



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



THE LIBRARY
OF
THE UNIVERSITY
OF CALIFORNIA

PRESENTED BY
PROF. CHARLES A. KOFOID AND
MRS. PRUDENCE W. KOFOID





(2)

My dear

Philadelphia

1845

Feb. 15th 1845



English

in the hands of
the people

QUINCY'S LEXICON-MEDICUM.

A NEW

MEDICAL DICTIONARY;

CONTAINING AN

EXPLANATION OF THE TERMS

IN

ANATOMY,
PHYSIOLOGY,
PRACTICE OF PHYSIC,
MATERIA MEDICA,

|| CHYMISTRY,
PHARMACY,
SURGERY,
MIDWIFERY,

AND THE VARIOUS BRANCHES OF NATURAL PHILOSOPHY CONNECTED
WITH MEDICINE.

**SELECTED, ARRANGED, AND COMPILED, FROM
THE BEST AUTHORS.**

"Nec araneorum sane texus ideo melior, quia ex se fila
gignunt, nec noster vilior quia ex alienis libamus ut apes."

JUST. LIPS. *Merit. Polit. Lib.* i, cap. i.

BY ROBERT HOOPER, M. D.

OF THE UNIVERSITY OF OXFORD, AND THE ROYAL COLLEGE OF PHYSICIANS OF LONDON;
PHYSICIAN TO THE ST. MARY-DE-BONE INFIRMARY, &c. &c.

PHILADELPHIA:

PUBLISHED BY M. CAREY & SON, BENJAMIN WARNER, AND EDWARD PARKER.

1817.

Griggs & Co. Printers.

K-R131

H6

B501

TO

WILLIAM SAUNDERS, M.D. F.R.S.

**FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS ;
OF THE ANTIQUARIAN AND OTHER SOCIETIES ;**

THIS WORK IS DEDICATED,

AS A MARK OF RESPECT AND ESTEEM,

BY HIS SINCERE FRIEND,

THE AUTHOR.

M372993

P R E F A C E.

WHEN Dr. Quincy published the first edition of his *Lexicon Medicum*, mathematical principles were generally adopted to explain the actions of the animal frame: hence we find in his work a continual recurrence to them. Since his time the functions of the animal economy and the knowledge of anatomy have received successive improvements, and the fashionable follies of mathematical explications have been reduced to their proper standard. To preserve the name which Dr. Quincy so deservedly obtained, and to render his work as useful as possible, such alterations and amendments were made in every following edition, as were suited to the doctrine of the times. It nevertheless has so happened, that his work, even in the thirteenth edition, contains very many of the absurdities of his day: The anatomical explanations are given in the language of the old schools, too often tedious, and abounding with every hypothesis; the physiology of the human body has been almost wholly overlooked; and all useful nosological descriptions omitted. Similar deficiencies and useless exuberances occur in every other department of the work.

PREFACE.

When, therefore, the present editor was solicited to undertake its revision, he thought he could not do a more acceptable office to the public, than almost wholly new model it. With this view he has been careful to collect such information as may render the work generally useful. Particular attention has been paid to the derivation of the terms, the anatomical description of the various parts, and the explanation of their functions; the diseases are considered according to the most approved nosological arrangement, and their symptoms and distinctions clearly enumerated: the *materia medica* and the preparations, especially those which enter the last edition of the *London Pharmacopœia*, have been amply considered; the improvements of modern Chymistry every where introduced, and the terms in Surgery, Midwifery, Medical Botany, and other Branches of Natural Philosophy, as far as connected with Medical Science, have been fully treated. In doing this, the editor has availed himself of the labours of the most eminent writers on the different branches of medicine, and has made such extracts, abridgments, translations, and selections, as the extent of the work would admit. It was his original intention to have given to each writer the merit of the particular description selected from his work; but having occasion to consult, frequently to abridge, and sometimes to alter various passages in works connected with his subject: and finding it difficult, and in many instances impossible to discover the original writer of several articles; and at the same time attended with no particular advantage, he prefers making a general acknowledgment of his obligations than to particularize the respective labours of each individual.

PREFACE.

The following have principally contributed to elucidate the several subjects. *Accum, Aiken, Albinus, Bell, Bergius, Blanchard, Burns, Burserius, Callisen, Castelli, Chaptal, Cooper, Cruickshank, Cullen, Denman, Duncan, Edinburgh Dispensatory, Edinburgh Encyclopædia, Editors of Motherby's Dictionary, Fourcroy, Green, Haller, Hunter, Innes, Latta, Lavoisier, Lewis, Linnæus, Meyer, Murray, Nicholson, Pott, Richerand, Richter, Saunders, Sauvage, Scarpa, Smith, Sæmmering, Swediaur, Symonds, Thomas, Thomson, Turton, Vaughan, Vossius, Willan, Willich's Encyclopædia, Wilson, Woodville.*

Deliusland Oct. 25th 1811
W. H. Bates

W. H. Bates

W. H. Bates

A NEW MEDICAL DICTIONARY.

A

A. **AA.** **ANA.** (From *ana*, which signifies of each.) A term in pharmacy.

It is never used but after the mention of two or more ingredients, when it implies, that the quantity mentioned of each ingredient should be taken; e. g. *℞. Potassæ nitratæ: Sacchari albi ℥℥. i. c.* Take the nitrate of potash and white sugar, of each one drachm.

ABAM. A term used by some ancient chymists for lead.

ABACTUS. *Abigeatus.* Among the ancient physicians, this term was used for a miscarriage, procured by art, or force of medicines, in contradistinction to *abortus*, which meant a natural miscarriage. The moderns know no such distinctions.

ABACUS. (From a Hebrew word, signifying dust.) A table for preparations, so called from the usage of mathematicians of drawing their figures upon tables sprinkled with dust.

ABAISIR. *Abasis.* *Spodium Arabum.* Ivory black; and also calcareous powder.

ABALIENATIO. A decay of the body, or mind.

ABALIENATUS. Corrupted. A part so destroyed as to require immediate extirpation; also the fault or total destruction of the senses, whether external or internal, by disease.

ABANET. (Hebrew, the girdle worn by the Jewish priests.) A girdle-like bandage.

ABANGA. *Ady.* The palm of the Island of St. Thomas, from which Thernal's restorative is prepared.

ABAPTISTA. (From *a*, priv. et *baptis*, to plunge.) *Abaptiston.* The shoulders of the old trepan. This term is employed by Galen, Fabricius ab Aquapendente, Scultetus, and others, to denote the conical saw with a circular edge, (otherwise called *modiolus*, or *terebra*), which was formerly used by surgeons to perforate the cranium.

ABAPTISTON. See *Abaptista*.

ABARNARAS. *Ovum ruffum.* A chemical term formerly used in the transmutation of metals, signifying *luna plena*, *magnes*, or *magnesia*.

ABARTAMEN. Plumbum, or lead.

ABARTICULATION. (From *ab*, and *articulus*, a joint.) That species of articulation which has evident motion. See *Diarthrosis*.

ABAS. (An Arabian word.) The scald-head; also epilepsy.

ABASIS. See *Abaisir*.

ABBREVIATION. The principal uses of medicinal abbreviations are in prescriptions; in which they are certain marks, or half words, used by physicians for despatch and conveniency when they prescribe, thus:—*℞* readily supplies the place of *recipe*—*h. s.* that of *hora somni*—*n. m.* that of *nux moschata*—*elect.* that of *electarium*, &c.; and in general all the names of compound medicines, with the several ingredients, are frequently wrote only up to their first or second syllable, or sometimes to their third or fourth, to make them clear and expressive. Thus *Croc. Anglic.* stands for *Crocus Anglicanus*—*Conf. Aromat.* for *Confectio Aromatica*, &c. A point being always placed at the end of such syllable shews the word to be incomplete.

ABDOMEN. (From *abdo*, to hide, because it hides the viscera. It is also derived from *abdere*, to hide, and *omentum*, the caul; and by others it is said to be only a termination, as from *lego*, *legumen*, so from *abdo*, *abdomen*.) The belly.

The abdomen is the largest cavity in the body, bounded superiorly by the diaphragm by which it is separated from the chest; inferiorly by the bones of the pubis and ischium; on each side by various muscles, the short ribs and *ossa ilii*, anteriorly by the abdominal muscles, and posteriorly by the vertebræ of the loins, the *os sacrum* and *os coccygis*. Internally it is invested by a smooth membrane called *peritoneum*, and externally by muscles and common integuments.

In the cavity of the abdomen are contained,

1. *Anteriorly and laterally.*
1. The epiploon. 2. The stomach.
3. The large and small intestines. 4. The mesentery. 5. The lacteal vessels. 6. The

pancreas. 7. The spleen. 8. The liver and gall-bladder.

Posteriorly, without the peritoneum, are,

1. The kidneys. 2. The supra-renal glands. 3. The ureters. 4. The receptaculum chyli. 5. The descending aorta. 6. The ascending vena cava.

3. *Inferiorly in the pelvis, and without the peritoneum,*

In men, 1. The urinary bladder. 2. The spermatic vessels. 3. The intestinum rectum.

In women, beside the urinary bladder and intestinum rectum, there are,

1. The uterus. 2. The four ligaments of the uterus. 3. The two ovaria. 4. The two Fallopian tubes. 5. The vagina.

The fore part of this cavity, as has been mentioned, is covered with muscles and common integuments, in the middle of which is the navel. It is this part of the body which is properly called abdomen; it is distinguished, by anatomists, into regions.

The posterior part of the abdomen is called the loins, and the sides the *Epitolic regions*.

Abdominal Hernia. See *Hernia Abdominalis*.

Abdominal muscles. See *Muscles*.

Abdominal ring. See *Annulus Abdominis*.

Abdominal regions. See *Regions*.

Abducens LABIORUM. A name given by Spigelius to the levator anguli oris. See *Levator anguli oris*.

Abducent nerves. See *Nervi abducentes*.

† *Abducent Muscles.* See *Abductor*.

ABDUCTIO. (From *abduco*, to draw away.)

A species of fracture, when a bone is divided transversely near a joint, so that each part recedes from the other. In Cælius Aurelianus it signifies a strain; and is mentioned as one of the causes of ischiadic and peoadic pains.

ABDUCTOR. (From *abduco*, to draw away.) *Abducens.* A name given to those muscles whose office is to pull back or draw the member to which it is affixed from some other, as the abductor pollicis draws the thumb from the fingers. The antagonists are called *adductores*, or *adductors*.

Abductor auricularis. See *Posterior auris*.

Abductor auris. See *Posterior auris*.

Abductor brevis alter. See *Abductor pollicis manus*.

ABDUCTOR INDICIS MANUS. *Abductor* of Douglas. *Semi-interosseus indicis* of Winslow. *Adductor indicis* of Cowper.

An internal interosseous muscle of the fore-finger, situated on the hand. It arises from the superior part of the metacarpal bone, and the os trapezium, on its inside, by a fleshy beginning, runs towards the metacarpal bone of the fore-finger, adheres to it, and is connected by a broad tendon to the superior part of the first phalanx of the fore-finger. Sometimes it arises by a

double tendon. Its use is to draw the fore-finger from the rest, towards the thumb, and to bend it somewhat towards the palm.

ABDUCTOR INDICIS PEDIS. An internal interosseous muscle of the fore-toe, which arises tendinous and fleshy, by two origins, from the root of the inside of the metatarsal bone of the fore-toe, from the outside of the root of the metatarsal bone of the great-toe, and from the os cuneiforme internum, and is inserted tendinous into the inside of the root of the first joint of the fore-toe. Its use is to pull the fore-toe inwards, from the rest of the small toes.

Abductor longus pollicis manus. See *Extensor ossis metacarpi pollicis manus*.

ABDUCTOR MEDI DIGITI PEDIS. An interosseous muscle of the foot, which arises tendinous and fleshy, from the inside of the root of the metatarsal bone of the middle toe internally, and is inserted tendinous into the inside of the root of the first joint of the middle toe. Its use is to pull the middle toe inwards.

ABDUCTOR MINIMI DIGITI MANUS. *Carpo-phalangien du petit doigt* of Dumas. *Extensor tertii internodii minimi digiti* of Douglas. *Hypothenar minor* of Winslow.

A muscle of the little finger, situated on the hand. It arises fleshy from the pisiform bone, and from that part of the *ligamentum carpi annulare* next it, and is inserted, tendinous, into the inner side of the upper end of the first bone of the little finger. Its use is to draw the little finger from the rest.

ABDUCTOR MINIMI DIGITI PEDIS. *Calcaneo-phalangien du petit doigt* of Dumas. *Adductor* of Douglas. *Parathenar major* of Winslow, by whom this muscle is divided into two, *Parathenar major* and *metatarsæus*. *Adductor minimi digiti* of Cowper.

A muscle of the little toe, which arises tendinous and fleshy, from the semicircular edge of a cavity on the inferior part of the protuberance of the os calcis, and from the rest of the metatarsal bone of the little toe, and is inserted into the root of the first joint of the little toe externally. Its use is to bend the little toe, and its metatarsal bone, downwards, and to draw the little toe from the rest.

ABDUCTOR OCULI. *Adductor* of Douglas and Winslow. *Orbito-fusus-scleroticus, orbito-ectus-scleroticus* of Dumas. *Rectus Adducens oculi* of Albinus. *Indignatorius*, or the scornful muscle. *Adducens Iracundus.* See *Rectus externus oculi*.

ABDUCTOR POLLICIS MANUS. *Scaphorus-phalangien du pouce* of Dumas. *Adductor pollicis manus*, and *Adductor brevis alter* of Albinus. *Adductor thenar Riolaui* of Douglas, (the *adductor brevis alter* of Albinus is the inner portion of this muscle.) *Adductor pollicis* of Cowper.

A muscle of the thumb situated on the hand. It arises by a broad tendinous and fleshy beginning, from the *ligamentum carpi annulare*, and from the *os trapezium*, and is inserted tendinous into the outer side of the root of the first bone of the thumb. Its use is to draw the thumb from the fingers.

ABDUCTOR POLLICIS PEDIS. *Caleaneo-phalangien du pouce* of Dumas. *Abductor* of Douglas. *Thenar* of Winslow. *Abductor pollicis* of Cowper.

A muscle of the great toe, situated on the foot. It arises fleshy, from the inside of the root of the protuberance of the *os calcis*, where it forms the heel, and tendinous from the same bone, where it joins the *os naviculare*; and is inserted tendinous into the internal sesamoid bone and root of the first joint of the great toe. Its use is to pull the great toe from the rest.

ABDUCTOR TERTII DIGITI PEDIS. An interosseous muscle of the foot, that arises tendinous and fleshy from the inside and the inferior part of the root of the metatarsal bone of the third toe; and is inserted tendinous into the inside of the root of the first joint of the third toe. Its use is to pull the third toe inwards.

ABEUS. (From *a*, neg. and *beus* firm.) *Abeus*. Weak, infirm, unsteady. A term made use of by Hippocrates de Signis.

ABEUS. See *Abeus*.

ABELMOSCHUS. (Arabian.) *Granum moschi*. *Moschus Arabum*. *Egyptia moschata*. *Bamid moschata*. *Alcea*. *Alcea Indica*. *Alcea Egyptiaca villosa*. *Abrette*. *Abelmosch*. *Abelmusk*. The seeds of a plant called the musk mallow, which have the flavour of musk. The plant *Hibiscus abelmoschus* of Linnaeus, is indigenous in Egypt, and in many parts of both the Indies. The best comes from Martinico. By the Arabians the seeds are esteemed cordial, and are mixed with their coffee, to which they impart their fragrance. In this country they are used by the perfumers.

Abelmosch. See *Abelmoschus*.

Abelmusk. See *Abelmoschus*.

ABERRATIO. (From *ab* and *erro*, to wander from.) *Lusus naturæ*. Dislocation.

ABESSI. (Arabian.) Filth. The alvine excrements.

ABESUM. Quick lime.

ABEVACUATIO. (From *ab*, dim. and *evacu*, to pour out.) A partial or incomplete evacuation of the peccant humours, either naturally or by art.

ABIES. (From *abeo*, to proceed, because it rises to a great height; or from *arbor*, a wild pear, the fruit of which its cones something resemble.) *Elate Thetela*. The fir. An evergreen tree. Linnaeus includes the abies in the genus *Pinus*. Botanists have enumerated several species: the four which follow, are the principal that afford materials for medicinal use.

1. *Pinus Picea*, the silver fir-tree, which affords the common turpentine.

2. *Pinus abies alba*, the Norway spruce fir-tree, which yields the Burgundy pitch.

3. *Pinus larix*, the common white larch-tree, from which is obtained the Venice turpentine.

4. *Pinus sylvestris*, the Scotch fir, which yields the *pice liquida*.

ABIES CANADENSIS. See *Balsamum Canadense*.

ABIGRATUS. See *Abactus*.

ABIOTOS. (From *a*, neg. and *bios*, to live.) A name given to hemlock, from its dead qualities. See *Centum*.

ABLACTATIO. (From *ab*, from, and *lact*, milk.) *Ab lactation*. The weaning of a child from the breast.

ABLATIO. (From *affero*, to take away.) The taking away from the body whatever is useless or hurtful; it comprehends all kinds of evacuations. Sometimes it signifies the subtraction of a part of the diet, with a medical view; and sometimes it expresses the interval betwixt two fits of a fever, or the time of remission.

Chymical *ablation* is the removal of any thing that is either finished or else no longer necessary in a process.

ABLUENTIA (*Abluentia*, sc. *medicamenta*, from *abluo*, to wash away.) *Abtergentia*. *Abluents*. Medicines which were formerly supposed to purify or cleanse the blood.

ABLUTION. (From *abluo*, to wash off.) A washing or cleansing either of the body or the intestines.

In chemistry it signifies the purifying of a body, by repeated effusions of a proper liquor.

ABOIT. An obsolete term of Arabic extraction for white lead.

ABOLITIO. (From *aboleo*, to destroy.) The separation or destruction of diseased parts.

ABORTION. (*Abortio*, from *aborior*, to be steril.) *Abortus*. *Amblous*. *Diaphthora*. *Ectrosis*. *Exumbroma*. *Examblois*. *Apopallens*. *Apopallens*. *Apophthora*.

Miscarriage, or the expulsion of the fetus from the uterus, before the seventh month, after which it is called premature labour. It most commonly occurs between the eighth and eleventh weeks of pregnancy, but may happen at a later period. In early gestation, the ovum sometimes comes off entire; sometimes the fetus is first expelled, and the placenta afterwards. It is preceded by flooding, pains in the back, loins, and lower part of the abdomen, evacuation of the water, shiverings, palpitation of the heart, nausea, anxiety, syncope, subsiding of the breasts and belly, pain in the inside of the thighs, opening and moisture of the os tinctæ.

ABORTIVES. (*Abortiva*, sc. *medicamenta*; from *aborior*, to be steril.) *Amblotica*. *Ectobica*.

Medicines capable of occasioning an

abortion, or miscarriage, in pregnant women. It is now generally believed, that the medicines which produce a miscarriage, effect it by their violent action on the system, and not by any specific action on the womb.

ABRASA. (From *abrado*, to shave off.) Ulcers attended with abrasion of part of their substance.

ABRASION. (*Abrasio*, from *abrado*, to tear off.) This word is generally employed to signify the destruction of the natural mucus of any part, as the stomach, intestines, urinary bladder, &c. It is also applied to any part slightly torn away by attrition, as the skin, &c.

ABRATHAN. Corrupted from *abrotanum*, southernwood. See *Abrotanum*.

ABRETTE. See *Abelmoschus*.

ABRIC. An obsolete Arabic term for sulphur.

ABROMA. (From *a*, neg. et *βρομα*, food; i. e. not fit to be eaten.) A tree of New South Wales, which yields a gum

ABROTANUM. (*Αβροτανον*, from *a*, neg. and *βροτος*, mortal; because it never decays: or from *αλεος*, soft, and *τερος*, extension; from the delicacy of its texture.) Common southernwood. *Abrotanum mas*.

Artemisia fruticosa, of Linnæus:—*foliis setaceis ramosissimis*. Class, *Syngenesia*. Order, *Polygamia superflua*. A plant possessed of a strong and, to most people, an agreeable smell; a pungent, bitter, and somewhat nauseous taste. It is supposed to stimulate the whole system, but more particularly the uterus. It is very rarely used unless by way of fomentation, with which intention the leaves are directed.

ABROTANUM MAS. See *abrotanum*.

ABROTONTES. (From *abrotanum*.) A wine mentioned by Dioscorides, impregnated with *abrotanum*, or southernwood, in the proportion of about one hundred ounces of the dried leaves, to about seven gallons of must.

ABSCIDENTIA. (From *abscedo*, to separate.) Decayed parts of the body, which, in a morbid state, are separated from the sound.

ABSCCESS. (From *abscedo*, to depart; because parts, which were before contiguous, become separated, or depart from each other.) *Abcessio* *Abcessus*. *Imposthuma*.

A collection of pus in the cellular membrane, or in the viscera, or in bones, preceded by inflammation.

ABSCISION. (*Abcissio*; from *ab*, and *scindo*, to cut.) *Apocope*. The taking away some morbid, or other part, by an edged instrument. The abscission of the prepuce makes what we call circumcision. *Abscission* is sometimes used by medical writers to denote the sudden termination of a disease in death, before it arrives at its decline. Celsus frequently uses the term *abscissa vox* to express a loss of voice.

ABSINTHIUM. (*Α-λ-σιν*, from *a*, neg. and *λσος*, pleasant: so called from the disagreeableness of the taste.) A genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia superflua*. Wormwood.

ABSINTHIUM COMMUNE. See *Absinthium vulgare*

ABSINTHIUM MARITIMUM. Sea wormwood, falsely called in our markets, Roman wormwood. *Artemisia maritima*.

Absinthium Ponticum of Linnæus:—*foliis multipartitis, tomentosis racemis cernuosis flosculis fœmineis ternis*. This plant grows plentifully about the sea-shore, and in salt marshes. The specific differences between it and the common wormwood, *absinthium vulgare*, are very evident. Its taste and smell are considerably less unpleasant than those of the common wormwood, and even the essential oil, which contains the whole of its flavour concentrated, is somewhat less ungrateful, and the watery extract somewhat less bitter than those of the common wormwood. Hence it is preferred, in those cases where the *Artemisia absinthium* is supposed to be too unpleasant for the stomach. A conserve of the tops of this plant is directed by the London Pharmacopœia.

ABSINTHIUM PONTICUM. See *Absinthium Maritimum*.

ABSINTHIUM VULGARE. Common wormwood. Falsely called in our markets *Absinthium Romanum*, or Roman wormwood. *Absinthium Ponticum* of Dioscorides and Pliny. Murray.

Artemisia Absinthium of Linnæus:—*foliis compositis multifidis, floribus subglobosis pendulis: receptaculo villosis*. Class, *Syngenum*. Order, *Polygamia superflua*. This plant is a native of Britain, and grows about rubbish, rocks, and sides of roads. The leaves of wormwood have a strong disagreeable smell: their taste is nauseous, and so intensely bitter as to be proverbial. The flowers are more aromatic and less bitter than the leaves, and the roots discover an aromatic warmth, without bitterness. This species of wormwood may be considered the principal of the herbaceous bitters. Its *virtus*, in the words of Bergius, is antiputredinosa, antacidia, anthelminthica, resolvens, tonica, spasmodica. And although it is now chiefly employed with a view to the two last-mentioned qualities, yet we are told of its good effects in a great variety of diseases, as intermittent fevers, hypochondriasis, obstructions of the liver and spleen, gout, calculi, scurvy, dropsy, worms, &c. See Woodville's Medical Botany. Cullen thinks it is possessed of a narcotic power, and that there is in every bitter, when largely employed, a power of destroying the sensibility and irritability of the nervous power.

Externally, wormwood is used in discu-

tient and antiseptic fomentations. This plant may be taken in powder, but it is more commonly preferred in infusion. The Edinburgh Pharmacopœia directs a tincture of the flowers, which is, in the opinion of Dr. Cullen, a light and agreeable bitter, and, at the same time, a strong impregnation of the wormwood.

Absorbing vessels. See *Absorbents*.

ABSORBENTS. *Absorbentia.*

1. Small, delicate, transparent vessels, which take up any fluid from the surface of the body, or of any cavity in it, and carry it to the thoracic duct, to be mixed with the blood. They are denominated according to the liquids which they convey, lacteals and lymphatics. See *Lacteals* and *Lymphatics*.

2. Medicines are so termed, which have no acrimony in themselves, and destroy acidities in the stomach and bowels; such are calcined magnesia, prepared chalk, oyster-shells, crab's claws, &c.

ABSORPTION. (From *absorbes*, to suck up.) A function in an animated body, arranged by physiologists under the head of natural actions. It signifies the taking up of substances applied to the mouths of absorbing vessels: thus the nutritious part of the food is absorbed from the intestinal canal by the lacteals: thus mercury is taken into the system by the lymphatics of the skin, &c. The principle by which this function takes place, is a power inherent in the mouths of the absorbents, a *vis insita*, dependent on the degree of irritability of their internal membrane by which they contract and propel their contents forwards.

ABSTENTIO. Cælius Aurelianus uses this word, to express a suppression, or retention. Thus, *abstentio stercorem*, a retention of the excrements, which he mentions as a symptom very frequent in a *satyriasis*. In a sense somewhat different, he uses the word *abstenta*, applying it to the pleura, where he seems to mean, that the humour of the inflamed pleura is prevented, by the adjacent bones, from extending itself.

ABSTERGENTS. (*Abstergentia*, *soilicet medicamenta*; from *abstergo*, to cleanse away.) Lotions, or any application that cleanses or clears away foulness. The term is seldom employed by modern writers.

ABSTRACTION. (From *abstraho*, to draw away.) A term employed by chymists in the process of humid distillation, to signify that the fluid body is again drawn off from the solid, which it had dissolved.

ABSTRACTIVUS. (From *abstraho*, to draw away.) Native spirit, not produced by fermentation.

ABUS. An obsolete term for the Egyptian lotus.

ABVACUATIO. (From *abvacuo*, to empty.) Local or morbid discharge. A large evacuation of any fluid, as of blood from a plethoric person.

ACACA. (From *α, neg.* and *κακα, bad.*) Diseases which are rather troublesome than dangerous.

ACACIA. (*Acacia*, from *ακαζω*, to sharpen.) The name of a genus of plants in the Linnæan system. The Egyptian thorn.

ACACIA GERMANICA. *Acacia nostras.* German acacia, or the German black-thorn or sloe-tree. *Acacia nostras.* *Succus pruni sylvestris.* The inspissated juice of the German wild sloe, *prunus spinosa*, or *prunus sylvestris spinosa* of Linnæus; now fallen into disuse.

ACACIE GUMMI. *Gummi acanthium.* *Gummi thebaicum.* *Gummi scorpioidis.* *Gum-lumac.* *Gummi senega*, or *senica.* *Acacia gum*, or *gum-arabic.* The gum of the Egyptian thorn.

Acacia vera, of Willdenow:—*spinis stipularibus patentibus, foliis bipinnatis; partibus extimis glandula intertinctis, spicis globosis pedunculatis.* Cairo and Alexandria were the principal marts for gum-arabic, till the Dutch introduced the gum from Senegal into Europe, about the beginning of the seventeenth century, and which now supplies the greater part of the vast consumption of this article.

The tree which yields the Senegal gum, grows abundantly on the sands, along the whole of the Barbary coast, and particularly about the river Senegal. There are several species, some of which yield a red astringent juice, but others afford only a pure, nearly colourless, insipid gum, which is the great article of commerce. These trees are from eighteen to twenty feet high, with thorny branches. The gum makes its appearance about the middle of November, when the soil has been thoroughly saturated with periodical rains. The gummy juice is seen to ooze through the trunk and branches, and, in about a fortnight, it hardens into roundish drops, of a yellowish white, which are beautifully brilliant where they are broken off, and entirely so when held in the mouth for a short time, to dissolve the outer surface. No clefts are made, nor any artificial means used by the Moors, to solicit the flow of the gum. The lumps of gum-senegal are usually about the size of partridge eggs, and the harvest continues about six weeks. This gum is a very wholesome and nutritious food; thousands of the Moors supporting themselves entirely upon it during the time of harvest. About six ounces is sufficient to support a man for a day; and it is besides, mixed with milk, animal broths, and other victuals.

The gum-arabic, or that which comes directly from Egypt and the Levant, only differs from the gum-senegal in being of a lighter colour, and in smaller lumps; and it is also somewhat more brittle. In all other respects, the two resemble each other perfectly.

Gum-arabic is neither soluble in spirit nor

ACETAS PLUMBI. Acetate of lead. A metallic salt composed of lead and acetic acid. See *Liquor plumbi acetatis*.

ACETAS ZINCI. A metallic salt composed of zinc and acetic acid. It is used by some as an astringent against inflammation of the eyes, urethra, and vagina, diluted in the same proportion as the sulphate of zinc.

Acetated vegetable Alkali. See *Acetas potassæ*.

Acetated volatile Alkali. See *Liquor acetatis ammoniæ*.

Acetic Acid. See *Acetum*.

ACETIFICATION. A term used by some chymists to denote the action or operation by which vinegar is made.

Acetat of Potash. See *Acetas potassæ*.

Acetat of Ammonia. See *Liquor ammoniæ acetatis*.

Acetat of Zinc. See *Acetas Zinci*.

ACETOSA. (From *acesco*, to be sour.) Sorrel. A genus of plants in some systems of botany.

ACETOSA VULGARIS. *Acetosa pratensis.* *Acetosa arvensis* Sorrel; sour-dock.

Rumex acetosus of Linnaeus:—*foliis oblongis sagittatis, floribus diœceis.* Class, *Hexandria*. Order, *Tryginia*. The leaves of this plant are sour, but not the root, which is bitter. It grows in the meadows and common fields.

ACETOSA ROMANA. *Acetosa rotundifolia hortensis.* Roman or garden sorrel.

Rumex scutatus or *helveticus*:—*foliis cordato-hastatis, ramis divergentibus, floribus hermaphroditis*, of Linnaeus. It is common in our gardens and in many places is known by the culinary name of Green-sauce.

ACETOSELLA. (From *acetosa*, sorrel; from the acidity of its leaves.) *Lujula*. *Alleluja*. Wood-sorrel.

Oxalis acetocella, of Linnaeus:—*foliis ternatis, scape uniflora, flore albo, capsulis pentagomis elasticis, radice squamoso-articulata.* Class, *Decandria*. Order, *Pentagynia*.—This plant grows wild in the woods, and flowers in April and May. The leaves are shaped like a heart, standing three together on one stalk. The acetocella is totally inodorous, but has a grateful acid taste, on which account it is used in sallads. Its taste is more agreeable than the common sorrel, and approaches nearly to that of the juice of lemons, or the acid of tartar, with which it corresponds in a great measure in its medical effects, being esteemed refrigerant, antiscorbutic, and diuretic. It is recommended by Bergius; in inflammatory, bilious, and putrid fevers. The principal use however of the acetosella is to allay inordinate heat and to quench thirst; for this purpose, a pleasant whey may be formed by boiling the plant in milk, which under certain circumstances may be preferable to the conserve directed by the London College, though an extremely grateful and useful medicine. Many have employed

the root of *Lujula*, probably on account of its beautiful red colour rather than for its superior efficacy. An essential salt is prepared from this plant, known by the name of essential salt of lemons, and commonly used for taking ink-stains out of linen. What is sold under the name of essential salt of lemons in this country, is said by some to consist of cream of tartar, with the addition of a small quantity of sulphuric acid. The leaves of sorrel when employed externally in the form of poultices, are powerful suppurants, particularly in indolent scrofulous humours.

Acetous Acid. Distilled vinegar. See *Acetum*.

Acetous fermentation. See *Fermentation*.

ACETUM. (From *acer*, sour.) Vinegar. A sour liquor obtained from many vegetable substances dissolved in boiling water, and from fermented and spirituous liquors, by exposing them to heat and contact with air; under which circumstances they undergo the acid fermentation, (see *Fermentation*.) and afford the liquor called vinegar.

Wine vinegar.—Let any quantity of vinous liquor be mixed with the acid and austere stalks of the vegetable from which wine was prepared. The whole must be frequently stirred and either exposed to the sun, or deposited in a warm place: after standing a few days it will ferment, become sour and in a fortnight it will be converted into vinegar.

Cyder vinegar, may be made by fermenting new cyder with the must of apples, in a warm room, or in the open air, where it should be exposed to the sun, and in the course of a week or nine days it will be fit for use.

Another method of preparing vinegar is that published by M. Heber: it consists in exposing a mixture of 72 parts of water, and 4 of rectified malt spirit in a temperature of from 70 to 80° of Fahrenheit, for about two months, at the expiration of which the acetous process will be effected.

Tarragon vinegar is manufactured by infusing one pound of the leaves of that vegetable (which has been gathered a short time before it flowers) in one gallon of the best vinegar, for the space of 14 days: when it should be strained through a flannel bag; and a drachm of isinglass dissolved in cyder must then be added, the whole be carefully mixed and decanted into bottles for a month. Thus the liquor will acquire a most exquisite flavour; it will become remarkably fine and almost colourless.

The utility of vinegar as a condiment for preserving and seasoning both animal and vegetable substances in various articles of food, is very generally known. It affords an agreeable beverage, when combined with water in the proportion of a table-spoonful of the former to half pint of the

latter. It is often employed as a medicine in inflammatory and putrid diseases, when more active remedies cannot be procured. Relief has likewise been obtained in hypochondrical and hysteric affections, in vomiting, fainting, and hiccup, by the application of vinegar to the mouth. If this fluid be poured into vessels and placed over the gentle heat of a lamp in the apartments of the sick, it greatly contributes to disperse foul or mephitic vapours, and consequently to purify the air.

Also as an external application, vinegar proves highly efficacious when joined with farinaceous substances, and applied as a cataplasm to sprained joints; it also forms an eligible lotion for inflammations of the surface, when mixed with alcohol and water in about equal proportions. Applied to burns and scalds, vinegar is said to be highly serviceable whether there is a loss of substance or not, and to quicken the exfoliation of carious bone. (Gloucester Infirmary.) Mixed with an infusion of sage, or with water, it forms a popular and excellent gargle for an inflamed throat, also for an injection to moderate the fluor albus. Applied cold to the nose in cases of hæmorrhage, also to the loins and abdomen in menorrhagia, particularly the profluvia after parturition, it is said to be very serviceable. An imprudent use of vinegar internally is not without considerable inconveniences. Large and frequent doses injure the stomach, coagulate the chyle, and produce not only leanness, but an atrophy. When taken to excess by females, to reduce a corpulent habit, tubercles in the lungs and a consumption have been the consequence.

Common vinegar consists of acetic acid combined with a large portion of water, and with this are in solution portions of gluten, mucilage, sugar, and attractive matter from which it derives its colour, and frequently some of the vegetable acids, particularly the malic and the tartaric.

Distilled with a gentle fire, in glass vessels, so long as the drops fall free from empyreuma, it affords the

ACIDUM ACETICUM.

Take of vinegar, a gallon.

Distil the acetic acid in a sand bath, from a glass retort into a receiver also of glass, and kept cold; throw away the first pint, and keep for use the six succeeding pints, which are distilled over.

In this distillation, the liquor should be kept moderately boiling, and the heat should not be urged too far, otherwise the distilled acid will have an empyreumatic smell and taste, which it ought not to possess. If the acid be prepared correctly, it will be colourless, and of a grateful, pungent, peculiar acid taste. One fluid-ounce ought to dissolve at least ten grains of carbonate of

lime (white marble.) This liquor is the *acetum distillatum*; the *acidum acetosum* of the London Pharmacopœia of 1767, and the *acidum aceticum* of the last (1809.)

When the acid of vinegar is greatly concentrated, that is, deprived of its water, it becomes the radical vinegar, or

CONCENTRATED ACID OF VINEGAR.

Distilled vinegar may be concentrated by freezing: the congelation takes place at a temperature below 28 degrees, more or less, according to its strength; and the congealed part is merely ice, leaving, of course, a stronger acid. If it be exposed to a very intense cold, equal to 38 degrees, it shoots into crystals; when the fluid part is withdrawn, the crystals liquefy, when the temperature rises, and the liquid is limpid as water, extremely strong, and has a highly pungent acetous odour. This is the pure acid of the vinegar, any foreign matter remaining in the uncongealed liquid.

Other methods are likewise employed to obtain the pure and concentrated acid. The process of Westendorf, which has been often followed, is to saturate soda with distilled vinegar, obtain the acetate by crystallization; and pour upon it, in a retort, half its weight of sulphuric acid. By applying heat, the acetic acid is distilled over; and, should there be any reason to suspect the presence of any sulphuric acid, it may be distilled a second time, from a little acetate of soda. According to Lowitz, the best way of obtaining this pure, is to mix three parts of the acetate of soda with eight of supersulphate of potass; both salts being perfectly dry, and in fine powder, and to distil from this mixture in a retort, with gentle heat.

It may also be obtained by distilling the verdigris of commerce, with a gentle heat. The concentrated acid procured by these processes, was supposed to differ materially from the acetous acid obtained by distilling vinegar; the two acids were regarded as differing in their degree of oxygenization, and were afterwards distinguished by the names of acetous and acetic acids. The acid distilled from verdigris was supposed to derive a quantity of oxygen from the oxide of copper, from which it was expelled. The experiments of Adet have, however, proved the two acids to be identical; the acetous acid, therefore, only differs from the acetic acid in containing more water, rendering it a weaker acid, and of a less active nature.

There exists, therefore, only one acid of vinegar, which is the acetic; and its compounds must be termed *acetates*; and the salts called *acetites* have no existence.

Acetic acid, when concentrated, has a fragrant and, at the same time, very penetrating smell, irritating the nostrils strongly. It is also so caustic, as to inflame the skin. Its acid taste is strong, even when

diluted with water; it is colourless, and has a specific gravity of 1.0626. The acid is capable of congelation; when it forms foliated arborescent crystals it is very volatile; its odour is diffused through the atmosphere, and, when exposed to it, gradually becomes weaker. By a moderate heat, it is converted into vapour; this vapour readily catches fire on the approach of a lighted taper. It combines with water in every proportion; and it combines readily with earthy, metallic, and alkaline bases, forming salts, which are acetates.

The compounds of the acid of vinegar, directed to be used by the new London Pharmacopœia, are *acetum colchici*, *acetum scilla*, *ceratum plumbi superacetatis*, *liquor ammonie acetatis*, *liquor plumbi acetatis*, *liquor plumbi acetatis dilutus*, *oxymel*, *oxymel scilla*, *potasse acetat*, and the *liquor ammonie acetatis*.

ACETUM AROMATICUM. Aromatic vinegar. A preparation of the Edinburgh pharmacopœia, thought to be an improvement of what has been named *thiours vinegar*.

"Take of the dried tops of rosemary;

The dried leaves of sage, of each four ounces.

Dried lavender flowers, two ounces.

Cloves, two drac ms;

Distilled vinegar, eight pounds.

Macerate for seven days, and strain the expressed juice through paper." Its virtues are antiseptic, and it is a useful composition to smell at in crowded courts of justice, hospitals, &c. where the air is offensive.

ACETUM DISTILATUM. See *acetum*.

ACETUM SCILLA. Lond. Pharm. Vinegar of squills. *S.* Squills recently dried, one pound; vinegar, six pints; proof spirit, half a pint. Macerate the squills with the vinegar in a glass vessel, with a gentle heat for twenty-four hours; then express the liquor and set it aside until the fœces subside. To the decanted liquor add the spirit. This preparation of squills is employed as an attenuant, expectorant, and diuretic, *gutt. xv. to lx.*

ACEDIA. Without hands.

ACHICOLUM. By this word Cælius Aurelianus, *Acut. lib. iii. cap. 17.* expresses the fornix, tholus, or sudatorium of the ancient baths, which was a hot room where they used to sweat.

ACHILLEA. (*ΑΧΙΛΛΕΙΑ*, from Achilles, who is said to have made his tents with it, or to have cured Telaphus with it.) The name of a genus of plants in the Linnæan system. Class *Syngenesia*. Order, *Polygamia superflua*. Milfoil.

ACHILLE AGERATUM. *Balsamita fœmina*. *Eupatorium Menues*. Maudlin or Maudlin tansey. This plant, the *ageratum* of the shops, is described by Linnæus as *Achillea foliis lanceolatis, obtusis, acutius-*

ratiss. It is esteemed in some countries as anthelmintic and alterative, and is given in hepatic obstructions. It possesses the virtues of tansey.

Achillea foliis pinnatis. See *Genipi verum*.

ACHILLEA MILLEFOLIUM. The systematic name of the milfoil. See *Millefolium*.

ACHILLEA PTARMICA. The systematic name of the sneezewort. See *Ptarmica*.

ACHILLIS TENDON. (So called, because, as fable reports, Thetis, the mother of Achilles, held him by that part when she dipped him in the river Styx, to make him invulnerable. Homer describes this tendon, and some writers suppose it was thus named by the ancients, from their custom of calling every thing *Achilleum*, that had any extraordinary strength or virtue. Others say it was named from its action in conducing to swiftness of pace, the term importing so much.) The strong and powerful tendon which is formed by the junction of the gastrocnemius and soleus muscles, and which extends along the posterior part of the tibia from the calf to the heel. When this tendon is unfortunately cut or ruptured, as it may be in consequence of a violent exertion, or spasm of the muscles, of which it is a continuation, takes place, the use of the leg is immediately lost, and unless the part be afterwards successfully united, the patient must remain a cripple for life. When the tendon has been cut, the division of the skin allows the accident to be seen. When the tendon has been ruptured, the patient hears a sound like that of the smack of a whip, at the moment of the occurrence. In whatever way the tendon has been divided, there is a sudden incapacity, or at least an extreme difficulty, either of standing or walking. Hence the patient falls down, and cannot get up again. Besides these symptoms there is a very palpable depression between the ends of the tendon: which depression is increased when the foot is bent and diminished, or even quite remove when the foot is extended. The patient can spontaneously bend his foot, none of the flexor muscles being interested. The power of extending the foot is still possible, as the peronei muscles, the tibialis posticus, and long flexors, remain perfect and may perform this motion. The indications are to bring the ends of the divided parts together, and to keep them so, until they have become firmly united. The first object is easily fulfilled by putting the foot into a state of complete extension; the second, namely, that of keeping the ends of the tendon in contact, is more difficult. It seems unnecessary to enumerate the various plans devised to accomplish these ends. The following is Desault's method: After the ends of the tendon had been brought into contact by moderate flexion of the knee,

and complete extension of the foot, he used to fill up the hollows on each side of the tendon with soft lint and compresses. The roller applied to the limb, made as much pressure on these compresses as on the tendon, and hence this part could not be depressed too much against the subjacent parts. Desault next took a compress about two inches broad, and long enough to reach from the toes to the middle of the thigh, and placed it under the foot, over the back of the leg and lower part of the thigh. He then began to apply a few circles of a roller round the end of the foot, so as to fix the lower extremity of the longitudinal compress: after covering the whole foot with the roller, he used to make the bandage describe the figure of 8, passing it under the foot and across the place where the tendon was ruptured, and the method was finished by encircling the limb upward with the roller as far as the upper end of the longitudinal compress.

ACHLYS. (Αχλυσ) Darkness, cloudiness. It is generally applied to a close, foggy air, or a mist.

Hippocrates, De Morbis Mulierum, lib. ii. signifies by this word condensed air in the womb.

Galen interprets it of those, who, during sickness, lose that usual lustre and loveliness observed about the pupil of the eye, during health.

Others express it by an ulcer on the pupil of the eye, or the scar left there by an ulcer.

It means also an opacity of the cornea; the same as the caligo cornea of Dr. Cullen.

ACHMADIUM. Antimony.

ACHMELLA. *Acmella. Achamella.* The herb and seeds of this plant, *Spilanthes achmella* of Linnaeus, are employed in cases of calculus of the kidneys and urinary bladder. The plant is very glutinous and bitter, and is given in infusion.

ACHNE. Chaff, scum or froth of the sea. A white mucus in the fauces, thrown up from the lungs, like froth; also a whitish mucilage in the eyes of those who have fevers, according to Hippocrates. It signifies also lint.

ACHOR. (αχας, qn. αχρας, from αχρν bran; according to Blanchard it is derived from α, priv. and χρας space, as occupying but a small compass.) *Lactumen: abas: acores: cerion: favus. Crusta lactea* of authors. The scald-head; so called from the branny scales thrown off it. A disease which attacks the hairy scalp of the head, for the most part of young children, forming soft and scaly eruptions. Dr. Willan, in his description of different kinds of pustules, defines the achor, a pustule of intermediate size between the phlyzadium and pyodacium which contains a straw-coloured fluid, having the appearance of and nearly the con-

sistence of strained honey. It appears most frequently about the head, and is succeeded by a dull white or yellowish scab. Pustules of this kind, when so large as nearly to equal the size of phlyzacia, are termed *ceria* or *favi*, being succeeded by a yellow, semi-transparent, and sometimes, cellular scab, like a honey-comb. The achor differs from the *favus* and *tinea* only in the degree, of virulence. It is called *favus* when the perforations are large; and *tinea* when they are like those which are made by moths in cloth; but generally by *tinea* is understood a dry scab on the hairy scalp of children, with thick scales and an offensive smell. When this disorder affects the face, it is called *crusta lactea* or milk scab. Mr. Bell, in his treatise on Ulcers, reduces the *tinea capitis* and *crusta lactea* to the same species of herpes, viz. the herpes pustulosus, differing only in situation.

ACHONISTOS. Inseparable. It is understood of accidents, symptoms, or signs, which are inseparable from the particular things. Thus, a pungent pain in the side is an inseparable symptom of the pleurisy.

ACHYRON. Useless. It is applied by Hippocrates to the limbs which, through weakness, are become useless.

ACHROIA. A paleness.

ACHYLOS. Deficient in bile.

ACHYRON, (αχρον.) This properly signifies bran or chaff, or straw.

Hippocrates, de Morbis Mulierum, most probably means by this word, bran. Achyron also signifies a straw, hair, or any thing that sticks upon a wall.

ACIA, (From αα, a point.) A needle with thread in it for chirurgic operations.

ACIRRS. It signifies weak, infirm, or faint, and in this sense it is used by Hippocrates, De Morb. lib. iv.

ACID. That which impresses upon the organs of taste a sharp or sour sensation. Acids are defined by modern chymists to be salts of a sour taste, changing the blue colour of various vegetable pigments to a red. The word *sour*, which is usually employed to denote the simple impression, or lively and sharp sensation produced on the tongue by certain bodies, may be regarded as synonymous to the word *acid*. The only difference which can be established between them is, that the one denotes a weak sensation, whereas the other comprehends all the degrees of force from the least perceptible to the greatest degree of causticity: thus we say that verjuice, gooseberries, or lemons, are *sour*; but we use the word *acid* to express the impression which the nitric, sulphuric, or muriatic acids make upon the tongue. The vegetable pigments usually employed to ascertain the presence of acids are tincture of turnsole or litmus, and syrup of violets. Acids readily combine with alka-

lis, earths, and metals, and form neutral salts. The characteristics, therefore, of an acid, are.

1. A peculiar taste termed acid.
2. Its changing blue vegetable juices red.
3. Combining with alkalis, earths, and metals.

Acids, according to the kingdom of nature in which they are found, are divided into mineral, vegetable, and animal.

The mineral acids as yet known, are the sulphuric or vitriolic, the nitric, muriatic, carbonic, boracic, fluoric, succinic, arsenic, molybdic, tungstic, and chromic.

The vegetable acids are, the acetic, oxalic, tartareous, pyrotartareous, gallic, citric, mallic, benzoic, pyroligneous, the succinic, pyromucous, camphric, and cortic.

Of the animal acids there are eight, viz. the phosphoric, lactic, saccholactic, formic, sebatic, prussic, bombic, and lithic, or uric.

Experiment proves that every acid consists of a peculiar body combined with the basis of oxygen gas: hence the origin of the word oxygen, which signifies the generation of acid, it being regarded as the *acidifying basis* or principle of acidity. The bodies which form the other constituents of acids, are regarded as the *acidifiable basis*; thus the principles of phosphoric acid are phosphorus and oxygen; those of carbonic acid, radical carbon and oxygen.

If an acid basis be perfectly saturated with oxygen, the acid, thus produced, is said to be perfect; but if the basis predominate, the acid is considered as imperfect. Modern chymists distinguish the former in Latin by the syllables *icum*, in English *ic*, and the latter in Latin by *osum*, and in English by *ous*: thus the perfect acid of nitre is called *acidum nitricum*, or nitric acid; the imperfect acid of nitre, *acidum nitrosum*, or nitrous acid. There are some cases where an acid is capable of combining with an excess of oxygen, in which case it is said to be *oxygenated*; and sometimes *super-oxygenated*. If the acidifiable basis be combined with oxygen, yet without showing any of the properties of an acid, the produce is then called an oxyd or oxyde: thus iron exposed to the air or water attracts the oxygen, and an oxyd of iron, the rust, is formed. The various acids employed medicinally are, the acetic, benzoic, tartaric, carbonic, citric, muriatic, oxygenated muriatic, nitric, nitrous, sulphuric and phosphoric.

Acid aërial. See *Carbonic acid*.

Acid acetic. See *Acetum*.

For the other Acids look to the word *Acidum*.

Acidifiable base. See *Acid*.

Acidifying base. See *Acid*.

ACIDIFICATION. The formation of an acid; also the impregnating of any thing with acid properties.

ACIDITY. *Aciditas*. Sourness.

ACIDS, ANIMAL. Those which are obtained from animals. See *Acid*.

ACIDS DULCIFIED. These are now called *Ethers*. See *Ether*.

ACIDS IMPERFECT. Those acids are so called in the chymical nomenclature, which are not fully saturated with oxygen. Their names are ended in Latin by *osum*, and in English by *ous*: e. g. *acidum nitrosum*, or nitrous acid.

ACIDS, MINERAL. Those acids which are found to exist in minerals, as the sulphuric, the nitric, &c. See *Acid*.

ACIDS, PERFECT. An acid is termed perfect in the chymical nomenclature, when it is completely saturated with oxygen. Their names are ended in Latin by *icum*, and in English by *ic*: e. g. *acidum nitricum*, or nitric acid.

ACIDS, VEGETABLE. Those which are found in the vegetable kingdom, as the citric, mallic, acetic, &c. See *Acid*.

ACIDULOUS WATER. Mineral waters, which contain so great a quantity of carbonic acid gas, as to render them acidulous, or gently tart to the taste. See *Mineral waters*.

ACIDUM ACETICUM. See *Acetum*.

ACIDUM ACETOSUM. See *Acetum*.

ACIDUM ÆTHEREUM. The sulphuric acid.

ACIDUM ALUMINOSUM. The sulphuric acid.

ACIDUM ARSENICUM. See *Arsenic*.

ACIDUM BENZOICUM. See *Benzoe*.

ACIDUM BORACICUM. See *Boracic acid*.

ACIDUM CARBONICUM. See *Carbonic acid*.

ACIDUM CATHOLICON. The acid of sulphur.

ACIDUM CITRICUM. See *Citric acid*.

ACIDUM MURIATICUM. See *Muriatic acid*.

ACIDUM NITRICUM. See *Nitric acid*.

ACIDUM NITROSUM. *Spiritus nitri fumans*, of the shops. The nitrous acid possesses the same properties as the nitric, but in a much inferior degree.

ACIDUM NITROSUM DILUTUM. This is the common *aqua fortis*. Diluted nitrous acid possesses the same properties as the nitric acid, but in an inferior degree.

ACIDUM PHOSPHORICUM. See *Phosphoric acid*.

ACIDUM PRIMOGENIUM. The sulphuric acid.

ACIDUM SUCCINICUM. See *Succinic acid*.

ACIDUM SULPHUREUM. The acid of sulphur.

ACIDUM SULPHURICUM. See *Sulphuric acid*.

ACIDUM SULPHURICUM DILUTUM. *Acidum vitriolicum dilutum*. *Spiritus vitrioli tenuis*.

ACIDUM TARTARICUM. See *Tartaric acid*.

ACIDUM VITRIOLICUM. See *Sulphuric acid*.

ACIDUM VITRIOLICUM DILUTUM. See *Acidum sulphuricum dilutum*.

Take of sulphuric acid, a fluidounce and half.

Distilled water, fourteen fluidounces and half. Add the water to the acid gradually, and mix.

ACIES. Steel.

ACINESIA. A loss of motion and strength.

ACINI BILIOSI. (*Acinus*, a grape-stone; so called from their supposed resemblance.) The small glandiform bodies of the liver, which separate the bile from the blood were formerly so called: they are now, however, more properly termed *penicilli*. See *Liver*.

ACINIIFORM TUNIC. *Tunica acinosa*. The coat of the eye called the *wren*, because the ancients, who dissected brutes, observed that, in them, it was usually of the colour of an unripe grape.

ACINUS. (A grape.) The glands which grow together in clusters are called by some *acid glandulosi*.

ACHASTICOS. A species of synochus, wherein the febrile heat continues of the same tenour to the end. *Actuarius*.

ACME. (From *αμην* a point.) The height or crisis of a disease. A term applied by physicians to that period or state of a disease in which it is at height. The ancients distinguished diseases into four stages: 1. the *arche*, the beginning or first attack. 2. *Anabasis*, the growth. 3. The *acme*, the height. 4. *Paracme*, or the decline of the disease.

ACHILLA. See *Achmella*.

ACNE. *Acna*, *acn*. A small pimple, or hard tubercle on the face. *Fæsius* says, that it is a small pustule or pimple, which arises usually about the time that the body is in full vigour.

ACNESTIA. (From *α*, priv. and *ανω*, to scratch.) That part of the spine of the back, which reaches from the metaphrenon, which is the part betwixt the shoulder-blades, to the loins. This part seems to have been originally called so in quadrupeds only, because they cannot reach it to scratch.

ACOX. (*Axon*) The sense of bearing. **ACOLIOS.** (From *α* priv. and *κολος*, the belly.) Without belly. It is applied to those who are so wasted, as to appear as if they had no belly. *Galen*.

ACORTUS. (*Acorus*) An epithet for honey, mentioned by Pliny: because it has no sediment, which is called *acorn*.

ACONION. (*Aconium*) A particular form of medicine among the ancient physicians, made of powders levigated, and probably like *collyria* for the disorders of the eyes.

ACONIVM. A little mortar.

ACONITUM. (Of this plant various derivations are given by etymologists; as, *acon* a whetstone or rock, because it is usually found in barren and rocky places: *a*, neg. and *con*, dust; because it grows without earth or on barren situations: *acon*, to sharpen; because it was used in medicines intended to quicken the sight: *acon*, *acn*, a dart; because they poison darts therewith: or, *aconizomas*, to accelerate; for it hastens death.) Aconite. Wolf's-bane. Monk's-hood.

1. A genus of plants in the Linnæan system. Class, *Polyandria*, *Trigyn*a.

2. The pharmacopœial name of the common, or blue, wolf's-bane. Monk's-hood. Aconite. *Camarum*. *Caniada*. *Cynocetanum*.

Aconitum napellus of Linnæus:—*foliorum lactuiss linearibus superne latioribus, lineâ exaratis*.

The aconite is cultivated in our gardens as an ornament, but is spontaneously produced in Germany, and some other northern parts of Europe. Every part of the plant is strongly poisonous, but the root is unquestionably the most powerful; and when first chewed, imparts a slight sensation of acrimony, but afterwards, an insensibility or stupor at the apex of the tongue and a pungent heat of the lips, gums, palate, and fauces are perceived, followed with a general tremor and sensation of chilliness. The juice applied to a wound, seemed to affect the whole nervous system; even by keeping it long in the hand, or on the bosom, we are told, unpleasant symptoms have been produced. The fatal symptoms brought on by this poison are, convulsions, giddiness, insanity, violent purgings, both upwards and downwards, faintings, cold sweats, and death itself. Dr. Stoerk appears to be the first who gave the wolf's-bane internally, as a medicine; and since his experiments were published, 1762, it has been generally and successfully employed in Germany and the northern parts of Europe, particularly as a remedy for obstinate rheumatisms; and many cases are related where this disease was of several years duration, and had withstood the efficacy of other powerful medicines, as mercury, opium, antimony, cicuta, &c. yet, in a short time, were entirely cured by the aconitum. Instances are also given us of its good effects in gout, serophulous swellings, venereal nodes, amaurosis, intermittent fevers, paralysis, ulceration, and scirrhus. This plant has been generally prepared as an extract or inspissated juice, after the manner directed in the Edinburgh and many of the foreign pharmacopœias: its efficacy is much diminished on being long kept. Like all virulent medicines, it should first be administered in

small doses. Stoerk recommends two grains of the extract to be rubbed into a powder, with two drams of sugar, and to begin with ten grains of this powder, two or three times a day. We find, however, that the extract is often given from one grain to ten for a dose; and Stoll, Scherckbecker, and others, increased this quantity considerably. Instead of the extract, a tincture has been made of the dried leaves, macerated in six times their weight of spirits of wine, and forty drops given for a dose. Some writers say that the napellus is not poisonous in Sweden, Poland, &c. but it should be noted that the napellus which is not poisonous, is the *Aconitum lycoctonum* of Linnaeus.

ACOPON. (From *a*, priv. and *xopw*, weariness.) It signifies originally whatever is a remedy against weariness, and is used in this sense by Hippocrates. Aph. viii. lib. ii. But in time, the word was applied to certain ointments.

ACOPA. According to Galen and Paulus Aegineta, the *Acopa Pharmaca* are remedies for indispositions of body which are caused by long or vehement motion. So are medicines against lassitudes.

ACOM. Acidity. It is sometimes used to express that sourness in the stomach contracted by indigestion, and from whence flatulencies and acid belching arise.

ACORDINA. An obsolete term for Indian tutty.

ACORIA. (From *a*, priv. and *xogos* to satiate.) Insatiability. In Hippocrates, it means a good appetite and digestion.

ACORITES VINUM. (From *axopw*, galangal.) A wine mentioned by Dioscorides, made with galangal, liquorice, &c. infused with wine.

ACORN. The fruit of the oak. Acorns were the food of the first ages; but when corn was cultivated, acorns were neglected. They are of little use with us, except for fattening hogs and other cattle and poultry. Among the Spaniards, the acorn, or *glans iberica*, is said to have long remained a delicacy, and to have been served up in the form of a dessert. In dearths, acorns have been sometimes dried, ground into meal, and baked as bread. Bartholin relates that they are used in Norway for this purpose. The inhabitants of Chio held out a long siege without any other food; and in a time of scarcity in France, A. D. 1709, they resorted to this food. But they are said to be hard of digestion, and to occasion headaches, flatulency, and colics. In Smoland, however, many instances occur, in which they have supplied a salutary and nutritious food. With this view they are previously boiled in water and separated from their husks, and then dried and ground; and the powder is

mixed with about one half, or one third of corn flour. A decoction of acorns is reputed good against dysenteries and colics; and a peassary of them is said to be useful in immoderate fluxes of the menes. Some have recommended the powder of acorns in intermittent fever; and in Brunswick, they mix it with warm ale, and administer it for producing a sweat in cases of erysipelas. Acorns roasted and bruised have restrained a violent diarrhoea. For other medical uses to which they have been applied, see Murray's Appar. Medic. vol. i. page 100.

From some late reports of the Academy of Sciences, at Petersburg, we learn that acorns are the best substitute to coffee that has been hitherto known. To communicate to them the oily properties of coffee, the following process is recommended. When the acorns have been toasted brown, add fresh butter in small pieces to them, while hot in the ladle, and stir them with care, or cover the ladle and shake it, that the whole may be well mixed. The acorns of the Holm oak are formed at Venice into cups about one inch and an half in diameter, and somewhat less in depth. They are used for dressing leather and instead of galls for dyeing woollen cloth black.

ACORTINUS. A lupin.

ACORUS. (*Ακωρον*, from *xopw*, the pupil; because it was esteemed good for disorders of the eyes.) The name of a genus of plants in the Linnæan system. Class, *Hexandria*. Order, *Digynia*. Sweet-flag. Sweet-rush.

ACORUS CALAMUS. The systematic name for the *calamus aromaticus*. See *Calamus aromaticus*.

ACORUS PALUSTRIS. See *Iris palustris*.

ACORUS VERUS. See *Calamus aromaticus*.

ACORUS VULGARIS. See *Iris palustris*.

ACOS. (From *ακοω*, to heal.) A remedy or cure.

ACOSMIA. (From *a*, neg. and *κοσμος*, beautiful.) Baldness; ill health; irregularity, particularly of the critical days of fevers.

ACOSTE. (From *ακωσθ*, barley.) An ancient food made of barley.

ACOUSTICA. (*Acoustica*, sc. *medicamenta*; *ακουω* from *ακουω* to hear.) Remedies which are employed with a view to restore the sense of hearing, when wanting or diminished. No internal remedies of this kind are known to produce any uniform effect.

Acoustic nerves. See *Auditory nerves*.

Acoustic duct. The external passage of the ear.

ACOUSTICS. That branch of general science which treats on the origin, propagation, and perception of sound.

ACBA. (Arab.) *Acrai nymphomania*. Ex.

cessive venereal appetite. The time of menstruation.

ACRACIA. (From *α, priv.* and *ακρας*, strength.) *Acrasia*. Debility, or impotence, from relaxation or lost tone of the parts. *Hippocrates*.

ACRAIPALA. (From *α, neg.* and *κραταια*, surfeit.) *Acraipala*. Remedies for the effects of a debauch.

ACRATISMA. (From *ακρατος*, unmixed wine.) A breakfast among the old Greeks, consisting of a morsel of bread, soaked in pure unmixed wine. The derivation of this word is the same as *Acrasia*, because the wine used on this occasion was not mixed with water.

ACRATONELLI. (From *ακρατος*, pure wine; and *μαλα*, honey.) *Mulsum*, or wine mixed with honey.

ACRE. (From *ακρας*, extreme.) The extremity of the nose.

ACRESA. (From *ακρας*, extreme.) *Acresia*. The extremities, i. e. the legs, arms, nose, and ears.

ACREPALOS. See *Acraipala*.

ACREBIA. (From *ακρις*, accurate.) An exact and accurate description and diagnosis, or distinction of diseases.

ACRID. (*Acris*.) A term employed in medicine to express a taste, the characteristic of which is pungency joined with heat.

ACRIMONY. (*Acrimonia*, from *acris*, acrid.) This term is used to express a quality in substances by which they irritate, corrode, or dissolve others. It has been supposed until very lately, there were acid and alkaline acrimonies in the blood, which produced certain diseases; and although the humoral pathology is nearly exploded, the term venereal acrimony and some others are still and must be retained.

ACRIS. Any fractured extremity.

ACRISIA. (From *α, priv.* and *ακρας*, to judge or separate.) A turbulent state of a disease, which will scarcely suffer any judgment to be formed thereof.

ACRITUS. (From *α, neg.* and *ακρις*, to judge.) Disease without regular crisis, the event of which is hazardous to judge.

ACROBUSTIA. (From *ακρας*, extreme, and *βω, to cover*.) The extremity of the prepuce.

ACROCHEIRIA. (From *ακρας*, extreme, and *χευ, a hand*.) An exercise among the ancients. Probably a species of wrestling, where they only held by the hands.

ACROCHEIRIBIA. (From *ακρας*, extreme, and *χευ, a hand*.) *Gorizus* says, it signifies the arm from the elbow to the ends of the fingers; *χευ* signifying the arm, from the scapula to the fingers' end.

ACROCHORDON. (From *ακρας*, extreme, and *χορδη, a string*.) Galen describes it as a round excrescence on the skin, with a slender base; and that it hath its name because of its situation on the surface of

the skin. The Greeks call that excrescence an *achrochordon*, where something hard concretes under the skin, which is rather rough, of the same colour as the skin, slender at the base, and broader above. Their size rarely exceeds that of a bean.

ACROCOLIA. (From *ακρας*, extreme, and *κολον, a limb*.) These are the extremities of animals, which are used in food, as the feet of calves, swine, sheep, oxen, or lambs, and of the broths of which, jellies are frequently made. *Castellus* from *Budæus* adds, that the internal parts of animals are also called by this name; in English *giblets*.

ACROLENION. *Castellus* says it is the same as *Olciranon*.

ACROMACHIA. (From *ακρας*, extreme, and *μαχη*, madness.) Total or incurable madness.

ACROMION. (From *ακρας*, extremity, and *μυος*, the shoulder.) A process of the scapula or shoulder-blade. See *Scapula*.

ACROMPHALUM. (*Ακρομφαλον*, from *ακρας*, extreme, and *μφαλον*, the navel.) *Acromphalon*. The tip of the navel.

ACROMPHALON. See *Acromphalium*.

ACRONIA. (From *ακρας*, the extremity.) The amputation of any extremity, as a finger or toe.

ACROPATHOS. (From *ακρας*, extreme, and *αθος*, a disease.) *Acropathos*. It signifies literally a disease at the top or superior part. *Hippocrates* in his treatise *De Superfizatione* applies it to the internal orifice of the uterus; and in *Prædict. lib. ii.* to cancers, which appear on the surface of the body.

ACROPATHUS. See *Acropathos*.

ACROPSIS. (From *ακρας*, the extremity, and *ωψ, the voice*.) Imperfect articulation, from a fault in the tongue.

ACROPOSTHIA. (From *ακρας*, extreme, and *σθη, the prepuce*.) The extremity of the prepuce; or that part which is cut off in circumcision.

ACROPSILON. (From *ακρας*, extreme, and *ψι, naked*.) The extremity of the denuded glans penis.

ACROPELOS. (From *ακρας*, the extremity, and *πελος*, black.) *Acrospelus*. The bromus *Dioscoridis*, or wild oat grass; so called because its ears, or tops, are often of a blackish colour.

ACROPELUS. See *Acrospelus*.

ACROTHERIA. (From *ακρας*, extreme.) The extreme parts of the body, as the hands, feet, nose, &c.

ACROTHERIASMUS. (From *ακροτηρες*, extremities, and this from *ακρας*, summus.) The amputation of an extremity.

ACROTHYMION. (From *ακρας*, extreme, and *θυμος*, thyme.) *Acrothymia*. *Acrothymum*. A sort of wart, described by *Celsus*, as hard, rough, with a narrow basis, and broad top; the top is of the colour of

thyme; it easily splits and bleeds. This tumour is also called *Thymus*.

ACETA. (From *aca*, to break.) *Acet.* The elder-tree, so called from its being easily broken. See *Sambucus*.

ACTIVA. The herb *Bunias* or *Napus*.

ACTINOMOLISMS (From *actin*, a ray, and *molle*, to cast out.) Irradiation. It is applied to the spirits, conveying the inclinations of the mind to the body: it is also called *Diradiatio*.

ACTION. (From *ago*, to act.) Any faculty, power, or function of the body, which, by physiologists are usually divided into vital, animal, or natural. The vital functions, or actions, are those which are absolutely necessary to life, and without which animals cannot exist; as the action of the heart, lungs, and arteries. The natural functions are those which are instrumental in repairing the several losses which the body sustains; digestion, and the formation of chyle, &c. fall under this head. The animal actions are those which we perform at will, as muscular motion, and all the voluntary motions of the body. Each part of the body is also said to have an action peculiar to itself.

ACTON WATER. A purging water procured from Acton, a village near London, where is a well that affords it. This is one of the strongest purging waters near London; and has been drank in the quantity of from one to three pints in a morning, against scorbutic and cutaneous affections. This medical spring is no longer resorted to by the public.

ACTUAL. This word is applied to any thing endued with a property or virtue which acts by an immediate power inherent in it: it is the reverse of potential; thus, a red-hot iron or fire is called an actual cauter, in contradistinction from caustics, which are called potential cauterics. Boiling water is actually hot; brandy, producing heat in the body, is potentially hot, though of itself cold.

ACTUATION. (From *ago*, to act.) That change wrought on a medicine, or any thing taken into the body, by the vital heat, which is necessary, in order to make it act and have its effect, is called its actuation.

ACUITAS. Acrimony.

ACUITIO. (From *acuo*, to sharpen.) The sharpening an acid medicine by an addition of something more acid; or in general, the increasing the force of any medicine, by an addition of something that hath the same sort of operation in a greater degree.

ACUXON. (From *a*, neg. and *uxon*, to well round: so called because its fruit is not involved in a cup, or sheath, like others.)

Acules. The fruit or acorn of the ilex, or scarlet oak.

ACULOS. See *Acodon*.

ACUMEN. A point. The extremity of a bone.

ACUPUNCTURA. (From *acus*, a needle, and *punctura*, a prick.) Acupuncture; bleeding performed by making many small punctures.

ACURES. *Plumbum*, or lead.

ACURON. (From *a*, neg. and *cura*, to happen.) A name of the *Alisma*: so called because it produces no effect if taken internally.

ACUSPASTORIS. A name of the *Scandix anthriscus*, the shepherd's needle, or Venus's comb. See *Scandix*.

ACUTE. *Morbus acutus.* A disease which is attended with violent symptoms, terminates in a few days, and is attended with danger. It is opposed to a chronic disease, which is slow in its progress, and not so generally dangerous.

ACUTENACULUM. (From *acus*, a needle, and *tenaculum*, a handle.) Heister calls the *portaguille* by this name. It is the handle for a needle, to make it penetrate easily when stitching a wound.

ACVISE. (From *a*, neg. and *vise*, to conceive.) In Vogel's nosology it signifies a defect of conception, or barrenness in women.

ACRAUS. (From *a*, priv. and *acus*, authority; so named from its little note in medicine.) The *Arnica montana*, or German leopard's-bane. See *Arnica*.

ADAMONIA. (From *a*, priv. and *saqueo*, a genius or fortune.) The restlessness and anxiety felt in acute fevers.

ADAISIS. Sal-ammoniac, or muriate of ammonia. See *Murias ammoniac*.

ADAMAS. (From *a*, neg. and *saqueo*, to conquer; as not being easily broken.) The adamant or diamond, the most precious of all stones, and which was formerly supposed to contain extraordinary cordial virtues.

ADAMPTA. *Adamitum.* A hard stone in the bladder.

Adam's Apple. See *Pomum Adami*.

ADAM'S NEEDLE. *Yucca gloriosa* of Linnaeus. The roots of this plant are thick and tuberos, and are used by the Indians instead of bread; being first reduced into a coarse meal. This, however, is only in times of scarcity.

ADARBS. (From *a*, neg. and *saqueo*, to acc.) A saltish concretion found about the reeds and grass in marshy grounds in Galatia, and so called because it hides them. It is used to clear the skin with, in leprosy, tetters, &c. Dr. Plott gives an account of this production in his *Natural History of Oxfordshire*. It was formerly in repute for cleansing the skin from freckles.

• **ADARBS.** An ammoniacal salt.

ADARBECK. Auripigmentum, or orpiment

Adarticulation. See *Arthrodia*.

ADDEPHAGIA. (From *ad*, abundantly, and *phagō*, to eat.) Insatiability. A voracious appetite. See *Bulimia*.

ADDITAMENTUM. (From *addo*, to add.) A term formerly employed as synonymous with *epiphysia*, but now only applied to two portions of sutures of the skull. See *Lambdoidal* and *Squamous Sutures*.

ADDITAMENTUM COLI. See *Appendix laci vermiformis*.

ADDUCTOR. (From *ad*, and *duco*, to draw.) A drawer or contractor. A name given to several muscles, whose office is to bring forwards or draw together those parts of the body to which they are annexed.

ADDUCTOR BREVIS FEMORIS. *Adductor femoris secundus* of Douglas. *Triceps secundus* of Winslow. A muscle, which, with the *adductor longus* and *magnus femoris* forms the *triceps adductor femoris*. It is situated on the posterior part of the thigh, arising tendinous from the os pubis near its joining with the opposite os pubis below, and behind the *adductor longus femoris*, and is inserted, tendinous and fleshy, into the inner and upper part of the *linea aspera*, from a little below the trochanter minor, to the beginning of the insertion of the *adductor longus*. See *Triceps adductor femoris*.

ADDUCTOR FEMORIS PRIMUS. See *Adductor longus femoris*.

ADDUCTOR FEMORIS SECUNDUS. See *Adductor brevis femoris*.

ADDUCTOR FEMORIS TERTIUS. See *Adductor magnus femoris*.

ADDUCTOR FEMORIS QUARTUS. See *Adductor magnus femoris*.

ADDUCTOR INDICIS PÉDIS. An external interosseous muscle of the fore-toe, which arises, tendinous and fleshy, by two origins, from the root of the inside of the metatarsal bone of the fore-toe, from the outside of the root of the metatarsal bone of the great-toe, and from the os cuneiforme internum. It is inserted, tendinous, into the inside of the root of the first joint of the fore-toe. Its use is to pull the fore-toe inwards from the rest of the small toes.

ADDUCTOR LONGUS FEMORIS. *Adductor femoris primus* of Douglas. *Triceps minimus* of Winslow. A muscle situated on the posterior part of the thigh, which, with the *adductor brevis*, and *magnus femoris*, forms the *triceps adductor femoris*. It arises by a pretty strong roundish tendon, from the upper and inferior part of the os pubis, and ligament of its synchondrosis, on the inner side of the pectineus, and is inserted along the middle part of the *linea aspera*. See *Triceps adductor femoris*.

ADDUCTOR MAGNUS FEMORIS. *Adductor femoris tertius et quartus* of Douglas. *Triceps magnus* of Winslow. A muscle which, with the *adductor brevis fe-*

moris, and the *adductor longus femoris*, forms the *Triceps adductor femoris*. It arises from the symphysis pubis, and all along the flat edge of the thyroid foramen, from whence it goes to be inserted into the *linea aspera* throughout its whole length. See *Triceps adductor femoris*.

ADDUCTOR MINIMI DIGITI PÉDIS. An internal interosseous muscle of the foot. It arises, tendinous and fleshy, from the inside of the root of the metatarsal bone of the little-toe. It is inserted, tendinous, into the inside of the root of the first joint of the little-toe. Its use is to pull the little-toe inwards.

ADDUCTOR OCULI. See *Rectus internus oculi*.

ADDUCTOR POLLICIS. See *Adductor pollicis manus*.

ADDUCTOR POLLICIS MANUS. *Adductor pollicis.* *Adductor ad minimum digitum.* A muscle of the thumb, situated on the hand, which arises, fleshy, from almost the whole length of the metacarpal bone that sustains the middle finger; from thence its fibres are collected together. It is inserted, tendinous, into the inner part of the root of the first bone of the thumb. Its use is to pull the thumb towards the fingers.

ADDUCTOR POLLICIS PÉDIS. *Abtithenar* of Winslow. A muscle of the great toe, situated on the foot: it arises, by a long thin tendon, from the os calcis, from the os cuboides, from the os cuneiforme externum, and from the root of the metatarsal bone of the second toe. It is inserted into the external os sesamoidem, and root of the metatarsal bone of the great toe. Its use is to bring this toe nearer to the rest.

ADDUCTOR PRESTATAE. A name given by Sanctonini to a muscle which he also calls *Levator prestatæ*, and which Winslow calls *Prestatæ superior*. Albimus, from its office, had very properly called it *compressor prestatæ*. See *Compressor prestatæ*.

ADDUCTOR TERTII DIGITI PÉDIS. An internal interosseous muscle of the foot, that arises, tendinous and fleshy, from the roots of the metatarsal bones of the third and little toe. It is inserted, tendinous, into the outside of the root of the first joint of the third toe. Its use is to pull the third toe outward.

ADDE. Soft milk, or butter-milk.

ADDECTA. See *Adectæ*.

ADDECTOS. *Adecta.* (From *a*, priv. and *decto*, to bite.) An epithet of those medicines which relieve from pain, by removing the uneasy situation caused by the stimulus of acrimonious medicines.

ADELPHIA. (*Ἀδελφία*, a relation.) Hippocrates calls diseases by this name that resemble each other.

ADEMONTA. (From *a*, priv. and *deimon*, a genus or divinity or fortune.) Hippocrates uses this word for uneasiness, restlessness,

or anxiety felt in acute diseases, and some hysterical fits.

ADEN. (*Aden*, a gland.) A gland. A bubo. See *Gland*.

ADENIFORM. (*Adeniformis*; from *aden*, a gland, and *forma*, resemblance.) Glandiform, or resembling a gland. A term sometimes applied to the prostate gland.

ADENDENTIS. An epithet applied to ulcers which eat and destroy the glands.

ADENOGRAPHY. (From *aden*, a gland, and *graphein*, to write.) A treatise on the glands. See *Gland*.

ADENOIDS. Glandiform: resembling a gland. An epithet applied also to the prostate gland.

ADENOLOGY. (From *aden*, a gland, and *logos*, a treatise.) The doctrine of the glands. See *Gland*.

ADENOUS ABSCESS. (*Abscessus adenosus*; from *aden*, a gland.) A hard glandular abscess, which suppurates slowly.

ADEPHAGIA. (From *aden*, abundantly, and *phagein*, to eat.) Insatiate appetite. See *Bulimia*.

ADEPS. Fat. An oily secretion from the blood into the cells of the cellular membrane. See *Fat*.

ADEPS ANSERINUS. Goose-grease.

ADEPS SUILLÆ. Hog's-lard.

ADEPTA MEDICINA. So Paracelsus calls that which treats of the diseases that are contracted by celestial operations, or communicated from heaven.

ADEPTA PHILOSOPHIA. Adept philosophy. It is that philosophy, whose end is the transmutation of minerals, and a universal remedy.

ADEPTA. (From *adipiscor*, to obtain.) Skilful alchemists. Such are called so as pretend to some extraordinary skill in chymistry; but these have too often proved either enthusiasts or impostors. The professors of the *Adepta Philosophia* are also called Adepts.

ADFLATUS. A blast: a kind of erysipelas.

ADMATODA. The Malabar nut-tree, which is a species of *Justicia*. It is used in India for expelling the dead fetus in an abortion, which it is said is the meaning of the word in the Zeylandic language.

ADHESION. (From *adherere*, to stick to.) The growing together of parts.

ADHESIVE INFLAMMATION. A term lately introduced into Surgery, to express that species of inflammation which terminates by an adhesion of the inflamed surfaces.

ADHESIVE PLASTER. A plaster made of common litharge plaster and resin, is so called because it is used for its adhesive properties. See *Emplastrum resine*.

ADIACHYTOS. (From *a*, neg. and *diachynein*, to diffuse, scatter, or be profuse.) Decent in point of dress. Hippocrates thinks the dress of a fop derogatory from the physi-

cian; though thereby he hides his ignorance, and obtains the good opinion of his patients.

ADIANTHUM. *Adiantum.* (*ad*, to, and *anthein*, to grow wet; so called because its leaves are not easily made wet.) Maidenhair. The name of a genus of plants in the Linnæan system. Class, *Cryptogamia*. Order, *Filices*.

ADIANTHUM CAPILLUS VENERIS. Maiden-hair. The leaves of this plant are somewhat sweet and austere to the palate, and possess mucilaginous qualities. A syrup, the *syrop de capillaire* is prepared from them, which is much esteemed in France. Orange-flower water and a proportion of honey, it is said, are usually added. It acts chiefly as a demulcent, sheathing the inflamed sides of the glottis.

ADIANTHUM AUREUM. The *Polytrichum commune* of Linnæus. It possesses, in an inferior degree, astringent virtues: and was formerly given in diseases of the lungs, and calculous complaints.

ADIAPHOROUS. A term which implies the same with neutral; and is particularly used of some spirits and salts, which are neither of an acid nor alkaline nature.

ADIANEUSTIA. (From the privative particle *a*, and *diarnein*, perspire. (A diminution or obstruction of natural perspiration, and that in which the ancients chiefly placed the cause of fevers.

ADIARRHÆA. (From *a*, priv. and *diarrhein*, to flow out or through.) A total suppression of all the necessary evacuations from the bowels.

ADIATHEOSUS. A spirit distilled from tartar.

ADIBAT. Mercury.

ADICK. (*Adica*.) A nettle.

ADIPOCIRE. (From *adeps*, fat, and *cera*, wax.) A substance that resembles soap, formed by a conversion of animal matter, placed under certain circumstances. Whole bodies have been found converted into this substance.

ADIPOSE MEMBRANE. (*Membrana adiposa*, from *adeps*, fat.) The fat collected in the cells of the cellular membrane. See *Fat*.

ADIPSAN. So the Greeks called medicines, &c. which abate thirst. Hippocrates applied this word to oxymel.

ADIPSIA. (From *a*, neg. and *diψa*, thirst.) A want of thirst. A genus of disease in the class *locales*, and order *dyssœremia* of Cullen's Nosology. It is mostly symptomatic of some disease of the brain.

ADIPOS. So the Greeks called the Egyptian palm-tree, whose fruit is said to be the *Myrobalans*. The tree is called *adiposa*, because its fruit quencheth thirst. Theophrastus calls this tree *Balanos*. *Adipos* is also a name for liquorice.

ADIRIZ. Ammoniacal salt.

ADJUTORIUM. (From *ad* and *juvō*, to help.) A name of the *humerus*, from its usefulness in lifting up the fore-arm.

ADJUVANTIA. Whatever assists in obviating disease.

ADNATA TUNICA. (*Adnata*, from *adnascor*, to grow to.) *Albuginea oculi*. *Tunica albuginea oculi*. This membrane is mostly confounded with the *conjunctiva*. It is, however, thus formed: five of the muscles which move the eye, take their origin from the bottom of the orbit, and the sixth arises from the edge of it; they are all inserted, by a tendinous expansion, into the anterior part of the *tunica sclerotica*; which expansion gives the whiteness peculiar to the fore-part of the eye. It lies betwixt the *sclerotica* and *conjunctiva*.

ADOC. Milk.

ADONION. (From *Adonis* the youth from whose blood it was feigned to have sprung.) *Adonium*. Southernwood.

ADOPTER. *Tubus intermedius*. A chymical instrument used to combine retorts to the cucurbits or matrasses in distillation, with retorts instead of receivers.

ADOR. A sort of corn, called also *spelta*.

ADOS. Water in which red-hot iron is extinguished.

AD PONDUS OMNIUM. The weight of the whole. These words are inserted in pharmaceutical preparations, or prescriptions, when the last ingredient ought to weigh as much as all the others put together.

ADRA RHIZA. Blancard says the root of the *Aristolochia* is thus named.

ADRACEN. The strawberry bay-tree. A species of *Arbutus*.

ADRAM. Fossil salt.

ADRARAGI. (Indian.) Garden-saffron.

ADROBOLON. (From *adros* large, and *bolos*, a globe, bole, or mass.) Indian *bdellium*, which is coarser than the Arabian.

Adstriction. Costiveness.

ADSTRINGENTS. See *Astringents*.

ADUSTION. An inflammation about the brain, and its membrane, with a hollow-ness of the eyes, a pale colour, and a dry body.

In Surgery, *adustion* signifies the same as cauterization, and means the application of any substance to the animal body, which acts like fire. The ancient surgeons, especially the Arabians, were remarkably fond of having recourse to *adustion* in local diseases: but the use of actual heat is very rarely admitted by the moderns.

ADVENTITIOUS. Any thing that accidentally, and not in the common course of natural causes, happens to make a part of another; as the glands in strumous cases are said to be *adventitious glands*, in distinction from those which are naturally produced. It is also used in opposition to hereditary; thus gout and scrofula are sometimes hereditary, and very often *adventitious*, they

having never before been known in the family.

ADY. *Abanga*. The palm of the island of St. Thomas, from which is prepared Thernel's restorative.

ADYNAMIA. (*Adynamia*: from *a*, priv. and *dynamis*, power.) A defect of vital power.

ADYNAMIÆ. The second order of the class *neuroses* of Cullen's Nosology; it comprehends *mycopie*, *dyspepsia*, and *hypochondriasis*.

ADYNAMON. (From *a*, neg. and *dynamis*, strength.) *Adynamum*. Among ancient physicians, it signified a kind of weak facitious wine, prepared from must, boiled down with water; to be given to patients to whom pure or genuine wine might be hurtful.

ÆNOÏA. (From *adnos*, modesty; or from *a*, neg. and *idō*, to see; as not being decent to the sight.) The pudenda, or parts of generation.

ÆDOROPHIA. (From *adnos*, pudenda, and *phos*, to break wind.) A term used by *Survages* and *Sagar*, to signify a flatus from the bladder, or from the womb, making its escape through the vagina.

ÆGAGROFILUS. (From *agros*, a wild goat, and *fila*, a ball.) *Ægagrophila*.

1. A ball found in the stomach of deer, goats, hogs, horned cattle, as cows, &c. It consists of hairs which they have swallowed from licking themselves. They are of different degrees of hardness, but have no medicinal virtues. Some rank these balls among the *Bezoars*. Hieronymus Velschius wrote a treatise on the virtues of this.

2. A species of *conserva* found in Wallenfennoor, from its resembling these concretions, is also so named.

ÆGIAS. A white speck on the pupil of the eye, which occasions a dimness of sight.

ÆGIDES. *Ægia*. A disorder of the eyes mentioned by Hippocrates. *Fœsius* thinks the disease consists of small cicatrices in the eye, caused by an afflux of corrosive humours upon the part. But in one passage of Hippocrates, *Fœsius* says it signifies small white concretions of humours which stick upon the pupil, and obscure the sight.

ÆGIDION. A collyrium or ointment for inflammations and defluxions of the eyes.

ÆGILOPS. Wild fescue grass. This plant is called *ægilops* from its supposed virtue in curing the disorder named *Ægilops*. It is a species of *Bromus* in the Linnean system.

ÆGIBETIA. Malabrian broom rape. A species of *Orobanchæ*.

ÆGIS. *Achlys*. A film on the eye.

ÆSOCHNAS. (From *as*, a goat, and *nos*, a horn: so called, because the pods were supposed to resemble the horns of a goat.) Pœnugreek. See *Trigonella Fœnum-græcum*, and *Bouccera*.

ÆGOLITHRON. (From *αἴς*, a goat, and *λίθος*, destruction; so named from the opinion of its being poisonous to goats.) Tournefort says it is the *Chamaerodendron*; now the *Azelea pontica* of Linnaeus.

ÆGOCYMON. (From *αἴς*, a goat, and *κύμη*, a hoof; because of the hardness of the seed.) Cromwell. See *Lithospermum*.

ÆGOPODIUM. (From *αἴς*, a goat, and *πῶς*, a foot; from its supposed resemblance to a goat's foot.) Goatweed. A genus of plants in the Linnaean system. Class, *Pentastria* Order, *Dignia*.

ÆGOPIDIUM. PODAGRARIA. (*Podagraria*, from its use in curing the *podagra*, or gout.) Goatweed. This plant is sedative, and was formerly applied to mitigate pains of gout, and to relieve piles, but not now employed. In its earlier state it is tender and esculent.

ÆGOPROSORON. (From *αἴς*, a goat, and *προσώρον*, a face; so called because goats are subject to defects in the eyes, or from having in it some ingredients named after the goat.) A name of a lotion for the eyes, when inflamed.

ÆGYLOPS. From *αἴς*, a goat, and *ὄψ*, an eye.) A disease so named from the supposition that goats were very subject to it.

The term means a sore just under the inner angle of the eye. The best modern surgeons seem to consider the ægylops only as a stage of the fistula lachrymalis. Paulus Ægineta calls it anchylops, before it bursts, and ægilops after. When the skin covering the lachrymal sac has been for some time inflamed, or subject to frequent returning inflammations, it most commonly happens that the puncta lachrymalia are affected by it; and the fluid, not having an opportunity of passing off by them, distends the inflamed skin, so that at last it becomes sloughy, and bursts externally. This is that state of the disease which is called perfect *aigylops*, or *ægylops*.

ÆGYPTIA MUSCATA. See *Hibiscus abelmoschus*.

ÆGYPTIACUM. A name given to different unguents of the detergent or corrosive kind. We meet with a black, a red, a white, a simple, a compound, and a magistral *Ægyptiacum*. The simple *Ægyptiacum*, which is that usually found in our shops, is a composition of verdigris, vinegar, and honey, boiled to a consistence. It is usually supposed to take its name from its dark colour, wherein it resembles that of the natives of Egypt. It is improperly called an unguent, as there is no oil, or rather fat, in it.

ÆGYPTIUM PHARMACUM AD AURES. Ætius speaks of this as excellent for detaching fetid ulcers of the ears, which he says it cures, where the patient were born with them.

ÆSOLUCA. (From *αἴς*, always and *λυκός*, sweet.) A sweetish wine, or must.

ÆIPATHEIA. (From *αἴς*, always, and *πάθος*, a disease.) Any disease of long duration.

ÆMBA. (From *αἴς*, brass, so called because it was formerly made of brass.) A catheter.

ÆOM. The apical marrow.

ÆONESIS. Fermentation. Sprinkling of the whole body.

ÆONION. The *sedum majus*, or common house-leek

ÆORA. (From *ἀνίστασθαι*, to lift up, to suspend on high.) Exercise without muscular action; as swinging. A species of exercise used by the ancients, and of which Ætius gives the following account. Gestation, while it exercises the body, the body seems to be at rest. Of the motion there are several kinds. First, swinging in a hammock, which, at the decline of a fever, is beneficial. Secondly, being carried in a litter, in which the patient either sits, or lies along. It is useful when the gout, stone, or such other disorder, attends, as does not admit of violent motions. Thirdly, riding in a chariot, which is of service in most chronic disorders; especially before the more violent exercises can be admitted. Fourthly, sailing in a ship, or boat. This produces various effects, according to the different agitation of the waters, and, in many tedious chronic disorders, is efficacious beyond what is observed from the most skilful administration of drugs. These are instances of a passive exercise.

ÆQUE. Equally. The same as *ana*.

AER. The fluid which surrounds the globe. See *Air and Atmosphere*.

ÆROS. An excrescence, or protuberance.

ÆMA. Darnel, or lolium.

ÆRITIS. The *Anagallis*, or pimpernell.

AEROLOGIA. (*Ἀερολογία*: from *αἴς*, the air, and *λόγος*, a discourse.) *Aerologie*.

Aerology. That part of medicine which treats of the nature and properties of air.

AEROLOGICE. See *Aerologia*.

AEROMELI. Honey, dew; also a name for manna.

AEROPHOBI. (From *αἴς*, air, and *φοβός*, fear.) According to Celsus Aurelianus, some phrenetic patients are afraid of a lucid and others of an obscure air: and these he calls *aerophobi*.

AEROPHOBIA. Fear of air, or wind. A symptom of the phrenitis; also a name of *Hydrophobia*.

ÆROSIS. The aerial vital spirit of the ancients.

ÆROSSUS LAPIS. So Pliny calls the *Lapis Calaminaris*, upon the supposition that it was a copper ore.

ÆRUGA. Verdigris.

ÆRUGO. (From *αἴς*, copper.) *Azager*. *Almáxadir*.

1. The rust of any metal, particularly of copper.

2. Verdigrise. See *Verdigrise*.

ÆRUGO PRÆPARATA. See *Subacetes cupri*.

ÆCHRONYTHESIS. The obscene language of the delirious.

ÆSCULUS. (*Æsculus*, from *esca*, food.) Horse-chestnut. The name of a genus of plants in the Linnæan system. Class, *Hep-tandria*. Order, *Monogynia*.

ÆSCULUS HIPPOCASTANUM. The systematic name for the hippocastanum. See *Hippocastanum*.

ÆSCAVUM. Auricalcum, or brass.

ÆSTÆTES. Freckles in the face; sun-burnings.

ÆSTHARA. Incineration, or burning of the flesh, or any other part of the body.

ÆSTUARIUM. A stove for conveying heat to all parts of the body at once. A kind of vapour bath. A vapour bath *Ambrose Percy* calls an instrument thus, which he describes for conveying heat to any particular part. *Palmaris, de morbis, contagiosis*, gives a contrivance under this name, for sweating the whole body.

ÆSTUATIO. The boiling up, or rather the fermenting of liquors when mixed.

ÆSTUS VOLATICUS. (From *æstus*, heat, and *volo*, to fly.) According to Vogel, synonymous with phlogosis. Sudden heat, or scorching, which soon goes off, but which for a time reddens the face.

ÆTHER. (*Æthyr*, a supposed fine subtle fluid.) *Liquor æthereus*. Ether. *Æther sulphuricus, nitrosus, muriaticus*, according to the acid from which it is formed combined with alcohol. A volatile liquor, obtained, by distillation, form a mixture of alcohol and a concentrated acid.

The medical properties of æther, when taken internally, are antispasmodic, cordial, and stimulant. Against nervous and typhoid fevers, all nervous diseases, but especially tetanic affections, soporose diseases from debility, asthma, palsy, spasmodic colic, hysteria, &c. it always enjoys some share of reputation. Regular practitioners seldom give so much as empirics, who sometimes venture upon large quantities, with incredible benefit. Applied externally, it is of service in the headach, toothach, and other painful affections. Thus employed, it is capable of producing two very opposite effects, according to its management; for, if it be prevented from evaporating, by covering the place to which it is applied closely with the hand, it proves a powerful stimulant and rubefacient, and excites a sensation of burning heat, as is the case with solutions of camphor in alcohol, or turpentine. In this way it is frequently used for removing pains in the head or teeth. On the contrary, if it be dropped on any part of the body, exposed freely to the air, its rapid evaporation produces an intense degree of cold; and as this is attended with a proportional diminution of bulk in the part

applied, in this way it has frequently contributed to the reduction of the intestine, in cases of strangulated hernia.

ÆTHEREA HERBA. The *Eryngium* was so called.

ÆTHEREAL OIL. An animal or vegetable oil, highly rectified, partaking, as it were, of the nature of æther.

ÆTHER SULPHURICUS. *S. Napthæ vitrioli. Ether Vitriolicus.* Sulphuric ether.

Take of rectified spirit,
Sulphuric acid, of each, by weight, a pound and a half.

Pour the spirit into a glass retort, then gradually add to it the acid, shaking it after each addition, and taking care that their temperature, during the mixture, may not exceed 120 degrees. Immerse the retort very cautiously to a sand bath, previously heated to 200 degrees, so that the liquor may boil as speedily as possible, and let the æther pass over into a tubulated receiver, to the tubulure of which another receiver is applied, and kept cold by immersion in ice, or water. Distil the liquor until a heavier part also begins to pass over, and appear under the æther in the bottom of the receiver. To the liquor which remains in the retort, pour on twelve fluidounces more of alcohol, and repeat the distillation in the same manner.

It is mostly employed as an excitant, nerve, antispasmodic, and diuretic, in cases of spasms, cardialgia, enteralgia, fevers, hysteria, cephalagia, and spasmodic asthma. The dose is from gr. xx to ʒij. Externally it cures toothach, and violent pains in the head. See *Æther*.

ÆTHER RECTIFICATUS. *Æther vitriolicus.*

Take of sulphuric æther, fourteen fluid-ounces.

Fused potash, half an ounce,

Distilled water, two fluidounces.

Dissolve the potash in the water, and add thereto the æther, shaking them well together, until they are mixed. Lastly, by means of a temperature about 200 degrees, distil over twelve ounces of rectified æther, from a large retort into a cooled receiver.

Sulphuric æther is impregnated with some sulphuric acid, as is evident in the smell, and with some ætherial oil: and these require a second process to separate them. Potash unites to the acid, and requires to be added in a state of solution, and in sufficient quantities, for the purpose of neutralizing it; and it also forms a soap with the oil. It is advantageous also to use a less quantity of water than exists in the ordinary solution of potash: and therefore the above directions are adopted in the last London Pharmacopœia. For its virtues, see *Æther*.

ÆTHIOPS. A term applied formerly to several preparations, because the powder becomes of a black colour, like the skin of an *Æthiopian*.

ÆTHIOPS ANTIMONIALIS. A preparation of antimony and mercury, once in high repute, and still employed by some practitioners in cutaneous diseases. A few grains are to be given at first, and the quantity increased as the stomach can bear it.

ÆTHIOPS MARTIALIS. A preparation of iron, formerly in repute, but now neglected.

ÆTHIOPS MINERAL. The substance heretofore known by this name, is called, by the London College, *Hydrargyrum cum sulphure*; by the Edinburgh, *Sulphuretum Hydrargyri nigrum*; and by that of Dublin, *Hydrargyrum sulphuratum nigrum*.

Æthmoid artery. See *Ethmoid artery*.

Æthmoid bone. See *Ethmoid bone*.

ÆTHNA. A chymical furnace.

ÆTHOCES. *Ætholices.* Superficial pustules in the skin, raised by heat, as boils, fiery pustules.

ÆTHUSA. (From *αἶσθα*, beggarly.) The name of a genus of plants of the Linnean system. Class, *Pentandria*. Order, *Digynia*.

ÆTHUSA MEUM. The systematic name of the meum of the Pharmacopœias. See *Meum Athamanticum*.

ÆTHYA. A mortar.

ÆTIOI PHLEBES. Eagle veins. The veins which pass through the temples to the head, were so called formerly by Rufus Ephesius.

ÆTIOLOGY. (*ἁιτιολογία*: from *αἴτιον*, a cause, and *λογος*, a discourse.) The doctrine of the causes of diseases.

ÆTOLION. *Ætolum.* The granum nidium.

ÆTONYCHUM. See *Lithospermum*.

AFFECTION. (This is expressed in Greek by *πάθος*: hence *pathema*, *passio*.) This term indicates any existing disorder of the whole body, or a part of it, as hysterics, colic, leprosy. Thus by adding a descriptive epithet to the term affection, most distempers may be expressed. We say febrile affection, cutaneous affection, &c. using the word affection synonymously with disease.

AFFINITY. (*Affinitas*, a proximity of relationship.) The term affinity is used indifferently with attraction. See *Attraction*.

Affinity of Aggregation. See *Attraction of Aggregation*.

Affinity of Composition. See *Attraction, chymical*.

Affinity, compound. When three or more bodies, on account of their mutual affinity, unite and form one homogeneous body, then the affinity is termed compound affinity or attraction: thus, if to a solution of sugar and water be added spirits of wine, these three bodies will form a homogeneous liquid by compound affinity. See *Attraction*.

Affinity, double. Double elective attraction. When two bodies, each consisting of

two elementary parts, come into contact, and are decomposed, so that their elements become reciprocally united, and produce two new compound bodies, the decomposition is then termed, decomposition by double affinity: thus, if we add common salt, which consists of muriatic acid and soda, to nitrate of silver, which is composed of nitric acid and silver, these two bodies will be decomposed; for the nitric acid unites with the soda, and the silver with the muriatic acid, and thus may be obtained two new bodies. The common salt and nitrate of silver therefore mutually decompose each other by what is called double affinity. See *Attraction*.

Affinity, intermediate. Appropriate affinity. Affinity of an intermedium—is, when two substances of different kinds, that show to one another no component affinity, do, by the assistance of a third, combine, and unite into a homogeneous whole: thus, oil and water are substances of different kinds, which, by means of alkali, combine and unite into an homogeneous substance: hence the theory of lixivia, of washing, &c. See *Attraction*.

Affinity, quiescent. Mr. Kirwan employs the term *Quiescent affinity* to mark that, by virtue of which, the principle of each compound of two bodies, decomposed by double affinity, adhere to each other; and *Divellent affinity*, to distinguish that by which the principles of one body unite and change order with those of the other: thus sulphate of potash or vitriolated tartar is not completely decomposed by the nitric acid or by lime, when either of these principles is separately presented; but if the nitric acid be combined with lime, this nitrate of lime will decompose the sulphate of potash. In this last case the affinity of the sulphuric acid with the alkali is weakened by its affinity to the lime. This acid, therefore, is subject to two affinities, the one which retains it to the alkali, called quiescent, and the other which attracts it towards the lime, called divellent affinity.

Affinity, divellent. See *Affinity quiescent*.

Affinity reciprocal. When a compound of two bodies is decomposed by a third; the separated principle being in its turn capable of decomposing the new combination.

Affinity, simple. Single elective Attraction. If a body consisting of two component parts, be decomposed on the approach of a third, which has a greater affinity with one of those component parts than with the other, then the decomposition is termed decomposition by simple affinity; for instance, if pure potash be added to a combination of nitric acid and lime, the union which existed between these two bodies will cease, because the potash combines with the nitric acid, and the lime being

disengaged is precipitated. The reason is, that the nitric acid has a greater affinity for the pure potash than for the lime, therefore it deserts the lime, to combine with the potash. When two bodies only enter into chymical union, the affinity, which was the cause of it, is also termed simple or single elective attraction; thus the solution of sugar and water is produced by simple affinity, because there are but two bodies. See *Attraction*.

AFIUM. *Affium*. An Arabic name for opium.

AFILATUS. (From *ad* and *flare*, to blow.) A vapour or blast. A species of erysipelas, which attacks people suddenly, so named upon the erroneous supposition that it was produced by some unwholesome wind blowing on the part.

AFUSIO. Pouring a liquor upon something; but sometimes it means the same as *suffusio*, a cataract.

After-birth. See *Placenta*.

AFIUM. An Arabic name for opium.

AGA CRETENSIS. The small Spanish milk-thistle.

AGALACTATIO. See *Agalactia*.

AGALACTIA. (*Ἀγαλακία*: from *a*, priv. and *γала*, milk.) *Agalactia*, *Agalactio*. *Agalactatio*. A defect of milk in childbirth.

AGALACTOS. (From *a*, priv. and *γала*, milk.) An epithet given to women who have no milk when they lie in.

AGALACTOS. See *Agalactia*.

AGALAXIS. See *Agalactia*.

AGALLOCHI VERI LIGNUM. See *Lignum aloes*.

AGALLUGUM. *Agallugum*. A name of the agallochum or aromatic aloe.

Agaric. See *Agaricus*.

AGARACOIDES. A species of *agaricus* or fungus.

AGARICUS. (*Ἀγαρίκος*: from *Agaria*, a town in Asia; or from *Agarus*, a river in Sarmatia, now Malowouda.) *Agaric*. The name of a genus of plants in the Linnæan system. Class, *Cryptogamia*. Order, *Fungi*.

AGARICUS CHIRURGORUM. *Agaricus quercus*. *Fungus igniarius*. *Boletus igniarius*. *Agaric* of the oak. Touchwood boletus. Female agaric. This fungus *Boletus igniarius* of Linnæus:—*acaulis pubinatus levis*, *peris tenuissimus*, has been much used by surgeons as an external styptic. Though still employed on the Continent, the surgeons in this country have not much confidence in it.

AGARICUS ALBUS. The plant known by this name in the pharmacopœias, is the *Boletus loricis* of Linnæus; so called from its being met with on old larch trees, in different parts of Europe. Several preparations, as troches, an extract, and pills, are ordered to be made with it in foreign pharmacopœias, which are administered against phthisical complaints.

AGALLUGUM. See *Agallugi*.

AGARICUS CAMPESTRIS. See *Mushroom*.

AGARICUS CHATARELLUS. A species of fungus, esteemed a delicacy by the French. Broiled with salt and pepper, it has much the flavour of a roasted cockle.

AGARICUS CINNAMOMUS. Brown Mushroom. A species of *agaricus*, of a pleasant smell. When broiled, it gives a good flavour.

AGARICUS DELICIOSUS. This fungus well seasoned and then broiled has the exact flavour of a roasted muscle. It is in season in September.

AGARICUS MUSCARIUS. Bug Agaric, so called from its known virtue in destroying bugs. This reddish fungus is the *Agaricus muscarius* of Linnæus:—*stipitatus, lamellis dimidiatis solitariis, stipite volutato, apice dilatato, basi ovato*. The use of this vegetable is not much known in this country. Haller relates that six persons of Lithuania perished at one time, by eating this kind of mushroom, and that in others it has caused delirium. It is employed externally to strumous, phagedenic, and fistulous ulcers, as an escharotic.

AGARICUS PIPERATUS. The plant thus named by Linnæus, is the pepper mushroom, also called pepper agaric. It is the *Fungus piperatus albus, lacteo-succo turgens* of Ray. *Fungus albus acris*. When freely taken, fatal consequences are related by several writers to have been the result. When this vegetable has even lost its acrid juice by drying, its caustic quality still remains.

AGARICUS PRATENSES. The Champignon of Hudson's Flora Anglica. This plant has but little smell, and is rather dry, yet when broiled and stewed, communicates a good flavour.

AGARICUS VIOLACEUS. Violet mushroom. This fungus requires much broiling, but when sufficiently done and seasoned, it is as delicious as an oyster. Hudson's bulbosus is only a variety of this.

AGE. The ancients reckoned six stages of life: *pueritia*, childhood, which is to the fifth year of our age;—*adulescentia*, youth, reckoned to the eighteenth, and youth properly so called, to the twenty-fifth year;—*juventus*, reckoned from the twenty-fifth to the thirty-fifth year;—*virilis ætas*, manhood, from the thirty-fifth to the fiftieth year;—*senectus*, old age, from fifty to sixty;—*crepita ætas*, decrepid age, which ends in death.

AGRESIA. (*Ἀγρῆσις*: from *a*, neg. and *γρομαι*, to beget.) Impotency in man. A term employed by Vægl. It is synonymous with *anaphrodisia* and *dyspermatismus*.

AGER. The common earth or soil.

AGER NATURÆ. The womb.

AGERATUS LAPIS. (*Ageratus* common.) A stone used by cobblers. It is ridiculously said to be discutient and gently astringent

If it possess any such virtues, it probably contains iron; a supposition countenanced by its being used in dyeing.

AGERATUM. (*Ἀγέραιος*: from *α*, priv. and *γέραιος*, senectus; never old, ever green; because its flowers preserve their beauty a long time.) See *Achillea ageratum*.

AGERS. (From *αγρός*, wicked; so called because it is generally the instrument of wicked acts.) The palm or hollow of the hand.

AGEUSTIA. (From *α*, neg. and *γεύωμαι*, guto, to taste.) *Agheustia*, *Apoguestia*, *Apogeusia*.

A defect or loss of taste. Callen ranks this as a genus of disease in the class *locales* and order *dysæsthesiæ*. The causes are fever or palsy, whence he forms two species: the latter he calls *organic*, arising from some affection in the membrane of the tongue, by which relishing things, or those which have some taste, are prevented from coming into contact with the nerves: the other *atomic*, arising without any affection of the tongue.

AGGLUTINANTIA. Adhesive medicines which heal by causing the parts to stick together.

AGGLUTINATIO. Agglutination. The adhesive union or sticking together of substances.

AGGLUTITIO. Obstruction in the œsophagus, or a difficulty in swallowing.

AGGREGATE GLANDS. (From *aggrego*, to assemble together.) An assemblage of glands, as those of the intestines.

Aggregation, affinity of. See *Attraction*.

AGHEUSTIA. See *Ageustia*.

AGIS. The thigh or femur.

AGITATORIA. Convulsive diseases, or those called clonic.

AGLACTATIO. Defect of milk.

AGLAXIS. See *Egides*.

AGLIA. *Aglium*. A shining tubercle or pustule on the face. White specks on the eye.

AGNA. *Agme*. A fracture.

AGNACAL. A tree, which, according to Ray, grows about the isthmus of Darien, and resembles a pear-tree, whose fruit is a great provocant to venery.

AGNATA. See *Adnata tunica*.

AGNINA MEMBRANA. (From *αγνός*, a lamb, and *membrana*, a membrane.) Aëtius calls one of the membranes which involve the fœtus by this name, which he derives from its tenderness. See *Ammios*.

AGNOIA. (From *α*, priv. and *γινώσκω*, to know.) Forgetfulness; a symptomatic affection in fevers.

AGNUS CASTUS. (From *αγνός*, a lamb; so called from the down upon its surface, which resembles that upon a lamb's skin; and *castus*, because the chaste matrons, at the feasts of Ceres, strewed them upon their beds, and lay upon them.) The plant bear-

ing this name in the pharmacopœias is the *Pistex agnus castus* of Linnaeus:—*foliis digitatis, serratis, stipulis verticillatis*. The seeds are the medicinal part, which have, when fresh, a fragrant smell and an acrid aromatic taste. Formerly they were celebrated as antaphrodisiacs; but experience does not discover in them any degree of such virtue, and some have ascribed to them an opposite one. They are now fallen into disuse.

AGNOER. The deduction or reasoning upon diseases from their symptoms and appearances. The order, state, or tenour of a disease or body.

AGOMPHIASIS. A looseness of the teeth.

AGONE. (*Ἄγων*: from *α*, neg. and *γινώσκω*, offspring.) Hyoscyamus or Henbane; so called because it was supposed to cause barrenness.

AGONIA. Sterility, impotence, agony.

AGONISTICUM. (*Ἀγωνιστικόν*, from *αγώνισμα*, to struggle.) A term used by ancient physicians to signify water extremely cold, which was directed to be given in large quantities, in acute erysipelatous fevers, with a view of overpowering or struggling with the febrile heat of the blood.

AGOROS. (From *α*, priv. and *γινώσκω*, or *γόνι*, an offspring; barren.) Hippocrates calls those women so who have not children, though they might have if the impediment were removed.

AGOSTOS. (From *αγείν*, to bring, or lead.) That part of the arm from the elbow to the fingers; also the palm or hollow of the hand.

AGRESTA. (*Ἄγρος*, wild.) Verjuice, which is made from the wild apple. The immature fruit of the vine.

AGRESTEA. A name for the common tartar.

AGRESTIS. In the works of some old writers it expresses an ungovernable malignity in a disease.

AGRIA. Holly. A malignant pustule, of which, the ancient surgeons describe two sorts; one which has been so called, is small, and casts a roughness or redness over the skin, slightly corroding it, smooth about its centre, spreads slowly, and is of a round figure; this sort is cured by rubbing it with the fasting spittle. The second ulcerates, with a violent redness and corrosion, so as to make the hair fall off; it is of an unequal form, and turns leprous; it is cured, by the application of pellitory of the wall in the manner of a poultice.

AGRIANPELOS. (From *αγρίος*, wild, and *αμπέλος*, a vine.) The wild vine, or white bryony. See *Bryonia*.

AGRIELEA. (From *αγρίος*, wild, and *ελεα*, the olive-tree.) The oleaster, or wild olive.

AGRIFOLIUM. (From *ακκ*, a prickle, and *φύλλον*, a leaf.) Aquifolium, or holly tree.

It should rather be called *acifolium* from its prickly leaves.

AGRIMONIA. (*Ἀγρίμον*: from *αγρός*, a

field, and *μωρος*, alone: so named from its being the chief of all wild herbs.) Agrimony.

1. The name of a genus of plants in the Linnaean system.

2. The pharmacopœial name of a plant; the common agrimony. *Agrimonia Eupatoria* of Linnaeus:—*foliis caulibus pinnatis, foliis undique serratis, omnibus minutis interstinctis, fructibus hispida*.

This plant is common in fields about hedges and shady places, flowering in June and July. It has been principally regarded in the character of a mild astringent and corroborant, and many authors recommend it as a deobstruent, especially in hepatic and other visceral obstructions. Chomel relates two instances of its successful use in cases where the liver was much enlarged and indurated. It has been used with advantage in hæmorrhagic affections, and to give tone to a lax and weak state of the solids. In cutaneous disorders, particularly in scabies, we have been told that it manifests great efficacy; for this purpose it was given infused with liquorice in the form of tea; but according to Alston it should be always exhibited in the state of powder. It is best used while fresh, and the tops, before the flowers are formed, possess the most virtue. Cullen observes that the agrimony has some astringent powers, but they are feeble; and pays little attention to what has been said in its favour.

AGRIMONIA EUPATORIA. (Called *Eupatori*; from *Eupator*, its inventor; or quasi *hepatorum*, *μωροτων*; from *μωρος*, the liver; because it is useful in disease of the liver.) The systematic name for the *Agrimonia* of the pharmacopœias. See *Agrimonia*.

Agrimony, hemp. The *Bidens tripartita* of Linnaeus.

AGRIOCARDANUM. (From *αγρος*, wild, and *καρδαμον*, the nasturtium.) Scitica cresses, or wild garden cress.

AGRIOCASTANUM. (From *αγρος*, wild, and *καστανον*, the chestnut.) Earth-nut or pig-nut.

AGRIOCINARA. (From *αγρος*, wild, and *κιναρα*, artichoke.) See *Cinara*.

AGRIOCOCINELA. (From *αγρος*, wild, *κωκκος*, a berry, and *μυλας*, an apple-tree.) The *prunus sylvestris*.

AGRIOMELA. The crab-apple.

AGRION. *Agriophyllum*. The peucedanum silaus, or hog's fennel, or sulphur wort.

AGRIOPHYLLON. See *Agrien*.

AGRIOPASTINACA. (From *αγρος*, wild, and *pastinaca*, a carrot.) Wild carrot, or parsnip.

AGRIORIGANUM. (From *αγρος*, wild, and *origανον*, marjoram.) Wild marjoram. See *Origanum*.

AGROSCELINUM. (From *αγρος*, wild, and *σκαλον*, paraley.) Wild-paraley. See *Hippocelsum*.

AGRIOSTARI. (From *αγρος*, wild, and *σταιν*, wild wheat.) A species of field corn called *Triticum creticum*.

AGRIPALMA. (From *αγρος*, wild, and *παλμα*, a palm tree.) *Agripalma gallia*. The herb mother-wort, or wild palm.

AGRIPALMA GALLIS. See *Agripalma*.

AGRIPE. Those children which are born with their feet foremost, are so called, because that was said to be the case with Agrippa the Roman, who was named *ad agro partu*, from his difficult birth. These births, though reckoned preter-natural, are often more safe and easy than the natural.

AGRIUM. An impure sort of natron, or soda. The purer sort was called *halmyrha*.

AGROX. A disease of the tongue peculiar to the Indians, in which it becomes extremely rough and chopped.

AGROMINA. Leeks, wild onions.

AGRYPNIA. (From *α*, priv. and *γρυπν*, sleep.) Watchfulness: want of sleep.

AGRYPNOCOMA. (From *αγρυπν*, without sleep, and *κομα*, a lethargy.) A lethargic kind of watchfulness, in which the patient is stupidly drowsy, and yet cannot asleep. A species of coma.

Ague. See *Febris Intermitiens*.

AGUS CAKE. The popular name for a hard tumour on the left side of the belly, lower than the false ribs in the region of the spleen, said to be the effect of intermittent fevers. However frequent it might have been formerly, it is now very rare, and although then said to be owing to the use of bark, it is now less frequent since the bark has been generally employed.

AGUS DROPS. This is a medicine sold for the cure of agues, composed of arseniate of potash in solution in water.

AGUS-FREE. A name given by some to *sassafras* on account of its supposed febrile virtue.

AGUS TREE. See *Laurus*.

AGUA. (From *a*, priv. and *γυν*, a member.) Paralytic debility. Where the use of the members is defective or lost.

AGUL. (Arab.) *Alhagi*. The Syrian thorn. The leaves are purgative.

AGUSTINE. A new earth discovered in the Saxon Beryl, or Beryl, of Georgien Stadt, a stone greatly resembling the Beryl of Siberia, by professor Tromsdorff of Erfurth in Germany, to which he has given the name of agustine on account of the property of forming salts which are nearly destitute of taste.

This earth is white and insipid; when moistened with water, it is somewhat ductile, but is not soluble in that fluid. Exposed to a violent heat, it becomes extremely hard, but acquires no taste. It combines with acids, forming salts, which have little or no taste. It does not combine either in the humid or dry way with alkalies,

or with their carbonates. It retains carbonic acid but feebly. It dissolves in acids equally well after having been hardened, by exposure to heat, as when newly precipitated. With sulphuric acid it forms a salt which is insipid, and scarcely soluble, but an excess of acid renders it soluble, and capable of crystallizing in stars. With an excess of phosphoric acid it forms a very soluble salt. With nitrous acid it forms a salt scarcely soluble.

ARUTIGUEPOOBI BRAZILIENSIS. (An Indian term.) Arrow-root: dartwort. Esculent and vulnerary, and used by the Indians to cure wounds made by arrows.

ASTION. See *Aguia*.

AGYRÆ (From *αγρεν* a crowd of people or a mob; or from *αγειν*, to gather together.) It formerly expressed certain strollers who pretended to strange things from supernatural assistances; but of late it is applied to all quack and illiterate dabblers in medicine.

AHALOTH. The Hebrew name of *lignum aloes*.

ACHAMELLA. See *Achmella*.

AROVAI THEVETICLUSH. A chesnut-like fruit of Brazil of a poisonous nature.

ARUSAL. Orpiment.

AJURAZAT. Lead.

AILNAD. An Arabian name for antimony.

AIMATEIA. A black bilious and bloody discharge from the bowels.

AIMORRHOIS. See *Hæmorrhœis*.

AIMORRHOEA. See *Hæmorrhagia*.

AIPATHEIA. (From *αι* always, and *παθος*, a disease.) A disease of long continuance.

AIPH. *Aipima cœtera.* *Aipipoca.* Indian words for Cassada. A poisonous root of India.

AIR. Common air. Atmospheric air. The word air seems to have been used at first to have denoted the atmosphere in general; but philosophers afterwards restricted it to the elastic fluid, which constitutes the greatest and the most important part of the atmosphere, excluding the water and the other foreign bodies which are occasionally found mixed with it. See *Atmosphere*.

Air is an elastic fluid, invisible indeed, but easily recognised by its properties. Its specific gravity, according to the experiments of Sir George Shuckburgh, when the barometer is at 30 inches, and the thermometer between 50 and 60 deg. is 0.0012, or 816 times lighter than water. One hundred cubic inches of air weigh 31 grains troy. But as air is an elastic fluid, and compressed at the surface of the earth by the whole weight of the incumbent atmosphere, its destiny diminishes according to its height above the surface of the earth. From the experiments of Pascal, Deluc,

General Roy, &c. it has been ascertained that the density diminishes in the ratio of the compression. Consequently the density decreases in a geometrical progression, while the heights increase in an arithmetical progression. Bouguer had suspected, from his observations made on the Andes, that at considerable heights the density of the air is no longer proportional to the compressing force: but the experiments of Saussure junior, made upon Mount Rose, have demonstrated the contrary.

Air is dilated by heat. From the experiments of General Roy and Sir George Shuckburgh, compared with those of Trembley, &c. it appears, that at the temperature of 60 deg. every degree of temperature increases the bulk of air about 1-82d part.

The specific caloric of air, according to the experiments of Dr. Crawford, is 1.79.

Although the sky is well known to have a blue colour, yet it cannot be doubted that air itself is altogether colourless and invisible. The blue colour of the sky is occasioned by the vapours which are always mixed with the air, and which have the property of reflecting the blue rays more copiously than any other. This has been proved by the experiments which Saussure made with his *cyanometer* at different heights above the surface of the earth. This consisted of a circular band of paper, divided into 51 parts, each of which were painted with a different shade of blue; beginning with the deepest mixed with black, to the lightest mixed with white. He found that the colour of the sky always corresponds with a deeper shade of blue, the higher the observer is placed above the surface; consequently, at a certain height, the blue will disappear altogether, and the sky appear black; that is to say, will reflect no light at all. The colour becomes always lighter in proportion to the vapours mixed with the air. Hence it is evidently owing to them.

The property which the air has of supporting combustion, and the necessity of it for respiration, are too well known to require any description.

For many ages, air was considered as an element, or simple substance. For the knowledge of its component parts, we are indebted to the labours of those philosophers in whose hands chymistry advanced with such rapidity during the last forty years of the eighteenth century.

Air is a compound of oxygen and nitrogen: but it becomes a question of considerable consequence to determine the proportion of these two ingredients, and to ascertain whether that proportion is in every case the same. Since nitrogen gas,

one of the component parts of that fluid, cannot be separated by any substance with which chymists are acquainted, the analysis of air can only be attempted by exposing it to the action of those bodies which have the property of absorbing its oxygen. By these bodies the oxygen gas is separated, and nitrogen gas is left behind, and the proportion of oxygen may be ascertained by the diminution of bulk; which, once known, it is easy to ascertain the proportion of nitrogen gas, and thus to determine the exact relative quantity of the component parts.

After the composition of the atmosphere was known to philosophers, it was taken for granted that the proportion of its oxygen varies in different times and in different places; and that upon this variation the purity or noxious qualities of air depended. Hence it became an object of the greatest importance to be in possession of a method of determining readily the quantity of oxygen in a given portion of air. Accordingly various methods were proposed, all of them depending upon the property which a variety of bodies possesses of absorbing the oxygen of the air, without acting upon its azot. These bodies were mixed with a certain known quantity of atmospheric air, in graduated glass vessels inverted over water, and the proportion of oxygen was determined by the diminution of bulk. These instruments received the name of *eudiometers*, because they were considered as measures of the purity of air. See *Eudiometer*.

It is considered as established by experiment, that air is composed of 0.22 of oxygen gas, and 0.78 of nitrogen gas by bulk. But as the weight of these two gases is not exactly the same, the proportion of the component parts by weight will differ a little: for as the specific gravity of oxygen gas is to that of nitrogen gas as 135 : 115, it follows that 100 parts of air are composed by weight of about 74 nitrogen gas
26, oxygen gas.

100

Having thus ascertained the nature and the proportion of the component parts of air, it remains only to inquire in what manner these component parts are united. Are they merely mixed together mechanically, or are they combined chymically? Is air a mechanical mixture, or a chymical compound? Philosophers seem at first to have adopted the former of these opinions, if we except Scheele, who always considered air as a chymical compound. But the supposition that air is a mechanical mixture, by no means agrees with the phenomena which it exhibits. If the two gases were only mixed together, as their specific gravity is different, it is scarcely possible that they would be uniformly mixed in every part of the atmosphere. Even Mr. Dalton's ingenious supposition, that they neither at-

tract nor repel each other, would not account for this equal distribution; for undoubtedly, on that supposition, they would arrange themselves according to their specific gravity. Since, therefore, air is in all places composed of the same ingredients, exactly in the same proportions, it follows that its component parts are not only mixed, but actually combined. When substances differing in specific gravity combine together, the specific gravity of the compound is usually greater than the mean. This holds also with respect to air. The specific gravity, by calculation, amounts only to 0.00119, whereas it actually is 0.0012; a difference by no means inconsiderable. But perhaps the specific gravity of nitrogen and oxygen gas can scarcely be considered as known with such precision as to entitle us to draw any consequence from this difference.

The difference between air and a mere mixture of its two component parts, has been demonstrated by the experiments of Morozzo and Humbolt. The artificial mixture is much more diminished by nitrous gas than air, even when the mixture contains less oxygen. It supports flame better and longer, and animals do not live in it the same time that they do in an equal portion of air, but longer.

The air is, therefore, to be considered as a chymical compound. Hence the reason that it is in all cases the same, notwithstanding the numerous decomposing processes to which it is subjected. The breathing of animals, combustion, and a thousand other operations, are constantly abstracting its oxygen, and decomposing it. The air thus decomposed or vitiated no doubt ascends in the atmosphere, and is again, by some unknown process or other, reconverted into atmospherical air. But the nature of these changes is at present concealed under an impenetrable veil. *Thompson.*

Air, alkaline. See *Ammonia*.

Air, atmospherical. See *Air*.

Air, azotic. See *Nitrogen gas*.

Air, fixed. See *Carbonic acid gas*.

Air, fluoric. See *Fluoric acid gas*.

Air, hepatic. See *Sulphuretted hydrogen gas*.

Air, inflammable. See *Hydrogen gas*.

Air, marine. See *Muriatic acid gas*.

Air, nitrous. See *Nitrous Oxide*.

Air, phlogisticated. See *Nitrogen gas*.

Air, phosphoric. See *Phosphoric acid gas*.

Air, sulphureous. See *Sulphuretted hydrogen gas*.

Air, vital. See *Oxygen gas*.

AISTHETERIUM. (From *αἰσθητός*, to perceive.) The sensorium commune, or common sensory, or seat, or origin of sensation. Cartesius and others say, it is the pineal gland; Willis says it is where the nerves of the external senses are terminated, which is about the beginning of the medulla ob-

longata, (or top of the spinal marrow,) in the corpus striatum.

AITMAD. Antimony.

AIX LA CHAPELLE. Called Aken by the Germans. *Thermæ Aquis-granensis*. A town in the south of France, where there is a sulphureous water, the most striking feature of which, and which is almost peculiar to it, is the unusual quantity of sulphur it contains; the whole, however, is so far united to a gaseous basis, as to be entirely volatilized by heat; so that none is left in the residuum after evaporation. In colour it is pellucid, in smell sulphureous, and in taste saline, bitterish, and rather alkaline. The temperature of these waters varies considerably, according to the distance from the source and the spring itself. In the well of the hottest bath, it is according to Lucas 136°, Monet 146°; at the fountain where it is drunk, it is 112°. This thermal water is much resorted to on the Continent, for a variety of complaints. It is found essentially serviceable in the numerous symptoms of disorders in the stomach and biliary organs, that follow a life of high indulgence in the luxuries of the table; in nephritic cases, which produce pain in the loins, and thick mucous urine with difficult micturition. As the heating qualities of this water are as decided as in any of the mineral springs, it should be avoided in cases of a general inflammatory tendency, in hectic fever and ulceration of the lungs: and in a disposition to active hæmorrhagy. As a hot bath, this water is even more valuable and more extensively employed than as an internal remedy. The baths of Aix la Chapelle may be said to be more particularly medicated than any other that we are acquainted with. They possess both temperature of any degree that can be borne, and a strong impregnation with sulphur in its most active forms, and a quantity of alkali which is sufficient to give it a very soft soapy feel, and to render it more detergent than common water. From these circumstances these baths will be found of particular service in stiffness and rigidity of the joints and ligaments, which is left by the inflammation of gout and rheumatism, and in the debility of palsy, where the highest degree of heat which the skin can bear is required. The sulphureous ingredient renders it highly active in almost every cutaneous eruption, and in general in every foulness of the skin; and here the internal use of the water should attend that of the bath. These waters are also much employed in the distressing debility which follows a long course of mercury and excessive salivation. Aken water is one of the few natural springs, that are hot enough to be employed as a vapour bath, without the addition of artificial heat. It is employed both in cases in which the hot bath is used, and is found to be a remarkably powerful auxiliary in curing some of the worst

species of cutaneous disorders. With regard to the dose of this water to be begun with, or the degree of heat to bathe in, it is in all cases best to begin with small quantities and low degrees of heat, and gradually increase them, agreeably to the effects and constitution of the patient. The usual time of the year for drinking these waters, is from the beginning of May to the middle of June, or from the middle of August to the latter end of September.

AIZOON. (From *ai* always, and *zoo* to live.) *Aizoon*. An evergreen aquatic plant, like the aloe said to possess antiscorbutic virtues.

AJAYA. (Indian.) A said used in the East Indies as a remedy for the cholera.

AL. The Arabian article which signifies *the*; it is applied to a word by way of eminence, as the Greek *o* is. The Easterns express the superlative by adding *God* thereto, as *the mountain of God*, for the highest mountain; and it is probable that *Al* relates to the word *Alla*, God: so *alchemy* may be *the chymistry of God*, or the most exalted perfection of chymical science.

ALA. A wing. The arm-pit, so called because it answers to the pit under the wing of a bird.

ALABARI. Lead.

ALFORMIS. Any thing like a wing

ALB AURIS. The upper part of the external ear.

ALB NASI. Two cartilages of the nose which form the nostrils.

ALB VESPERTILIONUM. That part of the ligaments of the womb, which lies between the tubes and the ovaria; so called from its resemblance to the wing of a bat.

ALB INTERNÆ MINORÆ. See *Nympha*.

ALAFI. *Alafor*. Alkaline.

ALLA PETHISIS. (From *alam*, blind, and *phok*, a wasting.) A consumption from a flux of humours from the head.

ALAMAD. *Alamed*. Antimony.

ALAMBIC. Mercury.

ALANBARLA. (Arab. bitter.) The bitter apple, or colocynth.

ALANTUTA. (Arab.) A vein between the chin and lower lip, which was formerly opened to prevent fetid breath.

ALAPOUTH. See *Bikimbi*.

ALARE EXTERNUM. A name of the external pterygoid muscle; so called because it takes its rise from the wing-like process of the sphenoid bone.

ALARIA OSSA. The wing-like processes of the sphenoid bone.

ALARIS VENÆ. The innermost of the three veins in the bend of the arm.

ALASALET *Alaset*. Ammoniacum.

ALASI. *Alafor*. An alkaline salt,

ALASTROB. Lead.

ALATAN. Litharge.

ALATERNUS. A species of rhamnus.

ALATL. Those who have prominent scapulae like the wings of birds.

ALAUBAT. Nitre.

ALBADAL. An Arabic name for the sesamoid bone of the first joint of the great toe.

ALBAGHIEL. *Albagiassi.* An Arabic name for the os sacrum.

ALBAGHIS STORA. So Avicenna names the lepra ischthyocis. Others call it lepra GRÆCORUM.

ALBAMENTUM. (From *albus*, white.) The white of an egg.

ALBANUM. Urinous salt.

ALBANA. (Chald.) The white leprosy.

ALBARAS. Arsenic. A white pustule.

ALBATIO. (From *alba*, to whiten.) *Alificatio.* The calcination or whitening of metals.

ALBERAS. (Arab.) White pustules on the face: also staphisagria, because its juice was said to remove these pustules.

ALBESTORE. Quick lime.

ALBETAD. Galbanum.

ALBI SUBLIMATI. Muriated mercury.

ALBICANTIA CORPORA. (From *albes*, to grow white.) The glands of a white colour which are usually called Willis's glands in the brain.

ALBIMENT. Orpiment. See *Auripigmentum*.

ALBINUM. See *Gnaphalium*.

ALBOR. Urine.

ALBORA. A sort of itch; or rather of leprosy. Paracelsus says, it is a complication of the morpew, serpigo, and leprosy. When cicatrices appear in the face like the serpigo, and then turn to small blisters of the nature of the morpew, it is the albora. It terminates without ulceration, but by fetid evacuations in the mouth and nostrils; it is also seated in the root of the tongue.

ALBORA. Quicksilver.

ALBOT. A crucible.

ALBOTAL. Turpentine.

ALBOTAR. Turpentine.

ALBOTAT. White lead.

ALBOTIK. Turpentine.

ALBOTIS. A cutaneous phlegmon or boil.

ALBUGINEA OCULI. (From *albus*, white.) See *Achna tunica*.

ALBUGINEA TESTIS. (*Albuginea*; from *albus*, white; so called on account of its white colour.) *Tunica albuginea testis.* The innermost coat of the testicle. It is a strong, white, and dense membrane, immediately covering the body or substance of the testicle. On its outer surface it is smooth, but rough and uneven on the inner.

ALBUGINOS HUMOUR. The aqueous humour of the eye.

ALBAEO OCULORUM. A white opacity of the cornea of the eyes. The Greeks named it *leucoma*; the Latins, *albugo*, *nebula* and *nubecula*; some ancient writers have called it *pterugium*, *janua oculi*, *onyx*, *unguis*, and

agides. It is a variety of Cullen's *Caligo Cornea*.

ALBUBAR. White lead.

ALBUM BALSANUM. The balsam of copaivi.

ALBUM GRÆCUM. The white dung of dogs. It was formerly applied as a discutient, to the inside of the throat, in quinseys, being first mixed with honey; medicines of this kind have long since justly sunk into disuse.

ALBUM OLUS. Lamb's lettuce, or corn-salad. The *Valeriana locusta* of Linnaeus.

ALBUMEN. *Albumena.* Albuminous matter. Coagulable lymph. Albumen is very abundant in the animal kingdom. It is the principal constituent part of the serum of the blood, and the lymphatic fluid. It forms the cheese in milk, and makes up the greater part of the white of eggs. It is composed of carbon, hydrogen, azot, oxygen, phosphorus, and somewhat of calcareous earth.

ALBUMEN OVI. *Alougo ovi.* *Albumen.* *alber ovi*, *ovi albus liquor*, *ovi candidum*, *albumentum*, *clavata.* The white of an egg.

ALCAHIST. An Arabic word to express an universal dissolvent, which was pretended to by Paracelsus and Helmont. Some say that Paracelsus first used this word, and that it is derived from the German words *al* and *geist*, i. e. *all spirit*. Van Helmont borrowed the word, and applied it to his invention, which he called the universal dissolvent.

ALCAOL. The solvent for the preparation of the philosopher's stone.

ALCALI. (Arab.) See *Alkali*.

ALCALIZATION. The impregnating any spiritous fluid with an alkali.

ALCRA INDICA. See *Hibiscus abelmoschus*.

ALCRA ÆSTHETIACA VILLOSA. See *Hibiscus abelmoschus*.

ALCRA ROSA. The systematic name for the malva arborea. See *Malva*.

ALCEA. (From *αλκω*, strength.) The name of a genus of plants in the Linnæan system. Class, *Monadelphæa*. Order *Polyandria*. Hollyhock.

ALCOAB. Sal almoniac, or muriat of ammoniac.

ALCANNA. (Indian word.) See *Alkanna*.

ALCEBAR. See *Agallochem*.

ALCEBRIS VIVUM. See *Sulphur vivum*.

ALCHABRIC. Sulphur vivum.

ALCHACHIL. Rosemary.

ALCHABITE. Quicksilver.

ALCHIKEN. This word occurs in the Theatrum Chemicum, and seems to signify that power in nature by which all corruption and generation are effected.

ALCHEMILLA. (So called because it was celebrated by the old alchemists.) *Ladies' mantle*.

1. The name of a genus of plants in the Linnæan system. Class *Tetrandria*. Order, *Monogynia*.

2. The parmacoporal name of a plant

called lady's mantle. *Alchemilla vulgaris*; *foliis lobatis* of Linnaeus. It was formerly esteemed as a powerful adstringent in hæmorrhages, fluor albus, &c. given internally.

ALCHIMILEC. (Heb.) The Egyptian meilot.

ALCHEMY. *Alchemia. Alchimia. Alkima.* That branch of chymistry which relates to the transmutation of metals into gold; the forming a panacea or universal remedy; an alkahest, or universal menstrum; a universal ferment; and many other absurdities.

ALCHIBRIC. See *Alkibric*.

ALCHIMILLA. See *Alchemilla*.

ALCHITRON. Oil of juniper; also the name of a dentifrice of Meusae.

ALCHUTE. See *Morum*.

ALCHYMY. Alchemy.

ALCHYLS. A speck on the pupil of the eye, somewhat obscuring vision.

ALCIMAD. Antimony.

ALCOS. Sal-ammoniac, or muriat of ammonia.

ALCOCALUM. (Perhaps Indian.) Artichoke, or cinara.

ALCOPOL. Antimony.

ALCOHOL. See *Alkohol*.

ALCOLA. (Heb.) The aphthæ, or thrush. Paracelsus gives this name to tartar, or excrement of urine, whether it appears as sand, mucilage, &c.

ALCOLITA. Urine.

ACLONE. Brass.

ALCOR. *Æs ustum*.

ALCTE. It is the name of a plant mentioned by Hippocrates. Fœsius thinks it is the elder.

ALCUBRITH. Sulphur.

ALCTONIUM. Bastard sponge, spongy plant-like substance, which is met with on the sea shore; it is of different shapes and colours. It is difficult to say what the Greeks called by this name. Dioscorides speaks of five sorts of it. They are calcined with a little salt, as dentifrice, and are used to remove spots on the skin.

Alder-tree. See *Alnus* and *Frangula*.

Alderberry bearing. See *Praugula*.

ALDER WINE. When well fermented, and having a proper addition of raisins in it, in its composition is frequently a rich and strong liquor; it keeps better than many of the other made wines, for a number of years, and was formerly supposed to possess many medical virtues; but these, experience does not seem to sanction: and the virtues of the alder, like those of many other simples formerly prized, have sunk into oblivion.

ALE. *Cerevisia. Liquor ceræis. Vinum hordeaceum.* Barley wine. A fermented liquor made from malt and hops, and chiefly distinguished from beer, made from the same ingredients, by the quantity of hops used therein; which is greater in beer, and therefore renders the liquor more bitter, and fitter for keeping. Ale, when well ferment-

ed, is a wholesome beverage, and seems only to disagree with those subject to asthma, or any disorder of the respiration or irregularity in the digestive organs. The old dispensaries enumerate several medicated ales, such as *cerevisia oxydrica*, for the eyes; *cerevisia antiarthritica*, against the gout; *cephalica, epileptica, &c.*

ALEARA. A cucurbit.

ALEBETHA. (From *alo*, to nourish.) Nourishing foods, or medicines.

ALEC. *Aléch.* Vitriol.

ALECHARITH. Mercury.

ALEIMA. (From *αλειν*, to anoint.) An ointment.

ALEION. (Ἄλειον, copious.) Hippocrates uses this word as an epithet for water.

ALEIPH. (From *αλειφ*, to anoint.) Any medicated oil.

ALEION. (From *αλε*, salt, and *ωαιον* oil.) Oil beat up with salt, to apply to tumours. Galen frequently used it.

ALEMA. (From *a* priv. and *λεμε*, hunger. Meat, food, or any thing that satisfies the appetite.

ALEMBO. (Some derive it from the Arabian particle *al*, and *ambic*, from *ambicare* to ascend. Avicenna declares it to be Arab.) Moorshead. A chymical utensil made of glass, metal, or earthen-ware, and adapted to receive volatile products from retorts. It consists of a body, to which is fitted a conical head, and out of this head descends laterally a beak to be inserted into the receiver.

ALEMBROTH. A chaldean word, importing the key of art. Some explained it by *sal mercurii*, or *sal philosophorum* and *artis*; others say it is named alembrot and *sal fusionis*, or *sal fixationis*. Alembroth desiccatum is said to be the *sal tartari*; hence this word seems to signify alkaline salt, which opens the bodies of metals by destroying their sulphurs, and promoting their separation from the ores. From analogy, it is supposed to have the same effect in conquering obstructions and attenuating viscid fluids in the human body. A peculiar earth, probably containing a fixed alkali, found in the island of Cyprus, has also this appellation; and a solution of the corrosive sublimate, to which the muriat of ammonia has been added, is called *sal alembroth*.

ALENZADAR. Crude sal ammoniac, or muriat of ammonia.

ALENZADAT. Crude sal ammoniac, or muriat of ammonia.

ALEPHNIS. A species of ash-tree which produces manna.

ALES. (From *αλε*, salt.) The name of a compound salt.

ALEUBON. (From *αλεω*, to grind.) Meal, **ALEXANDERS,** common. This plant, *Smyrniun olusatrum* of Linnaeus, was formerly cultivated for sallads. It is now superseded by celery.

ALEXANDERS, ROUND LEAVED. *Smyrniun perfoliatum* of Linnaeus. The blanched

stalks of this species are far preferable to those of common alexanders, and are esteemed as stomachic and nervine.

ALEXANDRIA. *Alexandrina.* The bay-tree, or laurel, of Alexandria.

ALEXANDRIUM. *Emplastrum viride.* A plaster described by Celsus, made with wax, alum, &c.

ALEXICACA. (From *αλεξω*, to drive away, and *κακον*, evil. *Alexicacum.* An antidote, or amulet, to resist poison.

ALEXIPHARMICA. (*Alexipharmica*, sc. *medicamenta*, fr. *ω αλεξω*, to expel, and *φαρμακον*, a poison.) *Antipharmica.* *Caco-alexiteria.* Medicines supposed to preserve the body against the power of poisons, or to correct or expel those taken. The ancients attributed this property to some vegetables, and even waters distilled from them. The term, however, is now disused.

ALEXIPYRETICUM. (From *αλεξω*, to drive away, and *πυρετρον*, fever.) A febrifuge. A remedy for fever.

ALEXIPYRETOS. *Alexipyretum.* The same as alexipyreticum.

ALEXIR. An elixir.

ALEXITERIA. Preservatives from contagion.

ALEXITERIUM. (From *αλεξω*, to expel, and *τηρεω*, to preserve.) A preservative medicine against poison, or contagion.

ALFACTA. Distillation.

ALFATIDE. Muriat of ammonia.

ALFASARA. *Alphasara.* Arabic terms for the vine.

ALFADAS. *Alfides.* Cernise.

ALPOL. Muriet of ammonia.

ALFUSA. Tutty.

ALGALI. A catheter. Also nitre.

ALGABAH. See *Anchilops*.

ALGAROTHE. (So called from Victorius Agaroth, a physician of Verone, and its inventor.) *Algarot.* *Algarothi.* *Mercurius rta.* *Pulvis Algarothi.* The antimonial part of the butter of antimony, separated from some of its acid by washing it in water. It is violently emetic in doses of two or three grains, and is preferred by many for making the emetic tartar.

ALGEDO. (From *αλγος*, pain.) A violent pain about the anus, perineum, testes, urethra, and bladder, arising from the sudden stoppage of a virulent gonorrhoea. A term very seldom used.

ALGEMA. (From *αλγω*, to be in pain.) *Algemedes.* *Algematodes.* Uneasiness, pain of any kind.

ALGERIA. *Algerie.* Lime.

ALGEROTH. See *Algaroth*.

ALGIBIC. Sulphur vitum.

ALGOR. A sudden chillness or rigor.

A term met with in Sauvage's and Sagar's Nosology.

ALGOSABEL. The Arabian-term for the *Daucus sylvestris*, or carrot.

ALGVADA. A white leprous eruption.

ALHAGI. (Arab.) A species of *Hedysarum*. The leaves are hot and pungent, the flowers purgative.

ALHANDALA. An Arabian name for colocyth, or bitter apple.

ALHASEF. (Arab.) *Alhasaf.* A sort of foetid pustule, called also *Hydroa*.

ALLA SQUILLA. (From *αλος*, belonging to the sea, and *σquilla*, a shrimp.) A prawn.

ALICA. (From *alo*, to nourish.) In general signification, a grain, a sort of food admired by the ancients; it is not certain whether it is a grain or a preparation of some kind thereof.

ALICES. (From *αλεξω*, to sprinkle.) Little red spots in the skin, which precede the eruption of pustules in the small-pox.

ALIENATIO MENTIS. (From *alieno*, to estrange.) *Delirium.* Estrangement of the mind.

ALIFORMES MUSCULI. Muscles so called from their supposed resemblance to wings. See *Pterugoideus*.

ALIMENTARY CANAL. Alimentary duct. A name given to the whole of those passages which the food passes through from the mouth to the anus. This duct may be said to be the true characteristic of an animal; there being no animal without it, and whatever has it, being properly ranged under the class of animals. Plants receive their nourishment by the numerous fibres of their roots, but have no common receptacle for digesting the food received, or for carrying off the excrements. But in all, even the lowest degree of animal life, we may observe a stomach and intestines, even where we cannot perceive the least formation of any organs of the senses, unless that common one of feeling, as in oysters.

ALIMENTARY DUCT. The alimentary canal. The thoracic duct is sometimes so called.

ALIMOS. Common liquorice.

ALIMUM. See *Arum*.

ALINDESIS. (*Ἀλινδῆσις*, from *αλινδνμι*, to be turned about.) A bodily exercise, which seems to be rolling on the ground, or rather in the dust, after being anointed with oil. Hippocrates says it hath nearly the same effect as wrestling.

ALIPENOS. (From *α*, neg. and *λιπνω*, to be fat.) *Alipenum.* *Alipantos.* An external remedy, without fat or moisture.

ALIPASMA. (From *αλειωω*, to anoint.) An ointment rubbed upon the body, to prevent sweating.

ALIPA. Remedies for wounds in the cheek, to prevent inflammation.

ALIPOW. A species of tureth, found near Mount Ceti, in Languedoc. It is a powerful purgative, used instead of senna, but much more active.

ALIPTRA. (From *αλειωω*, to anoint.) Servants who anointed the persons after bathing.

Alisanders. See *Smyrnum*.

ALISMA. (From *als*, the sea.) Water plantain. The name of a genus of plants in the Linnæan system. Class, *Hexandria*. Order, *Polygynia*.

ALISTELIS. (From *als*, the sea.) Muriat of ammonia.

ALITT. *Alith.* Asafetida.

ALKAPIAL. Antimony.

ALKAHEST. An imaginary universal menstruum, or solvent.

ALKAHEST GLAUBERI. Alkaline salts. See *Alcahest*.

ALKAHAT GLAUBERI. An alkali.

ALKALI. *Alkali*, in Arabic, signifies burnt; or from *al* and *kali*, i. e. the essence, or the whole of kali, the plant from which it was originally prepared, though now derived from plants of every kind.) *Alcali*, *alafi*, *alafar*, *alafort*, *calcadiis*. A term given to substances which possess the following properties: They are incombustible, and soluble in water; they possess an acrid, urinous taste. When mixed with siliceous substances, and exposed to an intense heat, they form a more or less perfect glass. They unite with another class of bodies called acids, and form new compounds, in which both the acid and alkaline properties are more or less lost. They render oils miscible with water. They change various blue vegetable pigments to green; red to violet, or blue, and yellow to brown. Blue pigments, that have been turned red with acids, are again restored by alkalies to their primitive colours. They emit light on the affusion of the dense acids when freed from water. They attract water and carbonic acid from the atmosphere. They unite to sulphur by fusion, and by means of water. They exert a great solvent power on the cellular membrane and animal fibre. They also corrode woollen cloth, and, if sufficiently concentrated, convert it into a sort of jelly.

They are only three kinds of alkalis at present known:

1. The mineral, called *soda*, in the new chymical nomenclature. See *Soda*.

2. The vegetable, called *potassa*, in the new chymical nomenclature. See *Potassa*.

3. *Ammonia*, or the caustic volatile alkali, is the third. See *Ammonia*.

To these, some chymists add barytes, and some other earths.

Alkalis are either *fixed*, that is, they are not reduced to the state of gas, by the most intense heat; or they are *volatile*, i. e. the common temperature is almost sufficient to change their state of aggregation; potash and soda are of the former kind; and, of the latter, ammonia is the only one known.

ALKALESCENT. Slightly alkaline.

ALKALI CAUSTIC. An alkali is so called when deprived of the carbonic acid it

contains for it then becomes more caustic and more violent in its action.

ALKALI FIXUM. Those alkalis are so called that emit no characteristic smell, and cannot be volatilized, but with the greatest difficulty. Two kinds of fixed alkalis have only hitherto been known, namely, potash and soda. See *Potash* and *Soda*.

Alkali, fossile. See *Soda*.

Alkali, mineral. (So called because it forms the basis of marine salts.) See *Soda*.

Alkali, vegetable. (So called because it abounds in many vegetables.) See *Potash*.

Alkali, volatile. (So called because it is volatile in opposition to the other alkalis, which are fixed.) See *Ammonia*.

ALKALINA. A class of substances described by Cullen as comprehending the substances otherwise termed *anacida*. They consist of alkalis, and also of substances, into which they enter in combination. The principal alkalines in use, are the carbonates and subcarbonates of soda, potash, and ammonia.

ALKALIZATION. (*Alcalizatio*, *onis*, f.) Alkalinization. The impregnating any thing with an alkaline salt, as spirit of wine, &c.

ALKAHAT. (*Alkanah*, a reed. Arab.) *Radix anchusæ*.

ALKANNA. See *Anchusa*.

ALKANNA VERA. *Alkauna Orientalis*. An Oriental plant; the *Lawsonia inermis*, *ramis inermibus*, of Linnæus; principally employed in its native place, as a dye. The root is the officinal part; which, however, is rarely met with in the shops. It possesses adstringent properties, and may be used as a substitute for the *anchusa*.

ALKASA. *Alkeal*. A crucible.

ALCANTHUM. Arsenic.

ALKANT. Quicksilver.

ALKEKENGI. (*Alkekengi*, Arab.) *Halicabaidro*. Winter cherry. This plant,

Physalis alkekengi of Linnæus:—foliis geminis integris acutis, caule herbaceo, inferne subramoso, is cultivated in our gardens. The berries are recommended as a diuretic, from six to twelve for a dose, in dropsical and calculous diseases.

ALKERMES. A term borrowed from the Arabs, denoting a celebrated remedy, of the form and consistence of a confection, whereof the kermes is the basis. See *Kermes*.

ALKERVA. (Arab.) Castor oil.

ALCOHOL. (An Arabian word, which signifies antimony: so called from the usage of the Eastern ladies to paint their eyebrows with antimony, reduced to a most subtle powder; which at last came to signify any thing exalted to its highest perfection.) *Alcohol*. *Alkol*. *Spiritus vinosus rectificatus*. *Spiritus vini rectificatus*. *Spiritus vini concentratus*. *Spiritus vini rectificatus*.

WINE. Alcohol is highly rectified spirit of wine, freed from all those aqueous particles which are not essential to it by duly performing rectification. In its purest state, it is quite colourless, and clear, of a strong and penetrating smell and taste; capable of being set on fire without a wick, and burning with a flame, without leaving a residue, and without smoke and soot. Alcohol is miscible with water in all proportions. It does not freeze in any degree of coldness. It is the direct menstruum or solvent of resins. It dissolves, also, the natural balsams. The resinous and various other parts of plants are also soluble in alcohol, hence it is made use of for extracting those parts, and for making the preparations called *elixirs, tinctures, essences*, &c. In England, alcohol is procured by distillation from molasses; in Scotland and Ireland, from an infusion of malt. This last, before its rectification, is termed *whiskey*. In the East-Indies, *arrack* is distilled from rice; in the West-Indies, *rum* from the sugar cane; and in France and Spain, *brandy* from wine; all these afford alcohol by distillation. On the human solids, alcohol acts as a most violent corrugator and stimulus.

ALKESOR. Camphire.

ALKI PLUMBI. Supposed to be acetat of lead.

ALKYMIA. Powder of basilisk.

ALLABOR. Lead.

ALLANTOIDES MEMBRANA. (*Allantoides*; from *alac*, a hog's pudding, and *oides*, likeness; because, in some brute animals, it is long and thick.) A membrane of the fœtus, peculiar to brutes, which contains the urine discharged from the bladder.

ALLERLUA. (Heb. *Praise the Lord*.) The acetosa, or wood-sorrel; so named from its many virtues. See *Acetocella*.

ALL-GOOD. *English mercury.* The vulgar name for the *Chenopodium bonus Henri-cus* of Linnæus; a plant which may be boiled for spinach, and which is in no degree inferior to it.

All-heal. See *Heracium* and *Stachys*.

ALLIARIA. (From *allium*, garlick; from its smell resembling garlick.) Jack of the hedge. Sauce-alone, or stinking hedge-mustard. The plant to which this name is given, in the Pharmacopœias, is the *Erysis mum alliaria*; *foliis cordatis* of Linnæus; it is sometimes exhibited in humid asthma and dyspnœa, with success. Its virtues are powerfully diaphoretic, diuretic, and antiscorbutic.

ALLICAR. Vinegar.

ALLICOA. Petroleum.

ALLIGATURA. A ligature, or bandage.

ALLIOTICUM. (From *allos*, to alter, or vary.) An alterative medicine, consisting of various antiscorbutics. *Galen*.

ALLIUM. (From *oleo*, to smell, because it stinks; or from *allos*, to avoid, as being unpleasant to most people.) Garlick.

1. The name of a genus of plants in the Linnæan system. Class, *Hexandria*. Order, *Monogynia*. Garlick.

2. The Pharmacopœial name of garlick. *Theriac rusticorum.* *Allium sativum* of Linnæus:—*caule planifolio bulbifero, bulbo composito, staminibus tricuspidatis*. This species of garlick, according to Linnæus, grows spontaneously in Sicily; but, as it is much employed for culinary and medicinal purposes, it has been long very generally cultivated in gardens. Every part of the plant, but more especially the root, has a pungent acrimonious taste, and a peculiarly offensive strong smell. This odour is extremely penetrating and diffusive; for, on the root being taken into the stomach, the alliaceous scent impregnates the whole system, and is discoverable in the various excretions, as in the urine, perspiration, milk, &c. Garlick is generally allied to the onion, from which it seems only to differ in being more powerful in its effects, and in its active matter, being in a more fixed state. By stimulating the stomach they both favour digestion, and, as a stimulus, are readily diffused over the system. They may, therefore, be considered as useful condiments with the food of phlegmatic people, or those whose circulation is languid and secretions interrupted; but with those subject to inflammatory complaints, or where great irritability prevails, these roots, in their acrid state, may prove very hurtful. The medicinal uses of garlick are various; it has been long in estimation as an expectorant in petuitous asthmas, and other pulmonary affections, *unattended* with inflammation. In hot bilious constitutions, therefore, garlick is improper: for it frequently produces flatulence, head-ach, thirst, heat, and other inflammatory symptoms. A free use of it is said to promote the piles in habits disposed to this complaint. Its utility as a diuretic in dropsies is attested by unquestionable authorities; and its febrifuge power has not only been experienced in preventing the paroxysms of intermittents, (Bergius says quartans have been cured by it; and he begins by giving one bulb, or clove, morning and evening, adding every day one more, till four or five cloves be taken at a dose: if the fever then vanishes, the dose is to be diminished, and it will be sufficient to take one, or even two cloves, twice a day, for some weeks;) but even in subduing the plague. Another virtue of garlick is that of an anthelmintic. It has likewise been found of great advantage in scorbutic cases, and in calculous disorders, acting in these, not only as a diuretic, but, in several instances, manifesting a lithontriptic power. That the juice of alliaceous plants in general, has considerable effects upon human calculi, is to be inferred by the experiments of Lobb; and we are abundantly warranted in asserting, that a decoction of the beads of

leeks, taken liberally, and its use persevered in for a length of time, has been found remarkably successful in calculous and gravelly complaints. The penetrating and diffusive acrimony of garlick, renders its external application useful in many disorders, as a rubefacient, and more especially as applied to the soles of the feet, to cause a revulsion from the head or breast, as was successfully practised and recommended by Sydenham. As soon as an inflammation appears, the garlick cataplasm should be removed, and one of bread and milk be applied, to obviate excessive pain. Garlick has also been variously employed externally, to tumours and cutaneous diseases; and, in certain cases of deafness, a clove, or small bulb of this root, wrapt in gauze or muslin, and introduced into the meatus auditorius, has been found an efficacious remedy. Garlick may be administered in different forms; swallowing the clove entire, after being dipped in oil, is recommended as the most effectual, or, where this cannot be done, by cutting it into pieces without bruising it, may be found to answer equally well, producing thereby no uneasiness in the fauces. On being beaten up, and formed into pills, the active parts of this medicine soon evaporate; this Dr. Woodville, in his Medical Botany, notices, on the authority of Culsen, who thinks that Lewis has fallen into a gross error, in supposing dried garlick more active than fresh. The syrup and oxymel of garlick, which formerly had a place in the British Pharmacopœias, are now expunged. It may be necessary to notice that, by some, the cloves of garlick are bruised and applied to the wrists, to cure agues, and to the bend of the arm, to cure the tooth-ache: when held in the hand, they are said to relieve hiccough; when beat with common oil into a poultice, they resolve sluggish humours; and, if laid on the navels of children, they are supposed to destroy worms in the intestines.

ALLIUM CÉPA. The systematic name for the *Cépa* of the shops. See *Cépa*.

ALLIUM FORBUM. The systematic name for the *Potrum* of the pharmacopœias. See *Potrum*.

ALLIUM SATIVUM. The systematic name for *Allium*. See *Allium*.

ALLIUM VICTORIALIS. The systematic name for the *Victoralis longa* of the pharmacopœias. See *Victoralis longa*.

ALLOCHOS. (From *αλλος*, another, and *αγο*, to speak.) Hippocrates uses this word to mean delirious.

ALLOESIS. (From *αλλος*, another.) Alteration in the state of a disease.

ALLOETICA. (From *αλλος*, another.) Alteratives. Medicines which change the appearance of the disease.

ALLOSINOSIS. (From *αλλος*, another, and *γινω*, to know.) Delirium; perversion of the judgment; incapability of distinguishing persons.

ALLOPHANTIS. (From *αλλος*, another, and *φασ*, to speak.) According to Hippocrates, a delirium, where the patient is not able to distinguish one thing from another.

ALLOTRIOPHAGIA. (From *αλλοτριος*, foreign, and *φας*, to eat.) A synonym of *pica*. See *Pica*. In Vogel's Nosology it signifies the greedily eating unusual things for food.

ALLOYS. By this word, chymists and artificers commonly understand any portion of base metal, or metallic mixture, which is added to combine metals by fusion into one seemingly homogeneous mass.

Allspice. See *Pimenta*.

ALMA. Water; and the first motion of a fetus to free itself from its confinement.

ALNABRI. A stone-like amber.

ALNAGRA. *Bolus cuprum*.

1. Red earth, or ochre, used by the ancients as an astringent.

2. Rulandus says it is the same as *Isio*.

3. In the Theatrum Chymicum it is a name for the white sulphur of the alchymists.

ALMARANDA. *Almakis*. Litharge.

ALMANDA CATHARTICA. A plant growing on the shores of Cayenne and Surinam, used by the inhabitants as a remedy for the colic; supposed to be cathartic.

ALMARCAN. (Arab.) Litharge of silver.

ALMACARIDA. Litharge of silver.

ALMARGEN. *Almaraga*. Coral.

ALMARKASITA. Mercury.

ALMARTAE. Powder of litharge.

AMATATICA. Copper.

ALMECASITE. *Almechasite*. Copper.

ALMBAILETU. A word used by Avicenna, to express a preternatural heat less than that of fever, and which may continue after a fever.

ALMENE. Sal lucidum, or sal gemmæ.

ALMIRA. Musk.

ALMISADIN. Verdigris, or muriat of ammonia.

ALMIZADAR. Muriat of Ammonia.

Almond, bitter. See *Amygdala*.

Almond, common. See *Amygdala*.

Almond, sweet. See *Amygdala*.

ALMONDS OF THE EARS. A popular name for the tonsils, which have been so called from their resemblance to an almond in shape. See *Tonsils*.

ALMONDS OF THE THROAT. A vulgar name for the tonsils.

ALNABATI. In Avicenna and Scorpion, this word means the *siliquadulcis*, a gentle laxative.

ALNEO. Stannum, or tin.

ALNERIO. Sulphur vivum.

ALNUS. (*Aino*, Ital.) The alder.

The pharmacopœial name of two plants, sometimes used in medicine, though rarely employed in the present practice.

Alnus rotundifolia; *ghumosa*; *viridis*; the common alder-tree, called *amendanus*.

Alnus nigra, vel *frangula*; the *rhamnus*

frangula of Linnaeus. The black alder, called also *alnus*,

All the parts of these trees are astringent and bitter. The bark is most astringent; a decoction of it has cured agues, and is often used to repel inflammatory tumours of the throat, by way of gargle. The inner yellow bark of the trunk, or root, given to ʒij., vomits, purges, and gripes; but, joined with aromatics, it operates more agreeably. An infusion or decoction in water inspissated to an extract, act yet more mildly than these. The berries of alder are purgative. They are not in use under their own name, but are often substituted for buck-thorn berries; to discover which, it should be observed, that the berries of the black alder have a black skin, a blue juice, and two seeds in each of them; whereas the buckthorn berries have a green juice, and commonly four seeds. The substitution of one for the other is not of material consequence, as the plants belong to the same genus, and the berries do not differ greatly.

Dr. Murray, of Göttingen, recommends from his own experience, the leaves of alder chopped in small pieces, and heated over the fire, as the best remedy with which he is acquainted for dispersing milk in the breasts.

Aloe. See *Aloes*.

ALOBABIA. (From *aloe*, the aloe.) Compound purging medicines so called from having aloes as the chief ingredient.

ALOPHARGINA. Medicines formed by a combination of aloes and aromatics.

ALOES. (From *alah*, a Hebrew word, signifying growing near the sea.) The Aloe.

1. A genus of plants of the Linnaean system. Class, *Hexandria*. Order, *Monogynia*.

2. The pharmacopoeial name of the inspissated juice of some of the aloe plants. *Fel naturæ*, nature's gall; so named from its intense bitterness. Aloes are distinguished into three species, *socotorine*, *hepatic*, and *caballine*; of which the two first are directed for official use in our pharmacopœia.

The 1st. *Aloes Succotorina* vel *Socotorina*. Succotorine aloes, is obtained from the *Aloe perfoliata* of Linnaeus:—*foliis caulibus dentatis, amplexicaulis vaginantibus, floribus corymbosis, cernuis, pedunculatis, subcylindricis*: it is brought over wrapt in skins, and is of a bright surface and in some degree pellucid; in the lump, of a yellowish red colour, with a purplish cast; when reduced into powder, it is of a golden colour. It is hard and friable in very cold weather; but in summer it softens very easily betwixt the fingers. It is extremely bitter and also accompanied with an aromatic flavour, but not so much as to cover its disagreeable taste. Its scent, though bitter, is rather agreeable, being somewhat similar to that of myrrh.

2. *Aloes hepatica*, vel *Barbadosensis*: the common or Barbadoes or hepatic aloes. The *perfoliata* of Linnaeus:—*floribus pedun-*

culatis, cernuis corymbosis, subcylindricis, foliis spinosis, confertis, dentatis, vaginantibus, planis, maculatis. The best is brought from Barbadoes in large gourd shells; an inferior sort in pots, and the worst in casks. It is darker coloured than the socotorine, and not so bright; it is also drier and more compact, though sometimes the sort in casks is soft and clammy. To the taste it is intensely bitter and nauseous, being almost wholly without that aroma which is observed in the socotorine. To the smell it is strong and disagreeable.

3. *Aloes caballina* vel *Guineensis*; Horse-aloes. This is easily distinguished from both the foregoing by its strong rank smell; in other respects it agrees pretty much with the hepatic, and is now not unfrequently sold in its place. Sometimes it is prepared so pure and bright as scarcely to be distinguishable by the eye, even from the socotorine, but its offensive smell betrays it; and if this also should be dissipated by art, its wanting the aromatic flavour of the finer aloes will be a sufficient criterion. This aloes is not admitted into the materia medica, and is not employed chiefly by farriers.

The general nature of these three kinds is nearly the same. Their particular differences only consist in the different proportions of gum to their resin, and in their flavour. The smell and taste reside principally in the gum, as do the principal virtues of the aloes. Twelve ounces of Barbadoes aloes yields nearly 4 ounces of resin, and 8 of gummy extract. The same quantity of socotorine aloes yields 3 ounces of resin and 9 of gummy extract.

Aloes is a well known stimulating purgative, a property which it possesses not only when taken internally, but also by external application. The cathartic quality of aloes does not reside in the resinous part of the drug, but in the gum, for the pure resin has little or no purgative power. Its medium dose is from 5 to 15 grains, nor does a larger quantity operate more effectually. Its operation is exerted on the large intestines, principally on the rectum. In small doses long continued, it often produces much heat, and irritation, particularly about the anus, from which it sometimes occasions a bloody discharge; therefore, to those who are subject to piles, or of an hæmorrhagic diathesis, or even in a state of pregnancy, its exhibition has been productive of considerable mischief; but on the contrary, by those of a phlegmatic constitution, or those suffering by uterine obstructions (for the stimulant action of aloes, it has been supposed, may be extended to the uterus,) and in some cases of dyspepsia, palsy, gout, and worms, aloes may be employed as a laxative with peculiar advantage. In all diseases of the bilious tribe, aloes is the strongest purge, and the best preparations for this purpose are the

pilula ex aloe cum myrrha, the *tinctura aloes*, or *extractum colocynthidis compositum*. Its efficacy in jaundice is very considerable, as it proves a succedaneum to the bile, which in that disease is defective either in quantity or quality. Aloes therefore may be considered as injurious where inflammation or irritation exist in the bowels or neighbouring parts, in pregnancy, or in habits disposed to piles, but highly serviceable in all hypochondriac affections, cachectic habits, and persons labouring under oppression of the stomach caused by irregularity: Aromatics correct the offensive qualities of aloes the most perfectly. The canella alba answers tolerably, and without any inconvenience; but some rather prefer the essential oils for this purpose. Dr. Cullen says, "If any medicine be entitled to the appellation of a *stomach purge*, it is certainly aloes. It is remarkable with regard to it, that it operates almost to as good a purpose in a small as in a large dose; that 5 grains will produce one considerable dejection, and 20 grains will do no more, except it be that in the last dose the operation will be attended with gripes, &c. Its chief use is to render the peristaltic motion regular, and it is one of the best cures in habitual costiveness. There is a difficulty we meet with in the exhibition of purgatives viz. that they will not act but in their full dose, and will not produce half their effect if given in half the dose. For this purpose we are chiefly confined to aloes. Neutral salts in half their dose will not have half their effect; although even from these, by large dilution, we may obtain this property; but besides them and our present medicine, I know no other which has any title to it except sulphur. Aloes sometimes cannot be employed. It has the effect of stimulating the rectum more than other purges, and with justice has been accused of exciting hæmorrhoidal swellings, so that we ought to abstain from it in such cases, except when we want to promote them. Aloes has the effect of rarefying the blood and disposing to hæmorrhagy, and hence it is not recommended in uterine fluxes. Fætid gums are of the same nature in producing hæmorrhagy, and perhaps this is the foundation of their emmenagogue power." Aloes is administered either simply in powders, which is too nauseous, or else in composition:—1. With purgatives, as soap, scammony, colocynth, or rhubarb. 2. With aromatics, as canella, ginger, or essential oils. 3. With bitters, as gentian. 4. With emmenagogues, as iron, myrrh, wine, &c. It may be exhibited in pills as the most convenient form, or else dissolved in wine, or diluted alcohol. The official preparations of aloes are the following—*Pilula Aloes*. *Pilula Aloes cum Assafœtidâ*. *Pil. Al. cum Colocynth.* *Pil. Al. cum Myrrh.* *T. Aloes.* *T. Al. Æth.* *T. Al. cum Myrrh.* *Vin. Aloes Soc.* *Pil.*

Aloes comp. *Pulv. Aloes cum Canell.* *Pulv. Al. cum Guaiac.* *Pulv. Al. cum Ferro.* *Tinct. Aloes comp.* *Ext. Colocynth. comp.* *Tinct. Benzoes. comp.* and some others.

ALLOTICS. Medicines wherein aloes is the chief or fundamental ingredient.

ALOGOTROPHIA. (From *αλογος*, disproportionate, and *τροφή*, to nourish.) Unequal nourishment, as in the rickets.

ALOHAR. (Arab.) *Alohec.* Mercury.

ALOES LIGNUM. See *Lignum Aloes*.

ALOMBA. (Arab.) *Alloc.* Plumbum, or lead.

ALOPECES. (From *αλωπεξ*, the fox.) The *psora* muscles are so called, by Fallopius and Vesalius, because in the fox they are particularly strong.

ALOPECIA. (from *αλωπεξ*, a fox; because the fox is subject to a distemper that resembles it: or, as some say, because the fox's urine will occasion baldness.) *Athrix depilis.* *Phalacroctis.* Baldness, or the falling off of the hair; when on the sinciput, *calvitium, calvitium*.

ALOSA. (From *αλωσα*, to take; because it is a ravenous fish.) The chad, *Clupea alosa* of Linnæus, whose flesh is by some commended as a restorative.

ALOSAT. *Alosohoc.* Quicksilver.

ALOSANTHO. (From *αλς*, salt, and *ανθος*, a flower.) *Alosanthum.* Flowers of salt.

ALPHARETICUM CHYMICUM. Raymond Lully hath given the world this alphabet, but to what end is difficult to say:

A significat Deum.

B ——— Mercurium

C ——— Salis Petram.

D ——— Vitriolum.

E ——— Menstruale.

F ——— Lunam claram.

G ——— Mercurium nostrum.

H ——— Salem purum.

I ——— Compositum Lune.

K ——— Compositum Solis.

L ——— Terram compositi Lune.

M ——— Aquam compositi Lune.

N ——— Ærem compositi Lune.

O ——— Terram compositi Solis.

P ——— Aquam compositi Solis.

Q ——— Ærem compositi Solis.

R ——— Ignem compositi Solis.

S ——— Lapidem Album

T ——— Medicinam corporis rubel.

U ——— Calorem fumi secreti.

X ——— Ignem siccum cineris.

Y ——— Calorem balnei.

Z ——— Separationem liquorum.

Z ——— Alembicum cum cucurbita.

ALPHANTIC. *Alphemic.* An Arabian word (signifying tender) for barley-sugar, or sugar-candy.

ALPHITA. (*Alplata*, the plural of *αλφον*, the meal of barley in general.) By Hippocrates this term is applied to barley-meal either toasted or fried. Galen says that *αριμα* is coarse meal, *αλφον* is fine meal, and *αλφοντα* is a middling sort.

ALPERTIDON. *Alphitidum*. It is when a bone is broken into small fragments like *Alphita*, i. e. bran.

ALPHONSIUM. The name of an instrument for extracting balls. It is so called from the name of its inventor Alphonso Ferrier, a Neapolitan physician. It consists of three branches, which separate from each other by their elasticity, but are capable of being closed by means of a tube in which they are included.

ALPHUS. (αλφός, from αλφαινω, to change; because it changes the colour of the skin.) *Vitiligo alba*. *Morphea alba*. *Lepra maculosa alba*. A species of leprosy, called by the ancients *vitiligo*, and which they divided into *alplus*, *melas*, and *leuce*. It is produced by a peculiar miasma, which is endemic to Arabia. See *Lepra*.

ALPINI BALSAMUM. Balm of Gilead.

ALRACHAS. Lead.

ALRATICA. A word used by Albucasis, to signify a partial or a total imperforation of the vagina. It is an Arabic word.

ALSAMACH. An Arabic name for the great hole in the os petrosus.

ALSINE. (From αλλος, a grove; so called because it grows in great abundance in woods and shady places.) The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Trigynia*. Chickweed.

ALSINE MEDIA. *Morus galline cunctunculus*. The name for the plant, called chickweed, which, if boiled tender, may be eaten like spinach, and forms also an excellent emollient poultice.

ALTAFOR. Camphire.

ALTERATIVES. (*Alterantia*, sc. *medicamenta*: from *altero*, to change.) Those remedies are so called, which are given with a view to re-establish the healthy functions of the animal æconomy, without producing any sensible evacuation.

ALTHÆA. (From αλθεω, to heal; so called from its supposed qualities in healing.) Marsh-mallow.

1. The name of a genus of plants of the Linnæan system. Class, *Monadelphia*. Order, *Polyandria*.

2. The pharmacopœial name of the marsh-mallow. *Althea officinalis* of Linneus:—*foliis simplicibus tomentosis*. Common marsh-mallow.

The mucilaginous matter with which this plant abounds, is the medicinal part of the plant: it is commonly employed for its emollient and demulcent qualities in tickling coughs, hoarseness, and catarrhs, in dysentery, and difficulty and heat of urine. It relaxes the passages in nephritic complaints, in which last case a decoction is the best preparation. Two or three ounces of the fresh roots may be boiled in a sufficient quantity of water to a quart, to which one ounce of gum-arabic may be added. The following is given where it is required that

large quantities should be used. An ounce of the dried roots is to be boiled in water enough to leave two or three pints to be poured off for use: if more of the root be used, the liquor will be disagreeably slimy. If sweetened, by adding a little more of the root of liquorice, it will be very palatable. The root had formerly a place in many of the compounds in the pharmacopœias, but now it is only directed in the form of syrup.

ALTHÆA OFFICINALIS. The systematic name of the marsh-mallow. See *Althea*.

ALTHANACA. *Althanacha*. Orpiment.

ALTHEBGEIUM. An Arabian name for a sort of swelling, such as is observed in cachectic and leuco-phlegmatic habits.

ALTHEXIS. (From αλθεω, to cure, or heal.) Hippocrates often uses this word to signify the cure of a distemper.

ALTIMIT. So Avicenna calls the *Laserpitium* of the ancients.

ALUD. (Arab.) Aloes.

ALUNDULS. Hollow spheres of stone, glass, or earthen-ware, with a short neck projecting at each end, by means of which one globe might be set upon the other. The uppermost has no opening at the top. They were used in former times for the sublimation of several substances.

Alum. See *Alumen*.

ALUMEN. (*Alum*. Arab.) *Assos*, *azub*, *aseb*, *elanula*. *Sulphas aluminæ acidulus*, *cum potassâ*. *Super-sulphas aluminæ et potassæ*. *Argilla virioliata*. Alum.

A neutral salt, formed by the combination of the earth called *alumine*, or pure clay, with sulphuric acid, and a little potash.

The alum of commerce, and that presented for medicinal purposes, is afforded by ores which are dug out of the earth for this purpose, and manufactured by first decomposing the ore, then lixiviating it, evaporating the lixiviums, and then crystallizing the alum, which affects the form of tetrahedral pyramids, applied to each other base to base; sometimes the angles are truncated.

The following kinds of alum are met with in the shops:

1. Ice or rock alum. *Alumen commune: alumen crystallinum, rupeum, factitium*. Common alum; fictitious alum: English alum. This is always in very large transparent masses, and derives its name from Rocca in Syria, now called Edessa, in which the earliest manufactory of this salt was established; or from the hardness and size the masses. This species is not very pure.

2. Roman alum. *Alumen Romanum: alumen rubrum rutium, rochi Galli*. Called rock alum by the French. This species, which is prepared in the territory of Civita-Vecchia, comes in lumps of the size of eggs, covered with a reddish efflorescence.

Alum, when first tasted, imparts a sweetness, but is soon felt to be strongly astringent; on account of which virtue it is of

very extensive use in medicine and surgery.

Internally it is used as a powerful astringent in cases of passive hæmorrhages from the womb, intestines, nose, and sometimes lungs. In bleedings of an active nature, i. e. attended with fever, and a plethoric state of the system, it is highly improper. Dr. Percival recommends it in the *colica pictorum* and other chronic disorders of the bowels, attended with obstinate constipation. See Percival's *Essays*. The dose advised in these cases, is from 5 to 20 grains, to be repeated every four, eight, or twelve hours. When duly persisted in, this remedy proves gently laxative, and mitigates the pain.

Alum is also powerfully tonic, and is given with this view in 10 grains of alum made into a bolus three times a day, in such cases as require powerful tonic and astringent remedies. Another mode of administering it, is in the form of whey made by boiling a drachm of powdered alum in a pint of milk, for a few minutes, and to be taken in the quantity of a tea-cup full three times a day. Dr. Cullen thinks it ought to be employed with other astringents in diarrhœas. In active hæmorrhages, as was observed, it is not useful, though a powerful medicine in those which are passive. It should be given in small doses, and gradually increased. It has been tried in the diabetes without success; though, joined with nutmeg, it has been more successful in intermittents given in a large dose, an hour or a little longer, before the approach of the paroxysm. In gargles, in relaxation of the uvula, and other swellings of the mucous membrane of the fauces, divested of acute inflammation, it has been used with advantage; also in every state of the cynanche tonsillaris. Externally alum is much employed by surgeons as a lotion for the eyes, and is said to be preferable to white vitriol, or acetated ceruse in the ophthalmia membranarum. From two to five grains dissolved in an ounce of rose water, forms a proper collyrium. It is also applied as a styptic to bleeding vessels, and to ulcers, where there is too copious a secretion of pus. It has proved successful in inflammation of the eyes, in the form of cataplasm, which is made by stirring or shaking a lump of alum in the whites of two eggs, till they form a coagulum, which is applied to the eye between two pieces of thin linen rag. This substance is also employed in the form of injection in cases of gleet or fluor albus.

When deprived of its humidity by exposure to the fire, by placing it in an earthen pan over a gentle fire, till it ceases to bubble, it is termed burnt alum, *alumen ustum*, and is sometimes employed by surgeons to destroy fungous flesh, and is a principle ingredient in most styptic powders. Alum is also applied to many purposes of life; in

this country, bakers mix a quantity with the bread, to render it white; this mixture makes the bread better adapted for weak and relaxed bowels; but in opposite states of the alimentary canal, this practice is highly pernicious. The official preparations of alum are: *Alumen purificatum*. Lond. *Sulphas aluminis exsiccata*. *Soluti sulphat. cupri ammon.* Eding. *Aq. alum. comp.* Lond.

ALUMEN CATTINUM. A name of potash.

ALUMEN USTRUM. See *Alumen*.

ALUMINE. *Alumina.* Earth of alum. Pure clay.

Aluminous earth derives its name from alum, of which it forms the base. It constitutes the lower strata of mountains and plains. It arrests the waters, and causes them to rise in springs to the surface of the earth. It enters into the natural composition of the schistus, and all these stones and earths called argillaceous, such as *potter's-clay*, *fuller's-earth*, *lapidolite*, *mica*, *corundum*, &c. Hitherto it has not been found pure any where, except in the garden of the public schools at Halle in Germany.

Properties of pure Alumina.—Alumine is white, and soft to the touch. It is insipid, adheres to the tongue, and occasions a sense of dryness in the mouth. When moistened with a small quantity of water, it forms a tenacious, ductile, kneadable paste. When heated to redness, it shrinks considerably in bulk, and at last becomes so hard as to strike fire with flint. After being ignited, it is no longer capable of being kneaded with water into a ductile mass. It recovers however this property by solution in an acid and precipitation. Alkalies dissolve it in the humid way, and form compounds decomposable by acids. It dissolves slowly in all acids. It possesses a powerful attraction for lime. The most intense heat of our furnaces is not able to melt it, but it becomes fusible when lime is added. Lavoisier has proved that it is capable of entering into a kind of fusion like paste, by the action of oxygen gas; it then cuts glass and resists the file. It absorbs water and carbonic acid from the atmosphere. By its mixture with water and silex it acquires great solidity. It does not unite with any combustible substance, but it becomes fused into coloured frits with metallic oxyda. Its specific gravity is 2. It is employed in a multitude of arts.

Method of obtaining pure alumine.—Take any quantity of alum of commerce, dissolve it in six parts of boiling distilled water, and add to this solution, when cold, liquid ammonia, till no further precipitate ensues. Then heat the whole nearly to the boiling point for a few minutes, and transfer it on a filter. In proportion as the fluid passes off, pour more water over the precipitate, until it passes tasteless. Let the precipitate obtained, while yet in a pasty state,

be transferred into a glass or Wedgwood's bason, and add to it muriatic acid in small quantities at a time, until the whole is dissolved. Then evaporate the solution, till a drop of it, when suffered to cool on a plate of glass, yields minute crystals: on letting it now cool, crystals of alum will be deposited. Remove these crystals by decanting the fluid, and renew the evaporation, until, on further cooling, no more crystals are formed. Nothing now but pure alumine remains in the solution; the fluid may therefore be decomposed by adding to it gradually liquid ammonia till no further precipitate ensues. The precipitate thus obtained, when well washed and dried, is pure alumine.

The process recommended in general by systematic writers for obtaining alumine, differs from this; it consists in decomposing a solution of alum of commerce by an excess of a carbonated alkali, washing the obtained precipitate, and exposing it to a sufficient heat to drive off the carbonic acid. This method however is imperfect, for if the alumine thus obtained be heated with charcoal, and a diluted acid is added to the mixture, sulphurated hydrogen gas will be liberated. It adheres to the tongue, and emits a peculiar odour when breathed upon. Sure signs that it is not pure.

It must be obvious that alumine cannot be obtained absolutely pure in this manner. For alum is a triple compound, consisting of alumine, potash, and sulphuric acid in excess. When this excess of acid is saturated, by adding to the solution an alkali, or even pure alumine, a highly insoluble salt (sulphate of alumine) is produced, differing from alum only in the proportion of its base. When we therefore gradually add to a solution of alum, a carbonated alkali, the first effect of the alkali is, to saturate the excess of the sulphuric acid; and the precipitate consists principally of the salt which is insoluble in water. A further quantity of the alkali effects instantly a decomposition of part of the salt, which, in proportion as it takes place, becomes mixed with the alumine: and it is thus covered from the further action of the alkali. This being the case, it is obvious that no subsequent washing can do more than separate the sulphate of potash, and therefore the residuum, instead of being pure alumine, contains also a variable proportion of true sulphate of alumine; the sulphuric acid of which becoming decomposed on heating it in contact with charcoal, accounts for the sulphurated hydrogen gas produced by the affusion of an acid. With the acids it is known to form more than twenty species of neutral salts. Of these only one is used in medicine and surgery, called *alum*, or *aluminous sulphate*. See *Alumen*.

ALUMINOUS WATERS. Waters impregnated with particles of alum.

ALUSAR. *Manna*.

ALVEARIUM. (From *alveare*, a beehive.) That part of the *meatus auditorius externus* is so called, which contains the wax of the ear.

ALVEOLI. (From *alveare*, a beehive; from their resemblance to its cells.) *Betrien*, *bothrien*; *frena*, *mortariolum*. The sockets of the teeth. There are usually sixteen of these alveoli, or sockets, in each jaw.

ALVEUS COMMUNIS. The common duct, or communication of the ampullæ of the membranaceous semicircular canals in the internal ear, is so termed by Scarpa.

ALVEUS AMPULNASCENS. Part of the duct conveying the chyle to the subclavian vein.

ALVIDUCA. (From *alvus*, the belly, and *duco*, to draw.) Purging medicines.

ALVIFLUXUS. (From *alvus*, and *fluo*, to flow.) A diarrhoea, or purging.

ALVUS. The belly, stomach and entrails.

ALYCE. (From *alvus*, to be anxious.) That anxiety which is attendant on low fevers.

ALYPIA. (From *α*, neg. and *λυω*, pain.) A gentle purgation of the humours without pain.

ALYTIAS. *Alypum*. A species of spurge, so called because it purges gently and without pain.

ALXANUS. (From *alvus*, to be restless.) Restlessness.

ALYSSUM. (From *α*, neg. and *λυω*, the bite of a mad dog: so called because it was foolishly thought to be a specific in the cure of the bite of a mad dog.) Mad-wort. The *Marrubium alysseon* of Linnæus, supposed by some to be diaphoretic.

ALYSSUM GALENI. The *marrubium*.

ALYSSUM PLINII. The *mollugo*.

ALYSSUM VERTICILLATUM. The *marrubium verticillatum*.

ALZEMAFOR. *Cinnabar*.

ALZUM. *Alidum*. *Alidrum*. The name of the tree which produces gum bdellium according to some ancient authors.

AMA. Together. A word used in composition.

AMALGAM. (From *αμα*, and *μαρτυρ*, to marry.) A substance produced by mixing mercury with a metal, the two being thereby incorporated.

AMANELIS. (From *αμα*, and *μαρμα*, an apple.) The hasted medlar of Hippocrates.

AMANITE. (From *α*, priv. and *μαρμα*, madness; so called, because they are eatable and not poisonous, like some others.) A tribe of fungus productions, called mushrooms, truffles, and morels, and by the French, *champignons*.

AMARA. (*Amara*, sc. *medicamenta*: from *amarus*, bitter.) Bitters.

The principal bitters used medicinally are: the pure bitters, *gentiana lutea*; *humulus lupulus*; and *quassia amara*: styptic bitters, *cinchona officinalis*; *crotan coarctilla*: *quassia*

amaroubia: and aromatic bitters, *artemesia abeynthium*; *anthemis nobilis*: *hyssopus*, &c.

AMARA DULCIS. See *Dulcemara*.

AMARACUS. (From *a*, neg. and *μαραω* to decay; because it keeps its virtues a long time.) Marjoram.

AMARANTHUS. (From the same.) The herb goldilocks.

AMARANTH ESCULENT. The leaves of the *amaranthus oleraceus* of Linnæus, and several other species, are eaten in India the same as cabbage is here.

AMARANTHUS OLERACEUS. See *Amaranth*, *esculent*.

AMATORIA FEBRIS. (From *amo*, to love.) See *Chlorosis*.

AMATORIA VENERICA. (From *amo*, to love, and *venerificium*, witchcraft.) Philters. Love-powders.

AMATORII. (*Amatorii*, sc. *musculi*.) A term given to the muscles of the eye, by which that organ is moved in ogling.

AMATEQUITL. (Indian.) *Unedo papyracea*. The *arbutus unedo* of Linnæus. A decoction of the bark of the root of this plant is commended in fevers.

AMAUROSIS. (*Αμαυρωσις*: from *αμαυρον*, to darken or obscure.) *Gutta serena*. *Amblyopia*. A total loss of sight without any visible injury to the eye, the pupil mostly dilated and immoveable. A genus of disease in the class *locales*, and order *dysæsthesiæ* of Cullen. It arises generally from compression of the optic nerves; *amaurosis compressionis*; from debility, *amaurosis atonica*; from spasm, *amaurosis spasmodica*; or from poisons, *amaurosis venenata*. See also *Gutta serena*.

Amber seed. See *Hibiscus abelmoschus*.

AMBE. (*Αμβη*, the edge of a rock; from *αμβαινω*, to ascend.) An old surgical machine for reducing dislocations of the shoulder, and so called, because its extremity projects like the prominence of a rock. Its invention is imputed to Hippocrates. The ambe is the most ancient mechanical contrivance for the above purpose, but is not at present employed.

AMELA. (Arab.) The cornered hazelnut, the bark of which is purgative.

AMBER. *Succinum*. A beautiful bituminous substance, of a yellow or brown colour, either transparent or opaque, which takes a good polish, and, after a slight rubbing, becomes so electric, as to attract straws and small bodies; hence it was called *electrum* by the ancients, and hence the word electricity. When powdered, it emits an agreeable smell. It is dug out of the earth at various depths, and often contains insects in high preservation; a circumstance which proves that it has been liquid. Amber is also found floating on the shores of the Baltic, and is met with in Italy, Sicily, Poland, Sweden, &c. From its colour or opacity it has been variously distinguished; thus white, orange, golden,

cloudy amber, &c. An oil is obtained from it, which, as well as its other preparations, is occasionally used in medicine against spasmodic diseases.

AMBERGRIS. (*Ambragrisæa*.) A concrete, bituminous substance, of a soft and tenacious consistence, marked with black and yellow spots, and of an agreeable and strong smell when heated or rubbed. It is found in very irregular masses, floating on the sea near the Molucca Islands, Madagascar, Sumatra, on the coast of Coromandel, Brazil, America, China, and Japan. Several American fishermen assured Dr. Schwediawer, that they often found this substance, either among the excrements of the *Physeter macrocephalus*, a species of whale, or in its stomach, or in a vessel near the stomach. The medical qualities of ambergris are stomachic, cordial, and antispasmodic. It is very seldom used in this country.

AMBLOSIA. (*Αμβλωσις*: from *αμβλω*, to cause abortion.) A miscarriage.

AMBLOTICA. (*Amblotica*, sc. *medicamentata*, *αμβλωτικα*: from *αμβλω*, to cause abortion.) Medicines which were supposed to occasion abortion.

AMBLYOTIA. (From *αμβλω*, dull, and *ωψ*, the eye.) Hippocrates means by this word, dimness, of sight to which old people are subject. Paulus Actuarius, and the best modern writers, seem to think that amblyopia means the same thing as the incomplete amaurosis. See *Gutta serena* and *Amaurosis*.

AMBLYOTISMUS. *Amblyotæ*. The same.

AMBO. (Indian.) The mango.

AMBON. (From *αμβανω*, to ascend.) Celsus uses this term to signify, the margin or tip of the sockets in which the heads of the large bones are lodged.

AMBONE. The same as ambe.

AMBRA. Amber. Also an aromatic gum.

AMBRA CINRACEA. (From *cineraceus*, of the colour of ashes.) Ambergris and grey amber.

AMBRA GRISCA. Ambergris.

AMBRAM. Amber.

AMBRETTE. See *Abelmoschus*.

AMBULATIVA. (From *ambulo*, to walk.) A species of herpes; so called because it walks or creeps as it were about the body.

AMBULO. (From *αμβελλω*, to cast forth.) *Fluxus furiosus*. A periodical flatulent disease, caused, according to Michalis, by vapours shooting through various parts of the body.

AMBUSTIO. (From *ambure*, to burn.) *Ambustiem*. A burn or scald.

AMELLA. The same as achmella.

AMENORRHEA. (From *a*, priv. and *ρρ*, a month, and *ρρ*, to flow.) A partial or total obstruction of the menses in women from other causes than pregnancy and old age. That this excrementitious discharge should be regular as to quantity and quality, and that it should observe the monthly period, is essential to health. When it is obstructed,

nature makes her efforts to obtain for it some other outlet. When these efforts of nature fail, the consequence may be pyrexia, pulmonic diseases, spasmodic affections, hysteria, epilepsy, mania, apoplexia, chlorosis, according to the general habit and disposition of the patient. Dr. Cullen places this genus in the class *locules*, and order *epischeses*. His species are, 1. *Emanatio mensium*: that is, when the menses do not appear so early as is usually expected. See *Chlorosis*. 2. *Suppressio mensium*, when, after the menses appearing and continuing as usual for some time, they cease without pregnancy, occurring. 3. *Amenorrhœa difficilis, vel Menorrhagia difficilis*, when the flux is too small in quantity, and attended with great pain, &c.

AMENTIA. (From *a*, priv. and *mens*, the mind.) Imbecility of intellect, by which the relations of things are either not perceived, or not recollected. A disease in the class *neuroses*, and order *semie* of Cullen. When it originates at birth, it is called *amentia congenita*, natural stupidity; when from the infirmities of age, *amentia senilis*, dotage or childishness; and when from some accidental cause, *amentia acquisita*.

American Balm. See *Balsamum Peruvianum*.

AMERICANUM TUBEROSUM. The potato. An America tuberosa root.

AMETHYSTA PHARMACA. (From *a*, neg. and *μῆδος*, wine.) Medicines which were said either to prevent or remove the effects of wine. *Galen*.

AMETHYSTUS. (From *a*, neg. and *μῆθος*, to be inebriated.) The amethyst. A precious stone, so called, because in former times, according to Plutarch, it was thought to prevent drunkenness. *Ruland in Lex.Chem.*

AMICULUM. A little short cloak. It is the same as the amnios, but anciently meant a covering for the pubes of boys, when they exercised in the gymnasium. *Rhodius*.

AMMI. (Αμμή from *αμμος*, sand; from its likeness to little gravel-stones.) The herb bishop's-weed, of which there are two sorts, the *ammi verum* and *vulgare*.

AMMI MAJUS. The systematic name for the *ammi vulgare* of the shops. See *Ammi vulgare*.

AMMI VERUM. The seeds of this plant, *Sison ammi* of Linnæus;—*foliis tripinnatis, radicalibus linearibus, caulibus setaceis, stipularibus longioribus*, have a grateful smell, somewhat like that of origanum, and were formerly administered as a carminative.

AMMI VULGARE. The seeds of this plant, *Ammi majus*, of Linnæus;—*foliis inferioribus pinnatis, lanceolatis, serratis; superioribus multifidis, linearibus*, are less powerful than those of the *Sison ammi*, but were exhibited with the same views.

AMIDUM. See *Amylum*.

AMINEUM. A wine produced in Ami-

neza, formerly a province of Italy; called also Salernum. Also a strong wine vinegar. *Galen* mentions *Amineum Neapolitanum*, and *Amineum Siculum*.

AMMIUM. *Ammiu*. Cinnabar.

AMMOCHOSIA. (From *αμμος*, sand, and *χύνω*, to pour.) A remedy for drying the body by sprinkling it with hot sand. *Oribasius*.

AMMONIA ACETATA. See *Liquor Ammonie acetatis*.

AMMONIA MURIATA. See *Muriate of ammonia*.

AMMONIA PREPARATA. See *Carbonas ammonie*.

AMMONIA. Ammonia-gas. The substance so called, is an aeriform or gaseous body.

Pure ammonia was long supposed to be a compound of hydrogen and nitrogen, rendered gaseous by the addition of caloric; but from the experiments of Mr. Davy on the alkalis, it appears to be a metallic oxyd.

Ammonia-gas has a strong and very pungent odour. It extinguishes flame, yet it increases the magnitude of the flame of a taper before extinction, producing a pale yellow colour round its edge. Animals cannot breathe it without death ensuing. It is lighter than atmospheric air, in the proportion of three to five. It tinges yellow vegetable colours brown, and blue ones green. It is rapidly absorbed by cold water; by ardent spirit, essential oils, ether, charcoal, sponge, bits of linen cloth, and all porous bodies.

When a piece of ice is brought in contact with this gas, it melts and absorbs the gas, while at the same time its temperature is diminished. It has no effect upon oxygen gas while cold; but when made to pass with it through an ignited tube, it detonates and becomes decomposed. The same is the case with common air. It is also decomposed by phosphorus at high temperatures.

It does not explode when mixed with hydrogen gas. Nitrogen gas has no effect upon it. Atmospheric air does not combine with it at common temperatures, but only mixes with and dilutes it. When made to pass through ignited charcoal, it forms with it a substance called prussic acid. If brought into contact with acid gases, both gases lose their gaseous form, and become concrete. It has no sensible action on earths, or on the *salino-terrene* substances. It combines readily with acids, and unites to sulphur, when both are in a state of vapour. It reduces oxyds of metals to their metallic state, and is decomposed by them. It is also decomposed by electrization, and by oxygenated muriatic acid gas, &c.—When exposed to the temperature of 46 degrees, it crystallizes, and when suddenly cooled down to 68 degrees, it assumes a gelatinous appearance, and has scarcely any odour.

Methods of obtaining Ammonia.—1. Mix together equal quantities of muriate of ammonia and quick-lime, separately powdered; introduce them into a gas-bottle or retort, apply the heat of a lamp, and receive the gas over mercury.

Explanation.—Muriate of ammonia consists of muriatic acid and ammonia; on adding lime to it, a decomposition takes place, the muriatic acid quits the ammonia and unites to the lime, in order to form muriate of lime, which remains in the retort, and the ammonia flies off in the state of gas.

Remark.—In order to obtain the gas in a state of purity, it is essentially necessary that a considerable quantity of the gas first disengaged, be suffered to escape, on account of the common air contained in the distilling vessel, and in the interstices of the ingredients.

2. Ammonia may likewise be obtained by heating the liquid ammonia of the shops (liquor of pure ammonia, *Pharm. Lond.*) in a retort placed in communication with the mercurial pneumatic trough.

In this process the ammonia contained in this liquid combines with caloric, assumes the form of ammonia-gas, and parts with the water to which it was united.

Remark.—The temperature of the fluid must not be carried so high as to cause the water to be converted into vapour, or, if this cannot well be avoided, a small vessel should be interposed between the retort and the receiver, which, when kept cool, may serve to condense the aqueous vapour which is formed, and cause the ammonia-gas to pass in a very pure and dry state.

Ammonia is likewise produced during the spontaneous decomposition of animal and vegetable substances; in these cases it did not pre-exist in them ready formed, but is generated by the union of the hydrogen and nitrogen contained in them.

In combination with water, this alkali forms a solution, or liquid ammonia, which is called, in the London pharmacopœia,

LIQUOR AMMONIÆ.

Take of muriate of ammonia,

Lime newly prepared, of each two pounds.

Water, a pint and a half.

Reduce the muriate of ammonia and the lime into powder separately; then mix them, and introduce them into a large glass retort, into which a pint of water has been previously poured. Having placed the retort in a sand bath, lute on a tubulated receiver, through which the ammonia may pass on into a third vessel containing half a pint of the water, and cooled. Then at first apply a gentle heat, and increase it by degrees, until the retort becomes red.

Great care and attention are necessary in every part of this process; the two salts are to be powdered separately, before they are mixed; for, if they be triturated toge-

ther, ammonia will be extricated, which should be prevented, until the means for its collection are adopted. The salts are to be shaken well together, rather than rubbed, and added to the water in the retort. The cold produced by the solution of the salt, will counteract the heat produced by the slacking of the lime, and a charge so made will be manageable until the receiver is fitted on, and the heat of the sand-bath applied. This heat need not be greater than 300 degrees, and should be very cautiously and slowly raised, to prevent the rapid ebullition and expansion during the extrication from gas of a charge of such density; and for the same reason, a large retort is directed. The ammonia rises immediately in the form of gas, and a portion of the water is therefore placed in a situation to condense it; in the subsequent stages, water will arise from the charge in the retort. The third vessel directed in the formula, may be either a common bottle, fitting moderately (for no great pressure is necessary,) to a straight tube issuing from the bottom of the receiver, and dipping below the surface of the water it contains; or Woulfe's apparatus may be used; but, with moderate attention, the simpler means will answer better, perhaps, than the more complex. In either case, the receiving-bottle must be kept cold by wet cloths, or ice, for the lower the temperature of the water, the greater quantity of ammonia-gas it will condense, and the condensation is accompanied by an increase of its heat. If two bottles, each containing half the quantity of water directed, be used, they will be most manageable, as they may be changed alternately, so as to prevent either from being overheated, and the contents of both may be mixed together at last. This preparation is colourless and transparent, with a strong peculiar smell; it parts with the ammonia in the form of gas, if heated to 130 degrees, and requires to be kept, with a cautious exclusion of atmospherical air, with the carbonic acid of which it readily unites: on this latter account, the propriety of keeping it small bottles instead of a large one, has been suggested. Water saturated with ammonia gas, has a less specific gravity than common water.

This is the *aqua ammonia pure* of the shops, and the *alkali volatile causticum*.

The preparations of ammonia in use are,

1. The carbonate of ammonia. See *Carbonas ammoniæ*, and *Liquor carbonatis ammoniæ*.

2. The acetate of ammonia. See *Liquor acetatis ammoniæ*.

3. The muriate of ammonia. See *Muriate of Ammonia*.

4. Ferrum ammoniatum.

5. Several tinctures and spirits, holding carbonate of ammonia in solution.

AMMONIACUM. (*Ammoniacum*: so called

from *Ammonia*, whence it was brought.) *Gum-ammoniac*. A concrete gummy-resinous juice, composed of little lumps, or tears, of a strong and somewhat ungrateful smell, and nauseous taste, followed by a bitterness. There has, hitherto, been no information had concerning the plant which affords this drug. It is imported here from Turkey, and from the East-Indies. Gum-ammoniacum is principally employed as an expectorant, and is frequently prescribed in asthma and chronic catarrh. Its dose is from 10 to 30 grains. It is given in the form of a pill, or diffused in water, and is frequently combined with squill, or tartrite of antimony. In large doses, it proves purgative. Externally, it is applied as a discutient, under the form of plaster, to white swellings of the knee, and to indolent tumours. The official preparations are: *Ammoniacum purificatum*: *Emplastrum ex ammoniaco cum hydragyro*: *Mistura ammoniacæ*.

AMMONIÆ CARBONAS. See *Carbonas ammoniæ*.

AMMONIÆ MURIAS. See *Muriate of ammonia*.

AMMONION. (From *αμμος*, sand.) *Ætius* uses this term to denote a collyrium of great virtue in many diseases of the eye, which was said to remove sand or gravel from the eyes.

AMMONIUM. The metal which with oxygen forms the alkali called ammonia. See *Ammonia*.

AMNESIA. (From *α, priv.* and *μνησις*, memory.) *Amnesia*. Forgetfulness; mostly a symptomatic affection.

AMNESTIA. See *Amnesia*.

AMNIOS. *Amnion*. (From *αμνος*, a lamb, or lamb's skin.) The soft internal membrane which surrounds the fœtus. It is very thin and pellucid in the early stage of pregnancy, but acquires considerable thickness and strength in the latter months. The amnios contains a thin watery fluid, in which the fœtus is suspended. In the abortion of the early months, we find the quantity of this fluid very great, in proportion to the whole ovum, and the amnios forms a delicate and almost gelatinous substance, and is a provision for the regular presentation of the head of the child; for now, the fœtus being suspended in the fluid, and hanging by the umbilicus, and the head and upper part of the body greatly preponderating, it takes that position with the head presenting to the orifice of the womb, which is necessary to natural and safe labour, the fœtus being prevented from shifting, in the latter months, by the closer embracing of the child with the uterus.

AMNIOTIC ACID. *Vauquelin* and *Buniva* have discovered a peculiar acid in the liquor of the amnios of the cow, to which they have given the name of amniotic acid.

It exists in the form of a white pulverulent powder. It is slightly acid, but sensibly reddens vegetable blues. It is difficultly soluble in cold, but readily soluble in boiling water, and in alcohol. When exposed to a strong heat, it exhales an odour of ammonia and of prussic acid. Assisted by heat, it decomposes carbonate of potash, soda, and ammonia. It produces no change in the solutions of silver, lead, or mercury, in nitric acid. Exposed to heat, it yields ammonia and prussic acid.

Amniotic acid may be obtained by evaporating the liquor of the amnios of the cow to a fourth part, and suffering it to cool; crystals of amniotic acid will be obtained in considerable quantity.

Whether this acid exists in the liquor of the amnios of other animals, is not yet known.

ANOMUM. (From an Arabian word, signifying a pigeon, whose foot it was thought to resemble.) The fruit of the *anomum verum*, True stone parsley. It is about the size of a grape, of a strong and grateful aromatic taste and penetrating smell. The seeds have been given as a carminative.

ANOMUM CARDAMOMUM. The systematic name for the *cardamomum minus*. See *Cardamomum minus*.

ANOMUM GRANUM PARADISI. The systematic name of the plant which affords the grains of paradise. See *Grana paradisi*.

ANOMUM ZINGIBER. The systematic name of the plant which affords ginger. See *Zingiber*.

AMORGE. See *Amurca*.

AMPELOSAGRIA. (From *αμπελος*, a vine.) See *Bryonia*.

AMPHIMERINOS. (From *αμφι*, about, *ημερα*, a day.) *Amphemerina*. A quotidian fever. A species of ague.

AMPHIARTHROSIS. (*Αμφιarthrosis*: from *αμφι*, both and *arthrosis*, an articulation; so called from its partaking both of diarthrosis and synarthrosis.) A mixed species of connexion of bones, which admits of an obscure motion, as is observed in the metacarpal and metatarsal bones, and the vertebræ.

AMPHIBIOUS. (From *αμφι*, *ambo*, and *βιος*, *vita*.) Animals are thus called, that live both on land and in the water. The *amphibious* animals, according to *Linnaeus*, are a class whose heart is furnished with one ventricle and one auricle, in which respiration is in a considerable degree voluntary.

AMPHIBLESTROINES. (From *αμφιβλετρον*, a net, and *ως*, a resemblance.) The retina, or net-like coat of the eye.

AMPHIBRANCHIA. (From *αμφι*, about, and *βραγχια*, the jaws.) The fauces or parts about the tonsils, according to *Hippocrates* and *Fæsius*.

AMPHICAUSTIS. (From *αμφι*, about, and *καυσος*, *puendum mulieb.*) A sort of wild barley growing about ditches. *Eusta-*

chius uses it to express the private parts of a woman.

AMPHIDEON. (From *αμφι*, on both sides, and *δαιω*, to divide.) *Amphideum*. *Amphidium*. The os tincæ, or mouth of the womb, which opens both ways, was so called by the ancients.

AMPHIDIARTHROSIS. The same as amphiarthrosis.

AMPHIMERINA. (From *αμφι*, about, and *μερα*, a day.) An intermitting fever of the quotidian kind.

AMPHIMETRIUM. (From *αμφι*, about, and *μετρα*, the womb) *Amphimetrium*. The parts about the womb. *Hippocrates*.

AMPHILEX. (From *αμφι*, about, and *πλεω*, to connect.) According to Rufus Ephesius, the part situated between the scrotum and anus, and which is connected with the thighs.

AMPHINEUMA. (From *αμφι*, about, and *πνευμα*, breath.) A difficulty of breathing. *Dyspnœa*. *Hippocrates*.

AMPHIPOLOS. (From *αμφι*, about, and *πολω*, to administer.) *Amphipolus*. One who attends the bed of a sick person, and administers to him. *Hippocrates*.

AMPHISMILA. (From *αμφι*, on both sides, and *σμιλα*, an incision-knife.) A dissecting knife, with an edge on both sides. *Galen*.

AMPULLA. (*αμβολλα*; from *αναβαλλω*, to swell out.) A bottle.

1. All bellied vessels are so called in chymistry, as bolt-heads, receivers, cucurbits, &c.

2. In anatomy this term is applied by Scarpa to the dilated portions of the membranaceous semicircular canals, just within the vestibulum of the ear.

AMPULLASCENS. (From *ampulla*, a bottle.) The most tumid part of Pecquet's duct is called alveus ampullascens.

AMPUTATIO. (From *amputo*, to cut off) *Ectome*. A surgical operation, which consists in the removal of a limb or viscus; thus we say a leg, a finger, the penis, &c. when cut off, are amputated; but when speaking of a tumour, or excrescence, it is said to be dissected out or removed.

AMULETUM. (From *αμμου*, a bond; because it was tied round the person's neck; or rather from *αμυνω*, to defend. An amulet, or charm; by wearing which the person was supposed to be defended from the admission of all evil; in particular, an antidote against the plague.

AMORCA. (From *αμυρω*, to press out.) *Amorça*. A small herb, whose expressed juice is used in dying. Also the sediment of the olive, after the oil has been pressed from it; recommended by Hippocrates and Galen as an application to ulcers.

AMUTICA. (From *αμυτω* to scratch.) Medicines that, by vellicating or scratching as it were, the bronchia, stimulate it to the discharge of whatever is to be thrown off the lungs.

ANYCHE. (From *αμυσσω*, to scratch.) A superficial laceration or exulceration of the skin: a slight wound. *Hippocrates*.—Scarification. *Galen*.

AMYTICA. (From *αμυσσω*, to vellicate.) Medicines which stimulate and vellicate the skin, according to Cælius Aurelianus.

AMYGDALA. (*Αμυδαλα*, from *αμυσσω*, to lacerate; so called, because after the green husk is removed from the fruit, there appear upon the shell certain fissures, as it were lacerations.) The almond.

1. The name of a genus of plants in the Linnæan system. Class, *Icosandria*. Order, *Monogynia*. The almond-tree.

2. The pharmacopœial name of the common almond. *Amygdalus communis* of Linnæus:—*foliis serratis infimis glandulosis, floribus sessilibus geminis*.

The almond is a native of Barbary. The same tree produces either bitter or sweet. Sweet almonds are more in use as food than medicine; but they are said to be difficult of digestion, unless extremely well comminuted. Their medicinal qualities depend upon the oil which they contain in the farinaceous matter, and which they afford on expression, nearly in proportion of half their weight. It is very similar to olive oil; perhaps rather purer, and is used for the same purposes. The oil thus obtained is more agreeable to the palate than most of the other expressed oils, and is therefore preferred for internal use, being generally employed with a view to obtund acrid juices, and to soften and relax the solids, in tickling coughs, hoarseness, costiveness, nephritic pains, &c. externally it is applied against tension and rigidity of particular parts, the milky solutions of almonds in watery liquors, usually called emulsions, possess, in a certain degree, the emollient qualities of the oil, and have this advantage over pure oil, that they may be given in acute or inflammatory disorders, without danger of the ill effects which the oil might sometimes produce by turning rancid. The official preparations of almonds, are the expressed oil and the emulsion; to the latter, the addition of gum-arabic is sometimes directed, which renders it a still more useful demulcent in catarrhal affections, stranguries, &c.

Bitter almonds yield a large quantity of oil, perfectly similar to that obtained from sweet almonds; but the matter remaining after the expression of the oil, is more powerfully bitter than the almond in its entire state. Great part of the bitter matter dissolves by the assistance of heat, both in water and rectified spirit; and a part arises also with both menstrua in distillation. Bitter almonds have been long known to be poisonous to various brute animals; and some authors have alleged that they are also deleterious to the human species; but the facts recorded upon this point appear to

want further proof. However, as the noxious quality seems to reside in that matter which gives it the bitterness and flavour, it is very probable, that when this is separated by distillation, and taken in a sufficiently concentrated state, it may prove a poison to man, as is the case with the common laurel, to which it appears extremely analogous. Bergius tells us, that bitter almonds, in the form of emulsion, cured obstinate intermittents, after the bark had failed. A simple water is distilled from bitter almonds, after the oil is pressed out, which possesses the same qualities, and in the same degree, as that drawn from cherry-stones. These afforded, formerly, the now exploded aqua cerasorum nigrorum, or black-cherry-water.

AMYGDALÆ. The almonds of the ears; from their supposed resemblance to almonds.

AMYGDALÆ ANARÆ. See *Amygdala*.

AMYGDALUS COMMUNIS. The systematic name of the plant which affords both sweet and bitter almonds. See *Amygdala*.

AMYGDALÆ DULCES. See *Amygdala*.

AMYGDALUS PERSICA. The systematic name of the common plum-tree. See *Persica*.

AMYLÆ. (From *amylum*, starch.) Any sort of chymical fæcula, or highly pulverized residuum.

AMYLEON. *Amylion*. Starch.

AMYLUM. (*Ἀμύλον*: from *a priv.* and *μύλος*, a mill; because it was formerly made from wheat without the assistance of a mill.) Starch. The fæcula of wheat, or starch of wheat. The white substance which subsides from the water that is mixed with wheaten flour. The starch-makers suffer it to remain in the water for a time after it has become acid, which makes it very white and soft to the touch, and scarcely sensible to the taste. As starch forms the greatest part of flour, it cannot be doubted but that it is the principal alimentary substance contained in our bread. In a medical point of view, it is to be considered as a demulcent; and, accordingly, it forms the principal ingredient of an official lozenge in catarrhs, and a mucilage prepared from it, often produces excellent effects, both taken by the mouth and in the form of a clyster, in dysenteries and diarrhœa, from irritation of the intestines. Milk and starch, with the addition of suet finely shred and incorporated by boiling, was the soup employed by Sir John Pringle, in dysenteries, where the mucous membrane of the intestines had been abraded. Externally, surgeons apply it as an absorbent in erysipelas.

ANTRIS ELEMIFERA. (From *a*, intensive, and *μύρον*, ointment, or balm; so called from its use, or smell. The systematic name of the plant from which it is supposed we obtain the resin called *gum elemi*. See *Elemi*.

ANTRIS OROBALSAMUM. The systematic name of the plant from which the balsam of Mecca is obtained. See *Balsamum Gileadense*.

AMYUM. (From *a*, priv. and *μύς*, muscle.) A limb so emaciated that the muscles scarcely appear.

ANA. *a*, or *aa*. In medical prescriptions it means "of each." See *A*.

ANABASIS. (From *αναβαίνω*, to ascend.) An ascension, augmentation, or increase of a disease, or paroxysm. It is usually meant of fevers. *Galen*. It also signifies equisetum.

ANABATICA. (From *αναβαίνω*, to ascend.) An epithet formerly applied to the synochus, or continual fever, when it increases in malignity.

ANABEXIS. (From *αναβέναι* to cough up.) An expectoration of matter by coughing.

ANABLEPSIS. (From *ανα* and *βλέπω*, to see again. The recovery of sight after it has been lost.

ANABLYSIS. (From *ανα* and *βλύω*, to gush out again.) Ebullition or effervescence.

ANABOLE. (From *αναβάλλω*, to cast up.) The discharge of any thing by vomit; also dilatation, or extension. *Galen*.

ANABROCHESIS. (From *ανα* and *βροχέω*, to resorb.) The reabsorption of matter.

ANABROCHISMOS. (From *αναβροχέω*, to reabsorb.) *Anabrochismus*. The taking up and removing the hair on the eye-lids, when they become troublesome. *Galen*. *Egineta*, and others.

ANABROSIS. (From *αναβροσέω*, to devour.) A corrosion of the solid parts, by sharp and biting humours. *Galen*.

ANACARDIUM OCCIDENTALE. (From *ανα*, without, and *καρδία*, a heart; without heart because the pulp of the fruit, instead of having the seed enclosed, as is usually the case, has the nut growing out of the end of it.) The cashew-nut. The oil of this nut is an active caustic, and employed as such in its native country; but neither it, nor any part of the fruit, is used medicinally in this country.

ANACARDIUM ORIENTALE. *Anacardium* or Malacca bean. The fruit, or nut, so called in the pharmacopœias, is of a shining black colour, heart-shaped, compressed, and about the size of the thumb-nail. It is the produce of the *Anicemia tomentosa*; *foliis cordato-ovatis, subtus tomentosis*, of Linnæus. It is now deservedly forgot in this country.

ANACATHARSIS. (From *ανα*, and *καθάρωμαι*, to purge up.) An expectoration of pus. It properly denotes a purgation by spitting, in which sense it stands contradistinguished from catharsis, or evacuation downwards. In this sense the word is used by Hippocrates and *Galen*. *Blancard* denotes, by this word, medicines which operate upwards, as vomiting. &c.

ANACATHARTICA. (From *ανακαθάρωμαι*, to purge upwards.) *Thoracica*. Medicines

which promote expectoration, or vomits which act upwards.

ANACHRON. Mineral alkali.

ANACLARIS. (From *ανακλαω*, to bend back.) A reflexion or recurvature of any of the members, according to Hippocrates.

ANACLISIS. (From *ανακλινω*, to recline.) A couch, or sick-bed. Hippocrates.

ANACOLIASMUS. (From *ανα*, and *κολω*, the bowels.) A gentle purge, which was sometimes used to relieve the lungs.

ANACOTHE. (From *ανακοχω*, to retard.) Delay in the administration of medicines; also slowness in the progress of a disease. Hippocrates.

ANACOLLEMA. (From *ανα*, and *κολλω*, to glue together.) A collyrium made of agglutinant substances, and stuck on the forehead. Galen.

ANACONCHYLISMOS. (From *ανακονχυλω*, to sound as a shell.) A gargarism so called, because the noise made in the throat is like the sound of a shell. Galen.

ANECTESIS. (From *ανακτιζω*, to renew.) Restoration of strength; recovery from sickness. Hippocrates.

ANACUPHISMA. (From *ανακυφω*, to lift up.) A kind of exercise mentioned by Hippocrates, which consists in lifting the body up and down, like our weigh-jolt.

ANACYCESIS. (From *ανακυκλω*, to mix.) The commixture of substances, or medicines, by pouring one upon another.

ANACYCLEON. (From *ανακυκλω*, to wander about.) *Anacycleus*. A mountebank or wandering quack.

ANACTRIOSIS. (From *ανα*, and *κριω*, authority.) By this word, Hippocrates means that gravity and authority which physicians should preserve among sick people and their attendants.

ANADIPLOSIS. (From *αναδιπλω*, to reduplicate.) A reduplication or frequent return of a paroxysm, or disease. Galen.

ANADOSIS. (From *ανα*, upwards, and *δοσμαι*, to give.) A vomit, or the distribution of alimen^t all over the body; or digestion.

ANADROME. (From *ανα*, upwards, and *δρομη*, to run.) A pain which runs from the lower extremities to the upper parts of the body. Hippocrates.

ANEDOS. (From *α*, priv. and *αιδω*, shame.) Shameless. Hippocrates uses this word metaphorically for without restraint, copious; and applies it to water rushing into the aspera arteria.

ANÆSTHESIA. (*Αναίσθησις*: from *α*, priv. and *αισθανομαι*, to feel.) Loss of the sense of touch. A genus of disease in the class *locales*, and order *dynæsthesiæ* of Cullen.

ANAGALLIS. (From *αναγαλλω*, to laugh; because, by curing the spleen, it disposes persons to be cheerful.) This plant *Anagallis arvensis*: *foliis indivisis, caule procumbente*, of Linnaeus, is small and delicately formed, and does not appear to possess any particular properties.

ANAGALLIS ARVENSI. The systematic name for the *anagallis* of the shops. See *Anagallis*.

ANAGARGALICTUM. (From *ανα*, and *γαργαζω*, the throat.) A gargarism, or wash for the throat.

ANAGARGARISTUM. The same.

ANAGLIFRE. (From *αναγλυφω*, to engrave.) A part of the fourth ventricle of the brain was formerly thus called, from its resemblance to a pen, or style.

ANAGNOSIS. (From *αναγνωσκω*, to know.) The persuasion, or certainty, by which medical men judge of a disease from its symptoms. Hippocrates.

ANAGRAPH. (From *αναγραφω*, to write.) A prescription, or receipt.

ANALEPSIA. (From *ανα*, and *λαμβάνω*, to take again.) A species of epilepsy, which proceeds from a disorder of the stomach, and with which the patient is apt to be seized very often and suddenly.

ANALENTIA. A fictitious term used by Paracelsus for epilepsy.

ANALEPSIS. (From *αναλαμβάνω*, to restore.) A recovery of strength after sickness. Galen.

ANALEPTICA. (From *αναλαμβάνω*, to recruit, or recover.) *Analeptics*. Restorative medicines; medicines, or food, which recover the strength which has been lost by sickness.

ANALOSIS. (From *αναλωσθαι*, to consume.) A consumption, or wasting.

ANALYSIS. (*Ανάλυσις*: from *αναλυω*, to resolve.) The resolution by chymistry, of any matter into its primary and constituent parts. The processes and experiments which chymists have recourse to, are extremely numerous and diversified, yet they may be reduced to two species, which comprehend the whole art of chymistry. The first is, *analysis*, or decomposition; the second, *synthesis*, or composition. In *analysis*, the parts of which bodies are composed, are separated from each other: thus, if we reduce cinnabar, which is composed of sulphur and mercury, and exhibit these two bodies in a separate state, we say we have decomposed, or analyzed cinnabar. But, if, on the contrary, several bodies be mixed together, and a new substance be produced, the process is then termed chymical composition, or *synthesis*: thus, if by fusion and sublimation, we combine mercury with sulphur, and produce cinnabar, the operation is termed chymical composition, or composition by synthesis.

ANAMNESIS. (From *αναμνησθαι* to remember.) Remembrance, or recollection of what has been done. Galen.

ANAMNESICA. (From the same.) Remedies for bad memory.

ANANAS. Called by the Brazilians *yayama*. The egg-shaped pine-apple. The plant which affords this fruit, is the *Bromelia ananas foliis ciliato-spinosis, mucronatis, spica comosa* of Linnaeus. It is used

principally as a delicacy for the table, and is also given with advantage as a refrigerant in fevers.

ANACE. (From *αναγκαζω*, to compel.) Necessity. It is applied to any desperate operation. *Hippocrates.*

ANAPHALANTIASIS. (From *αναφαλαντος*, bald.) A thinness of the hair upon the eye-brows. *Goræus.*

ANAPHORA. (From *αναφερω*, to bring up.) A person who spits blood. *Goræus.*

ANAPHORIXIS. (From *αναφορσσω*, to grind down.) The reducing of any thing to dust, or a very fine powder.

ANAPHRODISIA. (From *a*, priv. and *Αφροδισια*, the feast of Venus.) Impotence. A genus of disease in the class *locales*, and order *dysorexie* of Cullen. It either arises from paralysis, *anaphrodisia paralytica*; or from gonorrhœa, *anaphrodisia gonorrhœica*.

ANAPHROBOLI. (From *a*, neg. *αφρος*, froth, and *μελι*, honey.) Clarified honey.

ANAPLASIS. (From *αναπλασσω*, to restore again.) A restoration of flesh where it has been lost; also the reuniting a fractured bone. *Hippocrates.*

ANAPLEROSIS. (From *αναπληρωω*, to fill again.) The restitution, or filling up of wasted parts. *Galen.*

ANAPLEBOTICA. (From the same.) Medicines renewing flesh: incarnatives, or such medicines as fill up a wound so as to restore it to its original shape. *Galen.*

ANAPLEUSIS. (From *αναπλεωω*, to float upon.) The rotting of a bone, so that it drops off, and lies upon the flesh; exfoliation, or separation of a bone. *Hippocrates, Ægineta, &c.*

ANAPNEUSIS. (From *αναπνεωω*, to respire.) Respiration.

ANAPNOE. The same.

ANAPTOISIS. (From *αναπτωω*, to fall back.) A relapse.

ANAPTISIS. The same as *Anacatharsis*.

ANAREHESNIMIA. (From *ανα*, and *ρηγνυμι*, to break again.) *Anarthrosis*. A fracture; the fresh opening of a wound.

ANAREHGA. (From *ανα*, upwards, and *ρεω*, to flow.) A flux of humours from below upwards. *Schneider de Catarrho.*

ANAREUSOPIA. (From *ανα*, upwards, and *ρεωω*, to creep.) The same. *Hippocrates.*

ANAS DOMESTICA. (From *ανα*, to swim.) The tame duck. The flesh of this bird is difficult of digestion, and requires that warm and stimulating condiments be taken with it to enable the stomach to digest it.

ANASARCA. (From *ανα*, through, and *σαρξ*, flesh.) A species of dropsy from a serous humour, spread between the skin and flesh, or rather a general accumulation of lymph in the cellular system. Dr. Cullen ranks this genus of disease, in the class *Cachexie*, and the order *Intumescencia*. He enumerates the following species, viz. 1.

Anasarca serosa, as when the due discharge of serum is suppressed, &c. 2. *Anasarca oppilata*, as when the blood-vessels are considerably pressed, which happens to many pregnant women, &c. 3. *Anasarca exanthematica*, this happens after ulcers, various eruptive disorders, and particularly after the *erysipelas*. 4. *Anasarca anæmia*, happens when the blood is rendered extremely poor from considerable losses of it. 5. *Anasarca debilitum*, as when feebleness is induced by long illness, &c.

This species of dropsy shows itself at first with a swelling of the feet and ankles, toward the evening, which, for a time, disappears again in the morning. The tumefaction is soft and inelastic, and, when pressed upon by the finger, retains its mark for some time, the skin becoming much paler than usual. By degrees the swelling ascends upwards, and occupies the trunk of the body; and at last, even the face and eyelids appear full and bloated: the breathing then becomes difficult, the urine is small in quantity, high coloured, and deposits a reddish sediment; the belly is costive, the perspiration much obstructed, the countenance yellow, and a considerable degree of thirst, with emaciation of the whole body, prevails. To these symptoms succeed torpor, heaviness, a troublesome cough, and a slow fever. In some cases, the water oozes out, through the pores of the cuticle; in others, being too gross to pass by these, it raises the cuticle in small blisters; and sometimes the skin, not allowing the water to escape through it, is compressed and hardened, and is, at the same time, so much distended as to give the tumour a considerable degree of firmness.

ANASPAISIS. (From *ανα*, and *σπασσω*, to draw together.) Hippocrates uses this word to signify a contraction of the stomach.

ANASTTOS. (From *ανα*, upwards, and *σπωμαι*, to agitate.) *Anasytus*. Driven forcibly upwards. Hippocrates applies this epithet to air rushing violently upwards, as in hysteric fits.

ANASTALTICA. (From *ανασταλλω*, to contract.) Styptic or refrigerating medicines.

ANASTASIS. (From *αναστημι*, to cause to rise.) A recovery from sickness; a restoration of health. It likewise signifies a migration of humours, when expelled from one place and obliged to remove to another. *Hippocrates.*

ANASTOMOSIS. (From *ανα*, through, and *συνα*, a mouth.) The communication of vessels with one another.

ANASTOMOTICA. (From *ανα*, through, and *συνα*, the mouth.) Medicines which open the pores and mouths of the vessels, as cathartics, diuretics, deobstruents, and sudorifics.

ANATES. (From *ατες*, the buttocks.) A disease of the anus. *Forstus, &c.*

ANATOMY. (Ανατομή, or ανατομή: from *ana*, and *τεμνω*, to cut up.) *Anerotomy*. The dissection of the human body, to expose the structure, situation, and uses of every part.

ANATOMY, COMPARATIVE. Zootomy. The dissection of brutes, fishes, polypi, plants, &c. to illustrate, or compare them with the structure and functions of the human body.

ANATRESIS. (From *ana*, and *τρίγω*, to perforate.) A perforation like that which is made upon the skull by trepanning. *Galen*.

ANATRIBE. (From *ανατρίβω*, to rub.) Friction all over the body.

ANATRIPSIS. The same. *Moschion de Morb. Mulieb.* and *Galen*.

ANATRIS. *Antaris.* Mercury. *Ruland*.

ANATRON. (Arab. A lake in Egypt, where it was produced.) Soda, or fixed mineral alkali.

ANATROPE. (From *ανατρέπω*, to subvert. *Anatrophē.* *Anatrophā.* A relaxation, or subversion of the stomach, with loss of appetite and nausea. Vomiting. Indigestion. *Galen*.

ANATHUM. Soda.

ANAUDIA. (From *a*, priv. and *αὐδω*, the speech.) Dumbness; privation of voice; catalepsia. *Hippocrates*.

ANAXTRIS. (From *αναξυρίς*, the sole.) The herb sorrel; so called because its leaf is shaped like the sole of the shoe.

ANCHA. (Arab. to press upon, as being the support of the body.) The thigh. *Aricæana*, *Forestius*, &c.

ANCHILOPS. (From *αγγι*, near, and *ωψ*, the eye.) A disease in the inward corner of the eye, called also *Ægilops*. An incipient fistula lachrymalis.

ANCHYLE. See *Anchyle*.

ANCHORALIS PROCESSUS. (*Anchoralis*; from *αγκων*, the elbow.) See *Coracoïd process*.

ANCHOVY PEAR. This fruit, the produce of the *Grias cauliflora* of Linnæus, is eaten by the inhabitants of Jamaica, as a pleasant and refrigerant fruit.

ANCHUSA. (From *αγκυσ*, to strangle; from its supposed constringent quality; or, as others say, because it strangles serpents.) Alkanet.

1. The name of a genus of plants in the Linnæan system, Class, *Pentandria*. Order, *Monogynia*.

2. The name in some pharmacopœias for the alkanet root. The plant from which it is obtained, is the *anchusa tinctoria* of Linnæus, which grows wild in France, but is cultivated in our gardens. The root is externally of a deep purple colour. To oil, wax, turpentine, and alcohol, it imparts a beautiful deep red colour, for which purpose it is used. Its medicinal properties are scarcely perceptible.

ANCHUSA OFFICIALIS. The systematic

name for the *buglossum* of the shops. See *Buglossum*.

ANCHUSA TINCTORIA. The systematic name for the anchusa or alkanna of the pharmacopœias. See *Anchusa*.

ANCHYLOEISMIA. (From *αγκυλωμαι*, to bend.) Sagar uses this term to express a concretion, or growing together of the soft parts.

ANCHYLOSIS. (From *αγκυλωμαι*, to bend.) A stiff joint.

ANCI. Those who have a distorted elbow.

ANCIROMELE. See *Ancyromele*.

ANCINAR. BORAX.

ANCON. (From *αγκυλωμαι*, to embrace; *απο τῷ αγκυλωθεί τῷ ὅτι το ὅτι*; because the bones meeting, and there uniting, are folded one into another.) The elbow.

ANCONIUS. (*Anconus*, sc. *musculus*; from *αγκων*, the elbow.) *Anconius minor* of Winslow. *Anconius vel cubitalis Riolani* of Douglas. A small triangular muscle, situated on the back part of the elbow. It arises from the ridge and from the external condyle of the humerus, by a thick, strong, and short tendon: from this it becomes fleshy, and, after running about three inches obliquely backwards, it is inserted by its oblique fleshy fibres into the back part or ridge of the ulna. Its use is to extend the fore-arm.

ANCONIUS EXTERNUS. See *Triceps extensor cubiti*.

ANCONIUS INTERNUS. See *Triceps extensor cubiti*.

ANCONIUS MAJOR. See *Triceps extensor cubiti*.

ANCONIUS MINOR. See *Anconius*.

ANCONOID PROCESS. *Processus anconoides*. (From *αγκων*, the elbow.) A process of the cubit. See *Ulna*.

ANCTER. (*Αγκίτης*, a bond, or button.) A fibula, or button, by which the lips of wounds are held together. *Gorræus*.

ANCTERIASMUS. (From *αγκίτης*, a button.) The operation of closing the lips of wounds together by loops, or buttons. *Galen*.

ANCVBITUS. A disease of the eyes with a sensation of sand. *Joh. Anglic. Ro. Ang.*

ANCYLE. (From *αγκυλος*, crooked.) A species of contraction, called a stiff joint. *Galen*.

ANCYLOBLEPHARON. (From *αγκυλος*, a hook, and *βλεφαρον*, an eye-lid.) A disease of the eye, by which the eye-lids are closed together. *Ætius*.

ANCYLOGLOSSUM. (From *αγκυλος*, a hook, and *γλωσσα*, the tongue.) *Ancylium* of Ægineta. A contraction of the frænum of the tongue. Tongue-tied.

ANCYLOMELE. (From *αγκυλος*, crooked, and *μελλω*, a probe.) A crooked probe, or a probe with a hook. *Galen*, &c.

ANCYLOSIS. See *Anchylōsis*.

ANCYLOMUS. (From *αγκυλος*, a hook,

and *τρυακ*, to cut.) A crooked surgical knife, or bistoury. A knife for loosening the tongue. This instrument is no longer in use. *Egineta*, &c.

ANCTRA. (*Ἀγκυρα*, an anchor) A surgical hook. Epicharmus uses this word for the membrum virile, according to GORNEUS.

ANCTROIDES. (From *ἄγκυρα*, an anchor, and *ωδός*, a likeness.) A process of the scapula was so called, from its likeness to the beak of an anchor. It is the coracoid process. See *Scapula*.

ANCTRONELLE. See *Ancylomole*.

ANDRIA. A tree of Brazil, the fruit of which is bitter and astringent, and used as a vermifuge.

ANDRANATOMIA. *Andranatome*. (From *ανδρ*, a man, and *τομή*, to cut.) The dissection of the human body, particularly of the male. *M. Aur. Severinus, Zootome Democrit.*

ANDRAPODOCAPHELUS. (From *ἀνδραποδ*, a slave, and *καπηλός*, a dealer.) A crimp. Galen calls by this name the person whose office it was to anoint and slightly to wipe the body, to cleanse the skin from foulness.

ANDRIA. (From *ανδρ*, a man.) An hermaphrodite. *Bonnet.*

ANDROGÆTESIS. (From *ανδρ*, a man, and *κοίτη*, to cohabit with.) The venereal act; or the infamous act of sodomy. *Moschion*, &c.

ANDROGENUS. (From *ανδρ*, a man, and *γενν*, a woman.) An effeminate person. *Hipp.* An hermaphrodite.

ANDRONION. *Andronium*. A kind of plaster used by *Egineta* for carbuncles, invented by Andron.

ANDROTOMIA. *Androtome*. Human dissection, particularly of the male.

ANEBIUM. (From *ἀναβάναι*, to ascend.) The herb alkanet, so called from its quick growth.

ANEILESIS. (From *ἀνέλλω*, to roll up.) *Anilema*. An involution of the guts, such as is caused by flatulence and gripes. *Hippocrates.*

ANEMIA. (From *ἀνέμος*, wind.) Flatulence.

ANEMONE. (From *ἀνέμος*, wind; so named because it does not open its flowers till blown upon by the wind.) The wind flower. The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Polygynia*.

ANEMONE HEPATICA. The systematic name for the *hepatica nobilis* of the pharmacopœias. See *Hepatica nobilis*.

Anemone, meadow. See *Pulsatilla nigricans*.

ANEMONE NEMOROSA. The systematic name of the *ranunculus albus* of the pharmacopœias. See *Ranunculus albus*.

ANEMONE FRATENSIS. The systematic

name for the *pulsatilla nigricans* of the pharmacopœias. See *Pulsatilla nigricans*.

ANENCEPHALUS. (From *α*, priv and *κεφαλή*, the brain) A monster without brains. Foolish. *Galen de Hipp.*

ANEOS. A loss of voice and reason.

ANEPITHYMIA. (From *α*, priv. and *σθένος*, desire.) Loss of appetite.

ANERIC. *Anerit*. Sulphur vivum.

ANESIS. (From *ἀνέμω*, to relax.) A remission, or relaxation, of a disease, or symptom. *Ætius*, &c.

ANESUM. See *Anisum*.

ANETHUM. (*Ἄνηθος*: from *ἀνω*, afar, and *θω*, to run; so called because its roots run out a great way.) Fennel, dill, anet.

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order *Dygynia*.

2. The pharmacopœial name of the common dill, or anet.

Anethum graveolens of Linnæus:—*frutibus compressis*. This plant is a native of Spain, but cultivated in several parts of England. The seeds of dill are directed for use by the London and Edinburgh Pharmacopœias: they have a moderately warm, pungent taste, and an aromatic, but sickly smell. There is an essential oil, and a distilled water, prepared from them, which are given in flatulent colics and dyspepsia. They are also said to promote the secretion of milk.

ANETHUM FENICULUM. The systematic name for the *faniculum* of the shops. See *Feniculum*.

ANETHUM GRAVEOLENS. The systematic name for the *Anethum* of the shops. See *Anethum*.

ANETICA. (From *ἀνέμω*, to relax.) Pa-regorics; medicines which assuage pain, according to Andr. Tiraquell.

ANEURISMA. (*Ἀνευρυσμα*, from *ἀνευρυν*, to dilate.) An aneurism. A preternatural tumour formed by the dilatation of an artery. A genus of disease ranked by Cullen in the class *locales*, and order *tumores*. There are three species of aneurism: 1. The *true aneurism*, *aneurisma verum*, which is known by the presence of a pulsating tumour. The artery either seems only enlarged at a small part of its tract, and the tumour has a determinate border, or it seems dilated for a considerable length, in which circumstance the swelling is oblong, and loses itself so gradually in the surrounding parts, that its margin cannot be exactly ascertained. The first, which is the most common, is termed *circumscribed true aneurism*; the last, the *diffused true aneurism*. The symptoms of the circumscribed true aneurism, take place as follows: the first thing the patient perceives, is an extraordinary throbbing in some particular situation, and, on paying a little more attention, he discovers there a small pulsating tumour, which entirely dis-

appears when compressed, but returns again as soon as the pressure is removed. It is commonly unattended with pain or change in the colour of the skin. When once the tumour has originated, it continually grows larger, and at length attains a very considerable size. In proportion as it becomes larger, its pulsation becomes weaker, and, indeed, it is almost quite lost, when the disease has acquired much magnitude. The diminution of the pulsation has been ascribed to the coats of the artery losing their dilatable and elastic quality, in proportion as they are distended and indurated; and, consequently, the aneurismal sac being no longer capable of an alternate diastole and systole from the action of the heart. The fact is also imputed to the coagulated blood, deposited on the inner surface of the sac, particularly in large aneurisms, in which some of the blood is always interrupted in its motion. In true aneurisms, however, the blood does not coagulate so soon, nor so often as in false ones. Immediately such coagulated blood lodges in the sac, pressure can only produce a partial disappearance of the swelling. In proportion as the aneurismal sac grows larger, the communication into the artery beyond the tumour is lessened. Hence, in this state, the pulse below the swelling becomes weak and small, and the limb frequently cold and œdematous. On dissection, the lower continuation of the artery is found preternaturally small, and contracted. The pressure of the tumour on the adjacent parts, also produces a variety of symptoms, ulcerations, caries, &c. Sometimes an accidental contusion, or concussion, may detach a piece of coagulum from the inner surface of the cyst, and the circulation through the sac be obstructed by it. The coagulum may possibly be impelled quite into the artery below, so as to induce important changes. The danger of an aneurism arrives when it is on the point of bursting, by which occurrence the patient usually bleeds to death; and this is sometimes in a few seconds. The fatal event may generally be foreseen, as the part about to give way, becomes particularly tense, elevated, thin, soft, and of a dark purple colour. 2 The false or spurious aneurism, *aneurisma spurium*, is always owing to an aperture in the artery, from which the blood gushes into the cellular substance. It may arise from an artery being lacerated in violent exertions; but the most common occasional cause is a wound. This is particularly apt to occur at the bend of the arm, where the artery is exposed to be injured in attempting to bleed. When this happens, as soon as the puncture has been made, the blood gushes out with unusual force, and in a bright scarlet, irregular, interrupted current. It flows out, however, in an even and less ra-

pid stream when pressure is applied higher up than the wound. These last are the most decisive marks of the artery being opened; for blood often flows from a vein with great rapidity, and in a broken current, when the vessel is very turgid and situated immediately over the artery, which imparts its motion to it. The surgeon endeavours precipitately to stop the hæmorrhage by pressure; and commonly occasions a *diffused false aneurism*. The external wound in the skin is closed, so that the blood cannot escape from it. but innuates itself into the cellular substance. The swelling thus produced is uneven, often knotty, and extends upwards and downwards, along the tract of the vessel. The skin is also usually of a dark purple colour. Its size increases as long as the internal hæmorrhage continues, and, if this should proceed above a certain pitch, mortification of the limb ensues. 3. The *varicose aneurism*, *aneurisma varicosum*; this was first described by Dr W. Hunter. It happens when the brachial artery is punctured in opening a vein: the blood then rushes into the vein, which becomes varicose. Aneurisms may happen in any part of the body, except the latter species, which can only take place where a vein runs over an artery.

ANEURISMA SPURIUM. See *Aneurisma*.

ANEURISMA VARICOSUM. See *Aneurisma*.

ANEURISMA VERUM. See *Aneurisma*.

ANEXIS. (From *anexo*, to project.) A swelling, or protuberance.

ANGEILOLOGIA. (From *αγγιον*, a vessel, and *λογος*, a discourse.) A dissertation, or reasoning, upon the vessels of the body.

ANGIOTOMY. (From *αγγιον*, a vessel, and *τομη* to cut.) The dissection of the blood-vessels of an animal body; also the opening of a vein, or an artery.

ANGIOTOMUS. (From *αγγιον*, a vessel, and *τομη*, to cut.) A skilful dissector of the vessels.

ANGELICA. (So called from its supposed angelic virtues.) Angelica. 1. The name of a genus of plants in the Linnæan system. Class, *Penandria*. Order, *Digynia*.

2. The pharmacopœial name of the garden-angelica.

Angelica archangelica of Linnæus:—*foliorum impari lobato*. A plant, a native of Lapland, but cultivated in our gardens. The roots of angelica have a fragrant, agreeable smell, and a bitterish, pungent taste. The stalk, leaves, and seeds, which are also directed in the pharmacopœias, possess the same qualities, though in an inferior degree. Their virtues are aromatic and carminative. A sweetmeat is made, by the confectioners, of this root, which is extremely agreeable to the stomach, and is surpassed only by that of ginger.

ANGELICA ARCHANGELICA. The systematic name for the *angelica* of the shops. See *Angelica*.

Angelica, Garden. See *Angelica*.

ANGELICA SATIVA. See *Angelica sylvestris*.

ANGELICA SYLVESTRIS. *Angelica sativa.* Wild angelica. *Angelica sylvestris* of Linnaeus:—*foliis aequalibus ovato-lanceolatis serratis.* This species of angelica possesses similar properties to the garden species, but in a much inferior degree. It is only used when the latter cannot be obtained. The seeds, powdered and put into the hair, kill lice.

Angelica, wild. See *Angelica sylvestris*.

ANGELINE CORTEX. The tree from which this bark is procured is a native of Grenada. It has been recommended as an anthelmintic for children.

ANGELOCACOS. Myrobalans, or purging Indian plums.

ANGER. (From *angor*, anguish; because of their pain.) Bubbles in the groin. *Fallopian de Morbo Gallico.*

ANGILOSSUS. (From *αγκυρα*, a hook, and *γλωσσα*, the tongue.) A person who stammers.

ANGINA. (From *αγγω*, to strangle; because it is often attended with a sense of strangulation.) A soar throat. See *Cynanche*.

ANGINA MALIGNA. See *Cynanche maligna*.

ANGINA PAROTIDEA. See *Cynanche parotidea*.

ANGINA PECTORIS. An acute constrictory pain at the lower end of the sternum, inclining rather to the left side, and extending up into the left arm, accompanied with great anxiety. Violent palpitations of the heart, laborious breathings, and a sense of suffocation, are the characteristic symptoms of this disease.—It is found to attack men much more frequently than women, particularly those who have short necks, who are inclinable to corpulency, and who, at the same time, lead an inactive and sedentary life. Although it is sometimes met with in persons under the age of twenty, still it more frequently occurs in those who are between forty and fifty. In slight cases, and in the first stage of the disorder, the fit comes on by going up-hill, up-stairs, or by walking at a quick pace after a hearty meal; but as the disease advances, or becomes more violent, the paroxysms are apt to be excited by certain passions of the mind; by slow walking, by riding on horseback, or in a carriage, or by sneezing, coughing, speaking, or straining at stool. In some cases, they attack the patient from two to four in the morning, or whilst sitting or standing, without any previous exertion or obvious cause. On a sudden, he is seized with an acute pain in the breast, or rather at the extremity of the sternum, inclining to the left side, and extending up

into the arm, as far as the insertion of the deltoid muscle, accompanied by a sense of suffocation, great anxiety, and an idea that its continuance, or increase, would certainly be fatal. In the first stage of the disease, the uneasy sensation at the end of the sternum, with the other unpleasant symptoms, which seemed to threaten a suspension of life by a perseverance in exertion, usually go off upon the person's standing still, or turning from the wind; but, in a more advanced stage, they do not so readily recede, and the paroxysms are much more violent. During the fit, the pulse sinks in a greater degree, and becomes irregular; the face and extremities are pale, and bathed in a cold sweat, and, for a while, the patient is perhaps deprived of the powers of sense and voluntary motion. The disease having recurred more or less frequently during the space of some years, a violent attack at last puts a sudden period to his existence. Angina pectoris is attended with a considerable degree of danger; and it usually happens that the person is carried off suddenly. It mostly depends upon an ossification of the coronary arteries, and then we can never expect to effect a radical cure. During the paroxysms, considerable relief is to be obtained from fomentations, and administering powerful antispasmodics, such as opium and ether combined together. The application of a blister to the breast is likewise attended sometimes with a good effect. As the painful sensation at the extremity of the sternum often admits of a temporary relief, from an evacuation of wind by the mouth, it may be proper to give frequent doses of carminatives, such as peppermint, caraway, or cinnamon water. Where these fail in the desired effect, a few drops of oil of anise, on a little sugar, may be substituted.

With the view of preventing the recurrences of the disorder, the patient should carefully guard against passion, or other emotions of the mind: he should use a light, generous diet, avoiding every thing of a heating nature; and he should take care never to overload the stomach, or to use any kind of exercise immediately after eating. Besides these precautions, he should endeavour to counteract obesity, which has been considered as a predisposing cause; and this is to be effected most safely by a vegetable diet, moderate exercise at proper times, early rising, and keeping the body perfectly open. It has been observed that angina pectoris is a disease always attended with considerable danger, and, in most instances, has proved fatal under every mode of treatment. We are given, however, to understand, by Dr. Macbride, that of late, several cases of it have been treated with great success, and the disease readily removed, by inserting

a large issue in each thigh. These, therefore, should never be neglected. In one case, with a view of correcting, or draining off the irritating fluid, he ordered, instead of issues, a mixture of lime-water with a little of the spiritus juniperi comp. and an alterative proportion of Huxham's antimonial wine, together with a plain, light, perspirable diet. From this course the patient was soon apparently mended; but it was not until after the insertion of a large issue in each thigh, that he was restored to perfect health.

ANGINA TONSILLARIS. See *Cynanche tonsillaris*.

ANGINA TRACHEALIS. See *Cynanche trachealis*.

ANGIOLOGIA. (From *αἷμα*, a vessel, and *λογος*, a discourse.) The doctrine of the vessels of the human body.

ANGLICUS SUDOR. (From *Anglia*, England, and *sudor*, sweat.) The sweating sickness. *Sennertus*.

ANGOLAM. A very tall tree of Malabar, possessing vermifuge powers.

ANGORE. (From *αγγα*, to strangle.) A nervous sort of quinsey, or hysteric, suffocation, where the fauces are contracted and stopped up without inflammation.

ANGOR. Intense bodily pain. *Galen*.

ΑΝΓΟΣ. (*Αγγος*, a vessel.) A vessel; a collection of humours.

ANGUSTURÆ CORTEX. *Cortex cusparia*. A bark imported from Angustura, in South America. Its external appearances vary considerably. The best is not fibrous, but hard, compact, and of a yellowish brown colour, and externally of a whitish hue. When reduced into powder, it resembles that of Indian rhubarb. It is very generally employed as a febrifuge, tonic, and adstringent. While some deny its virtue in curing intermittents, by many it is preferred to the Peruvian bark: and has been found useful in diarrhoea, dyspepsia, and scrofula. It was thought to be the bark of the *Brucea antidyenterica*, or *ferruginea*. Wildenow suspected it to be the *Mugnolia plumieri*; but Humbolt and Bonpland, the celebrated travellers in South America, have ascertained it to belong to a tree not before known, and which they propose to describe by the name of *cusparia febrifuga*.

ANHELATIO. (From *anhelo*, to breathe with difficulty.) *Anhelitus*. Shortness of breathing.

ANICETON (From *a*, priv. and *νικη*, victory.) A name of a plaister invented by Crito, and so called because it was thought an infallible or invincible remedy for achores, or scald-head. It was composed of litharge, alum, and turpentine, and is described by *Galen*.

ANIMA. The thinking principle.

ANIMA ALOES. Refined aloes.

ANIMA ARTICULORUM. *Hermiodactylus*.

ANIMA HEPATIS. *Sal martis*.

ANIMA PULMONUM. The soul of the lungs. A name given to saffron on account of its use in asthma.

ANIMA RHABARBARI. The best rhubarb.

ANIMA SATURNI. A preparation of lead.

ANIMA VENERIS. A preparation of copper.

ANIMAL. An organized body endowed with life and voluntary motion.

ANIMAL ACTIONS. *Actiones animales*.

Those actions, or functions, are so termed which are performed through the means of the mind. To this class belong the external and internal senses, the voluntary action of muscles, voice, speech, watching, and sleep.

ANIMAL HEAT. Heat is essentially necessary to life. That of a man in health is from about 94 deg. to 100 deg. of Fahrenheit. It appears to depend upon the decomposition of the air in the lungs. See *Respiration*.

ANIMAL OIL. *Oleum animale*. An empyreumatic oil, obtained from the bones of animals, recommended as an anodyne and antispasmodic.

ANIME GUMMI. The substance which bears this name in the shops is a resin, the produce of the *Hymecia courbaril* of Linnaeus. It is seldom ordered in the practice of the present day, and is only to be met with in the collections of the curious.

ANIMI DELIQUIUM (From *animus*, the mind, and *delinquo*, to leave.) Fainting. See *Syncope*.

ANIMUS. This word is to be distinguished from *anima*; the former expresses the faculty of reasoning, and the latter the being in which that faculty resides.

ANINGA. A root which grows in the Antilla islands, and is used by sugar-bakers for refining their sugar.

ANISALPTOR. (From *anus*, the breech, and *scalpo*, to scratch.) The latissimus dorsi is so called, because it is the muscle chiefly instrumental in performing this office. *Burtholin*.

ANISOTACHYS. (From *ανισος*, unequal; and *ταχος*, quick.) A quick and unequal pulse. *Gorræus*.

ANISUM. (From *a*, neg. and *ισος*, equal.) Anise. *Anisum vulgare*. *Pimpinella anisa* of Linnaeus;—*foliis radicalibus trifidis incisiss*. A native of Egypt. Anise-seeds have an aromatic smell, and a pleasant, warm, and sweetish taste. An essential oil and distilled water are prepared from them, which are employed in flatulences and gripes, to which children are more especially subject; also in weakness of the stomach, diarrhoeas, and loss of tone in the primæ viæ.

ANISUM SILENSE. See *Anisum stellatum*.

ANISUM STELLATUM. *Anisum si-*

nense. Semen badian. The plant which affords these seeds is the *Illicium anisatum* of Linnæus. They are used with the same views as those of the *Pimpinella anisum*. The same tree is supposed to furnish the aromatic bark called *cortex anisi stellati*, or *cortex lavola*.

ANISUM VULGARE. See *Antrum*.

ANNUTES. (From *annuo*, to nod.) Some muscles of the head were formerly so called, because they perform the office of nodding, or bending the head downwards. *Cowper*, &c.

ANNULAR. (*Annularis*.) Like a ring; thus, annular bone, &c.

ANNULAR BONE. *Circulus ossæus*. A ring-like bone placed before the cavity of the tympanum in the fœtus.

Annular cartillages. See *Cricoid cartilages*.

ANNULARIS DIGITUS. The ring-finger. The one between the little and middle fingers.

ANNULARIS PROCESSUS. See *Pons varolii*.

ANO. (*Ana*, upwards; in opposition to *κατα*, downwards.) Upwards.

ANOCATHARTICA. (From *ana*, upwards, and *καθαρο*, to purge.) Emetics; medicines which purge upwards.

ANOCHILON. (From *ana*, upwards, and *χῆλος*, the lip.) The upper lip.

ANODIA. (From *a*, neg. and *δία*, the way.) Hippocrates uses this word for inaccuracy and irregularity in the description and treatment of a disease.

ANODYNA. See *Anodynes*.

ANODYNES. (*Anodyna*, sc. *medicamentia*. From *a*, priv. and *αδυν*, pain.) Those medicines are so termed which ease pain and procure sleep. They are divided into three sorts; paregorics, or such as assuage pain; hypnotics, or such as relieve by procuring sleep; and narcotics, or such as ease the patient by stupefying him.

ANODYNUM MINERALE. *Salprunella*.

ANODYNUM MARTIALE. *Ferrum ammoniacale* precipitated from water by potash.

ANOMALOUS. This term is often applied to those diseases whose symptoms do not appear with that regularity generally observed in diseases. A disease is also said to be anomalous, when the symptoms are so varied as not to bring it under the description of any known affection.

ANOMPHALOS. (From *a*, priv. and *ομφαλος*, the navel.) *Anomphalus*. Without a navel.

ANONYMUS. (From *a*, priv. and *ονυμα* a name.) Nameless. It was formerly applied to the cricoid muscle.

ANORCHIDES. (From *a*, priv. and *ορχις*, the testicle.) Children are so termed which come into the world without testicles. This is a very common occurrence. The testicles of many male infants at the time of birth are within the abdomen. The time of their descent is very uncertain, and

instances have occurred where they had not reached the scrotum at the age of ten and fifteen.

ANOREXIA. (From *a*, priv. and *ορεξ*, appetite.) A want of appetite, without loathing of food. Cullen ranks this genus of disease in the class *locales*, and order *dysorexia*; he believes it to be generally symptomatic, but enumerates two species, viz. the *anorexia humoralis* and the *anorexia atonica*.

ANOSMIA. (From *a*, neg. and *οσμη*, to smell.) A loss of the sense of smelling. This genus of disease is arranged by Cullen in the class *locales*, and order *dysæsthesie*. When it arises from a disease of the Schneiderian membrane, it is termed *anosmia organica*; and when from no manifest cause, *anosmia atonica*.

ANSER DOMESTICUS. The tame goose. The flesh of this bird is sometimes similar to that of the duck, and requires the assistance of spirituous and stimulating substances to enable the stomach to digest it. Both are very improper for weak stomachs.

ANSERINA. (From *anser*, a goose; so called, because geese eat it.) *Argentia*. Wild tansy, or goose-grass. This herb, *Potentilla anserina*, *foliis dentatis serratis, caule repente, pedunculis unifloris* of Linnæus, was formerly used as an astringent in laxity of the intestines and phthisical complaints, but is now fallen into disuse.

ANTACIDS. (*Antacida*, sc. *medicamenta*. From *αντι*, against, and *acidus*, acid.) Remedies which obviate acidity in the stomach. Their action is purely chymical, as they merely combine with the acid present, and neutralize it. They are only palliatives, the generation of acidity being to be prevented by restoring the tone of the stomach and its vessels. Dyspepsia and diarrhœa are the diseases in which they are employed. The principal antacids in use are the alkalies. *Liquoris potassæ*, gutt. xv. or from 5 to 15 gr. of carbonat of potash, or soda dissolved in water. The solution of soda called double soda-water, or potash supersaturated with carbonic acid, is more frequently used, as being more pleasant. Ammonia has been recommended as preferable to every other antacid, from 20 to 40 drops of the liquor ammoniæ in a cupful of water. The liquor calcis, or lime water, is likewise used to correct acidity, two or three ounces being taken occasionally. *Creta præparata* alone, or with the addition of a small quantity of any aromatic—*chelæ cancrorum præparata*, and *magnesia* also and its carbonate, are used for the same purpose.

ANTAGONIST MUSCLES. Counteracting muscles, or those muscles which have opposite functions. Such are flexor and extensor of any limb, the one of which contracts it, the other stretches it out; and also the abductors and adductors. Solitary muscles are

those without any antagonist, as the heart, &c.

ANTALGICA. (From *anti*, against, and *algos*, pain.) Anodynes. Remedies which relieve pain.

ANTALKALINES. (From *anti*, against, and *alkali*, an alkali.) Medicines which possess the power of neutralizing alkalis. All the acids are of this class.

ANTAPHRODISIACA. (From *anti*, against, and *aphrosis*, Venus.) Anti-venereals, or medicines which extinguish amorous desires. *Wedel. Amen. Med.*

ANTAPHRODITICA. The same.

ANTAPODOSIS. (From *antapodosis*, to reciprocate.) A vicissitude, or return of the paroxysm of fevers. *Hippocrates.* Called by Galen *epilepsis*.

ANTARTHRITICA. (From *anti*, against, and *arthra*, the gout.) Medicines which relieve or repel the gout.

ANTASTHMATICA. (From *anti*, against, and *asthma*, an asthma.) Remedies against asthma.

ANTATROPHICA. (From *anti*, against, and *tropha*, a consumption.) Medicines which relieve or restore consumption.

ANTECHESIS. (From *antechesis*, to resist.) A violent stoppage in the bowels, which resists all efforts to remove it. *Hippocrates.*

ANTELABIUM. (From *ante*, before, and *labium*, a lip.) The extremity of the lip.

ANTEMBASIS. (From *anti*, mutually, and *emba*, to enter.) A coalescence, or union of bone. *Galen.*

ANTEMETICA. (From *anti*, against, and *emetica*, to vomit.) Medicines which stop or prevent vomiting.

ANTENEASMUS. (From *anti*, against, and *enasmus*, implacable.) That species of madness in which the patient endeavours to destroy himself.

ANTEPHIALTICA. (From *anti*, against, and *ephialtes*, the night-mare.) Medicines which prevent the night-mare.

ANTEPILEPTICA. (From *anti*, against, and *epilepsis*, the epilepsy.) Remedies against the epilepsy, and other convulsive disorders.

ANTERIOR AURIS. One of the common muscles of the ear, situated before the external ear. It arises, thin and membranous, near the posterior part of the *zygoma*, and is inserted into a small eminence on the back of the helix, opposite to the concha, which it draws a little forwards and upwards.

ANTERIOR INTERCOSTAL NERVE. *Splanchnic nerve.* A branch of the great intercostal that is given off in the thorax.

ANTERIOR MALLEI. See *Laxator tympani*.

ANTHELIX. See *Antihelix*.

ANTHELMIA. (From *anti*, against, and *helmos*, a worm.) The herb Indian pink, or worm-grass, so called, because it was thought of great virtue in expelling worms. See *Spigelia Marylandica*.

ANTHELMINTICS. (*Anthelmintica*, sc.

medicamenta; from *anti*, against, and *helmos*, a worm.) Medicines which procure the evacuation of worms from the stomach and intestines. The greater number of them act mechanically, dislodging the worms, by the sharpness or roughness of their particles, or by their cathartic operation. Some seem to have no other qualities than those of powerful bitters, by which they either prove noxious to these animals, or remove that debility of the digestive organs, by which the food is not properly assimilated, or the secreted fluids poured in the intestines are not properly prepared; circumstances from which it has been supposed the generation of worms may arise. The principal medicines belonging to this class, are: Calomel, gamboge, *Geoffrea mermis*, *tanacetum*, *polypodium filix mas*, *spigelia Marylandica*, *artemesia santonica*, *olea Fropæa*, *stannum pulverisatum*, *ferri limaturæ*, and *dolichos pruriens*: which see under their respective heaps.

ANTHEMIS. (From *anthos*, *floreo*; because it bears an abundance of flowers.) Chamomile.

1. The name of a genus of plants in the Linnæan system. Class *Syngenesia*. Order, *Polygamia superflua*.

2. The name in the last London Pharmacopœia for chamomile. See *Chamamelum*.

ANTHEMIS COTULA. (*Cotula*, a dim. of *coa*, a whetstone; so called from its leaves resembling a whetstone.) The systematic name for the plant called *Cotula fetida* in the pharmacopœias. See *Cotula fetida*.

ANTHEMIS NOBILIS. (From *anthos*, a flower.) The systematic name for the *chamamelum* of the shops. See *Chamamelum*.

ANTHEMIS PYRETHRUM. The plant is so called from which we obtain the pyrethrum of the pharmacopœias. See *Pyrethrum*.

ANTHERA. (From *anthos*, a flower.)

1. A compound medicine used by the ancients; so called from its florid colour. *Galen. Fgineta*.

2. The male part of the fructification of plants.

ANTHOPHYLLI. (From *anthos*, a flower, and *phallos*, a leaf; so called from the fragrance of the flowers and the beauty of the leaves.) Cloves are so termed when they have been suffered to grow to maturity *G. Bauhin Pin*.

ANTHORA. (*Quasi antithora*, *antithora*: from *anti*, against, and *thora*, monkshood; so called because it is said to counteract the effects of the thora or monkshood.) A species of *Wolfsbane*. The root is the part of this plant (*Aconitum anthora*: *floribus pentagynis, foliorum laciniis linearibus* of Linnæus,) which is employed medicinally. Its virtues are similar to those of the *aconitum*. See *Aconitum*.

AUTHOS FLORES. The flowers of the *rosmarinus* are so termed in some pharmacopœias.

ANTHRACIA. See *Anthrax*.

ANTHRAX (From *αἰσθῆς*, a burning coal.) *Anthracia*. *Anthrocozium*. *Anthrocoma*. *Carbunculus*. A hard and circumscribed inflammatory tubercle like a boil, which sometimes forms on the cheek, neck, or back, and in a few days becomes highly gangrenous. It then discharges an extremely foetid sanies from under the black core, which, like a burning coal, continues destroying the surrounding parts. It is supposed to arise from a peculiar miasma, is most common in warm climates, and often attends the plague.

ANTHRACOSIS OCULI. A red, livid, burning, sloughy, very painful tumour, occurring on the eye-lids. *Agnetia*.

ANTHROPOGRAPHIA. (From *ἄνθρωπος*, a man, and *γραφω*, to write.) Description of man's structure.

ANTHROPOLOGIA. (From *ἄνθρωπος*, a man, and *λογος*, a discourse.) The description of man.

ANTHYPNOTICA. (From *ἄντη*, against, and *υπνος*, sleep.) Medicines which prevent sleep or drowsiness.

ANTHYPOCHONDRIACA. (From *ἄντη*, against, and *υποχονδρία*, the hypochondria.) Medicines adapted to cure low-spiritedness or disorders of the hypochondria.

ANTHYSTERICA. (From *ἄντη*, against, and *ύστερ*, the womb.) Uterines or medicines which relieve the hysterical passion. *Blanchard*.

ANTI. (*ἄντη*, against.) There are many names compounded with this word, as *antiasthmatica*, *antihysterica*, *antidysenteries*, &c. which signify medicines against the asthma, hysterics, dysentery, &c.

ANTIADÆS. (From *ἄντις*, to meet.) The tonsils are so called, because they answer one another. The mumps. *Nic Ples*.

ANTIAGRA. (From *ἄντις*, a tonsil, and *αἶψα*, a prey.) *Antiaagri*. A tumour of the tonsils. *Ulpian*, *Risland*, &c.

ANTIARTHRITICA. See *Antiarthritica*.

ANTICACHECTICA. (From *ἄντη*, against, and *κακία*, a cachexy.) Medicines against cachexy, or bad habit of body.

ANTICARDIUM. (From *ἄντη*, against, or opposite, and *καρδία*, the heart.) The hollow at the bottom of the breast, commonly called *scrobiculus cordis*, or pit of the stomach.

ANTICATARRAHIA. (From *ἄντη*, against, and *κατάρρως*, a catarrh.) Medicines which relieve a catarrh.

ANTICAUSTICA. (From *ἄντη*, against, and *καυσος*, a burning fever.) Remedies against burnings or fevers. We read, in *Corp. Pharm.* of *Juiken*, of a *symplicum anticausticum*.

ANTICHERIS. (From *ἄντη*, against, and *χρῆς*, the hand.) The thumb. *Galen*.

ANTICLIVUS. (From *ἄντη*, against, or opposite, and *κνήμη*, the calf of the leg.) That part of the tibia which is bare of flesh, and opposite the calf of the leg. The shin-bone. *Galen*.

ANTICOMICA. (From *ἄντη*, against, and

κομικα, the cholic.) Remedies against the cholic.

ANTIDIASTOLE. (From *ἄντη*, against, and *διαστολή*, to distinguish.) An exact and accurate distinction of one disease or symptom, from another.

ANTIDENICA. (From *ἄντη*, against, and *δενος*, circumgyration.) Medicines against a vertigo, or giddiness. *Blancard*.

ANTIDOTARIUM. (From *ἄντιδωτος*, an antidote.) A term used by former writers, for what we now call a dispensatory; a place where antidotes are prescribed and prepared. There are antidotaries extant of several authors, as those of *Nicholaus*, *Meuses*, *Myrepsus*, &c.

ANTIDOTUS. (From *ἄντη*, against, and *δωμι*, to give.) A preservative against sickness. A remedy. *Galen*.

ANTIDYSENTERICA. (From *ἄντη*, against, and *δυσεντερία*, a flux.) Medicines against a dysentery, or flux.

ANTIFEBRILIA. (From *ἄντη*, against, and *febris*, a fever.) A febrifuge, a remedy against fever.

ANTIHECTICA. [From *ἄντη*, against, and *ἡκτικός*, a hectic fever.) Remedies against a hectic fever.

ANTIHECTICUM POTERII. *Antimonium diaphoreticum Joviale*. A medicine invented by Poterius, formerly extolled as effectual in hectic fevers, but now disregarded. It is an oxyd of tin and chalybeated regulus of antimony, in consequence of their deflagration with nitre.

ANTIHELIX. (From *ἄντη*, against, and *ἑλὴξ*, the helix.) The inner circle of the auricle, so called from its opposition to the outer circuit called the helix.

ANTHELMINTICA. See *Anthelmintica*.

ANTHYSTERICA. (From *ἄντη*, against, and *ύστερ*, hysterics.) Medicines which prevent or relieve hysterics.

ANTILEPSIS. (From *ἄντιλαμβάνω*, to take hold of.) The securing of bandages or ligatures from slipping. *Hippocrates*.

ANTILOBUM. (From *ἄντη*, opposite, and *λοβός*, the bottom of the ear.) The tragus, or that part of the ear which is opposite the lobe.

ANTILOMICA. (From *ἄντη*, against, and *λοιμός*, the plague.) Remedies or preventives against the plague.

ANTILOPUS. The antelope. An African beast resembling a deer, whose hoofs and horns were formerly given in hysteric and epileptic cases.

ANTILYSsus. (From *ἄντη*, against, and *λύσσα*, the bite of a mad dog.) A medicine or remedy against the bite of a mad dog.

Antimonial powder. See *Pulvis antimoniæ*.

ANTIMONIALE. (From *antimonium*.) An antimonial, or composition in which antimony is a chief ingredient. A preparation of antimony.

ANTIMONIUM OXYDUM. See *Oxydum antimoniæ*.

ANTIMONII SULPHURETUM PRÆCIPITATUM. *Sulphur antimonii præcipitatum.* Precipitated sulphuret of antimony. This preparation of antimony appears to have rendered that called Kerme's mineral unnecessary. It is made thus:—

Take of sulphuret of antimony, in powder, two pounds:—of the solution of potash, four pints:—of distilled water, three pints.

Mix and boil the mixture over a slow fire, for three hours, stirring it well, and occasionally adding distilled water, so that the same measure may be preserved. Strain the solution forthwith through a double linen cloth; and while it is yet hot, drop in, gradually, as much sulphuric acid as may be required to precipitate the powder; then wash away the sulphate of potash, by hot water; dry the precipitated sulphuret of antimony, and reduce it to powder.

As an alterative and sudorific, it is in high estimation, and given in diseases of the skin and glands; and joined with calomel, it is one of the most powerful and penetrating alteratives we are in possession of.

ANTIMONIUM. See *Antimony*.

ANTIMONIUM CALCINATUM. See *Oxydum antimonii*.

ANTIMONIUM DIAPHORETICUM. An old name for oxyd of antimony.

ANTIMONIUM TARTARISATUM. *Tartarus emeticus.* *Tartarus emeticum.* *Tartarus antimonialis.* *Tartaris antimonii cum potassa.* *Tartarum stibiatum.* Tartar emetic is obtained by boiling an oxyd of antimony with acidulous tartrite of potash; the excess of tartarous acid dissolves the oxyd and a triple salt is obtained by crystallization. The London Pharmacopœia directs thus:

Take of oxyd of antimony, two ounces:—of supertartrite of potash, powdered, three ounces:—of distilled water, eighteen fluid-ounces.

To the water, whilst boiling in a glass vessel, add gradually the antimony and supertartrite of potash, previously mixed together, and continue to boil for half an hour; then filter the solution through paper, and evaporate it in a gentle heat, so that whilst it cools slowly, crystals may form.

Tartar emetic is the most useful of all the antimonial preparations. Its action is not dependent on the state of the stomach, and, being soluble in water, its dose is easily managed, while it also operates more speedily.

In doses of from one to three, four, or five grains, it generally acts powerfully as an emetic, and is employed whenever we wish to obtain the effects which result from full vomiting. As patients are differently affected by this medicine, the safest mode of exhibiting it is: *℞. Antimonii tartarizati, gr. iii. Aquæ distillatæ, ℥. v. Misce et cola. Dosis ℥ss. quadrante quâque horâ, donec supervenerit vomitus.*

For children, emetic tartar is not so safe an emetic as ipecacuanha powder: when

great debility of the system is present, even a small dose has been known to prove fatal to children. Sometimes it proves cathartic.

In smaller doses it excites nausea, and proves a powerful diaphoretic and expectorant. As an emetic it is chiefly given in the beginning of fevers and febrile diseases, when great debility is present, and in the advanced stages of typhoid fever its use is improper and even sometimes fatal. As a diaphoretic, it is given in small doses, of from an eighth to a quarter of a grain: and as an expectorant, in doses still smaller. Emetic tartar in small doses, combined with calomel, has been found a powerful yet safe alterative in obstinate eruptions of the skin. *℞. Antimonii tartarizati gr. iv. Hydragryi submuriatis, gr. xvi. Confectionis Rosæ gallicæ, q. s. Divide in pil. xxiv. Capiat i. mane nocteque ex thea sassafras.*

In the form of powder, or dissolved in water, it is applied by a pencil to warts and obstinate ulcers: it is also given in the form of clyster, with a view to produce irritation in soporose diseases, apoplexy, ileus, and hernia incarcerata. The powder mixed with any fluid, and rubbed on the scrobiculus cordis, excites vomiting. Another property which tartar emetic has, when rubbed on the skin, is that of producing a crop of pustules very like to the small-pox, and with this view it is used against rheumatic pains, white, and other obstinate swellings. The best antidote against the bad effects of too large a quantity of this and other antimonial preparations, is a decoction of the bark of cinchona.

ANTIMONIUM VITRIFACTUM. Glass of antimony. An oxyd of antimony, with a little sulphur.

ANTIMONY. (*Antimony.* The origin of this word is very obscure. The most received etymology is, from *anti*, against, and *monac*, a monk; because Valentine, by an injudicious administration of it, poisoned his brother monks.) *Antimonium.* *Stibium.* A metal found native, but very rarely; it has, in that state, a metallic lustre, and is found in masses of different shapes; its colour is white, between those of tin and silver. It generally contains a small portion of arsenic. It is likewise met with in the state of an oxyd, *antimonial ochre*. The most abundant ore of it is that in which it is combined with sulphur, the *grey ore of antimony*, or *sulphuret of antimony*. The colour of this ore is blueish, or steel-grey, of a metallic lustre, and often extremely beautifully variegated. Its texture is either compact, foliated, or striated. The striated is found both crystalized, massive, and disseminated: there are many varieties of this ore.

Properties of Antimony.—Antimony is a metal of a greyish white, having a slight bluish shade, and very brilliant. Its texture is lamellated, and exhibits plates crossing each other in every direction. Its surface

with herbarizations and foliage. Its specific gravity is 6.702. It is sufficiently hard to scratch all the soft metals. It is very brittle, easily broken and pulverizable. It fuses at 810° Fahr. It can be volatilized, and burns by a strong heat. When perfectly fused, and suffered to cool gradually, it crystallizes in octahedra. It unites with sulphur and phosphorus. It decomposes water strongly. It is soluble in alkaline sulphurets. Sulphureous acid, boiled upon antimony, is feebly decomposed. Nitric acid dissolves it in the cold. Muriatic acid scarcely acts upon it. The oxygenated muriatic acid gas inflames it, and the liquid acid dissolves it with facility. Arsenic acid dissolves it by heat with difficulty. It unites, by fusion, with gold and renders it pale and brittle. Platinum, silver, lead, bismuth, nickel, copper, arsenic, iron, cobalt, tin, and zinc, unite with antimony by fusion, and form with it compounds, more or less brittle. Mercury does not alloy with it easily. We are little acquainted with the action of alkalies upon it. Nitrate of potash is decomposed by it. It fulminates by percussion with oxygenated muriate of potash.

Methods of obtaining antimony—1. To obtain antimony, heat 32 parts of filings of iron to redness, and project on them, by degrees, 100 parts of antimony; when the whole is in fusion, throw on it, by degrees, 20 parts of nitrate of potash, and after a few minutes quiet fusion, pour it into an iron melting cone, previously heated and greased.

2. It may also be obtained by melting eight parts of the ore mixed with six of nitrate of potash, and three of acidulous tartaric of potash, gradually projected into a red-hot crucible, and fused.

To obtain perfectly pure antimony, Margraaf melted some pounds of the sulphureated ore in a luted crucible, and thus scorified any metals it might contain. Of the antimony thus purified, which lay at the bottom, he took sixteen ounces, which he oxidated cautiously, first with a slow, and afterwards with a strong heat, until it ceases to smell of sulphur, and acquired a grayish white colour. Of this gray powder he took four ounces, mixed them with six drachms of acidulous tartaric of potash, and three of charcoal, and kept them in fusion in a well covered and luted crucible, for one hour, and thus obtained a metallic button that weighed one ounce, seven drachms, and twenty grains.

The metal, thus obtained, he mixed with half its weight of desiccated carbonate of soda, and covered the mixture with the same quantity of the carbonate. He then melted it in a well covered and luted crucible, in a very strong heat, for half an hour, and thus obtained a button which weighed one ounce, six drachms, and seven grains, much whiter and more beautiful than the

former. This he again treated with one and a half ounce of carbonate of soda, and obtained a button, weighing one ounce, five drachms, and six grains. The button was still purer than the foregoing. Repeating these fusions with equal weights of carbonate of soda three times more, and an hour and a half each time, he at last obtained a button so pure as to amalgamate with mercury with ease, very hard, and in some degree malleable; the scoriz formed in the last fusion were transparent, which indicated that they contained no sulphur, and hence it is the obstinate adherence of the sulphur that renders the purification of this metal so difficult.

The preparations of antimony formerly in use were very many: those now directed to be kept are:—

1. Sulphuretum antimonii.
2. Oxydum antimonii.
3. Sulphuretum antimonii præcipitatum.
4. Antimonium tartarisatum.
5. Pulvis antimonialis.
6. Liquor antimonii tartarisati.

ANTIMORIS. (From *anti*, against, and *moros*, death, or disease.) A medicine to prolong life.

ANTINEPHRITICA. (From *anti*, against, and *nephr*, a disease of the kidneys.) Remedies against disorders of the kidneys. *Blancard*.

ANTI-DONTALGICUS. An insect described by Gerbi in a small work published at Florence 1794, so called from its property of allaying the tooth-ach. It is a kind of curculio found on a species of thistle, *carduus spinosissimus*. If twelve or fifteen of these insects in the state of larvæ, or when come to perfection, be bruised and rubbed slowly between the fore-finger and thumb until they have lost their moisture; and if the painful tooth where it is hollow, be touched with that finger, the pain ceases sometimes instantaneously. A piece of shamoy leather will answer the same purpose with the finger. If the gums are inflamed the remedy is of no avail. Other insects possess the property of curing the tooth-ach; such as the scarabeus ferrugineus of Fabricius; the coccinella septempunctata, or lady-bird; the chrysomela populi, and the chrysomela sanguinolenta. This property belongs to several kinds of the coleoptera.

ANTIPARALYTICA. (From *anti*, against, and *paralytic*, the palsy.) Medicines against the palsy.

ANTIPATHIA. (From *anti*, against, and *pathos*, an affection.) Antipathy. An aversion to particular objects.

ANTIPERISTALTIC. (From *anti*, against, and *peristaltic*, to contract.) Whatsoever obstructs the peristaltic motion of the intestines.

ANTIPERISTASIS. (From *anti*, against, and *peristasis*, to press.) A compression on all sides. *Theophrastus de igne*. Digitized by Google

ANTIPHARMACA. (From *ἀντί*, against, and *φάρμακον*, a poison.) The same as alexipharmaca. Remedies or preservatives against poison. *Dioscorides*.

ANTIPHLOGISTICA. (From *ἀντί*, against, and *φlogē*, to burn.) Antiphlogistics. A term applied to those medicines, plans of diet, and other circumstances, which tend to oppose inflammation, or which in other words, weaken the system by diminishing the activity of the vital power.

ANTIPHTHISICA. (From *ἀντί*, against, and *φθίσις*, a consumption.) Remedies against a consumption.

ANTIPHTHORA. (From *ἀντί*, against, and *φθορά*, corruption.) A species of wolfsbane which resists corruption.

ANTIPHTISICA. (From *ἀντί*, against, and *φύσας*, to blow.) Carminatives or remedies against wind.

ANTIPLEURITICA. (From *ἀντί*, against, and *πleuritis*, pleurisy.) Remedies against a pleurisy.

ANTIPODAGRICA. (From *ἀντί*, against, and *ποδάγρα*, the gout.) Medicines which relieve or remove the gout.

ANTIPTAXIA. (From *ἀντί*, against, and *πρᾶξις*, to work.) A contrariety of functions and temperaments in divers parts. Contrariety of symptoms.

ANTIPTYRETICA. (From *ἀντί*, against, and *πυρετός*, fever.) *Antifebrile*. Remedies against a fever.

ANTIQUARTANARIA. (From *ἀντί*, against, and *quartanum*, a quartan fever.) Remedies against quartan agues.

ANTIQUARTICUM. The same as Antiquartanaria.

ANTIARRHINUM. (*Ἀντίρρινος*: from *ἀντί*, against, and *ῥίς*, the nose; so called because it represents the nose of a calf.) Snap-dragon, or calf's-snout. The name of a genus of plants in the Linnæan system. Class, *Didynamia*. Order, *Angiospermia*.

ANTIRRHINUM LINARIA. The systematic name for the *linaria* of the pharmacopœias. See *Linaria*.

ANTISCOLICA. (From *ἀντί*, against, and *σκολῆξ*, a worm. Remedies against worms. Anthelmintics.

ANTISCORBUTICS. (*Antiscorbutica*, sc. *medicamenta*; from *ἀντί*, against, and *scorbutus*, the scurvy.) Medicines which cure the scurvy.

ANTISEPTICS. (*Antiseptica*, sc. *medicamenta*; from *ἀντί*, against, and *σῆναι*, to putrefy.) Those medicines which possess a power of preventing animal substances from passing into a state of putrefaction, and of obviating putrefaction when already begun. This class of medicine comprehends four orders.

1. *Tonic antiseptics*, as cinchona, angustura cortex, chamæmelum, &c. which are suited for every condition of body, and are, in general, preferable to other antiseptics, for those with relaxed habits.

2. *Refrigerating antiseptics*, as acids, which are principally adapted for the young, vigorous, and plethoric.

3. *Stimulating antiseptics*, as wine and alcohol, best adapted for the old and debilitated.

4. *Antispasmodic antiseptics*, as camphora and assafoetida, which are to be selected for irritable and hysterical habits.

ANTISPASIS. (From *ἀντί*, against, and *σπᾶσις*, to draw.) A revulsion. The turning the course of the humours, whilst they are actually in motion. *Galen*.

ANTISPASMODICS. (*Antispasmodica*, sc. *medicamenta*; from *ἀντί*, against, and *σπᾶσις*, a spasm.) Medicines which possess the power of allaying, or removing inordinate motions in the system, particularly those involuntary contractions which take place in muscles, naturally subject to the command of the will. Spasms may arise from various causes. One of the most frequent is a strong irritation, continually applied; such as dentition, or worms. In these cases, narcotics prove useful, by diminishing irritability and sensibility. Sometimes spasm arises from mere debility; and the obvious means of removing this is by the use of tonics. Both narcotics and tonics, therefore, are occasionally useful as antispasmodics, such as opium, camphor, and ether, in the one class, and zinc, mercury, and Peruvian bark, in the other. But there are farther, several other substances, which cannot be with propriety referred to either of these classes; and to these, the title of antispasmodics is more exclusively appropriated. The principal antispasmodics, properly so called, are moschus, castereum, oleum animale empyreumaticum, petroleum, ammonia, assafoetida, sagapenum, galbanum, valeriana, crocus, melaleuca leucadendron.

The narcotics, used as antispasmodics, are ether, opium, camphor.

Tonics used as antispasmodic, are cuprum, zincum, hydragryrus, cinchona.

ANTITRAXAR. (From *ἀντί*, against, and *τράχας*, the palm of the hand.) A muscle of the foot. See *Adductor pollicis pedis*.

ANTITRAGICUS. *Antitragus*. (*Antitragicus*, sc. *musculus*.) One of the proper muscles of the ear, whose use is to turn up the tip of the antitragus a little outwards, and to depress the extremity of the antithelix towards it.

ANTITRAGUS. (*Antitragus*, i. m. from *ἀντί*, and *τράχας*, the tragus.) An eminence of the outer ear, opposite to the tragus.

ANTIVENEREÆ. (From *ἀντί*, against, and *venereus*, venereal.) Medicines against the lues venerea.

ANTONI SANCTI IONIS. (So called because St. Anthony was supposed to cure it miraculously. In the Roman Missal, St. Anthony is implored as being the pre-

server from all sorts of fire.) St. Anthony's fire. See *Erysipelas*.

ANT-PHYLLON. (From *αντι*, against, and *φυλλον*, a leaf; so called because its leaves are opposite.) The male caryophyllus.

ANTRUM OF HIGMORE. (From the name of an anatomist, who gave the first accurate description of it.) *Antrum Highmorianum.* *Antrum genu Sinus maxillaris pituitarius.* *Antrum maxilla superioris.* Maxillary sinus. A large cavity in the middle of each superior maxillary bone, between the eye and the roof of the mouth, lined by the mucous membrane of the nose.

One or both antra are liable to several morbid affections. Sometimes their membranous lining inflames, and secretes pus. At other times, in consequence of inflammation, or other causes, various excrescences and fungi are produced in them. Their bony parietes are occasionally affected with exostosis, or caries. Extraneous bodies may be lodged in them, and it is even asserted that insects may be generated in them, and cause for many years, afflicting pains. Abscesses in the antrum are by far the most common. Violent blows on the cheek, inflammatory affections of the adjacent parts, and especially of the pituitary membrane lining the nostrils, exposure to cold and damp, and, above all things, bad teeth, may induce inflammation and suppuration in the antrum. The first symptom is a pain, at first imagined to be a tooth-ach, particularly if there should be a carious tooth at this part of the jaw. This pain, however, extends more into the nose than that usually does which arises from a decayed tooth; it also affects, more or less, the eye, the orbit, and the situation of the frontal sinuses. But even such symptoms are insufficient to characterize the disease, the nature of which is not unequivocally evinced till a much later period. The complaint is, in general, of much longer duration than one entirely dependent on a caries of the tooth, and its violence increases more and more, until at last a hard tumour becomes perceptible below the cheek-bone. The swelling by degrees extends over the whole cheek; but it afterwards rises to a point, and forms a very circumscribed hardness, which may be felt above the back-grinders. The symptom is accompanied by redness, and sometimes by inflammation and suppuration of the external parts. It is not uncommon also, for the outward abscess to communicate with that within the antrum. The circumscribed elevation of the tumour, however, does not occur in all cases. There are instances in which the matter makes its way towards the palate, causing the bones of the part to swell, and at length rendering them carious,

unless timely assistance be given. There are other cases, in which the matter escapes between the fangs and sockets of the teeth. Lastly, there are other examples, in which matter, formed in the antrum, makes its exit at the nostril of the same side, when the patient is lying with his head on the opposite one, in a low position. If this mode of evacuation should be frequently repeated, it prevents the tumour both from pointing externally, and bursting, as it would do if the purulent matter could find no other vent. This evacuation of the pus from the nostril is not very common.

ANTRUM UCCINOSUM. The cochlea of the ear.

ANTRUM PYLORI. The great concavity of the stomach approaching the pylorus.

ANTRUM MAXILLARE. See *Antrum of Higmore*.

Ante, acid of. See *Formic acid*.

ANTILION. (From *Antyllus*, its inventor.) An astringent application, recommended by Paulus Ægineta.

ANUS. (*Quasi onus*; as carrying the burden of the bowels.)

1. The fundament; the lower extremity of the great intestine, named the rectum, is so called; and its office is to form an outlet for the fæces. The anus is furnished with muscles which are peculiar to it, viz. the *sphincter*, which forms a broad circular band of fibres, and keeps it habitually closed, and the *levatoris ani*, which serve to dilate and draw it up to its natural situation, after the expulsion of the fæces. It is also surrounded, as well as the whole of the neighbouring intestine, with muscular fibres, and a very loose sort of cellular substance. The anus is subject to various diseases, especially piles, ulceration, abscesses, excrescences, prolapsus, imperforation in new-born infants.

2. The term *anus* is also applied to a small opening of the third ventricle of the brain, which leads into the fourth.

ANUS, ARTIFICIAL. An accidental opening in the parietes of the abdomen, to which opening some part of the intestinal canal leads, and through which the fæces are either wholly or in part discharged. When a strangulated hernia occurs, in which the intestine is simply pinched, and this event is unknown; when it has not been relieved by the usual means; or when the necessary operation has not been practised in time; the protruded part becomes gangrenous, and the fæces escape. But if the patient should be at last operated upon, his fæces are discharged through the wound, and the intestines are more easily emptied. In both cases, the excrement continues to be discharged from the artificial opening. In this way an artificial anus is formed, through which the excrement is evacuated during life.

ANTIDION. (From *αντι*, priv. and *διον*,

water; so called, because they who eat of it become thirsty.) A species of nightshade, according to Blancart.

ANIPRUTHNOS. (From *a*, neg. and *pruthnos*, hurtful.) Hippocrates, in his Precepts, uses this word to signify an accidental event, which cannot be charged on the physician, and for which he is not accountable.

AORTA. (From *aoz*, air, and *trapez* to keep; so called because the ancients supposed that only air was contained in it.) The great artery of the body, which arises from the left ventricle of the heart, forms a curvature in the chest, and descends into the abdomen. See *Artery*.

APALACHINE GALLIS. (From *απαλαχος*, to repel; because it is supposed to repel infection.) See *Cassine*.

APARTHROSIS. (From *απο* and *αρθρον*, a joint) Articulation.

APARINE. (From *grn*, a file; because its bark is rough, and rasps like a file.) *Philanthropus*. *Ampelocarpus*. *Omphalocarpus* *laxus*. *Aspurine*. *Asperula*. Goose-grass and cleaver's beers. Cleaver's. Goose-share. Hayrifi. This plant, which is common in our hedges and ditches, is the *Galium aparine* of Linnæus:—*foliis otoniis lanceolatis carinatis scabris retrosum aculeatis, geniculis venosis, fructu hispidis*. The expressed juice has been given with advantage as an aperient and diuretic in insipient dropsies; but the character in which it has of late been chiefly noticed, is that of a remedy against cancer. A tea-cup full internally, gradually increases to half a pint, two or three times a day, and the herb applied, in cataplasm, externally, has been said to cure cancers. Such beneficial results are not confirmed by the experience of others.

APELLA. (From *a*, priv. and *pellis*, skin.) Shortness of the prepuce. Galen gives this name to all whose prepuce, either through disease, section, or otherwise, will not cover the glans.

APERIENS PALPEBRARUM RECTUS. See *Levator palpebre superioris*.

APERIENTS. (*Aperientia*, sc. *medicamenta*; from *aperio*, to open.) Laxatives. Medicines which gently open the bowels.

APERISTATUS. (From *a*, neg. and *περιστοι*, to surround.) *Aperistaton*. An epithet used by Galen, of an ulcer which is not dangerous, nor surrounded by inflammation.

APERISTATION. See *Aperistatus*.

APERTOR OCULI. See *Levator palpebre superioris*.

APESIAL. (*Ἀπεσια*, *a*, f. *απεψια*: from *a*, priv. and *πεσσω*, to digest.) Indigestion. See *Dyspepsia*.

APRUTHNENUS. (From *απο* and *αδω*, straight.) A name formerly given to the intestinum rectum, or straight gut.

APEX. The extremity of a part; as the apex of the tongue, apex of the nose, &c.

APHANISUS. (From *αφανίζω*, to remove from the sight.) The removal, or gradual decay of a disorder.

APHERRIS. (From *αφαιρω*, to remove.) This term was formerly much used in the schools of surgery, to signify that part of the art which consists in taking off any diseased or preternatural part of the body.

APHERSENA. (From *απο*, and *εβα*, to boil.) A decoction.

APRESIS. (From *απαρμι*, to remit.) The remission or termination of a disorder.

APHISTE-IS. (From *αφιστημι*, to draw from.) An abscess.

APHODOS. (From *απο*, and *δω*, departure.) Excrement. The dejection of the body.

APHONIA. (*Aphonia*: from *a*, priv. and *φωνη*, the voice.) A suppression of the voice, without either syncope or coma. A genus of disease in the class *locales*, and order *dyscinesie* of Cullen.

When it takes place from a tumour of the fauces, or about the glottis, it is termed *aphonia gutturalis*;

When from a disease of the trachea, *aphonia trachealis*;

And when from a paralysis, or want of nervous energy, *aphonia atonica*;

APHORISM. (*Aphorismus*; from *απορρίζω*, to distinguish.) A maxim, or principle, comprehended in a short sentence.

APHRODISIA. (From *Αφροδιτη*, Venus.) An immoderate desire of venery.

APIRODISIACS. (*Aphrodisiaca*, sc. *medicamenta aphrodisiaca*; from *αφροδιτη*, venery.) Medicines which excite a desire for venery.

APHRODISIASTICON. (From *αφερο*, froth.) A troch so called by Galen, because it was given in dysenteries, where the stools were frothy.

APHRODISIUS MORBUS. (From *Αφροδιτη*, Venus.) The venereal disease.

APHTHA. See *Aphtha*.

APHITHÆ. (*Aphæ*: from *απθω*, to inflame.) The thrush. Frog, or sore mouth. *Aphtha lictucamen* of Sauvages. *Ulcera serpentia oris*, or spreading ulcers in the mouth, of Celsus. *Pustula oris*. *Alcola*. *Venicle gingivarum*. *Acacos Aphtha infantum*. A disease to which children are very subject. It appears in small white ulcers upon the tongue, gums, and around the mouth and palate, resembling small particles of curdled milk. When the disease is mild, it is confined to these parts; but when it is violent and of long standing, it is apt to extend through the whole course of the alimentary canal, from the mouth down to the anus; and so to excite severe purgings, flatulencies, and other disagreeable symptoms. The dis-

ease, when recent and confined to the mouth, may in general be easily removed; but when of long standing, and extending down to the stomach and intestines, it very frequently proves fatal.

The thrush sometimes occurs, as a chronic disease, but in warm climates and in those Northern countries where the cold is combined with a considerable degree of moisture, or where the soil is of a very marshy nature. It may in some cases, be considered as an idiopathic affection; but it is more usually symptomatic. It shews itself, at first, by an uneasy sensation, or burning heat in the stomach, which comes on by slow degrees, and increases gradually in violence. After some time, small pimples, of about the size of a pin's head, shew themselves on the tip and edges of the tongue; and these, at length, spread over the whole inside of the mouth, and occasion such a tenderness and rawness, that the patient cannot take any food of a solid nature; neither can he receive any vinous or spirituous liquor into his mouth, without great pungency and pain being excited; little febrile heat attends, with dry skin, pale countenance, small pulse, and cold extremities. These symptoms will probably continue for some weeks, the general health being sometimes better, and sometimes worse, and then the patient will be attacked with acid eructations, or severe purging, which greatly exhausts his strength, and produces considerable emaciation of the whole body. After a little time, these symptoms cease, and he again enjoys better health; but, sooner or later, the acrid matter shews itself once more in the mouth, with greater virulence than before, and makes frequent translations to the stomach and intestines, and so from these to the mouth again, until at last, the patient is reduced to a perfect skeleton. Elderly people, and persons with a shattered constitution, are more liable to its attacks. It is ranked by Cullen in the class *pyrexia*, and order *exanthemata*.

APIUM (From *apis*, Doric. *apios*, mild; or from *apes*, bees; because they are fond of it.

1. The name of a genus of plants in the Linnean system. Class, *Pentandria*. Order, *Digynia*.

2. The pharmacopœial name of the herb smallage. The root, seeds, and fresh plant; *Apium graveolens, foliolis caulinis, cuneiformibus, umbellis, sessilibus*, of Linneus, are aperient and carminative.

APIUM GRAVEOLENS. The systematic name for the *apium* of the pharmacopœias. See *Apium*.

APIUM PETROSELINUM. The systematic name for the *petroselinum* of the pharmacopœias. See *Petroselinum*.

APNEUSTIA. (From *a*, neg. and *pnos*, to

breathe.) A defect or difficulty of respiration, such as happens in a cold, &c. *Fæsius*.

APNOËA. The same. *Galen*.

APOCARNISUS. (From *apo*, and *καρνος*, smoke.) A fumigation.

APOCATHARSIS. (From *apo*, and *καθαίρω*, to purge.) An evacuation of humours; a discharge downward; but sometimes applied, with little discrimination, to vomiting.

APOCAULISIS. (From *αποκαυλίζω*, to break transversely.) A transverse fracture. *Hippocrates*.

APOCENOSIS. (From *apo*, and *κένω*, to evacuate.) A superabundant flux of blood, or other fluid, without pyrexia. The name of an order in the class *locales* of Cullen.

APOCOPE. (From *apo*, and *κοπέω*, to cut from.) Abscission, or the removal of a part by cutting it off.

APOCRISIS. (From *apo*, and *κρίνω*, to secrete from.) A secretion of superabundant humours. *Hippocrates*.

APOCRUSTINUM. *Apocrusticon*. (From *αποκρίνω*, to repel.) An astringent or repellent medicine. *Galen*.

APOCRUSTICON. See *Apocrustinum*.

APOCYISIS. (From *apo*, and *κυάω*, to bring forth.) Parturition, or the bringing forth of a child. *Galen*.

APODACHTYICA. (From *apo*, and *δακρυάω*, a tear.) Medicines which, by exciting tears, remove superfluous humours from the eyes, as onions, &c. *Pliny*.

APOGEUSIA. See *Aghæusia*.

APOGEUSIS. See *Aghæusia*.

APOBINOMESIS. (From *απονομομαι*, to be absent.) The remission or absence of a disease. *Hippocrates*.

APOBLAUTOSIS. (From *apo*, and *γλαυκός*, sky-coloured; so called because of its blueish appearance.) *Glaucoma*. A cataract of the eye. *Dioscorides*.

APOGONUM. (From *apo*, and *γονιμύω*, to beget.) A living fœtus in the womb. *Hippocrates*.

APOLEPSIS. (From *apo*, and *λαμβάνω*, to take from.) An interception, suppression, or retention of urine, or any other natural evacuation. *Hippocrates*.

APOLINOSIS. (From *apo*, and *ληνός*, flax.) The method of curing a fistula, according to Ægineta, by the application of raw flax.

APOLYSIS. (From *apo*, and *λύω*, to release.) The solution or termination of a disease. The removal of a bandage. *Eretianus*.

APOMAGNA. (From *apo*, and *μαγνέω*, to cleanse from.) Any thing used to cleanse and wipe away filth from sores, as a sponge, &c. *Hippocrates*.

APONATHENA. (From *apo*, neg. and *μαθάνω*, to learn.) *Hippocrates* expresses,

by this term, a forgetfulness of all that has been learnt.

APOMELI. (From *απο*, from, and *μελι*, honey.) An oxymel, or decoction, made with honey.

APONEUROSIS. (From *απο*, and *νευρον*, a nerve; from an erroneous supposition of the ancients, that it was formed by the expansion of a nerve.) A tendinous expansion. See *Muscle*.

AFONIA. (From *α*, priv. and *φωνε*, pain.) Freedom from pain.

AFONITROSIS. (From *απο*, and *νιτρον*, nitre.) The sprinkling an ulcer over with nitre.

AFOPALLESIS. (From *αποπαλλαι*, to throw off hastily.) An abortion, or premature expulsion of a foetus. *Hippocrates*.

AFOPEDASSIS. (From *απο*, and *σπιδαν*, to jump from.) A luxation.

AFOPHLEGMASIA. (From *απο*, and *φλεγμα*, phlegm.) A discharge of phlegm, or mucus.

AFOPHLEGMATICA. (From *απο*, and *φλεγμα*, phlegm.) *Apophlegmatizantia.* *Apophlegmatizantia.* Medicines which excite the secretion of mucus from the mouth and nose. Masticatories. *Er-rhines*.

AFOPHRAXIS. (From *απο*, and *φρασσα*, to interrupt.) A suppression of the menstrual discharge.

AFOPHTHARMA. (From *απο*, and *φθορα*, to corrupt.) A medicine to procure abortion.

AFOPHTHORA. (From *αποφθορα*, to be abortive.) An abortion.

AFOPHTAS. (From *αποφθα*, to proceed from.) Any thing which grows or adheres to another, as a wart to the finger.

AFOPHTADES. The ramifications of the veins and arteries. *Hippocrates*.

AFOPHYSIS. (From *αποφθα*, to proceed from.) *Appendix. Probole. Echphysis. Processus. Productio. Projectura. Protuberantia.* A process, projection, or protuberance, of a bone beyond a plain surface; as the nasal apophysis of the frontal bone, &c.

AFOPHTHEGMA. (From *αποφθεγγμαι*, to speak eloquently.) A short maxim, or axiom; a rule.

AFOPLECTA. A name formerly applied to the internal jugular vein; so called because, in apoplexies, it appears full and turgid. *Bartholin*.

AFOPLECTICA. (From *αποπληξια*, an apoplexy.) Medicines against an apoplexy.

APOPLEXIA. (From *απο*, and *πλησσω*, to strike or knock down; because persons, when seized with this disease, fall down suddenly.) Apoplexy. A sudden abolition, in some degree, of the powers of sense and motion, with sleep, and sometimes snoring; the respiration and motion

of the heart remaining. Cullen arranges it in the class *neuroses*, and order *comata*.

When it takes place from a congestion of blood, it is termed *apoplexia sanguinea*.

2. When there is an abundance of serum, as in persons of a cold temperament, *apoplexia serosa*.

3. If it arise from water in the ventricles of the brain, it is called *Apoplexia hydropthalica*.

4. If from a wound, *apoplexia traumatica*.

5. If from poisons, *apoplexia venenata*.

6. If from the action of suffocating exhalations, *apoplexia suffocate*.

7. If from passions of the mind, *apoplexia mentalis*.

8. And when it is joined with catalepsy, *apoplexia cataleptica*.

Apoplexy makes its attack chiefly at an advanced period of life; and most usually on those who are of a corpulent habit, with a short neck, and large head; and who lead an inactive life, make use of a full diet, or drink to excess. The immediate cause of apoplexy, is a compression on the brain, produced either by an accumulation of blood in the vessels of the head, and distending them to such a degree, as to compress the medullary portion of the brain; or by an effusion of blood from the red vessels, or of serum from the exhalants; which fluids are accumulated in such a quantity as to occasion compression. The former of these is called a *sanguineous* and the latter a *serous* apoplexy. These states, of over distinction and of effusion, may be brought on by whatever increases the afflux and impetus of the blood in the arteries of the head; such as violent fits of passion, great exertions of muscular strength, severe exercise, excess in venery, stooping down for any length of time, wearing any thing too tight about the neck, overloading the stomach, long exposure to excessive cold, or a vertice sun, the sudden suppression of any long-accustomed evacuation, the application of the fumes of certain narcotic and metallic substances, such as opium, alcohol, charcoal, mercury, &c. and by blows, wounds, and other external injuries: in short, apoplexy may be produced by whatever determines too great a flow of blood to the brain, or prevents its free return to it.

The young, and those of a full plethoric habit, are most liable to attacks of the sanguineous apoplexy; and those of a phlegmatic constitution, or who are much advanced in life, to the serous. Apoplexy is sometimes preceded by headach, giddiness, dimness of sight, loss of memory, faltering of the tongue in speaking, numbness in the extremities, drowsiness, stupor, and night-mare, all denoting an affection of the brain; but it more usually happens that,

without much previous indisposition, the person falls down suddenly, the countenance becomes florid, the face appears swelled and puffed up, the vessels of the head, particularly of the neck and temples, seem turgid and distended with blood; the eyes are prominent and fixed, the breathing is difficult, and performed with a snorting noise, and the pulse is strong and full. Although the whole body is affected with the loss of sense and motion, it nevertheless takes place often more upon one side than the other, which is called hemiplagia, and in this case, the side least affected with palsy is somewhat convulsed.

In forming an opinion as to the event, we must be guided by the violence of the symptoms. If the fit is of long duration, the respiration laborious and stertorous, and the person much advanced in years, the disease, in all probability, will terminate fatally. In some cases, it goes off entirely; but it more frequently leaves a state of mental imbecility behind it, or terminates in a hemiplegia, or in death. Even when an attack is recovered from, it most frequently returns again, after a short period of time, and in the end proves fatal. In dissections of apoplexy, blood is often found effused on the surface and in the cavities of the brain; and in other instances, a turgidity and distention of the blood-vessels are to be observed. In some cases, tumours have been found attached to different parts of the substance of the brain, and in others, no traces of any real affection of it could be observed.

APOPHYSIS. (From *ἀποφύω*, to suffocate.) A suffocation. *Mosclion*.

APOPSPHESIS. (From *ἀπο*, and *σφωω*, to emit wind.) The emission of wind by the anus or uterus, according to Hippocrates.

APOPSYCHIA. (From *ἀπο*, from, and *ψυχή*, the mind.) The highest degree of delirium, or fainting, according to Galen.

APOPTOSIS. (From *ἀποπίπτω*, to fall down.) A prolapsus, or falling down of any part, through relaxation. *Erelian*.

APOREXIS. (From *ἀπο*, and *αἰρώ*, to stretch out.) A play with balls, in the gymnastic exercises.

APORE. (From *α*, priv. and *αἶρος*, a duct. Restlessness, uneasiness, occasioned by the interruption of perspiration, or any stoppage of the natural secretions.

APOREHISIS. (From *ἀπορρίπτω*, to cast off.) Hippocrates uses this word to signify that kind of insanity where the patient tears off his clothes, and casts them from him.

APOKEPARNISMUS. (From *ἀπο* from, and *κεραυνός*, to strike with a hatchet.) *Deacatio*. A species of fracture, when part of a bone is chipped off. *Gorræus*.

APOCHASIS. *Aposchasius*. (From *ἀπο*,

and *σχαζω*, to scarify.) A scarification. Venesection. *Hippocrates*.

APOSITIA. *Apositios*. (From *ἀπο*, from, and *εἶδος*, food.) A loathing of food. *Galen*.

APOSPASMA. (From *ἀσπασμα*, to tear off.) A violent, irregular fracture of a tendon, ligament, &c. *Galen*.

APOSPHACELIS. (From *ἀπο*, and *σφακελος*, a mortification.) Hippocrates uses this word to denote a mortification of the flesh in wounds, or fractures, caused by too tight a bandage.

APOSTANIS. (From *ἀπο*, and *ιστημι*, to recede from.)

1. An abscess, or collection of matter.

2. The coming away of a fragment of bone, by fracture.

3. When a distemper passes away by some outlet, Hippocrates calls it an *apostasis* or *excretion*.

4. When the morbid matter, by its own weight, falls and settles on any part, an *apostasis* by settlement.

5. When one disease turns to another, an *apostasis* by metastasis.

APOSTAXIS. (From *ἀποσάω*, to distil from.) Hippocrates uses this word to express the defluxion or distillation of any humor, or fluid: as blood from the nose.

APOSTEMA. (From *ἀπιστῆμι*, to recede.) The term given by the ancients to abscesses in general. See *Abscessus*.

APOSTEMATIA. Those who, from an inward abscess, void pus downwards, are thus called by Aretæus.

APOSTRIGMA. (From *ἀποσθίγω*, fulsio.) Galen uses this word to denote a rest of a diseased part, a cushion.

APOSTOLORUM UNGENTUM. (From *ἀποστόλος*, an apostle.) *Dodecapharmacum*. The apostles' ointment; so called because it has twelve ingredients in it, exclusive of the oil and vinegar.

APOSTROME. (From *ἀπο* and *στροφή*, to turn from.) Thus Paulus Ægineta expresses an aversion for food.

APOSTRINGESIS. (From *ἀπο* and *σφίγγω*, a fistula.) The degeneracy of a sore into a fistula. *Hippocrates*.

APOSYMA. (From *ἀπο* and *σύνω*, to rub off.) An abrasion or desquamation of the bone, or skin. *Hippocrates*.

APOTANUSIS. (From *ἀπο* and *τείνω*, to extend.) An extension, or elongation, of any member or substance.

APOTELEISIS. (From *ἀπο* and *τέλειω*, a bog.) An expurgation of filth, or fæces.

APOTHECA. (*Αποθήκη*: from *ἀποθηκεύω*, to reposit.) A shop, or vessel, where medicines are sold or deposited.

APOTHECARIUS. (From *ἀπο*, and *θήκη*, *pono*, to put; so called from his employ being to prepare, and keep in readiness, the various articles in the *Materia Medica*, and to compound them for

the physician's use; or from *αποθήκη*, a shop.) An apothecary. In every European country, except Great Britain, the apothecary is the same as, in England, we name the druggist and chemist.

APOTHEKEIA. (From *apo* and *θεραπεύω*, to cure.) A perfect cure, according to Hippocrates.

APOTHEKEUTICA. (From *apo* and *θεραπεύω*, to heal.) Therapeutics; that part of medicine which teaches the art of curing disorders.

APOTHEKON. (From *apo* and *θεῖον*, heat.) An acrimonious pickle, with mustard, vinegar, and oil. *Galen.*

APOTHEKIS. (From *apo* and *τομή*, to replace.) The reduction of a dislocated bone, according to Hippocrates.

APOTHEKIS. (From *apo* and *θεῖον*, to press from.) The dregs or expressed juice of a plant.

APOTHEKIS. (From *apo* and *θεῖον*, to break.) Apocope. The taking away the splinters of a broken bone.

APOTHEKIS. (From *apo* and *τομή*, to bring forth.) Abortive; premature. *Hippocrates.*

APOTHEKIS. (From *apo* and *τομή*, to turn from.) A resolution or reversion of a suppurating tumour.

APOTHEKIS. (From *apo* and *τομή*, to avert.) An amulet, or charm, to avert diseases. *Plinius.*

APOTHEKIS. *Apothema.* (From *apo* and *θεῖον*, to boil.) A decoction.

APOTHEKIS. (From *apo* and *ζωή*, to separate.) The separation or removal of morbid parts. *Hippocrates.*

APOTHEKIS. (From *apo* and *ζωή*, fermented.) Fermented.

APPARATUS. (From *appareo*, to appear, or be ready at hand.) This term implies the preparation and arrangement of every thing necessary in the performance of an operation, or in the application of dressings. The apparatus varies according to circumstances. Instruments, machines, bandages, tapes, compresses, pledgets, dossils of lint, tents, &c. are parts of the apparatus, as well as any medical substances used. It is a rule in surgery to have the apparatus ready before beginning an operation. All preparations of this kind should not be made in the patient's room, when they can be avoided, nor any where in his presence, as it would agitate him, and render him timid and more restless in the operation.

APPARATUS MINOR. See *Lithotomy.*

APPARATUS MAJOR. See *Lithotomy.*

APPARATUS ALTUS. See *Lithotomy.*

APPENDICULA CECI VERMIFORMIS. A vermicular process, about four inches in length, and the size of a goose-quill, which hangs to the intestinum cæcum of the human body.

APPENDICULA EXFLOICA. *Appendices*

coli adipose. The small appendices of the colon and rectum, which are filled with adipose substance. See *Omentum.*

Apple, thorn. See *Stramenium.*

APPLE. The common crabtree, *pyrus malus* of Linnaeus, is the parent of all the vast variety of apples at present cultivated. Apples, in general, when ripe, afford a pleasant and easily digestible fruit for the table; but, when the stomach is weak, they are very apt to remain unaltered for some days, and to produce dyspepsia. Sour fruits are to be considered as unwholesome, except when boiled or baked, and rendered soft and mellow by the addition of sugar.

APRICOT. The fruit of the *Prunus armeniaca* of Linnaeus. When ripe, they are easily digested, and are considered as a pleasant and nutritious delicacy.

APYREXIA. (From *α*, priv. and *πύρεξια*, a fever.) Apyrexia. Without fever. The intermission of feverish heat.

AQUA. See *Water.*

AQUA AERIS FIXI. Water impregnated with fixed air. This is liquid carbonic acid, or water impregnated with carbonic acid; it sparkles in the glass, has a pleasant acidulous taste, and forms an excellent beverage. It diminishes thirst, lessens the morbid heat of the body, and acts as a powerful diuretic. It is also an excellent remedy in increasing irritability of the stomach, as in advanced pregnancy, and it is one of the best anti-emetics which we possess.

AQUA ALUMINIS COMPOSITA. Compound solution of alum, formerly called aqua aluminosa bateana. See *Liquor aluminis compositus.*

AQUA AMMONIÆ ACETATÆ. See *Liquor ammoniæ acetatis.*

AQUA AMMONIÆ FURÆ. See *Liquor ammoniæ.*

AQUA ANETHI. See *Anethum.*

AQUA CALCIS. See *Liquor calcis.*

AQUA CÆLESTIS. A preparation of copper.

AQUA CARUI. See *Carui.*

AQUA CINNAMO. See *Cinnamomum.*

AQUA CUPRI AMMONIATÆ. See *Liquor cupri ammoniaci.*

AQUA CUPRI VITRIOLATÆ COMPOSITA. This preparation of the Edinburgh Pharmacopoeia, is used externally, to stop hæmorrhages of the nose, and other parts. It is made thus:

℞. Cupri vitriolati, aluminis, sing. ℥ss. Aquæ puræ, ℥iv. Acid. vitriolici, ℥ij.

Boil the salts in water until they are dissolved; then filter the liquor, and add the acid.

AQUA DISTILLATA. Distilled water. This is made by distilling water in clean vessels, until about two-thirds have come over. In nature, no water is found perfectly pure. Spring or river water always

contains a portion of saline matter, principally sulphate of lime; and, from this impregnation, is unfit for a number of pharmaceutical preparations. By distillation, a perfectly pure water is obtained. The London College directs ten gallons of common water: of which, first distil four pints, which are to be thrown away; then distil four gallons. This distilled water is to be kept in glass vessels. See *Water*.

AQUA FENICULI. See *Feniculum*.

AQUA FORTIS. See *Acidum nitrosum dilutum*.

AQUA KALI PRÆPARATI. See *Liquor subcarbonatis potasse*.

AQUA KALI PURI. See *Liquor potasse*.

AQUA LITHARGYRI ACETATI. See *Liquor acetatis plumbi*.

AQUA LITHARGYRI ACETATI COMPOSITA. See *Liquor plumbi acetatis dilutus*.

AQUA MENTHE PIPERITÆ. See *Mentha piperita*.

AQUA MENTHE SATIVÆ. See *Mentha viridis*.

AQUA MENTHE VIRIDIS. See *Mentha viridis*.

AQUA PIMENTÆ. See *Pimenta*.

AQUA PULEGII. See *Pulegium*.

AQUA REGIA. *Aqua regalis* The acid now called the nitro-muriatic, was formerly called aqua regalis, because it was, at that time, the only acid that could dissolve gold. See *Nitro-muriatic acid*.

AQUA ROSÆ. See *Rosa centifolia*.

AQUA STYPTICA. A name formerly given to a combination of powerful astringents, viz. sulphate of copper, sulphate of alum, and sulphuric acid. It has been applied topically to check hæmorrhage, and largely diluted with water, as a wash in purulent ophthalmia.

AQUA ZINCI VITRIOLATI CUM CAMPHORA. Otherwise named *Aqua stercoraria camphorata*. This, when properly diluted, is a useful collyrium for inflammations of the eyes, in which there is a weakness of the parts. Externally it is applied by surgeons to scorbutic and phagedenic ulcerations.

AQUE DISTILLATÆ. Distilled waters. These are made by introducing vegetables, as mint, penny-royal, &c. into a still with water; and drawing off as much as is found to possess the properties of the plants. The London College orders the waters to be distilled from dried herbs, because fresh are not ready at all times of the year. Whenever the fresh are used, the weights are to be increased. But, whether the fresh or dried herbs are employed, the operator may vary the weight according to the season in which they have been produced and collected. Herbs and seeds, kept beyond the space of a year, are improper for the distillation of waters. To every gallon of these waters, five ounces, by measure, of proof spirit are to be added.

AQUE MINERALES. See *Waters, mineral*.

AQUE STILLATITIE SIMPLICES. Simple distilled waters.

AQUE STILLATITIE SPIRITUOSÆ. Spirituous distilled waters, now called only spiritus, as spiritus pulegii.

AQUADUCT OF FALLOPIUS. A canal in the petrous portion of the temporal bone, first accurately described by Fallopius.

Aquatic nut. See *Tribulus equalicus*.

AQUEOUS HUMOUR OF THE EYE. The very limpid watery fluid, which fills both chambers of the eye. See *Eye*.

AQUETTA. The name of a liquid poison, made use of by the Roman women, under the Pontificate of Alexander VII. It was prepared, and sold in drops, by Tophania, or Toffania, an infamous woman who resided at Palermo, and afterwards at Naples. From her, these drops obtained the name of *Aqua Toffania*, *Aqua della Toffania*, and also *Aqua di Napoli*. This poison is said, by some, to be a composition of arsenic, and by others opium and cantharides.

AQUROLIUM. (From *acus*, a needle, and *folium*, a leaf; so called on account of its prickly leaf.) The leaves of this plant, *Ilex aquifolium*; *foliis ovatis acutis spinatis*, of Linnaeus, have been known to cure intermittent fevers; and an infusion of the leaves, drank as tea, is said to be a preventive against the gout.

AQUILA. A chymical name formerly used for sal-ammoniac, mercurius præcipitatus, arsenic, sulphur, and the philosopher's stone.

AQUILA ALBA. One of the names given to calomel by the ancients. See *Sublimis hydrargyri*.

AQUILA ALBA PHILOSOPHORUM. *Aqua alba ganymedis*. Sublimed Sal-ammoniac.

AQUILA CÆLESTIS. A panacea, or cure for all diseases; a preparation of mercury.

AQUILA VENERIS. A preparation of the ancients, made with verdigrise and sublimed sal-ammoniac.

AQUILA, among the ancients, had many other epithets joined with it, as rubra, salutifera, volans, &c.

AQUILÆ VENÆ. (From *aquila*, an eagle.) Branches of the jugular veins, which are particularly prominent in the eagle.

AQUILÆ LIGNUM. Eagle-wood. It is generally sold for the agalochum.

AQUILEGIA. (From *aqua*, water, and *lego*, to gather; so called from the shape of its leaves, which retain water.) The herb columbine.

1. The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Pentagynia*.

2. The name, in the Pharmacopœia, for the columbine.

The seeds, flowers, and the whole plant, *Aquilegia vulgaris*; *nectaritis incurvis*, of Linnaeus, have been used medicinally,

the first in exanthematous diseases, the latter chiefly is an antiscorbutic. Though retained in several foreign pharmacopœias, their utility seems to be not allowed in this country.

AQUULA. (Diminutive of *aqua*.) A small quantity of very fine and limpid water. This term is applied to the pellucid water, which distends the capsule, of the crystalline lens, and the lens itself. Paulus Ægineta uses it to denote a tumour consisting of a fatty substance under the skin of the eyelid.

Arabic gum. See *Acacia gummi*.

ARACALAN. Amulets.

ARACA MIRA. (Indian.) A shrub growing in the Brazils, whose roots are diuretic and antidysergetic.

ARACHNE. (From *arag*, Heb. to weave; or from *araxus*, a spider.) The spider.

ARACHNOID MEMBRANE. (From *araxus*, a spider, and *ωσος*, likeness; so named from its resemblance to a spider's web.) A thin membrane of the brain, without vessels and nerves, situated between the dura and pia mater, and surrounding the cerebrum, cerebellum, medulla oblongata, and medulla spinalis. The term is also applied by some writers to the tunic of the crystalline lens and vitreous humour of the eye.

ARACK. (Indian.) An Indian spirituous liquor, prepared in many ways, often from rice; sometimes from sugar, fermented with the juice of cocoa-nuts; frequently from toddy, the juice which flows from the cocoa-nut tree by incision, and from other substances.

ARADOS. From *arados*, to be turbulent.) Hippocrates uses this term to signify a commotion in the stomach, occasioned by the fermentation of its contents.

ARMETICA. (From *arare*, to rarefy.) Things which rarefy the fluids of the body.

ARALIA. (From *ara*, a band in the sea; so called because it grows upon banks, near the sea.) The berry-bearing angelica. Of the several species of this tree, the roots of the nudicaulis, or naked-stalked, were brought over from North America, where it grows, and sold here for sarsaparilla.

ARANEA. (From *arare*, to knit together.) The spider.

ARBOR VITÆ. The tree of life.

1. The cortical substance of the cerebellum is so disposed, that, when cut transversely, it appears ramified like a tree, from which circumstance it is termed *arbor vite*.

2. The name of a tree, the leaves and wood of which were formerly in high estimation as resolvents, sudorifics, and expectorants, and were given in phthisical affections intermittent fevers, and dropsies. It is the *Thuya occidentalis*; *strobilis levibus*, *squamis obtusis*, of Linnæus.

ARBUTUS. The name of a genus of

plants in the Linnæan system. Class, *Decandria*. Order, *Monogynia*. The strawberry tree.

ARBUTUS UVA URSI. The systematic name for the officinal trailing arbutus. See *Uva ursi*.

ARCA ARCANORUM. The mercury of the philosophers.

ARCA CORDIS. The pericardium.

ARCANUM. (A secret.) A medicine whose preparation, or efficacy, is kept from the world, to enhance its value. With the chymists, it is a thing secret and incorporeal; it can only be known by experience, for it is the virtue of every thing, which operates a thousand times more than the thing itself.

ARCANUM CATHOLICUM. Bezoar, planetain, and colchicum.

ARCANUM DUPLIX. *Arcanum duplicatum.* A name formerly given to the combination of potash and sulphuric acid, more commonly called vitriolated tartar, and now sulphat of potash.

ARCANUM TARTARI. The acetate of potash.

ARCENTHOS. Juniper.

ARCHE. *Arx*, the beginning.) The first stage or attack of a disease.

ARCHÆUS. The universal archæus, or principle of Van Helmont, was the active principle of the material world; it means good health also.

ARCHE. (From *arx*, the beginning.) The earliest stage of a disease.

ARCHENDA. (Arab.) A powder made of the leaves of the ligustrum, to check the fetid odour of the feet. Detergent.

ARCHEOSTIS. White briony.

ARCHIMASIA. (From *arx*, the chief, and *maga*, Arab. meditation.) Chymistry, as being the chief of sciences.

ARCHITHOLUS. (From *arx*, the chief, and *tholos*, a chamber. The sudatorium, or principal room of the ancient baths.

ARCHUS. (From *arx*, an arch.) The anus; so called from its shape.

ARCHOPTOMA. (From *arx*, the anus, and *optoma*, to fall down.) A bearing down of the rectum, or prolapsus ani.

ARCTATIO. (From *arctis*, to make narrow.) *Arctitudo*. Narrowness.

1. A constipation of the intestines, from inflammation.

2. A preternatural straitness of the pendulum muliebree.

ARCTIUM. (From *arctos*, a bear; so called from its roughness.) The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia æqualis*. The burdock.

ARCTIUM LAPPA. (*Lappa*; *αρο το λακκον*, from its seizing the garments of passengers.) The herb clotbur, or burdock. The systematic name for the *barduna*. See *Bardana*.

ARCTURA. (From *arctis*, to straiten.)

AN INFLAMMATION of the finger, or toe, from a curvature of the nail.

ARCUALIA. (From *arcus*, a bow.) *Ar. curialis*. The satura coronalis is so named, from its bow-like shape; and, for the same reason, the bones of the sinciput are called arcualia ossia. *Bartholin*.

ARGUATIO. (From *arcus*, a bow.) A gibbosity of the fore-parts, with a curvature of the sternum of the tibia, or dorsal vertebra. *Avicenna*.

ARCULE. A dim. of *arca*, a chest.) The orbits or sockets of the eyes.

ARDAS. (From *ardus*, to defile.) Filth, excrement, or refuse. *Hippocrates*.

Ardent Spirit. See *Alcohol*.

ARDOR FEBRILIS. Feverish heat.

ARDOR URINÆ. *Dysuria*. Scalding of the urine. Difficulty and pain in making water, attended with a sense of heat in the urethra. It is a symptom of gonorrhœa, and some other affections.

ARDOR VENTRICULI. Heartburn.

AREA. An empty space. That kind of baldness where the crown of the head is left naked, like the tonsure of a monk.

ARECA INDICA. An inferior kind of nutmeg.

AREGON. (From *ageron*, to help.) A resolvent ointment; so called from its valuable qualities.

AREMAROS. Cinnabar.

ARENA. Sand, or gravel.

ARENAMEL. (From *arena*, sand; so called because it was said to be procured from sandy places.) *Arenamen*. Bole-armenia.

ARENATIO. (From *arena*, sand.) Saburation, or the sprinkling of hot sand upon the bodies of patients. *Andr. Baccus de Therm.*

ARENTES. (From *areo*, to dry up.) A sort of ancient cupping-glasses, used without scarifying.

AREOLA. (A dim. of *area*, a void space.) A small brown circle, which surrounds the nipples of females. During and after pregnancy, it becomes considerably larger.

ARETE. (*Agera*, virtue.) Hippocrates uses this word to mean corporeal or mental vigour.

ARETENOIDES. See *Arytenoides*.

AREUS. A pessary, invented by *Ægineta*.

AREFAR. *Arsay*. Arsenic. *Ruland*, &c.

ARGAL. Argol. Crude tartar, in the state in which it is taken from the inside of wine-vessels, is known in the shops by this name.

ARGASYLLIS. (From *Agros*, a serpent: which it is said to resemble.) The plant which was supposed to produce gum-ammoniac.

ARGEMA. (From *Agros*, white.) *Argemon*. A small white ulcer of the globe of the eye. *Eretianus*, *Galen*, &c.

ARGENTI NITRAS. See *Nitras argenti*.

ARGENTUM. Silver. See *Silver*.

ARGENTUM NITRATUM. See *Nitras argenti*.

ARGENTUM VIVUM. It was formerly, by some, called *argentum mobile*, and *argentum fusum*. See *Hydrargyrum*.

AREES. (From *agros*, white.) A serpent, with a whitish skin, deemed by Hippocrates exceedingly venomous.

AREILLA. (From *agros*, white.) White clay; argil; potter's earth.

AREILLA VITRIOLATA. Alum.

AREYRITIS. (From *argyros*, silver.) Litharge, or spume of silver. A kind of earth was formerly so named, which is taken from silver mines, and is bespangled with many particles of silver.

ARGYROCOME. (From *argyros* silver, and *κωμα*, hair.) A sort of cudweed, or gnaphalium, was so named, from its white silvery floscules.

ARGYROLIBANOS. The white olibanum.

ARGYROPHORA. An antidote, in the composition of which there is silver.

ARGYROTROPHEMA. (From *argyros* white, and *τροφημα*, food.) A white cooling food, made with milk. Milk diet. *Galen*.

ARHEUMATISTOS. (From *a*, neg. and *ρευματιζω*, to be afflicted with rheums.) Not being afflicted with gouty rheums.

ARICYMON. (From *aga* and *κυω*, to be quickly impregnated.) A woman who conceives quickly and often.

ARISTALTHÆA. (From *agros*, best, and *αλθα*, the althæa.) Althæa, or common marsh-mallow.

ARISTOLOCHIA. (From *agros*, good and *λοχι*, or *λοχη*, parturition; so called because it was supposed to be of sovereign use in disorders incident to child-birth.) 1. The name of a genus of plants in the Linnæan system. Class, *Gynandria*, Order, *Hexandria*. Birthwort.

2. The pharmacopœial name of the long-rooted birthwort.

Aristolochia longa of Linnæus:—*foliis cordatis, petiolatis, integerrimis, obtusiusculis; caule infirmo, floribus solitariis*. The root of this plant only is in use; it possesses a somewhat aromatic smell, and a warm bitterish taste, accompanied with a slight degree of pungency. The virtues ascribed to this root by the ancients were very considerable; and it was frequently employed in various diseases, but particularly in promoting the discharge of the *lochia*; hence its name. It is now very rarely used, except in gouty affections, as an aromatic stimulant.

ARISTOLOCHIA ANGUICIDA. Snake-killing birthwort.

Aristolochia:—*foliis cordatis, acuminatis, caule volubili, fruticoso; pedunculis solitariis; stipulis cordatis*, of Linnæus. The juice of the root of this plant has the property of so stupifying serpents, that they

may be handled with impunity. One or two drops are sufficient; and if more be dropt into the mouth, they become convulsed. So ungrateful is the smell of the root to those reptiles, that it is said they immediately turn from it. The juice is also esteemed as a preventive against the effects usually produced by the bite of venomous serpents.

ARISTOLOCHIA CLEMATITIS. (*Clematitis*, from *αἰμα*, a tendril; from its climbing up trees, or any thing it can fasten upon with its tendrils.) The systematic name of the *Aristolochia vulgaris* of some pharmacopœias. See *Aristolochia vulgaris*.

ARISTOLOCHIA FABACEA. The root of this plant, *Fumaria bulbosa* of Linnaeus:—*caule simplici, bracteis longitudine florum*; was formerly given to restore suppressed menses, and as an anthelmintic.

ARISTOLOCHIA LONGA. The systematic name for the aristolochia of our pharmacopœias. See *Aristolochia*.

ARISTOLOCHIA ROTUNDA. The root of this species of birthwort, *Aristolochia rotunda* of Linnaeus:—*foliis cordatis, subaeclibus, obtusis; caule infirmo; floribus solitariis*; is used indiscriminately with that of the *aristolochia longa*. See *Aristolochia*.

ARISTOLOCHIA SERPENTARIA. The systematic name for the *Serpentaria virginiana* of the pharmacopœias. See *Serpentaria virginiana*.

ARISTOLOCHIA TENUIS. See *Aristolochia vulgaris*.

ARISTOLOCHIA TRILOBATA. Three-lobed birthwort. The root, and every part of this plant, *Aristolochia trilobata* of Linnaeus:—*foliis trilobis, caule volubili, floribus maximis*; is diuretic, and is employed in America against the bite of serpents.

ARISTOLOCHIA VULGARIS. *Aristolochia tenuis*. An extract is ordered from this species, *Aristolochia clematitis* of Linnaeus: *foliis cordatis; caule erecto; floribus axillaribus confertis*; by the Wirtemberg Pharmacopœia, and the plant is retained in that of Edinburgh. It is esteemed as possessing antipodagric virtues.

ARISTOPHANEION. (From *Aristophanes*, its inventor.) The name of an ancient emollient plaister composed of wax, or pitch. *Goræus*.

ARMATURA. Hardness. The amnios or internal membrane which surrounds the fœtus.

ARNE. (From *αἶμα*, to adapt.) A junction of the lips of wounds; also the joining of the sutures of the head.

ARMILLA. (Dim. of *armus*, the arm.) The round ligaments which confine the tendons of the carpus.

ARMORACIA. (From *Armerica*, the country whence it was brought.) See *Raphanus rusticanus*.

ARMORACIA RADIX. Horse-radish root. See *Raphanus rusticanus*.

ARNICA. (*Agna*: from *agn*, a lamb; because of the likeness of the leaf of this plant to the coat of the lamb.) Leopard's-bane. *Arnica*.

1. The name of a genus of plants in the Linnaean system. Class, *Syngenesia*, Order, *Polygamia superflua*.

2. The pharmacopœial name of the *Deronicum Germanicum*. Mountain arnica. *Arnica montana* of Linnaeus:—*foliis ovatis integris; caulibus geminis oppositis*. The flowers of this plant are very generally employed on the Continent. Of the advantages derived from their use, in paralytic and other affections, depending upon a want of nervous energy, there are several proofs; and their extraordinary virtues, as a febrifuge and antiseptic, have been highly extolled by Dr. Collin, of Vienna. Much caution is necessary in regulating the dose, as it is a medicine very apt to produce vomiting, and much uneasiness of the stomach.

ARNICA MONTANA. The systematic name for the *arnica* of the pharmacopœias. See *Arnica*.

ARNICA SUEDENSIS. See *Conyza media*. **ARNOTTO.** (Spanish.) A curious shrub in Jamaica, the seeds of which are covered with a kind of wax, from which is made the Spanish arnotto.

AROMA. (From *αἶμα*, intensely, and *ωσ*, to smell.) *Spiritus rector*. Each plant has its characteristic smell. This odorant principle is called by the moderns, *aroma*. Water charged with aroma, is called the distilled water of the substance made use of; thus lavender and peppermint waters are water impregnated with the aroma of the lavender and peppermint.

AROMATICUS CORTEX. A name for *canella alba*.

AROMATICS. (*Aromatica*, sc. *medicamenta*; from *αἶμα*, an odour.) A term applied to all medicines which have a grateful spicy scent, and an agreeable pungent taste, as cinnamon bark, cardamoms, &c. Their peculiar flavour appears to reside in their essential oil, and arises in distillation either with water or spirit.

ARONATOPOLA. (From *αἶμα*, an odour and *ωσ*, to sell.) A druggist; a vender of drugs and spices.

ARQUEBUSADE. (A French word, implying good for a gun shot wound.) *Aqua scoloparia*. *Aqua vulneraria*. *Aqua catapultarium*. The name of a spirituous water, distilled from a farrago of aromatic plants.

ARRACK. A spirituous liquor distilled from rice, and drank, in the rice countries, as we do brandy in this Island. Its effects on the animal œconomy are the same.

ARRAPHUS. (From *α*, priv. and *αἶμα*, a suture.) Without suture. It is applied to the cranium when naturally without sutures.

ARRERA. (From *α*, neg. and *ρα*, to

flow.) The suppression of any natural flux, as the menses, &c.

ARROWHEAD. The roots of this plant, *Sagittaria sagittifolia* of Linnaeus, are said to be esculent, but it must be in times of very great scarcity.

ARROW-ROOT. *Indian arrow-root.* See *Maranta*.

ARSENIAE. (From *arsenicum*, arsenic.) An arseniate or arsenical salt. A salt formed by a combination of arsenic acid with different bases, as arseniate of ammonia, which is produced by the union of ammonia with arsenic acid. The only one used in medicine is the arseniate of potash. See *Liquor arsenicalis*.

ARSENIC. (From the Arabic term *Arsenek*; or from *agrow*, for *agrow masculus*; from its strong and deadly powers.) *Arsenicum crystallinum*, *risagallum*, *aquala*, *arfaz*, *aquila*, *zarnick*, *artaneck*. These names were all formerly applied to white arsenic.

Arsenic is a metal scattered, in great abundance, over the mineral kingdom. It is found in black heavy masses of little brilliancy, called *native arsenic*, (testaceous arsenic.) This exists in different parts of Germany. Mineralized by sulphur, it forms *sulphurized arsenic*, or *orpiment*. This mineral is met with in Italy, about Mount Vesuvius.—There are two varieties of this ore, which differ from each other in colour, occasioned by the different proportions of its component parts. The one is called *yellow sulphurized arsenic*, or *orpiment*; the other, *red sulphurized arsenic*, or *realgar*, (*ruby arsenic*;) both are met with in Hungary and different parts of Germany. The colour of the first ore is a lemon yellow, inclining sometimes to a green; the colour of the latter is a ruby red; it is more transparent than the former, and found in compact solid masses, sometimes crystallized in bright needles. Arsenic united to oxygen, constitutes the ore called *native oxide of arsenic*. This ore is scarce; it is generally found of an earthy appearance, or as an efflorescence, coating native, or metallic arsenic; its colour is a whitish grey; it is rarely met with crystallized. Arsenic exists likewise alloyed with cobalt, antimony, tin, copper, lead, and various other metals.

Properties. Arsenic is a brittle metal, and in the recent fracture, of a lively bright colour, between tin-white and lead gray; but, on exposure to the air, it soon loses its metallic lustre, and turns prismatic, dull, and at last black. Its specific gravity is between 8.310 and 5.763 according to its texture. Its hardness surpasses that of copper; but its ductility is so little, and its brittleness so great, that it is readily converted into a powder by the hammer. It is entirely volatilized when heated to 356° Fabr. It sublimes in close vessels, and then crystallizes in tetrahedra, or octahedra.

When heated with the access of air, it emits a strong smell of garlic, and burns with a blueish white flame. It combines with sulphur by fusion. It unites to phosphorus, and combines with most of the metals. It gives a white colour to copper, and renders many of the ductile metals brittle. When mixed with hyper-oxygenated muriate of potash, it detonates strongly by the stroke of a hammer. It is soluble in hydrogen gas by heat. It does not decompose water alone. It decomposes sulphuric acid by heat. The nitric and nitrous acids oxidate it rapidly. The muriatic acid attacks it with heat. The oxygenated muriatic acid when in a gaseous state, inflames it instantly. It is nearly unalterable by the fluoric, boracic, phosphoric, and carbonic acids. It unites with alkaline sulphurets, and hydro-sulphurets. It is a deadly poison.

Method of obtaining Arsenic. In order to obtain metallic arsenic, mix two parts of the white oxid of arsenic of commerce, with one of black flux (obtained by detoning one part of nitrate of potash with two of acidulous tartre of potash,) and put the mixture into a crucible, or melting-pot. Invert over this, another crucible, lute the two together with a little clay and sand, and apply gradually a red heat to the lower one. The oxid of arsenic will be reduced, and be found lining the upper crucible in small crystals of a metallic brilliancy.

The charcoal of the black flux takes in this process the oxygen from the white oxid, and forms carbonic acid gas; which flies off during the process, and the oxid becomes reduced to the metallic state. This reduction of the oxid is greatly facilitated by the alkali of the flux.

Remark.—In order to obtain arsenic in a state of absolute purity, the metal before obtained must be reduced to powder, dissolved by heat in nitro-muriatic acid, and then precipitated by immersing into the solution a plate of zinc. The arsenic is thus precipitated in a fine powder, and may be reduced to its metallic state, by exposing it in a covered crucible to a moderate heat.

If it be kept under water, its metallic brilliancy may be preserved. This effect is still better produced by alcohol.

Arsenic and its various preparations are the most active of all poisons. A nausea, sickness, and reaching, commonly ensue in half an hour after taking it, followed by violent vomitings, hiccups, and pains in the stomach and bowels; convulsions, and palsies of the limbs presently succeed, with intense heats, cold sweats, palpitations of the heart, extreme anxiety, prostration of strength, thirst and dryness of the mouth and throat; loss of reason, and at last death. If the quantity taken has been considerable, the stomach and intestines are often found, upon dissection, corroded.

or perforated; and the blood is fluid, though in general the patient expires before the action of the poison has proceeded to such a length. After death, the body runs into sudden putrefaction. When the quantity taken does not prove fatal, it occasions tremors, palsies, or lingering hectic, and in the end death. To detect the presence of arsenic, whether taken by design, the wickedness of others, or imprudence, it is necessary to throw on live coals, the contents of the stomach, when a garlicky smell is immediately obvious. On polished copper, if heated between its plates, a white spot is impressed, or in close vessels, the arsenic itself will be found sublimed in the upper parts. In the stomach, however, there are many substances, which may resemble or disguise the smell of arsenic, especially if the arsenic be in small quantities. We are therefore advised by Hahneman, to boil the contents of the stomach of the person supposed to be destroyed by this poison, in a large quantity of river water; to add to one third of the filtered liquor, hot and limpid lime-water; to another third, water saturated with hepatic gas; and to the remainder, a solution of copper, in pure aqua ammoniac. Each fluid is rendered turbid, if the suspected contents contain arsenic, and the sediment, thrown on live coals, emits the odour of garlic. The sediment of the lime-water is again dissolved by a recent solution of arsenic; the orange-coloured sediment, from the hepatic gas thrown on the coals, takes fire, and the smell of sulphur is observed previous to that of the garlic; while the yellow-green sediment of the copper is soluble in pure ammonia, and acids of every kind. Arsenic, however, is a valuable internal remedy, in its appropriate dose, viz. about one-eighteenth part of a grain. *Liquor arsenicalis*.

ARSENIOUS ACID. White arsenic. Oxyd of arsenic. The earliest chymists were embarrassed in the determination of the nature of the poisonous white substance known in commerce by the name of *white arsenic*. Subsequent experiments have shewn that this substance is metallic arsenic oxygenated in the first degree. The name of *arsenious acid* is therefore given to it. It is sometimes found in nature in sublimed crystals, in volcanoes; and in masses, or in stalactites among the ores of arsenic, cobalt, bismuth, and nickel.

It possesses a weak sub-acid taste, which slowly manifests itself. Though of but a feeble acidity, it sensibly reddens the tincture of cabbage and litmus. If placed on burning coals, or on a red-hot iron, it is volatilized in the form of a white vapour, which has a strong smell of garlic. It is slightly soluble in water. With phosphoric and boracic acids it fuses into glass. It decomposes the nitrates and the super-oxygenated muriate of potash. It unites with ma-

ny of the earths and alkalies, and forms saline compounds.

Methods of obtaining Arsenious Acid.—1. Pulverize arsenic, and put as much of it into a Florence flask as will fill it about one half, or less. Introduce a little tow or cotton, into the neck of the flask, and apply the heat of a lamp. A dense white smoke will be formed, and become precipitated on the internal sides of the flask. If the process be kept up till all the arsenic be oxidated, (which may be known by introducing a wire into the flask for a moment, which will become covered with a white crust, if the sublimation be not completed,) and the heat be then gradually augmented, the sublimed arsenious acid undergoes a sort of fusion, and an opake white mass, similar to that met with in commerce, is obtained.

2. The arsenious acid of the shops (or white arsenic) is chiefly obtained from arsenical ores of cobalt. These ores are thrown into a furnace, resembling a baker's oven, with a long flue, or chimney, either horizontal or winding, into which the fumes pass, and are condensed into a greyish or blackish powder. This is refined by a second sublimation, in close vessels, with a little alkali to arrest the impurities. As the heat is considerable, it melts the sublimed arsenious acid into those opake crystalline masses which are known in commerce by the name of white arsenic.

Arsenious acid, united with different bases, forms salts called *arsenites*.

ARSENIC ACID. This is arsenic fully oxygenated. It is always a product of art.

It is capable of existing in the solid state. It appears in the form of a white pulverulent matter. It attracts humidity from the air. It is soluble in water. The solution possesses a considerable acid taste. It may be evaporated to dryness, and even converted into glass. It is decomposable by all combustible bodies, and by many oxyds. It is soluble in some acids, but without change, or intimate combination.

Method of obtaining Arsenic Acid.—Take two ounces of white arsenic in powder, and put it into a tubulated retort; pour on it six or seven ounces of muriatic acid, and apply the heat of a lamp until the arsenious acid is dissolved. Then add three or four ounces of nitric acid, and heat it again gradually. An intestine motion now takes place, and much red vapour, or nitrous gas, is extricated. As soon as in the progress of the operation, the red vapours have ceased, an ounce of finely powdered arsenious acid is to be again added, and the solution effected as before, by a gentle ebullition; to this two ounces of nitric acid must be added, which will produce a second effervescence and discharge of red vapours: the distillation must now be continued to dryness, and the fire must be urged towards

the end, to such a degree, as to make the residual mass thoroughly red-hot. This mass is arsenic acid, which may either be preserved in that form, or be dissolved in boiling distilled water.

Arsenic acid, united to different bases, forms saline compounds, called *ARSENIATES*. The only one used in medicine is the arseniate of potash. See *Liquor arsenicalis*.

ARSENICI OXYDUM PREPARATUM. Prepared arsenious acid. It is sometimes used as a caustic.

ARSENICUM ALBUM. White arsenic. Rat's-bane. See *Arsenious acid*.

ARSENICAL CAUSTIC. A species of caustic said to possess useful properties, independent of those of destroying morbid parts to which it is applied. It is composed of two parts of levigated antimony to one of white arsenic. This is the caustic so extensively employed under the name of *arsenical caustic*, by the late Mr. Justamona, in his treatment of cancers.

Arsenical solution. See *Liquor arsenicalis*.

ARTEMISIA. (From a queen of that name, who first used it; or from *Agræus*, Diana; because it was formerly used in the diseases of women, over whom she presided.) The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia superflua*. Mugwort.

ARTEMISIA ABROTANUM. The systematic name for the *abrotanum* of the pharmacopœias. See *Abrotanum*.

ARTEMISIA ABSINTHIUM. The systematic name for the *Abysynthium vulgare* of the pharmacopœias. See *Abysynthium vulgare*.

ARTEMISIA JUDICIACA. The systematic name for the *Santonium* of the pharmacopœias. See *Santonium*.

ARTEMISIA MARITIMA. The systematic name for the *Abysynthium maritimum* of the pharmacopœias. See *Abysynthium maritimum*.

ARTEMISIA PONTICA. The systematic name for the *Abysynthium ponticum*. See *Abysynthium Ponticum*.

ARTEMISIA RUPESTRIS. The systematic name for the *genipi album* of the pharmacopœias. See *Genipi album*.

ARTEMISIA VULGARIS. Mugwort. This plant, *Artemisia, foliis pinnatifidis, planis, incis, subtus tomentosis, racemis simplicibus recurvatis floribus radio quinque-flo* of Linnæus, is slightly bitter, and, although in high esteem in former days, is now almost wholly forgotten. By beating and rubbing the dried tops of this plant, the Japanese prepare a soft substance, which they call *moxa*. See *Moxa*.

ARTEMONIUM. (From *Artemon*, its inventor.) A collyrium or wash for the eyes.

ARTERIA. See *Artery*.

ARTERIACA. (From *agrega*, an artery.) Medicines formerly used against disorders of the *aspera arteria*.

ARTERIA DIPOSA. The arteries which secrete the fat about the kidneys are so called. They are branches of the capsular and diaphragmatic, renal, and spermatic arteries.

ARTERIA VENOSA. The four pulmonary veins were so called by the ancients.

ARTERIOSUS DUCTUS. See *Ductus arteriosus*.

ARTERIOTOMY. (From *agrega*, an artery, and *τμήν*, to cut. The opening of an artery. This operation is only performed on the temporal artery.

ARTERY. (From *αἷμα*, air, and *ῥήμα*, to keep; so called because the ancients supposed that only air was contained in them.) *Arteria*. Arteries are membranous pulsating canals, which gradually become less as they proceed from the heart. They are composed of three membranes; a common or external, a muscular; and an internal one, which is very smooth. They originate from the heart; the pulmonary artery from the right ventricle, and the aorta from the left: the other arteries are all branches of the aorta. Their termination is either in the veins, or in capillary exhaling vessels, or they anastomose with one another. It is by their means that the blood is carried from the heart to every part of the body, for nutrition, preservation of life generation of heat, and the secretion of the different fluids. The action of the arteries, called the pulse, corresponds with that of the heart, and is effected by the contraction of their muscular, and great elasticity of their innermost, coat.

A Table of the Arteries.

All the arteries originate from the pulmonary artery and the aorta.

The *pulmonary artery* emerges from the right ventricle of the heart, soon divides into a right and left branch, which are distributed by innumerable branches through the lungs.

The *aorta* arises from the left ventricle of the heart, and supplies every part of the body with blood, in the following order:

- a. It first forms an *arch*;
- b. It then descends along the spine, and
- c. It divides into the two *iliacs*.

a. The *ARCH OF THE AORTA* gives off three branches.

I. The *arteria innominata*, which divides into the *right carotid* and *right subclavian*.

II. The *left carotid*.

III. The *left subclavian*.

I. The *carotids* are divided into *external* and *internal*.

The *external carotids* give off

1. The *thyroid*,
2. The *lingual*,
3. The *labial*,
4. The *inferior pharyngeal*,
5. The *occipital*,
6. The *posterior auris*,
7. The *internal maxillary*, from which the

opineus artery of the dura mater, the lower maxillary, and several branches about the palate and orbit arise,

2. The temporal.

The internal carotid affords

1. The *ophthalmic*,
2. The *middle cerebral*,
3. The *communicans*, which anastomoses with the vertebral.

II. The subclavian gives off the following branches :

1. The *internal mammary*, from which the *thymic*, *comes phrenici*, *pericardiac*, and *phrenico-pericardiac arteries* arise,
2. The *inferior thyroid*, which gives off the *tracheal*, *ascending thyroid*, and *transversalis humeri*.
3. The *vertebral*, which proceeds within the vertebrae, and forms within the cranium the *basiliary artery*, from which the *anterior cerebelli*, the *posterior cerebri*, and many branches about the brain are given off,
4. The *cervicalis profunda*,
5. The *cervicalis superficialis*,
6. The *superior intercostal*.
7. The *supra-scapular*.

As soon as the subclavian arrives at the arm-pit, it is called the *axillary artery*; and when the latter reaches the arm, it is called the *brachial*.

The *axillary artery* gives off,

1. *Four mammary arteries*,
2. The *sub-scapular*,
3. The *posterior circumflex*,
4. The *anterior circumflex*, which ramify about the shoulder-joint.

The *brachial artery* gives off,

1. *Many lateral branches*,
2. The *profunda humeri superior*,
3. The *profunda humeri inferior*,
4. The *great anastomosing artery*, which ramifies about the elbow-joint;

The *brachial artery* then divides, about the bend of the arm, into the *ulnar* and *radical arteries*, which are ramified to the ends of the fingers.

The *ulnar artery* gives off,

1. *Several recurrent branches*,
2. The *common interosseal*, of which the *dorsal*, *ulnar*, the *pulmaris profunda*, the *palmar arch*, and the *digitals*, are branches.

The *radial artery* gives off,

1. The *radial recurrent*,
2. The *superficialis volæ*, and then divides into the *palmaris profunda* and the *digitals*.

b. The DESCENDING AORTA gives off, In the breast,

1. The *bronchial*,
2. The *oesophageal*,
3. The *intercostals*,
4. The *inferior diaphragmatic* :

Within the abdomen,

1. The *cœliac*, which divides into three branches :
1. The *hepatic*, from which are given

off, before it reaches the liver,

- a. The *duodeno-gastric*, which sends off the *right gastro-epiploic* and the *pancreatico-duodenal*,
- β. The *pilorica superior hepatica* ;
2. The *coronaria ventriculi*,
5. The *splenic*, which emits the *great* and *small pancreaticæ*, the *posterior gastric*, the *left gastro-epiploic*, and the *vasa brevia* ;
2. The *superior mesenteric*,
3. The *emulgentæ*,
4. The *spermaticæ*,
5. The *inferior mesenteric*,
6. The *lumbar arteries*,
7. The *middle sacral*.

c. The aorta then bifurcates into the *iliacs*, each of which divides into *external* and *internal*.

The *internal iliac*, called also *hypogastric*, gives off,

1. The *lateral sacralæ*,
2. The *gluteal*,
3. The *ischiatricæ*,
4. The *pubical*, from which the *external hæmorrhoidal*, the *perineal*, and the *arterie penis* arise,
5. The *obturatoræ*.

The *external iliac* gives off, in the groin,

1. The *epigastric*,
2. The *circumflexa iliacæ* ;

It then passes under Poupart's ligament, and is called the *femoral artery* ; and sends off,

1. The *profunda*,
2. The *ramus anastomoticus magnus*, which runs about the knee-joint ;

Having reached the ham, where it gives off some small branches, it is termed the *popliteal*. It then divides into the *anterior* and *posterior tibial*.

The *tibialis antica* gives off,

1. The *recurrent*,
2. The *internal malleolar*,
3. The *external malleolar*,
4. The *tarsoal*,
5. The *metatarsæal*,
6. The *dorsales externa halices*.

The *posterior tibial* sends off,

1. The *nutricia tibiae*,
2. *Many small branches*,
3. The *internal plantar*,
4. The *external plantar*, from which an arch is formed, that gives off the *digitals* of the toes.

ARTHANITA. (From *arthos*, bread ; because it is the food of swine.) The herb sow-bread. See *Cyclamen*.

ARTHEMBOLUS. (From *arthron*, a joint, *emballon*, to impel.) An instrument for reducing luxated bones.

ARTHRITICA. (From *arthritic*, the gout.) 1. The herb ground-pine; so called because it was thought good against gouty disorders. 2. Remedies for the gout.

ARTHRITIS. (From *arthron*, a joint ; because it is commonly confined to the

joint.) The gout. Dr. Cullen, in his Nology, gives it the name of *podagra*, because he considers the foot to be the seat of idiopathic gout. It is arranged in the class *pyrexia* and order *phlegmasia*, and is divided into four species, the regular, atonic, retrocedent, and misplaced. The gout is a very painful disease, preceded usually by flatulency and indigestion, and accompanied by fever, pains in the joints of the hands and feet, particularly in that of the great-toe, and which returns by paroxysms, occurring chiefly in the spring and beginning of winter. The only disorder for which it can possibly be mistaken, is the rheumatism; and cases may occur wherein there may be some difficulty in making a just discrimination: but the most certain way of distinguishing them will be, to give due consideration to the predisposition in the habit, the symptoms which have preceded, the parts affected, the recurrences of the disease, and its connexion with other parts of the system. Its attacks are much confined to the male sex, particularly those of a corpulent habit, and robust body; but every now and then we meet with instances of it in robust females. Those who are employed in constant bodily labour, or who live much upon vegetable food, as likewise those who make no use of wine, or other fermented liquors, are seldom afflicted with the gout. The disease seldom appears at an earlier period of life than from five-and-thirty to forty; and, when it does, it may be presumed to arise from an hereditary disposition. Indolence, inactivity, and too free a use of tartareous wines, fermented liquors, and animal food, are the principal causes which give rise to the gout; but it may likewise be brought on by great sensuality and excess in venery, intense and close application to study, long want of rest, grief, or uneasiness of mind, exposure to cold, too free a use of accidulated liquors, a sudden change from a full to a spare diet, the suppression of any accustomed discharge, or by excessive evacuations; and that it sometimes proceeds from an hereditary disposition, is beyond all doubt, as females who have been remarked for their great abstemiousness, and youths of a tender age, have been attacked with it.

1. *Arthritis regularis*. A paroxysm of regular gout sometimes comes on suddenly, without any previous warning; at other times it is preceded by an unusual coldness of the feet and legs, a suppression of perspiration in them, and numbness, or with a sense of prickling along the whole of the lower extremities; and with these symptoms the appetite is diminished, the stomach is troubled with flatulency and indigestion, a degree of torpor and languor is felt over the whole body, great lassitude and fatigue are experienced after the least ex-

ercise, the body is costive and the urine pallid. On the night of the attack, the patient perhaps goes to bed in tolerable health, and, after a few hours, is awakened by the severity of the pain, most commonly in the first joint of the great-toe; sometimes, however, it attacks other parts of the foot, the heel, calf of the leg, or perhaps the whole of the foot. The pain resembles that of a dislocated bone, and is attended with the sensation as if cold water was poured upon the part; and this pain becoming more violent, is succeeded by rigors and other febrile symptoms, together with a severe throbbing and inflammation in the part. Sometimes both feet become swelled and inflamed, so that neither of them can be put to the ground; nor can the patient endure the least motion, without suffering excruciating pain. Towards morning, he falls asleep, and a gentle sweat breaks out, and terminates the paroxysm, a number of which constitutes what is called a fit of the gout; the duration of the fit will be longer or shorter, according to the disposition of the body to the disease, the season of the year, and the age and strength of the patient. When a paroxysm has thus taken place, although there is an alleviation of pain at the expiration of some hours, still the patient is not entirely relieved from it; and, for some evenings successively, he has a return both of pain and fever, which continue with more or less violence, until morning. The paroxysms, however, prove usually more mild every day, till at length the disease goes off either by perspiration, urine, or some other evacuation; the parts which have been affected becoming itchy, the cuticle falling off in scales from them, and some slight degree of lameness remaining. At first, an attack of gout occurs, perhaps, only once in two or three years; it then probably comes on every year, and, at length, it becomes more frequent, and is more severe, and of longer duration, each succeeding fit. In the progress of the disease, various parts of the body are affected, and translations take place from one joint, or limb, to another; and, after frequent attacks, the joints lose their strength and flexibility, and become so stiff as to be deprived of all motion. Concretions, of a chalky nature, are likewise formed upon the outside of the joints, and nephritic affections of the kidneys arise from a deposit of the same kind of matter in them which, although fluid at first, becomes dry and firm at last, and, when put into acids, is perfectly soluble.

2. *Arthritis atonica*. Atonic gout. It sometimes happens that, although a gouty diathesis prevails in the system, yet, from certain causes, no inflammatory affection of the joints is produced; in which case, the stomach becomes particularly affected, and

the patient is troubled with flatulency, indigestion, loss of appetite, eructations, nausea, vomiting, and severe pains; and these affections are often accompanied with much dejection of spirits, and other hypochondriacal symptoms. In some cases, the head is affected with pain and giddiness, and now and then with a tendency to apoplexy; and in other cases, the viscera of the thorax suffer from the disease, and palpitations, faintings, and asthma arise. This is what is called atonic gout.

3. *Arthritis retrograda*. Retrocedent gout. It sometimes happens that, after the inflammation has occupied a joint, instead of its continuing the usual time, and so going off gradually, it ceases suddenly, and is translated to some internal part. The term of retrocedent gout is applied to occurrences of this nature. When it falls on the stomach, it occasions nausea, vomiting, anxiety, or great pain; when on the heart, it brings on syncope; when on the lungs, it produces an affection resembling asthma; and, when it occupies the head, it is apt to give rise to apoplexy, or palsy.

4. *Arthritis aberrans*, or misplaced gout, is when the gouty diathesis, instead of producing the inflammatory affection of the joints, occasions an inflammatory affection of some internal part, and which appears from the same symptoms that attend the inflammation of those parts from other causes. All occurrences of this nature, as well as of the two former, are to be regarded as attacks of irregular gout and are to be guarded against as much as possible.

ARTHROCAE. (From *arthron*, a joint.) An ulcer of the cavity of the bone.

ARTHRODIA. (From *arthron*, to articulate.) A species of *diarthrosis*, or moveable connexion of bones, in which the head of one bone is received into the superficial cavity of another, so as to admit of motion in every direction, as the head of the humerus with the glenoid cavity of the scapula.

ARTHRODYNIA. (From *arthron*, a joint, and *dun*, pain.) Chronic pains in the joints, without pyrexia. It is one of the terminations of acute rheumatism. See *Rheumatismus*.

ARTHROPUOSIS. (From *arthron*, a joint, and *pus*, pus.) *Arthropyosis*. A collection of pus in a joint. It is, however, frequently applied to other affections, as *lumbago podagica*, &c.

ARTHROSIS. (From *arthron* to articulate, or join together.) Articulation.

Artichoke. See *Cynara*.

Artichoke, French. See *Cynara*.

ARTICHOKE, JERUSALEM. Although formerly in estimation for the table, this plant *Helianthus tuberosus* of Linnæus, is now neglected, it being apt to produce flatulency and dyspepsia.

ARTICULARIS. A name given to a dis-

ease which more immediately infests the *articuli*, or joints. The morbus articularis is synonymous with the Greek word arthritis, and our gout. A branch of the basilic vein is called *articularis vena*, because it passes under the joint of the shoulder.

ARTICULATION. (From *articulus*, a joint.) The skeleton is composed of a great number of bones, which are all so admirably constructed, and with so much affinity to each other, that the extremity of every bone is perfectly adjusted to the end of the bone with which it is connected; and this connexion is termed their articulation. Anatomists distinguish three kinds of articulation: the first they name *Diarthrosis*; the second, *Synarthrosis*; and the third *Amphiarthrosis*; which see, under their respective heads.

ARTISCUS. (From *agros*, bread.) A troch; so called because they are made like little loaves.

ARTOCREAS. (From *agros*, bread, and *zeugros*, fruit.) A nourishing food, made of bread and various meats, boiled together. *Galen*.

ARTOGALA. (From *agros*, bread, and *gala*, milk.) A cooling food, made of bread and milk.) A poultice.

ARTOMELI. (From *agros*, bread, and *mel*, honey.) A cataplasm made of bread and honey. *Galen*.

ARUM. (From the Hebrew word *jarom* which signifies a dart, so named because its leaves are shaped like a dart; or from *apa*, injury.)

1. The name of a genus of plants in the Linnæan system. Class, *Gynandria*. Order, *Polyandria*. *Arum*, or wake-robin.

2. The pharmacopœial name of the common *arum*, or wake-robin.

Arum maculatum of Linnæus:—*acaulis foliis hastatis integerrimis, spadice clavatis*. The root is the medicinal part of this plant, which, when recent, is very acrimonious; and, upon being chewed, excites an intolerable sensation of burning and pricking in the tongue, which continues for several hours. When cut in slices, and applied to the skin, it has been known to produce blisters. This acrimony, however, is gradually lost by drying, and may be so far dissipated by the application of heat, as to leave the root a bland farinaceous aliment. In this state, it has been made into a wholesome bread. It has also been prepared as starch. Its medicinal quality, therefore, resides wholly in the active volatile matter, and consequently the powdered root must lose much of its power, on being long kept. *Arum* is certainly a powerful stimulant, and, by promoting the secretions, may be advantageously employed in cachectic and chlorotic cases, in rheumatic affections, and in various other complaints of phlegmatic and torpid

constitutions; but more especially in a weakened or relaxed state of the stomach, occasioned by the prevalence of viscid mucus. If this root is given in powder, great care should be taken that it be young and newly dried, when it may be used in the dose of a scruple, or more, twice a day; but in rheumatisms, and other disorders requiring the full effect of this medicine, the root should be given in a recent state; and, to cover the insupportable pungency it discovers on the tongue, Dr. Lewis advises us to administer it in the form of emulsion, with gum-arabic and spermaceti, increasing the dose from ten grains to upwards of a scruple, three or four times a day. In this way, it generally occasioned a sensation of slight warmth about the stomach, and afterwards, in the remoter parts, manifestly promoted perspiration, and frequently produced a plentiful sweat. Several obstinate rheumatic pains were removed by this medicine. The root answers quite as well as garlic for cataplasms, to be applied on the feet in deliriums. The London College, in their pharmacopœia, 1788, ordered a conserve, in the proportion of half a pound of the fresh root to a pound and a half of double refined sugar, beat together in a mortar, which appears to be one of the best forms of exhibiting arum, as its virtues are destroyed by drying, and are not extracted by any menstruum. It may be given to adults in doses of a drachm.

ARUM MACULATUM. The systematic name for the arum of the pharmacopœias. See *Arum*.

ARYTENO-EPIGLOTTIDEUS. Innes. Albinus, *Aryteno-Epiglottici* of Winslow. A muscle composed of a number of fibres running between the arytenoid cartilage and epiglottis. It pulls the side of the epiglottis towards the external opening of the glottis, and when both act, they pull it close upon the glottis.

ARYTENOID CARTILAGE. *Cartilago arytenoidea.* The name of two cartilages of the larynx. See *Larynx*.

ARYTENOIDES. (From *apsana*, a funnel, and *ωει*, shape.) The name of some parts, from their being funnel-shaped.

ARYTENOIDEUS MAJOR. See *Arytenoideus transversus*.

ARYTENOIDEUS MINOR. See *Arytenoideus obliquus*.

ARYTENOIDEUS OBLIQUUS. Innes. Albinus, and Winslow. *Arytenoideus minor* of Douglas. A muscle of the glottis, which arises from the base of one arytenoid cartilage, and crossing its fellow, is inserted near the tip of the other arytenoid cartilage. It is a muscle that is occasionally wanting; but when present, and both muscles act, their use is to pull the arytenoid cartilages towards each other.

ARYTENOIDEUS TRANSVERSUS, of Innes, Albinus, Winslow. *Arytenoideus major* of Douglas. An azygos, or single muscle of the glottis, that arises from the side of one arytenoid cartilage, from near its articulation with the cricoid to near its tip. The fibres run across, and are inserted in the same manner into the other arytenoid cartilage. Its use is to shut the glottis, by bringing the two arytenoid cartilages, with their ligaments, nearer to each other.

ASAFETIDA. (From the Hebrew word *asa*, to heal.) *Hingisch* of the Persians, *Alhi* of the Arabians. By some thought to be the *σαλψιον*, vel, *σπος σαλψις* of Dioscorides, Theophrastus, and Hippocrates. *Laos* et *laserpitium* of the Latins. *Assafœtida* gum resin. The plant which affords this gum resin, is the *Ferula assafœtida* of Linnaeus:—*foliis alternatim sinuatis, obtusis.* Class, *Pentandria.* Order, *Digymia.* It grows plentifully on the mountains in the provinces of Chorasaa and Laar in Persia.

The process of obtaining it is as follows: the earth is cleared away from the top of the roots of the oldest plants; the leaves and stalks are then twisted away, and made into a covering, to screen the root from the sun; in this state the root is left for forty days, when the covering is removed, and the top of the root cut off transversely; it is then screened again from the sun for forty-eight hours, when the juice it exudes is scraped off, and exposed to the sun to harden. A second transverse section of the root is made, and the exudation suffered to continue for forty-eight hours, and then scraped off. In this manner it is eight times repeatedly collected in a period of six weeks. The juice thus obtained has a bitter, acrid, pungent taste, and is well known by its peculiar nauseous smell, the strength of which is the surest test of its goodness. This odour is extremely volatile, and of course the drug loses much of its efficacy by keeping. It is brought to us in large irregular masses, composed of various little shining lumps, or grains, which are partly of a whitish colour, partly reddish, and partly of a violet hue. Those masses are accounted the best which are clear, of a pale reddish colour, and variegated with a great number of elegant white tears. This concrete juice consists of two-thirds of gum and one-third of resin, its taste and smell residing in the resinous part. It yields all its virtues to alcohol. Triturated with water, it forms a milky-like mixture, the resin being diffused by the medium of the gum. Distilled with water, it affords a small quantity of essential oil. It is the most powerful of all the fœtid gums, and is a most valuable remedy. It is most commonly employed in hysteria, hypochondriasis, some symptoms of dyspepsia, flatulent

colics, and in most of those diseases termed nervous, but its chief use is derived from its antispasmodic effects; and it is thought to be the most powerful remedy we possess, for those peculiar convulsive and spasmodic affections, which often recur in the first of these diseases, both taken into the stomach and in the way of enema. It is also recommended as an emmenagogue, anthelmintic, antiasthmatic, and anodyne. Dr. Cullen prefers it as an expectorant to gum ammoniacum. Where we wish it to act immediately as an antispasmodic, it should be used in a fluid form, as that of tincture, from half a drachm to two drachms. When given in the form of a pill, or triturated with water, its usual dose is from 5 to 20 grs. When in the form of enema, two drachms are to be diffused in eight ounces of warm milk or water. It is sometimes applied externally as a plaster and stimulating remedy, and is much used in hysteria, hypochondriasis, dyspepsia, &c.

ASAPHATUM. (From *a*, neg. and *sapha*, clear.) An intercutaneous itch, generated in the pores, like worms with black heads: so called by reason of their minuteness: they are hardly visible.

ASAPHIA. (From *a*, neg. and *sapha*, clear.) A defect in utterance or pronunciation.

ASARABACCA. See *Asarum*.

ASARI FOLIA. Asarabacca leaves. The leaves of the *Asarum Europæum*. See *Asarum*.

ASARUM. (From *a*, neg. and *sapha*, to adorn, because it was not admitted into the ancient coronal wreaths.) Asarabacca.

1. The name of a genus of plants in the Linnæan system. Class, *Dodecandria*. Order, *Monogynia*.

2. The pharmacopœial name of the asarabacca.

Asarum Europæum of Linnæus, *foliis reniformibus obtusis binis*.

It is a native of England, but not very common. The leaves of this plant are extremely acrid, and are occasionally used, when powdered, as a sternutatory. For this purpose the leaves, as being less acrid than the roots, are preferred, and in moderate doses not exceeding a few grains, snuffed up the nose several evenings, produce a pretty large watery discharge, which continues for several days together, by which head-ache, tooth-ache, ophthalmia, and some paralytic and soporific complaints have been effectually relieved.

Prior to the introduction of ipecacuanha, the leaves and root of this plant were frequently employed on account of their emetic power: the dose of the dried leaves was 20 grains; of the dried roots 10 grains. As they were occasionally violent in their operation, they have fallen into disuse.

ASARUM EUROPEUM. The systematic

name of the asarabacca of the shops. See *Asarum*.

ASCALONITIS. A species of onion.

ASCARIDES. The plural of *ascaris*. See *Ascaris*.

ASCARIS. (From *arxus*, to move about; so called from its continued troublesome motion.) There are several kinds of worms distinguished by this term; but those which claim a place here as belonging only to the human body, are:—

1. *Ascaris vermicularis*, the thread or maw worm, which is very small and slender, not exceeding half an inch in length: they inhabit the rectum.

2. *Ascaris lumbricoides*, the long and round worm, which is a foot in length, and about the breadth of a goose-quill.

ASCENDENS OBLIQUUS. See *Obliquus ascendens*.

ASCIA. An axe or chissel. A simple bandage; so called from its shape in position. *Galen*.

ASCITES. (From *arxos*, a sack, or bottle: so called from its bottle-like protuberancy.) Dropsy of the belly. A tense, but scarcely elastic, swelling of the abdomen from accumulation of water. Cullen ranks this genus of disease in the class *cachexie*, and order *intumescencie*. He enumerates two species:—

1. *Ascites abdominalis*, when the water is in the cavity of the peritonæum, which is known by the equal swelling of the parietes of the abdomen.

2. *Ascites saccatus*, or encysted dropsy, in which the water is encysted, as in the ovary; the fluctuation is here less evident, and the swelling is at first partial.

Ascites is often preceded by loss of appetite, sluggishness, dryness of the skin, oppression at the chest, cough, diminution of the natural discharge of urine, and costiveness. Shortly after the appearance of these symptoms, a protuberance is perceived in the hypogastrium, which extends gradually, and keeps on increasing, until the whole abdomen becomes at length uniformly swelled and tense. The distension and sense of weight, although considerable, vary somewhat according to the posture of the body, the weight being felt the most on that side on which the patient lies, whilst at the same time, the distension becomes somewhat less on the opposite side. In general the practitioner may be sensible of the fluctuation of the water, by applying his left hand on one side of the abdomen, and then striking on the other side with his right. In some cases it will be obvious to the ear. As the collection of water becomes more considerable, the difficulty of breathing is much increased, the countenance exhibits a pale and bloated appearance, an immoderate thirst arises, the skin is dry and parched, and the urine is very scanty, thick, high coloured, and deposits a

lenticulous sediment. With respect to the pulse, it is variable, being sometimes considerably quickened, and at other times, slower than natural. The principal difficulty which prevails in ascites, is the being able to distinguish with certainty, when the water is in the cavity of the abdomen, or when it is in the different states of encysted dropsy. To form a just judgment, we should attend to the following circumstances: When the preceding symptoms give suspicion of a general hydropic diathesis; when at the same time, some degree of dropsy appears in other parts of the body; and when from its first appearance the swelling has been equally diffused over the whole belly, we may generally presume that the water is in the cavity of the abdomen. But when an ascites has not been preceded by any remarkable cachectic state of the system, and when, at its beginning, the tumour and tension had appeared in one part of the belly more than another, there is reason to suspect an encysted dropsy. Even when the tension and tumour of the belly have become general, yet if the system or the body in general appear to be little affected; if the patient's strength be little impaired; if the appetite continue pretty entire, and the natural sleep be little interrupted; if the menses in females continue to flow as usual; if there be yet no anasarca, or though it may have already taken place, if it be still confined to the lower extremities, and there be no leucophlegmatic paleness or sallow colour in the countenance; if there be no fever, nor so much thirst and scarcity of urine as occur in a more general affection: then according as more of these different circumstances take place, there will be the stronger grounds for supposing the ascites to be of the encysted kind.

ASX. A pustule like a millet seed.

ASGOS. *Asgen.* *Asogen.* Dragon's blood.

ASCLEPIAS. (From *Asclepias*, its discoverer; or from *Esculapius*, the god of medicine.) The herb swallow-wort. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*.

ASCLEPIAS VINETOXICUM. The systematic name for the *vincetoxicum* of the pharmacopœias. See *Vincetoxicum*.

ASCLEPIOS. (From *Asclepias*, its inventor.) A dried smegma and collyrium described by Galen.

ASCOMA. (From *asus*, a bottle.) The eminence of the pubes at the years of maturity; so called from its shape.

ASH. See *Fraxinus*.

ASODES. (From *asus*, to nauseate.) A nausea or loathing, or a fever with much sense of heat and nausea. *Arotens*.

ASIATICUM BALSAMUM. Balm of Gilead.

ASINUS. The ass. Its milk is much esteemed in medicine. See *Asses' milk*.

ASINUM LAC. *Asses' milk*.

ASITI. (From *a*, neg. and *erros*, food.)

Adria. Those are so called who take no food far want of appetite.

ASOGAN. (Indian) A tree growing in Malabar and the East Indies, whose juice is used against the colic.

ASPADIALIS. A suppression of urine from an imperforated urethra.

ASPALATHUM. The aromatic aloe.

ASPALATHI LIGNUM. See *Lignum aloes*.

ASPARAGUS. (*Asparagus*, a young shoot, before it unfolds its leaves.)

1. The name of a genus of plants in the Linnæan system. Class, *Hexandria*, Order, *Monogynia*. *Asparagus*.

2. The pharmacopœial name of the common sparage, or sparrow-grass.

Asparagus officinalis of Linnæus. The root has been esteemed as a diuretic. It is mostly employed as a food, but it contains very little nourishment.

ASPASIA. (From *a*, for *aqua*, together, and *asus*, to draw.) A constrictive medicine for the pudendum muliebre. *Captivac*.

ASPERA ARTERIA. (So called from the inequality of its cartilages.) See *Trachea*.

ASPERULA. (A diminutive of *asper*, the seeds being rough.) The name of a genus of plants in the Linnæan system. Class, *Tetrandria*. Order, *Monogynia*. Wood-roof.

ASPERULA ODORATA. The systematic name for the official *matriytha*. See *Matriytha*.

ASPHALITIS. A kind of trefoil: the last vertebra of the loins.

ASPHODELUS. (From *asus*, a serpent, and *delos*, fearful; because it destroys the venom of serpents; or from *asphalos*, ashes, because it was formerly sown upon the graves of the dead.) *Asphodel*.

1. The name of a genus of plants in the Linnæan system. Class, *Hexandria*. Order, *Monogynia*.

2. The pharmacopœial name of the daffodil, or branched asphodel.

Asphodelus ramosus, of Linnæus:—*caule nudo, foliis ensiformibus, cartilatis, levibus*. The plant was formerly supposed to be efficacious in the cure of sordid ulcers. It is now wholly laid aside.

ASPHODELUS RAMOSUS. The systematic name for the official *asphodelus*. See *Asphodelus*.

ASPHYXIA. (From *a*, priv. and *σφυα*, a pulse.) The state of the body, during life, in which the pulsation of the heart and arteries cannot be perceived. There are several species of asphyxia enumerated by different authors. See *Syncope*.

ASPIDISCUS. (From *aspid*, a buckler.) The sphincter muscle of the anus was formerly so called from its shape. *Culius Aurelianus*.

ASPLENIUM. (From *a*, priv. and *σπλην* the spleen; because it was supposed to remove disorders of the spleen.) The herb spleen-wort.

The name of a genus of plants in the Lin-

nean system. Class, *Cryptogamia*. Order, *Filices*.

ASPLENIUM RUTA MURARIA. The systematic name for the *ruta muraria* of the pharmacopœias. See *Ruta muraria*.

ASPLENIUM SCOLOPENDRIUM. The systematic name for the *scolopendrium* of the pharmacopœias. See *Scolopendrium*.

ASPLENIUM TRICHOMANES. The systematic name for the *trichomanes* of the pharmacopœias. See *Trichomanes*.

ASABA. A shrub found on the coast of Guinea, whose leaves are supposed to disperse buboes.

ASSAC. (Arab.) Gum ammoniacum.

ASSAFŒTIDA. See *Assafœtida*.

ASSALA. The nutmeg.

ASSANUS. A weight consisting of two drachms.

ASSARABACCA. See *Asarum*.

ASSARIUM, A Roman measure of twelve ounces.

ASTHETHROSIS. See *Astretation*.

ASSE. A loathing of food, from a conflux of humours. *Hippocrates*.

ASSES' MILK. This is preferred to cows' and other kinds of milk in phthisical cases, and where the stomach is weak; as containing less oleaginous particles, and being more easily converted into chyle.

ASSIMULATION. (*Assimilatio*, from *ad*, and *similis*, to make like to.) The conversion of the food into nutriment.

ASSISTENTES. (From *ad*, and *sisto*, to stand near.) A name of the prostate gland; so called because it lies near the bladder.

ASSODES. (From *ασωδω*, to nauseate, or from *ασωρε*, to burn.) *Assodes*. A continual fever attended with a loathing of food. Sauvages calls it *Tritæphyta assodes*; it is arranged by Cullen under the tertian remittents.

ASSOS. A name given formerly to alumen.

ASTACUS MARINUS. (From *a*, neg. and *στα*, to distil; so called from the hardness and dryness of its shell.) The lobster. The black tips of the claws of this fish, and of the sea-crab, and the stony concretions in the heads of the *astacus fluviatilis*, called crab's eyes, form some of the absorbent preparations of the shops.

ASTACUS FLUVIATILIS. The officinal crab, crevis, or cray-fish.

ASTAPIS. (From *στασις* *ura*, passa.) A raisin.

ASTARZOF. The name of an ointment of litharge, house-leek, &c. *Haracelous*.

ASTRACHILLOS. A malignant ulcer, by some called *araneus*.

ASTRANTIUM. (From *αστη*, a star.) *Astericum*. The herb pellitory: so called from its star-like form.

ASTHENIA. (From *a*, priv. and *σθεν*, strength.) Extreme debility. The asthenic diseases form one great branch of the Brunonian hypothesis.

ASTHENOLGY. (From *a*, priv. and *σθεν*, strength, and *λογος*, a treatise.) The doctrine of diseases arising from debility. The disciples of the Brunonian school, as they denominate themselves, maintain peculiar opinions on this subject.

ASTHMA. (From *ασμαζω*, to breathe with difficulty.) Difficult respiration, returning at intervals, with a sense of stricture across the breast, and in the lungs; a wheezing, hard cough, at first, but more free towards the close of each paroxysm, with a discharge of mucus, followed by a remission. It is ranked by Cullen in the class *neuroses*, and order *spasmi*. There are three species of asthma:—

1. *Asthma spontaneum*, when without any manifest cause.

2. *Asthma plethoricum*, when it arises from plethora.

3. *Asthma exanthematicum*, originating from the repulsion of some acrid humour.

Asthma rarely appears before the age of puberty, and seems to attack men more frequently than women, particularly those of a full habit, in whom it never fails, by frequent repetition, to occasion some degree of emaciation. In some instances, it arises from an hereditary predisposition, and in many others, it seems to depend upon a particular constitution of the lungs. Dyspepsia always prevails, and appears to be a very prominent feature in the predisposition. Its attacks are most frequent during the heats of summer in the dog-days, and in general commence at midnight. On the evening preceding an attack of asthma the spirits are often much affected, and the person experiences a sense of fullness about the stomach, with lassitude, drowsiness, and a pain in the head. On the approach of the succeeding evening, he perceives a sense of tightness and stricture across the breast, and a sense of straitness in the lungs, impeding respiration. The difficulty of breathing continuing to increase for some length of time, both inspiration and expiration are performed slowly and with a wheezing noise; the speech becomes difficult and uneasy, a propensity to coughing succeeds, and the patient can no longer remain in a horizontal position, being as it were threatened with immediate suffocation. These symptoms usually continue till towards the approach of morning, and then a remission commonly takes place; the breathing becomes less laborious and more full, and the person speaks and coughs with greater ease. If the cough is attended with an expectoration of mucus, he experiences much relief, and soon falls asleep. When he awakes in the morning, he still feels some degree of tightness across his breast, although his breathing is probably more free and easy, and he cannot bear the least motion, without rendering this more diffi-

cult and uneasy; neither can he continue in bed, unless his head and shoulders are raised to a considerable height. Towards evening, he again becomes drowsy, is much troubled with flatulency, in the stomach, and perceives a return of the difficulty of breathing, which continues to increase gradually, till it becomes as violent as on the night before. After some nights passed in this way, the fits at length moderate, and suffer more considerable remissions, particularly when they are attended by a copious expectoration in the mornings, and that this continues from time to time throughout the day; and the disease going off at last, the patient enjoys his usual rest by night, without further disturbance.

ASTITES. (From *ad*, and *sto*, to stand near.) A name given by the ancients to the prostate glands, because they are situated near the bladder.

ASTRAGALUS. (*Ἀσπράγγος*, a cockal, or die; because it is shaped like the die used in ancient games.) *Ballistæ os: artetrios: talus: quatrio: tetoreros: cardicula carvilla: diabebo: peza.*

1. The name of a bone of the *sermus*, upon which the tibia moves. Ankle-bone; also called the sling-bone, or first bone of the foot. It is placed posteriorly and superiorly in the tarsus, and is formed of two parts, one large, which is called its body, the other small, like a process. The part where these two unite is termed the neck.

2. The name of a genus of plants in the Linnæan system. Class, *Didelphia*. Order, *Decandria*. Milk-vetch.

ASTRAGALUS EXCAPUS. Stemless milk vetch. The root of this plant, *Astragalus aculeus excapus leguminibus lunatis, foliis villosis* of Linnæus, is said to cure confirmed syphilis, especially when in the form of nodes and nocturnal pains.

ASTRAGALUS TRAGACANTHA. The systematic name for the plant which affords the gum tragacanth. See *Tragacantha*.

ASTRANTIA VULGARIS. From *astrum*, a star; so called from the star-like shape of its flowers.) *Astrantia nigra*. The herb sanicle masterwort. A rustic purge.

ASTRAPE. (From *ασπράζω*, to corruscate.) Lightning. Galen reckons it among the remote causes of epilepsy.

ASTRICTA. (From *astringo*, to bind.) When applied to the belly, it signifies costiveness; thus *alvus stricta*.

Astringents. See *Astringents*.

ASTRONOMIA. (From *aster*, a star, and *nomos*, a law.) Astronomy, or the knowledge of the heavenly bodies. Hippocrates ranks this and astrology among the necessary studies of a physician.

ASUAR. Indian myrobalans, or purging nut.

ASUGAR. *Ærugo seris*, or verdigrise.

ASUOLI. Fuligo, or soot, an antispasmodic.

ATAO. Nitre.

ATAXIA. (From *a*, neg. and *taxis*, to order.) Want of regularity in the symptoms of a disease, or of the functions of an animal body.

ATAXIE. (Arab.) A tenesmus: a disease of the eyes.

ATAXIE. (Arab.) Removal of preternatural hairs growing under the natural ones on the eye-lids.

ATEBRAS. A chymical subliming vessel.

ATECNIA. (From *a*, neg. and *taxis*, to bring forth.) Venereal impotency: inability to procreate children.

ATHAMANTA CRETENSIS. The systematic name for the *daucus creticus* of the pharmacopœias. See *Daucus creticus*.

ATHAMANTA OREOSELINUM. The systematic name for the officinal *oreoselinum*. See *Oreoselinum*.

ATHANASIA. (From *a*, priv. and *θανος*, death; so called because its flowers do not wither easily.) The immortal plant. A name given to tansey: because when stuffed up the nose of a dead corpse, it is said to prevent putrefaction. See *Tanacetum*. It means also immortality. The name of an antidote of Galen, and another of Oribasius: it is the name also of a collyrium described by Ætius, and of many other compositions.

ATHANOR. (Arab.) A chymical digesting furnace.

ATHARA. (From *athap*, corn.) A panada, or pap for children, made of bruised corn.

ATHEKA. A plaster in much repute among the ancients.

ATHENATORIUM. A thick glass cover formerly used for chymical purposes.

ATHENIONIS CATAPOTIUM. The name of a pill in Celsus's writings.

ATHEKIPPOX. *Athenippum.* *Diasmyrnes.* The name of a collyrium.

ATHEROMA. (*Ἀθήρομα*, pulse, pap.) An encysted tumour that contains a soft substance of the consistence of a poultice.

ATHONOR. (Arab.) A chymical furnace.

ATHYMIA. (From *a*, neg. and *θυμος*, courage.) Pusillanimity. Despondence synonymous with melancholia.

ATINCAB. (Arab.) Borax.

ATLAS. (From *ἄταλα*, to sustain, because it sustains the head; or from the fable of Atlas, who was supposed to support the world upon his shoulders.) The name of the first cervical vertebra. This vertebra differs very much from the others. (See *Vertebra*.) It has no spinous process which would prevent the neck from being bent backwards, but in its place it has a small eminence. The great foramen of this is much larger than that of any other vertebra. Its body, which is small and thin, is nevertheless firm and hard. It is somewhat like a ring, and is distinguished into its *great arch*, which serves in the place of its body, and its *small posterior arch*. The atlas is joined superiorly to the head by

ginglymus; and inferiorly, to the second cervical vertebra, by means of the inferior oblique processes and the odontoid process by trochoides.

ATMOSPHERE. (From *ἀτμός*, vapour and *σφαῖρα*, a globe. The elastic invisible fluid which surrounds the earth to an unknown height and encloses it on all sides.

Neither the properties nor the composition of the atmosphere seem to have occupied much the attention of the ancients.

Aristotle considered it as one of the four elements, situated between the regions of water and fire, and mingled with two exhalations, the dry and the moist: the first of which occasioned thunder, lightning, and wind; while the second produced rain, snow, and hail.

The opinions of the ancients were vague conjectures, until the matter was explained by the sagacity of Hales, and of those philosophers who followed his career.

Boyle ventured beyond a doubt, that the atmosphere contained two distinct substances:—

1. An elastic fluid distinguished by the name of *air*.

2. Water in a state of vapour.

Besides these two bodies it was supposed that the atmosphere contained a great variety of other substances which were continually mixing with it from the earth, and which often altered its properties and rendered it noxious or fatal. Since the discovery of carbonic acid gas by Dr. Black, it has been ascertained that this elastic fluid always constitutes a part of the atmosphere.

The constituent parts of the atmosphere, therefore, are:—

1. Air.
2. Water.
3. Carbonic acid gas.
4. Unknown bodies.

For the properties, composition and account of the first, See *Air*.

2. *Water*.—That the atmosphere contains water, has been always known. The rain and dew which so often precipitate from it, the clouds and fogs with which it is often obscured, and which deposit moisture on all bodies exposed to them, have demonstrated its existence in every age. Even when the atmosphere is perfectly transparent, water may be extracted from it in abundance by certain substances. Thus if concentrated sulphuric acid be exposed to air, it gradually attracts so much moisture, that its weight is increased more than three times: it is converted into diluted acid, from which the water may be separated by distillation. Substances which have the property of abstracting water from the atmosphere, have received the epithet of *hygroscopic*, because they point out the presence of that water. Sulphuric acid, the fixed alkalies, muriat of lime, nitrat of lime, and in general

all deliquescent salts, possess this property. The greater number of animal and vegetable bodies likewise possess it. Many of them take water from moist air, but give it out again to the air when dry. These bodies augment in bulk when they receive moisture, and diminish again when they part with it. Hence some of them have been employed as *hygrometers* or measurers of the quantity of moisture contained in the air around them. This they do by means of the increase or diminution of their length, occasioned by the addition or abstraction of moisture. This change of length is precisely marked by means of an index. The most ingenious and accurate hygrometers, are those of Saussure and Deluc. In the first, the substance employed to mark the moisture is a human hair, which by its contractions and dilatations is made to turn round an index. In the second, instead of a hair, a very fine thin slip of whalebone is employed. The scale is divided into 1000. The beginning of the scale indicates extreme dryness, the end of it indicates extreme moisture. It is graduated by placing it first in air made as dry as possible by means of salts, and afterwards in air saturated with moisture. This gives the extremes of the scale, and the interval between them is divided into 100 equal parts.

The water, which constitutes a component part of the atmosphere, is chymically combined with air; but it exists in two different states. A small portion is held in solution in the state of water, but by far the greater proportion is in the state of an elastic fluid, whose specific gravity is to that of air as 10 to 12, and chymically combined with air in the same manner as one gas is combined with another. As the quantity of the water contained in the atmosphere varies considerably, it is impossible to ascertain its amount with any degree of accuracy.

3. *Carbonic acid gas*.—The existence of carbonic gas as a constituent part of the atmosphere, was observed by Dr. Black immediately after he had ascertained the nature of that peculiar fluid. If we expose a pure alkali or alkaline earth to the atmosphere, it is gradually converted into a carbonat by the absorption of carbonic acid gas. This fact, which had been long known, rendered the inference that carbonic acid gas existed in the atmosphere unavoidable as soon as the difference between a pure alkali and its carbonat had been ascertained to depend upon that acid. Not only alkalies and alkaline earths absorb carbonic acid when exposed to the air, but several of the metallic oxides also.

Carbonic acid gas not only forms a constituent part of the atmosphere near the surface of the earth, but at the greatest

heights which the industry of man has been able to penetrate. Saussure found it at the top of Mount Blanc, the highest point of the old continent; a point covered with eternal snow, and not exposed to the influence of vegetables or animals. Lime water diluted with its own weight of distilled water, formed a pellicle on its surface after an hour and three quarters exposure to the open air on that mountain; and slips of paper moistened with pure pot-ash, acquired the property of effervescing with acids after being exposed an hour and a half in the same place. This was at a height no less than 15,668 feet above the level of the sea. Humbolt has more lately ascertained the existence of this gas in air, brought by Mr. Garnerin from a height not less than 4280 feet above the surface of the earth, to which height he had risen in an air-balloon. This fact is a sufficient proof that the presence of carbonic acid in air does not depend upon the vicinity of the earth.

Now as carbonic acid gas is considerably heavier than air, it could not rise to great heights in the atmosphere unless it entered into combination with the air. We are warranted, therefore, to conclude that carbonic acid is not merely mechanically mixed, but it is chymically combined with the other constituent parts of the atmosphere. It is to the affinity which exists between carbonic acid and air that we are to ascribe the rapidity with which it disperses itself through the atmosphere, notwithstanding its great specific gravity. Fontana mixed 20,00 cubic inches of carbonic acid gas with the air of a close room, and yet half an hour after he could not discover the traces of carbonic acid in that air. Water impregnated with carbonic acid, when exposed to the air, very soon loses the whole of the combined gas. And when a phial full of carbonic acid gas is left uncorked, the gas, as Bergman first ascertained, very soon disappears, and the phial is found filled with common air.

It is owing to this strong affinity between air and carbonic acid gas, that it is so difficult to detect the presence of that gas in air by the common tests. Atmospheric air does not render lime water turbid, though agitated with it ever so long, or made to pass through it in ever so great a quantity. Neither has it any effect upon the most delicate vegetable blues. The great quantity of air with which it is combined, envelops it in such a manner that these bodies are not powerful enough to abstract it. We must employ for that purpose substances which have a very strong affinity for that acid, as the alkalis, milk of lime, &c. These substances detect its presence by acquiring the property of effervescing with acids.

The difficulty of separating this gas from air has hitherto prevented the possibility of

determining with accuracy the relative quantity of it in a given bulk of air; but from the experiments which have been made, we may conclude with some degree of confidence, that it is not very different from 0.01. From the experiments of Humbolt, it appears to vary from 0.005 to 0.01. This variation will by no means appear improbable, if we consider that immense quantities of carbonic acid gas must be constantly mixing with the atmosphere, as it is formed by the respiration of animals, by combustion, and several other processes which are going on continually. The quantity, indeed, which is daily formed by these processes is so great, that at first sight it appears astonishing that it does not increase rapidly. The consequence of such an increase would be fatal, as air containing 0.1 of carbonic acid extinguishes light and is noxious to animals. But there is reason to conclude, that this gas is decomposed by vegetables as rapidly as it forms.

4. *Bodies found in the atmosphere.*—From what has been advanced, it appears that the atmosphere consists chiefly of three distinct elastic fluids united together by chymical affinity; namely, air, vapour, and carbonic acid gas; differing in their proportions at different times and in different places; the average proportion of each is

98.6 air
1.0 carbonic acid
0.4 water

100.0

But besides these bodies which may be considered as the constituent parts of the atmosphere, the existence of several other bodies has been suspected in it. It is not meant in this place to include among those bodies electric matter, or the substance of clouds and fogs, and those other bodies which are considered as the active agents in the phenomena of meteorology, but merely those foreign bodies which have been occasionally found or suspected in air. Concerning these bodies, however, very little satisfactory is known at present, as we are not in possession of instruments sufficiently delicate to ascertain their presence. We can indeed detect several of them actually mixing with air, but what becomes of them afterwards we are unable to say.

1. Hydrogen gas is said to have been found in air situated near the crater of volcanoes, and it is very possible that it may exist always in a very small proportion in the atmosphere; but this cannot be ascertained till some method of detecting the presence of hydrogen combined with a great proportion of air be discovered.

2. Carbonated hydrogen gas is often emitted by marshes in considerable quantities during hot weather. But its presence

has never been detected in air; so that in all probability it is again decomposed by some unknown process.

3. Oxygen gas is emitted abundantly by plants during the day. There is some reason to conclude that this is in consequence of the property which plants have, of absorbing and decomposing carbonic acid gas. Now as this carbonic acid gas is formed at the expense of the oxygen of the atmosphere, as this oxygen is again restored to the air by the decomposition of the acid, and as the nature of atmospheric air remains unaltered, it is clear that there must be an equilibrium between these two processes; that is to say, all the carbonic acid formed by combustion must be again decomposed, and all the oxygen abstracted must be again restored. The oxygen gas which is thus continually returning to the air, by combining with it, makes its component parts always to continue in the same ratio.

4. The smoke and other bodies which are continually carried into the air by evaporation, &c. are probably soon deposited again, and cannot therefore be considered with propriety as forming parts of the atmosphere. But there is another set of bodies, which are occasionally combined with air, and which, on account of the powerful action which they produce on the human body, have attracted a great deal of attention. These are known by the name of *contagion*.

That there is a difference between the atmosphere in different places, as far as respects its effects upon the human body, has been considered as an established point in all ages. Hence some places have been celebrated as healthy, and others avoided as pernicious, to the human constitution. It is well known that in pits and mines the air is often in such a state as to suffocate almost instantaneously those who attempt to breathe it. Some places are frequented by peculiar diseases. It is known that those who are much in the apartments of persons ill of certain maladies, are extremely apt to catch the infection; and in prisons and other places, where crowds of people are confined together, when diseases once commence, they are wont to make dreadful havoc. In all these cases it has been supposed that a certain noxious matter is dissolved by the air, and that it is the action of this matter which produces the mischief.

This noxious matter is in many cases readily distinguished by the peculiarly disagreeable smell which it communicates to the air. No doubt this matter differs according to the diseases which it communicates, and the substance from which it has originated. Morveau lately attempted to ascertain its nature; but he soon found the chymical tests hitherto discovered al-

together insufficient for that purpose. He has put it beyond a doubt, however, that this contagious matter is of a compound nature, and that it is destroyed altogether by certain agents, particularly by those gaseous bodies which readily part with their oxygen. He exposed infected air to the action of various bodies, and he judged of the result by the effect which these bodies had in destroying the fetid smell of the air. The following is the result of his experiments.

1. Odorous bodies, such as benzoin, aromatic plants, &c. have no effect whatever.
2. Neither have the solutions of myrrh, benzoin, &c. in alcohol, though agitated in infected air.
3. Pyrolignous acid is equally inert.
4. Gunpowder, when fired in infected air, displaces a portion of it; but what remains still retains its fetid odour.
5. Sulphuric acid has no effect; sulphurous acid weakens the odour, but does not destroy it.
6. Acetous acid diminishes the odour, but its action is slow and incomplete.
7. Acetic acid acts instantly, and destroys the fetid odour of infected air completely.
8. The fumes of nitric acid, first employed by Dr. Carmichael Smith, are equally efficacious.
9. Muriatic acid gas, first pointed out as a proper agent by Morveau himself, is equally effectual.
10. But the most powerful agent is oxy-muriatic acid gas, first proposed by Mr. Cruickshanks, and now employed with the greatest success in the British navy and military hospitals.

Thus there are four substances which have the property of destroying contagious matter, and of purifying the air: but acetic acid cannot easily be obtained in sufficient quantity, and in a state of sufficient concentration to be employed with advantage. Nitric acid is attended with inconvenience, because it is almost always contaminated with nitrous gas. Muriatic acid and oxy-muriatic acid are not attended with these inconveniences; the last deserves the preference, because it acts with greater energy and rapidity. All that is necessary is to mix together two parts of salt with one part of the black oxide of manganese, to place the mixture in an open vessel in the infected chamber, and to pour upon it two parts of sulphuric acid. The fumes of oxy-muriatic acid are immediately exhaled, fill the chamber, and destroy the contagion.

ATROCHIA. (From *a*, neg. and *trachō*, off-spring; from *trachō*, to bring forth.) Inability, to bring forth children. Difficult labour.

ATONIC. Relaxed, diminution of strength, weakness, debility.

ATONY. (From *a*, neg. and *tonō*, to extend.) A defect of muscular power.

ATRAHILIARIE CAPSULE. See *Glandule renales*.

ATRABILIS. Black bile, or melancholy.

ATRACHELUS. (From α , priv. and $\tau\rho\alpha\chi\eta\lambda\omicron\varsigma$, the neck.) Short-necked.

ATRAGENE. *Viora*. *Clematis orthragene* of Theophrastus. The *Clematis vitalba* of Linnaeus. The traveller's joy: a common shrub in our hedges. It is said to have caustic qualities and to raise a blister when applied to the skin.

ATRAMENTUM SUTORIUM. A name of green vitriol.

ATRAIA. (From α , neg. and $\tau\rho\alpha\pi\alpha$, to perforate.) *Astria*. Imperforation. A disease where the anus or genitals have not their usual orifice.

ATRETARUM. (From α , neg. and $\tau\rho\alpha\omega$, to perforate.) A suppression of urine from the menses being retained in the vagina.

ATRICES. (From α , priv. and $\sigma\pi\epsilon\varsigma$, hair.) Small tubercles about the anus upon which hairs will not grow. *Vasellus*.

ATRICI. Small sinuses in the rectum, which do not reach so far up as to perforate into its cavity.

ATRIplex FETIDA. *Atriplex olida*. *Vulvaria*. *Garsenum*. *Raphex*. *Chenopodium fatidum*. *Bithum fatidum*. Stinking orach. The very fetid smell of this plant, *Chenopodium vulvaria*: *foliis integerrimis rhombeo-ovatis, floribus conglomeratis axillaribus*, of Linnaeus, induced physicians to exhibit it in hysterical diseases. It is now superseded by more active preparations.

ATRIplex HORTENSIS. The systematic name for the *atriplex sativa* of the pharmacopias. See *Atriplex sativa*.

ATRIplex SATIVA. The herb and seed of this plant, *Atriplex hortensis caule erecto herbaceo, foliis triangularibus*, of Linnaeus, have been exhibited medicinally as antiscorbutics, but the practice of the present day appears to have totally rejected them.

ATROPA. (From $\alpha\tau\rho\rho\epsilon$, the goddess of Destiny; so called from its fatal effects.) The deadly night-shade.

The name of a genus of plants in the Linnaean system. Class, *Pentandria*. Order, *Monogymia*.

ATROPA BELLADONNA. The systematic name for the *belladonna* of the pharmacopias. See *Belladonna*.

ATROPA MANDRAGORA. The systematic name for the plant which affords the *radic mandagore* of the pharmacopias. See *Mandragora*.

ATROPHIA. See *Atrophy*.

ATROPHY. (From α , neg. and $\tau\rho\alpha\sigma$, to nourish.) *Atrophia*. *Marasmus*. Nervous consumption. This disease is marked by a gradual wasting of the body, unaccompanied either by a difficulty of breathing, cough, or any evident fever, but usually attended with a loss of appetite and impaired digestion. It is arranged by Cullen

in the class *cachexia*, and order *macrures*. There are four species:—

1. When it takes place from too copious evacuations, it is termed *atrophia inanitionum*; by others called *tubes nutricum*;—*eductorior*;—*a sanguifluxu*, &c.

2. When from famine, *atrophia famelicorum*.

3. When from corrupted nutriment, *atrophia cacochymica*.

4. And when from an interruption in the digestive organs, *atrophia debilitum*.

The atrophy of children, is called *pald-atrophia*. The causes which commonly give rise to atrophy, are a poor diet, unwholesome air, excess in venery, fluor albus, severe evacuations, continuing to give suck too long, a free use of spirituous liquors, mental uneasiness, and worms; but it frequently comes on without any evident cause. Along with the loss of appetite and impaired digestion, there is a diminution of strength, the face is pale and bloated, the natural heat of the body is somewhat diminished, and the lower extremities are oedematous. Atrophy, arise from whatever cause it may, is usually very difficult to cure, and not unfrequently terminates in dropsy.

ATTENUANTS. (*Attenuantia*, sc. *medicamenta*; from *attenuo*, to make thin.) Diluents. Those substances are so termed, which possess a power of imparting to the blood a more thin and more fluid consistency than it had previous to their exhibition; such are, *aqua*, *serum lactis*, &c.

ATTOLLENS AURIS. (*Attollens*; from *attollo*, to lift up.) *Attollens auriculæ* of Albinus and Douglas; *Superior auris* of Winslow, and *Attollens auriculam* of Cowper. A common muscle of the ear, which arises, thin, broad, and tendinous, from the tendon of the occipito-frontalis, from which it is almost inseparable, where it covers the spongy process of the temporal muscle; and is inserted into the upper part of the ear, opposite to the antihelix. Its use is to draw the ear upwards, and to make the parts into which it is inserted, tense.

ATTOLLENS OCULI. One of the scoti-muscles which lie upon the upper part of the globe and pulls up the eye.

ATTONTUS MORBUS. (From *attono*, to surprise; so called because the person falls down suddenly.) *Attonitus stupor*. The apoplexy and epilepsy.

ATTRACTION. (From *attraho*, to attract. Affinity.)

The terms attraction, or affinity, and repulsion, in the language of modern philosophers, are employed merely as the expression of the general facts, that the masses or particles of matter have a tendency to approach to, or to recede from one another, and to unite to, or repel each other, under certain circumstances.

All bodies have a tendency or power to attract each other more or less, and it is this power which is called attraction.

Attraction is mutual, it extends to indefinite distances. All bodies whatever, as well as their component elementary particles, are endued with it. It is not annihilated, at how great a distance soever we suppose them to be placed from each other; neither does it disappear though they be arranged ever so near each other.

The nature of this reciprocal attraction, or at least the cause which produces it, is altogether unknown to us. Whether it be inherent in all matter, or whether it be the consequence of some other agent, are questions beyond the reach of human understanding; but its existence is nevertheless certain.

Proofs of attraction.

That the power of attraction really exists is obvious from the slightest view of the phenomena of nature. It is proved with mathematical certainty that the celestial bodies, which constitute the solar system, are urged towards each other by a force which preserves them in their orbits. It is further proved beyond any doubt, that this planetary attraction is possessed not only by the heavenly bodies as wholes, but that it also extends to the smaller particles of which they are formed, as may be evinced by means of the following experiments.

First.—If we place two or more globules of mercury on a dry glass or earthen plate, and push them gently towards each other, the globules will attract each other, and form one mass or sphere greater in bulk but precisely the same in nature.

Secondly.—If a plate of clean glass, perfectly dry, be laid on a large globule of mercury, the globule, notwithstanding the pressure applied to it, continues to preserve its spherical form; if we gradually charge the plate with weights carefully, the globule will be depressed and become thinner and thinner; but if we again remove the weights from the plate, the mercury will instantly recover its globular figure and push up the glass before it.

In both these experiments we see that there exists an attraction between the particles of mercury; in the first, the globules which are in contact with the plate of glass leave this substance completely, they attract each other and form a sphere greater in bulk. A mere inert fluid would in any case retain the figure it once possessed. It could not be endued with a globular form unless a real reciprocal attraction among its particles, took place, which in the latter experiment is still more striking, for it there is not only superior to gravitation, but actually overcomes an external force.

Thirdly.—If a glass tube of a fine bore

be immersed in water, contained in any vessel, the fluid will ascend to a certain height within the tube above its level, and its elevation in several tubes of different sizes will be reciprocally as the diameter of their bores.

This kind of attraction which takes place as well *in vacuo* as in the open air, has been called capillary attraction. It is this attraction which causes water to rise in sponge, cloth, sugar, sand, &c. for all these substances may be considered as fine tubes in which the fluid ascends.

Remark.—The ascension of fluids in glass tubes of a fine bore succeeds best when the inside of the tube has been previously moistened, which may conveniently be done by blowing through it with the mouth. And if the water be coloured with a little red or black ink, its ascension will be more obvious, particularly if the tube be held against a sheet of white paper.

Fourthly.—If two plates of glass previously wetted, be made to meet on one side, and be kept open at the other, at a small distance, by the interposition of a shilling, or any other thin substance, and then immersed in water, the fluid will ascend between the two plates unequally. Its upper surface will form a curve, in which the heights of the several points above the surface of the fluid will be to one another reciprocally, as their perpendicular distance from the line in which the plates meet. The ratio of this attraction is therefore as the squares of the increments with which the plates open.

Here then we have two other instances that an attraction prevails among the particles of bodies. For in both cases part of the fluid has left the contiguous mass, contrary to the laws of gravitation. It is drawn up as it were, or attracted by the tube or plate of glass.

Fifthly.—If we immerse a piece of tin, lead, bismuth, silver, or gold, in mercury, and draw it out again immediately, the mercury will attract the metal, and the latter will carry with it a portion of the former which will stick to it so obstinately as to be inseparable by mere friction.

There exists therefore an attraction between the different metals brought in contact with each other.

Sixthly.—If a small stick be dipped in water or any other fluid, and drawn out again, a drop will be found hanging at the end of it of a spherical form. The drop is spherical, because each particle of the fluid exerts an equal force in every direction, drawing other particles towards it on every side as far as its power extends.

Thus the very formation of drops obviously demonstrates that there must exist a cause which produces that effect. This

cannot be gravity, for agreeable to experience that is rather an obstacle to the formation of drops; since by the weight of the particles, large globules resting on solid bodies are flattened, and their regular spherical form prevented.

To explain this phenomenon there remains only the power of attraction, acting between the particles of the liquid body; for if it is supposed that the particles of a substance reciprocally attract each other with equal force, and their aptitude for being moved upon one another be great enough to overcome any impediment to their motion, it follows by the principles of mechanics, that the equilibrium of the attractive forces can only take place when the mass has received a globular form.

Hence it is that all liquid bodies assume a spherical figure when suffered to fall through the air, or form drops.

Division of attraction.

Though we are unable to discover the cause of the mutual attraction, experience has proved to us that this agency follows certain conditions or laws; for similar phenomena always present themselves, whenever the circumstances of experiment are the same.

Observation has taught us that attraction takes place between bodies of the same kind, and bodies of a different kind. The first is called *attraction of aggregation*, also *corpuscular attraction*; *molecular attraction*; and *attraction of cohesion*, or the *cohesive power*.

The latter is termed *chymical attraction*, *chymical affinity*, or *affinity of composition*.

ATTRACTION OF AGGREGATION.

Corpuscular attraction, or attraction of cohesion or aggregation, is that power by means of which the similar particles of bodies attract each other, and become united into one mass, without changing in the least the chymical properties they possessed before their union. The bodies may be in a solid, fluid, or æreiform state.

This attraction is different in different bodies. It is always in an inverse ratio to the power of repulsion, or the quantity of caloric interposed between the particles of the acting bodies.

It becomes obvious from this, that the agency of attraction of aggregation consists in a mere successive and constant accumulation of similar particles into one mass; and that it produces adherence of surface, or apparent contact in the ratio of the surface.

This force is inherent in all the particles of all bodies (caloric and light perhaps excepted;); we never find the particles of bodies in a detached state, but constantly in masses of greater or smaller magnitude, made up of an indefinite number of particles united together by virtue of the force of cohesion.

The simplest case of the exertion of the attraction of aggregation is that, where two bodies placed in mutual contact with each other form a direct union without changing their chymical properties: thus if different particles of sulphur be melted together, they form a uniform mass or whole, the particles of which are held together by virtue of the power of attraction of aggregation, but the properties of the body are not altered.

The same effect takes place when pieces of the same metal, or particles of resin, wax, &c. are united in a similar manner.

The force of this attraction in solid bodies may be measured by the weight necessary to demolish it. Thus if a rod of metal, glass, wood, &c. be suspended in a perpendicular direction, and weights be attached to its lower extremity till the rod is broken by them, the weight attached to the rod just before it broke is the measure of the cohesive force of the rod.

Laws of attraction of aggregation.

1. The agency of attraction of aggregation acts only at insensible distances; its force increases as the distance of the bodies presented to each other decreases, and as the surfaces of apparent contact are more numerous: thus, if we take two sections of a leaden ball, having each a flat and smooth surface, and press them forcibly together, they will cohere, and a considerable effort is necessary to force them asunder: so also two plates of glass wetted with a little water to fill up their inequalities, when laid together, will cohere; and two pieces of marble having each a flat, smooth, and well polished surface, when moistened and slipped upon each other with a gentle pressure, will unite, and a considerable force is required to separate them. But if the two substances placed together, be not sufficiently smooth or polished, it will be in vain to try to cause them to adhere together, for this reason that the particles touch each other only in a few points; whereas on the contrary the particles of the former flat and smooth surfaces touch each other in many points. It has been noticed that a silk-worm's thread can be interposed, but not two.

The pressure of the atmosphere has no influence on these experiments, for they succeed equally well in *vacuo* as in the open air.

It is on this account that carpenters when they intend to glue pieces of wood together, plane the surfaces perfectly smooth before they apply the glue: and that the surfaces of metals are scraped clean before they are soldered, &c.

Hence the attraction of aggregation always vanishes whenever the distance is measurable, and becomes exceedingly great whenever the distances is exceedingly diminished; but the particular rate which this power follows, is still unknown;

as we have no method of measuring either the distance at which it acts, or its relative intensity.

2. Attraction of aggregation acts differently in different bodies; according to the degree of force with which it acts between the particles of matter, the bodies appear under different forms.

It is on this account that rock-crystal, flint, diamond, and various other precious stones are extremely hard, for the attraction of aggregation unites the particles of these bodies with a great degree of force. Hence a considerable mechanical effort is necessary to disunite them.

In blocks of marble, chalk, lime-stone, &c. the particles are held together with a force considerably less. In these bodies it prevents all relative motion among the particles themselves, and hence the motion of one particle is followed by the motion of the whole mass; or if that is impossible, the cohesion is destroyed altogether, and the piece breaks.

The integrant parts of wax, tallow, suet, or lard, may be made to change their situations, with a less degree of force than the former.

In these substances the motion of one particle of the body is not necessarily followed by that of all the rest, neither does that motion destroy the cohesion, nor break them.

The particles of water, spirit, and ether, move or slide over each other very readily; hence their resistance is considerably less.

And lastly, vapours, the air of the atmosphere, and all the gasses, yield to the slightest possible impulse.

3. Attraction of aggregation may be annihilated by every effort which tends to separate the particles of bodies.

It need hardly be mentioned that all mechanical forces, such as grinding, cutting, filing, rasping, pounding, breaking, &c. are of this nature.

In all these cases the force applied must be more than equal to the force of the attraction; and as it was stated before, that the attraction of aggregation acts with different degrees of force between the particles of different bodies, so different degrees of force are necessary to destroy that attraction in different bodies; and hence it is that chalk is more easily reduced to powder than flint; wood is easier broken than lead; lead easier than iron, &c.

CHEMICAL AFFINITY.

Chymical affinity, or affinity of composition, is that power, by means of which the particles of bodies, whether simple or compound, attract each other so intimately as to produce a uniform whole, totally inseparable by mechanical efforts, and whose characteristic properties are often different, and sometimes contrary to those of its constituent parts.

It is obvious from this, that the particles of those bodies which are united by virtue of chymical affinity, form not a mere aggregate, but an entire new body, which can only be altered by the action of another chymical power.

In considering this kind of affinity, it will be necessary to state;—In what manner it takes place between the particles of different bodies;—In what proportion they are capable of combining;—Under what conditions;—With what degree of force they unite;—And what takes place when a variety of different substances are made to act upon each other at the same time, under certain circumstances and in different proportions.

Hence chymical affinity is of greater importance than affinity of aggregation, for it takes place in all the complex operations of chymistry.

Instances of chymical affinity.

To prove that chymical affinity acts differently from attraction of aggregation; that it takes place between the ultimate constituent parts of bodies; and that it produces substances possessing properties, frequently very different and sometimes contrary to those of the constituent parts, the following experiments may serve.

1. Put into a crucible placed in a coal fire, equal parts by weight of sulphur and mercury; stir the two substances together for a few minutes, and when the sulphur is melted, pour the contents out on a marble slab, or a piece of glass previously warmed and greased.

The substance obtained by this means is a *sulphuret of mercury*, in which the mercury and sulphur are united by virtue of chymical affinity; for the compound has neither the colour, the splendour, the inflammability, the volatility, nor the specific gravity of either of its constituent parts; nor can the sulphur and mercury be separated by mechanical means; they are therefore chymically united.

2. If we melt together two very malleable and ductile metals, for instance, tin and iron, in equal quantities, the compound produced will have totally lost the properties which its constituent parts possessed before their union, for the alloy formed will be a brittle metal which may easily be broken by the blow of a hammer.

3. Put two or three teaspoonfuls of an aqueous infusion of red cabbage or syrup of violets, into a wine glass of water, mix it well, and put half the mixture into another glass. By adding a few drops of sulphuric acid to one of the glasses and stirring it, the blue will be changed to a crimson; and by adding an alkali, for instance, potash, to the other glass, the blue fluid will be changed into a green.

If we drop carefully down the sides of the glass into the green obtained in this experiment, a few drops of sulphuric acid,

crimson will be perceived at the bottom, purple in the middle, and green at the top. On adding a little alkali to the other glass, containing the crimson, these colours will appear in an inverted order.

4. When equal parts of muriate of ammonia and slaked lime, both substances destitute of odour, are intimately blended in a stone mortar, a very pungent gas (ammonia) becomes evolved.

5. Water impregnated with ammonia and concentrated muriatic acid, both fluids of a strong odour, when mixed together in proper proportions, instantly lose their odour, and form a fluid void of smell, (muriate of ammonia.)

6. Into a saturated solution of muriate of lime, let fall gradually concentrated sulphuric acid, a quantity of pungent vapour will become disengaged, (muriatic acid gas,) and from the two fluids will thus be produced an almost solid compound, called sulphate of lime.

7. Let equal parts of fresh crystallized acetat of lead and acidulous sulphate of alumine and potash, be rubbed together intimately in a stone mortar, the saline mixture will soon become soft, and lastly fluid.

A like effect is produced by treating in a similar manner equal parts of crystallized nitrate of ammonia and sulphate of soda.

A solid alloy of mercury and bismuth, and another composed of lead and mercury, on being triturated together, instantly become fluid.

It is obvious from this, that when chymical combination takes place, the compound which is formed does not possess properties merely intermediate between those of its component parts, but has acquired others more or less new. This however does not hold good in all cases. There are various combinations in which the properties of bodies are only slightly altered.

Laws of chymical affinity.

Observation has shown that affinity of composition offers certain invariable phenomena, which being founded on a great number of facts are regarded by chymists as laws, and may be reduced under the following heads:

Law I.—Chymical affinity can exert its action between a number of bodies, simple or compound, and unite them chymically into one whole.

Law II.—The efficacy of chymical affinity is in an inverse ratio to that of attraction of aggregation.

Law III.—The agency of chymical affinity is influenced by temperature; its action is either accelerated, retarded, prevented, or rendered efficacious.

Law IV.—Chymical affinity is accompanied by a change of temperature at the instant of its action.

Law V.—The agency of chymical affinity existing between two or more bodies may be dormant, until it is called into action by the interposition of another body which frequently exerts no energy upon any of them in a separate state.

Law VI.—The ratio of the energy of chymical affinity acting between various bodies, is different in different substances.

Law VII.—The agency of chymical affinity is either limited, or unlimited in certain bodies; in other words, chymical affinity is capable of uniting bodies in definite, or in indefinite proportions.

Law VIII.—The energy of the chymical affinity of different bodies is modified in proportion to the ponderable quantities of the bodies placed within the sphere of action.

Such are the leading laws which regulate chymical affinity; they may be demonstrated by experiments.

1. Chymical affinity can exert its action between a number of bodies simple or compound, and unite them chymically into one whole;

There are an infinite variety of compounds, consisting of three, four, five or more simple substances in nature; and art can also effect combinations in which there are many simple bodies chymically united into one whole.

It frequently happens that various separate bodies presented to each other in a fluid, unite and form a single mass, which possesses all the characters of a homogeneous compound, and which retains these characters till its composition has been altered by chymical means.

A considerable number of triple salts are known, which consists of three different substances; for instance, the common alum of commerce consists of sulphuric acid united to alumine and potash or soda. The salt formerly called microcosmic salt, or phosphate of soda and ammonia, consists of phosphoric acid united to soda and ammonia, &c. When the oxygenated muriate of mercury is precipitated by the precise quantity of carbonate of soda which is requisite to effect its decomposition, the precipitate obtained contains muriatic acid, carbonic acid, and oxyd of mercury in excess,

It is a well known fact, that two, three, or more metals may be fused together so as to produce compounds whose properties are widely different from those of the constituent parts.

Melt together in an iron ladle or crucible, eight parts of bismuth, five of lead, and three of tin, the fusibility of the metals will thus be altered, for the alloy melts at 212° Fahr. A spoon or any other utensil formed of this compound will therefore melt in water kept boiling.

If in a similar manner an alloy be made of lead, tin, bismuth, and mercury, their proportions being two, three, five, and one, the compound produced melts at a heat even less than that of boiling water.

A composition of lead, zinc, and bismuth, in equal parts may be kept in fusion upon paper over a lamp.

II. The efficacy of chymical affinity is in an inverse ratio to that of corpuscular attraction:—

The cohesion of the particles of a body is owing to the mutual affinity existing between them. It is this force which must be overcome by the action of the substance which has a tendency to combine with those particles chymically. Chymical affinity therefore does not become *stronger* as the affinity of aggregation becomes *weaker*, it becomes only *more efficacious*; the absolute powers remain the same; the effect produced by that agency *increases*, because the resistance opposed to it *decreases*.

Remark.—It is from this law that it was formerly inferred that some or at least one of the bodies should be in a state of fluidity. This however is by no means necessary. It is in general true, that the weaker the attraction of aggregation is, the more easily chymical affinity takes place, as may be evinced by means of the following experiments:

Let any quantity of dry carbonate of soda and tartareous acid be mingled together, and put the mixture into a wine-glass, no chymical change will be produced; but if water be added, or either of the salts be previously dissolved, a violent effervescence ensues, and a chymical union is obtained.

The water added is of use merely to overcome the resistance which arises from the cohesion of the particles of the salts intended to be brought into the sphere of action, or to increase their mutual contact.

If we let fall a crystal, or lump of fluor spar (fluat of lime) into concentrated sulphuric acid, no sensible action will take place, both the sulphuric acid and the fluat of lime remain unaltered; but if the former be reduced to powder, and then brought into contact with the acid, a considerable action instantly takes place, the sulphuric acid unites to one of the constituent parts of the fluor spar, namely, to the lime, and its other constituent part, the fluoric acid, becomes disengaged in the state of white vapour, or fluoric acid gas.

If crystallized sulphate of alumine, or sulphate of soda, and acetate of lead are brought into contact with each other, the individuality of these bodies will not be destroyed, that is to say, no chymical change will take place; but if they be intimately rubbed together in a mortar, the

two solids will act upon each other and form a fluid.

It is obvious therefore that in order to facilitate chymical affinity, the attraction of aggregation must be broken; the bodies intended to be chymically united must not be presented to each other in their mass of contact, but mechanically divided, or reduced to the smallest molecule possible: hence liquids combine with more facility than solids, or even than a solid and a liquid, and in like manner vapours combine with rapidity and ease.

III. The agency of chymical affinity is influenced by temperature. Its action is either accelerated, retarded, prevented, or rendered efficacious:—

If we expose phosphorus in an open vessel to the action of the atmosphere, a chymical union will take place between the phosphorus and one of the constituent parts of the atmosphere, namely, the oxygen gas; the phosphorus will gradually (but very slowly) disappear and become converted into a fluid called phosphorus acid.

But if we heat the vessel containing the phosphorus, the latter will take fire, and become converted into a yellowish white substance, which in a short time is changed into an acid analogous to the former.

If equal quantities of muriate of ammonia and carbonate of magnesia are mixed with six or eight parts of water, and suffered to stand for some time exposed to the ordinary temperature of the atmosphere, a mutual decomposition of the two salts will take place. For if the mixture, and the fluid which passes, are left to evaporate spontaneously, muriate of magnesia, and carbonate of ammonia will be obtained. On the contrary,

If equal quantities of muriate of magnesia and carbonate of ammonia be exposed to a temperature of 200° in about four parts of water, the products obtained are, muriate of ammonia and carbonate of magnesia.

If muriate of soda and sulphate of magnesia be mixed together in any proportion, and exposed to a temperature below zero, they decompose each other, and muriate of magnesia and sulphate of soda are formed, but no decomposition takes place at a temperature above 30°.

Muriate of soda and acidulous sulphate of alumine and potash, exhibit precisely the same phenomena.

If ardent spirit and a solution of salt and water be mixed together, the compound formed is a real chymical union; but if we carefully heat the fluid, the caloric applied will be divided between the three ingredients according to their respective affinities; the union will be broken, for the ardent spirit will first be-

come volatilized, and the union of the salt and water remain unaltered. On increasing the temperature, the water will escape in the form of vapour, and the salt will be left behind.

There are numerous cases in which an increase of temperature is essentially necessary to determine bodies to unite. If mercury be exposed to oxygen gas at the common temperature of the atmosphere, the corpuscular attraction subsisting between its particles is sufficient to prevent combination. But if the mercury be heated to a certain degree, the force which kept its particles united will become annihilated, and it then combines with the oxygen which is present.

Again, if the oxyd of mercury thus formed be exposed to a higher degree of temperature, the union is demolished, and the quick-silver re-appears in its metallic state.

Hence it is obvious that the action of caloric favours the union of the oxygen and mercury, in consequence of the diminution of the mutual affinity of the parts; but at length, by augmenting this difference, it again breaks the union, or renders the combination impossible.

That increased temperature augments the power of chymical union, the solutions of salt and water afford instances of.

A larger quantity of salt is soluble in a given quantity of water at a high, than at a low temperature, and this larger quantity of salt is again separated by cooling.

IV. Chymical affinity is accompanied by a change of temperature at the instant of its action:—

When equal parts of concentrated sulphuric acid and ardent spirit are mingled together, the mixture in a few minutes becomes so hot as to render the vessel insupportable to the hands.

If four parts of sulphuric acid of commerce, and one part, by weight, of water, be mixed together, each at the temperature of 50°, the mixture immediately acquires a temperature of about 300°.

All the dense acids, ammonia, and ardent spirit, when mixed with water, have the property of raising its temperature remarkably: and the same is the case when alkalis are introduced into concentrated acids. On the contrary, in many instances cold is produced:—

Take one ounce and a half of muriate of ammonia and a like quantity of nitrate of potash; reduce each of these salts separately to a powder, and blend them intimately together: having done this, mix them gradually in a glass basin, or other thin glass vessel, with four ounces of water. The result will be, that the cold produced will sink a thermometer immersed in it, to 36° Fahr. A new edition of the same quantity of salts will cool it to 14°,

which therefore will freeze water in a glass tube that is immersed in it, without the use of snow or ice. If the water used in a first process be used to reduce other water and salts to the temperature of about 32°, and these be applied to the performance of a second experiment, the temperature may be lowered to 4° below 0°.

A number of experiments have lately been made to produce artificial cold by means of such freezing mixtures. The most complete act of this kind are those of Pepys, Lowitz and Walker.

V. The agency of chymical affinity between two or more bodies may lie dormant, until it is called into action by the interposition of another body, which frequently exerts no energy upon any of them in a separate state.

From this law originates what was formerly called disposing affinity, or that case in which two or more bodies are incapable of uniting, until the agency is called into action by the addition of a third body, which exerts no sensible affinity upon either of them. This may be proved in the following manner.

Water is a compound of hydrogen and oxygen; phosphorus is a simple body according to our present state of knowledge. If these be presented to each other, no chymical union will take place; but if we add to them an alkali, and then apply heat, the water will become decomposed; that is to say, part of the phosphorus will unite to the oxygen of the water, and form phosphoric acid, and the other part will be dissolved in the hydrogen gas and appear as phosphorated hydrogen.

Here the alkali acts as the substance requisite to favour the mutual action, or to give the disposing affinity.

If iron and water be brought into contact with each other no perceptible change will be produced; but if a little sulphuric acid be added to the water and iron, a violent effervescence will take place, the water will become decomposed, hydrogen gas will be evolved, and the iron become dissolved in the acid.

In this case the sulphuric acid is the condition necessary to accelerate the chymical action.

VI. The ratio of the energy of chymical affinity acting between various bodies, is different in different substances.

This is the most important law of chymical attraction. As beginners will find it rather difficult to understand what passes in this more complicated agency, they must remember, that the combination which is effected between two or more bodies by virtue of chymical affinity becomes broken whenever we present to the compound another body, which has an attraction to one of the constituent parts of the compound, superior to that attraction

by which they were held together: the bodies, therefore, between which the strongest attraction prevails, combine, and the rest are disengaged, thus :—

If muriatic acid be poured either on pure barytes, or on its carbonate, the barytes will be dissolved and the compound will be muriate of barytes, which compound is held together by the force of affinity existing between the muriatic acid and the barytes. On letting fall into this solution a few drops of sulphuric acid, an immediate change of principles takes place; the whole quantity of the muriatic acid which was combined with the barytes becomes disengaged, and the sulphuric acid unites to the barytes with a force equal to their affinity, *minus* that of the muriatic acid.

Again, if pure silver be dissolved in pure nitric acid, the silver will remain united to the acid, till another body is presented to it which has a greater force of attraction to one of the constituent parts of the compound; for instance, if mercury be added to this solution of silver, the mercury will be dissolved, and the silver becomes precipitated or disengaged. The supernatant fluid will then be a *solution of mercury in nitric acid*.

If to the before obtained solution of mercury in nitric acid, a piece of sheet lead be presented, the lead will be dissolved, and the mercury become precipitated. The fluid will then be a *solution of lead in nitric acid*.

If in this solution of lead, a thin slice of copper be suspended, the copper will be dissolved, and the lead will become disengaged. The fluid now is a *solution of copper in nitric acid*.

If in this solution of copper, a thin sheet of iron be kept immersed, the iron will be dissolved, and the copper become precipitated. The fluid now is a *solution of iron in nitric acid*.

If to this solution of iron, a piece of zinc be presented, the zinc will be dissolved, and the iron become precipitated. The solution then consists of *zinc and nitric acid*.

If to this solution of zinc in nitric acid some ammonia be gradually added, the ammonia will join to the acid, and the zinc will be precipitated. The solution will then be *nitrate of ammonia*.

If to this solution of nitrate of ammonia, some lime-water be added, the ammonia will become disengaged (and manifest itself by a pungent odour) and the solution will be *nitrate of lime*.

If to this solution of nitrate of lime some oxalic acid be added, the lime will be precipitated, and what now remains will be merely *nitric acid*.

We see from these experiments, that

different bodies have different degrees of affinity for one and the same substance, which can only be learnt from observation and experiments.

VII. The agency of chymical affinity is either limited or unlimited; in other words, chymical affinity is capable of uniting bodies in definite, or in indefinite proportions :—

Experience has convinced us that in all bodies there are certain precise limits of combinations beyond which their action cannot pass, namely a *minimum*, and a *maximum*; it remains still to be ascertained how bodies can combine within these limits.

If we attend to what is known at present, we are forced to acknowledge that this law comprehends several modifications, which may be arranged under the following classes.

1. Chymical affinity unites several bodies in any proportion whatsoever; their combination is therefore unlimited; for instance,

If water and ardent spirit be mingled together in any quantity, a chymical combination ensues; for the compound obtained has always a specific gravity different from the mean specific gravity of the fluids combined. Its bulk is likewise not the arithmetical mean of the fluids in a separate state.

The same is the case when liquid acids and water, or acids and ardent spirit, are combined together.

2. Chymical affinity combines several bodies to a certain extent or *maximum* only.

To this class belong all those bodies which are capable of saturation.

If we take a quantity of any of the dense acids diluted with water, for instance, sulphuric acid, and let fall into it a solution of an alkali, for example soda, by a little at a time, and examine the mixture after every addition of the alkali, we find for a considerable time it will exhibit the properties of an acid, it will have a sour taste, and convert vegetable blue colours into red; but if we continue to add greater quantities of soda, these acid properties will gradually diminish, and at last disappear altogether. At that point the combination is at an end, it has reached its *maximum* in this case; for if we continue to add more alkali, the mixture will gradually acquire alkaline properties; it will convert blue vegetables into green; it will have a urinous or alkaline taste, &c. These properties will become stronger, the greater the quantity of the soda which is added.

Again, take muriatic acid, and let fall into it gradually carbonate of lime, or magnesia; an effervescence will take place, for a chymical union ensues between the acid and the lime, or magnesia, and the

carbonic acid, the other constituent of these bodies, becomes disengaged. But if we continue the addition of the lime, or magnesia, until it produces no further effervescence, no chymical union will be obtained; the lime will fall to the bottom unaltered, for the combination is at its *maximum*.

It is on this account that water can only dissolve a certain quantity of salt; ardent spirit a certain quantity of resin, &c.

The union of oxygen and hydrogen belongs likewise to this class.

3. Chymical affinity is capable of uniting different bodies in two, three, or more proportions; each of these combinations produces compounds, possessing peculiar properties.

This peculiarity of combination is highly important.

It is owing to this circumstance that both nature and art produce substances of the same principles, only combined in different proportions, which possess peculiar properties, widely different from each other.

An instance of this law may be seen in the following experiment:

Introduce one ounce of copper filings into four ounces of muriatic acid, contained in a medicine-phial of eight ounces capacity, cork it well, and let it stand undisturbed; the acid will soon acquire a greenish colour, which becomes deeper in proportion as the copper becomes dissolved; but in a few days if the bottle be now and then agitated, the colour vanishes, and the solution at last becomes colourless.

If we now invert the bottle in mercury, or water, and remove the cork under that fluid, a quantity of the mercury will rush in: an evident proof that part of the air contained in the phial has disappeared.

If we examine the remaining air, we shall find that it is incapable of supporting flame, and that it is nearly deprived of all its oxygen. If we now open the phial, the solution becomes again green and colourless as before.

The *rationale* of these phenomena is this: The quantity of oxygen which is present in the confined quantity of air in the empty part of the phial, combines with the copper to a certain degree, which then becomes soluble in the acid, and exhibits the green solution.

This oxyd is gradually decomposed, more copper is dissolved, and the solution becomes colourless. If more oxygen be admitted, the solution becomes green again as before.

VIII. The energy of the chymical affinity of different bodies is modified in proportion to the ponderable quantities of the substances placed within the sphere of action.

It is obvious, from this, that the denomination of elective affinity is erroneous; since it supposes the union of one entire substance with another, in preference to a third. But this is not the case; a mere division of action takes place in instances of this kind; that is to say, the substances act according to the quantity existing within the sphere of activity. The excess of quantity is capable of compensating for the deficiency of the force of affinity.—When, therefore, a compound body of two substances is acted on by a third, that part of the compound which is the subject of combination, is divided between the two remaining, not only in proportion to their respective degrees of affinity, but also according to their ponderable quantities, so that by varying this in either, the effect produced will be varied.

Thus Berthollet has proved, that in all cases a large quantity of a body is capable of abstracting a portion of another, from a small portion of a third, how weak soever the affinity between the first and second of these bodies may be, and how strong soever the affinity between the second and third. Thus potash is capable of abstracting part of the acid from oxalate of lime, phosphate of lime, and carbonate of lime. Soda and lime decompose partially sulphate of potash. Nitric acid subtracts part of the base from oxalate of lime, &c.

The following experiment, advanced by Berthollet, will prove this more clearly.

If equal parts, by weight, of sulphate of barytes and potash be boiled, in a small quantity of water, to dryness, it will be found that the sulphuric acid has been divided between the two bases in the compound ratio of their mass, and their force of affinity. The greater part of the sulphate of barytes will be found undecomposed; a small quantity of barytes will be found at liberty; most of the potash will also be uncombined, but a certain portion will be united with the sulphuric acid which the barytes has lost, in the form of sulphate of potash.

It is not merely in the instance stated here that this division of one body between two others, according to their respective masses and affinities, takes place, there being scarcely any example to the contrary.

And as the affinities of bodies vary with their masses, it is obvious that, when we speak of the affinities of bodies, we ought to consider them as always acting in certain determinate proportions.

AUANTE. (From *auxina*, to dry.) A dry disease, proceeding from a fermentation in the stomach, described by Hippocrates de Morbis.

AUARSE. The same.

AUCIEN. (From *auxia*, to be proud.) The neck, which, in the posture of pride is made stiff and erect.

Auditory nerve. See *Nervus auditorius* and *Pertio mollis*.

Auditory passage. See *Meatus auditorius externus* and *internus*.

AUGUSTUM. An epithet given to several compound medicines.

AULISCOS. (From *αυλος*, a pipe.) A catheter, or clyster-pipe.

AULOS. The same.

AURA. (From *αω*, to breathe.) Any subtle vapour, or exhalation.

AURA EPILEPTICA. A sensation which is felt by epileptic patients, as if a blast of cold air ascended from the lower parts towards the heart and head.

AURA SEMINIS. The extremely subtle and vivifying portion of the semen virile, that ascends through the Fallopian tubes, to impregnate the ovum in the ovarium.

AURA VITALIS. So Helmont calls the vital heat.

AURANTII BACCÆ. Seville oranges. See *Aurantium*.

AURANTII CORTEX. See *Aurantium*.

AURANTIUM. (So called *ab aureo colore*, from its golden colour, or from *Aran-tium*, a town of Achaia.) *Aurantium hispaniense.* *Malus aurantia major.* *Malus aurantia.* *Aurantium vulgare.* *Malus aurantia vulgaris.* *Mala aurea.* *Chrysomela.* *Nerantia.* *Martianum pomum.* *Poma aurantia.* Seville orange. This plant is the *citrus aurantium* of Linnæus:—*petiolis alatis foliis acuminatis.* Class, *Polyadelphia.* Order, *Icosandria.* The China and Seville orange are both only varieties of the same species; the latter is met with in our pharmacopœias; and the *flowers, leaves, yellow rind, and juice*, are made use of for different medical purposes.

The *flowers, flores naphæ*, are highly odiferous, and are used as a perfume; they are bitter to the taste; they give their taste and smell both to water and to spirit, but most perfectly to rectified spirit of wine. The water which is distilled from these flowers, is called *aqua florum naphæ*. In distillation, they yield a small quantity of essential oil, which is called *oleum vel essentia neroli*: they are brought from Italy and France. Orange flowers were, at one time, said to be a useful remedy in convulsive and epileptic cases; but experience has not confirmed the virtues attributed to them.

The *leaves* have a bitterish taste, and yield, by distillation, an essential oil; indeed, by rubbing them between the fingers and the thumb, they manifest considerable fragrance. They have been applied for the same purposes as the flowers, but without success.

The *yellow rind* of the fruit, freed from the white fungous part, has a grateful aromatic flavour, and a warm, bitterish taste. Infused in boiling water, it gives out nearly

all its smell and taste; cold water extracts the bitter, but very little of the flavour. In distillation, a light, fragrant, essential oil rises, without the bitter. Its qualities are those of an aromatic and bitter. It has been employed to restore the tone of the stomach, and is a very common addition to combinations of bitters, used in dyspepsia. It has likewise been given in intermittents, in a dose of a drachm, twice or thrice a day. It is also much celebrated as a powerful remedy, in menorrhagia, and immoderate uterine evacuations. Its expressed oil is *essence of Bergamot*.

The *juice* of Seville oranges is a grateful acid, which, by allaying heat, quenching thirst, promoting various excretions, and diminishing the action of the vascular sanguiferous system, proves extremely useful in both ardent and putrid fevers; though the China orange juice, as impregnated with a larger proportion of sugar, becomes more agreeable, and may be taken in larger quantities. The Seville orange juice is particularly serviceable as an antiscorbutic, and alone will prevent or cure scurvy in the most apparently desperate circumstances. In dyspepsia, putrid bile in the stomach, both lemon and orange juice are highly useful.

AURANTIA CURASSAVENTIA. *Aurantium Curassavense.* Curassoa, or Curassao apples, or oranges. The fruit so called seem to be the immature oranges, that by some accident have been checked in their growth. They are a grateful aromatic bitter, of a flavour very different from that of the peel of the ripe fruit, and without any acid; what little tartness they have when fresh, is lost in drying. Infused in wine, or brandy, they afford a good bitter for the stomach. They are used to promote the discharge, in issues, whence their name of *issue peas*, and to give the flavour of hops to beer.

AURICULA. (Dim. of *auris*, the ear.) The external ear, upon which are several eminences and depressions, as the *helix*, *antihelix*, *tragus*, *antitragus*, *concha auriculae*, *schapha*, and *lobulus*.

AURICULA JUDÆ. *Fungus sambucinus Agaricus.* *Auricula forma.* Jew's ears. A membranaceous fungus, *Peziza auricula*; *concaua rugosa auriformis*, of Linnæus, which resembles the human ear. Its virtues are adstringent, and when employed, (by some its internal use is not thought safe,) it is made into a decoction, as a gargle for relaxed sore throats.

AURICULA MURIS. See *Pileolla*.

AURICULÆ CORDIS. The auricles of the heart. See *Heart*.

AURICULARIS. (*Auricularis*, sc. *digitus*: from *auris*, the ear.) The little finger; so called because people generally put it into the ear, when the hearing is obstructed.

AURIGA. (A wagoner. Lat.) A bandage for the sides; so called because it is made like the traces of a wagon horse. *Galen.*

AURIGO. (*Ab aures colore*: from its yellow colour.) The jaundice. See *Icterus*.

AURIFIGMENTUM. (From *aurum*, gold, and *pigmentum*, paint; so called from its colour and its use to painters. Yellow orpiment. See *Arsenic*.)

AURIS. (From *aura*, air, as being the medium of hearing.) The ear, or organ of hearing.) See *Ear*.

AURIS LEVATOR. See *Levator auris*.

AURISCALPIUM. (From *auris*, the ear, and *scalpo*, to scrape.) An instrument for cleansing the ear.

AURIUM SORDES. The wax of the ears.

AURIUM TINNITUS. A ringing noise in the ears.

AURUGO. The jaundice.

AURUM. Gold.

AURUM HORIZONTALE. Oil of cinnamon and sugar.

AURUM LEPROSUM. Antimony.

AURUM MUSIVUM. A preparation of tin, sulphur, sal-ammoniac, and quick-silver.

AURUM POTABILE. Gold dissolved and mixed with oil of rosemary, to be drank.

AURUS BRAZILIENSIS. *Calamus aromaticus*.

AUTHEMERON. (From *autos*, himself, and *emergo*, a day.) A medicine which gives relief, or is to be administered the same day.

AUTOLITHOTOMUS. One who cuts himself for the stone.

AUTOCRATHIA. The healing power of nature. *Hippocrates*.

AUTOPSIA. (From *autos*, himself, and *opsis* to see.) Ocular evidence.

AUTOPYROS. (From *autos*, itself, and *pyros*, wheat.) Bread made with the meal of wheat, from which the bran has not been removed. *Galen*.

AVENACU. A Molucca tree, of a caustic quality.

AVANGIS. *Avante*. Indigestion.

AVELLANA. (From *Abella*, or *avella*, a town in Campania, where they grew.) The hazel nut.

AVELLANA CATHARTICA. Barbadoes nuts. A purgative.

AVELLANA MEXICANA. Cocoa and chocolate nut.

AVELLANA PURGATRIX. Garden spurge.

AUXILIARI MUSCULI. The pyramidal muscles of the abdomen.

AVENA. (From *aveo*, to covet; because cattle are so fond of it.) The oat.

1. The name of a genus of plants in the Linnæan system. Class, *Triandria*. Order, *Dygyia*.

2. The pharmacopœial name of the oat.

Avena sativa of Linnæus. It is the seed which is commonly used, and called the

oat. There are two kinds of oats: the black and the white. They have similar virtues, but the black are chiefly sown for horses. They are less farinaceous, and less nourishing, than rice, or wheat; yet afford a sufficient nourishment, of easy digestion, to such as feed constantly on them. In Scotland, and some of the Northern counties of England, oats form the chief bread of the inhabitants. They are much used in Germany; but, in Norway, oat bread is a luxury, among the common people. Gruels, made with the flour, or meal, called oatmeal, digest easily, have a soft mucilaginous quality, by which they obtund acrimony, and are used for common drunk and food in fevers, inflammatory disorders, coughs, hoarseness, roughness, and exulceration of the fauces; and water-gruels answer all the purposes of Hippocrates' pisan. Externally, poultices, with oatmeal, vinegar, and a very little oil, are good for sprains and bruises. Stimulant poultices, with the grounds of strong beer, mixed up with oatmeal, are made for tumours, &c. of a gangrenous tendency.

AVENA SATIVA. The systematic name for the *avena* of the pharmacopœias. See *Avena*.

AVENÆ SEMINA. See *Avena*.

Avena, common. See *Caryophyllita*.

AVICENNIA TOMENTOSA. The systematic name for the plant which affords the *Anacardium orientale* of the pharmacopœias. See *Anacardium orientale*.

AVIGATO PEAR. This delicious fruit, the produce of the *Laurus persea* of Linnæus, when ripe, melts in the mouth like marrow, which it greatly resembles in flavour. It is supposed to be the most nutritious of all the tropical fruits, and grows in vast abundance in the West Indies and New Spain. The unripe fruit have but little taste; yet, being very salubrious, are often eaten with salt and pepper. The sailors, when they arrive at the Havannah, and those parts, purchase them in great quantities; and, chopping them into small pieces, with green capsicums and a little salt, regale themselves heartily with them. They are esteemed also for their antidiysenteric qualities, and are prepared in a variety of ways for the tables of the rich.

AXILLA. (*Axilla*, *atzil*, Heb. Scaliger deduces it from *ago*, to act; in this manner, *ago*, *axo*, *axa*, *axula*, *axilla*.) The cavity under the upper part of the arm, called the arm-pit.

AXILLARY ARTERIES. *Arterix axillares*. The axillary arteries are continuations of the subclavians, and give off, each of them, in the axilla, four mammary arteries, the subcapular, and the posterior and anterior circumflex arteries, which ramify about the joint.

AXILLARY NERVE. Articular nerve. A branch of the brachial plexus, and sometimes of the radial nerve. It runs outwards and backwards, around the neck of the humerus, and is lost in the muscles of the scapula.

AXILLARY VEINS. *Vene axillares.* The axillary veins receive the blood from the veins of the arm, and evacuate it into the subclavian vein.

AXIS. (From *ago*, to act.) See *Dentatus*.

AXUNGIA. (From *axis*, an axle-tree, and *unguo*, to anoint.) Hog's lard.

AXUNGIA CURATA. Purified hog's lard.

AXUNGIA DE MUMMIA. Marrow.

AZAC. (Arab.) Gum ammoniac.

AZAMAR. Native cinnabar. Vermilion.

AZED. A fine kind of camphire.

AZOT. (From *a*, priv. and *ζωω*, to live; because it is unfit for respiration.) See *Nitrogen*.

AZOT, GASEOUS OXYD OF. See *Nitrogen, gaseous oxyd of*.

AZOTH. An imaginary universal remedy.

AZUB. Alum.

AZURIUM. Quicksilver, sulphur, and sal-ammoniac.

AZYGES. (From *a*, priv. and *ζυγος*, a yoke.) The os phenoides was so called, because it has no fellow.

AZYGOS. (From *a*, priv. and *ζυγος*, a yoke; because it has no fellow.) Several single muscles, veins, bones, &c. are so called.

AZYGOS MORGAGNI. A muscle of the mouth.

AZYGOS PROCESSUS. A process of the os sphynoides.

AZYGOS UVULÆ. *Palato-staphilinus* of Douglas. *Staphilinus*, or *Epistaphilinus* of Winslow. A muscle of the uvula, which arises at one extremity of the suture which joins the palate bones, runs down the whole length of the velum and uvula, resembling an earth-worm, and adhering to the tendons of the circumflexi. It is inserted into the tip of the uvula. Its use is to raise the uvula upwards and forwards, and to shorten it.

AZYGOS VEIN. *Vena azygos. Vena sine pari.* This vein is situated in the right cavity of the thorax, upon the dorsal vertebrae. It receives the blood from the vertebral, intercostal, bronchial, pericardiac, and diaphragmatic veins, and evacuates it into the vena cava superior.

B

BABUZICARIUS. (From *babaζω*, to speak inarticulately.) The incubus, or nightmare; so called because, in this disorder, the person is apt to make an inarticulate and confused noise.

BACCA MONSPELIENSIS. See *Baccharis*.

BACCALIA. (*d baccharum copiosa*, because it abounds in berries.) The bay, or laurel-tree.

BACCE BERMUDENSES. See *Saponaria cucula*.

BACCE JUNIPERI. Juniper berries. See *Juniperus*.

BACCE LAURI. Laurel berries. See *Laurus*.

BACCE NORLANDICÆ. The fruit of the *Rubus arcticus* of Linnæus: *foliis alternatis, caule inermi unifloro*. They are recommended by Linnæus as possessing antiseptic, refrigerant, and antiscorbutic qualities.

BACCE PISCATORIÆ. See *Cocculus indicus*.

BACCHARIS. (From *bacchus*, wine; from its fragrance resembling that liquor.) *Bacchar. Bacca monspeliensis. Corryza tertia Dioscoridis, Eupatorium.* The

plant so called is the *Inula dysenterica* of Linnæus. Ploughman's spikenard. Great fleabane. It is sweet-scented, and the roots smell like cinnamon, and are said to powerfully emmenagogue, and the leaves moderately astringent.

BACHER'S PILLS. *Pilule tonice Bacheri.* A celebrated medicine in France, employed for the cure of dropsies. Their principal ingredient is the extract of melampodium, or black hellebore.

BACCHIA. (From *bacchus*, wine; because it generally proceeds from hard drinking and intemperance.) *Gutta roseacea*. A name given by Linnæus to a pimpled face.

BACULI. Is used, by some writers, for a particular kind of lozenges, shaped into little short rolls. Hildanus likewise uses it for an instrument in surgery.

BACOBÆ. The *Banana*.

BADIAGA. A kind of sponge usually sold in Russia, the powder of which is said to take away the livid marks of blows and bruises within a few hours. It is only described by Bauxbaum, and its nature is not properly understood.

BADIAN SEMEN. The seed of a tree which grows in China, and smells like aniseed. The Chinese (and Dutch, in imitation of them) sometimes use the badiane to give their tea an aromatic taste. See *Anisum stellatum*.

BADIZA AQUA. See *Bath water*.

BADRAHUM SEMEN. Indian aniseed.

BADUCCA. (Indian.) A species of caparis.

BANZHER. An antidote.

BÆOS. (*Bæos*.) In Hippocrates it means few; but in *P. Ægineta*, it is an epithet for a poultice.

BAGNIGGE WELLS. A saline mineral spring in London, resembling the Epsom mineral. In most constitutions, three half-pints is considered a full dose for purging.

BAGNIO. (From *bagno*, Ital.) A bathing or sweating-house.

BAHEI COYOLL. Ray takes it to be the *Arec*, or *Faufel*.

BAHEL SCHULLI. An Indian-tree; the *Genista spinosa Indica*; a decoction of the roots of which is diuretic. The leaves, boiled and sprinkled in vinegar, have the same effect, according to Ray.

BATAC. White lead.

BATAL. The mountain tree.

BALÆNA MACROCEPHALA. (*Balæna* from *βαλλω*, to cast, from its power in casting up water; and *μακροκεφαλος*; from *μακρος*, long, and *κεφαλη*, a head; from the length of its head.) The systematic name of a species of whale.

BALALINUM OLEUM. Oil of the ben-nut.

BALANOCASTANUM. (From *βαλανος*, a nut, and *καστανος*, a chestnut; so called from its tuberous root.) The *bunium bulbocastanum*, or earth-nut.

BALANOS. *Balanus*. (From *βαλλω*, to cast; because it sheds its fruit upon the ground.) 1. An acorn.

2. Hippocrates, in his *Treatise de Affectionibus*, expresses by it the oak.

3. Theophrastus uses it sometimes, to express any glandiferous tree.

4. From the similitude of form, this word is used to express suppositories and pessaries.

5. A name of the glans penis.

BALAUSTIUM. (From *βαλως*, various, and *αυω*, to dry; so called from the variety of its colours, and its becoming soon dry; or from *βλαστω*, to germinate.) *Balaustia*. A large rose-like flower, of a red colour, the produce of the plant from which we obtain the granatum. See *Granatum*.

BALBUTIES. (From *βαλβω*, to stammer; or from *babel*, Heb. to stammer.) A defect of speech; properly, that sort of stammering where the patient sometimes hesitates, and immediately after, speaks precipitately. It is the *Psyllismus Balbutiens* of Cullen.

BALISTA. (From *βαλλω*, to cast.) The astragulus, a bone of the foot, was formerly called *os balistæ*, because the ancients used to cast it from their slings.

Balmoney. See *Meum athamanticum*.

Balm. See *Melissa*.

Balm of Gilead. See *Moldavica*.

Balm of Mecca. See *Balsamum Gileadense*.

Balm, Turkey. See *Moldavica*.

BALLOTE. (From *βαλλω*, to send forth, and *ος*, *ωτος*, the ear; because it sends forth flowers like ears.) *Ballota*. Stinking horehound. A nettle-like plant. The *ballote nigra* of Linnæus.

BALNEUM. A bath, or bathing-house. See *Bath*.

BALNEUM ANIMALE. The wrapping any part of an animal, just killed, round the body, or a limb.

BALNEUM ARENÆ. A sand-bath for chymical purposes. See *Bath*.

BALNEUM CALIDUM. A hot-bath. See *Bath*.

BALNEUM FRIGIDUM. A cold bath. See *Bath*.

BALNEUM MARIE. *Balneum maris*. A warm-water bath. See *Bath*.

BALNEUM MEDICATUM. A bath impregnated with drugs.

BALNEUM SICCCUM. *Balneum cinereum*. A dry bath, either with ashes, sand, or iron filings.

BALNEUM SULPHUREUM. A sulphur bath.

BALNEUM TEPIDUM. A bath heated from 93 to 96 or 98 deg. of Fahrenheit's thermometer.

BALNEUM VAPORIS. A vapour bath.

BALLOON. (*Ballon* or *balon*, French) A large glass receiver in the form of a hollow globe. For certain chymical operations *ballons* are made with two necks, placed opposite to each other; one to receive the neck of a retort, and the other to enter the neck of a second *balloon*: this apparatus is called *enflated balloons*. Their use is to increase the whole space of the receiver, because any number of these may be adjusted to each other. The only one of these vessels which is generally used, is a small oblong *balloon* with two necks, which is to be luted to the retort, and to the receiver, or great *balloon*; it serves to remove this receiver from the body of the furnace, and to hinder it from being too much heated.

BALSAM. (*Balsamum*. From *baal samum*, Hebrew.) The term *balsam* was anciently applied to any strong-scented, natural vegetable resin of about the fluidity of treacle; inflammable, not miscible with water, without addition, and supposed to be possessed of many medical virtues. All the turpentine, the Peruvian

balsam, copaiba balsam, &c. are examples of natural balsams. Besides, many medicines compounded of various resins, or oils, and brought to this consistence, obtained the name of balsam. Latterly, however, the term has been restricted to those resins which contain the Benzoic acid. Of these only three are commonly known, the gum benzoin, balsam of Tolu, or Peru, and storax.

BALSAM APPLE, MALE. The fruit of the *momordica elaterium* of Linnæus. It is the fecula of the fruit of this plant, which affords the elaterium of the shops.

BALSAM, ARTIFICIAL. Compound medicines are thus termed which are made of a balsamic consistence and fragrance. They are generally composed of expressed or ethereal oils, resins, and other solid bodies, which give them the consistence of butter. The basis, or body of them, is expressed oil of nutmeg, and frequently wax, butter, &c. They are usually tinged with cinnamon and saffron.

Balsam Canary. See *Moldavica*.

Balsam of Canada. See *Balsamum Canadense*.

Balsam of Copaiva. See *Balsamum Copaivæ*.

BALSAM, NATURAL. A resin, which has not yet assumed the concrete form, but still continues in a fluid state, is so called, as common turpentine, balsamum copaiva, peruvianum, toluatum, &c.

Balsam, Peruvian. See *Balsamum Peruvianum*,

Balsam of sulphur. See *Balsamum sulphuris*.

Balsam of Tolu. See *Balsamum toluatum*.

Balsam Turkey. See *Moldavica*.

BALSAMATIO. (From *balsamum*, a balsam.) The embalming of dead bodies.

BALSAMEA. (From *balsamum*, balsam.) The balm of Gilead fir; so called from its odour. See *Balsamum canadense*.

BALSAMELEON. (From *balsamum*, balsam, and *leon*, oil.) Balm of Gilead, or true balsamum Judaicum.

BALSAM OLEUM. Balm of Gilead.

BALSAMICA. (*Balsamica*, sc. *medicamentis*; from *βάλανος*, balsam.) Balsamics. A term generally applied to substances of a smooth and oily consistence, which possess emollient, sweet, and generally aromatic qualities. Hoffman calls those medicines by this name, which are hot and acrid, and also the natural balsams, stimulating gums, &c. by which the vital heat is increased. Dr. Cullen speaks of them under the joint title of *balsamica et resinosa*, considering that turpentine is the basis of all balsams.

BALSAMIFERA BRAZILIENSIS. The balsam copaiba tree.

BALSAMIFERA INDICANA. The Peruvian balsam tree.

BALSAMITA MAS. (From *balsamum*.) *Balsamita major.* *Tanacetum hortense.* *Costus hortorum.* Costmary, or alecost. The plant which bears this name in the pharmacopœias, is the *Tanacetum balsamiti* of Linnæus:—*foliis ovatis, integris serratis*.

A fragrant-smelling herb, somewhat like that of mint; formerly esteemed as a corroborant, carminative, and emmenagogue.

BALSAMITA FEMINEA. See *Ageratum*.

BALSAMITA LUTEA. The *polygonum persicaria* of Linnæus. See *Persicaria*.

BALSAMITA MINOR. Sweet maudlin.

BALSAMITA MAJOR. See *Balsamita mas*.

BASAMUM. (From *baal samem*, Heb. the prince of oils.) A balsam. See *Balsam*.

BALSAMUM EGYPTIACUM. See *Balsamum Gileadense*.

BALSAMUM AMERICANUM. See *Balsamum Peruvianum*.

BALSAMUM ANODYNUM. A preparation made from tacamahacca, distilled with turpentine and soap liniment, and tincture of opium.

BALSAMUM ALPINUM. See *Balsamum Gileadense*.

BALSAMUM ANTIAONI. A remedy formerly applied to cancer.

BALSAMUM ARCEI. A preparation composed of gum-elemi and suet.

BALSAMUM ASIATICUM. See *Balsamum Gileadense*.

BALSAMUM BRAZILIENSE. See *Balsamum copaiba*.

BALSAMUM CANADENSE. Canada balsam. Balsam of Canadian fir. One of the purest turpentine, procured from the *Pinus balsamea* of Linnæus, and imported from Canada. For its properties, see *Turpentine*.

BALSAMUM CEPHALICUM. A distillation of oils, nutmeg, cloves, amber, &c.

BALSAMUM COMMENDATORIS. A composition of storax, benzoe, myrrh, aloes, &c.

BALSAMUM COPAIBÆ. (Many of the Americans call all odoriferous resins and sweet-scented gums, *copal*; and the word *iba*, or *iva*, is the name for a tree; hence *copaiba*.) *Balsamum Brazilianse.* *Balsamum copaibæ.* *Balsamum de copaibu.* *Balsamum copivi.* *Copaiba*.

Copaiba balsam is a yellow-resinous juice, of a moderately agreeable smell, and a bitterish biting taste, very permanent on the tongue. The tree which affords it, is the *Copaifera officinalis* of Linnæus. Class, *Decandria*. Order, *Mono-gynia*. It is obtained by making deep incisions near its trunk, when the balsam immediately issues, and, at the proper season, flows in such abundance, that sometimes, in three hours, twelve pounds have been procured. The older tree

afford the best balsam, and yield it two or three times in the same year. The balsam supplied by the young and vigorous trees, which abound with the most juice, is crude and watery, and is, therefore, accounted less valuable. While flowing from the tree, this balsam is a colourless fluid; in time, however, it acquires a yellowish tinge, and the consistence of oil; but, though by age it has been found thick, like honey, yet it never becomes solid, like other resinous fluids. By distillation in water, the oil is separated from the resin; and, in the former, the taste and smell of the balsam are concentrated. If the operation is carefully performed, about one-half of the balsam rises into the receiver, in the form of oil. The balsam unites with fixed and volatile oils, and with spirit of wine. It is given in all diseases of the urinary organs, when no inflammation is present. In gleets, and in gonorrhoea, it was once a favourite remedy, but is now disused. In diseases of the kidneys it is still employed, though less frequently than usual; and in hæmorrhoids it is occasionally trusted. The dose is from 20 to 30 drops, twice or three times a day, mixed with water, by means of an egg, or other mucilage. The balsam of copalva is occasionally adulterated with turpentine, but its virtues are not greatly injured by the fraud.

BALSAMUM EMBRYONUM. A preparation of aniseed.

BALSAMUM GENUINUM ANTIQUORUM. See *Balsamum Gileadense*.

BALSAMUM GILEADENSE. *Balsamum genuinum antiquorum. Balsameleon. Aegyptiacum balsamum. Balsamum Asiaticum. Balsamum Judicum. Balsamum Syriacum. Balsamum e Mecca Balsamum alpinum. Oleum balsami. Opobalsamum. Xyrobalsamum.* Balsam, or balm of Gilead. A resinous juice, obtained by making incisions into the bark of the *Amyris Gileadensis* of Linnaeus:—*foliis ternatis integerrimis, pedunculis unifloris lateralibus.* Class, *Octandria.* Order, *Monogynia.* The tree grows spontaneously, particularly near to Mecca, on the Asiatic side of the Red Sea. The juice of the fruit is termed *carpobalsamum* in the pharmacopœias, and that of the wood and branches *xylobalsamum*. The best sort is a spontaneous exudation from the tree, and is held in so high estimation by the Turks, that it is rarely, if ever, to be met with genuine among us. The medicinal virtues of the genuine balsam of Gilead, have been highly rated, undoubtedly with much exaggeration. The common balsam of Mecca is scarcely used; but its qualities seem to be very similar to those of the balsam of Tolu, with perhaps more acrimony. The dose is from 15 to 50 drops.

BALSAMUM GUAIAECINUM. Balsam of Peru and spirits of wine.

BALSAMUM GUIDONIS. The same as balsamum anodynum.

BALSAMUM HUNGARICUM. A balsam prepared from a coniferous tree on the Carpathian mountains.

BALSAMUM JUBAICUM. See *Balsamum Gileadense*.

BALSAMUM LOCATELLI. (*Locatelli*; so called from its inventor Lucatellus.) *Balsamum Lucatelli.* A preparation made of oil, turpentine, wax, and red saunders; now disused; formerly exhibited in coughs of long standing.

BALSAMUM MAS. The herb costmary. See *Balsamitana*.

BALSAMUM E MECCA. See *Balsamum Gileadense*.

BALSAMUM MEXICANUM. See *Balsamum Peruvianum*.

BALSAMUM NOVUM. A new balsam from a red fruit in the West Indies.

BALSAMUM ODORIFERUM. A preparation of oil, wax, and any essential oil.

BALSAMUM PERUVIANUM. (From its country, Peru.) *Putzochilli Indian, Mexican, and American balsam.* *Carbareiba* is the name of the tree from which, according to Piso and Ray, it is taken. It is the *Myroxylon peruiferum* of Linnaeus, which grows in the warmest provinces of South America, and is remarkable for its elegant appearance. Every part of the tree abounds with a resinous juice; even the leaves being full of transparent resinous points, like those of the orange-tree.

Balsam of Peru is of three kinds; or rather, it is one and the same balsam, having three several names: 1. The balsam of incision; 2. The dry balsam; 3. The balsam of lotion. The virtues of this balsam, as a cordial, pectoral, and restorative stimulant, and tonic, are by some thought to be very great. It is given with advantage from 5 to 10 or 15 drops for a dose, in dyspepsia, atonic gout, in consumptions, asthmas nephritic complaints, obstructions of the viscera, and suppressions of the menses. It is best taken dropped upon sugar. The yolk of an egg, or mucilage of gum-arabic, will, indeed, dissolve it; it may, by that way, be made into an emulsion; and it is less acrid in that form than when taken singly. It is often made an ingredient in boluses and electuaries, and enters into two of the official compositions: the tinctura balsami Peruviani compositi, and the trochisci glycyrrhyze. Externally, it is recommended as a useful application to relaxed ulcers, not disposed to heal.

BALSAMUM PERISICUM. A balsamum composed of storax, benzoe, myrrh, and aloes.

BALSAMUM RACKASIRA. This balsam, which is inodorous when cold, but of a smell approaching to that of Tolu balsam when heated, is brought from India in

gourd-shells. It is slightly bitter to the taste, and adheres to the teeth, on chewing. It is supposed to be one of the facitious balsams, and is scarcely ever prescribed in this country.

BALSAMUM SANECH. A facitious balsam, composed of tartar, dulcified by spirits of wine.

BALSAMUM SAPONACEUM. A name given to the preparation called opodeldoc.

BALSAMUM SATURNI. The remedy so named is prepared by dissolving the acetate of lead in oil of turpentine, and digesting the mixture till it acquires a red colour. This is found to be a good remedy for cleansing foul ulcers; but it is not acknowledged in our dispensatories.

BALSAMUM STYRACIS BENZOINI. Gum-benjoin is so called. See *Benzoinum*.

BALSAMUM SUCCINI. Oil of amber.

BALSAMUM SULPHURIS. A solution of sulphur in oil.

BALSAMUM SULPHURIS ANISATUM. Balsam of sulphur, and oil of aniseed.

BALSAMUM SULPHURIS BARBADENSE, Sulphur boiled with Barbadoes tar.

BALSAMUM SULPHURIS CRASSUM. Thick balsam of sulphur.

BALSAMUM SULPHURIS TEREBINTHINATUM. This is made by digesting the sulphur with oil of turpentine, and, in the latter, adding the oil of aniseed. They are now confined to veterinary medicine.

BALSAMUM SULPHURIS SIMPLEX. Sulphur boiled with oil.

BALSAMUM SYRIACUM. The balm of Gilead. See *Balsamum Gileadense*.

BALSAMUM TOLUTANUM. Balsam of Tolu. The tree *Toluifera balsamum* of Linnaeus, from which this balsam is procured, grows in South America, in the province of Tolu, behind Carthagena, whence we are supplied with the balsam, which is brought to us in little gourd-shells. The balsam is obtained by making incisions into the bark of the tree, and is collected into spoons, which are made of black wax, from which it is poured into proper vessels. It thickens, and in time becomes concrete: it has a fragrant odour, and a warm sweetish taste. It dissolves entirely in alcohol, and communicates its odour and taste to water, by boiling. It contains acid of benzoic. This is the mildest of all the balsams. It has been used as an expectorant; but its powers are very inconsiderable, and it is at present employed principally on account of its flavour somewhat resembling that of lemons. It is directed, by the pharmacopœias, in the *syrrupus Tolutanus*, *tinctura Tolutana*, and *syrrupus balsamicus*.

BALSAMUM TRAUMATICUM. Vulnerary balsam. A form of medicine prescribed in the London Dispensatory, intended to supply the place of the tincture commonly called *Fraser's balsam*, so famous for curing

old ulcers. The London College have named it *Tinctura Benzoini composita*.

BALSAMUM UNIVERSALE. A name given to the unguentum saturninum of old pharmacopœias.

BALSAMUM VERUM. See *Balsamum Gileadense*.

BALSAMUM VIRIDE. Linseed-oil, turpentine and verdigrise, mixed together.

BALSAMUM VITÆ HOFFMANNI. *Beaume de vie*. An artificial balsam, so named from its inventor, and composed of a great variety of the warmest and most grateful essential oils, such as nutmegs, cloves, lavender, &c. with balsam of Peru, dissolved in highly rectified spirit of wine; but it is now greatly abridged in the number of ingredients, and but little used.

BALZOINUM. The gum-benjamin.

BAMBALIO. (From *Bambano*, to speak inarticulately.) A person who stammers, or lisps.

BAMBOO. (Indian.) The young shoots of the *arundo bambos* of Linnaeus, which are prepared by the natives of both Indies with vinegar, garlic, pepper, &c. into a very excellent pickle, which promotes the appetite, and assists digestion.

BAMIA MOSCHATA. See *Abelmoschus*.

BAMIER. The name of a plant common in Egypt, the husk of which they dress with meat, and, from its agreeable flavour, make great use of it in their ragouts.

BAN ARBOR. The coffee-tree.

BANANA. (Indian.) *Bananeira*. *Ficoides*. *Ficus Indica*. *Musa fructu cucumerino breviori*. *Senoria*. *Paceira*. The Banana, or Plantain-tree. The most remarkable species of this genus of plants are,

1. The *paradisæica*, or plantain.
2. The *musa sapientum*, or banana-tree.

Both are among the most important productions of the earth. The first sort is cultivated in all the islands of the West-Indies, where the fruit serves the Indians for bread; and some of the white people also prefer it to most other things, especially to the yams and cassada bread. This tree is cultivated, on a very extensive scale in Jamaica; without the fruit of which, Dr. Wright says, the island would scarcely be habitable, as no species of provision would supply their place. Even flour, or bread itself, would be less agreeable, and less able to support the laborious negro, so as to enable him to do his business, or to keep in health. Plantains also fatten horses, cattle, swine, dogs, fowls, and other domestic animals. The leaves, being smooth and soft, are employed as dressings after blisters. The water from the soft trunk is astringent, and employed by some to check diarrhœas. Every other part of the tree is useful in different parts of rural economy. The leaves are used as napkins and tablecloths, and are food for hogs. The second sort, *musa sapientum*, or banana-tree, dif-

fers from the paradisica, in having its stalks marked with dark purple stripes and spots. The fruit is shorter, straighter, and rounder; the pulp is softer, and of a more luscious taste. It is never eaten green; but, when ripe, it is very agreeable, either eaten raw or fried in slices, as fritters, and is relished by all ranks of people in the West Indies. Both the above plants were carried to the West Indies from the Canary Islands; whither, it is believed, they had been brought from Guinea, where they grow naturally.

BANANERA. See *Banana*.

BANCIA. The *Elaphoboscum*, or wild parsnip.

BANDAGE. *Deligatio. Fascia.* An apparatus consisting of one or several pieces of linen, or flannel, and intended for covering or surrounding parts of the body for surgical purposes. Bandages are either simple or compound. The chief of the simple are the circular, the spiral, the uniting, the retaining, the expellent, and the creeping. The compound bandages used in surgery, are the T bandage, the suspensory one, the capistrum, the eighteen-tail bandage, and others to be met with in surgical treatises.

BANDURA. A plant which grows in Ceylon, whose root is said to be astringent.

BANGUS. *Bange.* A species of opiate in great use throughout the East, for its intoxicating qualities. It is the leaf of a kind of wild hemp, growing in the countries of the Levant, and made into powder, pills, or conserves.

BANICA. The wild parsnip.

BANILIA. See *Vanilla*.

BANILAS. See *Vanilla*.

BAOBAB. *Bahobab.* A species of the genus of plants called by Linnæus *Adansonia*. It grows mostly on the west coast of Africa, from the Niger to the kingdom of Benin. The bark is called *lalo*; the negroes dry it in the shade, then powder and keep it in little cotton bags, and put two or three pinches into their food. It is mucilaginous, and powerfully promotes perspiration. The mucilage obtained from this bark is a powerful remedy against the epidemic fevers of the country that produces these trees; so is a decoction of the dried leaves. The fresh fruit is as useful as the leaves, for the same purposes.

BAPTICA COCCUS. Kermes berries.

BAPTISTERIUM. (From *baptizo*, to immerge.) A bath, or repository of water, to wash the body.

BAPTISTRUM. (From *baptizo*, to dye.) A species of wild mustard; so called from its reddish colour.

BARAC. (From *borak*, Arabian, splendid.) *Baruch panis.* Nitre. According to Rulandus, nitrum salis.

BARAS. (Arabian.) In M. A. Severinus, it is synonymous with *Alphus*, or *Leuce*.

BARATHERUM. (Arabian.) Any cavity or hollow place.

BARBA. (From *barbarus*, because wild nations are usually unsheaven.) 1. The beard of man.

2. Some vegetables have the specific name of *barba*, whose ramifications are bushy, like a beard, as *barba jovis*, &c.

BARBA ARONIS. The arum.

BARBA CUPRIÆ. The ulmaria.

BARBA HIRCI. The tragopogon.

BARBA JOVIS. Jupiter's beard, or the silver bush. Also a name of the *sempervivum majus*, and of a species of *anthyllis*.

BARBADOES CHERRY. The fruit of the *Malphigia glabra* of Linnæus, resembling the inferior of our cherries.

Barbadoes nut. See *Ricinus major*.

BARBADOES TAR. (So named from the island from which it is chiefly procured.) The use of this article in medicine is limited to its external application, at times, in paralytic cases.

BARBARA. (From St. Barbary, who is said to have found its virtues.) The leaves of this plant, *Eriogonum barbarea: foliis lyrtis, extimo subrotundo* of Linnæus, may be ranked among the antiscorbutics.—They are seldom used in practice.

BARBARIA. *Barbaricum.* An obsolete term formerly applied to rhubarb.

BARBAROSSÆ PILULA. Barbarossa's pill. An ancient composition of quicksilver, rhubarb, diagridium, musk, amber, &c. It was the first internal mercurial medicine which obtained any real credit.

BARBARUM. The name of a plaster in Scribonius Largus.

BARBATINA. A Persian vermifuge seed.

BARBEL. *Barbo.* And oblong fish resembling the pike, the eating of whose roe often brings on the cholera morbus.

Barberry. See *Berberis*.

BARBOTA. The barbut. A small river-fish. It is remarkable for the size of its liver, which is esteemed the most delicate part of it.

BARDANA. (From *bardus*, foolish; because silly people are apt to throw them on the garments of passengers, having the property of sticking to whatever they touch) *Arctium. Betonica. Britannica. Naphis.* Burd. ck. The plant so called in the pharmacopœias, is *Arctium lappa: —foliis cordatis, inermibus, petiolatis*, of Linnæus. It grows wild in uncultivated grounds. The seeds have a bitterish sub-acrid taste; they are recommended as very efficacious diuretics, given either in the form of emulsion, or in powder, to the quantity of a drachm. The roots taste sweetish, with a slight austeritv and bitterness: they are esteemed aperient, diuretic, and sudorific; and are said to act without irritation, so as to be safely ventured upon in acute disorders. Decoctions

of them have of late been used, in rheumatic, gouty, venereal, and other disorders; and are preferred by some to those of sarsaparilla. Two ounces of the roots are to be boiled in three pints of water, to a quart; to this, two drachms of vitriolated Kali have been usually added. Of this decoction, a pint should be taken every day, in scorbutic and rheumatic cases, and when intended as a diuretic, in a shorter period.

BAREGE WATER. The small village of Barege, celebrated for its thermal waters, is situated on the French side of the Pyrenees, about half way between the Mediterranean and the Bay of Biscay. The hot springs are four in number. They have all the same component parts, but differ somewhat in their temperature, and in the quantity of sulphur, the hottest being most strongly penetrated with this active ingredient. The coolest of these waters raises Fahrenheit's thermometer to 73 deg.; the hottest to 120 deg. Barege waters are remarkable for a very smooth soapy feel; they render the skin very supple and pliable, and dissolve perfectly well soap and animal lymph; and are resorted to as a bath in resolving tumours of various kinds, rigidities, and contractions of the tendons, stiffness of the joints, left by rheumatic and gouty complaints, and are highly serviceable in cutaneous eruptions. Internally taken, this water gives considerable relief in disorders of the stomach, especially attended with acidity and heart-burn, in obstinate cholics, jaundice, and in gravel, and other affections of the urinary organs.

BARILLA. See *Barrilla*.

BARILLA. (*Barigla*, the place where it was formerly produced) *Bariglia*. *Bariller*. *Anatron*. *Natron*. *Anaton*. *Nitrum antiquorum*. *Aphronitrum*. *Baurach*. *Sal alkalinus fixus fossilis*. *Carbonas soda impurus*. *Sub-carbonas soda impurus*. Soda Barilla is the terra given, in commerce, to the impure mineral alkali, or imperfect carbonate of soda, imported from Spain and the Levant. It is made by burning to ashes different plants that grow on the seashore, chiefly of the genus *salsola* of Linnaeus, and is brought to us in hard porous masses, of a speckled brown colour. Kelp, a still more impure alkali, made in this country by burning various sea-weeds, is sometimes called British barilla. The marine plants, collected for the purpose of procuring barilla in this country, are the *salsola kali*, *salicornia Europaea*, *zostera maritima*, *triglochen maritimum*, *chenopodium maritimum*, *atriplex portulacoides* and *littoralis*, *plantago maritima*, *tamarix gallica*, *eryngium maritimum*, *sedum telephium*, *dipsacus fullonum*, &c. &c.

BARK. A term very frequently employed to signify, by way of eminence, Peruvian bark. See *Cinchona*.

Bark, Carribean. See *Cinchona Jamaicensis*.

Bark, Jamaica. See *Cinchona Jamaicensis*.

Bark, Peruvian. See *Cinchona*.

Bark, red. See *Cinchona rubra*.

Bark yellow. See *Cinchona flava*.

Barley. See *Hordeum*.

Barley, caustic. See *Cordilla*.

Barley, pearl. See *Hordeum perlatum*.

BARM. A name given to yeast.

BARNET WATER. It is of a purging kind, of a similar quantity to that of Epsom, and about half its strength.

BAROMETER. (From *bagos*, weight, and *metron*, measure.) An instrument to determine the weight of the air; it is commonly called a weather-glass.

BARONES. Small worms; called also *Nepones*.

BAROPTIS. A black stone, said to be an antidote to venomous bites.

BAROS. (*Bagos*.) Gravity.

1. Hippocrates uses this word to express by it, an uneasy weight in any part.

2. It is also the Indian name for a species of camphire, which is distilled from the roots of the true cinnamon-tree.

BARRENNESS. The same as sterility.

BARTHOLINIAN GLANDULE. See *Sublingual glands*.

BARTCOIA. (From *bagus*, heavy, and *akousis*, to hear.) Deafness, or difficulty of hearing.

BARYOCOCCALON. (From *bagus*, heavy, and *kokkalis*, a nut; because it gives a deep sound.) A name for the stramonium.

BARYPHONIA. (From *bagus*, dull, and *phona*, the voice.) A difficulty of speaking.

BARYTES. (From *bagus*, heavy; so called because it is very ponderous.) *Cauk*. *Calk*. *Terra ponderosa*. *Baryt*. *Ponderous earth*. *Heavy earth*.

Barytes does not exist pure in nature. It is always found in combination with sulphuric or carbonic acid. United with the sulphuric acid, it forms the mineral called *sulphate of barytes*, or *baroselenite*. It is found in Staffordshire, Derbyshire, &c. When united to carbonic acid, it is called *erated barytes*, or *carbonate of barytes*, found at Anglezark, near Chorley, in Lancashire. Both combinations are met with regularly crystallized and amorphous.

Pure barytes has a much stronger affinity than any other body for sulphuric acid; it turns blue tincture of cabbage green. It is entirely infusible by heat alone, but melts when mixed with various earths. Its specific gravity is 4.000. It changes quickly in the air, swells, becomes soft, and falls into a white powder, with the acquisition of about one-fifth of its weight. This slaking is much more active and speedy than that of lime. It combines with phosphorus, which compound decomposes water rapidly. It unites to sulphur by the

dry and humid way. It has a powerful attraction for water, which it absorbs with a hissing noise and consolidates it strongly. It is soluble in twenty times its weight of cold, and twice its weight of boiling water. Its crystals are long four-sided prisms of a satin-like appearance. It is a deadly poison to animals.

Method of obtaining pure Barytes.—1. Take native carbonate of barytes; reduce it to a fine powder and dissolve it in a sufficient quantity of diluted nitric acid; evaporate this solution till a pellicle appears, and then suffer it to crystallize in a shallow basin. The salt obtained is nitrate of barytes; expose this nitrate of barytes to the action of heat in a china cup, or silver crucible, and keep it in a dull red heat for at least one hour, then suffer the vessel to cool, and transfer the greenish solid contents, which are pure barytes, into a well stopped bottle. When dissolved in a small quantity of distilled water, and evaporated, it may be obtained in a beautiful crystalline form.

In this process the nitric acid, added to the native carbonate of barytes, unites to the barytes, and expels the carbonic acid, and forms nitrate of barytes; on exposing this nitrate to heat, it parts with its nitric acid, which becomes decomposed into its constituents, leaving the barytes behind.

2. Pure barytes may likewise be obtained from its sulphate. For this purpose, boil powdered sulphate of barytes in a solution of twice or three times its weight of carbonate of potash, in a Florence flask, for about two hours; filter the solution and expose what remains on the filter to the action of a violent heat.

In this case the sulphuric acid of the barytes unites to the potash, and the carbonic acid of the latter, joins to the barytes; hence sulphate of potash and carbonate of barytes are obtained. The former is in solution and passes through the filter; the latter is insoluble, and remains behind. From this artificial carbonate of barytes, the carbonic acid is driven off by heat.

BASAAL. (Indian.) The name of an Indian tree. A decoction of its leaves, with ginger, in water, is used as a gargle in disorders of the fauces. The kernels of the fruit kill worms. *Ray's Hist.*

BASALTES. (In the Æthiopic tongue, this word means iron, which is the colour of the stone.) A heavy and hard kind of stone, chiefly black, or green. It frequently contains iron, has a flinty hardness, is insoluble by acids, and is fusible by fire. The most remarkable property of this substance is its figure, being never found in strata, like other marbles, but always standing up in the form of regular angular columns, composed of a number of joints, one placed upon and nicely fitted to another, as if formed by the hands of a skilful architect. Some regard this fusible substance as a volcanic production, others

have supposed that it was formed of water. The Giant's Causeway, in the county of Antrim, in Ireland, and the rock of Pèreniere, near St. Santdoux, in Auvergne, are formed of these stones. The distinctive characters of basaltes are, a regular form, hardness sufficient to give fire with steel; and a cinereous, gray colour, inclining to black.

BASANITES. (From *basanizo* to find out.) A stone said by Pliny, to contain a bloody juice, and useful in diseases of the liver; also a stone upon which, by some, the purity of gold was formerly said to be tried, and of which medical mortars were made.

Base, acidifiable. See *Acid*.

Base, acidifying. See *Acid*.

BASIATTO. (From *basio*, to kiss.) Venereal connection between the sexes.

BASIATOR. See *Constrictor labiorum*.

BASIL. See *Basilicum*.

BASILARE OS. (*Basilaris*; from *basileus*, a king.) Several bones were so termed by the ancients; as the sphenoid and occipital bones.

BASILARIS ARTERIA. Basillary artery. An artery of the brain. So called because it lies upon the basillary process of the occipital bone. It is formed by the junction of the two vertebral arteries within the skull, and runs forwards to the sella turcica along the *pons varolii*, which it supplies, as well as the adjacent parts, with blood.

BASILARIS PROCESSUS. Basillary process. See *Occipital bone*.

BASILIARIS APOPHYSIS. The great apophysis of the os occipitis.

BASILICA MEDIANA. See *Basilica vena*.

BASILICA RUX. The walnut.

BASILICA VENA. The large vein that runs in the internal part of the arm, and evacuates its blood into the axillary vein. The branch which crosses, at the head of the arm, to join this vein, is called the *basilic median*. They may either of them be opened in the operation of blood-letting.

Basilicon ointment. See *Basilicum unguentum*.

BASILICUM. (From *basileus*, royal; so called from its great virtues.) *Ocimum*. Basil. The plant which bears this name in the pharmacopœias, is the *Ocimum basilicum* of Linnæus:—*foliis ovatis glabris; calycibus ciliatis*. It is supposed to possess nervine qualities, but is seldom employed but as a condiment to season high dishes, to which it imparts a grateful odour and taste.

BASILICUM UNGUENTUM. *Unguentum basilicum flavum*. An ointment popularly so called from its having the *ocimum basilicum* in its composition. It came afterwards to be composed of wax, resin, &c. and is now called *ceratum resinae flavæ*.

BASILICUS PULVIS. The royal powder.

A preparation formerly composed of calomel, rhubarb, and jalap. Many compositions, were, by the ancients, so called, from their supposed pre-eminence.

BASILION. An itchy ointment was formerly so called by Galen.

BASILIS. A name formerly given to collyriums of supposed virtues, by Galen.

BASILISCUS. (From *basilus*, a king.) The basilisk, or cockatrice, a poisonous serpent; so called from a white spot upon its head, which resembles a crown. Also the philosopher's stone, and corrosive sublimate.

BASIO-CERATO-CHONDRO-GLOSSUS. See *Hyaloglossus*.

BASIO-GLOSSUM. See *Hyaloglossus*.

BASIO-PHARYNGEUS. See *Constrictor pharyngis medius*.

BASIS. (From *basis*, to go: the support of any thing, upon which it stands or goes.)

1. This word is very frequently applied anatomically to the body of any part, or to that part from which the other parts appear, as it were, to proceed, or by which they are supported.

2. In pharmacy it signifies the principal ingredient.

BASIS CEREVIS. A term applied formerly to the palatum.

BASIS CORDIS. The broad part of the heart is so called, to distinguish it from the apex, or point.

BASIS COLICA. The name of a medicine in Scribonius Largus, compounded of aromatics and honey.

Bastard pleurisy. See *Peripneumonia notha*.

BATATAS. (So the natives of Peru call the potato, which is a native of that country, from our word potato.) A species of night-shade, *solanum tuberosum*, Linn. Potatoes were first brought into Europe by Sir Francis Drake, 1486, and planted in London. They are said to be natives of Peru.

BATH. *Balneum.* Baths are of several kinds.

1. A convenient receptacle of water, for persons to wash or plunge in, either for health or pleasure, is called a bath. These are distinguished into hot and cold; and are either natural or artificial. The natural hot baths are formed of the water of hot springs, of which there are many in different parts of the world; especially in those countries where there are, or have evidently been, volcanoes. The artificial hot baths consist either of water, or of some other fluid, made hot by art. The cold bath consists of water, either fresh or salt, in its natural degree of heat; or it may be made colder by art, as by a mixture of nitre, sal-ammoniac, &c. The chief hot baths in our country are those of Bath and Bristol, and those of Buxton and Matlock; which latter, however, are rather warm, or tepid, than hot. The use of these baths is found to be beneficial in diseases of the

head, as palsy, &c.; in cuticular diseases, as leprosy, &c.; obstructions and constipations of the bowels, the scurvy, and stone; and in many diseases of women and children. The cold bath, though popularly esteemed one of the most innocent remedies yet discovered, is not, however, to be adopted indiscriminately. On the contrary, it is liable to do considerable mischief in all cases of diseased viscera, and is not, in any case, proper to be used during the existence of costiveness. As a preventive remedy for the young, and as a general bracer for persons of a relaxed fibre, especially of the female sex, it often proves highly advantageous; and in general, the popular idea is a correct one, that the *glew* which succeeds the use of cold or temperate baths, is a test of their utility; while, on the other hand, their producing *chilliness*, head-ache, &c. is a proof of their being pernicious.

The Cold Bath.

The diseases and morbid symptoms, for which the cold bath, under one form or another, may be applied with advantage, are very numerous; and some of them deserve particular attention. One of the most important of its uses is in *ardent fever*; and, under proper management, it forms a highly valuable remedy in this dangerous disorder. It is highly important, however, to attend to the precautions which the use of this vigorous remedial process requires. "Affusion with cold water," Dr. Currie observes, "may be used whenever the heat of the body is steadily above the natural standard, when there is no sense of chilliness, and especially when there is no general nor profuse perspiration. If used during the cold stage of a fever, even though the heat be higher than natural, it brings on interruption of respiration, a fluttering, weak, and extremely quick pulse, and certainly might be carried so far as to extinguish animation entirely." The most salutary consequence which follows the proper use of this powerful remedy, is the production of profuse and general perspiration. It is this circumstance that appears to give so much advantage to a general effusion of cold water in fevers, in preference to any partial application. The cold bath is better known, especially in this country, as a general tonic remedy in various chronic diseases. The general circumstances of disorder for which cold bathing appears to be of service, according to Dr. Saunders, are a languor and weakness of circulation, accompanied with profuse sweating and fatigue, on very moderate exertion; tremors in the limbs, and many of those symptoms usually called nervous; where the moving powers are weak, and the mind listless and indolent; but, at the same time, where no permanent morbid obstruction, or visceral disease, is present. Such a state of body is often the conse-

quence of a long and debilitating sickness, or of a sedentary life, without using the exercise requisite to keep up the activity of the bodily powers. In all these cases, the great object to be fulfilled, is to produce a considerable reaction, from the shock of cold water, at the expense of as little heat as possible; and when cold-bathing does harm, it is precisely where the powers of the body are too languid to bring on reaction, and the chilling effects remain unopposed. When the patient feels the shock of immersion very severely, and, from experience of a pain, has acquired an insuperable dread of this application; when he has felt little or no friendly glow to succeed the first shock, but on coming out of the bath remains cold, shivering, sick at the stomach, oppressed with head-ache, languid, drowsy and listless, and verse to food and exercise during the whole of the day, we may be sure that the bath has been too cold, the shock too severe, and no reaction produced at all adequate to the impression on the surface of the body.

There is a kind of slow irregular fever, or rather febricula, in which Dr. Saunders has often found the cold bath of singular service. This disorder principally affects persons naturally of a sound constitution, but who lead a sedentary life, and at the same time are employed in some occupation which strongly engages their attention, requires much exertion of thought, and excites a degree of anxiety. Such persons have constantly a pulse rather quicker than natural, hot hands, restless nights, and an impaired appetite; but without any considerable derangement in the digestive organs. This disorder will continue for a long time, in an irregular way, never entirely preventing their ordinary occupation, but rendering it more than usually anxious and fatiguing, and often preparing the way for confirmed hypochondriasis. Persons in this situation, are remarkably relieved by the cold-bath, and for the most part, bear it well; and its use should also, if possible, be aided by that relaxation from business, and that diversion of the mind from its ordinary train of thinking, which are obtained by attending a watering-place. The Doctor also found cold bathing hurtful in chlorosis, and observes, that it is seldom admissible in those cases of disease in the stomach which are brought on by high living, and constitute what may be termed the true dyspepsia.

The topical application of cold water, or of a cold saturnine lotion, in cases of local inflammation, has become an established practice; the efficacy of which is daily experienced. Burns of every description will bear a most liberal use of cold water, or even of ice; and this may be applied to a very extensive inflamed surface, without even producing the ordinary effects of general chilling, which would be

brought on from the same application to a sound and healthy skin. Another very distressing symptom, remarkably relieved by cold water, topically applied, is that intolerable itching of the vagina, which women sometimes experience, entirely unconnected with any general cause, and which appears to be a kind of herpes confined to that part. Cold water has also been used topically in the various cases of strains, bruises, and similar injuries, in tendinous and ligamentous parts, with success; also in rigidity of muscles, that have been long kept at rest, in order to favour the union of bone, where there appears to have been no organic injury, but only a deficiency of nervous energy, and in mobility of parts, or at most, only slight adhesions, which would give way to a regular exercise of the weakened limb. Another very striking instance of the powerful effects of topical cold, in stimulating a part to action, is shown in the use of cold, or even iced water, to the vagina of parturient women, during the dangerous hemorrhages that take place from the uterus, on the partial separation of the placenta.

The Shower Bath.

A species of cold bath. A modern invention, in which the water falls, through numerous apertures, on the body. A proper apparatus for this purpose is to be obtained at the shops. The use of the shower bath applies, in every case, to the cold bath, and is of en attended with particular advantages. 1. From the sudden contact of the water, which, in the common cold bath, is only momentary, but which, in the shower bath may be prolonged, repeated, and modified, at pleasure; and, secondly, from the head and breast, which are exposed to some inconvenience and danger in the common bath, being here effectually secured, by receiving the first shock of the water.

The Tepid Bath.

The range of temperature, from the lowest degree of the warm bath to the highest of the cold bath, forms what may be termed the tepid. In general, the heat of water which we should term tepid, is about 90 deg. In a medicinal point of view, it produces the greatest effect in ardent fever, where the temperature is little above that of health, but the powers of the body weak, not able to bear the vigorous application of cold immersion. In cutaneous diseases, a tepid bath is often quite sufficient to produce a salutary relaxation, and perspirability of the skin.

The Warm Bath.

From 93 to 96 deg. of Fahrenheit, the warm bath has a peculiar tendency to bring on a state of repose, to alleviate any local irritation, and thereby induce sleep. It is, upon the whole, a safer remedy than the cold bath, and more peculiarly appli-

cable to very weak and irritable constitutions, whom the shock produced by cold immersion would overpower, and who have sufficient vigour of circulation for an adequate reaction. In cases of topical inflammation, connected with a phlogistic state of body, preceded by rigor and general fever, and where the local formation of matter is the solution of the general inflammatory symptoms, experience directs us to the use of the warm relaxing applications, rather than those which, by exciting a general reaction, would increase the local complaint. This object is particularly to be consulted when the part affected is one that is essential to life. Hence it is that in fever, where there is a great determination to the lungs, and the respiration appears to be locally affected, independently of the oppression produced by mere febrile increase of circulation, practitioners have avoided the external use of cold, in order to promote the solution of the fever; and have trusted to the general antiphlogistic treatment, along with the topically relaxing application of warm vapour, inhaled by the lungs. Warm bathing appears to be peculiarly well calculated to relieve those complaints that seem to depend on an irregular or diminished action of any part of the alimentary canal; and the state of the skin, produced by immersion in warm water, seems highly favourable to the healthy action of the stomach and bowels. Another very important use of the warm bath, is in herpetic eruptions, by relaxing the skin, and rendering it more pervious, and preparing it admirably for receiving the stimulant applications of tar ointment, mercurials, and the like, that are intended to restore it to a healthy state. The constitutions of children seem more extensively relieved by the warm bath than those of adults; and this remedy seems more generally applicable to acute fevers in them than in persons of a more advanced age. Where the warm bath produces its salutary operation, it is almost always followed by an easy and profound sleep. Dr. Saunders strongly recommends the use of the tepid warm bath, or even higher, in the true menorrhagia of females. In paralytic affections of particular parts, the powerful stimulus of heated water is generally allowed; and in these cases, the effect may be assisted by any thing which will increase the stimulating properties of the water, as, for instance, by the addition of salt. In these cases, much benefit may be expected from the use of warm sea-baths. The application of the warm bath topically, as in pediluvia, or fomentations to the feet, often produce the most powerful effects in quieting irritation in fever, and bringing on a sound and refreshing repose. The cases in which the warm bath is likely to be attended with danger, are particularly those

where there exists a strong tendency to a determination of blood to the head; and apoplexy has sometimes been thus brought on. The lowest temperature will be required for cutaneous complaints, and to bring on relaxation in the skin, during febrile irritation; the warmer will be necessary in paralysis; more heat should be employed on a deep-seated part than one that is superficial.

The Vapour Bath.

The vapour bath, called also *Balneum lacenicum*, though not much employed in England, forms a valuable remedy in a variety of cases. In most of the hot natural waters on the Continent, the vapour bath forms a regular part of the bathing apparatus, and is there highly valued. In no country, however, is this application carried to so great an extent as in Russia, where it forms the principal and almost daily luxury of all the people, in every rank; and it is employed as a sovereign remedy for a great variety of disorders. The Hon. Mr. Basil Cochrane has lately published a Treatise on the Vapour Bath, from which, it appears, he has brought the apparatus to such perfection, that he can apply it of all degrees of temperature, partially or generally, by shower, or by stream, with a great force or a small one; according to the particular circumstances under which patients are so variously placed, who require such assistance. See *Cochrane on Vapour Bath*. Connected with this article, is the *air-pump vapour-bath*; a species of vapour bath, or machine, to which the inventor has given this name. This apparatus has been found efficacious in removing paroxysms of the gout, and preventing their recurrence; in acute and chronic rheumatism, palsy, cutaneous diseases, ulcers, &c. It has also been proposed in chilblains, leprosy, yaws, tetanus, amenorrhœa, and dropsy.

II. When the vessels in which bodies are exposed to the action of heat, are not placed in immediate contact with the fire, but receive the required degree of heat by another intermediate body, such apparatus is termed a bath. These have been variously named, as dry, vapour, &c. Modern chymists distinguish three kinds:

1. *Balneum arenae*, or the sand bath. This consists merely of an open iron, or baked clay, sand-pot, whose bottom is mostly convex, and exposed to the furnace. Finely sifted sea-sand is put into this, and the vessel containing the substance to be heated, &c. in the sand bath, immersed in the middle.

2. *Balneum maris*, or the water bath. This is very simple, and requires no particular apparatus. The object is, to place the vessel containing the substance to be heated, in another, containing water; which last must be of such a nature as to

be fitted for the application of fire, as a common still, or kettle.

3. *The vapour bath.* When any substance is heated by the steam, or vapour, of boiling water, chymists say it is done by means of a vapour bath.

III. Those applications are called *dry baths*, which are made of ashes, salt, sand, &c. The ancients had many ways of exciting a sweat, by means of a dry heat; as by the use of hot sand, stove rooms, or artificial bagnios; and even from certain natural hot steams of the earth, received under a proper arch, or hot-house, as we learn from Celsus. They had also another kind of bath by insolation, where the body was exposed to the sun for some time, in order to draw forth the superfluous moisture from the inward parts; and to this day it is a practice, in some nations, to cover the body over with horse-dung, especially in painful chronic diseases. In New England, they make a kind of stove of turf, wherein the sick are shut up to bathe, or sweat. It was probably from a knowledge of this practice, and of the exploded doctrines of Celsus, that the noted empiric Dr. Graham drew his notions of the salutary effects of what he called *earth bathing*; a practice which, in the way he used it, consigned some of his patients to a perpetual mansion under the ground. The like name of *dry bath*, is sometimes also given to another kind of bath, made of kindled coals, or burning spirit of wine. The patient being placed in a convenient close chair, for the reception of the fume, which rises and provokes sweat in a plentiful manner; care being taken to keep the head out, and to secure respiration. This bath has been said to be very effectual in removing old obstinate pains in the limbs.

IV. *Medicated baths* are such as are saturated with various mineral, vegetable, or sometimes animal substances. Thus we have sulphur and iron baths, aromatic and milk baths. There can be no doubt that such ingredients, if duly mixed, and a proper temperature be given to the water, may, in certain complaints, be productive of effects highly beneficial. Water, impregnated with sulphate of iron, will abound with the bracing and sulphureous particles of that metal, and may be useful for strengthening the part to which it is applied, reinvigorating debilitated limbs, stopping various kinds of bleeding, restoring the menstrual and hæmorrhoidal discharges when obstructed, and, in short, as a substitute for the natural iron bath. There are various other medicated baths, such as those prepared with alum and quick-lime, sal-ammoniac, &c. by boiling them together, or separately, in pure rain water. These have long been reputed as eminently serviceable in paralytic, and all

diseases arising from nervous and muscular debility.

BATH WATERS. *Bathonia aqua Solis aque.* *Badlgue aque.* The city of Bath has been celebrated, for a long series of years, for its numerous hot springs, which are of a higher temperature than any in this kingdom, (from 112° to 116° ;) and, indeed, are the only natural waters which we possess that are at all hot to the touch; all the other thermal waters being of a heat below the animal temperature, and only deserving that appellation from being invariably warmer than the general average of the heat of common springs. By the erection of elegant baths, these waters are particularly adapted to the benefit of invalids, who find here a variety of establishments, contributing equally to health, convenience, and amusement. There are three principal springs in the city of Bath, namely, those called the *King's Bath*, the *Cross Bath*, and the *Hot Bath*; all within a short distance of each other, and emptying themselves into the river Avon, after having passed through the several baths. Their supply is so copious, that all the large reservoirs used for bathing are filled every evening with fresh water, from their respective fountains. In their sensible and medicinal properties, there is but a slight difference. According to Dr. Falconer, the former are—1. That the water, when newly drawn, appears clear and colourless, remains perfectly inactive, without bubbles, or any sign of briskness, or effervescence. 2. After being exposed to the open air, for some hours, it becomes rather turbid, by the separation of a pale yellow, ochery precipitate, which gradually subsides. 3. No odour is perceptible from a glass of the fresh water, but a slight pungency to the taste from a large mass of it, when fresh drawn; which, however, is neither fetid nor sulphureous. 4. When hot from the pump, it affects the mouth with a strong chalybeate impression, without being of a saline or pungent taste. And, fifthly, on growing cold, the chalybeate taste is entirely lost, leaving only a very slight sensation on the tongue, by which it can scarcely be distinguished from common hard spring-water. The temperature of the King's Bath water, which is usually preferred for drinking, is, when fresh drawn in the glass, above 116° deg.; that of the Cross Bath, 112° deg. But, after flowing into the spacious bathing vessels, it is generally from 100 to 106° deg. in the hotter baths, and from 92 to 94° deg. in the Cross Bath; a temperature which remains nearly stationary, and is greater than that of any other natural spring in Britain. A small quantity of gas is also disengaged from these waters, which Dr. Priestley first discovered to contain no more than

one-twentieth part of its bulk of fixed air, or carbonic acid. The chymical properties of the Bath waters, according to the most accurate analysers, Doctors Lucas, Falconer, and Gibbs, contain so small a proportion of iron, as to amount only to one-twentieth or one-thirty-eighth of a grain in the pint; and, according to Dr. Gibbs, fifteen grains and a quarter of siliceous earth in the gallon. Dr. Saunders estimates a gallon of the King's Bath water to contain about eight cubic inches of carbonic acid, and a similar quantity of air, nearly azotic, about eighty grains of solid ingredients, one-half of which probably consists of sulphat and muriat of soda, fifteen grains and a half of siliceous earth, and the remainder is selenite, carbonate of lime, and so small a portion of oxyd of iron as to be scarcely calculable. Hence he concludes, that the King's Bath water, is the strongest chalybeate; next in order, the Hot Bath water; and lastly, that of the Cross Bath, which contains the smallest proportions of chalybeate, gaseous and saline, but considerably more of the earthy particles; while its water, in the pump, is also two degrees lower than that of the others. It is likewise now ascertained, that these springs do not exhibit the slightest traces of sulphur, though it was formerly believed, and erroneously supported on the authority of Dr. Charleton, that the subtle aromatic vapour in the Bath waters, was a sulphureous principle, entirely similar to common brimstone.

With regard to the effect of the Bath waters on the human system, independent of their specific properties, as a medicinal remedy not to be imitated completely by any chymical process, Dr. Saunders attributes much of their salubrious influence to the natural degree of warmth peculiar to these springs, which, for ages, have preserved an admirable degree of uniformity of temperature. He thinks too, that one of their most important uses is that of an external application, yet supposes that, in this respect, they appear to differ little from common water, when heated to the same temperature, and applied under similar circumstances.

According to Dr. Falconer, the Bath water, when drunk fresh from the spring, generally raises, or rather accelerates the pulse, increases the heat, and promotes the different secretions. These symptoms, in most cases, become perceptible soon after drinking it, and will sometimes continue for a considerable time. It is, however, remarkable, that they are only produced in invalids. Hence we may conclude, that these waters not only possess heating properties, but their internal use is likewise attended with a peculiar stimulus, acting more immediately on the nerves.

One of the most salutary effects of the Bath water, consists in its action on the urinary organs, even when taken in moderate doses. Its operation on the bowels varies in different individuals, like that of all other waters, which do not contain any cathartic salt; but, in general, it is productive of costiveness: an effect resulting from the want of an active stimulus to the intestines, and probably also from the determination this water occasions to the skin, more than from any astringency which it may possess; for, if perspiration be suddenly checked during the use of it, a diarrhoea is sometimes the consequence. Hence it appears that its stimulant powers are primarily, and more particularly exerted in the stomach, where it produces a variety of symptoms, sometimes slight and transient, but, occasionally, so considerable and permanent, as to require it to be discontinued. In those individuals with whom it is likely to agree, and prove beneficial, the Bath waters excite, at first, an agreeable glowing sensation in the stomach, which is speedily followed by an increase both of appetite and spirits, as well as a quick secretion of urine. In others, when the use of them is attended with head-ache, thirst and constant dryness of the tongue, heaviness, loathing of the stomach, and sickness; or if they are not evacuated, either by urine or an increased perspiration, it may be justly inferred that their further continuance is improper.

The diseases for which these celebrated waters are resorted to, are very numerous, and are some of the most important and difficult of cure of all that come under medical treatment. In most of them, the bath is used along with the waters, as an internal medicine. The general indications, of the propriety of using this medicinal water, are in those cases where a gentle, gradual, and permanent stimulus is required. Bath water may certainly be considered as a chalybeate, in which the iron is very small in quantity, but in a highly active form; and the degree of temperature is in itself a stimulus, often of considerable powers. These circumstances again point out the necessity of certain cautions, which, from a view of the mere quantity of foreign contents, might be thought superfluous. Although, in estimating the powers of this medicine, allowance must be made for local prejudices in its favour, there can be no doubt but that its employment is hazardous, and might often do considerable mischief, in various cases of active inflammation; especially in irritable habits, where there exists a strong tendency to hectic fever; and even in the less inflammatory state of diseased and suppurating viscera; and, in general, wherever a quick pulse and dry tongue, indicate a de-

gree of general fever. The cases, therefore, to which this water are peculiarly suited, are mostly of the chronic kind; and by a steady perseverance in this remedy, very obstinate disorders have given way. The following, Dr. Saunders, in his *Treatise on Mineral Waters*, considers as the principal, viz. 1. Chlorosis, a disease which, at all times, is much relieved by steel, ~~and~~ will bear it, even where there is a considerable degree of feverish irritation, receives particular benefit from the Bath water; and its use, as a warm bath excellently contributes to remove that languor of circulation, and obstruction of the natural evacuations, which constitute the leading features of this common and troublesome disorder. 2. The complicated diseases, which are often brought on by a long residence in hot climates, affecting the secretion of bile, the functions of the stomach, and alimentary canal, and which generally produce organic derangement in some part of the hepatic system, often receive much benefit from the Bath water, if used at a time when suppurative inflammation is not actually present. 3. Another and less active disease of the biliary organs, the jaundice, which arises from a simple obstruction of the gall-ducts, is still oftener removed by both the internal and external use of these waters. 4. In rheumatic complaints, the power of this water, as Dr. Charleton well observes, is chiefly confined to that species of rheumatism which is unattended with inflammation, or in which the patient's pains are not increased by the warmth of his bed. A great number of the patients that resort to Bath, especially those that are admitted into the hospital, are affected with rheumatism in all its stages; and it appears, from the most respectable testimony, that a large proportion of them receive a permanent cure. (See *Falconer on Bath Water in Rheumatic Cases*.) 5. In gout, the greatest benefit is derived from this water, in those cases where it produces anomalous affections of the head, stomach, and bowels; and it is here a principal advantage to be able to bring, by warmth, that active local inflammation in any limb, which relieves all the other troublesome and dangerous symptoms. Hence it is that Bath water is commonly said to produce the gout; by which is only meant that, where persons have a gouty affection, shifting from place to place, and thereby much disordering the system, the internal and external use of the Bath water will soon bring on a general increase of action, indicated by a flushing in the face, fulness in the circulating vessels, and relief of the dyspeptic symptoms; and the whole disorder will terminate in a regular fit of the gout in the extremities, which is the crisis always to be wished for. 6. The colica pictonum, and

the paralysis, or loss of nervous power in particular limbs, which is one of its most serious consequences, is found to be peculiarly relieved by the use of the Bath waters, more especially when applied externally, either generally, or upon the part affected.

The quantity of water taken daily, during a full course, and by adults, is recommended by Dr. Falconer, not to exceed a pint and a half, or two pints: and in chlorosis, with irritable habits, not more than one pint is employed; and when the bath is made use of, it is generally two or three times a week, in the morning. The Bath waters require a considerable time to be persevered in, before a full and fair trial can be made. Chronic rheumatism, habitual gastritis, dyspepsia, from a long course of high and intemperate living, and the like, are disorders not to be removed by a short course of any mineral water, and many of those who have once received benefit at the fountains, find it necessary to make an annual visit to them, to repair the waste in health during the preceding year.

BATH, CAUTERES. A sulphureous bath near Barege, which raises the mercury in Fahrenheit's thermometer to 131 deg.

BATH, ST. SAUVEUR'S. A sulphureous and alkaline bath, in the valley adjoining Barege, the latter of which, raise Fahrenheit's thermometer as high as 131 deg. It is much resorted to from the South of France, and used chiefly externally, as a simple thermal water.

BATHNIS. (From *Bano*, to enter.) *Bathmus*. The seat, or base; the cavity of a bone, with the protuberance of another, particularly those at the articulation of the humerus and ulna, according to Hippocrates and Galen.

BATHONIE AQUE. Bath waters.

BATHRON. (From *Bano*, to enter.) *Bathrum*. The same as bathnis; also an instrument used in the extension of fractured limbs, called *scamnum*. *Hippocrates*. And described by Oribasius and Scultetus.

BATIA. A name formerly given to a retort.

BATINON-MORON. (From *Berus*, a bramble, and *moron*, raspberry. A raspberry.

BATRACHIUM. (From *Bargaxos*, a frog; so called from its likeness to a frog.) The herb crow's foot, or ranunculus.

BATRACHUS. (From *Bargaxos*, a frog; so called because they who are infected with it, croak like a frog.) An inflammatory tumour under the tongue.

BATTARIAMUS. (From *Battor*, a Cyrenian prince, who stammered.) Stammering; a defect in pronunciation. See *Psellismus*.

BATTATA VIRGINIANA. See *Potato*.

BATTATA PEREGRINA. The cathartic potato; perhaps a species of *ipomœa*. If about two ounces of them are eaten at bed-time, they greatly move the belly the next morning.

BATTARISMUS. Stammering with hesitation. The psellismus hæsitans of Cullen.

BAUDA. A vessel for distillation was formerly so called.

Baulmoney. See *Meum athamanticum*.

BAURACH. (Arab. *Bourach*.) A name formerly applied to nitre, or any salt; hence it is that borax took its name, which is also thus called, as well as the mineral fixed alkaline salt.

BAXANA. (Indian.) A poisonous tree growing near Ormuz; called by Ray, rabuxit.

Bay-cherry. See *Lauro-cerasus*.

Bay-leaves. See *Laurus*.

BAY-LEAVED PASSION-FLOWER. The plant so called is the *Passiflora laurifolia* of Linnæus. A native of Surinam, where the fruit grows to the size of a small lemon, which it greatly resembles. Its flavour is delicately acid, and much esteemed to quench thirst. It strengthens the stomach, and is a salutary fruit in gastric affections, fevers, &c.

BAY-SALT. A very pure salt, prepared from sea-water by spontaneous evaporation.

BAZCHER. A Persian word for antidote.

BDELLA. (From *βδελλα*, to suck.)

Bdellium. A horse-leech.

BDELLIUM. (From *bedallah*. Arab.)

Madeleon. Bolchon. Balchus. Called by the Arabians, mokel. A gum-like, very impure myrrh. It is one of the weakest of the deobstruent gums. It was sometimes used as a pectoral and an emmenagogue. Applied externally, it is stimulant, and promotes suppuration. It is never met with in the shops of this country.

BDELLUS. (From *βδελω*, to break wind.)

A discharge of the wind by the anus.

BDELTONIA. (From *βδελω*, to break wind.) Any filthy and nauseous odour.

BEAN. The common bean is the seed of the *Vicia faba* of Linnæus, a native of Egypt. There are many varieties. Beans are very wholesome and nutritious to those whose stomachs are strong, and accustomed to the coarser modes of living. In delicate stomachs they produce flatulency, dyspepsia, cardialgia, &c. especially when old. See *Legumina*.

Bean, French. See *Bean, kidney*.

BEAN, KIDNEY. This seed is often called the French bean; it is the pericarpium of the *phaseolus vulgaris* of Linnæus, which, when young and well boiled, is easy of digestion, and delicately flavoured. These are less liable to produce flatulency than peas. See *Legumina*.

Bean, Malacca. See *Anacardium orientale*.

Bean of Carthage. See *Bejuio*.

Bean, St. Ignatius. See *Nux vomica saprionis*.

BEARD. The hair growing on the chin and adjacent parts of the face, in adults of the male sex.

Beard's-weed. See *Acanthus*.

Beard's-foot. See *Helleboraster*.

Beard's whortleberry. *Uva ursi*.

BECCA. A fine kind of resin from the turpentine and mastich trees of Greece and Syria, formerly held in great repute.

BECCABUNGA. (From *bach bungen*, water-herb, German, because it grows in rivulets.) *Anagallis aquatica. Laver Germanicum. Veronica aquatica. Cepœa.* Water-pimpernel and brooklime. The plant which bears these names, is the *Veronica beccabunga* of Linnæus:—*racemis lateribus, foliis ovatis planis, caule repente.* It was formerly considered of much use in several diseases, and was applied externally to wounds and ulcers: but if it have any peculiar efficacy, it is to be derived from its antiscarbutic virtue. As a mild refrigerant juice, it is preferred where an acrimonious state of the fluids prevails, indicated by prurient eruptions upon the skin, or in what has been called the hot scurvy. To derive much advantage from it, the juice ought to be taken in large quantities, or the fresh plant eaten as food.

BECHA. See *Bechica*.

BECHICA. (From *βηχέ*, a cough.) *Bechita.* Medicines to relieve a cough. An obsolete term. The *trochisci bechici albi*, consist of starch and liquorice, with a small proportion of florentine orris made into lozenges, with mucilage of gum-tragacanth. They are a soft pleasant demulcent. The *trochisci bechici nigri*, consist chiefly of the juice of liquorice, with sugar and gum-tragacanth.

BECHION. (From *βηχέ*, a cough: so called from its supposed virtues in relieving coughs *Bechium.* The herb colt's foot, or tussilago.

BEQUIBA NUX. (Indian.) A large nut growing in Brazil, from which a balsam is drawn that is held in estimation in rheumatisms.

BEDEGUAR. (Arab.) *Bedeguar.* The *Carduus lacteus Syriacus* is so called, and also the *cynosbatus*, or *rosa canina*.

BEDEGSIAN. The name of the love-apples in Avicenna:

Bedetraw, lady's. See *Aparine*.

BEE. *Apis mellifica* of Linnæus. This insect was formerly exhibited, after being dried and powdered, internally, as a diuretic. It is to the industry of bees we are indebted for those valuable articles, honey and wax. See *Honey and Wax*.

Beech-tree. See *Fagus*.

BEE'S WAX. *Cera.* The production of the honey-comb: it is a hard compact substance, and of a clear yellow colour, much used for medical purposes externally.

in the composition of ointments, cerates, and plaisters.

Beet, red. See *Beta rubra*.

BEEF, WHITE. A variety of red beet. The juice and powder of the root are good to excite sneezing, and will bring away a considerable quantity of mucus.

BEMA. (From *βημα*, to cough.) A cough. Expectorated mucus, according to Hippocrates.

BEHEN ALBUM. (From *behen*, a finger, Arab.) *Jacea orientalis patula*. *Raphaniticoides lutea*. The true white behen of the ancients. *Centaurea behen* of Linnæus. The root possesses astringent virtues.

BEHEN OFFICINARUM. The spatting poppy—*Cucubalus behen* of Linnæus, formerly used as a cordial and alexipharmic.

BEHEN RUBRUM. *Limonium*. *Limonium majus*. Behen. Sea-lavender, or red behen. The *Statice limonium* of Linnæus. The roots possess astringent and strengthening qualities, but not in a very remarkable degree.

BEIDELBAR. *Beidellop*. A species of *Asclepias*, used in Africa as a remedy for fever and the bites of serpents. The caustic juice which issues from the roots when wounded, is used by the negroes to destroy venereal and similar swellings.

BEUTIO. *Habilla de Carthagend.* Bean of Carthage. A kind of bean in South America, famed for being an effectual antidote against the poison of all serpents, if a small quantity is eaten immediately. This bean is the peculiar product of the jurisdiction of Carthage.

BELA-AYE CORTEX. (Indian) *Belae*. A bark of Madagascar, said to be of considerable efficacy in the cure of diarrhoeas.

BELEMYOIDES. (From *βελαν*, a dart, and *ως*, form; so named from their dart-like shape.) *Belenoides*. *Beloides*. The styloid process of the temporal bone, and the lower end of the ulna, were formerly so called.

BELESON. (Indian.) *Belilia*. The *Mussaenda frondosa* of Linnæus, a decoction of which is, according to Ray, cooling.

BELLADONNA. (From *bella donna*, Italian, a handsome lady; so called because the ladies of Italy use it, to take away the too florid colour of their faces.) *Solanum melonocerasum*. *Solanum lethale*. *Atropa belladonna* of Linnæus:—*caule herbaceo, foliis ovatis integris*. This plant has been long known as a strong poison of the narcotic kind, and the berries have furnished many instances of their fatal effects, particularly upon children that have been tempted to eat them. The leaves were first used internally, to discuss scirrhus and cancerous tumours; and from the good effects attending their use, physicians were induced to employ them internally, for the same disorders; and there are a consi-

derable number of well authenticated facts, which prove them a very serviceable and important remedy. The dose, at first, should be small; and gradually and cautiously increased. Five grains are considered a powerful dose, and apt to produce dimness of sight, vertigo, &c.

BELLEGU. See *Myrobalani bellerici*.

BELLERESI. See *Myrobalani bellerici*.

BELVILES. See *Myrobalani bellerici*.

BELLERICE. See *Myrobalani bellerici*.

BELLIDIOIDES. (From *bellis*, a daisy, and *ως*, form.) See *Bellis major*.

BELLIS. (*à bello colore*, from its fair colour,) The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia superflua*. The daisy.

BELLIS MAJOR. *Euphthalmum majus*. *Leucanthemum vulgare*. *Bellikoides*. *Consolida media*. *Oculus bovis*. Ox-eye daisy. Maudin-wort. The pharmacopœal name for the *Chrysanthemum leucanthemum* of Linnæus:—*foliis amplexicaulibus, oblongis superne serratis, inferne dentatis*. The flowers and herb were formerly esteemed in asthmatic and phthisical diseases, but have now deservedly fallen into disuse.

BELLIS MINOR. *Bellis*. The common daisy. The *Bellis perennis* of Linnæus:—*scapo nudo, or bruise-wort*, was formerly directed in pharmacopœias by this name. Although the leaves and flowers are rather acrid, and are said to cure several species of wounds, they are never employed by modern surgeons.

BELLIS PERENNIS. The systematic name of the common daisy. See *Bellis minor*.

BELLOCLUS. (From *bellus*, fair, and *oculus*, the eye.) A precious stone, resembling the eye, and formerly supposed to be useful in its disorders.

BELLON. The Devonshire colic, or colica pictonum.

BELLONARIA. (From *Bellona*, the goddess of war.) An herb which, if eaten, makes people mad, and act outrageously, like the votaries of Bellona.

BELLUTTA TSJAMPACAM. (Indian.) A tree of Malabar, to which many virtues are attributed.

BELNUSCHUS. A name given to the *Abelmoschus*.

BELORRE. (Indian.) An evergreen plant of America, whose seeds purge moderately, but the leaves roughly.

BELOROIDES. *Beloides*. The same as *belemnoides*.

BELULCUM. (From *βελος*, a dart, and *ως*, to draw out.) A surgeon's instrument for extracting thorns, or darts.

BELUZZAR. *Beluzaar*. The Chaldean word for antidote.

BELZOR. See *Benzoinum*.

BELZOINUM. See *Benzoinum*.

BEMURINI. An Indian shrub used in gout.

BEN-TAMARA. (Arab.) The *fabia Egyptiaca*.

BEN. (Arab.) *Glans unguentaria*. *Ben nux.* *Balanus narepaica*. *Coatis*. The oily acorn, or ben-nut. A whitish nut, about the size of a small filberd, of a roundish triangular shape, including a kernel of the same figure, covered with a white skin. It is the fruit of the *Gustlandina moringa* of Linnæus:—*internis, foliis, subpinnatis, foliolis inferioribus ternatis*. They were formerly employed to remove obstructions of the primæ viz. The oil afforded by simple pressure, is remarkable for its not growing rancid in keeping, or, at least, not until it has stood for a number of years; and, on this account, it is used in extricating the aromatic principle of such odorous flowers as yield little or no essential oil in distillation. The unalterability of this oil would render it the most valuable substance for cerates, or liniments, were it sufficiently common. It is actually employed for this purpose in many parts of Italy.

BEN MAGNUM. Monardus calls by this name, the avellana purgatrix, which purges and vomits violently.

BEN TAMARA. The Egyptian bean.

BENATH. (Arab.) Small pustules produced by sweating in the night.

BENEDICT. (From *benedico*, to bless.) A specific name prefixed to many compositions and herbs on account of their supposed good qualities; as *benedicta herba*, *benedicta aqua*, &c.

BENEDICTA AQUA. Lime water was formerly so called; also, a water distilled from *serpyllum*, and, in Schroeder, it is the name for an emetic.

BENEDICTA AQUA COMPOSITA. Compound lime water.

BENEDICTA HERBA. See *Caryophyllata*.

BENEDICTA LAXATIVA. A compound of turbeth, scammony, and spurge, with some warm aromatics.

BENEDICTUM LAXATIVUM. Rhubarb, and sometimes the lenitive electuary.

BENEDICTUM LICTUM. A term applied to *Guaiacum*.

BENEDICTUM VINUM. Antimonial wine.

BENEDICTUS CARDUS. See *Carduus benedictus*.

BENEDICTUS LAPIS. A name for the philosopher's stone.

BENEOLENTIA. (From *bene*, well and *oleo*, to smell.) Sweet-scented medicines, as gums, &c.

BENG. A name given by the Mahomedans to the leaves of hemp, formed into pills, or conserve. They possess exhilarating and intoxicating powers.

BENGALÉ RADIX. (From *Bengal*, its native place.) See *Cassumuniar*.

BENGAL QUINCE. This fruit, which is the produce of the *Erateva marmelos* of Linnæus, of spontaneous growth in several

parts of India, is about the size of an orange, and covered with a hard bony shell, containing a yellow viscous pulp, of a most agreeable flavour; this is scooped out, and being mixed with sugar and orange, is brought to the tables of the grandes in India, who eat it as a great delicacy. It is also esteemed as a sovereign remedy against dysentery.

BENGALÉ INDORUM. (From *Bengal*, its native place.) See *Cassumuniar*.

BENGEI ERRI. A species of evergreen, Indian ricinus, which grows in Malabar.

Benit herb. See *Caryophyllata*.

BENIVI ARBOR. See *Benjaminum*.

Benjamin. See *Benzoinum*.

Benzinum flowers. See *Benzoic acid*.

BENZOAS. A benzoate. A salt formed by the union of benzoic acid, with an alkaline, earthy, or metallic base; as benzoate of alumine, &c.

BENZOE. See *Benzoinum*.

BENZOE AMYDALOIDES. See *Benzoinum*.

BENZOIC ACID. *Acidum benzoicum*.

Flores benzoes. *Flores benzoini.* *Benjamin flowers.*

This acid exists in several balsams, but principally in the concrete balsam, called benzoin. (See *Benzoinum*.) Chymists have obtained it from this balsam in various ways, either by sublimation, which gives beautiful foliated crystals, but requires to be repeated thrice, and pressed between bibulous paper after each sublimation, to obtain them white and free from any adherent essential oil: or, by forming some of its soluble compounds, and afterwards decomposing them, so as to precipitate the acid; or, by simply boiling the benzoin in water, which dissolves the acid, and, as it cools, allows it to separate again. The London Pharmacopœia directs it to be obtained thus:—Take of benzoin, a pound and a half: fresh lime, four ounces: water, a gallon and a half: muratic acid, four fluid-ounces. Rub together the benzoin and lime; then boil them in a gallon of the water, for half an hour, constantly stirring; and, when it is cold, pour off the liquor. Boil what remains, a second time, in four pints of water, and pour off the liquor as before. Mix the liquors, and boil down to half, then strain through paper, and add the muriatic acid gradually, until it ceases to produce a precipitate. Lastly, having poured off the liquor, dry the powder in a gentle heat; put it into a proper vessel, placed in a sand bath; and by a very gentle fire sublime the benzoic acid.

The Edinburgh Pharmacopœia forms a benzoate of soda, precipitates the acid by sulphuric acid, and afterwards crystallizes it by solution in hot water, which dissolves a larger quantity than cold.

Benzoic acid has a strong, pungent, aro-

matic, and peculiar odour. Its crystals are ductile, not pulverizable; it sublimes in a moderate heat, forming a white irritating smoke. It is soluble in about twenty-four times its weight of boiling water, which, as it cools, precipitates 19-20ths of what it had previously dissolved. It is soluble in alcohol.

Benzonic acid is very seldom used in the cure of diseases; but now and then it is ordered as a stimulant against convulsive coughs and difficulty of breathing. The dose is from one grain to five.

It combines with alkaline, metallic, and earthy bodies; and forms BENZOATES.

BENZOIFERA. *Benzoinum*.

BENZOINUM. (From the Arabic term *benzouh*.) *Bonjainum*. *Asa dulcis*. *Asa odorata*. *Liquor cyreniacus*. *Bulzoinum*. *Benzoin*. *Benjai*. *Benjua*. *Gumbenjamin*. This substance is classed, by modern chymists, among the balsams. There are two kinds of benzoin: *benzoe amygdaloides*, which is formed of white tears, resembling almonds, united together by a brown matter; and *common benzoin*, which is brown and without tears. The tree which affords this balsam, formerly called *Laurus benzoin*. *Benzosifera*. *Arbor benic*, is the *Styrax benzoin*, *foliis oblongis acuminatis, subus tomentosis, racemis compositis longitudine foliorum* of Dryander, from which it is obtained by incisions. The benzoin of the shops is usually in very large brittle masses. When chewed, it impresses very little taste, except that it impresses on the palate a slight sweetness; its smell, especially when rubbed or heated, is extremely fragrant and agreeable. It has rarely been used medicinally in a simple state, but its preparations are much esteemed against inveterate coughs and phthisical complaints, unattended with much fever; it has also been used as a cosmetic, and in the way of fumigation, for the resolution of indolent tumours. The acid of benzoin is employed in the *tinctura camphore composita*, and a tincture is directed to be made of the balsam.

BENTON FLORES. See *Benzonic acid*.

BENZOINI MAGISTERIUM. Magistery or precipitate of gum-benjamin.

BENZOINI OLEUM. Oil of benjamin.

BERBERIS. (*Berberi*, wild. Arab. used by Averrhoes and the officinal writers.)

1. The name of a genus of plants in the Linnaean system. Class, *Hexandria*. Order, *Monogynia*. The barberry, or pepperidge bush.

2. The pharmacopœical name for the common barberry, or pepperidge bush. *Oxyacantha Galeni*. *Spina acida Cræpina*.

This tree, *Berberis vulgaris* of Linnaeus: *pedunculis racemosis, spinis triplacibus*, is a native of England. The fruit or berries, which are gratefully acid, and moderately adstringent, are said to be of great

use in biliary fluxes, and in all cases where heat, acrimony, and putridity of the humours prevail. The filaments of this shrub possess a remarkable degree of irritability; for on being touched near the base with the point of a pin, a sudden contraction is produced, which may be repeated several times.

BERBERIS OELATINA. Barberries boiled in sugar.

BERBERIS VULGARIS. The systematic name for the berberis of the pharmacopœias. See *Berberis*.

BEREDINUM. An ointment.

BERENICE. (The city from whence it was formerly brought.) Amber.

BERENICUM. (From *bera*, to bring, and *nia*, victory.) A term applied by the old Greek writers to nitre, from its supposed power in healing wounds.

BERENI SCUM. Mugwort. See *Artemisia vulgaris*.

BERGAMOTE. A species of citron. *Citrus mella rosea* of Lamarck; and a variety of the *citrus medica* of Linnaeus. It was produced, at first, casually, by an Italian grafting a citron on a stock of a bergamot pear-tree; whence the fruit produced by this union, participated both of the citron tree and the pear-tree. The essence prepared from this fruit is called essence of bergamote and essentia de cedra.

BERIBERI. (An Hindostane word signifying a sheep.) *Berberia*. A species of palsy, common in some parts of the East Indies, according to Bontius. In this disease, the patients lift up their legs very much in the same manner as is usual with sheep. Bontius adds that this palsy is a kind of trembling in, which there is deprivation of the motion and sensation of the hands and feet, and sometimes of the body.

Bermudas berry. See *Saponaria nucula*.

BERNARVI. An electuary.

BERRIONIS. A name for colophony, or black rosin.

BERS. Formerly the name of an exhilarating electuary.

BERULA. An old name for brooklime.

BERULA GALLICA. Upright water parsnip.

BERYTTION. (From *Berytus*, its inventor. A collyrium described by Galen.

BES. An eight-ounce measure.

BESACHAR. An obsolete term for a sponge.

BESASA. Formerly applied to wild rue.

BESBAH. An old name for mace.

BESBENA. (Arab.) *Muscarum Fungus*. Probably a sponge, which is the nidus of some sorts of flies.

BESBAHEN. (Arab.) A redness of the external parts, resembling that which precedes the leprosy; it occupies the face and extremities. *Avicenna*.

BESRO. A name in Oribasius for *Saxifraga*.

BETA. (So called from the river *Betis*, in Spain, where it grows naturally; or, according to Blanchard, from the Greek letter *beta*, which it is said to resemble when turgid with seed.)

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*. The beet.

2. The pharmacopœial name of the common beet.

Beta vulgaris of Linnæus:—*floribus congestis*. The root of this plant is frequently eaten by the French; it may be considered as nutritious and antiscorbutic, and forms a very elegant pickle with vinegar. The root and leaves, although formerly employed as laxatives and emollients, are now forgotten. A considerable quantity of sugar may be obtained from the root of the beet. It is likewise said, that if beet roots be dried in the same manner as malt, after the greater part of their juice is pressed out, very good beer may be made from them. It is occasionally used to improve the colour of claret.

BETA VULGARIS. The systematic name for the beet of the pharmacopœias. See *Beta*.

BETELE. *Bethle. Betle. Betelle.* An oriental plant, like the tail of a lizard. It is chewed by the Indians, and makes the teeth black; is cordial and exhilarating, and in very general use throughout the East. It is supposed to be the long pepper.

BETONICA. (Corrupted from *Vettonica*, which is derived from the *Vettones*, an ancient people of Spain.)

1. The name of a genus of plants in the Linnæan system. Class, *Didynamia*. Order, *Gymnospermia*. Betony.

2. The pharmacopœial name for the wood betony.

Betonica purpurea. Vetonica cordi. Betonica officinalis of Linnæus:—*spica interrupta, corollarum labii lacinia intermedia emarginata*, common in our woods and heaths. The leaves and tops of this plant have an agreeable, but weak smell; and to the taste they discover a slight warmth, accompanied with some degree of astringency and bitterness. The powder of the leaves of betony, snuffed up the nose, provoke sneezing; and hence it is sometimes made an ingredient in sternutatory powders. Its leaves are sometimes smoked like tobacco. The roots differ greatly, in their quality, from the other parts; their taste is very bitter and nauseous; taken in a small dose, they vomit and purge violently, and are supposed to have somewhat in common with the roots of hellebore. Like many other plants, formerly in high medical estimation, betony is now almost entirely neglected. Antonius Musa, physician to the Emperor Augustus, filled a whole volume with enumerating its virtues, stating it as a remedy for no less than forty-seven disorders: and hence in

Italy the proverbial compliment, *You have more virtues than betony*.

BETONICA AQUATICA. *Scrophularia aquatica*. Greater waterfigwort. Waterbetony. The leaves of this plant, *Scrophularia aquatica* of Linnæus:—*foliis cordatis obtusis, petiolatis, decurrentibus; caule membranaceo angulato; racemis terminalibus*, are celebrated, as correctors of the ill flavour of senna. They were, also, formerly in high estimation against piles, tumours of the scrophulous nature, inflammations, &c.

BETONICA PAULI. A species of veronica.

BETONICA VULGARIS. The systematic name of the *betonica* of the pharmacopœias. See *Betonica*.

Betony, water. See *Betonica aquatica*.

BETULA. 1. The name of a genus of plants in the Linnæan system. Class, *Monoclea*. Order, *Tetrandria*. Alder and birch.

2. The pharmacopœial name of the white birch. *Betula alba*, of Linnæus:—*foliis ovatis, acuminatis, serratis*.

The juice, leaves, and bark of this tree, have been employed medicinally. If the tree be bored early in the spring, there issues, by degrees, a large quantity of limpid, watery, sweetish juice: it is said that one tree will afford from one to two gallons a day. This juice is esteemed as an antiscorbutic, deobstruent, and diuretic. The leaves and bark are used externally as resolvers, detergents, and antiseptics.

BETULA ALBA. The systematic name for the *betula* of the pharmacopœias. See *Betula*.

BETULA ALNUS. The systematic name for the *alnus* of the pharmacopœias. See *Alnus*.

BEX. (From *fluere*, to cough.) A cough.

BEXUSO. The root of the *Emathie Peruviana* of Casper Bauhin; one drachm of which is sufficient for a purge.

BEZAGUILLO. A name given to the white ipecacuanha, which the Spaniards bring from Peru, as the Portuguese do the brown from Brazil.

BEZAHAR. The fossile bezoar.

BEZETTA CORULEA. *Succia heliotropii. Lacmus seu torne. Lacca corulea. Litmus.* The juice of the *Croton tinctorium*; *foliis rhombeis repandis, capsulis pendulis, caule herbaceo*, of Linnæus. It is much used by chymists as a test. See *Testa*.

BEZOAR. (From *pa-zohar*, Persian, a destroyer of poison.) *Ispis bezardicus*. Bezoard. A preternatural or morbid concretion formed in the bodies of land-animals. Several of these kinds of substances were formerly celebrated for their medicinal virtues, and distinguished by the names of the countries from whence they came, or the animal in which they were found. They were considered as high alexipharmics, in so much so, that other medicines, possessed, or supposed to be possessed of alexipharmic powers, were called *bezoar-*

dies; and so efficacious were they once thought, that they were bought for ten times their weight in gold. These virtues, however, are in the present day justly denied them, as they produce no other effects than those common to the saline particles which they contain, and which may be given to greater advantage from other sources. A composition of bezoar with absorbent powders, has been much in repute, as a popular remedy for disorders in children, by the name of Gascoigne's powder and Gascoigne's ball; but the real bezoar was rarely, if ever, used for these, its price offering such a temptation to counterfeited it. Some have employed, for this purpose, a resinous composition, capable of melting in the fire and soluble in alcohol; but Newmann supposed that those nearest resembling it, were made of gypsum, chalk, or some other earth, to which the proper colour was imparted by some vegetable juice. We understand, however, that tobacco pipe clay, tinged with oxgall, is commonly employed, at least for the Gascoigne's powder; this giving a yellow tint to paper, rubbed with chalk, and a green to paper rubbed over with quicklime; which are considered as proofs of genuine bezoar, and which a vegetable juice would not effect.

BEZOAR BOVINUM. the bezoar from the ox.

BEZOAR GERMANICUM. The bezoar from the alpine goat.

BEZOAR HYSTRIS. *Lapis porcinus. Lapis malacensis. Pedro del porco.* The bezoar of the Indian porcupine. Said to be found in the gall-bladder of an Indian porcupine, particularly in the province of Malacca. This concrete differs from others; it has an intensely bitter taste; and on being steeped in water for a very little time, impregnates the fluid with its bitterness, and with aperient, stomachic, and, as it is supposed, with alexipharmic virtues. How far it differs in virtue from the similar concretions found in the gall-bladder of the ox, and other animals, does not appear.

BEZOAR OCCIDENTALE. Occidental bezoar. This concretion is said to be found in the stomach of an animal of the stag or goat kind, a native of Peru, &c. It is of a larger size than the oriental bezoar, and sometimes as large as a hen's egg; its surface is rough, and the colour green, greyish, or brown.

BEZOAR ORIENTALE. *Lapis bezoar orientalis.* Oriental bezoar stone. This concretion is said to be found in the pylorus, or fourth stomach of an animal of the goat kind, which inhabits the mountains of Persia. It is generally about the size of a kidney bean, of a roundish or oblong figure, smooth, and of a shining olive or dark greenish colour.

BEZOAR MICROCOSMICUM. The calculus found in the human bladder.

BEZOAR PORCINUM. See *Bezoar hystris*.

BEZOAR SIMIÆ. The bezoar of the monkey.

BEZOARDICA RADIX. See *Contragerva*.

BEZOARDICUM JOVIALE. *Bezoar* with tin. It differed very little from the *Antisepticum Poterii*.

BEZOARDICUM LUNALE. A preparation of antimony and silver.

BEZOARDICUM MARTIALE. A preparation of iron and antimony.

BEZOARDICUM MINERALE. A preparation of antimony, made by adding nitrous acid to butter of antimony.

BEZOARDICUM SATURNI. A preparation of antimony and lead.

BEZOARDICUS FULVIS. The powder of the oriental bezoar.

BEZOARTICUM MINERALE. An inert calx of antimony.

BEZOARTICUS SPIRITUS NITRI. The distilled acid of the bezoarticum minerale.

BEZOAR. A common chymical epithet.

BIZON. Wine of sun-raisons and sea-water.

BIBINELLA. See *Pimpinella*.

BIBITORIUS. (*Bibitorius*, sc. *musculus*; from *bibe*, to drink; because by drawing the eye inwards towards the nose, it causes those who drink to look into the cup.) See *Rectus internus oculi*.

BICEPS. (From *bis*, twice, and *caput*, a head.) Many muscles have this denomination, from their having two distinct heads, or origins.

BICEPS BRACHII. See *Biceps flexor cubiti*.

BICEPS CRURIS. See *Biceps flexor cruris*.

BICEPS CUBITI. See *Biceps flexor cubiti*.

BICEPS EXTENSUS. See *Triceps extensor cubiti*.

BICEPS FLEXOR CRURIS. *Biceps cruris* of Albinus. *Biceps* of Winslow, Douglas, and Cowper, and *Ichio-femoro-peronei* of Dumas. A muscle of the leg, situated on the hind part of the thigh. It arises by two distinct heads; the first, called *longus*, arises, in common with the semitendinosus, from the upper and posterior part of the tuberosity of the os ischium. The second, called *brevis*, arises from the linea aspera, a little below the termination of the glutæus maximus, by a fleshy acute beginning, which soon grows broader as it descends to join with the first head, a little above the external condyle of the os femoris. It is inserted, by a strong tendon, into the upper part of the head of the fibula. Its use is to bend the leg. This muscle forms what is called the outer hamstring; and, between it and the inner, the

nervus popliteus, arteria and vena poplitea, are situated.

BICEPS FLEXOR CUBITI. *Biceps brachii* of Albinus. *Coraco-radialis, seu biceps* of Winslow. *Biceps internus* of Douglas. *Biceps internus humeri* of Cowper. *Scapulo coracoradial* of Dumas. A muscle of the fore-arm, situated on the fore part of the *os humeri*. It arises by two heads. The first and outermost, called *longus*, begins tendinous from the upper edge of the glenoid cavity of the scapula, passes over the head of the *os humeri* within the joint, and, in its descent without the joint, is enclosed in a groove near the head of the *os humeri*, by a membranous ligament that proceeds from the capsular ligament and adjacent tendons. The second, or innermost head, called *brevus*, arises, tendinous and fleshy, from the coracoid process of the scapula, in common with the coracobrachialis muscle. A little below the middle of the fore-part of the *os humeri*, these heads unite. It is inserted by a strong roundish tendon into the tubercle on the upper end of the radius internally. Its use is to turn the hand supine, and to bend the fore-arm. At the bending of the elbow, where it begins to grow tendinous, it sends off an aponeurosis, which covers all the muscles on the inside of the fore-arm, and joins with another tendinous membrane, which is sent off from the triceps extensor cubiti, and covers all the muscles on the outside of the fore-arm, and a number of the fibres, from opposite sides, decussate each other. It serves to strengthen the muscles, by keeping them from swelling too much outwardly, when in action, and a number of their fleshy fibres take their origin from it.

BICEPS INTERNUS. See *Biceps flexor cubiti*.

BICORNIS. An epithet of certain peccatorals, or rather troches, described by Rhazes, which were made of liquorice, &c.

BICROS. A Portuguese name for the worms that get under the toe of the people in the Indies, which are destroyed by the oil of cashew nut.

BICORN. (*Bicornis*: from *bis*, twice, and *cornu* a horn.) An epithet sometimes applied to the *os hyoides*, which has two processes, or horns; and likewise, in former times, to muscles that had two terminations.

BICUSPIS. (*Bicuspis*: from *bis*, twice, and *cuspis*, a spear.) The name of those teeth which have double points, or fangs. See *Teeth*.

BIDENS. (From *bis*, twice, and *dens*, a tooth; so called from its being deeply serrated, or indented.)

1. The name of a genus of plants in the Linnean system. Class, *Syngeneria*. Order, *Polygamia equalis*.

BIFURCATED. (*Bifurcus*: from *bis*, twice, and *furca*, a fork.) A vessel, or nerve is said to bifurcate when it divides into two branches; thus the bifurcation of the aorta, &c.

BIGASTER. (*Bigaster*: from *bis*, twice, and *gaster*, a belly.) A name given to muscles which have two bellies.

BIBERNIUS. (From *bis*, double, and *hernia*, a disease so called.) Having a hernia, or rupture on each side of the scrotum.

BILABRE. A name of iron.

BILE. (*Bilis*. Nævius derives it from *bis*, twice, and *lis*, contention; as being supposed to be the cause of anger and dispute.) The gall. A bitter fluid, secreted in the glandular substance of the liver; in part flowing into the intestines, and in part regurgitating into the gall-bladder. The secretory organ of this fluid is the penicilli of the liver, which terminate in very minute canals, called biliary ducts. The biliary ducts exonerate their bile into the *ductus hepaticus*, which conveys it into the *ductus communis choledochus*, from whence it is in part carried into the duodenum. The other part of the bile regurgitates through the cystic duct (see *Gall-bladder*.) into the gall-bladder: for hepatic bile, except during digestion, cannot flow into the duodenum, which contracts when empty; hence it necessarily regurgitates into the gall-bladder. The branches of the *vena porta* contribute most to the secretion of bile; its peculiar blood, returning from the abdominal viscera, is supposed to be, in some respects, different from other venal blood, and to answer exactly to the nature of bile. It is not yet ascertained clearly whether the florid blood, in the hepatic artery, merely nourishes the liver, or whether, at the same time, it contributes a certain principle, necessary for the formation of bile. It has been supposed, by physiologists, that cystic bile was secreted by the arterial vessels of the gall-bladder; but the fallacy of this opinion is proved by making a ligature on the cystic duct of a living animal. From what has been said, it appears that there are, as it were, two kinds of bile in the human body: 1. *Hepatic bile*, which flows from the liver into the duodenum: this is thin, of a faint yellow colour, inodorous, and very slightly bitter, otherwise the liver of animals would not be eatable. 2. *Cystic bile*, which regurgitates from the hepatic duct into the gall-bladder, and there, from stagnating, becomes thicker, the aqueous part being absorbed by lymphatic vessels, and more acrid from concentration. Healthy bile is of a yellow, green colour; of a plastic consistence, like thin oil, and when very much agitated, it froths like soap and water: its smell is fatuous, somewhat like

must, especially the putrefying or evaporated bile of animals: its taste is bitter. The constituent principles of bile are: 1. *Water*, which constitutes the greatest part of bile. 2. *An albuminous principle*, precipitated by alcohol and mineral acids. 3. *A resinous principle*, obtained by evaporating a tincture made of alcohol and bile. 4. *A colouring principle*, which adheres to the resinous part, and gives the colour to bile. 5. *Soda*, in its caustic state: hence healthy bile does not effervesce with acids, and affords a neutral salt. 6. *A phosphated calx*. The primary uses of this fluid, so important in the animal economy, are: 1. *To extricate the chyle from the chyme*: thus chyle is never observed in the duodenum before the chyme has been mixed with the bile: and thus it is that oil is extricated from linen by the bile of animals. 2. By its *acridity* it excites the peristaltic motion of the intestines; hence the bowels are so inactive in people with jaundice. 3. It imparts a *yellow colour* to the excrements; thus the white colour of feces in jaundice, in which disease the flow of bile into the duodenum is entirely prevented. 4. It prevents the *abundance of mucus and acidity* in the primæ viæ; hence acid, pituitous, and verminous saburra are common from deficient or inert bile.

BILIARY DUCT. *Ductus biliaris*. The very vascular *glandules*, which compose almost the whole substance of the liver, terminate in very small canals, called *biliary ducts*, which at length form one trunk, the *ductus hepaticus*. Their use is to convey the bile, secreted by the liver, into the hepatic duct: this uniting with a duct from the gall-bladder, forms one common canal, called the *ductus communis choledochus*, which conveys the bile into the intestinal canal.

BILIMBI, (Indian.) *Biting-bing* of Boninus. The *malus indica*:—*fructus pentagono* of Europeans. It is carefully cultivated in the gardens of the East-Indies, where it flowers throughout the year. The juice of the root is cooling and drank as a cure for fevers. The leaves boiled and made into a cataplasm with rice, are famed in all sorts of tumours, and the juice of the fruit is used in almost all external heats, dipping linnen rags in it, and applying them to the parts. It is drank, mixed with arrack, to cure diarrhæas; and the dried leaves, mixed with betel leaves and given in arrack, are said to promote delivery. The ripe fruit is eaten as a delicacy, and the unripe made into a pickle for the use of the table.

BILIOUS. A term very generally made use of, to express diseases which arise from too copious a secretion of bile: thus bilious colic, bilious diarrhæa, bilious fever, &c.

BILIS See *Bile*.

BILIS ATRA. Black bile. The supposed cause among the ancients of melancholy.

BILIS CYSTICA. *Bilis fellea*. Cystic bile. The bile when in the gall-bladder is so called to distinguish it from that which is found in the liver.

BILIS HEPATICA. Hepatic bile. The bile when it has not entered the gall-bladder.

BILOBUS. (From *bis*, double, and *lobus*, the end of the ear.) Having two lobes, resembling the tips of ears.

BIMESTRIS. (From *bis*, twice, and *mensis*, month.) Two months old.

Bindweed, See *Convolvulus major*.

BINGALLS. See *Canumunar*.

BINGOLUS. (From *binus*, double, and *oculus*, the eye.) A bandage for securing the dressings on both eyes.

BINSICA. A disordered mind. *Helmant*.

BINSICA MORS. The binsical, or that death which follows a disordered mind.

BIOLITHIUM, (From *bios*, life, and *lithum*, a lamp.) Vital heat: an official nostrum.

BIOTZ. (From *bios*, life.) Life. Light food.

BIOTHANATI. (From *bios*, life, and *Thanatos*, death.) Those who die a violent death, or suddenly, as if there were no space between life and death.

BIPERMILLA. See *Pimpinella*.

BIPENELLA. See *Pimpinella*.

BIRA. Malt liquor or beer.

BIRAO. Stone parsley.

Birch tree. See *Betula*.

BIRDTONGUE. A name given to the seeds of the *Fraxinus excelsior* of Linnaeus.

BISSEK. (Heb. an aperture) A deep ulcer, or imposthume in the breast.

Birthwort, climbing. See *Aristolochia tenuis*.

BISCOCTUS. (From *bis*, twice, and *coccus*, to boil.) Twice dressed. It is chiefly applied to bread much baked, as biscuit.

BISCUTELLA. Mustard.

BISERMAS. A name formerly given to clay, or garden clay.

Bishop's weed. See *Animi*.

BISLINGUA. (From *bis*, twice, and *lingua*, a tongue; so called from its appearance of being double-tongued; that is, of having upon each leaf a less leaf.) The Alexandrian laurel.

BISMAVA. (From *vismaiva*, quasi *viscum* *malva*, from its superior viscosity.) The water or marsh-mallow.

BISMUTH. (*Bismut*, Germ.) *Bismuthum*.

A metal which is found in the earth in very few different states, more generally native or in the metallic state. Native *bismuth* is met with in solid masses, and also in small particles dispersed in and frequently deposited on different stones, at Schreeberg in Saxony, Sweden, &c. Sometimes it is crystallized in four-sided tables, or indistinct cubes. It exists combined with oxygen in the *oxyd of bismuth* (*bismuth ochre*), found in small particles, dispersed, of a blueish or yellowish-gray colour, needle-shaped and

capillary; sometimes laminated, forming small cells. It is also though more seldom united to sulphur and iron in the form of a sulphuret in the *marial sulphurated bismuth ore*. This ore has a yellowish-gray appearance, resembling somewhat the marial pyrites. And, it is sometimes combined with arsenic.

Properties.—Bismuth is of a silver-white colour inclining to red. It soon tarnishes and becomes iridescent. It is brittle and can easily be reduced to small particles. It is soft enough to be cut with a knife. It has a lamellated texture. Its specific gravity is 9.800. It requires less heat for fusion than any other metal, tin excepted, melting by a heat — 460 deg. Fahr. It can be volatilized by heat and escapes in the state of grayish-white vapour. It readily unites with mercury and with sulphur. When fused, it exhibits on cooling, cubical figures on the surface. It is soluble in sulphuric, nitric, and muriatic acids. The solution in nitric acid is decomposable by mere dilution with pure water. It inflames in oxygenated muriatic acid gas. It is capable of combining with the greatest number of the metals; and when in certain proportions, promotes their fusibility remarkably. It speedily becomes black by sulphurated hydrogen gas.

BISMUTHUM. (From *bismut*. German.) See *Bismuth*.

Bistort See *Bistorta*.

BISTORTA. (From *bis*, twice, and *torqueo*, to bend; so called from the contortions of its roots.) *Bistort*.

Polygonum bistorta of Linnæus:—*caule simplicissimo monostachio, foliis ovatis in petiolum decurrentibus*. This plant is a native of Britain. Every part manifests a degree of stipticity to the taste, and the root is esteemed to be one of the most powerful of the vegetable adstringents, and frequently made use of as such, in disorders proceeding from a laxity and debility of the solids, for restraining alvine fluxes, after due evacuations, and other preternatural discharges both serous and sanguineous. It has been sometimes given in intermitting fevers; and sometimes also, in small doses, as a corroborant and antiseptic, in acute malignant and colliquative fevers; in which intentions Peruvian bark has now deservedly superseded both these and all other adstringents. The common dose of bistort root in substance, is 15 or 20 grains: in urgent cases it is extended to a dram. Its astringent matter is totally dissolved both by water and rectified spirits.

BISTORTY. (*Bistoir*. French.) Any sm. II knife for surgical purposes.

BITUKOS. A Galenical plaister.

BITUMICI EMPLASTRUM. A plaister for the spleen.

Bitter apple. See *Colocynthis*.

BIT NOBEN. Salt of bitumen. A

white saline substance has lately been imported from India by this name, which is not a natural production, but a Hindoo preparation of great antiquity. It is called in the country *bit noben*, *padnoon*, and *soucheron* and popularly *khala mimur*, or black salt. Mr Henderson, of Bengal, conjectures it to be the *sal asphaltites*, and *sal sodomenus* of Pliny and Galen. This salt is far more extensively used in Hindostan than any other medicine whatever. The Hindoos use it to improve their appetite and digestion. They consider it as a specific for obstructions of the liver and spleen; and it is in high estimation with them in paralytic disorders, particularly those that affect the organs of speech, cutaneous affections, worms, old rheumatisms, and indeed all chronic disorders of man and beast.

BITUMEN. (*ἔλκευα*, *ἔλκευ*, pine; because it flows from the pine-tree; or, *quoddā tumeat ē terra*, from its bursting forth from the earth.) Bitumens are combustible, solid, soft, or fluid substances, whose smell is strong, acrid, or aromatic, composed of hydrogen and carbon with a contamination of earth and other substances in small proportions. They are found either in the internal part of the earth, or exuding through the clefts of the rocks, or floating on the surface of waters. Like oils they burn with a rapid flame. Natural historians have divided them into several genera; but modern chymists arrange them according to their chymical properties, and are only acquainted with six species, which are very distinct from each other: these are, naphtha, amber, asphaltos, jet, pit-coal, and petroleum.

BITUMEN BARBADENSE. See *Petroleum barbadense*.

BITUMEN JUDÆICUM. *Asphaltus*. Jews' pitch. A solid light bituminous substance, of a dusky colour on the outside, and a deep shining black within; of very little taste, and scarcely any smell, unless heated, when it emits a strong pitchy one. It is said to be found plentifully in the earth in several parts of Egypt, and floating on the surface of the Dead Sea. It is now wholly expunged from the catalogue of officinals of this country; but was formerly esteemed as a discutient, sudorific, and emmenagogue.

BITUMEN LIQUIDUM. See *Petroleum*.

BIVENTER (*Biventer*; from *bis*, twice, and *venter*, a belly.) A muscle is so termed, which has two bellies.

BIVENTER CERVICIS. A muscle of the lower jaw.

BIVENTER MAXILLÆ INFERIORIS. See *Digastricus*.

BIXA ORLEANA. The systematic name for the *terra orleana* of the pharmacopœias. See *Orleana*.

BLACCLE. The measles. *Rhazes*.

BLACKBERRY. The fruit of the common bramble, *Rubus fruticosus* of Linnæus. The berries are eaten in abundance by children, and are wholesome and gently aperient. Too large quantities, however, when the stomach is weak, produce vomiting and great distention of the belly, from flatus. See *Fruits, summer*.

Bladder. See *Urinary bladder*, and *Gall-bladder*.

Bladder, inflamed. See *Cystitis*.

Blade-bone. See *Scapula*.

BLASITAS. (From *blasus*.) A defect in speech, called stammering.

BLASUS. (From *blasia*, to injure.) A stammerer.

BLANCA. (*Blanc*, French.) A purging mixture; so called, because it was supposed to evacuate the white phlegmatic humours. Also white lead.

BLANCA MULIERUM. White lead.

BLASA. (Indian.) A tree, the fruit of which the Indians powder, and use to destroy worms.

BLASTEMA. (From *blasium*, to germinate.) A bud or shoot. Hippocrates uses it to signify a cutaneous pimple like a bud.

BLASTUM MOSYLITUM. Cassia bark kept with the wood.

BLATTA. (From *blasia*, to hurt.) A sort of beetle, or bookworm; so called from its injuring books and clothes: the kermes insect.

BLATTARIA LUTEA. (From *blatta*; so called, because, according to Pliny, it engenders the blatta.) The herb yellow moth-mullein.

BLECHON. (From *blaschomai*, to bleat; so called according to Pliny, because if sheep taste it they bleat.) The herb wild penny-royal.

Bleeding. See *Blood-letting* and *Hæmorrhage*.

Bleeding at the nose. See *Epistaxis*.

BLEMA. (From *blasia*, to inflict.) A wound.

BLENDE. A species of zinc ore, formed of zinc in combination with sulphur.

BLENA. (*Blena*.) *Blena*. Mucus, a thick excrementitious humour.

BLENNORRHOEA. (From *blena*, mucus, and *rho*, to flow.) *Gonorrhœa mucosa*. A gleet. An increased discharge of mucus from the urethra, arising from weakness.

BLENNORRHAGIA. (From *blena*, mucus, and *rho*, to flow.) The discharge of mucus from the urethra.

BLEPHARA. (*Quasi blennus opus*, as being the cover and defence of the sight.) The eyelids.

BLEPHARIDES. (From *blepharon*.) The hair upon the eyelids; also the part of the eyelids where the hair grows.

BLEPHAROPHTHALMIA. (From *blepharon*, the eyelid, and *ophthalmia*, a disease of the eye.) An inflammation of the eyelid.

BLEPHAROPTOSIS. (From *blepharon*, the eyelid, and *ptosis*, from *ptellein*, to fall.) A prolapse, or falling down of the upper eyelid, so as to cover the cornea.

BLEPHAROTIS. (From *blepharon*, the eyelid.) An inflammation of the eyelids.

BLEPHAROSTYX. (From *blepharon*, the eyelid, and *styx*, to scrape off.) A brush for the eyes. An instrument for cleansing or scraping off foul substances from the eye-lids.

BLEPHAROTYXIS. (From *blepharon*, the eyelid, and *styx*, to scrape off.) The cleansing of the eyelids. Inflammation of the eyelids.

Blessed thistle. See *Carduus benedictus*.

BLESTRISMUS. (From *blasia*, to throw about.) Pirenetic restlessness.

BLETA. A word used by Paracelsus to signify white, and applied to urine when it is milky, and proceeds from a disease of the kidneys.

BLETI. (*Bletus*, from *blasia*, to strike.) Those seized with dyspnoea or suffocation.

BLISTER. A topical application, when put on the skin raises the cuticle in the form of a vesicle, filled with a serous fluid. Various substances produce this effect on the skin; but the powder of the *lytta vesicatoria*, or blistering fly, is what operates with most certainty and expedition, and is now invariably made use of for the purpose.

When it is not wished to maintain a discharge from the blistered part, it is sufficient to make a puncture in the cuticle to let out the fluid; but when the case requires keeping up a secretion of pus, the surgeon must remove the whole of the detached cuticle with a pair of scissors, and dress the excoriated surface in a particular manner. Practitioners used formerly to mix powder of cantharides with ointment, and dress the part with this composition. But such a dressing not unfrequently occasioned very painful affections of the bladder, a scalding sensation in making of water, and very afflicting stranguries. The treatment of such complaints consists in removing every particle of the fly from the blistered part, making the patient drink abundantly of mucilaginous drinks, giving emulsions and some doses of camphor.

These objections to the employment of salves containing the *lytta*, for dressing blistered surfaces, led to the use of mezezon, euphorbium, and other irritating substances, which when incorporated with ointment, form very proper compositions for keeping blisters open, which they do without the inconvenience of irritating the bladder, like the blistering fly.—The favourite application however for keeping open blisters, is the powder of savine, which was brought into notice by Mr. Crowthen in his book on white swellings. The following is the form adopted by the London College: "Take of fresh

leaves of savine bruised, a pound; yellow wax, half a pound; prepared lard, two pounds. Having melted together the wax and lard, boil therein the savine leaves, and strain through a linen cloth." On the use of the savine cerate, immediately after the cuticle raised by the blister is removed, says Mr. Crowther, it should be observed that experience has proved the advantage of using the application lowered by a half or two-thirds of the unguentum cere. An attention to this direction will produce less irritation and more discharge, than if the savine cerate were used in its full strength. Mr. Crowther says also, that he has found fomenting the part with flannel, wrung out of warm water, a more easy and preferable way of keeping the blistered surface clean, and fit for the impression of the ointment, than scraping the part, as has been directed by others. An occasional dressing of unguentum resinæ flavæ, he has found a very useful application for rendering the sore free from an appearance of slough, or rather dense lymph, which has sometimes been so firm in its texture as to be separated by the probe, with as much readiness as the cuticle is detached after blistering. As the discharge diminishes, the strength of the savine dressing should be proportionably increased. The ceratum sabinz must be used in a stronger or weaker degree, in proportion to the excitement produced on the patient's skin.

BITUM FETIDUM. See *Atriplex fetida*.

BLOOD. *Sanguis.* A red homogeneous fluid, of a saltish taste, and somewhat urinous smell, and glutinous consistence, which circulates in the cavities of the heart, arteries, and veins. The quantity is estimated to be about twenty-eight pounds in an adult: of this, four parts are contained in the veins, and a fifth in the arteries. The colour of the blood is red; in the arteries it is of a florid hue, in the veins darker; except only the pulmonary veins, in which it is of a lighter cast. Physiology demonstrates, that it acquires this florid colour in passing through the lungs, and from the oxygen it absorbs. The blood is the most important fluid of our body. Some physicians and anatomists have considered it as alive, and have formed many ingenious hypotheses in support of its vitality. The temperature of this fluid is of considerable importance, and appears to depend upon the circulation and respiration. The blood of man, quadrupeds, and birds, is hotter than the medium they inhabit; hence they are termed animals of warm blood; whilst in fishes and reptiles, animals with cold blood, it is nearly of the temperature of the medium they inhabit. The microscope discovers that the blood contains a great number of round globules, which are seen floating about in a yellowish fluid, the serum. The blood also possesses remark-

able physical properties; its taste is saltish, and the smell of its halitus or vapour, when recently drawn, is somewhat urinous; it is of a plastic consistence, somewhat glutinous and adhesive. Chymical analysis of blood, by means of distillation, discovers, 1. A considerable quantity of *insipid water*, which very soon becomes putrid. 2. *Empyreumatic oil*. 3. *Ammoniacal spirit*. 4. *Carbon*, which remains behind, is very spongy, and with great difficulty incinerated. The ashes, however, consist of a small quantity of culinary salt, soda, phosphorated lime, and a very small portion of iron. While hot, and in motion, the blood remains constantly fluid, and red; when it cools, and is at rest, it takes the form of a fluid mass, which gradually and spontaneously separates into two parts; the one, which is red, and floating becomes of a darker colour, remains concrete, and is called the *cruur*, *crassamentum*, or *cake*; the other, which remains behind, the lower part of the vessel, is of a yellow greenish colour, and adhesive, and is called the *serum* or *lymph*.

The cruor forms more than one half of the blood; it is very plastic, thick, and, in consistence, like glutinous jelly. It soon putrefies in the temperature of the air; but, dried by a gentle heat, becomes a brittle, dark, red mass. It is insoluble in water; and, when boiled in it, is converted into a hard grumous mass, internally red. The surface of the cruor of the blood, after being exposed in a vessel to atmospheric air, becomes of a florid red colour; but the inferior surface, contiguous to the vessel, is of a deep black: the change of colour on the surface is owing to the oxygen of the atmosphere uniting with the blood. The cruor of the blood is composed of, 1. *Red globules*, which chymistry demonstrates consist of a fibrous gluten and oxydated iron. The experiments of the celebrated Rhades shew, that in twenty-five pounds of blood from the human body, near two drachms of the oxyd of iron were obtained. 2. The fibrous gluten of the cruor, which remains after washing the cruor of blood for a considerable time in cold water, and enclosed in a fine linen cloth; in which case the red globules are washed away. If the red water obtained in this experiment be evaporated, and then distilled to dryness, it leaves behind a carbon, exhibiting, when incinerated, a great quantity of iron, attractable by the magnet. From these experiments it would appear, that the redness of the globules is imparted from the oxydated iron, for which purpose a small quantity is sufficient.

The serum of the blood is a lymphatic fluid, almost inodorous; rather saltish to the taste; pellucid, and of a yellowish green colour; and rather of a plastic consistence. It forms scarcely one half of the

blood; and it contains, 1. A large portion of water; from forty-seven ounces of serum, forty-three of insipid water were yielded by distillation. 2. *Albuminous gluten*, like the white of an egg, obtained by boiling, or by stirring it with a stick, or by an admixture of alcohol or concentrated mineral acid. 3. *Jelly*. If equal parts of water and serum of the blood be coagulated by fire, that part of the serum which is not coagulated, upon being cooled, puts on the appearance of a tremulous jelly. 4. *Carbonated soda*, obtained by pouring a mineral acid upon recent diluted serum. 5. *Culinary salt*, found in the incinerated carbon of blood. The albuminous principle of the serum, more commonly called the coagulable lymph, appears to be of very considerable importance in the animal economy, both in diseased and healthy states of it: it affords, by analysis, carbon, azot, and hydrogen. The importance of the blood is very considerable; it distends the cavities of the heart and blood-vessels, and prevents them from collapsing; it stimulates to contraction the cavities of the heart and vessels, by which means the circulation of the blood is performed; it generates within itself animal heat, which it propagates throughout the body: it nourishes the whole body: and, lastly, it is that source from which every secretion of the body is separated.

Blood dragon's. See *Sanguis draconis*.

BLOOD-LETTING. Under this term is comprehended every artificial discharge of blood made with a view to cure or prevent a disease. Blood-letting is divided into *general* and *topical*. As examples of the former, *venesection* and *arteriotomy* may be mentioned; and of the latter, the application of *leeches*, *cupping-glasses*, and *scarification*.

Blood, spitting of. See *Hemoptysis*.

Blood, vomiting of. See *Hematemesis*.

Blood-stone. See *Hematites*.

Bloody flux. See *Dysentery*.

BOA (From *boe*, an ox.) A pustulous eruption like the small-pox, so called because it was cured, according to Pliny, by anointing it with hot ox-dung: also the name of a serpent of Calabria; and of the hydra.

BOCHTUM. A decoction of the woods prepared by a second boiling with fresh water.

BOCHIA. A subliming vessel.

BOCHUM. A swelling of the bronchial glands.

BODIES, COMBUSTIBLE. This term is given by chymists to all substances which, on account of their affinity with oxygen, are capable of burning.

BODIES, OASZOUS. See *Gas*.

BODIES, INFLAMMABLE. Chymists give this name to such bodies as burn with facility, and flame in an increased temperature; although, strictly speaking, all combustible

bodies are inflammable bodies: such are the diamond, sulphur, bitumens, &c.

BODIES, PHOSPHORESCENT. Bodies which produce light, though their temperature be not increased.

BODY. Corpus. The human body is divided by anatomists into the trunk and extremities. *i. e.* the head, and inferior and superior extremities, each of which have certain regions before any part is removed, by which the physician is enabled to direct the application of blisters and the like, and the situation of diseases is better described.

The head is distinguished into the hairy part and the face. The former has five regions, viz. the crown of the head or *vertex*, the fore part of the head or *sinciput*, the hind-part or *occiput*, and the sides, *partes laterales capitis*. In the latter are distinguished the region of the forehead, *frons*; temples, or *tempora*; the nose, or *nasus*; the eyes, or *oculi*; the mouth, or *os*; the cheeks, *buccae*; the chin, or *mentum*; and the ears, or *aurae*.

The trunk is distinguished into three principal parts, the neck, thorax, and abdomen. The neck is divided into the anterior region or *pars antica*, in which, in men, is an eminence called *pomum Adami*; the posterior region is called *nucha colli*; and the lateral regions, *partes laterales colli*.

The thorax is distinguished into the anterior region, in which are the *sternum* and *mammæ*, and at whose inferior part is a pit or hollow called *scrobiculus cordis*; a posterior region, called *dorsum*; and the sides, or *lateral thoracis*.

The abdomen is distinguished into an anterior region, properly the *abdomen*; a posterior region, called the loins, or *lumbi*; and lateral regions or flanks, called *lateral abdominis*. The anterior region of the abdomen being very extensive, is subdivided into the *epigastric*, *hypocondriac*, *umbilical*, and *hypogastric* regions, which are described under their respective names. Immediately below the abdomen is the *mons Veneris*, and at its sides the groins or *inguina*. The space between the organs of generation and the *anus*, or fundament, is called the *perineum*.

The superior extremity is distinguished into the shoulder, *summitas humeri*, under which is the arm-pit, called *axilla* or *fovea axillaris*; the *brachium*, or arm; the *antibrachium*, or fore-arm, in which anteriorly is the bend of the arm, where the veins are generally opened, called *flexura antibrachii*, and posteriorly the elbow, called *angulus cubiti*; and the hand, in which are the *carpus* or wrist, the back or *dorsum manus*, and the palm or *vola*.

The inferior extremity is divided into, 1. the region of the femur, in which are distinguished the *crua* or *regio-inchiadilla*, and outer and superior part; 2. the leg, in

which are the knee or *genu*, the bend or *cavum poplitæ*, and the calf or *sura*; 3. the foot, in which are the outer and inner ankle, or *malleolus externus* and *internus*, the back or *dorsum*, and the sole or *planta*.

BOE. (From *βοαι*, to exclaim.) Clamour, or moaning made by a sick person.

BOETHENA. (From *βοηθαι*, to assist.) A remedy.

BOETHEMATICA. (From *βοηθαι*, to assist.) Favourable symptoms.

Bog-bean. See *Trifolium paludosum*.

BOGIA GUMMI. Gamboge.

Bolsa tea. See *Tea*.

Bois de colasi. See *Quassi*.

Bolar earths. See *Bole*.

BOLE. (*Βολα*, a mass.) A friable earthy substance, uniting with water into a smooth paste, adhering to the tongue, and dissolving, as it were, in the mouth; of the argillaceous or clay kind, but more readily imbibing water than the clays strictly so called. Those used in medicine, are the Armenian and French boles. See *Bole Armenian*, and *Bolus Gallicus*. Many other bolar earths have been recommended for medicinal uses, and were formerly ranked amongst the officinals; as red boles from Armenia, Lemnos, Strigonium, Portugal, Tuscany, and Livonia; yellow boles from Armenia, Tockay, Silesia, Bohemia, and Blois; white boles from Armenia, Lemnos, Nocera, Eretria, Lamos, Chio, Malta, Tuscany, and Goltberg. Several of these earths have been commonly made into little cakes or flat masses, and stamped with certain impressions; from which circumstance they received the name of *terre sigillate*, or sealed earths.

BOLE, ARMENIAN. *Bolus Armenia*. Bole-armenic. A pale, but bright red coloured earth, which is occasionally mixed with honey, and applied to children's mouths when afflicted with aphthæ. It forms, like all argillaceous earths, a good tooth-powder, when mixed with some aromatic.

BOLETUS. (From *βολα*, a mass, or *βολη*, from its globular form.)

The name of a genus of plants in the Linnæan system Class, *Cryptogamia*. Order, *Fungi*. Boletus. Spunk.

BOLETUS CERV. The mushroom.

BOLETUS IGNIARIUS. The systematic name for the *agaricus* of the pharmacopœias. See *Agaricus*.

BOLETUS LARICIS. The systematic name for the officinal *agaricus albus*. See *Agaricus albus*.

BOLETUS PINI LARICIS. A species of *agaric*.

BOLETUS SUAVEVOLENS. The systematic name for the *fungus salicis* of the pharmacopœias. See *Fungus salicis*.

BOLIVUS. A voracious appetite, according to Avicenna; but most probably meant for bulimus.

BOLUS. (*Βολα*, a bole or bolus.) Any

medicine, rolled round, that is larger than an ordinary sized pea, and yet not too large to be swallowed.

BOLUS ALEXIPHARMICA. A preparation of contrayerva.

BOLUS EX ALUMINE. Alum, bark, and nutmeg.

BOLUS ARMENA. See *Bole Armenian*.

BOLUS ARMENA ALBA. The white armenian bole.

BOLUS ARMONIAC. See *Bole Armenian*.

BOLUS BLESSENSIS. Bole of Blois. See *Bole Armenian*.

BOLUS GALLICA. French bole. A pale red coloured bolar earth, variegated with irregular specks and veins of white and yellow. It is occasionally administered as an absorbent and antacid.

BOMBAX. *Gossypium*. The cotton-tree. The seeds of the cotton-tree, *Gossypium herbaceum* of Linnæus:—*foliis quinquelobis natis glandulosis, caule herbaceo*, are directed for medicinal use in some foreign pharmacopœias; and are administered in coughs, on account of the mucilage they contain. The cotton, the produce of this tree is well known for domestic purposes.

BOMBIAS. A bombiate. A salt formed by the union of the bombic acid with different bases; thus *bombias of alumine*, &c.

BOMBIC ACID. *Acidum bombycum*. Acid of the silk-worm. Silk-worms contain, especially when in the state of chrysalis, an acid liquor in a reservoir placed near the anus. It is obtained by expressing their juice in a cloth, and precipitating the mucilage by spirit of wine, and likewise by infusing the chrysalides in that liquor. This acid is very penetrating, of a yellow amber colour, but its nature and combinations are not yet well known.

BOMBUS. (*Βαμβος*.) A resounding noise, or ringing of the ears. Also, a sonorous expulsion of flatus from the intestines.

BON ARBOR. A name given to the coffee-tree.

BONA. *Boona*. The phaseolus, or kidney-bean.

BONDUCH INDORUM. Moluca or bezoar nut. The produce of the *Guilandina bonduc* of Linnæus. It possesses warm, bitter, and carminative virtues.

BONE. *Os*. Bones are hard, dry, and insensible parts of the body, of a whitish colour, and composed of a spongy, compact, or reticular substance. They vary much in their appearances, some being long and hollow, others flat and compact, &c. The greater number of bones have several processes and cavities, which are distinguished from their figure, situation, use, &c. Thus processes extended from the end of a bone, if smooth and round, are called *heads*; and *condyles*, when flattened either above or laterally. That part which is beneath the

head, and which exceeds the rest of the bone in smallness and levity, is called the neck. Rough, unequal processes are called *tubercles* or *tubercles*: but the longer and more acute, *spinous* or *styloid* processes, from their resemblance to a thorn. Thin broad processes with sharp extremities, are known by the name of *cristæ*, or *sharp edges*. Other processes are distinguished by their form, and called *alar* or *pterygoid*, *mandibular* or *mastoid*, *dentiform* or *odontoid*, &c. Others, from their situation, are called *superior*, *inferior*, *exterior*, and *interior*. Some have their name from their direction, as *oblique*, *straight*, *transverse*, &c.; and some from their use, as *trochanters*, *rotators* &c. *Furrows*, *depressions*, and *cavities*, are destined either for the reception of contiguous bones, to form an articulation with them, when they are called *articular cavities*, which are sometimes deeper, sometimes shallower; or they receive hard parts, but do not constitute a joint with them. Cavities serve also for the transmission and attachment of soft parts. Various names are given to them, according to the magnitude and figure of bones. If they be broad and large at the beginning, and not deep, but contracted at their ends, they are called *fossæ* or *pits*. *Furrows* are open canals, extending longitudinally in the surface of bones. A hollow, circular tube, for the most part of the same diameter from beginning to end, and more or less crooked, straight, long or short, is named a *canal*. *Foramina* are the apertures of canals, or they are formed of the excavated margins of two bones, placed against each other. If such be the form of the margin of a bone, as if a portion were taken out of it, it is called a *notch*.

With respect to the formation of bone, there have been various opinions. Physiologists of the present day assert that it is from a specific action of small arteries, by which ossific matter is separated from the blood, and deposited where it is required. The first thing observable in the embryo, where bone is to be formed, is a transparent *jelly*, which becomes gradually firmer, and is formed into *cartilage*. The cartilage gradually increases to a certain size, and when the process of ossification commences, vanishes as it advances. Cartilages, previous to the ossific action, are solid, and without any cavity; but when the ossific action of the arteries is about to commence, the absorbents become very active, and form a *small cavity* in which the bony matter is deposited; bone continues to be separated, and the absorbents model the mass into its required shape. The process of ossification is extremely rapid in utero: it advances slowly after birth, and is not completed in the human body till about the twentieth year. Ossification in the flat bones, as those of the skull, always begins from cen-

tral points, and the radiated fibres meet the radii of other ossifying points, or the edges of the adjoining bone. In long bones, as those of the arm and leg, the clavicle, metacarpal, and metatarsal bones, a *central ring* is formed in the body of the bone, the head and extremities being cartilage, in the centre of which ossification afterwards begins. The central ring of the body shoots its bony fibres towards the head and extremities, which extend towards the body of the bone. The head and extremities at length come so close to the body as to be merely separated by a cartilage, which becomes gradually thinner until the twentieth year. Thick and round bones, as those of the tarsus, carpus, sternum, and patella, are, at first, all cartilage: ossification begins in the centre of each. When the bones are deprived of their soft parts, and are hung together in their natural situation, by means of wire, the whole is termed an *artificial skeleton*; but when they are kept together by means of their ligaments, it is called a *natural skeleton*.—The uses of the bones are various, and are to be found in the account of each bone; it is, therefore, only necessary to observe, in this place, that they give shape to the body, contain and defend the vital viscera, and afford an attachment to all the muscles.

A Table of the Bones.

		No.
Bones of the Head.	Bones of the cranium or skull.	Frontal . . . 1
		Parietal . . . 2
		Occipital . . . 1
		Temporal . . . 2
		Ethmoid . . . 1
	Bones of the face.	Sphenoid . . . 1
		Superior maxil. . . 2
		Jugal . . . 2
		Nasal . . . 2
		Lachrymal . . . 3
	Dentes or teeth.	Palatine . . . 2
		Inferior spongy . . . 2
		Vomer . . . 1
	Bone of the tongue.	Inferior maxil. . . 1
		Incisores . . . 8
		Cuspidati . . . 4
Bones of the Trunk.	Bones of the ear, within the temporal bones.	Molares . . . 20
		Hyoides os . . . 1
		Malleus . . . 2
	The spine.	Incus . . . 2
		Stapes . . . 2
		Orbicular os . . . 2
	The thorax.	Cervical . . . 7
		Dorsal . . . 12
		Lumbar . . . 5
	The pelvis.	Sacrum . . . 1
		Coccygis os . . . 1
		Sternum . . . 1
	The pelvis.	Ribs . . . 24
		Innominate ossa . . . 2

Bones of the Upper Extrem.	The shoulder.	Clavicle	- - -	2
		Scapula	- - -	2
	The arm.	Humeri os	- - -	2
		Ulna	- - -	2
	The fore-arm.	Radius	- - -	2
		Naviculare os	- - -	2
		Lunare os	- - -	2
		Cuneiforme os	- - -	2
		Orbiculare os	- - -	2
	The hand.	Trapezium os	- - -	2
		Trapezoides os	- - -	2
		Magnum os	- - -	2
		Unciforme os	- - -	2
	Carpus or wrist.			
	Metacarpus			10
		Phalanges	- - - - -	28

Bones of the Low. Extr.	The thigh.	Femur	- - -	2
		Patella	- - -	2
	The leg.	Tibia	- - -	2
		Fibula	- - -	2
	The foot.	Calcaneus	- - -	2
		Astragalus	- - -	2
		Cuboides os	- - -	2
		Naviculare os	- - -	2
		Cuneiformia ossa	- - -	6
	Tarsus or instep.			
	Metatarsus			10
		Phalanges	- - - - -	28

Sesamond bones of the thumb and great toe, occasionally found } 8

Total 248

Bones, growth of. See Osteogony.

Bonebinder. See Osteocolla.

BONONIENSIS LAPIS. The Bononian stone. Called also *phosphorus bononiensis*, *phosphorus kircheri*, the light-carrier, or Bononian phosphorus. As a medicine the stone is caustic and emetic.

BONUS HENRICUS. (*Henricus*; so called because its virtues were detected by some one whose name was Henry.) *Tota bona*. *Lapathum unctuosum*. *Chenopodium*. English mercury. The plant to which this name is given in the pharmacopœias, is the *Chenopodium bonus Henricus*; *foliis triangulari-sagittatis integerrimis, specie compositis aphyllis axillaribus*, of Linnæus. It is a native of this country, and common in waste grounds from June to August. The young plant differs little from spinach when cultivated; and in many places the young shoots are eaten in spring like asparagus. The leaves of this plant are accounted emollient, and in this intention have been made an ingredient in decoctions for gylsters. They are applied by the common people to flesh wounds and sores under the notion of drawing and healing.

BORACIC ACID. *Acidum boracicum*. Sedative salt of Homberg. Acid of borax. Boracine acid. A concrete salt crystallized in small white scales, which may be obtained from borax, by adding concentrated sulphuric, the nitric, the muriatic, and even vegetable acids, to a hot solution of borax,

till the lixivium becomes somewhat acid: the solution is then to be cooled, when the acid will appear in the shape of bright scales. This acid in combination with alkalies, earths, and metallic oxyds, forms borates.

BORAS. A borate. A salt formed of boracic acid with an earthy, alkaline or metallic base: as borate of soda, &c.

Borage. See Borage.

BORAGO. (Formerly written *Corago*; from *cor*, the heart, and *ago*, to affect; because it was supposed to comfort the heart and spirits.) Borage.

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

2. The pharmacopœial name of the officinal borage. *Buglossum verum*. *Buglossum latifolium*. *Borago hortensis*.

The leaves and flowers of this plant, *Borago officinalis* of Linnæus:—*foliis omnibus alternis, calycibus patentibus*, are esteemed in some countries as refrigerant and cordial. A syrup is prepared from the leaves in France, and used in pleurisies and inflammatory fevers. Their principal use in this island is in that grateful summer beverage, known by the name of cool tankard.

BORAGO OFFICINALIS. The systematic name for the borage of the shops. See Borage.

BORAS SODÆ. Borate of soda. See *Sub boras sodæ* and *Borax*.

BORATE. See Borage.

BORAX. (*Borak*, Arab.) *Boras sodæ*, *sub-boras sodæ*. The obsolete synonyms are *Chrysocolla*, *capistrum auri*, *ancinar*, *boraxitron*, *anucar*, *antincar*, *tincal*, *amphitane*, *baurach*, *nitrum factitium*, *santerna*, and *nitrum nativum*. This salt consisting of boracic acid uniting with soda, the soda being slightly in excess, is brought from Thibet and Persia, where it is found in a native state. This native or crude borax is called *tincal*, *tincor*, *borech*, *pounza*, in the East Indies, and was formerly purified in Europe by the Venetians, when it was called refined or Venetian borax; but it is now prepared by the Dutch by solution in hot water, filtration, and careful crystallization. The particular process is not known. Its taste is cool: it is soluble in eighteen parts of cold and six of hot water. It is decomposed by several of the acids. Borax is rarely used internally in modern practice; and according to Murray it does not appear to possess any activity, although it is supposed by some to be, in doses of half a drachm or two scruples, diuretic and emenagogue. It is occasionally given in cardialgia as an antacid. Its solution is in common use as a cooling gargle, and to detach mucus, &c. from the mouth in putrid fever; and mixed with an equal part of sugar, it is used in the form of powder to remove the aphthous

rust from the tongue in children. The salts formed by the union of the acid of borax with different bases are called borates.

BORBORYGMUS. (From *βορβορυγμῶς*, to make a noise.) The rumbling noise occasioned by flatus in the intestines. It frequently precedes hysterical affections.

BOROZAIL. (Æthiop.) An epidemic disease of the Æthiopians, in appearance similar to the lues venerea.

BORRAGO. See *Borago*.

BORRI. (Indian.) *Borri-borri*. *Boberri*. The Indian name for turmeric; also an ointment used there, in which the roots of turmeric are a chief ingredient.

BOSA. An Egyptian word for a mass which is made of the meal of darnel, hempseed, and water. It is leavening.

BOSMOROS. (From *βοσμος*, to eat, and *μορος*, a part: because it is divided for food by the mill.) *Bosporas*. A species of meal.

BOTALE FORAMEN. A name formerly applied to the foramen ovale.

BOTANY. (*Botanica*, *βοτάνη*: from *βοτάνη*, an herb or grass.) That part of natural history which considers every thing respecting the natural history of vegetables.

BOTANICON. (From *βοτάνη*, an herb.) A plaster made of herbs, and described by Paulus Ægineta.

BOTHOR. (Arab.) Tumours: pimples in the face: also the small-pox or measles.

BOTHRION. (From *βοθρῖον*, a little pit.) *Botrium*. The alveolus or socket for the tooth: also an ulcerated cornea.

BOTIA. A name given to scrophula.

BOTIN. A name for turpentine.

BOTIUM. *Bocium*. Indurated bronchial glands.

BOTOTHINUM. The most evident symptom of disease.

BOTHRITIS. (From *βοθρῖον*, a bunch of grapes.) *Botrytes*. A sort of burnt cadmia, collected in the top of the furnace, and resembling a bunch of grapes.

BOTRIS. (*Βότρυς*, a cluster of grapes; so called because its seeds hang down like a bunch of grapes.) The oak of Jerusalem.

BOTRYS MEXICANA. *Botrys ambrosioides Mexicana*. *Chenopodium Mexicanum*. *Botrys Americana*. Mexican tea. Spanish tea and Artemisia botrys. A decoction of this plant, *Chenopodium ambrosioides* of Linnæus:—*foliis lanceolatis dentatis, racemis foliatis simplicibus*, is recommended in paralytic cases. Formerly the infusion was drunk instead of Chinese tea.

BOTRYS VULGARIS. *Botrys*. *Ambrosia*. *Artemisia chenopodium*. *Atriplex odorata*. *Atriplex sarcocolla*. Jerusalem oak. This plant, *Chenopodium botrys* of Linnæus:—*foliis oblongis sinuatis, racemis nudis multifidis*, was formerly administered in form of decoction in some diseases of the chest; as

humoral asthma, coughs, and catarrhs. It is now fallen into disuse.

BOTUS. *Botia*. *Botus barbatus*. A cucurbit of the chymists.

BOUBALIOS. See *Cucumis agrestis*, and *Pudendum muliebre*.

BOUBON. See *Bubo*.

BOUGIE. (French for wax candle.) *Candela cerea*. *Candela medicata*. *Catheters* of Swediaur. *Cerei medicati* of Le Dran. *Cereolus chirurgorum*. A term applied by surgeons to a long, slender instrument, that is introduced through the urethra into the bladder. Bougies made of the elastic gum are preferable to those made of wax. The caustic bougie differs from the ordinary one in having a thin roll of caustic in its middle, which destroys the stricture, or any part it comes in contact with. Those made of catgut are very seldom used, but are deserving of the attention of the surgeon. Bougies are chiefly used to overcome strictures in the urethra, and the introduction of them requires a good deal of address and caution. They should not be kept in the urethra so long at one time as to excite much pain or irritation. Before their use is discontinued, they should, if practicable, be carried the length of the bladder, in order to ascertain the extent of the strictures, taking care that this be performed not at once, but in a gradual manner, and after repeated trials; for much injury might arise from any hasty or violent efforts to remove the resistance that may present itself. There are bougies also for the œsophagus and rectum.

BOULIMUS. (From *βού*, greatly, and *λιμος*, hunger; or from *βουλομαι*, to desire.) A canine or voracious appetite.

BOVILLÆ. (From *bos*, an ox, because cattle were supposed subject to it.) The measles.

BOVINA FAMES. The same as bulimia.

BOVISTA *Crepitus lupi*. Puff ball. This is the *Lycoperdon bovista*;—*subrotundum, lacerato dehiscens*, of Linnæus, which when dry contains a powder used by the common people to stop the blood in recent cuts, &c.

Bax-tree. See *Buzus*.

BRACHERIUM. (From *brachiale*, a bracelet.) A truss or bandage for hernia; a term used by the barbarous Latin writers.

BRACHIEUS MUSCULUS. See *Brachialis*.

BRACHIEUS EXTERNUS. See *Brachialis externus*.

BRACHIEUS INTERNUS. See *Brachialis internus*.

BRACHIAL ARTERY. *Arteria brachialis*. The brachial artery is the continuation of the axillary artery, which, as it passes behind the tendon of the pectoralis major, receives the name of brachial. It runs down on the inside of the arm, over the musculus coraco-brachialis, and anconæus internus, and, along the inner

edge of the biceps, behind the vena basilica, giving out small branches as it goes along. Below the bend of the arm it divides into the cubitalis and radialis. Sometimes, though rarely, the *brachial artery* is divided from its origin into two large branches, which run down on the arm, and afterwards on the fore-arm, where they are called *cubitalis* and *radialis*.

BRACHIALE. The word means a bracelet: but the ancient anatomical writers apply this term to the carpus, the part on which the bracelet was worn.

BRACHIALIS. See *Brachialis internus*.

BRACHIALIS EXTERNUS. See *Triceps extensor cubiti*.

BRACHIALIS INTERNUS. *Brachieus* of Winslow. *Brachieus internus* of Cower, and *Humero-cubital* of Dumas. A muscle of the fore-arm, situated on the fore-part of the os humeri. It arises fleshy from the middle of the os humeri, at each side of the insertion of the deltoid muscle, covering all the inferior and fore-part of this bone, runs over the joint, and adheres firmly to the ligament; is inserted, by a strong short tendon, into the coronoid process of the ulna. Its use is to bend the fore-arm, and to prevent the capsular ligament of the joint from being pinched.

BRACHIO-CUBITAL LIGAMENT. *Ligamentum brachio cubitale.* The expansion of the lateral ligament, which is fixed in the inner condyle of the os humeri, runs over the capsular, to which it closely adheres, and is inserted like radii on the side of the great sigmoid cavity of the ulna; it is covered on the inside by several tendons, which adhere closely to it, and seem to strengthen it very considerably.

BRACHIO-RADIAL LIGAMENT. *Ligamentum brachio-radiale.* The expansion of the lateral ligament, which runs over the external condyle of the os humeri, is inserted round the coronary ligament, from thence all the way down to the neck of the radius, and also in the neighbouring parts of the ulna. Through all this passage it covers the capsular ligament, and is covered by several tendons adhering closely to both.

BRACHI OS. See *Humeri os*.

BRACHIUM. (*Brachion*, the arm.) The arm, from the shoulder to the wrist.

BRACHIUM MOVENS QUARTUM. See *Latisimus dorsi*.

BRACHUNA. According to Avicenna, a species of furor uterinus.

BRACHY-CHRONIUS. (From *Brachys*, short, and *chronos*, time.) A disease which continues but a short time.

BRACHYPNŒA. (From *Brachys*, short, and *pnœa*, to breathe.) Shortness and difficulty of breathing.

BRACHYS. (From *Brachys*, short.) A muscle of the scapula.

BRACHIUM. Copper. Verdigris.

BRADYPSYIA. (From *Bradys*, slow, and *psysia*, to concoct.) Weak digestion. See *Dyspepsia*.

BRASSAT. A name formerly applied to a ptisan of honey and water.

Brain. See *Cerebrum*.

Brain, little. See *Cerebellum*.

BRAN. *Furfur.* The husks or shells of wheat, which remain in the bolting machine. It contains a portion of the farinaceous matter and is said to have a laxative quality. Decoctions of bran, sweetened with sugar, are used by the common people, and sometimes with success, against coughs, hoarsenesses, &c.

BRANCA. (*Branca*, Span. a foot, or branch.) A term applied to some herba, which are supposed to resemble a particular foot; as *branca leonia*, lion's foot; *branca ursina*, bear's foot.

BRANCA LEONINA. See *Alchemilla*.

BRANCA URSINA. The plant which is directed by this name in foreign pharmacopœias, is the *Heracleum spondylium* of Linnæus:—*foliis pinnatifidis, levibus; floribus uniformibus*. In Siberia it grows extremely high, and appears to have virtues in the cure of dysentery, which the plants of this country do not possess. See also *Acanthus*.

BRANCA LEONIS. See *Alchemilla*.

BRANCHÆ. (From *Brancha*, to make moist.) *Branchi.* Swelled tonsils, or glandulous tumours, of the fauces, which secrete saliva.

BRANCHUS. (From *Brancha*, to moisten.) A defluxion of humours from the fauces.

BRANDY. *Spiritus Gallicus.* A colourless, slightly opaque, and milky fluid, of a hot and penetrating taste, and a strong and agreeable smell, obtained by distilling from wine. It consists of water, ardent spirit, and a small portion of oil, which renders it milky at first, and, after a certain time, colours it yellow. It is the fluid from which rectified or ardent spirit is obtained. Its peculiar flavour depends on the nature of the volatile principles, or essential oil, which come over along with it in the distillation, and likewise, in some measure upon the management of the fire, the wood of the cask in which it is kept, &c. It is said, that our rectifiers imitate the flavour of brandy, by adding a small proportion of nitrous ether to the spirit of malt, or molasses. The utility of brandy is very considerable, but, from its pleasant taste and exhilarating property, it is too often taken to excess. It gives energy to the animal functions; is a powerful tonic, cordial, and antispasmodic; and its utility with camphire, in gangrenous affections, is very great.

Branks. The name, in Scotland, for the mumps. See *Cynanche parotidea*.

Brankurine. See *Acanthus*.

BRASILIA. Brazil wood.

BRASILIENSE LIGNUM. See *Hama-*
zanthum.

BRASILIENSIS RADIX. The ipecacuan-
ha root is sometimes so called. See *Ipe-*
cacuanha.

BRASIUM. (From *βρασσα*, to boil.) Malt,
or germinated barley.

BRASMA. (From *βρασσα*, to boil.) The
unripe black pepper. Fermentation.

BRASMS. The same.

BRASS. *Æs*. A combination of copper
and zinc.

BRASSADELLA. *Brassatella*. Ophioglos-
sum, or the herb adder's tongue.

BRASSICA. (Varro says, *quasi præ-*
cisa: from *præscio*, to cut off; because it
is cut from the stalk for use; or from *αγρι-*
ον, a bed in a garden where they are
cultivated.) Crambe. Cabbage. Cole-
wort.

The name of a genus of plants in the
Linnean system. See *Brassica capi-*
tata.

BRASSICA ALBA. The white cabbage.

BRASSICA APIANA. Jagged or crimped
colewort.

BRASSICA CANINA. The mercurialis
sylvestris.

BRASSICA CAPITATA. Cabbage. There
are several varieties of cabbage, all of
which are generally hard of digestion, pro-
ducing flatulencies, and afford very little
nourishment. These inconveniences are
not experienced by those whose stomachs
are strong and accustomed to them. Few
vegetables run into a state of putrefaction
so quickly as cabbages; they ought, there-
fore, always to be used immediately after
cutting. In Holland and Germany there
is a method of preserving them, by cutting
them into pieces, and sprinkling salt and
some aromatic herbs among them: this
mass is put into a tub, where it is pressed
close, and left to ferment, when it is called
sour crout, or *sauer kraut*. These, and all
pickles of cabbage, are considered as
wholesome and antiscorbutic, from the
vinegar and spices they contain.

BRASSICA CONGYLODES. Turnip cab-
bage.

BRASSICA CUMANIA. Red colewort.

BRASSICA ERUCA. The systematic name
for the plant which affords the semen erucæ.
See *Eruca*.

BRASSICA ERUCASTRUM. See *Eruca*
sylvestris.

BRASSICA FLORIDA. Cauliflower. A va-
riety of the oleracea.

BRASSICA GONTICODES. Turnip cab-
bage.

BRASSICA LACUTURRIA. *Brassica lacu-*
terris. The savoy plant.

BRASSICA MARINA, *Κραμβή θαλασσια*,
Convolvulus maritimus. *Soldanella*. *Sol-*

danella. This plant, *Convolvulus soldan-*
ella of Linnæus:—*foliis reniformibus, pecten-*
catis unifloris, is a native of our coasts.
The leaves are said to be a drastic purge.
It is only used by the common people,
the pharmacopœias having now substituted
more safe and valuable remedies in its
place.

BRASSICA NAPUS. The systematic name
for the plant from which the *semen napi* is
obtained. See *Napi*.

BRASSICA OLERACEA. The systematic
name for the *brassica capitata* of the shops.
See *Brassica capitata*.

BRASSICA RAPA. The systematic name
for the plant whose root is called turnip.
See *Rapa*.

BRASSICA RUBRA. Red cabbage. Mr.
Watt finds that the red cabbage affords
a very excellent test both for acids and al-
kalis, in which it is super. or to litmus, be-
ing naturally blue, turning green with
alkalis, and red with acids.

BRASSICA SATIVA. The common gar-
den cabbage.

BRASSICA SABAUDA. The savoy plant.

BRASSIDELLA AHS. A way of curing
wounds, mentioned by Paracelsus, by ap-
plying the herb *Brassidella* to them.

BRATHU. (*Βραθυ*.) An old name for sa-
vine.

BREAD-FRUIT. The tree which af-
fords this, grows in all the Ladrone Islands
in the South Sea, in Otaheite, and now in
the West Indies. The bread-fruit grows up-
on a tree the size of a middling oak. The
fruit is about the size of a child's head, and
the surface is reticulated, not much unlike
the surface of a truffle. It is covered with
a thin skin, and has a core about the size
of a small knife. The eatable part is be-
tween the skin and the core: it is as white
as snow, and somewhat of the consistence
of new bread. It must be toasted before
it is eaten, being first divided into three or
four parts. Its taste is insipid, with a slight
sweetness, nearly like that of wheaton
bread and artichoke together. This fruit
is the constant food of the inhabitants all
the year, it being in season eight months.

BREAST. *Mamma*. The two globu-
lar projections, composed of common in-
tegments, adipose substance, and lacteal
glands and vessels, and adhering to the an-
terior and lateral regions of the thorax of
females. On the middle of each breast is
a projecting portion, termed the *papilla* or
nipple, in which the excretory ducts of the
glands terminate, and around which is a
coloured orb, or disc, called the *areola*.
The use of the breasts is to suckle new-born
infants.

Breast-bone. See *Sternum*.

BREMA. (From *βρεχω*, to moisten;
formerly so called because, in infants, and
sometimes even in adults, they are tender

and moist.) An old name for the parietal bones.

BREVIÀ. (From *brevis*, short.) A specific name of some parts whose termination is not far from their insertion, as *brevis vasa*, the branches of the splenic vein.

BREVIS MUSCULUS. A muscle of the scapula.

BREVIS CUBITI. A muscle of the forearm.

Brevis extensor pollicis pedis. See *Extensor brevis pollicis pedis*.

Brevis flexor pollicis pedis. See *Flexor brevis pollicis pedis*.

BREVIS PERONEUS. See *Peroneus brevis*.

Brevis pronator radii. See *Pronator radii brevis*.

BREYNIA. (An American plant named in honour of Dr. Brennius.) A species of capparid.

Briar, wild. See *Rosa canina*.

BRICUMUM. A name which the Gauls gave to the herb artemisia.

Brimstone. See *Sulphur*.

BRISTOL HOT-WELL. *Bristolensis aqua.* A pure, thermal or warm, slightly acidulated, mineral spring, situated about a mile below Bristol. The fresh water is inodorous, perfectly limpid, and sparkling, and sends forth numerous air bubbles when poured into a glass. It is very agreeable to the palate, but without having any very decided taste, at least none that can be distinguished by a common observer. Its specific gravity is only 1.00077, which approaches so near to that of distilled water, that this circumstance alone would shew that it contained but a very small admixture of foreign ingredients. The temperature of these waters, taking the average of the most accurate observations, may be reckoned at 74 deg.; and this does not very sensibly vary during winter or summer. Bristol water contains both solid and gaseous matter, and the distinction between the two requires to be attended to, as it is owing to the very small quantity of solid matter that it deserves the character of a very fine natural spring; and to an excess in gaseous contents, that it seems to be principally indebted for its medical properties, whatever they may be, independent of those of mere water, with an increase of temperature. From the different investigations of chymists, it appears that the principal component parts of the Hotwell water, are a large proportion of carbonic acid gas, or fixed air, and a certain portion of magnesia and lime, in various combinations, with the muriatic, vitriolic, and carbonic acids. The general inference is, that it is considerably pure for a natural fountain, as it contains no other solid matter than is found in almost all common spring water, and in less quantity.

On account of these ingredients, especially the carbonic acid gas, the Hotwell water is efficacious in promoting salutary hæmorrhages, in green sickness, as well as in the blind hæmorrhoides. It may be taken with advantage in obstructions, and weakness of the bowels, arising from habitual costiveness; and, from the purity of its aqueous part, it has justly been considered as a specific in diabetes, rendering the urinary organs more fitted to receive benefit from those medicines which are generally prescribed, and sometimes successful.

But the high reputation which this spring has acquired, is chiefly in the cure of pulmonary consumption. From the number of unsuccessful cases among those who frequent this place, many have denied any peculiar efficacy in this spring, superior to that of common water. It is not easy to determine how much may be owing to the favourable situation and mild temperate climate which Bristol enjoys; but it cannot be doubted that the Hotwell water, though by no means a cure for consumption, alleviates some of the most harassing symptoms of this formidable disease. It is particularly efficacious in moderating the thirst, the dry burning heat of the hands and feet, the partial night sweats, and the symptoms that are peculiarly hectic; and thus in the earlier stages of phthisis, it may materially contribute to a complete re-establishment of health; and even in the latter periods, mitigate the disease when the cure is doubtful, if not hopeless.

The sensible effects of this water, when drank warm and fresh from the spring, are a gentle glow of the stomach, succeeded sometimes by a slight and transient degree of headache and giddiness. By a continued use, in most cases it is diuretic, keeps the skin moist and perspirable, and improves the appetite and health. Its effects on the bowels are variable. On the whole, a tendency to costiveness seems to be the more general consequence of a long course of this medicinal spring, and therefore the use of a mild aperient is requisite. These effects, however, are applicable only to invalids, for healthy persons, who taste the water at the fountain, seldom discover any thing in it but a degree of warmth, which distinguishes it from the common element.

The season for the Hotwell is generally from the middle of May to October; but as the medicinal properties of the water continue the same throughout the year, the summer months are preferred merely on account of the concomitant benefits of air and exercise.

It should be mentioned, that another spring, nearly resembling the Hotwell, has

been discovered at Clifton, which is situated on the summit of the same hill, from the bottom of which the Hotwell issues. The water of Sion spring, as it is called, is one or two degrees colder than the Hotwell; but in other respects it sufficiently resembles it to be employed for all similar purposes.

BRITANNICA HERBA. See *Hydrolapathum*.

BRITISH OIL. A variety of the black species of petroleum, to which this name has been given as an empirical remedy.

BROCCOLI. *Brassica Italica*. As an article of diet, this may be considered as more delicious than cauliflower and cabbage. Sound stomachs digest broccoli without any inconvenience; but in dyspeptic stomachs, even when combined with pepper, &c. it always produces flatulency, and nauseous eructations.

BROCHOS. (*Βροχος*, a snare.) A bandage.

BROCHTHUS. (From *βροχθ*, to pour.) The throat; also a small kind of drinking-vessel.

BROCHUS. (*Βροχος*.) One with a prominent upper-lip, or one with a full mouth and prominent teeth.

BRODIUM. A term in pharmacy, signifying the same with *jusculum*, broth, or the liquor in which any thing is boiled. Thus we sometimes read of *Broodium Salis*, or a decoction of salt.

BROMA. (From *βρωμα*, to eat.) Food of any kind that is masticated, and not drank.

BROMA-THEON. (From *βρωμα*, to eat.) Mushrooms.

BROMATOLOGY. (*Bromatologia*: from *βρωμα*, food, and *λογος*, a discourse.) A discourse or treatise on food.

BROMELIA ANANAS. The systematic name of the plant which affords the *ananas*. See *Ananas*.

BROMELIA KARATAS. The systematic name of the plant from which we obtain the fruit called penguin, which is given in the Spanish West Indies to cool and quench thirst in fevers, dysenteries, &c. It grows in a cluster, there being several of the size of one's finger together. Each portion is clothed with a husk, containing a white pulpy substance, which is the eatable part; and if it be not perfectly ripe, its flavour resembles that of the pine-apple. The juice of the ripe fruit is very austere, and is made use of to acidulate punch. The inhabitants of the West Indies make a wine of the penguin, which is very intoxicating, and has a good flavour.

BROMION. (From *βρωμος*, the oat.) The name of a plaster, made with oat flour, mentioned by Paulus Aegineta.

BROMUS STERILIS. (From *βρωμα*, to eat.) The wild oat.

BRONCHIA. (From *βροχθ*, the throat.) See *Trachea*.

BRONCHIALES ARTERIE. Bronchial arteries. Branches of the aorta given off in the chest.

BRONCHIALES GLANDULE. Bronchial glands. Large blackish glands, situated about the bronchia and trachea, which secrete blackish mucus.

BRONCHOCELE. (From *βροχθ*, the windpipe, and *αλλω*, a tumour.) *Botium*. *Hernia gutturis* *Guttur tumidum*. *Tracheolophyena*. *Gossum*. *Excechebronchos*. *Gongrona*. *Botium*. *Hernia bronchialis*. *Tracheocele*. Derbyshire neck. This disease is marked by a tumour on the fore-part of the neck, and seated between the trachea and skin. In general it has been supposed principally to occupy the thyroid gland. We are given to understand that it is a very common disorder in Derbyshire; but its occurrence is by no means frequent in other parts of Great Britain, or in Ireland. Amongst the inhabitants of the Alps, and other mountainous countries bordering thereon, it is a disease very often met with, and is there known by the name of goitre. The cause which gives rise to it, is by no means certain, and the observations of different writers are of very little practical utility. Dr. Saunders controverts the general idea of the bronchocele being produced by the use of snow water. The swelling is at first without pain, or any evident fluctuation, when the disease is of long standing, and the swelling considerable, we find it in general a very difficult matter to effect a cure by medicine, or any external application; and it might be unsafe to attempt its removal with a knife, on account of the enlarged state of its arteries, and its vicinity to the carotids; but, in an early stage of the disease, by the aid of medicine a cure may be effected.

Although some relief has been obtained at times, and the disease probably somewhat retarded by external applications, such as blisters, discutient embrocations, and saponaceous and mercurial plaisters, still a complete cure has seldom been effected without an internal use of medicine; and that which has always proved the most efficacious, is burnt sponge. The form under which this is most usually exhibited, is that of a lozenge. *℞. spongiæ ustæ ℥ss. mucilag. Arab. gum. q. s. fiat trochiscus*. When the tumor appears about the age of puberty, and before its structure has been too morbidly deranged, a pill, consisting of a grain or two of calomel, must be given for three successive nights; and, on the fourth morning, a saline purge. Every night afterwards, for three weeks, one of the troches should, when the patient is in bed, be put under the tongue, suffered to dissolve gradually, and the solution swallowed. The disgust at first arising from this remedy soon wears off. The pills and the purge are to be repeated at the end of three

weeks, and the troches had recourse to as before; and this plan is to be pursued till the tumour is entirely dispersed. Some recommend the burnt sponge to be administered in larger doses. Sulphurated potash dissolved in water, in the proportion of 30 grains to a quart daily, is a remedy which has been employed by Dr. Richter with success, in some cases, where calcained sponge failed. The natron preparatum being the basis of burnt sponge, is now frequently employed instead of it, and, indeed, it is a more active medicine.

BRONCHOTOMY. (*Bronchotomia*.—from *βρογχος*, the wind-pipe, and *τομή*, to cut.) Tracheotomy. Laryngotomy. This is an operation in which an opening is made into the larynx, or trachea, either for the purpose of making a passage for the air into and out of the lungs, when any disease prevents the patient from breathing through the mouth and nostrils, or of extracting foreign bodies, which have accidentally fallen into the trachea; or, lastly, in order to be able to inflate the lungs, in cases of sudden suffocation, drowning, &c. Its practicableness, and little danger, are founded on the facility with which certain wounds of the wind-pipe, even of the most complicated kind, have been healed, without leaving any ill effects whatever, and on the nature of the parts cut, which are not furnished with any vessel of consequence.

BRONCHOS. (*βρογχος*, the wind-pipe.) A catarrh; a suppression of the voice from a catarrh.

BRONCHUS. (From *βρω*, to pour.) The wind-pipe. The ancients believed that the solids were conveyed into the stomach by the œsophagus, and the fluids by the bronchia; whence its name.

Brooklime Speedwell. See *Beccabunga*.

Broom, Common. See *Genista*.

BRUCEA. (So named by Sir Joseph Banks in honour of Mr. Bruce, the traveller in Abyssinia, who first brought the seeds thence into England.) The name of a genus of plants in the Linnæan system.

BRUCEA ANTIDYSSENTERICA. The systematic name of the plant from which it was supposed we obtained the angustura bark. See *Angustura cortex*.

BRUCEA FERRUGINEA. This plant is also supposed to afford the angustura bark. See *Angustura cortex*.

Bruscewort. See *Saponaria*.

BRUNELLA. See *Prunella*.

BRUNNER'S GLANDS. *Brunneri glandule.* Peyer's glands. The muciparous glands, situated between the villous and cellular coat of the intestinal canal; so named after Brunner, who discovered them.

BRUVUS. An erysipelatous eruption.

BRUVUS. See *Ruscus*.

BRUTA. (Arab.) Instinct. Savine.

BRUTIA. An epithet for the most resinous kind of pitch, therefore used to make the *Olum Picinum*. The *Pix Brutia* was so called from Brutia, a country in the extreme parts of Italy, where it was produced.

BRUTINO. Turpentine.

BRUTOXON. The name of an ointment used by the Greeks.

BRUTUA. See *Pareira brava*.

BRUXANELI. (Indian.) A tall tree in Malabar, whose bark is diuretic, according to Ray.

BRUXANUS. (From *βρυξ*, to make a noise.) A peculiar kind of noise, such as is made by gnashing or grating the teeth; or, according to some, a certain kind of convulsion affecting the lower jaw, and striking the teeth together, most frequently observed in such children as have worms.

BRYONIA. (From *βρω* to abound, from its abundance.) Bryony.

1. The name of a genus of plants in the Linnæan system. Class, *Diœcia*. Order, *Syngœnia*.

2. The pharmacopœial name of the white bryony. *Vitis alba sylvestris*. *Agrostis*. *Ampelæ*. *Archeustris*. *Echetrosis* of Hippocrates. *Bryonia aspera*. *Cedrostris*. *Chelidonium*. *Labrusca*. *Melastrium*. *Ophrostaphylon*. *Psilothrum*. *Bryonia alba* of Linnæus:—*foliis palmatis utrinque callosa scabris*.

White bryony, or wild vine, is a very common plant in woods and hedges. The root has a very nauseous biting taste, and disagreeable smell. Bergius states the virtues of this root to be purgative, hydragogue, emmenagogue, and diuretic; the fresh root emetic. This powerful and irritating cathartic, though now seldom prescribed by physicians, is said to be of great efficacy in evacuating serous humours, and has been chiefly employed in hydropical cases. Instances of its good effects in other chronic diseases are also mentioned, as asthma, mania, and epilepsy. In small doses, it is reported to operate as a diuretic, and to be resolvent and deobstruent. In powder, from ℥j. to a drachm, it proves strongly purgative; and the juice, which issues spontaneously, in doses of a spoonful or more, has similar effects, but is more gentle in its operation. An extract prepared by water acts more mildly, and with greater safety than the root in substance, given from half a drachm to a drachm. It is said to prove a gentle purgative, and likewise to operate powerfully by urine. Of the expressed juice, a spoonful acts violently both upwards and downwards; but cream of tartar is said to take off its virulence. Externally, the fresh root has been employed in cataplasms, as a resolvent and discutient; also in ischiadic and other rheumatic affections.

BRYONIA ALBA. The systematic name of the white bryony plant. See *Bryonia*.

BRYONIA MEXICANICA FLORIDANS. A name given to the jalap root.

BRYONIA NIGRA Black bryony, or vine.

The *Tamus communis* of Linnæus.

BRYONIA PERUVIANA. Jalap.

Bryony, black. See *Bryonia nigra*.

Bryony, white. See *Bryonia*.

BRETHION. (*Brethun.*) A malagma so called, and described by Paulus Ægineta.

BRETON. (From *breu*, to pour out.) A kind of ale, or wine, made of barley.

BUBASTECORDIUM. (From *bubastus* and *cor*, the heart.) A name formerly given to artemisia, or mugwort.

BUBO. (From *bucor*, the groin; because they most frequently happen in that part.) Modern surgeons mean, by this term, a swelling of the lymphatic glands, particularly of those of the groin and axilla. The disease may arise from the mere irritation of some local disorder, when it is called *sympathetic bubo*; from the absorption of some irritating matter, such as the venereal poison; or from constitutional causes, as in the pestilential bubo, and scrophulous swellings, of the inguinal and axillary glands.

BUBON. (From *bucor*, the groin, or a tumour to which that part is liable, and which it was supposed to cure.) The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*.

BUBON GALBANUM. The systematic name of the plant which affords the official galbanum. See *Galbanum*.

BUBON MACEDONICUM. The systematic name of the plant which affords the *semen petroselinæ Macedonici* of the shops. See *Petroselinum Macedonicum*.

BUBONIUM. (From *bucor*, the groin.) A name of the golden starwort; so called because it was supposed to be efficacious in diseases of the groin.

BUBONOCELE. (From *bucor*, the groin, and *anas*, a tumour.) *Hernia inguinalis*. Inguinal hernia, or rupture of the groin. A species of hernia, in which the bowels protrude, at the abdominal ring. See *Hernia*.

BUCCA. (Heb.) The cheek. The hollow inner part of the cheek, that is inflated by the act of blowing.

BUCCACRATON. (From *bucca*, or *bucella*, that is, a morsel of bread soaked in wine, which served in old times for a breakfast.) Paracelsus calls by the name of *Buccella*, the carneous excrescence of the polypus in the nose, because he supposes it to be a portion of flesh parting from the bucca, and insinuating itself into the nose.

BUCCAL GLANDS. (*Glandule buccinæ*: from *bucca*, the cheek.) The small glands of the mouth, under the cheek, which assist in secreting saliva into that cavity.

BUCEA. (From *bucca*, the cheek; as much as can be obtained at one time with

in the cheeks.) A mouthful; a morsel; a polypus of the nose.

BUCCELLATON. (From *buccella*, a morsel.) A purging medicine, made up in the form of a loaf; consisting of scammony, &c. put into fermented flour, and then baked in an oven.

BUCELLA. See *Buccæa*.

BUCCELLATIO. (From *buccellatus*, cut into small pieces.) *Bucellatio*. A method of stopping an hæmorrhage, by applying small pieces of lint to the vein, or artery.

BUCGINATOR (*Musculus buccinator*. So named from its use in forcing the breath to sound the trumpet; from *bucinator*, a trumpet.) *Retractor anguli oris* of Albinus, and *alveolo maxillare* of Dumas. The trumpeter's muscle. The buccinator was long thought to be a muscle of the lower jaw, arising from the upper alveoli, and inserted into the lower alveoli, to pull the jaw upward; but its origin and insertion, and the direction of its fibres, are quite the reverse of this. For this large flat muscle, which forms in a manner the walls of the cheek, arises chiefly from the coronoid process of the lower jaw-bone, and partly also from the end of the alveoli, or socket process of the upper-jaw, close by the pterygoid process of the sphenoid bone: it goes forward with direct fibres, to be implanted into the corner of the mouth; it is thin and flat, covers in the mouth, and forms the walls of the cheek, and is perforated in the middle of the cheek by the duct of the parotid gland. These are its principal uses:—it flattens the cheek, and so assists in swallowing liquids;—it turns, or helps to turn, the morsel in the mouth, while chewing, and prevents it from getting without the line of the teeth;—in blowing wind-instruments, it both receives and expels the wind;—it dilates like a bag, so as to receive the wind in the cheeks; and it contracts upon the wind, so as to expel the wind, and to swell the nose. In blowing the strong wind-instruments, we cannot blow from the lungs, for it distresses the breathing, we reserve the air in the mouth, which we keep continually full; and from this circumstance, as mentioned above, it is named buccinator, from blowing the trumpet.

BUCULA. (Dim. of *bucca*, the cheek.) The fleshy part under the chin.

BUCFHALON, RED-FRUITED. The plant so called, is the *Trophis Americanus* of Linnæus. Its fruit is a kind of rough red berry, which is eaten by the inhabitants of Jamaica, although its flavour is by no means pleasant.

BUCERAS. (From *buc*, an ox, and *eras*, a horn; so called from the horn-like appearance of its seed.) *Buceros*. Fenugreek seed. See *Fenugracum*.

Buck-bean. See *Trifolium paudoum*.

Buck-thorn. See *Spina cervicæ*.

BUCKWHEAT. The *Polygonum fagopyrum* of Linnæus. The grain of this plant constitutes the principal food of the inhabitants of Russia, Germany, and Switzerland.

BUCK-WHEAT, EASTERN. The *Polygonum divaricatum* of Linnæus. The roots, reduced into a coarse meal, are the ordinary food of the Siberians.

BUCRANION. (From *βuc*, an ox, and *κρανιον*, the head; so called from its supposed resemblance to a calf's snout.) The antirrhinum, or snap-dragon plant.

BUCRON. The hymen, according to Piræus.

BUGANTIA. Chilblains.

Bugle. See *Primella*.

Bugloss. See *Buglossum*.

BUGLOSSUM. (From *βuc*, an ox, and *γλωσσα*, a tongue; so called from the shape and roughness of its leaf.) *Buglossa*. *Buglossum angustifolium majus*. *Buglossum vulgare majus*. *Buglossum sativum*. Official bugloss, or alkanet. This plant, *Anchusa officinalis* of Linnæus:—*foliis lanceolatis strigosis, spicis secundis imbricatis, calycibus quinquepartitis*, was formerly esteemed as a cordial in melancholic and hypochondriacal diseases. It is seldom used in modern practice, and then only as an aperient and refrigerant.

BUGLOSSUM SYLVESTRIS. The stone bugloss.

BUGULA. (A dim. of *buglossa*.) See *Consolida media*.

BULBOCASTANUM. (From *βολβοc*, a bulb, and *καστανον*, a chestnut; so called from its bulbous appearance.) *Agriocastanum*. *Nucula terrestris*. *Bulbocastaneum*. *Bulbocastanum majus et minus*. Earth-nut. Hawk-nut. Kipper-nut, and pig-nut. This plant, the *Bunium bulbocastanum* of Linnæus, has a root as large as a nutmeg; hard, tuberous and whitish; is eaten raw, or roasted. It is sweetish to the taste, nourishing, and supposed to be of use against stranguary and bloody urine.

BULBOCAVERNOSUS. (*Bulbocavernosus*, sc. *musculus*: so called from its origin and insertion) See *Accelerator urine*.

BULBONACH. (Germ.) The *Lunaria rediviva* of Linnæus. Satin and honesty. It is said, by Ray, to be a warm diuretic.

BULBUS EXCELENTUS. Such bulbous roots as are commonly eaten are so called.

BULBUS VOMITORIUS. *Muscari*. *Hyacinthus muscari*, of Linnæus. Musk. Grape-flower. Emetic and diuretic, according to Ray.

Bulge-water tree. The *Geoffroya Jamaicensis*.

BULIMIA. (From *βu*, a particle of excess, and *ιμος*, hunger.) *Bulimiasis*, *Boulimos*, *Bulimus*. *Bolimos* of Avicenna. *Fames canina*. *Appetitus caninus*. *Phagedæna*. *Adephagia*. *Bupeina lyncæria*. Insatiable hunger, or canine appetite.

Dr. Cullen places this genus of disease in the class *locales*, and order *dysoresia*; and distinguishes three species. 1. *Bulimia helvorum*; in which there is no other disorder of the stomach, than an excessive craving of food. 2. *Bulimia cyncepalis*; in which there is a frequent desire of food, and the sense of hunger is preceded by swooning. 3. *Bulimia emetica*, also *cyncexia*; in which an extraordinary appetite for food is followed by vomiting. The real causes of this disease are, perhaps, not properly understood. In some cases, it has been supposed to proceed from an acid in the stomach, and in others, from a superabundance of acid in the gastric juice, and from indigested sordes, or worms. In most instances, some consider it as depending more frequently on monstrosity than disease. An extraordinary and well-attested case of this disease, is related, in the third volume of the Medical and Physical Journal, of a French prisoner, who, in one day, consumed of raw cow's udder, 4 lbs. raw beef, 10 lbs. candles, 2 lbs.; total, 16 lbs.; besides 5 bottles of porter.

BULIMIA ADEPHAGIA. A voracious appetite.

BULIMIA CANINA. A voracious appetite, with subsequent vomiting.

BULIMIA CARDIALGICA. A voracious appetite, with heartburn.

BULIMIA CONVULSORUM. A voracious appetite, with convulsions.

BULIMIA EMETICA. A voracious appetite, with vomiting.

BULIMIA HELVORUM. Gluttony.

BULIMIA LUNARIO. Gluttony.

BULIMIA SYNCOPALIS. A voracious appetite, with fainting, from hunger.

BULIMIA VERMINOSA. A voracious appetite, from worms.

BULIMIASIS. See *Bulimia*.

BULIMUS. See *Bulimia*.

BULITHUM. (From *βuc*, an ox, and *λιθος*, a stone.) A bezoar, or stone, found in the kidneys, or gall, or urinary bladder, of an ox, or cow.

BULLA. (A bubble.) A clear vesicle, which arises from burns, or scalds; or other causes.

BULLACE. The fruit so called, is the produce of the *Prunus insitia* of Linnæus, which grows wild in our hedges. There are two varieties of bullace, the red and the white, which are used with the same intentions as the common damsons.

BULLOSA FEBRIS. An epithet applied to the vesicular fever, because the skin is covered with little vesicles, or blisters. See *Pemphigus*.

BUNIUM. (From *βuvoc*, a little hill; so called from the tuberosity of its root.)

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*.

2. The name of the wild parley.

BURSUS VINUM. (From *bunium*, wild parsley.) A wine made of bunium and must.

BURUM. **BULBOCASTANUM.** The systematic name of a plant whose root is called the pig-nut. See *Bulbocastanum*.

BUNIOS. A species of turnip.

BURINA. (From *bu*, a particle of magnitude, and *ina*, hunger.) A voracious appetite.

BUPHAGOS. (From *bu*, a particle of excess, and *φαγεω*, to eat.) The name of an antidote which created a voracious appetite in Marcellus Empiricus.

BUPHTHALNUM. (From *βου*, an ox, and *οφθαλμος*, an eye; so called from its flowers, which are supposed to resemble an eye.) The herb ox-eye daisy. See *Bellis major*.

BUPHTHALNUM CRITICUM. Pellitory of Spain. See *Pyrethrum*.

BUPHTHALNUM GERMANICUM. The common ox-eye daisy.

BUPHTHALNUM MAJUS. Great, or ox-eye daisy. See *Bellis major*.

BUPHTHALMUS. (From *βου*, an ox, and *οφθαλμος*, an eye; so named from its large appearance, like an ox's eye.) Ox-eye.

1. Diseased enlargement of the eye.

2. Houseleek.

BUPLEURUM. (From *βυ*, large, and *πλευρα*, a rib; so named from its having large rib-like filaments upon its leaves.)

1. The name of a genus of plants in the Linnean system.

2. The pharmacopœial name of the *Bupleium Bupleioides*. The herb hare's-car. See *Perfoliata*.

BUPLEURUM ROTUNDIFOLIUM. The systematic name of the plant called perfoliata, in the pharmacopœias. See *Perfoliata*.

Burdock. See *Bardana*.

Burgundy pitch. See *Pix Burgundica*.

BURAC. (Arab.) **BORAX.** It also means any kind of salt.

BURIS. According to Avicenna, a scirrhous hernia, or hard abscess.

BURINA. Pitch.

Burnet saxifrage. See *Pimpinella*.

BURNING. *Brenning.* An ancient medical term, denoting an infectious disease, got in the stews by conversing with lewd women, and supposed to be the same with what we now call the venereal disease.

BURRHU SPIRITUS MATRICARIS. *Burru's spirit*, for disorders of the womb. A compound of myrrh, olibanum, amber, and spirit of wine.

Burst hartshorn. See *Cornu ustum*.

Burst sponge. See *Spongia usta*.

BURSA. A bag

1. The scrotum.

2. A herb called *Thlaspi bursa pastoris*, from the resemblance of its seminal follicles to a triangular purse.

BURSALOGY. (From *bursa*, a bag,

and *λογος*, a discourse.) The doctrine of the *bursæ mucosæ*.

BURSÆ MUCOSÆ. Mucous bags, composed of proper membranes, containing a kind of mucous fat, formed by the exhaling arteries of the internal coat. They are of different sizes and firmness, and are connected by the cellular membrane with articular cavities, tendons, ligaments, or the periosteum. The use of the *bursæ mucosæ* is to secrete, and contain a substance to lubricate tendons, muscles, and bones, in order to render their motion easy.

A Table of all the Bursæ Mucosæ.

In the Head.

1. *A bursa of the superior oblique muscle of the eye, situated behind its trochlea in the orbit.*

2. *The bursa of the digastricus, situated in the internal surface of its tendons.*

3. *A bursa of the circumflexus, or tensor palati, situated between the hook-like process of the sphenoid bone and the tendon of that muscle.*

4. *A bursa of the sterno-hyoidous muscle, situated between the os hyoides and larynx.*

About the Shoulder-joint.

1. *The external acromial, situated under the acromion, between the coracoid process, deltoid muscle, and capsular ligament.*

2. *The internal acromial, situated above the tendon of the infra-spinatus and teres major: it often communicates with the former.*

3. *The coracoid bursa, situated near the root of the coracoid process: it is sometimes double and sometimes triple.*

4. *The claviculæ bursa, found where the clavicle touches the coracoid process.*

5. *The subclavian bursa, between the tendon of the subclavicularis muscle and the first rib.*

6. *The coraco-brachial, placed between the common origin of this muscle and the biceps, and the capsular ligament.*

7. *The bursa of the pectoralis major, situated under the head of the humerus, between the internal surface of the tendon of that muscle and another bursa placed on the long head of the biceps.*

8. *An external bursa of the teres major, under the head of the os humeri, between it and the tendon of the teres major.*

9. *An internal bursa of the teres major, found within the muscle where the fibres of its tendon diverge.*

10. *A bursa of the latissimus dorsi, between the tendon of this muscle and the os humeri.*

11. *The humero-bicipital bursa, in the vagina of the tendon of the biceps.*

There are other *bursæ mucosæ* about the humerus, but their situation is uncertain.

Near the Elbow-joint.

1. *The radio-bicipital, situated between*

the tendon of the biceps, brachialis, and anterior tubercle of the radius.

2. *The cubito-radial*, between the tendon of the biceps, supinator brevis, and the ligament common to the radius and ulna.

3. *The anconeal bursa*, between the olecranon and tendon of the anconeus muscle.

4. *The capitulo-radial bursa*, between the tendon common to the extensor carpi radialis brevis, and extensor communis digitorum and round head of the radius. There are occasionally other bursæ, but as their situation varies, they are omitted.

About the inferior part of the Fore-arm and Hand.

On the inside of the Wrist and Hand.

1. A very large bursa, for the tendon of the flexor pollicis longus.

2. Four short bursæ on the fore-part of the tendons of the flexor sublimis.

3. A large bursa behind the tendon of the flexor pollicis longus, between it and the fore-part of the radius, capsular ligament of the wrist and os trapezium.

4. A large bursa behind the tendons of the flexor digitorum profundus, and on the fore-part of the end of the radius, and fore-part of the capsular ligament of the wrist. In some subjects it communicates with the former.

5. An oblong bursa between the tendon of the flexor carpi radialis and os trapezium.

6. A very small bursa between the tendon of the flexor carpi ulnaris and os pisiforme.

On the back part of the Wrist and Hand.

7. A bursa between the tendon of the abductor pollicis longus and the radius.

8. A large bursa between the two extensores carpi radiales.

9. Another below it, common to the extensores carpi radiales.

10. A bursa, at the insertion of the tendon of the extensor carpi radialis.

11. An oblong bursa, for the tendon of the extensor pollicis longus, and which communicates with 9.

12. A bursa, for the tendon of the extensor pollicis longus, between it and the metacarpal bone of the thumb.

13. A bursa between the tendons of the extensor of the fore, middle, and ring fingers.

14. A bursa for the extensors of the little finger.

15. A bursa between the tendon of the extensor carpi ulnaris and ligament of the wrist.

There are also bursæ mucosæ between the musculi lumbricales and interossei.

Near the Hip-joint.

On the fore-part of the joint.

1. The *ileo puberal*, situated between the iliacus internus, psoas magnus, and the capsular ligament of the head and the femur.

2. The *pectineal*, between the tendon of the pectineus and the thigh bone.

3. A small bursa of the gluteus medius muscle, situated between it and the great trochanter, before the insertion of the pyramiformis.

4. A bursa of the gluteus minimus muscle between its tendon and the great trochanter.

5. The *gluteo-fascial*, between the gluteus maximus and vastus externus.

On the posterior part of the Hip-joint.

6. The *tubero-ischiatic bursa*, situated between the obturator internus muscle, the posterior spine of the ichium, and its tuberosity.

7. The *obturator bursa*, which is oblong, and found between the obturator internus and gemini muscles, and the capsular ligament.

8. A bursa of the semi-membranosus, under its origin and the long head of the biceps femoris.

9. The *gluteo-trochanteral bursa*, situated between the tendon of the psoas muscle and the root of the great trochanter.

10. Two *gluteo-femoral bursæ*, situated between the tendon of the gluteus maximus and os femoris.

11. A bursa of the quadratus femoris, situated between it and the little trochanter.

12. The *iliac bursa*, situated between the tendon of the iliacus internus and the little trochanter.

Near the Knee-joint.

1. The *supra-gemal*, which adheres to the tendons of the vastus and cruralis and the fore-part of the thigh-bone.

2. The *infra gemal bursa*, situated under the ligament of the patella, and often communicating with the above.

3. The *anterior gemal*, placed between the tendon of the sartorius, gracilis and semitendinosus and the internal and lateral ligament of the knee.

4. The *posterior gemal*, which is sometimes double, and is situated between the tendons of the semi-membranosus, the internal head of the gastrocnemius, the capsular ligament, and internal condyle.

5. The *popliteal*, conspicuous between the tendon of that muscle, the external condyle of the femur, the semilunar cartilage, and external condyle of the tibia.

6. The *bursa of the biceps cruris*, between the external part of the tendon, the biceps cruris, and the external lateral ligament of the knee.

In the Foot.

On the back, side, and hind part of the Foot.

1. A bursa of the *tibiæ anticus*, between its tendon, the lower part of the tibia, and capsular ligament of the ankle.

2. A bursa between the tendon of the extensor pollicis pedis longus, the tibia and capsular ligament of the ankle.

3. A bursa of the *extensor digitorum communis*, between its tendons, the tibia, and ligament of the ankle.

4. *A large bursa*, common to the tendons of the peronei muscles.

5. *A bursa of the peroneus brevis*, proper to its tendon.

6. *The calcaneal bursa*, between the tendo Achillis and os calcis.

In the Sole of the Foot.

1. *A bursa for the tendon of the peroneus longus.*

2. *A bursa* common to the tendon of the flexor pollicis pedis longus, and the tendon of the flexor digitorum pedis communis longus profundus.

3. *A bursa of the tibialis posticus*, between its tendon, the tibia, and astragalus.

4. *Five bursae for the flexor tendons*, which begin a little above the first-joint of each toe, and extend to the root of the third phalanx, or insertion of the tendons.

BURSALIS MUSCULUS. (From its resemblance to a bursa, or purse.) See *Obturator externus et internus*.

BUSKINUM. (From *bus*, great, and *skinos*, parsley.) A large species of parsley.

BUSII SPIRITUS, BEZOARDICUS. The bezoardic spirit of Bussius, an eminent physician at Dresden. A distillation of ivory, sal-ammoniac, amber, &c.

Butchersbroom. See *Ruscus*.

BUTIRA. A synonym for gutta rosacea.

BUTIRO. Turpentine.

BUTIRON. See *Iris palustris*.

BUTTER. (*Butyrum*: from *bus*, a cow, and *tuos, conguum*, or cream.) A concrete and soft substance, of a yellow colour, approaching more or less to that of gold, and of a mild, agreeable taste. It melts by a gentle heat, and becomes solid by cooling. Fresh butter is nourishing, and relaxing, but it readily becomes sour, and, in general, agrees with few stomachs. Rancid butter is one of the most unwholesome and indigestible of all food.

Butter-bar. See *Petasites*.

Butter-flower. See *Ranunculus*.

BUTTER-MILK. The thin and sour milk which is separated from the cream by churning it into butter.

Butterwort. See *Pinguicula*.

BUTUA. See *Pariera brava*.

BUTYRUM. See *Butter*.

BUTYRUM ANTIMONII. See *Murias antimoni*.

BUXTON WATERS. *Buxtonienses aque.* Warm mineral springs, which rise in the village of Buxton, in Derbyshire. They have been long celebrated for their medicinal properties. With respect to sensible properties, the Buxton-water cannot be distinguished from common spring water, when heated to the same temperature. Its temperature in the gentleman's bath, is invariably 82°. The principal peculiarity in the appearance of this spring, is a large quantity of elastic vapour, that rises, and forms bubbles, which pass through

the water, and break as soon as they reach the surface. The air of these bubbles was ascertained, by Dr. Pearson, to consist of azotic gas, mixed with a small proportion of atmospheric air. Buxton water is frequently employed, both internally and externally; one of which methods often prove beneficial, when the other would be injurious; but, as a bath alone, its virtues may not be superior to those of tepid common water. As the temperature of 82° is several degrees below that of the human body, a slight shock of cold is felt on the first immersion into the bath; but this is almost immediately succeeded by a pleasant glow over the whole system. It is therefore proper for very delicate and irritable habits. The cases which derive most benefit from the external use of Buxton waters, are those in which a loss of action, and sometimes of sensation, affects particular limbs, in consequence of long continued or violent inflammation, or external injury. Hence the chronic rheumatism, succeeding the acute, and where the inflammation has been seated in particular limbs, is often wonderfully relieved by this bath. The internal use of the water has been found to be of considerable service in symptoms of defective digestion, and derangement of the alimentary organs. A judicious use of this simple remedy, will often relieve the heartburn, flatulency, and sickness; it will increase the appetite, animate the spirits, and improve the health. At first, however, it sometimes occasions a diarrhæa, which is rather salutary than detrimental; but costiveness is a more usual effect, especially in sluggish habits. It also affords great relief when taken internally, in painful disorders of the bladders and kidneys; and has likewise been recommended in cases of gout; but when taken for these complaints, the addition of some aromatic tincture is recommended. In all cases of active inflammation, the use of these waters should be carefully avoided, on account of their supposed heating properties. A full course consists of two glasses, each containing one-third of a pint, before breakfast; which quantity should be repeated between breakfast and dinner. In chronic cases a long residence on the spot is requisite to ensure the desired effect.

BUXUS. (From *βυζα*, to become hard.) The box tree.

1. The name of a genus of plants in the Linnæan system. Class, *Monoclea*. Order, *Triandria*.

2. The pharmacopœial name of the *Buxus sempervirens* of Linnæus, possess a very strong nauseous, bitter taste, and aperient virtues. They are occasionally exhibited, in form of decoction, amongst the lower orders of people, in cases of dropsy and asthma, and worms. As much as will lay

upon a shilling, of the common dwarf box, dried and powdered, may be given at bedtime, every night, to an infant.

BUXUS SEMPERVIRENS. The systematic name of the *buxus* of the pharmacopœias. See *Buxus*.

BYARUS. A plexus of blood vessels in the brain.

BYNG. A Chinese name for green tea.

BYRETHURUM. (*Boretta*, Ital. or *barette*, Fr. a cap.) *Byrethrus*. An odoriferous cap, filled with cephalic drugs, for the head.

BYRBA. (*Bugra*, leather.) A leather skin, to spread plaisters upon.

BYRUM. (From *byr*, to hide, and *aym*, the neck.) Morbid stiffness of the neck.

BRASUS. (Heb.) A woolly kind of moss. *Pudendum maliebre*. A kind of fine linen.

BYTROS. (*Butor*, deep.) An epithet used by Hippocrates for the bottom of the stomach.

BYZEN. (From *byz*, to rush together.) In a heap; throngingly. Hippocrates uses this word to express the hurry in which the menses flow in an excessive discharge.

C.

C, In the Chemical alphabet, means nitre.

CAA-APIA. (Indian.) A Brazil root, which, chewed, has nearly the effects of ipecacuanha. It is the *Dorstenia Brasiliensis* of Willdenow. The Brazilians cure the wounds from poisoned darts with the juice of this root, which they pour into the wound.

CAA-ATATA. (Indian.) A bitter plant of Brazil, very powerfully cathartic and emetic. It resembles the euphrasia. *Ray*.

CAACICA. (Indian.) A Brazilian herb applied in cataplasms against venomous bites; called also *colubrina Lusitanica*. *Ray*.

CAACO. The name of a species of sensitive plant, whose root is used by the natives of America as an antidote to several poisons.

CAARTIMAY. *Senecio Brasiliensis*. A decoction of the plant thus called, is used as a wash to cure the itch. Its systematic name is unknown. *Ray*.

CAAGITURO. (Indian.) *Frutex baccifer Brasiliensis*. A shrub of Brazil, whose leaves are applied to ulcers, as desiccative.

CAA-OFIA. (Indian.) *Arbuscula gum-mifera Brasiliensis*. *Hypericum bacciferum* of modern naturalists. The name of a tree in the Brazils, whose bark emits a juice, when wounded, which, in a dried state, resembles gamboge, except that it is rather of a darker colour.

CAAFERA. See *Pareira brevia*.

CAAPONGA. (Indian.) The Brazilian name for crithmum; also called *Trifolia spica*. *Crithmum maritimum non spinosum*. *Inula crithmoides* of Linnaeus. The leaves and young stalks are pickled for the use of the table, they are gently diuretic.

CAABORA. (Indian.) The name of a tree which grows in the Brazils. A decoction of its leaves promotes perspiration, and is given in the cure of the venereal disease. *Ray*.

CABALISTICA ARS. *Cabala*. *Cabula*. *Kabala*. The cabalistic art. It is derived from the Hebrew word, signifying to receive by tradition. It is a term that hath been anciently used, in a very mysterious sense, amongst divines; and since, some enthusiastic philosophers and chemists have transplanted it into medicine, importing by it somewhat magical: but such unmeaning terms are now justly rejected.

Caballine ulcers. See *Aloes*.

Cabbage. See *Brassica*.

Cabbage-bark tree. See *Cortex Geoffree Jamaicensis*.

Cabbalistic art. See *Cabalistica ars*.

CABUREIBA. *Caburiba*. A name of the *Balsamum Peruvianum*. *Ray* thinks it is the tree which affords that balsam.

CACASOGA. (From *zaxos*, excrement; and *ayo*, to expel.) Cathartics. Ointments which, being rubbed on the fundus, procure stools, according to Paulus Aegineta.

CACALIA. (From *zaxos*, bad, and *nas*, exceedingly; because it is mischievous to the soil on which it grows. (*Cacamum*. The herb wild chervil, or wild carraways, formerly said to be pectoral.

CACAMOTIOTLANO QUILONI. (Indian.) *Batatas peregrina*. The purging potatoe.

CACANUM. See *Cocalia*.

CACAO. *Cocoa*. *Cocos*. *Cacavifera*. *Quahol*. *Cucata*. The cocoa or chocolate nut of Virginia and Jamaica.

CACAPRONIA. (From *zaxos*, bad, and *qaw*, the voice.) Defective articulation.

CACARI. See *Cacao*.

CACATORIA FEBRIS. (From *caco*, to go to stool.) An epithet given by Sylvius to a kind of intermittent fever, attended with copious stools.

CACCIONTE. A sort of pill recommended by Baglivi against dysenteries; its basis is catechu.

CACHEXIA. (From *zaxos*, bad, and *exis*, a habit.) A bad habit of body, known

by a depraved, or vitiated state of the solids and fluids.

CACHEXIÆ. A class of diseases in Cullen's nosology, embracing three orders, viz. *marcoræ*, *intumescentiæ*, and *impetiginosæ*.

CACHERIA UTERINA. The fluor albus is sometimes so called.

CACHERIA ICTERICA. The jaundice, or a disposition thereto.

CACHIAN. The *bathalmum verum*.

CACHELI. A little stone, or pebble. Galen says, that the cachleces, heated in the fire and quenched in whey, become astringents and useful in dysenteries.

CACHINATIO. (From *cachino*, to laugh aloud.) A tendency to immoderate laughter, as in some hysteric and maniacal affections.

CACHORE. A name for catechu.

CACHOS. (Indian.) A shrub which the Indians use as a diuretic, and to expel concretions from the kidneys.

CACHUNDE. A medicine highly celebrated among the Chinese and Indians, made of several aromatic ingredients, perfumes, medicinal earths, and precious stones. They make the whole into a stiff paste, and form out of it several figures, according to their fancy, which are dried for use. These are principally used in the East Indies, but are sometimes brought over to Portugal. In China, the principal persons usually carry a small piece in their mouths, which is a continued cordial, and gives their breath a very sweet smell. It is highly esteemed as a medicine in nervous complaints; and it is reckoned a pro-longer of life, and a provocative to venery; the two great intentions of most of the medicines used in the East.

CACHERYS. Galen says it sometimes means parched barley. In Linneus's botany, it is the name of a genus of which he enumerates three species.

CACHERYS ODONTALGICA. A plant, the root of which may be substituted for that of the pyrethrum.

CACHERIA. (*Kachurima*.) An imperfect metal, or an immature metalline ore, according to Paracelsus.

CACOLEXTERIUM. (From *κακος*, bad, and *αλεξιπνευ*, to preserve.) An antidote to poison or infectious diseases. Alexipharmics.

CACCOLIA. (From *κακος*, and *κολος*, bile.) An indisposition, or disease of the bile.

CACOCYLLA. (From *κακος*, bad, and *χυλη*, the chyle. Indigestion or depraved chylification.

CACOCYHIA. (From *κακος*, bad, and *χυμος*, juice, or humour.) A diseased or depraved state of the humours.

CACOCNEMUS. (From *κακος*, bad, and *αἷμα*, the leg.) Having a natural defect in the tibia.

CACOCORREA. (From *κακος*, bad, and *κορρεω*, to purge, or cleanse.) A medicine which purges off the vitiated humours.

CACODÆMON. (From *κακος*, bad, and *δαίμων*, a spirit.) An evil spirit, or genius, which was supposed to preside over the bodies of men, and afflict them with certain disorders. The night-mare.

CACODIA. (From *κακος*, bad, and *αἶσθησις*, to smell.) A defect in the sense of smelling.

CACOSTHES. (From *κακος*, ill, and *σθες*, a word which when applied to diseases, signifies a quality, or a disposition.) Hippocrates applied this word to malignant and difficult distempers. Galen and some others, express by it an incurable ulcer, that is rendered so through the acrimony of the humours flowing to it. Linnaeus and Vogel use this term much in the same sense with Galen, and describe the ulcer as superficial spreading, weeping, and with callous edges.

CACOPATHIA. (From *κακος*, bad, and *παθω*, affection.) An ill affection of the body, or part.

CACOPHONIA. (From *κακος*, bad, and *φωνη*, the voice.) A defect in the organs of speech; a bad pronunciation.

CACOPRAGIA. (From *κακος*, bad, and *πραγναι*, to perform.) Diseased chylopoietic viscera.

CACORHYTHUS. (From *κακος*, bad, and *ρυθμος*, order.) A disordered pulse.

CACORIS. (From *κακος*, bad.) A bad disposition of body.

CACOSTIA. (From *κακος*, and *σιση*, food.) An aversion to food, or nausea.

CACOPHYXIA. (From *κακος*, bad, and *φυξις*, pulse.) A disorder of the pulse.

CACOSTOMACHUS. (From *κακος*, bad, and *στομαχος*, the stomach.) A bad or disordered stomach; also food which the stomach rejects.

CACOSTOMUS. (From *κακος*, bad, and *στος*, a mouth.) Having a bad formed, or disordered mouth.

CACOTHYRIA. (From *κακος*, ill, and *θυρις*, the mind.) Any vicious disposition of the mind; or a diseased mind.

CACOTROPHIA. (From *κακος*, ill, and *τροφη*, nutriment.) A vitiated nourishment; a wasting of the body, through a defect of nutrition.

CACTUS. The name of a genus of plants in the Linnæan system. Class, *Icosandria*. Order, *Monogynia*. The melon-thistle.

CACTUS OPUNTIA. The systematic name of the plant bearing the epithet opuntia in the pharmacopœias. See *Opuntia*.

CACUBALUS. (From *κακος*, evil, and *βαλλω*, to cast out; so named because it was thought to be efficacious in expelling poisons.) The berry-bearing chickweed. See *Cucubalus*.

CACALA. The Arabian term for cardamoms.

CACUMEN. The top or point.

CADAVÉR. (From *cado*, to fall; because the body, when deprived of life, falls to the ground.) A carcase. A body deprived of life.

CADNIA. (Heb.) *Chlimia. Catimia.* A name given to the lapis calaminaris. See *Zinc*.

CADNIA METALLICA. A name given, by the Germans, to cobalt.

CADUCA. (From *cado*, to fall down.) See *Decidua*.

CADUCUS MORBUS. (From *cado*, to fall down.) The epilepsy, or falling sickness.

CACITAS. (From *cæcus*, blind.) Blindness. See *Cægo*, and *Amareosis*.

CÆCUM. From *cæcus*, blind. (The cæcum, or blind gut: so called from its being perforated at one end only. The first portion of the large intestines, placed in the right iliac region, about four fingers' breadth in length. It is in this intestine that the ileum terminates by a valve, called the valve of the cæcum. The *appendicula cæci vermiformis* is also attached to it. See *Intestines*.)

CÆROS. (*Kæros*.) Hippocrates, by this word, means the opportunity or moment in which whatever is to be effected should be done.

CÆSARIAN OPERATION. (So called because Julius Cæsar is said to have been extracted in this manner.) *Hysterotomia Hysterotomacæcia*. The operation for extracting the fœtus from the uterus, by dividing the integuments of the abdomen and the uterus.

There are three cases in which this operation may be necessary.—1. When the fœtus is perceived to be alive, and the mother dies, either in labour or in the last two months. 2. When the fœtus is dead, but cannot be delivered in the usual way, from the deformity of the mother, or the disproportionate size of the child. 3. When both the mother and the child are living, but delivery cannot take place, from the same causes as in the second instance. Both the mother and the child, it accounts can be credited, have often lived after the Cæsarian operation, and the mother even borne children afterwards. Heister gives a relation of such success, in his *Institutes of Surgery*, and others. In England, the Cæsarian operation has almost always failed. Mr. James Barlow, of Charley, Lancashire, succeeded, however, in taking a fœtus out of the uterus by this bold proceeding, and the mother was perfectly restored to health.

CÆSARES. *Cæsones*. Children who are brought into the world by the Cæsarian operation.

CÆTECHU. See *Catechu*.

CÆF. (Arab.) *Cufa. Caffa*. A name given by the Arabs to camphire.

CABASTRUM. A barbarous term used by

Paracelsus, to express the morbid matter which generates diseases.

CAJAN. *Cajan*. The *Phasolus creticus* of Linnaeus. A decoction of the leaves restrains the hæmorrhoids when excessive. *Ray*.

CAJEPUT OIL. *Oleum cajepute. Oleum Wittnebianum. Oleum volatile melaleuca. Oleum cajeputi*. The tree which affords this oil, by distillation of its leaves, was supposed to be the *Melaleuca leucodendron* of Linnaeus, but it appears from the specimens of the tree producing the true oil, sent home from India, by Mr. Christopher Smith, that it is another species which is therefore named *Melaleuca cajuputi*. Thunberg says cajeput oil has the appearance of inflammable spirit, is of a green colour, and so completely volatile, that it evaporates entirely, leaving no residuum; its odour of the camphoraceous kind, with a terebinthinate admixture. Goetz says it is limpid, or rather yellowish. It is a very powerful medicine, and in high esteem in India and Germany, in the character of a general remedy in chronic and painful diseases: it is used for the same purposes for which we employ the officinal æthers, to which it seems to have a considerable affinity; the cajeput, however, is more potent and pungent; taken into the stomach, in the dose of five or six drops, it heats and stimulates the whole system, proving, at the same time, a very certain diaphoretic, by which probably the good effects, it is said to have in dropsies and intermittent fevers, are to be explained. For its efficacy in various convulsive and spasmodic complaints, it is highly esteemed. It has also been used both internally and externally, with much advantage, in several other obstinate disorders; as palsies, hypochondriacal and hysterical affections, deafness, defective vision, tooth ache, gout, rheumatism, &c. The dose is from two to six, or even twelve drops.

CALABA. The Indian mastich-tree. *Cataphyllum inophyllum* of Linnaeus, a native of America, accounted vulnerary, resolvent and anodyne.

CALAGUALE RADIX. *Calaguale radix*. The root so called is knotty, and somewhat like that of the polyphyd tribe. It has been exhibited internally at Rome, with success, in dropsy; and it is said to be efficacious in pleurisy, contusions, abscesses, &c. It was first used in America, where it is obtained; and Italian physicians have since written concerning it, in terms of approbation.

CALAMAGROSTIS. (From *καλαμος*, a reed, and *αγροστις*, a sort of grass.) Sheer grass. Reed grass.

CALAMBAC. (Indian.) The agallochum, or aromatic aloë.

CALANACORUS. Indian reed.

CALAMERON. (From *καλαμος*, a reed.)

A sort of fracture which runs along the bone, in a straight line, like a reed, but is lunated in the extremity.

CALAMINA PRÆPARATA. Prepared calamine. Burn the calamine, and reduce it to powder; then let it be brought into the state of a very fine powder, in the same manner that chalk is directed to be prepared. See *Calamine*.

CALAMINE. (From *calamus*, a reed; so called from its reed-like appearance.) *Cadmia*. *Cuthnda*. *Cadmia lapidosa erosa*. *Cadmia fossilis*. *Calamina*. *Lapis calaminarius*. An ore of zinc. A sort of stone, or mineral, containing zinc and carbonic acid, united with a portion of iron, and sometimes other substances. It is very heavy, moderately hard and brittle, of a gray, yellowish, red, or blackish brown; found in quarries of considerable extent, in several parts of Europe, and particularly in this country, in Derbyshire, Gloucestershire, Nottinghamshire, and Somersetshire; as also in Wales. The calamine of England is, by the best judges, allowed to be superior in quality to that of most other countries. It seldom lies very deep, being chiefly found in clayey grounds, near the surface. In some places it is mixed with lead ores. This mineral, or semimetal, is an article in the materia medica: but, before it comes to the shops, it is usually roasted, or calcined, to separate some arsenical or sulphureous particles which, in its crude state, it is supposed to contain, and in order to render it more easily reducible into a fine powder. In this state, it is employed in collyria, for weak eyes, for promoting the cicatrization of ulcers, and healing excoriations of the skin. It is the basis of an official cerate, called *Ceratum calaminæ*, by the London College, formerly called *ceratum lapidis calaminaris*, *ceratum epuloticum*; and *ceratum carbonis zinci impuri* by the Edinburgh College. These compositions form the cerate which Turner strongly recommends for healing ulcerations and excoriations, and which have been popularly distinguished by his name. The collyria in which the prepared calamine has been employed, have consisted simply of that substance added to rose-water, or elder-flower water.

Calamint, common. See *Calamintha*.

Calamint, mountain. See *Calamintha magno flore*.

CALAMINTHA. (From *καλος*, beautiful, or *καλαμος*, a reed, and *μινθος*, mint.) Common calamint. *Calamintha montana*. *Calamintha vulgaris*. *Calamintha officinarum*. *Melissa calamintha* of Linnæus:—*pedunculis axillaribus, dichotomis, longitudine foliorum*. This plant smells strongly, like wild mint, though more agreeable; and is often used by the common people, in form of tea, against weakness of the

stomach, flatulent colic, uterine obstructions, hysteria, &c.

CALAMINTHA ANGELICA. Field calamint. Spotted calamint. *Calamintha pulegiis odore*. *Nepeta agrestis*. It is the *Melissa nepeta* of Linnæus. It was formerly used as an aromatic.

CALAMINTHA HUMILIOR. The ground-ivy.

CALAMINTHA MAGNO FLORE. *Calamintha montana*. Mountain calamint. This plant, *Melissa grandiflora* of Linnæus, has a moderately pungent taste, and a more agreeable aromatic smell than the common calamint, and appears to be more eligible as a stomachic.

CALAMINTHA MONTANA. See *Calamintha*.

CALAMUS. A word of Arabian derivation.

1. A general name denoting the stalk of any plant.

2. The name of a genus of plants in the Linnæan system. Class, *Hexandria*. Order, *Monogynia*.

CALAMUS AROMATICUS. (From *kalam*, Arab.) *Acorus verus*. *Calamus odoratus*. *Calamus vulgaris*. *Thiringa*. *Jacurantatinga*. *Typha aromatica*. *Clava rugosa*. Sweet-flag or acorus. *Acorus calamus scapi mucrone longistimo foliaceo* of Linnæus. The root of this plant has been long employed medicinally. It has a moderately strong aromatic smell, and a warm, pungent, bitterish taste; and is deemed useful as a warm stomachic. Powdered, and mixed with some absorbent, it forms a useful and pleasant dentifrice.

CALAMUS AROMATICUS ASIATICUS. The *Acorus calamus* of Linnæus.

CALAMUS ODORATUS. See *Calamus aromaticus*.

CALAMUS ROTANG. The systematic name of the plant from which we obtain the Dragon's blood. See *Sanguis draconis*.

CALAMUS SCRIPTORIUS. A kind of canal at the bottom of the fourth ventricle of the brain, so called from its resemblance to a writing pen.

CALAMUS VULGARIS. See *Calamus aromaticus*.

CALATHIANA. (From *καλαθος*, a twig basket; so called from the shape of its flowers.) The herb marsh-gentian, or *Gentiana pneumonanthe* of Linnæus.

CALBIANUM. The name of a plaster in Myrepsus.

CALCADIRUM. Vitriol.

CALCADIS. An Arabian name for white vitriol and alkali.

CALCANEUM. (From *calx*, the heel.) *Calcar pterna*. *Os calcis*. The largest bone of the tarsus, which forms the heel. It is situated posteriorly under the astragalus, is very regular, and divided into a body and processes. It has a large tuberosity or knob, projecting behind to form the heel. A *sinuous cavity*, at its fore part

which in the fresh subject is filled with fat, and gives origin to several ligaments. Two *prominences*, at the inner and fore part of the bone, with a pit between them, for the articulation of the under and fore part of the astragalus. A *depression*, in the external surface of the bone near its forepart, where the tendon of the peronæus longus runs. A large *cavity*, at the inner side of the bone, for lodging the long flexors of the toes, together with the vessels and nerves of the sole. There are two *prominences*, at the under and back part of this bone, that give origin to the aponeurosis and several muscles of the sole. The anterior surface of the os calcis is concave, for its articulation with the os cuboides, and it is articulated to the astragalus by ligaments.

CALCANTHUM. (From *χαλκος*, brass, and *ανθος*, a flower; i. e. flowers of brass.) *Calcanthos*. *Copperas*. *Vitriol*.

CALCAR. (From *calx*, the heel; also from *caleo*, to heat.) The heel-bone; also the furnace of a laboratory.

Calcareous earth. See *Calx*.

CALCARIUS FLOS. The Larkspur.

CALCARIUS LAPIS. Limestone.

CALCATAR. A name for vitriol.

CALCATON. White arsenic. *Troches of arsenic*.

CALCATRIPPA. See *Consolida media*.

CALCENA. *Calcenonius*. *Calcetius*. *Paracelsus* uses these words to express the tartarous matter in the blood; or that the blood is impregnated with tartarous principles.

CALCES, METALLIC. Metals which have undergone the process of calcination, or combustion; or any other equivalent operation.

CALCEUM EQUINUM. (From *calceus*, a shoe, and *equus*, a horse; so called from the figure of its leaf.) The herb tussilago, or colt's foot.

CALCHITHEOS. (From *καλχιος*, purple.) *Verdigrise*.

CALCHOIDES. (From *χαλξ*, a chalk-stone, and *ωδω*, fort.) *Calchoidea ossicula*. A name of the cuneiform bones.

CALCIDICIUM. The name of a medicine in which arsenic is an ingredient.

CALCIFRAGA. (From *calx*, a stone, and *frango*, to break; so named from its supposed property of breaking the human calculus.) Breakstone. The herb spleenwort, or scolopendrium, in *Scribonius Largus*.

CALCINATION. Oxydation. The fixed residues of such matters as have undergone combustion are called cinders, in common language, and calces, but now more commonly oxyds, by chymists; and the operation, when considered with regard to these residues, is termed calcination. In this general way, it has likewise been applied to bodies not really combustible, but only

deprived of some of their principles by heat. Thus we hear of the calcination of chalk, to convert it into lime by driving off its carbonic acid and water; of gypsum, or plaster stone, of alum, of borax, and other saline bodies, by which they are deprived of their water of crystallization; of bones which lose their volatile parts by this treatment, and of various other bodies.

CALCINATUM. *Cinificatum*. Terms applicable to calcined substances.

CALCINATUM MAJUS. It is whatsoever is dulcified by the chymical art, which was not so by nature; such as dulcified mercury, lead, and the like substances, which are very speedily consolidated.

CALCINATUM MAJUS POTERII. Mercury dissolved in aqua-fortis, and precipitated with salt water. *Poteries* used it in the cure of ulcers.

CALCINATUM MINUS. Any thing which is sweet by nature, and speedily cures, as sugar, manna, tamarinds &c.

CALCINONIA. See *Calcena*.

CALCIS AQUA. See *Liquor calcis*.

CALCIS VIVI FLORES. The pellicle on lime water.

CALCIS OS. See *Calcanseum*.

CALCITARI. Alkaline salt.

CALCITTA. *Vitriol*.

CALCITEOSA. *Litharge*.

CALCITHOS. *Verdigrise*.

CALCITRAPA. *Carduus stellatus*. *Jacea ramossissima, stellata, rupina*. Common star-thistle. Star-knapweed. The plant thus called in the pharmacopœias, is the *Centaurea calcitrapa* of Linnaeus:—*calycibus subduplicato-spinosis, sessilibus; foliis pinnatifidis, linearibus dentatis; caule piloso*; every part of which is bitter. The juice, or extract, or infusion, are said to cure intermitents, and the bark of the root, and the seeds, have been recommended in nephritic disorders, and in suppression of urine. It scarcely differs, in its effects, from other bitters, and is now little used.

CALCITRAPA OFFICINALIS. *Carduus stellatus lutea*. *Carduus solstitialis*. *Jacea stellata*. *Jacea lutea capite spinosa minor*. *Leucanthæ vetterum*. St. Barnaby's thistle. The *Centaurea solstitialis* of Linnaeus. It is commended as an anticteric, anti-cachectic, and lithontriptic, but is, in reality only a weak tonic.

CALCITREA. *Vitriol*.

CALCOIDEA OSSICULA. The cuneiform bones.

CALCOTAR. *Vitriol*.

CALCIFRAGUS. (From *calculus*, a stone, and *frango*, to break.) Having the power to break calculi, or stones in the human body.

1. A synonym of lithontriptic. See *Lithontriptics*.

2. A name sometimes applied to scolopendrium, or the pimpernel, from its supposed virtue.

CALCULUS. (Dim. of *calx*, a lime-stone.) *Calculus humanus.* *Bezoar microcormicum.* Gravel. Stone. In English we understand by *gravel*, small sand-like concretions, or stones, which pass from the kidneys through the ureters in a few days; and by *stone*, a calculous concretion in the kidneys or bladder, of too large a size to pass, without great difficulty. Similar concretions are found occasionally in other cavities, or passages. When a disposition to form minute calculi or gravel exists, we often find nephritic paroxysms, as they are called, (see *Nephritis*.) which consist of pain in the back, shooting down through the pelvis to the thighs; sometimes a numbness in one leg, and a retraction of either testicle in men, symptoms arising from the irritation of a stone passing through the ureters, as these cross the spermatic cord, on the nerves passing to the lower extremities. These pains, often violent, are terminated by the painful discharge of small stones through the urethra, and the patient is for a time easy. What, however, is meant by the stone is a more serious and violent disease. It is singular that these discharges of small gravel do not usually terminate in stone. Many have experienced them during a long life, without any more serious inconvenience: while the latter is a disease chiefly of the young, and depending on circumstances not easily explained. If the stone attacks persons more advanced in age, it is often the consequence of paroxysms of gout, long protracted, and terminating imperfectly.

When once a stone has acquired a moderate size, it usually occasions the following symptoms:—frequent inclination to make water, excessive pain in voiding; it drop by drop, and sometimes a sudden stoppage of it, if discharged in a stream; after making water, great torture in the glans penis, which lasts one, two, or three minutes; and in most constitutions, the violent straining makes the rectum contract and expel its excrements; or, if it be empty, occasions a tenesmus, which is sometimes accompanied with a prolapsus ani. The urine is often tinged with blood, from a rupture of the vessels, and sometimes pure blood itself is discharged. Sometimes the urine is very clear, but frequently there are great quantities of slimy sediment deposited at the bottom of it, which is only a preternatural separation of the mucilage of the bladder, but has often been mistaken for pus. The stone is a disease to which both sexes and all ages are liable; and calculi have even been found in the bladders of very young children, nay of infants only six months old.

Women seem less subject to this complaint than men, either owing to constitutional causes, or to the capaciousness, shortness, and straightness of their urethra, al-

lowing the calculi to be discharged while small, together with the urine.

Chymical analysis of urinary calculi.

It is only since the time of Scheele that we have become acquainted with the nature of urinary calculi, this subject having been quite in the dark before that great chymist discovered, in the year 1776, a peculiar acid (the lithic acid) in them, and at the same time found them to contain no lime, a circumstance which was soon after confirmed by the experiments of Bergman. From this period the chymists bestowed a particular attention upon the examination of urinary concretions, as appears from the writings of Dobson, Percival, Falconer, Achard, Hartenkeit, Tychsen, Link, Titius, Walther, Garner, Brugnatelli, Pearson, and several others, some of whom confirmed the discovery of Scheele, while others contradicted, and others enlarged it.

But we are particularly indebted to Fourcroy and Vauquelin, who, since 1786, had turned their attention on this subject, for having made many experiments, by which great light is thrown on the nature of urinary concretions. The following are the interesting results of their chymical inquiries.

The Seat and Physical Properties of Urinary Calculi.

Calculi are found in different parts of the urinary system, in the pelvis of the kidney, in the ureters, in the bladder and urethra: but as they, for the most part, originate in the kidney, the calculi renales make the nucleus of the greatest number of urinary stones. The *calculi renales* differ greatly with respect to their external qualities; for the most part, however, they consist of small, concrete, roundish, smooth, glossy, and crystalline bodies, of a red-yellow colour, like that of wood, and so hard as to admit of polishing. On account of their minuteness, they easily pass through the urinary passages in form of gravel, which being sometimes of a rough surface, cause several complaints on their passage. But in some instances they are of too great a size to be able to pass along the ureters; in which case they increase in the kidneys, sometimes to a great size. Calculi renales of this kind are generally of a brown, dark red, or black colour, and surrounded with several strata of coagulated blood and pus; they have also been observed of a yellow, reddish, and lighter colour: and some consisting of a homogeneous stony mass: but white or grey calculi renales are very rarely to be met with. Amongst the great number that were examined, one or two only were found of a grey or blackish colour, and a composition similar to those which generally bear the name of mulberry-like stones.

The stones in the ureters, which on passing into the ureters, are prevented by their size from descending into the bladder,

frequently increase very much; they, however, rarely occur; their colour is white, and they consist of phosphat of lime.

The stones in the bladder are the most frequent urinary concretions that have been principally examined; they draw their first origin from the kidneys, whence they descend into the bladder, where they increase; or they immediately originate and increase in the bladder; or they arise from a foreign body that by chance has got into the bladder, which not unfrequently happens, particularly in the female sex. Concretions of this kind differ greatly in their respective physical qualities and external form, which, however, is generally spherical, oval, or compressed on both sides; and sometimes, when there are several stones in the bladder, they have a polyedrous or cubical form; their extremities are frequently pointed or roundish, but they are very seldom found cylindrical, and more rarely with cylinders at their ends.

There is a great variety in the size of the calculi, and likewise in their colour, which is materially different, according to their respective nature and composition. They occur, 1. of a yellowish colour, approaching nearly to red, or brown; such stones consist of lithic acid. 2. Gray, or more or less white; these stones always contain phosphats of earths. 3. Dark gray, or blackish; stones of this colour have oxalats of earths. Many stones shew brown or gray spots, on a yellow or white ground, generally raised on the surface, and consisting of oxalat of lime, which is enclosed in lithic acid, when the ground-colour of the stone is of a wood colour, or in phosphat of lime, when it is white. These spots are, in general, only to be observed in the middle of the stone, or at one of its extremities.

All that is here stated, is the result of observations on more than 600 calculi; and different other colours, that are said to have been observed, either arise from heterogeneous substances, or are merely variations of the above colours. Their surface is smooth and polished in some, in others only smooth, and in others uneven, and covered with rough or smooth corpuscles, which are always of a yellow colour; in some, the surface is partly smooth and partly rough. The white ones are frequently even and smooth, half transparent, and covered with shining crystals, that generally indicate phosphat of ammonia with talc; or they are faint, and consist of minute grains; or rough, in which case they consist of phosphat of lime. The brown and dark gray stones are, from their similarity to mulberries, called mulberry-stones, and being frequently very rugged, they cause the most pain of all.

On examining the specific weight of urinary calculi in more than 500 specimens, it was found to be, in the lightest, 1213:1000,

in the heaviest, as 1976:1000. Their smell is partly strong, like urine or ammonia, partly insipid, and terrecous; especially, the white ones, which are like sawed ivory or rasped bone.

The internal texture of calculi is but seldom guessed from their external appearance, particularly when they exceed the size of a pigeon's egg. On breaking them, they generally separate into two or three strata, more or less thick and even, which prove that they are formed by different precipitations, at different times. In the middle, a nucleus is generally seen, of the same mass as the rest. When the place they are broken at is finely streaked, and of a yellow or reddish colour, the lithic acid predominates; but when they are half transparent, luminous like spar, they have ammoniacal magnesia in them, and phosphat of lime and then they are brittle and friable; but when they are so hard as to resist the instrument, of a smooth surface, and a smell like ivory, they contain saccharic lime. It frequently happens, that the exterior stratum consists of white phosphat of earth, while the nucleus is yellow lithic acid, or oxalat of lime, covered sometimes with a yellow stratum of lithic acid, in which case the nucleus appears radiant; but when it consists of lithic acid, and is covered with white phosphate of earth, it is roundish, oval, and somewhat crooked. These concretions have very seldom three strata; namely, on the outside a phosphat of salt, towards the inside lithic acid, and quite withinside an oxalat of lime; but still rarer these substances occur in more strata, or in another order, as before-mentioned.

Stones of the urethra are seldom generated in the urethra itself; however, there are instances of their having been formed in the fossa navicularis, by means of foreign bodies that have got into the urethra. We also very frequently observe stony concretions deposited between the glans and prepuce. All the concretions produced in the inside and outside the urethra consist of phosphats of earths, which are easily precipitated from the urine. There are likewise stones in the urethra, which have come out of the bladder, having been produced there, or in the kidneys; and they generally possess the properties of stones of the kidneys.

The different constituent Particles of Urinary Calculi.

It has been mentioned before, that Scheele found a peculiar acid in the urinary concretions, and likewise that phosphat of lime was discovered in them. The identity of the lithic acid, however, was much doubted by modern chymists, particularly by Dr. Pearson, who asserted that it was merely an oxyd, whereby he gave rise to the discoveries which Fourcroy and Vauquelin have since made on this subject, because they were induced to repeat the ex-

périments, in order to examine whether the lithic acid were really an acid. Their endeavours were fully rewarded, as they not only found the lithic acid and phosphat of lime in the different calculi, but also five other substances, viz. the lithat of ammonia, oxalat of lime, siliceous earth, phosphat of ammoniacal magnesia, and an animal matter.

1. *Of the Lithic or Uric Acid.*

The acid discovered by Mr. Scheele, in the urinary concretions, was styled lithic acid; or, according to Dr. Pearson's Researches, uric acid, which, after Scheele, has the following properties. It is insipid, without smell, hard, crystallizable, not soluble in cold water, and in boiling water only in several thousand times greater quantity. The solution, after having become cool, deposits the acid in form of minute yellow needles, easily soluble in the lie of fixed alkalis, out of which, however, it is precipitated by all acids, even the carbonic acid, except the sulphuric and muriatic acid, which have no effect on it. Concentrated nitric acid, on dissolving it, obtains a red colour. On distilling the lithic acid, it yields a small quantity of sublimed, undecomposed acid, very little oil and water crystallized carbonat of ammonia, carbonic acid, and a very black coal, which, however, contains neither alkali nor lime. Besides these properties, it possesses still others. On rubbing it with concentrated lie of kali or natron, it immediately forms a saponaceous, thick, and pulpy mass, which is very soluble in water, when supersaturated with alkali, but little soluble when only saturated with it. The saturated combinations have little taste, are not crystallizable, and when diluted with water, the muriatic acid precipitates the uric acid in form of small needle-like, shining, somewhat yellowish crystals. Ammonia receives very little of it, which combination is almost quite indissoluble. Lime-water has likewise very little effect on it, and the carbonats of alkalis none at all. On being dissolved in nitric acid, a part of the lithic acid is changed into oxalic acid. The red colour which appears after this combination, is said to prove, according to Pearson, that substance to be merely an oxyd, but it arises from a peculiar animal matter. When oxygenated muriatic acid is brought in contact with lithic acid, the colour of it grows pale, it puffs up, becomes soft and gelatinous, and at last obtains the consistency of a milky liquor; from which process only one-sixtieth of a white, light animal substance remains, and a quantity of carbonic acid evolves itself under continual slow effervescence. The liquor yields muriat of ammonia, oxalat of ammonia, both in crystals, free muriatic and malic acid; consequently the oxygenated muriatic acid separates the uric acid into ammonia, carbonic acid, oxalic acid, and malic acid, where-by we observe that the oxygenated muriatic

acid changes the uric acid, first into ammonia and malic acid, but on the addition of more acid, into oxalic acid; and when still more acid is added, into water and carbonic acid. The remaining white substance is the same, from which the red colour originates that appears on the combination of the uric acid with nitric acid, and which imparts the cubical form to the muriat of ammonia, obtained by the evaporation of the liquor. It remains now to be stated what is observed in the distillation of that acid, by which it yields, not only carbonat of ammonia, but also carbonic gas, very little oil, Prussic acid, partly in form of gas, partly in fluid form, a considerable quantity of coal that contains no salt, and a little water. The productions thus obtained have the smell of bitter almonds. The results of these inquiries manifestly shew, that the lithic acid is really a distinct acid from all others, consisting of azote, carbon, hydrogen, and oxygen. This peculiar acid is an excrementitious substance, which is carried off by the urine, and, at the forming of calculi, combines itself with a coloured animal matter, from which also it probably originates by a process still unknown.

2. *Of the Lithat of Ammonia.*

This substance seems to have been unknown before, or at least not properly discerned from the uric acid; and, though Scheele has observed it, he was ignorant of its particular nature. It is easily to be distinguished, by the small even strata in which it is formed, by its colour, that looks like milk coloured with coffee, and by its forming but small calculi. It dissolves in the lees of kali and natron like the lithic acid, but with the characteristic difference that it discharges ammonia, a phenomenon already observed by Scheele. It is more soluble in cold as well as warm water, than the lithic acid. It is in the same way affected by acids, except that a greater quantity is required for changing it. It is generally mixed with phosphat of ammoniacal magnesia, because it seems only to take place after a sufficient quantity of ammoniacal magnesia has been formed, to saturate the phosphat of kali and the free uric acid.

3. *Of the Phosphat of Lime.*

The existence of this substance had hitherto been but inaccurately determined, every substance which was not lithic acid being formerly comprised by the name of phosphat of lime. It occurs in small friable strata, which break in scales, or splints, of a grey white colour, and are faint, opaque, without any smell or taste, and crystallized in aluminous or spar-like form; instead of strata, it is frequently composed of friable grains, that slightly cohere, and has many holes and pores, like a spongy texture. It never forms a calculus by itself, being in a calculus always united with

animal gelatinous matter; on account of which circumstance it becomes black by exposing it to a strong heat, and burns to ash, exhaling the odour of burned bones; it yields water, oil, carbonat of ammonia, and a carbonaceous residuum. Being calcined white, it only leaves lime, and phosphat of lime, without any water of crystallization. It is not soluble in cold water, but in boiling water a part of its saline dissolves, spreading an animal odour. All acids, except the boracic and carbonic acid, dissolve it, leaving on the bottom of the vessels transparent spots of animal matter. These solutions are all precipitated by alkalis, but without any deposition, the precipitation remaining phosphat of lime. On treating the phosphat of lime with concentrated nitric acid, a thick pulpy mass of acid sulphat and phosphat of lime will be obtained, on which the alkalis, as well as carbonat of alkalis, have no effect. We never could find acid phosphat of lime, as Brugnatelli pretends to have observed.

Of the Phosphat of Ammoniacal Magnesia.

It consists of scaly, half-transparent, white, and coherent strata; can be sawed without crumbling, and reduced to a fine, white and white powder. It is of a sweetish, lipid taste, somewhat soluble, and crystallized in rhomboids, or thick laminas, persed in the cavities of other calculous substances; and it is frequently found on the surface of other calculi. It consists, betwixt its strata, a gelatinous substance, but less than the phosphat of lime which account it also blackens by being heated. Though it be but little soluble in water, yet it dissolves in such a quantity as to be capable of crystallizing by slow evaporation. Acids dissolve it more quickly than they do the phosphat of lime. Weak sulphuric acid entirely dissolves it, forming sulphat of ammoniacal magnesia. In diluted muriatic or nitric acid, it disappears more quickly than phosphat of lime. Ammonia, by which that it is made turbid, only precipitates small particles of magnesia. The lees of fixed alkalis disengage from it ammonia, without forming with it a solution; and, depriving it of the phosphoric acid, leave the magnesia behind.

5. Of the Oxalat of Lime.

It is, according to our observations, only found in the mulberry-like calculi, in combination with a coloured animal matter, and consist of strata covered with pointed, reddish, rough or smooth protuberances; outside it appears of a dark or brown colour, but internally it is grey, frequently with white streaks, of a solid texture, and may be polished like ivory; it breaks in scales, or in the shape of shells; and, on being pounded, or sawed, it exhales an animal odour, like semen. It is the heaviest

of all calculous substances, and the only one which yields one-third of lime by calcination. It dissolves with difficulty in acids and is precipitated unaltered by alkalis from nitric acid. The fixed alkalis decompose it when they are impregnated with carbonic acid, and when it is pulverized, and the solution heated, whereby carbonat of lime and oxalat of alkalis are obtained.

The great quantity of animal matter which constantly adheres to this oxalat of lime is very characteristic, it imparts the brown, reddish, blackish colour to the above kind of stones, and likewise the fine and solid texture. This substance may be obtained by putting small pieces of these stones into diluted nitric acid, whereby it appears of the same colour, and becomes soft and spongy. The great hardness of this kind of calculous substance, most probably arises from the intimate connexion of its particles, produced by the combination of the oxalat of lime with animal matter, in the same way as lime obtains a great degree of solidity by its combination with albuminous matter, of which, and of a peculiar matter of urine, that animal substance seems to consist.

6. Of the Siliceous Earth.

Amongst 600 calculi that were examined, there were only two which contained this earth; both had the texture of mulberry-like stones, though of a lighter colour, and by being calcined, lost one-third of their weight, without giving free lime; heated with acids they lost nothing, but when melted with four times as much of alkali, they yielded siliceous earth by being treated with muriatic acid. They contained phosphat of lime, and an animal matter similar to that which is united with the oxalat of lime. They were hard, difficult to be sawed and pulverized, and the powder made scratches in metal. On being burnt, they emit an animal odour; they imparted nothing to the boiling water, and to the acids a little phosphat of lime, which difficultly separates from the siliceous earth. Alkalis, either pure, or combined with carbonic acid, did not affect them, merely depriving them of a part of their animal matter. Their essential character consists in their being fusible and vitrifiable with fixed alkalis.

7. Of the Animal Matter.

All the six substances just examined, which constitute the urinary calculi of the human species, are always combined with an animal matter, as appears from its being burnt to coal from the productions it yields by distillation, from its stench on being burnt, and from the cellulous membranous floccula which remain when pieces of calculi are dissolved in diluted acids. This animal matter has been frequently, and with good reason, considered as the basis of all urinary concretions, like as in bones

the gelatinous matter, the first basis of the bones, forms an organic texture, in the interstices of which the phosphat of lime is deposited. It is very remarkable, that the different constitutory particles of urinary calculi are combined with a dissimilar animal matter, which are sometimes albuminous, sometimes gelatinous, sometimes composed of both, and frequently united with the matter of urine. Thus the lithic acid, or the lithat of ammonia, contains a third of albuminous matter, combined with the matter of urine, the phosphats of earths, albuminous matter, gelatine in form of membranes, and laminas, or tela cellulosa; the oxalat of lime, a spongy, yet more solid texture, of the colour of albumen, and the siliceous earth, a similar substance. On the whole, the animal matter seems to unite and join together all the acid and saline particles of urinary concretions.

The Classification of Urinary Stones.

The old classification of urinary calculi, made according to their figure and their size, cannot at present, where we have acquired so accurate a knowledge of their internal nature, be retained, as they ought rather to be classed according to their constituent particles; however, no regard is to be had to the animal matter, as being found in all urinary concretions, and having no influence on their respective difference. On comparing the results of the analyses of more than 600 stones, Fourcroy was induced to bring them under three genera; *the first* of which comprehends such stones as are merely composed of one substance, besides the animal matter; *the second* contains urinary concretions, consisting of two substances, besides the animal matter; and *the third* comprises all those which are formed by more than three calculous substances. These three genera comprehend about twelve species, namely, the first genus three, the second seven, and the third two; but it must be remembered that the number of the genera, as well as of the species, is determined after the observations hitherto made, and may consequently be increased in future.

1. *The first species* of urinary concretions consists of lithic acid, and stones of this kind most frequently occur, as there were, amongst 600, about 150. They are easily distinguished by their reddish or high yellow colour, much resembling that of wood, by their brittle, radiant-like, homogeneous, and fine texture, and by their perfect solubility in the lies of fixed alkalis, without disengaging the smell of ammonia. Their size varies from the bigness of a pea to that of a duck's egg, &c. and their figure is roundish, spheroid, compressed, oval, oblong, &c. the surface polished like marble, but frequently rough and watry; of a crimson light red, yellowish, light brown colour, but never white, gray, or black; their strata differ in num-

ber and thickness, and are frequently of a smooth surface. The specific weight of these stones is from 1,276, to 1,786, but generally more than 1,500. The urinary concretions in the kidneys are mostly of this species.

2. *The second species* is composed of lithate of ammonia, and differs from the former by disengaging ammonia on their being dissolved in the lies of fixed alkalis. Concretions of this kind are generally small, of a pale or gray colour, and consist of fine strata, easily separable from each other; they mostly contain a nucleus, which is easily separated from the strata that cover it. Their figure is generally oblong, compressed like almonds, and of a smooth surface, which is frequently crystalline. Their specific weight varies from 1,225 to 1,720. They are entirely soluble in water, particularly when previously pulverized. All acids, principally the muriatic acid, deprive them of the ammonia, leaving the pure lithic acid behind. They are frequently found covered with a thin stratum of lithic acid. Amongst 600 calculi there were but few of this kind.

3. *The third species*, consisting of oxalat of lime, are easily to be distinguished by the protuberances and inequality of their surface, whence they have got the appellation of mulberry-like stones; by their hardness, gray colour, solid texture, their polish like ivory, in the inside, and their particular smell on being sawed, which resembles that of semen. A peculiar characteristic, which distinguishes them from all others, consists in their leaving lime after the calcination, in their being with difficulty soluble in acids and not soluble in alkalis, and, at last, in their being only decomposed by the lees of carbonats of alkali. They weigh from 1,428 to 1,976, and their size varies from that of a calculus renalis to the bigness of an egg, or more; their figure is generally spherical or spheroid. They often make the nucleus of other stones, in which case they belong to another species. In 300 stones they bore the proportion of one-fourth or one-fifth.

4. *Stones of this species* contain lithic acid and phosphat of earth, but in a separate state. Their surface is white, cretaceous, brittle, and half transparent, as it either consists of phosphat of lime, or of phosphat of ammoniacal magnesia, the kernel being formed by lithic acid; thus both constituents are exactly separate from each other. They were found in the proportion of one-twelfth amongst the stones that were examined, and they grow bigger than any of the rest, as they appear from the size of an egg to that of the whole bladder, even when extended. They generally have an oval form, often pointed at one end, of a smooth surface, which, however, is frequently covered with crystals of phosphat of ammoniacal mag-

nesia. Sometimes the lithic acid in the middle is alternately covered with phosphat of lime, and phosphat of ammoniacal magnesia. The specific weight of these stones is extremely variable.

5. The fifth species of calculi contains, likewise, lithic acid and phosphats of earth, but intimately mixed with each other. Of these stones a great many varieties are observed, depending on the proportionable quantity of their constituent particles, as well as on the strata in which they lie above one another. The chief constituents, the phosphats of earths, are separated in different strata, but sometimes so intimately mixed with each other, that it is impossible to distinguish them with the eye; and the analysis could only shew their difference. From this circumstance arise the variety in the colour, figure, and number of the strata. The colour, however, is generally gray, but frequently variegated like marble, sometimes like soap. Their figure is irregular, oval, or globular, and the surface mostly brittle, cretaceous, or whitish, as to make us believe that they only consist of phosphat of lime. The polyedrous stones generally belong to this species, when they have the appearance of being worn away by rubbing. They make about one-fifth of the stones that were examined. Their specific weight varies extremely, the least being 1,213, the greatest 1,739.

6. This species is constituted by lithat of ammonia and phosphat of earth, *i. e.* of lime and ammoniacal magnesia; and resembles in its external appearances the fourth species. One of the constituents, generally the lithat of ammonia, makes the nucleus, while a mixture of the two others, but rarely one by itself, forms the crust. Sometimes, however, the nucleus contains also the phosphats, and the crust a little lithat of ammonia, which, even in some varieties, is mixed with pure lithic acid. The strata in stones of this kind are more easily separable, and always smaller than those of the fourth species. Their specific weight is 1,312 to 1,761; and they are more rarely met with than most of the rest. Amongst 600 there were only twenty of this kind,

7. Stones of the seventh species consist likewise of lithat of ammonia and phosphat of earths, but intimately mixed with each other. They are of a paler colour, much lighter than the first species, and disengage a great deal of ammonia on their being treated with kali. We found them only in the proportion of one-fortieth amongst the stones which we have analysed. They never grow so large as the two former.

8. The constituent particles of the eighth species are phosphat of lime and phosphat of ammoniacal magnesia. The pure white colour, the friability, their being insoluble in alkalis, and their easy solubility even in

weak acids, constitute the chief characteristics of this sort of stones, of which about 60 were found amongst 600: sometimes they are of an enormous size, of irregular form, rarely round, but frequently of an uneven surface, and resembling an incrustation. Their texture is formed of white brittle strata, sometimes interwoven with solid half-transparent crystals of ammoniacal magnesia. The crusts formed on foreign bodies that happened to penetrate into the bladder, belong to this species; the specific weight of which is 1,138 to 1,473.

9. This species of calculi contains oxalat of lime, but externally uric acid, in more or less quantity, and are only to be distinguished by the nucleus from the first species. The proportion of both constituents, and the specific weight, vary extremely, the latter being 1,341 to 1,754. Sometimes the nucleus, consisting of oxalat of lime, is only covered on one side with uric acid, and discernible on the other by protuberances with which the surface is variegated; which variety, however, seldom occurs.

10. Stones of this species have, in their centre, oxalat of lime, surrounded by phosphat of earths; the kernel is gray, or brown, and radiant-like, the crust white and cretaceous; their size and figure differ extremely, and their specific weight is from 1,168 to 1,752. They amount to one-fifth of the 600 stones that were examined.

11. This species contains stones composed of three or four calculous substances, namely, of oxalat of kali, phosphat of earths, and of uric acid, either pure or combined with ammonia. They rarely occur; and amongst 600 stones only ten or twelve were observed. They often consist of three distinct strata, *viz.* in the interior, of oxalat of lime; in the middle of lithat of ammonia; and the exterior, of phosphats of earths, which are frequently mixed with uric acid or lithat of ammonia, all which are distinguished on their being sawed through. This species comprehends three varieties; the first of which consists of oxalat of lime, uric acid, and phosphats of earths; the second contains lithat of ammonia, combined with pure uric acid, and the two other constituents; the third has, besides these two substances, free uric acid and lithat of ammonia, mixed with the phosphats of earths. We forbear to mention other varieties of this species, as being less remarkable and instructive.

12. The last species of calculi is of a very complicated composition. The siliceous earth seems to have taken the place of the oxalat of lime; it is mixed with uric acid and lithat of ammonia, and covered by phosphats of earths. Stones of this kind are the rarest of all, and there were only two amongst 600.

The Causes of the Generation of Urinary Calculi.

To inquire into the causes by which

which urinary concretions are produced, is both interesting and useful, however attended with the greatest difficulties. The writings of medical authors are full of conjectures and hypotheses with regard to this subject, on which nothing could be ascertained before we had acquired an accurate knowledge of the nature of urinary concretions. It is owing to this circumstance that the most enlightened physicians acquiesced in ascribing the immediate cause of them to a superabundance of terreous matter in the urine; and Boerhaave, as well as, particularly, Van Swieten, imagined that the urine of all men contained calculous matter in the natural state, and that, for the generation of stones, a nucleus was only required, to attract it. That this may be the case, in some instances, is proved by frequent experience; but stones produced by foreign bodies, that have accidentally got into the urethra or bladder, are always white and composed of phosphat of earth, and seldom or never covered with lithic acid, a substance which is observed to form the stones that most frequently occur; but even in these the nucleus consists of a substance formed in the body itself, as a particle descended from the kidneys, &c. which must, therefore, have necessarily originated in a peculiar internal cause. A superabundance of uric acid in stony patients, and its more copious generation than in a sound state, though it seems to be one of the principal and most certain causes, is by no means satisfactory, as it only explains the precipitation of stony matter from the urine, but not why it unites in strata. A coagulating substance is required for separating, attracting, and, as it were, agglutinating the condensable particles that are precipitated. This substance is undoubtedly the animal matter which we have constantly found in all calculous masses, and which seems to constitute the basis of stones, like the membranous gelatina that of bones. It is known that the urine of calculous patients is generally muddy, ductile, in threads, slimy, and as if mixed with albumen, which quality it obtains at the moment when the ammonia is disengaged, or on the addition of kali that separates it from the acid in which it was dissolved; and in all cases of superabundance of lithic acid the urine contains a great quantity of that animal matter, which promotes the precipitation of it, and attracts and unites the particles thus separated. Hence it appears, that every thing capable of increasing the quantity of that pituitous gluten in the urine, may be considered as the remote cause of the formation of calculi. And the old ideas on pituitous temperaments, or superabundant pituita, &c. which were thought to dispose people to a calculus, seems to be connected with the late discoveries on the nature of urinary stones. Though the animal matter appears

to be different in different calculi, yet certain, that every calculous substance contains an animal gluten, from which it crete and solid state arises; whence may fairly state the superabundance of substance as the chief and principal of the formation of calculi.

There are, however, other causes, seem to have a particular influence on nature of urinary stones, and the str which they are formed; but it is extremely difficult to penetrate and to explain. We are, for instance, entirely ignorant the manner in which urinary stones formed from the oxalat of lime; tho from their occurring more frequently children than in adults, we might be titled to ascribe them to a disposition, a cause considered by Boerhaave the general source of a great number diseases incident to the infantile age. opinion seems to be proved by the id Bonhomme, physician at Avignon, o oxalic or saccharic acid, as the cause mollities ossium in the rickets; by this being discovered in a species of saliva Brugnatelli; and, lastly, by an observation of Turgais, who found this acid i urine of a child diseased with worms. but rarely observe saccharic acid in human body, which appears to be adventitious, and by which the a matter is rendered coagulable, and sited, or precipitated, with the oxal lime; or the oxalic acid decompose phosphat of lime, and forms an insol combination, incapable of being any kept dissolved in the urine. It is, ever, extremely difficult to determine far the constitution of the body is covered with that particular disposition i urine, of precipitating sometimes phosphat of lime mixed with oxalat of lime, times phosphat of ammoniacal magn either by itself or mixed with lithic &c. &c. Who can explain the reason of 600 stones, there were only two in siliceous earth could be traced? Still difficult is it to explain the causes the above substances precipitate either once or in different strata; but it may suffice to have shewn how many observations and experiments are required, and accurate attention and perseverance necessary, in order to throw light on difficult a subject.

CALCULUS BILIARIS. See *Gall-stone*.

CALDARIUM. (From *calo*, to make A vessel in the baths of the ancient bold hot water.

CALIFACIENTE. (*Calificientia*, sc. *camenta*: from *calidus*, warm, and *fac* make. (Medicines, or other substances which excite a degree of warmth in parts to which they are applied; as *spiritus vini*, &c. They belong to the of stimulants.

CALCIBULA. (*Quid angustis calc*
U

i. e. mensibus, floreat; so called because it flowers every month.) Marigold.

1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia necessaria*.

2. The pharmacopœial name of the *Calendula sativa*. *Chrysanthemum. Sponsa solis. Caltha, vulgaris*. Single marigold. Garden marigold. The flowers and leaves of this plant, *Calendula officinalis* of Linnæus:—*seminibus cymbiformibus, muricatis, incurvatis omnibus*, have been exhibited medicinally: the former, as aperients in uterine obstructions and icteric disorders, and as diaphoretics in exanthematous fevers; the latter, as gentle aperients, and to promote the secretions in general.

CALENDULA ALPINA. The *Arnica montana* of Linnæus. See *Arnica*.

CALENDULA ARVENSIS. The wild marigold. The *Caltha arvensis* of Linnæus. It is sometimes preferred to the former. Its juice is given, from one to four ounces, in jaundice and cachexia; and the leaves are commended as a salad for children afflicted with scrophulous tumours.

CALENDULA OFFICINALIS. The systematic name of the single marigold plant. See *Calendula*.

CALENDULA PALUSTRIS. The *Caltha palustris* of Linnæus. Common single marsh marigold. It is said to be caustic and deleterious: but this may be questioned.

CALENTURE. A febrile delirium, said to be peculiar to sailors, wherein they imagine the sea to be green fields, and will throw themselves into it if not restrained. Bonetus gives an account of it; also Dr. Oliver and Dr. Stubbs. It is probably a species of phrenitis.

CALISTUM. (Indian.) A tree which grows in Malabar, whose bark, made into an ointment with butter, cures convulsions from wounds, and heals ulcers. The juice of the bark cures the aphthæ, and, taken inwardly, the dysentery. *Ray*.

CALI. (Arab.) The same as kali.

CALICHAFA. The spina alba, or white-thorn.

CALIDUM. In medical language, it is commonly used with the adjective *animale*, or *innatum*, for animal heat, or the vis vitæ.

CALISTA. (From *καλός*, a nest, which it somewhat resembles.) *Calliæte*. A fungus growing on the juniper-tree.

CALIGO. A disease of the eye, known by diminished or destroyed sight; and by the interposition of a dark body between the object and the retina. It is arranged by Cullen in the class *locales*, and order *dysæsthesiæ*. The species of caligo are distinguished according to the situation of the interposed body: thus *caligo lentis*, *caligo corneæ*, *caligo pupillæ*, *caligo humorum*, and *caligo palpebrarum*.

CALIGO LENTIS. *Glaucoma Woulhousei*. The true cataract. See *Cataract*.

CALIGO CORNEÆ. An opacity of the cornea. See *Caligo*.

CALIGO PUPILLÆ. *Synchysis. Amyosis*. Blindness from obstruction in the pupil. See *Caligo*.

CALIGO HUMORUM. *Glaucoma Vogelii*. Blindness from a fault in the humours of the eye. See *Caligo*.

CALIGO PALPEBRARUM. Blindness from a disorder in the eyelids. See *Caligo*.

CALIHACHA. The cassia-lignea, or cassia-tree of Malabar.

CALIMIA. The lapis calaminaris.

CALIX. (*Κάλυξ*, *icis*, m.; from *καλυπτω*, to cover.) *Calyx*.

1. The term calix is given to the membrane which covers the papillæ in the pelvis of the human kidney.

2. The name of the case, or sheath, in which the flower of plants is concealed before it expands.

CALLEUM. (From *καλλυνω*, to adorn.) *Calleon*. The gills of a cock, which, Galen says, is food not to be praised or condemned.

CALLENA. A kind of salt-petre.

CALLI. Nodes in the gout. *Galen*.

CALLIA. (From *καλός*, beautiful.) A name of the chamomile.

CALLIBLEPHARA. (From *καλός*, good, and *βλεφαρον*, the eyelid.) Medicines, or compositions, appropriated to the eyelids.

CALLICREAS. (From *καλός*, good, and *κρεας*, meat; so named from its delicacy as food.) The pancreas, or sweetbread.

CALLIGONUM. (From *καλός*, beautiful, and *γονυ*, a knot, or joint; so named from its being handsomely jointed, like a cane.) The polygonum, or knot-grass.

CALLIOMARCHUS. The Gaulish name, in Marcellus Empiricus, for tussilago, or colt's-foot.

CALLON. A kind of night-shade.

CALLIPHYLLUM. (From *καλός*, beauty, and *φυλλον*, a leaf.) The herb adiantum, or maidenhair. See *Adiantum*.

CALLISTRUTHIA. (From *καλός*, good, and *τρυθός*, a sparrow: because it was said to fatten sparrows.) A fig mentioned by Pliny, of a good taste.

CALLITRICUM. (From *καλός*, beauty, and *τριχ*, hair; so named because it has the appearance of long, beautiful hair, or, according to Littleton, because it nourishes the hair, and makes it beautiful.) The herb maidenhair.

CALONX. (From *καλός*, fair.) Hippocrates uses this word, to signify that decency and gravity of character and deportment which it is necessary that all medical men should be possessed of.

CALLOSITAS. Callosity, or preternatural hardness.

CALLOUS. A surgical term, signifying hardened or indurated; thus the callous edges of ulcers.

CALLUS. *Callum.*

1. The bony matter deposited between the divided ends of broken bones, about the fourteenth day after the fracture.

2. A preternatural hardness, or induration, of any fleshy parts.

CALOCATANUS. (From *καλος*, beautiful, and *καταν*, a cup; so called from the beauty of its flower and shape.) The papaver rhæas, or wild poppy.

CALOMELANUS TURQUETI. So Riverius calls a purgative medicine, composed of calomel and scammony.

CALOMELAS. (From *καλος*, good, and *μελας*, black; from its virtues and colour. The preparation called Æthiop's mineral, or *hydrargyrus cum sulphure*, was formerly and properly so named. But calomel now means a white preparation of sublimed mercury.) See *Sublimatus hydrargyri*.

CALORIC. (*Caloricum*; from *calor*, heat.) Heat. Igneous fluid.

Heat and cold are perceptions of which we acquire the ideas from the senses; they indicate only a certain state in which we find ourselves independent of any exterior object. But as these sensations are for the most part produced by bodies around us, we consider them as causes, and judging by appearances, we apply the terms *hot*, or *cold*, to the substances themselves; calling these bodies *hot*, which produce in us the sensation of heat, and those *cold*, which communicate the contrary sensation.

This ambiguity, though of little consequence in the common affairs of human life, has led unavoidably to confusion and perplexity in philosophical discussions. It was to prevent this, that the framers of the new nomenclature adopted the word *caloric*, which denotes that which produces the sensation of heat.

Theories of Heat.

Two opinions have long divided the philosophical world concerning the nature of heat.

1. The one is: that the cause which produces the sensation of heat, is a real, or distinct substance, universally pervading nature, penetrating the particles or pores of all bodies, with more or less facility, and in different quantities.

This substance, if applied to our system in a greater proportion than it already contains, warms it, as we call it, or produces the sensation of heat; and hence it has been called *caloric* or *calorific*.

2. The other theory concerning heat is; that the cause which produces that sensation, is *not* a separate or self-existing substance; but that it is merely like gravity, a property of matter; and that it consists in a specific or peculiar motion, or vibration of the particles of bodies.

The arguments in favour of the first theory have been principally deduced from the evolution, and absorption of heat

during chemical combinations; those of the latter, are chiefly founded on the production of heat by friction. For it has been observed, that whatever is capable of producing motion in the particles of any mass of matter, excites heat. Count Rumford and Professor Davy have paid uncommon attention to this fact, and proved, that heat continues to be evolved from a body subjected to friction, so long as it is applied, and the texture or form of the body be not altered.

All the effects of heat, according to this theory, depend therefore entirely on the vibratory motion of the particles of bodies. According as this is more or less intense, a higher or lower temperature is produced; and as it predominates over—is nearly equal—or inferior to the attraction of cohesion, bodies exist in the gaseous, fluid, or solid state.

Different bodies are susceptible of it in different degrees, and receive and communicate it with different celerity. From the generation, communication and abstraction of this repulsive motion, under these laws, all the phenomena ascribed to heat are explicable.

Each of these theories has been supported by the most able philosophers, and given occasion to the most important disputes in which chemists have been engaged; and have contributed in a very particular manner to the advancement of the science. The obscurity of the subject, however, is such, that both parties have been able to advance the most plausible arguments.

Setting aside all enquiries concerning the merits of these different doctrines, we shall confine ourselves to the general effects, which heat produces on different bodies. For the phenomena which heat presents, and their relation to each other, may be investigated with sufficient precision, though the materiality, or immateriality of it, may remain unknown to us.

Nature of Heat.

Those who consider heat as matter, assert that caloric exists in two states, namely, in combination, and in a radiant state, or at liberty.

In the first state it is not sensible to our organs, nor indicated by the thermometer: it forms a constituent part of the body; but it may be brought back to the state of radiant or sensible heat. In this state it affects animals with the sensation of heat. It therefore has been called sensible or free heat, or fire; and is synonymous with uncombined caloric, thermometrical caloric, caloric of temperature, interposed caloric, &c. expressions now pretty generally superseded.

From the diversity of opinions among chemists respecting the nature of caloric, several other expressions have been intro-

duced, which it is proper to notice. For instance, by *specific heat* is understood, the relative quantities of caloric contained in equal weights of different bodies at the same temperature. *Latent heat* is the expression used to denote that quantity of caloric which a body absorbs when changing in its form. It is, however, more properly called *caloric of fluidity*. The disposition, or property, by which different bodies contain certain quantities of caloric, at any temperature, is termed their *capacity for heat*. By the expression of *absolute heat*, is understood the whole quantity of caloric which any body contains.

Methods of exciting and collecting Heat.

Of the different methods of exciting heat, the following are the most usual:

1. *Production of Heat by Percussion or Collision.*

This method of producing heat is the simplest, and therefore it is generally made use of in the common purposes of life for obtaining fire.

When a piece of hardened steel is struck with a flint, some particles of the metal are scraped away from the mass, and so violent is the heat which follows the stroke, that it melts and vitrifies them. If the fragments of steel are caught upon paper, and viewed with a microscope, most of them will be found perfect spherules, and very highly polished. Their sphericity demonstrates that they have been in a fluid state, and the polish upon their surface, shews them to be vitrified.

No heat, however, has been observed to follow the percussion of liquids, nor of the softer kind of bodies which yield to a slight impulse.

2. *Production of Heat by Friction.*

Heat may likewise be excited by mere friction. This practice is still retained in some parts of the world. The natives of New Holland are said to produce fire in this manner, with great facility, and spread it in a wonderful manner. For that purpose, they take two pieces of dry wood; one is a stick, about eight or nine inches long, and the other piece is flat; the stick they point into an obtuse point at one end, and pressing it upon the other piece, they turn it very nimbly, by holding it between both hands, as we do a chocolate-mill, often shifting their hands up, and then moving down upon it, in order to increase the pressure as much as possible. By this method, they get fire in a few minutes, and from the smallest spark they increase it with great speed and dexterity.

If the irons at the axis of a coach-wheel are applied to each other, without the interposition of some unctuous matter to keep them from immediate contact, they will become so hot when the carriage runs swiftly along, as to set the wood on fire; and the fore-wheels, being smallest, and

making most revolutions in a given time, will be most in danger.

The same will happen to mill-work, or to any other machinery.

It is no uncommon practice in this country, for blacksmiths to use a plate of iron as an extemporaneous substitute for a tinder-box; for it may be hammered on an anvil till it becomes red-hot, and will fire a brimstone match. A strong man, who strikes quick, and keeps turning the iron so that both sides may be equally exposed to the force of the hammer, will perform this in less time than would be expected.

If, in the coldest season, one dense iron plate be laid on another, and pressed together by a weight, and then rubbed upon each other by reciprocal motions, they will gradually grow so hot as, in a short time, to emit sparks, and at last become ignited.

It is not necessary that the substances should be very hard; a cord rubbed backwards and forwards swiftly against a post or a tree will take fire.

Count Rumford and Professor Pictet have made some very ingenious and valuable experiments concerning the heat evolved by friction.

3. *Production of Heat by Chemical Action.*

To this belongs the heat produced by combustion. There are, besides this, many chemical processes wherein rapid chemical action takes place, accompanied with a development of heat, or fire ~~and~~ flame.

4. *Solar Heat.*

It is well known that the solar rays, when collected by a mirror, or lens, into a focus, produce the most astonishing effects.

Dr. Herschell has discovered that there are rays emitted from the sun, which have not the power of illuminating or producing vision; and that these are the rays which produce the heat of the solar light.

Consequently, heat is emitted from the sun in rays, but these rays are not the same with the rays of light.

5. *Production of Heat by the Electric Spark, and by Galvanism.*

The effects of electricity are too well known in this point of view, to need any description.

Galvanism has of late become a powerful instrument for the purpose of exciting heat. Not only easily inflammable substances, such as phosphorus, sulphur, &c. have been fired, but likewise gold, silver, copper, tin, and the rest of the metals, have been burnt by means of galvanism.

General Effects of Heat.

Expansive property of Heat.—This is the first and most obvious effect which heat produces on bodies. Experience has taught us that, at all times, when bodies become hot, they increase in bulk. The bodies experience a dilatation which is greater in proportion to the accumulation of caloric.

or, in other words, to the intensity of the heat. This is a general law, which holds good as long as the bodies have suffered no change either in their combination or in the quantity of their chemical principles.

This power which heat possesses, consists, therefore, in a constant tendency to separate the particles of bodies. Hence philosophers consider heat as the *repulsive power* which acts upon all bodies whatever, and which is in constant opposition to the power of attraction.

The phenomena which result from these mutual actions, seem, as it were, the secret springs of nature. Heat, however, does not expand all bodies equally, and we are still ignorant of the laws which it follows.

1. *Expansion of Fluid Bodies by Heat.*

Take a glass globe, with a long slender neck (called a bolt head); fill it up to the neck with water, ardent spirit, or any other fluid which may be coloured with red or black ink, in order to be more visible, and then immerse the globe of the instrument in a vessel of hot water; the included fluid will instantly begin to mount into the neck. If it be taken out of the water and brought near the fire, it will ascend more and more, in proportion as it becomes heated; but upon removing it from the source of heat, it will sink again: a clear proof that caloric dilates it, so as to make it occupy more space when hot than when cold. These experiments may, therefore, serve as a demonstration that heat expands fluid bodies.

It appears that liquids of the least density expand most, with the same temperature. Thus hydrogen gas dilates more with the same degree of heat than atmospheric air; atmospheric air more than sulphuric ether; ether more than ardent spirit; ardent spirit more than oil; oil more than water; water more than acids, and acids more than mercury. But if we compare the periods of time necessary for each fluid to acquire the *maximum* of rarefaction it is susceptible of, there is no law to guide us yet known.

2. *Expansion of Aeriform Bodies by Heat.*

Take a bladder partly filled with air, the neck of which is closely tied, so as to prevent the inclosed air from escaping, and let it be held near a fire. The air will soon begin to occupy more space, and the bladder will become gradually distended; on continuing the expansion of the air, by increasing the heat, the bladder will burst with a loud report.

3. *Expansion of Solid Bodies by Heat.*

If we take a bar of iron, six inches long, and put it into a fire till it becomes red-hot; and then measure it in this state accurately, it will be found 1-90th of an inch longer than it was before; that is, about 120th part of the whole. That the metal is proportionally expanded in

breadth, will be seen by trying to pass it through an aperture which it fitted exactly when cold, but which will not admit it when red hot. The bar is, therefore, increased in length and diameter.

To discover the minutest changes of expansion by heat, and the relative proportions thereof, instruments have been contrived, called *Pyrometers*, the sensibility of which is so delicate as to shew the expansion from 1-50000 to 1-100000 of an inch.

It is owing to this expansion of metals, that the motion of time-pieces is rendered erroneous; but the ingenuity of artists has discovered methods of obviating this inaccuracy, by employing the greater expansion of one metal to counteract the expansion of another; this is effected in what is called the grid-iron pendulum. Upon the same principle a particular construction of watches has been contrived.

The expansion of metals is likewise one of the principal reasons that clocks and watches vary in winter and summer, when worn in the pocket, or exposed to the open air, or when carried into a hotter or a colder climate. For the number of the vibrations of the pendulum are always in the sub-duplicate ratio of its length, and as the length is changed by heat and cold, the times of vibration will be also changed. The quantity of alteration, when considered in a single vibration, is exceedingly small, but when they are often repeated, it will be very sensible. An alteration of one-thousandth part in the time of a single vibration of a pendulum which beats seconds, will make a change of eighty-six whole vibrations in twenty-four hours.

As different metals expand differently with the same degree of heat; such musical instruments, therefore, whose parts are to maintain a constant true proportion, should never be strung with different metals. It is on this account that harpsichords, &c. are out of tune by a change of temperature.

Bodies which are brittle, or which want flexibility, crack or break, if suddenly heated or cooled. This likewise depends upon the expansive force of heat, stretching the surface to which it is applied, while the other parts, not being equally heated, do not expand in the same ratio, and are therefore torn asunder or break. Hence thin vessels stand heat better than thick ones.

Measurement of Heat.

Upon the expansive property of heat, which we have considered before, is founded its artificial measurement. Various means have been therefore employed to assist the imperfection of our sensations in judging of the different degrees of heat, for our feelings unaided afford but very inaccurate information concerning this matter; they indicate the presence of heat, only when the

bodies presented to them are *hotter* than the actual temperature of our organs of feeling. When those bodies are precisely of the same temperature with our body, which we make the standard of comparison, we then are not sensible of the presence of heat in them. When their temperature is less hot than that of our bodies, their contact gives us what is called the sensation of cold.

The effects of heat upon material bodies in general, which are easily visible to us, afford more precise and determinate indications of the intensity, than can be derived from our feelings alone. The ingenuity of the philosopher and artist has therefore furnished us with instruments for measuring the relative heat or temperature of bodies. These instruments are called *Thermometers* and *Pyrometers*. By these, all degrees are measurable, from the slightest, to that of the most intense heat.

1. Nature of the Thermometer.

A thermometer is a hollow tube of glass, hermetically sealed, and blown at one end in the shape of a hollow globe. The bulb and part of the tube are filled with mercury, which is the only fluid which expands equally. When we immerse the bulb of the thermometer in a hot body, the mercury expands, and of course *rises* in the tube; but when we plunge it into a cold body, the mercury contracts, and of course *falls* in the tube.

The rising of the mercury indicates, therefore, an increase of heat; its falling, a diminution of it; and the quantity which it rises or falls, denotes the proportion of increase or diminution. To facilitate observation, the tube is divided into a number of equal parts, called degrees.

Further, if we plunge a thermometer ever so often into melting snow or ice, it will always stand at the same point. Hence we learn that *snow* or *ice* always begins to melt at the same temperature.

If we plunge a thermometer repeatedly into water kept boiling, we find that the mercury rises up to a certain point. This is therefore the point at which water always boils, provided the pressure of the atmosphere be the same.

There are four different thermometers used at present in Europe, differing from each other in the number of degrees into which the space between the freezing and boiling points is divided. These are Fahrenheit's, Reaumur's, Celsius's, and Delisle's.

The thermometer uniformly used in Britain, is Fahrenheit's; in this the freezing point is fixed at 32° —the boiling point, at 212° above 0° —or the part at which both the ascending and descending series of numbers commence.

In the thermometer which was first con-

structed by Reaumur, the scale is divided into a smaller number of degrees upon the same length, and contains not more than 80° between the freezing and the boiling points. The freezing point is fixed in this thermometer precisely at 0° , the term between the ascending and the descending series of numbers 100 is the number of the degrees between the freezing and the boiling points in the scale of Celsius; which has been introduced into France, since the revolution, under the name of the Centigrade thermometer; and the freezing point is in this, as in the thermometer of Reaumur, fixed at 0° . One degree on the scale of Fahrenheit, appears, from this account, to be equal to $4\frac{1}{9}$ ths of a degree on that of Reaumur, and to $5\frac{9}{16}$ ths of a degree on that of Celsius.

The space in Delisle's thermometer between the freezing and boiling points is divided into 150° , but the graduation begins at the boiling point, and increases towards the freezing point. The boiling point is marked 0, the freezing point 150° . Hence $180^{\circ} F = 150^{\circ} D$, or $6^{\circ} F = 5^{\circ} D$. To reduce the degrees of Delisle's thermometer under the boiling point to those of Fahrenheit; we have $F = 212 - 6.5 D$; to reduce those *above* the boiling point $F = 212 - 6.5 D$. Upon the knowledge of this proportion it is easy for the student to reduce the degrees of any of these thermometers into the degrees of any other of them.

2. Nature of the Pyrometer.

To measure those higher degrees of heat to which the thermometer cannot be applied, there have been other instruments invented by different philosophers: these are called *pyrometers*. The most celebrated instrument of this kind, and which has been adopted into general use, is that invented by the late ingenious Mr. Wedgwood.

This instrument is also sufficiently simple. It consists of two pieces of brass fixed on a plate, so as to be $\frac{6}{10}$ ths of an inch asunder at one end, and $\frac{3}{10}$ ths at the other; a scale is marked upon them, which is divided into 240 equal parts, each $\frac{1}{10}$ th of an inch; and with this his gauge, are furnished a sufficient number of pieces of baked clay, which must have been prepared in a red heat, and must be of given dimensions. These pieces of clay, thus prepared, are first to be applied cold, to the rule of the gauge, that there may no mistake take place in regard to their dimensions. Then any one of them is to be exposed to the heat which is to be measured, till it shall have been completely penetrated by it. It is then removed and applied to the gauge. The difference between its former and its present dimensions, will shew how much it has shrunk; and will consequently indicate to what degree the

intensity of the heat to which it was exposed, amounted.

High temperatures can thus be ascertained with accuracy. Each degree of Wedgwood's pyrometer is equal to 130° of Fahrenheit's.

Exceptions to the Expansion by Heat.

Philosophers have noticed a few exceptions to the law of heat expanding bodies. For instance; water, when cooled down to the freezing point, instead of contracting on the farther deprivation of heat, actually expands.

Another seeming exception is manifested in alumine, or clay; others occur in the case of cast-iron, and various other metals. Alumine contracts on being heated, and cast-iron, bismuth, &c. when fully fused, are more dense than when solid; for, as soon as they become so, they decrease in density, they expand in the act of cooling, and hence the sharpness of figures upon iron which has been cast in moulds, compared to that of other metals.

Some philosophers have persuaded themselves that these exceptions are only *apparent*, but not really true. They say when water freezes, it assumes a crystalline form, the crystals cross each other and cause numerous vacuities, and thus the ice occupies more space. The same is the case with fused iron, bismuth, and zinc. The contraction of clay is owing to the loss of water, of which it loses a part at every increased degree of temperature hitherto tried; there is therefore a loss of matter; and a reduction of volume must follow.

Mr. Tilloch has published a brief examination of the received doctrines respecting heat and caloric, in which these truths are more fully considered, together with many other interesting facts relative to the received notions of heat.

Equal Distribution of Heat.

If a number of bodies of different temperatures are placed in contact with each other, they will all at a certain time acquire a temperature, which is the mean temperature of the different substances; the caloric of the hottest body will diffuse itself among those which are heated in a less degree, till they have all acquired a certain temperature. Thus, if a bar of iron which has been made red-hot be kept in the open air, it does not retain the heat which it had received, but becomes gradually colder and colder, till it arrives at the temperature of the bodies in its neighbourhood. On the other hand; if we cool down the iron bar by keeping it for some time covered with snow, and then carry it into a warm room, it does not retain its low temperature, but becomes gradually hotter, till it acquires the temperature of the room. It is therefore obvious, that in the one instance the temperature is lowered, and in the other it is raised.

These changes of temperature occupy a

longer or a shorter time, according to the nature of the body, but they always take place at last. This law itself is, indeed, familiar to every one: when we wish to heat a body, we carry it towards the fire; when we wish to cool it, we surround it by cold bodies.

Propagation of Heat.

We have seen, that when bodies of higher temperature than others are brought into contact with each other, the heat is propagated from the first to the second, or the colder body deprives the warmer of its excess of heat.—We shall now see that some bodies do so much more quickly than others. Through some bodies caloric passes with undiminished velocity, through others its passage is prodigiously retarded.

This disposition of bodies of admitting, under equal circumstances, the refrigeration of a heated body within a shorter or a longer time, is called the *power of conducting heat*; and a body is said to be a *better* or *worse conductor of heat*, as it allows the refrigeration to go on quicker or slower. Those bodies, therefore, which possess the property of letting heat pass with facility, are called *good conductors*; those through which it passes with difficulty, are called *bad conductors*, and those through which it does not pass at all, are called *non conductors*: thus we say, in the common language, some bodies are *warm*, or capable of preserving warmth; and from this arises the great difference in the sensation excited by different bodies, when applied at the same temperature to our organs of feeling. Hence, if we immerse our hand in mercury, we feel a greater sensation of cold than when we immerse it in water, and a piece of metal appears to be much colder than a piece of wood, though their temperatures, when examined by means of the thermometer, are precisely the same.

It is probable that all solids conduct heat in some degree, though they differ very much in their conducting power. Metals are the best conductors of heat; but the conducting powers of these substances are by no means equal. Stones seem to be the next best conductors. Glass conducts heat very slowly; wood and charcoal still slower; and feathers, silk, wool, and hair, are still worse conductors than any of the substances yet mentioned.

The best conductors of electricity and galvanism are also the best conductors of heat.

Experiment.—Take a number of straight wires, of equal diameters and lengths, but of different metals; for instance, gold, silver, copper, iron, &c.; cover each of them with a thin coat of wax, or tallow, and plunge their extremities into water, kept boiling, or into melted lead. The melting of the coat of wax will shew that caloric is more quickly transmitted through some metals than others.

It is on this account also, that the end of a glass rod may be kept red-hot for a long time, or even melted, without any inconvenience to the hand which holds the other extremity; though a similar metallic rod, heated in the same manner, would very soon become too hot to be held.

Liquor and Aeriform Bodies convey Heat by an actual Change in the Situation of their Particles.

Count Rumford was the first who proved that fluids in general, and aeriform bodies, convey heat on a different principle from that observed in solids. This opinion is pretty generally admitted, though various ingenious experiments have been made by different philosophers to prove the contrary. In water, for instance, which is, in the strictest sense of the word, a perfect non-conductor of heat, the Count has proved that caloric is propagated only in consequence of the motion which is occasioned in the insulated and solitary particles of that fluid.

All fluids are considered, strictly speaking, in a similar respect as non-conductors of caloric. They can receive it, indeed, from other substances, and can give it to other substances, but no particle can either receive it from or give it to another particle. Before a fluid, therefore, can be heated or cooled, every particle must go individually to the substance from which it receives or to which it gives out caloric. Heat being, therefore, only propagated in fluids, in consequence of the internal motion of their particles, which transport the heat; the more rapid these motions are, the more rapid is the communication of heat. The cause of these motions is the change in the specific gravity of the fluid, occasioned by the change of temperature, and the rapidity is in proportion to the change of the specific gravity of the liquid by any given change of temperature. The following experiment may serve to illustrate this theory.

Take a thin glass tube, eight or ten inches long, and about an inch in diameter. Pour into the bottom part, for about the depth of one inch, a little water coloured with Brazil-wood, or litmus, and then fill up the tube with common water, extremely gently, so as to keep the two strata quite distinct from each other. Having done this, heat the bottom part of the tube over a lamp; the coloured infusion will then ascend, and gradually tinge the whole fluid; the water in the upper part of the tube may be made to boil, but the colouring matter will remain at the bottom undisturbed. The heat cannot act downwards to make it ascend.

By thus being able to make the upper part of a fluid boil without heating the bottom part, water may be kept boiling

for a considerable time in a glass tube over ice, without melting it.

Other experiments, illustrating the same principle, may be found in Count Rumford's excellent *Essays*, especially in *Essay the 7th*; 1797.

To this indefatigable philosopher we are wholly indebted for the above facts: he was the first who taught us that air and water were nearly non-conductors. The results of his experiments, which are contained in the above *Essay*, are highly interesting; they also shew that the conducting power of fluids is impaired by the admixture of fibrous and glutinous matter.

Count Rumford proved that ice melted more than 80 times slower, when boiling-hot water stood on its surface, than when the ice was placed to swim on the surface of the hot water. Other experiments shewed that water, only eight degrees of Fahrenheit above the freezing point, or at the temperature of forty degrees, melts as much ice, in any given time, as an equal volume of that fluid at any higher temperature, provided the water stands on the surface of the ice. Water, at the temperature of 41°, is found to melt more ice, when standing on its surface, than boiling water.

It becomes further evident, from the Count's ingenious experiments, that of the different substances used in clothing, hares' fur and gilder-down are the warmest; next to these, beavers' fur, raw silk, sheep's wool, cotton wool, and lastly, lint, or the scrapings of fine linen. In fur, the air interposed among its particles is so engaged as not to be driven away by the heat communicated thereto by the animal body; not being easily displaced, it becomes a barrier to defend the animal body from the external cold. Hence it is obvious that those skins are warmest which have the finest, longest, and thickest fur; and that the furs of the beaver, otter, and other like quadrupeds, which live much in the water, and the feathers of water-fowl, are capable of confining the heat of those animals in winter, notwithstanding the coldness of the water which they frequent. Bears, and various other animals, inhabitants of cold climates, which do not often take the water, have their fur much thicker on their backs than on their bellies.

The snow which covers the surface of the earth in winter, in high latitudes, is doubtless designed as a garment to defend it against the piercing winds from the polar regions, which prevail during the cold season.

Without dwelling farther upon the philosophy of this truth, we must briefly remark that the happy application of this law, satisfactorily elucidates some of the most interesting facts of the economy of nature.

Theory of Caloric of Fluidity, or Latent Heat.

There are some bodies which, when submitted to the action of caloric, dilate to such a degree, and the power of aggregation subsisting among their particles is so much destroyed and removed to such a distance by the interposition of caloric, that they slide over each other in every direction, and therefore appear in a fluid state. This phenomenon is called *fusion*. Bodies thus rendered fluid by means of caloric are said to be *fused*, or *melted*; and those that are subject to it, are called *fusible*.

The greater number of solid bodies may, by the application of heat, be converted into fluids. Thus metals may be fused; sulphur, resin, phosphorus, may be melted; ice may be converted into water; &c.

Those bodies which cannot be rendered fluid by any degree of heat hitherto known, are called *fixed*, or *infusible*.

If the effects of heat under certain circumstances, be carried still further than is necessary to render bodies fluid, vaporization begins; the bodies then become converted into the vaporous or *gaseous state*. Vaporization, however does not always require a previous fusion. Some bodies are capable of being converted into the vaporous state, without previously becoming fluid, and others cannot be volatilized at any temperature hitherto known.

Fluidity is therefore by no means essential to any species of matter, but always depends on the presence of a quantity of caloric. Solidity is the natural state of all bodies, and there can be no doubt that every fluid is capable of being rendered solid by a due reduction of temperature; and every solid may be fused by the agency of caloric, if the latter does not decompose them at a temperature inferior to that which would be necessary for their fusion.

Caloric of Fluidity.

Dr. Black was the first who proved that, whenever caloric combines with a solid body, the body becomes heated only, until it is rendered fluid: or that, whenever it has acquired the fluid state, its temperature remains stationary, though caloric is continued to be added to it. The same is the case when fluids are converted into the æriform or vaporous state.

From these facts, the laws of latent heat have been inferred. The theory may be illustrated by means of the following experiments.

If a lump of ice, at a low temperature, suppose at 22°, be brought into a warm

room, it will become gradually less cold, as may be discovered by means of the thermometer. After a very short time, it will reach the temperature of 32°, (the freezing point); but there it stops. The ice then begins to melt; but the process goes on very slowly. During the whole of that time its temperature continues at 32°; and as it is constantly surrounded by warm air, we have reason to believe that caloric is constantly entering into it; yet it does not become hotter till it is changed into water. Ice, therefore, is converted into water by a quantity of caloric uniting with it.

It has been found by calculation, that one pound of ice in melting absorbs 140° of caloric, the temperature of the water produced still remaining at 32°.

This fact may be proved in a direct manner.

Take one pound of ice, at 32 degrees reduced to a coarse powder; put it into a wooden bowl, and pour over it gradually, one pound of water, heated to 172 deg.; all the ice will become melted, and the temperature of the whole fluid, if examined by a thermometer, will be 32 deg.; 140 deg. of caloric are therefore lost, and it is this quantity which was requisite to convert the ice into water.

This caloric has been called *latent caloric*, or rather caloric of fluidity, because its presence is not measurable by the thermometer.

Dr. Black has also ascertained, by experiment, that the fluidity of melted wax, tallow, spermaceti, metals, &c. is owing to the same cause; and Ladrani proved that this is the case with sulphur, alum, nitrate of potash, &c.

We consider it, therefore, as a general law, that whenever a solid is converted into a fluid, it combines with caloric, and that is the cause of fluidity.

On the sudden transition of solids into fluids, is founded the well known *Production of Artificial Cold, by Means of Frigorific Mixtures.*

A number of experiments have been lately made by different philosophers, in order to produce artificial cold. And as these methods are often employed in chymistry, with a view to expose bodies to the influence of very low temperatures, we shall enumerate the different substances which may be made use of for that purpose, and the degrees of cold which they are capable of producing. We are indebted for them to Pepys, Walker, and Lowitz.

A TABLE OF FREEZING MIXTURES.

<i>Mixtures.</i>		<i>Thermometer Sinks</i>	
Muriate of ammonia	5 parts	From 50° to 10°	
Nitrate of potash	5		
Water	16		
Muriate of ammonia	5	From 50° to 4°	
Nitrate of potash	5		
Sulphate of soda	8		
Water	16		
Sulphate of soda	3 parts	From 50° to 3°	
Diluted nitric acid	2		
Sulphate of soda	8 parts	From 50° to 0°	
Muriatic acid	5		
Snow	1 part	From 32° to 0°.	
Muriate of soda	1		
Snow, or pounded ice	2 parts	From 0° to —5°.	
Muriate of soda	1		
Snow, or pounded ice	1 part	From —5° to —18°.	
Muriate of soda	5		
Muriate of ammonia and			
Nitrate of potash	5		
Snow, or pounded ice	12 parts	From —18° to —25°.	
Muriate of soda	5		
Nitrate of ammonia	5		
Snow, and		From 0° to —46°.	
Diluted nitric acid			
Muriate of lime	3 parts	From 32° to —50°.	
Snow	2		
Potash	4 parts	From 32° to —51°.	
Snow	3		
Snow	2 parts	From —10° to —56°.	
Diluted sulphuric acid	1		
Diluted nitric acid	1		
Snow	1 part	From 20° to —60°.	
Diluted sulphuric acid	1		
Muriate of lime	2 parts	From 0° to —66°.	
Snow	1		
Muriate of lime	3 parts	From —40° to —73°.	
Snow	1		
Diluted sulphuric acid	10 parts	From —68° to —91°.	
Snow	8		
Nitrate of ammonia	1 part	From 50° to 4°.	
Water	1		
Nitrate of ammonia	1 part	From 50° to 3°.	
Carbonate of soda	1		
Water	1		
Sulphate of soda	6 parts	From 50° to 10°.	
Muriate of ammonia	4		
Nitrate of potash	2		
Diluted nitric acid	4	From 50° to 14°.	
Sulphate of soda	6 parts		
Nitrate of ammonia	5	From 50° to 12°.	
Diluted nitric acid	4		
Phosphate of soda	9 parts	From 50° to 21°.	
Diluted nitric acid	4		
Phosphate of soda	9 parts		
Nitrate of ammonia	6	From 50° to 3°.	
Diluted nitric acid	4		
Sulphate of soda	5 parts		
Diluted sulphuric acid	4		

Management of the preceding Mixtures for producing Cold.

To produce the effects before stated, the salts must be reduced to powder, and contain their full quantity of water of crystallization. The vessel in which the freezing mixture is made should be very thin, and just large enough to hold it, and the materials should be mixed together as expeditiously as possible, taking care to stir the mixture at the same time with a rod of glass or wood.

In order to obtain the full effect, the materials ought to be first cooled to the temperature marked in the table, by introducing them into some of the other frigorific mixtures, and then mingling them together in a similar mixture. If, for instance, we wish to produce -46° , the snow and diluted nitric acid ought to be cooled down to 0° , by putting the vessel which contains each of them into the fifth freezing mixture in the above table, before they are mingled together. If a more intense cold be required, the materials to produce it are to be brought to the proper temperature by being previously placed in the second freezing mixture.

This process is to be continued till the required degree of cold has been procured.

Conversion of Solids and Fluids into the Aëriform or Gaseous State.

We have seen before, that, in order to render solids fluid, a certain quantity of caloric is necessary, which combines with the body, and therefore cannot be measured by the thermometer; we shall now endeavour to prove, that the same holds good in respect to the conversion of solids or fluids into the vaporous or gaseous state.

Take a small quantity of carbonate of ammonia, introduce it into a retort, the neck of which is directed under a cylinder filled with mercury and inverted in a basin of the same fluid. On applying heat to the body of the retort, the carbonate of ammonia will be volatilized, it will expel the mercury out of the cylinder, and become an invisible gas, and would remain so, if its temperature was not lowered.

The same is the case with benzoic acid, camphire, and various other substances.

All fluids may by the application of heat be converted into an aëriform elastic state.

When we consider water in a boiling state, we find that this fluid, when examined by the thermometer, is not hotter, after boiling several hours, than when it began to boil, though to maintain it boiling, a brisk fire must necessarily be kept up. What then, we may ask, becomes of the wasted caloric? It is not perceptible in the water, nor is it manifested by the steam; for the steam, if not compressed,

upon examination is found not to be hotter than boiling water. The caloric is therefore absorbed by the steam, and although what is so absorbed is absolutely necessary for the conversion of water into the form of steam, it does not increase its temperature, and is therefore not appreciable by the thermometer.

This conclusion is further strengthened by the heat given out by steam on its being condensed by cold. This is particularly manifested in the condensation of this fluid in the process of distilling, where, upon examining the refrigerator, it will be found that a much greater quantity of caloric is communicated to it, than could possibly have been transmitted by the caloric which was sensibly acting before the condensation. This may be easily ascertained by observing the quantity of caloric communicated to the water in the refrigerator of a still, by any given quantity of liquid that passes over.

1. The boiling points of different fluids are influenced by atmospheric pressure.

The boiling point, or the conversion of fluids into gases, always takes place at certain temperatures, which is different in different fluids, provided the pressure of the atmosphere be the same.

Put any quantity of sulphuric ether into a Florence flask, suspend a thermometer in it, and hold the flask over an Argand's lamp, the ether will immediately begin to boil, and the thermometer will indicate 98° , if the ether has been highly rectified.

If highly rectified ardent spirit is heated in a similar manner, the thermometer will rise to 176° , and there remain stationary.

If water is substituted, it will rise to 212° .

If strong nitrous acid of commerce be made use of, it will be found to boil at 248° ;—sulphuric acid at 546° ;—and mercury and linseed-oil at 600° , &c.

2. The boiling point of fluids is retarded by pressure.

Mr. Watt heated water under a strong pressure to 400° . Yet still, when the pressure was removed, only part of the water was converted into vapour, and the temperature of this vapour, as well as that of the remaining fluid, was no more than 212° . There was therefore 188° of caloric suddenly lost. This caloric was carried off by the steam. Now as only about one-fifth of the water was converted into steam, that steam must contain not only its own 188° , but also the 188° lost by each of the other four parts; that is to say, it must contain 188×5 , or about 940° . Steam, therefore, is water combined with at least 940° of caloric, the presence of which is not indicated by the thermometer.

3. When pressure is removed from the surface of bodies, their conversion into the gaseous state is greatly facilitated, or their boiling point is lowered.

In proof of this the following experiments may serve:

Let a small bottle be filled with highly rectified sulphuric ether, and a piece of wetted bladder be tied over its orifice around its neck. Transfer it under the receiver of an air-pump, and take away the super-incumbent pressure of the air in the receiver. When the exhaustion is complete, pierce the bladder by means of a pointed sliding wire, passing through a collar of leather which covers the upper opening of the receiver. Having done this, the ether will instantly begin to boil, and become converted into an invisible gaseous fluid.

Take a small retort or Florence flask, fill it one half, or less, with water, and make it boil over a lamp; when kept briskly boiling, for about five minutes, cork the mouth of the retort as expeditiously as possible, and remove it from the lamp.

The water, on being removed from the source of heat, will keep boiling for a few minutes, and when the ebullition begins to slacken, it may be renewed by dipping the retort into cold water.

The water during boiling becomes converted into vapour; this vapour expels the air of the vessel, and occupies its place; on diminishing the heat, it condenses; when the retort is stopped, a partial vacuum is formed; the pressure becomes diminished, and a less degree of heat is sufficient to cause an ebullition.

For the same reason water may be made to boil under the exhausted receiver at 94° Fahr. or even at a much lower degree; alcohol at 56°, and ether at—20°.

On the conversion of fluids into gases is founded the following experiment, by which water is frozen by means of sulphuric ether.

Take a thin glass tube, four or five inches long, and about two or three-eighths of an inch in diameter, and a two-ounce bottle furnished with a capillary tube fitted to its neck. In order to make ice, pour a little water into the tube, taking care not to wet the outside, nor to leave it moist. Having done this, let a stream of sulphuric ether, fall through the capillary tube upon that part of it containing the water, which by this means will be converted into ice in a few minutes, and this it will do even near a fire, or in the midst of summer.

If the glass tube containing the water be exposed to the brisk thorough air, or free draught of an open window, a large quantity of water may be frozen in a shorter time; and if a thin spiral wire be introduced previous to the congelation of the

water, the ice will adhere to it, and may thus be drawn out conveniently.

A person might be easily frozen to death during very warm weather, by merely pouring upon his body, for some time, sulphuric ether, and keeping him exposed to a thorough draught of air.

Artificial Refrigeration.

The cooling or refrigeration of rooms in the summer season by sprinkling them with water, becomes likewise obvious on this account.

The method of making ice artificially in the East-Indies depends on the same principle. The ice-makers at Benares dig pits in large open plains, the bottom of which they strew with sugar-canes or dried stems of maize or Indian corn. Upon this bed they place a number of unglazed pans, made of so porous an earth that the water penetrates through their whole substance. These pans are filled towards evening in the winter season with water that has boiled, and left in that situation till morning, when more or less ice is found in them, according to the temperature of the air; there being more formed in dry and warm weather, than in that which is cloudy, though it may be colder to the human body.

Every thing in this process is calculated to produce cold by evaporation; the beds on which the pans are placed suffer the air to have free passage to their bottoms; and the pans, constantly oozing out water to their external surface, are cooled by the evaporation of it.

In Spain, they use a kind of earthen jars, called *buxaros*, which are only half-baked, the earth of which is so porous, that the outside is kept moist by the water which filters through it, and though placed in the sun, the water in the jar becomes as cold as ice.

It is a common practice in China to cool wine or other liquors by wrapping the bottle in a wet cloth, and hanging it up in the sun. The water in the cloth becomes converted into vapour, and thus cold is produced.

The Blacks in Senegambia have a similar method of cooling water by filling tanned leather bags with it, which they hang up in the sun; the water oozes more or less through the leather, so as to keep the outward surface wet, which by its quick and continued evaporation cools the water remarkably.

The winds on the borders of the Persian Gulph are often so scorching, that travellers are suddenly suffocated, unless they cover their heads with a wet cloth; if this be too wet, they immediately feel an intolerable cold, which would prove fatal if the moisture was not speedily dissipated by the heat.

If a cold vessel is brought into a warm room, particularly where many people are

assembled, the outside of it will soon become covered with a sort of dew.

Before some changes of weather, the stone pavements, the walls of a house, the balustrades of staircases and other solid objects, feel clammy and damp.

In frosty nights, when the air abroad is colder than the air within, the dampness of this air, for the same reason, settles on the glass panes of the windows, and is there frozen into curious and beautiful figures.

Thus *fogs* and *dews* take place, and in the higher regions *clouds* are formed from the condensed vapour. The still greater condensation produces *mists* and *rain*.

Capacity of Bodies for containing Heat.

The property which different bodies possess, of containing at the same temperature, and in equal quantities, either of mass or bulk, unequal quantities of heat, is called their capacity for heat. The capacities of bodies for heat are therefore considered as great or small in the ratio as their temperatures are either raised or lowered by the addition, or diminished by the deprivation of equal quantities of heat.

In homogeneous bodies, the quantities of caloric which they contain are in the ratio of their temperature and quantity of mass; for instance, when equal quantities of water, oil, or mercury, of unequal temperatures, are mingled together, the temperature of the whole will be the *arithmetical* mean between the temperatures of the two quantities that had been mixed together. It is a self-evident truth that this should be the case, for the particles of different portions of the same substance being alike, their effects must be equal. For instance:

Mix a pound of water at 172° , with a pound at 32° , half the excess of heat in the hot water will quit it to go over into the colder portion; thus the hot water will be cooled 70° , and the cold will receive 70° of temperature; therefore $172 - 70$, or $32 + 70 = 102$ will give the heat of the mixture. To attain the arithmetical mean very exactly, several precautions however are necessary.

When heterogeneous bodies of different temperatures are mixed together, the temperature produced is never the arithmetical mean of the two original temperatures.

In order to ascertain the comparative quantities of heat of different bodies, equal weights of them are mingled together; the experiments for this purpose being in general more easily executed than those by which they are compared from equal bulks.

Thus, if one pound of mercury heated to 110° Fahr. be added to one pound of water of 44° , the temperature of the blended fluids will not be changed to 77° , as it would be if the surplus of heat were divided among those fluids in the

proportion of their quantities. It will be found on examination to be only 47° .

On the contrary; if the pound of mercury be heated to 44° and the water to 110° , then on stirring them together the common temperature will be 107° .

Hence if the quicksilver loses by this distribution 63° of caloric, an equal weight of water gains only 3° from this loss of 63° of heat. And on the contrary, if the water loses 3° , the mercury gains 63° .

When, instead of comparing the quantities of caloric which equal *weights* of different bodies contain, we compare the quantities contained in equal *volumes*, we still find that the same difference takes place. Thus it is found by experiment, that the quantity of caloric necessary to raise the temperature of a given volume of water any number of degrees, is, to that necessary to raise an equal volume of mercury, the same number of degrees as 2 to 1. This is therefore the proportion between the comparative quantities of caloric which these two bodies contain, estimated by their volumes; and similar differences exist with respect to every other kind of matter.

From the nature of the experiments by which the quantities of caloric which bodies contain are ascertained, it is evident that we discover merely the *comparative*, not the *absolute* quantities. Hence water has been chosen as a standard, to which other bodies may be referred; its capacity is stated as the arbitrary term of 1000, and with this the capacities of other bodies are compared.

It need not be told that pains have been taken to estimate on these experiments that portion of heat which diffuses itself into the air or into the vessel where the mercury and water are blended together. As however such valuations cannot be made with complete accuracy, the numbers stated above are only an approximation to truth.

Radiation of Caloric.

Caloric is thrown off or radiates from heated bodies in right lines, and moves through space with inconceivable velocity. It is retarded in its passage by atmospheric air, by colourless fluids, glass, and other transparent bodies.

If a glass mirror be placed before a fire, the mirror transmits the rays of light, but not the rays of heat.

If a plate of glass, talc, or a glass vessel filled with water be suddenly interposed between the fire and the eye, the rays of light pass through it, but the rays of caloric are considerably retarded in its passage; for no heat is perceived until the interposed substance is saturated with heat, or has reached its *maximum*. It then ceases to intercept the rays of caloric, and allows

them to pass as freely as the rays of light.

It has been lately shewn by Dr. Herschel that the rays of caloric are refrangible, but less so than the rays of light; and the same philosopher has also proved by experiment, that it is not only the rays of caloric emitted by the sun, which are refrangible, but likewise the rays emitted by common fires, by candles, by heated iron, and even by hot water.

Whether the rays of caloric are differently refracted, in different mediums, has not yet been ascertained. We are certain, however, that they are refracted by all transparent bodies which have been employed as burning glasses.

The rays of caloric are also reflected by polished surfaces, in the same manner as the rays of light.

This was long ago noticed by Lambert, Suessure, Sheele, Pictet, and lately by Dr. Herschel.

Professor Pictet placed two concave metallic mirrors opposite to each other at the distance of about twelve feet. When a hot body, an iron bullet for instance, was placed in the focus of the one, and a mercurial thermometer in that of the other, a substance radiated from the bullet; it passed with incalculable velocity through the air, it was reflected from the mirrors, it became concentrated, and influenced the thermometer placed in the focus, according to the degree of its concentration.

An iron ball, two inches in diameter, heated so that it was not luminous in the dark, raised the thermometer not less than ten and a half degrees of Reaumur's scale, in six minutes.

A lighted candle occasioned a rise in the thermometer nearly the same.

A Florence flask, containing two ounces and three drachms of boiling water, raised Fahrenheit's thermometer three degrees. He blackened the bulb of his thermometer, and found that it was more speedily influenced by the radiation than before, and that it rose to a greater height.

M. Pictet discovered another very singular fact; namely, the *apparent radiation of cold*. When, instead of a heated body, a Florence flask full of ice or snow is placed in the focus of one of the mirrors, the thermometer placed in the focus of the other immediately descends, and ascends again whenever the cold body is removed.

This phenomenon may be explained on the supposition, that from every body at every temperature caloric radiates, but in less quantity as the temperature is low; so that in the above experiment, the thermometer gives out more caloric by radiation, than it receives from the body in the opposite focus, and therefore its temperature is lowered. Or, as Pictet has supposed,

when a number of bodies near to each other have the same temperature, there is no radiation of caloric, because in all of them it exists in a state of equal tension; but as soon as a body at an inferior temperature is introduced, the balance of tension is broken, and caloric begins to radiate from all of them, till the temperature of that body is raised to an equality with theirs. In the above experiment, therefore, the placing the snow or ice in the focus of the mirror causes the radiation of caloric from the thermometer, and hence the diminution of temperature which it suffers.

These experiments have been lately repeated by Dr. Young and Professor Davy, at the theatre of the Royal Institution. These gentlemen inflamed phosphorus by reflected caloric; and proved that the heat thus excited was very sensible to the organs of feeling.

It is, therefore evident that caloric is thrown off from bodies in rays which are invisible, or incapable of exciting vision, but which are capable of exciting heat.

These invisible rays of caloric are propagated in right lines, with extreme velocity, and are capable of the laws of reflection and refraction.

The heating agency however is different in the different coloured rays of the prismatic spectrum. According to Dr. Herschel's experiments, it follows inversely the order of the refrangibility of the rays of light. The least refrangible possessing it in the greatest degree.

Sir Henry Englefield has lately made a series of experiments on the same subject, from which we learn that a thermometer, having its ball blackened, rose when placed in the *blue* ray of the prismatic spectrum in 3' from 55° to 56°; in the *green*, in 3' from 45° to 58°; in the *yellow*, in 3' from 56° to 62°; in the *full red*, in 2 1-2' from 56° to 72°; in the *confines of the red*, in 2' from 58° to 73 10-20; and *quite out of the visible light*, in 2' from 61° to 79°.

Between each of the observations, the thermometer was placed in the shade so long, as to sink it below the heat to which it had risen in the preceding observation, of course its rise above that point could only be the effect of the ray to which it was exposed. It was continued in the focus long after it had ceased to rise; therefore the heats given are the greatest effects of the several rays on the thermometer in each observation. A thermometer placed constantly in the shade near the apparatus, was found scarcely to vary during the experiments.

Sir Henry made other experiments with thermometers with naked balls, and with others whose balls were painted white, for which we refer the reader to the interesting

paper of the Baronet, from which the above experiments are transcribed.

The coloured rays emitted from the sun and combustible bodies, since they excite heat and vision, must consist of a mixture of heat-making rays, and rays of light.

And as the rays of heat and light accompany each other when emitted from luminous bodies, the velocity with which the rays of caloric move must be equal to that of light, and hence its particles must be equally minute. They differ however in this particular, that the rays of light produce the sensation of vision, and possess certain chemical properties, whilst in those of caloric the peculiar agency of heat resides.

CALORIMETER. An instrument by which the whole quantity of absolute heat existing in a body in chemical union can be ascertained.

CALTHA. (Καλθε, corrupted from καλχα, yellow, from whence, says Vossius, come calthula, caldula, caledula, calendula.) Marsh marigold.

1. The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Polygynia*.

2. The pharmacopœial name of the herb marigold, so called from its colour. See *Calendula arvensis*.

CALTHA PALUSTRIS. The marsh marigold. The young buds of this plant make, when properly pickled, very good substitutes for capers.

CALTHA VULGARIS. See *Calendula arvensis*.

CALTHULA. The caltha is so called.

CALTROPS. A name of the *Trapa natans* of Linnæus, whose fruit is said to be nutritious and demulcent, and to be useful in diarrhœas from abraded bowels, and against calculus.

CALUMBA. The name now adopted by the London college of physicians for the columbo. See *Columbo*.

CALVA. (From *calvus*, bald.) The scalp or upper part of the cranium or top of the head; so called because it often grows bald first.

CALVARIA. (From *calvus*, bald.) The upper part of the cranium which becomes soon bald. It means all above the orbits, temples, ears and occipital eminence.

CALVITIA. (From *calvus*, bald.) *Calvitium*. Baldness; want or loss of hair, particularly upon the sinciput.

CALX. (Καλακ, to burn. Arab.)

1. Chalk. Limestone.

2. Lime. *Calx viva*. The London College direct it to be prepared thus: Take of limestone one pound. Break it into small pieces and heat it in a crucible in a strong fire for an hour, or until the carbonic acid is entirely driven off, so that on the addition of acetic acid, no bubbles of gas shall be extricated. Lime may be made by the same process from shells previously washed in

boiling water, and cleared from extraneous matters. See *Lime*.

CALX ANTIMONII. See *Oxydum antimoni*.

CALX OUM KALI FURO. The preparation formerly called by this name is now termed, in the London pharmacopœia, potassæ eum calce.

CALX HYDRARGYRI ALBA. See *hydrargyri præcipitatus albus*.

CALX VIVA. See *Calx*.

CALYPTER. (From καλυπτω, to hide.) A carnicous excrescence covering the hemorrhoidal vein.

CAMARA. (καμαρα, a vault.) *Camarium*. The fornix of the brain: also the vaulted part of the auricle of the heart.

CAMARIUM. (From Καμαρα, a vault.) See *Camara*.

CAMAROMA. (From καμαρα, a vault.) *Camarois*, *Camaratia*. A fracture of the skull, in the shape of an arch or vault.

CAMBING. A tree of the Molucca islands, whose bark has been recommended in dysenteries.

CAMBIREA. So Paracelsus calls the venereal bubo.

CAMBIVM. (From *cambio*, to exchange.) That nutritious humour which is changed into the matter of which the body is composed.

CAMBODIA. See *Gambogia*.

CAMBOGIA GUTTA. See *Gambogia*.

CAMBOGIUM. (From the province of Cambodia, whence it was brought.) See *Gambogia*.

CAMBRO-BRITANNICA. See *Charsamorius*.

CAMBUCA. *Cambuta membrata*. So Paracelsus calls the venereal cancer. Also by some it is described as a bubo, an ulcer, an abscess on the pudenda: also a boil in the groin.

CAMBU. The wild American myrtle of Piso and Margrave, which is said to be astringent.

Camel's hay. See *Fucus odoratus*.

CAMERA. The chambers or cavities of the eye are termed camera.

CAMERATIO. See *Camaroma*.

CAMES. *Camet*. Silver.

CANIERA. See *Canella alba*.

CANINUS. A furnace and its chimney. In Rulandus it signifies a bell.

CANISIA FORTUS. (From the Arabic term *kandash*, an under garment.) The shirt of the fatus. It is frequently put for the chorion;

Camomille. See *Chamæmelum*.

Camomille, stinking. See *Cotula fetida*.

CANOMILLA. Corrupted from *chamæmelum*.

CANORUM. (καμνορον, quia homines, naxo moras, perimat; because, if eaten, it brings men to a miserable end.) A species of monkhood. See *Aconitum*.

CAMPANA. A bell. In Chemistry, a receptacle like a bell, for making sulphuric acid; thus the oleum sulphuris per campanam.

CAMPANULA. (From *Campana*, a bell, named from its shape.) The bell-flower.

The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

CAMPE. (From *καμπη*, to bend.) A flexure or bending. It is also used for the ham, and a joint, or articulation.

Campeachy wood. See *Lignum campechense*.

CAMPICHENSE LIGNUM. See *Lignum Campechense*.

Camphire. See *Camphora*.

Camphor. See *Camphora*.

CAMPHORA. (*Camphura*, Arab. The ancients by camphor meant what now is called asphaltum, or Jews' pitch; *καμψη*.) *Camphura*. *Caf.* *Cafar.* *Ligatura veneris.* *Caphora*. *Capur.* *Alkhor.* *Alteor.* *Camphire.* *Camphor.* A peculiar concrete substance prepared by distillation from the *Laurus camphora* of Linnæus;—*foliis triplinerviis lanceolato-ovatis*, a tree indigenous to Japan, where it grows abundantly. The camphire is found to lodge every where in the interstices of the fibres of the wood, pith, and knots of the tree. The crude camphire, exported from Japan, appears in small greyish pieces, and is intermixed with various extraneous matters; in this state it is received by the Dutch, and purified by a second sublimation; it is then formed into loaves, in which state it is sent to England. When pure, it is white, pellucid, somewhat unctuous to the touch; of a bitterish, aromatic, acid taste, yet accompanied with a sense of coolness; of a fragrant smell, and approaching to that of rosemary, but much stronger. It is totally volatile and inflammable, soluble in vinous spirits, oils, and the mineral acids; not in water, fixed nor volatile alkaline liquors, nor in acids of the vegetable kingdom. The use of this important medicine, in different diseases, is very considerable. It has been much employed, with great advantage, in fevers of all kinds, particularly in nervous fevers attended with delirium and much watchfulness. The experienced Werlhoff has witnessed its utility in several inflammatory diseases, and speaks highly in favour of its refrigerant qualities. The benefit derived from it in putrid fevers, where bark and acids are contra-indicated, is remarkable. In spasmodic and convulsive affections it is also of much service, and even in epilepsy. In chronic diseases this medicine is likewise employed; and against rheumatism, arthritis, and mania, we have several accounts of its efficacy. Nor is it less efficacious when applied externally in certain diseases: it dissipates inflammatory tumours in a short time; and its antiseptic quality, in resisting and during gangrene, is very considerable. Another property peculiar to this medicine must not, however, be omitted; the power it possesses of obviating the stranguy that is produced by cantharides, when sprinkled over a blister.

The preparations of camphor are, *spiritus camphoratus*, *oleum camphoratum*, *linimentum camphoræ*, *tinctura opii camphorata*, and the *mustura camphorata*. Camphor dissolved in acetic acid, with some essential oils, forms the aromatic vinegar.

CAMPHORÆ FLORES. The subtile substance which first ascends in subliming camphor. It is nothing more than the camphor.

CAMPHORÆ FLORES COMPOSITI. Camphor sublimed with gum benzoin.

CAMPHOROSMA. (From *camphora*, and *osma*, to smell; so called from its smelling of camphire.) The camphor-smelling plant.

1. The name of a genus of plants in the Linnæan system. Class, *Tetrandria*. Order, *Monogynia*.

2. The pharmacopœial name of the camphorala. See *Camphorala*.

CAMPHORASMA. (From *camphora*: so called from its camphor-like smell.) Balm of Gilend. See *Moldavica*.

CAMPHORATA. *Chamepeuce.* *Camphorata hirsuta.* *Camphorasma Monspelica.* Stinking ground-pine. This plant, *Camphorosma Monspelienis* of Linnæus;—*foliis hirsutis linearibus*, took its name from its smell resembling so strongly that of camphor: it has been exhibited internally, in form of decoction, in dropsical and asthmatic complaints, and by some is esteemed in fomentations against pain. It is rarely or ever used in modern practice.

CAMPHORAS. A salt formed by the union of the camphoric acid with different bases: thus, *camphorat of allumin*, *camphorat of ammoniac*, &c.

CAMPHORATUM OLEUM. A mixture of olive oil, two parts, with one of camphor: of use in inflammatory swellings of the throat, if mixed with a proper cataplasm, and applied to it. In ascites, when the abdomen is much distended, if rubbed on freely every night and morning, it is supposed to be useful.

CAMPHORIC ACID. *Acidum camphoricum.* If nitric acid be distilled several times (six or eight) from camphor, a crystallized salt is obtained, called the acid of camphor, which reddens syrup of violets and the tincture of turnsole. Its taste is bitter, and it differs from oxalic acid in not precipitating lime from the muriatic acid. The union of this acid with different bases forms what is called *camphorates*, none of which have yet been used medicinally.

CAMPHOROSMA MONSPELIENSIS. The systematic name of the plant called camphorata in the pharmacopœias. See *Camphorata*.

CAMPTER. (From *καμπη*, to bend.) An inflexion or incurvation.

CAMPULUM. (From *καμπη*, to twist about.) A distortion of the eyelids or other parts.

CAMPULOTIS. (From *καμπη*, bent.

A preternatural incurvation, or recurvation of a part. A distortion of the eyelids.

CAMPYLUM. See *Campylotis*.

CANABIL. A sort of medicinal earth.

CANABINA AQUATICA. See *Bidens*.

CANABIS INDICA. See *Banque* and *Cannabis*.

Canada Balsam. See *Balsamum Canadense*.

CANABIS PEREGRINA. See *Cannabis*.

CANADENSIS. (Brought from *Canada*.) A name of the copaiva and other balsams.

CANALES SEMICIRCULARES. Three semicircular canals placed in the posterior part of the labyrinth of the ear. They open by five orifices into the vestibulum. See *Ear*.

CANALICULUS. (Dim. of *canalis*, a channel.) That blood-vessel, which in a fetus is situated between the pulmonary artery and the aorta, but in the adult is extinct, is called the canaliculus arteriosus. The same as *canalis arteriosus*.

CANALIS. (From *χωρος*, an aperture, or rather from *canna*, a reed.) A Canal. A hollow round instrument like a reed, for embracing and holding a broken limb. The hollow of the spine. Also it is specifically applied to many parts of the body; as *canalis venosus*.

CANALIS ARTERIOSUS. *Canalis Botalli.* A blood-vessel peculiar to the fetus, disappearing after birth; through which the blood passes from the pulmonary artery into the aorta.

CANALIS NASALIS. A canal going from the internal canthus of the eye downwards into the nose: it is situated in the superior maxillary bone, and is lined with the pituitary membrane continued from the nose.

CANALIS PETITIANUS. A triangular cavity, naturally containing a moisture, between the two laminae of the hyaloid membrane of the eye, in the anterior part, formed by the separation of the anterior lamina from the posterior. It is named after its discoverer, M. Petit.

CANALIS SEMISPETROS. The half bony canal of the ear.

CANALIS VENOSUS. A canal peculiar to the fetus, disappearing after birth, that conveys the maternal blood from the *porta* of the liver to the ascending *vena cava*.

Canary balm. See *Melissa Turcica*.

CANCANUM GRÆCORUM. See *Courbaril*.

CANCELLI. Lattice-work; generally applied to the reticular substance in bones.

CANCELLUS. (From *cancer*, a crab.) The wrong heir. Bernard the hermit. A species of cray-fish supposed to cure rheumatism, if rubbed on the part.

CANCER. (From *καρκινος*, a crab; so called by the ancients, because it exhibited large blue veins like crab's claws.)

1. The name of a disease likewise called *Carcinoma*, *carcinos* by the Greeks, *Lupus* by the Romans, because it eats away the flesh like a wolf. Dr. Cullen places this genus of disease in the class *locales*, and order *tumores*. He defines it a painful scirrhus tumour, terminating in a fatal ulcer. Any part of the body may be the seat of cancer, though the glands are most subject to it. It is distinguished according to its stages into *occult* and *open*; by the former is meant its scirrhus state, which is a hard tumour that sometimes remains in a quiet state for many years. When the cancerous action commences in it, it is attended with frequent shooting pains: the skin that covers it becomes discoloured, and ulceration sooner or later takes place; when the disease is denominated *open* cancer. Mr. Pearson says, "When a malignant scirrhus or a warty excrescence, hath proceeded to a period of ulceration, attended with a constant sense of ardent and occasionally shooting pains, is irregular in its figure, and presents an unequal surface; if it discharges sordid, sanious or fetid matter; if the edges of the sore be thick, indurated, and often exquisitely painful, sometimes inversed, at other times retorted, and exhibit a serrated appearance; and should the ulcer in its progress be frequently attended with hæmorrhage, in consequence of the erosion of blood-vessels; there will be little hazard of mistake in calling it a cancerous ulcer." In men, a cancer most frequently seizes the tongue, mouth, or penis; in women, the breasts, or the uterus, particularly about the cessation of their periodical discharges; and in children, in the eyes. The following description of Scirrhus and Cancer, from the above writer, will serve to elucidate the subject. A hard unequal tumour that is indolent and without any discoloration in the skin is called a scirrhus; but when an itching is perceived in it, which is followed by a pricking, shooting or lancinating pain, and a change of colour in the skin, it is usually denominated a cancer. It generally is small in the beginning and increases gradually; but though the skin changes to a red or livid appearance, and the state of the tumour from an indolent to a painful one, it is sometimes very difficult to say when the scirrhus really becomes a cancer, the progress being quick or slow according to concurring causes. When the tumour is attended with a peculiar kind of burning, shooting pains, and the skin hath acquired the dusky purple or livid hue, it may then be deemed the malignant scirrhus or *confirmed cancer*. When thus far advanced in women's breasts, the tumour sometimes increases speedily to a great size, having a knotty unequal surface, more glands becoming obstructed, the nipple sinks in, turgid veins are conspicuous, ramifying around and resembling a crab's claw. These are the

characteristics of an occult cancer on the external parts; and we may suspect the existence of one internally when such pain and heat as has been described, succeed in parts where the patient hath before been sensible of a weight and pressure, attended with obtuse pain. A cancerous tumour never melts down in suppuration like an inflammatory one: but when it is ready to break open, especially in the breast, it generally becomes prominent in some minute point, attended with an increase of the peculiar kind of burning, shooting pain, felt before at intervals, in a less degree and deeper in the body of the gland. In the prominent parts of the tumour, in this state, a corroding ichor sometimes transudes through the skin, soon forming an ulcer; at other times a considerable quantity of a thin lymphatic fluid tinged with blood from eroded vessels is found on it. Ulcers of the cancerous nature discharge a thin, fetid, acrid sanies, which corrodes the parts, having thick dark-coloured retorted lips; and fungous excrescences frequently rise from these ulcers, notwithstanding the corrosiveness of the discharge. In this state they are often attended with excruciating, pungent, lancinating, burning pains, and sometimes with bleeding.

Though a scirrhus may truly be deemed a cancer, as soon as a pain is perceived in it, yet every painful tumour is not a cancer; nor is it always easy to say whether a cancer is the disorder or not: irregular hard lumps may be perceived in the breast; but on examining the other breast, where no uneasiness is perceived, the same kind of tumours are sometimes found, which renders the diagnostic uncertain. Yet in every case, after the cessation of the catamenia, hard unequal tumours in the breast are suspicious; nor, though without pain, are they to be supposed indolent or innoxious.

2. The name of a crab-fish, from which the *chele cancerorum*, *oculi cancerorum*, or *lapi-des cancerorum* are produced. The shell-fish so called is the *Cancer astacus* of Linnæus: the official preparations are nevertheless obtained also from the *cancer gammarus*, *marurus* and *pagurus* of Linnæus. Crab's claws and crab's eyes, as they are called, which are cerebral concretions, are of a calcareous quality and possess antacid virtues. They are exhibited with their compounds in pyrosis, diarrhoea, and in infantile convulsions from acidity.

CANCER MUNDITORUM. Chimney-sweep-er's cancer.

CANCERYS. *Cachrys*. *Libanotis*. Galen says it sometimes means parched barley.

CANCERENA. Paracelsus uses this word of gangrenæ.

CANCERUM CHELE. Crab's claws. See *calcis durior*.

CANCERUM OCULI. See *Carbonas cal-*

CANCERUM ORIS. (From *cancer*, a spreading ulcer.) Canker of the mouth; called also *splithæ serpentes*, *gangræna oris*, &c. See *Aphtha*.

CANDELA. (From *candeo*, to shine.) A candle.

CANDELA FUMALIS. A candle made of odoriferous powders and resinous matters, to purify the air and excite the spirits.

CANDELA REGIA. See *Candelaria*.

CANDELARIA. (From *candela*, a candle, so called from the resemblance of its stalks to a candle.) The herb mullein. See *Verbascum*.

Candy carrot. See *Daucus Creticus*.

CANELA. Sometimes used by the ancients for cinnamon, or rather cassia.

CANELLA. (*Canella*, dim. of *canna*, a reed; so named because the pieces of bark are rolled up in the form of a reed.) The name of a genus of plants in the Linnæan system. Class, *Dodecanthia*. Order, *Monogymia*. The canella-tree.

CANELLA ALBA. The pharmacopœial name of the laure-leaved canella *Cortex Winteranus spurius*. *Canella cubana*. *Canella alba* of Linnæus. The tree which produces the bark so called, is a native of the West-Indies. It is brought into Europe in long quills, somewhat thicker than cinnamon; their taste is moderately warm, aromatic, and bitterish; and of an agreeable smell, somewhat resembling that of cloves. *Canella alba* has been supposed to possess considerable medicinal powers in the cure of the scurvy and some other complaints. It is now merely considered as a useful and cheap aromatic, and is chiefly employed for the purpose of correcting, and rendering less disagreeable the more powerful and nauseous drugs; with which view it is used in the *tinctura amara*, *vinum amarum*, *vinum rhei*, &c. of the Edinburgh Pharmacopœia.

CANELLA CUBANA. See *Canella alba*.

CANELLA CUURDO. The true cinnamon-tree.

CANELLE MALABARICÆ CORTEX. See *Cassia lignea*.

CANELIFERA MALABARICA. See *Cassia lignea*.

CANEON. (From *cana*, because it was made of split cane.) A sort of tube or instrument, mentioned by Hippocrates, for conveying the fumes of antihysterical drugs into the womb.

CANICA. A spice used in the Island of Cuba, probably the pimento; or from some of the species of myrrhs.

CANICE. (From *canis*.) Coarse meal was so called by the ancients, from *canis*, a dog, because it was food for dogs. Hence *panis caniceus*, very coarse bread.

CANICIDA. (From *canis*, a dog, and *cædo*, to kill; so called because dogs are destroyed by eating it.) The herb dog's bane or aconitum.

CANICIDIUM. (From *canis*, a dog, and *cedo*, to kill.) The anatomical dissection of living dogs.

CANINA BRASSICA. The *mercurialis sylvestris*.

CANINA LINGUA. The cynoglossum.

CANINA MALUS. The mandragora.

CANINA RABIES. The hydrophobia.

CANINE. Whatever partakes of, or has any relation to the nature of a dog:

Canine appetite. See *Bulimia*.

Canine madness. See *Hydrophobia*.

CANINE TEETH. *Dentes canini.* *Cynodontes.* *Cuspidati* of Mr. John Hunter; because they have the two sides of their edge sloped off to a point, and this point is very sharp or cuspidated. *Cohimellares* of Varro and Pliny. The four eye-teeth are so called from their resemblance to those of the dog. They are situated, two in each jaw, on the side of the four middle or incisor-teeth. Their fangs are longer than those of the incisores, and therefore, from the fangs of those in the upper jaw being supposed to extend the greatest part of the way to the eye, they have been called the *eye-teeth*.

CANINUS. (*Caninus*, sc. *musculus*; because it arises near the canine or eye-tooth.) See *Levator anguli oris*.

CANINUS SENTIS. (From *canis*, a dog, and *sentis*, a thorn; from its being prickly like a thorn.) See *Cynosbatus*.

CANIRAM. (Indian.) See *Nux vomica*.

CANIRUBUS. (From *canis*, and *rubus*, a bramble.) See *Cynosbatus*.

CANIS. A dog. The white dung of this animal called *album græcum*, was formerly in esteem, but now disused. This term was also applied to the frænum of the penis.

CANUS INTERFECTOR. Indian caustic barley or cevadilla.

CANIS PONTICUS. See *Castor*.

CANNA. (Heb.) A reed or hollow cane: also a name of the fibula, from its resemblance to a reed.

CANNA FISTULA. See *Cassia fistula*.

CANNA INDICA. The *Sagittaria alexipharmica*.

CANNA MAJOR. The tibia.

CANNA MINOR CRURIS. A name formerly applied to the fibula.

CANNABINA. (From *canna*, a reed; named from its reed-like stalk.) So Tournefort named the *Datisca*.

CANNABIS. (From *kana*, a reed. *Kanah* are foul springs, wherein hemp, &c. grow naturally. Or from *kanaba*, from *kanah*, to mow. Arab.) Hemp.

1. The name of a genus of plants in the Linnæan system. Class, *Diœcia*. Order, *Pentandria*.

2. The pharmacopœial name of the *Cannabis sativa* of Linnæus. It has a rank smell of a narcotic kind. The effluvia from the fresh herb are said to affect the eyes and

head, and that the water in which it has been long steeped is a sudden poison. Hemp-seeds, when fresh, afford a considerable quantity of oil. Decoctions and emulsions of them have been recommended against coughs, ardor urinæ, &c. Their use, in general, depends on their emollient and demulcent qualities. The leaves of an oriental hemp, called *bang* or *bangue*, and by the Egyptians *assis*, are said to be used in Eastern countries, as a narcotic and aphrodisiac. See *Bangue*.

CANNABIS SATIVA. The systematic name of the hemp plant. See *Cannabis*.

CANNACORUS RADICE CROCEA. See *Curcuma*.

CANNULA. (Dim. of *canna*, a reed.) The name of a surgical instrument. See *Cunula*.

CANON. (*Κανον*.) A rule or canon, by which medicines are compounded.

CANONIAL. (*Κανονιαι*.) Hippocrates in his book, *De Aëre*, &c. calls those persons thus, who have straight, and not prominent beillies. He would intimate that they are disposed, as it were by a straight rule.

CANOPICON. (From *καναριον*, the flower of the elder.) A sort of spurge named from its resemblance; also a collyrium, of which the chief ingredient was elder-flowers.

CANOPITE. The name of a collyrium mentioned by Celsus.

CANOPUM. (*Καναρον*.) The flower or bark of the elder-tree, in Paulus Ægineta.

CANTABRICA. *Convolvulus minimus spice foliis.* *Convolvulus linaria folio.* *Convolvulus Cantabrica* of Linnæus. Lavender-leaved bindweed. Pliny says it was discovered in the time of Augustus, in the country of the Cantabri in Spain; whence its name. It is anthelmintic and actively cathartic.

CANTABRUM. (From *kanta*, Heb.) In Cælius Aurelianus it signifies bran or fur.

CANTACON. Garden saffron.

CANTARA. The plant which bears the St. Ignatius's bean.

CANTARI FIGULINI. Earthen cucurbits.

CANTHARIS. (*Cantharis*, pl. *cantharides*; from *κανθαρος*, a beetle, to whose tribe it belongs.) *Musca Hispanica* *Lytta venicatoria* of Linnæus. The blistering fly. Spanish fly. The importance of these flies, by their stimulant, corrosive, and epispastic qualities, in the practice of physic and surgery, is very considerable; indeed, so much so, as to induce many to consider them as the most powerful medicine in the materia medica. These flies have a green shining gold body, and are common in Spain, Italy, France, and Germany. The largest come from Italy, but the Spanish cantharides are generally preferred. When applied on the skin, in the form of a plaster, it soon raises a blister full of serous matter, and thus

relieves inflammatory diseases, as phrenitis, pleuritis, hepatitis, phlegmon, bubo, myositis, arthritis, &c. The tincture of these flies is also of great utility in several cutaneous diseases, rheumatic affections, sciatic pains, &c. but ought to be used with much caution. See *Blister*.

CANTHUS. Sugar-candy.

CANTHUS. (*Karbox*, the iron binding of a cart-wheel. Dr. Turton, in his glossary, supposes, from its etymology, that it originally signified the circular extremity of the eye-lid.) The angle or corner of the eye, where the upper and under eye-lids meet. That next the nose is termed the internal or greater canthus, and the other, the external or lesser canthus.

CANTION. An epithet for sugar.

CANTUARIENSIS AQUA. Canterbury water is strongly impregnated with iron, sulphur, and carbonic acid gas; recommended in disorders of the stomach, in gouty complaints, jaundice, diseases of the skin and chlorosis.

CANULA. (Dim. of *canna*, a reed.) A tube adapted to a sharp instrument, with which it is thrust into a cavity or tumour, containing a fluid; the perforation being made, the sharp instrument is withdrawn, and the banula left, in order that the fluid may pass through it.

CANUSA. Crystal.

CAOUTCHOUC. See *Indian rubber*.

Copaiva balsam. See *Balsamum Copaiba*.

CAPELINE. (From *capeline*, a woman's hat, or bandage, French.) A double-headed roller put round the head.

CAPELLA. A cupel or test.

Caper-bush. See *Capparis*.

CAPETUS. (*Karidos*, per aphæresin, pro *καυτός*: from *καυτός*, to dig.) Hippocrates means by this word a foramen, which is impervious and needs the use of a surgical instrument to make an opening; as the anus of some new-born infants.

CAPHORA. (Arab.) Camphire.

C-PHURA BARON INDORUM. A name for camphire.

CAPHURE OLEUM. An aromatic essential oil distilled from the root of the cinnamon-tree.

CAPILLARES VERMICULI. See *Crinones* and *Dracunculi*.

CAPILLARY VESSELS. (*Vasa capillaria*; from *capillus*, a little hair; so called from their resemblance to hairs or fine threads.) The very small ramifications of the arteries, which terminate upon the external surface of the body, or on the surface of internal cavities.

CAPILLATIO. (From *capillus*, a hair.) A capillary fracture of the cranium.

CAPILLUS. (Quasi *capitis pilus*, the hair of the head.) The hair. Small, cylindrical, transparent, insensible, and elastic filaments, which arise from the skin, and are fastened in it by means of small roots.

The human hair is composed of a spongy, cellular texture, containing a coloured liquid, and a proper covering. Hair is divided into two kinds: *long*, which arises on the scalp, cheek, chin, breasts, of men, the anterior parts of the arms and legs, the arm-pits, groins, and pelvis: and *short*, which is softer than the long, is present over the whole body, except only the palms of the hand and sole of the foot. The hair originates in the adipose membrane from an oblong membranous bulb, which has vessels peculiar to it. The hair is distinguished by different names in certain parts: as, *capillus*, on the top of the head; *crinis*, on the back of the head; *circinnus*, on the temples; *cilium*, on the eye-lids; *supercilium*, on the eyebrows; *vibrissa*, in the nostrils; *barba*, on the chin; *pappus*, on the middle of the chin; *mustax*, on the upper lip; *pilus*, on the body.

CAPILLUS VENERIS. See *Adiantum*.

CAPILLUS VENERIS CANADENSIS. The *Adiantum Canadense*.

CAPILENIUM. (From *caput*, the head, and *plenus*, full.) A catarrh. It is a barbarous word; but Baglivi uses it to signify that continual heaviness or disorder in the head, which the Greeks call *Carebaria*, *nagbagia*.

CAPISTRATIO. (From *capistrum*, a bridle; so called because the præpuce is restrained as it were with a bridle.) See *Phymosis*.

CAPISTRUM. (From *caput*, the head.) A bandage for the head is so called. In Vogel's Nosology it is the same as *Trismus*.

CAPITAL. The upper part of an alembic; likewise called the head.

CAPITALIA. (From *caput*, the head.) *Cephalics*: medicines which relieve disorders of the head.

CAPITELLUM. The head or seed vessels, frequently applied to mosses, &c. Some say it signifies soapy water, others say it is a lixivium.

CAPITILEVIUM. (From *caput*, the head, and *lavo*, to wash. A lotion or bath for the head.

Capitis obliquus inferior et major. See *Obliquus inferior*.

Capitis par tertium Follopii. See *Complexus minor*.

Capitis posticus. See *Rectus major capitis*.

Capitis rectus. See *Rectus minor capitis*.

CAPITULUM. (Dim. of *caput*, the head.) An alembic. In anatomy, a small head or protuberance of a bone, received into the concavity of another bone.

CAPIVI. (Indian.) A tree of Brazil, which affords the drug called balsam of capivi. See *Balsamum copaiba*.

CAPNELUM. (From *capnos*, smoke, and *oleum*, oil; so named from its smoky exhalations when exposed to heat.) In Galen's works, it is said to be a resin.

CAPNUS. (From *καπνος*, a smoke.) A jasper of a smoky colour. Also a kind of vine which bears white and part black grapes.

CAPNISTOM. (From *καπνος*, smoke.) A preparation made of spices and oil, by kindling the spices and fuming the oil.

CAPNITIS. (From *καπνος*, smoke; so called from its smoky colour.) Tutty.

CAPNOIDES. (From *καπνος*, fumitory, and *οειδης*, likeness.) A species of fumitory.

CAPNOS. *καπνος*. Fumitory: so called, says Blanchard, because its juice, if applied to the eyes, produces the same effect and sensations as smoke.

CAPNO MOLAGO. The Piper Indicum.

CAPPA. (*α-capite*, from the head; so called from its supposed resemblance.) The herb monkshood.

CAPPARIS. (From *cabar*, Arab or *απαρ* to *καππαρις* *απαρ*, from its curing madness and melancholy.) The caper plant.

1. The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Menogynia*.

2. The pharmacopœial name of the *Capparis spinosa* of Linnæus:—*pedunculis sociatis unifloris, stipulis spinosis, foliis annuis, capsulis ovalibus*. The buds or unexpanded flowers of this plant, are in common use as a pickle, which is said to possess antiscorbutic virtues. The bark of the root was formerly in high esteem as a deobstruent.

CAPPARIS SPINOSA. The systematic name of the caper plant. See *Cupparis*.

CAPREOLARIS. (From *capreolus*, a tendril.) *Capreolatus*. Resembling in its contortions, or other appearance, the tendrils of a vine; as the spermatic vessels.

CAPREOLATUS. See *Capreolaris*.

CAPREOLUS (Dim. of *caprea*, a tendril.) It means the helix or circle of the ear, from its tendril-like contortion. Dr. Turton suggests its derivation from *caper*, a goat, whose horn its contortions somewhat resemble.

CAPRICORNUS. Lead.

CAPRICUS. (From *caper*, a goat, and *ficus*, a fig; because they are a chief food of goats.) The wild fig-tree.

CAPRIZANS. Is by Galen and others used to express an inequality in the pulse, when it leaps, and, as it were, dances in uncertain strokes and periods.

CAPSELLA. (Dim. of *capsea*, a chest, from its resemblance.) A name in Marcellus Empiricus for viper's bugloss.

CAPSICUM. (From *καρσιν*, to bite, on account of its effect on the mouth.)

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*. Guinea pepper.

2. The pharmacopœial name of the *Capicum annuum* of Linnæus. What is generally used under the name of Cayenne pepper, is an indiscriminate mixture of the powder

of the dried pods of many species of capsicums, but especially of the capsicum minimum, or bird pepper, which is the hottest of all. These peppers have been chiefly used as condiments. They prevent flatulence from vegetable food and give warmth to the stomach, possessing all the virtues of the oriental spices, without producing those complaints of the head which the latter are apt to occasion. An abuse of them, however, gives rise to visceral obstructions, especially of the liver. In the practice of medicine, there can be little doubt that they furnish us with one of the purest and strongest stimulants which can be introduced into the stomach. Dr. Adair who first introduced them into practice, found them useful in the cachexia Africana, which he considers as a most frequent and fatal predisposition to disease among the slaves. Dr. Wright says, that in dropsical and other complaints where chalybeates are indicated, a minute portion of powdered capsicum forms an excellent addition and recommends its use in lethargic affections. This pepper has also been successfully employed in a species of cynanche maligna, which proved very fatal in the West Indies, resisting the use of Peruvian bark, wine, and other remedies commonly employed. In tropical fevers, coma and delirium are common attendants, and in such cases cataplasms of capsicum have a speedy and happy effect. They reddens the parts, but seldom blister unless when kept on too long. In ophthalmia from relaxation, the diluted juice of capsicum is found to be a valuable remedy. Dr. Adair gave six or eight grains for a dose, made into pills; or else he prepared a tincture by digesting half an ounce of the pepper in a pound of alcohol, the dose of which was one or two drachms, diluted with a sufficient quantity of water. A *tinctura capsaci* is now for the first time introduced into the London pharmacopœia.

CAPSULA. (Dim. of *capsea*, a chest or case.) A term given by anatomists to any membranous production enclosing a part of the body like a bag; as the capsular ligaments, the capsule of the crystalline lens, &c.

CAPSULE ATRABILARIS. See *Renal capsules*.

CAPSULE RENALES. See *Renal capsules*.

CAPSULAR LIGAMENT. (*Capsularis*; from *capsea*, a bag.) *Ligamentum capsulare*. The ligament which surrounds every movable articulation, and contains the synovia like a bag.

CAPSULE OF GLYSSON. *Capsula communis, Glyssonii, Vagina portæ, Vagina Glyssonii*. A strong tunic, formed of cellular texture, which accompanies the vena portæ, and its most minute ramifications throughout the whole liver.

CAPULUM. (From *καμπερα*, to bend.) A contortion of the eye-lids, or other parts.

CAPUL. (Arab.) Camphor.

CAPUT. (From *capio*, to take; because from it, according to Vāro, the senses take their origin.) The head, cranium or skull. It is situated above or upon the trunk, and united to the cervical vertebrae.

For its bones, see *Skull*. It is distinguished into skull and face. On the skull are observed *vertex* or crown; *sinciput*, or fore-part; *occiput*, or hinder part; and the *temples*. The parts distinguished on the face are well known; as the forehead, nose, eyes, &c. The arteries of the head are branches of the carotids; and the veins empty themselves into the jugulars.

CAPUT GALLINAGINIS. *Verumontanum*. A cutaneous eminence in the urethra of men, before the neck of the bladder, somewhat like the head of a cock in miniature, around which the seminal ducts, and the ducts of the prostate gland, open.

CAPUT MORTUUM. A fanciful term, much used by the old chymists, but now entirely rejected. It denoted the fixed residue of operations. As the earlier chymists did not examine these, they did not find any inconvenience in one general term to denote them: but the most slender acquaintance with modern chymistry must show that it is utterly impracticable to denote, by one general term, all the various matters that remain fixed in certain degrees of heat.

CAPUT OBSTIPUM. The wry neck, mostly a spasmodic complaint.

CAPUT PURGIA. (A barbarous word, from *caput*, the head, and *purgo*, to purge.) Medicines which purge the head. Errhines. Masticatories.

CAPRYDION. (From *καρυγος*, burnt.) *Capyrion* A medicated cake, much baked.

CAPRYON. See *Capyridion*.

CARABE. (Persian.) Amber.

CARABE FUNERUM. A name given to bitumen.

CARABUS. A genus of insects of the beetle kind. Two species, the chrysoccephalus and ferruginous, have been recommended for the tooth-ach. They must be pressed between the fingers, and then rubbed on the gum and tooth affected.

CARACOSMOS. A name of the sour mare's milk, so much admired by the Tartars.

CARAGUATA. The common aloe of Brazil.

CARANEA, Caragna, Caranne gummi. (Spanish.) *Brazilia*. A concrete resinous juice, that exudes from a large tree, of which we have no particular account. It is brought from New Spain and America,

in little masses, rolled up in leaves of figs; externally and internally it is of a brownish colour, variegated with irregular white streaks. When fresh, it is soft and tenacious, but becomes dry and friable by keeping. Pure caranna has an agreeable aromatic smell, especially when heated, and a bitterish slightly pungent taste. It was formerly employed as an ingredient in vulnerary balsams, strengthening, discutient, and suppurating plaisters; but its scarcity has caused it to be forgotten.

CARA SCHULLI. (Indian) *Frutex Indica spinosa*. An Indian shrub, like the caper-bush. A decoction of the root proves diuretic. *Ray*.

Carawayseed. See *Carum*.

CARBASUS. (*Καρβασος*.) Scribonius Largus uses this word for lint.

CARBO. (*Carbah*, Heb. burnt, or dried.) Coal. In medicine and chymistry, it is commonly understood to mean charcoal, and receives its name from its mode of preparation, which is by burning pieces of light wood into a dry black coal.

CARBO LIGNI. Charcoal. As an external application, powdered charcoal has been recommended in the cure of gangrene, from external causes, and all descriptions of fetid ulcers. Meat which has acquired a mawkish or even putrid smell, is found to be rendered perfectly sweet by rubbing it with powdered charcoal.

CARBON. (From *carbo*, coal.) The chymical name of charcoal. It is the black residue of vegetables, which have suffered a complete decomposition of their volatile principles by fire. Charcoal is black, brittle, sonorous and light. It is placed among simple bodies, because no experiment has hitherto shown the possibility of decomposing it. It exists in the animal, vegetable, and mineral kingdom. When it is required to procure carbon in a state of great purity, it must be dried by strong ignition in a closed vessel. The diamond when burnt in oxygen gas forms charcoal. Charcoal is therefore considered to be an oxyd of diamond, and the diamond pure carbon.

CARBON, GASEOUS OXYD OF. Gaseous oxyd of carbon was first described by Dr. Priestley, who mistook it for a hydro-carbonate. With the true nature of it we have been only lately acquainted. It was first proved to be a peculiar gas, by Mr. Cruikshank, of Woolwich, who made it known to us as such, in April 1801, through the medium of Nicholson's Journal for that month. Several additional properties of this gas were soon afterwards noticed by Desormes, Clement and others. Gaseous oxyd of carbon forms an intermediate substance between the pure hydro-carbonates and carbonic acid gas; but not being possessed of acid properties, Mr. Cruikshank has called

it conformable to the rules of the chymical nomenclature, *gaseous oxyd of carbon*, for it consists of oxygen and carbon rendered gaseous by caloric.

Though the gaseous oxyd of carbon has some of the properties peculiar to the common hydro-carbonates, the following characteristic properties sufficiently prove that none of those at present known are similar to it. We are, therefore, entitled to consider it as a peculiar gas.

Properties.—Gaseous oxyd of carbon is considerably lighter than any of the hydro-carbonates. It is lighter than common air, in the proportion of 22 to 23. When mingled with common air, and ignited, it does not explode, but burns with a lambent blue flame, and the product is carbonic acid. It is very little absorbable by water; it is void of taste and odour. A mixture of 20 parts of gaseous oxyd of carbon and 8 of oxygen gas, fired over mercury, by electricity, diminishes to a volume equal to about 18 or 19 parts, which is carbonic acid gas. It contains neither water nor the basis of that fluid. It is exceedingly noxious: animals die in it instantly; when breathed for a few minutes only, it produces giddiness and faintings. Neither light, heat, nor electricity have any effect upon it. When equal quantities of gaseous oxyd of carbon and hydrogen gas are passed through a red-hot glass tube, the tube is lined with charcoal, water is formed, and an excess of hydrogen makes its escape. If a piece of iron be put into the tube, it is oxydated, but not converted into steel. Neither nitrogen gas nor sulphur have any action on it even at high temperatures. It is capable of dissolving a minute quantity of charcoal, and increases in bulk. It dissolves phosphorus and acquires the property of burning with a yellow flame. The alkalis have no effect on this gas. It is not altered when passed with ammonia through an ignited tube. When the red oxyd of mercury is heated in it, a commencement of reduction takes place. Neither sulphuric, nitric, nor nitro-muriatic acids, alter it, when passed with it through a red-hot tube. Four parts of oxygenated muriatic acid gas left with one of carbonic acid gas, decompose it completely. Nitrous gas has no effect upon it. When mixed with sulphurated hydrogen gas, and passed through a red-hot tube, sulphur is deposited, and sulphurated hydrogen gas remains mixed with gaseous oxyd of carbon.

Methods of obtaining Gaseous Oxyd of Carbon.—Gaseous oxyd of carbon may be obtained by a decomposition of carbonic acid at high temperatures, by means of various fixed substances which have a considerable affinity to oxygen. This may be done by distilling a mixture of charcoal with any of the metallic oxyds, or by ex-

posing to a strong red heat, a mixture of carbonate of lime or barytes, and filings of iron, zinc, &c.

The method of obtaining the gaseous oxyd of carbon in a state of purity, recommended by Mr. Cruikshank, is the following:

1. Take one part of chalk, previously exposed to a low red heat, for about ten minutes, mix it with an equal quantity of perfectly dry filings of zinc; let the mixture be introduced into a retort, and expose it to a heat gradually increased. As soon as the retort becomes of a dull red heat, gas will be disengaged in great abundance. The gas which comes over first is carbonic acid gas, but as soon as the retort becomes thoroughly ignited, pure gaseous oxyd of carbon is liberated in a prodigious quantity, which may be collected in the usual manner over water.

In this process, a decomposition of the carbonic acid of the chalk takes place in its nascent state. The zinc robs the carbonic acid of part of its oxygen at a high temperature, and becomes to a certain degree oxydated. The carbonic acid, by being thus deprived of part of its oxygen, becomes converted into a new inflammable gas, which is the gaseous oxyd of carbon.

Carbonaceous acid. See *Carbonic acid*.

CARBONAS. A carbonate. A neutral salt, formed by the union of carbonic acid with an alkaline, earthy, or metallic base. The carbonates employed in medicine are some of them perfect and some imperfect.

The imperfect carbonates in use are—

1. The subcarbonas potassæ.
2. The subcarbonas sodæ.
3. The subcarbonas sodæ exsiccata.
4. The subcarbonas plumbi.

The perfect carbonates are—

1. The carbonas ammoniæ.
2. The liquor carbonatis ammoniæ.
3. The carbonas potassæ.
4. The carbonas sodæ.
5. The creta præparata.
6. The carbonas magnesiæ.
7. The carbonas ferri.

CARBONAS AMMONIÆ. Carbonate of ammonia. This preparation was formerly called *ammonia præparata*, and *sal volatilis salis ammoniaci*, and *sal volatilis*. It is made thus:—take of muriate of ammonia, a pound; of prepared chalk, dried, two pounds. Reduce them separately to powder; then mix them together, and sublime in a heat gradually raised, till the retort becomes red.

This salt possesses nervine and stimulating powers, and is highly beneficial in the dose of from two to eight grains, in nervous affections, debilities, flatulency, from acidity and dyspepsia.

CARBONAS CALCIS. Carbonate of lime. Several of these are used in medi-

cine; the purest and best are the creta preparati, chelz cancrorum, testz ostrearum, testz ovorum, and oculi cancrorum.

CARBONAS PLUMB. See *Cerussa*.

CARBONAS POTASSÆ. Carbonate of potash. This preparation, which has been long known by the name of *Kali eratum*, appeared in the last London pharmacopœia, for the first time. It is made thus:—Take of subcarbonate of potash, made from tartar, a pound; carbonate of ammonia three ounces; distilled water, a pint. Having previously dissolved the subcarbonate of potash in the water, add the carbonate of ammonia: then, by means of a sand-bath, apply a heat of 180° for three hours, or until the ammonia shall be driven off; lastly, set the solution by, to crystallize. The remaining solution may be evaporated in the same manner, that crystals may again form when it is set by.

This process was invented by Bertholet. The potash takes the carbonic acid from the ammonia, which is volatile, and passes off in the temperature employed. It is, however, very difficult to detach the ammonia entirely. Potash is thus saturated with carbonic acid, of which it contains double the quantity that the subcarbonate of potash does; it gives out this proportion on the addition of muriatic acid, and may be converted again into the sub-salt, by heating it again a short time, to redness. It is less nauseous to the taste than the subcarbonate; it crystallizes, and does not deliquesce. Water, at the common temperature, dissolves one-fourth its weight, and at 212°, five-sixths; but this latter heat detaches some of the carbonic acid.

The carbonate of potash is generally used for the purpose of imparting carbonic acid to the stomach, by giving a scruple in solution with a table spoon-ful of lemon-juice, in the act of effervescing.

CARBONAS SODÆ. Carbonate of soda. Take of the subcarbonate of soda, a pound; of the carbonate of ammonia, three ounces; of distilled water, a pint. Having previously dissolved the soda in the water, add the ammonia: then, by means of a sand-bath, apply a heat of 180° for three hours, or until the ammonia is driven off. Lastly, set the solution by, to crystallize. The remaining solution may, in the same manner, be evaporated, and set by, that crystals may again form.

This salt which is called also *aërated soda* and *natron*, bears to the subcarbonate of soda, the same relation that the carbonate of potash does to its subcarbonate. It is prepared in the same way, possesses the same comparative advantages, and contains double the quantity of carbonic acid.

CARBONAS MAGNESIÆ. *Magnesia alba.* The carbonate of magnesia. This

preparation is variously prepared. The college of physicians of London direct it thus:—Take of sulphate of magnesia, of subcarbonate of potash, of each a pound; water, three gallons. Dissolve the subcarbonate of potash in three pints of the water, and strain. Dissolve also the sulphate of magnesia separately, in five pints of the water, and strain; then add the rest of the water to the latter solution, apply heat, and, when it boils, pour in the former solution, stirring them well together; next, strain through a linen cloth; lastly, wash the powder repeatedly with boiling water, and dry it upon bibulous paper, in a heat of 200 deg. .

Carbonate of magnesia is esteemed as an aperient and antacid, and is given against constipation, flatulency, acidity of the stomach, and its effects. The dose is from ten grains to a drachm.

CARBONAS FERRI. *Ferrum precipitatum.* Carbonate of iron. This preparation is made by decomposing the sulphate of iron by the subcarbonate of soda, thus—Take of the sulphate of iron, eight ounces; subcarbonate of soda, ten ounces; boiling water, a gallon. Dissolve the sulphate of iron and subcarbonate of soda separately, each in four pints of water; next mix the solutions together, and set it by, that the precipitated powder may subside; then, having poured off the supernatant liquor, wash the carbonate of iron with hot water, and dry it upon bibulous paper, in a gentle heat.

This salt is a subcarbonate of iron, and not a perfect carbonate. It is substituted for the *rubigo ferri* of the former pharmacopœias. It is much esteemed as a mild chalybeate, and is given in the dose of from five grains to a scruple.

CARBONATED HYDROGEN GAS, LIGHT. Light carbonated hydrogen gas is hydrogen gas holding charcoal in solution. There are several combinations of this kind of gas obtained by different processes, which differ in their properties, and in the proportion of their constituent principles.

Properties.—Light carbonated hydrogen gas has a fœtid odour. It is neither absorbed nor altered by water. It is inflammable, and burns with a denser and deeper coloured flame than hydrogen gas. It is unalterable by acids or alkalis, and by water. Its specific gravity is greater than that of hydrogen gas, or that of common air. Its combustion with a due proportion of oxygen gas, is productive of water and carbonic acid. When passed through melted sulphur, it becomes converted into sulphurated hydrogen gas, and charcoal is deposited. Electrization dilates it permanently to a little more than twice its original bulk. The air thus expanded, re-

quires a greater quantity of oxygen to decompose it, than the same quantity of gas not diluted by electricity; 100 cubic inches of pure carbonated hydrogen gas weigh from 16 to 24 grains.

Light carbonated hydrogen gas may be obtained from animal, vegetable, or mineral substances. Nature produces it ready formed in marshes and ditches, on the surface of putrid water, in burying-places, common sewers, and in those situations where putrid animal and vegetable matters are accumulated. It is also generated in the intestinal canal of living animals.

1. Light carbonated hydrogen gas may be plentifully procured from most stagnant waters: to do this, fill a wide-mouthed bottle with the water, and keep it inverted therein, with a funnel in its neck; then, with a stick, stir the mud at the bottom, just under the funnel in the bottle, so as to let the bubbles of air, which rise from the mud, enter into the bottle; when, by thus stirring the mud in various places, and catching the air in the bottle, it is filled, it must be corked under water.

2. It may be also obtained during the distillation of animal and vegetable matters. For instance:

Let shavings of wood, or saw-dust, be put into a retort, and begin the distillation with a gentle heat, increasing it gradually, till the retort becomes red hot; a great quantity of gas will be liberated, which may be caught over water. On examining this gas, it will be found to consist of carbonic acid gas and carbonated hydrogen gas. In order to obtain the latter in a state of purity, the whole must be mixed with lime-water, or with a caustic alkaline solution. The carbonic acid gas will be absorbed, and the carbonated hydrogen gas left behind, in a pure state.

The production of this gas in this manner, is the result of a partial analysis of the wood. It proves that wood contains solid hydrogen, carbon, and oxygen. When the intensity of the heat has reached a certain degree, a part of the charcoal unites with part of the oxygen and produces carbonic acid, which, by means of caloric, is melted into the gaseous state, and forms carbonic acid gas; at the same time, a part of the hydrogen of the wood combines with another portion of carbon and caloric, and forms carbonated hydrogen gas.

Remark.—The flame of burning wood, &c. is the inflamed carbonated hydrogen gas, liberated on the application of caloric to such bodies.

3. Charcoal is in general made use of for obtaining light carbonated hydrogen gas. For this purpose, put some moistened charcoal into an earthen retort, apply heat, and increase it till the retort becomes ig-

nited; gas will be evolved, consisting partly of carbonic acid gas, and partly of light carbonated hydrogen gas, which may be separated as before.

In this case a decomposition of the water takes place, by means of the charcoal. The oxygen forsakes its hydrogen, and unites to part of the charcoal, at this temperature, and forms carbonic acid gas, in conjunction with caloric; the liberated hydrogen assisted by caloric, dissolves another portion of the charcoal, and forms with it light carbonated hydrogen gas.

4. Light carbonated hydrogen gas may be formed in a direct manner, by dissolving charcoal in hydrogen gas. This may be effected by directing the rays of the sun collected in the focus of a mirror, upon a small piece of charcoal placed on mercury, in a receiver filled with hydrogen gas.

CARBONATED HYDROGEN GAS, HEAVY. This gas was first brought into notice by a society of Dutch chymists, consisting of Deiman, Troostwyk, Bond, and Laurenburgh. They observed in this gas the particular property, that when it was combined with oxygenated muriatic acid gas, in a certain proportion, the elastic form of both fluids became destroyed, and an oil was produced; for which reason they called it *Olefiant gas*.

Properties.—Heavy carbonated hydrogen gas is not absorbed or altered by water. Its weight, compared with common air, is as 909 to 1000. It has a disagreeable fœtid odour, different from that of light carbonated hydrogen gas. It burns with a strong compact flame, similar to that of a resinous oil. When mixed with oxygenated muriatic acid gas, its bulk is diminished, and an oil is formed. When the mixture of these two gases is fired, a quantity of charcoal is immediately deposited, in the form of fine soot. Sulphuric, sulphureous, nitric, nitrous, and muriatic acids do not act upon it; neither does nitrous gas, nor any of the fixed alkalies. Ammonia adds to its volume without occasioning any other change. Phosphorus heated in it, even to fusion does not affect it. When made to pass through an ignited glass tube, it does not diminish in volume, but loses the property of forming oil with oxygenated muriatic acid gas. Electric shocks passed through it, dilate and likewise deprive it of this property. When passed through an ignited porcelain tube, it affords hydrogen gas mingled with carbonic acid, and carbon is deposited. When passed through a tube with sulphur in fusion, sulphurated hydrogen gas is obtained, and charcoal deposited. When burnt with oxygen gas, or when passed through a red-hot tube, filled with oxyd of manganese, carbonic acid gas is formed.

Preparation.—Heavy carbonated hydrogen gas is obtained by decomposing alcohol by sulphuric acid, at high temperatures. It is also obtained in abundance when alcohol or ether is passed through a red-hot earthen tube. Sulphuric ether mixed with sulphuric acid, and subjected to heat, also affords it, but in a less pure state.

The Dutch chymists observed, that if the vapour of ardent spirit or ether be made to pass through a glass tube, over the component parts of the earthen tube, namely, alumine and silex, this gas was also produced; or by passing it through a red-hot tube of pipe-clay.

In order to obtain this gas the following method may serve:

Let four parts of concentrated sulphuric acid, and one of highly rectified ardent spirit, be mingled together gradually in a glass retort; heat will be developed, the mixture will become brown, and heavy carbonated hydrogen gas will be extricated without the application of external heat. When a moderate heat is applied, the action is very violent, and the gas is liberated very copiously, and may be received over water.

The gas obtained, is always mixed with a considerable quantity of sulphureous acid gas, from which it may be freed by agitating it in contact with lime-water, or a solution of potash.

Remark.—In this operation the heat ought to be regulated with great care, and the retort holding the mixture ought to be very capacious, otherwise the matter will be forced over into the receiver. The heat of a candle, or lamp, is sufficient.

CARBONIC ACID GAS. *Acidum Carbonicum.* Fixed air. Carbonaceous acid. Aërial acid. Carbonic acid gas is the first elastic aëriform fluid that was known. We find that the ancients were in some measure acquainted with it. Van Helmont called it the gas of Must, or of the vintage, or *gas sylvestre*.

We are indebted to Dr. Black of Edinburgh for the knowledge of some of the most remarkable properties of this fluid. In the year 1755 he discovered the affinity between this gas and alkalies: and Bergman, in 1772, proved that it was an acid.

Properties.—Carbonic acid gas is invisible. It extinguishes flame. It is fatal to animal life. It exerts powerful effects on living vegetables. Its taste is pungent and acid. Its energy, as an acid, is but feeble, although distinct and certain. Neither light or caloric seem to produce any distinct effect upon it, except that the latter dilates it. It mixes without combining with oxygen gas. It unites with water slowly. These two fluids, after considerable agitation, at last combine, and form a sub-acid fluid. The

colder the water, and the greater the pressure applied, the more carbonic acid gas will be absorbed. The water impregnated with it, sparkles upon agitation; it has a pungent acidulous taste, and reddens tincture of litmus. Heat again disengages the gas from the water. This gas precipitates lime, strontia, and barytes, from their solutions in water. It is greedily attracted by all the alkalies. It undergoes no alteration by light. Its specific weight is to that of atmospheric air, as 1500 to 1000. It may be poured out of one vessel into another. It is not acted upon by oxygen, nor is it altered by any of the simple combustible bodies at common temperatures; but phosphorus, iron, and lime, are capable of decomposing it, when assisted by heat.

Methods of obtaining Carbonic Acid Gas.

Of all the gases, carbonic acid gas is that, perhaps, which is diffused in the greatest abundance throughout nature. It is found in three different states:—1st, In that of gas; 2dly, In that of mixture; and 3dly, In that of combination. The various processes for obtaining it are the following:

1. Put into a common glass-bottle, or retort, a little marble, chalk, or limestone, and pour on it sulphuric acid, diluted with about six times its weight of water, an effervescence will ensue, and carbonic acid gas will be liberated, which those who have an opportunity may collect over mercury; but a mercurial apparatus is not absolutely necessary, since the gas may be collected over water, if it is to be used immediately when procured.

In this instance the carbonic acid is disengaged from the state of combination, and reduced to the aëriform state of gas. The marble, lime-stone, or chalk, consists of this acid and lime; on presenting to it sulphuric acid, a decomposition takes place, the sulphuric acid has a greater affinity to the lime than the carbonic acid has; it therefore unites to it, and forms sulphate of lime, disengaging at the same time, the carbonic acid in the state of gas, at the temperature of our atmosphere.

Remark.—Carbonic acid gas may, in this manner, be disengaged from all its combinations with alkalies; by using indifferently any other dense acid, possessing a superior affinity to the alkali in the common acceptance of the word.

2. It may likewise be obtained from the same substances by the action of caloric.

For this purpose, reduce marble, or chalk, to powder; introduce it into a gun-barrel, which must be placed across a furnace; adapt a bent tube to its lower extremity, and insert it below a receiver in the pneumatic apparatus. Maintain a strong heat, till the barrel is brought to a state of

ignition, and at that temperature carbonic acid gas will be liberated in abundance.

In this case, a decomposition of the marble or carbonate of lime takes place, on account of the action of caloric, which at a high temperature breaks the affinity of the carbonic acid and lime; it unites with the first, and leaves the lime behind in that state which is generally called quick-lime.

3. Carbonic acid gas may also be obtained by burning charcoal in oxygen gas.

Take a bell-glass, filled with oxygen gas, resting inverted in a basin of mercury; pass up into it some bits of new-made charcoal, with some touch-paper affixed to them; set fire to them by means of a lens collecting the sun's rays, and carbonic acid will be produced by the combustion of the charcoal.

Carbonic acid gas is often found occupying the lower parts of mines, caverns, tombs, and such other subterraneous places as contain materials for producing it. It is called choke, or chalk-damp. The grotto del Cane, near Naples, has long been famous for the quantity of carbonic acid gas produced there, which runs out at the opening like a stream of water. The quantity of carbonic acid gas generated in this cavern, is so great, that a dog, or any other animal, is immediately killed if his nose be thrust into it.

The carbonic acid, existing naturally in the state of gas, may be collected by filling bottles with water and emptying them into the atmosphere of this gas; the gas takes the place of the water, and fills the bottles, which must then be corked.

Carbonic acid gas is likewise formed during fermentation; on account of its great weight, it occupies the apparently empty space, or upper part of the vessel, in which the fermenting process is going on. It may in this case, be collected in a manner similar to that above.

Carbonic acid gas is also obtained during the reduction of metallic oxids, and during the deflagration of nitrates with combustible bodies. This gas is much esteemed in the cure of typhus fevers, and irritability and weakness of stomach producing vomiting. Against the former diseases it is given by administering yeast, bottle porter, and the like; and for the latter it is disengaged from the carbonated alkali by lemon juice in a draught given while effervescing.

CARBUNCLE. (Dim. of *carbo*, a burning coal.) *Carbo. Rubinus verus. Codebella. Erythema gangrenosum. Granatistrum. Pruna. Percus ignis* of Avicenna. An inflammatory tumour which soon becomes gangrenous. See *Anthrax*.

CARBUNCULUS. See *Carbuncle*.

CARCARUS. *Carcarus.* (From *κακαρος*, to resound.) A kind of fever in which the

patient has a continual horror and trembling, with an unceasing sounding in his ears.

CARCAS. The Barbadoes nut-tree, the *Cataputia*.

CARCAZ. (From *κακα*, a head.) A species of poppy, with a very large head.

CANCER. Paracelsus means by it, a remedy proper for restraining the disorder by motions of body and mind, as in curing the *chorea Sancti Viti*.

CANCRESIUS. (*Κακρησιος.*) A name of some bandages noticed by Galen, and described by Oribasius. Properly it is the top of a ship's mast.

CARCINOMA. (From *καρκινος*, a cancer, and *μασ*, to feed upon.) See *Cancer*.

CARCINOS. (*Καρκινος*, a cancer.) See *Cancer*.

CARDAMANTICA. (From *καρδαμω*, the nasturtium.) A species of asiatic cresses.

CARDAMELEUM. A medicine of no note, mentioned by Galen.

CARDAMINE. (From *καρδα*, the heart; because it acts as a cordial and strengthener, or from its having the taste of cardamum, that is, nasturtium, or cress.) Cuckoo-flower.

1. The name of a genus of plants in the Linnæan system. Class, *Tetradynamia*. Order, *Siliquosa*.

2. The pharmacopœial name of the common lady's smock, or cuckoo flower. *Cardamine pratensis* of Linnæus:—*foliis pinnatis, foliis radicalibus subrotundis, caulibus lanceolatis*. This plant is also called *Cardamantica. Nasturtium aquaticum. Culi flos. Iberis sophia*. It is the flower of this plant which has a place in the *materia medica*, upon the authority of Sir George Baker, who has published five cases, two of *chorea Sancti Viti*, one of spasmodic asthma, and a case of spasmodic affections of the lower limbs, wherein the *flores cardamine* were supposed to have been successfully used. A variety of virtues have been given to this plant, which do not deserve the attention of practitioners.

CARDAMINE PRATENSIS. The systematic name of the plant called *cardamine* in the pharmacopœias. See *Cardamine*.

CARDAMINES FLORES. See *Cardamine*.

CARDAMOMUM. (From *καρδαμω*, and *μασ*, because it partakes of the nature, and is like both the cardamum and amomum.) The cardamom seed, or grains of Paradise.

CARDAMOMUM MAJUS. A large, brown, somewhat triangular husk, the thickness of one's thumb, and pyramidal. The seeds resemble the *grana paradisi*, their virtues are similar to those of the *cardamomum minus*.

CARDAMOMUM MEDIUM. The seeds correspond, in every respect, with the lesser, except in size, they being twice as long, but no thicker than the *cardamomum minus*.

CARDAMOMUM MINUS. Official cardamom. *Anomum repens*, seu le cardamome de la côte de Malabar, of Sonnerat. *Elettaria cardamomum*, of Maton in Act. Soc. Lin. Class, *Menandria*. Order, *Monogymia*. The seeds of this plant are imported in their capsules or husks, by which they are preserved, for they soon lose a part of their flavour when freed from this covering. On being chewed they impart a glowing aromatic warmth, and grateful pungency; they are supposed gently to stimulate the stomach, and prove cordial, carminative, and antispasmodic, but without that irritation and heat which many of the other spicy aromatics are apt to produce. Simple and compound spirituous tinctures are prepared from them, and they are ordered as a spicy ingredient in many of the official compositions.

CARDAMOMUM PINKERATUM. The grains of Paradise.

CARDAMOMUM SIBERIENSE. The Anisum Indicum.

CARDAMUM. (From καρδία, the heart; because it comforts and strengthens the heart.) Garden cresses.

CARDIA. (From καρδία, the heart; so the Greeks called the heart.) The superior opening of the stomach.

CARDIACA. (From καρδία, the heart.)

1. Cordials. See *Cordials*.

2. The pharmacopœial name of motherwort. So named from the supposed relief it gives in faintings and disorders of the stomach. *Agripalmia gallica*. *Marrubium Cardiacum crispum*. *Leonurus cardiaca* of Linnaeus:—*foliis caulibus lanceolatis, trilobis*. The leaves of this plant have a disagreeable smell and a bitter taste, and are said to be serviceable in disorders of the stomach of children, to promote the uterine discharge, and to allay palpitations of the heart.

CARDIACA CONFECTIO. See *Confectio aromatica*.

CARDIACA PASSIO. The cardiac passion. Ancient writers frequently mention a disorder under this name, but the moderns always speak of it as a syncope.

CARDIACUS MORBUS. A name by which the ancients called the typhus fever.

CARDIALGIA. (From καρδία, the cardia, and αλγος, to be pained.) Pain at the stomach. The heartburn. Dr. Cullen ranks it as a species of dyspepsia. Heartburn is an uneasy sensation in the stomach, with anxiety, a heat more or less violent, and sometimes attended with oppression, faintness, an inclination to vomit, or a plentiful discharge of clear lymph, like saliva. This pain may arise from various and different causes; such as *status*; from sharp humours, either acid, bilious, or rancid; from worms gnawing and vellicating the coats of the stomach; from *acid* and *pungent* food, such as spices, aromatics, &c.; as also from *rheumatic* and *gouty humours*, or *sinfeits*; from too free a use of tea, or

watery fluids relaxing the stomach, &c.; from the natural mucus being abraded, particularly in the upper orifice of the stomach.

CARDIALGIA INFLAMMATORIA. Inflammation in the stomach.

CARDIALGIA SPUTATORIA, i. e. pyrosis. See *Pyrosis*.

CARDIMELECH. (From καρδία, the heart, and μελεχ, Heb. a governor.) A fictitious term in Dolzuz's Encyclopædia, by which he would express a particular active principle in the heart, appointed to what we call the vital functions.

CARDINOMA. A name for Cardialgia.

Cardinal flowers, blue. See *Lobelia*.

CARDINAMENTUM. (From cardo, a hinge.) A sort of articulation like a hinge.

CARDIOGMUS. (From καρδιον, to have a pain in the stomach.) The same as *Cardialgia*. Also an aneurism in the aorta, near the heart, which occasions pain in the præcordia.

CARDIOTHUS. (From καρδία, the heart, and ογκος, a tumour.) An aneurism in the heart, or in the aorta near the heart.

CARDIOTROTUS. (From καρδία, the heart, and τρωω, to wound.) One who hath a wound in his heart.

CARDITIS. (From καρδία, the heart.) Inflammation of the heart. It is a genus of disease arranged by Cullen in the class *pyrexia*, and order *phlegmasia*. It is known by pyrexia, pain in the region of the stomach, great anxiety, difficulty of breathing, cough, irregular pulse, palpitation, and fainting, and the other symptoms of inflammation.

CARDO. (A hinge.) The articulation called *Ginglymus*; also the second vertebra of the neck.

CARDONET. A wild artichoke, esculent.

CARDONIUM. So Paracelsus calls wine medicated with herbs.

CARDOPATIA. The low carline thistle, said to be diaphoretic.

CARDUUS. (d *carere*, quasi aptus carende lanæ, being fit to tease wool; or from καρδία, to abrade; so named from its roughness, which abrades and tears whatever it meets with.) The thistle, or teasel. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia equalis*.

CARDUUS ACANTHUS. The bear's breech.

CARDUUS ALEUTIS. The artichoke.

CARDUUS BENEDICTUS. *Cnicus sylvestris*. Blessed or holy thistle. *Centaurea benedicta*; *calycibus duplicato spinosis lanatis involucratiss, foliis semis decurrentibus denticulato-spinosis* of Linnaeus. Class, *Syngenesia*. Order, *Polygamia frustranea*. This exotic plant, a native of Spain and some of the Archipelago islands, obtained the name of *Benedictus*, from its being supposed to possess extraordinary medicinal virtues. In loss of appetite, where the stomach was injured by irregularities, its good effects

have been frequently experienced. It is a powerful bitter tonic and astringent. Bergius considers it as antacid, corroborant, stomachic, sudorific, diuretic, and ecceproptic. Chamomile flowers are now generally substituted for the *carduus benedictus*, and are thought to be of at least equal value.

CARDUUS HÆMORRHOIDALIS. (So called because it is said to relieve the pains of the hæmorrhoids, if beat into a poultice and applied.) Also called *carduus vinearum repens*, *sonchi folio*, *cirsium arvense*, *ceanothos*. The common creeping way thistle. *Serratula arvensis* of Linnaeus.

CARDUUS LACTEUS. See *Carduus Marie*.

CARDUUS LACTUS SYRIACUS. The Spanish milk-thistle. Stomachic and anodyne.

CARDUUS MARIÆ. *Carduus albis maculis notatus vulgaris*, C. B. Common milk-thistle, or Lady's thistle. The seeds of this plant, *Carduus marianus*; *foliis amplexicaulibus*, *hastato-pinnatifidis*, *spinosis*; *calycibus aphyllis*; *spinis canaliculatis*, *duplicato-spinosis*, of Linnaeus, and the herb have been employed medicinally. The former contains a bitter oil, and are recommended as relaxants. The juice of the latter is said to be salutary in dropsies, in the dose of four ounces; and, according to Millar, to be efficacious against pungent pains.

CARDUUS MARIANUS. The systematic name of the officinal *Carduus marie*.

CARDUUS SATIVUS. The artichoke.

CARDUUS SOLSTITIALIS. The calcitrapa officinalis.

CARDUUS TORRENTOSUS. The woolly thistle. The plant distinguished by this name in the pharmacopœias, is the *Oenopodium acanthium* of Linnaeus:—*calycibus squarrosis*; *squamis patentibus*; *foliis ovato-oblongis, sinuatis*. Its expressed juice has been recommended as a cure for cancer, either applied by moistening lint with it, or mixing some simple farinaceous substance, so as to form a poultice, which should be in contact with the disease, and renewed twice a-day.

CAREBRARIA. (From *careo*, the head, and *baros*, weight.) A painful and uneasy heaviness of the head.

CARENUM. (From *careo*, the head.) Galen uses this word for the head.

CARENUM VINUM. Strong wine.

CAREUM. From *Carea*, the country whence they were brought.) The caraway.

CAREX. (From *careo*, not *quia riribus careat*, but because, from its roughness, it is fit *ad carendum*, to card, tease, or pull.) Sedge. The name of a genus of plants in the Linnaean system. Class, *Monocotyledon*. Order, *Triandria*.

CAREX ARENARIA. The systematic name of the officinal *sarsaparilla Germanica*. See *Sarsaparilla Germanica*.

CARICA. (From *Carica*, the place

where they are cultivated.) *Ficus*. *Ficus vulgaris*. *Ficus communis*. *Sium* of the Greeks. The fig. The plant which affords this fruit is the *Ficus Carica* of Linnaeus. Fresh figs, are, when completely ripe, soft, succulent, and easily digested, unless eaten in immoderate quantities, when they are apt to occasion flatulency, pain of the bowels, and diarrhoea. The dried fruit, which is sold in our shops, is pleasanter to the taste, and more wholesome and nutritive. They are directed in the *decoctum hordei compositum*, and in the *electuarium lenitivum*. Applied externally, they promote the suppuration of tumours; hence they have a place in maturing cataplasms; and are very convenient to apply to the gums, and, when boiled with milk, to the throat.

CARICA PAPAYA. Papaw tree. Every part of the papaw tree, except the ripe fruit, affords a milky juice, which is used, in the Isle of France, as an effectual remedy for the tape-worm. In Europe, however, whether it has been sent in the concrete state, it has not answered, perhaps from some change it had undergone, or not having been given in a sufficient dose.

CARIUM. (From *Caricus*, its inventor.) *Carycum*. An ointment for cleansing ulcers, composed of hellebore, lead, and cantharides.

CARIES. (From *carah*, Chald.) Rotteness, or mortification of the bones.

CARINA. The cassada bread.

CARINA. A name formerly applied to the back-bone.

CARIUM TERRA. Lime.

CARIVILLANDI. A name of *sarsaparilla* root.

CARLINA. (From *Carolus*, Charles the Great, or Charlemagne; because it was believed that an angel shewed it to him, and that, by the use of it, his army was preserved from the plague.) Carline thistle. The name of a genus of plants in the Linnaean system. Class, *Syngenesia*. Order, *Polygamia equalis*. The official name of two kinds of plants. See *Chamæmelen album* and *Carlina gummiifera*.

CARLINA ACAULIS. The systematic name of the *chamæmelen album*.

CARLINA GUMMIIFERA. *Carduus pinea*. *Ixine*. Pine thistle. This plant is the *Attractylis gummiifera* of Linnaeus. The root, when wounded, yields a milky, viscous juice, which concretes into tenacious masses, at first whitish, resembling wax, when much handled growing black; it is said to be chewed with the same views as mastich.

Carline thistle. See *Chameleon album*.

CARLO SANCTO RADIX. St. Charles's root; so called by the Spaniards, on account of its great virtues. It is found in Mechoachan, a province in America. Its bark hath an aromatic flavour, with a bitter acrid taste. The root itself consists of

slender fibres. The bark is sudorific, and strengthens the gums and stomach.

CARMEN. (A verse; because charms usually consisted of a verse.) A charm; an amulet.

CARMES. (The Carmelite friars, Fr.) Carmelite water; so named from its inventors; composed of baum, lemon-peel, &c.

CARMINANTIA. See *Carminatives*.

CARMINATIVES. (*Carminativa*, sc. *medicamenta*: from *carmen*, a verse or charm; because practitioners, in ancient times, ascribed their operation to a charm or enchantment.) A term applied to those substances which allay pain, and dispel flatulencies of the primæ viæ. The principal carminatives are the semina cardamomi, anisi et carui; olea essentialia carui, anisi et juniperi; confectio aromatica; pulvis aromaticus; tinctura cardamomi; tinctura cinnamomi composita; zinziber; tonics, biters, and astringents.

CARNABADIUM. Caraway seed.

CARNÆ COLUMNÆ. The fleshy pillars or columns in the cavities of the heart. See *Heart*.

CARNICULA. (Dim. of *caro*, *carnis*, the flesh.) The fleshy substance which surrounds the gums.

CANTIFORNIS. (From *caro*, flesh, and *forma*, likeness.) Having the appearance of flesh. It is commonly applied to an abscess where the flesh surrounding the orifice is hardened and of a firm consistence.

CARO. Flesh. The red part or belly of a muscle; also the pulp of fruit.

CARO ADNATA. The recent swelled testicle.

CAROLINA. See *Carlinæ*.

CAROPI. The amomum verum.

CARORA. The name of a vessel that resembles an urinal.

CAROSIS. See *Carus*.

CAROTA. See *Daucus*.

CAROTIDÆ ARTERIÆ. See *Carotides*.

CAROTID ARTERY. (From *caro*, to cause to sleep; so called because, if tied with a ligature, they cause the animals to be comatose, and have the appearance of being asleep. The carotids are two considerable arteries that proceed, one on each side of the cervical vertebræ, to the head, to supply it with blood. The right carotid does not arise immediately from the arch of the aorta, but is given off from the arteria innominata. The left arises from the arch of the aorta. Each carotid is divided into external and internal, or that portion without and that within the cranium. The external gives off eight branches to the neck and face, viz. *anteriorly*, the superior thyroideal, the sublingual, the inferior maxillary, the external maxillary; *posteriorly*, the internal maxillary, the occipital, the external auditory, and the temporal. The internal carotid or cerebral artery, gives off four branches within the cavity of the cranium; the

anterior cerebral, the posterior, the central artery of the optic nerve, and the internal orbital.

CAROUN. The caraway seed.

CARPASUS. (So named *καρπας το καρον* because it makes the person who eats it appear as if he was asleep.) An herb, the juice of which was formerly called opocarpason, opocarpathon, or opocalpason: according to Galen it resembles myrrh; but is esteemed highly poisonous.

CARPATICUS BALSAMUS. *Oleum Germanis. Carpathicum.* This balsam is obtained both by wounding the young branches of the *Pinus cembra* of Linnæus: —*foliis quinis, levibus*, and by boiling them. It is mostly diluted with turpentine, and comes to us in a very liquid and pellucid state, rather white.

CARPENTARIA. (From *Carpentarius*, a carpenter; and so named from its virtues in healing cuts and wounds made by a tool.) A vulnerary herb; but not properly known what it is.

CARPHALEUS. (From *καρφα*, to exsiccate.) Hippocrates uses this word to mean *dry*, opposed to *moist*.

CARPUS. (From *καρπη*, a straw.) In Hippocrates it signifies a mote, or any small substance. A pustule of the smallest kind. Also the herb fenugreek.

CARPIA. (From *carpo*, to pluck off, as lint is from linen cloth.) Lint. See *Lint*.

CARPISNUS. The wrist.

CARPOBALSAMUM. (From *καρπος*, fruit, and *βασανισμος*, balsam.) See *Balsamum Gileadense*.

CARPOLOGAI. (From *carpo*, to pluck or pull gently.) Picking the clothes, as in dangerous fevers.

CARPUS. (*Καρπος*, the wrist.) The wrist, or carpus. It is situated between the fore arm and hand. See *Bones*.

Carrot. See *Daucus*.

Carrot, candy. See *Daucus Creticus*.

Carrot poultice. See *Cataplasma dauci*.

CARTHAMUS. (From *καθαρος*, to purge.) 1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygomia equalis*.

2. The pharmacopœial name of the saffron flower, *Carthamus tinctorius* of Linnæus: —*foliis ovatis, integris, serrato-aculeatis*: called also *Cnicus*, *Crocus Saracenicus*, *Carthamum officinarum*, *Carduus sativus*. The plant is cultivated in many places on account of its flowers, which are used as a yellow dye. The seeds, freed from their shells, have been celebrated as a gentle cathartic in a dose of one or two drachms. They are also supposed to be diuretic and expectorant; particularly useful in humoral asthma, and similar complaints. The *carthamus lanatus* is considered in France, as a febrifuge and sudorific. The dried flowers are frequently mixed with saffron, to adulterate it.

CARTHAMUS TINCTORIUS. The systematic name of the safflower plant. See *Carthamus*.

CARTHUSIANUS. (From the Monks of that order, who first invented it.) A name of the precipitated sulphur of antimony.

CARTILAGE. (Quasi *cartilago*; from *caro*, *carnis*, flesh.) A white elastic, glistening substance, growing to bones, and commonly called *gristle*. Cartilages are divided, by anatomists into *obducent*, which cover the moveable articulations of bones; *inter-articular*, which are situated between the articulations, and *uniting* cartilages, which unite one bone with another. Their use is to lubricate the articulations of bones, and to connect some bones by an immoveable connexion.

CARTILAGO ANNULARIS. See *Cartilago cricoidea*.

CARTILAGO ARYTÆNOIDEA. See *Larynx*.

CARTILAGO CRICOIDEA. The cricoid cartilage belongs to the larynx, and is situated between the thyroid and arytenoid cartilages and the trachea; it constitutes, as it were, the basis of the many annular cartilages of the trachea.

CARTILAGO ENSIFORMIS. *Cartilago xyphoidea*. Ensiform cartilage. A cartilage shaped somewhat like a sword or dagger, attached to the lowermost part of the sternum, just at the pit of the stomach.

CARTILAGO SCUTIFORMIS. See *Thyroid cartilage*.

CARTILAGO THYROIDEA. See *Thyroid cartilage*.

CARTILAGO XYPHOIDEA. See *Cartilago ensiformis*.

CARUL. (*Caruia*, Arabian.) The caraway. See *Carum*.

CARUM. (*Καρός*: so named from *Caria*, a province of Asia.) The caraway.

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

2. The pharmacopœial name of the caraway plant. It is also called *Carui*. *Cuminum pratense*. *Caros*. *Caruon*. And is the *Carum carui* of Linnæus. The seeds are well known to have a pleasant spicy smell, and a warm aromatic taste; and, on this account, are used for various economical purposes. They are esteemed to be carminative, cordial, and stomachic, and recommended in dyspepsia, flatulencies, and other symptoms attending hysterical and hypochondriacal disorders. An essential oil and distilled water are directed to be prepared from them by the London college.

CARUM CARUI. The systematic name for the plant whose seeds are called caraway. See *Carum*.

CARUNCLE. (Diminutive of *caro*, flesh.) *Caruncula*. A little fleshy excrescence; as the *caruncula myrtiformes*, *caruncula lachrymales*, &c.

CARUNCULA LACHRYMALIS. A long conoidal gland, red externally, situated in the internal canthus of each eye, before the union of the eyelids. It appears to be formed of numerous sebaceous glands, from which many small hairs grow. The hardened smegma observable in this part of the eye in the morning, is separated by this caruncle.

CARUNCULE CUTICULARES ALB. The nymphæ.

CARUNCULE MAMILLARES. The extremities of the tubes in the nipple.

CARUNCULE MYRTIFORMES. When the hymen has been lacerated by attrition, there remains in this place two, three, or four caruncles, which have received the name of myrtiform.

CARUNCULÆ PAPILLARES. The protuberances within the pelvis of the kidney, formed by the papillous substance of the kidney.

CARUNCULOSA ISCHURIA. A suppression of urine, from caruncles in the urethra.

CARUON. See *Carum*.

CARUS. (*Καρός*: from *καρᾶ*, the head, as being the part affected.) *Caros*. *Carosis*.

1. Insensibility and sleepiness, with quiet respiration.

2. A profound sleep, without fever. A lethargy.

3. The name of the caraway-seed.

CARYA. The cassia lignea.

CARYEDON. (From *καρύα*, a nut.) *Carydon*. A sort of fracture, where the bone is broken into small pieces, like the shell of a cracked nut.

CARYEDON. See *Caryedon*.

CARYOCOSTIUM. (From *καρυον*, the caryophyllus, and *κοστιον*, composed of the costus.) An electuary, named from its ingredients.

CARYOPHYLLATA. (From *καρυοφυλλον*, the caryophyllus; so named because it smells like the caryophyllus, or clove July flower.) *Herba benedicta*. *Caryophyllus vulgaris*. *Garyophilla*. *Janamunda*. Avens, or herb bennet. The root of this plant, *Geum urbanum* of Linnæus:—*floribus erectis, fructibus globosis villosis, aristis uncinatis nudis, foliis lyratis*, has been employed as a gentle styptic, corroborant, and stomachic. It has a mildly austere, somewhat aromatic taste, and a very pleasant smell, of the clove kind. It is also esteemed, on the continent, as a febrifuge.

CARYOPHYLLOIDES CORTEX. See *Cul-lawan*.

CARYOPHYLLUS. (*Καρυοφυλλον*: from *καρυον*, a nut, and *φυλλον*, a leaf; so named because it smells like the leaves of the Indian nut, or clove-tree.) The clove-tree.

The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Monogynia*.

CARYOPHYLLUM AROMATICUM.

The clove. The tree which affords this spice is the *Eugenia caryophyllata* of Willdenow, which grows in the East Indies, the Moluccas, &c. The clove is the unexpanded flower, or rather the calyx; it has a strong agreeable smell, and a bitterish, hot, not very pungent, taste. The oil of cloves, commonly met with in the shops, and received from the Dutch, is highly acrimonious, and sophisticated. Clove is accounted the hottest and most acrid of the aromatics; and, by acting as a powerful stimulant to the muscular fibres, may, in some cases of atonic gout, paralysis, &c. supersede most others of the aromatic class; and the foreign oil, by its great acrimony, is also well adapted for several external purposes: it is directed by several pharmacopœias, and the clove itself enters many official preparations.

CARYOPHYLLUM RUBRUM. *Tunica. Vetonica. Betonica. Cornaria. Caryophyllus hortensis.* Clove pink. Clove gilliflower. Clove July flower. This fragrant plant, *Dianthus caryophyllus* of Linnaeus:—*floribus solitariis, squamis calycinis subovatis, brevissimis, corollis crenatis*, grows wild in several parts of England; but the flowers, which are pharmaceutically employed, are usually produced in gardens: they have a pleasant aromatic smell, somewhat allied to that of clove spice: their taste is bitterish and sub-adstringent. These flowers were formerly in extensive use, but are now merely employed in form of syrup, as a useful and pleasant vehicle for other medicines.

CARYOPHYLLUS AROMATICUS AMERICANUS. The piper Jamaicensis.

CARYOPHYLLUS HORTENSIS. The caryophyllum rubrum.

CARYOPHYLLUS VULGARIS. The caryophyllata.

CARYOTIS. (From *raguor*, a nut.) *Caryota*. Galen uses this word to mean a superior sort of dates, of the shape of a nut.

CASCARILLA. (Dim. of *cascara*, the bark, or shell. Span.) *Chocarilla Eluteria. Eluteria.* The tree that affords the cascarilla bark, is the *Croton cascarilla* of Linnaeus. The bark comes to us in quills, covered upon the outside with a rough, whitish matter, and brownish on the inner side, exhibiting, when broken, a smooth, close, blackish brown surface. It has a lightly agreeable smell, and a moderately bitter taste, accompanied with a considerable aromatic warmth. It is a very excellent tonic, adstringent, and stomachic, and is deserving of a more general use than it has hitherto met with.

CASCHU. See *Catechu*.

Cashew nut. See *Anacardium occidentale*.

CASHOO. An aromatic drug of Hindostan, said to possess pectoral virtues.

CASIA. See *Cassia*.

CASMINARIS. The cassamunia of Bengal.

CASSA. (Arab.) The thorax, or breast.

CASSADA. See *Cassava*.

CASSAMUM. The fruit of the balsam tree.

CASSAVA. *Cassada. Cacavi. Cassare. Pain de Madagascar. Ricinus minor. Maniot. Yucca. Maniibar. Aipi. Aipima coxera. Aipipoca. Janipha.* The plant so called is the *Jatropha manihot* of Linnaeus. The leaves are boiled, and eaten as we do spinach. The root abounds with a milky juice, and every part, when raw, is a fatal poison. It is remarkable that the poisonous quality is destroyed by heat: hence the juice is boiled with meat, pepper, &c. into a wholesome soup, and what remains after expressing the juice, is formed into cakes, or meal, the principal food of the inhabitants. This plant, which is a native of three quarters of the world, is one of the most advantageous gifts of Providence, entering into the composition of innumerable preparations of an economical nature.

Cassada roots yield a great quantity of starch, called tapioca, exported in little lumps by the Brazilians, and now well known to us as diet for sick and weakly persons.

CASSADA. See *Cassava*.

CASSIA. (From the Arabic *katsia*, which is from *katsa*, to tear off; so called from the act of stripping the bark from the tree.) The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Monogynia*. Cassia and Seneca belong to this genus. See *Cassia fistularis*, and *Senna*.

CASSIA CARYOPHYLLATA. Clove bark tree. The bark of this tree, *Myrtus caryophyllata* of Linnaeus:—*pedunculis trifido-multifloris, foliis ovatis*, is a warm aromatic, of the smell of clovespice, but weaker, and with a little admixture of the cinnamon flavour. It may be used with the same views as cloves, or cinnamon.

CASSIA FISTULA. The systematic name of the purging cassia. See *Cassia fistularis*.

CASSIA FISTULARIS. *Cassia nigra. Cassia fistula. Alexandrina. Chaiarxambar. Canna. Cussia solutiva. Tlai. Kiem.* Purging cassia. This tree, *Cassia fistula* of Linnaeus:—*foliis quinquejugis ovatis acuminatis glabris, petiolis eglandulatis*, is a native of both Indies. The pods of the East-India cassia are of less diameter, smoother, and afford a blacker, sweeter, and more grateful pulp, than those which are brought from the West-Indies. Those pods which are the heaviest, and in which the seeds do not rattle on being shaken, are commonly the best, and contain the most pulp, which is the part medicinally employed, and to be obtained in the manner described in the pharmacopœias. The best pulp is of a bright shining black colour, and of a sweet taste, with a slight degree of acidity. It

has been long used as a laxative medicine, and being gentle in its operation, and seldom disturbing the bowels, is well adapted to children, and to delicate or pregnant women. Adults, however, find it of little effect, unless taken in a very large dose, as an ounce or more; and, therefore, to them this pulp is rarely given, but usually conjoined with some of the brisker purgatives. The official preparation of this drug, is the *confectio cassiæ*; it is also an ingredient in the *confectio sennæ*.

CASSIÆ ARABICÆ. The pulp of cassia.

CASSIÆ FLORES. What are called cassia flowers in the shops, are the flowers of the true cinnamon tree, *Laurus cinnamomum* of Linnæus. They possess aromatic and adstringent virtues, and may be successfully employed in decoctions, &c. in all cases where cinnamon is recommended. See *Cinnamomum*.

CASSIA LIGNEA. *Cortex canella Malabarica. Cassia lignea Malabarica. Xyl-cassia. Canella Malabarica et Javensis. Karva. Canella Cubana. Arbor Judaica. Cassia canella. Canellifera Malabarica. Cortex crassol. Cinnamomum Malabaricum. Cakacha canela.* Wild cinnamon-tree. Malabar cinnamon-tree, or cassia lignea tree. Cassia lignea is the bark of the *Laurus cassia* of Linnæus:—*foliis triplinerviis lanceolatis*, whose leaves are called *folia malabathri* in the shops. The bark and leaves abound with the flavour of cinnamon, for which they may be substituted; but in much larger doses, as they are considerably weaker.

Cassia, purging. See *Cassia fistularia*.

CASSIA SENNA. The systematic name of the plant which affords senna. See *Senna*.

CASSIÆ PULPA. See *Cassia fistularia*.

CASSOB. An obsolete term for kali.

CASSOLETA. Warm fumigations described by Marcellus.

CASSONADA. Sugar.

CASUMINUM. (Of uncertain derivation, perhaps Indian.) *Casuminar. Casmina. Risagon. Bengale Indorum.* The root, occasionally exhibited under one of these names, is brought from the East Indies. It comes over in irregular slices of various forms, some cut transversely, others longitudinally. The cortical part is marked with circles of a dusky brown colour: the internal part is paler, and unequally yellow. It possesses moderately warm, bitter, and aromatic qualities, and a smell like ginger. It is recommended in hysterical, epileptic, and paralytic affections.

CASTANEA. (*Castanea, nescio*: from *Castana*, a city in Thessaly, whence they were brought.) *Lopina. Mota. Glans Jovis Theophrasti.* Jupiter's acorn, and Sardinian acorn. The common chestnut. The fruit of this plant, *Fagus castanea* of

Linnæus:—*foliis lanceolatis, acuminato-seratis, subtus nudis*, are much esteemed as an article of luxury, after dinner. Toasting renders them more easy of digestion; but, notwithstanding, they must be considered as improper for weak stomachs. They are moderately nourishing, as containing sugar, and much farinaceous substance.

CASTANEA EQUINA. The horse-chestnut. See *Hippocastanum*.

Castanea flore albo. Coffee.

CASTLE-LEOD WATERS. A sulphureous spring in Rosshire, celebrated for the cure of cutaneous diseases and foul ulcers.

CASTOR FIBER. The generic name of the beaver. See *Castoreum Russicum*.

Castor. See *Castoreum Russicum*.

Castor oil. See *Ricinus*.

Custer, Russian. See *Castoreum Russicum*.

CASTOREUM RUSSICUM. (*Castoreum*, from *castor*, the beaver, *quasi castor*; from *castor*, the belly; because of the largeness of its belly; or *a castrando*, because he is said to castrate himself in order to escape the hunters.) *Castoreum Russian castor.* A peculiar concrete substance, obtained from the *Castor fiber* of Linnæus, or beaver, an amphibious quadruped inhabiting some parts of Prussia, Russia, Germany, &c.; but the greatest number of these animals is met with in Canada. The name of *castoreum* is given to two bags, situated in the inguinal regions of the male beaver, which contain a very odorous substance, soft, and almost fluid when recently cut from the animal, but which dries, and assumes a resinous consistence in process of time. This substance has an acrid, bitter and a nauseous taste; its smell is strong, aromatic, and even fetid. It is used medicinally, as a powerful antispasmodic in hysterical and hypochondriacal affections, and in convulsions, in doses of from 10 to 30 grains. It has also been successfully administered in epilepsy and tetanus. It is occasionally adulterated with dried blood, gum-ammoniacum, or galbanum, mixed with a little of the powder of castor, and some quantity of the fat of the beaver.

CASTORIUM. See *Castoreum Russicum*.

CASTRATION. *Celotomia. Orchotomia.* A chirurgical operation, by which a testicle is removed from the body.

CASTRENSIS. (From *castra*, a camp.) A name applied to those diseases with which soldiers, encamped in marshy places, are afflicted.

CATABASIS. (From *καταβασις*, (to descend.) A descent or operation downwards.

CATABIBASIS. (From *καταβίβας*, to come to descend.) An exclusion, or expulsion of the humours downwards.

CATABLACTUM. (From *καταβλάσκειν*, to be useless.) Hippocrates uses this word

to signify carelessness and negligence in the attendance on, and administration to the sick.

CATABLEMA. (From *καταβάλλω*, to throw round.) The outermost fillet, which secures the rest of the bandages.

CATABRONCHESIS. (From *κατα*, and *βρογχος* the throat; or *καταβρογχίζω*, to swallow.) The act of swallowing.

CATACAUMA. (From *καίω*, to burn.) A burn, or scald.

CATACAUSIS. (From *καίω*, to burn.) The act of combustion, or burning.

CATACCLIVENUS. (From *κατακλινόμεαι*, to lie down.) Keeping the bed, from the violence of a disease.

CATACECHRAMENUS. (From *κατακεχαρμνυμι*, to reduce to small particles.) Broken into small pieces. It is used of fractures.

CATACEBRASTICA. (From *κατακεκραννυμι*, to mix together.) Medicines which obtund the acrimony of humours, by mixing with them and reducing them.

CATACHLIDESIS. (From *καταχλιδάω*, to indulge in delicacies.) A gluttonous indulgence in sloth and delicacies, to the generation of diseases.

CATACHRYSTON. (From *καταχρίω*, to anoint.) An unguent, or ointment.

CATACHRYMA. An ointment.

CATACLASIS. (From *κατακλάω*, to break, or distort.) Distorted eyelids.

CATACLEIS. (From *κατα*, beneath, and *κλαίω*, the clavicle.) The subclavicle, or first rib, which is placed immediately under the clavicle.

CATACLINES. (From *κατακλινω*, to lie down.) One who, by disease, is fixed to his bed.

CATACLISIS. (From *κατακλινω*, to lie down.) A lying down. It means also incurvation.

CATACLYSMA. (From *κατακλύω*, to wash.) A clyster.

CATACLYSMUS. (From *κατακλύω*, to wash.) An embrocation. A dashing of water upon any part.

CATACHEKMONOS. (From *κατα*, and *κρημνός*, a precipice.) Hippocrates means, by this word, a swollen and inflamed throat, from the exuberance of the parts.

CATACBUSIS. (From *καταβύω*, to drive back.) A revulsion of humours.

CATADOULESIS. (From *καταδούλω*, to enslave.) The subduing of passions, as in a phrensy, or fever.

CATAEZESIS. (From *καταερίζω*, to repel.) A revulsion or rushing back of humours, or wind in the intestines.

CATÆONISIS. (From *καταϊονω*, to irrigate.) Irrigation by a plentiful affusion of liquor on some part of the body.

CATAGMA. (From *κατα*, and *αγω*, to break.) A fracture. Galen says a solution of the bone is called catagma, and *elcos* is a solution of the continuity of the flesh: that when it happens to a cartilage, it has

no name, though Hippocrates calls it *catagma*.

CATASMATICA. (From *κατασμα*, a fracture.) Catagmatics. Remedies proper for cementing broken bones, or to promote a callus.

CATAGOGE. (From *καταγομαι*, to abide.) The seat or region of a disease or part.

CATASTIOSIS. (From *καταστίνω*, to debilitate.) An imbecility and enervation of the strength and limbs.

CATALEPSIS. (From *καταλαμβάνω*, to seize, to hold.) *Catoche*. *Catochus*. *Congelatio*. *Detentio*. *Encatalepsis*; and by Hippocrates *uphonia*; by Antigenes *anautidia*; by Cælius Aurelianus *apprehensio*, *oppressio*; by *comprehensio*. *Apoplexia cataleptica* of Cullen. A sudden suppression of motion and sensation, the body remaining in the same posture that it was in when seized.

Dr. Cullen says, he has never seen the catalepsy except when counterfeited; and is of opinion, that many of those cases related by other authors, have also been counterfeited. It is said to come on suddenly, being only preceded by some languor of body and mind, and to return by paroxysms. The patients are said to be for some minutes, sometimes (though rarely) for some hours, deprived of their senses, and all power of voluntary motion; but constantly retaining the position in which they were first seized, whether lying or sitting; and if the limbs be put into any other posture during the fit, they will keep the posture in which they are placed. When they recover from the paroxysm, they remember nothing of what passed during the time of it, but are like persons awakened out of a sleep.

CATALOTICA. (From *καταλωω*, to grind down.) Medicines to soften and make smooth the rough edges and crust of cicatrices.

CATALYSIS. (*Καταλυσις*: from *καταλύω*, to dissolve or destroy.) It signifies a palsy, or such a resolution as happens before the death of the patient; also that dissolution which constitutes death.

CATAMARASMUS. (From *καταμαρῖνω*, to grow thin.) An emaciation, or resolution of tumours.

CATANASSESIS. (From *καταμασσομαι*, to manducate.) The grinding of the teeth, and biting of the tongue; so common in epileptic persons.

CATAMENIA. (From *κατα*, according to, and *μήν* the month.) *Menses*. The monthly discharge of blood from the uterus of females, between the ages of 14 and 45. Many have questioned whether the catamenial discharge arose from a mere rupture of vessels, or whether it was owing to a secretory action. There can be little doubt of the truth of the latter. The secretory organ is composed of the arterial

vessels situated in the fundus of the uterus. The dissection of women who have died during the time of their menstruating, proves this. Sometimes, though very rarely, women, during pregnancy, menstruate; and when this happens, the discharge takes place from the arterial vessels of the vagina. During pregnancy and lactation, when the person is in good health, the catamenia, for the most part, cease to flow. The quantity a female menstruates at each time, is very varied; depending on climate, and a variety of other circumstances. It is commonly, in England, from five to six ounces: it rarely exceeds eight. Its duration is from three to four, and sometimes, though rarely, five days. With respect to the nature of the discharge, it differs very much from pure blood; it never coagulates, but is sometimes grumous, and membranes like the decidua are formed in difficult menstruations: in some women it always smells rank and peculiar; in others it is inodorous. The use of this monthly secretion is to render the uterus fit for the conception and nutrition of the fetus; therefore girls rarely conceive before the catamenia appear, and women rarely after their entire cessation; but very easily soon after menstruation.

CATANANCE. Succory.

CATANIPHTHIS. (From *κατανιψω*, to wash.) Washed, or scoured. It is used by Hippocrates of a diarrhoea washed and cleansed by boiled milk.

CATANTELEMA. (From *κατασπασσω*, to pour upon.) A lotion by infusion of water, or medicated fluids.

CATANTELEUSIS. A medicated fluid.

CATAPASMA. (From *κατασπασσω*, to sprinkle.) *Catapastum. Consersio. Epipaston. Pusma sympasmata. Asperio. Aspergines.* The ancient Greek physicians meant by this, any dry medicine reduced to powder, to be sprinkled on the body. Their various uses may be seen in Paulus of Eginæ, lib. vii. cap. xiii.

CATAPAUSIS. (From *καταπαύω*, to rest, or cease.) That rest or cessation from pain which proceeds from the resolution of uneasy tumours.

CATAPLATES. (From *κατα*, against, and *πλατή*, a shield.) This word means a sling, a granado, or battery; and is also used to signify the medicine which heals the wounds and bruises made by such an instrument.

CATAPHORA. (From *καταφερο*, to make sleepy.) *Coma somnolentum.* A preternatural propensity to sleep. A mild apoplexy.

CATAPHORA ARTERITICA. Apoplexy from gout.

CATAPHORA COMA. Sanguineous apoplexy.

CATAPHORA EXANTHEMATICA. A lethargy in eruptive diseases.

CATAPHORA HYDROCEPHALICA. Serous apoplexy.

CATAPHORA SCORBUTICA. Apoplectic symptoms in scurvy.

CATAPHORA TIMOR. A lethargic disposition.

CATAPHRACTA. (From *καταφρασσω*, to fortify.) A bandage on the thorax.

CATAPLASMA. (From *καταπλασσω*, to spread like a plaster.) A poultice. The following are among the most useful.

CATAPLASMA ALUMINIS. This application was formerly used to inflammation of the eyes, which was kept up from weakness of the vessels; it is now seldom used, a solution of alum being mostly substituted.

CATAPLASMA ACETOSÆ. Sorrel poultice; *℞. Acetosz, ℥bj.* To be beaten in a mortar into pulp. A good application to scorbutic ulcers.

CATAPLASMA ERATUM. See *Cataplasma fermenti.*

CATAPLASMA CICUTÆ. Hemlock poultice. *℞. Herbz cicutæ exfoliatæ, ʒij, Aquæ fontanæ, ℥bj.* To be boiled till only a pint remains, when as much linseed meal as necessary is to be added. This is an excellent application to many cancerous and scrophulous ulcers, and other malignant ones; frequently producing great diminution of the pain of such diseases, and improving their appearance. Justamond preferred the herb fresh bruised.

CATAPLASMA CUMINI. This is a warm and stimulating poultice, and was formerly much used as an irritating antiseptic application to gangrenous ulcers, and the like. It is now seldom ordered.

CATAPLASMA DAUCI. Carrot poultice, *℞. Radicis dauci recentis, ℥bj.* Bruise it in a mortar into pulp. Some, perhaps with reason, recommend the carrots to be first boiled. The carrot poultice is employed as an application to ulcerated cancers, scrophulous sores of an irritable kind, and various inveterate malignant ulcers.

CATAPLASMA FERMENTI. Yeast cataplasm. Take of flour a pound; yeast half a pint. Mix and expose to a gentle heat, until the mixture begins to rise. This is a celebrated application in cases of sloughing and mortification.

CATAPLASMA LINI. Linseed poultice, *℞. Farinz lini, ℥ss. Aquæ serventis, ℥jss.* The powder is to be gradually sprinkled into the water, while they are quickly blended together with a spoon. This is the best and most convenient of all emollient poultices for common cases, and has, in a great measure, superseded the bread and milk one, so much in use formerly.

CATAPLASMA LITHARGYRI ACETATI. *℞. Aquæ lithargyri acetati, ʒj. Aquæ distill. ℥bj. Micæ panis, q. s. Misce.* Practitioners who place much confidence in the virtues of lead, often use this poultice in cases of inflammation.

CATAPLASMA QUERCUS MARINI. This is prepared by bruizing a quantity of the

marine plant, commonly called sea-tang, which is afterwards to be applied by way of a poultice. Its chief use is in cases of scrophula, white swellings, and glandular tumours more especially. When this vegetable cannot be obtained in its recent state, a common poultice of sea-water and oat-meal has been substituted by the late Mr-Hunter, and other surgeons of eminence.

CATAPLASMA SINAPEOS. See *Cataplasma sinapis*.

CATAPLASMA SINAPIS. Mustard cataplasm. Take of Mustard-seed, linseed, of each powdered half a pound; boiling vinegar, as much as is sufficient. Mix until it acquires the consistence of a cataplasm.

CATAPLEXIS. (From *πλησσειν*, to strike.) Any sudden stupefaction, or deprivation of sensation, in any of the members, or organs.

CATAPOISIS. (From *καταπινομαι*, to swallow down.) According to Aretaeus, it signifies the instruments of deglutition. Hence also catapodium.

CATAPOTIUM. (*Καταποτιον*) A pill.

CATAPSYXIS (From *ψυχειν*, to refrigerate.) A refrigeration without shivering, either universal, or of some particular part. A chilliness, or, as Vogel defines it, an uneasy sense of cold in a muscular or cutaneous part.

CATAPTOSIS. (From *καταπτωω*, to fall down.) It implies such a falling down as happens in apoplexies; or the spontaneous falling down of a paralytic limb.

CATAPUTIA. (From *καταπυτω*, to have an ill savour; or from the Italian, *cacapuzza*, which has the same meaning; so named from its foetid smell.) Spurge.

CATAPUTIA MAJOR. See *Ricinus*.

CATAPUTIA MINOR. The plant so called in the pharmacopoeias, is the *Euphorbia lathyris*; *umbella, quadrifida, dichotoma, foliis oppositis integerrimis*, of Linnæus. The seeds possess purgative properties; but if exhibited in an over dose, prove drastic and poisonous; a quality peculiar to all the *euphorbia*.

CATABRACT. (*Cataracta*; from *καταβασις*, to confound or disturb; because the sense of vision is confounded, if not destroyed.) The *Cálogo lentis* of Cullen.

Hippocrates calls it *γλαυχμα*. Galen, *σπηγμα*. The Arabians, *gutta opaca*. Celsus, *suffusio*. A species of blindness, arising almost always from an opacity of the crystalline lens, or its capsule, preventing the rays of light passing to the optic nerve. It commonly begins with a dimness of sight; and this generally continues a considerable time before any opacity can be observed in the lens. As the disease advances, the opacity becomes sensible, and the patient imagines there are particles of dust, or motes, upon the eye, or in the air, which are called *musce volitantes*. This opacity gradually increases till the person either becomes entirely blind, or

can merely distinguish light from darkness. The disease commonly comes on rapidly, though sometimes its progress is slow and gradual. From a transparent state, it changes to a perfectly white, or light gray colour. In some very rare instances, a black cataract is found. The consistence also varies, being at one time hard, at another entirely dissolved. When the opaque lens is either more indurated than in the natural state, or retains a tolerable degree of firmness, the case is termed, a *firm* or *hard* cataract. When the substance of the lens seems to be converted into a whitish or other kind of fluid, lodged in the capsule, the case is denominated a *milky* or *fluid* cataract. When the substance is of a midling consistence, neither hard nor fluid, but about as consistent as a thick jelly, or curds, the case is named a *soft* or *caseous* cataract. When the anterior or posterior layer of the crystalline capsule becomes opaque, after the lens itself has been removed from this little membranous sac, by a previous operation, the affection is named a *secondary membranous cataract*. There are many other distinctions made by authors. Cataract is seldom attended with pain; sometimes, however, every exposure to light creates uneasiness, owing probably to the inflammation at the bottom of the eye. The real cause of cataract is not yet well understood. Numbers of authors consider it as proceeding from a preternatural contraction of the vessels of the lens, arising from some external violence, though more commonly from some internal and occult cause. The cataract is distinguished from gutta serena, by the pupils in the latter being never affected with light, and from no opacity being observed in the lens. It is distinguished from hypopyon, staphyloma, or any other disease in the fore part of the eye, by the evident marks which these affections produce, as well as by the pain attending their beginning. But it is difficult to determine when the opacity is in the lens, or in its capsule. If the retina (which is an expansion of the optic nerve in the inside of the eye) be not diseased, vision may, in most cases, be restored, by either depressing the diseased lens, or extracting it entirely, which is termed couching.

CATARREUMA. (From *καταρρεω*, to flow from.) A catarrh, or defluxion of humours.

CATARREXIS. (From *καταρρεω*, to pour out.) A violent and copious eruption, or effusion, joined with *καλας*: it is a copious evacuation from the belly, and sometimes alone it is of the same signification. In Vogel's Nosology, it is defined, a discharge of pure blood from the intestines, such as takes place in dysentery.

CATARREXUS. (From *καταρρεω*, to flow from.) A word applied to diseases proceeding from a discharge of phlegm.

CATARROPSA PYTHATA. (*Katarrōpsa pythata*.) Tubercles tending downward; or, as Galen relates, those that have their apex on a depending part.

CATARROPOS NOUROS. (*Katarrōpos nouros*.) A remission of the disease, or its decline, and opposed to the paroxysm.

CATARRHUS. (From *καταρρῆω*, to flow down.) *Coryza.* A catarrh. An increased secretion of mucus from the membranes of the nose, fauces, and bronchiz, with fever, and attended with sneezing, cough, thirst, lassitude, and want of appetite. It is a genus of disease in the class *pyrexia*, and order *profusiva*, of Cullen. There are two species of catarrh, viz. *catarrhus à frigore*, which is very common, and is called a cold in the head; and *catarrhus à contagione*, the influenza, which sometimes seizes a whole city. Catarrh is also symptomatic of several other diseases. Hence we have the *catarrhus rubescens*; *scissis variolosa*, *verminosa*, *calculosa*, *phthisica*, *hysterica*, *à dentitione*, *gravidarum*, *metastolicarum*, &c.

CATARRHUS À FRIGORE. Catarrh from cold.

CATARRHUS EKLINSULANUS. Mumps, or cynanche parotidæa.

CATARRHUS À CONTAGIO. The influenza.

CATARRHUS SUPPOCATIVUS. The croup, or cynanche trachealis.

CATARRHUS VESICÆ. Strangury, with discharge of mucus.

CATARTISMUS. (From *καταρτίζω*, to make perfect.) According to Galen, it is a translation of a bone from a preternatural to its natural situation.

CATASARCA. (From *κατα*, and *σαρξ*, flesh.) The same as *Anasarca*.

CATASBERTIS. (From *κατα*, and *σβεννυμι*, to extinguish.) The resolution of tumours without suppuration.

CATASCHASMUS. (From *κατασχαζω*, to scarify.) Scarification.

CATASISIA. (From *κατα*, and *σειω*, to shake.) A concussion.

CATASPASMA. (From *κατασπασμα*, to draw backwards.) A revulsion or retraction of humours, or parts.

CATASTASMOUS. (From *κατα*, and *στέλλω*, to distil.) This is the name which the Greeks, in the time of Celsus, had for a distillation.

CATASTALTICUS. (From *κατασταλλω*, to restrain, or *σπασμα*, to contract.) It signifies styptic, astringent, repressing.

CATASTASIS. (*Καταστασις*.) The constitution, state, or condition of any thing.

CATATASIS. (From *κατατάσσω*, to extend.) In Hippocrates it means the extension of a fractured limb, or a dislocated one, in order to replace it. Also the actual replacing it in a proper situation.

CATAXIS. (From *καταξω*, to break.) A fracture. Also a division of parts by an instrument.

CATECHOMENUS. (From *κατα*, to resist.) Resisting and making ineffectual the remedies which have been applied or given.

CATECHU. *Catechu.* It is said, that, in the Japanese language, *kate* signifies a tree, and *chu*, juice.) *Terra Japonica.* Japan earth. An extract prepared in India, it was supposed till lately, from the juice of the *Mimosa catechu* of Linnæus:—*epinis stipularibus, foliis bipinnatis multijugis, glandulis purtialium singulis, spicis axillaribus geminis seu ternis pedunculatis*; by boiling the wood and evaporating the decoction by the heat of the sun. But the shrub is now ascertained to be an acacia, and is termed *Acacia catechu*. In its purest state, it is a dry, pulverable substance, outwardly of a reddish colour, internally of a shining dark brown, tinged with a reddish hue; in the mouth it discovers considerable adstringency, succeeded by a sweetish mucilaginous taste. It may be advantageously employed for most purposes where an adstringent is indicated, and is particularly useful in alvine fluxes, where astringents are required. Besides this, it is employed also in uterine profusiva, in laxity and debility of the viscera in general; and it is an excellent topical adstringent, when suffered to dissolve leisurely in the mouth, for laxities and ulcerations of the gums, aphthous ulcers in the mouth, and simular affections. This extract is the basis of several formulæ in our pharmacopœias, particularly of a tincture: but one of the best forms under which it can be exhibited, is that of a simple infusion in warm water with a proportion of cinnamon, for by this means it is at once freed of its impurities and improved by the addition of the aromatic. Fourcroy says that catechu is prepared from the seeds of a kind of palm, called areca.

CATEADION. (From *κατα*, and *αια*, a blade of grass.) An instrument, having at the end a blade of grass, or made like a blade of grass, which was thrust into the nostrils to provoke a hæmorrhage when the head ached. It is mentioned by Aretæus.

CATELLUS. (Dim. of *catulus*, a whelp.) A young whelp. Also a chymical instrument called a cupel, which was formerly in the shape of a dog's head.

CATHERESIS. (From *κατα*, to take away.) The subtraction or taking away any part or thing from the body. Sometimes it means an evacuation, and Hippocrates uses it for such. A consumption of the body, as happens without manifest evacuation.

CATHERETICA. (From *κατα*, to take away.) Medicines which consume or remove superfluous flesh.

CATHARMA. (From *καθαίρω*, to remove.) The excrements, or humours, purged off from the body.

CATHARMUS. (From *καθαίρω*, to remove.) A purgation of the excrements, or humours. A cure by incantation, or the royal touch.

CATHARSIA. (From *καθαίρω*, to purge.) Cathartics, having a purging property.

CATHARSIS. (From *καθαίρω*, to take away.) A purge, or purgation of the excrements, or humours, either medically or naturally.

CATHARTICS. (*Cathartica*, sc. *medicamenta*: *καθάρματα*: from *καθαίρω*, to purge.) Those medicines which, taken internally, increase the number of alvine evacuations. The different articles referred to this class of medicines are divided into five orders:

1. *Stimulating cathartics*, as jalap, aloes, and bitter apple, which are well calculated to discharge accumulations of serum, and are mostly selected for indolent and phlegmatic habits, and those who are hard to purge.

2. *Refrigerating cathartics*, as Glauber's salts, Epsom salts, sal polychrest, and cremor tartar. These are better adapted for plethoric habits, and those with an inflammatory diathesis.

3. *Adstringent cathartics*, as rhubarb and damask roses, which are mostly given to those whose bowels are weak and irritable, and subject to diarrhoea.

4. *Emollient cathartics*, as manna, malva, castor oil, and olive oil, which may be given in preference to other cathartics, to children and the very aged.

5. *Narcotic cathartics*, as tobacco, hyoscyamus, and digitalis. This order is never given but to the very strong and indolent, and to maniacal patients, as their operation is very powerful.

Murray, in his *Materia Medica*, considers the different cathartics under the two divisions of laxatives and purgatives; the former being mild in their operation, and merely evacuating the contents of the intestines; the latter being more powerful, and even extending their stimulant operation to the neighbouring parts. The following he enumerates among the principal laxatives:—Manna, *Cassia fistularis*, *Tamarindus Indica*, *Ricinus communis*, Sulphur, Magnesia. Under the head of purgatives, he names *Cassia senna*, *Rhæum palmatum*, *Convolvulus jalapa*, *Helleborus niger*, *Brionia alba*, *Cucumis colocynthis*, *Momordica elaterium*, *Rhamnus catharticus*, *Aloe perfoliata*, *Convolvulus scammonia*, *Gambogia gutta*, *Submurias hydrargyri*, Sulphas magnesæ, Sulphas sodæ, Sulphas potassæ, Supertartris potassæ, Tartris potassæ, Tartris potassæ et sodæ, Phosphas sodæ, Murias sodæ, Terebinthina veneta, *Nicotiana tabacum*.

CATHARTICUS SAL. See *Sulphas magnesæ*, and *Sulphas sodæ*.

CATHARTICUS HISPANICUS SAL. A kind of Glauber's salt, produced near Madrid, from some springs.

CATHARTICUS GLAUBERI SAL. See *Sulphas sodæ*.

CATHEDRA. (From *καθίζωμαι*, to sit.) The anus, or rather, the whole of the buttocks, as being the part on which we sit.

CATHERETICA. (From *καθαίρω*, to remove.) Corrosives. Medicines which, by corrosion, remove superfluous flesh.

CATHETER. (From *καθίμι*, to thrust into.) A long and hollow tube, that is introduced by surgeons into the urinary bladder, to remove the urine, when the person is unable to pass it. Catheters are either made of silver or of the elastic gum. That for the male urethra is much longer than that for the female, and so curved, if made of silver, as to adapt itself to the urethra.

CATHETERISMUS. (From *καθίμι*, a catheter.) The term given by P. Egrieta to the operation of introducing the catheter.

CATHIDRYSIS. (From *καθίδω*, to place together.) The reduction of a fracture. The operation of setting a broken bone.

CATHMIA. A name for litharge.

CATHODOS. (From *κατα*, and *ὄδος*.) A descent of humours.

CATHOLCEUS. (From *κατα*, and *ὀλκω*, to draw over.) An oblong fillet, made to draw over and cover the whole bandage of the head.

CATHOLICON. (From *κατα*, and *ὀλκω*, universal.) A panacea, or universal medicine. A term formerly applied to medicines that were supposed to purge all the humours.

CATHYPIA. (From *κατα*, and *υπνος*, sleep.) A profound but unhealthy sleep.

CATIAS. (From *καθίμι*, to place in.) An incision knife, formerly used for opening an abscess in the uterus, and for extracting a dead fœtus.

CATILLUS. See *Catellus*.

CATINUM ALUMEN. A name given to potash.

CATINUS. (*Καταγών*.) A crucible.

Catmini. See *Nepeta*.

CATOCATHARTICA. (From *κατα*, downwards, and *καθαίρω*, to purge.) Medicines that operate by stool.

CATOCHÆ. (From *κατεχω*, to detain.) See *Cataleptis*.

CATOCHEILUM. (From *κατα*, beneath, and *χῆλος*, the lip.) The lower lip.

CATOCCHUS. (From *κατεχω*, to detain.) A catalepsy. Also a tetanus or spasmodic disease in which the body is rigidly held in an upright posture.

CATOCCHUS CERVINUS. Tetanus, particularly affecting the neck.

CATOCCHUS DIVENUS. An occasional tetanus.

CATOCCHUS HOLOTONICUS. Another name for tetanus.

CATOMISEUS. (From *κατα*, below, and *αμος*, the shoulder.) By this word, P. Ægine expresses a method of reducing a luxated shoulder, by raising the patient over the shoulder of a strong man, that, by the weight of the body the dislocation may be reduced.

CATOPHIS. (From *κατατομαι*, to see clearly.) An acute and quick perception. The acuteness of the faculties which accompanies the latter stages of consumption.

CATOPTR. (From *κατα*, and *στρομα*, to see, and, by metaphor, to probe.) A probe. An instrument called a *speculum ani*.

CATORCHITES. (From *κατα*, and *ορχις* the orchis.) A wine in which the orchis root has been infused.

CATORETICA. (From *κατα*, downwards, and *ρεω*, to flow.) *Catoteretica*. *Catoterica*. Medicines which purge by stool.

CATORETICA. See *Catoteretica*.

CATULOTICA. (From *καταλυω*, to cicatrize.) Medicines that cicatrize wounds.

CATUTRIPALI. A name of the piper longum.

CAUCALIS. (From *καυκων*, a cup; or from *δευκαλις*, the daucus.) Bastard parsley, so named from the shape of its flower. Also, the wild carrot.

CAUCALOIDES. (From *caucalis*, and *ειδος*, a likeness; from its likeness to the flower of the caucalis.) The patella is sometimes so called.

CAUDA. (From *cado*, to fall; because it hangs or falls down behind.) A tail.

1. The tail of animals.

2. A name formerly given to the os coccygis, that being in tailed animals the beginning of the tail.

3. A fleshy substance, protuberating from the lips of the vagina of the pudendum muliebri, and resembling a tail, according to Ætius.

4. Many herbs are also named cauda, with the affixed name of some animal, whose tail the herb is supposed to be like; as *cauda equina*, horse-tail; *cauda muris*, mouse-tail; and in many other instances.

CAUDA EQUINA. The spinal marrow, at its termination about the second lumbar vertebra, gives off a large number of nerves, which, when unravelled, resemble the horse's tail; hence the name. See also *Equisetum*.

CAUDATIO. (From *cauda*, a tail.) An elongation of the clitoris.

CAUL. The English name for the omentum. See *Omentum*.

CAULEDON. (From *καυλος*, a stalk.) A transverse fracture, when the bone is broken, like the stump of a tree.

CAULIFLOWER. A species of brassica, whose flower is cut before the fructification expands. The observations which have been made concerning cabbages are applicable here. Cauliflower is, however, a fit more delicious vegetable.

CAULIS. (Kalab. A Chaldean word.)

1. The stem or stalk of a plant.

2. A cabbage.

3. It means too the penis of a man.

CAULIS FLORIDA. Cauliflower.

CAULODES. (From *καυλος*, the cabbage.) The white or green cabbage.

CAULOTON. (From *καυλος*, a stem: because it grows upon a stalk.) A name given to the beet.

CAUMA. (From *καω*, to burn.) The heat of the body or the heat of the atmosphere, in a fever.

CAUNGA. A name of the areca.

CAUSIS. (From *καω*, to burn.) A burn; or rather, the act of combustion, or burning.

CAUSODES. (From *καω*, to burn.) A term applied by Celsus to a burning fever.

CAUSOMA. (From *καω*, to burn.) An ardent or burning heat and inflammation. A term used by Hippocrates.

CAUSTIC ALKALI. The pure alkalis are so called. See *Alkali*.

Caustic barley. See *Cevadilla*.

CAUSTICS. (*Caustica*, sc. *medicamenta*; from *καω*, to burn; because they always produce a burning sensation.) See *Escharotics*.

CAUSTICUM AMERICANUM. The *cevadilla*.

CAUSTICUM ANTIMONIALE. See *Muriate of antimony*.

CAUSTICUM COMMUNE FORTIUS. See *Potassa cum calce*.

CAUSTICUM LUNARE. See *Nitras argenti*.

CAUSUS. (From *καω*, to burn.) A highly ardent fever. According to Hippocrates, a fiery heat, insatiable thirst, a rough and black tongue, complexion yellowish, and the saliva bilious, are its peculiar characteristics. Others also are particular in describing it; but whether ancients or moderns, from what they relate, this fever is no other than a continued ardent fever in a bilious constitution. In it the heat of the body is intense: the breath is particularly fiery; the extremities are cold; the pulse is frequent and small; the heat is more violent internally than externally, and the whole soon ends in recovery or death.

CAUSUS, ENDEMIAL. The name given, by Dr. Mosely, to the yellow fever of the West Indies.

CAUTERY. (From *καω*, to burn.) Cauteries were divided, by the ancients, into *actual* and *potential*; but the term is now given only to the red-hot iron, or *actual cautery*. This was formerly the only means of preventing hæmorrhages from divided arteries, till the invention of the ligature. It was also used in diseases, with the same view as we employ a blister. *Potential cautery* was the name by which *kali purum*, or *potassa*, was distinguished in the former dispensaries of Edinburgh,

Surgeons understand, by this term, any caustic application.

CAVA. The name of a vein, and also of the pudendum muliebre. See *Veins*.

CAVERNA. (From *cavus*, hollow.) A cavern. Also the name of the pudendum muliebre.

CAVIARE. *Caviarium.* A food made of the hard roes of sturgeon, formed into cakes, and much esteemed by the Russians.

CAVICULA. (Dim. of *cavilla*.) See *Ca-villa*.

CAVILLA. (From *cavus*.) The ankle, or hollow of the foot.

CAVITAS. (From *cavus*, hollow.) Any cavity, or hollowness. The auricle of the heart was formerly called the *cavitas innominata*, the hollow without a name.

Cayenne pepper. See *Piper Indicum*.

CAZABI. See *Cassala*.

CEASMA. (From *caeo*, to split, or divide.) *Cœmus.* A fissure, or fragment.

CEBER. (Arab.) The *agalochumi*. Also the *capparis*.

CEBIPIRA. (Indian.) A tree which grows in Brazil, decoctions of whose bark are used in baths and fomentations, to relieve pains in the limbs, and cutaneous diseases.

CEDAR. See *Cedrinum lignum*.

CEDMA. (From *cedo*, to disperse.) A defluxion, or rheumatic affection, scattered over the parts about the hips.

CEDRA, ESSENTIA DE. See *Bergamotte*

CEDRINUM LIGNUM. Cedar of Lebanon. *Pinus cedrus* of Linnæus. An odorous wood, more fragrant than that of the fir, but possessing similar virtues.

CEDRITES. (From *cedrus*, the cedar-tree.) Wine in which the resin which distills from the cedar-tree has been steeped.

CEDRIUM. Cedar. It is also a name for common tar, in old writings.

CEIDROKELA. The fruit of the citron-tree.

CEIDRONELLA. Turkey baum.

CEIDROSTIA. (From *cedrus*, the cedar-tree.) A name of the white bryony, which smells like the cedar.

CEIDRUS. (From *Kedron*, a valley where they grew abundantly.) The *Pinus cedrus* of Linnæus, or the cedar-tree.

CEIDRUS AMERICANA. The arbor vitæ.

CEIDRUS BACCIFERA. The sylvine.

CEIRIA. (From *ceiro*, to abrade.) The tape-worm; so called from its excoriating and abrading the intestines.

Calandine. See *Chetidonium majus*.

CELASTRUM. (From *celao*, a dart or pole, which it represents.) *Ceanothus Americanus* of Linnæus. Some noted Indians depend more on this than on the lobelia, for the cure of syphilis, and use it in the same manner as lobelia.

CÊLE. (From *celao*.) A tumour caused by the protusion of any soft part. Hence the compound terms *hydrocele*, *bubonocœle*.

CELEBY. The English name for a variety of the apium graveolens.

Celiac artery. See *Celiac artery*.

CELLIS. (From *celo*, to burn.) A spot or blemish upon the skin, particularly that which is occasioned by a burn.

CELLA TURCICA. See *Sella turcica*.

CELLULA. (Dim. of *cella*, a cell.) A little cell or cavity.

CELLULÆ MASTOIDEÆ. See *Temporal bone*.

CELLULAR MEMBRANE. *Membrana cellulosa* *Tela cellulosa*. *Panniculus adiposus.* *Membrana adiposa, pinguedinosa, et reticularis.* The cellular structure of the body, composed of laminae and fibres variously joined together, which is the connecting medium of every part of the body. It is by means of the communication of the cells of this membrane, that the butchers blow up their veal. The cellular membrane is, by some anatomists, distinguished into the reticular and adipose membrane. The former is evidently dispersed throughout the whole body, except the substance of the brain. It makes a bed for the other solids of the body, covers them all, and unites them one to another. The adipose membrane consists of the reticular substance, and a particular apparatus for the secretion of oil, and is mostly found immediately under the skin of many parts, and about the kidneys.

CELOTOMIA. (From *celo*, hernia, and *tomia*, to cut.) The operation for hernia.

CELSA. A term of Paracelsus's, to signify what is called the beating of the life in a particular part.

CEMENTERIUM. A crucible.

CENCHRANIS. (From *κενχρος*, millet.) A grain or seed of the fig.

CENCHRIBUS. A species of Herpes that resembles *κενχρος*, or millet.

CENHANGIA. (From *κενο*, to empty, and *αγγος*, a vessel.) The evacuation of blood, or other fluids, from their proper vessels.

CENISDAM. *Ceniplam.* *Cenigotam.* *Cenipolam.* The name of an instrument anciently used for opening the head in epilepsies.

CENIOTENIUM. A purging remedy, formerly of use in the venereal disease, supposed to be mercurial.

CENOSIS. (From *κενο*, empty.) Evacuation. It must be distinguished from *Catharsis*. *Cenosis* imports a general evacuation; *Catharsis* means the evacuation of a particular humour, which offends with respect to quality.

CENTAUREA. (So called from *Chiron*, the centaur, who is said to have employed one of its species to cure himself of a wound accidentally received, by letting one of the arrows of Hercules fall upon his foot.)

The name of a genus of plants in the Linnæan system, of the Order *Polygamia frustanea*. Class, *Syngenesia*.

CENTAURIA BEHEN. The systematic name of the official *behen albuma*. See *Behen album*.

CENTAURIA BENEDICTA. The systematic name of the blessed thistle. See *Cardus benedictus*.

CENTAURIA CALCITRAPA. The systematic name of the calcitrapa. See *Calcitrapa*.

CENTAURIA CYANUS. The systematic name of the plant which affords the *flores cyani*. See *Cyanus*.

CENTAURIODES. The gratiola.

CENTAURIUM MAGNUM MAJUS. The great centaury.

CENTAURIUM. (From *centaurus*, a centaur; so called because it was feigned that Chiron cured Hercules's foot, which he had wounded with a poisonous arrow, with it.) *Centaureum minus vulgare*. *Centaureum parvum*. *Centaureum minus*. *Centaury*. *Chironia centaureum* of Linnaeus:—*caeruleis quinquefidis infundibuliformibus, caule dichotomo, pistillo simplici*. This plant is justly esteemed to be the most efficacious bitter of all the medicinal plants indigenous to this country. It has been recommended, by Cullen, as a substitute for gentain, and by several is thought to be a more useful medicine. The tops of the centaury plant are directed for use by the colleges of London and Edinburgh, and are most commonly given in infusion; but they may also be taken in powder, or prepared into an extract.

Centaury. See *Centaureum*.

CENTAURIUM MINUS. See *Centaureum*.

CENTIMORBIA (From *centum*, a hundred, and *morbus*, a disease.) Nummularia, or moneywort; named from its supposed efficacy in the cure of a multitude of disorders.

CENTINODIA. (From *centum*, a hundred, and *nodus*, a knot.) The herb *Polygonum*; so called from its many knots, or joints.

CENTIPEDES. (From *centum*, a hundred, and *pes*, a foot.) Woodlice, named from the multitude of their feet.

CENTRATIO. (From *centrum*, a centre.) The concentration and affinity of certain substances to each other. Paracelsus expresses by it the degenerating of a saline principle, and contracting a corrosive and exulcerating quality. Hence *Centrum Salis* is said to be the principle and cause of ulcers.

CENTRIUM. (From *centrum*, to prick.) A plaster recommended by Galen against stiches and pricks in the side.

CENTRUM. (From *centrum*, to point or prick.) The middle point of a circle. In chymistry it is the residence or foundation of matter. In medicine, it is the point in which its virtue resides. In anatomy, the middle point of some parts is so

named, as *centrum nervorum* the middle or tendinous part of the diaphragm.

CENTRUM OVALE. When the two hemispheres of the brain are removed on a line with the level of the *corpus callosum*, the internal medullary part presents a somewhat oval centre; hence it is called *centrum ovale*. Vieussenius supposed all the medullary fibres met at this place.

CENTRUM TENDINOSUM. The tendinous centre of the diaphragm is so called. See *Diaphragm*.

CENTRUM NERVEUM. The centre of the diaphragm is so called. See *Diaphragm*.

CENTURNODIA. (From *centum*, a hundred, and *nodus*, a knot; so called from its many knots, or joints.) Common knot-grass. This plant, *Polygonum aviculare* of Linnaeus, is never used in this country; it is said to be useful in stopping hæmorrhages, diarrhæas, &c. but little credit is to be given to this account.

CENTUNCULUS. Bastard pimpernel.

CEPA. (From *capere*, a woolcard, from the likeness of its roots.) The onion. This bulbous root belongs to the *Allium cepa* of Linnaeus:—*scapo nudo inferne verticillis longioribus, foliis teretibus*. Dr. Cullen says, onions are acrid and stimulating, and possess very little nutriment. With bilious constitutions they generally produce flatulency, thirst, head-ach, and febrile symptoms; but where the temperament is phlegmatic, they are of infinite service, by stimulating the habit and promoting the natural secretions, particularly expectoration and urine. They are recommended in scorbutic cases, as possessing antiscorbutic properties. Externally, onions are employed in suppurating poultices, and suppression of urine in children is said to be relieved by applying them, roasted, to the pubes.

CEPÆA. A species of onion which used to be esteemed for salads in spring; but is now disregarded.

CEPHALÆA. (From *κεφαλή*, the head.) The flesh of the head, which covers the skull. Also a long continued pain of the cerebrum, and its membranes.

CEPHALALGIA. (From *κεφαλή*, the head, and *αλγος*, pain.) *Cephalæa*. The head-ach. It is symptomatic of very many diseases, but is rarely an original disease itself. When mild, it is called cephalalgia; when inveterate, cephalæa. When one side of the head only is affected, it takes the names of *hemicephalia*, *migrana*, *hemipalgia*, and *megrim*; in one of the temples only, *crotophæa*; and that which is fixed to a point, generally in the crown of the head, is distinguished by the name of *clavus*.

CEPHALALGIA CATARRHALIS. Head-ach, with catarrh, from cold.

CEPHALALGIA INFLAMMATORIA. Phrenitis, or inflammation of the brain.

CEPHALALGIA SPASMODICA. The sick head-ach. A species of indigestion.

CEPHALARTICA. (From *κεφαλη*, the head, and *artē*, to make pure.) Medicines which purge the head.

CEPHALE. (*Κεφαλη*.) The head.

CEPHALIC VEIN. (*Vena cephalica*; so called, because the head was supposed to be relieved by opening it.) The anterior vein of the arm, that receives the cephalic of the thumb.

CEPHALICA. (From *κεφαλη*, the head.) Cephalics. Such remedies as are adapted for the cure of disorders of the head. Of this class are the snuffs, which produce a discharge from the mucous membrane of the nose, &c.

CEPHALICA FOLLIES. A branch from the cephalic vein, sent off from about the lower extremity of the radius, running superficially between the thumb and the metacarpus.

CEPHALICUS FULVIS. A powder prepared from *asarum*.

CEPHALINE. (From *κεφαλη*, the head.) The head of the tongue. That part of the tongue which is next the root and nearest the fauces.

CEPHALITIS. (From *κεφαλη*, the head.) Inflammation of the head. See *Phrenitis*.

CEPHALORRHOEA. (From *κεφαλη*, the head, and *ρως*, a disease.) This term is applied to the febris hungarica, in which the head is principally affected.

CEPHALO-PHARYNGEUS. (From *κεφαλη*, the head, and *φαρυγξ*, the throat.) A muscle of the pharynx, otherwise named *constrictor pharyngis inferior*; which see.

CEPHALOPONTIA. (From *κεφαλη*, the head, and *πονος*, pain.) Head-ach; heaviness of the head.

CEPINI. Vinegar.

CEPULA. Large myrobalans.

CERA. Wax. Bees' wax. A solid concrete substance, collected from vegetables by bees; and extracted from their combs after the honey is got out, by heating and pressing them. With rectified spirit it forms, by the assistance of heat, a gelatinous liquid. It is perfectly insoluble in watery liquors. When melted, it assumes the appearance of oil, and in this state is easily combined with oils and liquid fats. It is very inflammable, and burns totally away. In the state in which it is obtained from the combs, it is called yellow wax, *cera flavo*; and this, when new, is of a lively yellow colour, somewhat tough, yet easy to break; by age it loses its fine colour, and becomes harder and more brittle. Yellow wax, after being reduced into thin cakes, and bleached by a long exposure to the sun and open air, is again melted, and formed into round cakes, called virgin's wax, or white wax, *cera alba*.

The chief medicinal use of wax, is in plasters, unguents, and other like external applications, partly for giving the requisite consistence to other ingredients, and partly on account of its own emollient quality.

CERA ALBA. See *Cera*.

CERA DICARDO. The carduus pinea.

CERA FLAVA. Yellow wax. See *Cera*.

CERNE. (From *κερας*, a horn.) So Rufus Ephesius calls the cornua of the uterus.

CERANITES. (From *κερανισμι*; to temper together.) A name formerly applied to a pastel, or troch, by Galen.

CERAS. (*Κερας*, a horn.) A wild sort of parsnip is so named from its shape.

CERASA NIGRA (*Κερασε*, the cherry-tree; from *κεραση*, a town in Pontus, whence Lucullus first brought them to Rome; or from *ang*, the heart; from its resemblance to it in shape and colour.) *Eucosmia*, because cherries keep the body open. The black cherry. The ripe fruit of the *Prunus avium* of Linnaeus:—*umbellulis sessilibus, foliis ovato-lanceolatis, subtus pubescentibus, conduplicatis*. The flavour of these is esteemed by many, and if not taken in too large quantities, they are extremely salutary. A gum exudes from the tree, whose properties are similar to those of gum arabic.

CERASA RUBRA. *Cerasa sativa*, or *anglica*. The red cherry. The ripe fruit of the *Prunus cerasus* of Linnaeus:—*umbellulis subpedunculatis, foliis ovato-lanceolatis, glabris, conduplicatis*. This species possesses a pleasant, acidulated, sweet flavour, and is extremely proper in fevers, scurvy, and bilious obstructions. Red cherries are mostly eaten as a luxury, and are very wholesome, except to those whose bowels are remarkably irritable.

CERASIATUM. (From *cerasus*, a cherry.) A purging medicine in Libavius; so called because the juice of cherries is an ingredient.

CERASTUS. *Craetes*. (From *cerasus*, a cherry.) The name of two ointments in Mesuë.

CERASMA. (From *κερανισμι*, to mix.) A mixture of cold and warm water, when the warm is poured into the cold.

CERASUS. (From *Κερασειον* a town in Pontus, from whence Lucullus is said first to have brought them to Rome.) The cherry-tree.

CERATE. (From *cerum*, wax.) *Ceratum* The obsolete synonyms, are *cerelea*, *ceromata*, *ceronia*, *cerota*. *Ceratomalgamata*. Cerates take their name from the wax which enters into their composition, and to which they owe their consistence, which is intermediate between that of plasters and that of ointments; though no very definite rule

for this consistence is, in fact, either given or observed.

CERATIA. (From *apac*, a horn, which its fruit is supposed to resemble.) *Ceratium*. *Ceratomia*. The siliqua dulcis.

CERATIA DIPHYLLUS. See *Courbaril*.

CERATO-GLOSSUS. (From *apac*, a horn, and *glossa*, a tongue.) A muscle, so named from its shape and insertion into the tongue. See *Hyoglossus*.

CERATO-HYOIDEUS. (From the *os hyoides*.) See *Stylo-hyoides*.

CERATOIDES. (From *apac*, the genitive of *apac*, a horn.) See *Cornea*.

CERATO-MALGAMA. A cerate.

CERATONIA SILLIQUA. The systematic name of the plant which affords the sweet pod. See *Siliqua dulcis*.

CERATUM. See *Cerate*.

CERATUM COMMUNE. Common cerate. Take of olive oil, six fluid ounces; yellow wax, four ounces. Having melted the wax, mix in the oil.

CERATUM ALBUM. See *Ceratum cetacei*.

CERATUM CALAMELANOS. \mathfrak{z} Calom. \mathfrak{z} j. Cerat. calam. \mathfrak{z} ss. Misce. Some practitioners are partial to this as a dressing for chancre.

CERATUM CALAMINAE. Formerly called *ceratum lapidis calaminaris*, and *ceratum epuloticum*. Calamine cerate. Take of prepared calamine, yellow wax, of each half a pound; olive oil, a pint. Mix the oil with the melted wax; then remove it from the fire, and, as soon as it begins to thicken, add the calamine, and stir it constantly, until the mixture becomes cold. A composition of this kind was first introduced under the name of Turner's cerate. Its virtue is desiccative, epulotic, and is well calculated to promote the cicatrization of ulcers.

CERATUM CETACEL. *Ceratum spermatis ceti*. *Ceratum album*. Spermaceti cerate. Take of spermaceti, half an ounce; white wax, two ounces; olive oil, 4 fluid-ounces. Add the oil to the spermaceti and wax, previously melted together, until the mixture becomes cold. This cerate is cooling and emollient, and applied to excoriations, &c. and may be applied with advantage to all ulcers, where no stimulating substance can be applied, being extremely mild and unctuous.

CERATUM CICUTAE. Hemlock cerate. \mathfrak{z} . unguenti cicutae lb. \mathfrak{z} j. Spermatis ceti \mathfrak{z} j. Cerat. alb. \mathfrak{z} iii. Misce. One of the formulæ of St. Bartholomew's hospital, occasionally applied to cancerous, scrophulous, phagedenic, herpetic, and other inveterate sores.

CERATUM CITRINUM. See *Ceratum resinæ*.

CERATUM EPULOTICUM. See *Ceratum calamine*.

CERATUM LAPIDIS CALAMINARIS. See *Ceratum calamine*.

CERATUM LITHARGYRI ACETATI COMPOSITUM. See *Ceratum plumbi compositum*.

CERATUM LITTE. *Ceratum cantharidis*. Cerate of blistering fly. Take of spermaceti cerate, six drachms; blistering flies, very fine powder, a drachm. Having softened the cerate by heat, add the flies, and mix them together.

CERATUM PLUMBI SUPERCACETATIS. *Unguentum ceruæ acetate*. Cerate of superacetate of lead. Take of superacetate of lead, powdered, two drachms; white wax two ounces; olive oil, a pint. Dissolve the wax in seven fluid-ounces of oil; then gradually add thereto the superacetate of lead, separately rubbed down with the remaining oil, and stir the mixture with a wooden slice, until the whole has united. This cerate is cooling and desiccative.

CERATUM PLUMBI COMPOSITUM. *Ceratum lithargyri acetati compositum*. Compound cerate of lead. Take of solution of lead, two fluid-ounces and a half; yellow wax, four ounces; olive oil, nine fluid ounces; camphor, half a drachm. Mix the wax previously melted, with eight fluid ounces of oil; then remove it from the fire, and, when it begins to thicken, add gradually the solution of acetate of lead, and constantly stir the mixture with a wooden slice, until it gets cold. Lastly, mix in the camphor, previously dissolved in the remainder of the oil. Its virtues are cooling, desiccative, resolvent against chronic rheumatism, &c. &c.; and as a proper application to superficial ulcers, which are inflamed.

CERATUM RESINÆ. *Ceratum resinæ flavæ*. *Ceratum citrinum*. Resin cerate. Take of yellow resin, yellow wax, of each a pound; olive oil, a pound. Melt the resin and wax together, over a slow fire; then add the oil, and strain the cerate, while hot, through a linen cloth. Digestive.

CERATUM SABINÆ. Savine cerate. Take of fresh leaves of savine, bruised, a pound; yellow wax, half a pound; prepared lard, two pounds. Having melted together the wax and lard, boil therein the savine leaves, and strain through a linen cloth. This article is of late introduction, for the purpose of keeping up a discharge from blistered surfaces. It was first described, by Mr. Crowther, and has since been received into extensive use, because it does not produce the inconveniences that follow the constant application of the common blistering cerate. A thick white layer forms daily upon the part, which requires to be removed, that the cerate may be applied immediately to the surface from which the discharge is to be made.

CERATUM SAPONIS. Soap cerate. Take of hard soap, eight ounces; yellow wax, ten ounces; semi-vitreous oxyd of lead, powdered, a pound; olive oil, a pint; vinegar, a gallon. Boil the vinegar, with the oxyd of lead, over a slow fire, constantly stirring,

until the union is complete; then add the soap, and boil it again in a similar manner, until the water is entirely evaporated; then mix in the wax, previously melted with the oil. Resolvent; against arophulous tumours, &c. It is a convenient application in fractures, and may be used as an external dressing for ulcers.

CERATUM SPERMATISCONTI. See *Ceratum crotali*.

CERBERUS. (Κεραυρ.) A fanciful name given to the compound powder of scammony, because, like the dog Cerberus, it has three heads, or principal ingredients, each of which is eminently active.

CERCIVALEX. (From *κερα*, to make a noise.) A wheezing, or bubbling noise, made by the trachea, in breathing.

CERCINUS. (From *κερα*, to wheeze.) Wheezing.

CERCINOSUS. (From *κερα*, to wheeze.) One who labours under a dense breathing, accompanied with a wheezing noise.

CERCINODES. The same as *cercinodes*.

CERCIS. (From *κερα*, to shriek.) This word literally means the spoke of a wheel, and has its name from the noise which wheels often make. In anatomy it means the radius, a bone supposed to be like a spoke. Also a pebble, from its shape.

CERCOSIS. (From *κερα*, a tail.) A polypus of the uterus. It is sometimes applied to an enlargement of the clitoris.

CERA. (From *cera*, wax.) The cerumen aurium, or wax of the ear.

CEREALIA. (Solemn feasts to the goddess Ceres.) All sorts of corn, of which bread or any nutritious substance is made, come under the head of *cerealia*, which term is applied by bromatologists as a genus.

CEREBELLA URINA. (Paracelsus thus distinguishes urine which is whitish, of the colour of the brain, and from which he pretended to judge of some of its distempers.

CEREBELLUM. (Dim. of *cerebrum*.) The little brain. A somewhat round viscus, of the same use as the brain; composed, like the brain, of a cortical and medullary substance, divided by a septum into a right and left lobe, and situated under the tentorium, in the inferior occipital fossæ. In the cerebellum are to be observed the *crura cerebelli*, the fourth ventricle, the *valvula magna cerebri*, and the *pro tuberantia vermiciformis*.

CEREBRUM. (Quasi *cerebrum*; from *κερα*, the head.) The brain. A large round viscus, divided superiorly into a right and left hemisphere, and inferiorly into six lobes, two anterior, two middle, and two posterior; situated within the cranium, and surrounded by the dura and pia mater, and tunica arachnoidea. It is composed of a cortical substance, which is external; and a medullary, which is internal. It has three

convolutes, called *ventricles*; two anterior, or lateral, which are divided from each other by the *septum lœidam*; and in which is the *choroid plexus*, formed of blood-vessels; the third ventricle is a space between the thalami nervorum optico-rum. The principal prominences of the brain are, the *corpus callosum*, a medullary eminence, conspicuous upon laying aside the hemispheres of the brain; the *corpus striatum*, two striated protuberances, one in the anterior part of each lateral ventricle; the *thalami nervorum optico-rum*, two whitish eminences behind the former, which terminate in the optic nerves; the *corpora quadrigemina*, four medullary projections called by the ancients, *nates* and *testes*; a little cerebrine tubercle lying upon the nates, called the *pineal gland*; and lastly, the *crura cerebri*, two medullary columns which proceed from the basis of the brain to the *medulla oblongata*. The cerebral arteries are branches of the carotid and vertebral arteries. The veins of the head are called *sinusæ*, which return their blood into the internal jugulars. The use of the brain is to give off nine pairs of nerves, through whose means the various senses are performed, and muscular motion excited.

CEREBRUM ELONGATUM. The medulla oblongata.

CEREPOLIUM. (A corruption of *chærophyllyum*.) *Chærophyllyum*. *Cerefolium*. Chervil. This plant, *Scandix cerefolium* of Linnæus:—*seminibus nitidis, ovato-subulatis; umbellis erectissimis, lateratibus*, is a salubrious culinary herb, sufficiently grateful both to the palate and stomach, slightly aromatic, gently aperient, and diuretic.

CEREPOLIUM HISPARTUM. The plant called by us Sweet-cicely.

CEREPOLIUM SYLVESTRIS. See *Cerefolium*.

CEREI MEDICATI. See *Bougies*.

CERELUM. (From *κερα*, wax, and *ελαιον*, oil.) A cerate, or liniment, composed of wax and oil. Also the oil of tar.

CERVISIE FERMENTUM. Yeast.

CERVISIA. (From *cereus*, corn, of which it is made.) Ale. Beer. Any liquor made from corn.

CERVISIE CATAPLASMA. Into the grounds of strong beer stir as much oatmeal as will make it of a suitable consistence. This is sometimes employed as a stimulant and antiseptic to mortified parts.

CERRA. (From *cereus*, soft, taper.) *Cerræ*. The flat worms which breed in the intestines.

CERION. (From *κερα*, a honey-comb.) A kind of achor.

CEROMA. (From *κερα*, wax.) *Ceromum*. Terms used by the ancient physicians for an unguent, or cerate, though originally applied to a particular compo-

sition which the wrestlers used in their exercises.

CEROPISSUS. (From *κερος*, wax, and *αμρα*, pitch.) A plaister composed of pitch and wax.

CEROTUM. (*Κερωτον*.) A cerate.

CERUMEN AURIS. (*Cerumen*; dim. of *cera*, wax.) *Aurium ceras.* *Marmorata aurium.* *Cypsel.* *Cypselus fugile.* The waxy secretions of the ears, situated in the meatus auditorius externus.

GERUSSA. (Arab.) Ceresse, or white lead. A subcarbonate of lead. See *Subcarbonas plumbi*.

CERUSSA ACETATA. See *Super-acetas plumbi*.

CERVI SPINA. See *Rhamnus catharticus*.

CERVICAL. (*Cervicalis*; from *cervix*, the neck.) Belonging to the neck; as cervical nerves, cervical muscles, &c.

CERVICAL ARTERIES. *Arterie cervicales.* Branches of the subelavians.

CERVICAL VERTEBRE. The seven uppermost of the vertebrae, which form the spine. See *Vertebra*.

CERVICARIA. (From *cervix*, the neck; so named because it was supposed to be efficacious in disorders and ailments of the throat and neck.) The herb throatwort.

CERVIX. (Quasi *cerebri vis*; as being the channel of the spinal marrow.) The neck. That part of the body which is between the head and shoulders. The cervix uteri is the neck of the uterus; or that part of it which is immediately above or beyond the os tunc. This term is also applied to other parts, as cervix vesicæ, ossis, &c.

CERYTHES. (From *κερος*, betony.) Wine impregnated with betony.

CESTRUM. (From *κερα*, a dart; so called from the shape of its flowers, which resemble a dart; or because it was used to extract the broken ends of darts from wounds.) The herb betony.

CATERACH. (Blanchard says this word is corrupted from *Pteryga*, *πτερυγες*, q. v. as pteryga, cetergya, and ceterach.) *Scelopendria vera.* *Doradilla?* Spicewort. Miltwaste. This small bushy plant, *Asplenium ceterach* of Linnaeus; *frondibus pinnatifidis, lobis alternis confusius obtusis*, grows upon old walls and rocks. It has an herbaceous, mucilaginous, roughish taste, and is recommended as a pectoral. In Spain it is given, with great success, in nephritic and calculous diseases.

CEVABILLA. (Dim. of *cevada*, barley. Spanish.) *Cevadilla Hispanorum.* *Sevadilla.* *Sabadilla.* *Hordeum causticum.* *Canis intersector.* Indian caustic barley. The plant whose seeds are thus denominated, is a species of *veratrum*; they are powerfully caustic, and are administered with very great success as a vermifuge. They

are also diuretic and emetic. The dose to a child, from two to four years old, is two grains; from hence to eight, five grains; from eight to twelve, ten grains.

Cayenne pepper. See *Capicum*.

CHKA. A Chinese name for tea.

CHEROPOLIUM. See *Cerefolium*.

CHEROPHYLLUM. (*Χαροφυλλον*; from *χαρο*, to rejoice, and *φυλλον*, a leaf; so called from the abundance of its leaves.) *Chervi*.

1. The name of a genus of plants in the Linnaean system. Class, *Pentandria*. Order, *Digynia*.

2. The pharmacopœial name of some plants. See *Cicentaria* and *Cerefolium*.

CHEROPHYLLUM SYLVESTRIS. See *Cicentaria*.

CHACARILLE CORTEX. See *Cascarilla*.

CHETA. (From *χεω*, to be diffused.) The human hair.

CHALASIS. (From *χαλαω*, to relax.) Relaxation.

CHALASTICA. (From *χαλαω*, to relax.) Medicines which relax.

CHALAZION. (From *χалаζα*, a hailstone.) *Chalaza.* *Chalazium Grands.* An indolent, moveable tubercle, on the margin of the eyelid, like a hailstone. A species of hordeolum. It is that well-known affection of the eye, called a sty, or stian. It is white, hard, and encysted, and differs from the *criske*, another species, only in being moveable. Writers mention a division of Chalazion into scirrhus, cancerous, cystic, and earthy.

CHALIBANE. (*Καλαβη*.) Galbanum.

CHALCANTHUM. (From *χαλκας*, brass, and *ανθος*, a flower.) Vitriol; or rather, vitriol calcined red. The flowers of brass.

CHALCETON. A species of pimpinella.

CHALCIDEUM OS. The os cuneiforme of the tarsus.

CHALK. *Creta.* A carbonate of lime. Pure chalk is a neutral salt, formed by the union of the cretaceous acid with lime. It is much used as an absorbent and antacid, to stop diarrhoeas, accompanied with acidity. See *Carbonas calcis*.

CHALK STONES. A name given to the concretions of calcareous matter in the hands and feet of people violently afflicted with the gout.

CHALICRATUM. (From *χαλκ*, an old word that signifies pure wine, and *κρατουμε*, to mix.) Wine mixed with water.

CHALINOS. *Chalinus.* That part of the cheeks, which, on each side, is contiguous to the angles of the mouth.

CHALYBEATE. (*Chalybeata*, ac, *medicamenta*; from *chalybs*, steel.) Of or belonging to iron. A term given to any medicine into which iron enters; as chalybeate mixture, pills, waters, &c.

CHALYBEATE WATERS. Any mineral water which abounds with steel, or

iron; such as the waters of Tunbridge, Spa, Pyrmont, Cheltenham, Scarborough, and Hartel; and many others.

CHALYBIS RUBICO PRÆPARATA. See *Carbonas ferri*.

CHALYBS. (From *Chalybes*, a people in Pontus, who dug iron out of the earth.) *Acies*. Steel. The best, hardest, finest, and the closest-grained forged iron. As a medicine, it differs not from iron.

CHALYBS TARTARISATUS. The *ferrum tartaribatum*.

CHAMBALANUS. (From *χαμαί*, on the ground, and *βάλανος*, a nut.) Wood peas. Earth nuts.

CHAMBUXUS. (From *χαμαί*, on the ground, and *βύξ*, the box-tree.) The dwarf box-tree.

CHAMCEDRUS. (From *χαμαί*, on the ground, and *κεδρος*, the cedar tree.) *Chamaedrys*. A species of dwarf abrotanum.

CHAMECISUS. (From *χαμαί*, on the ground, and *κισσός*, ivy.) Ground ivy.

CHAMEDRYS. (From *χαμαί*, the ground, and *δρυς*, the oak; so called from its leaves resembling those of the oak.) *Chamedrys minor repinis, vulgaris. Quercula calamandrina. Trixago: Chamedrops* of Paulus Ægineta and Oribasius. This plant, creeping germander, small germander, and English treacle: *Teucrium chamaedrys* of Linnaeus:—*foliis cuneiformi-ovatis, incisis, crenatis, petiolatis; floribus ternis; caulis procumbentibus, subpilosis*, has a moderately bitter and somewhat aromatic taste. It is in high repute amongst the ancients in intermittent fevers, rheumatism, and gout; and where an aromatic bitter is wanting, germander may be administered with success. The best time for gathering this herb is when the seeds are formed, and the tops are then preferable to the leaves. When dry, the dose is from ʒss to ʒj. Either water or spirit will extract their virtue; but the watry infusion is more bitter. This plant is an ingredient in the once celebrated powder called from the Duke of Portland.

CHAMEDRYS INCANA MARITIMA. The *marum syriacum*.

CHAMEDRYS FRUTESCENS. A name for *teucrium*.

CHAMEDRYS PALUSTRIS. A name given to *scordium*.

CHAMEDRYS SPURIA. A name given to *veronica*.

CHAMELEA. (From *χαμαί*, on the ground, and *ελαια*, the olive-tree.) *Chamelea*. The herb widow-wail, or *Daphne alpina* of Linnaeus. A sort of dwarf olive-tree, said to be purgative in the dose of ʒij. The mezercon is also so called, because it has leaves like the olive-tree.

CHAMELEAGNUS. (From *χαμαί*, on the ground, and *ελαιαγνος*, the wild olive.) The *myrtus brabantica*.

CHAMELENA. (From *χαμαί*, on the ground, and *ελαια*, ivy.) The ground-ivy.

CHAMELEON. (From *χαμαί*, on the ground, and *λεων*, a lion, i. e. dwarf lion.) The chameleon, an animal supposed to be able to change its colour at pleasure. Also the name of many thistle, so named from the variety and uncertainty of their colours.

CHAMELEON ALBUM. *Carlina. Cardopatiium.* Carline thistle. *Carlina acaulis* of Linnaeus:—*caule unifloro, flore breviori*. The root of this plant is bitter, and said to possess diaphoretic and anthelmintic virtues. It is also extolled by foreign physicians in the cure of acute, malignant, and chronic disorders.

CHAMELEON VERUM. The distaff thistle.

CHAMELEUCE. (From *χαμαί*, on the ground, and *λευκός*, the herb colt's-foot.) Tuisslago, or colt's-foot.

CHAMELINUM. (From *χαμαί*, on the ground, and *λεων*, flax.) *Linum catharticum*, or purging of flax.)

CHAMEMELUM. (From *χαμαί*, on the ground, and *μαλόν*, an apple; because it grows upon the ground, and has the smell of an apple.) *Chamemelum nobile. Chamomilla romana. Euanthemum* of Galen. *Anthemis* of the last London pharmacopœia. Common chamomile. *Anthemis nobilis* of Linnaeus:—*foliis pinnato-compositis linearibus acutis subvillosis*. Both the leaves and flowers of this indigenous plant have a strong, though not ungrateful smell, and a very bitter, nauseous taste: but the latter are the bitterer, and considerably more aromatic. They possess tonic and stomachic qualities, and are much employed to restore tone to the stomach and intestines, and as a pleasant and cheap bitter. They have been long successfully used for the cure of intermittents, as well as of fevers of the irregular nervous kind, accompanied with visceral obstructions. The flowers have been found useful in hysterical affections, flatulent or spasmodic colics, and dysentery; but, from their laxative quality, Dr. Cullen tells us they proved hurtful in diarrhœas. A simple infusion is frequently taken to excite vomiting, or for promoting the operation of emetics. Externally, they are used in the *decoctum pro fomento*, and are an ingredient in the *decoctum pro enema*.

CHAMEMELUM CANARIENSE. The *Chrysanthemum frutescens* of Linnaeus.

CHAMEMELUM CHRYSANTHEMUM. The *bupthalmum germanicum*.

CHAMEMELUM FETIDUM. The *Anthemis cotula* of Linnaeus.

CHAMEMELUM FLORE PLENO. *Chamemelum nobile flore multiplici*. Double chamomile. A variety of the *anthemis nobilis*. See *Chamemelum*.

CHAMÆMELUM NOBILE. See *Chamæmelum*.

CHAMÆMELUM VULGARE. *Chamamilla nostras.* *Leucanthemum* of Dioscorides. Common wild corn, or dog's chamomile. The plant directed under this name in the pharmacopœia, is the *Marricaria chamamilla* of Linnæus:—*receptaculis conicis; radiis prætenuibus; squamis calycinis, marginibus aequalibus.* Its virtues are similar to those of the chamæmelum, but in a much inferior degree.

CHAMÆMORUS. (From *χαμας*, on the ground, and *μορος*, the mulberry tree.) *Chamerubus foliis rubis Anglicæ.* *Rubus palustris humilis.* *Vaccinium Lencutrense.* *Rubus alpinus humilis.* *Anglicus* Cloude, berries, and knot berries. The ripe fruit of this plant, *Rubus chamæmorus* of Linnæus—*foliis simplicibus lobatis caule internodioso unifloro*, is prepared into a jam: and is recommended to allay thirst, &c. in fevers, phthisical diseases, hæmoptysis, &c. As an antiscorbutic, it is said to excel the scurvy-grass and vegetables of that tribe in common use.

CHAMÆPICE. (From *χαμας*, on the ground, and *πικρα*, the pine-tree.) Camphorata, or stinking ground-pine, formerly said to be antirheumatic.

CHAMÆPITYS. (From *χαμας*, the ground, and *πυτος*, the pine-tree.) *Arthritica.* *Arthretica Ajuga.* *Abiga.* *Iva arthritica.* *Eulocyron.* *Ionia.* *Sideritis.* Common ground-pine. This low hairy plant, *Teucrium chamæpitys* of Linnæus:—*foliis trifidis, linearibus, intergerminis; floribus sessilibus, lateralibus, solitariis; caule diffusis*, has a moderately bitter taste, and a resinous, not disagreeable smell, somewhat like that of the pine. The tops, or leaves, are recommended as aperients and corroborants of the nervous system, and said to be particularly serviceable in female obstructions and paralytic disorders.

CHAMÆPITYS MOSCHATA. *Iva moschata mœspeliensis.* *Chamæpitys anthyllis.* The *Teucrium iva* of Linnæus. French ground pine. It is weaker, but of similar virtues to chamæpitys.

CHAMÆPLION. A name in Oribasius for crysimum, or hedge mustard.

CHAMÆRAPANUM. So Paulus Ægineta calls the upper part of the root of the apium.

CHAMÆRAPANUS. From *χαμας*, on the ground, and *ραπανος*, the radish.) The upper part of the root of apium, according to P. Ægineta. The smallage, or parsley. Also dwarf radish.

CHAMÆRIPES. The palma minor.

CHAMÆRODENDRON. (From *χαμας*, on the ground, and *ροδον*, the rose laurel.) The *Azalea pontica* of Linnæus.

CHAMÆRUBUS. (From *χαμας*, on the ground, and *rubus*, the bramble.) The chamæmorus.

CHAMÆSPARTIUM. (From *χαμας*, on the ground, and *σπαστον*, Spanish broom.) The genista tinctoria.

CHAMBERS. The space between the capsule of the crystalline lens and the cornea of the eye, is divided by the iris into two spaces called chambers; the space before the iris is termed the anterior chamber; and that behind it, the posterior. They are filled with an aqueous fluid.

CHAMONILLA NOSTRAS. See *Chamæmelum vulgare*.

CHAMONILLA ROMANA. See *Chamæmelum*.

CHANCER. (From *καρκινος*, cancer.) A sore which arises from the direct application of the venereal poison to any part of the body. Of course it mostly occurs on the genitals. Such venereal sores as break out from a general contamination of the system, in consequence of absorption, never have the term chancre applied to them.

CHAMANTIA SIGNA. So Paracelsus calls those prognostics that are taken from observations of the air; and the skill of doing this, the same author calls *Chuamancia*.

CHAOSDA. Paracelsus uses this word as an epithet for the plague.

CHARABE (Arab.) A name given to amber. See *Succinum*.

CHARABRA. (From *χαγαρρο*, to excavate.) The bowels, or sink of the body.

CHARAMAIS. Purging hazel-nut.

CHARANTIA. The momordica elaterium. *Charcoal.* See *Carbon*.

CHARDONS. The *Cinara spinosa*.

CHARISTOLOCHIA. (From *χαρε*, joy, and *λοχη*, the flux of women after child-birth; so named from its supposed usefulness to women in childbirth.) The plant mugwort, or artemisia.

CHARME. (From *χαμας*, to rejoice.) *Charmis.* A cordial antidote mentioned by Galen.

CHARPIE. (French.) Scraped linen, or lint.

CHARTA. (Chald.) Paper. The amnios, or interior fetal membrane, was called the *charta virginica*, from its likeness to a piece of fine paper.

CHARTREUX, POUDBE DE. (So called because it was invented by some friars of the Carthusian order.) A name of the kermes mineral.

CHASME. (From *χαμας*, to gape.) *Chasmus.* Ossification. Gaping.

Chaste tree. The *Agnus castus*.

CHATE. The cucumis *Ægyptia*.

Check-bone. See *Jugale os*.

CHEESE. The coagulum of milk. When prepared from rich milk, and well made, it is very nutritious in small quantities: but mostly indigestible when hard and ill prepared, especially to weak stomachs.

CHELOCACE. (From *χαλας*, a lip, and

μακρυ, an evil.) The lip-evil. A swelling of the lips, or canker in the mouth.

CHEIMELTON. (From χιμα, winter.) Chilblains.

CHEIRANTHUS. (From χηρ, a hand, and ανθος, a flower; so named from the likeness of its blossoms to the fingers of the hand.) The name of a genus of plants in the Linnæan system. Class, *Tetradynamia*. Order, *Silicosa*. The wall flower.

CHEIRANTHUS CHEIRI. The systematic name of the wall-flower. See *Cheiri*.

CHEIRAPsia. (From χηρ, the hand, and σκωμαι, to scratch.) The act of scratching; particularly the scratching one hand with another, as in the itch.

CHEIRI. (*Cheiri*, Arab.) *Leucolum lateum*. *Viola lutea*. Common yellow wall-flower. The flowers of this plant, *Cheiranthus cheiri* of Linnæus:—*foliis lanceolatis, acutis, glabris; ramis angulatis; caule fruticoso*, are recommended as possessing nervine and deobatruent virtues. They have a moderately strong, pleasant smell, and a nauseous, bitter, somewhat pungent taste.

CHEIRIATER. (From χηρ, the hand, and ιατρος, a physician. A surgeon whose office it is to remove maladies by operations of the hand.

CHEIRIUMA. (From χηρζομαι, to labour with the hand.) Handling. Also a manual operation.

CHEIRIXIS. (From χηρζομαι, to labour with the hand. The art of surgery.

CHEIRONOMIA. (From χηρνομαι, to exercise with the hands.) An exercise mentioned by Hippocrates, which consisted of pectulations with the hands, like our dumb-bells.

CHELA. (From χηλη, *forceps*; from χη, to take.) A forked probe, for drawing a polypus out of the nose. Fissures in the feet, or other places.

CHELE CANCERUM. See *Cancer*.

CHELIDON. The bend of the arm.

CHELIDONIUM MAJUS. (From χελιδων the swallow. It is so named from an opinion that it was pointed out as useful for the eyes by swallows, who are said to open the eyes of their young by it; or because it blossoms about the time when swallows appear.) *Papaver corniculatum, luteum*. Tetterwort, and great colandine. The herb and root of this plant, *Chelidonium majus* of Linnæus:—*pedunculis umbellatis*, have a faint, unpleasant smell, and a bitter, acrid, durable taste, which is stronger in the roots than the leaves. They are aperient and diuretic, and recommended in icterus; when not accompanied with inflammatory symptoms. The chelidonium should be administered with caution, as it is liable to irritate the stomach and bowels. Of the dried root, from ℥ss to ʒj is a dose; of the fresh root, infused in water, or wine, the dose may be about ʒss. The decoction of the fresh root is used in

droopy, cachexy, and cutaneous complaints. The fresh juice is used to destroy warts and films in the eyes; but, for the latter purpose, it is diluted with milk.

CHELIDONIUM MINUS. *Serophylaria minor*. *Chelidonia retundifolia minor*. *Curcuma hemorrhoidalis herba*. *Ranunculus vernus*. Lessercelandine, and pilewort. The leaves and root of this plant, *Ranunculus ficaria; foliis cordatis angulatis petiolatis, caule unifloro*, of Linnæus, are used medicinally. The leaves are deemed antiscorbutic, and the root reckoned a specific, if beat into cataplasma, and applied to the piles.

CHELONÆ. (χελωνæ.) The tortoise. An instrument for extending a limb, and so called because, in its slow motions, it represents a tortoise. This instrument is mentioned in Oribasius.

CHELONTION. (From χελωνæ, the tortoise; so called from its resemblance to the shell of a tortoise.) A hump, or gibbosity in the back.

CHELTENHAM WATER. One of the most celebrated purging waters in England and the reputation of it is daily increasing, as it possesses both a saline and chalybeate principle. When first drawn it is clear and colourless, but somewhat brisk; has a saline, bitterish, chalybeate taste. It does not keep, nor bear transporting to any distance; the chalybeate part being lost by precipitation of the iron, and in the open air it even turns fetid. The salts, however, remain. Its heat, in summer, was from 50° to 55°, or 59°, when the medium heat of the atmosphere was nearly 15° higher. On evaporation, it is found to contain a calcareous earth, mixed with ochre and a purging salt. A general survey of the component parts of this water, according to a variety of analyses, shows that it is decidedly saline, and contains much more salt than most mineral waters. By far the greater part of the salts are of a purgative kind, and therefore an action on the bowels is a constant effect, notwithstanding the considerable quantity of selenite and earthy carbonates which may be supposed to have a contrary tendency. Cheltenham water is, besides, one of the strongest chalybeates we are acquainted with. The iron is suspended entirely by the carbonic acid, of which gas the water contains about an eighth of its bulk; but, from the abundance of earthy carbonates, and oxyd of iron, not much of it is uncombined. It has, besides, a slight impregnation of sulphur, but so little, as to be scarcely appreciable, except by very delicate tests. The sensible effects produced by this water, are generally, on first taking it, a degree of drowsiness, and sometimes head-ach, but which soon go off spontaneously, even previous to the operation on the bowels. A moderate dose acts powerfully, and speedily, as a cathartic, without occa-

sioning griping, or leaving that faintness and languor which often follow the action of the rougher cathartics. It is principally on this account, but partly too from the salutary operation of the chalybeate, and perhaps the carbonic acid, that the Cheltenham water may be, in most cases, persevered in, for a considerable length of time, uninterruptedly, without producing any inconvenience to the body; and during its use, the appetite will be improved, the digestive organs strengthened, and the whole constitution invigorated. A dose of this water, too small to operate directly on the bowels, will generally determine pretty powerfully to the kidneys. As a purge, this water is drank from one to three pints; in general, from half a pint to a quart is sufficient. Half a pint will contain half a drachm of neutral purging salts, four grains of earthy carbonate, and selenite, about one-third of a grain of oxyd of iron; together with an ounce in bulk of carbonic acid, and half an ounce of common air, with a little sulphurated hydrogen. Cheltenham water is used, with considerable benefit, in a number of diseases, especially of the chronic kind, and particularly those called bilious: hence it has been found of essential service in the cure of glandular obstruction, and especially those that affect the liver, and the other organs connected with the functions of the alimentary canal. Persons who have injured their biliary organs, by a long residence in hot climates, and who are suffering under the symptoms either of excess of bile or deficiency of bile, and an irregularity in its secretion, receive remarkable benefit from a course of this water, judiciously exhibited. Its use may be here continued, even during a considerable degree of debility; and from the great determination to the bowels it may be employed with advantage to check the incipient symptoms of dropsy, and general anasarca, which so often proceed from an obstruction of the liver. In scrophulous affections, the sea has the decided preference; in painful affections of the skin, called scorbutic eruptions, which make their appearance at stated intervals, producing a copious discharge of lymph, and an abundant desquamation, in common with other saline purgative springs, this is found to bring relief; but it requires to be persevered in for a considerable time, keeping up a constant determination to the bowels, and making use of warm bathing externally. The season for drinking the Cheltenham water is during the whole of the summer months.

CHILTA. (*χάλυς*, a shell.) The breast is so called, as resembling, in shape and office, the shell of some fishes.

CHELYSTON. (From *χάλυς*, the breast.) A dry, short cough, in which the muscles of the breast are very sore.

CHEMA. A measure mentioned by the

Greek physicians, supposed to contain two small spoonfuls.

CHEMIA. (Arab.) *Chimia*. Chymistry.

CHEMICAL APPARATUS. A general expression, denoting the instruments, vessels, machinery, furniture, and utensils of a laboratory.

CHEMISTRY. (*χημια*, and sometimes *χημια*: *chamia*, from *chama*, to burn, Arab. this science bring the examination of all substances by fire.) The learned are not yet agreed as to the most proper definition of chymistry. Boerhaave seems to have ranked it among the arts. According to Macquer, it is a science, whose object is to discover the nature and properties of all bodies by their analyses and combinations. Dr. Black says, it is a science which teaches, by experiments, the effects of heat and mixture on bodies; and Fourcroy defines it a science which teaches the mutual actions of all natural bodies on each other. "Chymistry," says Jacquin, "is that branch of natural philosophy which unfolds the nature of all material bodies, determines the number and properties of their component parts, and teaches us how those parts are united, and by what means they may be separated and recombined." Mr. Heron defines it, "That science which investigates and explains the laws of that attraction which takes place between the minute component particles of natural bodies." The objects to which the attention of chymists is directed, comprehend the whole of the substances that compose the globe.

CHEMOSIS. (From *χῆμα*, to gape; because it gives the appearance of a gap, or aperture.) Inflammation of the conjunctive membrane of the eye, in which the cellular structure, or white of the eye, is distended with a florid fluid, and elevated above the margin of the transparent cornea. In Cullen's Nosology, it is a variety of the ophthalmia membranarum, or an inflammation of the membranes of the eye.

CHENOPODIO-MORUS. (From *chenopodium* and *μωρος*, the mulberry; so called because it is a sort of chenopodium, with leaves like a mulberry.) The herb mulberry-blight, or strawberry-spench.

CHENOPODIUM. (From *χην*, a goose, and *πους*, a foot; so called from its supposed resemblance to a goose's foot.) The herb chenopody, goose's foot. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*.

CHENOPODIUM AMEROSIOIDES. The systematic name of the Mexican tea-plant. See *Botrys Mexicana*.

CHENOPODIUM ANTHELMINTICUM. The seeds of this plant, *Chenopodium anthelminticum* of Linnæus:—*foliis ovato-oblongis dentatis, racemis aphyllis*, though in great esteem in America for the cure of worms,

are never exhibited in this country. They are powdered and made into an electuary, with any proper syrup, or conserve.

CHENOPODIUM BOTRYS. The systematic name of the Jerusalem oak. See *Botrys vulgaris*.

CHENOPODIUM BONUS HENRICUS. The systematic name of the English mercury. See *Bonus henricus*.

CHENOPODIUM VULVARIA. The systematic name for the stinking orach. See *Atriplex fatida*.

CHENAS (From *χένω* to pour out.) The struma, or scrophula.

CHEREPOLIUM. See *Cherophyllum*.

CHERMES. (Arab.) A small berry, full of insects like worms: the juice of which was formerly made into a confection, called confectio alkerimes, which has been long disused. Also the worm itself.

CHERMES MINERALIS. See *Sulphuretum ammonii præcipitatum*.

CHERNIBIUM. *Chernibion*. In Hippocrates it signifies an urinal.

CHERONIA. (From *Χερον*, the Centaur.) See *Centaurium*.

Cherry. See *Cerasa nigra* and *Cerasa rubra*.

Cherry, bay. The *Lauro-cerasus*.

Cherry, laurel. The *Lauro-cerasus*.

Cherry, winter. The *Alkekegi*.

CHERVILLUM. See *Cerefolium*.

Chestnut, horse. See *Hippocastanum*.

CHESIS. (From *χένω*, to pour out.) Li-quation. Infusion.

CHEVA-TRE. A double-headed roller, applied by its middle below the chin; then running on each side, it is crossed on the top of the head; then passing to the nape of the neck is there crossed: it then passes under the chin, where crossing, it is carried to the top of the head, &c. until it is all taken up.

CHEZANANCE. (From *χίζω*, to go to stool, and *ανάγκη*, necessity.) It signifies any thing that creates a necessity to go to stool; but, in *P. Æ. ineta*, it is the name of an ointment, with which the anus is to be rubbed, for promoting stools.

CHIA. (From *Χίος*, an island where they were formerly propagated.) A sweet fig of the island of Chio, or Scio. Also an earth from that island, formerly used in fevers.

CHIACUS. (From *Χίος*, the island of Scio.) An epithet of a collyrium, whose chief ingredient was wine of Chios.

CHIADUS. In Paracelsus it signifies the same as furunculus.

Chian pepper. See *Capsicum*.

Chian turpentine. See *Chio turpentine*.

CHIASEMS (From *χιάζω*, to form like the letter X, *chi*.) The name of a bandage, whose shape is like the Greek letter X, *chi*.

CHIASTOS. The name of a cruciate bandage in Orribasius; so called from its resembling the letter X, *chi*.

CHIASTRE. The name of a bandage for the temporal artery. It is a double-headed roller, the middle of which is applied to the side of the head, opposite to that in which the artery is opened and, when brought round to the part affected, it is crossed upon the compress that is laid upon the wound, and then, the continuation is over the coronal suture, and under the chin; then crossing on the compress, the course is, at the first, round the head, &c. till the whole roller is taken up.

CHIBOU. A spurious species of gum-elemi, spoken of by the faculty of Paris, but not known in England.

CHIBUR. Sulphur.

CHICHINA. Contracted from China *chinz*. See *Cinchona*.

CHICOS. *Chirces*. The affectio bovina, or distemper of black cattle.

Chicken pox. See *Varicella*.

Chickweed. See *Aloine media*.

CHILBLAIN. *Pernio*. An inflammation of the extreme parts of the body, from the application of cold; attended with a violent itching, and soon forming a gangrenous ulcer.

CHILI, BALSAM DE. Salmon speaks, but without any proof, of its being brought from Chili. The Barbadoes tar, in which are mixed a few drops of the oil of aniseed, is usually sold for it.

CHILIODYNAMON. (From *χίλιον*, a thousand, and *δυναμις*, virtue.) An epithet of the herb *Potimonium*. In Dioscorides, this name is given on account of its many virtues.

CHILON. (*Χίλιον*.) An inflamed and swelled lip.

CHILPELAGUA. A variety of capsicum.

Chilterspin. A species of capsicum.

CHIMETHLON. A chilblain.

CHIMIA. See *Chemia*.

CHINIATER. (From *chemia*, chymistry, and *ιατρος*, a physician.) A physician who makes the science of chymistry subservient to the purposes of medicine.

CHINOLEA LAXA. Paracelsus means, by this word, the sublimed powder which is separated from the flowers of saline ores.

CHINA. (So named from the country of China, from whence it was brought.) *China orientalis.* *Sankia.* *Guoquara.* *Smilax aspera Chinensis.* China root. It is obtained from the *Smilax china* of Linnaeus:—caule aculeato, teretiusculo; foliis inermis, ovato-cordatis, quinque nerviis. It was formerly in esteem, as sarsaparilla now is, in the cure of the venereal disease, and cutaneous disorders.

CHINA CHINE. A name given to the Pervian bark, a native of some parts in China.

CHINA OCCIDENTALIS. *China spuria nodosa.* *Smilax pseudo-China.* *Smilax Indica spinosa.* American or West-Indian China. The root is chiefly brought from

Jamaica, in large round pieces, full of knots. In scrophulous disorders, it has been preferred to the oriental kind. In other cases it is of similar but inferior virtue.

CHINA SUPPOSITA. *Senecio madraspatanus.* *Senecio pseudo-China* of Linnaeus. Bastard China. It grows in Malabar. The root greatly resembles the China root in appearance and qualities.

CHINCHINA. See *Cinchona*.

CHINCHINA CARIBEA. See *Cinchona Caribaea*.

CHINCHINA DE SANTA FE. There are several species of bark sent from Santa Fè: but neither their particular nature, nor the trees which afford them, are yet accurately determined.

CHINCHINA JAMAICENSIS. See *Cinchona Caribaea*.

CHINCHINA RUBRA. See *Cinchona oblongifolia*.

CHINCHINA DE ST. LUCIA. St. Lucia bark is collected from the *Anchona floribunda* of Swartz:—*floribus paniculatis glabris, capsulis turbinatis levibus, foliis ellipticis acuminatis glabris*; it has an adstringent, bitter taste, somewhat like ginsain. It is recommended in intermittents, putrid dysentery, and dyspepsia: it should always be joined with some aromatic.

Chincough. See *Pertussis*.

CHINENSE. The aurantium sinense, or Chinese orange.

CHINESE SMILAX. See *China*.

CHIO TURPENTINE. *Terebinthina de Chio.* Cyprus turpentine. Chian turpentine. This substance is classed among the resins. It is procured by wounding the bark of the trunk of the *Pistachia terebinthus* of Linnaeus. The best Chio turpentine is about the consistence of honey, very tenacious, clear, and almost transparent; of a white colour, inclining to yellow, and a fragrant smell, moderately warm to the taste, but free from acrimony and bitterness. Its medicinal qualities are similar to those of the turpentine. See *Turpentine*.

CHIOLI. In Paracelsus it is synonymous with furunculus.

CHIQUE. A name for the worms which get into the toes of the negroes, and which are destroyed by the oil which flows out of the cashew nut-shell.

CHIRAGRA. (From *χυσ*, the hand, and *αγρα*, a seizure.) The gout in the joints of the hand. See *Arthritis*.

CHIRONES. (From *χυσ*, the hand.) Small pustules on the hand and feet, enclosed in which is a troublesome worm.

CHIRONIA. (From *Chiron*, the Centaur, its inventor.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*. *Centaurium*. See *Centaurium*.

2. (From *χυσ*, the hand.) An affection

of the hand, where it is troubled with chirones.

CHIRONIUM. (From *Χιρων*, the Centaur, who is said to have been the first who healed them.) A malignant ulcer, callous on its edges, and difficult to cure.

CHIROTHECA. (From *χυσ*, the hand, and *τίθημι*, to put.) A glove of the scarf-skin, with the nails, which is brought off from the dead subject, after the cuticle is loosened by putrefaction, from the parts under it.

CHIRURGIA. (From *χυσ*, the hand, and *εργον*, a work; because surgical operations are performed by the hand.) Chirurgery, or surgery.

CHITON. (*χιτων*.) A coat or membrane.

CHIU. (From *Χιος*, the island where it was produced.) An epithet of wine made at Scio.

CHILASMA. (From *χλιαστω*, to make warm.) A warm fomentation, called also *thermasma*.

CHLOASMA. (From *χλασσω*, to become green.) Chlorosis.

CHLOROSIS. (From *χλωρος*, green pale; from the yellow-greenish look those have who are affected with it.) *Febris alba. Febris amatoria. Icterus albus.*

The green sickness. A genus of disease in the class *cachexia*, and order *impetiginæ* of Cullen. It is a disease which affects young females who labour under a suppression of the menses. Heaviness, listlessness to motion, fatigue on the least exercise, palpitations of the heart, pains in the back, loins, and hips, flatulency and acidiities in the stomach and bowels, a preternatural appetite for chalk, lime, and various other absorbents, together with many dyspeptic symptoms, usually attend on this disease. As it advances in its progress, the face becomes pale, or assumes a yellowish hue; the whole body is flaccid, and likewise pale; the feet are affected with œdematous swellings; the breathing is much hurried by any vigorous exertion of the body; the pulse is quick, but small; and the person is apt to be affected with many of the symptoms of hysteria. To procure a flow of the menses, proves in some cases a very difficult matter; and where the disease has been of long standing, various morbid affections of the viscera are often brought on, which at length prove fatal. Dissections of those who have died of chlorosis, have usually shewn the ovaria to be in a scirrhus, or dropsical state. In some cases, the liver, spleen, and mesenteric glands, have likewise been found in a diseased state.

CHNUS. (From *χναω*, to grind, or rasp.) Chaff. Bran. Also fine wool, or lint, which is, as it were, rasped from lint.

CHROKE DAMP. The name given by miners to a noxious air, occasionally found in the bottom of mines and pits. It is

heavier than common air, therefore lies chiefly at the bottom of the pits; it extinguishes flame, and is noxious to animals. It is probably carbonic acid. See *Carbonic acid*.

CHOANA. (From *χωα*, to pour out.) It is properly a funnel, but is used to signify the infundibulum of the kidney and brain.

CHOANUS. (*χοανα*, a funnel.) A furnace made like a funnel, for melting metals.

CHOCOLATE. (Dr. Alston says this word is compounded of two Indian words, *choco*, sound, and *atte*, water; because of the noise made in its preparation.) An article of diet prepared from the cocoa-nut; highly nourishing, particularly when boiled with milk and eggs. It is frequently recommended as a restorative in cases of emaciation and consumption.

CHONICIS. The trepan, so called by Galen and P. Aegineti, from *χωνικος*, the nave of a wheel.

CHORADES. (From *χοιρος*, a swine.) *Chorades*. The same as *scrofula*.

CHORADOLETHRON. (From *χοιρος*, a swine, and *αλεσθω*, destruction; so named from its being dangerous if eaten by hogs.) Hogbane. A name in *Ætius* for the *Xanthium*, or louse-bur.

CHOIRAS. (From *χοιρος*, a swine; so called because hogs are diseased with it.) The scrophula.

CHOLADES. (From *χολη*, the bile.) So the smaller intestines are called, because they contain bile.

CHOLAGO. The same as *cholus*.

CHOLAGOGA. (From *χολη*, bile, and *αγω*, to evacuate.) *Chologon*. By *chologogues*, the ancients meant only such purging medicines as expelled the internal feces, which resembled the cystic bile in their yellow colour, and other properties.

CHOLAS. (From *χολη*, the bile.) All the cavity of the ilium is so called, because it contains the liver which is the strainer of the gall.

CHOLE. (*χολη*) The bile.

CHOLEDOCHUS DUCTUS. (*Choledochus*; from *χολη*, bile, and *αγωγος*, to receive; receiving or retaining the gall.) *Ductus communis choledochus*. The common biliary duct, which conveys both cystic and hepatic bile into the intestinum duodenum.

CHOLEGON. The same as *chologoga*.

CHOLERA. (From *χολη*, bile, and *ρωα*, to flow.) *Diarrhœa cholericæ*. *Fellina passio Cholera*. A genus of disease arranged by Cullen in the class *neuroses*, and order *spasmi*. It is a purging and vomiting of bile, with anxiety, painful gripings, spasms of the abdominal muscles, and those of the thighs. There are two species of this genus: 1. *Cholera spontanea*, which happens, in hot seasons, without any manifest cause. 2. *Cholera accidentalis*, which

occurs after the use of food that digests slowly, and irritates. In warm climates it is met with in all seasons of the year, and its occurrence is very frequent: but in England, and other cold climates, it is apt to be most prevalent in the middle of summer, particularly in the month of August; and the violence of the disease has usually been observed to be greater in proportion to the intenseness of the heat. It usually comes on with soreness, pain, distension, and flatulency in the stomach and intestines, succeeded quickly by a severe and frequent vomiting, and purging of bilious matter, heat, thirst, a hurried respiration, and frequent but weak and fluttering pulse. When the disease is not violent these symptoms, after continuing for a day or two, cease gradually, leaving the patient in a debilitated and exhausted state; but where the disease proceeds with much violence there arises great depression of strength, with cold clammy sweats, considerable anxiety, a hurried and short respiration, and hiccup, with a sinking, and irregularity of the pulse, which quickly terminate in death; an event that not unfrequently happens within the space of 24 hours.

CHOLERICA. (From *χολη*, the cholera.) Medicines which relieve the cholera. Also a bilious flux of the bowels, without pain or fever.

CHOLICELE. (From *χολη*, bile, and *κληη*, a tumour.) A swelling formed by the bile morbidly accumulated in the gall-bladder.

CHOLOMA. (From *χολος*, lame, or maimed.) Galen says that, in Hippocrates, it signifies any distortion of a limb. In a particular sense, it is taken for a halting or lameness in the leg.

CHONDROGLOSSUS. (From *χονδρος*, a cartilage, and *γλωσσα*, the tongue.) A muscle so named from its insertion, which is in the basis or cartilaginous part of the tongue. See *Hyoglossus*.

CHONDROLOGY. (*Chondrologia*; from *χονδρος*, a cartilage, and *λογος*, a discourse.) A discourse or treatise on cartilages.

CHONDRO-PHARYNGEUS. (From *χονδρος*, a cartilage, and *φαρυγξ*, the upper part of the fauces.) A muscle so named because it rises in the cartilaginous part of the tongue, and is inserted in the pharynx.

CHONDROS. (*Χονδρος*.) A food of the ancients, the same as *alica*. Also any grumous concretion, and a cartilage.

CHONDROSYNDESMUS. (From *χονδρος*, a cartilage, and *συνδεω*, to tie together.) A cartilaginous ligament.

CHONDROS. See *Chondros*.

CHONE. (*Χωνη*.) The infundibulum.

CHOXA. (*Χοξα*.) A region. Galen, in his book *De Usu Partium*, expresses by it particularly the cavities of the eyes; but,

in others of his writings, he intimates by it any void space.

CHORDA. A cord. A tendon. A painful tension of the penis in venereal disease. See *Chordee*. Sometimes the intestines are called chordæ.

CHORDA MAGNA. A name of the *tendo Achillis*.

CHORDA TYMPANI. A branch of the seventh pair of nerves that passes through the tympanum

CHORDÆ TENDINÆ. The tendinous and cord-like substances which connect the *cornu columæ* of the ventricles of the heart to the auricular valves.

CHORDÆ WILLISII. The small fibres which cross the sinuses of the dura mater. They are so termed, because Willis first described them

CORDAPUS. (From *χορδή*, a cord, and *πύσις*, to knit) A sort of painful cholæ, where the intestines appear to be twisted into knots, like pieces of string.

CHORDEE. (*Chordé* French.) A spasmodic contraction of the penis, that sometimes attends gonorrhœa, and is often followed by a hæmorrhage.

CHOREA SANCTI VITI (*Chorea*, *χορεία*) from *χορός*, a chorus which of old accompanied dancing. It is called St. Vitus's dance, because some devotees of St. Vitus exercised themselves so long in dancing, that their intellects were disordered, and could only be restored by dancing again at the anniversary of St. Vitus.) St. Vitus's dance. Convulsive motions of the limbs, as if the person were dancing. It is a genus of disease arranged by Cullen in the class *neuroses*, and order *spasmi*. These convulsive motions, most generally, are confined to one side, and affecting principally the arm and leg. When any motion is attempted to be made, various fibres of other muscles act which ought not; and thus a contrary effect is produced from what the patient intended. It is chiefly incident to young persons of both sexes, and makes its attack from between the age of ten and fifteen, occurring but seldom after that of puberty.

By some practitioners it has been considered rather as a paralytic affection than as a convulsive disorder and has been thought to arise from a relaxation of the muscles, which, being unable to perform their functions in moving the limbs, shake them irregularly by jerks. *Chorea sancti Viti* is occasioned by various irritations, as teething, worms, offensive smells, poisons, &c. It arises likewise in consequence of violent affections of the mind, as horror, fear, and anger. In many cases it is produced by general weakness; and in a few, it takes place from sympathy, at seeing the disease in others.

The fits are sometimes preceded by a coldness of the feet and limbs, or a kind of tingling sensation, that ascends like cold

air up the spine, and there is a flatulent pain in the left hypochondrium, with obstinate costiveness. At other times, the accession begins with yawning, stretching, anxiety about the heart, palpitations, nausea, difficulty of swallowing, noise in the ears, giddiness and pains in the head and teeth; and then come on the convulsive motions.

These discover themselves at first by a kind of lameness, or instability of one of the legs, which the person draws after him in an odd and ridiculous manner; nor can he hold the arm of the same side still for a moment; for if he lays it on his breast, or any other part of his body, it is forced quickly from thence by an involuntary motion. If he is desirous of drinking, he uses many singular gesticulations before he can carry the cup to his head, and it is forced in various directions, till at length he gets it to his mouth; when he pours the liquor down his throat with great haste, as if he meant to afford amusement to the bystanders. Sometimes various attempts at running and leaping take place, and at others, the head and trunk of the body are affected with convulsive motions. In many instances, the mind is affected with some degree of fatuity, and often shews the same causeless emotions, such as weeping and laughing, which occur in hysteria. When this disease arises in children, it usually ceases before the age of puberty; and in adults, is often carried off by a change from the former mode of living. Unless it passes into some other disease, such as epilepsy, it is never attended with danger.

CHORION. (From *χαίρω*, to escape; because it always escapes from the uterus with the fœtus.) Shaggy chorion. The external membrane of the fœtus in utero.

CHOROID MEMBRANE. (*Membrana chorioidea*; from *χορῖον*, the chorion, and *ωειδής*, resemblance.) The second tunic of the eye, lying immediately under the sclerótica, to which it is connected by vessels. The true knowledge of this membrane, is necessary to a perfect idea of the iris and uvea. The tunica chorioidea commences at the optic nerve, and passes forwards, with the sclerotic coat, to the beginning of the cornea transparens, where it adheres very firmly to the sclerotic membrane, by means of a cellular membrane, in the form of a white fringe, called the *ciliary circle*. It then recedes from the sclerótica and cornea and ciliary circle, directly downwards and inwards, forming a round disk, which is variously coloured; hence blue, black eyes, &c. This coloured portion, reflected inwards, is termed the *iris*, and its posterior surface is termed *uvea*. The choroid membrane is highly vascular, and its external vessels are disposed like stars, and termed *vasa vorticosa*. The internal surface of this membrane is

covered with a black pigment, called the pigment of the choroid membrane.

CHOROID PLEXUS. *Plexus choroidea.* A plexus of blood-vessels, situated in the lateral ventricles of the brain.

CHOROID TUNIC. See *Choroid membrane.*

CHRISIS (From *χρῖω*, to anoint.) Anunction, or anointing of any part.

Christma rose. See *Helleborus niger.*

CHRISTUM. (From *χρῖω*, to anoint.) An unguent, or ointment of any kind.

CHROMAS. A chromate, or salt, formed by the union of earthy, metallic, or alkaline bases, with chromic acid; a chromate of lead, &c.

CHROMATISMUS. (From *χρῶμα*, colour, to colour.) The morbid discoloration of any of the secretions, as of the urine, or blood.

CHROME. (From *χρῶμα*, colour; because its primary combinations impart its colour to all secondary ones.) A white metal, inclining to a grey, very brittle, and crystallizable at an elevated temperature, in feathered filaments on the surface. Its internal fracture presents in some parts close grains, in other parts needles crossing each other. It is an ingredient in the fossil, known by that name in Siberia.

Natural History.—This metal, which is extremely scarce, and exists only in the state of a metallic oxid, was discovered by Vauquelin. He found it in an ore called red-lead ore of Siberia, or *chromate of lead*. The colour of this ore is red, with a shade of yellow; when reduced to powder, it is of a bright orange. Chrome has likewise been found in combination with iron, alumine, and silex, (*chromate of iron and alumine*,) in the department of Var, in France. It is met with in irregular masses. Its colour is brown; it has very little metallic lustre. Pontier has lately found chrome combined with iron (*chromate of iron*,) in a quarry near Gussin in the road to Cavalaire. It sometimes forms large masses. The emerald of Peru and spinel ruby owe their colours to this metal.

Properties.—Chrome is obtained in small agglutinated masses of a white colour, inclining to yellow; it is very hard, extremely brittle and refractory, and crystallizes into needles.

Exposed to the heat of a blow-pipe, it is covered with a lilac-coloured crust, which becomes green on cooling. Heated by the same apparatus with borax, it does not melt; but a part, after being oxidated, is dissolved in this salt, and communicates to it a very beautiful green colour. Acids have only a weak action on this metal. The nitric is the only acid which produces any remarkable change, it converts it into an oxid. It is capable of combining with three different portions of oxygen, and forming three different oxida. It has not yet been combined with any combustible

body. It does not appear to decompose water. It is unalterable by the alkalies. The other properties of this metal are not yet ascertained, neither are its uses known; perhaps it may afford beautiful and durable colours to the painter or the enameller.

Method of obtaining Chrome. Chrome is obtained from its native combinations, by decomposing them by the alkaline carbonates, precipitating the chromic acid, and heating it strongly in a crucible.

The following method is recommended by Vauquelin. Seventy-two parts of chromic acid are to be introduced into a charcoal crucible, placed within another of porcelain, filled with charcoal dust. The apparatus is then to be put in a furnace, and subjected to a very strong heat. Metallic chrome will then be found in the charcoal crucible. From seventy-two parts, Vauquelin obtained forty-three of metal.

CHROMIC ACID. This is obtained by decomposing the chromate of lead by potash, and treating the chromate of potash with nitric or muriatic acid. It is of an orange-red colour, and a pungent metallic taste; very soluble in water, and crystallizes by gentle evaporation.

CHRONIC (From *χρῶς*, time.) A term applied to diseases which are of long continuance, and mostly without fever. It is used in opposition to the term acute. See *Acute*.

CHRUPIA. (From *χρῶς*, colour, and *εἶναι*, sight.) *Visus coloratus.* A disease of the eyes, in which the person perceives objects of a different colour from their natural.

CHRYSANTHEMUM. (From *χρῶς*, gold, and *ἄνθος*, a flower.) 1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia*. Sun flower, or marigold.

2. Many herbs are so called whose flowers are of a bright yellow colour.

CHRYSANTHEMUM LEUCANTHEMUM. The systematic name of the great ox eye-daisy. See *Bellis major*.

CHRYSE. (From *χρῶς*, gold.) The name of a yellow plaster.

CHRYSSELETRUM. (From *χρῶς*, gold, and *ἄμπελον*, amber.) Amber of a golden yellow colour.

CHRYSIPPEA. (From *Chrysis*, its inventor. A herb enumerated by Pliny.

CHRYSITIS. (From *χρῶς*, gold.) Litharge. The yellow foam of lead. Also the herb yarrow, from the golden colour of its flower.

CHRYDOLANUS. (From *χρῶς*, gold, and *δολανος*, a nut; so named because of its colour, which, before it is dried, is yellow.) The nutmeg.

CHRYSOCOLLA. (From *χρῶς*, gold, and *κόλλα*, cement.) Gold solder.

CHRYSOCOMA. (From *χρῶς*, gold, and *κομη*, hair; so called from its golden, hair-

like appearance.) The herb milfoil, or *yarrow*.

CHRYSOSONIA. (From *χρυσος*, gold, and *συνα*, to become.) The tincture of gold.

CHRYSOLOCHANON. (From *χρυσος*, gold, and *λαχανον*, the olus; so named for its having a yellow leaf, and a flower like the olus.) The herb orach, or atriplex.

CHRYSOSPENIUM. (From *χρυσος*, gold, and *σπένειν*, spikenwort.) The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Digynia*. Golden saxifrage.

CHRYZULCUS. (From *χρυσος*, gold, and *αλα*, to take away.) The *qua regia* is so called, as having the property to dissolve gold.

CHYLARIA. (From *χυλος*, chyle.) A discharge of a whitish mucus urine, of the colour and consistence of chyle.

CHYLE. (*Χυλος*. *Chylus*.) The milk like liquor observed some hours after eating, in the lacteal vessels of the mesentery, and in the thoracic duct. It is separated by digestion from the chyme, and is that fluid substance from which the blood is formed.

The chyle is absorbed by the mouths of the lacteal vessels, which are in the greatest number in the jejunum and ilium, whilst the fæces of the chyme, with the bile, are propelled into the large intestines. The chyle of the human body smells like milk; has a sweetish taste, a white colour, and a consistence thinner than blood and milk. Its specific gravity is lighter than that of the blood, and hence it is that chyle is occasionally seen swimming on the blood, if a vein be opened some hours after eating. The quality of the chyle is similar to that of milk; for, like it, it coagulates and ascends; but sometimes its nature is altered from bad digested food or medicines: thus the chyle becomes blue, from eating indigo; yellow, from the yolk of eggs, &c. The quantity of chyle depends upon that of the ingesta, and their greater or less nourishing power: from five or six pounds of food, very little more than two pounds of chyle are elaborated.

The constituent principles of chyle, are, 1. *Water*, which forms its greatest part. 2. *Oily cream*, which chymistry teaches to be hydrogen and carbon. 3. *Cheese*, which, by the vis vitæ, is formed by the carbon and azot of the ingested food. 4. *Earth*, which is obtained from lacteal calculi, that are occasionally found in the receptaculum chyli and lacteals. 5. *Animal lymph*, which is mixed with the gastric and enteric juices.

The nutritive principles of vegetables, are starch; an albuminous principle, oil, vegetable gluten, and sugar. The nutritive principles of animal substances, are oil, jelly, and animal gluten; and hence the reason why the chyle, separated from vegetables, is of the same nature with that prepared from animal ingesta, that the

principles of both are dissolved into their elements, which are the same in animal and vegetable food: thus the *cream* of the chyle is formed of carbon and hydrogen; and the *cheese* of the chyle, from the carbon and azot of both animal and vegetable substances.

The chyle is mixed with the albuminous and gelatinous lymph in the thoracic duct, which receives them from the lymphatics.

The uses of the chyle are, 1. To supply the matter from which the blood and other fluids of our body are prepared; from which fluids the solid parts are formed. 2. By its ascescent nature, it somewhat restrains the putrescent tendency of the blood; hence the dreadful puridity of the humours from starving; and thus milk is an excellent remedy against scurvy. 3. By its very copious aqueous latex, it prevents the thickening of the fluids, and thus renders them fit for the various secretions. 4. The chyle secreted in the breasts of puerperal women, under the name of milk, forms the most excellent nutriment of all aliments for new-born infants.

CHYLIFICATION. (From *chylus*, and *φο*, to become.) *Chylification*. The process, carried on in the small intestines, and principally in the duodenum, by which the chyle is separated from the chyme.

CHYLISMA. (From *χυλος*, juice.) An expressed juice.

CHYLOPOETIC (*Chylopoeticus*; from *χυλος*, chyle, and *ποιω*, to make.) Chylopoetic. Any thing connected with the formation of chyle; thus chylopoetic viscera, chylopoetic vessels, &c.

CHYLOSIS. (From *χυλαω*, to express the juice from any thing.) Chylification, or the changing the food into chyle.

CHYLOSTAGMA (From *χυλος*, juice, and *σταω*, to distil.) The distillation or expression of any juice, or humid part from its dry one.

CHYLOSTAGMA DIAPHORETICUM MINDEBERI. A distillation of Venice treacle and mithridate.

CHYME. (*Chymus*; from *χυμος*, which signifies humour or juice.) The ingested mass of food, that passes from the stomach into the duodenum, and from which the chyle is prepared in the small intestines by the admixture of the bile, &c.

CHYMIA. Chymistry.

CHYMIATER. A chymical physician. See *Chimicater*.

CHYMIATRIA. (From *χυμα*, chymistry, and *ιατρον* to heal.) The art of curing diseases by the application of chymistry to the uses of medicine.

CHYMOSIS. See *Chemosis*.

CHYLEN RADIX. A cylindrical root, of the thickness of a goose-quill, brought from China. It has a bitterish taste, and imparts a yellow tinge to the salva. The Chinese hold it in great estimation as a stomachic, infused in wine.

CYSEA. (From *χυν*, to pour out.) Fusion, or the reduction of solid bodies into fluid by heat.

CYTLON. (From *χυν*, to pour out.) An anointing with oil and water.

CIBALIS VISTULA. An obsolete term for the œsophagus.

CIBATIO. (From *cibus*, food.) In chymistry it means incorporation: also the taking of food.

CIBUR. An obsolete term for sulphur.

CICATRISANTIA. (*Cicatrissantia*, sc. remedia; from *cicatrigo*, to skin over.) Such applications as dispose wounds and ulcers to dry up and heal, and to be covered with a skin.

CICATRIX. (From *cicatrigo*, to heal up or skin over.) A seam or scar upon the skin after the healing of a sore or ulcer.

CICER. (A plant so called. The Cicerones had their name from this pulse, as the Pisones had from the pisum or pea, and the Lentuli from the lens or lentil.)

1. The name of a genus of plants in the Linnæan system. Class, *Diadelphia*. Order, *Decandria*. The vetch.

2. The pharmacopœial name of the common cich or ciches. *Erebinthus*. *Cicer arietinum* of Linnæus:—*foliis serratis*. The seeds have been employed medicinally, but are now fallen into disuse. In some places they are toasted, and used as coffee; and in others, ground into a flour for bread. The colour of the aryllus of the seed is sometimes white, red, or black: hence the distinction into *cicer album*, *rubrum*, and *nigrum*.

CICERARIETINUM. The systematic name of the cicer plant.

CICERA. (From *cicer*, the vetch.) A small pill of the size of a vetch.

CICERA TABTARI. Small pills composed of turpentine and cream of tartar, of the size of a vetch.

CICHORIUM. (Originally, according to Pliny, an Egyptian name, and adopted by the Greeks. It is written sometimes *Κίχουρον*: whence Horace has—*cichoreæ, leuœque malvæ*: sometimes *Κίχουρον*, or *Κίχουρον*. It is supposed to have this name, *αρχαίο το δέντρο χίχουρον*, from its creeping through the fields.—Others derive it from *αίχμα, introitus*; on account of its being so readily found, or so common.) Succory.

1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia equalis*.

2. The pharmacopœial name of the wild chicory. This plant, called *Cichoreum*, *Cichorium scytostre vel officinarum*, is the *Cichorium intybus* of Linnæus:—*floribus geminis, seedilibus; foliis runcinatis*.

It belongs to the same family with the garden endive, and by some botanists has been supposed to be the same plant in its uncultivated state; but the endive commonly used as sallad is an annual, or at most a biennial plant, and its parent is now

known to be the *cichorium endivia*. Wild succory or cichory, abounds with a milky juice, of a penetrating bitterish taste, and of no remarkable smell or particular flavour: the roots are more bitter than the leaves or stalks, and these much more so than the flowers. By culture in gardens and by blanching, it loses its bitterness, and may be eaten early in the spring in sallads. The roots, if gathered before the stem shoots up, are also eatable, and when dried may be made into bread. The roots and leaves of this plant are stated by Lewis to be very useful aperients, acting mildly and without irritation, tending rather to abate than to increase heat, and which may therefore be given with safety in hectic and inflammatory cases. Taken freely, they keep the belly open, or produce a gentle diarrhœa; and when thus continued for some time, they have often proved salutary in the beginning obstructions of the viscera, in jaundices, cachexies, hypochondriacal and other chromical disorders. A decoction of this herb, with others of the like kind, in whey, and rendered purgative by a suitable addition of polychrest salt, was found a useful remedy in cases of biliary calculi, and promises advantage in many complaints requiring what have been termed attenuants, and resolvents. The virtues of succory, like those of dandelion, reside in its milky juice; and we are warranted, says Dr. Woodville, in asserting that the expressed juice of both these plants, taken in large doses frequently repeated, has been found an efficacious remedy in phthisis pulmonalis, as well as the various other remedies above mentioned. The milky juice may be extracted by boiling in water, or by pressure. The wild and the garden sorts are used indifferently. If the root is cut into small pieces, dried, and roasted, it resembles coffee, and is sometimes a good substitute for it.

CICHORIUM ENDIVIA. The systematic name of the endive. See *Endivia*.

CICHORIUM INTYBUS. The systematic name of the wild chicory. See *Cichorium*.

Chichory. See *Cichorium*.

Chichory, wild. See *Cichorium*.

CICINDELLA. (A dim. of *candelis*: i. e. a little candle; so called from its light.) The glow-worm. Some think them anodyne, other lithontriptic, though as the editor of Motherby's Dictionary justly observes, probably neither.

CICINUM OLEUM. (From *καμ*, the ricinus.) An oil, obtained by boiling the bruised seeds of the *Jatropha curcas* of Linnæus. It is somewhat similar in its properties to castor oil. See *Ricinus*.

CICLA. A name for the beta alba.

CICUTA. (*Quasi cecuta*, blind; because it destroys the sight of those who use it. *Cicuta* signifies the internode, or space between two joints of a reed: the hollow stem of any plant which the shepherds used for making their rural pipes.

Est mihi disparibus septem conjuncta cicutis festula.—Virgil.) Hemlock.

1. The name of a genus of plants in the Linnean system. Class, *Pentandria*. Order, *Digynia*.

2. The name, in most pharmacopœias, of the common hemlock, or *Conium maculatum* of Linnæus:—*seminibus striatis*. It is called by some *camaran*; by others *abitos*; and according to Erotian, *cambeison* is an old Sicilian word for *cicuta*. *Cicuta major fetida*. *Conium* of the last London pharmacopœia.

Hemlock is found in every part of England, and is distinguished from those plants which bear some remembrance to it, by the spotted stem. It is generally believed to be a very active poison. In a very moderate dose it is apt to occasion sickness and vertigo; in a larger quantity it produces anxiety, cardialgia, vomiting, convulsions, coma, and death. Baron Stoerk was the first who brought hemlock into repute as a medicine of extraordinary efficacy; and although we have not in this country any direct facts, like those mentioned by Stoerk, proving that inveterate scirrhus cancers, ulcers, and many other diseases hitherto deemed irremediable, were completely cured by the *cicuta*; we have however the testimonies of several eminent physicians, shewing that some complaints, which had resisted other powerful remedies, yielded to hemlock; and that even some disorders, which, if not really cancerous, were at least suspected to be of that tendency, were greatly benefited by this remedy. In chronic rheumatisms, some glandular swellings, and in various fixed and periodical pains, the *cicuta* is now very generally employed; and from daily experience, it appears in such cases to be a very efficacious remedy. It has also been of singular use in the whooping-cough. Nor is it less efficacious when applied externally; a poultice made of oatmeal and the expressed juice, or a decoction of the extract, when the former cannot be obtained, allays the most excruciating torturous pains of a cancer, and thus gives rest to the distracted patient.

The proper method of administering *conium* internally, is to begin with a few grains of the powder or inspissated juice, and gradually to increase the dose until a giddiness affects the head, a motion is felt in the eyes as if pressed outwards, with a slight sickness and trembling agitation of the body. One or more of these symptoms are the evidence of a full dose, which should be continued until they have ceased, and then after a few days the dose may be increased; for little advantage can be expected but by a continuance of the greatest quantity the patient can bear. In some constitutions even small doses greatly offend, occasioning spasms, heat and thirst; in

such instances it will be of no service. As the powder of the dried leaves has been thought to act, and may be depended upon with more certainty than the extract, the following direction should be observed in the preparation; gather the plant about the end of June, when it is in flower; pick off the little leaves, and throw away the leafstalks: dry the small selected leaves in a hot sun, or in a tin or pewter dish before the fire. Preserve them in bags made of strong brown paper, or powder them and keep the powder in glass phials where the light is excluded; for light dissipates the beautiful green colour very soon, and thus the medicine loses its appearance, if not its efficacy: this mode is recommended by Dr. Withering. The extract should also be made of the plant gathered at this period. From 10 to 20 grains of the powder may be taken twice or thrice a day.

CICUTA AQUATICA. *Cicutaria virosa*. *Sium majus alterum angustifolium*. *Sium cruce folio*. Long-leaved water hemlock and cowbane. This plant, *Cicuta virosa* of Linnæus:—*umbellis oppositifoliis; petiolis marginatis obtusis*, is seldom employed medicinally in the present day. It is an active poison, and often eaten by mistake for the wild smallage, the *Apium graveolens* of Linnæus; when it produces tremors, vertigo, a violent burning at the stomach, epilepsy, convulsions, spasms of the jaw, a flowing of blood from the ears, tumefaction of the abdomen, and death.

CICUTA VIROSA. The systematic name of the water hemlock. See *Cicuta aquatica*.

CICUTARIA. (*Cicutaria*; from *cicuta*, hemlock.) Bastard hemlock. This plant, *Cherophyllum sylvestre* of Linnæus:—*caule lavi striato; geniculis tumidiusculis*, is often mistaken for the true hemlock. It may with great propriety be banished from the list of officinals, as it possesses no remarkable property. See *Cherophyllum*.

CICUTARIA AQUATICA. A name for the phellœdrium aquaticum.

CICUTARIA VIROSA. See *Cicuta aquatica*.

CINONIUM. See *Cydonium*.

CILIAR LIGAMENT. (From *cileo*, to move about.) *Ligamentum ciliare*. The circular portion that divides the choroid membrane from the iris, and which adheres to the sclerotic membrane. It appears like a white circular ring. See *Choroid membrane*.

CILIARE LIGAMENTUM. See *Choroid membrane*.

CILIARIS MUSCULUS. That part of the musculus orbicularis palpebrarum which lies nearest the cilia, considered by Riolan as a distant muscle.

CILIUM. (From *cileo*, to move about.) The hair on the eyelid or eyelash.

CILIARY PROCESSES. The white folds

at the margin of the uvea in the eye, covered with a black matter, which proceeds from the uvea to the crystalline lens, upon which they lie.

CILLO. (From *cilium*, the eyelid.) One who is affected with a spasm or trembling of the eyelids.

CILLOSI. (From *cilium*, the eyelid.) A spasmodic trembling of the eyelids.

CIMEX. (From *cupias*, to inhabit; so called because they infect houses.) The wall-louse or bug. *Cimex domesticus*. Six or seven are given inwardly to cure the ague, just before the fits come on, and have the same effect with every thing nauseous and disgusting.

CIMOLIA ALBA. (From *Κίμολος*, Cimolus, an island in the Cretan sea, where it is procured.) Tobacco-pipe clay. Its virtues are similar to those of the bolar earths; but it is never administered medicinally.

CIMOLIA PURPURESCENS. Fullers-earth. A bolar earth, of a grayish brown colour.

CINA CINÆ. See *China chinæ*.

CINÆ SEMEN. See *Santonicum*.

CINARA. (From *κινω*, to move; *quasi movet ad venerem vel urinam*.) Artichoke.

1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia equalis*.

2. The pharmacopœial name for the common artichoke. *Alloccalum*. *Articocalus*. *Artischocæ levis*. *Costus nigra*. *Carduus sativus non spinosus*. *Cinara hortensis*. *Scolymus sativus*. *Carduus domesticus capite majore*. *Carduus altiss.* The species thus called, and which is used in medicine, is the *Cinara scolymus* of Linnæus: — *foliis subspinosis pinnatis indivisisque calycinis squamis oratis*. A native of the southern parts of Europe, but cultivated here for culinary purposes. The leaves are bitter, and afford, by expression, a considerable quantity of juice, which, when strained, and mixed with an equal quantity of white wine, has been given successfully in dropsies, the dose of 3 or 4 table-spoonfuls every night and morning, but it is very uncertain in its operation.

CINCHONA. (Geoffroy states that the use of this bark was first learned from the following circumstance:—Some cinchona trees being thrown by the winds into a pool of water, lay there till the water became so bitter, that every body refused to drink it. However, one of the neighbouring inhabitants being seized with a violent paroxysm of fever, and finding no other water to quench his thirst, was forced to drink of this, by which he was perfectly cured. He afterwards related the circumstance to others, and prevailed upon some of his friends, who were ill of fevers, to make use of the same remedy, with whom it proved equally successful. The use of this excellent remedy, however, was very little known till about the year 1638, when

a signal cure having been performed by it on the Spanish viceroy's lady, the Countess del Cinchon, at Lima, and it came into general use, and hence it was distinguished by the appellation of *cortex cinchonæ*, and *pulvis comitessæ*, or the Countess's powder. On the recovery of the Countess, she distributed a large quantity of the bark to the Jesuits, in whose hands it acquired still greater reputation, and by them it was first introduced into Europe, and thence called *cortex*, or *pulvis jesuiticus*, *pulvis patrium*; and also Cardinal del Lugo's powder, because that charitable prelate bought a large quantity of it at a great expense, for the use of the religious poor at Rome.) *Cortex china*, *chinu* or *chinchina*; *kina kina*, or *kinkina*; and *quina quina*, or *quinquina*.

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*. *Cinchona*, or Peruvian bark-tree.

2. The pharmacopœial name of several kinds of barks; the trees affording which grow wild in the hilly parts of Peru; the bark is stripped from the branches, trunk and root, and dried. Three kinds of it are now in use.

CORTX CINCHONÆ CORDIFOLIÆ.

The plant which affords this species is the *Cinchona cordifolia* of Zee: the *Cinchona officinalis* of Linnæus: the *Cinchona macrocarpa* of Willdenow. Heart-leaved cinchona.

The bark of this tree is called *yellow bark*, because it approaches more to that colour than either of the others do, is in flat pieces, not convoluted like the pale, nor dark-coloured like the red; externally smooth, internally of a light cinnamon colour, friable and fibrous; has no peculiar odour different from the others, but a taste incomparably more bitter, with some degree of astringency.

CORTX CINCHONÆ LANCEFOLIÆ.

This species is obtained from the *Cinchona lancefolia* of Zee. Lance-leaved cinchona. This is the *Quilled bark*, which comes in small quilled twigs, breaking close and smooth, friable between the teeth, covered with a rough coat of a brownish colour, internally smooth and of a light brown; its taste is bitter and slightly astringent; flavour slightly aromatic, with some degree of mustiness.

CORTX CINCHONÆ OBLONGIFOLIÆ.

This kind is procured from the *Cinchona oblongifolia* of Zee. Oblong-leaved cinchona. This bark is the *red bark*: it is in large thick pieces, externally covered with a brown rugged coat, internally more smooth and compact, but fibrous; of a dark-red colour; taste and smell similar to that of the *cinchona lancefolia cortex*, but the taste rather stronger.

From the general analysis of bark, it

appears to consist, besides the woody matter which composes the greater part of it, of gum, resin, gallic acid, of very small portions of tannin and essential oil, and of several salts, having principally lime for their basis. Sequin also supposed the existence of gelatin in it, but without sufficient proof. Cold water infused on pale bark for some hours acquires a bitter taste, with some share of its odour; when assisted by a moderate heat, the water takes up more of the active matter; by decoction, a fluid, deep-coloured, of a bitter styptic taste, is obtained, which, when cold, deposits a precipitate of resinous matter and gallic acid. By long decoction, the virtues of the bark are nearly destroyed, owing to the oxygenation of its active matter. Magnesia enables water to dissolve a larger portion of the principles of a bark, as does lime, though in an inferior degree. Alcohol is the most powerful solvent of its active matter. Brandy and other spirits and wines afford also strong solutions, in proportion to the quantity of alcohol they contain. A saturated solution of ammonia is also a powerful solvent; vinegar is less so even than water. By distillation, water is slightly impregnated with the flavour of bark; it is doubtful whether any essential oil can be obtained.

The action of menstrua on the red bark is nearly the same, the solutions only being considerably stronger, or containing a larger quantity of resinous matter and of the astringent principle.

The analysis of the yellow bark shows that its active principles are more concentrated than in either of the others, affording to water, alcohol, &c. tinctures much stronger, both in bitterness and astringency, especially in the former principle.

From the general analysis of these barks, it appears that they consist of nearly the same proximate principles, which vary in their proportions; the most active component parts are the resin, extractive matter and the gallic acid, and these in combination probably constitute the tonic quality of bark. In the best pale bark this active matter amounts to about one-eighth.

The red bark has been considered as superior to the pale, the yellow is represented, apparently with justice, as being more active than either of the others.

The effects of Peruvian bark are those of a powerful and permanent tonic, so slow is its operation, that its stimulating property is scarcely perceptible by any alteration in the state of the pulse, or of the temperature of the body. In a large dose, it occasions nausea and head-ache; in some habits it operates as a laxative; in others it occasions costiveness. It is one of those medicines, the efficacy of which, in removing

disease, is much greater than could be expected, *a priori*, from its effects on the system in a healthy state.

Intermittent fever is the disease, for the cure of which bark was introduced into practice, and there is still no remedy which equals it in power.

The disputes respecting the mode of administering it are now settled. It is given as early as possible, with perhaps the previous exhibition of an emetic, to evacuate the stomach; it is repeated in the dose of one scruple or half a drachm every second or third hour, during the interval of the paroxysm; and it may even be given during the hot fit, but it is then more apt to excite nausea.

In remittent fever it is given with equal freedom, even though the remission of the fever may be obscure.

In some forms of continued fever which are connected with debility, as in typhus, cynanche maligna, confluent small-pox, &c. it is regarded as one of the most valuable remedies. It may be prejudicial, however, in those diseases, where the brain, or its membranes are inflamed, or where there is much irritation, marked by subsultus tendinum, and convulsive motions of the extremities; and in pure typhus it appears to be less useful in the beginning of the disease than in the convalescent stage.

Even in fevers of an opposite type, where there are marks of inflammatory action, particularly in acute rheumatism, bark has been found useful after blood-letting. In erysipelas, in gangrene, in extensive suppuration and venereal ulceration, the free use of bark is of the greatest advantage.

In the various forms of passive hæmorrhagy, in many other diseases of chronic debility, dyspepsia, hypochondriasis, paralysis, rickets, scrophula, dropsy, and in a variety of spasmodic affections, epilepsy, chorea, and hysteria, it is administered as a powerful and permanent tonic, either alone, or combined with other remedies suited to the particular case.

Its usual dose is half a drachm. The only inconvenience of a larger dose is its sitting uneasy on the stomach. It may therefore, if necessary, be frequently repeated, and in urgent cases may be taken to the extent of an ounce, or even two ounces, in twenty-four hours.

The powder is more effectual than any of the preparations: it is given in wine, in any spirituous liquor; or, if it excite nausea, combined with an aromatic. The cold infusion is the least powerful, but most grateful; the decoction contains much more of the active matter of the bark, and is the preparation generally used when the powder is rejected; its dose is from two to four ounces. The spirituous tincture, though containing still more of the bark, cannot

be extensively used on account of the menstruum, but is principally employed, occasionally, and in small doses of two or three drachms, as a stomachic. The extract is a preparation of considerable power, when properly prepared, and is adapted to those cases, where the remedy requires to be continued for some time. It is then given in the form of pill, in a dose from five to fifteen grains.

Bark is likewise sometimes given in the form of enema; one scruple of the extract, or two drachms of the powder, being diffused in four ounces of starch mucilage. The decoction is also sometimes applied as a fomentation to ulcers.

CINCHONA CARIBÆA. The systematic name of the Caribæan bark-tree. It grows in Jamaica, where it is called the sea-side beech. According to Dr. Wright, the bark of this tree is not less efficacious than that of the cinchona of Peru, for which it will prove an useful substitute: but by the experiments of Dr. Skeete, it appears to have less astringent power.

CINCHONA FLORIBUNDA. The systematic name of the plant which affords the Saint Lucé bark. Dr. Withering considers this bark as greatly inferior to that of the other species of this genus. In its recent state it is considerably emetic and cathartic, properties which in some degree it retains on being dried; so that the stomach does not bear this bark in large doses, and in small ones its effects are not such as to give it any peculiar recommendation.

CINCHONA OFFICINALIS. The name of the officinal Peruvian bark. See *Cinchona*.

CINCHONA SANCTA FÉ. Several species of cinchona have been lately discovered at Sancta Fé, yielding barks both of the pale and red kind; and which, from their sensible qualities, are likely upon trial to become equally useful with those produced in the kingdom of Peru.

CINCHONA RUBRA. See *Cinchona*.

CINCHONA FLAVA. See *Cinchona*.

CINCINNUS. The hair on the temples. See *Capillus*.

CINCLESIS. (From *κινεῖν*, to move.) *Cinismus*. An involuntary nictitation or winking. *Vogel*.

CINERARIUM. (From *cinis*, ashes.) The ash-hole of a chymical instrument.

CINERES. (plur. of *cinis*, ashes.) Ashes.

CINERES CLAVELLATI. (*Clavellatus*; from *clarus*, a wedge. The name of *cineres clavellati* originated from the little wedges or billets into which the wood was cut to make potash.) *Alumen cutinum*. *Sal alkali fixum*. *Cineres russici*. *Kali Potassa gastrinum*. *Kali impurum*. Impure potash or pearl ash. The name adopted in the new London pharmacopœia is *potassa impura*. It is from this salt the various preparations of potash are made.

The ancients called the ashes of burnt-wood *lix*, from whence the modern word *lixivia*. The English name *potash* is from the pots in which the *lixivium* was boiled.

CINERES RUSSICI. See *Cineres clavellati*. **CINERITIOUS.** (From *cinis*, ashes.) Of the colour of ashes. A name applied to the cortical substance of the brain, from its resemblance to an ash-colour.

CINERITIUM. (From *cinis*, ashes.) A cupel or test; so named from its being commonly made of the ashes of vegetables or bones.

CINERULAN. A name for *spodium*.

CINETUS. An epithet formerly applied to the diaphragm.

CINGULARIA. (From *cingulum*, a girdle; because it grows in that shape.) The *lycopodium*.

CINGULUM. (From *cingo*, to bind.) A girdle or belt about the loins.

CINGULUM MERCURIALE. A mercurial girdle, called also *cingulum sapientie*, and *cingulum stultitie*. It was an invention of Rulundus's; different directions are given for making it, but the following is one of the neatest: "Take three drams of quicksilver; shake it with two ounces of lemon-juice until the globules disappear; then separate the juice and mix with the extinguished quicksilver half the white of an egg; gum-dragon finely powdered, a scruple; and spread the whole on a belt of flannel.

CINGULUM SANCTI JOHANNIS. A name of the *artemisia*.

CINIFICATUM. A name for *calcinatum*.

CINNABAR. (Pliny says the Indians call by this name a mixture of the blood of the dragon and elephant, and also many substances which resemble it in colour, particularly the *minium*.) *Minium purum*. *Minium Græcorum*. *Magnes eplépsia*. *Atzemafor*. *Amisyon*. *Azamar*. Vitruvius calls it *anthrax*. A red mineral substance composed of mercury combined with sulphur. It is either native or factitious. The native is an ore of quicksilver moderately compact, and of an elegant striated red colour. This kind of cinnabar, artificial cinnabar, employed as a factitious cinnabar, is a mixture of mercury and sulphur sublimed, and thus reduced to a fine red substance. The best is of a high colour and full of needle-like spiculae. See *Sulphuretum hyargyri rubrum*. Cinnabar is often employed as a mild mercurial, and as an alterative. Hoffman greatly recommends it as a sedative and antispasmodic. Others deny that cinnabar taken internally has any medicinal quality; and their opinion is grounded on the insolubility of it in any menstruum. It is found in the Dutchy of Deuxponts, in the Palatinate, in Spain, South America, &c. It is called native vermilion, and cinnabar in flowers.

CINNABAR FACTITIA. See *Sulphuretum hydrargyri rubrum*.

CINNABAR NATIVA. See *Cinnabar*.

CINNABARIS GRÆCORUM. The sanguis draconis and cinnabar.

CINNAMOMUM. (From *kinamon*, Arab.) Cinnamon. The tree which affords the true cinnamon, which is its inner bark, is the *Laurus cinnamomum* of Jacquin:—*foliis trinerviis ovato-oblongis: nervis versus apicem evanescentibus*. Cinnamon bark is one of the most grateful of the aromatics; of a fragrant smell, and a moderately pungent, glowing, but not fiery taste, accompanied with considerable sweetness, and some degree of astringency. It is one of the best cordial, carminative, and restorative spices we are in possession of, and is generally mixed with the diet of the sick. The essential oil, on account of its high price, is seldom used: a tincture, simple and spirituous water, are directed to be kept in the shops. The watery infusion of cinnamon is given with advantage to relieve nausea and check vomiting.

Cinquefoil. See *Pentaphyllum*.

CION. (*Kion*, a column, from *κίον*, to move.) The uvula was formerly so named from its pyramidal shape: also an enlargement of the uvula.

CIONIS. (From *κίον*, the uvula.) A diseased enlargement and painful swelling of the uvula.

CIRCEÆA. (From *Circe*, the enchantress; so named from the opinion, that it was used by Circe in her enchanted preparations.) Enchanter's nightshade.

1. The name of a genus of plants in the Linnæan system. Class, *Diandria*. Order, *Monogynia*.

2. The name in some pharmacopœias for the *Circea lutiana*, which is now fallen wholly into disuse.

CIRCOCELE. (From *κίρκος*, varix, or a dilatation of a vein, and *κύλα*, a tumour.)

Varicocele. A varicose distension and enlargement of the spermatic veins; and whether considered on account of the pain, or on account of the wasting of the testicle, which now and then follows, it may truly be called a disease. It is frequently mistaken for a descent of a small portion of omentum. The uneasiness which it occasions, is a kind of pain in the back, generally relieved by suspension of the scrotum. It has been resembled to a collection of earthworms. It is most frequently confined to that part of the spermatic process, which is below the opening in the abdominal tendon; and the vessels generally become rather larger as they approach the testes. There is one sure method of distinguishing between a circocele and omental hernia: place the patient in a horizontal posture, and empty the swelling by pressure upon the scrotum; then put the fingers firmly upon the upper part of the abdomi-

nal ring, and desire the patient to rise; if it is a hernia, the tumour cannot re-appear, as long as the pressure is continued at the ring: but if a circocele, the swelling returns with increased size, on account of the return of blood into the abdomen being prevented by the pressure.

CIRCOS. (From *κίρκος*, to roll up.) A ring. It is sometimes used for the sphincter muscle, which is round like a ring.

CIRCULATION. (*Circulatio*; from *circulo*, to compass about.) A vital action performed by the heart in the following manner: the blood is returned into the right auricle of the heart by the descending and ascending *venæ cavæ*, which, when distended, contracts and sends its blood into the right ventricle; from the right ventricle it is propelled through the pulmonary artery, to circulate through, and undergo a change in, the lungs, being prevented from returning into the right auricle by the closing of the valves, which are situated there for that purpose. Having undergone this change in the lungs, it is brought to the left auricle of the heart by the four pulmonary veins, and from thence it is evacuated into the left ventricle. The left ventricle, when distended, contracts, and throws the blood through the aorta to every part of the body, to be returned by the veins into the two *venæ cavæ*. It is prevented from passing back from the left ventricle into the auricle by a valvular apparatus; and the beginning of the pulmonary artery and aorta is also furnished with similar organs, to prevent its returning into the ventricles.—(See *Heart*.) It is by means of this important action, that every part of the body lives, becomes warm, and is nourished, the various secretions separated, and the chyle converted into blood. In the fœtus the blood passes from the umbilical veins, partly into the vena portæ, and partly through the *canalis venosus*, into the ascending cava. The lungs being contracted, a very small quantity circulates through them, and the greater part flows through the *canalis arteriosus* and *foramen ovale* to the left side of the heart, and into the aorta, and is carried back by the umbilical arteries to the placenta.

CIRCULATOR. (From *circulo*, to compass about.) A wandering practitioner in medicine. A quack. A mountebank.

CIRCULATORIUM. (From *circulo*, to move round.) A chemical digesting vessel in which the fluid performs a circulatory motion.

CIRCULUS. (Dim. of *circus*, a circle.) A circle or ring. Any part of the body which is round or annular, as *circulus oculi*. A round chemical and chirurgical instrument, sometimes called *abbreviatorium* by the old chymists.

CIRCULUS ARTERIOSUS IRIDIS. The

artery which runs round the iris and forms a circle, is so termed.

CIRCULUS QUADREPLEX. A bandage.

CIRCUMCAULALIS. A name of the adnata of the eye.

CIRCUMCISION. (*Circumcisio*, from *circumcindo*, to cut about.) The cutting off the prepuce from the glans penis; an ancient custom, still practised amongst the Jews.

CIRCUMFLEXUS. (*Circumflexus*, sc. *musculus*.) *Tensor palati* of Innes. *Circumflexus palati mollis* of Albinus. *Sphenosalphingo-staphylinus*, seu *staphylinus externus* of Winslow. *Musculus tube nose* of Valsalva. *Palato-salphingus* of Douglass. *Pterigo-staphylinus* of Cowper, and *Petrosalphingo-staphylin* of Dumas. This muscle arises from the spinous process of the sphenoid bone, behind the foramen ovale, which transmits the third branch of the fifth pair of nerves, from the Eustachian tube, not far from its osseous part; it then runs down along the pterygoideus internus, passes over the hook of the internal plate of the pterygoid process by a round tendon, which soon spreads into a broad membrane. It is inserted into the velum pendulum palati, and the semilunar edge of the os palati, and extends as far as the suture which joins the two bones. Generally some of its posterior fibres join with the constrictor pharyngis superior, and palato-pharyngæus. Its use is to stretch the velum, to draw it downwards, and to a side towards the hook. It hath little effect upon the tube, being chiefly connected to its osseous part.

CIRCUMGYRATIO. (From *circumgyro*, to turn round.) Circumgyration, or the turning a limb round in its socket.

CIRCUMLITIO. (From *circumlino*, to anoint all over.) A medicine used as a general unction or liniment to the part.

CIRCUMSSALIS. (From *circum*, about, and *os*, a bone.) Surrounding a bone as the periosteum does; or surrounded by a bone.

CIRCUS. (From *carka*, to surround. Chald.) A circle or ring. A circular bandage, called also plinthius lacqueus.

CIRRESIA. (From *supra*, to mix.) An union of separate things.

CIRSIUM ARVENSE. (From *supra*, a vein or swelling of a vein, which this herb was supposed to heal.) The *carduus hæmorrhoidalis*.

CIRSOCELE. See *Circocèle*.

CIRSOIDES. (From *supra*, a varix, and *similis*, likeness.) Resembling a varix: an epithet applied by Rufus Ephesius to the upper part of the brain.

CIRSOS. (From *supra*, to dilate.) A varix or preternatural distension of any part of a vein.

CISSA. (From *ciroa*, a glutinous bird.) A depraved appetite proceeding from previous gluttony and voracity.

CISSAMPELOS. (From *ciroa*, ivy, and *amela*, the vine.) The name of a genus of plants in the Linnæan system. Class, *Dioccia*. Order, *Monadelphia*. The wild vine with leaves like ivy.

CISSAMPELOS PARIERA. The systematic name of the *pariera brava*. See *Pariera brava*.

CISSINUM. (From *ciroa*, ivy.) The name of a plaster mentioned by Ægineta.

CISTA. (From *ciroa*, to deposit.) A cyst.

CISTERNA. (From *cista*, a cyst.) The fourth ventricle of the brain is so called from its cavity; also the lacteal vessels in women.

CISTUS. (*Kistoc*, the derivation of which is uncertain.) The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Monogynia*. The *Cistus*.

CISTUS CRETICUS. (From *kis*, Heb.) *Cisthortia*. *Cissarus*. *Dorycnium*. The systematic name of the plant from which the laudanum of the shops is obtained. See *Ladanum*.

CISTUS HUMILIS. A name of *parnassia* or white liverwort.

CISTUS LADANIFERA. See *Ladanum*.

CISTUS LEDON. A name of *Ledum palustre*. See *Rosmarinus*.

CITHARUS. (From *ciroa*, a harp.) The breast is sometimes so named from its shape.

CITRAGO. (From *citrus*, a citron, so called from its citron-like smell.) *Citraria*. *Melissa* or *baum*.

CITRAS. (From *citrus*, the citron.) A citrate. A salt formed by the union of the citric acid, or acid of lemons, with different bases; as the *citrate of alumina*, *citrate of ammonia*, *citrate of potash*.

CITREA. See *Limon*.

CITREUM. (From *citrus*.) *Citron*. *Malus medica*. *Malus citria*. The citron-tree. *Citrus medica* of Linnæus. Its fruit is called *Cedromela*, which is larger and less succulent than the lemon; but in all other respects the citron and lemon trees agree. The lemon is a variety only of the *citrus medica*. The citron juice when sweetened with sugar is called by the Italians *cigre di cedre*.

CITRIC ACID. *Acidum citricum*. The juice of lemons. The citric acid may be obtained pure in concrete crystals, by the following method: Saturate boiling lemon-juice with pulverized chalk. The acid forms with lime a salt that is scarce soluble, and the mucilaginous and extractive substances remain dissolved in the supernatant liquor; the precipitate is to be washed with lukewarm water, till it ceases to deepen in colour; it dissolves nearly as well as sulphate of lime; it is then to be treated with as much sulphuric acid as is requisite to saturate the chalk, diluted in ten parts of water; and this mixture is to be boiled

for a few minutes. Afterwards, it must be cooled and filtered; the sulphate of lime remains on the filter, and the liquor affords a crystallized acid by evaporation. See *Limon*.

CITRINATIO. Complete digestion.

CITRINULA. (A Dim. of *citrus*.) A small citron.

Citron. See *Limon*.

Citrus, Sicilian. See *Citrullus*.

CITRULLUS. *Angura.* *Jace brasiliensis.* *Tetraguria.* Sicilian citrul, or water-melon. The seeds of this plant, *Cucurbita citrullus* of Linnæus — *foliis multipartitis*, were formerly used medicinally, but now only to re-produce the plant. Water-melon is cooling, and somewhat nutritious; but so soon begins to ferment, as to prove highly noxious to some stomachs, and bring on spasms, diarrhæas, cholera morbus, colics, &c.

CITRUS. 1. The name of a genus of plants in the Linnæan system. Class, *Polyadelphia.* Order, *Icosundria.*

2. The name of the lemon. See *Limon*.

CITRUS AURANTIUM. The systematic name of the orange tree. See *Aurantium*.

CITRUS MEDICA. The systematic name of the lemon-tree. See *Limon*.

CITTA. A voracious appetite.

CITROSIS. See *Chlorosis*.

Civet-cat. See *Zibethum*.

CIVETTA. (From *sebet*, Arab.) Civet. An unctuous odoriferous drug now only used by perfumers.

Clap. See *Gonorrhæa*.

CLARET. (*Claretum*; from *claro*, to be clear.) A French wine, that may be given with great advantage, as a tonic and antiseptic, where red port wine disagrees with the patient; and in typhoid fevers of children and delicate females, it is far preferable as a common drink.

CLARETUM. See *Claret*. Also a wine impregnated with spices and sugar, called by some *Vinum Hippocraticum*. A *Claretum purgatorium*, composed of a vinous infusion of glass of antimony in cinnamon water with sugar, is mentioned by Schroeder.

CLARIFICATIO. The depuration of any thing.

Clary. A name for the *horminum*.

CLASIS. (From *κλασ*, to break.) *Clasma.* A fracture.

CLAUSTRUM. (From *claudo*, to shut.) *Cleithrum gutturis*. Any aperture which has a power of contracting itself, or closing its orifice by any means, as the passage of the throat.

CLAUSTRUM VIRGINITATIS. The hymen.

CLAUSURA. (From *claudo*, to shut.) An imperforation of any canal or cavity in the body. Thus *clausura uteri* is a preternatural imperforation of the uterus; *clausura tubarum Fallopiarum*, a morbid imperforation of the Fallopian tubes, men-

tioned by Ruysch as one cause of infecundity.

CLAVATIO. (From *clava*, a club.) A sort of articulation without motion, where the parts are, as it were, driven in with a hammer, like the teeth in the sockets. See *Gomphosis*.

CLAVELLATUS. (From *clavus*, a wedge.) A specific name of potash, or cineres clavellati; and so named from the little wedges, or billets, into which the wood was cut to make it.

CLAVICLE. (Dim. of *clavis*; so called from its resemblance to an ancient key.) Collar-bone. The clavicle is placed at the root of the neck, and at the upper part of the breast. It extends across, from the tip of the shoulder to the upper part of the sternum; it is a round bone, a little flattened towards the end, which joins the scapula; it is curved like an Italic S, having one curve turned out towards the breast; it is useful as an arch, supporting the shoulders, preventing them from falling forwards upon the breast, and making the hands strong antagonists to each other; which, without this steadying, they could not have been.

1. The thoracic end, that next the sternum, or what may be called the inner head of the clavicle, is round and flat, or button-like; and it is received into a suitable hollow on the upper piece of the sternum. It is not only like other joints surrounded by a capsule or purse; it is further provided with a small moveable cartilage, which, like a friction-wheel in machinery, saves the parts and facilitates the motions, and moves continually as the clavicle moves.

2. But the outward end of the clavicle is flattened, as it approaches the scapula, and the edge of that flatness is turned to the edge of the flattened acromion, so that they touch but in one single point. This outer end of the clavicle, and the corresponding point of the acromion, are flattened and covered with a crust of cartilage; but the motion here is very slight and quite insensible; they are tied firmly by strong ligaments; and we may consider this as almost a fixed point; for there is little motion of the scapula upon the clavicle; but there is much motion of the clavicle upon the breast, for the clavicle serves as a shaft, or axis, firmly tied to the scapula, upon which the scapula moves and turns, being connected with the trunk only by this single point, viz. the articulation of the clavicle with the breast-bone.

CLAVICULUS. See *Clavicle*.

CLAVICULA. See *Clavicle*.

CLAVIS. (From *claudo*, to shut.) The same as clavicle.

CLAVUS. A nail. The sensation resembling the driving a nail into the head. A fixed pain in the forehead, which may

be covered by one's thumb, giving a sensation like as if a nail were driven into the part. When connected with hysterics, it is called *clavus hystericus*. This term is also applied to *corns*, from their resemblance to the head of a nail; and to an artificial palate, or diseased uterus.

CLAVUS HYSTERICUS. See *Clavus*.

CLAVUS OCULORUM. A staphyloma, or tumour on the eyelids.

Clay. See *Alumina*.

Cleavers. See *Aparine*.

CLIDION. *Clidion*. The epithet of a pastil, described by Galen and Paulus Aegineta; and it is the name also of an epithem described by Aëtius.

CLIDOMA. (From *κλιδω*, to close.) A pastil, or torch. Also the clavicula.

CLIDOMASTOIDEUS. (From *κλιδωμα*, the clavicle, and *μαστοιδευς*, the mastoideus muscle.) See *Sterno-cleidomastoideus*.

CLISAGRA. (From *κλυς*, the clavicle, and *αγρα*, a prey.) The gout in the articulation of the clavicles.

CLITHRON. (From *κλυθω*, to shut.) See *Clastrum*.

CLEMATIS. (From *κλημα*, a tendril; so named from its climbing up trees, or any thing it can fasten upon with its tendrils.) The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Polyginia*.

CLEMATIS RECTA. The systematic name of the upright virgin's-bower. See *Flamula Jovis*.

CLEMATIS VITALBA. The systematic name of the traveller's-joy. See *Vitalba*.

CLEMATITIS. The same as *clematis*.

CLEONIS COLLYRIUM. The name of a collyrium described by Celsus.

CLEONIS GLUTEN. An astringent formula of myrrh, frankincense, and white of egg.

CLEPSIDRA. (From *κλεπτα*, to conceal, and *ιδρα*, water.) Properly, an instrument to measure time by the dropping of water through a hole, from one vessel to another; but it is used to express a chymical vessel, perforated in the same manner. It is also an instrument mentioned by Paracelsus, contrived to convey suffumigations to the uterus, in hysterical cases.

CLIBANUS. (Quasi *καλχανος*: from *κλυθω*, to conceal.) A portable furnace, or still, in which the materials to be wrought on are shut up.

CLIMACTER. (From *κλιμαζω*, to proceed gradually.) The progression of the life of man. It is usually divided into periods of seven years.

CLIMAX. (From *κλιμαζω*, to proceed.) A name of some antidotes, which, in regular proportion, increased or diminished the ingredients of which it was composed. e. g. *Chamedryos* ʒijj. *Centaurei* ʒij. *Hyperici* ʒj.

Climbing birthwort. See *Aristolochia tenuis*.

CLINICAL. (*Clinicus*; from *κλινω*, a bed.) Any thing concerning a bed: thus clinical lectures, notes, a clinical physician, &c.; which mean lectures given at the bed side, observations taken from patients when in bed, a physician who visits his patients in their bed, &c.

CLINOID. (*Clinoideus*; from *κλινω*, a bed, and *ειδος*, resemblance.) Resembling a bed. The four processes surrounding the sella turcica of the sphenoid bone are so called, of which two are anterior, and two posterior.

CLINOMASTOIDEUS. A corruption of *cleidomastoideus*.

CLISSUS. A chymical term denoting mineral compound spirits; but antimony is considered as the basis clissii. A spirit of antimony is called *clissus*.

CLITORIDIS MUSCULUS. See *Erector clitoridis*.

CLITORIS. (From *κλυω*, to enclose, or hide; because it is hid by the labia pudendorum.) *Columella.* A small glandiform body, like a penis in miniature, and, like it, covered with a præpuce, or fore-skin. It is situated above the nymphæ, and before the opening of the urinary passage of women. Anatomy has discovered, that the clitoris is composed, like the penis, of a cavernous substance, and of a glans, which has no perforation, but is, like that of the penis, exquisitely sensible. The clitoris is the principal seat of pleasure: during coition it is distended with blood, and after the venereal orgasm it becomes flaccid and falls. Instances have occurred, where the clitoris was so enlarged as to enable the female to have venereal commerce with others; and, in Paris, this fact was made a public exhibition of to the faculty. Women thus formed appear to partake, in their general form, less of the female character, and are termed hermaphrodites. The clitoris of children is larger, in proportion, than in full-grown women: it often projects beyond the external labia at birth.

CLITORISMIUS. (From *κλιτορις*.) A morbid enlargement of the clitoris.

CLONIC. (From *κλονω*, to move to and fro.) *Spasmi clonici.* See *Convulsion*.

CLONODES. (From *κλονω*, to agitate.) A strong unequal pulso.

Clove bark. See *Cassia caryophyllata*.

Clove gilliflower. See *Caryophyllum rubrum*.

Clove July flower. See *Caryophyllum rubrum*.

Clove pink. See *Caryophyllum rubrum*.

Clooc. See *Caryophyllum aromaticum*.

CLUNESIA. (From *clunes*, the buttocks.) *Proctalga.* An inflammation of the buttocks.

CLUSIA. (So called in memory of

Charles Clusius, an eminent botanist.) The name of a genus of plants in the Linnæan system. Class, *Polygamia*. Order, *Monocla*. Balsam-tree.

CLUTIA KLUTERIA. The systematic name of the tree which was supposed to afford the cascarilla bark.

CLUYTIA. (Named after Cluyt, and generally spelt *clutius*.) The name of a genus of plants in the Linnæan system. Class, *Dioecia*. Order, *Gymandria*.

CLUDON. (*Kludon*.) A fluctuation and flatulency in the stomach.

CLYPEALIS. (From *clypeus*, a shield.) Formed like a shield, hence cartilago clypealis.

CLYSMUS. *Clyma*. (From *κλύω*, to wash.) A glyster.

CLYSUS. *Clytus*. A term anciently used by the chymists for medicines made by the re-union of different principles, as oil, salt, and spirit, by long digestion; but it is not now practised, and the term is almost lost.

CLYSUS ANTIMONII. *Clysus mineralis*. A weak acid of sulphur.

CLYSTER. (*Clysterium*. From *κλύω*, to cleanse.) A glyster.

CNEMIA. Any part connected with the tibia.

CNEMODACTYLEUS. (From *κνήμη*, the tibia, and *δακτύλος*, a finger, or toe.) A muscle whose origin is in the tibia, and whose insertion is in the toes. Its office is to elevate the toes. See *Extensor digitorum longus*.

CNERIS. (From *κνῆμι*, to scratch.) *Cnema*. *Cnemosis*. A painful itching of any part.

CNICILION. (From *κνίκος*, cnicus, et *λαϊον*, oil.) Oil made of the seeds of cnicus. Its virtues are the same with those of ricinus, but in an inferior degree.

CNICUS. (From *κνῆμι*, to scratch.) The plant used by Hippocrates by this name, is supposed to be the carthamus; but modern botanists exclude it from the species of this plant.

CNICUS SYLVESTRIS. The *Carduus benedictus*.

CNIDI COCCI. See *Coccognidia*.

CNIDI GRANA. See *Coccognidia*.

CNIDOSIS. (From *κνήμη*, the nettle.) An itching sensation, such as is perceived by the nettle. A dry ophthalmia.

CNIPOTES. An itching.

CNISKOS. See *Cneris*.

CNYNA. (From *κνῆμι*, to scrape, or grate.) In Hippocrates it signifies a rasure, puncture, or vellication: also the same as *cnemosis*, or *cnesis*.

COAGULANTIA. (From *coagulo*, to incassate, or curdle.) Such medicines as coagulate the blood and juices flowing from it.

COAGULABLE LYMPH. *Lympha coagulabilis*. This substance has a great

affinity to the white of eggs. It is a component part of the serum of the human blood. It may be obtained in considerable quantities by stirring the serum about with a stick, when it adheres to its sides. In certain diseased actions it is separated from the blood, and is often found in very considerable quantities in the circumscribed cavities of the body. It has neither taste nor smell; it always possesses a white and opaque colour; is of a glutinous consistence, and, if dried by a gentle heat, becomes horny. Its presence is detected by an admixture of the diluted mineral acids. See also *Albumen*.

COAGULATION. (*Coagulatio*; From *con*, and *ago*, to drive together.) The separation of the glutinous or viscid particles, contained in any fluid, from the more thin and not coagulable particles: thus, when milk curdles, the coagulable particles form the curd; and when acids are thrown into any fluid containing coagulable particles, they form what is called a *coagulum*.

COAGULUM. A term applied frequently to blood and other fluids, when they assume a jelly-like consistency.

COAGULUM ALUMINIS. This is made by beating the white of eggs with a little alum, until it forms a coagulum. It is recommended as an efficacious application to relaxations of the conjunctive membrane of the eye.

COALTERNE FEBRES. (From *con*, and *alternus*, alternate.) Fevers mentioned by Belini, which he describes as two fevers affecting the same patient, and the paroxysm of one approaching as that of the other subsides.

COARCTATIO. (From *coarcto*, to straiten.) The contraction, or diminution of any thing. Applied to the pulse, it means a lessening in number.

COARTICULATION. (From *con*, and *articulatio*, an articulation.) That sort of articulation which has manifest motion.

COBALT. *Cadmia metallica*. A metal that has never been found pure in nature. We meet with it almost always either in the state of an oxyd; alloyed with other metals in the form of a sulphuret; or combined with an acid.

Cobalt in the state of an oxyd forms the *black cobalt ore*. This ore is found in Germany, either in powder of a black or grey colour, or in compact masses. In the last form, it exhibits at its fracture, rose coloured spots. There are several varieties of this ore.

Cobalt, alloyed with other metals, forms the *dull white cobalt ore*. In this ore, which occurs either amorphous or crystallized, cobalt is united to iron and arsenic. The colour of this ore, when fresh broken, is white or blueish-grey, sometimes with a shade of red. It has a metallic lustre. Its texture is compact. Cobalt, united to

sulphur, forms the *white Cobalt ore*. It is met with in masses, or crystallized in cubes, dodecahedra, and octahedra. Its colour is a tin-white, sometimes reddish-yellow. Cobalt combined with arsenic acid forms the *red cobalt ore*, arseniate of cobalt. It is found in masses of various shapes. Its colour is red, inclining to orange.

Cobalt, when in a pure state, is of a steel-grey colour, with a tinge of red, and a fine close grain. It has a granulated fracture, and is easily broken and pulverized. Its specific weight is between 7.700 and 7.811. It requires a very intense heat for its fusion, nearly equal to that necessary to melt cast-iron. When heated in contact with the air, it oxydates before fusion. Phosphorus renders it very fusible, and converts it into a phosphuret. It unites to sulphur with difficulty, but very well with the alkali sulphurets by fusion. When alloyed with metals it renders them granulated, rigid, and brittle. It is attacked by the greater number of the acids, and unites with the boracic acid. Its solutions in different acids become green when heated; and from this property, it is used as an ink, which, when written with on paper, is invisible, but becomes visible when gently heated, and disappears when cold. It takes fire in oxygenated muriatic acid gas. It colours glass of a fine blue. It unites with platina, gold, iron, nickel, copper and arsenic, by fusion; but silver, lead, bismuth, and mercury, refuse to unite with it in the dry way. In its purest state, it is not only obedient to the magnet, but, if we may trust to the accuracy of some experiments made by Kohl and Wenzel, it may even receive a magnetical attractive power.

Nitrate of potash oxydates cobalt readily. It detonates by the blow of a hammer, when mixed with oxygenated muriate of potash. It produces fine colours in porcelain, enamels, artificial gems, &c.

COBHAM WATERS. Weak saline purging waters at Cobham.

COBRA DE CAPELO. (From *cobra*, the head, or covering. Span.) *Crotalus horridus* of Linnaeus. The rattle-snake; the stone out of whose head is said to be an antidote to the poison of venomous animals.

COCCA CNIDIA. See *Grana cnidia*.

COCCARIUM. (From *coccos*, a berry.) A very small pill.

COCCINELLA. (Dim. from *coccus*, a berry; from its resemblance to a berry.) *Coccinella*. *Ficus India grana*. *Scarabeolus hemisphericus*. *Cochinifera cochinnilla*. *Coccus Americanus*. *Cochinelle*. *Coccus Indicus tinctorius*. Cochineal. The female of a species of insect called *Coccus cacti*, that is found on, and collected in South America, from the *Opuntia*, or Indian fig-

tree. It possesses stimulating qualities, and is ordered by the College in the *tinctura cardamomi composita* and *cinchone composita*; but most probably, on account of the beautiful red colour which it imparts to them.

COCCO-BALSAMUM. The fruit of the true balsam.

COCCONIDIA. *Grana cnidia*. *Cocci cnidii*. The seeds of the *Daphne mezereum* are so termed. They are violently purgative. See *Mezereum*.

COCCOS. See *Grana cnidia*.

COCCULI INDI AROMATICI. The pipet Jamaicense.

COCCULUS INDUS. (Dim. of *anusac*, a berry.) *Coccus Indicus*. *Coccule officinarum*. *Cocci Orientales*. The berry so called is rugous and kidney-shaped, and contains a white nucleus; it is the produce of the *Menispermum cocculus*; *foliis cordatis, retusis mucronatis*; *caule lacero*, of Linnaeus. It is brought from Malabar and the East Indies. It is poisonous if swallowed, bringing on a nausea, fainting and convulsion. Mixed with paste it stupifies fishes, so that they will lie on the water, and not attempt to escape from the hand that takes them. The berries possess an inebriating quality; and are supposed to impart that power to most of the London porter.

COCCUM BAPHICUM. A name for cheremes.

COCCUS. The name, in entomology, of a tribe of insects.

COCCUS CACTI. The systematic name of the cochineal animal. See *Coccinella*.

COCCYGEUS. (*Coccygeus*, sc. *musculus*; from *κωκυξ*: because it is inserted into the *coccyx*.) *Ischio-coccygien* of Dumas. A muscle of the os coccygis, situated within the pelvis. It arises tendinous and fleshy, from the spinous process of the ischium, and covers the inside of the sacro-ischiatic ligament: from this narrow beginning it gradually increases to form a thin fleshy belly, interspersed with tendinous fibres. It is inserted into the extremity of the os sacrum, and near the whole length of the os coccygis, laterally. Its use is to support and move the os coccygis forwards, and to tie it more firmly to the sacrum.

COCCYGIS OS. (From *κωκυξ*, the cuckoo, whose bill it is said to represent.) *Cauda*. *Ossis sacri acumen*. *Coccyx*. This bone is a small appendage to the point of the sacrum, terminating this inverted column with an acute point, and found in very different conditions in the several stages of life. In the child, it is merely cartilage, and we can find no point of bone; during youth, it is ossifying into distinct bones, which continue moveable upon each other till manhood; then the separate bones gradually unite with each other, so as to form one conical bone,

with bulgings and marks of the pieces of which it was originally composed; but still the last bone continues to move upon the joint of the sacrum, till, in advanced years, it is at last firmly united; later in women than in men, with whom it is often fixed at twenty or twenty-five. It is not, like the os sacrum, flat, but of a roundish form, convex without, and concave inwards; forming with the sacrum the lowest part of the pelvis behind. It has no holes like the sacrum; has no communication with the spinal canal, and transmits no nerves; but points forwards to support the lower parts of the rectum; thus it contracts the lower opening of the pelvis, so as to support effectually the rectum, bladder, and womb; and yet continues so moveable in women, as to recede in time of labour, allowing the head of the child to pass.

COCYX. (Κουκῦξ, the cuckoo.) See *Coccygis os*. Also the part in which the os coccygis is placed.

COCHIA. (From κοχια, to turn or make round.) An ancient name of some officinal pills.

Cochineal. See *Goccinella*.

COCHLEA. (From κοχλια, to turn round.) A cavity of the internal ear, resembling the shell of a snail, in which are observed, the *modiolus*, or *nucleus*, extending from its basis to the apex, the *scala tympani*, *scala vestibuli*, and *apical lamina*.

COCHLEA TERRESTRIS. See *Limax*.

COCHLEARE. (From *cochlea*, a cockle, whose shell its bowl represents.) A spoonful. In prescriptions it is sometimes abbreviated thus, *coch*. *Cochleare magnum*, is a table-spoon; *cochleare medium*, a dessert or pap-spoon; and *cochleare minimum*, a tea-spoon.

COCHLEARIA. (From *cochleare*, a spoon; so called from its resemblance.) The name of a genus of plants in the Linnean system. Class, *Tetradynamia*. Order, *Sticulea*.

COCHLEARIA ARMORACIA. The systematic name of the horse-radish. See *Raphanus rusticanus*.

COCHLEARIA HORTENSIS. Lemon scurvy-grass. This indigenous plant, *Cochlearia officinalis* of Linnæus:—*foliis radicalibus cordato-subrotundis; caulibus oblongis subsinuatis*, is cultivated in gardens for its medicinal qualities. Its expressed juice has been long considered as the most effectual of the scorbutic plants.

COCHLEARIA OFFICINALIS. The systematic name of the lemon-scurvy-grass. See *Cochlearia hortensis*.

COCHONE. (From κοχον, to turn round.) Galen explains this to be the juncture of the ischium, near the seat, or breech; whence, says he, all the adjacent parts about the seat are called by the same name. Hesiychius says, that *cochone* is the

part of the spine which is adjacent to the os sacrum.

COCOS. (So called from the Portuguese *coco*, or *coquen*, the three holes at the end of the cocoa-nut shell, giving it the resemblance of a monkey's head.) The name of a genus of plants in the Linnean system. Class, *Monocla*. Order, *Hexandria*. The cocoa-nut tree.

Cocos nycifera. The systematic name of the plant whose fruit is the cocoa-nut.

Cocos butyracea. The systematic name of the plant which affords the palm oil.

COCTION. (From *coquo*, to boil.) Concoction. Digestion. In a medical sense, signifies that alteration, whatever it be, or however occasioned, which is made in the crude matter of a distemper, whereby it is either fitted for a discharge, or rendered harmless to the body. This is often brought about by nature; that is, by the vis vitæ, or the disposition or natural tendency of the matter itself, or else by proper remedies, which may so alter its bulk, figure, cohesion, or give it a particular determination, so as to prevent any farther ill effects, or drive it quite out of the body. And, that time of a disease wherein this action is performing, is called its state of coction.

COCUSTU. The name for courbaril.

CODOCLE. (From *codon*, a bulb, and *ocle*, a tumour.) A bubo.

CONOGA PALA. See *Goness cortex*.

CŒCALIS. (From *cæcum*, the blind gut, through which it runs.) A vein, being a branch from the concave side of the vena mesaraica.

CŒLA. (From *καλος*, hollow.) The hollow pits above, and sometimes below the eyes. The hollow parts at the bottom of the feet.

CŒLIA. (From *καλος*, hollow.) A cavity in any part of the body. The belly. The womb.

CELIAC ARTERY. (*Cæliacus*, belonging to the belly; from *καλος*, the belly.) *Arteria cæliaca*. The first branch given off from the aorta in the cavity of the abdomen. It sends branches to the diaphragm, stomach, liver, pylorus, duodenum, omentum, and spleen.

CELIAC PASSION. (*Cæliacus*, belonging to the belly; from *καλος*, the belly.) *Cælica Chylosa*. *Cælica lactea*. There are very great differences among physicians, concerning the nature of this disease. Sauvages says it is a chronic flux, in which the aliment is discharged half digested. Dr. Cullen considers it as a species of diarrhœa, and mentions it in his third and fourth species, under the terms *mucosa*, *chylosa*, *lactea*; making the purulenta only symptomatic. See *Diarrhœa*. It is at-

tended with great pains in the stomach, resembling the pricking of pins; rumbling and flatulency in the intestines; white stools, because deprived of bile; while the patient becomes weak and lean.

CŒLOMA. (From *καλος*, hollow.) An ulcer in the tunica cornea of the eye.

CŒLOSTOMIA. (From *καλος*, hollow, and *σoma*, the mouth.) A defect in speaking, when a person's speech is obscured by sounding as if his voice proceeded from a cavern.

CŒNOLOGIA. (From *κοινος*, common, and *λογος*, discourse.) A consultation, or common consideration of a disease, by two or more physicians.

CŒNOTAS. (From *κοινος*, common.) The physicians of the methodic sect asserted that all diseases arose from relaxation, stricture, or a mixture of both. These were called *cœnotas*, viz. what diseases have in common.

CŒRULEUS LAPIS. The sulphate of copper. See *Sulphas cupri*.

CŒTE. (From *κυμα*, to lie down.) A bed, or couch, for a sick person.

COFFĒA. (The true name is *Caffē*, from *Caffa*, the province in South America, where the tree grows spontaneously in great abundance.) The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Mongynia*. The coffee-tree.

COFFĒE. (From *kofuah*, a mixing together, Hebr.; so called from the pleasant position which is made from its berry.) *Jasminum Arabicum*. *Choava*. The seeds of the *Coffea Arabica*; *floribus quinquefidis, dispermis* of Linnæus. The coffee-tree is cultivated in Arabia, Persia, the East Indies, the Isle of Bourbon, and several parts of America. Good Turkey Coffee is by far the most salutary of all liquors drunk at meal-time. It possesses nerve and adstringent qualities, and may be drunk with advantage at all times, except when there is bile in the stomach. It is said to be a good antidote against an over-dose of opium, and to relieve obstinate spasmodic asthma. For the latter purpose, the coffee ought to be of the best Mocha, newly burnt, and made very strong, immediately after grinding it. Sir John Pringle commonly ordered one ounce for a dose; which is to be repeated fresh, after the interval of a quarter or half an hour; and which he directed to be taken without milk or sugar.

If coffee be drunk warm within an hour after dinner, it is of singular use to those who have head-ach, from weakness in the stomach, contracted by sedentary habits, close attention, or accidental drunkenness. It is of service when the digestion is weak; and persons afflicted with the sick head-ach are much benefited by its use, in

some instances, though this effect is by no means uniform. Coffee is often imitated, by roasting rye with a few almonds.

COFFĒE ARABICA. The plant which affords coffee. See *Coffee*.

COHESION. (From *con*, and *hæreo*, to stick together.) *Vis cohesionis*. *Vis adhesionis*. *Vis attractionis*. That force in the particles of matter whereby they are connected in such a way that they resist any attempt towards their removal or separation. It is a species of attraction. See *Attraction*.

COHOBATION. (A term invented by Paracelsus.) *Cohobatio*. *Cohobium*. *Cohoph*. The ancient chymists use this term to signify the distillation of a fluid poured afresh upon a substance of the same kind as that upon which it was before distilled, and repeating this operation several times, to make it more efficacious. For this purpose, the vessel called pelican was employed.

COHOL. (*Cohol*. Heb.) Castellus says this word is used in Avicenna, to express dry collyria for the eyes, in fine powder.

COLICA. (From *κολλη*, the bowels.) A sudden swelling of the belly from wind.

COLLOSTOMIA. (From *καλος*, hollow, and *σoma*, the mouth.) A defect of speaking, from the palate, or through the nose.

COINDICANTIA. (From *con*, and *indico*, to indicate.) Signs, or symptoms, are called coincident, when, besides the usual incidental appearances, there occur others, as age, habit, season, &c.

COIRA. A name for the terra Japonica.

COITUS. (From *coeo*, to go together.) The conjunction of the male and female in the act of procreation.

COLA. (From *καλος*, a joint.) The joints.

COLATORIA LACTEA. Astruc says they were formerly called glands, and are situated in the third and internal tunic of the uterus, and, that they are vesiculo-vascular bodies.

COLATORIUM. (From *colo*, to strain.) A strainer of any kind.

COLATURE. (From *colo*, to strain.) A filtered or strained liquor.

COLCAQUAHUITL. An American plant, commended in palsies and uterine disorders, according to Ray.

COLCESTRENSIS AQUA. Colchester water. This mineral water is of the bitter purging kind, similar to that of Epsom, but not so strong.

COLCHICUM. (From *Colchis*, a city of Armenia; where this plant is supposed to have been common.) 1. The name of a genus of plants in the Linnæan system. Class, *Hexandria*. Order, *Trigynia*. Meadow-saffron.

2. The pharmacopœial name of the common meadow-saffron. *Colchicum autumnale* of Linnæus:—*foliis planis lan-*

colicis arvensis. A native of England. The sensible qualities of the fresh root are very various, according to the place of growth and season of the year. In autumn it is almost inert, but in the beginning of summer highly acrid; hence some have found it to be a corrosive poison, whilst others have eaten it in considerable quantity, without experiencing any effect. When it is possessed of acrimony, this is of the same nature with that of garlic and some other plants, and it is entirely destroyed by drying. The German physicians have celebrated its virtues as diuretic, in hydrothorax and other dropsies; and in France it continues to be a favourite remedy; but it is, nevertheless, in this country unsuccessful, and at best a very uncertain remedy. The expressed juice is used, in Alsace, to destroy vermin in the heads of children. The officinal preparations of colchicum are, Syr. Colch. Autum. Edin. Pharm. The oxymel colchici of the former London pharmacopœia is now omitted, and the acetum colchici ordered in its room; the College observing that the honey may easily be added extemporaneously, if it be thought requisite. The following is the formula of the present London Pharm. for preparing the acetum colchici, or vinegar of meadow-saffron: Take of meadow-saffron root sliced, an ounce; acetic acid, a pint; proof spirit, a fluid-ounce. Macerate the meadow-saffron root in the vinegar, in a covered glass vessel, for twenty-four hours; then press out the liquor and set it by, that the feculencies may subside; lastly, add the spirit to the clear liquor. The dose is from ʒss to ʒjss.

COLCHICUM AUTUMNALE. The systematic name of the common meadow-saffron. See *Colchicum*.

COLCHICUM ILLYRICUM. The plant supposed to afford the *hermodactylus*. See *Hermodactylus*.

COLCHICUM ZEGLANICUM. See *Zedaria*.

COLOTHAR VITRIOLI. *Chalcitis*. The remains of calcined martial vitriol.

COLD. A privation of heat. It is nothing positive, but somewhat of the negative kind. The human body contains within itself, as long as it is living, a principle of warmth: if any other body, being in contact with it, does at the same time impart to it more caloric or heat than it obtains from the human body, it is said to be warm; but if it receives from the human body more heat than it remits, it is said to be cold.

A cold is a popular name also for a catarrhus affection of the nostrils, throat, and fauces. See *Catarrhus*.

COLD AFFUSION. A process formerly practised by physicians, but lately introduced by Dr. Currie, of Liverpool,

in the treatment of typhus fever, and which appears to possess a uniformity of success, which we look for in vain in almost any other branch of medical practice. The remedy consists merely in placing the patient in a bathing-tub, or other convenient vessel, and pouring a pailful of cold water upon his body; after which he is wiped dry, and again put to bed. It should be noted,

First, That it is the *low contagious fever* in which the cold affusion is to be employed. The first symptoms of which are a dull head-ach, with restlessness and shivering; pains in the back, and all over the body, the tongue foul, with great prostration of strength; the head-ach becoming more acute, the heat of the body, by the thermometer, 102° to 105° or more; general restlessness, increasing to delirium, particularly in the night.

Secondly, That it is in the *early stage* of the disease we must employ the remedy; and generally in the *state of the greatest heat and exacerbation*.

Thirdly, It is *affusion, not immersion*, that must be employed.

Since the first publication of Dr. Currie's work, the practice of affusion has been extended throughout England; and its efficacy has been established in some stages of the disease, from which the author had originally proscribed the practice of it. One of the cautionary injunctions which had been given for the affusion of cold water in fever was, *never to employ it in cases where the patient had a sense of chilliness upon him*, even if the thermometer, applied to the trunk of the body, indicated a preternatural degree of heat. In his last edition of Reports, however, Dr. Currie has given the particulars of a case of this description, in which the cold affusion was so managed as to produce a successful event.

In fevers arising from, or accompanied by, *topical inflammation*, his experience does not justify the use of cold affusion; though in a great variety of these cases, the warm affusion may be used with advantage. "And," says he, "though I have used the cold affusion in some instances, so late as the twelfth or fourteenth day of contagious fever, with safety and success, yet it can only be employed, at this advanced period, in the instances in which the heat keeps up steadily above the natural standard, and the respiration continues free. In such cases I have seen it appease agitation and restlessness, dissipate delirium, and, as it were, snatch the patient from impending dissolution. But it is in the *early stages* of fever (let me again repeat) that it ought always to be employed, if possible; and where, without any regard to the heat of the patient, it is had recourse to in the last stage of fever, after every other remedy has failed, and the case appears desperate, (of

which I have heard several instances,) can it appear surprising that the issue should sometimes be unfavourable?"

Numerous communications from various practitioners, in the West and East Indies, in Egypt and America, also shew the efficacy of affusion in the raging fevers of hot countries.

COLIC. (From *καλός*, a stalk.) *Colis*. The penis.

Colewort. See *Brassica*.

COLI DEXTRUM LIGAMENTUM. Where the mesentery changes its name for that of mesocolon, (near the extremity of the ileum,) the particular l. mina, which is turned to the right side, forms a small transverse fold which is thus named.

COLI SINISTRUM LIGAMENTUM. It is a contraction of the mesocolon, a little below the left kidney.

COLICA. (From *καλόν*, colon, the name of one of the intestines.) The colic. The appellation of colic is commonly given to all pains in the abdomen, almost indiscriminately; but, from the different causes and circumstances of this disorder, it is differently denominated. When the pain is accompanied with a vomiting of bile, or with obstinate costiveness, it is called a *bilious colic*: if flatus causes the pain, that is, if attended with temporary distention, relieved by the discharge of wind, it takes the name of *flatulent* or *windy colic*; when accompanied with heat and inflammation, it takes the name of *inflammatory colic*, or *enteritis*. When this disease arises to a violent height, and is attended with a stercoraceous vomiting, obstinate costiveness, and an evacuation of fæces by the mouth, it is called *pseudo iliaca*, or *iliac passion*.

Dr. Cullen places this genus of disease in the class *neuroses*, and order *spasmi*; and defines it pain of the abdomen, particularly round the umbilicus, attended with vomiting and costiveness. He enumerates seven species.

1. *Colica spasmodica*, with retraction of the navel, and spasm of the muscles of the belly.

2. *Colica pictorum*. This is called from the place where it is endemial; the Pictou, the Surinam, the Devonshire colic; from its victims, the plumbers' and the painters' colic; from its symptoms, the dry belly-ach, the nervous and spasmodic colic. It has been attributed to the poison of lead, and this is undoubtedly the cause, when it occurs to glaziers, painters, and those employed in lead works; but, though this is one, it is by no means the only cause. In Devonshire it certainly more often arises from the early cider, made of harsh, unripe fruit, and in the West Indies from new rum. The characteristics of this disease are—obstinate costiveness, with a vomiting of an acrid or porraceous bile, pains about

the region of the navel, shooting from thence to each side with excessive violence, strong convulsive spasms in the intestines, and a tendency to a paralysis of the extremities. It is occasioned by a long continued costiveness; by an accumulation of acrid bile; by cold, applied either to the extremities or to the belly itself; by a free use of unripe fruits, and by great irregularity in the mode of living. From its occurring frequently in Devonshire, and other cider counties, it has been supposed to arise from an impregnation of lead, received into the stomach; but this seems to be a mistake, as it is a very prevalent disease in the West Indies likewise, where no cider is made, and where there is only a very small quantity of lead in the mills employed to extract the juice from the sugar-canes. One or other of the causes just enumerated may justly be said always to give rise to this species of colic.

The disease comes on gradually, with a pain at the pit of the stomach, extending downwards to the intestines, accompanied with eructations, slight sickness at the stomach, thirst, anxiety, obstinate costiveness, and a quick contracted pulse. After a short time, the pains increase considerably in violence; the whole region of the belly is highly painful to the touch; the muscles of the abdomen are contracted into hard irregular knots, or lumps; the intestines themselves exhibit symptoms of violent spasm, insomuch that a glyster can hardly be injected, from the powerful contraction of the sphincter ani; and there is constant restlessness, with a frequent vomiting of an acrid or porraceous matter, but more particularly after taking either food, or medicine.

Upon a farther increase of the symptoms, or their not being quickly alleviated, the spasms become more frequent, as well as violent; the costiveness proves invincible, and an inflammation of the intestines ensues, which soon destroys the patient by gangrene. In an advanced stage of the disease, it is no uncommon occurrence for dysuria to take place, in a very high degree.

The dry belly-ach is always attended with some degree of danger; but which is ever in proportion to the violence of the symptoms, and the duration of the disease. Even when it does not prove fatal, it is too apt to terminate in palsy, and to leave behind it contractions of the hands and feet, with an inability in their muscles to perform their office; and in this miserable state of existence the patient lingers out many wretched years.

Dissections of this disease usually shew the same morbid appearances as in common colic, only in a much higher degree.

3. *Colica stercorea*, which happens

from obstinate and long-continued costiveness.

4. *Colica accidentalis*, called also *cholera sicca*, from acrid undigested matters.

5. *Colica meconialis*, in infants, from a retention of meconium.

6. *Colica callosa*, from a sensation of a stricture in some part of the colon, and frequently of previous flatulence, gradually passing off; the habit costive, or faeces liquid, and in small quantity.

7. *Colica calculosa*, from calculi formed in the intestines, attended with a fixed hardness in some part of the abdomen. It is distinguished by the previous discharge of calculi.

8. *Colica flatulenta* may be added to these species. It is distinguished by a sudden fullness, with pain and constipation, relieved by a discharge of wind from the mouth, or anus.

The colic is distinguished from inflammation of the intestines by the pain being *wringing*, and not of a burning sensation; by the *spasmodic contraction* of the abdominal muscles; by the *absence* or *trifling* degree of fever; by the *state* of the pulse, and by the *diminution* of pain upon pressure.

The flatulent and inflammatory colic are thus distinguished from each other:—In the flatulent colic, the pain comes on by fits, flies from one part of the bowels to another, and is much abated by a discharge of wind, either upwards or downwards; but in the inflammatory colic the pain remains equable, and fixed and settled in one spot; the vomitings are severe, and frequently bilious; the belly is obstinately bound, and the pulse quick and feverish.

The colic should be distinguished from a fit of the gravel; stones passing through the ureters; rheumatic pains in the muscles of the belly; a beginning dysentery; the blind piles; and from a stone passing through the gall-duct. Gravel in the kidney produces often colic pains, not easily distinguishable; but when stones pass through the ureters, the testicle on that side is often retracted, the leg is benumbed, a pain shoots down the inside of the thigh; symptoms occasioned by the stone passing through the ureter over the spermatic chord, or the sacro-sciatic nerve. Rheumatic pains in the muscles of the belly rarely affect so accurately the umbilical region, but dart in various directions, to the chest, or to the pelvis, and are attended with soreness, not confined to the abdomen. A beginning dysentery differs little from colic. The pain from the blind piles is confined to the rectum: and that from a stone in the gall-duct is felt in the pit of the stomach, occasionally shooting through the body to the back.

COLICA ACCIDENTALIS. Colic from cruelties in the bowels.

COLICA ARTERIA SINISTRA. The lower mesenteric artery.

COLICA ARTERIA SUPERIOR. The upper mesenteric artery.

COLICA BILIOSA. Colic from excess of bile.

COLICA CALCULOSA. Colic from stony matters in the intestines.

COLICA CALLOSA. Colic from hardened and obstinate strictures.

COLICA DAMNONIORUM. Colic peculiar to Devonshire. See *Colica*.

COLICA FEBRICOSA. Colic with fever.

COLICA FLATULENTA. Colic from wind.

COLICA GRAVIDARUM. Colic in pregnant women.

COLICA HYSTERICA. Hysteric colic.

COLICA LACTANTIUM. Colic peculiar to nurses.

COLICA LAPTONICA. Colic peculiar to Laplanders.

COLICA MECONIALIS. Colic from meconium in infants.

COLICA MESENTERICA. Colic from diseased mesentery.

COLICA NERVOSA. The nervous colic.

COLICA PANCREATICA. Colic from diseased pancreas.

COLICA PHLOGISTICA. Colic with inflammation.

COLICA PICTONUM. See *Colica*.

COLICA PITUITOSA. The spasmodic colic.

COLICA PLETHORICA. The inflammatory colic.

COLICA PLUMBARIORUM. The colic of lead-workers.

COLICA PULSATILIS. The inflammatory colic.

COLICA SATURNINA. The Devonshire colic. See *Colica*.

COLICA SCIRRHOSEA. The colic from scirrhus tumours.

COLICA SPASMODICA. The spasmodic colic.

COLICA STERCOREA. Colic from retained faeces.

COLICA VENA. A branch of the upper mesenteric artery.

COLICA VENA RECTA. The vein of the colon.

COLICA VERMINOSA. The colic from worms.

COLICE. The colic.

COLIFORMIS. (From *cola*, a strainer, and *forma*, a likeness; so called from its having many perforations, like a strainer.) *Coliforme os*. A name formerly given to the os cribriform.

COLIPHUM. (From *κωλον*, a limb, and *φωμ*, strongly.) A kind of bread given to wrestlers. It was made of flour and bran together, and was thought to make men athletic.

COLIS. The same as coles.

COLLAPsus. (From *collabor*, to shrink

down.) A wasting or shrinking of the body, or strength.

COLLATENNA. A specific vulnerary.

COLLATERALES. So Spigelius calls the erecitors penis, from their collateral order of fibres.

COLLETICA. (From *κolla*, glue.) Conglutinating medicines.

COLLICIE. (From *colligo*, to collect.) The union of the ducts, which convey the humours of the eyes from the puncta lachrymalia to the cavity of the nose.

COLLICULUM. (Dim. of *collis*, a hill.) The nymph, or prominence, without the vagina of women.

COLLIGAMEN. (From *colligo*, to tie together.) A ligament.

COLLIQUAMENTUM. (From *colliquo*, to melt.) A term first made use of by Dr. Harvey, in his application of it to the first rudiments of an embryo in generation.

COLLIQUATIVE. (From *colliquo*, to melt.) Any excessive evacuation is so called which melts down, as it were, the strength of the body: hence colliquative perspiration; colliquative diarrhæa, &c.

COLLISIO. (From *collido*, to beat together.) A contusion.

COLLIX. (From *κολον*, food.) A troch, or lozenge.

COLLOBOMA. (From *κολλαω*, to glue together.) The growing together of the eyelids.

COLLODES. (From *κολλα*, glue.) Glutinous.

COLLUM. (From *καλόν*, a member, as being one of the chief; or dim. of *columna*, as being the pillar and support of the head.) The neck.

COLLUTORIUM. (From *colluo*, to wash.) A gargarism or wash for the mouth.

COLLUVIES. (From *colluo*, to cleanse.) Filth. Excrement. The discharge from an old ulcer.

COLLYRI. (*Κολλυρι*, a little round cake; so called from its likeness to a cake.) A bump, or knob, which rises after a blow.

COLLYRIUM. (From *καλυω*, to check, and *ρυς*, a defluxion; because it stops the defluxion.) Any medicine was formerly so called which was applied with that intention. The term is now only given to fluid applications for the eyes, or eye-waters.

COLOBOMA. (From *κολλαω*, to glue together.) The growing together of the eyelids; also the want of a particular member of the body.

COLOBOMATA. In Celsus this word is expressed by *curta*. Both the words signify a deficiency in some part of the body, particularly the ears, lips, or alæ of the nostrils.

COLOCASIA. (From *κολος*, food, and

κατα, to adorn; so called from its use as a food, and the custom of wearing its flowers in wreaths.) The faba *Ægyptia*.

COLOCYNTHIS. (From *καλος*, the colon, and *κινω*, to move; because of its great purging powers.) *Alhandula* of the Arabians. *Colocynthis medulla*. Colocynthida. Bitter apple. Bitter gourd. Bitter cucumber. The fruit which is the medicinal part of the *Cucumis colocynthis*; *foliis multifidis, pomis globosis glabris*, is imported from Turkey. Its spongy membranous medulla, or pith, is directed for use; it has a nauseous, acrid, and intensely bitter taste; and is a powerful irritating cathartic. In doses of ten and twelve grains, it operates with great vehemence, frequently producing violent gripes, bloody stools, and disordering the whole system. It is recommended in various complaints, as worms, mania, dropsy, epilepsy, &c.; but is seldom resorted to, except where other more mild remedies have been used without success, and then only in the form of the *extractum colocynthis compositum*, and the *pilule ex colocynthide cum aloë* of the pharmacopœias.

COLOMBO. See *Columbo*.

COLON. (*Κολον*, quasi *καλον*: from *καλος*, hollow; so called from its capacity, or from its generally being found empty, and full of wind, in dissection.) The ascending portion of the large intestine is so called. It proceeds towards the liver, by the name of the *ascending portion of the colon*; and having reached the liver, forms a *transverse arch* across to the other side. The colon then descends, forming what is termed its *sigmoid flexure*, into the pelvis, where the gut is called rectum. See *Intestines*.

COLOPHONIA. (*Κολοφωνα*, the city from whence it was first brought) *Resina nigra*. The black resin which remains in the retort, after distilling the common resin with a strong fire. Paracelsus seems to mean by it what is now prescribed by the name of *terebinthina cocta*: but the ancients, and particularly Galen, seemed to understand by it a soft kind of mastich, from *Chio*, probably the same as our *Chio turpentine*.

COLOSTRUM. (From *κολος*, food, or *κολλωμαι*, to agglutinate; so called, either because it is the first food of the young, or from its being at that time peculiarly glutinous.) Is the first milk in the breasts after delivery, according to some authors; but Bartholme applies it to an emulsion made by the solution of turpentine with the yolk of an egg.

COLOTOMES. (From *καλωτης*, a lizard, and *μοις*, likeness.) Variegated like the skin of a lizard. Hippocrates applied it to the excrements.

COLOQUINTIDA. See *Colocynthis*.

COLOCELE. (From *καλος*, the vagina,

and *κωλη*, a tumour.) A hernia forced into the vagina.

COLPOPTOSIS. (From *κολπος*, the vagina, and *πτωσις*, to fall down.) A bearing down of the vagina.

Colt's foot. See *Tussilago*.

COLUMBER HERUS. (*Quod colit umbram*, because it delights in shade.) The systematic name of a viper. See *Vipera*.

COLUMBRINA VIRGINEANA. See *Serpentaria*.

COLUMBRINUM LIGNUM. (*Colubrinus*; from *coluber*; so called from the snake-like contortions of its roots.) This species of snake-wood is brought from America. It is solid, ponderous, acrid, extremely bitter, and inodorous; its bark is of a ferruginous colour, covered with cineritious spots.

Columbine. See *Aquilegia*.

COLUMBA. See *Columbo*.

COLUMBO. *Colombo. Calombo. Calombo.* The root formerly so called is now termed *Calumba* in the London pharmacopœia. It is imported from *Colomba*, in Ceylon, in circular, brown knobs, wrinkled on the outer surface, yellowish within, and consisting of cortical, woody, and medullary lamina. Its smell is aromatic; its taste pungent, and nauseously bitter. From Dr. Percival's experiments on the root, it appears that rectified spirit of wine extracts its virtues in the greatest perfection. The watery infusion is more perishable than that of other bitters. An ounce of the powdered root, half an ounce of orange-peel, two ounces of brandy, and fourteen ounces of water, macerated 12 hours without heat, and then filtered through paper, afford a sufficiently strong and tolerably pleasant infusion. The extract made first by spirit and then with water, and reduced by evaporation to a pilular consistence, is found to be equal, if not superior, in efficacy to the powder. As an antiseptic, *Calumba* root is inferior to the bark; but as a corrector of putrid bile, it is much superior to the bark; whence also it is probable that it would be of service in the West-India yellow fever. It also restrains alimentary fermentation, without impairing digestion; in which property it resembles mustard. It does not appear to have the least heating quality, and therefore may be used in phthisis pulmonalis, and in hectic cases, to strengthen digestion. It occasions no disturbance, and agrees very well with a milk diet, as it abates flatulence, and is indisposed to acidity. The London, Edinburgh, and Dublin Colleges, direct a tincture of *Calumba* root. The dose of the powdered root is half a drachm, which, in urgent cases, may be repeated every third or fourth hour.

COLUMBORE. See *Columbo*.

COLUMELLA. (Dim. of *columna*, a column.) See *Uvula*, and *Clitoria*.

COLUMBIUM. Mr. Hatchett describes the ore from which this metal is obtained, as being of a dark brownish grey externally, and more inclining to an iron grey internally; the longitudinal fracture he found lamellated; and the cross fracture had a fine grain. Its lustre was vitreous, slightly inclining in some parts to metallic; moderately hard and very brittle. The colour of the streak, or powder, was dark chocolate brown. The particles were not obedient to the magnet. Its specific gravity, at a temperature of 65° Fahr. Mr. Hatchett found to be 5918.

A series of accurate experiments made by its discoverer, prove that this ore consists of iron, combined with a new metallic acid, which constitutes more than three-fourths of the whole.

The smallness of the quantity Mr. Hatchett had to operate upon, has hitherto prevented us from seeing the metal in its metallic state; but the accuracy with which the properties of its acid have been investigated, leave no doubt of its being different from any of the acidifiable metals hitherto known.

COLUMNELLARIS. (From *columella*, a little column.) A name of the *dens caninus*.

COLUMNA. A column, or pillar. Many parts of the body, which in their shape or office resemble columns, are so named; as *columnæ carnæ*, &c.

COLUMNA NASI. The lowest and fleshy part of the nose, which forms a part of the septum.

COLUMNA ORIS. A name for the uvula.

COLUMNÆ CARNÆ. *Columnæ cordis.* See *Carnæ columnæ*.

COLUMPIUM. (*Πασι το κώλων τὸν ἐνν*; because it prevents a defluxion.) A tent to thrust into a sore, to prevent a defluxion of humours.

COMA. (From *καμν*, or *καμν*, to lie down.) A propensity to sleep. This word anciently meant any total suppression of the powers of sense; but now it means a lethargic drowsiness. The *coma vigil* is a disease where the patients are continually inclined to sleep, but cannot.

COMA SOMNOLENTUM. Is when the patient continues in a profound sleep; and, when awakened, immediately relapses, without being able to keep open his eyes.

COMATA. (*Κοματα*; from *coma*) A diminution of the powers of voluntary motion, with sleep, or the senses impaired. It is an order of the class *neuroses* of Cullen's Nosology.

COMATOSE. Having a strong propensity to sleep.

COMBUSTIO. (From *comburo*, to burn.) A burn, or scald.

COMIPER. See *Cubæ*.

COMBUSTION. (From *comburo*, to burn.) Burning. Among the various operations.

tions of chymistry, none acts a more conspicuous part than combustion; and in proportion to its utility in the science, the necessity of thoroughly investigating its nature and mode of action becomes more obvious to the philosophical chymist.

Lavoisier's Theory of Combustion.

Lavoisier's theory of combustion is founded upon the absorption of oxygen by a combustible body.

Taking this for granted, it follows that combustion is only the play of affinity between oxygen, the matter of heat, and a combustible body.

When an *incombustible* body (a brick for instance) is heated, it undergoes no change, except an augmentation of bulk and temperature: and when left to itself, it soon regains its former state. But when a *combustible* body is heated to a certain degree, in the open air, it begins to become on a sudden intensely hot, and at last emits a copious stream of caloric and light to the surrounding bodies. During this emission, the burning body gradually wastes away. It either disappears entirely, or its physical properties become totally altered. The principal change it suffers is that of being no longer capable of combustion. If either of these phenomena, namely, the emission of heat and light, and the waste of substance be wanting, we do not say that a body is undergoing combustion, or that it is burning. It follows, therefore, that every theory of combustion ought to explain the following facts:

1. Why a burning body is consumed, and its individuality destroyed.
2. Why, during the progress of this alteration, heat and light are emitted.

For the elucidation of these objects, Lavoisier's theory has laid down the following laws:

1. Combustion cannot take place without the presence of oxygen, and is more rapid in proportion to the quantity of this agent in contact with the inflamed body.
2. In every act of combustion, the oxygen present is consumed.
3. The weight of the products of every body after combustion, corresponds with the weight of the body before combustion, *plus* that of the oxygen consumed.
4. The oxygen absorbed by the combustible body may be recovered from the compound formed, and the weight regained will be equal to the weight which disappeared during the combustion.
5. In every instance of combustion, light and heat, or fire, are liberated,
6. In a limited quantity of air, only a certain quantity of the combustible body can be burnt.
7. The air, wherein a body has been burnt, is rendered unfit for containing combustion, or supporting animal life.

Though every case of combustion re-

quires that light and heat should be evolved, yet this process proceeds very differently in different circumstances; hence the terms *ignition*, or glowing heat; *inflammation*, or accension; and *detonation*, or explosion.

Ignition takes place when the combustible body is not in an aëriiform state.

Charcoal, pyrophorus, &c. furnish instances of this kind.

It seems as if the phenomenon of glowing was peculiar to those bodies which require a considerable quantity of oxygen to become converted into the gaseous state.

The disengagement of caloric and light is rendered more evident to the senses in the act of

Inflammation, or accension. Here the combustible substances are more easily converted into an elastic or aëriiform state. Flame, therefore, consists of the inflammable matter in the act of combustion in the gaseous state. When all circumstances are favourable to the complete combustion of the products, the flame is perfect; if this is not the case, part of the combustible body, capable of being converted into the gaseous state, passes through the luminous flame unburnt, and exhibits the appearance of smoke. Soot, therefore, always indicates an imperfect combustion. Hence a common lamp smokes, an Argand's lamp yields no smoke.

This degree of combustion is very accurately exemplified in the

Flame of Candles.—When a candle is first lighted, which must be done by the application of actual flame, a degree of heat is given to the wick, sufficient to destroy the affinity of its constituent parts; part of the tallow is instantly melted, volatilized, and decomposed; its hydrogen takes fire, and the candle burns. As this is destroyed by combustion, another portion melts, rises and supplies its place, and undergoes a like decomposition. In this way combustion is maintained. The tallow is liquified as it comes into the vicinity of the flame, and is, by the capillary attraction of the wick, drawn up to supply the place of what is decomposed; the unmelted tallow, by this means, forms a kind of cup.

The congeries of capillary tubes which form the wick is black, because the charcoal of the cotton becomes predominant, the circum-ambient air is defended by the flame from oxidating it; it therefore remains, for a considerable time, in its natural state; but when the wick, by the continual consumption of tallow, becomes too long to support itself in a perpendicular position, its upper extremity projects nearly out of the cone of the flame, and there forms a support for an accumulation of soot, which is produced by the imperfect combustion. A candle in this situation, affords scarcely one-tenth of the light it

can give, and tallow candles, on this account, require continual snuffing.

But if the candle be made of wax, the wick does not long occupy its place in the middle of the flame; its thinness makes it bend on one side, when its length is too great for its vertical position; its extremity comes then into contact with the air, and is completely burnt, or decomposed, except so much of it as is defended by the continual afflux of the melted wax. This small wick, therefore, performs the office of snuffing itself. The difficult fusibility of wax enables us to use a thinner wick for it than can be used for tallow, which is more fusible. But wax being a substance which contains much more oxygen than tallow, or oil, the light it affords is not so luminous.

Detonation is an instantaneous combustion, accompanied with a loud report; it takes place in general when the compound resulting from the union of two or more bodies, occupies less space than the substances did before their union; a vacuum is therefore formed, and the surrounding air rushing in from all sides to fill it up is the cause of the report.

Gun-powder, fulminating gold, silver, and mercury; oxygenated muriate of potash; and various other explosive compounds, are capable of producing very loud detonations.

With respect to the disengagement of light and caloric.

By the older chymists, it was universally supposed that the light and heat emitted during combustion, proceeded from the inflammable body; and this opinion would indeed appear unquestionable, while the composition of the atmosphere was imperfectly known. The burning body appeared luminous and felt hot, and no other agent was supposed to be concerned; the conclusion that the light and heat were evolved from the burning substance, was therefore unavoidable. But when the nature of the atmosphere was ascertained, and when it became evident that part of the air was absorbed during combustion, the former conclusion fell to the ground; for, when two bodies exert a mutual action on each other, it becomes *a priori* equally probable that the products may be derived from either of them; consequently, the light and heat evolved might proceed either from the one or the other. Whether they proceed from the atmosphere, or from the combustible body, they must be separated at the part where the combination takes place; that is, upon the surface of the burning body itself; and consequently it appeared luminous and heated, while the air being invisible escaped observation.

When the laws of heat became known,

at least when it was ascertained that bodies in the æriform state contain at the same temperature, and in equal quantities, either of mass or bulk, unequal quantities of heat, the conclusion became probable, that the caloric evolved in combustion proceeded rather from the oxygen gas of the atmosphere, than from the combustible body; since the former contains a much larger quantity than the latter. The caloric evolved was therefore supposed to be derived from the *condensation* of the oxygen gas in the new combination into which it entered.

Though *approaching* to the truth, this explanation is not strictly true. It is not merely from the oxygen gas being *condensed* that the caloric is evolved, because, in many cases of combustion, the product still exists in the gaseous state, and in others, the quantity of caloric evolved bears no proportion to the degree of condensation. Philosophers ascribed this to a change of capacity; for, in different bodies, the difference in the proportion of the capacities before and after combustion, is by no means uniform; and hence the difference in the quantities of caloric extricated in various cases of combustion.

This being premised, it remains to explain the origin of the light emitted during combustion; for although we take it for granted that the caloric is evolved from the oxygen gas, we cannot infer that the light has the same origin.

It is very probable that light is a constituent part of inflammable bodies; for it is frequently evolved in combinations when the oxygen is merely *transferred* from one inflammable substance to another. In those cases it must proceed from the inflammable body. The accension of oils by the affusion of acids, the combustion of metals with sulphur, furnish instances of the kind.

It seems, therefore, probable, that the light is derived from the inflammable substance; and that the oxygen combining with the bases of these substances, disengages the light.

It may be concluded then, that light enters into the composition of all combustible bodies; but as we are unable to separate the light, so as to obtain these bodies pure, we treat of them as simple bodies.

According to this theory, the combustion of phosphorus in oxygen gas is therefore the effect of a double affinity. The basis of the oxygen gas unites with the phosphorus, to form phosphoric acid; and the light disengaged from the phosphorus, together with part of that of the oxygen gas, produces the vivid flame.

The quantity of light emitted by different bodies is supposed to depend on the quantity contained in them, and on the

proportion in which it is united to caloric.

Such is the theory of combustion of Lavoisier, modified by Gren, Leonardi, Richter, and formerly taught in this capital by Dr. Crichton.

Thomson's Theory of Combustion.

Though the preceding theory of combustion is simple and beautiful, it appears from what we are now going to state, to be by no means satisfactory.

It has misled chymists, by confining the term combustion to the act of oxygenation, and considering that all bodies, during their combustion, combine with oxygen, without at the same time recollecting, that this latter effect may take place without any of the phenomena usually attendant on combustion; and that, though certainly all combustion pre-supposes the combination of oxygen with a base, yet this combination may be, and repeatedly is, effected where no combustion can possibly take place. Nothing can be more evident than the difference which, in numberless instances, prevails between the act of oxygenation in bodies and that of combustion, in as much as neither the phenomena attending on, nor the results arising from them, are the same. That a distinction therefore should be made between these processes is obvious; and it is on this account that Dr. Thomson has offered a theory, which considers this subject in a new point of view, and which bids fair to enable us to estimate the phenomena of combustion much better than has hitherto been done.

According to Dr. Thomson's theory, all the bodies concerned in combustion are either, 1. *Combustibles*—2. *Supporters of Combustion*—3. *Incombustibles*.

1. **COMBUSTIBLE BODIES** are those substances which are said, in common language, to *burn*. During the combustion, they appear to emit light and heat, and, at the same time, gradually waste away. When this change has reached its *maximum*, the process of combustion is at an end.

The class of combustibles is very numerous; but all the bodies belonging to it may be sub-divided into three sets, namely:

1. Simple combustibles. 2. Compound combustibles 3. Combustible oxyds.

Simple Combustibles.

1. Sulphur.
2. Phosphorus.
3. Diamond.
4. Hydrogen gas.
5. All the metals.

Compound Combustibles.

The *compound combustibles* consist of compounds, formed by the simple combustibles uniting together two and two, and are of course much more numerous than the simple combustibles. They may be arranged under the five following heads:

1. Sulphurets.

2. Phosphurets.

3. Carburets.

4. Alloys.

5. Sulphurated, phosphorated, and carbonated hydrogen.

The *combustible oxyds* are either simple, having a single base, or compound, having more than one base. All the simple combustible oxyds are by combustion converted into acids.

The compound combustible oxyds are by far the most numerous.

II. The **SUPPORTERS OF COMBUSTION** are bodies which are not of themselves, strictly speaking, capable of undergoing combustion, but which are absolutely necessary for the process; for no combustible body can burn unless some one or other of them be present. Whenever they are excluded, combustion ceases. All the supporters of combustion known at present are the following six:

1. Oxygen gas.
2. Air.
3. Gaseous oxyd of nitrogen.
4. Nitrous gas.
5. Nitric acid.
6. Oxygenated muriatic acid.

There are indeed certain substances besides these, which possess nearly the same properties; these shall be afterwards enumerated under the title of *partial supporters*.

They all contain one common principle, namely, *oxygen*.

III. The **INCOMBUSTIBLE BODIES** are neither capable of undergoing combustion themselves, nor of supporting the combustion of those bodies that are; they are therefore not immediately connected with combustion. At present we are acquainted with about thirteen incombustible bodies, not reckoning the compounds which they are capable of forming with each other. Those are,

1. Nitrogen gas.
2. The alkalis.
3. The earths.

Some of the alkalis and earths possess certain properties in common with combustibles, and are capable of exhibiting phenomena somewhat analogous to combustion; which will be described afterwards under the title of *semi-combustion*.

III. In every case of combustion, there must therefore be present a *combustible* body, and a *supporter* of combustion. During combustion, the combustible always unites with the oxygen of the supporter. *It is this combination which occasions the apparent waste and alteration of the combustible.* The new compound thus formed is a *product of combustion*. Every product of combustion is either, 1. *water*, 2. *an acid*, or 3. *a metallic oxyd*. It is true, indeed, that other bodies sometimes make their appearance during combustion, but

these will be found, upon examination, not to be products, nor to have undergone combustion.

Thus one of the two characteristic marks which distinguish combustion, namely, the *apparent waste and alteration of the combustible body*, has been fully explained. For the explanation of it we are indebted to Lavoisier, as stated before.

But though the combination of the combustible with oxygen be a constant part of combustion, yet the facility with which combustibles burn is not proportional to their parent affinity for oxygen.

Phosphorus, for instance, burns more readily than charcoal; yet charcoal is capable of abstracting oxygen from phosphorus, and of course has a greater affinity for it. The combustible oxyds take fire more readily than some of the simple combustibles; thus charcoal burns more easily than diamond; alcohol, ether, and oils, are exceedingly combustible, whereas all the metals require very high temperatures when the supporter is air.

This greater combustibility of combustible oxyds is probably owing to the weaker affinity by which their particles are united. Hence they are more easily settled than homogeneous particles, and of course combine more readily with oxygen; those simple combustibles which melt easily, or which are in the state of elastic fluids, are also very combustible, because the cohesion between their particles is easily overcome.

It is owing to the same inferiority in the cohesion of heterogeneous particles, that some of the compound supporters occasion combustion in circumstances when the combustibles would not be acted on by simple supporters.

Thus phosphorus burns in air at the common temperature; but it does not burn in oxygen gas unless its temperature be raised. In oxygenated muriatic acid gas, phosphorus burns rapidly at the common temperature of the air, and so do several of the metals; though they cannot be made to burn in air, except at a very high temperature. Thus also oils burn rapidly when mixed with nitrous acid. Nitrous gas and the gaseous oxyd of nitrogen, constitute exceptions to this rule.

IV. None of the *products* of combustion are combustible, according to the definition of combustion here given. This want of combustibility is not owing to their being saturated with oxygen; for several of them are capable of combining with an *additional dose* of it. But, during this combination, no caloric or light is ever emitted; and the compound formed differs essentially from a *product* of combustion; for, by this additional dose of oxygen, the *product* is converted into a *supporter*. Hence we see that *combustion ought not to be confounded with*

the combination of a body with oxygen, as was done formerly.

Combustion, indeed, cannot take place without the combination of oxygen; but oxygen may combine with bodies in different proportions without the phenomena of combustion; and the *product obtained* is capable of becoming converted into a *supporter of combustion*; for instance, if lead be melted, and kept so for some time, it becomes covered with a white pellicle, or white *oxyd of lead*, a product consisting of oxygen and lead; but if this white oxyd is suffered to be heated longer, it absorbs an additional quantity of oxygen, and becomes converted into a yellow powder, called *yellow oxyd of lead*. If this yellow oxyd be again exposed to heat, it absorbs still more oxygen, and becomes converted into *red oxyd of lead*. All these oxyds are therefore *supporters*. When the *supporters* thus formed by the combination of oxygen with *products*, are made to support combustion, they do not lose all their oxygen, but only the additional dose which constituted them supporters. Of course they are again reduced to their original state of products of combustion. Hence it follows, that they owe their properties as supporters, not to the *whole* of the oxygen which they contain, but to the *additional dose* which constituted them supporters. We may therefore call them *partial supporters*, indicating by the term, that part only of their oxygen is capable of supporting combustion, and not the whole.

All the partial supporters with which we are acquainted, contain a metallic basis; for metallic oxyds are the only products at present known, capable of combining with an additional dose of oxygen. It is a circumstance highly deserving attention, that when metals are capable of combining with several doses of oxygen, the product, or oxyd formed by combustion is seldom or never that which contains a *maximum* of oxygen.

Thus it is evident that several of the products of combustion are capable of combining with oxygen. *The incombustibility of products, therefore, is not owing to their want of affinity for oxygen, but to some other cause.*

5. No product of combustion is capable of *supporting* combustion. This is not occasioned by any want of affinity to combustible bodies; for several of them are capable of combining with an additional dose of their basis. But by this combination, they *lose* their properties as products, and are converted into *combustibles*. The process, therefore, differs essentially from combustion. Thus sulphuric acid, a *product* of combustion, by combining with an additional dose of sulphur, or its oxyd, is converted into *sulphureous acid*, a substance which, from several of its properties, the

doctor concludes to be combustible. Thus also phosphoric acid, a product of combustion, is capable of combining with phosphorated hydrogen, and of forming *phosphorous* acid a combustible body. When this last acid is heated in contact with a supporter, it undergoes combustion; but it is only the additional dose of the combustible which burns, and the whole is converted into phosphoric acid. Hence we see that it is not the whole basis of these compounds which is combustible, but merely the additional dose. The compounds, therefore, formed by the union of a product and combustible, may be termed *partial combustibles*; indicating by the name, that a part only of the base is capable of undergoing combustion. Since the products of combustion are capable of combining with oxygen, but never exhibit the phenomena of combustion, except when they are in the state of partial combustibles, combustible bodies must contain a substance which they lose in burning, and to which they owe their combustibility; for, after they have lost it, they unite to oxygen *without* exhibiting the phenomena of combustion.

Though the products of combustion are not capable of supporting combustion, they not unfrequently part with their oxygen just as supporters do, give it out to combustibles, and convert them into products; but during this process, no heat nor light is ever evolved. Water, for instance, gives out its oxygen to iron, and converts it into *black oxyd*, a product; and sulphuric acid gives out its oxygen to phosphorus, and converts it into phosphoric acid. Thus we see that the oxygen of products is capable of converting combustibles into products, just as the oxygen of supporters; but during the combustion of the last only, are heat and light emitted. The oxygen of supporters then contains something which the oxygen of products wants.

6. Whenever the whole of the oxygen is abstracted from products, the combustibility of their base is restored as completely as before combustion; but no substance is capable of abstracting the whole of the oxygen, except a *combustible*, or a *partial combustible*. Water, for instance, is a product of combustion, whose base is hydrogen. To restore the combustibility of the hydrogen, we have only to mix water with iron or zinc filings, and an acid; the metal is oxydated, and the hydrogen gas is evolved as combustible as ever. But no substance, except a combustible, is capable of separating hydrogen gas from water, by combining with its oxygen. Thus we see that combustibles are capable of restoring the combustibility of the bases of products; but they themselves lose their combustibility by the process, and are converted into products. Combustibility, therefore, may

be thrown at pleasure from one body to another.

From these facts it is obvious, that the products of combustion may be formed without combustion; but in these cases a new combustible is always evolved. The process is merely an interchange of combustibility; for the combustible is converted into a product only by means of a product. Both the oxygen and the base of the product having undergone combustion, have lost something which is essential to combustion. The process is merely a double decomposition. The product yields its oxygen to the combustible, while at the same time the combustible gives out something to the base of the product; the combustibility of that base then is restored by the loss of its oxygen, and by the restoration of something which it receives from the other combustible thus converted into a product.

There is indeed another method of forming the products of combustion without actual combustion in certain cases; but the phenomena are much more complicated. This method is to expose them to the action of some of the supporters dissolved in water; especially nitric acid. Thus most of the metallic oxyds may be formed without combustion by the action of that acid on the metals. But, in that case, a new supporter is always evolved, namely, nitrous gas; ammonia, a new combustible, is also usually formed; and, not unfrequently, the *product* is converted into a *partial supporter*.

7. No supporter can be produced by combustion, or by any equivalent process. As all the supporters, except oxygen gas, consist of oxygen combined with a base, it follows as a consequence, that oxygen may combine with a base without losing that ingredient, which occasions combustion. The act of combination of oxygen with a base, therefore, is by no means the same with combustion. If we take a view of the different supporters, we shall find that all of them which can be obtained artificially, are procured either from other supporters or by the agency of electricity.

I. OXYGEN gas may be procured from nitric acid and oxygenated muriatic acid, two supporters; and from several of the partial supporters, as the black oxyd of manganese, the red oxyds of lead and of mercury. The action of heat is always necessary; but the process is very different from combustion.

II. AIR, as far as is known at present, cannot be formed artificially. The gas, indeed, which comes over during part of the usual distillation of nitrate of potash and sulphuric acid, to obtain nitrous acid, resembles air very closely. But it is obtained from a supporter.

III. THE GASEOUS OXYD OF NITROGEN

has hitherto been only procured from nitrous gas and nitric acid (nitrate of ammonia,) both of which are supporters.

IV. NITROUS GAS can only be procured by the decomposition of nitric acid, a supporter.

V. OXIGENATED MURIATIC ACID can be formed by combining muriatic acid with the oxygen of the black oxyd of manganese, the red oxyds of lead, iron, mercury; all of which are partial supporters.

VI. NITRIC ACID is formed spontaneously upon the surface of the earth, by processes with which we are but imperfectly acquainted; but which certainly have no resemblance to combustion. Its oxygen is probably furnished by the air, which is a supporter; at least, it has been observed, that nitrogen and oxygen, at high temperatures, are capable of forming nitric acid.

This formation of nitric acid by means of electricity, has been considered as a combustion, but for what reason it is not easy to say: the substance acted upon is not a combustible with a supporter, but a supporter alone. Electricity is so far from being equivalent to combustion, that it sometimes acts in a manner diametrically opposite; *unburning*, if we may use the expression, a substance which has already undergone combustion, and converting a *product* into a *combustible* and a *supporter*. Thus it decomposes water, and converts it into oxygen and hydrogen gas; therefore it must be capable of supplying the substances which the oxygen and combustible lose when they combine by combustion, and form a product.

8. Several of the supporters and partial supporters are capable of combining with combustibles, without undergoing decomposition, or exhibiting the phenomena of combustion. In this manner, the yellow oxyd of gold and the white oxyd of silver combine with ammonia; the red oxyd of mercury with oxalic acid; and oxygenated muriatic acid with ammonia. Thus also nitrate of potash and oxygenated muriate of potash may be combined, or at least intimately mixed with several combustible bodies, as in gun-powder, fulminating powder, &c. In all these compounds, the oxygen of the supporter and the combustible retain the ingredients which render them susceptible of combustion; hence the compound is still combustible. And in consequence of the intimate combination of the component parts, the least alteration is apt to destroy the equilibrium which subsists between them; the consequence is, combustion and the formation of a new compound. Hence these compounds burn with amazing facility, not only when heated, but when triturated, or struck smartly with a hammer. They have therefore received the name of *detonating* or *fulminating* bodies. Thus we have fulminating gold,

fulminating silver, fulminating mercury, fulminating powder, &c.

9. Such are the properties of the combustibles, the supporters, and the products; and such the phenomena which they exhibit when made to act upon each other.

If we compare together the *supporters* and the *products*, we shall find that they resemble each other in many respects. Both of them contain oxygen, as an essential constituent part; both are capable of converting combustibles into products; and several of both combine with combustibles and with additional doses of oxygen. But they differ from each other in their effects on combustibles. The former only produce combustion; whereas the products convert combustibles into products without combustion. Now, as the ultimate change produced upon combustibles by both these sets of bodies is the same, and as the substance which combines with the combustibles is in both cases the same, namely oxygen, we must conclude that this oxygen in the supporters contains something which the oxygen of the products wants, something which separates during the passage of the oxygen from the product to the combustible, and occasions the combustion, or emission of fire, which accompanies this passage. The oxygen of supporters then contains some ingredient which the oxygen of products wants. Many circumstances concur to render it probable that this ingredient is *caloric*.

The *combustibles* and the *products* also resemble each other. Both of them contain the same or a similar base; both frequently combine with combustibles, and likewise with oxygen; but they differ essentially in the phenomena which accompany their combination with oxygen. In the one case, *fire is emitted*, in the other not. If we recollect that no substance but a combustible is capable of restoring combustibility to the base of a product, and that at its doing so it always loses its own combustibility; and if we recollect farther, that the base of a product does not exhibit the phenomena of combustion even when it combines with oxygen, we cannot avoid concluding, that all combustibles contain an ingredient which they lose when converted into products, and that this loss contributes to the fire which makes its appearance during the conversion. Many circumstances contribute to render it probable that this ingredient is *light*.

If we suppose that the oxygen of supporters contains caloric as an essential ingredient, and that light is a component part of all combustibles, the phenomena of combustion above enumerated, numerous and intricate as they are, admit of an easy and obvious explanation. The component parts of the oxygen of supporters are two; namely, 1. a base, 2. caloric. The com-

ponent parts of combustibles are likewise two; namely, 1. a base, 2. light. During combustion, the base of the oxygen combines with the base of the combustible, and forms the product; while, at the same time, the caloric of the oxygen combines with the light of the combustible, and the compound flies off in the form of fire. Thus combustion is a double decomposition; the oxygen and combustible divide themselves each into two portions, which combine in pairs; the one compound is the *product*, and the other the *fire*, which escapes.

Hence the reason that the oxygen of products is unfit for combustion. It wants its caloric. Hence the reason that combustion does not take place when oxygen combines with products, or with the base of supporters. These bodies contain no light. The caloric of the oxygen of course is not separated, and no fire appears. And this oxygen still retaining its caloric, is capable of producing combustion whenever a body is presented which contains light, and whose base has an affinity for oxygen. Hence also the reason why a combustible alone can restore combustibility to the base of a product. In all such cases, a double decomposition takes place. The oxygen of the product combines with the base of the combustible, while the light of the combustible combines with the base of the product.

But the application of this theory to all the different phenomena described above, is so obvious, that it is needless to give any more examples. Let us rather inquire, with the author, into the evidences which can be brought forward in its support.

10. As caloric and light are always emitted during combustion, it follows that they must have previously existed in the combustible, the supporter, or in both.

That the oxygen of the supporters contains either one or both of these substances, follows incontrovertibly from a fact already mentioned, namely, that the oxygen of products will not support combustion, while that of supporters will. Hence the oxygen of supporters must contain something which the oxygen of the products wants, and this something must be caloric, or light, or both.

That the oxygen of some of the supporters at least contains caloric, as an ingredient, has been proved, in a satisfactory manner, by the experiments of Crawford, Lavoisier, and La Place. Thus the temperature of hot blooded animals is maintained by the decomposition of *air*. Now if the oxygen of one supporter contains caloric, the same ingredient must exist in the oxygen of every supporter, because all of them are obviously in the same state. Hence we conclude that the oxygen of every supporter contains caloric as an essential ingredient.

The light emitted during combustion, must either proceed from the combustible or the supporter. That it proceeds from the combustible, must appear pretty obvious, if we recollect that the colour of the light emitted during combustion varies, and that this variation usually depends, not upon the supporter, but upon the combustible. Thus charcoal burns with a red flame, sulphur with a blue or violet, zinc with a greenish white, &c.

The formation of combustibles in plants, obviously requires the presence and agency of light. The leaves of plants emit oxygen gas, when exposed to the sun's rays, but never in the shade, or in the dark.

Besides vegetation, we are acquainted with two other methods of *unburning* products, or of converting them into products and combustibles, by exposing them, in certain circumstances, to the agency of *fire*, or of *electricity*. The oxydes of lead, mercury, &c. when heated to redness, are decomposed, oxygen gas is emitted, and the pure metal remains behind. In this case, the necessary caloric and light must be furnished by the fire; a circumstance which explains why such reductions always require a red heat. When carbonic acid is made to pass repeatedly over red-hot charcoal, it combines with a portion of charcoal, and is converted into gaseous oxyd of carbon. If this gas be a combustible oxyd, the base of the carbonic acid and its oxygen must have been supplied with light and caloric from the fire; but if it be a *partial combustible*, it is merely a compound of carbonic acid and charcoal: which of the two it is, remains still to be ascertained.

Electricity decomposes water, and converts it into oxygen gas and hydrogen gas; it must therefore supply the heat and the light which these bodies lost when converted into a product.

These facts, together with the exact correspondence of the theory given above with the phenomena of combustion, render it so probable, that Dr. Thomson has ventured to propose it as an additional step towards a full explanation of the theory of combustion. Every additional experiment has served to confirm it more and more. It even throws light upon the curious experiments of the accension of metals with sulphur, which succeed *in vacuo*, under mercury, in nitrogen gas, &c.

Dr. Thomson has noticed, that the same emission of caloric and light, or of *fire*, takes place when melted sulphur is made to combine with potash, or with lime, in a crucible or glass tube, and likewise when melted phosphorus is made to combine with lime heated to redness. He supposes that in all probability, barytes and strontia exhibit the same phenomenon when combined with melted sulphur or phosphorus; and

perhaps some of the metals when combined with phosphorus.

The phenomena Dr. Thomson explains thus: The sulphur and phosphorus are in the melted state, and therefore contain caloric as an ingredient; the alkalies, earths, and metals, which produce the phenomenon in question, contain light as an essential ingredient. The sulphur, or phosphorus combines with the base of the metal, earth, or alkali; while at the same time, the caloric, to which the sulphur or phosphorus owed its fluidity, combines with the *light* of the metal, earth or alkali; and the compound flies off under the form of *fire*.

Thus the process is exactly the same with: combustion, excepting as far as regards the product. The melted sulphur, or phosphorus, acts the part of the *supporter*, while the metal, earth, or alkali, occupies the place of the *combustible*. The first furnishes caloric, the second light, while the base of each combines together. Hence we see that the base of sulphurets and phosphurets resembles the base of products in being destitute of light; the formation of these bodies exhibiting the separation of fire like *combustion*, but the product differing from a product of combustion in being destitute of oxygen, Dr. Thomson distinguishes the process by the title of *semi-combustion*; indicating by the term, that it possesses one half of the characteristic marks of combustion, but is destitute of the other half.

The only part of this theory which requires proof is, that light is a component part of the earths and alkalies. But as potash and lime are the only bodies of that nature, which we are certain to be capable of exhibiting the phenomena of semi-combustion, the proofs must of necessity be confined to them. That lime contains light as a component part, has been long known. Meyer and Pelletier observed long ago, that when water is poured upon lime, not only heat but light is emitted. Light is emitted also abundantly when sulphuric acid is poured upon magnesia, or upon lime, potash, or soda, freed from the water of crystallization. In all these cases, a *semi-combustion* takes place. The water and the acid being solidified, give out *caloric*, while the lime or potash gives out *light*.

That lime during its burning combines with light, and that light is a component part of lime, is demonstrated by the following experiment, for which we are indebted to Scheele.

Fluor spar (fluat of lime) has the property of phosphorescing strongly when heated, but the experiment does not succeed twice with the same specimen. After it has been once heated sufficiently, no subsequent heat will cause it to phosphorate. Now phosphorescence is merely the emis-

sion of light; light of course is a component part of fluor spar, and heat has the property of separating it. But the phosphorescing quality of the spar may be again recovered to it, or, which is the same thing, the light which the spar had lost may be restored by the following process:

Decompose the fluat of lime by sulphuric acid, and preserve the fluoric acid separate. Boil the sulphate of lime thus formed with a sufficient quantity of carbonate of soda; a double decomposition takes place; sulphate of soda remains in solution, and carbonate of lime precipitates. Ignite this precipitate in a crucible, till it is reduced to lime, and combine it with the fluoric acid to which it was formerly united. The fluor spar thus regenerated, phosphoresces as at first. Hence the lime, during its ignition, must have combined with light.

That potash contains light, may be proved in the same manner as the existence of that body in lime. Now as potash is deprived of its carbonic acid by lime, the Doctor supposes that the process must be a double decomposition; namely, that the base of the lime combines with carbonic acid, while its light combines with the potash.

These remarks on semi-combustion might easily be much enlarged upon. For it is obvious, that whenever a liquid combines with a solid containing light, and the product is a solid body, something analogous to semi-combustion must take place. Hence the reason why water increases the violence of combustion when thrown sparingly into a common fire, &c.

COMEDONES. (From *comedo*, a glutton.) A sort of worms which eat into the skin and devour the flesh.

Comfrey. See *Symphitum*.

COMISDI. The gum arabic.

COMISTE. The epilepsy. This name arose from the frequency of persons being seized with this disorder while in the assemblies called *Comitia*.

COMITISSA. (A countess.) Some preparations are distinguished by this name, as *pulvis Comitissæ de Cantia*, the Countess of Kent's powder.

COMMAGENUM. (From *Commagene*, a place in Syria, whence it was brought.) Syrian ointment. mentioned by Galen.

COMMANDUCATIO. (From *commanduco*, to eat.) The act of mastication, or chewing.

COMMANSUM. (From *commando*, to eat.) A masticatory. A medicine put into the mouth chewed to promote a discharge of phlegm, or saliva.

COMMENDATORIUS. (From *commendo*, to recommend.) An epithet of the traumatic balsam, *Tinctura Benzoes composita*, from its singular virtues and usefulness.

COMMI. Gum. When alone it signifies

gum Arabic. The *καμμη λαικον* mentioned by Hippocrates in his *De Morb. Mulieb.* is gum Arabic.

COMMISSURA (From *committo*, to join together.) A suture, juncture, or joint. A term applied in anatomy to the corners of the lips, where they meet together; and also to certain parts of the brain which go across and join one hemisphere to the other.

COMMISSURA ANTERIOR CEREBRI. The white nerve-like substance which crosses the anterior part of the third ventricle of the brain, immediately above the infundibulum, and between the anterior crura of the fornix; uniting one hemisphere of the brain with the other.

COMMISSURA MAGNA CEREBRI. The *corpus callosum* of the brain is so termed by some writers.

COMMISSURA POSTERIOR CEREBRI. A white nerve-like substance, which passes from one hemisphere of the brain across to the other, immediately over the opening of the aqueduct of Sylvius, in the posterior part of the third ventricle of the brain, and above the *corpora quadrigemina*.

COMMUNICANT. (From *communico*, to make partake.) A term applied, by Bellini, to fevers of two kinds afflicting the same person, wherein as one goes off the other immediately succeeds.

COMPAGES. (From *compingo*, to put together.) A suture, or joint. A commissure.

COMPARATIVE ANATOMY. *Anatomia comparativa.* Zootomy. The dissection of brute animals and fishes, to compare them with the human body.

CONFERA. See *Cubeba*.

COMPLETION. A term used by the ancient writers in various acceptations; but latterly it signifies only the same as *Plethora*.

COMPLEXUS. (*Complexus*, sc. *musc.* from *complector*, to comprise.) *Complexus seu biventer cervicis* of Albinus. *Dorso trachelon occipital* of Dumas. A muscle situated on the back part of the neck, that draws the head backwards, and to one side; and when both act, they draw the head directly backward. It arises from the transverse processes of the seven superior vertebrae of the back, and four inferior of the neck, by as many distinct tendinous origins; in its ascent, it receives a fleshy slip from the spinous process of the first vertebrae of the back: from these different origins it runs upwards, and is every where intermixed with tendinous fibres. It is inserted, tendinous and fleshy, into the inferior edge of the protuberance in the middle of the *os occipitis*, and into a part of the curved line that runs forwards from that protuberance. It draws the head backwards.

COMPRESSION. (From *comprimo*, to

press together.) By this term, surgeons express a diseased state of the body, which is the effect of something pressing upon the brain. It should be distinguished from concussion and inflammation. When the brain is compressed either by bone, extravasated blood, or any other fluid, there is a general insensibility, the eyes are half open, the pupil dilated and motionless, even when a candle is brought near the eye; the retina is insensible; the limbs relaxed; the breathing stertorous; the pulse slow, and, according to Mr. Abernethy, less subject to intermission than in cases of concussion. Nor is the patient ever sick, when the pressure on the brain and the general insensibility are considerable; for the very action of vomiting betrays an irritability in the stomach and oesophagus.

COMPRESSOR NARIS. (*Compressor*; from *comprimo*, to press together) *Reneus vel nasalis* of Douglas. *Transversarius vel myrtiliformis* of Winslow. *Dilatator alarum nasi* of Cowper; and *Maxillo narial* of Dumas. A muscle of the nose, that compresses the ala towards the septum nasi, particularly when we want to smell acutely. It also corrugates the nose, and assists in expressing certain passions. It arises, by a narrow beginning, from the root of the ala nasi externally, and spreads into a number of thin, separate fibres, which run up along the cartilage in an oblique manner towards the back of the nose, where it joins with its fellow, and is inserted into the narrow extremity of the *os nasi*, and nasal process of the superior maxillary bone.

COMPURETIO. (From *compungo*, to prick.) A puncture.

CONARIUM. (From *conus*, a cone.) The pineal gland is so named, from its conical shape. See *Pineal gland*.

CONCAUSA. (From *con*, with, and *causa*, a cause.) A cause which co-operates with another in the production of a disease.

CONCENTRANTIA. (From *concentro*, to concentrate.) Absorbents of acids are so called, because they remove the obstructions which keep asunder the affinities between the two powers.

CONCENTRATION. (From *con*, and *centrum*, having the same centre.) The volatilizing of part of the water of fluids in order to improve their strength. The matter to be concentrated, therefore, must be of superior gravity to water. This operation is performed on some acids, particularly the sulphuric and phosphoric. It is also employed in solutions of alkalis and neutral salts.

CONCEPTION. (*Conceptio*; from *concipio*, to conceive.) The impregnation of the ovulum in the female ovarium by the subtle prolific aura of the semen virile. In order to have a fruitful coition, it is necessary that the semen be propelled into the uterus, or vagina, so that its fe-

condensing vapour shall be conveyed through the Fallopian tubes to the ovarium: it is also necessary that there be a certain state of the ovarium of the female in order to impregnate it; which is, that the ovum shall be mature, and embraced by the fimbriae of the Fallopian tube, to convey that vivifying principle to the ovum. See *Generation*.

CONCHA. (*Concha*, *αρχή*, a liquid measure amongst the Athenians.) A term applied by anatomists to several parts of the body, as the hollow of the ear, the spongy bones of the nose, &c.

CONCHA AURICULÆ. See *Auricula*.

CONCHA AURIS. The hollow part of the cartilage of the outer ear.

CONCHÆ NARIUM. (*Concha*, a shell.) The turbinated portion of the ethmoid bone, and the inferior spongy bones of the nose, which are covered by the Schneiderian membrane, are so termed.

CONCHUS. (From *αρχή*, a shell; so named from their likeness to a shell.) The cranium, and the cavity of the eye.

CONCIDENTIA. (From *concide*, to decay.) A decrease of bulk in the whole or any part of the body. A diminution of a tumour.

CONCOAGULATIO. (From *con*, and *coagulo*, to coagulate together.) The coagulation, or chrySTALLIZATION of different salts, first dissolved together in the same fluid.

CONCOCTIO. From *concoquo*, to digest.) Digestion. That operation of nature upon morbid matter which renders it fit to be separated from the healthy fluids.

CONCREMATIO. (From *con*, and *crema*, to burn together.) The same as calcination.

CONCRETION. (From *concreresco*, to grow together.)

1. The condensation of any fluid substance into a more solid consistence.

2. The growing together of parts which, in a natural state, are separate.

CONCURSUS. (From *concurro*, to meet together.) The congeries or collection of symptoms which constitute and distinguish the particular disease.

CONCUSSION. From *concutio*, to shake together.) Concussion of the brain. Various alarming symptoms, followed sometimes by the most fatal consequences, are found to attend great violence offered to the head; and upon the strictest examination, both of the living and the dead; neither fissure, fracture, nor extravasation of any kind can be discovered. The same symptoms and the same events are met with, when the head has received no injury at all *ab externis*, but has only been violently shaken; nay, when only the body, or general frame, has seemed to have sustained the whole violence. The symptoms attending a concussion, are generally in proportion to the degree of violence which

the brain itself has sustained; and which, indeed, is cognizable only by the symptoms. If the concussion be very great, all sense and power of motion are immediately abolished, and death follows soon; but between this degree and that slight confusion (or stunning, as it is called,) which attends most violences done to the head, there are many stages. The following is Mr. Abernethy's description of the symptoms of concussion, and which, he is of opinion may be properly divided into three stages.

The first is, that state of insensibility and derangement of the bodily powers which immediately succeeds the accident. While it lasts, the patient scarcely feels any injury that may be inflicted on him. His breathing is difficult, but in general without stertor; his pulse intermitting, and his extremities cold. But such a state cannot last long; it goes off gradually, and is succeeded by another, which is considered as the *second* stage of concussion. In this, the pulse and respiration become better, and though not regularly performed, are sufficient to maintain life, and to diffuse warmth over the extreme parts of the body. The feeling of the patient is now so far restored, that he is sensible of his skin being pinched; but he lies stupid and inattentive to slight external impressions. As the effects of concussion diminish, he becomes capable of replying to questions put to him in a loud tone of voice, especially when they refer to his chief suffering at the time, as pain in the head, &c.; otherwise he answers incoherently, and as if his intention was occupied by something else. As long as the stupor remains, the inflammation of the brain seems to be moderate; but as the former abates, the latter seldom fails to increase; and this constitutes the *third* stage, which is the most important of the series of effects proceeding from a concussion.

These several stages vary considerably in their degree and duration; but more or less of each will be found to take place in every instance where the brain has been violently shaken. Whether they bear any certain proportion to each other or not, is not known; indeed this will depend upon such a variety of circumstances in the constitution, the injury, and the after-treatment, that it must be difficult to determine.

To distinguish between an extravasation and a commotion by the symptoms only, Mr. Pott says, is frequently a very difficult matter; sometimes an impossible one. The similarity of the effects in some cases, and the very small space of time which may intervene between the going off of the one and the accession of the other, render this a very nice exercise of the judgment. The first stunning, or deprivation of sense,

whether total or partial, may be from either, and no man can tell from which; but when these first symptoms have been removed, or have spontaneously disappeared, if such patient is again oppressed with drowsiness, or stupidity, or total or partial loss of sense, it then becomes probable that the first complaints were from commotion, and that the latter are from extravasation; and the greater the distance of time between the two, the greater is the probability not only that an extravasation is the cause, but that the extravasation is of the limpid kind, made gradatim, and within the brain.

Whoever seriously reflects on the nature of these two causes of evil within the cranium, and considers them as liable to frequent combination in the same subject, and at the same time considers that, in many instances, no degree of information can be obtained from the only person capable of giving it (the patient), will immediately be sensible how very difficult a part a practitioner has to act in many of these cases, and how very unjust it must be to call that ignorance which is only a just diffidence arising from the obscurity of the subject, and the impossibility of attaining materials to form a clear judgment.

CONDENSATION. (From *condenso*, to make thick.) A contraction of the pores of the skin, by means of astringent or cooling medicines. A thickening of any fluid.

CONDIMENTUM. (From *condio*, to preserve, or season.) A preserve, or sweetmeat.

CONDUCTIO. (From *conduco*, to draw along.) In Cælius Aurelianus, it is a spasm, or convulsion, drawing the muscles out of their proper positions.

CONDUCTOR. (From *conduco*, to lead, or guide.) A surgical instrument, whose use is to direct the knife in certain operations. It is more commonly called a director.

CONDYLE. (From *κονδυ*, an ancient cup, shaped like a joint.) A rounded eminence of a bone in any of the joints.

CONDYLOMA. (From *κονδυλος*, a tubercle, or knot.) *Sarcoma*. A soft, wart-like excrescence, that appears about the anus and pudendum of both sexes. There are several species of condylomata, which have received names from their appearances, as *ficus*, *crystæ*, *thymus*, from their resemblance to a fig, &c.

CONESION. In Hippocrates it imports the *Cicuta*. It is said to be thus named (from *κονειν*, to turn round,) because it produces a vertigo in those who take it inwardly.

Conessi bark. See *Conessi cortex*.

CONESSI CORTEX. *Codagopala.* *Cortex profluvi.* The bark of the *Nerium antidysentericum* of Linæus:—*foliis ova-*

lis, acuminatis, postolatis. It grows on the coast of Malabar. It is of a dark black colour externally, and generally covered with a white moss, or scurf. It is very little known in the shops; has an austere, bitter taste; and is recommended in diarrhæas, dysenteries, &c. as an adstringent.

CONFECTIO. (From *conficio*, to make up.) A confection. In general it means any thing made up with sugar. This term, in the new London Pharmacopœia, includes those articles which were formerly called electuaries and conserves, between which there do not appear to be sufficient grounds to make a distinction.

CONFECTIO AMYGDALÆ. Confection of almond. Take of sweet almonds, an ounce; Acacia gum powdered, a drachm; refined sugar, half an ounce. The almonds having been previously macerated in water, and their external coat removed, beat the whole together, until they are thoroughly incorporated. It has been objected to the almond mixture, which is an article of very general use, that it requires considerable time for its extemporaneous preparation, and that it spoils and cannot be kept when it is made. This will be obviated by the present form, which does keep for a sufficient length of time, and rubs down into the mixture immediately.

CONFECTIO AROMATICA. This preparation was formerly called *Confectio cardaca.* *Confectio Raleighana.* Take of cinnamon bark, nutmegs, of each two ounces; cloves, an ounce; cardamom seeds, half an ounce; saffron dried, 2 ounces; prepared shells, 16 ounces; refined sugar powdered, two pounds; water, a pint. Reduce the dry substances, mixed together, to very fine powder; then add the water, and gradually mix the whole, until it is incorporated. This preparation is much simplified by the London college. It is an excellent medicine, possessing stimulant, antispasmodic, and adstringent virtues; and is exhibited with these views to children and adults, in a vast variety of diseases, mixed with other medicines. It may be given in doses of 10 grains to an ounce.

CONFECTIO AURANTII. *Conserva corticis exterioris aurantii hispalensis.* *Conserva flavedinis corticis aurantium.* Take of fresh external rind of oranges, separated by rasping, a pound; refined sugar, three pounds. Bruise the rind with a wooden pestle, in a stone mortar; then, after adding the sugar, bruise it again, until the whole is thoroughly incorporated. This is well calculated to form the basis of a tonic and stomachic confection, and may be given alone in doses of from 2 to 5 drachms, twice, or three times a-day.

CONFECTIO CARMACA. See *Confectio aromatica*.

CONJECTIO CASSIAE. *Electuarium cassiae.* *Electuarium e cassia.* Confection of cassia. Take of fresh cassia pulp, half a pound; manna, two ounces; tamarind pulp, an ounce; syrup of roses, half a pound. Bruise the manna; melt it in the syrup by a water-bath; then mix in the pulps, and evaporate down to a proper consistence. This is a very elegant, pleasant, and mild aperient for the feeble, and for children. Dose from 2 drachms to an ounce.

CONJECTIO OPII. *Confectio opiata.* *Philonium Londinense.* *Philonium Romanum.* Confection of opium. Take of hard opium powdered, six drachms; long pepper, an ounce; ginger-root, two ounces; carraway-seed, three ounces; syrup, a pint. Rub together the opium and the syrup previously heated; then add the remaining articles reduced to powder, and mix. To the credit of modern pharmacy, this is the only one that remains of all those complicated and confused preparations called mithridate, theriaca, &c.; it more nearly approximates, in its composition, the philonium than any other, and may be considered as an effectual substitute for them in practice. This very warm and stimulating confection is admirably calculated to relieve diarrhoea, spasms of the stomach and bowels, and is frequently ordered as a nervine, stimulant, and adstringent, in doses of from 10 grs. to half a drachm. About 36 grains contains one of opium.

CONJECTIO ROSAE CANINAE. *Conserua cynosbati.* *Conserua fructus cynosbati.* Conserve of hips. Confection of dog-rose. Take of dog-rose pulp, a pound; refined sugar powdered, twenty ounces. Rub them together until they are thoroughly incorporated. This preparation is cooling and adstringent; it is seldom given alone, but mostly joined to some other medicine, in the form of linctus, or electuary.

CONJECTIO ROSAE GALLICAE. *Conserua rosae.* *Conserua rosarum rubrarum.* Conserve of red rose. Take of the petals of the red rose, before it is expanded, and without the claws, a pound; refined sugar, three pounds. Bruise the petals in a stone mortar; then, having added the sugar, beat them again together, until they are thoroughly incorporated. This is an excellent subastringent composition. Rubbed down with water, it forms an excellent drink, with some lemon juice, in hemorrhagic complaints; it may also be given with vitriolated zinc, in the form of an electuary.

CONJECTIO RUTAE. *Electuarium e baculis lauri.* Confection of rue. Take of rue leaves dried, carraway seeds, bay berries, of each an ounce and a half; sagapenum, half an ounce; black pepper, two drachms; clarified honey, sixteen ounces. Rub the dry articles together, into a very fine powder; then add the honey, and mix

the whole. Its use is confined to chylsters.

CONJECTIO SCAMMONIÆ. *Electuarium scammonii.* *Electuarium e scammonio.* *Electuarium caryocostinum.* Confection of scammony. Take of scammony gum resin powdered, an ounce and a half; cloves bruised, ginger-root powdered, of each 6 drachms; oil of carraway, half a drachm; syrup of roses, as much as is sufficient. Rub the dry articles together, into very fine powder; next rub them again whilst the syrup is gradually added; then add the oil of carraway, and mix the whole well together. This is a strong stimulating cathartic, and calculated to remove worms from the primæ viæ, with which view it is mostly exhibited. Dose from ʒss to ʒj.

CONJECTIO SENNÆ. *Electuarium sennæ.* *Electuarium lenitivum.* Confection of senna. Take of senna leaves, eight ounces; figs, a pound; tamarind pulp, pulp of prunes, cassia pulp, of each half a pound; coriander seeds, four ounces; liquorice root, three ounces; refined sugar, two pounds and a half. Powder the senna leaves with the coriander seeds, and separate, by sifting ten ounces of the mixed powder. Boil the remainder with the figs and the liquorice-root, in four pints of water, until it be reduced to half; then press out and strain the liquor. Evaporate the liquor, until a pint and a half only remains of the whole; then add the sugar, to make syrup. Lastly, mix the pulps gradually with the syrup, and, having added the sifted powder, mix the whole together. This is a mild and elegant aperient, well adapted for pregnant women, and those whose bowels are easily moved. Dose, ʒss to ʒss.

CONFERVA. (From *confervee*, to knit together.) 1. The name of a genus of plants in the Linnean system. Class, *Cryptogamia*. Order, *Alga*.

2. A kind of moss: named from its use formerly in healing broken bones.

CONFERVA HELMINTHOCORTOS. See *Corallina coriacea*.

CONFERVA RIVALIS. This plant, *Conferua rivalis* of Linneus:—*filamentis simplicissimis equalibus longissimis*, has been recommended in cases of spasmodic asthma, phthisis, &c. on account of the great quantity of vital air it contains.

CONFIRMANTIA. (From *con*, and *firmo*, to strengthen.) Restoratives; also medicines which fasten the teeth in their sockets.

Confluent small-pox. See *Variola*.

CONFLUXIO. It is much used by Hippocrates, and his interpreter Galen, in the same sense as we use consent and transpirable, from a notion that parts at a distance have mutual consent with one another, and that they are all perspirable by many subtle streams. Paracelsus, according to his way, expressed the former by confederation.

CONFORMATION. (From *conforme*, to shape or fashion.) The natural shape and form of any thing, also a description of some diseases which arise from a bad formation of the parts.

CONFORTANTIA. (From *conforto*, to strengthen.) Cordial medicines. Strengtheners.

CONFORTATIVA. The same.

CONFUSIO. (From *confundo*, to mix together.) A confusion, or disorder in the eyes, proceeding from a rupture of the membranes, which include the humours, by which means they are all confounded together.

CONGELATI. (From *congelio*, to freeze.) *Congelatici.* Persons afflicted with a catalepsy are so called, by which all sensation seems to be taken away.

CONGELATION. (From *congelio*, to freeze.) That change of liquid bodies which takes place when they pass to a solid state, by losing the caloric which kept them in a state of fluidity.

CONGELATIVA. (From *congelio*, to congel.) Medicines that inspissate humours, and stop fluxions and rheuma.

CONSERVA. (From *con*, and *genuo*, of the same kind.) Of the same kind; concurring in the same action. It is usually said of the muscles.

CONGESTION. (From *congero*, to amass.) A collection of blood or fluid; a swelling which rises gradually, and ripens slowly, in opposition to that which is soon formed, and soon terminated.

CONGLOBATE GLAND. (From *con-globo*, to gather into a ball.) *Glandula conglobata.* Lymphatic gland. Globate gland. A round gland formed of a contortion of lymphatic vessels, connected together by cellular structure, having neither a cavity nor any excretory duct: such are the mesenteric, inguinal, axillary glands, &c. See *Glands*.

CONGLOMERATE GLAND. (From *conglomerare*, to heap upon one.) *Glandula conglomerata.* A gland composed of a number of glomerate glands, whose excretory ducts all unite into one common duct: such are the salival, parotid glands, &c.

CONGLUTINANTIA. (From *conglutino*, to glue together.) Healing medicines; and such as unite parts disjointed by accident.

CONIS. (Κοκ.) Dust, fine powder, ashes, a nit in the hair, scurf from the head; and sometimes it signifies lime.

CONIUM. (From *conia*, dust, according to Linnæus, or from *novum, circumago*, on account of its inebriating and poisonous quality.) Hemlock.

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*.

2. The pharmacopœial name of the official hemlock. See *Cicuta*.

CONIUM MACULATUM. The systematic name for the *cicuta* of the pharmacopœias. See *Cicuta*.

CONJUNCTIVE MEMBRANE. *Membrana conjunctiva.* The thin, transparent, delicate membrane, that lines the internal superficies of one eyelid, and is reflected from thence, over the anterior part of the bulb, to the edge of the other eyelid. That portion which covers the transparent cornea cannot, without much difficulty, be separated from it. Inflammation of this membrane is called *ophthalmia*.

CONNATUS. (From *con*, and *nascor*, to grow together.) Used much by Hippocrates for what is born with a person; the same with *congenita*.

CONNEXION. See *Articulation*.

CONNUTRITUS. (From *con*, and *nutrior*, to be nourished with.) It is what becomes habitual to a person from his particular nourishment, or what breaks out into a disease in process of time, which gradually had its foundation in the first aliments, as from sucking a distempered nurse, or the like.

CONQUASSATIO. Conquassation. In pharmacy it is a species of comminution, or an operation by which moist concrete substances, as recent vegetables, fruits, the softer parts of animals, &c. are agitated and bruised, till, partly by their proper succulence, or by an effusion of some liquor, they are reduced to a soft pulp.

Consent of parts. See *Sympathy*.

CONSERVA. (From *conservo*, to keep.) A conserve. A composition of some recent vegetable and sugar, beat together into an uniform mass of the consistence of honey; as conserve of hips, orange peel, &c. Conserves are called confections in the last edition of the London Pharmacopœia. See *Confectio*.

CONSERVA ABSINTHII MARITIMI. See *Absinthium maritimum*.

CONSERVA ARI. This is occasionally exhibited as a stimulant and diuretic. See *Arum*.

CONSERVA AURANTII HISPALENSIS. See *Confectio aurantii*.

CONSERVA CYNOSBATI. See *Confectio rose canine*.

CONSERVA LULULÆ. A preparation of wood-sorrel, possessing acid, cooling, and antiseptic qualities. See *Acetocella*.

CONSERVA MENTHÆ. This preparation of mint is given occasionally as a stomachic, in sickness and weakness of the stomach. See *Mentha viridis*.

CONSERVA PRUNI SYLVESTRIS. Astringent virtues are ascribed to this medicine, which is now seldom used but in private formulæ.

CONSERVA ROSÆ. This conserve, rubbed down with water, to which is added

some lemon-juice, forms an excellent drink in hemorrhagic complaints. See *Confectio rose gallica*.

CONSERVA SCILLÆ. A preparation of squills, which affords an excellent basis for an electuary, possessing expectorant and diuretic qualities.

CONSISTENTIA. (From *consisto*, to abide.) The state or acme of a disease. The appearance or state of the humours and excrements.

CONSOLIDA. (So called, *quia consolidandi et conglutinandi vi pollet*; named from its power and use in agglutinating and joining together things broken.) Comfrey.

CONSOLIDA AUREA. *Aurea cordis*. A name of the chamæscistus.

CONSOLIDA MAJOR. See *Symphitum*.

CONSOLIDA MEDIA. *Bugula*. Upright bugloss. Middle consoud. This plant, *Ajuga pyramidalis* of Linnaeus — *tetragono-pyramidalis, villosa, foliis radicalibus maximis*, possesses subadstringent and bitter qualities: and has been recommended in *phthisis, apthæ, and cynanche*.

CONSOLIDA MINOR. See *Prunella*.

CONSOLIDA REALIS. *Calceatrippa*. Many virtues are attributed to this plant, *Delphinium consolida*, of Linnaeus: — *nectaritis monophyllis, caule subultrio*. The flowers are bitter, and a water distilled from them is recommended in ophthalmia. The herb has been administered in calculous cases, obstructed menses, and visceral diseases.

CONSOLIDA SARACENICA. See *Virga aurea*.

Consoud. See *Symphitum*.

Consoud middle. See *Consolida media*.

CONSTIPATION. (From *constipo*, to crowd together.) *Obstipatio*. A person is said to be constive when the alvine excrements are not expelled daily, and when the feces are so hardened as not to receive their form from the impression of the rectum upon them.

CONSTRICIVA. (From *constringo*, to bind together.) *Styptics*.

CONSTRUCTOR. (From *constringo*, to bind together.) A name given to those muscles which contract any opening of the body.

CONSTRUCTOR ALÆ NASÆ. See *Depressor labii superioris aequæ nasi*.

CONSTRUCTOR ANI. See *Sphincter ani*.

CONSTRUCTOR ISTHMI FAUCIUM. *Glosso-staphylinus* of Winslow, Douglas, and Cowper; and *Glosso staphilin* of Dumas. A muscle situated at the side of the entry of the fauces, that draws the *velum pendulum palati* towards the root of the tongue, which it raises at the same time, and with its fellow contracts the passage between the two arches, by which it shuts the opening of the fauces.

CONSTRUCTOR LABIORUM. See *Orbicularis oris*.

CONSTRUCTOR MUSCULUS. See *Buccinator*.

CONSTRUCTOR ORIS. See *Orbicularis oris*.

CONSTRUCTOR PALPEBRARUM. See *Orbicularis palpebrarum*.

CONSTRUCTOR PHARYNGIS INFERIOR. *Crico-pharyngeus*. *Thyro-pharyngeus* of Douglass and Winslow. *Crico-thyropharyngien* of Dumas. A muscle situated on the posterior part of the pharynx. It arises from the side of the thyroid cartilage, near the attachment of the sterno-hyoidæus and thyro-hyoidæus muscles; and from the cricoid cartilage, near the crico-thyroidæus; it is inserted into the white line, where it joins with its fellow, the superior fibres running obliquely upwards, covering nearly one-half of the middle constrictor, and terminating in a point: the inferior fibres run more transversely, and cover the beginning of the œsophagus. Its use is to compress that part of the pharynx which it covers, and to raise it with the larynx a little upwards.

CONSTRUCTOR PHARYNGIS MEDIUS. *Hyo-pharyngeus* and *cephalo-pharyngeus* of Douglas and Winslow. *Chondro-pharyngeus* of Douglas. *Syndesmo-pharyngeus* of Winslow. *Cephalo-pharyngeus* of Winslow and Douglas. *Hyo-glossobasi pharyngien* of Dumas. A muscle situated on the posterior part of the pharynx. It arises from the appendix of the os hyoides, from the corner of that bone, and from the ligament which connects it to the thyroid cartilage; the fibres of the superior part, running obliquely upwards, and covering a considerable part of the superior constrictor, terminate in a point; and is inserted into the middle of the cuneiform process of the os occipitis, before the foramen magnum, and joined to its fellow at a white line in the middle part of the pharynx. This muscle compresses that part of the pharynx which it covers, and draws it and the os hyoides upwards.

CONSTRUCTOR PHARYNGIS SUPERIOR. *Glosso-pharyngeus*. *Mylo-pharyngeus*. *Pterygo-pharyngeus* of Douglas and Winslow, and *Pterigo syndesmo staphili pharyngien* of Dumas. A muscle situated on the posterior part of the pharynx. It arises above, from the cuneiform process of the os occipitis, before the foramen magnum, from the pterygoid process of the sphenoid bone, from the upper and under jaw, near the roots of the last dentes molares, and between the jaws. It is inserted in the middle of the pharynx. Its use is to compress the upper part of the pharynx, and to draw it forwards and upwards.

CONSTRUCTOR VESICÆ URINARIÆ. See *Detrusor urine*.

CONSTRUCTORES PHARYNGEI. Muscles of the œsophagus.

CONTRACTORIUM. Diseases attended with constriction, or spasmodic diseases.

CONSTRINGENTIA. (From *constringo*, to bind together.) Astringent medicines.

CONSUMPTION. (From *consumo*, to waste away.) See *Phthisis*.

CONTABESCENTIA. (From *contabesco*, to pine or waste away.) An atrophy, or nervous consumption.

CONTAGION. (From *contingo*, to meet or touch each other.) *Effluvia*. *Miasma*. *Virus*. *Lues*. Infection. The very subtle particles arising from putrid or other substances, or from persons labouring under contagious diseases, which communicate the disease to others; thus the contagion of putrid fever, the effluvia of dead animal or vegetable substances, the miasma of bogs and fens, the virus of small-pox, lues venerea, &c. &c. There does not appear to be any distinction between contagious and infectious diseases. Would it not be proper to apply the term *contagious* to those which are communicated by contact only, as the venereal disease, itch, &c.; and *infectious*, to those which are caught through the medium of the atmosphere, &c. without contact, as putrid fever, &c.

CONTENSIO. (From *contineo*, to restrain.) It is sometimes used to express a tension or stricture.

CONTINENS FEBRIS. A continual or continent fever, which proceeds regularly in the same tenor, without either intermission or remission. This rarely if ever happens.

CONTINUA FEBRIS. (From *continuo*, to persevere.) A continued fever, attended with exacerbations and slight remissions, but no intermission; sometimes called *assidua*.

CONTORSIO. (From *contorqueo*, to twist about.) A contorsion, or twisting. In medicine this word has various significations, and is applied to iliac passion, to luxation of the vertebrae, head, and back, &c.

CONTRA-APERTURA. (From *contra*, against, and *aperio*, to open.) A counter-opening. An opening made opposite to the one that already exists.

CONTRACTILITY. Expansive elasticity. A property in bodies, the effect of the cohesive power, by which their particles resume their former propinquity when the force ceases which was applied to separate them.

CONTRACTION. (From *contraho*, to draw together.) *Contractura*. *Beriberia*. A rigid contraction of the joints. It is a genus of disease in the class *locales*, and order *dyscinetice* of Cullen. The species are,

1. *Contractura primaria*, from a rigid contraction of the muscles, called also *obstipitas*; a word that, with any other

annexed, distinguishes the variety of the contraction. Of this species be forms four varieties. 1. *Contractura ab inflammatione*, when it arises from inflammation. 2. *Contractura a spasma*, called also *tonic spasm* and *cramp*, when it depends upon spasm. 3. *Contractura ab antagonistas paraliticos*, from the antagonist muscles losing their action. 4. *Contractura ab acrimonia irritante*, which is induced by some irritating cause.

2. *Contractura articularis*, originating from a disease of the joint.

CONTRAFISSURA. (From *contra*, against, and *fendo*, to cleave.) A crack in the skull, opposite to the part on which the blow was given.

CONTRAHESSIA. (From *contraho*, to contract.) Medicines which shorten and strengthen the fibres. Astringents are the only medicines of this nature.

CONTRA-INDICATION. (*Contraindicatio*; from *contra*, against, and *indico*, to shew.) A symptom attending a disease, which forbids the exhibition of a remedy which would otherwise be employed; for instance, bark and acids are usually given in putrid fevers; but if there be difficulty of breathing, or inflammation of the side, or of any viscus, they are contra-indications to their use.

CONTRA-LUNARIS. (From *contra*, and *luna*, the moon.) An epithet given by Diemerbroeck to a woman who conceives during the menstrual discharge.

CONTRA-SEMIN. See *Santonicum*.

CONTRE COU. A species of fracture of the skull, called in Latin *contra-fissura*, in which the fracture happens in that part of the bone opposite to where the blow was received.

CONTRAYERVA RADIX. See *Contrayerva*.

CONTRAYERVA. (From *contra*, against, and *yerva*, a herb. Span. *i. e.* a herb good against poison.) *Drakena*. *Cyperus longus, odoratus, peruvianus*. *Bezoardica radix*. *Dorstenia contrayerva* of Linnaeus.

The contrayerva was first brought into Europe about the year 1581, by Sir Francis Drake, whence its name *Drakena*. It is the root of a small plant found in Peru, and other parts of the Spanish West-Indies. Dr. Houlston observes, that the roots of different species of *dorstenia* are promiscuously gathered and exported for those of the contrayerva, and, as all the species bear a great resemblance to each other they are generally used for medical purposes in this country. The tuberos parts of these roots are the strongest, and should be chosen for use. They have an agreeable aromatic smell; a rough, bitter, penetrating taste; and when chewed, they give out a sweetish kind of acrimony.

It is diaphoretic and antiseptic; for-

merly used in low nervous fevers,¹ and those of the malignant kind; though taken freely, it does not produce much heat. It is, however, now seldom used, though, with the Peruvian bark in decoction, it is occasionally employed in ulcerated sore throats, as a gargle.

Dr. Cullen observes, that this and serpentaria are powerful stimulants; and both have been employed in fevers in which debility prevailed. However, he thinks, wine may always supersede the stimulant powers of these medicines; and that debility is better remedied by the tonic and antiseptic powers of cold and Peruvian bark, than by any stimulants.

By the assistance of heat, both spirit and water extract all its virtues; but they carry little or nothing in distillation; extracts made by inspissating the decoction, retain all the virtues of the root.

The London College forms the compound powder of contrayerva by combining five ounces of contrayerva root with a pound and a half of prepared shells. This powder was formerly made up in balls, and called *lapis contrayervae*, employed in the decline of ardent fevers, and through the whole course of low and nervous ones. The radix serpentaria virginianensis, in all cases, may be substituted for the contrayerva.

CONTRAYERVA ALBA. *Contrayerva Germanorum.* A name for asclepias.

CONTRAYERVA NOVA. Mexican contrayerva. This is the root of the *Peoralea pentaphylla* of Linnæus. It was introduced into Europe after the former, and is brought from Guiana as well as from Mexico. It is but little if any thing inferior to contrayerva.

CONTRAYERVA VIRGINIANA. See *Serpentaria*.

CONTRITIO. The same as comminution.

CONTUSION. (From *contundo*, to knock together.) A bruise, or contused wound.

CONVALESCENCE. That space from the departure of a disease, to the recovery of the strength lost by it.

CONVALLARIA. (From *convallis*, a valley; named from its abounding in valleys and marshes.) The name of a genus of plants in the Linnæan system. Class, *Hexandria*. Order, *Monogynia*.

CONVALLARIA MAJALIS. The systematic name of the lily of the valley. See *Lilium convallium*.

CONVALLARIA POLYGONATUM. The systematic name of Solomon's seal. See *Sigillum Salomonis*.

CONVOLUTA SUPERIORA OSSA. The superior turbinated bones of the nose.

CONVOLUTA INFERIORA OSSA. The lower turbinated bones of the nose.

CONVOLVULUS. (From *convolvo*, to roll together; so named from its spiral

shape, and its twisting round other trees and shrubs.)

1. A name for the *Iliaca passion*.

2. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*, which affords the Jalapa, mechoacana, turbith, and scammony. The whole genus usually abounds with plants containing a milky juice strongly cathartic and caustic.

CONVOLVULUS AMERICANUS. The jalap root.

CONVOLVULUS CANTABRICA. A name for the cantabrica.

CONVOLVULUS COLUMBINUS. The pariera brava.

CONVOLVULUS JALAPA. The systematic name of the jalap plant. See *Jalapium*.

CONVOLVULUS MAJOR ALBUS. The juice of this plant, *Convolvulus sepium* of Linnæus:—*foliis sagittatis postice truncatis pedunculis tetragonis, unifloris*, is violently purgative, and given in dropsical affections. A poultice of the herb, made with oil, is recommended in white swellings of the knee-joint.

CONVOLVULUS MARITIMUS. The brassica maritima, or sea colewort.

CONVOLVULUS MECOACAN. See *Mechoacanæ radix*.

CONVOLVULUS SCAMMONIA. The systematic name of the scammony plant. See *Scammonium*.

CONVOLVULUS SEPIUM. See *Convolvulus major albus*.

CONVOLVULUS SOLDANELLA. The systematic name of the sea convolvulus. See *Brassica marina*.

CONVOLVULUS SYRIACUS. A name for the scammonium.

CONVOLVULUS TURPETHUM. The systematic name of the turbith plant. See *Turpethum*.

CONVULSION. (*Convulsio*; from *convello*, to pull together.) *Hieranosos. Distentio nervorum.* Clonic spasm. A diseased action of muscular fibres, known by alternate relaxations, with violent and involuntary contractions of the muscular parts, without sleep. Cullen arranges convulsion in the class *neuroses*, and order *spasmi*. Convulsions are universal or partial, and have obtained different names, according to the parts affected, or symptoms; as the *risus sardonicus*, when the muscles of the face are affected; St Vitus's dance, when the muscles of the arm are thrown into involuntary motions with lameness and rotations. The hysterical epilepsy, or other epilepsies, arising from different cause, are convulsive diseases of the universal kind; the muscles of the globe of the eye, throwing the eye into involuntary distortions in defiance to the direction of the will, are instances of partial convulsion. The muscles principally

affected in all species of convulsions, are those immediately under the direction of the will; as those of the eyelids, eye, face, jaws, neck, superior and inferior extremities. The muscles of respiration, acting both voluntarily and involuntarily, are not unfrequently convulsed; as the diaphragm, intercostals, &c. The more immediate causes of convulsions are, 1. Either mental affection, or any irritating cause exciting a greater action in the arterial system of the brain and nerves. 2. An increase of nervous energy, which seems to hold pace or be equi-potent with the increased arterial energy, excited in the brain. 3. This increased energy, conveying its augmented effects, without the direction of the will, to any muscles destined to voluntary motion, over-irritates them. 4. The muscles, irritated by the increased nervous energy and arterial influx, contract more forcibly and involuntarily by their excited vis insita, conjointly with other causes, as long as the increased nervous energy continues. 5. This increased energy in the nervous system may be excited either by the mind, or by any acrimony in the blood, or other stimuli sufficiently irritating to increase the arterial action, nervous influence, and the vires insitæ of muscles. 6. After muscles have been once accustomed to act involuntarily, and with increased action, the same causes can readily produce the same effects on those organs. 7. All parts that have muscular fibres may be convulsed. 8. The sensations in the mind most capable of producing convulsions, are timidity, horror, anger, great sensibility of the soul, &c.

CONVULSIO ABDOMINIS. Convulsion of the muscles of the belly.

CONVULSIO CANINA. A wry mouth.

CONVULSIO CEREALIS. Cereal convulsion, is a singular disorder of the spasmodic convulsive kind, not common to this country, but mentioned by Cælius under this title, from the peculiar tingling and formication perceived in the arms and legs. *Motus spasmodicus* of Hoffman. It is endemic in some places in Germany, but more a rural than urbanical disorder, said to arise from the use of spoiled corn.

CONVULSIO CLONICA. Convulsion with alternate relaxation.

CONVULSIO GRAVIDARUM. Convulsion of pregnant women.

CONVULSIO HABITUALIS. The chorea St. Viti.

CONVULSIO HEMITONOS. Convulsion approaching to tetanus.

CONVULSIO AB INANITIONE. Convulsion from inanition.

CONVULSIO INDICA. Tetanus.

CONVULSIO INTERMITTENS. Convulsion occurring in paroxysms.

CONVULSIO NEPHRALGICA. Convulsion from stones in the kidneys.

CONVULSIO AB ORANISMO. Convulsion from self pollution.

CONVULSIO RAPHANIA. Spasmodic painful disease of the joints.

CONVULSIO TONICA. Common or permanent convulsion.

CONVULSIO UTERI. Abortion.

CONYZA. (From *zōis*, dust; because its powder is sprinkled to kill fleas in places where they are troublesome.) The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia superflua*.

CONYZA ETHIOPICA. A name for the elichrysium.

CONYZA CERULEA. The herb flea-bane. This acrid plant is exhibited on the continent in some diseases of the chest.

CONYZA MAJOR. *Conyza major vulgaris.* *Mas theophrasti.* *Conyza major* of Dioscorides. Greater flea-bane.

CONYZA MEDIA. *Arnica Suedensis.* *Inula dysenterica* of Linnæus:—*foliis amplexicaulibus, cordato-oblongis; caule villosa, paniculato, squamis calycinis, setaceis.* An acrid, subaromatic plant, possessing antidyenteric virtues. It is sometimes called *arnica spuria*.

CONYZA MINOR. *Conyza minor flore globoso.* *Pulicaria.* *Conyza minima et media.* Small flea-bane. The *inula dysenterica*. The chief use of all the flea-banes is to destroy fleas and gnats, by burning. They are occasionally used as an antipsoric remedy.

COOPERFORIA. (From *co operio*, to cover over.) *Cartilago thyroidea.* Called also *abicum*.

COOSTRUM. The centre of the diaphragm.

COPAIBA. See *Balsamum copaiba*.

COPAIFERA. (From *Copaiva*, the Indian name, and *fero*, to bear.) The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Digynia*.

COPAIFERA OFFICINALIS. The systematic name of the plant from which the *Copaiva* balsam is obtained. See *Balsamum Copaivæ*.

COPAL. (The American name of all clear odoriferous gums.) Gum copal. This resinous substance is imported from Guinea, where it is found in the sand on the shore. It is of a yellow colour, faintly glistening, imperfectly transparent, and apt to break with a conchoidal fracture. It is tasteless, and, while cold, inodorous. It is used, dissolved in rectified spirit of wine, in laxities of the gums, with the same views as mastich.

Copaiva balsam. See *Balsamum Copaivæ*.

COPELLA. See *Cupella*.

COPHER. A name for camphor.

COPHOS. (*Kopos*, dumb.) Deaf or dumb. Also a dulness in any of the senses.

COPHOSIS. (From *copos*, deaf.) A difficulty of hearing. It is often symptomatic of some disease. See *Dyscoëcia*.

COPPER. (*Quasi* *as* *Cyprium*; so named from the island of Cyprus, whence it was formerly brought.) A metal found in the earth, in various states. It is found native, (*native copper*), possessing the red colour, malleability, and many of its other properties; it is, however, not quite pure, but generally mixed with a minute portion of gold, or silver.

This ore is found of an indeterminate figure, in solid and compact masses; sometimes in plates and threads, which assume a variety of forms. It crystallizes in cubes. It then is flexible. It has much metallic lustre. It is found in many parts of Europe.

Copper ores are very numerous. Copper combined with oxygen, forms *oxyd of copper*, or the *earthy copper ore*, (*mountain blue*.) United to carbonic acid, it constitutes the hepatic copper ores, (*mountain green*.) &c. The compact ore of this kind is termed *malachite*. It generally exhibits a very fine grass-green, emerald green, or apple-green colour. It is found in solid masses of an indeterminate shape. It has often a beautiful satin-like appearance, or silky lustre.

Copper also exists mineralized by the muriatic acid, sulphuric acid, arsenic acid, &c. Copper mineralized with sulphur is called *vitreous copper ore*. Its colour is generally lead-gray. Combined with sulphur and iron, it forms the *azure copper ore*, and all the varieties of copper pyrites. Mineralized with sulphur, arsenic, iron, and zinc, it constitutes the *brown or blende copper ores*, of which there are many varieties. Copper mines are abundant in Britain, Germany, &c.

Properties of Copper.—Pure copper is of a rose-red colour, very sonorous, very tenacious, ductile, and malleable; of a considerable compactness; moderately hard and elastic. Its texture is granulated, and subject to blisters. It crystallizes in quadrilateral pyramids. Its specific gravity is between 7.788 and 8.584. When rubbed, it emits a disagreeable odour. It melts at 27° of Wedgwood's pyrometer. At a higher temperature, it burns with a beautiful green flame. It is a good conductor of caloric, of electricity, and of galvanism. Exposed to the air it becomes brown, and at last green, by absorbing carbonic acid. When heated, it turns blue, yellow, violet, and brown. It readily fuses with phosphorus and unites to sulphur, when finely divided by mere trituration. It does not decompose water at the temperature of ignition. It is acted on by the greater number of the acids. Nitric acid acts on copper with great vehemence. Sulphuret of potash combines with it in the dry and in the humid way. It is capable of alloying with the greater number of the metals. With zinc it forms the compound metals

called brass, pinchbeck, and others: with tin it forms bell-metal and bronze. It unites to the earths merely in vitrification. Liquid ammonia causes it to oxydate quickly when air is admitted. It decomposes muriate of ammonia, and red sulphurated oxyd of mercury, by heat. It is poisonous to the human constitution.

Method of obtaining Copper.—Copper is procured from its ores, by different processes according to the nature of those ores. If they contain much sulphur, after being pounded and washed, they are roasted in the open air to dispel the sulphur. The ore is afterwards roasted once or twice more, and is melted in an open fire into a mass, called a *mat of copper*. In this state it still contains a large quantity of sulphur, which the workmen continue to expel by repeated roastings and fusion, till the metal acquires a certain degree of purity, and is called black copper, which is somewhat malleable, but still contains sulphur, iron, and in general some other impurities. In order to get entirely rid of these, the copper is hastily fused with three times its weight of lead. The lead unites with the copper, and expels the iron; and the rest of the metals which happen to be mixed with the copper are thus expelled. The copper is afterwards refined, by keeping it heated in crucibles for a considerable time, so that it may throw up all the foreign substances it still contains in the form of scoriae. It is examined from time to time by immersing iron rods into it, which become coloured with a small quantity of copper, and its purity is judged of by the brilliant redness of these specimens.

COPPERAS. A name given to blue, green, and white vitriol.

COPRAGOSA. (From *κωρος*, dung, and *αγω*, to bring away.) *Copragagum*. The name of a gently purging electuary, mentioned by Rulandus.

COPRIEXESIS. (From *κωρος*, excrement, and *εγω*, to vomit.) A vomiting of feces.

COPROCRITICA. (From *κωρος*, excrement, and *αγω*, to separate.) Mild cathartic medicines.

COPROPHORIA. (From *κωρος*, excrement, and *αγω*, to bring away.) A purging.

COPROS. (*Κωρος*.) The feces, or excrements.

COPROSTASIA. (From *κωρος*, feces, and *στασις*, to remain.) Costiveness, or a constriction of the belly.

COPARTION. (*Κορτή*, a small cake.) *Copartarium*. A medicine in the shape of a very small cake, directed for disorders of the aspera arteria and lungs, and for many other intentions, by the ancients.

COPTE (*Κορτή*, a small cake.) It was the form of a medicine used by the ancients; also a cataplasm generally made of

vegetable substances, and applied externally to the stomach, and internally on many occasions.

COPULA. (*Quasi compula*; from *compello*, to restrain.) A name for a ligament.

COQUENTIA. (From *coquo*, to digest.) Medicines which promote concoction.

COR.

1. The heart. See *Heart*.

2. Gold.

3. An intense fire.

CORACINE. (From *κοράξ*, a crow; so named from its black colour.) A name for a lozenge, quoted by Galen from Asclepiades.

CORACO-BRACHIALIS. (From *κοράξ*, a crow, and *βραχίων*, the arm.) *Coraco-humeral* of Dumas. *Coraco-brachius*. A muscle, so called from its origin and insertion. It is situated on the humerus, before the scapula. It arises, tendinous and fleshy, from the fore-part of the coracoid process of the scapula, adhering, in its descent, to the short head of the biceps; inserted, tendinous and fleshy, about the middle of the internal part of the os humeri, near the origin of the third head of the triceps, called *brachialis externus*, where it sends down a thin, tendinous expansion to the internal condyle of the os humeri. Its use is to raise the arm upwards and forwards.

CORACO-HYOIDEUS. (*Coraco-hyoideus*, sc. *musculus*, *κοράξο-υοειδής*: from *κοράξ*, a crow, and *υοειδής*, the bone called *hyoides*.) See *Omo-hyoideus*.

CORACOID. (*Coracoides*; from *κοράξ*, a crow, and *οὐδός*, resemblance; because it is shaped like the beak of a crow.) A name given to a process on the upper and anterior part of the scapula.

Coral. See *Corallium*.

CORALLINA. (Dim. of *corallium*; from *κορη*, a daughter, and *αλς*, the sea; because it is generated in the sea.) *Muscus maritimus*. *Corallina officinalis*. *Corallina alba*. Sea coralline, and white wormseed. A marine production, resembling a small plant without leaves, consisting of numerous brittle cretaceous substances, friable betwixt the fingers, and crackling between the teeth. Powdered, it is administered to children as an anthelmintic.

CORALLINA CORSICANA. *Helminthocorton*. *Conferva helmintho-cortici*. *Corallina rubra*. *Corallina melito-corton*. *Lemitho-corton*. Corsican worm weed. *Fucus helmintho-corton* of De La Tourette. This plant has gained great repute in destroying all species of intestinal worms. Its virtues are extolled by many; but impartial experimentalists have frequently been disappointed of its efficacy. The Geneva Pharmacopœia directs a syrup to be made of it.

CORALLINA MELITO-CORTON. See *Corallina corsicana*.

CORALLINA RUBRA. See *Corallina corsicana*.

Coralline. See *Corallina*.

Coralline, Corsican. See *Corallina Corsicana*.

CORALLIUM ALBUM. A hard, white, calcareous, brittle substance; the nidus of the *Madrepora oculata*. Class, *Vermes*. Order, *Lithophyta*. It is sometimes exhibited as an absorbent earth.

CORALLIUM RUBRUM. (From *κορη*, a daughter, and *αλς*, the sea; so named because it is generated in the sea.) *Acma*. *Azur*. The red coral is mostly employed medicinally. It is a hard, brittle, calcareous substance, resembling the stalk of a plant, and is the habitation of the *Isis nobilis*. Class, *Vermes*. Order, *Zoophyta*. When powdered, it is exhibited as an absorbent earth to children; but does not appear to claim any preference to common chalk.

CORALLODENDRON. (From *κοράλλον*, coral, and *δένδρον*, a tree; resembling in hardness and colour a piece of coral.) The coral-tree of America; antivenereal.

CORALLOIDES SEPTUOL. Tooth or coralwort; corroborant.

CORALLOIDES FURGUS. (From *κοράλλον*, coral, and *ωδός*, likeness.) *Erotulus*. *Clavaria coralloides* of Linnaeus. It is said to be corroborant and astrigent.

COCHURON. (From *κορη*, the pupil of the eye, and *κορεω*, to purge; so called because it was thought to purge away rheum from the eyes.) The herb pimpinell, or chickweed.

CORDA. See *Chordee*.

CORDA TYMPANI. The portio dura of the seventh pair of nerves, having entered the tympanum, sends a small branch to the stapes, and another more considerable one, which runs across the tympanum from behind forwards, passes between the long leg of the incus and the handle of the malleus, then goes out at the same place where the tendon of the anterior muscle of the malleus enters. It is called *corda tympani*, because it crosses the tympanum as a cord crosses the bottom of a drum. Dr. Monro thinks that the *corda tympani* is formed by the second branch of the fifth pair, as well as by the portio dura of the seventh.

CORDÆ WILLISII. See *Dura mater*.

CORDIALS. Medicines are generally so termed, which possess warm and stimulating properties, and that are given to raise the spirits.

CORDIA MYXA. *Sebastina*. The systematic name of the Sebesten plant. See *Sebesten*.

CORDINEMA. (From *κορη*, the head, and *δυνας*, to move about.) A head-ach, attended with a vertigo.

CORDOLIUM. (From *cor*, the heart, and *dolor*, pain.) A name formerly applied to cardialgia, or heart-burn.

CORE. (*Κορη*.) The pupil of the eye.

COREMATA. (From *κορεω*, to cleanse.) Medicines for cleansing the skin.

Coriander. See *Coriandrum*.

CORIANDRUM. (From *coron*, a pupil, and *and*, a man; because of its roundness, like the pupil of a man's eye; or probably so called from *coron*, *cimezz*, a bug, because the green herb, seed and all, stinks intolerably of bugs.) *Coriander*.

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digymia*.

2. The pharmacopœial name of the official coriander. *Cassiope*. *Corianon*. The *Coriandrum sativum* of Linnæus:—*fructibus globosis*. This plant is a native of the South of Europe, where, in some places, it is said to grow in such abundance as frequently to choke the growth of wheat and other grain. From being cultivated here as a medicinal plant, it has for some time become naturalized to this country, where it is usually found in corn-fields, the sides of roads, and about dunghills. Every part of the plant, when fresh, has a very offensive odour, but, upon being dried, the seeds have a tolerable grateful smell, and their taste is moderately warm and slightly pungent. They give out their virtue totally to rectified spirit, but only partially to water. In distillation with water, they yield a small quantity of a yellowish essential oil, which smells strongly and pretty agreeably of the coriander.

Dioscorides asserts, that the seeds, when taken in a considerable quantity, produce deleterious effects; and, in some parts of Spain and Egypt, where the fresh herb is eaten as a cordial, instances of fatuity, lethargy, &c. are observed to occur very frequently; but these qualities seem to have been unjustly ascribed to the coriander; and Dr. Withering informs us, that he has known six drachms of the seeds taken at once, without any remarkable effect. These seeds, and indeed most of those of the umbelliferous plants, possess a stomachic and carminative power. They are directed in the infusum amarum, the infusum sennæ tartarizatum, and some other compositions of the pharmacopœias; and, according to Dr. Cullen, the principal use of these seeds is, "that infused along with senna, they more powerfully correct the odour and taste of this than any other aromatic that I have employed, and are, I believe, equally powerful in obviating the griping that senna is very ready to produce."

CORIANDRUM SATIVUM. The systematic name of the plant called *coriandrum* in the pharmacopœias. See *Coriandrum*.

CORIANON. See *Coriandrum*.

CORIS. (From *coron*, to cleave, or cut; so called because it was said to heal wounds.) The herb hypericum.

CORIS LUTEA. *Coris legitima cretica*. The hypericum saxatile, or bastard St. John's wort.

CORIS MONAPHELIENSIS. This plant is intensely bitter and nauseous, but apparently an active medicine, and employed, it is said, with success in syphilis.

CORK. The bark of the *Quercus ruber* of Linnæus, formerly employed as an astringent, but now disused. It affords an acid.

CORN. *Clavus*. A hardened portion of cuticle, produced by pressure: so called because a piece can be picked out like a corn of barley. Corns are sometimes connected with the periosteum.

CORNACHINI PULVIS. Scammony, anti-mony, and cream of tartar.

CORNEA OPAKA. The sclerotic membrane of the eye is so called, because it is of a horny consistence and opaque. See *Sclerotic membrane*.

CORNEA TRANSPARENS. *Sclerotica ceratoidea*. The transparent portion of the sclerotic membrane, through which the rays of light pass, is so called, to distinguish it from that which is opaque. See *Sclerotic membrane*.

CORNESTA. A chemical retort.

Cornflower. See *Cyanus*.

CORNICULA. (From *cornu*, a horn.) A cupping instrument, made of horn.

CORNICULARIS. (From *cornu*, a horn.) Shaped like a horn; the coracoid process.

CORN-SALLAD. This is the *Valeriana locusta* of Linnæus. It is cultivated in our gardens, and eaten amongst the early sallads. It is a very wholesome succulent plant, possessing anti-scorbutic and gently aperient virtues.

CORNU AMMONIS. *Cornu arietis*. When the pes hippocampi of the human brain is cut transversely through, the cortical substance is so disposed as to resemble a ram's horn. This is the true cornu ammonis, though the name is often applied to the *pes hippocampi*.

CORNU ARIETIS. See *Cornu ammonis*.

CORNU. *Cornu cervi*. Hartshorn. The horns of several species of stag, as the *cervus aloe*, *cervus dama*, *cervus elaphus*, and *cervus tarandus*, are used medicinally. Boiled, they impart to the water a nutritious jelly, which is frequently served at table. Hartshorn jelly is made thus:—Boil half a pound of the shavings of hart's-horn, in six pints of water, to a quart; to the strained liquor add one ounce of the juice of lemon, or of Seville orange, four ounces of mountain wine, and half a pound of sugar; then boil the whole to a proper consistence. The chief use of the horns is for calcination, and to afford the *liquor volatilis cornu cervi* and carbonate of ammonia.

CORNU CERVİ CALCINATUM. See *Cornu ustum*.

CORNU USTUM. Burnt hartshorn shavings possess absorbent, antacid, and astringent properties, and are given in form of decoction.

tion, as a common drink in diarrheas, pyrosis, &c.

CORNUA. Warts. Horny excrescences, which mostly form on the joints of the toes. Similar diseased productions have been known to arise on the head, and other parts.

CORNUA UTERI. *Plectene*. In comparative anatomy, the horns of the womb; the womb being in some animals triangular, and its angles resembling horns.

CORNUMUSA. A retort.

CORNUS. 1. The name of a genus of plants in the Linnæan system. Class, *Tetrandria*. Order, *Monogymia*.

2. The pharmacopœial name of the cornel tree. *Cornus sanguinea* of Linneus. The fruit is moderately cooling and astringent.

CORNUTA. (From *cornu*; from its resemblance to a horn.) A retort.

CORONA CILIARIS. The ciliar ligament.

CORONA GLANDIS. The margin of the glans penis.

CORONA IMPERIALIS. A name for crown-imperial. The Turks use this plant as an emetic. The whole plant is considered poisonous.

CORONA REGIA. The melilotus.

CORONA SOLIS. Sun-flower. Called also *chimalati*. The *Helianthus annuus* of Linneus. It has been noticed as heating, and an agreeable food. The seeds are made into bread.

CORONA VENERIS. Venereal blotches on the forehead are so termed.

CORONAL SUTURE. (From *corona*, a crown, or garland; so named because the ancients wore their garlands in its direction.) *Sutura coronalis*. *Sutura arcuata*. The suture of the head, that extends from one temple across to the other, uniting the two parietal bones with the frontal.

CORONARIUS STOMACHICUS. Part of the eighth pair of nerves.

CORONARY VESSELS. *Vasa coronaria*. The arteries and veins of the heart and stomach. The term coronary is here given from *corona*, a crown, surrounding any part in the manner of a crown.

CORONARY LIGAMENTS. (From *corona*, a crown.) Ligaments uniting the radius and ulna. The term ligamentum coronarium is also applied to a ligament of the liver.

CORONE. (*Korona*, a crow; so named from its supposed likeness to a crow's bill.) The acute process of the lower jaw bone.

CORONOID. (*Coronoides*; from *zagon*, a crow, and *oides*, likeness.) Processes of bones are so called, that have any resemblance to a crow's beak, as *coronoides apophysis ulne*, *coronoides apophysis maxille*.

CORONORUS. (From *zagon*, a carrion crow, and *rus*, foot; the plant being said to resemble a crow's foot.) *Coronopodium*. *Cornu*. *Cerbinum*. *Stella terre*. *Planta*

go. Buck's-horn plantain. The *Plantago coronopus* of Linneus. Its medical virtues are the same as those of the other plantains.

CORPORA ALBICANTIA. Corpora albicantia Willisii.

CORPORA CAVERNOSA CLITORIDIS. Two hollow crura, forming the clitoris.

CORPORA CAVERNOSA PENIS. Two spongy bodies that arise, one from each ascending portion of the ischium, and form the whole bulk of the penis above the urethra, and terminate obtusely behind its glans. See *Præis*.

CORPORA FIMBRIATA. The flattened terminations of the posterior crura of the fornix of the brain, which turn round into the inferior cavity of the lateral ventricle, and end in the *pedes hippocampi*.

CORPORA LOBOSA. Part of the cortical part of the kidney.

CORPORA NERVO-SPONGIOSA. The corpora cavernosa penis.

CORPORA NERVOSA. The corpora cavernosa clitoridis.

CORPORA OLIVARIA. The two external prominences of the medulla oblongata, that are shaped somewhat like an olive.

CORPORA PYRAMIDALIA. Two internal prominences of the medulla oblongata, which are more of a pyramidal shape than the former.

CORPORA QUADRIGEMINA. See *Tubercula quadrigemina*.

CORPORA STRIATA. So named from their appearance. See *Cerebrum*.

CORPUS ANNULARE. A synonym of *pons Varolii*. See *Pons Varolii*.

CORPUS. The body. Many parts and substances are also distinguished by this name: as *corpus callosum*, *corpus luteum*, &c. See also *Body*.

CORPUS CALLOSUM. *Commissura magna cerebri*. The white medullary part joining the two hemispheres of the brain, and coming into view under the falx of the dura mater when the hemispheres are drawn from each other. On the surface of the *corpus callosum* two lines are conspicuous, called the *raphe*.

CORPUS GLANDULOSUM. The prostrate gland.

CORPUS LUTEUM. The granulous papilla which is found in that part of the ovary of females, from whence an ovum had proceeded; hence their presence determines that the female has been impregnated; and the number of the *corpora lutea* corresponds with the number of impregnations. It is, however, asserted by a modern writer, that *corpora lutea* have been detected in young virgins, where no impregnations could possibly have taken place.

CORPUS MUCOSUM. See *Retis mucosum*.

CORPUS PAMPINIFORME. (*Pampiniformis*; from *pampinus*, a tendril, and *forma*, likeness, resembling a tendril.) *Corpus pyramidale*. Applied to the spermatic chord, and thoracic duct; also to the plexus of veins surrounding the spermatic artery in the cavity of the abdomen.

CORPUS RETICULARE. See *Rete mucosum*.

CORPUS SESAMOIDEUM. A little prominence at the entry of the pulmonary artery.

CORPUS SPONGIOSUM URETHRÆ. *Substantia spongiosa urethræ.* *Corpus spongiosum penis.* This substance originates before the prostate gland, surrounds the urethra and forms the *bulb*; then proceeds to the end of the corpora cavernosa, and terminates in the *glans penis*, which it forms.

CORPUS VARICOSUM. The spermatic chord.

CORRAGO. (From *cor*, the heart; it being supposed to have a good effect in comforting the heart.) See *Borago*.

CORRE. (From *corra*, to shave.) The temples. That part of the jaws where the beard grows, and which is used to shave.

CORROBORANTS. (*Corroborantia*, sc. *medicamenta*.) Medicines, or whatever gives strength to the body, as bark, wine, beef, cold bath, &c. &c. See *Tonica*.

Corrosive sublimate. See *Oxyurias hydrogryi*.

CORROSIVES. (*Corrosiva*, sc. *medicamenta*; from *corrodo*, to eat away.) See *Escharotica*.

CORRUGATOR SUPERCILII. (From *corrugo*, to wrinkle.) *Musculus supercilii* of Winslow. *Musculus frontalis verus*, sc. *corrugator cæterii* of Douglas, and *Cutaneus superciliorum* of Dumas. A small muscle situated on the forehead. When one muscle acts, it is drawn towards the other, and projects over the inner canthus of the eye. When both muscles act, they pull down the skin of the forehead and make it wrinkle, particularly between the eyebrows.

CORTEX. This term is generally, though improperly, given to the Peruvian bark. It applies to any rind or bark.

CORTEX ANGELINÆ. The bark of a tree growing in Grenada. A decoction of it is recommended as a vermifuge. It excites tormina, similar to jalap, and operates by purging.

CORTEX ANGUSTURÆ. See *Angustura cortex*.

CORTEX ANTISCORBUTICUS. The canella alba.

CORTEX AROMATICUS. The canella alba.

CORTEX BELA-AYE. See *Bela-aye cortex*.

CORTEX CANELLÆ MALABRICÆ. See *Cassia lignea*.

CORTEX CARDINALIS DE LUGO. The Peruvian bark was so called, because the

Cardinal Lugo had testimonials of above a thousand cures performed by it in the year 1653.

CORTEX CEREBRI. The cortical substance of the brain.

CORTEX CHINÆ REGIUS. See *Cinchona*.

CORTEX CHINÆ SURINAMENSIS. This bark is remarkably bitter, and preferable to the other species in intermittent fevers.

CORTEX CHINCHINÆ. See *Cinchona*.

CORTEX NEUTHERIÆ. See *Coccarilla cortex*.

CORTEX GEOFFROYI & JAMAICENSIS. Bulge-water-tree bark. The bark of the *Geoffroya Jamaicensis*, *inermis foliolis lanceolatis*, of Swartz. It is principally used in Jamaica, and with great success, as a vermifuge.

CORTEX LAVOLA. The bark bearing this name is supposed to be the produce of the tree which affords the *animum stellatum*. Its virtues are similar.

CORTEX MASSELANICUS. See *Winteranus cortex*.

CORTEX MASSOY. The produce of New Guinea, where it is beaten into a pulsatious mass with water, and rubbed upon the abdomen to allay tormina of the bowels. It partakes of the smell and flavour of cinnamon.

CORTEX PATRUM. The Peruvian bark.

CORTEX PERUVIANUS. See *Cinchona*.

CORTEX PERUVIANUS FLAVUS. See *Cinchona*.

CORTEX PERUVIANUS RUBER. See *Cinchona*.

CORTEX POCCKEBÆ. This bark is sent from America; and is said to be serviceable in diarrhoea, dysenteries, and hepatic fluxes.

CORTEX QUASSIÆ. See *Quassia*.

CORTEX WINTERANUS. See *Winteranus cortex*.

CORTICAL. Cerebrous substance. The external substance of the brain is of a darker colour than the internal, and surrounds the medullary substance, as the bark does the tree; hence it is termed *cortical*. See also *Kidney*.

CORTUSA. The plant self-heal; bear's ear; sanicle. It is expectorant.

CORU CANARICA. A quince-like tree of Malabar; it is antydysenteric.

CORYLUS. (Derivation uncertain; according to some, from *cupra*, a walnut.) 1. The name of a genus of plants in the Linnæan system. Class, *Monocia*. Order, *Polymardia*.

2. The pharmacopœial name of the hazel-tree. The nuts of this tree are much eaten in this country; they are hard of digestion, and often pass the bowels very little altered; if however, they are well chewed, they give out a nutritious oil. An oil is also obtained from the wood of this tree, *Corylus avellana* of Linnaeus:—*stipulis ovatis, obtusis*, which is efficacious

against the tooth-ach, and is said to kill worms.

CORYLUS AVELLANA. The hazel-nut tree. See *Corylus*.

CORYPH. (Κορυφή.) The vertex of the head. The inner parts of the nails.

CORYZA. (Coryza, κορυζα: from *κορυ*, the head, and *ζω*, to boil.) An increased discharge of mucus from the nose. See *Catarrh*.

CORYZA CATARRHALIS. A catarrh from cold.

CORYZA FEBRICOSA. A catarrh with fever.

CORYZA PNEUMATORRHOÏA. A catarrh, with much discharge of mucus.

CORYZA PURULENTA. A catarrh, with discharge of matter.

CORYZA VARIOLOSA. A catarrh accompanying small-pox.

CORYZA VIRULENTA. A catarrh, with discharge of acrid mucus.

COSCULIA. The grains of kermes.

COSMETIC. A term applied to remedies against blotches and freckles.

COSMOS. *Rythmus*. A regular series. In Hippocrates it is the order and series of critical days.

COSMIS. *Cossi*. A worm that breeds in wood; also a little tubercle in the face, like the head of a worm.

COSMUM. A malignant ulcer of the nose mentioned by Paracelsus.

COSTA. (*A custodiendo*; because the ribs surround and defend the vital parts.) A rib. The ribs are four-and-twenty in number, twelve on each side of the thorax. See *Ribs*.

COSTA PULMONARIA. *Costa herba*. A name of the herb hawkweed.

COSTO-HYOIDEUS. (From *costa*, a rib, and *hyoideus*, belonging to the hyoidal bone.) A muscle so named from its origin and insertion. See *Omo-hyoideus*.

COSTUS. (From *kasta*, Arabian.) The name of a genus of plants in the Linnæan system. - Class, *Monandria*. Order, *Mono-gynia*.

COSTUS AMARUS. See *Costus arabicus*.

COSTUS ARABICUS. *Costus indicus*, *amarus*, *dulcis*, *orientalis*. Sweet and bitter costus. The root of this tree, *Costus arabicus* of Linnæus, possesses bitter and aromatic virtues, and is considered as a good stomachic. Formerly there were two other species, the *bitter* and *sweet*, distinguished for use. At present, the Arabic only is known, and that is seldom employed. It is, however, said to be stomachic, diaphoretic, and diuretic.

COSTUS CORTICOSUS. The canella alba.

COSTUS HORTORUM MINOR. The ageratum.

COSTUS NIGRA. The artichoke.

COSTYLE. (Κοτύλη, the name of an old measure.) The socket of the hip-bone.

COTABONIUM. A word coined by Paracelsus, implying a liquor into which all bodies, and even their elements, may be dissolved.

COTIS. (From *κοτήν*, the head.) The back part of the head; sometimes the hollow of the neck.

COTULA FETIDA. (*Cotula*, dim. of *cos*, a whetstone, from the resemblance of its leaves to a whetstone; or from *κόλλη*, a hollow.) *Chamæmelum fatidum*. Mayweed. Stinking camomile. This plant, *Anthemis cotula* of Linnæus:—*receptaculis conicis, paleis sataceis, seminibus nudis*, has a very disagreeable smell; the leaves, a strong, acrid, bitterish taste; the flowers, however, are almost insipid. It is said to have been useful in hysterical affections, but is very seldom employed.

COTYLOID CAVITY. (*Cotyloides*; from *κοτύλη*, the name of an old measure, and *οἶος*, resemblance.) The acetabulum. See *innominatum os*.

COTYLOIDES. See *Cotyloid cavity*.

COUCHING. A surgical operation that consists in removing the opaque lens out of the axis of vision, by means of a needle, constructed for the purpose. There are two couching needles, which now seem to be preferred to all others; the one used by Mr. Hey, and that employed by Professor Scarpa.

Couch-grass. See *Græmen caninum*.

COUGH. *Tussis*. A sonorous concussion of the thorax, produced by the sudden expulsion of the inspired air.

COUM. The meadow-saffron.

COUNTER-OPENING. *Contra-apertura*. An opening made in any part of an abscess opposite to one already in it. This is often done in order to afford a readier egress to the collected pus.

COUT DE SOLEIL. An erysipelas from a scorching sun.

COURAP. (Indian.) A distemper very common in Java, and other parts of the East-Indies, where there is a perpetual itching and discharge of matter. It is a herpes on the axilla, groins, breast, and face.

COURBAIL. The tree which produces the gum anime. See *Anime*.

COURONDI. An evergreen tree of India, said to be antidyseric.

COURROY MOELLI. A shrub of India, said to be antivenomous.

COUSCOS. An African food, much used about the river Senegal. It is a composition of the flower of millet, with some flesh, and what is there called lalo.

Cowhage. See *Dolichos*.

Cow-itch. See *Dolichos*.

COWPER'S GLANDS. (*Cowperi glandule*; named from Cowper, who first described them.) Three large muciparous glands of the male, two of which are situated before the prostate gland under the

accelerator muscles of the urine, and the third more forward, before the bulb of the urethra. They excrete a fluid, similar to that of the prostate gland, during the venereal orgasm.

COFFERI GLANDULÆ. See *Compton's glands*.

COVOLAM. The *Cratæva marmelos* of Linnæus, whose fruit is astringent whilst unripe; but when ripe, of a delicious taste. The bark of the tree strengthens the stomach, and relieves hypochondriac languors.

COXA. The ischium is sometimes so called, and sometimes the os coceygi.

COXEDIX. (From *coxa*, the hip.) The ischium; the hip-joint.

CRABLOUSE. A species of pediculus which infests the axillæ and pudenda.

CRAB-TAWS. A name in Jamaica for a kind of ulcer on the soles of the feet, with callous lips, so hard that it is difficult to cut them.

CRAMBE. (*Κραμβή*, the name given by Dioscorides, Galen, and others, to the cabbage; the derivation is uncertain.) The name of a genus of plants in the Linnæan system. Class, *Tetradynamia*. Order, *Siliculosa*. Cabbage.

CRAMBE MARITIMA. The systematic name for the sea-coal. See *Sea-coal*.

CRAMP. (From *krampen*, to contract. Germ.) A spasm of a muscle or muscles.

Cranesbill, bloody. See *Geranium sanguinum*.

CRANIUM. (*Κρανίον*, quasi *caput*; from *caput*, the head.) The skull, or superior part of the head. (See *Caput*.)

CRANTERES. (From *crantæ*, to perform.) A name given to the sapientizæ dentes and molares, from their office of masticating the food.

CRAPULA. (*Κραμπυλα*.) A surfeit; drunkenness.

CRASIS. (From *κρασύνω*, to mix.) Mixture. A term applied to the humours of the body, when there is such an admixture of their principles as to constitute a healthy state: hence, in dropsies, scurvy, &c. the crasis, or healthy mixture of the principles of the blood, is destroyed.

CRASPEDON. (*Κρασπεδόν*, the hem of a garment; from *κρίνω*, to hang down.) A relaxation of the uvula, when it hangs down in a thin, long membrane, like the hem of a garment.

CRASSAMENTUM. (From *crassus*, thick.) See *Blood*.

CRASSULA. (From *crassus*, thick: so named from the thickness of its leaves.) See *Faba crassa*.

CRATEGUS. (From *κρατός*, strength: so called from the strength and hardness of its wood.) The wild service-tree, whose virtues are astringent.

CRATICULA. (From *crates*, a hurdle.)

The bars or grate which covers the ash-hole in a chymical furnace.

Cream of tartar. See *Supertartarus potassæ*.

CREMASTER. (From *κρεμαίνω*, to suspend.) A muscle of the testicle, by which it is suspended, and drawn up and compressed, in the act of coition. It arises from Poupart's ligament, passes over the spermatic cord, and is lost in the cellular membrane of the scrotum, covering the testicles.

CRENSTUS. (From *κρεμνός*, a precipice, or shelving place.) The lips of an ulcer is so called. Also the labium pudendi.

CREMOR. Cream. Any substance floating on the top, and skimmed off.

CREPITUS. (From *crepo*, to make a noise.) A puff or little noise: the crackling made by the joints when there is a defect of synovia.

CREPITUS LUTÆ. See *Muriæta*.

Cress, water. See *Nasturtium aquaticum*.

CRETA PRÆPARATA. Prepared chalk is a carbonate of lime, and possesses antacid qualities: it is exhibited in form of electuary, mixture, or bolus, in pyrosis, cardialgia, acidities of the primæ viæ, rachitis, crusta lactea, &c. and is an antidote against white arsenic. See *Carbonas calcæ*.

Cretaceous acid. See *Carbonic acid*.

Crete, dittany of. See *Dictamnus creticus*.

CRIBRIFORMIS. (From *cribrum*, a sieve, and *forma*, likeness; because it is perforated like a sieve.) See *Ethmoid bone*.

CRICO. Names compounded of this word belong to muscles which are attached to the cricoid cartilage.

CRICO-ARYTÆNOIDEUS LATERALIS. *Crico-lateri-arythenoidien* of Dumas. A muscle of the glottis, that opens the *rima* by pulling the ligaments from each other.

CRICO-ARYTÆNOIDEUS POSTICUS. *Crico-creti-arythenoidien* of Dumas. A muscle of the glottis, that opens the *rima glottidis* a little, and by pulling back the arytenoid cartilage, stretches the ligament so as to make it tense.

CRICO-PHARYNGÆUS. See *Constrictor pharyngis inferior*.

CRICO-THYROIDÆUS. *Crico-thyroidien* of Dumas. The last of the second layer of muscles between the os hyoides and trachea, that pulls forwards and depresses the thyroid cartilage, or elevates and draws backwards the cricoid cartilage.

CRICOIDES. (From *κρίκος*, a ring, and *οἶδος*, resemblance.) A round ring like cartilage of the larynx is called the cricoid. See *Larynx*.

CRIMONES. (From *κρίνον*, bran.) A term applied to urine, which deposits a sediment like bran.

CRINATUS. (From *κρίνον*, the lily.) A term

given to a suffumigation mentioned by P. Ægineta, composed chiefly of the roots of lilies.

CRINIS. The hair. See *Capillus*.

CRINOMYRON. (From *κρινος*, a lily, and *μυρον*, ointment.) An ointment composed chiefly of lilies.

CRINODES. (From *crinis*, the hair.) *Comedones.* Collections of a sebaceous fluid in the cutaneous follicles upon the face and breast, which appear like black spots, and when pressed out, look like small worms, or, as they are commonly called, maggots.

CRIOGENES. An epithet for certain troches, mentioned by P. Ægineta, and which he commends for cleansing sordid ulcers.

CRIPSORCHIS. (From *κρυπτός*, to conceal, and *ορχις*, a testicle.) Having the testicle concealed, or which is not yet descended into the scrotum.

CRISIS. (From *κρίνω*, to judge.) The judgment. The sudden change of symptoms in acute diseases, from which the recovery or death is prognosticated or judged of.

CRISFATURA. (From *crispo*, to curl.) A spasmodic contraction, or curling of the membranes and fibres.

CRISTA. (*Quasi cerista*; from *κρῆς*, a horn, or *carista*: from *κρῆς*, the head, as being on the top of the head.) Any thing which has the appearance of a crest or comb upon the head of a cock, as *crista colitidis*, the nympha. Also a tubercle about the anus; so called from its form.

CRISTA GALLI. An eminence of the ethmoid bone, so called from its resemblance to a cock's comb. See *Ethmoid bone*.

CRITHMUM. See *Crithmum*.

CRITERION. (From *κρίνω*, to judge.) The same as *crisis*.

CRITHE. (*Κριθν.*) Barley. A sty or tumour on the eyelid, in the shape and of the size of a barley-corn.

CRITHMUM. (From *κρῆν*, to secrete; so named from its supposed virtues in promoting a discharge of the urine and menses.) Samphire, or sea-fennel.

CRITHODES. (From *κριθν*, barley, and *οδος*, resemblance.) Resembling a barley-corn. It is applied to small protuberances.

CRITICAL. Determining the event of a disease. Many physicians have been of opinion, that there is something in the nature of fevers which generally determines them to be of a certain duration; and, therefore, that their terminations, whether salutary or fatal, happen at certain periods of the disease, rather than at others. These periods, which were carefully marked by Hippocrates, are called *critical days*. The critical days, or those on which we suppose the termination of continued fevers especi-

ally to happen, are the third, fifth, seventh, ninth, eleventh, fourteenth, seventeenth, and twentieth.

CRUCIDIXIS. (From *κρῖναι*, to gather wool.) *Fleclatio*. A fatal symptom in some diseases, where the patient gathers up the bed-clothes, and seems to pick up substances from them.

CROCUM. (From *κρόκος*, saffron.) Oil of saffron, or a mixture of oil, myrrh and saffron.

CROCONES. (From *κρόκος*, saffron; so called from the quantity of saffron they contain.) A name of some old troches.

CROCOMAGMA. (From *κρόκος*, saffron, and *μαγμα*, the thick oil or drega.) A troch made of oil of saffron and spices.

CROCUS. (*Κρόκος* of Theophrastus.) The story of the young Crocus, turned into this flower, may be seen in the fourth book of Ovid's *Metamorphoses*. Some derive this name from *κρῖον*, or *κρῖον*, a thread; whence the stamens of flowers are called *κρῖνες*. Others, again, derive it from *Corticus*, a city and mountain of Cilicia, and others from *crocin*. (Chald.) Saffron.

1. The name of a genus of plants in the Linnæan system. Class, *Triandria*. Order, *Monogynia*. Saffron.

2. The pharmacopœial name of the prepared stigmata of the *Crocus sativus* of Linnaeus:—*spatha univocali rutilicali, corollæ tubo longissimo*. Saffron has a powerful, penetrating, diffusive smell, and a warm, pungent, bitterish taste. Many virtues were formerly attributed to this medicine, but little confidence is now placed in it. The Edinburgh College directs a tincture, and that of London a syrup of this drug.

3. A term given by the older chymists to several preparations of metallic substances, from their resemblance: thus, *crocus martis, crocus veneris*.

CROCUS ANTIMONII. *Crocus metallorum*. This preparation is a sulphurated oxyd of antimony, and therefore called *oxydum stibii sulphuretum* in the new chymical nomenclature. It possesses emetic and drastic cathartic powers, producing a violent diaphoresis afterwards.

CROCUS GERMANICUS. See *Carthamus*.

CROCUS INDICUS. See *Curcuma*.

CROCUS MARTIS. Green vitriol exposed to fire till red.

CROCUS METALLORUM. See *Crocus antimonii*.

CROCUS OFFICINALIS. See *Crocus*.

CROCUS SARACENICUS. See *Carthamus*.

CROCUS SATIVUS. See *Crocus*.

CROCUS VENERIS. Copper calcined to a red powder.

CROMMYON. (*Παρα το τα κρῖος μύνη*, because it makes the eyes wink.) An onion.

CROMMYOXYSSEMA. (From *κρῖον*, an onion, and *ξύς*, acid, and *μύνη*, to break out.) An acid eruption accompanied with a taste resembling onions.

CROTAPHITA ARTERIA. The tendon of the temporal muscle.

CROTAPHITES. (*Crotaphites*, sc. *musculus*; from *κρότος*, the temple.) See *Temporalis*.

CROTAPHIUM. (From *κρότος*, to pulsate; so named from the pulsation which in the temples is eminently discernible.) *Crotaphos*. *Crotaphus*. A pain in the temples.

CROTAPHOS. See *Crotaphium*.

CROTAPHUS. See *Crotaphium*.

CROTCHET. A curved instrument with a sharp hook to extract the fœtus.

CROTON. (From *κρότος*, to beat.)

1. An insect called a tick, from the noise it makes by beating its head against wood.

2. A name of the ricinus or castor-oil-berry, from its likeness to a tick.

3. The name of a genus of plants in the Linnean system. Class, *Monocotyledon*. Order, *Monadelphæa*.

CROTON BENZOË. Gum-Benjamin was formerly so called.

CROTON CASCARILLÆ. The systematic name of the plant which affords the Cascarilla bark. See *Cascarilla*.

CROTON LACCIFERUM. The systematic name of the plant upon which gum-lac is deposited.

CROTON TIGLIUM. The tree which affords the pavaia wood and tiglii seeds. See *Lignum pavaia*, and *Tiglia grana*.

CROTON TINCTORIUM. The systematic name of the lacmus plant. See *Bezetta cerulea*.

CROTONE. (From *κρότος*, the tick.) A fungus on trees produced by an insect like a tick; and by metaphor applied to tumours and small fungous excrescences on the periosteum.

Croup. See *Cynanche*.

CROUSIS. (From *κρούω*, to beat, or pulsate.) Pulsation.

CROUSMATA. (From *κρούω*, to pulsate.) Rheums or defluxions from the head.

Crowsfoot. See *Ranunculus*.

Crowsfoot cranesbill. See *Geranium batrachionæ*.

CRUCIAL. Some parts of the body are so called when they cross one another, as the crucial ligaments of the thigh.

CRUCIALIS. Cross-like. Mugweed or crosswort.

CRUCIBLE. (*Crucibulum* from *crucio*, to torment; so named, because, in the language of old chymists, metals are tormented in it, and tortured, to yield up their powers and virtues.) A chemical vessel made mostly of earth to bear the greatest heat. They are of various shapes and composition.

CRUDITAS. (From *crudus*, raw.) It is applied to undigested substances in the stomach and humours in the body unprepared for concoction.

CRUENON. (From *κρῦον*, a torrent.) A

medicine mentioned by Ætius, and named from the violence of its operation as a diuretic.

CRUOR. The red part of the blood. See *Blood*.

CRURA. The plural of *crus*, a leg or root; applied to some parts of the body, from their resemblance to a leg or root: thus, *crura cerebri*, *crura cerebelli*, the *crura* of the diaphragm, &c. &c.

CRURA CLITORIDIS. See *Clitoris*.

CRURA MEDULLÆ OBLONGATÆ. The roots of the medulla oblongata.

CRURÆUS. (From *crus*, a leg; so named, because it covers almost the whole foreside of the upper part of the leg, or thigh.) *Cruralis*. A muscle of the leg, situated on the forepart of the thigh. It arises, fleshy, from between the two trochanters of the os femoris, but nearer the lesser, firmly adhering to most of the forepart of the os femoris; and is inserted, tendinous, into the upper part of the patella, behind the rectus. Its use is to assist the vasti and rectus muscles in the extension of the leg.

CRURAL. Belonging to the crus, leg, or lower extremity.

CRURAL HERNIA. Femoral hernia: A tumour under the groin, and in the upper part of the thigh, arising from a protusion of part of an abdominal viscus under Poupert's ligament. See *Hernia cruralis*.

CRURALIS. See *Cruræus*.

CRUSTA. A shell; a scab; the scum or surface of a fluid.

CRUSTA LACTEA. A disease that mostly attacks some part of the face of infants at the breast. It is known by an eruption of broad pustules, full of a glutinous liquor, which form white scabs when they are ruptured. It is cured by mineral alteratives.

CRUSTA VILLOSA. The inner coat of the stomach and intestines has been so called.

CRUSTELA. (Dim. of *crusta*, a shell.) An ecchymosis or discoloration of the flesh from a bruise, where the skin is entire and covers it over like a shell.

CRUSTUMINATUM. (From *Crustuminum*, a town where they grew.)

1. A kind of Catherine pear.

2. A rob or electuary made of this pear and apples boiled up with honey.

CRYMONES. (From *κρύος*, cold.) An epithet for a fever, wherein the external parts are cold.

CRYPTORCHIS. (From *κρυπτός*, to conceal, and *ὄρχις*, a testicle.) When the testicles are hid in the belly, or have not descended into the scrotum.

CRYPTÆ. (From *κρυπτός*, to hide.) The little rounded appearances at the end of the small arteries of the cortical substance of the kidneys, that appear as if formed by the artery being convoluted upon itself.

CRYPTOPTICA ISCHURIA. A suppression of urine from a retraction of the penis within the body.

CRYSORCHIA. (*Κρυσορχις*.) A retraction or retrocession of one of the testicles, the same as crypsorchis.

CRYSTALLI. Eruptions about the size of a lupine, white and transparent, which sometimes break out all over the body. They are also called *Crystallina*, and by the Italians *Taroli*. They are probably the pemphigus of modern writers.

CRYSTALLINE LENS. (*Lens crystallina*—*crystallina*, from its crystal-like appearance.) A lentiform pellucid part of the eye, enclosed in a membranous capsule, called the capsule of the crystalline lens, and situated in a peculiar depression in the anterior part of the vitreous humour. Its use is to transmit and refract the focus of the rays of light to the vitreous humour.

CRYSTALLINUM. (From *κρυσταλλος*, a chrysal; so called from its transparency.) White arsenic.

CRYSTALLIZATION. (From *crystallus*, a chrysal.) *Crystallization*. A property by which crystallizable bodies tend to assume a regular form, when placed in circumstances favourable to that particular disposition of their particles. Almost all minerals possess this property, but it is most eminent in saline substances. The circumstances which are favourable to the crystallization of salts, and without which it cannot take place, are two. 1. Their particles must be divided and separated by a fluid, in order that the corresponding faces of those particles may meet and unite. 2. In order that this union may take place, the fluid which separates the ingredient parts of the salt must be gradually carried off, so that it may no longer divide them.

CRYSTALLUS. (From *κρυος*, cold, and *σπασω*, to contract : i. e. contracted by cold into ice.) Chrysal. The ancients supposed that chrysalts were water intensely frozen. It also means an eruption over the body of white transparent pustules. See *Crystalli*.

CTEDONES. (From *κτενω*, a rake.) The fibres are so called from their pectinated course.

CTES. (*Κτες*.) A comb or rake. *Cteses*, in the plural number, implies those teeth which are called incisores, from their likeness to a rake.

CUBEBA. (From *cubabah*, Arab) *Piper caudatum*. *Cumamus*. Cubebs. The dried berries of the *Piper cubeba* of Linnaeus :—*foliis oblique ovatis, seu oblongis venosis acutis, spica solitaria pedunculata oppositifolia, fructibus pedicellatis*. They are of an ash-brown colour, generally wrinkled, and resembling pepper, but furnished each with a slender stalk. They are a warm spice, of a pleasant smell, and moderately pungent

taste; imported from Java; and may be exhibited in all cases where warm spicy medicines are indicated, but they are inferior to pepper.

Cubebs. See *Cubeb*.

CUBITAL ARTERY. *Arteria cubitalis, Arteria ulnaris*. A branch of the brachial that proceeds in the fore-arm, and gives off the recurrent and inter-osseals, and forms the palmary arch, from which arise branches going to the fingers, called digitalis.

CUBITAL NERVE. *Nervus cubitalis. Nervus ulnaris*. It arises from the brachial plexus, and proceeds along the ulna.

CUBITALIS MUSCULUS. An extensor muscle of the fingers.

CUBITUS EXTERNUS. An extensor muscle of the fingers.

CUBITUS INTERNUS. A flexor muscle of the fingers.

CUBITI PROFUNDA VENA. A vein of the arm.

CUBETUS. (From *cubo*, to lie down; because the ancients used to lie down on that part at their meals.) The fore-arm, or that part between the bend of the arm, including the elbow and wrist.

CUBOIDES OS. (From *κυβος*, a cube or die, and *ωδς*, likeness.) A tarsal bone of the foot, so called from its resemblance.

Cuckoo flower. See *Gardamine*.

CUCULLARIN. (*Cucullaris* sc. *musculus*; from *cucullus*, a hood: so named, because it is shaped like a hood.) See *Trapezius*.

CUCULLUS. A hood. An odoriferous cap for the head.

Cucumber. See *Cucumis*.

Cucumber, bitter. See *Colocynthis*.

Cucumber, squirting. See *Cucumis agrestis*.

Cucumber, wild. See *Cucumis agrestis*.

CUCUMIS. (*Quasi curvimeres*, from their curvature. The cucumber.

1. The name of a genus of plants in the Linnæan system. Class, *Monococcia*. Order, *Syngenesia*. The cucumber.

2. The pharmacopœial name of the garden cucumber, *Cucumis sativus* of Linnaeus :—*foliorum angulatis rectis; pomis oblongis scabris*. It is cooling and aperient, but very apt to disagree with bilious stomachs. It should always be eaten with pepper and oil. The seeds were formerly used medicinally.

CUCUMIS AGRESTIS. *Cucumis asiaticus. Cucumis xylocrostris. Elaterium officinarum. Boubalios. Guarberba erba.* Wild or squirting cucumber. *Momordica elaterium* of Linnaeus :—*pomis hispido cirrhis nullis*. The dried juice of this plant is the elaterium of the shops. It has neither smell nor taste, and is the most powerful cathartic in the whole materia medica. Its efficacy in dropsies is said to be considerable; it, however, requires great caution in the ex-

hibition. From the eighth to the half of a grain should be given at first, and repeated at proper intervals until it operates.

CUCUMIS ASININUS. See *Cucumis agrestis*.

CUCUMIS COLOCYNTHIS. The systematic name for the official bitter apple. See *Colocynthis*.

CUCUMIS MELO. The systematic name of the melon plant. See *Melo*.

CUCUMIS SATIVUS. The systematic name of the cucumber plant. See *Cucumis*.

CUCUMIS SILVESTRIIS. See *Cucumis agrestis*.

CUCURHA. A hood. An odoriferous cap for the head composed of aromatic drugs.

CUCURBITA. (*A carvitate*, according to Scaliger; the first syllable being doubled, as in *cacula*, *populus*, &c.)

1. The name of a genus of plants in the Linnæan system. Class, *Monocotyledon*. Order, *Syngenesia*. The pumpkin.

2. The pharmacopœial name of the common pumpkin or gourd. The seeds of this plant, *Cucurbita pepo*; *foliis lobatis, pomis levibus*, are used indifferently with those of the *Cucurbita lagenaria*; *foliis subangulatis, tomentosis, basi subtus biglandulosis; pomis lignosis*. They contain a large proportion of oil, which may be made into emulsions; but is superseded by that of sweet almonds.

3. A chymical distilling vessel shaped like a gourd.

CUCURBITA CITRULLUS. The systematic name of the water-melon plant. See *Citrullus*.

CUCURBITA LAGENARIA. The systematic name of the bottle-gourd plant. See *Cucurbita*.

CUCURBITA PEPO. The systematic name of the common pumpkin. See *Cucurbita*.

CUCURBITINUS. A species of worm, so called from its resemblance to the seed of the gourd. See *Tania*.

CUCURBITULA. (A diminutive of *cucurbita*, a gourd; so called from its shape.) A cupping-glass.

CUCURBITULA CRUENTA. A cupping-glass with scarification to procure blood.

CUCURBITULA CUM FERRO. A cupping-glass with scarification to draw out blood.

CUCURBITULA SICCA. A cupping-glass without scarification.

CURMA. (From *cur*, to carry in the womb.) The conception, or rather, as Hippocrates signifies by this word, when the complete rudiments of the fœtus are formed.

CURUSIO. A sort of stranguary, or rather heat of urine.

CULLAWAN CORTEX. *Cullitawan*. *Cortex caryophyllides*. The bark of the *Laurus cullawan* of Linnæus:—*foliis tripli-*

nerolis oppositis. It very much resembles cinnamon in appearance and properties.

CULINARY. (*Culinarius*, from *culina*, a kitchen.) Any thing belonging to the kitchen, as culinary salt, culinary herbs.

CULTER. (From *colo*, to cultivate.) A knife or shear. The third lobe of the liver is so called from its resemblance.

CULUS. (From *καλος*.) The anus or fundament.

CUMAMUS. See *Cubeba*.

Cumin seeds. See *Cuminum*.

CUMINUM. (From *cum*, to bring forth; because it was said to cure sterility.) *Cuminum*. *Faniculum orientale*.

1. The name of a genus of plants in the Linnæan system. Class, *Heptandria*. Order, *Digymia*. The cumin plant.

2. The pharmacopœial name of the *Cuminum cyminum* of Linnæus. A native of Egypt and Ethiopia, but cultivated in Sicily and Malta, from whence it is brought to us. The seeds of cumin which are the only part of the plant in use, have a bitterish taste, accompanied with an aromatic flavour, but not agreeable. They are generally preferred to other seeds for external use in discussing indolent tumours, as the encysted scrophulous, &c. and give name both to a plaster and cataplasm in the pharmacopœias.

CUMINUM ETHIOPICUM. A name for the *ammi verum*.

CUMINUM CYMINUM. The systematic name of the cumin plant. See *Cuminum*.

CUNEALIS SUTURA. The suture by which the os sphenoides is joined to the os frontis.

CUNEIFORM. (*Cuneiformis*; from *cuneus*, a wedge, and *forma*, likeness.) Some parts of the body are so called, being shaped or fixed in, like a wedge: such are the sphenoid bone, and some bones of the wrist and tarsus.

CUNEOLOUS. (From *cuneo*, to wedge.) A crooked tent to put into a fistula.

CUPEL. (*Kuppel*, German.) *Copella*, *Catellus cinereus*. *Cinertium*. *Patella domesticæ*. *Testa probatrix exploratrix*, or *domicastica*. A chymical instrument, which suffers the baser metals to pass through it, when exposed to heat, and retains the pure metal. This process is termed cupellation.

CUPELLATION. (From *kuppel*, German.) The purifying of perfect metals by means of an addition of lead, which at a due heat becomes vitrified and promotes the vitrification and calcination of such imperfect metals as may be in the mixture, so that these last are carried off in the fusible glass that is formed, and the perfect metals are left nearly pure. The name of this operation is taken from the vessels made use of, which are called cupels.

CURROS. (Κεφος.) Light, when applied to aliments, imports their being easily digested; when to distempers, that they are mild.

CUPRESSUS. (So called, *αὐτὸ τὴν αὐτὴν παρὰ τὴν αὐτὴν*, because it produces equal branches.) Cypress.

1. The name of a genus of plants in the Linnæan system. Class, *Monocla.* Order, *Monadelphia.* The cypress-tree.

2. The pharmacopœial name of the *Cupressus sempervirens* of Linnæus:—*foliis imbricatis squamis quadrangulis*; called also *cyparissius*. Every part of the plant abounds with a bitter, aromatic, terebinthinate fluid; and is said to be the remedy against intermittents. Its wood is extremely durable, and constitutes the cases of Egyptian mummies.

CUPRESSUS SEMPERVIRENS. The systematic name of the cupressus of the shops. See *Cupressus*.

CUPRI AMMONIATI AQUA. *Aqua sapharina.* "Take of lime water, one pint; sal ammoniac, one drachm: let them stand together in a copper vessel until the ammonia is saturated."

CUPRI RUBIGO. Verdigris or rust of copper.

CUPRUM. (*Quasi ex Cyprum*: so called from the island of Cyprus, whence it was formerly brought.) See *Copper*.

CUPRUM AMMONIACALE. *Cuprum ammoniacum.* Blue vitriol and prepared ammonia. An ammoniacal sulphat of copper.

CUPRUM VITRIOLATUM. See *Sulphas cupri*.

CURA AVENACEA. A decoction of oats and succory roots, in which a little nitre and sugar were dissolved, was formerly used in fevers, and was thus named.

CURCAS. The Barbadoes nut; a drastic purge.

CURCULIO. (From *karkarah.* Heb.) The throat; the aspera arteria.

CURCUM. The large celandine; deobstruent.

CURCUMA. (From the Arabic *curcum*, or *hercum*.) Turmeric.

1. The name of a genus of plants in the Linnæan system. Class, *Monandria.* Order, *Monogynia.*

2. The pharmacopœial name of the turmeric-tree. *Curcuma longa* of Linnæus:—*foliis lanceolatis; nervis lateralibus numerosissimis*; called also, *Crocus Indicus.* *Terra marita.* *Cassiacorus radice croce.* *Curcuma longa.* *Curcuma rotunda.* *Mayella.* *Kua kaha* by the Indians. The Arabians call every root of a saffron colour by the name of *curcim*. The root of this plant is imported here in its dried state from the East Indies, in various forms. Externally it is of a pale yellow colour, wrinkled, solid, ponderous, and the inner substance of a deep saffron or gold colour: its odour is somewhat fragrant; to the taste it is bitterish, slightly acid, exciting a moderate degree of warmth in the mouth, and on being chewed it tinges the saliva yellow.

It is an ingredient in the composition of *Curry powder*, is valuable as a dyeing drug, and furnishes a chymical test of the presence of uncombined alkalies. It is now very seldom used medicinally, but retains a place in our pharmacopœias.

CURCUMA LONGA. The turmeric plant. See *Curcuma*.

CURCUMA ROTUNDA. See *Curcuma*.

CURMI. (From *azga*, to mix.) Ale. A drink made of barley, according to Dioscorides.

Currants. See *Ribes*.

CURSUMA. *Curturna.* The *Ranunculus ficaria* of Linnæus.

CURVATOR COCCYIS. A muscle bending the coccyx.

CURSUTA. (Corrupted from *casuta*, *kanuth*, Arab.) The root of the *Gentiana purpurea* of Linnæus.

CUSCUTA. (According to Linnæus, a corruption from the Greek *καρύς*, or *καρύς*, which is from the Arabic *Chesuth*, or *Chasuth*.) Dodder.

1. The name of a genus of plants in the Linnæan system. Class, *Tetrandria.* Order, *Digynia.*

2. The pharmacopœial name of dodder of thyme. See *Epithymum*.

CUSCUTA EPITHYMIUM. The systematic name of dodder of thyme. See *Epithymum*.

CUSCUTA EUROPEA. The systematic name of a species of dodder of thyme. See *Epithymum*.

CUSPIDATUS. (From *cuspis*, a point.) See *Teeth*.

CUSPARIA. The general name given by Messrs. Humboldt and Boupland to the tree from which we obtain the *Augustura* bark. See *Augustura*.

CUSPIS. (From *caipa*, Chald. a shell, or bone, with which spears were formerly pointed.) The glans penis was so called, from its likeness to the point of a spear. Also a bandage.

CUSTOS OCULI. An instrument to fix the eye during an operation.

CUTANEBULI. (From *cutis*, the skin.) Cutaneous worms; scorbutic itching.

CUTANEUS MUSCULUS. The platysma myoides.

CUTANEOUS. (From *cutis*, the skin.) Belonging to the skin.

CUTICLE. (*Cuticula*, dim. of *cutis*, the skin.) *Epidermis.* Scarf-skin. A thin, pellucid, insensible membrane, of a white colour, that covers and defends the true skin, with which it is connected by the hairs, exhaling and inhaling vessels, and the rete mucosum.

CUTIS. *Dermis.* The true skin. A thick, fibrous, vascular, and nervous membrane, that covers the whole external surface of the body, and is the situation of the organ of touch, exhalation, and inhalation.

CUTIS ANSERINA. (*Anserina*; from

anser, a goose.) The rough state the skin is sometimes thrown into from the action of cold, or other cause, in which it looks like the skin of the goose.

CUTIS VERA. The true skin under the cuticle.

CYANUS. (*Κυανος*, azurean, or sky-blue; so called from its colour.) Blue-bottle. Corn-flower. The flowers of this plant, *Centaurea cyaneus* of Linnæus:—*calycibus serratis; foliis linearibus, integerrimis, infamis dentatis*, were formerly in frequent use; but their antiphlogistic, antispasmodic, cordial, aperient, diuretic, and other properties are now, with great propriety, forgotten.

CYAR. (From *κυω*, to pour out.) The lip of a vessel. The eye of a needle; and the orifice of the internal ear, from its likeness to the eye of a needle.

CYASMA. Spots on the skin of pregnant women.

CYATHISCUS. (From *κυαθος*, a cup.) The hollow part of a probe, formed in the shape of a small spoon, as an ear-picker.

CYBITOS. See *Cubitus*.

CYBITUM. See *Cubitus*.

CYBITUS. See *Cubitus*.

CYBOIDES. See *Cuboides*.

CYCEUM. From *κυκαα*, to mix.) *Cyceon*. A mixture of the consistence of pap.

CYCLIA. (From *κυκαα*, to mix.) So called from the mixture of the ore with lead, by which litharge is made.

CYCLAMEN. (From *κυκλος*, circular; either on account of the round form of the leaves, or of the roots.) Cyclamen.

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

2. The pharmacopœial name of the sow-bread. See *Arthanita*.

CYCLAMEN EUROPEUM. The systematic name of the sow-bread. See *Arthanita*.

CYCLISCUS. (From *κυκλος*, a circle.) An instrument in the form of a half moon, formerly used for scraping rotten bones.

CYCLISMUS. (From *κυκλος*, a circle.) A lozenge.

CYCLOPHORIA. (From *κυκλος*, a circle, and *φορεω*, to bear.) The circulation of the blood, or other fluids.

CYCLOPIOX. (From *κυκλωω*, to surround, and *οφθαλμος*, the eye.) The white of the eye.

CYCLOS. A circle. Hippocrates uses this word to signify the cheeks, and the orbits of the eyes.

CYCLUS METASYNCRITITUS. It is a long protracted course of remedies, persisted in with a view of restoring the particles of the body to such a state as is necessary to health.

CYDONIUM MALUM. (From *Cydon*, a town in Crete, where they grew.) The quince. The tree which affords this fruit is the *Pyrus cydonia* of Linnæus:—*foliis integerrimis, floribus solitariis*. Quince seeds

are directed by the London College to be made into a mucilage, which is recommended in apthous affections, and excoriations of the mouth and fauces.

CYEMA. (From *κυω*, to bring forth.) Parturition.

CYLICHRIS. (From *κυλεξ*, a cup.) A gallipot or vessel of any kind to hold medicines in.

CYLOSIS. (From *κυλλω*, to make lame.) A tibia or leg bending outwards.

CYLINDRUS. (From *κυλλω*, to roll round.) A cylinder. A tent for a wound, equal at the top and bottom.

CYLLUS. (From *κυλλω*, to make lame.) In Hippocrates, it is one affected with a kind of luxation, which bends outwards, and is hollowed inward. Such a defect in the tibia is called *Cylosis*, and the person to whom it belongs is called, by the Latins, *Varus*, and is opposed to *Valgus*.

CYMATODES. Is applied by Galen and some others to an unequal fluctuating pulse.

CYMA. (From *κυμας*, hollow.) A boat, or pinnace. A bone of the wrist is so called, from its supposed likeness to a skiff.

CYMINALIS. Gentian.

CYMINUM. See *Cuminum*.

CYNANCHE. (From *κυων*, a dog, and *ανω*, to suffocate, or strangle; so called from dogs being said to be subject to it.) Sore throat. A genus of disease in the class *pyrexia*, and order *phlegmasie* of Cullen. It is known by pain and redness of the throat, attended with a difficulty of swallowing and breathing. The species of this disease are—

1. *Cynanche tracheitis*; called also *cynanche laryngea*. *Suffocatio stridula angina perniciosa*. *Asthma infantum*. *Cynanche stridula*. *Morbus strangulatorius*. *Catarhus suffocatus Barbadensis*. *Angina polyposa sive membranacea*. The croup. A disease that mostly attacks infants, who are suddenly seized with a difficulty of breathing and a crouping noise; it is an inflammation of the mucous membrane of the trachea, that induces the secretion of a very tenacious coagulable lymph, which lines the trachea and bronchia, and impedes respiration. The croup does not appear to be contagious, whatever some physicians may think to the contrary; but it sometimes prevails epidemically. It seems, however, peculiar to some families; and a child having once been attacked, is very liable to its returns. It is likewise peculiar to young children, and has never been known to attack a person arrived at the age of puberty.

The application of cold seems to be the general cause which produces this disorder, and therefore it occurs more frequently in the winter and spring, than in the other seasons. It has been said, that it is most

prevalent near the sea coast; but it is frequently met with in inland situations, and particularly those which are marshy.

Some days previous to an attack of the disease, the child appears drowsy, inactive, and fretful; the eyes are somewhat suffused and heavy; and there is a cough, which, from the first, has a peculiar shrill sound; this, in the course of two days, becomes more violent and troublesome, and likewise more shrill. Every fit of coughing agitates the patient very much; the face is flushed and swelled, the eyes are protuberant, a general tremor takes place, and there is a kind of convulsive endeavour to renew respiration at the close of each fit. As the disease advances, a constant difficulty of breathing prevails, accompanied with a swelling and inflammation in the tonsils, uvula, and velum pendulum palati; and the head is thrown back, in the agony of attempting to escape suffocation. There is not only an unusual sound produced by the cough, (something between the yelping and barking of a dog,) but respiration is performed with a hissing noise, as if the trachea was closed up by some slight spongy substance. The cough is generally dry; but if any thing is spit up, it has either a purulent appearance, or seems to consist of films resembling portions of a membrane. Where great nausea and frequent retchings prevail, coagulated matter of the same nature is brought up. With these symptoms, there is much thirst, and uneasy sense of heat over the whole body, a continual inclination to change from place to place, great restlessness, and frequency of the pulse.

In an advanced stage of the disease, respiration becomes more stridulous, and is performed with still greater difficulty, being repeated at longer periods, and with greater exertions, until at last it ceases entirely.

The croup generally proves fatal by suffocation, induced either by spasm affecting the glottis, or by a quantity of matter blocking up the bronchiæ; but when it terminates in health, it is by a resolution of the inflammation, by a ceasing of the spasms, and by a free expectoration of the matter exuding from the trachea, or of the crusts formed there.

The disease has, in a few instances, terminated fatally within twenty-four hours after its attack; but it more usually happens, that where it proves fatal, it runs on to the fourth or fifth day. Where considerable portions of the membranous films, formed on the surface of the trachea, are thrown up, life is sometimes protracted for a day or two longer than would otherwise have happened.

Dissections of children who have died of the croup have always shown a preternatural membrane, lining the whole internal

surface of the upper part of the trachea, which may always be easily separated from the proper membrane. There is likewise usually found a good deal of mucus, with a mixture of pus, in the trachea and its ramifications.

2. *Cynanche tonsillaris*. The inflammatory quincy, called also *angina inflammatoria*. In this complaint the inflammation principally occupies the glands, such as the tonsils; but often extends through the whole mucous membrane of the fauces, so as essentially to interrupt the speech, respiration, and deglutition of the patient.

The causes which usually give rise to it are, exposure to cold, either from sudden vicissitudes of weather, from being placed in a partial current of air, wearing damp linen, sitting in wet rooms, or getting wet in the feet; all of which may give a sudden check to perspiration. It principally attacks those of a full and plethoric habit, and is chiefly confined to cold climates, occurring usually in the spring and autumn; whereas the ulcerated sore throat chiefly attacks those of a weak irritable habit, and is most prevalent in warm climates. The former differs from the latter likewise in not being contagious. In many people their seems to be a particular tendency to this disease; as from every considerable application of cold it is readily induced.

An inflammatory sore throat discovers itself by a difficulty of swallowing and breathing, accompanied by a redness and tumour in one or both tonsils, dryness of the throat, foulness of the tongue, lancinating pains in the parts affected, a frequent but difficult excretion of mucus, and some small degree of fever. As the disease advances, the difficulty of swallowing and breathing becomes greater, the speech is very indistinct, the dryness of the throat and thirst increase, the tongue swells and is incrustated with a dark fur, and the pulse is full and frequent. In some cases, a few white sloughy spots are to be observed on the tonsils. If the inflammation proceeds to such a height as to put a total stop to respiration, the face will become livid, the pulse will sink, and the patient will quickly be destroyed.

The chief danger arising from this species of quincy is, the inflammation occupying both tonsils, and proceeding to such a degree as to prevent a sufficient quantity of nourishment for the support of nature from being taken, or its occasioning suffocation; but this seldom happens, and its usual termination is either in resolution or suppuration. When proper steps are adopted, it will in general readily go off by the former.

Where the disease has proved fatal by suffocation, little more than a highly inflamed state of the parts affected, with some morbid phenomena in the head, have been observed on dissection.

3. *Cynanche pharyngea*. This species is so called when the pharynx is chiefly affected. Dr. Wilson, in his treatise on Febrile Diseases, includes in his definition of cynanche tonsillaris, that of cynanche pharyngea. These varieties of cynanche differ considerably when they are exquisitely formed. But the one is seldom present in any considerable degree without being attended with more or less of the other. Dr. Cullen declares, indeed, that he never saw a case of true cynanche pharyngea; that is, a case in which the inflammation was confined to the pharynx; it constantly spread in a greater or less degree to the tonsils and neighbouring parts. Besides the mode of treatment is, in almost every instance, the same in both cases. And if we admit the cynanche pharyngea to be a distinct variety, we must admit another, the cynanche œsophagea; for the inflammation frequently attacks the œsophagus, and is sometimes even confined to it.

4. *Cynanche parotidea*. The mumps. A swelling under the jaw, extending over the neck; an inflammation of the parotid gland, rendering deglutition difficult, declining the fourth day. Epidemic and contagious.

5. *Cynanche maligna*. The malignant, putrid, or ulcerous sore throat. Called also *cynanche gangrenosa*. *Angina ulcerosa*. *Febris epidemica cum angina ulcusculosa*. *Angina epidemica*. *Angina gangrenosa*. *Angina suffocativa*. *Angina maligna*. This disease is readily to be distinguished from the inflammatory quincy, by the soreness and white specks which appear in the fauces, together with the great debility of the system, and small fluttering pulse, which are not to be observed in the former. In the inflammatory sore throat there is always great difficulty of swallowing, a considerable degree of tumour, with a tendency in the parts affected to suppurate, and a hard, full pulse. Moreover in the former affection the disease is seated principally in the mucous membrane of the mouth and throat; whereas in the latter the inflammation chiefly occupies the glandular parts.

The putrid sore throat often arises from a peculiar state of the atmosphere, and so becomes epidemical; making its attacks chiefly on children, and those of a weak relaxed habit. It is produced likewise by contagion, as it is found to run through a whole family, when it has once seized any person in it; and it proves often fatal, particularly to those in an infantile state.

It usually makes its attack with cold shiverings, anxiety, nausea, and vomiting, succeeded by heat and restlessness, debility and oppression at the chest. The face looks flushed, the eyes are red, and a

stiffness is perceived in the neck, with a hoarseness of voice, and soreness in the throat; and, upon viewing the internal fauces, there appears a fiery redness in every part, with some slight degree of swelling in the tonsils, which, however, is by no means so great as to impede either respiration or deglutition.

The inflammation, after a short time, takes a peculiar termination; for, upon further inspection into the throat, a number of white specks, or sloughs, are to be observed on the tonsils and uvula, the breath is highly offensive, the tongue is covered with a thick brown fur, and the inside of the lips are beset with vesicles, containing an acrid matter, which, falling on the corners of the mouth and other parts, occasions excoriations. With these symptoms there is likewise a coryza, which pours out a thin acrid matter, excoriating the nostrils. A purging often attends also, particularly in infants, and a thin acrid matter flows from the anus, excoriating this and the neighbouring parts.

From the first attack of the complaint, there is a considerable degree of fever, with a small, frequent and irregular pulse; and every evening there occurs a manifest exacerbation, and in the morning some slight remission, together with general loss of strength, and debility. In some cases the brain is affected with delirium, or coma.

About the second or third day, large patches of a scarlet or fiery red colour, make their appearance about the face and neck, which, by degrees, become dispersed over every part of the body, even to the extremities of the fingers, which feel swelled and stiff. These eruptions, after continuing for about four days, then depart without producing any remission of the symptoms.

In bad cases, the sloughs corrode deeper and deeper, and spread throughout the whole of the alimentary tube, so as to terminate at last in gangrene; and the symptoms of irritation continuing to increase, together with a severe purging, the patient is at length cut off; which event happens usually before the seventh day, and in some cases so early as on the third.

Where there is a great increase of the evening paroxysm of fever, with vast debility, irregularity in the pulse, much fetor of breath, and a livid appearance in the ulcers, with a purging, or hæmorrhage, the disease will certainly terminate fatally; but when the fever is moderate, and of a less putrid nature, and suffers a remission on the appearance of the efflorescence on the skin; and when this remission increases as it proceeds, together with a falling off of the cuticle in scales, it promises fair to terminate in a return of health.

From dissections it appears that, in this

disease, the fauces are inflamed, suppurated and gangrenous; and that the trachea and larynx are likewise in a state of inflammation, and lined with a viscid fetid matter. In many instances, the inflammatory affection extends to the lungs themselves. Large swellings of the lymphatic glands about the neck, occasioned by an absorption of the acrid matter poured out in the fauces, are now and then to be found. The same morbid appearances which are to be met with in putrid fever, present themselves in other parts of the body.

CYNANCHE A DEGLUTITIS. Quinoy from hard substances swallowed.

CYNANCHE A DYSENTERIA. Quinoy from dysentery.

CYNANCHE ANGINOSA. The inflammatory quinoy.

CYNANCHE ARTHRITICA. Quinoy from gout.

CYNANCHE EPIDEMICA. The cynanche maligna.

CYNANCHE GANGRENOZA. The cynanche maligna.

CYNANCHE HEPATICA. Quinoy from a disease of the liver.

CYNANCHE LARYNGEA. The cynanche trachealis.

CYNANCHE MALIGNA. See *Cynanche*.

CYNANCHE PAROTIDEA. See *Cynanche*.

CYNANCHE PHARYNGEA. See *Cynanche*.

CYNANCHE PRUNELLA. Common sore throat.

CYNANCHE PURPURE-PAROTIDEA. A cynanche maligna, or malignant sore throat.

CYNANCHE STRIDULA. The croup. See *Cynanche*.

CYNANCHE THYMICA. Sore throat from an enlargement of the thyroid gland.

CYNANCHE TONSILLARIS. See *Cynanche*.

CYNANCHE TRACHEALIS. See *Cynanche*.

CYNANCHE ULCEROSA. The malignant sore throat.

CYNANCHICA. (From *κυνάχην*, the quinoy.) Medicines which relieve a quinoy.

CYNANTHROPIA. (From *κυν*, a dog, and *άνθρωπος*, a man.) It is used by Bellini, De Morbis Capitis, to express a particular kind of melancholy, when men fancy themselves changed into dogs, and imitate their actions.

CYNARA SCOLYMUS. See *Cinara*.

CYNCHIS. (Κύκκιν.) A vessel of any kind to hold medicines in.

CYNOCOCTANUM. (From *κυν*, a dog, and *κόκκος*, the herb coctanum.) A species of aconitum, said to destroy dogs if they eat it.

CYNOCRAME. (From *κυν*, a dog, and *ράμν*, cabbage; a herbe of the cabbage tribe, with which dogs are said to physic themselves.) Dog's mercury. *Mercurialis perennis* of Linnaeus. A poisonous plant very common in our hedges. It produces vomiting and purging, and the person then

goes to sleep, from which he does not often awake.

CYNOCTYSIS. (From *κυν*, a dog, and *κύστις*, the cytismus; so named because it was said to cure the distemper of dogs.) The dog-rose. See *Cynosbatus*.

CYNODACTOS. (From *κυν*, a dog, and *δάκν*, to bite.) So Dioscorides calls a person bit with a mad dog.

CYNODISMION. (From *κυν*, a dog, and *δισμ*, to bind; so named because in dogs it is very discernible and strong.) A ligature by which the prepuce is bound upon the glans. Sometimes it signifies the lower part of the prepuce.

CYNODONTES. (Κυνόδοντες: from *κυν*, a dog, and *όντις*, a tooth.) The canine teeth.

CYNOGLOSSUM. (From *κυν*, a dog, and *γλῶσσα*, a tongue; so named from its supposed resemblance.) Hound's tongue.

1. The name of a genus of plants in the Linnaean system. Class, *Pentandria*. Order, *Monogynia*.

2. The pharmacopœial name of the *Lingua canina* of some authors. *Cynoglossum, officinale* of Linnaeus:—*staminibus corollæ brevioribus; foliis late lanceolatis, tomentosis, sessilibus*. It possesses narcotic powers, but is seldom employed medicinally. Acids are said to counteract the ill effects from an overdose more speedily than any thing else, after clearing the stomach.

CYNOGLOSSUM OFFICINALE. The systematic name for hound's tongue. See *Cynoglossum*.

CYNOLOPHUS. (From *κυν*, a dog, and *λόφος*, a protuberance; so called because in dogs they are peculiarly eminent.) The asperities and prominences of the vertebrae.

CYNOLISSA. (From *κυν*, a dog, and *λύσσα*, madness.) Canine madness.

CYNOMORIUM. The name of a genus of plants in the Linnaean system. Class, *Monacia*. Order, *Menandria*.

CYNOMORIUM COCCINEUM. The systematic name of the *fungus melitensis*. See *Fungus melitensis*.

CYNOPASTUM. (From *κυν*, a dog, and *πάσκειν*, to attract.) See *Cynosbatus*.

CYNOREXIA. (From *κυν*, a dog, and *ῥέξις*, appetite.) A voracious or canine appetite. See *Bulimia*.

CYNOSBATOR. See *Cynosbatus*.

CYNOSBATUS. (From *κυν*, a dog, and *βέλος*, a thorn; so called because dogs are said to be attracted by its smell.) *Cynorrhodon*. *Cynosbatus*. The dog-rose, or wild brier, or hip tree. *Rosa canina* of Linnaeus:—*germinibus ovatis pedunculisque glabris, caule petiolisque aculeatis*. The fruit of this tree, called heps, or hips, has a sourish taste, and obtains a place in the London pharmacopœias in the form of conserve. It is seldom employed but to give form to more active remedies, in pills, boluses, linctuses, &c.

CYONORIA. (From *κυν*, a fœtus, and

page, to bear.) **Gestation.** The pregnancy of a woman.

CYPARISSUS. See *Cyperus*.

CYPERUS. (From *κυμαος*, a little round vessel which its roots are said to resemble.) *Cyperus*. The name of a genus of plants in the Linnæan system. Class, *Triandria*. Order, *Monogymia*.

CYPERUS LONGUS. The pharmacopœial name of the English galangale. *Cyperus longus* of Linnæus:—*culmotriquetro folioso, umbella folioso supra-decomposita; pedunculis nudis, spicis alternis*. The smell of the root of this plant is aromatic, and its taste warm, and sometimes bitter. It is now totally fallen into disuse.

CYPERUS ROTUNDUS. This species, the round cyperus, *Cyperus rotundus* of Linnæus:—*culmo triquetro subnudo, umbella decomposita; spicis alternis linearibus*, is generally preferred to the former, being a more gratefully aromatic bitter. It is chiefly used as a stomachic.

CYPHOMA. (From *κυρτω*, to bend.) A gibbosity, or curvature of the spine.

CYPHOSIS. An incurvation of the spine.

Cypress spurge. See *Eaula minor*.

CYPERINUM OLEUM. Flowers of cypress, calamus, cardamoms, &c. boiled in olive oil.

CYPRUM. (From *Κυπρος*, Cyprus, an island where it is said formerly to have abounded.) Copper.

CYPRUS. The cypress-tree, or Eastern privet; so called from the island of Cyprus, where it grew abundantly.

CYPSILIS. (From *κυψαλα*, a bee-hive.) The aperture of the ear; the ear-wax.

CYRNEISIS. (From *κυμανα*, to mix.) A mixture, or composition.

CYRTOMA. (From *κυρτω*, curved.) An unnatural convex tumour; tympanites.

CYRTONOSUS. (From *κυρτω*, curved, and *νοσος*, a disease.) The rickets, or curved spine.

CYSIARUS. (From *κυρος*, the anus.) The intestinum rectum is so called, because it reaches to the anus.

CYSOTIS. (From *κυρος*, the anus.) An inflammation of the anus.

CYSTOLITHUS. (From *κυστις*, the bladder, and *λιθος*, a stone.) The stone in the bladder.

CYSTHUS. (*Κυθος*.) The anus.

CYSTIC. Belonging to the urinary or gall bladder.

CYSTIC DUCT. *Ductus cysticus.* The membranous canal that conveys the bile from the hepatic duct into the gall-bladder.

CYSTICA. (From *κυστις*, the bladder.) Remedies for diseases of the bladder.

CYSTIDES. (From *κυστις* a bag.) Encysted tumours.

CYSTIPHLOGIA. (From *κυστις*, the bladder, and *φλεγω*, to burn.) An inflammation in the bladder.

CYSTIRRHAGIA. (From *κυστις*, the blad-

der, and *ρηνω*, to flow.) A discharge of blood from the bladder.

CYSTIS. (*Κυστις*, a bag.) The bladder; any receptacle of morbid humours. See *Urinary bladder*.

CYSTIS CHOLEDOCHA. See *Gall-bladder*.

CYSTIS FELLEA. See *Gall-bladder*.

CYSTITIS. (From *κυστις*, the bladder.) Inflammation of the bladder. A genus of disease arranged by Cullen in the class, *pyrexie*, and order *phlegmasie*. It is known by great pain in the region of the bladder, attended with fever and hard pulse, a painful discharge of urine, and a frequent desire to make water. This is rarely a primary disease, and when it occurs, the above character of it will readily point it out. There is frequently also nausea and vomiting, and, in some cases, delirium. It most generally arises in consequence of inflammation of the adjacent parts, or from calculi in the bladder.

CYSTOCELE. (From *κυστις*, the bladder, and *κελω*, a tumour.) An hernia formed by the protusion of the urinary bladder.

CYSTOLITHICUS. (From *κυστις*, the bladder, and *λιθος*, a stone.) A suppression of urine from a stone in the bladder, is called *ischuria cystolithica*.

CYSTOPHLEGICUS. (From *κυστις*, the bladder, and *φλεγω*, to burn.) A suppression of urine from an inflammation of the bladder, was formerly called *ischuria cystophlegmatica*.

CYSTOPHLEGMATICA. (From *κυστις*, the bladder, and *φλεγμα*, phlegm.) A suppression of urine, from too much matter or mucus in the bladder, was called *ischuria cystophlegmatica*.

CYSTOPROCTICA. (From *κυστις*, the bladder, and *προκτω*, the anus, or rectum.) A suppression of urine from pain in the bladder, caused by wind, inflammation of the rectum, hardened feces, &c. is called *ischuria cystoproctica*.

CYSTOPTOSIS. (From *κυστις*, the bladder, and *πτωσις*, to fall.) A protusion of the inner membrane of the bladder, through the urethra.

CYSTOSPASTICUS. (From *κυστις*, the bladder, and *σπασμα*, a spasm.) A suppression of urine, from a spasm in the sphincter of the bladder, was called *ischuria cystospastica*.

CYSTOSPITICUS. (From *κυστις*, the bladder, and *πυον*, pus.) A suppression of urine, from purulent matter in the bladder, was called *ischuria cystospitica*.

CYSTOTROMBOIDES. (From *κυστις*, the bladder, and *θρομβος*, a coagulation of blood.) A suppression of urine, from a concretion of grumous blood in the bladder, was called *ischuria cystothromboides*.

CYSTOTOMIA. (From *κυστις*, the bladder, and *τομη*, to cut.) The operation of cutting or piercing the bladder.

CYTHON. An eye-wash.

CYTINUS. (From *cyo*, to produce; so called from its fecundity.) The bud or flower of the pomegranate.

CYTINUS HYPOCISTIS. The plant from whose fruit the *succus hypocistidis* is obtained. See *Hypocistia*.

CYTISO-GENISTA. Common heath. See *Genista*.

CYTXENUS. A plaster for wounds of the nerves.

CYEMER. A painful swelling of the wrists.

D.

D. This letter signifies vitriol in the old chemical alphabet.

DACNERUS. (From *daeno*, to bite.) Biting. Pungent. An epithet for a sharp collyrium, or eye-wash, composed of burnt copper, pepper, cadmia, myrrh, and opium.

DACRYDIUM. (From *daugu*, a tear.) The inspissated juice of scammony. It is in small drops, and therefore called a tear.

DACRYGELOSIS. (From *dauguo*, to weep, and *gelo*, to laugh.) A species of insanity, in which the patient weeps and laughs at the same time.

DACRYODUS. (From *dauguo*, to weep.) A sanious ulcer. A weeping sore.

DACTROMA. (From *dauguo*, to weep.) A union of one or more of the puncta lachrymalia, causing an effusion of tears.

DACTYLETHRA. (From *daulolos*, a finger.) A species of bougies shaped like a finger, to excite vomiting.

DACTYLETUS. (From *daulolos*, the date.) The hermodactylus.

DACTYLUS. (From *daulolos*, a finger.) A round pastil troche, or lozenge, shaped like a finger.

DACTYLUS. (A finger; so called from the likeness of its fruit to a finger.) The date. The oblong fruit of the *Phoenix dactylifera* of Linnæus:—*frondibus pinnatis; foliis ensiformibus complicatis*. Before they are ripe, dates are rather rough and astringent; but when perfectly matured, they are much of the nature of the fig. See *Carica*. Senegal dates are most esteemed, they having a more sugary, agreeable flavour than those of Egypt and other places.

DÆDIUM. (From *dæu*, a torch.) A small torch or candle. A bougie.

DAMNATUS. (From *damno*, to condemn.) *Caput mortuum*. The dry useless feces left in a vessel after the moisture has been distilled from it, is called *terra damnata*.

Daisy, common. See *Bellis minor*.

Daisy, ox eye. See *Bellis major*.

Damask rose. See *Rosa centifolia*.

DEMONOMANIA. (From *daimon*, a demon, and *mania*, madness.) That species of melancholy, where the patient supposes himself to be possessed of devils.

DAMSON. The fruit of the *Prunus damascena*, which, when perfectly ripe, affords a wholesome article for pies, tarts, &c. gently opening the body; but when

damsons are not perfectly mature, they produce cholicky pains, diarrhœa, and convulsions in children.

Dandelion. See *Taraxacum*.

Dandrif. See *Pityriasis*.

Dane-wort. See *Ébulus*.

DAPHNE. (*Daphne, daφn*: from *da*, to burn, and *φν*, a noise; because of the noise it makes when burnt.) The name of a genus of plants in the Linnæan system. Class, *Octandria*. Order, *Monogynia*. The laurel, or bay-tree.

Daphne, flux-leaves. See *Thymelæa*.

DAPHNE EXIDIMUM. The systematic name of the tree which affords the garou. See *Thymelæa*.

DAPHNE LAUREOLA. The systematic name of the spurge laurel. See *Laureola*.

DAPHNE MEZERIUM. The systematic name of the mezereum. See *Mezereum*.

DAPHNELÆON. (From *daφn*, the laurel, and *laion*, oil.) The oil of bay-berries.

DAPHNITIS. (From *daφn*, the laurel.) A sort of cassia resembling the laurel.

DAPHNOIDES. (From *daφn*, the laurel, and *ωδo*, a likeness.) The herb spurge laurel.

DARSIN. (From *darzin*, Arab.) The grosser sort of cinnamon.

DARSIS. (From *daφo*, to excoriate.) An excoriation.

DARTOS. (From *daφo*, to excoriate; so called from its raw and excoriated appearance. The part so called, under the skin of the scrotum, is by some anatomists considered as a muscle, although it appears to be no more than a condensation of the cellular membrane lining the scrotum. It is by means of the dartos that the skin of the scrotum is corrugated and relaxed.

DASYNA. (From *daφu*, rough.) A scabby roughness of the eye-lids.

DASTS. (*daφu*, rough.) A dry, parched tongue. Difficult respiration.

Date plum, Indian. See *Indiun date plum*.

Date. See *Dactylus*.

DATURA. (Blanchard says it is derived from the Indian word *datiro*, o which he knows not the meaning.) Th name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

DATURA STRAMONIUM. The systematic name of the thorn apple. See *Stramonium*.

DAUCUS VIRID. Wild carrot-seeds steeped in must.

DAUCUS. (*Are* or *daus*, from its relieving the cholic, and discussing flatulencies.) The carrot.

1. The name of a genus of plants in the Linnean system. Class, *Pentandria*. Order, *Digynia*.

2. The pharmacopœial name of the garden carrot. *Daucus sylvestris*. *Pastinaca sylvestris tenuifolia officinarum*. The cultivated root of the *Daucus carota* of Linnaeus:—*seminibus hispida, petiolis subtus nervosis*, scraped and applied in the form of a poultice, it is an useful application to phagedenic ulcers, and to cancers and putrid sores. The seeds, which obtain a place in the materia medica, have a light aromatic smell, and a warm acrid taste, and are esteemed for their diuretic qualities, and for their utility in calculous and nephritic complaints, in which an infusion of three spoonfuls of the seeds in a pint of boiling water, has been recommended; or the seeds may be fermented in malt liquor, which receives from them an agreeable flavour, resembling that of lemon peel. The boiled root is said, by many, to be difficult of digestion; but this is the case only when the stomach is weak. It contains a considerable quantity of the saccharine principle, and is very nutritious.

DAUCUS ALSATICUS. The oreoselinum pratense.

DAUCUS ANNUUS MINOR. The caucalis atrisculus.

DAUCUS CAROTA. The systematic name of the carrot plant. See *Daucus*.

DAUCUS CRETICUS. *Myrrhus annua*. Candy Carrot. The seeds of this plant, *Athamanta cretensis* of Linnaeus:—*foliis linearibus planis, hirsutis; petalis bipartitis; seminibus oblongis, hirsutis*, are brought from the isle of Candy; they have an aromatic smell, and a slightly biting taste; and are occasionally employed as carminatives and diuretics in diseases of the primæ viæ and urinary passages.

DAUCUS SATIVUS. A variety of the *daucus carota*, whose seeds are preferred by some practitioners.

DAUCUS REPENS. Common chervil.

DAUCUS SYLVESTRIS. Wild carrot, or bird's nest. The seeds of the wild plant are said to be more efficacious than those of the garden carrot; they possess demulcent and aromatic qualities, and are given, in infusion, or decoction, in calculous complaints.

Dead nettle. See *Labium album*.

Deadly nightshade. See *Beladonna*.

DEAFNESS. It is occasioned by any thing that proves injurious to the ear, as loud noises from the firing of cannon, violent colds, particularly affecting the head, inflammation or ulceration of the membrane, hard wax, or other substances inter-

rupting sounds; too great a dryness, or too much moisture in the parts; or by atony, debility, or paralysis of the auditory nerves. In some instances it ensues in consequence of preceding diseases, such as fever, syphilis, &c. and in others it depends upon an original defect in the structure or formation of the ear. In the last instance the person is usually not only deaf but likewise dumb. See *Paracusis*.

DEARTICULATIO. (From *de*, and *articulus*, a joint.) Articulation admitting evident motion.

DEASCIATIO. (From *de*, and *ascio*, to chip, as with a hatchet.) A bone splintered on its side.

DECATRON. (From *deka*, ten, and *muon*, an ointment.) An aromatic ointment, mentioned by Oribasius, containing ten ingredients.

DECIDENTIA. (From *decido*, to fall down.) *Cataptosis*. Any change prolonging acute diseases.

DECIDUA. (*Decidua*, sc. *membrana*; from *decido*, to fall down.) *Membrana decidua*. A very thin and delicate membrane or tunica, which adheres to the gravid uterus, and is said to be a reflexion of the chorion, and, on that account, is called *decidua reflexa*. The tunica decidua comes away after delivery, in small pieces, mixed with the *lochia*.

DECIMANUS. (From *decem*, ten, and *mane*, the morning.) Returning every tenth day, applied to some erratic fevers.

DECLIVIS. (From *de*, and *clivis*, a hill.) Declining, descending. A name of an abdominal muscle, because of its posture.

DECOCTUM. (From *decoquo*, to boil.) A decoction. Any medicine made by boiling in a watery fluid. In a chemical point of view, it is a continued ebullition with water, to separate such parts of bodies as are only soluble at that degree of heat. The following are among the most approved decoctions.

DECOCTUM ALBUM. See *Decoction cornu*.

DECOCTUM ALOES COMPOSITUM. Compound decoction of aloes. Take of extract of liquorice, half an ounce; subcarbonate of potash, two scruples; extract of spiked aloe powdered, myrrh powdered, saffron stigmata, of each a drachm; water, a pint. Boil down to twelve fluid ounces, and strain; then add compound tincture of cardamoms, four fluid ounces. This decoction, now first introduced in the London Pharmacopœia, is analogous to an article in very frequent use, invented by the late Dr. Devalingin, and sold under the name of *beaume de vie*. By the proportion of tincture, which is added, it will keep unchanged for any length of time.

DECOCTUM ALTHÆÆ. Decoction of marsh mallows. Take of dried marsh mallow roots, ℥iv; raisins of the sun stoned, ℥jij;

water, ℥viij. Boil to five pounds; place apart the strained liquor, till the feces have subsided, then pour off the clear. This preparation, directed in the Edinburgh Pharmacopœia, may be exhibited as a common drink in nephralgia, and many diseases of the urinary passages with advantage.

DECOCTUM ANTHEMIDIS. See *Decoctum chamemeli*.

DECOCTUM ASTRAGALI. Take of the root of the astragalus excapus, ℥j; distilled water, ℔vj. These are to be boiled, till only a quart of fluid remain. The whole is to be taken, a little warmed, in the course of 24 hours. This remedy was tried very extensively in Germany, and said to evince very powerful effects, as an antisyphilitic.

DECOCTUM BARDANÆ. Take of bardana root, ℥vj; of distilled water, ℔vj. These are to be boiled till only two quarts remain. From a pint to a quart in a day is given, in those cases where sarsaparilla and other remedies that are called alterative are supposed to be requisite.

DECOCTUM CHAMEMELI. Chamomile decoction. Take of chamomile flowers, ℥j; caraway seeds, ℥ss; water, ℔ssv. Boil fifteen minutes, and strain. A very common and excellent vehicle for tonic powders, pills, &c. It is also in very frequent use for fomentation and clysters.

DECOCTUM CINCHONÆ. Decoction of cinchona, commonly called decoction of Peruvian bark. Take of lance-leaved cinchona bark bruised, an ounce; water, a pint. Boil for ten minutes, in a vessel slightly covered, and strain the decoction while hot. According to the option of the practitioner, the bark of either of the other species of cinchona, the cordifolia, or yellow, or the oblongifolia, or red, may be substituted for the lancifolia, or quilled; which is here directed. This way of administering the bark is very general, as all the other preparations may be mixed with it, as necessity requires. It is a very proper fomentation for prolapsus of the uterus and rectum.

DECOCTUM CORNU. *Decoctum album.* Decoction of hartshorn. Take of burnt and prepared hartshorn, two ounces by weight; gum-arabic, 6 drachms by weight; distilled water, three pints. Boil, constantly stirring, to two pints, and strain. This decoction, which is omitted in the last edition of the London Pharmacopœia, is a much weaker absorbent than the julep ecreta, but is much more agreeable to most people. It forms an excellent drink in fevers attended with diarrhœa, and acidities of the primæ viæ.

DECOCTUM CYDONIÆ. *Mucilago seminis cydonii mali.* *Mucilago seminum cydoniorum.* Decoction of quince seeds. Take of quince seeds, two drachms; water, a pint. Boil over a gentle fire for ten minutes, then

strain. This decoction, in the new London Pharmacopœia, has been removed from among the mucilages, as being less dense than either of the others, and as being employed in larger doses, like other mucilaginous decoctions. In addition to gum, it contains other constituent parts of the seeds, and is, therefore, more apt to spoil than common mucilage, over which it possesses no other advantages than that it is more grateful, and sufficiently thin, without further dilution, to form the bulk of any liquid medicine. Its virtues are demulcent. Joined with syrup of mulberry, and a little borax, it is useful against aphthæ of the mouth and fauces.

DECOCTUM DAPHNIS MEZEREI. Decoction of mezereon. Take of the bark of mezereon, ℥j; liquorice root bruised, ℥ss; water, ℔vj. Boil it, with a gentle heat, down to two pounds, and strain it. From four to eight ounces of this decoction may be given four times a day, in some obstinate venereal and rheumatic affections. It operates chiefly by perspiration.

DECOCTUM DULCAMARÆ. Decoction of woody nightshade. Take of woody nightshade stalks, newly gathered, ℥j; distilled water, ℔jss. These are to be boiled away to a pint, and strained. The dose is half an ounce to two ounces, mixed with an equal quantity of milk. This remedy is employed in inveterate cases of scrophula; in cancer and phagedæna; in lepra and other cutaneous affections; and in anomalous local diseases, originating in venereal lues.

DECOCTUM GROSSARÆ INERMIS. Decoction of cabbage-tree plant. Take of bark of the cabbage-tree powdered, ℥j; water, ℔vj. Boil it, with a gentle fire, down to one pound, and strain. This is a powerful anthelmintic. It may be given in doses of one table-spoonful to children, and four to adults. If disagreeable symptoms should arise from an over-dose, or from drinking cold water during its action, we must immediately purge with castor-oil, and dilute with acidulated drinks.

DECOCTUM GUIACI OFFICINALIS COMPOSITUM. *Decoctum liquorum.* Compound decoction of guaiacum, commonly called decoction of the woods. Take of guaiacum raspings, ℥jjj; raisins stoned, ℥jj; sassafras root, liquorice, each ℥j; water, ℔x. Boil the guaiacum and raisins, with the water, over a gentle fire, to the consumption of one half; adding, towards the end, the sassafras and liquorice. Strain the liquorice without expression. This decoction possesses stimulant and diaphoretic qualities, and is generally exhibited in rheumatic and cutaneous diseases, which are dependent on a vitiated state of the humours. It may be taken by itself, to the quantity of a quarter of a pint, twice or thrice a day, or used as an assistant in a

course of mercurial or anatomical alteratives; the patient, in either case, keeping warm, in order to promote the operation of the medicine.

DECOCTUM HELLEBORI ALBI. Decoction of hellebore. Take of the root of white hellebore powdered, by weight, $\mathfrak{z}\mathfrak{j}$; distilled water, two pints; rectified spirits of wine, $\mathfrak{z}\mathfrak{j}\mathfrak{j}$ by weight. Boil the water, with the root, to one pint; and the liquor being cold and strained, add to it the spirit. This decoction, in the last London Pharmacopœia is called decoctum veratri. It is, however, a very efficacious application externally, as a wash, in tinea capitis, lepra, psora, &c. When the skin is very tender and irritable, it should be diluted with an equal quantity of water.

DECOCTUM HORDEI. *Decoctum hordei distichi, Apua hordeata.* Take of pearl barley, $\mathfrak{z}\mathfrak{j}\mathfrak{j}$; water, four pints and a half. First wash away any adhering extraneous substances with cold water; next, having poured upon the barley half a pint of water, boil for a few minutes. Let this water be thrown away, and add the remainder of the water boiling; then boil down to two pints and strain. Barley water is a nutritive and softening drink, and the most proper of all liquors in inflammatory diseases. It is an excellent gargle in inflammatory sore throats, mixed with a little nitre.

DECOCTUM HORDEI COMPOSITUM. *Decoctum pectorale.* Compound decoction of barley. Take of decoction of barley, two pints; figs sliced, $\mathfrak{z}\mathfrak{j}\mathfrak{j}$; liquorice root, sliced and bruised, $\mathfrak{z}\mathfrak{s}$; raisins stoned, $\mathfrak{z}\mathfrak{j}\mathfrak{j}$; water, a pint. Boil down to two pints, and strain. From the pectoral and demulcent qualities of this decoction, it may be administered as a common drink in fevers and other acute disorders, in catarrh, and several affections of the chest.

DECOCTUM HORDEI GUMI GUMMI. Barley water, $\mathfrak{lb}\mathfrak{j}\mathfrak{j}$; gum arab. $\mathfrak{z}\mathfrak{j}$. The gum is to be dissolved in the barley decoction whilst warm. It then forms a suitable diluent in strangury, dysury, &c. for the gum, finding a passage into the bladder in an unaltered state, mixes with the urine, and prevents the action of its neutral salts on the urinary canal.

DECOCTUM LICHENIS. Decoction of liverwort. Take of liverwort, one ounce; water, a pint and a half. Boil down and strain. The dose is from $\mathfrak{z}\mathfrak{j}$ to $\mathfrak{z}\mathfrak{iv}$.

DECOCTUM LOBELIS. Take a handful of the roots of the lobelia syphilitica; distilled water, $\mathfrak{lb}\mathfrak{x}\mathfrak{j}\mathfrak{j}$. These are to be boiled in the usual way, till only four quarts remain. The very desirable property of curing the venereal disease has been attributed to this medicine; but it is not more to be depended on than guaiacum, or other vegetable substances, of which the same thing has been alleged. The effects of

this decoction are purgative; and the manner of taking it, as described by Swediaur, is as follows. The patient is to begin with half a pint twice a day. The same quantity is then to be taken four times a day, and continued so long as its purgative effect is not too considerable. When the case is otherwise, it is to be discontinued for three or four days, and then had recourse to again till the cure is completed. As this is a remedy on the old system, and not admitted into our pharmacopœias, little confidence ought to be placed in it.

DECOCTUM LUSITANICUM. Take of sliced sarsaparilla, lignum sassafras, lignum santalum rubrum, official lignum guaiacum, of each one ounce and a half; of the root of mezereon, coriander seed, of each half an ounce; distilled water, ten pounds. These are to be boiled till only half the fluid remains. The dose is a quart or more in a day.

"Take of sliced sarsaparilla, lignum santalum rubrum, lignum santalum citrinum, of each $\mathfrak{z}\mathfrak{j}\mathfrak{s}$; of the root of glycyrrhiza and mezereon, of each $\mathfrak{z}\mathfrak{j}\mathfrak{j}$; of lignum rhodii, official lignum guaiacum, and lignum sassafras, of each $\mathfrak{z}\mathfrak{s}$; of antimony, $\mathfrak{z}\mathfrak{j}$; distilled water, $\mathfrak{lb}\mathfrak{x}$." These ingredients are to be macerated for 24 hours, and afterwards boiled, till the fluid is reduced to half its original quantity. From one to four pints are given daily.

The late Mr. Hunter notices this and also the following formula, in his Treatise on the Venereal Disease.

"Take of sliced sarsaparilla, of the root of China, of each $\mathfrak{z}\mathfrak{j}$; walnut peel dried, $\mathfrak{x}\mathfrak{x}$; antimony, $\mathfrak{z}\mathfrak{j}\mathfrak{j}$; pumice stone, powdered, $\mathfrak{z}\mathfrak{j}$; distilled water, $\mathfrak{lb}\mathfrak{x}$. The powdered antimony and pumice stone are to be tied in separate pieces of rag, and boiled along with the other ingredients." This last decoction is reckoned to be the genuine Lisbon diet drink, whose qualities have been the subject of so much encomium.

DECOCTUM MALVE COMPOSITUM. *Decoction pro enemate. Decoction commune pro clystere.* Compound decoction of mallows. Take of mallows dried, an ounce; chamomile flowers dried, half an ounce; water, a pint. Boil for a quarter of an hour, and strain. A very excellent form for an emollient clyster. A variety of medicines may be added to answer particular indications.

DECOCTUM MEZEREI. See *Decoction daphnes mezerei*.

DECOCTUM PAPAVERIS. *Decoction pro fomento, Fetus communis.* Decoction of poppy. Take of white poppy capsules bruised, $\mathfrak{z}\mathfrak{iv}$; water four pints. Boil for a quarter of an hour, and strain. This preparation possesses antiseptic properties, and may be directed with advantage in spæculus, &c.

DECOCTUM PRO ENEMATE. See *Decoctum mafae compositum*.

DECOCTUM PRO PONENTO. See *Decoctum papaveris*.

DECOCTUM QUERCUS. Decoction of oak bark. Take of oak bark, \mathfrak{zj} ; water, two pints. Boil down to a pint, and strain. This astringent decoction has lately been added to the Lond. Pharm. and is chiefly used for external purposes. It is a good remedy in prolapsus ani, and may be used also in some cases as an injection.

DECOCTUM SARSAPARILLÆ. Decoction of sarsaparilla. Take of sarsaparilla root, sliced, \mathfrak{ziv} ; boiling water, four pints. Macerate for four hours, in a vessel lightly covered, near the fire; then take out the sarsaparilla and bruise it. After it is bruised, put it again into the liquor, and macerate it in a similar manner for two hours more; then boil it down to two pints and strain.

This decoction is much extolled by some practitioners, in phthisis, and to restore the strength after a long course of mercury.

DECOCTUM SARSAPARILLÆ COMPOSITUM. Compound decoction of sarsaparilla. Take of decoction of sarsaparilla, boiling, 4 pints; sassafras root sliced, guaiacum wood shavings, liquorice root bruised, of each an ounce; mezereon root bark, \mathfrak{zjjj} . Boil for a quarter of an hour, and strain. The alterative property of the compound is very great; it is generally given after a course of mercury, where there have been nodes and indolent ulcerations, and with great benefit. The dose is from half a pint to a pint in twenty-four hours.

DECOCTUM SENEÆ. Decoction of senega. Take of senega root, \mathfrak{zj} ; water, two pints. Boil down to a pint, and strain. This is now first introduced in the London Pharm. as being a useful medicine especially in affections of the lungs, attended with debility and inordinate secretion.

DECOCTUM ULMÆ. Decoction of elm bark. Take of fresh elm bark bruised, four ounces; water, four pints. Boil down to two pints, and strain. This may be employed with great advantage as a collyrium in chronic ophthalmia. It is given internally in some cutaneous eruptions.

DECOCTUM VERATRI. See *Decoctum hellobori albi*.

DECOLLATIO. (From *decollo*, to behead.) The loss of a part of the skull.

DECOMPOSITION. A separation of parts. See *Analysis*.

DECORTICATION. (From *de*, from, and *cortex*, bark.) The stripping of any thing of its bark, husk, or shell: thus almonds, and the like, are decorticated, that is, deprived of their pellicle, when ordered for medicinal purposes.

DECREPITATION. (From *decrepo*, to crackle.)

A kind of crackling noise, which takes place in bodies when heated: it is peculiar to some kinds of salts; which, from a state of solution, are crystallized so rapidly, that the crystals formed burst into minute pieces.

DECUSSION. (From *decutio*, to divide.) When nerves, or muscular fibres, cross one another, they are said to decussate each other.

DECUSSUMUM. (From *decusso*, to divide.) An instrument to depress the dura mater, after trepanning.

DEFENSIVA. (From *defendo*, to preserve.) Cordial medicines, or such as resist infection.

DEFERENS. (From *defero*, to convey; because it conveys the semen to the vesiculæ seminales.) See *Vas deferens*.

DEFLAGRATION. (From *deflagro*, to burn.) Calcination. A chymical term, chiefly employed to express the burning or setting fire to any substance; as nitre, sulphur, &c.

DEFLUXION. (From *defluis*, to run off.) *Defluxio*. A falling down of humours from a superior to an inferior part. Many writers mean nothing more by it than inflammation.

DEGLUTITION. (From *deglutis*, to swallow down.) A natural action, by which the masticated bolus or a fluid is conveyed from the mouth into the fauces, and from thence through the œsophagus into the stomach.

DEGMUS. (From *degnus*, to bite.) A biting pain in the orifice of the stomach.

DEJECTIO ALVINA. Discharge of excrement by stool.

DEJECTORIA. (From *dejicio*, to cast out.) Purging medicines.

DEMOSIS. (From *demo*, to exaggerate.) An enlargement of the supercilia.

DELACHRYMATIVA. (From *de*, and *lachryma*, a tear.) Medicines which dry the eyes, first purging them of tears.

DELAPSO. (From *delabor*, to slip down.) A falling down of the anus, uterus, or intestines.

DELETERIOUS. (*Deleterius*; from *deleas*, to hurt, or injure.) Those substances are so called which are of a poisonous nature.

DELIQUESCENCE. Deliquation, or the gradually melting down of crystallized salts, from exposure to the air.

DELIQUIUM ANIMI. (*Deliquium*; from *delinquo*, to leave.) See *Syncope*.

DELIRIUM. (From *deliro*, to rave.) A febrile symptom, consisting in the persons acting or talking unreasonably. It is to be carefully distinguished from an alienation of the mind, without fever.

DELOCATIO. (From *de*, from, and *locus* a place.) A dislocation, or putting any part out of its proper place.

DELPHINIUM. (From *δελφίνος*, the dolphin.) Larkspur; so called from the likeness of its flower to the dolphin's head. The name of a genus of plants in the Linnean system. Class, *Polyandria*. Order, *Trigynia*.

DELPHINIUM CONSOLIDA. The systematic name of the *consolida regalis*. See *Consolida regalis*.

DELPHINIUM STAPHYSAGNIA. The systematic name of stave-acre. See *Staphysagria*.

DELPHYS. *Δελφύς*. The uterus, or pudendum muliebre.

DELTA. (The Greek letter, Δ) The external pudendum muliebre is so called, from the triangular shape of its hair.

DELTOIDES. (From *δελτα*, the Greek letter Δ, and *ομοίος*, a likeness; shaped like the Greek delta.) *Sous-acromio-clavi-humeral* of Damas. A muscle of the superior extremity, situated on the shoulder. It arises exactly opposite to the trapezius, from one third part of the clavicle, from the acromion and spine of the scapula, and is inserted, tendinous, into the middle of the os humeri, which bone it lifts up directly; and it assists with the supra-spinatus and coracobrachialis in all the actions of the humerus, except the depression; it being convenient that the arm should be raised and sustained, in order to its moving on any side.

DEMENTIA. (From *de*, and *mens*, without mind.) Madness. Delirium. Absence of intellect.

DEMULCENTS. (*Demulcentia*, sc. *medicamenta*; from *demulceo*, to soften.) Medicines suited to obviate and prevent the action of acrid and stimulant matters; and that not by correcting or changing their acrimony, but by involving it in a mild and viscid matter, which prevents it from acting upon the sensible parts of our bodies, or by covering the surface exposed to their action.

Where these substances are directly applied to the parts affected, it is easy to perceive how benefit may be derived from their application. But where they are received by the medium of the stomach, into the circulating system, it has been supposed that they can be of no utility, as they must lose that viscosity on which their lubricating quality depends. Hence it has been concluded that they can be of no service in gonorrhœa, and some similar affections. It is certain, however, says J. Murray, in his *Elements of Materia Medica and Pharmacy*, that many substances which undergo the process of digestion, are afterwards separated, in their entire state, from the blood, by particular secreting organs, especially by the kidneys; and it is possible, that mucilaginous substances, which are the principal demulcents, may be separated in this manner. There can be no doubt, however, but that a great

share of the relief demulcents afford, in irritation or inflammation of the urinary passages, is owing to the large quantities of water in which they are diffused, by which the urine is rendered less stimulating from dilution. In general, demulcents may be considered merely as substances less stimulating than the fluids usually applied.

Catarrh, diarrhœa, dysentery, calculus, and gonorrhœa, are the diseases in which demulcents are employed. As they are medicines of no great power, they may be taken in as large quantities as the stomach can bear.

The particular demulcents may be reduced to the two divisions of mucilages and expressed oils. The principal demulcents are the acacia vera, *astragalus tragacantha*, *linum usitatissimum*, *althœa officinalis*, *malva sylvestris*, *glycyrrhiza glabra*, *cycas circinalis*, *orchis mascula*, *maranta arundinacea*, *triticum hybernum*, *ichthyocolla*, *olea Europœa*, *amygdalus communis*, *cetaceum*, and *cera*.

DENDROBIANUS. (From *δένδρον*, a tree, and *οὐκάρω*, frankincense.) The herb rosemary, or frankincense tree.

DENS. (*Quasi edens*; from *edo*, to eat, or from *δεν*, *δένος*.) A tooth. See *Teeth*.

Many herbs have this specific name, from their fancied resemblance to the tooth of some animal; as *dens leonis*, the dandelion; *dens canis*, dog's tooth, &c.

DENS LEONIS. See *Taraxacum*.

DENTAGRA. (*Dentagra*, *δενταγρα*; from *δεν*, a tooth, and *αγρα*, a seizure.) The tooth-ach; also an instrument for drawing the teeth.

DENTARIA. (*Dentaria*, from *dens*, a tooth; so called because its root is denticulated.) *Dentillaria*. Tooth-wort. This plant is to be distinguished from the pelletory of Spain, which is also called dentaria. It is the *Plumbago Europœa* of Linnæus:—*foliis amplexicaulibus, lanceolatis scabris*. The root was formerly esteemed, prepared in a variety of ways, as a cure for the tooth-ach, arising from caries.

DENTARAGA. (From *δεν*, a tooth, and *αγραζω*, to fasten upon.) An instrument for drawing of teeth.

DENTATA. See *Dentatus*.

DENTATUS. (From *dens*, a tooth; from its tooth-like process.) *Dentata*. *Epistropheus*. The second vertebra of the neck. It differs from the other cervical vertebrae, by having a tooth-like process at the upper part of the body. See *Vertebra*.

DENTELLARIA. (From *dentella*, a little tooth; so called because its root is denticulated.) The herb tooth-wort. See *Dentaria*.

DENTES INCISORES. See *Teeth*.

DENTES CANINI. See *Canine teeth*.

DENTES LACTEI. The milk-teeth. See *Teeth*, and *Dentition*.

DENTES MOLARES. See *Teeth*.

DENTIDUCUM. (From *dens*, a tooth, and *duco*, to draw.) An instrument for drawing of teeth.

DENTIFRICE. (From *dens*, a tooth, and *frico*, to rub.) A medicine to clean the teeth.

DENTI-CALPIUM. (From *dens*, a tooth, and *scalpo*, to scrape.) An instrument for scaling teeth.

DENTITION. (From *dentio*, to breed teeth.) The breeding or cutting of the teeth. The first dentition takes place about the sixth or seventh month, and the teeth are termed the *primary* or *milk* teeth. About the seventh year, these fall out, and are succeeded by others, which remain during life, and are called the *secondary* or *perennial* teeth. The last dentition takes place between the ages of twenty and five-and-twenty, when the four last grinders appear; they are called *dentes sapientie*. See also *Teeth*.

DENTODUCUM. See *Dentiducum*.

DENUATIO. (From *denudo*, to make bare.) A laying bare the bone.

DEOBSSTRUENTS. (*Deobstruentia*, sc. *medicamenta*; from *de*, and *obstruo*, to obstruct.) Medicines that are exhibited with a view of removing any obstruction.

DEOPFILANTIA. (From *de*, and *oppilo*, to stop.) *Deoppilativa*. Medicines which remove obstructions; deobstruent or aperitive medicines.

DEPARTITIO. (From *de*, and *partior*, to divide.) Separating metals.

DEPERDITIO. (From *deperdo*, to lose.) Abortion, or the undue loss of the fœtus.

DEPETIGO. (From *de*, and *petigo*, a running scab.) A ring-worm, or tetter. A scurf, or itch, where the skin is rough.

DEPHLEGMATIO. (From *de*, and *phlegma*, phlegm.) The operation of rectifying or freeing spirits from their watery parts.

DEPILATORY. (*Depilatoria*, sc. *unguenta*; from *de*, of, and *pilus*, the hair.) Any application which removes the hairs from any part of the body; thus, a pitch cap pulls the hairs of the head out by the roots.

DEPLUMATIO. (From *de*, and *pluma*, a feather.) A disease of the eyelids, which causes the hair to fall off.

DEPREHENSIO. (From *deprehendo*, to catch unawares.) The epilepsy is so called, from the suddenness with which persons are seized with it.

DEPRIMIO. (From *deprimo*, to press down.) Depression. When the bones of the skull are forced inwards by fracture, they are said to be depressed.

DEPRESSOR. (From *deprimo*, to press down.) Several muscles are so termed, because they depress the part on which they act.

DEPRESSOR ALÆ NASI. See *Depressor labii superioris alæque nasi*.

DEPRESSOR ANGULI ORIS. *Triangularis* of Winslow. *Depressor labiorum communis* of Douglas. *Depressor labiorum* of Cowper. *Sous-maxillo-labial* of Dumas. A muscle of the mouth and lip, situated below the under lip. It arises, broad and fleshy, from the lower edge of the lower jaw, near the chin; and is inserted into the angle of the mouth, which it pulls downwards.

DEPRESSOR LABII INFERIORIS. *Quadratus* of Winslow. *Depressor labii inferioris proprius* of Douglas and Cowper. *Mentonier labial* of Dumas. A muscle of the mouth and lip, that pulls the under lip and skin of the side of the chin downwards, and a little outwards.

DEPRESSOR LABII SUPERIORIS ALÆQUE NASI. *Depressor alæ nasi* of Albinus. *Incisivus medius* of Winslow. *Depressor labii superioris proprius* of Douglas. *Constrictores alarum nasi, ac depressores labii superioris* of Cowper. *Maxillo-alveoli nasal* of Dumas. A muscle of the mouth and lip, situated above the mouth, that draws the upper lip and ala nasi downwards and backwards. It arises, thin and fleshy, from the superior maxillary bone, immediately above the joining of the gums, with the two incisor teeth and cuspidatus; from thence it runs upwards, and is inserted into the upper lip and root of the ala of the nose.

DEPRESSOR LABII SUPERIORIS PROPRIUS. See *Depressor labii superioris alæque nasi*.

DEPRESSOR LABIORUM COMMUNIS. See *Depressor anguli oris*.

DEPRESSOR OCULI. See *Rectus inferior oculi*.

DEPRIMENS. See *Rectus inferior oculi*.

DEPURANTIA. (From *depure*, to make clean.) Medicines which evacuate impurities.

DEPURATION. The freeing a liquor or solid body from its foulness.

DEPURATORIUS. (From *de*, and *purus*, pure.) It is applied to fevers, which terminate in perspiration.

DERIS. (*Δερσις*: from *δερν*, to excoriate.) The skin.

DERIVATIO. (From *derivo*, to drain off.) The doctrines of derivation and revulsion, talked of by the ancients, are now, in their sense of the terms, wholly exploded. Derivation means the drawing away any disease from its original seat to another part.

DERMA. (*Δερμα*.) The skin.

DERMATODES. (From *derma*, skin, and *ωδωρ*, a likeness.) Resembling skin, or leather, in its consistence. It is applied to the dura mater.

DERMATOLOGIA. (From *derma*, the skin, and *λογος*, a discourse.) A discourse or treatise on the skin.

DESTRON. (From *δερν*, skin.) The omentum, or peritonæum, is so named, from its skin-like consistence.

DESCENSORIVM. (From *descendo*, to move downwards.) A vessel in which the distillation by descent is performed.

DESCENSUS. (From *descendo*, to move downwards.) The same chymists call it a distillation *per descensum*, by descent, when the fire is applied at the top and round the vessel whose orifice is at the bottom.

DESICCATIVA. (From *desicco*, to dry up.) Such medicines as, being applied outwardly, dry up the humours and moisture running from a wound.

DESIPIENTIA. (From *desipio*, to dote.) A defect of reason. Symptomatic phrenzy.

DESINE. (From *deso*, to bind up.) A bandage, or ligature.

DESMIDION. (From *desmion*, a handful.) A small bundle, or little bandage.

DESMOS. (From *deso*, to bind up.) A bandage. An inflammatory stricture of a joint, after luxation.

DESPUMATION. (From *despumo*, to clarify.) The clarifying a fluid, or separating its foul parts from it.

DESQUAMATION. (From *desquamo*, to scale off.) The separating of lamina, or scales, from a bone. Exfoliation.

DESQUAMATORIVM. (From *desquamo*, to scale off.) A trepan, or instrument to take a piece out of the skull.

DESTILLATION. See *Distillation*.

DESUDATIO. (From *desudo*, to sweat much.) An unnatural and morbid sweating.

DETENTIO. (From *detineo*, to stop, or hinder.) Epilepsy is so called, from the suddenness with which the patient is seized.

DETERGENTS. (From *detergo*, to wipe away.) Medicines which cleanse and remove such viscid humours as adhere to and obstruct the vessels. Also such applications as clear away foulness from ulcers.

DETONATION. (From *detono*, to make a noise.) Explosion.

DETRACTOR. (From *detracto*, to draw.) applied to a muscle, whose office is to draw the part to which it is attached.

DETRAHENS QUADRATUS. See *Platysma myoides*.

DETRUSOR URINÆ. (From *Detrudo*, to thrust out.) The name of a muscle whose office is to squeeze out the urine. The muscular coat of the urinary bladder was formerly so called.

DEUTERI. (From *δευτερος*, second; because it is discharged next after the fetus.) The secundines, or after-birth.

DEUTEROPATHIA. (From *δευτερος*, second, and *pathos*, a suffering.) An affection or suffering by consent, where a second part suffers, from consent, with the part originally affected, as where the stomach is disturbed through a wound in the head.

Devil's dung. See *Assafœtida*.

DIABERUS. (From *διὰ βῆσαν*, to strengthen; so called, as affording the chief support to the foot.) The ankle-bone.

DIABETES. (From *δια*, through, and *βαίνω*, to pass.) An immoderate flow of urine. A genus of disease in the class *neuroses*, and order *spasmi* of Cullen. There are two species of this complaint: 1. *Diabetes scrobus*, in which there is a superabundant discharge of limpid urine, of its usual urinary taste. 2. *Diabetes mellitus*, in which the urine is very sweet, and contains a great quantity of sugar. Great thirst, with a voracious appetite, gradual emaciation of the whole body, and a frequent discharge of urine, containing a large proportion of saccharine and other matter, which is voided in a quantity far exceeding that of the aliment or fluid introduced, are the characteristics of this disease. Those of a shattered constitution, and those who are in the decline of life, are most subject to its attacks. It not unfrequently attends on hysteria, hypochondriasis, dyspepsia, and asthma; but it is always much milder when symptomatic, than when it appears as a primary affection.

Diabetes may be occasioned by the use of strong diuretic medicines, intemperance of life, and hard drinking; excess in venery, severe evacuations, or by any thing that tends to produce an impoverished state of the blood, or general debility. It has, however, taken place, in many instances, without any obvious cause.

That which immediately gives rise to the disease, has ever been considered as obscure, and various theories have been advanced on the occasion. It has been usual to consider diabetes as the effect of relaxation of the kidneys, or as depending on a general colliqution of the fluids. Dr. Rilcher, professor of medicine in the university of Goettengen, supposes the disease to be generally of a spasmodic nature, occasioned by a stimulus acting on the kidneys; hence a *secretio aucta urinæ*, and sometimes *perversa*, is the consequence. Dr. Darwin thinks that it is owing to an inverted action of the urinary branch of the lymphatics; which doctrine, although it did not escape the censure of the best anatomists and experienced physiologists, met, nevertheless, with a very favourable reception, on its being first announced. The late Dr. Cullen offered it as his opinion, that the proximate cause of this disease might be some fault in the assimilatory powers, or in those employed in converting alimentary matters into the proper animal fluids, which theory has since been adopted by Dr. Dobson, and still later by Dr. Rollo, surgeon general to the royal artillery. The liver has been thought, by some, to be the chief source of the disease; but diabetes is hardly ever attended with any affection of this organ, as has been

proved by frequent dissections; and when observed, it is to be considered as accidental.

The primary seat of the disease is, however far from being absolutely determined in favour of any hypothesis yet advanced, and, from the most attentive consideration of all the circumstances, the weight of evidence appears to induce the majority of practitioners to consider diabetes as depending on a primary affection of the kidneys.

Diabetes sometimes comes on slowly and imperceptibly, without any previous disorder; and it now and then arises to a considerable degree, and subsists long without being accompanied with evident disorder in any particular part of the system; the great thirst which always, and the voracious appetite which frequently occur in it, being often the only remarkable symptoms; but it more generally happens, that a considerable affection of the stomach precedes the coming on of the disease; and that, in its progress, besides the symptoms already mentioned, there is a great dryness in the skin, with a sense of weight in the kidneys, and a pain in the ureters, and the other urinary passages.

Under a long continuance of the disease, the body becomes much emaciated, the feet œdematous, great debility arises, the pulse is frequent and small, and an obscure fever, with all the appearance of hectic, prevail.

The urine in diabetes, from being at first insipid, clear, and colourless, soon acquires a sweetish or saccharine taste, its leading characteristic; and, when subjected to experiment, a considerable quantity of saccharine matter is to be extracted from it.

In some instances, the quantity of urine is much greater than can be accounted for from all the sources united. Cases are recorded, in which 25 to 30 pints were discharged in the space of a natural day, for many successive weeks, and even months; and in which the whole ingesta, as was said, did not amount to half the weight of the urine. To account for this overplus, it has been alleged that water is absorbed from the air by the surface of the body; as also that an extraordinary quantity of water is compounded in the lungs themselves.

Dissections of diabetes have usually shewn the kidneys to be much affected. In some instances they have been found in a loose flabby state, much enlarged in size and of a pale ash colour; in others, they have been discovered much more vascular than in an healthy state, approaching a good deal to what takes place in inflammation, and containing, in their infundibula, a quantity of whitish fluid, somewhat resembling pus, but without any sign of ulcera-

tion whatever. At the same time that these appearances have been observed in their interior, the superficial veins on their surface were found to be much fuller of blood than usual, forming a most beautiful net-work of vessels, the larger branches of which exhibited an absorbent appearance. In many cases of dissection, the whole of the mesentery has been discovered to be much diseased, and its glands remarkably enlarged; some of them being very hard, and of an irregular texture; others softer, and of a uniform spherical shape. Many of the lacteals have likewise been seen considerably enlarged. The liver, pancreas, spleen, and stomach, are in general perceived to be in a natural state; when they are not so, the occurrence is to be considered as accidental. The bladder, in many cases, is found to contain a considerable quantity of muddy urine.

DIABETES HYSTERICUS. Large discharge of urine in hysterical women.

DIABOLUS METALLORUM. Tin.

DIABOTANUM. (From *δια* and *βοτανη*, a herb.) A plaster made of herbs.

DIACADMIAS. (From *δια*, and *καδμια*, cadmia.) The name of a plaster whose basis is cadmia.

DIACALAMINTHES. (From *δια*, and *καλαμινθος*, calamint.) The name of an antidote, whose chief ingredient is calamint.

DIACARCINUM. (From *δια* and *καρκινος*, a crab.) The name of an antidote prepared from the flesh of crabs and crayfish.

DIACARYON. (From *δια*, and *καρυον*, a nut.) Rob of nuts, or wall nuts.

DIACASSIA. (From *δια*, and *κασσια*, cassia.) Electuary of cassia.

DIACASTORIUM. (From *δια*, and *καστωρ*, castor.) An antidote whose basis is castor.

DIACATHOLICON. (From *δια*, and *καθολικος*, universal.) The name of a purge, so called from its general usefulness.

DIACENTAURIUM. (From *δια*, and *κενταυριον*, centaur.) The Duchess of Portland's powder is so called, because its chief ingredient is centaur.

DIACENTROTUM. From *δια*, and *κεντρον*, to prick.) A collyrium, so called from its pungency and stimulating qualities.

DIACHALCITIS. (From *δια* *χαλκισ*, chalcitis.) A plaster whose chief ingredient is chalcitis.

DIACHALSIIS. (From *δια* *χαλασ*, to be relaxed.) A relaxation. The opening of the sutures of the head.

DIACHIRISMUS. (From *δια*, and *χειρ*, the hand.) Any operation performed by the hand.

DIACHELIDONIUM. (From *δια*, and *χελιδονιον*, celandine.) A plaster whose chief ingredient was the herb celandine.

DIACHOREMA. (From *δια* *χορηξω*, to

separate from.) *Diachoresis*. Any excretion or excrement, but chiefly that by stool.

DIACHORESIS. See *Diachorema*.

DIACHERISTA. (From *δια*, and *χρησ*, to anoint.) Medicines to anoint sore or bruised parts.

DIACHERYSUM. (From *δια*, and *χρυσος*, gold.) A plaster for fractured limbs; so named from its yellow colour.

DIACHYLUM. (From *δια*, and *χυλος*, chyle.) The plaster of this name was formerly made of certain juices, but it now means an emollient digestive plaster.

DIACHYSIS. (From *δια*, and *χυω*, to pour out.) Fusion or melting.

DIACHYTICA. (From *διαχυνω*, to dissolve.) Medicines which discuss and dissolve tumours.

DIACINEMA. (From *δια*, and *κινω*, to move.) A slight dislocation.

DIACISSUM. (From *δια*, and *κισσος*, ivy.) An application composed of ivy-leaves.

DIACLASIS. (From *δια*, and *κλαω*, to break.) A small fracture.

DIACLYSMA. (From *διακλυω*, to wash out.) A gargism, or wash for the mouth.

DIACOCCTHELOX. (From *δια*, and *κακκυμυλλος*, a plum.) An electuary made of prunes.

DIACODIUM. (From *δια*, and *καρδια*, a poppy head.) A composition made of the heads of poppies.

DIACOLCYNTHIS. (From *δια*, and *κολοκυνθικη*, the colocynth.) A preparation whose chief ingredient is colocynth.

DIACOMMA. (From *διακομω*, to cut through.) *Diacoephe*. A deep cut or wound.

DIACOPE. See *Diacomma*.

DIACOPRÆGIA. (From *δια*, *κοπρος*, dung, and *αιξ*, a goat.) A preparation with goat's dung.

DIACORALLUM. (From *δια*, and *κοραλλον*, coral.) A preparation in which coral is a chief ingredient.

DIACRISIS. (From *διακρινω*, to distinguish.) The distinguishing diseases one from another by their symptoms.

DIACROCIVM. (From *δια*, and *κροκος*, saffron.) A collyrium in which is saffron.

DIACURCUNA. (From *δια*, and *κυρκμα*, turmeric.) An antidote in which is turmeric or saffron.

DIACYDONIUM. (From *δια*, and *κυδωνια*, a quince.) Marmalade of quinces.

DIADAPHNIDIOM. (From *δια*, and *δαφνικη*, the laurel-tree.) A drawing-plaster in which were bay-berries.

DIADEMA. (From *διαδωω*, to surround.) A diadem or bandage to put round the head.

DIADEXIS. (From *διαδεχομαι*, to transfer.) *Diadoche*. A transposition of humours from one place to another.

DIADOCHES. See *Diadexis*.

DIADOSIS. (From *διαδωδωμαι*, to distribute.) The remission of a disorder.

DIARESIS. (From *διαρρω*, to divide or

separate.) A solution of continuity of the soft parts of the human body.

DIARETICA. (From *διαρρω*, to divide.) *CORTOSIVE* medicines.

DIÆTA. (From *διαίτω*, to nourish.) Diet; food. It means also the whole of the non-naturals. See *Diet*.

DIAGLAUCIUM. (From *δια*, and *γλαυκων*, the blue juice of a herb.) An eye-water made of the purging thistle.

DIAGNOSIS. (From *διαγνωσκω*, to discern or distinguish.) The science which delivers the signs by which a disease may be distinguished from another disease; hence those symptoms which distinguish such affections are termed *diagnostic*.

DIAGRYDIUM. Corrupted from *diacrydium* or scammony.

DIATHERMODACTYLUM. (From *δια*, and *αθηροδωδωλος*, the hermodactyl.) A purging medicine whose basis is the hermodactyl.

DIATRIZON. (From *δια*, and *τρις*, the lily.) An antidote in which is the root of the lily.

DIATUM. (From *δια*, and *τις*, a violet.) A pastel whose chief ingredient is violets.

DIALACCA. (From *δια*, and *λακκα*.) An antidote in which is the lacca.

DIALAGOURM. (From *δια*, and *λαγος*, a hare.) A medicine in which is the dung of a hare.

DIALEMMMA. (From *δια*, and *λεπω*, to leave.) The remission of a disease.

DIALEPSIS. (From *διαλεπω*, to leave a space.) An intermission. Also a space left between a bandage.

DIALIBANUM. (From *δια*, and *λιβανον*, frankincense.) A medicine in which frankincense is a chief ingredient.

DIALOES. (From *δια*, and *αλω*, the aloe.) A medicine chiefly composed of aloes.

DIALTHEA. (From *δια*, and *αλθαια*, the mallow.) An ointment composed chiefly of mallows.

DIALYSIS. (From *διαλυω*, to dissolve.) A solution of continuity, or a destruction of parts.

DIALYSES. An order in the class *locales* of Cullen's nosology.

DIALYTICA. (From *διαλυω*, to dissolve.) Medicines which heal wounds and fractures.

DIAMARGARITON. (From *δια*, and *μαργαριτι*, pearl.) An antidote in which pearls are the chief ingredient.

DIANASSEMMA. (From *δια*, and *μασσιμω*, to chew.) A masticatory, or substance put into the mouth and chewed to excite a discharge of the saliva.

DIAMBRA. (From *δια*, and *αμβρα*, amber.) An aromatic composition in which was ambergris.

DIAMELOX. (From *δια*, and *μελον*, a quince.) A composition in which are quinces.

DIAMISTOS. (From *δια*, and *μισω*, misy.) A composition in which misy is an ingredient.

DIAMOND. The diamond, which was well known to the ancients, is principally found in the western peninsula of India, on the coast of Coromandel, in the kingdoms of Golconda and Visapour, in the island of Borneo, and in the Brazils.

They are generally found bedded in yellow ochre, or in rocks of free-stone, or quartz, and sometimes in the beds of running waters. When taken out of the earth they are incrustated with an exterior earthy covering, under which is another, consisting of carbonate of lime.

In the Brazils, it is supposed that diamonds might be obtained in greater quantities than at present, if the sufficient working of the diamond mines was not prohibited, in order to prevent that diminution of their commercial value, which a greater abundance of them might occasion.

Brazilian diamonds are, in commercial estimation, inferior to the oriental ones.

In the rough, diamonds are worth two pounds sterling the carat, or four grains, provided they are without blemish. The expense of cutting and polishing amounts to about four pounds more. The value however is far above what is now stated when they become considerable in size.

The usual method of calculating the value of diamonds is by squaring the number of carats, and then multiplying the amount by the price of a single carat; thus supposing one carat to be 2*l*. a diamond of 8 carats is worth 128*l*. being $8 \times 8 \times 2$.

The famous Pigot diamond weighs 188 1-8th grains.

Physical Properties of Diamond.

Diamond is always crystallized, but sometimes so imperfectly, that at first sight it might appear amorphous. The figure of diamond when perfect, is an eight-sided prism. There are also cubical, flat, and round diamonds. It is the oriental diamond which crystallizes into octohedra, and exhibits all the varieties of this primitive figure. The diamond of Brazil crystallizes into dodecahedra.

The texture of the diamond is lamellated, for it may be split or cleft with an instrument of well-tempered steel, by a swift blow in a particular direction. There are however some diamonds which do not appear to be formed of *laminae*, but of *distended* and interwoven fibres, like those of knots in wood. These exceed the others greatly in hardness, they cannot be cut or polished, and are therefore called by the lapidaries *diamonds of nature*.

The diamond is one of the hardest bodies known. It resists the most highly-tempered steel file, which circumstance renders it necessary to attack it with diamond powder. It takes an exquisite and lasting polish. It has a great refractive power, and hence its lustre, when cut into the form of a regular solid, is uncommonly great. The usual colour of diamonds is a light gray, often

inclining to yellow, at times lemon colour, violet, or black, seldomer rose-red, and still more rarely green or blue, but more frequently pale brown. The purest diamonds are perfectly transparent. The colourless diamond has a specific gravity which is in proportion to that of water as 3.512 to 1.000, according to Brisson. This varies however considerably. When rubbed it becomes *positively* electric, even before it has been cut by the lapidary.

Diamond is not acted upon by acids, or by any chymical agent, oxygen excepted; and this requires a very great increase of temperature to produce any effect.

The diamond burns by a strong heat, with a sensible flame, like other combustible bodies, attracting oxygen and becoming wholly converted into carbonic acid gas during that process.

It combines with iron by fusion, and converts, it like common charcoal, into steel; but diamond requires much more oxygen to burn in than common charcoal does, and even then it consumes but slowly, and ceases to burn the instant its temperature is lowered.

It is considered by modern chymists as *pure crystallized carbon*. See *Carbon*.

DIAMORON. (From *dia*, and *μωρον*, a mulberry.) A preparation of mulberries.

DIAMOSCRUM. (From *dia*, and *μοσχος*, musk.) An antidote in which musk is a chief ingredient.

DIAMOTOSIS (From *dia*, and *μολος*, lint.) The introduction of lint into an ulcer or wound.

DIANA. (A name of the moon.) The chymical name for silver from its white shining appearance.

DIANANCASMUS. (From *dia*, and *ανακατα*, to force.) The forcible restoration of a luxated part into its proper place. An instrument to reduce a distorted spine.

DIANTHUS. (From *Δις*, *Δις*, Jove, and *ανθος*, a flower; so called from the elegance and fragrance of its flowers.) The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Digynia*.

DIANTHUS CARYOPHYLLUS. The systematic name of the clove-pink. See *Caryophyllum rubrum*.

DIAPASMA. (From *διαπασσω*, to sprinkle.) A medicine reduced to powder and sprinkled over the body, or any part.

DIAPHEDESIS. (From *διαφενδω*, to leap through.) The transudation or escape of blood through the coats of an artery.

DIAPESMA. (From *διαπυρνω*, to close together.) A surgical instrument for closing together broken bones.

DIAPENTE. (From *dia*, and *πεντε*, five.) A medicine composed of five ingredients.

DIAPHANOUS. (From *dia*, through, and *φανω*, to shine.) A term applied to any substance which is transparent; as the hyaloid membrane covering the vitreous

humour of the eye, which is as transparent as glass.

DIAPHENICUM. (From *δια*, and *φενίξ* a date.) A medicine made of dates.

DIAPHORA. (From *διαφορά*, to distinguish.) The distinction of diseases by their characteristic marks and symptoms.

DIAPHORESIS. (From *διαφύω*, to carry through.) Perspiration or increased cutaneous secretion.

DIAPHORETICS. (*Diaphoretica*, sc. *medicamenta*; from *διαφύω*, to carry through.) Medicines which, from being taken internally, increase the discharge by the skin. When this is carried so far as to be condensed on the surface, it forms sweat; and the medicines producing it are named sudorifics. Between diaphoretics and sudorifics there is no distinction; the operation is in both cases the same, and differs only in degree from augmentation of dose, or employment of a-sistant means. This class of medicines comprehends five orders: 1. *Pungent diaphoretics*, as the *volatile salts* and *essential oil*, which are well adapted for the aged; those in whose system there is little sensibility; those who are difficultly affected by other diaphoretics; and those whose stomachs will not bear large doses of medicines. 2. *Calefacient diaphoretics*, such as *serpentaria*, *contrayerva*, and *guaiacum*: these are given in cases where the circulation is low and languid. 3. *Stimulant diaphoretics*, as antimonial and mercurial preparations, which are best fitted for the vigorous and plethoric. 4. *Antispasmodic diaphoretics*, as *opium*, *musk*, and *camphire*, which are given to produce a diaphoresis, when the momentum of the blood is increased. 5. *Diluent diaphoretics*, as water, whey, &c. which are best calculated for that habit in which a predisposition to sweating is wanted; and in which no diaphoresis takes place, although there be evident causes to produce it.

DIAPHRAGMA (From *δια*, and *φάρω*, to divide.) *Septum transversum.* The midriff, or diaphragm. A muscle that divides the thorax from the abdomen. It is composed of two muscles; the first and superior of these arises from the sternum, and the ends of the last ribs on each side. Its fibres, from this semi-circular origination, tend towards their centre, and terminate in a tendon, or aponeurosis, which is termed the *centrum tendinosum*. The second and inferior muscle comes from the vertebræ of the loins by two productions, of which that on the right side comes from the first, second, and third vertebræ of the loins; that on the left side is somewhat shorter, and both these proportions join and make the lower part of the diaphragm, which joins its tendons with the tendon of the other, so that they make but one muscular partition. It is covered by the pleura on its upper side, and by the

peritonæum on the lower side. It is pierced in the middle, for the passage of the vena cava; in its lower part for the œsophagus, and the nerves, which go to the upper orifice of the stomach; and betwixt the productions of the inferior muscle, passes the aorta, the thoracic duct, and the vena azygos. It receives arteries and veins called phrenic or diaphragmatic, from the cava and aorta; and sometimes on its lower part two branches from the vena adiposa, and two arteries from the lumbares. It has two nerves which come from the third vertebra of the neck, which pass through the cavity of the thorax, and are lost in its substance. In its natural situation the diaphragm is convex on the upper side towards the breast, and concave on its lower side towards the belly: therefore, when its fibres swell and contract, it must become plain on each side, and consequently the cavity of the breast is enlarged to give liberty to the lungs to receive air in inspiration; and the stomach and intestines are pressed for the distribution of their contents; hence the use of this muscle is very considerable; it is the principal agent in respiration, particularly in inspiration; for when it is in action the cavity of the thorax is enlarged, particularly at the sides, where the lungs are chiefly situated; and as the lungs must always be contiguous to the inside of the thorax and upper side of the diaphragm the air rushes into them, in order to fill up the increased space. In expiration it is relaxed and pushed up by the pressure of the abdominal muscles upon the viscera of the abdomen; and at the same time that they press it upwards, they pull down the ribs, by which the cavity of the thorax is diminished, and the air suddenly pushed out of the lungs.

DIAPHRAGMATITIS. (From *διαφράγμα*, the diaphragm.) *Diaphragmitis. Paraphrenitis.* An inflammation of the diaphragm. See *Paraphrenitis*.

DIAPHRAGMATITIS. See *Paraphrenitis*.

DIAPHTHORA. (From *διαφθω*, to corrupt.) An abortion where the fœtus is corrupted in the womb.

DIAPHYLACTICA. (From *διαφυλάσσω*, to preserve.) Medicines which resist putrefaction or prevent infection.

DIAPHYSIS. (From *διαφύω*, to divide.) An interstice, or partition between the joints.

DIAPISSELEUM. (From *δια*, and *πισσελαιον*, the oil of pitch, or liquid pitch.) A composition in which is liquid pitch.

DIAPLASIS. (From *διαπλάσσω*, to put together.) The replacing a luxated or fractured bone in its proper situation.

DIAPLASMA. (From *διαπλάσσω*, to anoint.) An unction or fomentation applied to the whole body or any part.

DIAPHNE. (From *διανναι*, to blow through

or pass gently as the breath does.) An involuntary and insensible discharge of the urine.

DIAPHOE. (From *διαφω*, to breathe through.) The transpiration of air through the pores of the skin.

DIAPHOREA. (From *διαφω*, to transpire.) Diaphoretics or medicines which promote perspiration.

DIAPHOREMA. (From *διαφω*, to be in doubt.) Nervous anxiety.

DIAPHORON. (From *δια*, and *πορος*, autumnal fruits.) A composition in which are several autumnal fruits, as quinces, medlars, and services.

DIAPHRASUM. (From *δια*, and *πρασιν*, horehound.) A composition of horehound.

DIAPHRUNUM. (From *δια*, and *πρυν*, a prune.) An electuary of prunes.

DIAPHROBICUM. (From *δια*, and *σφοδ*, the itch or scurvy.) A medicine for the itch or scurvy.

DIAPTERNES. (From *δια*, and *πτερν*, the heel.) A composition of cow heels and cheese.

DIAPTEROSIS. (From *δια*, and *πτερον*, a feather.) The cleaning the ears with a feather.

DIAPTEMA. (From *δια*, and *πυσ*, pus.) A suppuration or abscess.

DIAPTEMATA. (From *διαπτενμα*, a suppuration.) Suppurating medicines.

DIAPYPTICA. (From *διαπτενμα*, a suppuration.) See *Diaptemata*.

DIARHOCHA. (From *δια*, and *ρως*, a space.) The space between the foldings of a bandage.

DIARIUS. (From *dies*, a day.) A term applied to fevers which last but one day.

DIAROMATICUM. (From *δια*, and *αρωματιζω*, an aromatic.) A composition of spices.

DIARRHAGE. (From *διαρρηγνμι*, to break asunder.) A fracture of the temple bones.

DIARRHODOMELI. (From *διαρροδον*, a rose, and *μελι*, honey.) Scammony, agaric, pepper and honey.

DIARRHODON. (From *δια*, and *ροδον*, a rose.) A composition of roses.

DIARRHŒA. (From *διαρρω*, to flow through.) A purging. It is distinguished by frequent stools with the natural excrement, not contagious, and seldom attended with pyrexia. It is a genus of disease in the class *neuroses*, and order *spasmi* of Cullen, containing the following species: 1. *Diarrhœa crapulosa*. The feculent diarrhœa, from *crapulus*, one who overloads his stomach. 2. *Diarrhœa biliosa*. The bilious, from an increased secretion of bile. 3. *Diarrhœa mucosa*. The mucous, from a quantity of slime being voided. 4. *Diarrhœa hepaticæ*. The hepatic, in which there is a quantity of serous matter, somewhat resembling flesh, voided; the liver being primarily affected. 5. *Diarrhœa lenterica*. The lenteric; when the food passes

unchanged. 6. *Diarrhœa cœliaca*. The cœliac passion: the food passes off in this affection in a white liquid state like chyle.

7. *Diarrhœa verminea*. Arising from worms.

DIARTHROSIS. (From *διαρθρω*, to articulate.) A moveable connexion of bones. This genus has five species, viz. *enarthrosis*, *arthrodia*, *ginglymus*, *trochoides*, and *amphiarthrosis*.

DIASAPONIUM. (From *δια*, and *σαπων*, soap.) An ointment of soap.

DIASATTIRIUM. (From *δια*, and *σαττιρον*, the orchis.) An ointment of the orchis-root.

DIASCILLIUM. (From *δια*, and *σκυλλα*, the squill.) Oxyinel and vinegar of squills.

DIASCINCUS. (From *δια*, and *σινκε*, the crocodile.) A name for the mithridate, in the composition of which there was a part of the crocodile.

DIASCORDIUM. (From *δια*, and *σχοδρον*, the water germander.) Electuary of scordium.

DIASENA. (From *δια*, and *sena*.) A medicine in which is senna.

DIASMYRNUM. (From *δια*, and *σμιρνη*, myrrh.) A collyrium containing myrrh.

DIABOSTICA. (From *διασσω*, to preserve.) Medicines which preserve health.

DIASPERMATUM. (From *δια*, and *σπέρμα*, seed.) A medicine composed chiefly of seeds.

DIASPHAGE. (From *διασφαζω*, to separate.) *Diaphaxis*. The interstice between two veins.

DIASPHYXIS. (From *δια*, and *σφυζω*, to strike.) The pulsation of an artery.

DIASTASIS. (From *διαστημι*, to separate.) *Diastema*. A separation.

1. A separation of the ends of bones.
2. (From *διασσω*, to distend.) The extension of a fractured limb, in order to reduce it.

DIASTETUMA. (From *δια*, and *στυπ*, fat.) An ointment of the fat of animals.

DIASTEMA. See *Diastasis*.

DIASTOLE. (From *δια*, and *στέλλω*, to stretch.) The dilatation of the heart and arteries.

DIASTOMOSIS. (From *διασσω*, to dilate.) Any dilatation, or dilating instrument.

DIASTREMA. (From *διαστροφω*, to turn aside.) *Diastrophe*. A distortion of any limb or part.

DIASTROPE. See *Diastrema*.

DIATECOLITHUM. (From *δια*, and *τεκολιθος*, the Jew's stone.) An antidote containing lapis judaicus.

DIATERESIS. (From *δια*, and *τερεω*, to perforate.) A perforation or aperture.

DIATERETICA. (From *δια*, and *τερεω*, to preserve.) Medicines which preserve health and prevent disease.

DIATESSARON. (From *δια*, and *τεσσαρες*, four.) A medicine compounded of four simple ingredients.

DIATETIUM. (From *dia*, and *trētē*, a grasshopper.) A medicine in the composition of which were grasshoppers.

DIATHESIS. (From *diathesis*, to dispose.) Any particular state of the body: thus, in inflammatory fever, there is an inflammatory diathesis, and, during putrid fever, a putrid diathesis.

DIATHESMUS. (From *diathesis*, to run through.) A rupture through which some fluid escapes.

DIATRAGACANTHUM. (From *dia*, and *τραγακάνθη*, tragacanth.) A medicine composed of gum tragacanth.

DIATRUM. (From *dia*, and *τρεῖς*, three.) A medicine composed of three simple ingredients.

DIAXYLALOE. (From *dia*, and *ξύλαλον*, the ligni aloes.) A medicine in which is lignum aloes.

DIAXOMA. (From *diastomum*, to surround; because it surrounds the cavity of the thorax.) The diaphragm.

DIAZOSTER. (From *diastomum*, to surround; because when the body is girded, the belt usually lies upon it.) A name of the twelfth vertebra of the back.

DICENTETUM. (From *dia*, and *κίωσις*, to stimulate.) A pungent stimulating collyrium.

DICHESTERES. (From *diachein*, to divide, because they divide the foot.) A name of the fore-teeth.

DICHOPHYLLA. (From *diχα* double, and *φυον*, to grow.) A distemper of the hairs, in which they split and grow forked.

DICROTIC. (*Dicroticus*, sc. *pulsus*; from *dia*, twice, and *κρούω*, to strike.) A term given to a pulse in which the artery rebounds after striking, so as to convey the sensation of a double pulsation.

DICTAMNITES. (From *dictamnus*, dittany.) A wine medicated with dittany.

DICTAMNUS. (From *Dictamnus*, a city in Crete, on whose mountains it grows.) The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Monogynia*. Dittany.

DICTAMNUS ALBUS. White fraxinella, or bastard dittany *Fraxinella*. *Dictamnus albus*; *foliis pinnatis, caule simpliciter*, of Linnæus. The root of this plant is the part directed for medicinal use; when fresh, it has a moderately strong, not disagreeable, smell. Formerly it was much used as a stomachic, tonic, and alexipharmic, and was supposed to be a medicine of much efficacy in removing uterine obstructions and destroying worms; but its medicinal powers became so little regarded by modern physicians, that it had fallen almost entirely into disuse, till Baron Stœrck brought it into notice, by publishing several cases of its success, viz. in tertian intermittents, worms (lunatic) and menstrual suppressions. In all these cases, he

employed the powdered root to the extent of a scruple twice a day. He also made use of a tincture, prepared of two ounces of the fresh root digested in 14 ounces of spirit of wine; of this, 22 to 50 drops two or three times a day, were successfully employed in epilepsies, and, when joined with steel, this root, we are told, was of great service to chlorotic patients. The dictamnus undoubtedly, says Dr. Woodville, is a medicine of considerable power; but notwithstanding the account of it given by Stœrck, who seems to have paid little attention to its *modus operandi*, we may still say with Haller "*nondum autem vires pro dignitate exploratus est*," but is now fallen into disuse.

DICTAMNUS CRETICUS. Dittany of Crete. *Origanum creticum*. *Onitis*. The leaves of this plant, *Origanum dictamnus*; *foliis inferioribus tomentosis, apicibus mutantibus*, of Linnæus, are now rarely used; they have been recommended as emmenagogue and alexipharmic.

DIDYMÆA. (From *didymos*, double.) A cataplasm; so called by Galen, from the double use to which he put it.

DIDYMI. (From *didymos*, double.) Twins. An old name of the testicles, and two eminences of the brain, from their double protuberance.

DROBOLIUM. (From *dia*, and *βάλλω*, to cast out.) A medicine causing an abortion.

DIELECTRON. (From *diez*, and *ελεκτρον*, amber.) A name of a troche, in which amber is an ingredient.

DIERVILLA. (Named in honour of Mr. Dierville, who first brought it from Arcadia.) The young branches of this species of honeysuckle, *Lonicera diervilla racemis terminalibus, foliis serratis* of Linnæus, are employed in North America as a certain remedy in gonorrhœa and suppression of urine. It has not yet been exhibited in Europe.

DIET. *Dieta*. The dietetic part of medicine is no inconsiderable branch, and seems to require a much greater share of regard than it commonly meets with. A great variety of diseases might be removed by the observance of a proper diet and regimen, without the assistance of medicine, were it not for the impatience of the sufferers. However, it may on all occasions come in as a proper assistant to the cure, which sometimes cannot be performed without a due observance of the non-naturals. That food is, in general, thought the best and most conducive to long life, which is most simple, pure, and free from irritating, and such as approaches nearest to the nature of our own bodies in a healthy state, or capable of being easiest converted into their substance by the vis vitæ, after it has been duly prepared by

the art of cookery; but the nature, composition, virtues, and uses of particular aliments can never be learnt to satisfaction, without assistance of practical chymistry.

DIET DRINK. An alterative decoction employed daily in considerable quantities, at least from a pint to a quart. The decoction of sarsaparilla and mezereon, the Lisbon diet drink, is the most common and most useful.

DIETETICS. That part of medicine which considers the way of living with relation to food, or diet, suitable to any particular case.

DIXODOS. (From *dis*, and *ἄδος*, a way to pass out.) *Diodos*. In Hippocrates it means evacuation by stool.

DIFFLATIO. (From *difflo*, to blow away.) Perspiration,

DIGASTRICUS. (*Digastricus*, sc. *musc.* from *dis*, twice, and *γαστρῆς*, a belly.) *Biventer maxillæ* of Albinus. *Mastoido hygenien* of Dumas. A muscle so called from its having two bellies, situated externally between the lower jaw and *os hyoides*. It arises, by a fleshy belly, from the upper part of the processus mastoideus, and descending, it contracts into a round tendon, which passes through the stylohyoidæus, and an annular ligament which is fastened to the *os hyoides*; then it grows fleshy again, and ascends towards the middle of the edge of the lower jaw, where it is inserted. Its use is to open the mouth by pulling the lower jaw downwards and backwards; and when the jaws are shut to raise the larynx, and consequently the pharynx, upwards, as in deglutition.

DIGERENTIA. (From *digero*, to digest.) Medicines which promote the secretion of proper pus in wounds and ulcers.

DIGESTION. (*Digestio*, from *digero*, to dissolve.)

1. An operation in which such matters as are intended to act slowly on each other, are exposed to a slow heat, continued for some time.

2. The change that the food undergoes in the stomach, by which it is converted into chyme. The circumstances necessary to effect a healthy digestion of the food are: 1. A certain degree of heat of the stomach. 2. A free mixture of saliva with the food in the mouth. 3. A certain quantity of healthy gastric juice. 4. The natural peristaltic motion of the stomach. 5. The pressure of the contraction and relaxation of the abdominal muscles and diaphragm. From these circumstances, the particles of the food are softened, dissolved, diluted, and intimately mixed into a soft pap, called chyme, which passes through the pylorus of the stomach into the duodenum. The fluid which is termed gastric juice, is separated by the minute arteries opening into the cavity of the stomach.—

See *Gastric Juice*. From various experiments of physiologists, it is ascertained that the gastric juice reduces the aliments into an uniform pap or paste, even out of the body; that it acts in the same manner after death; and that it is the chief agent in the process of digestion.

Animals only are invested with organs of digestion; every one, from man to the poly-pus, presents an alimentary canal differently formed; the existence of a digestive apparatus, then, could be given as an essential character of animal life. In man this consists of a long canal extending from the mouth to the anus, into which open the excretory ducts of various glands, situated in the vicinity, that secrete liquors, necessary to alter, liquefy, and animalize alimentary matter.

It would be useless to recapitulate the hypotheses formed to explain digestion; they may be reduced to *coction, fermentation, trituration, putrefaction, and maceration* of the food received into the cavity of the stomach. Physiologists are generally agreed, at present, in considering digestion in the stomach as a solution of the aliment by the gastric juice. This liquid, copiously poured on the internal surface of the stomach, when this viscus is irritated by the presence of food, is the production of arterial exhalation; it is neither an acid nor alkali, and seems to be of a nature nearly analogous to saliva; the gastric juice producing great properties of solutions, penetrates into the alimentary matter on all sides, separates and divides its particles, combines with it, changes its composition, and impresses qualities very different from those it possessed before this mixture. In fact, if a mouthful of wine or food be returned from the stomach some minutes after it has been received, the odour, taste, and all the qualities, both physical and chymical, of these substances, are so altered, that we can with difficulty distinguish them; and vinous liquors, more or less acid, are no longer susceptible of spirituous fermentation. The energy of the power of the gastric juice, perhaps exaggerated by some physiologists, is sufficient to reduce to a soft mass the hardest bones, on which certain animals subsist: it is very probable that its chymical composition is different and variable, and that it is acid, alkaline, or saponaceous, according to the nature of the aliment. Although gastric juice is the most powerful agent of digestion in the stomach, its dissolvent power has need of assistance from the action of several secondary causes, as heat, which seems to augment and concentrate itself in the epigastric region. So long as the exertion of the stomach continues, there is a sort of intestine fermentation, which should not be, in the full sense, compared to the motion by which fermentative and putre-

scent substances are decomposed ; there is also a moderate and peristaltic motion of the muscular fibres of the stomach which press the aliment on all sides, and perform a slight trituration, while the gastric moisture softens and macerates the food before it is dissolved ; it may then be affirmed that the process of digestion is at the same time chymical, vital, and mechanical : the authors, therefore, of various theories to explain this function, have erred by attributing to one cause only, as heat, fermentation, putrefaction, trituration, maceration, and the gastric juice, that which is the aggregate result of all those causes united.

The aliment remains a greater or less time in the stomach, agreeably to the facility or difficulty of the necessary changes taking place. Gossé of Geneva, has proved on himself, that the animal and vegetable fibre, the white of an egg boiled, white and tendinous parts, paste kneaded with butter, sebaceous substances, and those things which are not fermented, or very little fermentative, make greater resistance to the gastric juice, than the gelatinous parts of vegetables and animals, fermented bread, &c. that the latter class of substances require only an hour for their complete dissolution, while the digestion of the former was not completed at the end of several hours.

During the time of digestion, both orifices of the stomach are closed ; no gas, disengaged from the aliment, ascends through the œsophagus, unless in cases of bad digestion : slight chills are felt ; the pulse becomes quicker and stronger, and the powers of life seem diminished in some organs, to be carried to the seat of the digestive process. The parieties of the stomach soon begin to act : their circular fibres contract in different parts of its extent ; these peristaltic oscillations, at first vague and uncertain, become more regular, and are directed from above downwards, from left to right, that is, from the cardia, towards the pylorus ; its longitudinal fibres also contract, and thus approximate both terminations. In these different motions the stomach becomes parallel with the pylorus, and the angle formed by the duodenum is almost totally obliterated, which renders the passage of food easier. It has been remarked, that digestion proceeds better during sleep, when we lie on the right side than on the left, and this difference has been attributed to the compression made by the liver on the stomach. It should rather be considered, that on the right side, the passage of food is accelerated by its own gravity, the situation of the stomach is naturally oblique, from left to right, and becomes more so in consequence of changes induced by food.

The aperture of the pylorus is furnished with a muscular ring, covered by a duplication of mucous membrane ; this kind of sphincter keeps it closed during the time of digestion in the stomach, and does not give passage to the aliment until it has undergone a very material alteration. The pylorus, possessing a peculiar and extremely delicate sensibility, may be considered as a sort of vigilant centinel, that prevents any thing from passing that has not suffered proper changes. Many authors quoted by Haller, have been well aware that the food did not pass from the stomach successively in the same manner as it was received, but agreeably to its greater or less facility of digestion.

It would appear that there is a real selection of food in the stomach, for those aliments that admit of an easy digestion are directed towards the pylorus, which gives passage to them ; while, on the contrary, such as are not sufficiently digested are not permitted to pass, but kept back in the stomach. This delicate feeling, which we attribute to the pylorus, this exquisite sense, by which it exerts a kind of choice on the food that passes through, may be perhaps objected to ; pieces of money, however, or other extraneous indigestible bodies, remain a longer or shorter time in the stomach before they go into the intestines, and present themselves several different times at the orifice of the pylorus, and do not get through till after it has been accustomed to their contact. It is the same with the gastric system as with a secreting gland ; and in the same manner the commencement of excretory ducts, possessed of a sort of *elective* sensibility, do not receive the secreted liquor before it has undergone necessary preparations in the glandular parenchyma ; so the pylorus, which may be considered the excretory duct of the stomach, does not admit the food nor suffer it to pass into the intestines, until it has been sufficiently elaborated by the action of this organ.

In proportion as the stomach becomes empty, the spasm of the skin ceases, a moderate heat succeeds the shiverings, the pulse becomes more evident and elevated, the quantity of insensible perspiration increases ; digestion then produces a general motion, analogous to a feverish paroxysm ; and this digestive fever, described also by the ancients, is most easy to be observed in women of great sensibility. Nothing positive can be established on the duration of digestion in the stomach. The aliments go out of the stomach with more or less celerity, in proportion as they offer a greater or less resistance to those powers which serve to dissolve them, and agreeably to the energy and strength of the stomach and activity of the gastric juice : five hours, however,

may be considered the ordinary time of their presence.

The action of the parietics of the stomach ceases when this viscus is entirely liberated from the aliments that were in its cavity, but not before; the gastric juice, the secretion of which is not augmented by any stimulus, is no longer poured out by its arteries; and the parietics which come into contact with each other, are only lubricated by the mucous copiously secreted by its internal coat.

DIGESTIVES. (*Digestiva*, sc. *medicamentaria*; from *digero*, to dissolve.) A term applied by surgeons to those substances which, when applied to an ulcer or wound, promote suppuration: such are the *ceratum resine flavae*, *unguentum elemi*, warm poultices, fomentations, &c.

DIGESTIVUM SAL SYLVII. See *Murias potasse*.

DIGITALIS. (From *digitus*, a finger; because its flower represents a finger.) 1. The name of a genus of plants in the Linnean system. Class, *Didynamia*. Order, *Angiospermia*. Fox-glove.

2. The pharmacopœial name of the common fox-glove. *Digitalis purpurea* of Linneus:—*calycidis foliolis ovatis acutis, corollis obtusis, labio superiore integro*. The leaves of this plant have a bitter nauseous taste, but no remarkable smell; they have been long used externally to ulcers and scrophulous tumours with considerable advantage. When properly dried, their colour is a lively green. They ought to be collected when the plant begins to blossom, to be dried quickly before the fire, and preserved unpowdered.

Of all the narcotics, digitalis is that which diminishes most powerfully the actions of the system; and it does so without occasioning any previous excitement. Even in the most moderate dose, it diminishes the force and frequency of the pulse, and, in a large dose, reduces it to a great extent, as from 70 beats to 40 or 35 in a minute, occasioning, at the same time, vertigo, indistinct vision, violent and durable sickness, with vomiting. In a still larger quantity, it induces convulsions, coldness of the body, and insensibility; symptoms which have sometimes terminated fatally. As a narcotic, fox-glove has been recommended in epilepsy, insanity, and in some acute inflammatory diseases. Lately it has been very extensively employed in phthisis, and the beneficial effects which it produces in that disease, are probably owing to its narcotic power, by which it reduces the force of the circulation through the lungs and general system. It is administered so as to produce this effect. One grain of the powdered leaves, or ten drops of the saturated tincture, may be given night and morning. This dose is increased one half every second day, till its action on the sys-

tem becomes apparent. As soon as the pulse begins to be diminished, the increase of dose must be made with more caution; and, whenever nausea is induced, it ought rather to be reduced, or, if necessary, intermitted for a short time. If the sickness become urgent, it is best relieved by stimulants, particularly large doses of brandy, with aromatics. The tincture has been supposed to be the best form of administering digitalis, when the remedy is designed to act as a narcotic: it is also more manageable in its dose and more uniform in its strength, than the dried leaves.

Besides its narcotic effects, digitalis acts as one of the most certain diuretics in dropsy, apparently from its power of promoting absorption. It has frequently succeeded where the other diuretics have failed. Dr. Withering has an undoubted claim to this discovery; and the numerous cases of dropsy related by him, and other practitioners of established reputation, afford incontestable evidence of its diuretic powers, and of its practicable importance in the cure of those disorders. From Dr. Withering's extensive experience of the use of the digitalis in dropsies, he has been able to judge of its success by the following circumstances:—"It seldom succeeds in men of great natural strength, of tense fibre, of warm skin, of florid complexion, or in those with a tight and cordy pulse. If the belly in ascites be tense, hard, and circumscribed, or the limbs in anasarca solid and resisting, we have but little hope. On the contrary, if the pulse be feeble, or intermitting, the countenance pale, the lips livid, the skin cold, the swollen belly soft and fluctuating, the anasarcaous limbs readily pitting under the pressure of the finger, we may expect the diuretic effects to follow in a kindly manner." Of the inferences which he deduces, the fourth is, "that if it (digitalis) fails, there is but little chance of any other medicine succeeding." Although the digitalis is now generally admitted to be a very powerful diuretic, yet it is but justice to acknowledge that this medicine has more frequently failed than could have been reasonably expected from a comparison of the facts stated by Dr. Withering. The dose of the dried leaves in powder, is from one to three grains twice a day. But if a liquid medicine be preferred, a drachm of the dried leaves is to be infused for four hours, in half a pint of boiling water, adding to the strained liquor an ounce of any spirituous water. One ounce of this infusion, given twice a day, is a medium dose. It is to be continued in these doses till it either acts upon the kidneys, the stomach, the pulse, (which as has been said, it has a remarkable power of lowering,) or the bowels.

The administration of this remedy re-

quires to be conducted with much caution. Its effects do not immediately appear; and when the doses are too frequent, or too quickly augmented, its action is concentrated so as to produce frequently the most violent symptoms. The general rules are, to begin with a small dose, to increase it gradually, till the action is apparent on the kidneys, stomach, intestines, or vascular system; and immediately suspending its exhibition, when its effects on any of these parts take place.

The symptoms arising from too large a dose of digitalis are, extreme sickness, vertigo, indistinct vision, incessant vomiting, and a great reduction of the force of the circulation, terminating sometimes in syncope or convulsions. They are relieved by frequent and small doses of opium, brandy, aromatics, and strong bitters, and by a blister applied to the region of the stomach.

DIGITUM. (From *digitus*, a finger.) A contraction of the finger-joint. A paronychia, or whitlow, or other sore upon the finger.

DIGITUS. (From *directus*, to direct.) A finger.

DIGITUS MANUS. A finger. The fingers and thumb in each hand consist of fourteen bones, there being three to each finger; they are a little convex and round towards the back of the hand, but hollow and plain towards the palm, except the last, where the nails are. The order of their dispositions is called first, second, and third *phalanx*. The first is longer than the second, and the second longer than the third. What has been said of the fingers, applies to the toes also.

DIGITUS PEDIS. A toe. See *Digitus manus*.

DIGLOSSUM. (From *duo*, double, and *glossa*, a tongue; so called because above its leaf there grows a lesser leaf, like two tongues.) The *laurus alexandrina*. Galen speaks of a man born with two tongues.

DIGNOSCO. (From *dignosco*, to distinguish.) See *Diagnosis*.

DIGNATON. (From *duo*, and *anima*, blood.) An antidote in which is the blood of many animals.

DIALON. (From *duo*, and *aleo*, salt.) A plaster prepared with salt and nitre, adapted to foul ulcers.

DIAPTES. (From *Zeus*, *duo*, Heaven, and *aptes*, to fall: i. e. falling as rain.) An epithet applied by Hippocrates to semen, when it is discharged like a sudden shower of rain.

DILATATIO. (From *dilato*, to enlarge.) Dilatation, or enlargement. Diastole.

DILATOR. (From *dilato*, to enlarge.) The name of some muscles whose office is to open and enlarge parts.

DILATORES ALAARUM NASI. See *Levator*.

DILATORIUM. (From *dilato*, to enlarge.)

A surgical instrument for enlarging any part. A speculum oris.

Dill. See *Anethum*.

DILUENTS. (*Diluentia*, sc. *medicamenta*; from *diluo*, to wash away.) Those substances which increase the proportion of fluid in the blood. It is evident that this must be done by watery liquors. Water is indeed, properly speaking, the only diluent. Various additions are made to it, to render it pleasant, and frequently to give it a slightly demulcent quality. But these are not sufficiently important to require to be noticed, or to be classed as medicines.

Diluents are merely secondary remedies. They are given in acute inflammatory diseases, to lessen the stimulant quality of the blood. They are used to promote the action of diuretics in dropsy, and to favour the operation of sweating.

DINICA. (From *diuo*, giddiness.) Medicines which relieve a giddiness.

DIREA. (From *diro*, to turn round.) A vertigo, or giddiness.

DIOCRES. The name of a lozenge.

DIODOS. (From *duo*, and *odo*, the way through.) Evacuation by stool.

DIERANTHEM. (From *duo*, and *anthen*, the flower of the vine.) A remedy said to be good for cholera morbus, in which was the flower of the vine-tree.

DIOENUS. (From *diuo*, to persecute.) A distressing palpitation of the heart.

DIONYSISCUS. (From *diuo*, Bacchus, was of old represented as having horns.) Certain bony excrescences, near the temples, were called dionysisci.

DIONYSIOPHYLLA. (From *diuo*, Bacchus, and *phylla*, a nymph.) A herb which, if bruised, smells of wine, and yet resists drunkenness.

DIAPORUM. (From *duo*, and *porum*, autumnal fruits.) A medicine composed of ripe fruits for quincy.

DIOPTRA. (From *dioptra*, to see through.) *Dioptron*. Speculum ani oris, or uteri. Also the lapis specularis.

DIOPTRICA. (From *dioptra*, to see through.) Dioptrics, or doctrine of the refraction of light.

DIOPTRISUM. (From *dioptra*, to see through.) Dilatation of any natural passage.

DIORUBUM. (From *duo*, and *agros*, a vetch.) A medicine, in the composition of which there are vetches.

DIOREXOSIS. (From *duo*, and *exor*, the serum.) *Diorosis*. A dissolved state of the blood. A conversion of the humours into serum and water.

DIORTHROSIS. (From *dioptra*, to direct.) The reduction of a fracture.

DIOSCOREA. (Named in honour of Dioscorides.) The name of a genus of plants in the Linnæan system. Class, *Diacia*. Order, *Hexandria*.

DIOSCOREA ALATA. See *Yam*.

DIOSCOREA BULBIFERA. See *Yam*.

DIOSCOREA SATIVA. See *Yam*.

DIOSCURI. (i. e. *Διοσ Κυριοι*, the sons of Jupiter, or Castor and Pollux.) The parotids were so named from their twin-like equality in shape and position.

DIOSEPYROS LOTUS. See *Indian date plum*.

DIOXELÆUM. (From *δια*, *οξύς*, acid, and *αἶμα*, oil.) A medicine composed of oil and vinegar.

DIOXOS. (From *δια*, and *οξύς*, acid.) A collyrium composed chiefly of vinegar.

DIPLESIASMUS. (From *διπλοος*, to double.) The re-exacerbation of a disease.

DIPLOE. (From *διπλοος*, to double.) *Meditullum*. The spongy substance between the two tables of the skull.

DIPLOPIA. (From *διπλος*, double, and *οπταμαι*, to see.) *Virus duplicatus*. A disease of the eye, in which the person sees an object double or triple. Dr. Cullen makes it a variety of the second species of pseudoblepsia, which he calls mutans, in which objects appear changed from what they really are: and the disease varies according to the variety of the remote cause.

DIPYROS. (From *δις*, twice, and *πναι*, to breathe.) An epithet for wounds which are perforated quite through, and admit the air at both ends.

DIPSACUS. (From *διψα*, thirst; so called from the concave situation of its leaves, which hold water, by which the thirst of the traveller may be relieved.) *Diapacum*.

1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia*. The teasel.

2. A diabetes, from the continual thirst attending it.

DIPYRENUM. (From *δις*, twice, and *πυρ*, a berry.) A berry, or kernel: a probe with two buttons.

DIPYRITES. (From *δις*, twice, and *πυρ*, fire.) *Dipyros*. An epithet given by Hippocrates to bread twice baked, and which he recommended in dropsies.

DIRECTOR. (From *dirigo*, to direct.) A hollow instrument for guiding an incisor-knife; also the name of a muscle which lifts up the penis.

DIRECTORES PENIS. (From *dirigo*, to direct.) The same as *erectores penis*.

DIRINGA. A name, in the isle of Java, for the *calamus aromaticus*.

DISCESSUS. (From *discedo*, to depart.) The separation of any two bodies, before united by chymical operation.

DISCIFORMIS. (From *discus*, a quoit, and *forma*, likeness.) Resembling a disk, or quoit, in shape. It is applied to the knee-pan.

DISCORDES. (From *δισκος*, a quoit.) Resembling a disk, or quoit, in shape. It is applied to the crystalline humour of the eye.

DISCROMEN. A small roller. A term applied to the diaphragm.

DISCUTIENTS. (*Discutientia*, sc. *medicamenta*; from *discutio*, to make in pieces.) *Discusoria*. *Diachytica*. A term in surgery applied to those substances which possess a power of repelling or resolving tumours.

DISEASE. *Morbus*. Any alteration from a perfect state of health is a disease. A disease is variously termed: when it pervades the whole system, and does not depend on any other disease, as an inflammatory fever, for instance, it is called a *general disease*, to distinguish it from inflammation of the eye, or any other viscus, which is a *partial* or *local* one; and when it does not depend on another disease, it is termed an *idiopathic disease*, which may be either general or partial, to distinguish it from a *symptomatic affection*, which depends upon another disease, and is produced by consent of parts. See also *Endemic*, *Epidemic*, *Sporadic*, &c.

DISLOCATION. (From *disloco*, to put out of place.) *Luxation*. The secession of a bone of a moveable articulation from its natural cavity.

DISPENSARY. (*Dispensarium*, from *dispendo*, to distribute.) The shop, or place, in which medicines are prepared. Also the name of an institution, in which the poor are supplied with medicines and advice.

DISPENSATORY. (*Dispensatorium*; from *dispendo*, to distribute.) *Antidotarium*. A book which treats of the composition of medicines.

DISSECTION. (From *disseco*, to cut asunder.) The cutting to pieces of any part of an animal, or vegetable, for the purpose of examining its structure.

DISEPTUM. (From *dissepio*, to inclose round.) The diaphragm, or membrane, which divides the cavity of the thorax from the abdomen.

DISSOLVENTIA. (From *dissolvo*, to loosen.) Medicines which loosen and dissolve morbid concretions in the body. In chymistry it means *menstrua*.

DISSOLUTUS. (From *dissolvo*, to loosen.) Loose. An epithet applied to the dysentery, or *morbus dissolutus*.

DISTENTIO. (From *distendo*, to stretch out.) Distention, or dilatation. A convulsion.

DISTICHTA. See *Distichiasis*.

DISTICHIASIS. (From *διστιχια*: from *δις*, double, and *στιχ*, a row.) *Distichiasis*. *Distichia*. A disease of the eyelash, in which there is a double row of hairs, the one row growing outwards, the other inwards towards the eye.

DISTILLATION. (From *distillo*, to drop little by little.) *Alaecta*. *Catastagnos*. A chymical process, very similar to evaporation, instituted to separate the volatile from the fixed principles by means of heat. Distillatory vessels are either alembics or retorts; the former consist of an

inferior vessel, called a cucurbit, designed to contain the matter to be examined, and having an upper part fixed to it, called the capital, or head. In this last, the vapours are condensed by the contact of the surrounding air, or, in other cases, by the assistance of cold water surrounding the head, and contained in a vessel called the refrigeratory. From the lower part of the capital proceeds a tube, called the nose, beak, or spout, through which the vapours, after condensation, are, by a proper figure of the capital, made to flow into a vessel called the receiver, which is usually spherical. These receivers have different names, according to their figure, being called mattresses, balloons, &c. Retorts are a kind of bottle of glass, pottery, or metal, the bottom being spherical, and the upper part gradually diminishing into a neck, which is turned on one side.

DISTORTION. (From *distorqueo*, to wrest aside.) *Distortio*. A term applied to the eyes, when a person seems to turn them from the object he would look at, and is then called squinting, or strabismus. It also signifies the bending of a bone preternaturally to one side; as distortion of the spine, or vertebrae.

DISTORTOR. (From *distorqueo*, to wrest aside.) A muscle, whose office is to draw the mouth awry.

DISTORTOR ORIS. (From *distorqueo*, to wrest aside.) The zygomaticus minor.

DISTRICHIASIS. See *Distichiasis*.

DISTRIX. (From *δύω*, double, and *ἄκρῃ*, the hair.) A disease of the hair, when it splits and divides at the end.

Dittander. See *Nasturtium hortense*.

Dittany, bastard. See *Dictamnus albus*.

Dittany of Crete. See *Dictamnus creticus*.

Dittany, white. See *Dictamnus albus*.

DIURESIS. (From *διῶ*, through, and *πῶς*, to flow.) An increased secretion of urine. It is also applied to a diabetes.

DIURETICS. (*Diuretica*, sc. *medicamenta*, *διυρητικά*: from *διῶ*, a discharge of urine.) Those medicines or substances are so called which, when taken internally, augment the flow of urine from the kidneys. It is obvious that such an effect will be produced by any substance capable of stimulating the secreting vessels of the kidneys. All the saline diuretics seem to act in this manner. They are received into the circulation; and passing off with the urine, stimulate the vessels, and increase the quantity secreted.

There are other diuretics, the effect of which appears not to arise from direct application, but from an action excited in the stomach, and propagated by nervous communication to the secreting urinary vessels. The diuretic operation of squill,

and several other vegetables, appears to be of this kind.

There is still, perhaps, another mode in which certain substances produce a diuretic effect; that is, by promoting absorption. When a large quantity of watery fluid is introduced into the circulating mass, it stimulates the secreting vessels of the kidneys, and is carried off by urine. If, therefore, absorption be promoted, and if a portion of serous fluid, perhaps previously effused, be taken up, the quantity of fluid secreted by the kidneys will be increased. In this way digitalis seems to act: its diuretic effect, it has been said, is greater when exhibited in dropsy than it is in health.

On the same principle, (the effect arising from stimulating the absorbent system,) may probably be explained the utility of mercury in promoting the action of several diuretics.

The action of these remedies is promoted by drinking freely of mild diluents. It is also influenced by the state of the surface of the body. If external heat be applied, diuresis is frequently prevented, and diaphoresis produced. Hence the doses of them should be given in the course of the day, and the patient, if possible, be kept out of bed.

The direct effects of diuretics are sufficiently evident. They discharge the watery part of the blood; and, by that discharge, they indirectly promote absorption over the whole system.

Dropsy is the disease in which they are principally employed; and when they can be brought to act, the disease is removed with less injury to the patient than it can be by exciting any other evacuation. Their success is very precarious, the most powerful often failing; and, as the disease is so frequently connected with organic affection, even the removal of the effused fluid, when it takes place, only palliates without effecting a cure.

Diuretics have been likewise occasionally used in calculous affections, in gonorrhœa, and with a view of diminishing plethora, or checking profuse perspiration.

Murray, in his *Elements of Materia Medica*, classes the super-tartrate of potash, or cream of tartar, and nitrate of potash, or nitre, the muriate of ammonia, or crude salammoniac, potash, and the acetate of potash, or kali acetatum, among the *saline* diuretics; and selects the following from the *vegetable* kingdom:—*scilla maritima*, *digitalis purpurea*, *nicotiana tabacum*, *solanum dulcamara*, *lactuca virosa*, *colchicum autumnale*, *gratiola officinalis*, *spartium scoparium*, *juniperus communis*, *copaifera officinalis*, *pinus balsamea*, and *pinus larix*; and the *lytta vesicatoria* from the *animal* kingdom.

In speaking of particular diuretics, Dr.

Cullen says, the diuretic vegetables mentioned by writers are of very little power, and are employed with very little success. Of the umbellatae, the medicinal power resides especially in their seeds; but he never found any of them very efficacious. The semen dauci sylvestris has been commended as a diuretic; but its powers as such are not very remarkable. In like manner some of the *plantae stellatae* have been commended as diuretics; but none of them deserve our notice, except the *rubia tinctorum*, the root of which passes so much by the kidneys as to give its colour to the urine. Hence it may fairly be supposed to stimulate the secretories; but Dr. Cullen found its diuretic powers did not always appear, and never to any considerable degree; and as in brute animals it has always appeared hurtful to the system, he does not think it fit to be employed to any extent in human diseases. The bardana, lithospermum, ononis, asparagus, enula campana, are all substances which seem to pass, in some measure, by the kidneys; but their diuretic powers are hardly worth notice.

The principal articles included by Dr. Cullen, in his catalogue of diuretics, are dulcamara, digitalis, scilla; some of the alliacea and siliquosa; the balsams and resins; cantharides, and the diuretic salts.

DIVAPORATIO. Evaporation.

DIVARICATION. The crossing of any two things: thus when the muscular or tendinous fibres intersect each other at the different angles, they are said to divaricate.

DIVERSORIUM. (From *divertere*, to resort to.) The receptaculum chyli.

DIVERTICULUM. A mal-formation or diseased appearance of a part, in which a portion goes out of the regular course; and thereby forms a diverticulum, or deviation from the usual course. It is generally applied to the alimentary canal.

DIVERTICULUM NUCKII. The opening through which the round ligaments of the uterus pass. Nuck asserted that it remained open a long time after birth; to these openings he gave the name of *diverticula*.

DIVINUS. A pompous epithet of many compositions, from their supposed excellence.

DIVULSIO. (From *divello*, to pull asunder.) Urine, with a ragged and uneven sediment.

DOCIMASTIC ART. *Arts docimasticae*. The art of examining fossils, in order to discover what metals, &c. they contain.

Dock-cresses. See *Lampyris*.

Dock, sour. See *Acetosa*.

Dock, water. See *Hydrolapathum*.

Dodder of thyme. See *Epithymum*.

DODECACTYLUS. (From *dodua*, twelve, and *actylus*, a finger; so named, be-

cause its length is about the breadth of twelve fingers.) The duodenum, an intestine so called. It must be observed, that at the time this name was given, anatomy consisted in the dissection of brutes; and the length was therefore probably adjudged from the gut of some animal, and not of man.

DODECAPHARMACUM. (From *dodua*, twelve, and *pharmacum*, a medicine.) An ointment consisting of twelve ingredients, for which reason it was called the ointment of the twelve apostles.

DODECATHEON. (From *dodua*, twelve, and *theon*, to put.) An antidote consisting of twelve simples.

Dog-rose. See *Cynosbatus*.

DOU'S BANE, SYRIAN. This plant, *Aclepias syriaca* of Linnaeus, is particularly poisonous to dogs, and also to the human species. Boiling appears to destroy the poison in the young shoots, which are then said to be esculent, and flavoured like asparagus.

Dog's-grass. See *Gramen caninum*.

Dog's mercury. See *Cynocrambe*.

Dog-stones. See *Satyrion*.

DOGMA. (From *dodua*, to be of opinion.) An opinion founded on reason and experience.

DOLICHOS. (From *doluxos*, long: so called from its long shape.) 1. The name of a genus of plants in the Linnaean system. Class, *Diadelphia*. Order, *Decandria*.

2. The pharmacopoeial name of the cowhage. *Dolichos pruriens* of Linnaeus:—*volubilis*, *leguminibus racemosis*, *valvulis subcarinatis hirtis*, *pedunculis ternis*. The pods of this plant are covered with sharp hairs which are the parts employed medicinally in form of electuary, as anthelmintics. The manner in which these hairy spiculae act, seems to be purely mechanical: for neither the tincture, nor the decoction possess the least anthelmintic power.

DOLICHOS SOJA. The plant which affords the soy. It is much cultivated in Japan, where it is called *daidzei*: and where the pods supply their kitchens for various purposes; but the two principal are, with a sort of butter, termed *miso*, and a pickle called *soeju*.

DOLICHOS FRUENS. The systematic name of the cowhage. See *Dolichos*.

Dolor faciei. See *Tic Doloureux*.

Doloureux tic. See *Tic doloureux*.

DORONICUM. (From *dorongi*, Arab.)

Leopard's bane. See *Arnica*.

DORONICUM GERMANICUM. See *Arnica*.

DORONICUM PARDALIANUM. The systematic name of the Roman leopard's bane. See *Doronicon romanum*.

DORONICUM ROMANUM. Roman leopard's bane. *Doronicon pardalianches*; *foliis cordatis*, *obtusis*, *denticulatis*; *radicali-*

bus petiolatis; caulibus amplexicaulibus, of Linnæus. The root of this plant, if given in a full dose, possesses poisonous properties; but instances are related of its efficacy in epileptical and other nervous diseases.

DORSAL. Belonging to the back.

DORSALIS NERVI. The nerves which pass out from the vertebra of the back.

DORSI SPINALIS. See *Spinalis dorsi*.

DORSTENIA. (Named in honour of Dr. Dorsten.) A name of the contrayerva.

DORSTENIA DRAKENA. The systematic name for the contrayerva.

DORSTENIA HOUSTONII. See *Contrayerva*.

DOETHIEN. A name for furunculus.

DOVERI PULVIS. See *Pulvis ipecacuanhe compositus*.

Dove's foot. The geranium columbinum.

DRABA. (From *dravō*, to seize; so called from its sudden effect upon the nose of those who eat it.)

1. The name of a genus of plants in the Linnæan system. Class, *Tetradynamia*. Order, *Siliculosa*.

2. A name of the lepidium, or Arabian mustard, and Turkey cresses.

DRACO SYLVESTRIS. See *Parmica*.

DRACOCEPHALUM. (From *drakon*, a dragon, and *kephala*, a head.) The name of a genus of plants in the Linnæan system. Class, *Didynamia*. Order, *Gymnospermia*.

DRACOCEPHALUM CANARIENSE. The systematic name of the balm of Gilead. See *Moldavia*.

DRACONIS SANGUIS. See *Sanguis draculæ*.

DRACONTIUM. (From *drakon*, a dragon; so called because its roots resemble a dragon's tail.) *Dracunculus polyphyllus*. *Colubrina dracontia*. *Eroa de Sancta Maria*. *Gigarus serpentaria*. *Arum polyphyllum*. Dragon's wort. Many-leaved arum. This plant is the *Arum dracunculus* of Linnæus. Its roots and leaves are extremely acrimonious, more so than the arum maculatum, with which it agrees in medicinal virtues.

DRACUNCULUS. (From *drakon*, a serpent.) The Guinea worm; called also vermiculus capillaris. These animalcules are common in both Indies, in most parts of Africa, occasionally at Genoa, and other hot countries. These worms resemble the common worm, but are much larger; commonly found in the legs, but sometimes in the muscular part of the arms. They principally affect children, and their generation is not unlike that of the broad worms of the belly, hence their name tape-worm. While they move under the skin, they create no trouble; but, in length of time, the place near the dracunculus suppurates, and the animal puts forth its head. If it be drawn, it excites considerable uneasiness especially if drawn so

forcibly as to break it; for the part left within creates intolerable pain. These worms are of different lengths. In the Edin. Med. Essays, mention is made of one that was three yards and a half in length.

DRAGACANTHA. See *Tragacantha*.

Dragant gum. See *Tragacantha*.

Dragon's blood. See *Sanguis draconis*.

Dragon's wort. See *Dracontium*.

DRAKENA. See *Contrayerva*.

DRASTICA. (*Drastica*, sc. *medicamenta drastica*, active, brisk; from *dravō*, to effect.) A term generally applied to those medicines which are very violent in their action; thus, drastic purges, emetics, &c.

DRESDENSIS PULVIS. An oleo saccharum, containing the oil of cinnamon.

DROMA. The name of a plaster described by Myrepsus.

DRORACISMUS. (From *dravō*, to remove.)

Dropax. A stimulant plaster of pitch, wax, &c. to take off hair.

DROFAX. See *Drofacismus*.

DROPSY. A collection of a serous fluid in the cellular membrane; in the viscera and the circumscribed cavities of the body. See *Ascites*, *Anasarca*, *Hydrocephalus*, *Hydrothorax*, *Hydrocele*.

Dropsy of the belly. See *Ascites*.

Dropsy of the brain. See *Hydrocephalus*.

Dropsy of the cellular membrane. See *Anasarca*.

Dropsy of the chest. See *Hydrothorax*.

Dropsy of the ovaria. See *Ascites*.

Dropsy testicle. See *Hydrocele*.

Dropwort. See *Enanthe*, and *Filipendula*.

Dropwort hemlock. See *Enanthe*.

Dropwort water. See *Enanthe*.

DROSERA. (From *drosera*, dewy; which is from *droso*, dew; drops hanging on the leaves like dew.) The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Hexagynia*. Sun-dew.

DROSERA ROTUNDFOLIA. The systematic name of the sun-dew. See *Ros solis*.

DROSIOBOTANUM. (From *droso*, dew, and *botan*, a herb; so called from its being covered with an aromatic dew.) The herb betony. See *Betonica*.

DROSMELI. (From *droso*, dew, and *mel*, honey.) Honey-dew. Manna.

Dry belly ach. See *Colica pictonum*.

DUCTILITY. A property by which bodies are dilated by repeated or continued pressure. It is peculiar to metals.

Ducts, biliary. See *Biliary ducts*.

DUCTUS ARTERIOSUS. A great in-osculation found only in the fœtus, and very young children, betwixt the pulmonary artery and the aorta. In adults it is closed up.

DUCTUS AD NASUM. See *Canalis nasalis*.

DUCTUS AURIS PALATINUS. The eustachian tube.

DUCTUS BILIARIS. See *Choledochus ductus*.

DUCTUS COMMUNIS CHOLEDOCHUS. See *Choledochus ductus*.

DUCTUS HEPATICUS. See *Hepatic duct*.

DUCTUS LACHRYMALIS. See *Lachrymal ducts*.

DUCTUS LACTIFERI. *Ductus galactiferi.* The excretory ducts of the glandular substance composing the female breast. The milk passes along these ducts to the nipple.

DUCTUS PANCREATICUS. The pancreatic duct. It is white and small, and arises from the sharp extremity of the pancreas runs through the middle of the gland towards the duodenum, into which it pours its contents by an opening common to it and the *ductus communis choledochus*.

DUCTUS SALIVALES. The excretory ducts of the salivary glands, which convey the saliva into the mouth.

DUCTUS STENONIS. The Stenonian duct, which was so called after its discoverer *Stena*. It arises from all the small excretory ducts of the parotid gland, and passes transversely over the masseter muscle, penetrates the buccinator, and opens into the mouth.

DUCTUS THORACICUS. See *Thoracic duct*.

DUCTUS VENOSUS. When the vena cava passes the liver in the fœtus, it sends off the ductus venosus which communicates with the sinus of the vena portæ; but, in adults, becomes a flat ligament.

DUCTUS WARTHONIANUS. The excretory duct of the maxillary glands; so named after its discoverer.

DULCACIDUM. (From *dulcis*, sweet, and *acidus*, sour.) An oxymel. A medicine composed of a sweet and sour ingredient.

DULOAMARA. (From *dulcis*, sweet, and *amarus*, bitter.) *Solanum scamens. Glycyphoros, sive amaradulcis. Solanum hignosum. Ξεγυρως* of Theophrastes. Woody nightshade. *Solanum dulcamara* of Linnaeus:—*caule inermi frutescente flexuoso, foliis superioribus hastatis racemis cymosis.* The roots and stalks of this nightshade, upon being chewed, first cause a sensation of bitterness, which is soon followed by a considerable degree of sweetness; and hence the plant obtained the name of bittersweet. The berries have not yet been applied to medical use; they seem to act powerfully upon the primæ viæ, exciting violent vomiting and purging. Thirty of them were given to a dog, which soon became mad, and died in the space of three hours; and, upon opening his stomach, the berries were discovered to have undergone no change by the powers of digestion; there can, therefore, be little doubt of the deleterious effects of these berries: and,

as they are very common in the hedges, and may be easily mistaken, by children, for red currants, which they somewhat resemble, this circumstance is the more worthy of notice. The stipites, or younger branches, are directed for use in the Edinburgh Pharm. and they may be employed either fresh or dried, making a proportionate allowance in the dose of the latter for some diminution of its powers by drying. In autumn, when the leaves are fallen, the sensible qualities of the plant are said to be the strongest; and, on this account, it should be gathered in autumn rather than spring. *Dulcamara* does not manifest those narcotic qualities which are common to many of the nightshades; it is, however, very generally admitted to be a medicine of considerable efficacy. Murray says it promotes all the secretions; Haller observes, that it partakes of the milder powers of the nightshade joined to a resolvent and saponaceous quality; and the opinion of Bergius seems to coincide with that of Murray:—"Virtus: pellens urinam, sudorem, menses, lochia, sputa; mundificans." The diseases in which we find it recommended by different authors, are extremely various; but Bergius confines its use to rheumatism, retentio menisium, et lochiorum. *Dulcamara* appears also by the experiments of Razoux and others, to have been used with advantage in some obstinate cutaneous affections. Dr. Cullen says, "We have employed only the stipites, or slender twigs of this shrub, but as we have collected them, they come out very unequal, some parcels of them being very mild and inert, and others of them considerably acrid. In the latter state, we have employed a decoction of them in the cure of rheumatism, sometimes with advantage, but at other times without any effect. Though the *dulcamara* is here inserted in the catalogue of diuretics, it has never appeared to us as powerful in this way; for, in all the trials made here, it has hardly ever been observed to be in any measure diuretic." This plant is generally given in decoction, or infusion, and to prevent its exciting nausea, it is ordered to be diluted with milk, and to begin with small doses, as large doses have been found to produce very dangerous symptoms. Razoux directs the following: ꝑ. Stipitum dulcam. rec. drac ss in aquæ font. unc. 16 coquantur ad unc 8. This was taken in the dose of three or four drachms, diluted with an equal quantity of milk, every four hours. Linnaeus directs two drachms, or half an ounce of the dried stipites, to be infused half an hour in boiling water, and then to be boiled ten minutes; and of this decoction he gives two tea-cups full morning and evening. For the formula of this plant according to the London Pharm. see *Decoctum dulcamare*.

Dung, devil's. See *Asafoetida*.

Duo. (*Δύο*, two.) Some compositions consisting of two ingredients, are distinguished by this term, as *pilulæ ex duobus*.

DUODENUM. (From *duodenus*, consisting of twelve: so called because it was supposed not to exceed the breadth of twelve fingers; but as the ancients dissected only animals, this does not hold good in the human subject.) The first portion of the small intestines. See *Intestines*.

DUPLEXANA. (From *duplex*, double.) A name of the double tertian fever.

DURA MATER. (From *durus*, hard, and *mater*, a mother; called *dura*, from its comparative hardness with the *pia mater*, and *mater*, from its being supposed to be the source of all the other membranes.)

Dura meninx. *Dermatodes.* A thick and somewhat opaque and insensible membrane, formed of two layers, that surrounds and defends the brain, and adheres strongly to the internal surface of the cranium. It has three considerable processes, the falxiform, the tentorium, and the septum cerebelli; and several sinusses, of which the longitudinal, lateral, and inferior longitudinal, are the principal. Upon the external surface of the dura mater, there are little holes, from which emerge fleshy-coloured papillæ, and which, upon examining the skull-cap will be found to have corresponding foveæ. These are the external glandulæ Pacchioni. They are in number from ten to fifteen on each side, and are chiefly lateral to the course of the longitudinal sinus. The arteries which supply this membrane with vessels for its own nourishment, for that of the contiguous bone, and for the perpetual exudation of the fluid, or halitus rather, which moistens or bedews its internal surface, may be divided into anterior, middle, and posterior. The first proceeds from the ophthalmic and ethmoidal branches; the second from the internal maxillary and superior pharyngeal; the posterior from the occipital and vertebral arteries.

The principal artery of the dura mater, named by way of distinction, the great artery of the dura mater, is derived from the internal maxillary artery, a branch of the external carotid. It is called the spinalis, or spheno-spinalis, from its passing into the head through the spinous hole of the sphenoid bone, or meningeæ media, from its relative situation, as it rises in the great middle fossa of the skull. This artery, though it sometimes enters the skull in two branches, usually enters in one considerable branch, and divides, soon after it reaches the dura mater, into three or four branches, of which the anterior is the largest; and these spread their ramifications beautifully upon the dura mater, over all that part which is opposite to the anterior, middle, and posterior lobes of the

brain. Its larger trunks run upon the internal surface of the parietal bone, and are sometimes for a considerable space buried in its substance. The extreme branches of this artery extend so as to inosculate with the anterior and posterior arteries of the dura mater, and through the bones, (chiefly parietal and temporal bones.) They inosculate with the temporal and occipital arteries. The meningeal artery has been known to become aneurismal, and distended at intervals; it has formed an aneurism, destroying the bones and causing epilepsy.

DURA MENINX. Before the time of Galen, the term *meninx* was common to all the membranes of the body; afterwards it was appropriated to those of the brain. See *Dura mater*.

Dwale. See *Belladonna*.

Dwarf elder. See *Ebulus*.

DYOTA. (From *δύο*, two, and *ωτ*, *ωτος*, an ear.) A chymical instrument with two ears, or handles.

DYSÆSTHESIA. (From *δυσ*, difficulty, and *αἰσθησις*, to feel or perceive. Impaired feeling.

DYSANAGOGUS. (From *δύο*, with difficulty, and *αναγω*, to subdue.) Viscid expectoration.

DYSCATAPOTIA. (From *δυσ*, and *καταποτιν*, to drink.) A difficulty of swallowing liquids, which Dr. Mead thinks a more proper term than that used for canine madness, viz. hydrophobia; as it is more particularly descriptive of the affection under which the unhappy patients labour; for in reality, they dread water from the difficulty of swallowing it.

DYSCINESIA. (From *δυσ*, bad, and *κίνησις*, to move.) Bad or imperfect motion.

DYSCINESIÆ. An order in the class *locales* of Cullen's nosology; embracing disease in which the motion is impeded, or depraved, from an imperfection of the organ.

DYSCOROSIA. (From *δυσ*, with difficulty, and *ακοησις*, to be deaf.) A defect in the sense of hearing.

DYSCRASIA. (From *δυσ*, with difficulty, and *κρασις*, to mix. A bad habit of body.

DYSECOEA. (From *δυσ*, difficult, and *εξω*, hearing.) *Cophosis*. Deafness. Hearing diminished, or destroyed. A genus of disease in the class *locales*, and order *dysæsthesiæ* of Cullen, containing two species: *Dysecoëa organica*, which arises from wax in the meatus, injuries of the membrane, or inflammation and obstruction of the tube: *Dysecoëa atonica*, when without any discernible injury of the organ.

DYSECLIA. (From *δυσ*, with difficulty, and *ελκος*, an ulcer.) An ulcer difficult to heal.

DYSEMETUS. (From *δυσ*, with difficulty, and *εμεσις*, to vomit.) A person not easily made to vomit.

DYSENTERY. (From *δυσ*, difficulty, and *ένεα*, the bowels.) *Dysentēria*. *Dis-solutus morbus*. *Diarrhœa carnosæ*. The flux. A genus of disease in the class *pyrexia*, and order *profusio* of Cullen's nosology. It is known by contagious pyrexia; frequent griping stools; tenesmus; stools, chiefly mucous, sometimes mixed with blood, the natural fæces being retained or voided in small, compact, hard substances, known by the name of scybala; loss of appetite, and nausea. It occurs chiefly in summer and autumn, and is often occasioned by much moisture succeeding quickly intense heat, or great drought; whereby the perspiration is suddenly checked, and a determination made to the intestines. It is likewise occasioned by a use of unwholesome and putrid food, and by noxious exhalations and vapours; hence it appears often in armies encamped in the neighbourhood of low marshy grounds, and proves highly destructive; but the cause which most usually gives rise to it, is a specific contagion; and when it once makes its appearance, where numbers of people are collected together, it not unfrequently, spreads with great rapidity. A peculiar disposition in the atmosphere seems often to predispose, or give rise to the dysentery, in which case it prevails epidemically.

It frequently occurs about the same time with autumnal intermittent and remittent fevers, and with these it is often complicated.

The disease, however, is much more prevalent in warm climates than in cold ones; and in the months of August, September, and October, which is the rainy season of the year in the East Indies, it is very apt to break out and to become very general among the negroes on the different plantations in the colonies. The body having been rendered irritable by the great heat of the summer, and being exposed suddenly to much moisture with open pores, the blood is thereby thrown from the exterior vessels upon the interior, so as to give rise to dysenteries.

An attack of dysentery is sometimes preceded by loss of appetite, costiveness, flatulency, sickness at the stomach, and a slight vomiting, and comes on with slight chills, succeeded by heat in the skin, and frequency of the pulse. These symptoms are in general the forerunners of the griping and increased evacuation which afterwards occur.

When the inflammation begins to occupy the lower part of the intestinal tube, the stools become more frequent, and less abundant; and, in passing through the inflamed parts, they occasion great pain, so that every evacuation is preceded by a severe griping, as also a rumbling noise.

The motions vary both in colour and consistence, being sometimes composed of

frothy mucus streaked with blood, and at other times of an acrid watery humour, like the washings of meat, and with a very fetid smell. Sometimes pure blood is voided; now and then lumps of coagulated mucus, resembling bits of cheese, are to be observed in the evacuations, and in some instances a quantity of purulent matter is passed.

Sometimes what is voided consists merely of a mucous matter, without any appearance of blood, exhibiting that disease which is known by the name of dysentery alba, or morbus mucosus.

Whilst the stools consist of these various matters, and are voided frequently, it is seldom that we can perceive any natural fæces among them, and when we do, they appear in small hard balls, called scybala, which being passed, the patient is sure to experience some temporary relief from the griping and tenesmus.

It frequently happens, from the violent efforts which are made to discharge the irritating matters, that a portion of the gut is forced beyond the verge of the anus, which, in the progress of the disease, proves a troublesome and distressing symptom; as does likewise the tenesmus, there being a constant inclination to go to stool, without the ability of voiding any thing, except perhaps a little mucus.

More or less pyrexia usually attends with the symptoms which have been described, throughout the whole of the disease, where it is inclined to terminate fatally, and is either of an inflammatory or putrid tendency. In the other case, the febrile state wholly disappears after a time, while the proper dysenteric symptoms probably will be of long continuance.

When the symptoms run high, produce great loss of strength, and are accompanied with a putrid tendency and a fetid and involuntary discharge, the disease often terminates fatally in the course of a few days; but when they are more moderate, it is often protracted to a considerable length of time, and so goes off at last by a gentle perspiration, diffused equally over the whole body; the fever, thirst, and griping then ceasing, and the stools becoming of a natural colour and consistence. When the disease is of long standing, and has become habitual, it seldom admits of an easy cure; and when it attacks a person labouring under an advanced stage of scurvy, or pulmonary consumption, or whose constitution has been much impaired by any other disorder, it is sure to prove fatal. It sometimes appears at the same time with autumnal intermittent and remittent fevers, as has been observed, and is then more complicated and difficult to remove.

Upon opening the bodies of those who die of dysentery, the internal coat of the intestines (but more particularly of the

colon and rectum) appears to be affected with inflammation and its consequences, such as ulceration, gangrene, and contractions. The peritonæum and other coverings of the abdomen, seem likewise, in many instances, to be affected by inflammation.

DYSEMPULOTICUS. (From *δυσ*, with difficulty, and *εμυλον*, to cicatrize.) *Dysempulotus*. An inveterate ulcer difficult to be healed.

DYSESTHESIA. The name of an order of diseases in the class *locales* of Cullen's nosological arrangement; embracing those diseases in which the senses are injured or destroyed by the imperfections of the organs.

DYSHÆMORRHOIS. (From *δυσ*, with difficulty, and *αιμορροια*, the piles.) Suppression of bleeding piles.

DYSLOCHIA. (From *δυσ*, difficulty, and *λοχεια*, the lochia.) A suppression of the lochia.

DYSMENORRHEA. (From *δυσ*, with difficulty, and *μηνορροια*, the menses.) A difficult or painful menstruation, accompanied with severe pains in the back, loins, and bottom of the belly.

DYSODOR. (From *δυσ*, bad, and *οδω*, to smell.) A bad smell. Fœtid. Hippocrates applies it to a fœtid disorder of the small intestines. Also the name of a malagma and acopon in Galen and Paulus Aegineta.

DYSOPIA. (From *δυσ*, bad, and *οψω*, an eye.) *Paropsis*. Difficult sight. Sight depraved, requiring one certain quantity of light, one particular distance, or one position. A genus of disease in the class *locales*, and order *dysæsthesiæ* of Cullen, containing the five following species: 1. *Dysopia tenebrarum*, called also *amblyopia crepuscularis*, requiring objects to be placed in a strong light. 2. *Dysopia luminis*, likewise termed *amblyopia meridiana*, objects only discernible in a weak light. 3. *Dysopia distitorum*, in which distant objects are not perceived. 4. *Dysopia proximorum*, or *dysopia amblyopia*, in which objects too near are not perceived. 5. *Dysopia lateralis*, called also *amblyopia luscorum*, in which objects are not seen, unless placed in an oblique position.

DYSOREXIA. (From *δυσ*, bad, and *ορεξις*, appetite.) A bad or depraved appetite.

DYSOREXIAE. The name of an order in the class *locales* of Cullen's nosology, which he divides into two sections, *appetitus erronei* and *deficientes*.

DYSPEPSIA. (From *δυσ*, bad, and *πεψω*, to concoct.) *Apepsia*. Indigestion. Dr. Cullen arranges this genus of disease in the class *neuroses*, and order *adynamia*. It chiefly arises in persons between thirty and forty years of age, and is principally to be met with in those who devote much time to study, or who lead either a very sedentary or irregular life. A great singu-

larity attendant on it is, that it may, and often does, continue a great length of time, without any aggravation or remission of the symptoms.

Great grief and uneasiness of mind, intense study, profuse evacuations, excess in venery, hard drinking, particularly of spirituous liquors, and of tea, tobacco, opium, and other narcotics, immoderate repletion, and over distention of the stomach, a deficiency in the secretion of the bile or gastric juice, and the being much exposed to moist and cold air, when without exercise, are the causes which usually occasion dyspepsia.

A long train of nervous symptoms generally attend on this disease, such as a loss of appetite, nausea, heart-burn, flatulency, acid eructations, a gnawing in the stomach when empty, a sense of constriction and uneasiness in the throat, with pain in the side, or sternum, so that the patient at times can only lay on his right side; great costiveness, habitual chilliness, paleness of the countenance, languor, unwillingness to move about, lowness of spirits, palpitations, and disturbed sleep.

The number of these symptoms varies in different cases, with some being felt only in part; in others being accompanied even with additional ones, equally unpleasant, such as severe transient pains in the head and breast, and various affections of the sight, as blindness, double vision, &c.

Dyspepsia never proves fatal, unless when, by a very long continuance it produces great general debility and weakness; and so passes into some other disease, such as dropsy: but it is at all times very difficult to remove, but more particularly so in warm climates.

The morbid appearances to be observed on dissections of this disease, are principally confined to that part of the stomach which is called the pylorus; which is often found either in a contracted, scirrhus, or ulcerated state. In every instance the stomach is perceived to be considerably distended with air.

DYSPERMATISMUS. (From *δυσ*, bad, and *σπέρμα*, seed.) *Agnesia*. Slow, or impeded emission of semen during coition, insufficient for the purpose of generation. A genus of disease in the class *locales*, and order *epischæses* of Cullen. The species are: 1. *Dyspermatismus urethralis*, when the obstruction is in the urethra. 2. *Dyspermatismus nodosus*, when a tumour is formed in either corpus cavernosum penis. 3. *Dyspermatismus preputialis*, when the impediment is from a straightness of the orifice of the prepuce. 4. *Dyspermatismus mucosus*, when the urethra is obstructed by a viscid mucus. 5. *Dyspermatismus hypertonicus*, when there is an excess of erection of the penis. 6. *Dyspermatismus epilepticus*, from epileptic fits coming on during

coition. 7. *Dyspermatus ephractodes*, from a want of vigour in the genitals. 8. *Dyspermatus refluxus*, in which the semen is thrown back into the urinary bladder.

DYSPHAGIA. (From *δύς*, with difficulty, and *φαγῶς*, to eat.) A difficulty of deglutition.

DYSPHONIA. (From *δύς*, bad, and *φωνή*, the voice.) A difficulty of speaking.

DYSPPNEA. (From *δύς*, difficult, and *πνέω*, to breathe.) *Dyspnoea*. Difficult respiration, without sense of stricture, and accompanied with cough through the whole course of the disease. A genus of disease in the class *neuroses*, and order *spasmi* of Cullen. He distinguishes eight species. 1. *Dyspnea catarrhalis*, when with a cough there are copious discharges of viscid mucus, called also *asthma catarrhale*, *pneumodes*, *pneumonicum*, and *pituitosum*. 2. *Dyspnea sicca*, when there is a cough without any considerable discharge. 3. *Dyspnea aerea*, when the disease is much increased by slight changes of the weather. 4. *Dyspnea terrea*, when earthy or calculous matters are spit up. 5. *Dyspnea aquosa*, when there is a scarcity of urine and oedematous feet, without any symptoms of a dropy in the chest. 6. *Dyspnea pinguedinosa*, from corpulency. 7. *Dyspnea thoracica*, when parts surrounding the chest are injured or deformed. 8. *Dyspnea extrinseca*, from manifest external causes.

DYSPNOEA. See *Dyspnea*.

DYSRACHITIS. The name of a plaster in Galen.

DYSTHYMIA. (From *δύς*, bad, and *θυμός*, mind.) Insanity.

DYSTOCHIA. (From *δύς*, with difficulty, and *τέλει*, to bring forth.) Difficult labour or childbirth.

DYSTOECHIASIS. (From *δύς*, bad, and *σύνταξις*, order.) An irregular disposition of the hairs in the eyelids.

DYSURIA. (From *δύς*, difficult, and *ουρά*, urine.) *Stillicidium*. *Ardor urinae*. *Culbicio*. A suppression or difficulty in discharging the urine. A total suppression is called ischuria; a partial suppression, dysuria; and this may be with or without heat. When there are frequent, painful, or uneasy urgings to discharge the urine, and it passes off only by drops, or in very small quantities, the disease is called stranguy. When a sense of pain, or heat, attends the discharge, it

passes with difficulty, and is styled *heat* of the urine. The dysuria is acute, or chronic. Dr. Cullen places this disease in the class *locales*, and order *epischees*, containing six species: 1. *Dysuria ardens*, a sense of heat, without any manifest disorder of the bladder. 2. *Dysuria spasmodica*, from spasm. 3. *Dysuria compressionis*, from a compression of the neighbouring parts. 4. *Dysuria phlogistica*, from violent inflammation. 5. *Dysuria calculosa*, from stone in the bladder. 6. *Dysuria mucosa*, from an abundant secretion of mucus. The causes which give rise to these diseases are, an inflammation of the urethra, occasioned either by venereal sores, or by a use of acrid injections, tumour, ulcer of the prostate gland, inflammation of the kidneys, or bladder, considerable enlargements of the hemorrhoidal veins, a lodgment of indurated faeces in the rectum, spasm at the neck of the bladder, the absorption of cantharides, applied externally or taken internally, and excess in drinking either spirituous or vinous liquors; but particles of gravel, sticking at the neck of the bladder, or lodging in the urethra, and thereby producing irritation, prove the most frequent cause. Gouty matter falling on the neck of the bladder, will sometimes occasion these complaints.

In dysury, there is a frequent inclination to make water, with a smarting pain, heat, and difficulty in voiding it, together with a sense of fulness in the region of the bladder. The symptoms often vary, however, according to the cause which has given rise to it. If it proceeds from a calculus in the kidney or ureter, besides the affections mentioned, it will be accompanied with nausea, vomiting, and acute pains in the loins and region of the ureter and kidney of the side affected. When a stone in the bladder, or gravel in the urethra, is the cause, an acute pain will be felt at the end of the penis, particularly on voiding the last drops of urine, and the stream of water will either be divided into two, or be discharged in a twisted manner, not unlike a cork-screw. If a scirrhus of the prostate gland has occasioned the suppression or difficulty of urine, a hard indolent tumour, unattended with any acute pain, may readily be felt in the perineum, or by introducing the finger in ano.

E.

EAR. *Auris*. The organ of hearing is situated at the side of the head, and is divided into external and internal ear. The *auricula*, commonly called the ear, constitutes the external, and contains several eminences and depressions, as the *helix*, *an-*

tihelix, *tragus*, *antitragus*, *concha auriculae*, *scapha*, and *lobulus*. The external auditory passage, containing the wax, proceeds from its middle down to the membrane of the tympanum, which divides the external from the internal parts of this organ. Behind

the *membrana tympani* is an irregular cavity, the cavity of the tympanum, in which are four little bones, the *malleus*, *incus*, *stapes*, and *os orbiculare*; and four openings one of the Eustachian tube, mastoid sinus, fenestra ovalis, and fenestra rotunda. The tympanum is terminated by the labyrinth. The labyrinth is the remaining part of the internal ear, consisting of the *cochlea vestibulum*, and *semicircular canals*. The arteries of the ear are the external and internal auditory. The veins empty themselves into the external jugulars. The muscles of the ear are divided into three classes: the common, proper, and internal. The common muscles are, the *attollens aurem*, *anterior auris* and *retrahentes auris*, which move the whole ear. The proper are, *helicis major*, *helicis minor*, *tragicus*, *antitragicus*, and *transversus auris*; these affect the parts only to which they are connected. The muscles of the internal ear are, *laxator tympani*, *tensor tympani*, and *stapedius*, which belong to the *ossicula auditus*. The nerves of the external ear are branches of the *nervus auditorius durus*, and those of the internal ear, are branches of the *nervus auditorius mollis*.

EARITES. Hematites, or blood-stone.

EARTH. *Terra*. Though there seems to be an almost infinite variety of earthy substances scattered on the surface of this globe, yet when we examine them with a chymical eye, we find, not without surprise, that all the earth and stones which we tread under our feet, and which compose the largest rocks, as well as the numerous different specimens which adorn the cabinets of the curious, are composed of a very few simple or elementary earths, in number no more than seven: viz.—*Silex*, *alumine*, *glucine*, *zircon*, *agustine*, *ytria*, and *magnesia*.

These are all the simple earths hitherto known: none of them have yet been decomposed, nor has the smallest proof been brought that they are compound; we must therefore consider them as simple bodies, which nature presents to us completely formed, though one or more of them enters into the composition of a great many bodies. They have a variety of properties which are common to all:—they are dry, incombustible bodies. They are insoluble in water and alcohol, or nearly so. They have little or no taste. Their specific gravity does not exceed 4.9. When perfectly pure, they assume the form of a white powder, harsh to the touch. They are infusible. They are capable of combining with acids, when they form neutral salts. They are likewise disposed to unite with the alkalis, with sulphur, and phosphorus; with metallic oxyds, and with each other, either by fusion or solution in water.

Every one of these characters is not perhaps rigorously applicable to each of these

bodies; but they all possess a sufficient number of them to render it useful to arrange them under one class.

Stones differ from earths principally in cohesion and hardness, and therefore are included under the same general name.

Earth, absorbent. See *Absorbents*.

Earth, aluminous. Earth which contains alumina. See *Alumene*.

Earth, animal calcareous. This term is applied to crab's claws, &c. which are calcareous earth, and obtained from the animal kingdom.

Earth, argillaceous. See *Alumine*.

EARTH-BATH. A remedy recommended by some writers on the continent, as a specific in consumption. In this country it produced to the patients very distressing sensations of cold; in some it seemed to be productive of bad effects; and it does not appear that, in any consumptive cases, good effects were ever derived from its use.

Earth, bolar. See *Bole*.

EARTH, FULLERS'. *Cimelia purpureascens*. A compact bolar earth, commonly of a greyish colour. It is sometimes applied by the common people to inflamed breasts, legs, &c. with a view of cooling them.

Earth, heavy. See *Barytes*.

Earth, Japan. See *Catechu*.

Earth, mineral calcareous. Those calcareous earths which are obtained from the mineral kingdom. The term is applied in opposition to those obtained from animals.

Earth-nut. See *Pig-nut*.

EARTH, SEALED. *Terra sigillata*. Little cakes of bolar earths, which are stamped with impressions. They were formerly in high estimation as absorbents, but now fallen into disuse.

EARTH-WORM. *Lambricus terrestris*. *Vermis terrestris*. These insects are supposed to possess a diuretic and antispasmodic virtue, with which views they are occasionally employed in foreign countries.

EAR-WAX. *Cerumen aurium*. A waxy secretion found in the meatus auditorius externus, into which it is separated by the glands around that canal.

EATON'S STYPTIC. French brandy highly impregnated with calcined green vitriol. A remedy for checking hæmorrhages.

EAU-DE-LUCE. See *Spiritus ammonia succinatus*.

EAU-DE-RABEL. This is composed of one part of sulphureous acid to three of rectified spirit of wine. It is much used in France, when diluted, in the cure of gonorrhœas, leucorrhœas, &c.

EBEL. The seeds of sage, or of juniper.

EBENUM. Indian ebony. It is supposed to be opthalmic.

EBESMECH. A name in Langius for quicksilver.

EBISCUS. The *Hibiscus*, or marsh mallow.

EBRIECATUM. (From *ebrio*, to be drunk.) By this term Paracelsus expresses loss of sense by drunkenness.

EBRIECATUM CÆLESTE. By this term Paracelsus means that kind of enthusiasm which is affected by many heathen priests.

EBSENECH. A name in Languis for quicksilver.

EBULLITION. (From *ebullio*, to bubble up.) *Ebullitio*. Boiling. This consists in the change which a fluid undergoes from a state of liquidity to that of an æri-form fluid, or gas, in consequence of the application of heat, which dilates and converts it into vapour.

EBULUS. (From *ebullio*, to make boil; so called because of its supposed use in purifying the humours of the body.) *Chamaeacte*. *Sambucus humilis*. *Sambucus herbacea*. Dwarf elder, or dane-wort. The root, interior bark, leaves, flowers, berries, and seeds of this herbaceous plant, *Sambucus ebulus*; *cymistridis*, *stipulis foliaceis*, *caule herbaceo*, of Linnaeus, have all been administered medicinally, in moderate doses, as resolvents and deobstruents, and, in larger doses, as hydragogues. The plant is chiefly employed by the poor of this country, amongst whom it is in common use as a purgative, but Dr. Cullen speaks of it as a violent remedy.

ECBOLICA. (From *εκβαλλω*, to cast out.) Medicines which were formerly said to cause abortion.

ECBOLIOS. (From *εκβαλλω*, to cast out.) Miscarriage.

ECBRASATA. (From *αεζαζω*, to be very hot.) *Ecchymata*. Painful fiery pimples in the face, or surface of the body.

ECBRASMUS. (From *αεζαζω*, to become hot.) Fermentation.

ECBYRSOMATA. (From *αε*, and *βυρσο*, the skin.) Protuberances of the bones at the joints, which appear through the skin.

ECCATHARTICA. (From *εκκαθαζω*, to purge outwards.) According to Gorræus, eccathartics are medicines which open the pores of the skin; but in general they are understood to be deobstruents. Sometimes expectorants are thus called, and also purgatives.

ECCHYLOMA. (From *αε*, and *χυλος*, juice.) An extract.

ECCHYMATA. (From *αχυω*, to pour out.) See *Ecbrasmata*.

ECCHYMOMA. (*Εκχυμομα*; from *αχυω*, to pour out.) *Ecchymosis*. Sometimes called *crustula* and *sugillatio*. Extravasation. A black and blue swelling, either from a bruise or extravasation of blood. A genus of disease in the class *locales*, and order *tumores* of Cullen.

ECCHYMOMA ARTERIOSUM. The false aneurism.

ECCHYMOISIS. See *Ecchymoma*.

ECCLISIS. (From *εκκληνω*, to turn aside.) A luxation or dislocation.

ECCOPE. (From *εκκοβω*, to cut off.) The cutting off any part.

ECCORFUS. (From *εκκοψω*, to cut off.) An ancient instrument, the raspatory, used in trepanning.

ECCOPROTICA. (From *αε*, and *κοπρω*, dung.) Opening medicines, whose operation is very gentle; such as manna, *sepea*, &c.

ECCRINOCRITICA. (From *αεζω*, to secrete, and *κρηνω*, to judge.) Judgments formed from the secretions.

ECCRINOLOGIA. *Ecclinologica*. (From *αεζω*, to secrete, and *λογος*, a discourse.) The doctrine of secretions.

ECCRISIS. (From *αεζω*, to secrete.) A secretion of any kind.

ECCYMOSIS. See *Ecchymoma*.

ECDORA. (From *εκδωω*, to excoriate.) An excoriation; and particularly used for an excoriation of the urethra.

ECDORIA. (From *εκδωω*, to excoriate.) Medicines which excoriate and burn through the skin.

ECHECOLLON. (From *εχω*, to have, and *κολλα*, glue.) *Echecollum*. Any topical glutinous remedy.

ECHETROISIS. So Hippocrates calls the white brisny.

ECHINIDES. In Hippocrates it is mentioned as what he used for purging the womb with.

ECHINOPHTHALMIA. (From *εχινος*, a hedge-hog, and *οφθαλμα*, an inflammation of the eye.) An inflammation of the hairy part of the eyelids, where the hairs bristle out like the quills of an echinus, or hedge-hog.

ECHINOPODIUM. (From *εχινος*, a hedge-hog, and *πους*, a foot; so named because its flowers resemble the foot of an urchin.) A species of broom, or genista.

ECHINOPUS. (From *εχινος*, as beset with prickles.) *Crocodilium*. *Acanthaluca*. *Seabiosa cardiifolia*. *Spherocephala elation*. Globe thistle. *Echinops sphaerocephalus* of Linnaeus. It is raised in our gardens. The root and seeds are moderately diuretic, but not used.

ECHIUUM. (From *εχμη*, a viper; so called because it was said to heal the stings of vipers.) The name of a genus of plants in the Linnaean system. Class, *Pentandria*. Order, *Monogynia*. Viper's bugloss. Antiepileptic.

ECHIUUM ÆGYPTIACUM. Wall bugloss; vulnerary, sudorific.

ECHOS. (*Εχος*; sound.) In Hippocrates it signifies the same as the *tinnitus aurium*, or noise in the ears.

ECHYSIS. (From *εχυω*, to pour out.) A fainting, or swooning.

ECLAMPSIA. (From *εκλαμπω*, to shine.) See *Eclampsia*.

ECLAMPSIS. (From *εκλαμπω*, to shine.) *Eclampsia*. It signifies a splendour, bright-

ness, effulgence, flashing of light, scintillation. It is a flashing light, or those sparkings which strike the eyes of epileptic patients. Celsus Aurelianus calls them *circuli ignei*, scintillations, or fiery circles. Though only a symptom of the epilepsy Hippocrates puts it for epilepsy itself.

ECCLECTICA. (From *εκλεκτα*, to elect.) Archigenus and some others selected from all other sects what appeared to them to be the best and most rational; hence they were called *Eclectics*, and their medicine *Eclectic medicine*.

ECCLECTOS. (From *εκλεκτω*, to lick up.) A linctus, or soft medicine to be licked up.

ECCLESMA. (From *εκλεκτω*, to lick.) Is a form of medicine made by the incorporation of oils and syrups, and which is to be taken upon a liquorice stick; the same as *Linctus*.

ECLISIS. (From *εκλυω*, to dissolve.) *Eclectos*. An universal faintness.

ECHASMA. (From *εκαθιστω*, to form together.) A mass of substances kneaded together.

ECHPEISMENOS. (From *εκαπνιστω*, to press out.) An epithet for ulcers with protuberating lips.

ECHPHRACTIC. (From *εκαφρασσω*, to remove obstructions.) Are such medicines as incide and render more thin tough humours, so as to promote their discharge.

ECHPHRACTICA. (From *εκαφρασσω*, to remove obstructions.) Deobstruent medicines.

ECHPHRAXIS. (From *εκαφρασσω*, to remove obstruction.) A diaphoresis; an opening of the pores.

ECHPHIAS. (From *εκ*, and *φωα*, to produce.) An appendix, or excrescence. Some call the appendicula vermiciformis thus.

ECHPHYS. Flatus from the bladder through the urethra, and from

ECHPHYSISIS. (From *εκαφυσσω*, to breathe through.) A quick expulsion of the air from the lungs.

ECHPHYSIS. (From *εκαφω*, to produce.) An apophysis, or appendix. A process.

ECHPIESMA. (From *εκαπνιστω*, to press out.) A fracture of the skull, in which the bones press inwardly.

ECHPIESMOS. (From *εκαπνιστω*, to press out.) A disorder of the eye, in which the globe is almost pressed out of the socket by an afflux of humours.

ECHPIEROMA. (From *εκαπνισσω*, to fill.) In Hippocrates they are hard balls of leather, or other substances, adapted to fill the arm-pits, while by the help of the heels, placed against the balls, and repressing the same, the luxated os humeri is reduced into its place.

ECHPLEXIA. (From *εκαπνισσω*, to terrify or astonish.) A stupor, or astonishment, from sudden external accidents.

ECPIROX. (From *εκαπνισσω*, to breathe.) Expiration; that part of respiration in which the air is expelled from the lungs.

ECPIROMA. (From *εκαπνισσω*, to fall out.)

1. A luxation of a bone.
2. The exclusion of the secundines.
3. Speaking of corrupt parts, it signifies a falling off.

4. An hernia in the scrotum.

5. A falling down of the womb.

ECPICTICA. (From *εκαπνισσω*, to condense.) *Incrassants*. Medicines that render the fluids more solid.

ECPIEMA. (From *εκ*, and *πυω*, pus.) A copious collection of pus or matter, from the suppuration of a tumour.

ECPIESMA. (From *εκαπνισσω*, to break.) A rupture.

ECPIEXIS. (From *εκαπνισσω*, to break.) A rupture. Hippocrates expresses by it a rupture or laceration of the womb.

ECPIETHMOS. (From *εκ*, and *γυμνος*, harmony.) A term applied to the pulse, and signifies that it is disorderly or irregular.

ECPIOS. (From *εκαπνισσω*, to flow out.) An efflux, or the course by which any humour which requires purging is evacuated.

ECPIULLA. The French name for scrophula.

ECPIUIS. (From *εκαπνισσω*, to flow out.) In Hippocrates it is an efflux of the semen before it receives the conformation of a fœtus, and therefore is called an efflux, to distinguish it from abortion.

ECSARCOMA. (From *εκ*, and *σαρξ*, flesh.) A fleshy excrescence.

ECSTASIS. (*Εκστασις*: from *εξιστημι*, to be out of one's senses.) An exstasy, or trance. In Hippocrates it signifies a delirium. Dr. Cullen ranks it as a kind of apoplexy. See *Exstasis*.

ECTHELYSIS. (From *εκαθελω*, to render effeminate.) Softness. It is applied to the skin and flesh, when lax and soft, and to bandages, when not sufficiently tight.

ECTHLYMA. (From *εκαθελω*, to press out against.) An ulceration caused by pressure of the skin.

ECTHLYPSIS. (From *εκαθελω*, to press out against.) Elision, or expression. It is spoken of swelled eyes, when they dart forth sparks of light.

ECTHYMA. (From *εκαθελω*, to break out.) A pustule, or cutaneous eruption.

ECTHYMATA. (From *εκαθελω*, to break out.) Pimples, pustules, or cutaneous eruptions.

ECTILLOTICA. (From *εκτελλω*, to pull out.) Medicines which eradicate tubercles, or corns, or destroy superfluous hair.

ECTOPIA. (From *εκτοπος*, out of place.) Displaced.

ECTOPIÆ. Parts displaced. An order in the class *locales* of Cullen's nosology.

ECTRAPELOGASTROS; (From *εκτραπηνωμαι*,

to degenerate, and *γαστήρ*, a belly.) One who has a monstrous belly, or whose appetite is voraciously large.

ΕΚΤΡΙΜΜΑ. (From *αλείω*, to rub off.) An attrition, or galling. In Hippocrates it is an excoriation of the skin about the os sacrum.

ΕΚΤΡΟΠΗ. (From *αλγίζω*, to divert, pervert, or invert.) It is any duct by which the humours are diverted and drawn off. In *P. Egineta* it is the same as *Ectropium*.

ΕΚΤΡΟΠΙΟΣ. (From *αλγίζω*, to invert.) An epithet for any medicine that makes the blind piles appear outwardly.

ECTROPIUM. (From *αλγίζω*, to evert.) An eversion of the eyelids so that their internal surface is outermost.

There are two species of this disease; one produced by an unnatural swelling of the lining of the eyelids, which not only pushes their edges from the eyeball, but also presses them so forcibly, that they become everted; the other arising from a contraction of the skin, covering the eyelid, or of that in the vicinity, by which means the edge of the eyelid is first removed for some distance from the eye, and afterwards turned completely outward, together with the whole of the affected eyelid.

The morbid swelling of the lining of the eyelids, which causes the first species of ectropium, arises mostly from a congenial laxity of this membrane, afterwards increased by obstinate chronic ophthalmies, particularly of a scrophulous nature, in relaxed, unhealthy subjects; or else the disease originates from the small-pox affecting the eyes.

While the disease is confined to the lower eye-lid, as it most commonly is, the lining of this part may be observed rising in the form of a semilunar fold, of a pale red colour, like the fungous granulations of wounds and intervening between the eye and eyelid, which latter it in some measure everts. When the swelling is afterwards occasioned by the lining of both the eyelids, the disease assumes an annular shape, in the centre of which the eyeball seems sunk, while the circumference of the ring presses and everts the edges of the two eye-lids, so as to cause both great uneasiness and deformity. In each of the above cases, on pressing the skin of the eye-lids with the point of the finger, it becomes manifest that they are very capable of being elongated, and would readily yield, so as entirely to cover the eye-ball, were they not prevented by the intervening swelling of their membranous lining.

Besides the very considerable deformity which the disease produces, it occasions a continual discharge of tears over the cheek, and what is worse, a dryness of the eye-ball,

frequent exasperated attacks of chronic ophthalmia, incapacity to bear the light, and lastly, opacity and ulceration of the cornea.

The second species of ectropium, or that arising from a contraction of the integuments of the eyelids, or neighbouring parts, is not unfrequently a consequence of puckered scars, produced by the confluent small-pox, deep burns, or the excision of cancerous or encysted tumours, without saving a sufficient quantity of the skin; or lastly, the disorder is the effect of malignant carbuncles, or any kind of wound attended with much loss of substance. Each of these causes is quite enough to bring on such a contraction of the skin of the eyelids as to draw the parts towards the arches of the orbits, so as to remove them from the eye-ball and turn their edges outward. No sooner has this circumstance happened, than it is often followed by another one equally unpleasant, namely a swelling of the internal membrane of the affected eyelids, which afterwards has a great share in completing the eversion. The lining of the eye-lids, though trivially everted, being continually exposed to the air, and irritation of extraneous substances, soon swells, and rises up like fungus. One side of this fungus-like tumour, covers a part of the eye-ball, the other pushes the eye-lid so considerably outward, that its edge is not unfrequently in contact with the margin of the orbit. The complaints induced by this second species of ectropium are the same as those brought on by the first; it being noticed, however, that in both cases, whenever the disease is very inveterate, the fungous swelling of the inside of the eye-lids becomes hard, coriaceous, and as it were callous.

Although, in both species of ectropium, the lining of the eye-lids, seems equally swollen, yet the surgeon can easily distinguish to which of the two species the disease belongs. For, in the first, the skin of the eyelids, and adjoining parts, is not deformed with scars; and by pressing the everted eyelid with the point of the finger, the part would with ease cover the eye, were it not for the intervening fungous swelling. But in the second species of ectropium, besides the obvious cicatrix and contraction of the skin of the eye-lids, or adjacent parts; when an effort is made to cover the eye with the everted eyelid, by pressing upon the latter part with the point of the finger, it does not give way so as completely to cover the globe, or only yields, as it ought to do, for a certain extent; or it does not move in the least from its unnatural position, by reason of the integuments of the eye-lids having been so extensively destroyed, that their margin has become adherent to the arch of the orbit.

ECTROISIS. (*Barbaric*: from *αἰσχρονομία*, to miscarry.) A miscarriage.

ECTROISIS. (From *αἰσχρονομία*, to miscarry.) An abortion.

ECTROTICA. (From *αἰσχρονομία*, to miscarry.) *Ectrotica*. Medicines which cause abortion.

ECOTYLOICA. See *Ecaltolica*.

ECTYTOTICA. See *Ectrotica*.

ECZEMA. (From *εἰς*, to boil out.)

Eczema. A hot painful eruption, or pustule. Mr. Pearson calls the erythema mercuriale, *eczema mercuriale*.

EDBEPHUS. Prognosis from the nature of elements.

EDERA TRIIFOLIA. The poison-tree of America.

EDES. A name for amber.

EDESSENUM. *Polarium*. An eye-water of tragacanth, arabic, acacia, opium, &c.

EDITE. Amber.

EDIC. *Edich*. *Edir*. An old name for iron.

EDRA. A fracture; also the lower part of the rectum.

EDULCORANTIA. (From *edulco*, to make sweet. *Edulcorants*. Sweeteners. Medicines which absorb the vicious humours of the body, sweeten the fluids, and deprive them of their acrimony.

EFFERVESCENCE. (From *effervesco*, to grow hot.) *Effervescentia*. That agitation which is produced by mixing an acid and an alkali together. A small degree of ebullition.

EFFIDES. An old name for ceruss.

EFFILA. Freckles.

EFFLORESCENCE. (From *effloresco*, to blow as a flower.) *Efflorescentia*.

1. A preternatural redness of the skin.

2. In chymistry it means that phenomenon which takes place upon crystals, producing a white powder when exposed to air.

EFLUVIUM. (From *effluo*, to spread abroad.) See *Contagium*.

EFFRACTURA. (From *effringo*, to break down. *Eclipsma*. A species of fracture, in which the bone is much depressed by the blow.

EFFUSION. (From *effundo*, to pour out.) *Effusio*. In surgery it means the escape of any fluid out of the vessel, or viscous, naturally containing it, and its lodgment in another cavity, in the cellular substance, or in the substance of parts. Effusion also sometimes signifies the natural secretion of fluids from the vessels; thus surgeons frequently speak of the coagulable lymph being effused on different surfaces.

EGEREIS. (From *egere*, to carry out.) *Egeosis*. An excretion, or evacuation.

EGG. *Ovum*. The eggs of poultry are chiefly used as food: the different parts are likewise employed in pharmacy and in medicine. The calcined shell is esteemed as an absorbent. The oil of the egg is softening, and is used externally to burns

and chaps. The yolk of the egg renders oil miscible with water, and is triturated with the same view with resinous and other substances. Raw eggs have been much recommended as a popular remedy for jaundice.

EGEREISIS. (From *egere*, to watch.) A watchfulness. A morbid want of sleep.

EJACULANTIA. (From *ejaculo*, to cast out.) *Ejaculatoria*. The vessels which convey the seminal matter secreted in the testicles to the penis. These are the epididymis, and the vasa deferentia; the vesiculæ seminales are the receptacles of the semen.

EJECTIO. (From *ejicio*, to cast out.) *Ejectio*. The discharging of humours or excrements.

EILANIS. (From *εἰλω*, to involve.) A membrane involving the brain.

EILENA. (From *εἰλω*, to form convolutions.) In Hippocrates it signifies painful convolutions of the intestines from flatulence. Sometimes it signifies a covering. Vogel says it is a fixed pain in the bowels, as if a nail was driven in.

EILEON. (From *εἰλω*, to wind.) *Goræus* says it is a name of the intestinum ileum.

EILINOS. (From *εἰλω*, to form convolutions.) The iliac passion.

EISBOLE. (From *εἰς*, into, and *βολα*, to cast.) It signifies strictly an injection, but is used to express the access of a distemper, or of a particular paroxysm.

EISPHON. (From *εἰς*, into, and *φω*, to breathe.) Inspiration of air.

ELA CALLI. An Indian cathartic shrub, the *Euphorbia nerifolia* of Linnæus.

ELÆAGNOM. (From *ελαιον*, oil, and *αγνος*, chaste.) The agnus castus was formerly so called.

ELÆOSACCHARUM. (From *ελαιον*, oil, and *σακχαρον*, sugar.) A mixture of essential oil with sugar.

ELÆOSELINUM. (From *ελας*, a lake, and *ελαιον*, parsley.) Water parsley.

ELAIS GUINEENSIS. A species of palm which grows spontaneously on the coast of Guinea, but is much cultivated in the West Indies. From this tree is obtained the palm oil which is considered as an emollient and strengthener of all kinds of weakness of the limbs. It also is recommended against bruises, strains, cramps, pains, swellings, &c.

ELAMBICATIO. A method of analyzing mineral waters.

ELANULA. An old name for alum.

ELAPHOBOSCON. (From *ελαφος*, a stag, and *βοσκα*, to eat; so called, because deer eat them greedily.) The wild parsnip. See *Pastinaca*.

ELAPHOSCORODON. (From *ελαφος*, the stag, and *σκορδα*, garlic.) Stag's or viper's garlic.

ELAQUR. Red vitriol.

ELAS MARIS. Butir lead.

ELASMA. (From *elasma*, a lamina or plate of any kind.) A term used to express a cleft r-pipe.

Elastic fluid. See *Gas*.

Elastic gum. See *Indian-rubber*.

ELASTICITY. A force in bodies, by which they endeavour to restore themselves to the posture from whence they were displaced by any external force. To solve this property, many have recourse to the universal law of nature, attraction, by which the parts of solid and firm bodies are caused to cohere together: whereby, when hard bodies are struck or bent, so that the component parts are a little moved from one another, but not quite disjoined or broken off, nor separated so far as to be out of the power of attracting force, by which they cohere together, they certainly must, on the cessation of the external violence, spring back with a very great velocity to their former state; but in this circumstance the atmospherical pressure will account for it as well: because such a violence, if it be not great enough to separate the constituent particles of a body far enough to let in any foreign matter, must occasion many vacuola between the separated surfaces, so that upon the removal they will close again by the pressure of the aerial fluid upon the external parts, i. e. the body will come again into its natural posture. The included air, likewise, in most bodies, gives that power of resiliation upon their percussion.

If two bodies perfectly *elastic* strike one against another, there will be or remain in each the same relative velocity as before, i. e. they will recede with the same velocity as they met together with. For the compressive force, or the magnitude of the stroke in any given bodies, arises from the relative velocity of those bodies, and is proportional to it: and bodies perfectly *elastic* will restore themselves completely to the figure they had before the shock; or, in other words, the restitutive force is equal to the compressive, and therefore must be equal to the force with which they came together, and consequently they must by elasticity recede again from each other with the same velocity. Hence, taking equal times before and after the shock, the distances between the bodies will be equal: and therefore the distances of times from the common centre of gravity will, in the same times, be equal. And hence the laws of percussion of bodies perfectly elastic are easily deduced.

ELATERIUM. (From *elatus*, to stimulate or agitate: so named from its great purgative qualities.) See *Cucumis agrestis*.

ELATERIA. A name for the cascarrilla bark.

ELATINE. (From *elatus*, smaller, being

the smaller species.) Fluellin, or female speedwell. *Antirrhinum elatine* of Linnaeus. The leaves of this plant have a roughish bitter taste, but no smell. It was formerly much used against scurvy and old ulcerations, but now wholly forgotten.

ELATITE. Bloodstone.

ELCOSIA. (From *elus*, an ulcer.) A disease attended with fetid, carious, and chronic ulcers. The term is seldom used.

Elder. See *Sambucus*.

Elder dwarf. See *Ebulus*.

Elecampane. See *Enula campana*.

ELECTRICITY. (*Electricitas*, from *electrum*, amber, from *elektron*, the sun, because of its bright shining colour; or from *elao*, to draw, because of its magnetic power.) A property which certain bodies possess when rubbed, heated, or excited, whereby they attract remote bodies, and frequently emit sparks or streams of light. The ancients first observed this property in amber, which they called *electrum*, and hence arose the word electricity. The efficacy of electricity in the cure of several diseases has been supported by many very respectable authorities, especially in paralytic diseases. It considerably augments the circulation of the blood, and excites the action of the absorbents.

ELECTRODES. (From *elektron*, amber.)

An epithet for stools which shine like amber.

ELECTRUM MINERALIS. The tincture of metals. It is made of tin and copper, to which some add gold, and double its quantity of martial regulus of antimony, melted together; from these there results a metallic mass, to which some chymists have given the name of *electrum mineralis*. This mass is powdered and detonated with nitre and charcoal to a kind of scoria; it is powdered again whilst hot, and then digested in spirit of wine, whence a tincture is obtained of a fine red colour.

ELECTUARIUM. An electuary. The London Pharmacopoeia refers those articles which were formerly called electuaries to confections and conserves.

ELECTUARIUM ANTIMONII. *p.* Electuarii senex, one ounce: guaiaci gummi, hydragry cum sulphure antimonii ppti. sing. half an ounce; syrupi simplicis q. s. misce. Of this electuary from a dram to about two drams is given twice a day, in those cutaneous diseases which go under the general name of scorbatic. It is usually accompanied with the decoctions of elm, bark, or sarsaparilla.

ELECTUARIUM CASSIE. See *Confectio cassie*.

ELECTUARIUM CATECHU. *Confectio Japonica.* Electuary of catechu, commonly called Japonic confection. Take of mimosa catechu, four ounces; kino, three ounces; cinnamon, nutmeg each, one ounce; opium diffused in a sufficient quantity of Spanish white wine one drachm and a

half; syrup of dried roses boiled to the consistence of honey, two pounds and a quarter. Reduce the solids to powder, and having mixed them with the opium and syrup, make them into an electuary. A very useful astringent, and perhaps the most efficacious way of giving the catechu to advantage. Ten scruples of this electuary contain one grain of opium.

ELECTUARIUM CINCHONÆ CUM NATRO. ℞ natri ppti ʒij; pulveris cinchonæ unc; mucilaginis gummi arabici q. s. misce. In this composition mucilage is preferred to syrup, on account of its covering the taste of the bark much more advantageously. It should, for this purpose, however, be made thin, otherwise it will increase the bulk of the electuary too much.

This remedy will be found an excellent substitute for the burnt sponge, whose powers as a remedy in scrophula, are known solely to depend on the proportion of natron contained in it. The dose is two drachms, twice or thrice a day.

ELECTUARIUM OPIATUM. See *Confectio opii*.

ELELISPHACOS. (From *ελεῖν*, to distort, and *εφαξ*, sage; so named from the spiral coiling of its leaves and branches.) A species of sage.

ELEMBRAT. An obsolete term for alkaline salts.

ELEMENTS. Radicals. First principles. The minutest particles of any substance, which can no further be divided or decomposed by chymical analysis. Many substances cannot be farther decomposed by the chymist into constituent parts, but this does not entitle the ranking them among the elements. Though they are as yet decomposed, it does not follow that they are *undecomposable*; as, perhaps, neither our senses nor our instruments will ever reach those substances which by their nature admit of no sort of decomposition. The bodies which are known to us at present, however, as simple substances, amount to forty-one; some of these may be sensibly exhibited in their simple state, uncombined with other matters: these are termed *ostensible, producible, simple substances*, to distinguish them from those whose existence or presence is only inferred from facts, and are called *unostensible, unproducible, simple substances*. The following is a list of simple substances at present known.

Unproducible, simple Substances.

- | | |
|-----------------------------------|----------------------|
| 1. Phlogiston, or basis of light. | 10. Boracic radical. |
| 2. Oxygen. | 11. Rad. of gold. |
| 3. Hydrogen. | 12. ——— platina. |
| 4. Azotic. | 13. ——— silver. |
| 5. Carbonic. | 14. ——— mercury. |
| 6. Sulphur. | 15. ——— lead. |
| 7. Phosphor. | 16. ——— copper. |
| 8. Muriatic. | 17. ——— iron. |
| 9. Fluoric. | 18. ——— tin. |
| | 19. ——— zinc. |

- | | |
|----------------------|---------------------|
| 20. Rad. of bismuth. | 26. Rad. of molybd. |
| 21. ——— antim. | 27. ——— wolfranc. |
| 22. ——— nickel. | 28. ——— uranium. |
| 23. ——— cobalt. | 29. ——— titanium. |
| 24. ——— arsenic. | 30. ——— tellurium. |
| 25. ——— mangan. | 31. ——— chromic. |

Producible, ostensible, simple Substances.

- | | | |
|----------------|----------------|----------|
| 32. Caloric. | 37. S. rontian | } Earth. |
| 33. Siliceous. | 38 Argillac. | |
| 34. Calcar. | 39. Glucine. | } Alk. |
| 35. Magnesia. | 40. Vegetab. | |
| 36. Ponder. | 41. Mineral. | |

The alkalis have lately been decomposed.

See *Potassa* and *Soda*.

ELEMI. (It is said this is the Ethiopian name) Gum elemi. The parent plant of this resin is supposed to be the *Amryis elemifera* of Linnæus:—*foliis terribus quinatopinnatisque subtus tomentosis*. Elemi is brought here from the Spanish West Indies: it is most esteemed when softish, somewhat transparent, of a pale whitish colour, inclining a little to green, and of a strong, though not unpleasant smell. It is only used in ointments and plasters, and is a powerful digestive.

ELEMI UNQUENTUM. See *Unguentum elemi compositum*.

ELEMIFERA CURASSAVICA ARBOR. The gum elemi-tree.

ELENGI. A tree of Malabar. Cardiac.

ELEOCHRYSUM (From *ελεος*, the sun, and *χρυσος*, gold; so called from their shining yellow appearance) *Elio-chrysum*. Goldilocks. See *Elio-chrysum*.

ELEOSILINUM. (From *ελεος*, a lake, and *σιν*, par-ley.) See *Apium*.

ELEPHANTIA. (From *ελεphas*, an elephant; so called from the great enlargement of the body in this disorder.) A species of anasarca.

ELEPHANTIA ARABUM. In Dr. Cullen's nosology it is synonymous with elephantiasis. The term is, however, occasionally confined to this disease when it affects the feet.

ELEPHANTIASIS. (From *ελεphas*, an elephant; so named from the legs of people affected with this disorder growing scaly, rough, and wonderfully large, at an advanced period, like the legs of an elephant.) *Elephas. Elephantia. Lazari morbus vel malum. Phenicæus morbus* A disease that attacks the whole body, but mostly affects the feet, which appear somewhat like those of the elephant. It is known by the skin being thick, rough, wrinkly, unctuous, and void of hair, and mostly without the sense of feeling. It is said to be contagious. Cullen makes it a genus of disease in the class *cachexia*, and order *impetiginæ*.

Elephantiasis has generally been supposed to arise in consequence of some slight attack of fever, on the cessation of which the morbid matter falls on the leg, and occasions a distention and tumefaction of the limb, which is afterwards overspread

with uneven lumps, and deep fissures. By some authors it has been considered as a species of leprosy; but it often subsists for many years without being accompanied with any of the symptoms which characterize that disease.

It sometimes comes on gradually, without much previous indisposition; but more generally, the person is seized with a coldness and shivering, pains in the head, back, and loins, and some degree of nausea. A slight fever then ensues, and a severe pain is felt in one of the inguinal glands, which, after a short time, becomes hard, swelled, and inflamed. No suppuration, however, ensues; but a red streak may be observed running down the thigh from the swelled gland to the leg. As the inflammation increases in all the parts, the fever gradually abates, and perhaps, after two or three days continuance, goes off. It however, returns again at uncertain periods, leaving the leg greatly swelled with varicose turgid veins, the skin rough and rugged, and a thickened *membrana cellulosa*. Scales appear also on the surface, which do not fall off, but are enlarged by the increasing thickness of the membranes; uneven lumps, with deep fissures, are formed, and the leg and foot become at last of an enormous size.

A person may labour under this disease many years, without finding much alteration in the general health, except during the continuance of the attacks; and perhaps the chief inconvenience he will experience is the enormous bulky leg which he drags about with him. The incumbrance has, indeed, induced many who have laboured under this disease to submit to an amputation; but the operation seldom proves a radical cure, as the other leg frequently becomes affected.

Hillary observes, that he never saw both legs swelled at the same time. Instances where they have alike acquired a frightful and prodigious size, have, however, frequently fallen under the observation of other physicians.

ELIPHANTINUM EMPLASTRUM. A plaster described by Oribasius. Celsus describes one of the same name, but very different in qualities.

ELEPHAS. (*Elephas*, the elephant.) The disease called elephantiasis; also *aqua fortis*.

ELERSNA. An obsolete term for black lead.

ELESMATIS. An old term for burnt lead.

ELIETARI FRIMUM. See *Anemum verum*.

ELIUTHERIA BARK. See *Cascarilla*.

ELIUTHERIA CORTEX. See *Cascarilla*.

ELEVATIO. (From *elevo*, to lift up.) Elevation. Sublimation.

ELEVATOR. (From *elevo*, to lift up.) A muscle is so called whose office is to lift

up the part to which it is attached. Also a chirurgic instrument *elevatorium*, with which surgeons raise any depressed portion of bone, but chiefly those of the cranium.

ELEVATOR LABII INFERIORIS PROPRIUS. See *Levator labii inferioris*.

ELEVATOR LABII SUPERIORIS PROPRIUS. See *Levator labii superioris alicque nasi*.

ELEVATOR LABIORUM. See *Levator anguli oris*.

ELEVATOR NASI ALARUM. Muscles of the ala of the nose.

ELEVATOR OCULI. See *Levator oculi*.

ELEVATOR PALPEBRÆ SUPERIORIS. See *Levator palpebræ superioris*.

ELEVATOR SCAPULÆ. See *Levator scapulae*.

ELEVATORIUM. (From *elevo*, to lift up.) An instrument to raise a depression in the skull.

ELIBANUM. See *Olibanum*.

ELICHRYSUM. (From *elao*, the sun, and *chrysos*, gold; so called from their shining yellow appearance.) *Stachas citrina*. Goldilocks. This small downy plant is the *Gnaphalium stachas* of Linnaeus. The flowers are warm, pungent, and bitter, and said to possess aperient and corroborant virtues.

ELIDION. Mastich; a mixture of brass.

ELISMA. An old name for a linctus.

ELIOMELI. (From *elao*, and *melas*, honey.) A sweet purging oil like honey.

ELIOSSELINUM. See *Eleoselinum*.

ELITHROIDES. The vaginal coat of the testicle.

ELIXATIO. (From *elizo*, to boil.) The act of seething, or boiling.

ELIXIR. (From *elektar*, an Arabic word signifying quintessence.) A term formerly applied to many preparations similar to compound tinctures. It is now very little employed.

ELIXIR OF HEALTH. *Elixir salutis*. A term formerly applied to what is now called compound tincture of senna. See *Tinctura sennæ compoſita*.

ELIXIR PAREGORICUM. Paregoric elixir. See *Tinctura opii camphorata*.

ELIXIR PROPRIETATIS. A preparation of aloes.

ELIXIR SACRUM. See *Tinctura rhæi cum aloë*.

ELIXIR SALUTIS. See *Tinct. sennæ comp.*

ELIXIR STOMACHICUM. Stomachic elixir. See *Tinctura gentianæ compoſita*.

ELIXIVATIO. (From *elizo*, to boil, or from *elivium*, lye.) The extraction of a fixed salt from vegetables, by an affusion of water.

ELIIBORUM. See *Helleborus albus*.

ELM. See *Ulmus*.

ELMINTHES. (From *elao*, to involve, from its contortions.) Worms.

Elm-leaved sumach. See *Sumach*.

ELMDA. (From *elao*, a swamp.) A

term given to a sweating fever, from its great moisture.

ELONGATIO. (From *elongo*, to lengthen out.) An imperfect luxation, where the ligament is only lengthened, and the bone not put out of its socket.

ELUTRIATIO. (From *elutriatio*, to cleanse.) Washing over. It is the pouring a liquor out of one vessel into another, in order to separate the subsiding matter from the clear and fluid part.

ELUVIES (From *eluo*, to cleanse.) The effluvia from a swampy place. Also the humour discharged in fluor albus.

ELUXATIO. (From *eluxo*, to put out of joint.) A luxation, or dislocation.

ELYNASTROSTIS. (From *ελυστρος*, the herb panic, and *αγροσ*, wild.) Wild panic.

ELYMUS. (*Ελμυος*.) The herb panic.

ELYTROCELE. (From *ελυτρος*, the vagina, and *κελ*, a tumour.) A hernia in the vagina.

ELYTROIDES. (*Elytroides*; from *ελυτρος*, a sheath, and *ειδος*, form.) Like a sheath. The tunica vaginalis is so called by some writers, because it includes the testes like a sheath.

ELYTRON. (From *ελυα*, to involve.) The vagina. A sheath. The membranes which involve the spinal marrow are called *elytra*, *ελυτρα*.

EMARGINATIO. (From *emargino*, to cleanse the edges.) The cleansing of the edges of wounds from scurf and filth.

EMASCLATUS. (From *emasculo*, to render impotent.) Having the testicles in the belly, and not fallen into the scrotum.

EMAMMA. (From *εμβασμα*, to immerge in.) A medicated pickle to dip the food in.

EMBOLE. (From *εμβολα*, to put in.) The reduction or setting of a dislocated bone.

EMBOLUM. (From *εμβολα*, to cast out; so named because it ejects the semen.) The penis.

EMBRISMA. (From *εμβριση*, to make wet.) A fluid application to any part of the body.

EMBROCATIO ALUMINIS. ꝑ. *Aluminis* ʒij. *Aceti spiritus vinosi tenuioris*, sing. lbs̄ss. For chilblains and diseased joints.

EMBROCATIO AMMONIÆ. ꝑ. *embrocationis ammoniæ acetatæ cum saponē* ʒij. *Aquæ ammoniæ puræ* ʒij. For sprains and bruises.

EMBROCATIO AMMONIÆ ACETATÆ CAMPHORATÆ. ꝑ. *solutionis saponis cum camphora, aquæ ammoniæ acetatæ* sing. ʒj. *Aquæ ammoniæ puræ* ʒss. For sprains and bruises. It is also frequently applied to disperse chilblains which have not suppurated. It is said to be the same as *sters' opodeldoc*.

EMBROCATIO AMMONIÆ ACETATÆ. ꝑ. *aquæ ammoniæ acetatæ. Solutionis sa-*

ponis sing. ʒj. M. For bruises with inflammation.

EMBROCATIO CANTHARIDIS CUM CAMPHORA. ꝑ. *cantharidia. Spiritus camphoræ* sing. ʒj. M. This may be used in any case in which the object is to stimulate the skin. The absorption of cantharides, however, may bring on a strangury.

EMBROCATIO. (From *εμβριση*, to moisten, or soak in.) *Embroche. Embrocatio.* A fluid application to rub any part of the body with. Many use the term, however, as synonymous with liniment. The following embrocations are noticed in the *Pharmacopœia Chirurgicalis*.

EMBROCHE. See *Embrocation*.

EMBRYO. (From *εμβρυον*, to bud forth.) The *fœtus in utero* is so called before the fifth month of pregnancy, because its growth resembles that of the budding of a plant.

EMBRYOTHLASTES. (From *εμβρυον*, the fœtus, and *θλαω*, to break.) *Embryorectes.* A chirochet, or instrument for breaking the bones of a dead fœtus to promote its delivery.

EMBRYOTOMY. (From *εμβρυον*, a fœtus, and *τομω*, to cut.) *Embryotomia.* The separating of any part of the fœtus whilst *in utero*, to extract it.

EMBRULCUS. (From *εμβρυον*, a fœtus, and *ελκω*, to draw.) A blunt hook, or forceps, for drawing the child from the womb.

EMERUS. Scorpion senna. A laxative.

EMESIA. (From *εμεω*, to vomit.) *Emesma. Emetia.* The act of vomiting. Medicines which cause vomiting.

EMETICS. (*Emetica, sc. medicamenta;* from *εμεω*, to vomit.) Substances capable of exciting vomiting, independent of any effect arising from the mere quantity of matter introduced into the stomach, or of any nauseous taste or flavour.

The susceptibility of vomiting is very different in different individuals, and is often considerably varied by disease.

Emetics are employed in many diseases.

When any morbid affection depends upon, or is connected with, over distention of the stomach, or the presence of acrid, indigestible matters, vomiting gives speedy relief. Hence its utility in impaired appetite, acidity in the stomach, in intoxication, and where poisons have been swallowed.

From the pressure of the abdominal viscera in vomiting, emetics have been considered as serviceable in jaundice, arising from biliary calculi obstructing the ducts.

The expectorant power of emetics, and their utility in catarrh and phthisis, have been ascribed to a similar pressure extended to the thoracic viscera.

In the different varieties of febrile affections, much advantage is derived from exciting vomiting, especially in the very commencement of the disease. In high inflam-

matory fever it is considered as dangerous, and in the advanced stage of typhus it is prejudicial.

Emetics, given in such doses as only to excite nausea, have been found useful in restraining hamorrhage.

Different species of dropsy have been cured by vomiting, from its having excited absorption. To the same effect, perhaps, is owing the dispersion of swelled testicle, bubo, and other swellings, which have occasionally resulted from this operation.

The operation of vomiting is dangerous, or hurtful, in the following cases: where there is determination of the blood to the head, especially in plethoric habits; in visceral inflammation; in the advanced stage of pregnancy; in hernia and prolapsus uteri; and wherever there exists extreme general debility. The frequent use of emetics weakens the tone of the stomach. An emetic should always be administered in the fluid form. Its operation may be promoted by drinking any tepid diuretic, or bitter infusion.

The individual emetics may be arranged under those derived from the vegetable, and those from the mineral kingdom. From the vegetable kingdom are numbered ipecacuanha, scilla maritima, anthemis nobilis, sinapis alba, asarum Europæum, nicotiana tabacum. From the mineral kingdom, antimony, sulphat of zinc and copper, and the subacetat of copper. To these may be added ammoniac and its hydro-sulphuret.

EMETOCATHARTICUS. (From *emetos*, to vomit, and *kathartikos*, to purge.) Purging both by vomit and stool.

EMINENTIE QUADRIGEMINE. See *Tubercula quadrigemina*.

EMMENAGOGUES. (*Emmenagoga*, sc. *medicamenta, emmenagoga*: from *emmenae*, the menses, and *ago*, to move.) Those medicines that possess a power of promoting that monthly discharge of blood by the uterus, which, from a law of the animal æconomy, should take place in certain conditions of the female system. The articles belonging to this class may be referred to four orders:

1. *Stimulating emmenagogues*, as *hydrargyrite* and *antimonial preparations*, which are principally adapted for the young, and those with peculiar insensibility of the uterus.

2. *Irritating emmenagogues*, as *aloes*, *savine*, and *Spanish flies*: these are to be preferred in torpid and chlorotic habits.

3. *Tonic emmenagogues*, as *ferruginous preparations*, *cold bath*, and *exercise*, which are advantageously selected for the lax and phlegmatic.

4. *Antispasmodic emmenagogues*, as *assafoetida*, *castor*, and *pediluvia*: the constitutions to which these are more especially

suited are the delicate, the weak, and the irritable.

EMMENEA. (From *em*, in, and *men*, a month.) The menstrual flux.

EMOLLIENTS. (*Emollientia*, sc. *medicamenta*; from *emollio*, to soften.) Those substances which possess a power of relaxing the living and animal fibre, without producing that effect from any mechanical action. The different articles belonging to this class of medicines may be comprehended under the following orders:

1. *Humectant emollients*, as *warm water* and *tepid vapours*, which are fitted for the robust and those in the prime of life.

2. *Relaxing emollients*, as *althea*, *malva*, &c. These may be employed in all constitutions, while, at the same time, they do not claim a preference to others from any particular habit of body.

3. *Lubricating emollients*, as *bland oils*, *fat*, and *tard*. The same observation will hold of this order as was made of the last mentioned.

4. *Atonic emollients*, as *opium* and *pediluvia*: these are applicable to any constitution, but are to be preferred in habits where the effects of this class are required over the system in general.

EMPERIA. (From *em*, and *pega*, to endeavour.) Professional experience.

EMPHROMENUS. (From *emphro*, to bear.) Urine, or other substance, which has a sediment.

EMPHRACTICA. (From *emphrakto*, to obstruct.) Medicines which applied to the skin, shut up the pores.

EMPHYSEMA. (From *emphusos*, to inflate.) *Pneumatosus*. Air in the cellular membrane. In general it is confined to one place; but in a few cases it spreads universally over the whole body, and occasions a considerable degree of swelling. It sometimes arises spontaneously, which is, however, a very rare occurrence, or comes on immediately after delivery, without any evident cause; but it is most generally induced by some wound or injury done to the thorax, and which affects the lungs; in which case, the air passes from these, through the wound, into the surrounding cellular membrane, and from thence spreads over the whole body.

Emphysema is attended with an evident crackling noise, and elasticity upon pressure; and sometimes with much difficulty of breathing, oppression, and anxiety.

We are to consider it as a disease by no means unattended with danger; but more probably from the causes which give rise to it, than any hazard from the complaint itself.

EMPIRIC. (*Empiricus, empiricus*: from *em* in, and *pega*, experience.) One who practises the healing art upon experience, and not theory. This is the true meaning of the word empiric: but it is now applied, in a very opposite sense, to those who de-

viate from the line of conduct pursued by scientific and regular practitioners, and vend nostrums, or sound their own praise in the public papers.

EMPLASTICA. (From *εμπλαστον*, to obstruct.) Medicines which, spread upon the skin, stop the pores.

EMPLASTRUM. (From *εμπλαστον*, to spread upon.) A plaster. Plasters are composed of unctuous substances, united either to powders or metallic oxyds, &c. They ought to be of such a consistence as not to stick to the fingers when cold, but to become soft, so as to be spread out, in a moderate degree of heat, and in that of human body, to continue tenacious enough to adhere to the skin. They owe their consistence either to metallic oxyds, especially those of lead, or to wax, resin, &c. They are usually kept in rolls wrapped in paper, and spread, when wanted for use, upon thin leather; if the plaster be not of itself sufficiently adhesive, it is to be surrounded at its margin by a boundary of resin plaster.

EMPLASTRUM AMMONIACI. Take of purified ammoniacum, five ounces; acetic acid half a pint. Dissolve the ammoniacum in the acid, then evaporate the liquor in an iron vessel, by means of a water-bath, constantly stirring it, until it acquires a proper consistence. This plaster is now first introduced in the London Pharmacopœia; it adheres well to the skin, without irritating it, and without producing inconvenience by its smell.

EMPLASTRUM AMMONIACI CUM HYDRARGYRO. "Take of purified ammoniacum, a pound; purified mercury, three ounces; sulphurated oil, a drachm." Rub the mercury with the sulphurated oil until the globules disappear; then add by degrees the ammoniacum, previously melted, and mix the whole together. This mixture of ammoniacum hydrargyrus and sulphur, is said to possess resolvent virtues; and the plaster is recommended with this view to be applied to nodes, tophi, indurated glands, and tumours.

EMPLASTRUM ASEFETIDE. *Emplastrum antihystericum.* Plaster of *asafoetida*. Take of plaster of semi-vitrified oxyd of lead, *asafoetida*, each two parts; galbanum, yellow wax, each one part. This plaster is said to possess anodyne and antispasmodic virtues. It is, therefore, occasionally directed to be applied to the umbilical region in hysterical cases.

EMPLASTRUM CANTHARIDIS. See *Emplastrum lytte*.

EMPLASTRUM CERÆ. Wax plaster. *Emplastrum attrahens.* Take of yellow wax, prepared suet, of each three pounds; yellow resin, a pound. Mix them together and strain. This is a gently drawing preparation, calculated to promote a moderate discharge from the blistered surface,

with which intention it is mostly used. Where the stronger preparations irritate, this will be found in general to agree.

EMPLASTRUM CUMINI. Cumin plaster, "Take of cumin-seeds, caraway-seeds, bay-berries, of each three ounces; dried pitch, three pounds; yellow wax, three ounces." Having melted the dried pitch and wax together, add the remaining articles, previously powdered, and mix. A warm stomachic plaster, which, when applied to the stomach, expels flatulency. To indolent scrophulous tumours, where the object is to promote suppuration, this is an efficacious plaster.

EMPLASTRUM GALBANI COMPOSITUM. Compound Galbanum plaster, formerly called *emplastrum lithargyri compositum* and *diachylon magnum cum gummi*. Take of galbanum gum resin, eight ounces. Lead plaster, three pounds; common turpentine, ten drachms; resin of the spruce fir, three ounces. Having melted the galbanum gum resin with the turpentine, mix in first the powdered resin of the spruce fir, and then the lead plaster, previously melted by a slow fire, and mix the whole. This plaster is used as a warm digestive and suppurative, calculated to promote maturation of indolent or scirrhus tumours, and to allay the pains of sciatica, arthrodynia, &c.

EMPLASTRUM HYDRARGYRI. Mercurial plaster. *Emplastrum lithargyri cum hydrargyro.* "Take of purified mercury, three ounces; sulphurated oil, a fluid drachm; lead plaster, a pound." Rub the mercury with the sulphurated oil, until the globules disappear; then add by degrees the lead plaster, melted, and mix the whole.

EMPLASTRUM LADANI COMPOSITUM. This may be used with the same intentions as the cumin plaster, to which it is in no way superior, though composed of more expensive materials. Formerly it was considered as a very elegant stomach plaster, but is now disused.

EMPLASTRUM LITHARGYRI. *Diachylon*, "Take of semi-vitrified oxyd of lead, in very fine powder, five pounds; olive oil, a gallon; water, two pints." Boil them with a slow fire, constantly stirring until the oil and litharge unite, so as to form a plaster. Excoriations of the skin, slight burns, and the like may be covered with this plaster; but it is in more general use, as a defensive, where the skin becomes red from lying a long time on the part,

EMPLASTRUM LITHARGYRI COMPOSITUM. See *Emplastrum Galbani compositum*.

EMPLASTRUM LITHARGYRI CUM RESINA. See *Emplastrum resine*.

EMPLASTRUM LYTTE. Blistering fly-plaster. *Emplastrum cantharides.* *Emplastrum vesicatorium.* Take of blistering flies, in very fine powder, a pound; wax

plaster, a pound and a half; prepared fat, a pound. Having melted the plaster and fat together, and removed them from the fire, a little before they become solid sprinkle in the blistering flies, and mix the whole together. See *Blister* and *Cantharides*.

EMPLASTRUM OPII. Plaster of opium. "Take of opium, powdered, half an ounce; resin of the spruce fir, powdered, three ounces; lead plaster, a pound." Having melted the plaster, mix in the resin of the spruce fir and opium, and mix the whole. Opium is said to produce somewhat, though in a smaller degree, its specific effect when applied externally.

EMPLASTRUM PICIS COMPOSITUM. Compound pitch plaster. *Emplastrum picis Burgundicæ.* "Take of dried pitch, two pounds; resin of spruce fir, a pound; yellow resin, yellow wax, of each four ounces; expressed oil of nutmegs, an ounce." Having melted together the pitch, resin, and wax, add first the resin of the spruce fir, then the oil of nutmegs, and mix the whole together. From the slight degree of readiness this stimulating application produces, it is adapted to gently irritate the skin, and thus relieve rheumatic pains. Applied to the temples it is sometimes of use in pains of the head.

EMPLASTRUM PLUMBI. Lead plaster *Emplastrum lithargyri* *Emplastrum commune.* *Diachylon simplex.* This plaster is of great importance, as forming the basis, by additions to which many other plasters are prepared. See *Emplastrum lithargyri*.

EMPLASTRUM RESINÆ. Resin plaster. *Emplastrum lithargyri cum resina.* "Take of yellow resin, half a pound; lead plaster, three pounds. Having melted the lead plaster over a slow fire, add the resin in powder, and mix." The adhesive, or sticking plaster, is chiefly used for keeping on other dressings, and for retaining the edges of recent wounds together.

EMPLASTRUM SAPONIS. Soap plaster. Take of hard soap sliced, half a pound; lead plaster, three pounds. Having melted the plaster, mix in the soap; then boil it down to a proper consistence. Discutient properties are attributed to this elegant plaster, with which view it is applied to lymphatic and other indolent tumours. It forms an admirable defensive and soft application, spread on linen, to surround a fractured limb.

EMPLASTRUM THURIS COMPOSITUM. Compound frankincense plaster. Take of frankincense, half a pound; dragon's blood, three ounces; litharge plaster, two pounds. To the melted lead plaster add the rest powdered. This plaster is said to possess strengthening, as well as adhesive powers. By keeping the skin firm, it may give tone to the relaxed muscles it surrounds, but cannot, in any other way, impart more

strength than the common adhesive plaster.

EMPNEUMATOSIS. (From *em*, in, and *pnos*, to blow.) An inflation of the stomach, or any other viscus.

EMPORIUM. (From *emporion*, to negotiate.) A mart. The brain is so called, as being the place where all rational and sensitive transactions are collected.

EMPRION. (From *em*, and *prion*, a saw.) Serrated. An epithet of a pulse, in which the artery at different times is unequally distended.

EMPROSTHOTONOS. (From *emprosthen*, before, or forwards, and *tonos*, to draw.) A clonic spasm of several muscles, so as to keep the body in a fixed position and bent forward. Cullen considers it as a species of tetanus. See *Tetanus*.

EMPTYSIS. (From *empton*, to spit out.) A discharge of blood from the mouth and fauces.

EMPYEMA. (From *em*, within, and *pus*, pus.) A collection of pus in the cavity of the thorax. It is one of the terminations of pleuritis. There is reason for believing that matter is contained in the cavity of the chest, when, after a pleurisy, or inflammation in the thorax, the patient has a difficulty of breathing, particularly on lying on the side opposite the affected one; and when an œdematous swelling is externally perceptible.

EMPYEMATA. (From *em*, and *pus*, pus.) Suppurating medicines.

EMPYREUMA. (From *empyreuma*, to kindle, from *pyr*, fire.) The offensive smell that distilled waters and other substances receive from being exposed too much to fire.

EMPYREUMATIC. (*Empyreumatica*; from *empyreuma*, to kindle.) Smelling as it were burnt; thus empyreumatic oils are those distilled with a great heat, and impregnated with a smell of the fire.

EMULGENT. (*Emulgentia*; from *emulgeo*, to melt out; applied to the veins and arteries which go from the aorta and vena cava to the kidneys, because the ancients supposed they strained, and, as it were, milked the serum through the kidneys.) The vessels of the kidneys are so termed. The emulgent artery is a branch of the aorta. The emulgent vein evacuates its blood into the ascending cava.

EMULSIO AMYGDALÆ COMMUNIS. Almond emulsion. Take of almonds, one ounce; water, two pounds and a half. Beat the blanched almonds in a stone mortar, gradually pouring on them the water; then strain off the liquor. It possesses cooling and demulcent properties.

EMULSIO ARABICA. This is made in the same manner as the almond emulsion, only adding two ounces, while beating the almonds mucilage, of gum arabic. This cooling and demulcent emulsion, ordered

in the Edinburgh Pharmacopœia, may be drank ad libitum to mitigate ardor urinæ, whether from the venereal virus or any other cause. In difficult and painful micturition, and strangury, it is of infinite service.

EMULSIO CAMPHORATA. "Take of camphor, one scruple; sweet almonds, blanch'd, two drachms; double refined sugar, one drachm; water, six ounces." This is to be made in the same manner as the common emulsion. It is calculated for the stomachs of those who can only bear small quantities of camphire.

EMULSION. (*Emulsio*; from *emulgio*, to milk.) A soft and somewhat oily medicine, resembling milk.

Emulsion, almond. See *Emulsio amygdale communis*.

Emulsion, Arabic. See *Emulsio arabica*.

Emulsion, camphorated. See *Emulsio camphorata*.

Emulsion of asafetida. See *Mistura asafetide*.

Emulsion of gum-ammoniac. See *Mistura ammoniaci*.

EMUNCTORY. (From *emungo*, to drain off.) The excretory ducts of the body are so termed; thus the exhaling arteries of the skin constitute the great emunctory of the body.

ENEMA. (From *en*, and *aima*, blood.) *Enemas*. So Hippocrates and Galen call such topical medicines as are appropriated to bleeding wounds.

ENUREMA. (From *en*, and *ouros*, to lift up.) The pendulous substance which floats in the middle of the urine.

ENAMEL. See *Teeth*.

ENANTHESIS. (From *en*, and *anthe*, to meet.) The near approach of ascending and descending vessels.

ENARTHROSIS. (From *en*, in, and *arthron*, a joint.) The ball and socket-joint. A species of diarthrosis, or moveable connexion of bones, in which the round head of one is received into the deeper cavity of another, so as to admit of motion in every direction; as the head of the os femoris with the acetabulum of the os innominatum. See *Articulation*.

ENCANTHIS. (From *en*, and *kanthos*, the angle of the eye.) A disease of the caruncula lachrymalis, of which there are two species. *Encanthis benigna*, and *Encanthis maligna seu inveterata*.

The encanthis, at its commencement, is nothing more than a small, soft, red, and sometimes rather livid excrescence, which grows from the caruncula lachrymalis, and, at the same time, from the neighbouring semimular fold of the conjunctiva. This excrescence, on its first appearance, is commonly granulated, like a mulberry, or is of a ragged and fringed structure. Afterwards, when it has acquired a certain size, one part of it represents a granulated

tumour, while the rest appears like a smooth, whitish, or ash-coloured substance, streaked with varicose vessels, sometimes advancing as far over the conjunctiva, covering the side of the eye next to the nose, as where the cornea and sclerotica unite.

The encanthis keeps up a chronic ophthalmia, impedes the action of the eyelids, and prevents, in particular, the complete closure of the eye. Besides, partly by compressing and partly by displacing the orifices of the puncta lachrymalis, it obstructs the free passage of the tears into the nose. The inveterate encanthis is ordinarily of a very considerable magnitude; its roots extend beyond the caruncula lachrymalis and semimular fold to the membranous lining of one or both eyelids. The patient experiences very serious inconvenience from its origin and interposition between the commissure of the eye-lids, which it necessarily keeps asunder on the side towards the nose. Sometimes the disease assumes a cancerous malignancy. This character is evinced by the dull red, and, as it were leaden colour of the excrescence; by its exceeding hardness, and the lancing pains which occur in it, and extend to the forehead, the whole eye-ball and the temple, especially when the tumour has been slightly touched. It is also shewn, by the propensity of the excrescence to bleed, by the partial ulcerations on its surface, which emit a fungous substance, and a thin and exceedingly acrid discharge.

ENCATALEPSIS. (From *en*, and *καταληψω*, to leave.) A catalepsy.

ENCATHISMA. (From *en*, and *καθιμι*, to sit in.) A semicupium. A bath for half the body.

ENCAUMA. (From *en*, in, and *καω*, to burn.) *Encausis*. A pustule produced from a burn.

ENCAUSIS. (From *en*, and *καω*, to burn.) A burn, or scald.

ENCEPHALOCELE. (From *encephalon*, the brain, and *cele*, a tumour.) A rupture of the brain.

ENCEPHALON. (From *en*, in, and *κεφαλη*, the head.) *Encephalum*. By some writers the cerebrum only is so called; and others express by this term the contents of the cranium.

ENCERIS. (From *en*, and *κερος*, wax.) A roll of wax for making plasters.

ENCEROSIS. (From *en*, and *κερος*, to wax.) The covering of a plaster with wax.

ENCHARAXIS. (From *en*, and *καρσσω*, to scarify.) A scarification.

ENCHEIRESIS. (From *en*, and *χειρ*, the hand.) *Encheiria*. Galen uses this word as part of the title to one of his works, which treats of dissection. The word imports the manual treatment of any subject

ENCHIRIA. See *Encheiresis*.

ENCHILOMA. See *Enchyloma*.

ENCHONDRIUS. (From *ω*, and *χονδρος*, a cartilage.) A cartilage.

ENCHRISTA. (From *ενχρησμι*, to anoint.) Unguents. Ointments.

ENCHYLOMA. (From *ω*, and *χυλος*, juice.) An inspissated juice. An elixir, according to Lemery.

ENCHYMA. (From *ω*, and *χυν*, to infuse.) An infusion. A sanguineous plethora.

ENCHYMATA. (From *ενχυνω*, to infuse.) Injections for the eyes and ears.

ENCHYMONA. (From *ω*, and *χυν*, to pour in.) In the writings of the ancient physicians, it is a word by which they express that sudden effusion of blood into the cutaneous vessels, which arises from joy, anger, or shame; and in the last instance is what we usually call blushing.

ENCHYMOISIS. (*ενχυμοσις*.) Blushing; also an extravasation of blood, which makes the part appear livid. Thus, but improperly, it is synonymous with *Ecchymosis*.

ENCHYSIS. See *Enchyma*.

ENCHYSMA. (From *ω*, and *κλυσω*, to cleanse out.) A clyster.

ENGELIUM. (From *ω*, within, and *καλας*, the belly.) The abdominal viscera.

ENGOLISMUS. (From *ενκολισμος*, to insinuate.) An uterine injection.

ENCHRANIUM. (From *ω*, within, and *κρανιον*, the skull.) The cerebrum; the whole contents of the skull.

ENCHRASICHOLUS. (From *ω*, in, and *καρκα*, the head; and *χολα*, bile; because it is said to have the gall in its head.) The anchovy.

ENCHRES. *Ενχρε*. A cake of meal, oil, and honey.

ENCHYMON. (From *ω*, and *κυν*, to conceive.) Pregnancy.

ENCHYSIS. (From *ω*, and *κυν*, to bring forth.) Parturition.

ENCYSTED. A term applied to those tumours which consist of a fluid or other matter, enclosed in sac or cyst.

ENCHYSIS. (From *ω*, in, and *κυστις*, a bag.) A wen. A hard tumour.

ENDEMIC. (*Endemicus*; from *ω*, in, and *δημος*, people.) A disease is so termed that is peculiar to a certain class of persons, or country; thus struma is endemic to the inhabitants of Derbyshire and the Alps; scurvy to seafaring people; and the plica polonica is met with in Poland.

ENDESIS. (From *ω*, and *δεω*, to tie up.) A ligature. A bandage.

Endive. See *Endivia*.

ENDIVIA. (*Quasi cundo via, quia parvisim naciatur*; named from the quickness of its growth.) *Endiva*. Endive. This plant, *Cichorium endivia*; *floribus solitariis, pedunculatis; foliis integris, crenatis*, of Linnæus, is an extremely wholesome salad, possessing bitter and anodyne qualities.

ENDEOSIS. (From *ω* and *διδωμι*, to give.)

A remission, particularly of febrile disorders.

ENELLASMENUS. (From *εναλλάττω*, to interchange.) An epithet applied to the union of the joints of the vertebrae.

ENEMA. (From *ενεμι*, to inject.) Clyster. Injection. *Lavement*. A clyster. A well-known form of conveying both nourishment and medicine to the system, under certain morbid circumstances. The former takes place where obstruction of the passage to the stomach is so great as to render access to that organ impossible, such as occurs in lock jaw, diseased œsophagus, &c. By this means the body can be supported for a few weeks, till an attempt is made at effecting a cure. It is composed, in such cases, of animal broths, gruels made of farinaceous seeds, mucilages, &c. As a form of medicine, clysters are no less useful; and, according to the intention with which they are prescribed, they are either of an emollient, anodyne, or purgative nature. The following forms are in general use.

ENEMA ANODYNUM. Take of starch jelly, half a pint; tincture of opium, forty to sixty drops. Mix. The whole to be injected by means of a pewter clyster-syringe, in cases of dysentery or violent purging, and pain in the bowels.

ENEMA ANTISPASMODICUM. Take of tinct. of asafoetida, half an ounce; tincture of opium, forty drops, Mix. For spasmodic affections of the bowels.

ENEMA LAXATIVUM. Take of Epsom salt, two ounces; dissolve in three quarters of a pint of warm gruel, or broth, with an ounce of fresh butter, or sweet oil.

ENEMA NICOTIANÆ. Take of the leaves of tobacco, two drachms; boiling water, one pound. The tobacco to be infused in the water for the space of ten minutes. Employed in cases of strangulated hernia.

ENEMA NUTRIENS. Take of strong beef tea, twelve ounces; thicken with hartshorn shavings, or arrow-root.

ENEMA TEREBINTHINÆ. Take of common turpentine, half an ounce; the yolk of one egg, and half a pound of gruel. The turpentine, being first incorporated with the egg, is to be added to the gruel. This clyster is generally used, and with great good effect, in violent fits of the stone.

ENERESIS. (From *ενεωδω*, to adhere to.) A compression. A tight ligature

ENERGY. (*Energia*; from *ενεργω*, to act.) Action. The degree of force exercised by any power: thus, nervous energy, muscular energy, &c.

ENEURESIS. See *Enuresis*.

ENGALACTUM. (From *ω*, and *γαλα*, milk; so called because it is eaten by nurses to increase their milk.) The herb saltwort.

ENCASTRIMYTUS. (From *ω*, in, and *κατρω*, the belly, and *μυθωμαι*, to discourse.) A

Ventriloquist; one who appears to speak from his belly.

ENGISOMA. (From *ἔγγειον*, to approach.) *Camarosia*. An instrument for making the parts of the broken clavicle meet. Also a fracture of the cranium.

English mercury. See *Mercurialis*.

ENGLOTTO-SASTOR. (From *εν*, *γλωττιν*, the tongue; and *σαστις*, the belly. A ventriloquist.

ENGOMPHOSIS. (From *εν*, and *γομφος*, a nail.) That species of articulation which resembles a nail, driven into wood, as a tooth in its socket.

ENGONTON. (From *εν*, and *γωνια*, an angle.) The flexure, or angle made by the bending of a joint.

ENIXUM PARACELSI. The caput mortuum of the spirit of nitre, joined with vitriolic acid.

ENNEAPHARMACUM. (From *εννα*, nine, and *φαρμακον*, a medicine. A medicine composed of nine simple ingredients.

ENNEAPHYLLUM. (From *εννα*, nine, and *φυλλον*, a leaf; because its flower consists of nine leaves.) A name for belleboraster, or bear's foot.

ENRITHMUS. (From *εν*, priv. and *ριθμος*, number.) An irregular pulse.

ENS MARRII. A name antiently given to the oxyde of iron, which arises in sublimation, with twice its quantity of sal-ammoniac. Medical practice does not at present place this practical preparation in a higher rank of estimation than other oxydes of iron.

ENS PRIMUM SOLARE. Antimony.

ENS VENERIS. The ens veneris is in many dispensatories called by this name.

ENSIFORM. (*Ensiformis*, sc. *cartilago*; from *ensis*, a sword, and *forma*, resemblance.) Sword-like. A term applied to a cartilage. See *Cartilago ensiformis*.

ENSTACTUM. (From *εν*, and *σταζειν*, to instil.) A liquid medicine, which is applied stillatim, or drop by drop.

ENTATICA. (From *εντασσειν*, to strain.) Provocatives: medicines which excite venereal inclination.

ENTERA. (From *εντος*, within. The bowels. Hippocrates calls by this name the bags in which were formerly enclosed medicines for fomentations.

ENTERADENES. (From *εντερον*, an intestine, and *αδενη*, gland.) The intestinal glands.

ENTERENCHTA. (From *εντερον*, the bowels, and *εγχυω*, to infuse into.) An instrument for administering clysters. A clyster-pipe.

ENTERITIS. (*Enteritis*: from *εντερον*, an intestine.) Inflammation of the intestines. It is a genus of disease in the class *pyrexia*, and order *phlegmasia* of Cullen, and is known by the presence of pyrexia, fixed pain in the abdomen, costiveness, and vomiting. The causes of enteritis are much

the same as those of gastritis, being occasioned by acrid substances, indurated faeces, long continued and obstinate costiveness, spasmodic cholic, and a strangulation of any part of the intestinal canal; but another very general cause is the application of cold to the lower extremities, or to the belly itself. It is a disease which is most apt to occur at an advanced period of life, and is very liable to a relapse.

It comes on with an acute pain, extending in general over the whole of the abdomen; but more especially round the navel, accompanied with eructations, sickness at the stomach, a vomiting of bilious matter, obstinate costiveness, thirst, heat, great anxiety, and a quick and hard small pulse. After a short time, the pain becomes more severe, the bowels seem drawn together by a kind of spasm, the whole region of the abdomen is highly painful to the touch, and seems drawn together in lumpy contractions; invincible costiveness prevails, and the urine is voided with great difficulty and pain.

The inflammation continuing to proceed with violence, terminates at last in gangrene; or abating gradually, it goes off by resolution.

Enteritis is always attended with considerable danger, as it often terminates in gangrene in the space of a few hours from its commencement; which event is marked by the sudden remission of pain, sinking of the pulse, shrinking of the features, and distention of the belly; and it frequently proves fatal likewise, during the inflammatory stage. If the pains abate gradually, if natural stools be passed, if an universal sweat, attended with a firm equal pulse, comes on, or if a copious discharge of loaded urine, with the same kind of pulse, takes place, a resolution and favourable termination may be expected.

Dissections of this disease shew that the inflammation pervades the intestinal tube to a very considerable extent; that adhesions of the diseased portion to contiguous parts are formed; and that, in some cases, the intestines are in a gangrenous state, or that ulcerations have formed. They likewise shew that, besides obstinate obstructions, intususception, constrictions, and twistings, are often to be met with; and that, in most cases, the peritoneum is more or less affected, and is perceived, at times, to be covered with a layer of coagulable lymph.

ENTEROCELE. (From *εντερον*, an intestine, and *κελη*, a tumour.) *Hernia intestinalis*. Every hernia may be so called that is produced by the protusion of an intestine, whether it is in the groin, navel, or elsewhere.

ENTERO-EPIPOCLE. (From *εντερον*, an intestine, *επιπλοον*, the epiploon, and *κελη*,

αλλη, a tumour.) A rupture formed by the protrusion of part of an intestine, with a portion of the epiploon.

ENTERO-HYDROCELE. (From *εντερον*, an intestine, *υδωρ*, water, and *αλλη*, a tumour.) This must mean a common scrotal hernia, with a good deal of water in the hernial sac; or else a hernia congenita, (in which the bowels descend into the tunica vaginalis testis,) attended with a collection of fluid in the cavity of this membrane.

ENTEROMPHALUS. (From *εντερον*, an intestine, and *ομφαλος*, the navel.) An umbilical hernia, produced by the protrusion of a portion of intestine.

ENTEROPHYTUM. (From *εντερον*, an intestine, and *φυλον*, a plant.) The sea-chitlerling; a plant which grows in the form of a gut.

ENTERORAPHIA. (From *εντερον*, an intestine, and *ραφω*, a suture.) A suture of the intestines, or the sewing together the divided edges of an intestine.

ENTEROSCHOCOLE. (From *εντερον*, an intestine, and *σχων*, the scrotum, and *αλλη*, a rupture.) *Hernia scrotalis*, or rupture of the intestines into the scrotum.

ENTHEMATA. (From *ενθηναι*, to put in.) Anti-inflammatory styptics.

ENTHELASIS. A contusion, with the impression of the instrument by which it happened.

ENTROPIUM. (From *εν*, and *τροπον*, to turn.) A disease of the eyelids, occasioned by the eyelashes and eyelid being inverted towards the bulb of the eye.

ENTYPOSIS. (From *εντυπον*, to make an impression) The acetabulum, or concave bone of the shoulder.

ENULA CAMPANA. (A corruption of *henula*, or *Helenium*, from *Helene*, the island where it grew.) *Helenium*. Common inula, or elecampane. *Enula helenium* of Linnaeus:—*foliis amplexicaulibus ovatis rugosis subtus tomentosis, calycum squamis ovatis*. This plant, though a native of Britain, is seldom met with in its wild state, but mostly cultivated. The root, which is the part employed medicinally, in its recent state, has a weaker and less grateful smell than when thoroughly dried; and kept for a length of time, by which it is greatly improved, its odour then approaching to that of Florentine orris. It was formerly in high estimation in dyspepsia, pulmonary affection, and uterine obstructions, but is now fallen into disuse.

ENULON. (From *εν*, and *υλος*, the gums.) The internal flesh of the gums, or that part of them which is within the mouth.

ENURESIS. (From *ενωρεω*, to make water.) An inconinency or involuntary flow of urine. This disease usually proceeds either from relaxation or a paralytic affection of the sphincter of the bladder,

induced by various debilitating causes, as too free a use of spirituous liquors, manutrition, and excess in venery; or it arises from compression on the bladder, from a diseased state of the organ, or from some irritating substance contained in its cavity. It is arranged in the class *locales*, and order *apoceneses* of Cullen, and contains two species:—1. *Enuresis atonica*, the sphincter of the bladder having lost its tone from some previous disease: 2. *Enuresis ab irritatione, vel compressione vesicae*, from an irritation or compression of the bladder.

EPACMASTICUS. (From *επι*, and *ακμαζω*, to increase.) It is applied to fever which is still increasing in malignity.

EPACME. (From *επακμαζω*, to increase.) The increase, or exacerbation of a disease.

EPAGOGIUM. (From *επαγω*, to draw over.) The præpuce, that part of the penis which is drawn over the glans, according to Dioscorides.

EPANADIDOTES. (From *επαναδιδωμι*, to increase.) A term applied to fevers which continue to increase in their degree of heat.

EPANADIPLOSIS. (From *επαναδιπλωω*, to reduplicate.) The reduplication of a fit of a semiterian fever; that is, the return of the cold fit before the hot fit is ended.

EPANASTASIS. (From *επι*, and *ανισταμι*, to excite.) A tubercle, or small pustule upon the skin.

EPANCYLOTUS. (From *επι*, and *αγκυλος*, crooked.) A sort of crooked bandage in Oribasius.

EPARMA. (From *επαρτω*, to elevate.) *Eparsis*. Any kind of tumour, but frequently applied to the parotis.

EPARSIS. See *Eparma*.

EPASMATICA FEBRIS. A fever is so called by Bellini, and others, while it is in its increase.

EPENCRAIUS. (From *επι*, *εν*, in, and *κρανιον*, the skull.) The name of the cerebellum.

EPHREBUM. (From *επι*, and *ηλη*, the groom.) The hair upon the pubes.

EPHEDRA. (From *εφεζομαι*, to sit upon.) *Ephedrana*. The buttocks. Also a species of horse-tail.

EPHEDRANA. See *Ephedra*.

EPHELICIS. (From *επι*, upon, and *υλος*, an ulcer.) The crust of an ulcer; hardened purulent expectoration.

EPHELIS. (From *επι*, and *ηλος*, the sun.) A broad, solitary, or aggregated spot, attacking most commonly the face, back of the hand, and breast, from exposure to the sun.

EPHEMERA. (From *επι*, upon, and *ημερα*, a day.) A fever which begins, is perfectly formed, and runs through its course, in the space of twelve hours.

EPHEMERIDES. (From *εφημερις*, an almanack; so called because, like the moon's age, they may be foretold by the almanack.) Diseases which return at particular times of the moon.

EPHIALTES. (From *επαιλλωμαι*, to leap upon; so called because it was thought a demon leaped upon the breast.) Incubus, or night-mare.

EPHIALTIA. (From *εφιάλτης*, the night-mare; so called because it was said to cure the night-mare.) The herb piony.

EPIDROSIS. (From *επιδρῶς*, to perspire.) *Sudatio.* *Mador.* A violent and morbid perspiration. A genus of disease in the class *locales*, and order *apoceneses* of Cullen.

EPHIPPIUM. (A saddle, which it is thought to resemble.) See *Sella turcica*.

EPHODOS. (From *επι*, and *οδος*, a way.) In Hippocrates it hath three significations: 1. The ducts, or passages, by which the excrements of the body are evacuated. 2. The periodical attack of a fever, from the common use of it to express the attack of thieves. 3. The access of similar or dissimilar things, which may be useful or hurtful to the body.

EPHIALTES. See *Ephialtes*.

EPHALUS. (From *επι*, gently, and *αλγος*, to heat.) *Epihalos.* An ardent fever, in which both heat and cold are felt in the same part at the same time. Galen defines it to be a fever in which the patient labours under a preternatural heat and a coldness at the same time. The ancient Latins call it *Quercera*.

EPIBOLE. (From *επιβαλλω*, to press upon.) The night-mare, or ephialtes.

EPICANTHIS. (From *επι*, and *καθος*, the angle of the eye.) The angle of the eye.

EPICARPIUM. (From *επι*, upon, and *καρπος*, the wrist.) A topical medicine applied to the wrist.

EPICAUMA. (From *επι*, and *καυω*, to burn.) *Encauma.* *Epicautis.* A burn, or scald.

EPICAUSIS. See *Epicauma*.

EPICERAS. (From *επι*, and *κερας*, a horn; so called because its pods are shaped like a horn.) The herb fænugreek.

EPICERASTICA. (From *επι*, and *καταμιμναι*, to mix.) Medicines which, by mixing with acrimonious juices, temper them and render them less troublesome. Emollients.

EPICHEIRESIS. (From *επι*, and *χειρ*, the hand.) A manual operation.

EPICHOLUS. (From *επι*, and *χολη*, the bile.) Bilious.

EPICHORDIS. (From *επι*, upon, and *χορδη*, a gut.) The mesentery.

EPICHRIOIS. (From *επι*, upon, and *χορδη*, a region.) The same as epidermis.

EPICHELIS. (From *επι*, upon, and *οφθαλμος*, the eyelid.) The upper eyelid.

EPICOLIC. (*Epicolica*; from *επι*, upon, and *κολον*, the colon.) Upon the colon. That

part of the abdomen which lies over the head of the cæcum and sigmoid flexure of the colon, called the epicolic region.

EPICOROSIS. (From *επι*, and *κορος*, deaf.) A total deafness.

EHICRANIUM. (From *επι*, and *κρανιον*, the cranium.) The common integuments, aponeurosis, and muscular expansion which lie upon the cranium.

EPICRANIUS. See *Occipito-frontalis*.

EPICRISIS. (From *επι*, and *κρίσις*, to temper.) A critical evacuation of bad humours, an attemperation of bad ones. When a cure is performed in the alterative way, it is called *per Epicrisin*.

EPICRISIS. (From *επι* *κρίσις*, to judge from.) A judgment of the termination of a disease from present symptoms.

EPICENTUM. (From *επι*, about, and *κεντρον*, the pubes.) The parts above and about the pubes.

EPICYMA. (From *επι*, upon, and *κυω*, to conceive.) *Epiyesis* Superfætation. Superimpregnation.

EPICYTIS. See *Epicyma*.

EPIDEMIC. (*Epidemicus*; from *επι*, upon, and *δημος*, the people.) A contagious disease is so termed, that attacks many people at the same season, and in the same place; thus putrid fever, plague, dysentery, &c. are often epidemic.

EPIDENDRUM. (From *επι*, upon, and *δενδρον*, a tree; because all this genus of plants grow parasitically on the trunk or branches of other trees.) The name of a genus of plants in the Linnæan system. Class, *Gynandria*. Order, *Monandria*.

EPIDENDRUM VANILLA. The systematic name of the vanelloe plant. See *Vanilla*.

EPIDERIS. (From *επι*, and *δερμα*, the skin.) The clitoris.

EPIDERMIS. (From *επι*, upon, and *δερμα*, the true skin.) The scarf-skin. See *Cuticle*.

EPIDESIS. (From *επι*, upon, and *δεω*, to bind.) A bandage to stop a discharge of blood.

EPIDESMUS. (From *επι*, upon, and *δεω*, to bind.) A bandage by which splints, bolsters, &c. are secured.

EPIDIDYMIS. (From *επι*, upon, and *ιδυμος*, a testicle.) A hard, vascular, oblong substance, that lies upon the testicle, formed of a convolution of the *vas deferens*. It has a thick end, which is convex, and situated posteriorly; and a thin end, which is rather flat, and situated inferiorly. The epididymis adheres to the testicle by its two extremities only. for its middle part is free, forming a bag, to which the tunica vaginalis of the testicle is attached.

EPIDONIS. (From *επιιδοναι*, to grow upon.) A preternatural enlargement of any part.

EPIDROME. (From *επιιδρωμα*, to run upon.) An flux of humours.

EPIGASTRIC. (*Epigastricus*, *επιγαστρικος*; from *επι*, upon, or above, and *γαστηρ*,

the stomach.) That part of the abdomen that lies over the stomach, is called the epigastric region; it reaches from the pit of the stomach to an imaginary line above the navel, supposed to be drawn from one extremity of the last of the false ribs to the other. Its sides are called hypochondria, and are covered by the false ribs, between which lies the epigastrium.

EPIGASTRIUM. (From *epi*, upon, or above, and *gaster*, the belly.) The region or part immediately over the stomach.

ΕΠΙΓΕΝΝΕΜΑ. (From *επιγενναι*, to generate upon.) *Epigenesis*. The fur on the tongue. An accessory symptom.

ΕΠΙΓΕΝΝΕΣΙΣ. See *Epigenemad*.

ΕΠΙΕΙΝΟΜΕΝΑ. (From *επινομιαι*, to succeed or supervene.) Galen says, they are those symptoms which naturally succeed; or may be expected in the progress of a disease; but Pæsius says, they are accessions of some other affection to diseases, which never happen but in stubborn and malignant diseases.

ΕΠΙΓΛΩΤΤΗΝ. (From *epi*, upon, and *γλωττις*, the tongue; so called because a lesser leaf grows above the larger in the shape of a tongue.) The Alexandrian laurel.

EPIGLOTTIS. (From *epi*, upon, and *γλωττις*, the tongue.) The cartilage at the root of the tongue that falls upon the glottis or superior opening of the larynx. Its figure is nearly oval; it is concave posteriorly, and convex anteriorly. Its apex or superior extremity is loose, and is always elevated upwards by its own elasticity. While the back of the tongue is drawn backwards in swallowing, the epiglottis is put over the aperture of the larynx, hence it shuts up the passage from the mouth into the larynx. The base of the epiglottis is fixed to the thyroid cartilage, the os hyoides, and the base of the tongue, by a strong ligament.

ΕΠΙΓΛΩΤΤΗΝ. (From *επιγλωττις*, the epiglottis, which it resembles in shape.) An instrument mentioned by Paracelsus for elevating the eyelids.

ΕΠΙΓΛΩΤΤΙΣ. (From *epi*, upon, and *γλωττις*, the buttocks.) The superior parts of the buttocks.

ΕΠΙΓΟΝΑΤΙΣ. (From *epi*, upon, and *γονυ*, the knee.) The patella or knee pan.

ΕΠΙΓΟΝΙΔΕΣ. (From *epi*, and *γονυ*, the knee.) The muscles inserted into the knees.

ΕΠΙΓΟΝΟΝ. (From *επινομιαι*, to proceed upon.) A superciliation.

ΕΠΙΛΕΨΙΑ. See *Epilepsia*.

EPILEPTIA. Corrupted from *epilepsia*.

EPILEPSY. (From *επιλαμβάνω*, to seize upon; so called, from the suddenness of its attack.) It is also called falling sickness, from the patient suddenly falling to the ground on an attack of this disease. By the ancients it was termed, from its affect-

ing the mind, the most noble part of the rational creature, the sacred disease. It consists of convulsions with sleep, and usually froth issuing from the mouth. It is a genus of disease in the class *neuroses*, and order *spasmi* of Cullen, and contains nine species: 1. *Epilepsia traumatica*, arising from an injury of the head: 2. *Epilepsia à dolore*, from pain: 3. *Epilepsia verminosa*, from the irritation of worms: 4. *Epilepsia à veneno*, from poisons: 5. *Epilepsia exanthematica*, from the repulsion of cutaneous eruptions: 6. *Epilepsia à cruditate ventriculi*, from crudities of the stomach: 7. *Epilepsia ab inanitione*, from debility: 8. *Epilepsia aërina*, from hysterical affections: 9. *Epilepsia ex onanismo*, from onanism.

Epilepsy attacks by fits, and after a certain duration goes off, leaving the person most commonly in his usual state; but sometimes a considerable degree of stupor and weakness remain behind, particularly where the disease has frequent recurrences. It is oftener met with among children than grown persons, and boys seem more subject to its attacks than girls. Its returns are periodical, and its paroxysms commence more frequently in the night than in the day, being somewhat connected with sleep. It is a disease sometimes counterfeited, in order to extort charity or excite compassion.

Epilepsy is properly distinguished into sympathetic and idiopathic, being considered as sympathetic, when produced by an affection in some other part of the body, such as acidities in the stomach, worms, teething, &c. an idiopathic when it is a primary disease, neither dependent on, nor proceeding from any other.

The causes which give rise to epilepsy, are blows, wounds, fractures, and other injuries done to the head by external violence, together with lodgments of water in the brain, tumours, concretions and polypi. Violent affections of the nervous system, sudden frights, fits of passion, great emotions of the mind, acute pains in any part, worms in the stomach or intestines, teething, the suppression of long accustomed evacuations, too great emptiness or repletion, and poisons received into the body, are causes which likewise produce epilepsy. Sometimes it is hereditary, and at others it depends on a predisposition arising from a mobility of the sensorium, which is occasioned either by plethora, or a state of debility.

An attack of epilepsy is now and then preceded by a heavy pain in the head, dimness of sight, noise in the ears, palpitations, flatulency in the stomach and intestines, weariness, and a small degree of stupor, and in some cases, there prevails a sense of something like a cold vapour or aura arising up to the head; but it more generally happens that the patient falls

down suddenly without much previous notice; his eyes are distorted or inverted, so as that only the whites of them can be seen; his fingers are closely clenched, and the trunk of his body, particularly on one side, is much agitated; he foams at the mouth, and thrusts out his tongue, which often suffers great injury, from the muscles of the lower jaw being affected; he loses all sense of feeling, and not unfrequently voids both urine and feces involuntarily.

The spasms abating he recovers gradually; but on coming to himself, feels very languid and exhausted, and retains not the smallest recollection of what has passed during the fit.

When the disease arises from an hereditary disposition, or comes on after the age of puberty, or where the fits recur frequently, and are of long duration, it will be very difficult to effect a cure; but when its attacks are at an early age, and is occasioned by worms, or any accidental cause, it may in general be removed with ease. In some cases, it has been entirely carried off by the occurrence of a fever, or by the appearance of a cutaneous eruption. It has been known to terminate in apoplexy, and in some instances to produce a loss of the powers of the mind, and to bring on idiotism.

The appearances usually to be observed on dissection are, serous and sanguineous effusion, a turgid tense state of the vessels of the brain without any effusion, a dilatation of some particular part of the brain, excrescences, polypi, and hydatides adhering to it, and obstructing its functions, and likewise ulcerations.

EPILOBIUM. (From *επι λοβιον*, a violet or beautiful flower, growing on a pod.) The name of a genus of plants in the Linnean system. Class *Octandria*. Order, *Monogynia*.

EPILOBIUM ANGUSTIFOLIUM. Rose-bay-willow herb. The young tender shoots cut in the spring, and dressed as asparagus, are little inferior.

EPIMEDIUM. The plant barren-wort.

EPIMORIUM. (From *επι*, and *μοριον*, to divide.) A term applied to an unequal pulse.

EPIMYLIS. (From *επι*, and *μυλα*, the knee.) The patella or knee bone.

EPINACUCUS. (From *επι νανος*, to nod or incline.) An unequal pulse.

EPINOTIUM. (From *επι*, upon, and *νωτος*, the back.) The shoulder blade.

EPINOTIS. (From *επι*, and *νυξ*, night.) A pustule, which rises in the night, forming an angry tumour on the skin of the arms, hands, and thighs, of the size of a lupine, of a dusky red, and sometimes of a livid and pale colour, with great inflammation and pain. In a few days breaks, and sloughs away.

EPIACTIS. (From *επι ακτις*, to coagulate.) A plant mentioned by Dioscorides; and so named because its juice was said to coagulate milk.

EPIPAROXYSMUS. (From *επι*, upon, and *παροξυσμος*, a paroxysm.) An unusual frequency of febrile exacerbation.

EPIFASTUM. (From *επι*, upon, and *πασα*, to sprinkle.) Any powdered drug sprinkled on the body.

EPIPECHYS. (From *επι*, above, and *πυξ*, the cubit.) That part of the arm above the cubit.

EPITHELOGISMA. (From *επι*, upon, and *φλεγω*, to inflame.) Violent inflammation, or burning heat in any part, attended with pain; tumour, and redness: also a name given by Hippocrates to the shingles.

EPIPHORA. (From *επι ροη*, to carry forcibly.) The watery eye. An involuntary flow of tears. A superabundant flowing of a serous or aqueous humour from the eyes. A genus of disease in the class *locales*, and order *apocremes*, of Cullen. The humour which flows very copiously from the eye in epiphora appears to be furnished, not only by the lachrymal gland, but from the whole surface of the conjunctive membrane, Meibomius's glands, and the caruncula lachrymalis; which increased and morbid secretion may be induced from any stimulus seated between the globe of the eye and lids, as sand, acrid fumes, and the like; or it may arise from the stimulus of active inflammation; or from the acrimony of scrophula, measles, small-pox; or from general relaxation. The disease may also arise from a more copious secretion of tears than the puncta lachrymalia can absorb, or, as it is most common, from an obstruction in the lachrymal canal, in consequence of which the tears are prevented from passing freely from the eye into the nose.

EPIPHYSIS. (From *επι*, upon, and *φυω*, to grow.) Any portion of bone growing upon another, but separated from it by a cartilage.

EPIPLAEMA. (From *επι*, upon, and *πλεσσω*, to spread.) A poultice; also a name for an application of wheat meal, boiled in hyalazum, to wounds.

EPIPLOCELE. (From *επι πλοον*, the omentum, and *κελη*, a tumour.) An ommental hernia. A rupture produced by the protusion of a portion of the omentum. See *Hernia*.

EPIPOLOMISTES. (From *επι πλοον*, the omentum, and *πομω*, to carry.) One who has the omentum morbidly large.

EPIPLOIC APPENDAGES. See *Appendicula epiploica*.

EPIPLOITIS. (From *επι πλοον*, the omentum.) An inflammation of the process of the peritoneum, that forms the epiploon or omentum. See *Peritonitis*.

EPIPOLOPEALON. (From *επι πλοον*, the

omentum, and *ομφαλος*, the navel.) An omental hernia protruding at the navel.

EPIPLOON. (From *επιπλοον*, to sail over, because it is mostly found floating, as it were, upon the intestines.) See *Omentum*.

EPIPLOSCHOECELE. (From *επιπλοον*, the omentum, *σχημα*, the scrotum, and *κηλη*, a tumour or hernia.) A rupture of the omentum into the scrotum, or a scrotal hernia containing omentum.

EPITOLASIS. (From *επιπλάζω*, to swim on the top.) A fluctuation of humours. A species of chymical sublimation.

ΕΠΙΤΟΜΑ. (From *επι*, upon, and *τομή*, a lid.) An instrument to cover the shoulder in a luxation.

ΕΠΙΤΟΜΑ. (From *επιπύρρειν*, to harden.) An indurated tumour in the joints; a callous concretion, a tophus, a tophaceous callus molesting the joints.

ΕΠΙΤΥΞΙΣ. (From *επιτίσσω*, to close up.) A spasmodic closing of the lips.

ΕΠΙΦΥΞΙΣ. (From *επι*, upon, and *πύρεξιν*, to be feverish.) A rapid exacerbation in a fever.

ΕΠΙΡΙΓΗΣΙΣ. (From *επι*, and *ρίζω*, to become cold.) An unusual degree of cold, or repetition of rigors.

ΕΠΙΡΡΗΟΞ. (From *επι*, upon, and *ρην*, to flow.) An influx or afflux of humours to any part.

ΕΠΙΣΑΝΚΙΔΙUM. (From *επι*, upon, and *σάρξ*, the flesh.) An anasarca, or dropsy spread between the skin and flesh.

ΕΠΙΣΧΕΣΕΣ. (From *επισχέω*, to restrain.) A suppression of excretions. It is an order in the class *locales* of Cullen's nosology.

ΕΠΙΣΧΙUM. (From *επι*, upon, and *ισχον*, the hip-bone.) The os pubis.

ΕΠΙΣΚΟΡΑΛΙΣ. (From *επισκοπος*, a bishop, or mitred dignitary.) Resembling a bishop's mitre. It is applied to a valve at the orifice between the left auricle and ventricle of the heart, called the mitral valve.

ΕΠΙΣΠΑΣΜΟΣ. (From *επισπασω*, to draw together.) A quick inspiration of the breath.

ΕΠΙΣΠΑΣΤΙΚΟΣ. (*Epispastica*, sc. *medicamenta*; from *επισπασω*, to draw together.) Those substances which are capable, when applied to the surface of the body, of producing a serous or puriform discharge, by exciting a previous state of inflammation. The term, though comprehending likewise issues and setons, is more commonly restricted to blisters—those applications which, exciting inflammation on the skin, occasion a thin serous fluid to be poured from the exhalants, raise the cuticle, and form the appearance of a vesicle. This effect arises from their strong stimulating power, and to this stimulant operation and the pain they excite, as to be ascribed the advantages derived from them in the

treatment of disease. The evacuation they occasion is too inconsiderable to have any effect.

It is a principle sufficiently established with regard to the living system, that where a morbid action exists, it may often be removed by inducing an action of a different kind in the same or neighbouring part. On this principle is explained the utility of blisters in local inflammation and spasmodic action, and it regulates their application in pneumonia, gastritis, hepatitis, phrenitis, angina, rheumatism, colic, and spasmodic affections of the stomach; diseases in which they are employed with the most marked advantage.

A similar principle exists with respect to pain; exciting one pain often relieves another. Hence blisters often give relief in tooth-ach, and some other painful affections.

Lastly, blisters, by their operation, communicate a stimulus to the whole system, and raise the vigour of the circulation. Hence, in part, their utility in fevers of the typhoid kind, though in such cases they are used with still more advantage to obviate or remove local inflammation.

ΕΠΙΣΦΗΡΙΑ. (From *επι*, and *σφαίρα*, a sphere; so called from the spherical shape of the brain.) The windings of the exterior surface of the brain; or the winding vessels upon it.

ΕΠΙΣΤΑΓΜΟΣ. (From *επι*, and *σταζω*, to trickle down.) A catarrh.

ΕΠΙΣΤΑΦΙΛΙΝΟΣ. (From *επι*, and *σταφυλη*, a parnip, from their resemblance to a carrot.) See *Urtica*.

ΕΠΙΣΤΑΞΙΣ. (From *επισταζω*, to distil from.) Bleeding at the nose, with pain, or fullness of the head. A genus of disease arranged by Cullen in the class *pyrexiae*, and order *hæmorrhagia*.

Persons of a sanguine and plethoric habit and not yet advanced to manhood, are very liable to be attacked with this complaint: females being much less subject to it than males, particularly after menstruation.

Epistaxis comes on at times without any previous warning; but at others, it is preceded by a pain and heaviness in the head, flushing in the face, heat and itching in the nostrils, a throbbing of the temporal arteries and a quickness of the pulse. In some instances a coldness of the feet, and shivering over the whole body, together with a costive belly, are observed to precede an attack of this hæmorrhage.

This complaint is to be considered as of little consequence; when occurring in young persons, being never attended with any danger; but when it arises in those who are advanced in life, flows profusely and returns frequently, it indicates too great fullness of the vessels of the head, and not

unfrequently precedes apoplexy, palsy, &c. and therefore, in such cases, is to be regarded as a dangerous disease. When this hemorrhage arises in any putrid disorder, it is to be considered as a fatal symptom.

EPISTHOTONOS. (From *επιστροφή*, for wards, and *τόνος*, to extend.) A spasmodic affection of muscles drawing the body forwards. See *Tetanus*.

EPISTOMION. (From *επι*, upon, and *στόμα*, a mouth.) A stopper for a bottle. Also a vent-hole of a furnace, called the register.

EPISTROPHALUS. (From *επι*, upon, and *στροφή*, to turn about.) It is applied to the first vertebra of the neck, because it turns about upon the second as upon an axis, which therefore was so called by the ancients. Some, though improperly, called the second thus. It is also written *Epistrophea* and *Epistrophis*.

EPISTROPHUS. (From *επιστροφή*, to invert.) An inversion of any part, as when the neck is turned round. Also a return of a disorder which has ceased.

EPISTROPHEUS. (From *επιστροφή*, to turn round, because the head is turned upon it.) *Epistropheus*. The second cervical vertebra. See *Dentatus*.

EPISTROPHIS. See *Epistrophe*.

EPITASIS. (From *επι*, and *τάσις*, to extend.) The beginning and increase of a paroxysm or disease.

EPITHELIUM. The cuticle on the red part of the lips.

EPITHEMA. (From *επι*, upon, and *θέμα*, to apply.) A term sometimes applied to a lotion, fomentation, or some external application.

EPITHEMATIUM. The same. A plaster.

EPITHESES. (From *επι*, and *θέσις*, to cover or lay upon.) The rectification of crooked limbs by means of instruments.

EPITHYMUM. (From *επι*, upon, and *θυμός*, the herb thyme.) *Cuscuta*. Dodder of thyme. A parasitical plant, possessing a strong disagreeable smell, and a pungent taste, very durable in the mouth. Two kinds are recommended in melancholia, as cathartics, viz. *Cuscuta epithymum* of Linnæus:—*foliis sessilibus, quinquefidis, bracteis obvallatis, and cuscuta europæa; floribus sessilibus*.

EPOTHE. (From *επι*, over, and *οθή*, a song.) *Epodos*. The method of curing distempers by incantation.

EPOMIS. (From *επι*, upon, and *ομος*, the shoulder.) The acromion, or upper part of the shoulder.

EPOMPHALIUM. (From *επι*, upon, and *ομφαλός*, the navel.) An application to the naval.

EPSOM SALT. A purging salt, formerly obtained by boiling down the mineral water, found in the vicinity of Epsom in Surrey. It is at present prepared from

sea water, which, after being boiled down, deposits an uncrystallized brine, that consists chiefly of muriated magnesia, and is sold in the shops under the name of sal. catharticus amarus, or bitter purging salt. See *Sulphas Magnese*.

EPSOM WATER. *Aqua Epsomensis*. This water evaporated to dryness leaves a residuum, the quantity of which has been estimated from an ounce and a half in the gallon to five drachms and one scruple. Of the total residuum, by far the greater part, about four or five sixths, is sulphate of magnesia mixed with a very few muriats, such as that of lime, and probably magnesia, which render it very deliquescent, and increase the bitterness of taste, till purified by repeated crystallizations. There is nothing sulphureous or metallic ever found in this spring. The diseases in which it is employed are similar to those of Sedlitz water. There are many other of the simple saline springs that might be enumerated, all of which agree with that of Epsom, in containing a notable proportion of some purging salt. This, for the most part, is either Epsom or Glauber's salt, or often a mixture of both, such as Acton, Kilburne, Bagnigge Wells, Dog and Duck, St. George's Fields, &c.

EPULIS. (From *επι*, and *υαλ*, the gums.) A small tubercle on the gums. It is said sometimes to become cancerous.

EPULOTICA. (*Epulotica*, sc. *medicamentosa*; from *επύλωσις*, to cicatrize.) A term given by surgeons to those applications which promote the formation of skin.

EQUISETUM. (From *equus*, a horse, and *seta*, a bristle: so named from its resemblance to a horse's tail.)

1. The name of a genus of plants in the Linnæan system. Class, *Cryptogamia*. Order, *Filices*.

2. The pharmacopœial name of the *Cauda equina*. Horse-tail, mare's-tail. The plant directed for medicinal purposes under this name is the *Hippuris vulgaris* of Linnæus. It possesses astringent qualities, and is frequently used by the common people as tea in diarrhœas and hæmorrhages. The same virtues are also attributed to the *Equisetum arvense, fluviatile, limosum*, and other species, which are directed indiscriminately by the term *Equisetum*.

EQUISETUM ARVENSE. See *Equisetum*.

EQUUS ASINUS. The systematic name of the animal called an ass; the female affords a light and nutritious milk.

ERANTHEMUS. (From *ερα*, the spring, and *ανθος*, a flower; so called because it flowers in the spring.)

EREBINTHUS. (*Erebthos*.) The vetch.

ERECTOR CLITORIDIS. First muscle of the clitoris of Douglas. *Ichio-cavernosus* of Winslow, and *Ichio-clitoridien* of Dumas. A muscle of the clitoris

that draws it downwards and backwards, and serves to make the body of the clitoris more tense, by squeezing the blood into it from its crus. It arises from the tuberosity of the ischium, and is inserted into the clitoris.

ERECTOR PENIS. *Iechio-cavernosus* of Winslow, and *Iechio-cavernosus* of Du-mas. A muscle of the penis that drives the urine or semen forwards, and, by grasping the bulb of the urethra, pushes the blood towards the corpus cavernosum and the glans, and thus distends them. It arises from the tuberosity of the ischium, and is inserted into the sides of the cavernous substance of the penis.

ERETHISMUS. (From *ἐρεθίζω*, to excite or irritate.) Increased sensibility and irritability. It is variously applied by modern writers.

Mr. Pearson has described a state of the constitution produced by mercury acting on it as a poison. He calls it the mercurial erethismus, and mentions that it is characterized, by great depression of strength, anxiety about the præcordia, irregular action of the heart, frequent sighing, trembling, a small, quick, sometimes intermitting pulse, occasional vomiting, a pale contracted countenance, a sense of coldness; but the tongue is seldom furred, nor are the vital and natural functions much disturbed. In this state any sudden exertion will sometimes prove fatal.

EROSTERIUM. (From *εἶρον*, work.) A laboratory: that part of the furnace in which is contained the matter to be acted upon.

ERICA. (From *εἶρα*, to break; so named from its fragility, or because it is broken into rods to make besoms of.) The name of a genus of plants in the Linnæan system. Class, *Octandria*. Order, *Monogymia*. Heath.

ERICERUM. (From *εἶρα*, heath.) A medicine in which heath is an ingredient.

ERIGERUM. (From *εἶρ*, the spring, and *γῆρας*, old, so called, because in the spring it has a white blossom like the hair of an old man.) *Erigeron*. Groundsel. This very common plant, *Senecio vulgaris* of Linnæus, is frequently applied bruised to inflammations and ulcers; as a refrigerent and antiscorbutic.

EROSION. (From *ερεδο*, to gnaw off.) *Erosio*. This word is very often used in the same sense as ulceration, viz. the formation of a breach or chasm in the substance of parts, by the action of the absorbents.

EROTOMANIA. (From *εἶρα*, love, and *μανία*, madness.) That melancholy, or madness, which is the effect of love.

ERPEX. (From *εἶρα*, to creep; so named from their gradually increasing in size.) The shingles, or herpes.

ERRATIC. (*Erraticus*, from *erre*, to wander.) Wandering. Irregular. A term occasionally applied to pains, or any disease which is not fixed, but moves from one part to another, as gout, rheumatism, &c.

ERRHINE. (*Errhina*, *ec. medicamenta*; *errhina*, from *ε*, in, and *ρῆ*, the nose.) By errhines are to be understood those medicines, which, when topically applied to the internal membrane of the nose, excite sneezing, and increase the secretion, independent of any mechanical irritation. The articles belonging to this class may be referred to two orders.

1. *Stimulatory errhines*, as *nicotiana*, *hol-luborus*, *euphorbium*, which are selected for the torpid, the vigorous, but not plethoric; and those to whom any degree of evacuation would not be hurtful.

2. *Evacuating errhines*, as *asarum*, &c. which is calculated for the phlegmatic and infirm.

ERROR LOCAL. Boerhaave is said to have introduced this term, from the opinion that the vessels were of different sizes, for the circulation of blood, lymph, and serum; and that when the larger sized globules were forced into the lesser vessels by an error of place, they were obstructed. But this opinion does not appear to be well grounded.

ERUCA. (From *erugo*, to make smooth, so named from the smoothness of its leaves; or from *erre*, to burn, because of its biting quality.) Garden rocket. Roman rocket. Rocket gentle. The seeds of this plant, *Brassica eruca*; *foliis hyaritis*, *caule hirsuto aliquis glabris*, of Linnæus, and of the wild rocket, have an acrid taste, and are eaten by the Italians in their pickles, &c. They are said to be good aperients and antiscorbutics, but are esteemed by the above-mentioned people for their supposed aphrodisiac qualities.

ERUCA SILVESTRIS. The wild rocket, *Brassica erucastrum* of Linnæus. See *Eruca*.

ERTHEMA. (From *ερθεω*, to make red.) A fiery red tumour, or pustules on the skin.

ERVUM. (*Quasi aruum*, a field, because it grows wild in the fields: or from *erre*, to pluck out, because it is diligently plucked from corn.)

1. The name of a genus of plants in the Linnæan system. Class, *Diadelphia*. Order, *Decandria*.

2. The pharmacopœial name of tare. *Orobis*. The plant ordered by this name is the *Ervum crœlia*; *germinibus undatoplicatis*, *foliis imparipinnatis*, of Linnæus. In times of scarcity the seeds have been made into bread, which is not the most salubrious. The meal was formerly amongst the resolvent remedies by way of poultice.

ERYON KUELLA. The systematic name of the Grobos. See *Eryon*.

ERYON LENS. The systematic name of the lentil. See *Lens*.

ERYNGIUM. (From *eryssa*, to eructate.)

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*.

2. The pharmacopœial name of the sea eryngo, or holly. *Eryngium maritimum*; *foliis radicalibus subrotundis plicatis spinosis, capitulis pedunculatis, paleis tricuspidatis*, of Linnæus. The root of this plant is directed for medicinal use. It has no particular smell, but to the taste it manifests a grateful sweetness; and, on being chewed for some time, it discovers a light aromatic warmth or pungency. It was formerly celebrated for its supposed aphrodisiac powers, but it is now very rarely employed.

ERYNGIUM CAMPESTRE. The root of this plant, *Eryngium campestre*; *foliis radicalibus, amplexicaulibus, pinnatis-lanceolatis*, of Linnæus, is used in many places for that of the sea eryngo. See *Eryngium*.

Eryng. See *Eryngium*.

Eryngo, sea. See *Eryngium*.

Eryngo leaved lichen. See *Lichen islandicus*.

ERYSIMUM. From *eruo*, to draw, so called from its power of drawing and producing blisters; others derive it *ero tu equum*, because the leaves are much cut; others from *eripio*, precious.)

1. The name of a genus of plants in the Linnæan system. Class, *Tetradynamia*. Order, *Silicquosa*.

2. The pharmacopœial name of the hedge mustard. This plant, *Erpsimum officinale*; *siliquis spica adpressa, foliis runcinatis*, of Linnæus, was formerly much used for its expectorant and diuretic qualities, which are now forgotten. The seeds are warm and pungent, and very similar to those of mustard in their sensible effects.

ERISIMUM ALLIARIA. The systematic name of Jack in the hedge. See *Alliaria*.

ERISIMUM BARBAREA. The systematic name of the barbarea of the shops. See *Barbarea*.

ERYSIMUM OFFICINALE. The systematic name of the hedge mustard. See *Erysimum*.

ERYSIPELAS. (From *eruo*, to draw, and *epi*, adjoining; named from the neighbouring parts being affected by the eruption.) *Ignis sacer*. The rose, or St. Anthony's fire. A genus of disease in the class *pyrexia*, and order *exanthemata* of Cullen. It is known by synocha of two or three days continuance, with drowsiness, and sometimes with delirium; pulse commonly full and hard; then erythema of the face, or some other part, with continuance

of synœchia, tending either to abscess or gangrene. There are two species of this disease, according to Cullen: 1. *Erysipelas vesiculosum*, with large blisters; 2. *Erysipelas phlyctenodes*, the shingles, or an erysipelas with phlyctenæ, or small blisters.

This disease is an inflammatory affection, principally of the skin, when it makes its appearance externally, and of the mucous membrane, when it is seated internally; and is more liable to attack women and children, and those of an irritable habit, than those of a plethoric and robust constitution.

It is remarkable that erysipelas sometimes returns periodically, attacking the patient once or twice in the year, or even once every month, and then by its repeated attacks it often gradually exhausts the strength, especially if he be old and of a bad habit.

When the inflammation is principally confined to the skin, and is unattended by any affection of the system, it is then called erythema; but when the system is affected it is named erysipelas.

Every part of the body is equally liable to it, but it more frequently appears on the face, legs, and feet, than any where else, when seated externally; and it occurs often in warm climates than phlegmonous inflammation.

It is brought on by all the causes that are apt to excite inflammation, such as injuries of all kinds, the external application of stimulants, exposure to cold, and obstructed perspiration; and it may likewise be occasioned by a certain matter generated within the body, and thrown out on its surface. A particular state of the atmosphere seems sometimes to render it epidemical, as we often find the scarlatina anginosa, which is a species of internal erysipelas, prevail as such.

In slight cases, where it attacks the extremities, it makes its appearance with a roughness, heat, pain, and redness of the skin, which becomes pale when the finger is pressed upon it, and again returns to its former colour, when it is removed. There prevails likewise a small febrile disposition, and the patient is rather hot and thirsty. If the attack is mild, these symptoms will continue only for a few days, the surface of the part affected will become yellow, the cuticle or scarf-skin will fall off in scales, and no further inconvenience will perhaps be experienced; but if the attack has been severe, and the inflammatory symptoms have run high, then there will ensue pains in the head and back, great heat, thirst, and restlessness; the part affected will slightly swell: the pulse will become small and frequent; and about the fourth day, a number of little vesicles, con-

taining a limpid, and, in some cases, a yellowish fluid, will arise. In some instances, the fluid is viscid, and instead of running out, as generally happens when the blister is broken, it adheres to and dries upon the skin.

In unfavourable cases, these blisters sometimes degenerate into obstinate ulcers, which now and then become gangrenous. This, however, does not happen frequently; for although it is not uncommon for the surface of the skin, and the blistered places to appear livid or even blackish, yet this usually disappears with the other symptoms of the complaint.

The period at which the vesicles shew themselves is very uncertain. The same may be said of the duration of the eruption. In mild cases it often disappears gradually, or is carried off by spontaneous sweating. In some cases it continues without shewing any disposition to decline for twelve or fourteen days, or longer.

The trunk of the body is sometimes attacked with erysipelatous inflammation, but less frequently so than the extremities. It is not uncommon, however, for infants to be attacked in this manner a few days after birth; and in these it makes its appearance about the genitals. The inflamed skin is hard, and apparently very painful to the touch. The belly often becomes uniformly tense, and sphacelated spots sometimes are to be observed. From dissections made by Dr. Underwood, it appears, that in this form of the disease, the inflammation frequently spreads to the abdominal viscera.

Another species of erysipelatous inflammation, which most usually attacks the trunk of the body, is that vulgarly known by the name of *shingles*, being a corruption of the French word *ceintle*, which implies a belt. Instead of appearing an uniform inflamed surface, it consists of a number of little pustules, extending round the body a little above the umbilicus, which have vesicles formed on them in a short time. Little or no danger ever attends this species of erysipelas.

When erysipelas attacks the face, it comes on with chilliness, succeeded by heat, restlessness, thirst, and other febrile symptoms, with a drowsiness or tendency to coma or delirium, and the pulse is very frequent and full. At the end of two or three days, a fiery redness appears on some part of the face, and this extends at length to the scalp, and then gradually down the neck, leaving a tumefaction in every part the redness has occupied. The whole face at length becomes turgid, and the eyelids are so much swelled as to deprive the patient of sight. When the redness and swelling have continued for some time, blisters of different sizes, containing a thin colourless

acid liquor, arise on different parts of the face, and the skin puts on a livid appearance in the blistered places: but in those not affected with blisters, the cuticle, towards the close of the disease, falls off in scales.

No remission of the fever takes place on the appearance of the inflammation on the face; but, on the contrary, it is increased as the latter extends, and both will continue probably for the space of eight or ten days. In the course of the inflammation, the disposition to coma and delirium are sometimes so increased as to destroy the patient between the seventh and eleventh days of the disease. When the complaint is mild, and not marked by a fatal event, the inflammation and fever generally cease gradually, without any evident crisis.

If the disease arises in a bad habit of body, occupies a part possessed of great sensibility, is accompanied with much inflammation, fever, and delirium, and these take place at an early period, we may suppose the patient exposed to imminent danger. Where translations of the morbid matter take place, and the inflammation falls on either the brain, lungs, or abdominal viscera, we may entertain the same unfavourable opinion. Erysipelas never terminates in suppuration, unless combined with a considerable degree of phlegmonous inflammation, which is, however, sometimes the case; but in a bad habit, it is apt to terminate in gangrene, in which case there will be also great danger. When the febrile symptoms are mild, and unaccompanied by delirium or coma, and the inflammation does not run high, we need not be apprehensive of danger.

Where the disease has occupied the face, and proves fatal, inflammation of the brain, and its consequences, are to be met with on dissection.

ERYTHEMA. (From *ερυθος*, red.) A morbid redness of the skin, as is observed upon the cheeks of hectic patients after eating, and the skin covering bubo, phlegmon, &c. Erysipelas is so called when the inflammation is principally confined to the skin.

ERYTHRODANUM. (From *ερυθος*, red, so called from the colour of its juice.) See *Rubia*.

ERYTHROIDES. (From *ερυθος*, red, and *ωδρ*, a likeness, so called from its colour.) A name given to the tunica vaginalis testis.

ERYTHRONIUM. (From *ερυθος*, red, so called from the red colour of its juice.) A species of *satyrium*.

ERYTHROXYLUM. (From *ερυθος*, red, and *ξυλον*, wood, so named from its colour.) Logwood.

ERYTHRUS. (From *ερυθος*, red, so named from the red colour of its juice.) The herb *her' sumach*.

ESAPHE. (From *ωσφαι*, to feel.) The touch; or feeling the mouth of the womb, to know its state.

ESCHAR. (*Eschara*, from *ωσχα*, to scab over.) *Eschara*. The portion of flesh that is destroyed by the application of a caustic.

ESCHAROTICS. (*Escharotica*, sc. *medicamenta*, *ωσχαλμα*, from *ωσχα*, to scab over.) Caustics. Corrosives. A term given by surgeons to those substances which possess a power of destroying the texture of the various solid parts of the animal body to which they are directly applied. The articles of this class of substances may be arranged under two orders.

1. *Eroding escharotics*, as blue vitriol, alumen ustum, &c.

2. *Caustic escharotics*, as *lapis infernalis*, *argentum nitratum*, *oleum vitrioli*, *acidum nitrosum*, &c.

ESCULENT. An appellation given to such plants, or any part of them, that may be eaten for food.

ESOX LUCIUS. The systematic name of fish of the class *pisces*, and order *abdominales*, from whose liver an oil spontaneously is separated, which is termed in some pharmacopœias *oleum lucii piscis*. It is used in some countries by surgeons, to destroy spots of the transparent cornea.

ESSENCE. Several of the volatile or essential oils are so called by the perfumers.

ESSENTIAL OIL. See *Oil*.

ESSERA. (Arab. A humour.) A species of cutaneous eruption, distinguished by broad, shining, smooth, red spots, mostly without fever, and differing from the nettle rash in not being elevated. It generally attacks the face and hands.

ESULA. (From *erus*, eating, because it is eaten by some as a medicine.) Spurge.

ESULA MAJOR. The officinal plant ordered by this name in some pharmacopœias is the *Euphorbia palustris* of Linnæus:—*umbella multifida, bifida, involucrellis ovatis, foliis lanceolatis, ramis sterilibus*. The juice is exhibited in Russia as a common purge; and the plant is given, in some places, in the cure of intermittents.

ESULA MINOR. *Tithymalis cyparissius*. Cypress spurge. This, like most of the spurges, is very acrimonious, inflaming the eyes and œsophagus after touching them. It is now fallen into disuse, whatever were its virtues formerly, which, no doubt, amongst some others, was that of opening the bowels, for, amongst rustics, it was called poor man's rhubarb.

ETHER. See *Æther*.

ETHER, ACETIC. Acetic naphtha. An ethereal fluid, drawn over from an equal admixture of alcohol and acetic acid, distilled with a gentle heat from a glass retort in a sand-bath. It has a grateful

smell, is extremely light, volatile, and inflammable.

ETHER, MURIATIC. Marine ether. Marine ether is obtained by mixing and distilling alcohol with extremely concentrated muriatic of tin. It is stimulant, antiseptic, and diuretic.

ETHER, NITROUS. Nitric naphtha. This is only a stronger preparation than the spiritus ætheris nitrosi of the London Pharmacopœia; it is produced by the distillation of two parts of alcohol to one part and an half of fuming nitric acid.

ETHER, SULPHURIC. See *Æther sulphuricus*.

ETHER, VITRIOLIC. See *Æther sulphuricus*.

ETHERIAL OIL. Any highly rectified essential oil may be so termed.

Ethiops antimonial. See *Sulphuretum hydrargyri stibiatum nigrum*.

Ethiops mineral. See *Hydrargyrum cum sulphure*.

Ethiops per se. See *Oxydum hydrargyri nigrum*.

ETHMOIDES. See *Ethmoid bone*.

ETHMOID BONE. (*Os ethmoides*; from *σμεος*, a sieve, and *ωδης*, form; because it is perforated like a sieve.) *Os ethmoidæum*. *Os ethmoides*. Cribriform bone. A bone of the head. This is, perhaps, one of the most curious bones of the human body. It appears almost a cube, not of solid bone, but exceedingly light, spongy, and consisting of many convoluted plates, which form a net-work, like honey-comb. It is curiously enclosed in the os frontis, betwixt the orbitary processes of that bone. One horizontal plate receives the olfactory nerves, which perforate that plate with such a number of small holes, that it resembles a sieve; whence the bone is named cribriform, or ethmoid bone. Other plates dropping perpendicularly from this one, receive the divided nerves, and gives them an opportunity of expanding into the organ of smelling; and these bones, upon which the olfactory nerves are spread out, are so much convoluted as to extend the surface of this sense very greatly, and are named spongy bones. Another flat plate lies in the orbit of the eye; and being very smooth, by the rolling of the eye, it is named the os planum, or smooth bone. So that the ethmoid bone supports the forepart of the brain, receives the olfactory nerves, forms the organ of smelling, and makes a chief part of the orbit of the eye; and the spongy bones, and the os planum, are neither of them distinct bodies, but parts of this ethmoid bone.

The *cribriform plate* is exceedingly delicate and thin; lies horizontally over the root of the nose; and fills up neatly the space betwixt the two orbitary plates of

the frontal bone. The olfactory nerves, like two small flat lobes, lie out upon this plate, and, adhering to it, shoot down like many roots through this bone, so as to perforate it with numerous small holes, as if it had been dotted with the point of a pin, or like a nutmeg-grater. This plate is horizontal; but its processes are perpendicular, one above, and three below.

1. The first perpendicular process is what is called *crista galli*, a small perpendicular projection, somewhat like a cock's comb, but exceedingly small, standing directly upwards from the middle of the cribriform plate, and dividing that plate into two; so that one olfactory nerve lies upon each side of the *crista galli*; and the root of the *falx*, or *septum*, betwixt the two hemispheres of the brain, begins from this process. The *foramen cæcum*, or blind hole of the frontal bone, is formed partly by the root of the *crista galli*, which is very smooth, and sometimes, it is said, hollow, or cellular.

2. Exactly opposite this, and in the same direction with it, i. e. perpendicular to the æthmoid plate, stands out the *nasal plate* of the æthmoid bone. It is sometimes called *azygous*, or single process of the æthmoid, and forms the beginning of that *septum*, or partition, which divides the two nostrils. This process is thin but firm, and composed of solid bone; it is commonly inclined a little to one side, so as to make the nostrils of unequal size. The *azygous* process is united with the *vomer*, which forms the chief part of the partition; so that the *septum*, or partition of the nose, consists of the *azygous* process of the æthmoid bone above, of the *vomer*, below, and of the cartilage in the fore or projecting part of the nose; but the cartilage rots away, so that whatever is seen of this *septum* in the skull, must be either of the æthmoid bone or *vomer*.

3. Upon either side of the *septum*, there hangs down a *spongy bone*, one hanging in each nostril. They are each rolled up like a scroll of parchment; they are very spongy; are covered with a delicate and sensible membrane; and when the olfactory nerves depart from the cribriform plate of the æthmoid bone, they attach themselves to the *septum*, and to these upper spongy bones, and expend upon them, so that the convolutions of these bones are of material use in expanding the organ of smelling, and detaining the odorous effluvia till the impression be perfect. Their convolutions are more numerous in the lower animals, in proportion as they need a more acute sense. They are named *spongy* or *turbinated bones*, from their convolutions resembling the many folds of a turban.

The spongy bones have a great many honeycomb-like cells connected with

them, which belong also to the organ of smell, and which are useful perhaps by detaining the effluvia of odorous bodies, and also by reverberating the voice. Thus, in a common cold, while the voice is hurt by an affection of these cells, the sense of smelling is almost lost.

4. The *orbital plate* of the æthmoid bone, is a large surface, consisting of a very firm plate of bone, of a regular square form; exceedingly smooth and polished; it forms a great part of the socket for the eye, lying on its inner side. When we see it in the detached bone, we know it to be just the flat side of the æthmoid bone; but while it is incased in the socket of the eye, we should believe it to be a small square bone; and from this, and from its smoothness, it has got the distinct name of *os planum*.

The cells of the æthmoid bone, which form so important a share of the organ of smell, are arranged in great numbers along the spongy bone. They are small neat cells, much like a honey-comb, and regularly arranged in two rows, parted from each other by a thin partition; so that the *os planum* seems to have one set of cells attached to it, while another regular set of cells belongs in like manner to the spongy bones. There are thus twelve in number, opening into each other, and into the nose.

These cells are frequently the seat of venereal ulcers; and the spongy bones are the surface where polypi often sprout up. And from the general connections and forms of the bone, we can easily understand how the venereal ulcer, when deep in the nose, having got to these cells, cannot be cured, but undermines all the face; how the venereal disease, having affected the nose, soon spreads to the eye, and how even the brain itself is not safe. We see the danger of a blow upon the nose, which, by a force upon the *septum*, or middle partition, may depress the delicate cribriform plate, so as to oppress the brain with all the effects of a fractured skull, and without any operation which can give relief. And we also see the danger of pulling away polypi, which are firmly attached to the upper spongy bone.

ETHOX. (From *etho*, to eat, as containing the receptacles of the food.) The hypogastrium.

EUANTHEMUM. (From *eu* well, and *anthem*, a flower; so named from the beauty of its flowers.) The chamomile.

EUAPHIUM. (From *eu*, well, and *aphi*, the touch; so called because its touch was supposed to give ease.) A medicine for the piles.

EUGENIA. The name of a genus of plants in the Linnæan system. Class, *Isocandria*. Order, *Monogynia*.

EUDIOMETER. An instrument by

which the quantity of oxygen and nitrogen in atmospherical air can be ascertained. They are all founded upon the principle of decomposing common air by means of a body which has a greater affinity with the oxygen. See *Eudiometry*.

EUDIOMETRY. The method of ascertaining the purity of atmospherical air.

No sooner was the composition of the atmosphere known, than it became an inquiry of importance to find out a method of ascertaining, with facility and precision, the relative quantity of oxygen gas contained in a given bulk of atmospherical air.

The instruments in which the oxygen gas of a determined quantity of air was ascertained, received the name of *Eudiometers*, because they were considered as measurers of the purity of air. They are, however, more properly called *Oximeters*.

The eudiometers proposed by different chymists, are the following:

1. *Priestley's Eudiometer.*

The first eudiometer was made in consequence of Dr. Priestley's discovery, that when nitrous gas is mixed with atmospherical air over water, the bulk of the mixture diminishes rapidly; in consequence of the combination of the gas with the oxygen of the air, and the absorption of the nitric acid thus formed by the water.

When nitrous gas is mixed with nitrogen gas, no diminution takes place: but when it is mixed with oxygen gas in proper proportions, the absorption is complete. Hence it is evident, that in all cases of a mixture of these two gases, the diminution will be proportional to the quantity of the oxygen. Of course it will indicate the proportion of oxygen in air; and by mixing it with different portions of air, it will indicate the different quantities of oxygen which they contain, provided the component parts of air be susceptible of variation.

Dr. Priestley's method was to mix together equal bulks of air and nitrous gas in a low jar, and then transfer the mixture into a narrow graduated glass tube about three feet long, in order to measure the diminution of bulk. He expressed this diminution by the number of hundredth parts remaining. Thus, suppose he had mixed together equal parts of nitrous gas and air, and that the sum total was 200 (or 2.00:) suppose the residuum when measured in the graduated tube to amount to 104 (or 1.04,) and of course that 96 parts of the whole had disappeared, he denoted the purity of the air thus tried by 104.

This method of analyzing air by means of nitrous gas is liable to many errors. For the water over which the experiment is made may contain more or less carbonic acid, atmospherical air, or other heterogeneous substance. The nitrous gas is not always of the same composition, and is partly absorbed by the nitrous acid which

is formed; the figure of the vessel, and many other circumstances are capable of occasioning considerable differences in the results.

Fontana, Cavendish, Ladrioni, Magellan, Von Humbolt, and Dr. Falconer, have made series of laborious experiments to bring the test of nitrous gas to a state of complete accuracy; but, notwithstanding the exertions of these philosophers, the methods of analyzing air by means of nitrous gas are liable to so many anomalies, that it is unnecessary to give a particular description of the different instruments invented by them.

2. *Scheele's Eudiometer.*

This is merely a graduated glass cylinder, containing a given quantity of air, exposed to a mixture of iron filings and sulphur, formed into a paste with water. The substances may be made use of in the following manner:

Make a quantity of sulphur in powder, and iron filings, into a paste with water, and place the mixture in a saucer, or plate, over water, on a stand raised above the fluid; then invert over it a graduated bell-glass, and allow this to stand for a day or two. The air contained in the bell-glass will gradually diminish, as will appear from the ascent of the water.

When no further diminution takes place, the vessel containing the sulphuret must be removed, and the remaining air will be found to be nitrogen gas, which was contained in that quantity of atmospherical air.

In this process the moistened sulphuret of iron has a great affinity to oxygen, it attracts and separates it from the atmospherical air, and the nitrogen gas is left behind; the sulphur, during the experiment, is converted into sulphuric acid, which unites to the alkali and forms sulphate of potash.

The air which is exposed to moistened iron and sulphur, gradually becomes diminished, on account of its oxygen combining with a portion of the sulphur which becomes converted into sulphuric acid, and its nitrogen remains behind. The quantity of oxygen contained in the air examined becomes thus obvious, by the diminution of bulk, which the volume of air submitted to examination, has undergone.

The only error to which this method is liable is, that the sulphuric acid which is formed acts on the iron and produces hydrogen gas, which joins to the nitrogen remaining after the absorption, and occasions an incorrect result; and hence it is that the absorption amounts in general to 0.27 parts, although the true quantity of oxygen is no more than from 0.21 to 0.23.

3. *De Marti's Eudiometer.*

De Marti obviated the errors to which the method of Scheele was liable. He

availed himself. for that purpose, of a hydrogenated sulphuret, formed by boiling sulphur and liquid potash, or lime-water, together. These substances, when newly prepared, have the property of absorbing a minute portion of nitrogen gas; but they lose this property when saturated with that gas, which is easily effected by agitating them for a few minutes in contact with a small portion of atmospheric air.

The apparatus is merely a glass tube, ten inches long, and rather less than half an inch in diameter, open at one end, and hermetically sealed at the other. The close end is divided into one hundred equal parts, having an interval of one line between each division. The use of this tube is to measure the portion of air to be employed in the experiment. The tube is filled with water; and by allowing the water to run out gradually, while the tube is inverted, and the open end kept shut with the finger, the graduated part is exactly filled with air. These hundred parts of air are introduced into a glass bottle, filled with liquid sulphuret of lime previously saturated with nitrogen gas, and capable of holding from two to four times the bulk of the air introduced. The bottle is then to be closed with a ground glass stopper, and agitated for five minutes. After this, the stopper is to be withdrawn, while the mouth of the phial is under water; and for the greater accuracy, it may be closed and agitated again. Lastly, the air is to be again transferred to the graduated glass tube, in order to ascertain the diminution of its bulk.

4. *Humbolt's Eudiometer.*

Consists in decomposing a definite quantity of atmospheric air, by means of the combustion of phosphorus, after which, the portion of gas which remains must be measured.

Take a glass cylinder, closed at the top, and whose capacity must be measured into sufficiently small portions by a graduated scale fixed on it. If the instrument be destined solely for examining atmospheric air, it will be sufficient to apply the scale from the orifice of the cylinder down to about half its length, or to sketch that scale on a slip of paper pasted on the outside of the tube, and to varnish it over with a transparent varnish.

This half of the eudiometrical tube is divided into fifty equidistant parts, which in this case indicate hundredth parts of the whole capacity of the instrument.

Into this vessel, full of atmospheric air, put a piece of dry phosphorus, (one grain to every twelve cubic inches,) close it airtight, and heat it gradually, first the sides near the bottom, and afterwards the bottom itself. The phosphorus will take fire and burn rapidly. After every thing is cold, invert the mouth of the eudiometer

tube into a basin of water, or mercury, and withdraw the cork. The water will ascend in proportion to the loss of oxygen gas the air has sustained, and thus its quantity may be ascertained.

Analogous to this is

5. *Seguin's Eudiometer.*

Which consists of a glass tube, of about one inch in diameter, and eight or ten inches high, closed at the upper extremity. It is filled with mercury, and kept inverted in this fluid in the mercurial trough. A small bit of phosphorus is introduced into it, which on account of its specific gravity being less than that of mercury, will rise up in it to the top. The phosphorus is then melted by means of a red-hot poker, or burning coal applied to the outside of the tube. When the phosphorus is liquified, small portions of air destined to be examined, and which have been previously measured in a vessel graduated to the cubic inch, or into grains, are introduced into the tube. As soon as the air which is sent up reaches the phosphorus, a combustion will take place and the mercury will rise again. The combustion continues till the end of the operation; but, for the greater exactness, Mr. Seguin directs the residuum to be heated strongly. When cold it is introduced into a small vessel, whose capacity has been ascertained at the same time as that of the preceding. The difference of the two volumes gives the quantity of the oxygen gas contained in the air subjected to examination.

6. *Bertholet's Eudiometer.*

Instead of the rapid combustion of phosphorus, Bertholet has substituted its spontaneous combustion, which absorbs the oxygen of atmospheric air completely; and, when the quantity of air operated on is small, the process is accomplished in a short time.

Bertholet's apparatus consists of a narrow graduated glass tube, containing the air to be examined, into which is introduced a cylinder, or stick of phosphorus, supported upon a glass rod, while the tube stands inverted in water. The phosphorus should be nearly as long as the tube. Immediately after the introduction of the phosphorus, white vapours are formed which fill the tube; these vapours gradually descend, and become absorbed by the water. When no more white vapours appear, the process is at an end, for all the oxygen gas which was present in the confined quantity of air, has united with the phosphorus; the residuum is the quantity of nitrogen of the air submitted to examination.

This eudiometer, though excellent of the kind, is nevertheless not absolutely to be depended upon; for, as soon as the absorption of oxygen is completed, the nitrogen gas exercises an action upon the phosphorus, and thus its bulk becomes

increased. It has been ascertained, that the volume of nitrogen gas is increased to 1.40th part; consequently the bulk of the residuum, diminished by 1.40, gives us the bulk of the nitrogen gas of the air examined; which bulk, subtracted from the original mass of air, gives us the proportion of oxygen gas contained in it. The same allowance must be made in the eudiometer of Seguin.

7. Davy's Eudiometer.

Until very lately, the preceding processes were the methods of determining the relative proportions of the two gases which compose our atmosphere.

Some of these methods though very ingenious, are so extremely slow in their action, that it is difficult to ascertain the precise time at which the operation ceases. Others have frequently involved inaccuracies, not easily removed.

The eudiometer of Mr. Davy is not only free from these objections, but the result it offers is always constant; it requires little address, and is very expeditious; the apparatus is portable, simple and convenient.

Take a small glass tube, graduated into one hundred equi-distant parts; fill this tube with the air to be examined, and plunge it into a bottle, or any other convenient vessel, containing a concentrated solution of green muriate or sulphate of iron, strongly impregnated with nitrous gas. All that is necessary to be done is, to move the tube in the solution a little backwards and forwards; under these circumstances, the oxygen gas contained in the air will be rapidly absorbed, and condensed by the nitrous gas in the solution, in the form of nitrous acid.

N. B. The state of the greatest absorption must be marked, as the mixture afterwards emits a little gas which would alter the result.

This circumstance depends upon the slow decomposition of the nitrous acid (formed during the experiment,) by the green oxyd of iron, and the consequent production of a small quantity of æriform fluid (chiefly nitrous gas); which, having no affinity with the red muriate, or sulphate of iron, produced by the combination of oxygen, is gradually evolved and mingled with the residual nitrogen gas.

The impregnated solution with green muriate, is more rapid in its operation than the solution with green sulphate. In cases when these salts cannot be obtained in a state of absolute purity, the common sulphate of iron of commerce may be employed. One cubic inch of moderately strong impregnated solution, is capable of absorbing five or six cubic inches of oxygen, in common processes; but the same quantity must never be employed for more than one experiment.

In all these different methods of analyzing air, it is necessary to operate on air of a determinate density, and to take care that the residuum be neither more condensed nor dilated than the air was when first operated on. If these things are not attended to, no dependance whatever can be placed upon the result of the experiments; how carefully soever they may have been performed. It is, therefore, necessary to place the air before and after the examination, into water of the same temperature. If this, and several other little circumstances have been attended to, for instance, a change in the height of the barometer, &c. we find that air is composed of 0.22 of oxygen gas, and 0.78 of nitrogen gas by bulk. But as the weight of these two gases is not exactly the same, the proportion of the component parts by weight will differ a little; for as the specific gravity of oxygen gas is to that of nitrogen gas as 134:115, it follows that 100 parts of air are composed by weight of about 74 nitrogen gas, and 26 oxygen gas.

The air of this metropolis, examined by means of Davy's eudiometer, was found, in all the different seasons of the year to contain 0.21 of oxygen; and the same was the case with air taken at Islington and Highgate; in the solitary cells in Cold-bath-fields prison, and on the river Thames. But the quantity of water contained in a given bulk of air from these places, differed considerably.

EUGENIA. (So named by Micheli, in compliment to Prince Eugene of Savoy, who sent him from Germany almost all the plants described by Clusius.) The name of a genus of plants in the Linnæan system. Class, *Icosandria*. Order, *Monogynia*.

EUGENIA JAMBOS. The systematic name of the Malabar plum-tree. See *Malabar plum*.

EUEKUS. (From *eu*, well, and *kus*, the earth; so called, because of its fertility.) The uterus.

EULE. (From *εὐλαζω*, to putrefy.) A worm bred in foul and putrid ulcers.

EUNUCHIUM. (From *εὐνοχος*, an eunuch; so called because it was formerly said to render those who eat it impotent, like an eunuch.) The lettuce.

EUPATORIOPHALACRON. (From *εὐρατοριον*, agrimony, and *φαλακρον*, bald.) A species of agrimony with naked h ads.

EUPATORIUM. (From *Eupator*, its inventor; or *quasi hepatorium*, from *επα*, the liver; because it was said to be useful in diseases of the liver.) 1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia equalis*.

2. The pharmacopœial name of *Eupatorium Arabicum*. Hemp agrimony. This very bitter and strong-smelling plant, is the *Eupatorium cannabinum*; *foliis digitatis*, of

Linnaeus. Its juice proves violently emetic and purgative, if taken in sufficient quantity, and promotes the secretions generally. It is recommended in dropsies, jaundices, agues, &c. and is in common use in Holland, amongst the lower orders, as a purifier of the blood in old ulcers, scurvy, and anaesca.

EUPATORIUM ARABICUM. See *Eupatorium*.

EUPATORIUM CANNABINUM. The systematic name of the hemp agrimony. See *Eupatorium*.

EUPATORIUM MESUES. See *Ageratum*.

EUPESIA. (From *eu*, well, and *πείω*, to concoct.) A good digestion.

EUPETIC. (*Eueptica*; from *eu*, good, and *πείω*, to digest.) Substances are so called that are easy to digest.

EUPHORBIA ANTHORUM. The Linnaean name of a plant supposed to produce the *Euphorbium*.

EUPHORBIA CANARIENSIS. In the Canary islands this species of spurge affords the gum euphorbium.

EUPHORBIA CYPARISSIAS. The systematic name of the cypress spurge. See *Enula minor*.

EUPHORBIA LATHYRIS. The systematic name of the plant, which affords the lesser cataputia seeds. See *Cataputia minor*.

EUPHORBIA OFFICINARUM. The systematic name of the plant which affords the euphorbium in the greatest abundance. See *Euphorbium*.

EUPHORBIA PALUSTRIS. The systematic name of the greater spurge. See *Enula major*.

EUPHORBIA PARALIAS. See *Tithymalus paralias*.

EUPHORBIA. The name of a genus of plants in the Linnaean system. Class, *Dodecandria*. Order, *Trigynia*.

EUPHORBUM. (From *Euphorbus*, the physician of king Juba, in honour of whom it was named.) An inodorous gum-resin, in yellow tears, which have the appearance of being worm-eaten; said to be obtained from several species of *Euphorbia*, but principally from the *Euphorbia officinarum*; *aculeata nuda multangularis*, *aculeis germinatis*, of Linnaeus; it is imported from Ethiopia, Libya, and Mauritania. It contains an active resin, and is very seldom employed internally, but, as an ingredient, it enters into many resolvent and discutient plasters.

EUPHRASIA. (Corrupted from *Euphrosyne*, *υπερσυν*, from *ευφρον*, joyful; so called because it exhilarates the spirits.)

1. The name of a genus of plants in the Linnaean system. Class, *Didynamia*. Order, *Angiospermia*.

2. The pharmacopoeial name of eye-bright. This beautiful little plant, *Euphrasia officinalis*; *foliis ovatis, lineatis, urgute dentatis*, of Linnaeus, has been greatly es-

teemed by the common people, as a remedy for all diseases of the eyes; yet, notwithstanding this, and the encomiums of some medical writers, is now wholly fallen into disuse. It is an ingredient in the British herb-tobacco.

EUPHRASIA OFFICINALIS. The systematic name of the eye-bright. See *Euphrasia*.

EUSTACHIAN TUBE. *Tuba Eustachiana*. The tube so called was discovered by the great Eustachius. It begins, one in each ear, from the anterior extremity of the tympanum, and runs forwards and inwards in a bony canal, which terminates with the petrous portion of the temporal bone. It then goes on, partly cartilaginous, and partly membranous, gradually becoming larger, and at length ends behind the soft palate. Through this tube the air passes to the tympanum.

Eustachian valve. See *Valvula Eustachii*.

EUTHYFORIA. (From *ευθυ*, straight, and *πείω*, to pass into.) *Euthyporos*: An extension made in a straight line, to put in place a fracture, or dislocation.

EVAPORATION. The volatilization of a fluid, by means of heat, with access of air, in order to diminish its fluidity, to obtain any fixed salts it may hold in solution, or to diminish the quantity of a residuum. In this manner sea water is evaporated, and the salt obtained, and decoctions made into extracts.

EVERICULUM. (From *everro*, to sweep away.) A sort of spoon, used to clear the bladder from gravel.

EXACERBATION. (From *exacerbesco*, to become violent.) An increase of febrile symptoms.

EXAREISIS. (From *εξαίρω*, to remove.) One of the divisions of surgery, adopted by the old surgeons; the term implies the removal of parts.

EXALMA. (From *εξαλλομαι*, to leap out.) Hippocrates applies it to the starting of the vertebrae out of their places.

EXAMBLOMA. (From *εξμβλομα*, to miscarry.) An abortion.

EXAMBLOSIS. An abortion.

EXANASTOMOSIS. (From *εξαναστομω*, to relax or open.) The opening of the mouths of vessels, to discharge their contents; also the meeting of the extremities of the veins and arteries.

EXANTHEMA. (From *εξανθηω*, to spring forth, to bud.) *Exanthema*. An eruption of the skin. Cullen makes *exanthemata* an order in the class *pyrexia*. It includes diseases, beginning with fever, and followed by an eruption on the skin.

EXANTHEMATA. (The plural of *exanthema*.) The name of an order of diseases in Cullen's Nosology, which embraces eruptive fevers.

EXAN-ISMIA. See *Exanthema*.

EXANTHROPIA. (From *εξ*, without, and

αυδωμος, a man, i. e. having lost the faculties of a man.) A species of melancholy, where the patient fancies himself some kind of brute.

EXARAGMA. (From *εξαγασθαι*, to break.) A fracture.

EXARMA. (From *εξασθαι*, to lift up.) A tumour; a swelling.

EXARTEMA. (From *εξασθαι*, to suspend.) An amulet, or charm, hung round the neck.

EXARTHREMA. (From *εξαρθρεω*, to put out of joint.) *Exarthroma*. *Exarthrosis*. A dislocation, or luxation.

EXARTHROMA. See *Exarthrema*.

EXARTHROSIS. See *Exarthrema*.

EXARTICULATIO. (From *ex*, out of, and *articulus*, a joint.) A luxation. The dislocation of a bone from its socket.

EXCIPULUM. (From *excipio*, to receive.) A chymical receiver.

EXCITABILITY. See *Excitement*.

EXCITEMENT. A term introduced into medicine by Dr. Browne. Animals, differ from themselves in their dead state, or from any other inanimate matter, in this property alone; they can be affected by external agents, as well as by certain functions peculiar to themselves in such a manner, that the phenomena peculiar to the living state can be produced. This proposition extends to every thing that is vital in nature, and therefore applies to vegetables.

The external agents are reducible to heat, diet, and other substances taken into the stomach, blood, the fluids secreted from the body and air.

The functions of the system itself, producing the same effect, are muscular contraction, sense, or perception, and the energy of the brain in thinking, and in exciting passion and emotion. These affect the system in the same manner as the other agents; and they arise both from the other and from themselves.

If the property which distinguishes living from dead matter, or the operation of either of the two sets of powers be withdrawn, life ceases. Nothing else than the presence of these is necessary to life.

The property on which both sets of powers act Dr. Browne names *Excitability*, and the powers themselves exciting powers. The word body, means both the body simply so called, and also as endued with an intellectual part, a part appropriated to passion and emotion, or a soul: the usual appellation in medical writings is system.

The effects common to all the exciting powers, are sense, motion, mental exertion, and passion. Their effects being the same, it must be granted, that the operation of all their powers is the same.

The effects of the exciting powers acting upon the excitability, Dr. Browne, denominates *excitement*.

EXCITING CAUSE. Occasional cause.

Procataretic cause. Remote cause. That which when applied to the body, excites a disease. The exciting, or remote causes of diseases, are either external or internal.

EXCORIATION. (From *excorio*, to take off the skin.) *Excoriatio*. An abrasion of the skin.

EXCREMENT. (From *excerno*, to separate from.) The alvine fæces.

EXCRESCENCE. (From *exresco*, to grow from.) *Excrecentia*. Any preternatural formation of flesh, on any part of the body, as wens, warts, &c.

EXCRETION. (From *excerno*, to separate from.) *Excretio*. This term is applied to the separation or secretion of those fluids from the blood of an animal, that are supposed to be useless, as the urine, perspiration, and alvine fæz.

EXFOLIATION. (From *exfolio*, to cast the leaf.) *Exfoliatio*. The separation of a dead piece of bone from the living.

EXFOLIATIVUM. (From *exfolio*, to shed the leaf.) A raspatory or instrument for scraping exfoliating portions of bone.

EXISCHIOS. (From *εξ*, out of, and *ισχιον*, the ischium.) A luxation of the thigh-bone.

EXITURA. (From *exeo*, to come from.) A running abscess.

EXITUS. (From *exeo*, to come out.) A prolapsus, or falling down of the womb or anus.

EXOCHAS. (From *εξ*, without, and *εχσθαι*, to have.) *Exoche*. A tubercle on the outside of the anus.

EXOCHÆ. See *Exochas*.

EXOCYSTÆ. See *Exocystis*.

EXOCYSTIS. (From *εξ*, without, and *κυστις*, the bladder.) *Exocyste*. A prolapsus of the inner membrane of the bladder.

EXOMPHALUS. (From *εξ*, out, and *ομφαλος*, the navel.) *Exomphalos*. An umbilical hernia. See *Hernia*.

EXONCHOMA. (From *εξ*, and *ονχος*, a tumour.) A large prominent tumour.

EXOPHTHALMIA. (From *εξ*, out, and *οφθαλμος*, the eye.) A swelling or protrusion of the bulb of the eye, to such a degree that the eyelids cannot cover it. It may be caused by inflammation, when it is termed *exophthalmia inflammatoria*; or from a collection of pus in the globe of the eye, when it is termed the *exophthalmia purulenta*; or from a congestion of blood within the globe of the eye, *exophthalmia sanguinea*.

EXOSTOSIS. (From *εξ*, and *οστος*, a bone.) *Hyperostosis*. A morbid enlargement, or hard tumour of a bone. A genus of disease arranged by Cullen in the class *locales*, and order *tumores*. The bones most frequently affected with exostosis, are those of the cranium, the lower jaw, sternum, humerus, radius, ulna, bones of the carpus, the femur, and tibia. There is, however, no bone of the body which may

not become the seat of this disease. It is not uncommon to find the bones of the cranium affected with exostosis, in their whole extent. The ossa parietalia sometimes become an inch thick.

The exostosis, however, mostly rises from the surface of the bone, in the form of a hard round tumour, and venereal exostoses, or nodes, are observed to arise chiefly on compact bones; such of these as are only superficially covered with soft parts, as, for instance, the bones of the cranium, and the front surface of the tibia.

EXPECTORANTS. (*Expectorantia*, sc. *medicamenta*, from *expecto*, to discharge from the breast.) Those medicines which increase the discharge of mucus from the lungs. The different articles referred to this class may be divided into the following orders: 1. *Nauseating expectorants*, as squills ammoniacum, and garlic, which are to be preferred for the aged and phlegmatic. 2. *Stimulating expectorants*, as marrubium, which is adapted to the young and irritable, and those easily affected by expectorants. 3. *Antispasmodic expectorants*, as vesicatories, pediluvium, and watery vapours; these are best calculated for the plethoric and irritable, and those liable to spasmodic affections. 4. *Irritating expectorants*, as fumes of tobacco and acid vapours. The constitutions to which these are chiefly adapted, are those past the period of youth, and those in whom there are evident marks of torpor, either in the system generally, or in the lungs in particular.

EXPIRATION. (From *expiro*, to breathe.) *Expiratio*. That part of respiration in which the air is thrust out from the lungs. See *Respiration*.

EXPRESSED OILS. Such oils as are obtained by pressing the substance containing them, as olives, which give out olive oil, almonds, &c.

EXSUCCATIO. (From *ex*, out of, and *succus*, humour.) An ecchymosis, or extravasation of humours, under the integuments.

EXTENSOR. (From *extendo*, to stretch out.) A term given to those muscles whose office it is to extend any part; the term is in opposition to flexor.

EXTENSOR BREVIS DIGITORUM PEDIS. *Extensor brevis* of Douglas. *Calcano phalangium commune* of Dumas. A muscle of the toes situated on the foot. It arises fleshy and tendinous from the fore and upper part of the os calcis, and soon forms a fleshy belly, divisible into four portions, which send off an equal number of tendons that pass over the upper part of the foot under the tendons of the extensor longus digitorum pedis, to be inserted into its tendinous expansion. Its office is to extend the toes.

EXTENSOR CARPI RADIALIS BRE-

VIOR. *Radialis externus brevior* of Albinus. *Radialis secundus* of Winslow. An extensor muscle of the wrist, situated on the fore-arm. It arises tendinous from the external condyle of the humerus, and from the ligament that connects the radius to it, and runs along the outside of the radius. It is inserted by a long tendon into the upper and back part of the metacarpal bone of the middle finger. It assists in extending and bringing the hand backward.

EXTENSOR CARPI RADIALIS LONGIOR. *Radialis externus longior* of Albinus. *Radialis externus primus* of Winslow. An extensor-muscle of the carpus, situated on the fore-arm, that acts in conjunction with the former. It arises thin, broad, and fleshy, from the lower part of the external ridge of the os humeri, above its external condyle, and is inserted by a round tendon into the posterior and upper part of the metacarpal bone that sustains the fore-fingers.

EXTENSOR CARPI ULNARIS. *Ulnaris externus* of Albinus and Winslow. It arises from the outer condyle of the os humeri, and then receives an origin from the edge of the ulna: its tendon passes in a groove behind the styloid process of the ulna to be inserted into the inside of the basis of the metacarpal bone of the little finger.

EXTENSOR DIGITORUM COMMUNIS. *Cum extensore proprio auricularis* of Albinus. *Extensor digitorum communis manus* of Douglas and Winslow. *Extensor digitorum communis*, seu *digitorum tensor* of Cowper, and *Epichondylo-rusphalangettien commune* of Dumas. A muscle situated on the fore-arm, that extends all the joints of the fingers. It arises from the external protuberance of the humerus: and at the wrist it divides into three flat tendons, which pass under the annular ligament, to be inserted into all the bones of the fore, middle, and ring fingers.

EXTENSOR DIGITORUM LONGUS. See *Extensor longus digitorum pedis*.

EXTENSOR INDICIS. See *Indicator*.

EXTENSOR LONGUS DIGITORUM PEDIS. *Extensor longus peroneo tibialis phalangitien commune* of Dumas. A muscle situated on the leg, that extends all the joints of the four small toes. It arises from the upper part of the tibia and fibula, and the interosseous ligament; its tendon passes under the annular ligament, and then divides into five, four of which are inserted into the second and third phalanges of the toes, and the fifth goes to the basis of the metatarsal bone. This last Winslow reckons a distinct muscle, and calls it *Peroneus brevis*.

EXTENSOR LONGUS POLLICIS PEDIS. See *Extensor proprius pollicis pedis*.

EXTENSOR MAGNUS. See *Gastrocnemius internus*.

EXTENSOR MAJOR POLLCIS MANUS. See *Extensor secundi internodii*.

EXTENSOR MINOR POLLCIS MANUS. See *Extensor primi internodii*.

EXTENSOR OSSIS METACARP POLLCIS MANUS. *Abductor longus pollicis manus* of Albinus. *Extensor primi internodii* of Douglas. *Extensor primus pollicis* of Winslow. *Extensor primi internodii pollicis* of Cowper. *Cubito-radialis metacarpien du pouce* of Dumas. It arises fleshy from the middle and posterior part of the ulna, from the posterior part of the middle of the radius, and from the interosseous ligament, and is inserted into the os trapezium, and upper part of the metacarpal bone of the thumb.

EXTENSOR POLLCIS PRIMUS. See *Extensor primi internodii*.

EXTENSOR POLLCIS SECUNDUS. See *Extensor secundi internodii*.

EXTENSOR PRIMI INTERNODII. *Extensor minor pollicis manus* of Albinus. This muscle, and the *Extensor ossis metacarpi pollicis manus*, are called *Extensor pollicis primus* by Winslow. *Extensor secundi internodii* by Douglas. *Extensor secundi internodii ossis pollicis* of Cowper. *Cubito-susphalangien du pouce* of Dumas. A muscle of the thumb, situated on the hand, that extends the first bone of the thumb obliquely outwards. It arises fleshy from the posterior part of the ulnar, and from the interosseous ligament, and is inserted tendinous into the posterior part of the first bone of the thumb.

EXTENSOR PROPRIUS POLLCIS PEDIS. *Extensor longus* of Douglas. *Extensor pollicis longus* of Winslow and Cowper. *Peroneo susphalangien du pouce* of Dumas. An exterior muscle of the great toe, situated on the foot. It arises by an acute, tendinous and fleshy beginning, some way below the head and anterior part of the fibula, along which it runs to near its lower extremity, connected to it by a number of fleshy fibres, which descend obliquely, and form a tendon, which is inserted into the posterior part of the first and last joint of the great toe.

EXTENSOR SECUNDI INTERNODII. *Extensor majus pollicis manus* of Albinus. *Extensor pollicis secundus* of Winslow. *Extensor tertii internodii* of Douglas. *Extensor internodii ossis pollicis* of Cowper. *Cubito-susphalangietien du pouce* of Dumas. A muscle of the thumb, situated on the hand, that extends the last joint of the thumb obliquely backwards. It arises tendinous and fleshy from the middle part of the ulna, and the interosseous ligament; it then forms a tendon, which runs through a small groove at the inner and back part of the radius, to be inserted into the last bone of the thumb. Its use is to extend the last phalanx of the thumb obliquely backwards.

EXTENSOR SECUNDI INTERNODII INDICIS PROPRIUS. See *Indicator*.

EXTENSOR TARSII MINOR. See *Plantaris*.

EXTENSOR TARSII SUBALIS. See *Gastrocnemius internus*.

EXTENSOR TERTII INTERNODII MINIMI DIGITI. See *Abductor minimi digiti manus*.

EXTENSOR TERTII INTERNODII INDICIS. See *Prior indicis*.

EXTENSUS MALLEI. See *Laxator tympani*.

EXTIRPATION. (From *extirpo*, to eradicate.) *Extirpatio*. The complete removal or destruction of any part, either by cutting instruments, or the action of caustics.

EXTRACTION. (From *extraho*, to draw out.) *Extractio*. The taking extraneous substances out of the body. Thus bullets and splinters are said to be *extracted* from wounds; stones from the urethra, or bladder.

Surgeons also sometimes apply the term *extraction* to the removal of tumours out of cavities, as, for instance, to the taking of cartilaginous tumours out of the joints. They seldom speak of extracting any diseased original part of the body; though they do so in one example, viz. the cataract.

EXTRACT. (*Extractum*, from *extraho*, to draw out.) The generic term *extract* is used pharmaceutically, in an extensive sense, and includes all those preparations from vegetables which are separable by the agency of various liquids, and afterwards obtained from such solutions, in a solid state, by evaporation of the menstruum. It also includes those substances which are held in solution by the natural juices of fresh plants, as well as those to which some menstruum is added at the time of preparation. Now, such soluble matters are various, and mostly complicated; so that chymical accuracy is not to be looked for in the application of the term. Chymists, however, have affixed this name to one peculiar modification of vegetable matter, which has been called *extractive*, or *extract*, or *extractive principle*; and, as this forms one constituent part of common extracts, and possesses certain characters, it will be proper to mention such of them as may influence its pharmaceutical relations. The extractive principle has a strong taste, differing in different plants: it is soluble in water, and its solution speedily runs into a state of putrefaction, by which it is destroyed. Repeated evaporations and solutions render it at last insoluble, in consequence of its combination with oxygen from the atmosphere. It is soluble in alcohol, but insoluble in ether. It unites with alumine, and if boiled with neutral salts thereof, preci-

pitates them. It precipitates with strong acids, and with the oxyds from solutions of most metallic salts, especially muriate of tin. It readily unites with alkalies, and forms compounds with them, which are soluble in water. No part, however, of this subject has been hitherto sufficiently examined.

In the preparation of all the extracts, the London Pharmacopœia requires that the water be evaporated as speedily as possible, in a broad, shallow dish, by means of a water-bath, until they have acquired a consistence proper for making pills; and, towards the end of the inspissation, that they should be constantly stirred with a wooden rod. These general rules require minute and accurate attention, more particularly in the immediate evaporation of the solution, whether prepared by expression or decoction, in the manner as well as the degree of heat by which it is performed, and the promotion of it by changing the surface by constant stirring, when the liquor begins to thicken, and even by directing a strong current of air over its surface, if it can conveniently be done. It is impossible to regulate the temperature over a naked fire, or if it be used, to prevent the extract from burning; the use of a water-bath is, therefore, absolutely necessary, and not to be dispensed with, and the beauty and precision of extracts so prepared, will demonstrate their superiority.

EXTRACTIVE. See *Extract*.

EXTRACTUM ACONITI. Extract of aconite. "Take of aconite leaves, fresh, a pound; bruise them in a stone mortar, sprinkling on a little water; then press out the juice, and, without any separation of the sediment, evaporate it to a proper consistence." The dose is from one grain to five grains. For its virtues, see *Aconitum*.

EXTRACTUM ALOES. Extract of aloes. "Take of extract of spike aloes, powdered, half a pound; boiling water, four pints." Macerate for three days in a gentle heat, then strain the solution, and set it by, that the dregs may subside. Pour off the clear solution, and evaporate it to a proper consistence. The dose, from v to xv grs. See *Aloes*.

EXTRACTUM ANTHEMIDIS. Extract of chamomile, formerly called *extractum chamœmeli*. "Take of chamomile flowers, dried, a pound. Water, a gallon." Boil down to four pints, and strain the solution while it is hot, then evaporate it to a proper consistence. The dose is x grs. to a scruple. For its virtues, see *Chamamelum*.

EXTRACTUM BELLADONNÆ. Extract of belladonna. "Take of deadly night-shade leaves, fresh, a pound." Bruise them in a stone mortar, sprinkling on a little water; then press out the juice, and without any previous separation of the sediment, evapo-

rate it to a proper consistence. The dose is from one to five grains. For its virtues, see *Belladonna*.

EXTRACTUM CINCHONÆ RESINOSUM. Resinous extract of bark. "Take of lance-leaved cinchona bark, a pound; rectified spirit, four pints." Macerate for four days, and strain. Distil the tincture in the heat of a water-bath, until the extract has acquired a proper consistence. This is considered by many as much more grateful to the stomach, and, at the same time, producing all the effects of bark in substance, and, by the distillation of it, is intended that the spirit which passes over shall be collected and preserved. The dose is from ten grains to half a drachm. See *Cinchona*.

EXTRACTUM COLOCYNTHIDIS. Extract of colocynth. "Take of colocynth pulp, a pound; water, a gallon." Boil down to four pints, and strain the solution while it is hot, and evaporate it to a proper consistence. The dose is from five to thirty grains. For its virtues, see *Colocynthis*.

EXTRACTUM COLOCYNTHIDIS COMPOSITUM. Compound extract of colocynth. "Take of colocynth pulp, sliced, six drachms; extract of spike aloes, powdered, an ounce and half; scammony gum-resin powdered, half an ounce; cardamom-seeds powdered, a drachm; hard soap, three drachms; boiling water, two pints." Macerate the colocynth pulp in the water, for four days, in a gentle heat; strain the solution, and add to it the aloes, scammony, and soap; then, by means of a water-bath, evaporate it to a proper consistence, constantly stirring, and about the end of the inspissation, mix in the cardamom-seeds. The dose, from five to thirty grains.

EXTRACTUM CONII. Extract of hemlock, formerly called *succus cicutæ spissatus*. "Take of fresh hemlock, a pound." Bruise it in a stone mortar, sprinkling on a little water; then press out the juice, and, without any separation of the sediment, evaporate it to a proper consistence. The dose from five grains to a scruple.

EXTRACTUM ELATERII. Extract of elaterium. "Cut the ripe, wild cucumbers into slices, and pass the juice, very gently expressed, through a very fine hair sieve, into a glass vessel; then set it by for some hours, until the thicker part has subsided. Pour off, and throw away the thinner part, which swims at the top. Dry the thicker part which remains in a gentle heat." The dose from half a grain to three grains. For its virtues, see *Cucumis agrestis*.

EXTRACTUM GENTIANÆ. Extract of gentian. "Take of gentian root, sliced, a pound; boiling water, a gallon." Macerate for twenty-four hours, then boil down to four pints: strain the hot liquor, and evaporate it to a proper consistence. Dose, from ten to thirty grains. See *Gentiana*.

EXTRACTUM GLYCYRRHIZÆ. Extract of

liquorice. "Take of liquorice-root, sliced, a pound; water, boiling, a gallon." Macerate for twenty-four hours, then boil down to four pints; strain the hot liquor, and evaporate it to a proper consistence. Dose, from one drachm to half an ounce. See *Glycyrrhiza*.

EXTRACTUM HEMATOXYLI. Extract of logwood, formerly called *extractum ligni campechensis*. "Take of logwood, powdered, a pound; water, boiling, a gallon." Macerate for twenty-four hours, then boil down to four pints; strain the hot liquor, and evaporate it to a proper consistence. Dose, from ten grains to half a drachm. For its virtues, see *Lignum campechense*.

EXTRACTUM HUMULI. Extract of hops. "Take of hops, half a pound; water, boiling, a gallon." Boil down to four pints; strain the hot liquor, and evaporate it to a proper consistence. This extract is said to produce a tonic and sedative power combined. The dose is from five grains to one scruple. See *Lupulus*.

EXTRACTUM HYOSCYAMI. Extract of henbane. "Take of fresh henbane leaves, a pound." Bruise them in a stone mortar, sprinkling on a little water; then press out the juice, and, without separating the feculencies, evaporate it to a proper consistence. Dose, from five to thirty grains. For its virtues, see *Hyosciamus*.

EXTRACTUM JALAPÆ. Extract of jalap. "Take of jalap-root, powdered, a pound; rectified spirit, four pints; water, ten pints." Macerate the jalap-root in the spirit for four days, and pour off the tincture; boil the remaining powder in water, until it be reduced to two pints; then strain the tincture and decoction separately, and let the former be distilled and the latter evaporated, until each begins to grow thick. Lastly, mix the extract with the resin, and reduce it to a proper consistence. Let this extract be kept in a soft state, fit for forming pills, and in a hard one, so that it may be reduced to powder. The dose, from ten to twenty grains. For its virtues, see *Jalapium*.

EXTRACTUM OPII. Extract of opium, formerly called *extractum thebaicum*. *Opium colatum*. "Take of opium, sliced, half a pound; water, three pints." Pour a small quantity of the water upon the opium, and macerate it for twelve hours, that it may become soft; then, adding the remaining water gradually, rub them together until the mixture be complete. Set it by, that the feculencies may subside; then strain the liquor, and evaporate it to a proper consistence. Dose, from half a grain to five grains.

EXTRACTUM PAPAVERIS. Extract of white poppy. "Take of white poppy capsules, bruised, a pound; water, boiling, a gallon." Macerate for twenty-four hours,

then boil down to four pints; strain the hot liquor, and evaporate it to a proper consistence. Six grains are about equivalent to one of opium. Dose, from half a grain to five grains. For its virtues, see *Papaver album*.

EXTRACTUM RHEI. Extract of rhubarb. "Take of rhubarb root, powdered, a pound; proof spirit, a pint; water, seven pints." Macerate for four days in a gentle heat, then strain, and set it by, that the feculencies may subside. Pour off the clear liquor, and evaporate to a proper consistence. This extract possesses the purgative properties of the root, and the fibrous and earthy parts are separated; it is, therefore a useful basis for pills, as well as given separately. Dose, from ten to thirty grains. See *Rhubarbarum*.

EXTRACTUM SARSAPARILLÆ. Extract of sarsaparilla. "Take of sarsaparilla-root, sliced, a pound; water, boiling, a gallon." Macerate for twenty-four hours, then boil down to four pints; strain the hot liquor, and evaporate it to a proper consistence. In practice this is much used, to render the common decoction of the same root stronger and more efficacious. Dose, from ten grains to a drachm. For its virtues, see *Sarsaparilla*.

EXTRACTUM SATURNI. See *Liquor acetatis plumbi*.

EXTRACTUM TARAXACI. Take of dandelion root, fresh and bruised, a pound; water, boiling, a gallon. Macerate for twenty-four hours; boil down to four pints, and strain the hot liquor through a woollen cloth; then evaporate it to a proper consistence. Dose, from ten grains to a drachm. For its virtues, see *Taraxacum*.

EXTRAVASATION. (From *extra*, without, and *vas*, a vessel.) *Extravasatio*. A term applied by surgeons to fluids, which are out of their proper vessels or receptacles. Thus, when blood is effused on the surface, or in the ventricles of the brain, it is said that there is an extravasation. When blood is poured from the vessels into the cavity of the peritoneum, in wounds of the abdomen, surgeons call this accident *extravasation*. The urine is also said to be *extravasated*, when, in consequence of a wound, or of sloughing, or ulceration, it makes its way into the cellular substance, or among the abdominal viscera. When the bile spreads among the convolutions of the bowels, in wounds of the gall-bladder, it is also a species of extravasation.

EYE. *Oculus*. The parts which constitute the eye are divided into external and internal. The external parts are—1. The *eyebrows*, or *supercilia*, which form arches of hair above the orbit, at the lower part of the forehead. Their use is to prevent the sweat falling into the eyes, and for moderating the light above.

2. The eyelashes, or *cilia*, are the short hairs that grow on the margin of the eyelids; they keep external bodies out of the eyes, and moderate the influx of light.

3. The eyelids, or *palpebre*, of which, one is superior or upper, and the other inferior or under; where they join outwardly, it is called the *external canthus*; inwardly, towards the nose, the *internal canthus*; they cover and defend the eyes.

The margin of the eyelids, which is cartilaginous, is called *tarsus*.

In the *tarsus*, and internal surface of the eyelids, small glands are situated, called *glandule Meibomiane*, because Meibomius discovered them; they secrete the oily mucilaginous fluid, which prevents the attrition of the eyes and eyelids, and facilitates their motions.

4. The lachrymal glands, or *glandule lachrymales*, which are placed in the external canthus, or corner of the eyes, in a little fovea of the os frontis.

From these glands six or more canals issue, which are called lachrymal ducts, or *ductus lachrymales*, and they open in the internal superficies of the upper eyelids.

5. The lachrymal caruncle, or *carunculus lachrymalis*, which is situated in the internal angle, or canthus of the eyelids.

6. *Puncta lachrymalia*, are two callous orifices or openings, which open in the internal angle of the tarsus of the eyelids; the one in the superior, the other in the inferior eyelid.

7. The lachrymal ducts, or *canales lachrymales*, are two small canals, which proceed from the lachrymal points into the lachrymal sac.

8. The lachrymal sac, or *saccus lachrymalis*, is a membranous sac, which is situated in the internal canthus of the eye.

9. The nasal duct, or *ductus nasalis*, is a membranous canal, which goes from the inferior part of the lachrymal sac through the bony canal below, and a little behind, into the cavity of the nose, and opens under the inferior spongy bone into the nostrils.

10. The conjunctive membrane, or *membrana conjunctiva*, which, from its white colour, is called *albuginea*, or white of the eye, is a membrane which lines the internal superficies of the eyelids, and covers the whole fore-part of the globe of the eye; it is very vascular, as may be seen in inflammations.

The bulb or globe of the eye is composed of eight membranes, or coverings, two chambers, or *camere*, and three humours, improperly so called.

The membranes of the globe of the eye are, four in the hinder or posterior part of the bulb, or globe, viz. *sclerotica*, *choroidea*, *retina*, and *hyaloidea*, or *arachnoidea*; four in the fore or anterior part of the bulb, viz.

cornea *transparens*, *iris*, *uvea*, and *capsula* of the crystalline lens.

The *membrana sclerotica*, or the sclerotic or horny membrane, which is the outermost, begins from the optic nerve, forms the spherical or globular cavity, and terminates in the circular margin of the transparent cornea.

The *membrana choroidea*, or *choroidea*, is the middle tunic of the bulb, of a black colour, beginning from the optic nerve, and covering the internal superficies of the sclerotic, to the margin of the transparent cornea. In this place it secedes from the tornea, and defects transversely and inwardly, and in the middle forms a round perforated foramen. This circular membrane of the choroidea in the anterior surface is called *iris*, in the posterior superficies, *uvea*.

The round opening in the centre is called the *pupil*, or *pupilla*. This foramen, or round opening, can be dilated, or contracted, by the moving powers of almost invisible muscular fibres.

The *membrana retina*, is the innermost tunic, of a white colour, and similar to mucus, being an expansion of the optic nerve, chiefly composed of its medullary part. It covers the inward superficies of the choroidea, to the margin of the chrySTALLINE lens, and there terminates.

The *chambers*, or *camere*, of the eyes are:

1. *Camera anterior*, or fore-chamber; is an open space, which is formed anteriorly, by the hollow superficies of the *cornea* *transparens*, and posteriorly, by the superficies of the *iris*.

2. *Camera posterior*, is that small space which remains anteriorly from the *tunica uvea*, and *pupilla*, or pupil; posteriorly from the anterior superficies of the chrySTALLINE lens.

Both these chambers are filled with aqueous humour. The humours of the eye, as they are called, are in number three:

1. The *aqueous humour*, which fills both chambers.

2. The *crystalline lens*, or humour, is a pellucid body, about the size of a lentil, which is included in an exceeding fine membrane, or *capsula*, and lodged in a concave fovea of the vitreous humour.

3. The *vitreous humour* is a pellucid, beautifully transparent substance, which fills the whole bulb of the eye behind, the chrySTALLINE lens. Its external superficies is surrounded with a most pellucid membrane, which is called *membrana hyaloidea*, or *arachnoidea*. In the anterior part is a fovea, or bed, for the chrySTALLINE lens, in which the lens is seated.

The connection of the bulb is made anteriorly, by means of the conjunctive membrane, with the inner surface of the eyelids, or *palpebra*; posteriorly, by the

adhesion of six muscles of the bulb and optic nerve, with the orbit.

The optic nerve, or *nervus opticus*, perforates the sclerotica and choroides, and then constitutes the retina, by spreading itself on the whole posterior superficies of the internal globe of the eye.

The muscles by which the eye is moved in the orbit, are six; much adesa surrounds them, and fills up the cavities in which the eyes are seated. The arteries are the internal orbital, the central, and the ciliary arteries. The veins empty themselves into the external jugulars. The nerves are the optic, and branches from the third, fourth, fifth, and sixth pair.

The use of the eye is to form the organ of vision.

Externally, the globe of the eye and the transparent cornea, are moistened with a most limpid fluid, called *lachryma*, or tears; the same pellucid subtle fluid exactly fills all the pores of the transparent cornea; for, deprived of this fluid, and being exposed to the air, that coat of the eye becomes dry, shrivelled, and cloudy, impeding the rays of light.

Eye-bright. See *Euphrasia*.

EYE-BROW. *Supercilium.* A layer of short hair, which lies thick upon the low part of the frontal bone, or the superior prominent part of the orbit.

EYE-LID. *Palpebra.* The semi-lunar moveable production of the skin which covers the eye when shut. It is distinguished into upper and under eye-lid.

F.

F. or **ft.** In a prescription these letters are abbreviations of *fiat*, or *fiant*, let it or them be made; thus *f. bolus*, let the substance or substances prescribed be made into a bolus.

FABA. See *Bean*.

FABA CRASSA. *Telephium.* *Fabaria crassula.* *Anacampteros.* The plant which bears these names in various pharmacopœias, is the orphine, *Sedum telephium* of Linnaeus: —*foliis planiusculis serratis, corymbo foliose, caule erecto.* It was formerly ranked as an antiphlogistic, but now forgotten.

FABA ÆGYPTIACA. *Cyamos Ægyptiacus.* *Nymphaea Indica.* The pontic, or Egyptian bean. The fruit of the *nymphaea nelumbo* of Linnaeus, which grows on marshy grounds in Egypt, and some of the neighbouring countries. It is eaten either raw or boiled, and is a tonic and astringent.

FABA FEBRIFUGA. See *Faba Indica*.

FABA INDICA. *Faba sancti ignatii.* *Faba febrifuga.* The seeds of a gourd-like fruit, the produce of the *Ignatia amara* of Linnaeus. They are of a roundish figure, very irregular and uneven, about the size of a middling nutmeg, semi-transparent, and of a hard, horny texture. They have a very bitter taste, and no considerable smell. They are said to be used in the Philippine islands in all diseases, acting as a vomit and purgative. Infusions are given in the cure of intermittents, &c.

FABA MAJOR. The Turkey or garden bean. See *Bean*.

FABA MINOR. *Equina frasa.* The horse-bean. It differs no otherwise from the garden bean than in being less.

FABA PECHURIM. *Faba pichurim.* *Faba pechuris.* An oblong oval, brown, and

ponderous seed supposed to be the produce of a *Laurus*, brought from the Brazils. Their smell is like that of musk, between it and the scent of sassafras. They are exhibited as carminatives in flatulent colics, diarrhœas, and dysenteries.

FABA PURGATRIZ. See *Ricinus*.

FABA SANCTI IGNATII. See *Faba indica*.

FABA SUILLA. See *Hyosciamus*.

FABARIA. (From *faba*, a bean, which it resembles.) Orphine. See *Faba crassa*.

FACE. The lower and anterior part of the cranium, or skull.

FACIAL. Belonging to the face; as facial nerve, &c.

FACIAL NERVE. *Nervus facialis.* *Portio dura* of the auditory nerve. These nerves are two in number, and are properly the eighth pair: but are commonly called the seventh, being reckoned with the auditory, which is the portio mollis of the seventh pair. They arise from the fourth ventricle of the brain, pass through the petrous portion of the temporal bone to the face, where they form the *pes anserinus*, which supplies the integuments of the face and forehead.

FACIES HIPPOCRATICA. That particular disposition of the features which immediately precedes the stroke of death is so called, because it has been so admirably described by Hippocrates.

FACIES RUBRA. See *Gutta serena*.

FACTITIOUS. A term applied to any thing which is made by art, in opposition to that which is native, or found already made in nature.

FACULTY. *Facultas.* The power or ability by which any action is performed.

FÆCES. The plural of *fec*. The alvine excretions.

FÆCULA. (Diminutive of *fax*.) A substance obtained by bruising or grinding certain vegetables in water. It is that part which, after a little, falls to the bottom. The fæcula of plants appears to be only a slight alteration of mucilage, for it differs from mucilage in no other respect than being insoluble in cold water, in which it falls with wonderful quickness. There are few plants which do not contain fæcula; but the seeds of gramineous and leguminous vegetables, and all tuberos roots contain it most plentifully.

FÆX. The alvine excretions are called *fæces*.

FAGARA. The name of a genus of plants in the Linnæan system. Class, *Tetrandria*. Order, *Monogymia*.

FAGARA MAJOR. (From *fagus*, the beech, which it resembles) *Custana Luzonia*. *Cubebis*. *Fagara pierota* of Linnaeus. A plant found in the Philippine islands. The berries are aromatic, and, according to Ayicenna, heating, drying, good for cold, weak stomachs, and astringent to the bowels.

FAGARA OCTANDRA. The systematic name of the plant which affords *tacamahaca*. See *Tacamahaca*.

FAGOPYRUM. (From *φαγος*, the beech, and *φυρον*, wheat; because its seeds were supposed to resemble the mast, i. e. fruit of beech.) See *Polygonum fagopyrum*.

FAGOTRITICUM. See *Polygonum fagopyrum*.

FAGUS. (From *φαγε*, to eat; its nut being one of the first fruits used by man.)

1. The name of a genus of plants in the Linnæan system. Class, *Monæcia*. Order, *Polyandria*.

2. The pharmacopœial name of what is also called *Oxya*, *Balanda*, *Valanida*. The beech-tree. The fruit and interior bark of this tree, *Fagus sylvatica*; *foliis ovatis, obsolete serratis*, of Linnaeus, are occasionally used medicinally, the former in obstinate head-achs, and the latter in the cure of hectic fever. The oil expressed from beech-nuts is supposed to destroy worms; a child may take two drachms of it night and morning; an adult, an ounce. The poor people in Silesia use this oil instead of butter.

FAGUS CASTANEA. The systematic name of the chesnut-tree. See *Castanea*.

FAGUS SYLVATICA. The systematic name of the beech-tree. See *Fagus*.

Fainting. See *Syncope*.

FAIRBURN WATER. A sulphureous water in the county of Ross, in the north of England.

FALCIFORM PROCESS. (*Falciformis*; from *falx*, a scythe, and *forma*, resemblance.) The *falx*. A process of the dura mater, that arises from the *grista galli*, separates the hemispheres of the brain, and terminates in the tentorium.

FALSELLA. Contorted joint, used for compresses.

Falling sickness. See *Epilepsia*.

Fallopiæ tube. See *Tuba Fallopiæ*.

Fallopiæ ligament. See *Poupart's ligament*.

FALX. See *Falciform process*.

FANES CANINA. See *Bulimia*.

FANIGERATISSIMUM EMPLASTRUM. (From *famigeratus*, renowned; from *fama*, fame, and *gero*, to bear; so named from its excellence.) A plaster used in intermittent fevers, made of aromatic, irritating substances, and applied to the wrists.

FARFARA. (From *farfurus*, the white poplar; so called because its leaves resemble those of the white poplar.) See *Tussilago*.

FABINA. (From *far*, corn, of which it is made.) Meal, or flour. A term given to the pulverulent and glutinous part of wheat, and other seeds, which is obtained by grinding and sifting. It is highly nutritious, and consists of gluten, starch, and mucilage. See *Wheat*.

FARINACEA. (From *farina*, flour.) This term includes all those substances employed as aliment, called *cerealia*, *legumina*, and *nucæ oleosæ*.

FARINACEOUS. A term given to all articles of food which contain *farina*. See *Farina*.

FARINARIUM. See *Alica*.

FARNEUS. (From *far*, corn.) Scurfy. An epithet of urine, where it deposits a branny sediment.

FASCIA. (From *fascis*, a bundle; because, by means of a band, materials are collected into a bundle.) *Ligatio*. *Ligatura*. *Alligatura*. A bandage, fillet, or roller.

2. The aponeurotic expansions of muscles, which bind parts together, are termed *fasciæ*.

FASCIA LATA. A thick and strong tendinous expansion, sent off from the back, and from the tendons of the glutei and adjacent muscles, to surround the muscles of the thigh. It is the thickest on the outside of the thigh and leg, but towards the inside of both becomes gradually thinner. A little below the trochanter major, it is firmly fixed to the *linea aspera*; and, farther down, to that part of the head of the tibia that is next the fibula, where it sends off the tendinous expansion along the outside of the leg. It serves to strengthen the action of the muscles, by keeping them firm in their proper places when in action, particularly the tendons that pass over the joints where this membrane is thickest.

FASCIALIS. (*Fascialis*, sc. *musculus*.) See *Tensor vaginæ femoris*.

FASCIATIO. (From *fascia*, a fillet.) The binding up any diseased or wounded part with bandages.

FASCICULUS. (From *fascis*, a bundle.) A handful.

FAT. *Adeps.* A concrete oily matter contained in the cellular membrane of animals, of a white or yellowish colour, with little or no smell, nor taste. It differs in all animals in solidity, colour taste, &c. and likewise in the same animal at different ages. In infancy it is white, insipid, and not very solid; in the adult it is firm and yellowish, and in animals of an advanced age, its colour is deeper, its consistence various, and its taste in general stronger. Fat meat is nourishing to those that have strong digestive powers. It is used externally, as a softening remedy, and enters into the composition of ointments and plasters.

FATUITAS. (From *fatuus*, silly.) Foolishness. A synonym of *Amentia*.

FAUCES. (*Faux*, pl. *fauces*.) *Isthmion.* *Amphibranchia.* A cavity behind the tongue, palatine arch, uvula, and tonsils: from which the pharynx and larynx proceed.

FAUEL. *Terra japonica*, or catechu.

FAVASO AUSTRALIS. (From *favus*, a honey-comb, from its resemblance to a honey-comb.) A species of bastard sponge.

FAVUS. A honey-comb. A species of achor, or foul ulcer.

FEBRES. An order in the class *pyrexie* of Cullen, characterized by the presence of pyrexia, without primary or local affection.

FEBRICULA. (Dim. of *febris*, a fever.) A term employed to express a slight degree of symptomatic fever.

FEBRIFUGA. (From *febrim fugare*, to drive away a fever.) The plant fever-few; lesser centaury.

FEBRIFUGE. (*Febrifuga*, from *febris*, a fever, and *fugo*, to drive away.) A medicine that possesses the property of abating the violence of any fever.

FEBRIFUGUM CERATII. Regulus of antimony.

FEBRIFUGUM OLEUM. Febrifuge oil. The flowers of antimony, made with sal-ammoniac and antimony sublimed together, and exposed to the air, when they deliquesce.

FEBRIFUGUS PULVIS. Febrifuge powder. The Germans give this name to the pulvis stypticus Helvetii. In England, a mixture of oculi cancrorum and emetic tartar, in the proportions of half a drachm and two grains, has obtained the same name; in fevers it is given in doses of gr. iii. to iv.

FEBRIFUGUS SAL. Regenerated marine salt.

FEBRIFUGUS SPIRITUS DOMINI CLUTTON. Mr. Clutton's febrifuge spirit. An imperfect ether, which is sometimes given diluted in water, as a common drink in fevers.

FEBRIS. A fever. A disease characterized by an increase of heat, an accelerated pulse, a foul tongue and an impaired state of the functions of the body.

FEBRIS ALBA. See *Chlorosis*.

FEBRIS AMPHIMERINA. A quotidian, or remittent fever.

FEBRIS ANGINOSA. The scarlatina anginosa.

FEBRIS APHTHOSEA. An aphthous fever.

FEBRIS ARDENS. A burning inflammatory fever.

FEBRIS ASSODES. A tertian fever, with extreme restlessness.

FEBRIS AUTUMNALIS. An autumnal or bilious fever.

FEBRIS BULLOSA. The pemphigus, or vesicular fever.

FEBRIS CACATORIA. An intermittent, with diarrhoea.

FEBRIS CARCERUM. The prison fever.

FEBRIS CATARRHALIS. The catarrhal fever.

FEBRIS CHOLERICA. A fever with diarrhoea.

FEBRIS CONTINUA. A continued fever. A division of the order *febres*, in the class *pyrexie* of Cullen. Continued fevers have no intermission, but exacerbations come on twice in one day. The genera of continued fever are: 1. *Synocha*, or inflammatory fever, known by increased heat; pulse frequent, strong, and hard; urine high coloured; senses not much impaired. See *Synocha*; 2. *Typhus*, or putrid-tending fever, which is contagious, and is characterized by moderate heat; quick, weak, and small pulse; senses much impaired, and great prostration of strength. Typhus has four varieties, viz. 1. *Typhus petechialis*, typhus with petechiæ; 2. *Typhus mitior*, the nervous fever; 2. *Typhus gravior*, the putrid fever; 4. *Typhus icterodes*, the yellow fever. See *Typhus*.

FEBRIS LODES. A fever with continual and profuse sweating.

FEBRIS EPIDIAL. A fever with a continual sense of cold.

FEBRIS Erysipelatosa. See *Erysipelas*.

FEBRIS EXANTHEMATICA. Fever with eruptions.

FEBRIS FLAVA. The yellow fever.

FEBRIS HECTICA. A genus of disease in the class *pyrexie* and order *febres* of Cullen. It is known by exacerbations at noon, but chiefly in the evening, with slight remissions in the morning, after nocturnal sweats; the urine depositing a surfuraceous-lateritious sediment; appetite good; thirst moderate. Hectic fever is symptomatic of chlorosis, scrophula, phthisis, diseased viscera, &c.

FEBRIS HUNGARICA. A species of tertian fever.

FEBRIS HYDRODES. A fever with profuse sweats.

FEBRIS INFLAMMATORIA. See *Synocha*.

FEBRIS CASTRENSIS. A camp fever, generally typhus.

FEBRIS INTERMITTENS. An intermittent fever, or ague. A division of the order *febres* of Cullen, in the class *pyrexia*. Intermittent fevers are known by cold, hot, and sweating stages, in succession, attending each paroxysm, and followed by an intermission or remission. There are three genera of intermitting fevers, and several varieties.

1. *Intermittens quotidiana*. A quotidian ague. The paroxysms return in the morning, at an interval of about twenty-four hours.

2. *Intermittens tertiana*. A tertian ague. The paroxysms commonly come on at mid-day, at an interval of about forty-eight hours.

3. *Intermittens quartana*. A quartan ague. The paroxysms come on in the afternoon, with an interval of about seventy-two hours. The tertian ague is most apt to prevail in the spring, and the quartan in autumn.

Of the quotidian, tertian, and quartan intermittents, there are several varieties and forms; as the double tertian, having a paroxysm every day, with the alternate paroxysms, similar to one another. The double tertian, with two paroxysms every other day. The triple tertian, with two paroxysms on one day, and another on the next. The double quartan, with two paroxysms on the first day, none on the second and third, and two again on the fourth day. The double quartan, with a paroxysm on the first day, another on the second, but none on the third. The triple quartan, with three paroxysms every fourth day. The triple quartan, with a paroxysm every day, every fourth paroxysm being similar.

When these fevers arise in the spring of the year, they are called vernal; and when in the autumn, they are known by the name of autumnal. Intermittents often prove obstinate, and are of long duration, in warm climates: and they not unfrequently resist every mode of cure, so as to become very distressing to the patient; and by the extreme debility which they thereby induce, often give rise to other chronic complaints.

It seems to be pretty generally acknowledged, that marsh miasma, or the effluvia arising from stagnant water, or marsh ground, when acted upon by heat, is the most frequent exciting cause of this fever. In marshes, the putrefaction of both vegetable and animal matter is always going forward, it is to be presumed; and hence it has been generally conjectured, that vegetable and animal putrefaction imparted a peculiar quality to the watery particles of the effluvia arising from thence.

We are not yet acquainted with all the circumstances, which are requisite to render marsh miasma productive of intermittents; but it may be presumed that a moist atmosphere has a considerable influence in promoting its action. A watery poor diet, great fatigue, long watching, grief, much anxiety, exposure to cold, lying in damp rooms or beds, wearing damp linen, the suppression of some long-accustomed evacuation, or the recession of eruptions, have been ranked among the exciting causes of intermittents; but it is more reasonable to suppose that these circumstances act only by inducing that state of the body, which predisposes to these complaints. By some, it has been imagined that an intermittent fever may be communicated by contagion; but this supposition is by no means consistent with general observation.

One peculiarity of this fever is, its great susceptibility of a renewal from very slight causes, as from the prevalence of an easterly wind, or from the repetition of the original exciting cause. It would appear likewise, that a predisposition is left in the habit, which favours the recurrence of the complaint. In this circumstance, intermittents differ from most other fevers, as it is well known that, after a continued fever has once occurred, and been removed, the person affected is by no means so liable to a fresh attack of the disorder, as one in whom it had never taken place.

We have not yet attained a certain knowledge of the proximate cause of an intermittent fever, but a deranged state of the stomach and primæ viæ is that which is most generally ascribed.

Each paroxysm of an intermittent fever is divided into three different stages, which are called the *cold*, the *hot*, and the *sweating stages* or *fits*.

The *cold* stage commences with languor, a sense of debility and sluggishness in motion, frequent yawning and stretching, and an aversion to food. The face and extremities become pale, the features shrink, the bulk of every external part is diminished, and the skin over the whole body appears constricted, as if cold had been applied to it. At length the patient feels very cold, and universal rigours come on, with pains in the head, back, loins and joints, nausea, and vomiting of bilious matter; the respiration is small, frequent and anxious; the urine is almost colourless; sensibility is greatly impaired; the thoughts are somewhat confused; and the pulse is small, frequent, and often irregular. In a few instances, drowsiness and stupor have prevailed in so high degree as to resemble coma or apoplexy; but this is by no means usual.

These symptoms abating after a short time, the second stage commences with

an increase of heat over the whole body, redness of the face, dryness of the skin, thirst, pain in the head, throbbing in the temples, anxiety and restlessness; the respiration is fuller and more free, but still frequent; the tongue is furred, and the pulse has become regular, hard and full. If the attack has been very severe, then perhaps delirium will arise.

When these symptoms have continued for some time, a moisture breaks out on the forehead, and by degrees becomes a sweat, and this, at length, extends over the whole body. As this sweat continues to flow, the heat of the body abates, the thirst ceases, and most of the functions are restored to their ordinary state. This constitutes the third stage.

It must, however, be observed, that in different cases these phenomena, and their mode of succession, may prevail in different degrees, that the series of them may be more or less complete, and that the several stages, in the time they occupy, may be in different proportions to one another.

Such a depression of strength has been known to take place on the attack of an intermittent, as to cut off the patient at once; but an occurrence of this kind is very uncommon.

Patients are seldom destroyed in intermittents from general inflammation, or from a fulness of the vessels either of the brain or of the thoracic viscera, as happens sometimes in a continued fever; but when they continue for any length of time, they are apt to induce other complaints, such as a loss of appetite, flatulency, scirrhus of the liver, dropsical swellings, and general debility, which in the end now and then prove fatal. In warm climates, particularly, intermittents are very apt to terminate in this manner, if not speedily removed; and, in some cases, they degenerate into continued fevers. When the paroxysms are of short duration, and leave the intervals quite free, we may expect a speedy recovery; but when they are long, violent, and attended with much anxiety and delirium, the event may be doubtful. Relapses are very common to this fever at the distance even of five or six months, or even a year; autumnal intermittents are more difficult to remove than vernal ones, and quartans more so than the other types.

Dissections of those who have died of an intermittent, shew a morbid state of many of the viscera of the thorax and abdomen; but the liver, and organs concerned in the formation of bile, as likewise the mesentery, are those which are usually most affected.

FEBRIS LACTEA. Milk fever.

FEBRIS LENTA. A slow fever.

FEBRIS LENTICULARIS. A species of petechial fever.

FEBRIS MALIGNA. The malignant fever.

FEBRIS MILIARIA. The miliary fever.

FEBRIS MORBILLORA. The measles.

FEBRIS NERVOSA. *Febris lenta nervosa.* The nervous fever. A variety of *typhus mitior* of Cullen, but by many considered as a distinct disease. It mostly begins with loss of appetite, increased heat and vertigo; to which succeed nausea, vomiting, great languor, and pain in the head, which is variously described, by some like cold water pouring over the top, by others a sense of weight. The pulse, before, little increased, now becomes quick, feeble, and tremulous; the tongue is covered with a white crust, and there is great anxiety about the præcordia. Towards the seventh or eighth day, the vertigo is increased, and tinnitus aurium, copiosis, delirium, and a dry and tremulous tongue, take place. The disease mostly terminates about the fourteenth or twentieth day.

FEBRIS NOSOCOMICONUM. The fever of hospitals.

FEBRIS PALUSTRIS. The marsh fever.

FEBRIS PESTILENS. The plague.

FEBRIS PETECHIALIS. Fever with purple spots.

FEBRIS PUTRIDA. See *Typhus gravior*.

FEBRIS SCARLATINA. Scarlet fever.

FEBRIS SUDATORIA. Sweating fever.

FEBRIS SYNOCHA. See *Synocha*.

FEBRIS TYPHOIDES. See *Typhus*.

FEBRIS VARIOLOSA. A variolous or small-pox fever.

FEBRIS URTICARIA. Fever with nettle-rash.

FEBRIS VESICULOSA. See *Erysipelas*.

FECULA. See *Fecula*.

FEL. See *Bile*.

FEL NATURÆ. See *Aloes*.

Fel-wort. So called from its bitter taste like bile. See *Gentiana*.

FELLICULUS. The gall-bladder.

FELLIFLVA PASSIO. A name given to cholera morbus.

Felen. See *Paronychia*.

FERN. (*Quasi ferimen*; from *fero*, to bear; so called because it is the chief support of the body.) The thigh.

FEMORALIS ARTERIA. A continuation of the external iliac along the thigh, from Poupart's ligament to the ham.

FEMORIS OS. See *Femur*.

FEMUR. *Os femoris.* The thigh-bone. A long cylindrical bone, situated between the pelvis and tibia. Its upper extremity affords three considerable processes; these are, the head, the trochanter major, and trochanter minor. The head, which forms about two thirds of a sphere, is turned inwards, and is received into the acetabulum of the os innominatum, with which it is articulated by enarthrosis. It is covered by a cartilage, which is thickest in its middle part, and thin at its edges, but which is

wanting in its lower internal part, where a round spongy fossa is observable, to which the strong ligament, usually, though improperly, called the *round one*, is attached. This ligament is about an inch in length, flattish, and of a triangular shape, having its narrow extremity attached to the fossa just described, while its broader end is fixed obliquely to the rough surface near the inner and anterior edge of the acetabulum of the os innominatum, so that it appears shorter internally and anteriorly than it does externally and posteriorly.

The head of the os femoris is supported obliquely, with respect to the rest of the bone, by a smaller part, called the *cervez*, or *neck*, which, in the generality of subjects, is about an inch in length. At its basis we observe two oblique ridges, which extend from the trochanter major to the trochanter minor. Of these ridges, the posterior one is the most prominent. Around this neck is attached the capsular ligament of the joint, which likewise adheres to the edge of the cotyloid cavity, and is strengthened anteriorly by many strong ligamentous fibres, which begin from the lower and anterior part of the ilium, and, spreading broader as they descend, adhere to the capsular ligament, and are attached to the anterior oblique ridge at the bottom of the neck of the femur. Posteriorly and externally, from the basis of the neck of the bone, a large unequal protuberance stands out, which is the *trochanter major*. The upper edge of this process is sharp and pointed posteriorly, but is more obtuse anteriorly. A part of it is rough and unequal, for the insertion of the muscles; the rest is smooth, and covered with a thin cartilaginous crust, between which and the tendon of the glutæus maximus that slides over it, a large bursa mucosa is interposed. Anteriorly, at the root of this process, and immediately below the bottom of the neck, is a small process called *trochanter minor*. Its basis is nearly triangular, having its two upper angles turned towards the head of the femur and the great trochanter, while its lower angle is placed towards the body of the bone. Its summit is rough and rounded. These two processes have gotten the name of *trochanters*, from the muscles that are inserted into them being the principal instruments of the rotatory motion of the thigh. Immediately below these two processes the body of the bone may be said to begin. It is smooth and convex before, but is made hollow behind by the action of the muscles. In the middle of this posterior concave surface is observed a rough ridge, called *linea aspera*, which seems to originate from the trochanters, and extending downwards, divides at length into two branches, which terminate in the tuberosities near the condyles.

At the upper part of it, blood-vessels pass to the internal substance of the bone, by a hole that runs obliquely upwards.

The lower extremity of the os femoris is larger than the upper one, and somewhat flattened, so as to form two surfaces, of which the anterior one is broad and convex, and the posterior one narrower and slightly concave. This end of the bone terminates in two large protuberances, called *condyles*, which are united before, so as to form a pulley, but are separated behind by a considerable cavity, in which the crural vessels and nerves are placed secure from the compression to which they would be otherwise exposed in the action of bending the leg. Of these two condyles, the external one is the largest; and when the bone is separated from the rest of the skeleton, and placed perpendicularly, the internal condyle projects less forwards, and descends nearly three-tenths of an inch lower than the external one: but in its natural situation, the bone is placed obliquely, so that both condyles are then nearly on a level with each other. At the side of each condyle, externally, there is a tuberosity, the situation of which is similar to that of the condyles of the os humeri. The two branches of the *linea aspera* terminate in these tuberosities, which are rough, and serve for attachment of ligaments and muscles.

Fennel. See *Feniculum*.

Fennel, hog's. See *Peucedanum*.

FENESTRA OVALIS. An oblong or elliptical foramen, between the cavity of the tympanum and the vestibulum of the ear. It is shut by the stapes.

FENESTRA ROTUNDA. A round foramen, leading from the tympanum to the cochlea of the ear. It is covered by a membrane in the fresh subject.

Fenugreek. See *Fenugrecum*.

FERINE. (*Ferinus*, sc. *morbus*, savage or brutal.) A term occasionally applied to any malignant or noxious disease.

FERMENTATION. *Fermentatio*. A spontaneous commotion in a vegetable substance, by which its properties are totally changed. There are several circumstances required in order that fermentation may proceed: such are, 1. A certain degree of fluidity: thus, dry substances do not ferment at all. 2. A certain degree of heat. 3. The contract of air. Chymists, after Boerhaave, have distinguished three kinds of fermentation: the *spirituous*, which affords ardent spirit; the *acetous*, which affords vinegar, or acid; and the *putrid* fermentation, or putrefaction, which produces volatile alkali. The conditions necessary for spirituous fermentation are, 1. A saccharine mucilage. 2. A degree of fluidity slightly viscid. 3. A degree of

heat between 55 and 65 of Fahrenheit.

4. A large mass, in which a rapid commotion may be excited. When these four conditions are united, the spirituous fermentation takes place, and is known by the following characteristic phenomena: 1. An intestine motion takes place. 2. The bulk of the mixture then becomes augmented. 3. The transparency of the fluid is diminished by opaque filaments. 4. Heat is generated. 5. The solid parts mixed with the liquor rise and float in consequence of the disengagement of elastic fluid. 6. A large quantity of carbonic acid gas is disengaged by bubbles. All these phenomena gradually cease in proportion as the liquor loses its sweet and mild taste, and it becomes brisk, penetrating, and capable of producing intoxication. In this manner wine, beer, cider, &c. are made. All bodies which have undergone the spirituous fermentation are capable of passing on to the acid fermentation; but although it is probable that the acid fermentation never takes place before the body has gone through the spiritous fermentation, yet the duration of the first is frequently so short and imperceptible, that it cannot be ascertained. Besides the bodies which are proper for spirituous fermentation, this class includes all sorts of *fecula* boiled in water. The conditions required for the acid fermentation are, 1. A heat from 20 to 25 degrees of Fahrenheit. 2. A certain degree of liquidity. 3. The presence of atmospheric air. 4. A moderate quantity of fermentable matter. The phenomena which accompany this fermentation, are an intestine motion, and a considerable absorption of air. The transparent liquor becomes turbid, but regains its limpidity when fermentation is over. The fermented liquor now consists, in a great measure, of a peculiar acid, called the acetic acid, or vinegar. Not a vestige of spirit remains, it being entirely decomposed; but the greater the quantity of spirit in the liquor, previous to the fermentation, the greater will be the quantity of true vinegar obtained. See also *Putrefaction*.

FERMENTUM. (*Quasi fermentum* from *ferveo* to work) yeast.

Fern, male. See *Filix*.

Fern, female. See *Filix femina*.

FERRAMENTUM. An instrument made of iron.

FERRI CARBONAS. Carbonate of iron; formerly called *chalybis rubigo preparata* and *ferris rubigo*. "Take of sulphate of iron, eight ounces; subcarbonate of soda, ten ounces; boiling water, a gallon." Dissolve the sulphate of iron and subcarbonate of soda separately, each in four pints of water; then mix the solutions together and set it by, that the precipitated powder may subside; then, having poured off the

supernatant liquor, wash the carbonate of iron with hot water, and dry it upon bibulous paper in a gentle heat. It possesses corroborant and stimulating properties, and is exhibited with success in leucorrhœa, ataxia, asthenia, chlorosis, dyspepsia, rachites, &c. &c. Dose from two to ten grains.

FERRI LIMATURA PURIFICATA. Purified steel filings. These possess tonic, astringent, and deobstruent virtues, and are calculated to relieve chlorosis and other diseases in which steel is indicated, where crudity in the primæ viæ abounds.

FERRI RUBIGO. See *Ferri carbonas*.

FERRI SUBCARBONAS. See *Ferri carbonas*.

FERRI SULPHAS. Sulphate of iron; formerly called *sal martis*, *vitriolum martis*, *vitriolum ferri*, and lately *ferrum vitriolatum*. Green vitriol. "Take of iron, sulphuric acid, of each by weight, eight ounces; water, four pints." Mix together the sulphuric acid and water in a glass vessel, and add thereto the iron; then after the effervescence has ceased, filter the solution through paper, and evaporate it until chrysalis form as it cools. Having poured away the water, dry these upon bibulous paper. This is an excellent preparation of iron, and is exhibited, in many diseases, as a styptic, tonic, astringent and anthelmintic. Dose from one grain to five grains.

FERRUM AMMONIATUM. Ammoniated iron; formerly known by the name of *flores martiales*; *flores salis ammoniaci martiales*; *ens martis*; *ens veneris* Boyle: *sal martis muriaticum sublimatum*, and lately by the title of *ferrum ammoniacale*. "Take of carbonate of iron, muriate of ammonia, of each a pound." Mix them intimately, and sublime by immediate exposure to a strong fire: lastly, reduce the sublimed ammoniacal iron to powder. This preparation is astringent and deobstruent, in doses from 3 to 15 grs. or more in the form of bolus or pills, prepared with some gum. It is exhibited in most cases of debility, in chlorosis, asthenia menorrhagia, intermittent fevers, &c. This or some other strong preparation of iron, as the *Tinct. ferri muriatis*, Mr Cline is wont to recommend in scirrhus affections of the breast.

FERRUM TARTARIZATUM. Tartarized iron. A tartrate of potash and iron; formerly called *tartarus chalybeatus*; *mare solubilis*; *ferrum potabile*. "Take of iron, a pound; supertartrate of potash, powdered, two pounds; water, a pint." Rub them together, and expose them to the air in a broad glass vessel for eight days, then dry the residue in a sand bath, and reduce it to a very fine powder. Add to this powder a pint more water, and expose it for eight days longer, then dry it, and reduce it to a very fine powder. Its virtues are astringent and tonic, and it forms in solution an excellent tonic fomentation to con-

tusions, lacerations, distortions, &c. Dose from ten grains to half a drachm.

FEBRA. The measles.

* **FERULA.** The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*.

FERULA AFRICANA GALBANIFERA. The galbanum plant.

FERULA ASSAFETIDA. The systematic name of the assafetida plant. See *Assafetida*.

FERULA MINOR. Allheal of Esculapius; this plant is said to be detergent.

FERULACCA. The ferula galbanifera.

Fever. See *Febris*.

Feverfew. See *Matricaria*.

FIBER. (From *fiber*, extreme, because it resides in the extremities of lakes and rivers.) The beaver. The female beaver. *Castor fiber* of Linnæus, it has two excretory follicles near the anus, filled with an unctuous substance called castor. See *Castoreum*.

FIBRE. *Fibra.* An anatomical term for a very simple filament composed of earthy particles, connected together by an intermediate gluten. It is owing to the different arrangements of the fibres that the cellular structure, membranes, muscles, vessels, nerves, and, in short, every part of the body, except the fluids, are formed.

Fibre muscular. See *Muscular fibre*.

FIBRINE. The coagulable lymph is so termed by the French.

FIBULA. (*Quasi figillata*; from *figo*, to fasten; so named because it joins together the tibia and the muscles.) A long bone of the leg, situated on the outer side of the tibia, and which forms, at its lower end, the outer ankle. Its upper extremity is formed into an irregular head, on the inside of which is a slightly concave articulating surface, which, in the recent subjects, is covered with cartilage, and receives the circular flat surface under the edge of the external cavity of the tibia. This articulation is surrounded by a capsular ligament, which is farther strengthened by other strong ligamentous fibres, so as to allow only a small motion backwards and forwards.—Externally, the head of the fibula is rough and protuberant, serving for the attachment of ligaments, and for the insertion of the biceps cruris muscle.—Immediately below it, on its inner side, is a tubercle, from which a part of the gastrocnemius internus has its origin. Immediately below this head the body of the bone begins. It is of a triangular shape, and appears as if it were slightly twisted at each end, in a different direction. It is likewise a little curved inwards and forwards. This curvature is in part owing to the action of muscles: and in part perhaps to the carelessness of nurses.—Of the three angles of the bone, that which is turned towards the tibia is the most prominent, and serves for

the attachment of the interosseous ligament, which, in its structure and uses, resembles that of the fore-arm, and, like that, is a little interrupted above and below. The three surfaces of the bone are variously impressed by different muscles. About the middle of the posterior surface is observed a passage for the medullary vessels, slanting downwards. The lower end of the fibula is formed into a spongy, oblong head, externally rough and convex, internally smooth, and covered with a thin cartilage, where it is received by the external triangular depression at the lower end of the tibia. This articulation, which resembles that of its upper extremity, is furnished with a capsular ligament, and farther strengthened by ligamentous fibres, which are stronger and more considerable than those before described. They extend from the tibia to the fibula, in an oblique direction, and are more easily discernible before than behind. Below this the fibula is lengthened out, so as to form a considerable process, called *malleolus externus*, or the outer ankle. It is smooth, and covered with cartilage on the inside, where it is contiguous to the astragalus, or first bone of the foot. At the lower and inner part of this process there is a spongy cavity, filled with fat; and a little beyond this, posteriorly, is a cartilaginous groove, for the tendons of the peroneus longus and peroneus brevis, which are here bound down by the ligamentous fibres that are extended over them.

The principal uses of this bone seem to be, to afford origin and insertion to muscles, and to contribute to the articulation of the leg with the foot.

FICARIA. (From *ficus*, a fig, so called from its likeness.) The herb figwort, or pile-wort, the *ranunculus ficaria* of Linnæus.

FICATIO. (From *ficus*, a fig.) Tubercles near the anus and pudenda.

FICOIDEA. *Ficoides.* Resembling a fig. A name of the houseleek, which is emollient.

FICUS. 1. A fleshy excrescence about the anus, in figure resembling a fig.

2. The name of a genus of plants in the Linnæan system. Class, *Polygamia*. Order, *Diœcia*. The fig-tree.

FICUS CARICA. The systematic name of the fig-tree. See *Carica*.

FICUS INDICA. See *Lacca*.

FIDICINALES. (*Fidicinalis*, sc. *musculus*.) See *Lumbricalis*.

Figs. See *Carica*.

Figwort. See *Picaria*.

FILAGO. Cud or cotton-weed; an astringent.

FILAMENT. (*Filamentum*; from *filum*, a thread.) A term applied in anatomy to a small thread-like portion adhering to any part, and frequently synonymous with fibre. See *Fibre*.

FILLLUM. (From *filum*, a thread, because it resembles a string.) The frænum of the penis.

FILSTRUM. (From *filum*, a thread; named from its string-like appearance.) The frænum of the tongue.

FILICULA. (Dim. of *filix*, fern; a small sort of fern; also from *filum*, a thread, which it resembles.) Common maiden-hair. See *Adiantum capillus Veneris*.

FILIPENDULA. (From *filum*, a thread, and *pendeo*, to hang; so named because the numerous bulbs of its roots hang, as it were, by small threads.) *Saxifraga rubra*. Drop wort. The root of this plant, *Spiræa filipendula*; *foliis pennatis, foliolis uniformibus serratis, caule herbaceo, floribus corymbosis* of Linnaeus, possesses adstringent, and it is said lithontriptic virtues. It is seldom used in the practice of the present day.

FILIPENDULA AQUATICA. Water-drop-wort.

FILIUS ANTE PATREM. Any plant whose flower comes out before the leaf, as colts-foot.

FILIX. (From *filum*, a thread; so called from its being cut, as it were, in slender portions, like threads.) *Pteris*. *Blancnon* *Oribasii*. *Lonchitis*. Male polypody, or fern. *Polypodium filix mas* of Linnaeus. The root of this plant has lately been greatly celebrated for its effects upon the *tænia oculi superficialibus*, or broad tape-worm. Madame Noufer acquired great celebrity by employing it as a specific. This secret was thought of such importance by some of the principal physicians at Paris, who were deputed to make a complete trial of its efficacy, that it was purchased by the French king, and afterwards published by his order. The method of cure is the following:—after the patient has been prepared by an emollient glyster, and a supper of panada, with butter and salt, he is directed to take in the morning, while in bed, a dose of two or three drachms of the powdered root of the male fern. The powder must be washed down with a draught of water, and, two hours after, a strong cathartic, composed of calomel and scammony, is to be given, proportioned to the strength of the patient. If this does not operate in due time, it is to be followed by a dose of purging salts, and if the worm be not expelled in a few hours, this process is to be repeated at proper intervals. Of the success of this, or a similar mode of treatment, in cases of *tænia*, there can be no doubt, as many proofs in this country afford sufficient testimony; but whether the fern-root or the strong cathartic is the principal agent in the destruction of the worm, may admit of a question; and the latter opinion, Dr. Woodville believes, is the more generally adopted by physicians. It appears, however, from some experiments made in

Germany, that the *tænia* has, in several instances, been expelled by the repeated exhibition of the root, without the assistance of any purgative.

FILIX ACULEATA. Spear-pointed fern. The *Polypodium aculeatum* of Linnaeus.

FILIX FEMINA. The female fern. The plant which is thus called, in the pharmacopœias, is not the *Polypodium filix femina*, but the *Pteris aquilina*; *frondibus supradecompositis, foliis pinnatis, pinnis lanceolatis, infimis pinnatifidis, superioribus minoribus* of Linnaeus. The root is esteemed as an anthelmintic, and is supposed to be as efficacious in the destroying the tape-worm as the root of the male fern.

FILIX FLORIDA. The *Osmunda regalis* of Linnaeus. The flowering fern. *Emmenagogue*.

FILIX MAS. See *Filix*.

FILTRATION. (*Filtratio*: from *filtrum*, a strainer.) A method of rendering fluids clear, by passing them through a porous solid, as the filtering stone, compact close linen, woollen cloths, or porous paper, which is generally used for this purpose, as a lining to a funnel, or other such vessel. Filtration is also performed on a principle somewhat different, as by immersing one end of a porous substance, as a piece of list, skain of cotton, or slip of thick paper, or other such substance, moistened in its whole length in the fluid, and allowing the other end of it to hang down, over the outside of the vessel. The fluid in this depending part drains out by its own gravity, and is supplied by capillary attraction from the portion next within the vessel, which is supplied in the same manner from the surface of the fluid, till the whole passes over, unless too deep, the list, &c. appearing to act as syphons.

FILTRUM. A filter, straining or filtering instrument.

FILUM ARSENICALE. Corrosive sublimate.

FIMBRIÆ. (*Quasi finibria*; from *finis*, the extremity.) The extremities of the Fallopian tubes. See *Uterus*.

Finkle. See *Feniculum*.

Fine leaved water hemlock. See *Feniculum aquaticum*.

Finochio. The Italian name of the sweet azorian fennel.

Fir-tree. See *Abies*.

Fir, balsam of Gilead. See *Balsamea*.

Fir, Canada. See *Balsamum Canadense*.

Fir, Norway spruce. See *Pinus abies*.

Fir, Scotch. See *Pinus sylvestris*.

Fir, silver. See *Pinus picea*.

FIRE. *Ignis*. A very simple and active element, the principal agent in nature to balance the power and natural effect of attraction. The most useful acceptance of the word fire comprehends *heat* and *light*. There have been several theories proposed respecting fire, but no one as yet is gene-

rally received. It is therefore, at present, only known by its effects, namely, light, heat, rarefaction. See *Caloric* and *Light*.

FIRMISIMUM MINERALIUM. Antimony.

Fish-glue. See *Ichthyocollium*.

FISSURA MAGNA SYLVII. The anterior and middle lobes of the cerebrum on each side are parted by a deep narrow sulcus, which ascends obliquely backwards from the temporal ala of the os sphenoides, to near the middle of the os parietale, and this sulcus is thus called.

FISSURE. *Fissura.* That species of fracture in which the bone is slit, but not completely divided.

Fistic-nut. See *Pistachio-nut*.

FISTULA. (*Quasi funiculus*; from *fundo*, to pour out; or from its similarity to a pipe, or reed.) *Eligii morbus.* A term in surgery, applied to a long and sinuous ulcer that has a narrow opening, and which sometimes leads to a larger cavity; and has no disposition to heal. No technical term has been more misapplied; and no mis-interpretation of a word has had worse influence in practice than this. Many simple healthy abscesses, with small openings, have too often been called *fistulous*; and, the treatment pursued has in reality at last rendered them so, and been the only reason of their not having healed.

FISTULARIA. (From *fistula*, a pipe; so called because its stalk is hollow.) A name for stavesacre.

Five-leaved grass. See *Pentaphyllum*.

Fixed air. See *Carbonic acid*.

FIXED BODIES. Chymists give this name to those substances, which cannot be caused to pass by a strong rarefaction from the liquid state to that of an elastic fluid.

Flag, sweet. See *Calamus aromaticus*.

Flag, yellow water. See *Iris palustris*.

FLAMMULA. (Dim. of *flamma*, a fire, named from the burning pungency of its taste.) *Surrecta alba.* Small water crow-foot, or spear-wort. The roots and leaves of this common plant, *Ranunculus flammula*; *foliis ovatis-lanceolatis, petiolatis, caule declinato* of Linnæus; taste very acrid and hot, and, when taken in a small quantity, produce vomiting, spasms of the stomach, and delirium. Applied externally, they vesicate the skin. The best antidote, after clearing the stomach, is cold water acidulated with lemon-juice, and then mucilaginous drinks.

FLAMMULA JOVIS. *Flammula jovis*, so called from the burning pungency of its taste.) Upright virgin's bower. *Clematis recta*; *foliis pinnatis, foliolis ovato lanceolatis integerrimis, caule erecto, floribus pentapetalis tetrapetalisque* of Linnæus. More praises have been bestowed upon the virtue which the leaves of this plant are said to possess, when exhibited internally, as an antiveneræal, by foreign physicians, than its trials in this country can justify. The

powdered leaves are sometimes applied externally to ulcers, as an escharotic.

FLATULENT. Windy.

Flax-leaved daphne. See *Thymelea*.

Flax, purging. See *Linum Catharticum*.

Flax, spurge. See *Thymelea*.

Flea wort. See *Prylimum*.

FLEKEN. (From *flecto*, to incline downwards.) *Flegmen.* A tumour about the ancles.

FLEURESIN. Gout.

FLEXOR. The name of several muscles, whose office it is to bend parts into which they are inserted.

Flexor accessorius digitorum pedis. See *Flexor longus digitorum pedis*.

FLEXOR BREVIS DIGITORUM PEDIS, PERFORATUS, SUBLIMIS. *Flexor brevis digitorum pedis perforatus* of Albinus. *Flexor brevis* of Douglas. *Flexor digitorum brevis, sive perforatus pedis* of Winslow. *Perforatus seu flexor secundus internodii digitorum pedis* of Cowper, and *calcaneus phalangeitien communis* of Dumas. A flexor muscle of the toes, situated on the foot. It arises by a narrow, tendinous, and fleshy beginning, from the inferior protuberance of the os calcis. It likewise derives many of its fleshy fibres from the adjacent aponeurosis, and soon forms a thick belly, which divides into four portions. Each of these portions terminates in a flat tendon, the fibres of which decussate, to afford a passage to a tendon of the long flexor, and afterwards re-uniting, are inserted into the second phalanx of each of the four lesser toes. This muscle serves to bend the second joint of the toes.

FLEXOR BREVIS MINIMI DIGITI PEDIS. *Parathenar minor* of Winslow. This little muscle is situated along the inferior surface and outer edge of the metatarsal bone of the little toe. It arises tendinous from the basis of that bone, and from the ligaments that connect it to the os cuboides. It soon becomes fleshy, and adheres almost the whole length of the metatarsal bone, at the anterior extremity of which it forms a small tendon, that is inserted into the root of the first joint of the little toe. Its use is to bend the little toe.

FLEXOR BREVIS POLLICIS MANUS. *Flexor secundus internodii* of Douglas. *Thenar* of Winslow. *Flexor primi et secundus ossis pollicis* of Cowper, and *Carpophalangin du pouce* of Dumas. This muscle is divided into two portions by the tendon of the flexor longus pollicis. The outermost portion arises tendinous from the anterior part of the os trapezoides and internal annular ligament. The second, or innermost, and thickest portion, arises from the same bone, and likewise from the os magnum, and os cuneiforme. Both these portions are inserted tendinous into the sesamoid bones of the thumb. The use of this muscle is to bend the second joint of the thumb.

FLEXOR BREVIS POLLICIS PE-

DMS. *Flexor brevis* of Douglas. *Flexor brevis pollicis* of Cowper, and *Tarso-phalangien du ponce* of Dumas. A muscle of the great toe, that bends the first joint of the great toe. It is situated upon the metatarsal bone of the great toe, arises tendinous from the under and anterior part of the os calcis, and from the under part of the os cuneiforme externum. It soon becomes fleshy, and divisible into two portions, which do not separate from each other till they have reached the anterior extremity of the metatarsal bone of the great toe, where they become tendinous, and then the innermost portion unites with the tendon of the abductor, and the outermost with that of the abductor pollicis. They adhere to the external os sesamoideum, and are finally inserted into the root of the first joint of the great toe. These two portions, by their separation, form a groove, in which passes the tendon of the flexor longus pollicis.

FLEXOR CARPI RADIALIS. *Radialis internus* of Albinus and Winslow, and *Epirochlo-metacarpien* of Dumas. This is a long thin muscle, situated obliquely at the inner and anterior part of the fore-arm, between the palmaris longus and the pronator teres. It arises tendinous from the inner condyle of the os humeri, and, by many fleshy fibres, from the adjacent tendinous fascia. It descends along the inferior edge of the pronator teres, and terminates in a long, flat, and thin tendon, which afterwards becomes narrower and thicker, and, after passing under the internal annular ligament, in a groove distinct from the other tendons of the wrist, it spreads wider again, and is inserted into the fore and upper part of the metacarpal bone that sustains the fore-finger. It serves to bend the hand, and its oblique direction may likewise enable it to assist in its pronation.

FLEXOR CARPI ULNARIS. *Ulnaris internus* of Winslow and Albinus. *Epirochli-cubito-carpiei* of Dumas. A muscle situated on the cubit or fore-arm, that assists the former in bending the arm. It arises tendinous from the inner condyle of the os humeri, and, by a small fleshy origin, from the anterior edge of the olecranon. Between these two portions, we find the ulnar nerve passing to the fore-arm. Some of its fibres arise likewise from the tendinous fascia that covers the muscles of the fore-arm. In its descent, it soon becomes tendinous, but its fleshy fibres do not entirely disappear till it has reached the lower extremity of the ulna, where its tendon spreads a little, and, after sending off a few fibres to the external and internal os pisiforme, is inserted into the os pisiforme.

FLEXOR LONGUS DIGITORUM PEDIS, PROFUNDUS, PERFORANS. *Perforans seu flexor profundus* of Douglas,

Flexor digitorum longus, et se perforans pedis, and *perforans seu flexor tertii internodii digitorum pedis* of Cowper, and *Tibio phalangettien* of Dumas. A flexor muscle of the toes, situated along the posterior part and inner side of the leg. It arises fleshy from the back part of the tibia, and, after running down to the internal ankle, its tendon passes under a kind of annular ligament, and then through a sinuosity at the inside of the os calcis. Soon after this it receives a small tendon from the flexor longus pollicis pedis, and about the middle of the foot it divides into four tendons, which pass through the slits of the flexor brevis digitorum pedis, and are inserted into the upper part of the last bone of all the lesser toes. About the middle of the foot, this muscle unites with a fleshy portion, which, from the name of its first describer, has been usually called *massa carnea Jacobi Sylvii*: it is also termed *Flexor accessorius digitorum pedis*. This appendage arises by a thin fleshy origin, from most part of the sinuosity of the os calcis, and likewise by a thin tendinous beginning from the anterior part of the external tubercle of that bone; it soon becomes all fleshy, and unites to the long flexor just before it divides into its four tendons. The use of this muscle is to bend the last joint of the toes.

FLEXOR LONGUS POLLICIS MANUS. *Flexor longus pollicis* of Albinus, *Flexor tertii internodii* of Douglas. *Flexor tertii internodii seu longissimus pollicis* of Cowper, and *radio-phalangettien du ponce* of Dumas. This muscle is placed at the side of the flexor longus digitorum pedis, profundus, perforans, and is covered by the extensores carpi radiales. It arises fleshy from the anterior surface of the radius, immediately below the insertion of the biceps, and is continued down along the oblique ridge, which serves for the insertion of the supinator brevis, as far as the pronator quadratus. Some of its fibres spring likewise from the neighbouring edge of the interosseous ligament. Its tendon passes under the internal annular ligament of the wrist, and after running along the inner surface of the first bone of the thumb, between the two portions of the flexor brevis pollicis, goes to be inserted into the last joint of the thumb, being bound down in its way by the ligamentous expansion that is spread over the second bone. In some subjects we find a tendinous portion arising from the inner condyle of the os humeri, and forming a fleshy slip that commonly terminates near the upper part of the origin of this muscle from the radius. The use of this muscle is to bend the last joint of the thumb.

FLEXOR LONGUS POLLICIS PEDIS. This muscle is situated along the posterior part of the leg. It arises tendinous and fleshy a little below the head of

the fibula, and its fibres continue to adhere to that bone almost to its extremity. A little above the heel it terminates in a round tendon, which, after passing in a groove formed at the posterior edge of the astragalus, a distal internal and lateral part of the os calcis, in which it is secured by an annular ligament, goes to be inserted into the last bone of the great toe, which it serves to bend.

FLEXOR OSSIS METACARPI POLLICIS: *Seu opponens pollicis* of Innes. *Opponens pollicis manus* of Albinus. *Flexor primi internodii* of Douglas. *Antithenar et sic semi-interosseus pollicis* of Winslow, and *carpo-phalangien du pouce* of Dumas. This muscle is situated under the abductor brevis pollicis, which it resembles in its shape. It arises tendinous and fleshy from the os scaphoides, and from the anterior and inner part of the internal annular ligament. It is inserted tendinous and fleshy into the under and anterior part of the first bone of the thumb. It serves to turn the first bone of the thumb upon its axis, and at the same time to bring it inwards, opposite to the other fingers.

FLEXOR PARVUS MINIMI DIGITI. *Abductor minimi digiti*, *Hypothenar Riolani* of Douglas. *Hypothenar minimi digiti* of Winslow, and *second-carpo-phalangien du petit doigt* of Dumas. The situation of this muscle is along the inner surface of the metacarpal bone of the little finger. It arises tendinous and fleshy from the hook-like process of the unciform bone, and likewise from the anterior surface of the adjacent part of the annular ligament. It terminates in a flat tendon, which is connected with that of the abductor minimi digiti, and inserted into the inner and anterior part of the upper end of the first bone of the little finger. It serves to bend the little finger, and likewise to assist the abductor.

FLEXOR PROFUNDUS PERFORANS. *Profundus* of Albinus. *Perforans* of Douglas. *Perforans vulgo profundus* of Winslow. *Flexor tertii internodii digitorum manus, vel perforatus manus* of Cowper, and *cubito-phalangietien commun.* of Dumas. This muscle is situated on the fore-arm, immediately under the perforatus, which it greatly resembles in its shape. It arises fleshy from the external side, and upper part of the ulna, for some way downwards, and from a large portion of the interosseous ligament. It splits into four tendons a little before it passes under the annular ligament of the wrist, and these pass through the slits in the tendons of the flexor sublimis, to be inserted into the fore and upper part of the third or last bone of all the four fingers, the joint of which they bend.

FLEXOR SUBLIMIS PERFORATUS. This muscle, which is the *perforatus* of

Cowper, Douglas, and Winslow, is by Albinus and others named *sublimis*. It has gotten the name of *perforatus* from its tendons being perforated by those of another flexor muscle of the finger, called the *perforans*. They who give it the appellation of *sublimis*, consider its situation with respect to the latter, and which, instead of *perforans*, they name *perfundus*. It is a long muscle, situated most commonly at the anterior and inner part of the forearm, between the palmaris longus and the flexor carpi ulnaris; but, in some subjects, we find it placed under the former of these muscles, between the flexor carpi ulnaris and the flexor carpi radialis. It arises, tendinous and fleshy, from the inner condyle of the os humeri, from the inner edge of the coronoid process of the ulna, and from the upper and fore part of the radius, down to near the insertion of the pronator teres. A little below the middle of the forearm, its fleshy belly divides into four portions, which degenerate into as many round tendons, that pass all together under the internal annular ligament of the wrist, after which they separate from each other, become thinner and flatter, and running along the palm of the hand, under the aponeurosis palmaris, are inserted into the upper part of the second bone of each finger. Previous to this insertion, however, the fibres of each tendon decussate near the extremity of the first bone, so as to afford a passage to a tendon of the perforans. Of these four tendons, that of the middle finger is the largest, that of the fore-finger the next in size, and that of the little-finger the smallest. The use of this muscle is to bend the second joint of the fingers.

Flexor tertii internodii. See *flexor longus pollicis manus*.

FLORES BENZOES. See *Benzoic acid*.

FLORES MARTIALES. See *Ferrum ammoniatum*.

FLORES SALIS AMMONIACI. See *Carbonas ammoniac*.

FLORES SULPHURIS. See *Sulphur*.

FLORES SULPHURIS LOTI. When sublimed sulphur is boiled and washed in water, it forms the *flores sulphuris loti* of the pharmacopoeias. Analogous to this preparation is the *sulphur precipitatum*; they possess eccoprotic, diaphoretic, and antispasmodic virtues, and are administered in obstipation, where there are piles, colica pictonum, worm cases, to diminish salivations, &c.

FLOWERS. A term formerly employed by chymists to the fine parts which are sublimated from certain bodies, as the flowers of benjamin, sulphur, zinc, &c. - *Flower-de-luce.* See *Iris nostras*.

Flowers of Benjamin. See *Benzoic acid*.

FLUAT. A salt formed by the fluoric acid, combined with different bases: thus, *fluat of alumina, fluat of ammonia, &c.*

FLUCTUATION. A term used by surgeons, to express the undulations of a fluid; thus, when pus is formed in an abscess, or when water accumulates in the abdomen, if the abscess or abdomen be lightly pressed with the fingers, the motion of fluctuation may be distinctly felt.

Fuellin. See *Elatine*.

FLUID. A fluid is that, the constituent principles of which so little attract each other, that, when poured out, it drops *guttatim*, and adapts itself, in every respect, to the form of the vessel containing it.

FLUIDS OF THE BODY. This term is often applied to the blood, and other humours of which the body is composed. The drying of any part of the human body demonstrates that by far the greater part consists of fluid. The quantity of fluid in a man of one hundred and sixty pounds weight, is estimated at one hundred and thirty-five pounds. The fluids of the human body are divided into, 1. *Crude*, or those which have not yet entirely put on the animal nature, as the chyme and chyle. 2. *Sanguineous*; to this is referred the blood, or the cruor of the blood. 3. *Lymphatic*, which are those of the lymphatic vessels, and the nutritious jelly. 4. *Secreted*; to this head are referred all those separated from the blood, which are very numerous. 5. *Excrementitious*, which are eliminated from the body, as the alvine faeces, urine, cutaneous and pulmonary perspirable matter.

The secreted humours are divided into, 1. *Lacteal*, which are white; as the milk, juice of the prostate and thymus glands. 2. *Aqueous*, as the aqueous humour of the eye. 3. *Mucous*, as the mucus of the nostrils and primæ viæ. 4. *Albuminous*, as the serum of the blood. 5. *Oleous*, as the oil of the adipose membrane. 6. *Bilious*, as the bile and wax of the ears.

The fluids of the human body are also divided from their motion into, 1. *Circulating*, which continually circulate in the vessels. 2. *Commorant*, which circulate with a slow motion, as the oil of the adipose membrane and male semen. 3. *Stagnant*, which remain for a certain time in any receptacle, as cystic bile, urine, and the alvine faeces.

FLUOR ALBUS. The *Leucorrhæa*.

FLUOR SPAR. Vitreous spar. Sparry flour. Derbyshire spar. A species of salt which abounds in nature, formed by the combination of the fluoric acid with lime. It is called spar, because it has the sparry form and fracture; fluor, because it melts very readily; and vitreous, because it has the appearance of glass, and may be fused into glass of no contemptible appearance.

FLUORIC ACID. The acidifiable base of fluoric acid is unknown; for no experi-

ments have yet been made by which it can be exhibited. It is merely from general inference, that we may conjecture it to consist of a simple acidifiable basis united to oxygen.

This acid is to be obtained by putting one part of finely pulverized fluuate of lime into a leaden or tin retort, and pouring upon it two parts and a half of sulphuric acid. Lute the retort to a receiver of the same metal, containing one part of water, and apply a gentle heat. The fluoric acid gas, disengaged, will be absorbed by the water, and form liquid fluoric acid, which must be kept in well closed leaden or tin bottles, or phials, coated within with wax or varnish.

The distinguishing property of fluoric acid is, its power of dissolving and volatilizing silex. Its odour resembles muriatic acid. On being exposed to a moist atmospheric air, it emits white fumes. Its action upon all the inflammable substances is very feeble; it does not afford oxygen to them. It has no action upon most of the metals, but it dissolves many of their oxids.

Fluoric acid, united to different bases, forms saline compounds called *FLUATS*.

FLUORIC ACID GAS. The most remarkable property of this gas is, its power of dissolving silex, and keeping it suspended in the gaseous state; it therefore dissolves glass, crystals, and various precious stones. It is heavier than atmospheric air. It does not maintain combustion, nor can animals breathe it. It is absorbed by water, and forms with it liquid fluoric acid. It has a penetrating odour, approaching nearly to that of muriatic acid gas. It corrodes animal and vegetable matters. Light has no effect upon it. It emits white fumes, in contact with moist atmospheric air. It is very sour, and therefore reddens blue vegetable colours. It precipitates lime-water. With ammonia it unites and forms a concrete body. It has no action upon platina, gold, silver mercury, tin, lead, antimony, cobalt, nickel, and bismuth; but it attacks iron, arsenic, and manganese.

Method of obtaining Fluoric Acid Gas.

1. Fluoric acid gas may be obtained by decomposing fluuate of lime (Derbyshire spar) by means of sulphuric acid.

For that purpose, put one part of powdered fluuate of lime into a leaden or tin retort, and pour over it two or three parts of concentrated sulphuric acid. A violent action takes place instantly, and fluoric acid gas is extricated, which must be collected over mercury, either in a leaden or tin vessel, or in a glass receiver, covered within with a thick coat of varnish and wax. When no more gas is produced spontaneously, the action of the acid must be assisted by a gentle heat.

Fluate of lime consists of fluoric acid, united to lime; on adding sulphuric acid, in the above process a decomposition takes place. The sulphuric acid having a greater attraction for the lime than the fluoric acid, unites to it and forms sulphate of lime, disengaging at the same time the fluoric acid, which takes the gaseous state at the instant of its extrication, and appears under the form of fluoric acid gas.

2. Fluoric acid gas may likewise be obtained from liquid fluoric acid, by heat.

Flux. This word is mostly employed for *dyenteria sanguinea*.

FLUXION. A term mostly applied by chymists, to signify the change of metals, or other bodies, from the solid into the fluid state, by the application of heat. See *Fusion*.

Fly, Spanish. See *Cantharis*.

FOCILE MAJUS. The ulna and the radius are occasionally denominated by the barbarous appellations of *focele majus* and *minus*; the tibia and fibula in the leg are also so called.

FOCUS. A lobe of the liver.

FODINA. (From *fodio*, to dig.) A quarry. The labyrinth of the ear.

FENICULATUM LIGNUM. A name for *sassafras*.

FENICULUM. (*Quasi fanum oculorum*, the hay or herb good for the sight; so called because it is thought good for the eyes.) Fennel.

FENICULUM AQUATICUM. Water fennel. Fine-leaved water hemlock. The plant which bears this name in the pharmacopœias is the *Phellandrium aquaticum*; *foliorum ramificationibus divaricatis* of Linnaeus. It possesses vertiginous and poisonous qualities, which are best counteracted by acids, after clearing the primæ viæ. The seeds are recommended by some, in conjunction with Peruvian bark, in the cure of pulmonary phthisis.

FENICULUM ALPINUM. The herb spig-nel.

FENICULUM ANNUUM. Royal cummin.

FENICULUM DULCE. Common fennel. *Anethum feniculum*, *fructibus ovatis* of Linnaeus. The seeds and roots of this indigenous plant are directed by the colleges of London and Edinburgh. The seeds have an aromatic smell, and a warm sweetish taste, and contain a large proportion of essential oil. They are stomachic and carminative. The root has a sweet taste, but very little aromatic warmth, and is said to be pectoral and diuretic.

FENICULUM MARINUM. Samphire.

FENICULUM ORIENTALE. Cuminum.

FENICULUM PORCINUM. See *Peucedanum*.

FENICULUM SINENSE. Aniseed.

FENICULUM SYLVESTRIS. Bastard spig-nel.

FENICULUM TOMATOSUM. French Bitter-wort.

FENICULUM VULGARE. *Feniculum germanicum.* *Maruthram.* Common fennel, or fenchel. A variety of the *Anethum feniculum*. See *Feniculum dulce*.

FENUM CAMELORUM. See *Juncus sub-ratus*.

FENUM GRÆCUM. (*Fenum*, hay, and *græcus*, belonging to Greece; because, in Greece, it grew in the meadows, like hay.) *Buccras.* *Ægoceras.* Fenugreek. *Trigonella fenum græcum* of Linnaeus, *leguminibus scabulis strictis erectiusculis subul-catis acuminatis, caule erecto*. A native of Montpellier. The seeds are brought to us from the southern parts of France and Germany; they have a strong disagreeable smell, and an unctuous farinaceous taste, accompanied with a slight bitterness. They are esteemed as assisting the formation of pus, in inflammatory tumours; and the meal, with that intention, is made into a poultice with milk.

FENUM SYLVESTRIS. Wild fenugreek.

FESTABULUM. (From *fæto*, to become putrid.) An encysted abscess, or foul ulcer.

FÆTUS. (From *fæo*, to bring forth, according to Vossius.) *Epicyema.* *Epigonion*. The child enclosed in the uterus of its mother, is called a *fætus* from the fifth month after pregnancy until the time of its birth. The internal parts peculiar to the *fætus*, are the thymus gland, *canalis venosus*, *canalis arteriosus*, *foramen ovale*, and the *membrana pupillaris*. Besides these peculiarities, there are other circumstances in which the *fætus* differs from the adult. The lungs are black and collapsed, and sink in water; the liver is very large; all the glands, especially the thymus and suprarenal, and the vermiform process of the cæcum, are also considerably larger in proportion. The teeth of the *fætus* are hid within their sockets; the great intestines contain a substance called *meconium*; the *membrana tympani* is covered with a kind of mucous membrane, and the bones in many places are cartilaginous.

FOLIATA TERRA. (From its resemblance to *folium*, a leaf.) Sulphur, also the acetas potassæ.

FOLIUM ORIENTALE. See *Senna*.

FOLLICLE. *Folliculus*; dim. of *folia*, a bag.) See *Folliculose gland*.

FOLLICULOSE GLAND. Follicle. One of the most simple species of gland, consisting merely of a hollow vascular membrane or follicle, and an excretory duct; such are the muciparous glands, the sebaceous, &c.

FOLLICULUS FELLIS. The gall-bladder.

FOMENTATION. *Fomentatio*. A sort of partial bathing, by applying hot flannels to any part, dipped in medicated

decoctions, whereby steams are communicated to the diseased parts, their vessels are relaxed, and their morbid action sometimes removed.

FONNA VENTRICULI. Hypochondriacism.

FONITES. A term mostly applied as synonymous with contagion.

FONS PULSATILIS. See *Fontanella*.

FONTANELLA. (Dim. of *fons*, a fountain.) *Fons pulsatilis.* The parietal bones and the frontal bones do not coalesce until the third year after birth, so that, before this period, there is an obvious interstice, commonly called *mould*, and scientifically the *fontanel*, or *fons pulsatilis*. There is also a lesser space, occasionally, between the occipital and parietal bones, termed the *posterior fontanel*. These spaces between the bones are filled up by the dura mater and the external integuments, so that during birth, the size of the head may be lessened; for, at that time, the bones of the head, upon the superior part, are not only pressed nearer to each other, but they frequently lap over one another, in order to diminish the size during the passage of the head through the pelvis.

FONTANELLA ANTERIOR. See *Fontanella*.

FONTANELLA POSTERIOR. See *Fontanella*.

FONTICULUS. (Dim. of *fons*, an issue.) An artificial ulcer formed in any part, and kept discharging by introducing daily a pea, covered with any digestive ointment.

FORAMEN. (From *foro*, to pierce.) A little opening.

FORAMEN CÆCUM. A single opening in the basis of the cranium between the ethmoid and the frontal bone, that gives exit to a small vein.

FORAMEN LACERUM IN BASI CRANII. See *Foramina lacera*.

FORAMEN LACERUM ORBITALE SUPERIUS. A large opening between the greater and lesser wing of the parietal bone through which the third, fourth, first branch of the fifth and the sixth pair of nerves, and the ophthalmia artery pass.

FORAMEN OPTICUM. The hole transmitting the optic nerve.

FORAMEN OVALE. The opening between the two auricles of the heart of the fœtus. See also *Innominatum os*.

FORAMEN OF WINSLOW. An opening in the omentum. See *Omentum*.

FORAMINA LACERA. A pair of foramina in the basis of the cranium, through which the internal jugular veins and the eighth pair of accessory nerves pass.

FORAMINULUM OS. The ethmoid bone.

FORCEPS. (*Quasi ferriceps*, as being the iron with which we seize any thing hot, from *ferrum*, iron, and *capio*, to take.)

Placens. A surgical instrument with which extraneous bodies or other substances are extracted. Also an instrument occasionally used by men midwives to bring the head of the fœtus through the pelvis.

Foreskin. See *Prepuce*.

FORMIAT. *Formias.* A salt produced by the union of the formic acid with different bases: thus, *formiat of alumina*, *formiat of ammonia*, &c.

FORMICA. (*Quod ferat micas*, because of his diligence in collecting small particles of provision together.) 1. The ant, or pismire. This industrious little insect, *Formica rufa* of Linnæus, contains an acid juice and gross oil, which were supposed to possess aphrodisiac virtues. The crystalles of this animal are said to be diuretic and carminative, and by some recommended in the cure of dropsy.

2. The name of a black wart with a broad base, and cleft superficies, because the pain attending it resembles the biting of an ant.

3. A varicose tumour on the anus and glans penis.

FORMICA MILIARIS. Any herpetic eruption.

FORMIC ACID. *Addum formicum.* The acid of ants was known to Tragus, Bauhin, Fisher, Etmüller, Hoffman, and many others. It is obtained chiefly from the red ant, *Formica rufa* of Linnæus, by distilling them in a retort, and by washing them in boiling water. When rectified, and rather concentrated, it has a penetrating smell, and is corrosive; and its taste is so agreeable, when greatly diluted with water, that it has been proposed to be used instead of vinegar.

FORMIX. The herpes exedens.

FORMULA. (Dim. of *forma*, a form.) A little form of prescription, such as physicians direct in extemporaneous practice, in distinction from the greater forms in pharmacopœias, &c.

FORFAX. *Furnus.* A furnace. A chemical instrument. See *Furnace*.

FORNIX. (*Formix*, an arch or vault.) A part of the corpus callosum in the brain is so called, because, if viewed in a particular direction, it has some resemblance to the arch of an ancient vault. It is the medullary body, composed of two anterior and two posterior crura, situated at the bottom and inside of the lateral ventricle, over the third ventricle, and below the septum lucidum.

FORTIS AQUA. A weak nitrous acid.

FOSSA. (From *fodis*, to dig.) *Fovea.* A little depression or sinus. The pudendum muliebre.

FOSSA AMYNTÆ. A double-headed roller for the face.

FOSSA MAGNA. The great groove of the ear, the pudendum muliebre.

FOSSA NAVICULARIS. The cavity at the U U

entrance of the pudendum muliebre: also the great groove of the ear.

FOSSA OVALIS. The depression in the right auricle of the human heart, which in the fetus opened into the other auricle, forming the foramen ovale.

FOSSA PITUITARIA. The fossa of the sella turcica of the sphenoid bone.

FOSSIL. (*Fossilis*, from *fodio*, to dig.) Any thing dug out of the earth.

FOSILLUS. The bone of the leg.

FOVEA. (From *fodio*, to dig.) A little depression. The pudendum muliebre. A partial sweating-bath.

Fox-glove. See *Digitalis*.

FOX-GLOVE, EASTERN. The seeds of this plant, *Sesamum orientale* of Linnæus, are in much esteem in South Carolina, where they are called *oily grain*, they are made into soups and puddings after the manner of rice. Toasted over the fire, they are mixed with other ingredients, and stewed into a delicious food. The fresh seed affords a considerable quantity of a warm pungent oil, otherwise not unpalatable. In a year or two the pungency leaves it, when the oil is used for salad, &c. The seed of the *Sesamum indicum* are used in the same manner.

FRACTURE. (*Fractura*; from *frango*, to break.) *Catagma. Cassis. Clasma. Agme.* A Fracture is a solution of a bone into two or more fragments. A *simple fracture* is when the bone only is divided. A *compound fracture* is a division of the bone, with a laceration of the integuments, the bone mostly protruding. A fracture is also termed *transverse, oblique, &c.* according to its direction.

FRÆNULUM. (Dim. of *frenum*, a bridle.) The cutaneous fold, under the apex of the tongue, that connects the tongue to the infralingual cavity. It is sometimes, in infancy, so short as to prevent the child from sucking, when it is necessary to cut it, in order to give more room for the motion of the tongue.

FRÆNUM. The membranous fold which connects the præpuce to the inferior part of the glans penis.

FRAGA. (From *frago*, to smell sweet. The strawberry. See *Fragaria*.)

FRAGARIA. (From *frago*, to smell sweet.) The strawberry.

1. The name of a genus of plants in the Linnæan system. Class, *Icosandria*. Order, *Polygynia*.

2. The pharmacopœial name of the strawberry. *Fraga*. The mature fruit of the *Fragaria vesca, fragellis reptantibus* of Linnæus, was formerly recommended in gouty and calculous affections, in consequence, it would appear, of its efficacy in removing tartar from the teeth, which it is said to do very effectually.

FRAGARIA STABILLIS. *Barren strawberry.* Astrigent.

FRAGARIA VESCA. The systematic name of the strawberry. See *Fragaria*.

FRAGILITAS OSSUM. *Frailty of the bones.* Brittleness of the bones.

FRASMEN. *Fragmentum.* A splinter of a bone.

FRAMBŒSIA. (From *framboise*, Fr. for a raspberry.) The yaws. A genus of disease arranged by Cullen in the class *cachexie*, and order *impetigines*. It is somewhat similar in its nature to the lues venerea, and is endemic to the Antilla islands. It appears with excrescences, like mulberries, growing out of the skin in various parts of the body, which discharge an ichorous fluid.

FRANGULA. (From *frango*, to break, so called because of the brittleness of its branches.) Black alder. This officinal tree is the *Rhamnus frangula* of Linnæus:—*inermis floribus monogynis hermaphroditis, foliis integerrimis*. The berries and bark are used medicinally as strong purgatives. The former are often substituted for those of the buckthorn; the latter, which is the internal bark, and of a yellow colour, is mostly employed by the common people in dropsical and other disorders.

Frankincense. See *Thus*.

FRAXINELLA. (From *fraxinus*, the ash, so called because its leaves resemble those of the ash.) See *Dictamnus albus*.

Fraxinella, white. See *Dictamnus albus*.

FRAXINUS. (*A fragore*, from the noise its seeds make when shaken by the wind; or from *qæsar*, a hedge, because of its use in forming hedges.) The ash.

1. The name of a genus of plants in the Linnæan system. Class, *Polygamia*. Order, *Diœcia*. The ash.

2. The pharmacopœial name of the ash-tree, called also *brumeli* and *bumelia*. The bark of this tree, *Fraxinus excelsior* of Linnæus:—*foliis serratis floribus upetalis*, when fresh, has a moderately strong bitterish taste. It possesses resolvent and diuretic qualities, and has been successfully exhibited in the cure of intermittents. The seeds are occasionally exhibited medicinally as diuretics, in the dose of a drachm. In warm climates, a species of manna exudes from this species of fraxinus.

FRAXINUS EXCELSIOR. The systematic name of the ash-tree. See *Fraxinus*.

FRAXINUS ORNUS. The systematic name of the tree from which manna flows. See *Manna*.

FRAXINUS ROTUNDIFOLIA. The systematic name of a tree which affords manna. See *Manna*.

FRENA. The sockets of the teeth.

FRIGIDARIUM. (From *frigidus*, cold.) The cold bath.

FRIGORANA. A putrid fever.

FRONS. The forehead. The part between the eyebrows and the hairy scalp.

Frontal bone. See *Os frontis*.

Frontal sinus. See *Os frontis*.

FRONTALIS. See *Occipitis frontalis*.

FRONTALIS VERUS. See *Corrugator supercilli*.

FRONTIS OS. See *Os frontis*.

FRUCTUS HORREI. See *Fruits, summer*.

FRUITS, SUMMER. *Fructus horrei.* Under this term physicians comprehend strawberries, cherries, currants, mulberries, raspberries, and the like. They possess a sweet sub-acid taste, and are exhibited as dietetic auxiliaries, as refrigerants, antiseptics, attenuants, and aperients. Formerly they were exhibited medicinally in the cure of putrid affections, and to promote the alvine and urinary excretions. Considering them as an article of diet, they afford little nourishment, and are liable to produce flatulencies. To persons of a bilious constitution and rigid fibres, and where the habit is disposed naturally, or from extrinsic causes, to an inflammatory or putrescent state, their moderate, and even plentiful, use is salubrious; by those of a cold inactive disposition, where the vessels are lax, the circulation languid, and the digestion weak, they should be used very sparingly. The juices extracted from these fruits, by expression, contain their active qualities, freed from their grosser indigestible matter. On standing, the juice ferments and changes to a vinous or acetous state. By proper addition of sugar, and by boiling, their fermentive power is suppressed, and their medicinal qualities preserved. The juices of these fruits, when purified from their feculencies by settling and straining, may be made into syrups, with a due proportion of sugar, in the usual way.

FRUMENTACEOUS. A term applied to all such plants as have a conformity with wheat, either with respect to their fruit, leaves, or ears.

FUCUS. The name of a genus of plants in the Linnæan system. Class, *Cryptogamia*. Order, *Alga*.

FUCUS DIGITATUS. This fucus grows upon stones and rocks in the sea near the shore. It has several plain, long leaves or sinuses springing from a round stalk, in the manner of fingers when extended. It affords soda.

FUCUS ESCULENTUS. Edible fucus. Mr. Hudson has made this a distinct species, but Linnæus included it under his *saccharinus*. It grows plentifully in the sea, near the shores of Scotland, and also those of Cumberland. It has a broad, plain, simple, sword-shaped leaf, springing from a pinnated stalk.

FUCUS HELMINTOCORPTON. See *Corralina coriicana*.

FUCUS PALMATUS. Handed fucus. This grows in the sea, and consists of a thin lobed leaf like a hand.

FUCUS SACCHARINUS. Sea-belts; so

called from the supposed resemblance of its leaves to a belt or girdle. It grows upon rocks and stones by the sea shore. The leaves are very sweet, and when washed and hanged up to dry will exude a substance like that of sugar, from whence it was named.

FUCUS VESICULOSUS. The systematic name of the sea oak. See *Quercus marina*.

FULIGO. (*Quasi fumidigo*, from *fumus*, smoke.) *Arazos. Asper. Ascoli.* Soot: Wood soot, *fuligo ligni*, or the condensed smoke from burning wood, has a pungent, bitter, and nauseous taste, and is resolved by chymical analysis into a volatile alkaline salt, and empyreumatic oil, a fixed alkali, and an insipid earth. The tincture prepared from this substance, *tinctura fuliginis*, is recommended as a powerful antispasmodic in hysterical affections.

FULMINATION. A quick and lively explosion of bodies, such as takes place with fulminating gold, fulminating powder, and in the combustion of inflammable gas and vital air.

FUMARIA. (From *fumus*, smoke, from its juice, when dropped into the eye, producing the same sensations as smoke.)

1. The name of a genus of plants in the Linnæan system. Class, *Diadelphica*. Order, *Decandria*. Fumitory.

2. The pharmacopœial name of the common fumitory. *Fumia terre. Capnos. Herba melancholicifuga.* The leaves of this indigenous plant, *Fumaria officinalis* of Linnæus:—*pericarpium monocarpium racemosum, caule diffuso*, are directed for medicinal use by the Edinburgh college; they are extremely succulent, and have no remarkable smell, but a bitter, somewhat saline taste. The infusion of the dried leaves, or the expressed juice of the fresh plant, is esteemed for its property of clearing the skin of many disorders of the leprous kind.

FUMARIA RUBROSA. See *Aristolochia Jabacea*.

FUMARIA OFFICINALIS. The systematic name of the fumitory. See *Fumaria*.

FUMIGATION. The application of fumes, either from metallic or other preparations, to particular parts of the body, as those of the mercurial kind to venereal sores, &c.

Fumitory. See *Fumaria*.

FUMUS ALBUS. Mercury.

FUMUS CITRINUS. Sulphur.

FUMUS DUPLEX. Sulphur and mercury.

FUMUS RUBENS. Orpiment.

FUNCTION. See *Action*.

FUNGUS. 1. Proud flesh. A term in surgery to express any luxuriant formation of flesh.

2. The name of an order of plants in the Linnæan system, belonging to the *Cryptogamia* class.

FUNGUS IGNARIUS. See *Agaricus*.

FUNGUS LARICIS. See *Agaricus albus*.

FUNGUS MELITENSIS. This is impro-

perly called a fungus, it being the *Cinomorium coccineum* of Linnæus, a small plant which grows only on a little rock adjoining Malta. A drachm of the powder is given for a dose in dysenteries and hæmorrhages, and with remarkable success.

FUNGUS ROSACEUS. See *Bedeguar*.

FUNGUS SALICIS. The willow fungus. The species of fungus ordered in some pharmacopœias by this name is the *Boletus suævolens*; *acaulis superne leviss*, *salicibus*, of Linnæus, and the *Boletus albus* of Hudson. When fresh, it has a suburinous smell, and at first an acid taste, followed by a bitter. It is seldom used at present, but was formerly given in phthical complaints.

FUNGUS SAMBUCINUS. See *Auricula Jude*.

FUNGUS VINOSUS. The dark cobweb-like fungus, which vegetates in dry cellars, where wine, ale, and the like, are kept.

FUNICULUS UMBILICALIS. (*Funiculus*; dim. of *funis*, a cord.) See *Umbilical cord*.

FUNIS. A rope or cord.

FUNIS UMBILICALIS. See *Umbilical cord*.

FURCALA. The clavicle or collar-bone.

FURCELLA INFERIOR. The ensiform cartilage.

FURFUR. Bran. A disease of the skin, in which the cuticle keeps falling off in small scales like bran.

FURFURACEOUS. (*Furfuraceus*, from *furfur*, bran.) A term applied to the sediment deposited in the urine of persons afflicted with fever, of a reddish or whitish matter, which generally appears within an hour or two after the urine is passed, and

only falls in part to the bottom, the urine remaining turbid.

FURNACE. *Furnus*. The furnaces employed in chemical operations are of three kinds: 1. The *evaporatory furnace*, which has received its name from its use; it is employed to reduce substances into vapour by means of heat, in order to separate the more fixed principles from those which are more ponderous, and were mixed, suspended, compounded, or dissolved in the fluid. 2. The *reverberatory furnace*, which name it has received from its construction, being appropriated to distillation. 3. The *forge furnace*, in which the current of air is determined by bellows.

FUROR UTERINUS. (From *furo*, to be mad, and *uterus*, the womb.) *Acra*. *Brachura*. *Estromania*. *Arascon*. *Aratum*. See *Nymphomania*.

FURUNCULUS. (From *furo*, to rage; so named from its heat and inflammation before it suppurates.) *Dothetis* of Paracelsus. *Chiadus*. *Chieti*. A boil. An inflammation of a subcutaneous gland, known by an inflammatory tumour that does not exceed the size of a pigeon's egg.

FUSION. (*Fusio*; from *fundo*, to pour out.) A chymical process, by which bodies are made to pass from the solid to the fluid state, in consequence of the application of heat. The chief objects susceptible of this operation are salts, sulphur, and metals. Salts are liable to two kinds of fusion: the one, which is peculiar to saline matters, is owing to water, and is called *aqueous fusion*; the other, which arises from the application of fire, is known by the name of *igneous fusion*.

G.

GABIANUM OLEUM. See *Petroleum rubrum*.

GABIREA. A fatty kind of myrrh, mentioned by Dioscorides.

GALACTIA. (From γάλα, milk.) *Galactirrhœa*. An excess or overflowing of the milk.

GALACTINA. (From γάλα, milk.) Aliment prepared of milk.

GALACTIRRHŒA. (From γάλα, milk, and ρεω, to flow.) See *Galactia*.

GALACTODES. (From γάλα, milk.) In Hippocrates it signifies both milk warm, and a milky colour.

GALACTOPHORA MEDICAMENTA. (From γάλα, milk, and φεω, to carry.) Medicines which increase the milk.

GALACTOPHOROUS DUCTS. (*Ga-*

lactophorus; from γάλα, milk, and φεω, to carry, because they bring the milk to the nipple.) The excretory ducts of the glands of the breasts of women, which terminate in the papilla, or nipple, are so called.

GALACTOPOETICA. (From γάλα, milk, and ποω, to make.) Milk-making, an epithet applied to the faculty of making milk.

GALACTOPOSIA. (From γάλα, milk, and ποω, to drink.) The method of curing diseases by a milk diet.

GALANGA. (Perhaps its Indian name) Galangal. The roots of this plant are used medicinally; two kinds are mentioned in the pharmacopœias; the greater galangal *Kemferi galanga* of Linnæus, and the smaller galangal, the root of the *Maranta galanga*;

causino simplicifolia lanceolata subsessilibus, of Linnæus. The dried root is brought from China, in pieces from an inch to two in length, scarce half so thick, branched, full of knots and joints, with several circular rings of a reddish brown colour on the outside, and brownish within. It has an aromatic smell, not very grateful, and an unpleasant, bitterish, hot, biting taste. It was formerly much used as a warm stomachic bitter, and generally ordered in bitter infusions. It is now, however, seldom employed.

GALANGA MAJOR. See *Galanga*.

GALANGA MINOR. See *Galanga*.

Galangal. See *Galanga*.

Galangal, English. See *Cyperus*.

GALBANUM. (From *chalbanah*, Heb.) *Albetad*, *Chalbane*, *Gesor*. The plant is also named *Ferula Africana*; *Oreoselinum Africamum*; *Anisum fruticosum galbaniferum*; *Anisum Africamum frutescens*; and *Ayberzat*. Galbanum is the gummy-resinous juice, obtained partly by its spontaneous exudation from the joints of the stem of the Lovage-leaved bubon, *Bubon galbanum* of Linnæus:—*foliis rhombeis dentatis striatis glabris umbellis paucis*; but more generally, and in greater abundance, by making an incision in the stalk, a few inches above the root, from which it immediately issues, and soon becomes sufficiently concrete to be gathered. It is imported into England from Turkey and the East Indies, in large, softish, ductile, pale-coloured masses, which, by age, acquire a brownish yellow appearance: these are intermixed with distinct whitish tears, that are the most pure part of the mass. Galbanum has a strong unpleasant smell, and a warm, bitterish, acrid taste; "like the other gummy resins it unites with water, by trituration into a milky liquor, but does not perfectly dissolve, as some have reported, in water, vinegar or wine. Rectified spirit takes up much more than either of these menstrua, but not the whole: the tincture is of a bright golden colour. A mixture of two parts of rectified spirit, and one of water, dissolves all but the impurities, which are commonly in considerable quantity. In distillation with water, the oil separates and rises to the surface, in colour yellowish, in quantity one-twentieth of the weight of the galbanum. Galbanum, medicinally considered, may be said to hold a middle rank between assafoetida and ammoniacum; but its fœtidness is very inconsiderable, especially when compared with the former; it is therefore accounted less antispasmodic, nor are its expectorant qualities equal to those of the latter; it, however, is esteemed more efficacious than either in hysterical disorders. Externally it is often applied, by surgeons, to expedite the suppuration of inflammatory and indolent tumours, and,

by physicians, as a warm stimulating plaster. It is an ingredient in the *pilule galbani composite*, the *emplastrum galbani compositum* of the London Pharmacopœia, and in the *emplastrum ad clavos pedum* of the Edinburgh.

GALBEI. *Galbeum*. A sort of ornamental and medical bracelets worn by the Romans.

GALBULUS. (From *galbus*, yellow.) When the skin of the body is naturally yellow.

GALDA SUMMI. This is a gum-resin, mentioned by old writers, but totally forgot in the present day, and not to be obtained. Externally, it is of a brown colour, but white within, of a hard lamellated structure, and smells and tastes somewhat like elemi. When burnt it gives out an agreeable odour. It was formerly used as a warm, stimulating medicine, and applied in plasters as a strengthener.

GALEA. (From *γᾱλᾱ*, a cat, of the skin of which it was formerly made.) A helmet. In anatomy, the amnios is so called, because it surrounds the fœtus like a helmet. In surgery, it is a bandage for the head. A species of head-ach is so called, when it surrounds the head like a helmet.

GALEANTHROPIA. (This term seems to be from *γᾱλᾱ*, a cat, and *ἄνθρωπος*, a man.) It is a species of madness, in which a person imagines himself to be a cat, and imitates its manners.

GALEGA. (From *γᾱλᾱ*, milk; so named because it increases the milk of animals which eat it.) The name of a genus of plants in the Linnæan system. Class, *Dialphæa*. Order, *Decandria*.

2. The pharmacopœial name of the *Ruta praria*. Goats rue. From the little smell and taste of this plant, *Galega officinalis* of Linnæus: *leguminibus strictis, erectis; foliis lanceolatis, striatis, nudis*, it may be supposed to possess little virtues. In Italy the leaves are eaten amongst salads.

GALEGA OFFICINALIS. The systematic name of the goats rue. See *Galega*.

GALEZZ. A species of senna. The *casia tora* of Linnæus.

GALENA. (From *γᾱλᾱν*, to shine.) The name of an ore formed by the combination of lead with sulphur.

GALENIC MEDICINE. That practice of medicine which conforms to the rules of Galen, and runs much upon multiplying herbs and roots in the same composition, though seldom torturing them any otherwise than by decoction. It is opposed to chymical medicine, which by the force of fire, and a great deal of art, fetches out the virtues of bodies, chiefly mineral, into a small compass.

GALENTUM. (From *γᾱλᾱν*, galena.) A cataplasm, in the composition of which was the galena. In Paulus Ægineta it is considered as anodyne.

GALEODULON. See *Lamium album*.

GALBOREIA. (From *galon*, good, and *o-fa*, vision; so called because it was thought good for the sight; or from *galu*, a cat, and *o-fa*, aspect; the flowers gaping like the open mouth of that animal.) *Lamium rubrum*, *Urtica iners magna fetidissima* *Stachys fetida*. Hedge nettle. See *Lamium album*.

GALERICULUM APONEUROTICUM. The tendinous expansion which lies over the pericranium.

GALIUM. (From *gala*, milk; some species having the property of coagulating milk.) 1. The name of a genus of plants in the Linnean system. Class, *Tetrandria*. Order, *Monogynia*.

2. The pharmacopoeial name of the herb cheese rennet, or ladies bedstraw. The tops of this plant, *Galium verum*; *edlis octomis*, *linearibus*, *sulcatis*; *ramis floriferis*, *brevibus*, of Linnaeus, were long used as an efficacious medicine in the cure of epilepsy, but, in the practice of the present day, they are abandoned. Indeed, from the sensible qualities of the plant, little can be expected. The leaves and flowers possess the property of curdling milk; it is on that account styled cheese rennet.

3. A name for madder.

GALIUM ALBUM. The greater ladies bedstraw. * This herb, *Galium mollugo-folids octomis*, *ovato-linearibus*, *suberratis*, *patentissimis*, *mucronatis*; *caule flaccido*, *ramis patentibus*, of Linnaeus, with its flowers, are used medicinally. Five ounces, or more, of the expressed juice, taken every evening upon an empty stomach, is said to cure epilepsy.

GALIUM APARINE. The systematic name of the gobse-grass. See *Aparine*.

GALIUM MULLUGO. The systematic name of the greater ladies bedstraw. See *Galium album*.

GALIUM VERUM. The systematic name of the true ladies bedstraw. See *Galium*.

Gall. See *Bile*.

GALL SICKNESS. A popular name for the remitting fever occasioned by marsh miasmata, in the Netherlands, and which proved so fatal to thousands of the English soldiers after the capture of Walcheren in the year 1809. Dr. Lind informs us, that at Middleburg, the capital of Walcheren, a sickness generally reigns towards the latter end of August, or the beginning of September, which is always most violent after hot summers. It commences after the rains which fall in the end of July; the sooner it begins the longer it continues, and it is only checked by the coldness of the weather. Towards the end of August and the beginning of September, it is a continual burning fever, attended with a vomiting of bile, which is the *gall sickness*. This fever, after continuing three or four days, intermits, and

assumes the form of a double tertian; leaving the patient in a fortnight, or perhaps sooner. Strangers, that have been accustomed to breathe a dry, pure air, do not recover so quickly. Foreigners, in indigent circumstances, such as the Scots and German soldiers, who were garrisoned in the adjacent places, were apt, after those fevers, to have a swelling in the legs, and a dropsy; of which many died.

These diseases are the same with the double tertians common within the tropics. Such as are seized with the gall sickness, have at first some flushes of heat over the body, a loss of appetite, a white, foul tongue, a yellow tinge in the eyes, and a pale colour of the lips. Such as live well, drink wine, and have warm clothes, and a good lodging, do not suffer so much during the sickly season as the poor people; however, these diseases are not infectious, and seldom prove mortal to the natives.

Sir John Pringle observes, that the prevailing epidemic of autumn, in all marshy countries, is a fever of an intermitting nature, commonly of a tertian form, but of a bad kind; which, in the dampest places and worst seasons, appears as a double tertian, a remitting, or even an ardent fever. But, however these may vary in their appearance, according to the constitution of the patient, and other circumstances, they are all of a similar nature. For though, in the beginning of the epidemic, when the heat, or rather the putrefaction in the air, is the greatest, they assume a continued or a remitting form; yet, by the end of autumn, they usually terminate in regular intermittents.

But although, in the gall-sickness, there is both a redundancy and a deprivation of the bile, still the disease cannot, with justice, be said to originate wholly from that cause. It is certain, however, that the disease may be continued, and the symptoms aggravated, by an increased secretion and putrefaction of the bile, occasioned by the fever. In proportion to the coolness of the season, or the height and dryness of the ground, this disease is milder, remits and intermits more freely, and removes further from the nature of a continued fever. The higher ranks of people in general are the least liable to the diseases of the marshes; for such countries require dry houses, apartments raised above the ground, moderate exercise, without labour, in the sun, or evening damps; a just quantity of fermented liquors, plenty of vegetables, and fresh meats. Without such helps, not only strangers, but the natives themselves, are sickly, especially after hot and close summers. The hardiest constitutions are very little excepted more than others; and hence the British in the Netherlands have always been subject to fevers.

By this disease the British troops were harassed throughout the war, from 1743 to 1747. It appeared in the month of August, 1743: the paroxysms came on in the evening, with great heat, thirst, a violent head-ache, and often a delirium. These symptoms lasted most of the night, but abated in the morning, with an imperfect sweat; sometimes with an hæmorrhage of the nose, or looseness. The stomach, from the beginning, was disordered, with a nausea and sense of oppression; frequently with a bilious and offensive vomiting. If evacuations were either neglected, or too sparingly used, the patient fell into a continued fever, and sometimes grew yellow, as in jaundice. When the season was further advanced, this fever was attended with a cough, rheumatic pains, and sizzly blood. The officers, being better accommodated than the common men, and the cavalry, who had cloaks to keep them warm, were not so subject to it; and others who belonged to the army, but lay in quarters, were least of all effected; and the less in proportion to their being exposed to heats, night damps, and the other fatigues of the service. In this manner did the remitting fever infest the army for the remaining years of the war: and that exactly in proportion to their distance from the marshy places, of which we have several notable instances in Pringle's observations.

GALLA. (From *Gallus*, a river in Bythenia.) *Nux galla*. *Galla maxima orbiculata*. The gall-nut. The production of the *quercus cerris* of Linnaeus. By this name is usually denoted any protuberance, tubercle, or tumour, produced by the puncture of insects on plants and trees of different kinds. These galls are of various forms and sizes, and no less different with regard to their internal structure. Some have only one cavity, and others a number of small cells, communicating with each other. Some of them are as hard as the wood of the tree they grow on; whilst others are soft and spongy; the first being termed gall-nuts, and the latter berry-galls, or apple-galls.

The gall is thus produced:—the cynips *quercus folii*, an insect of the fly kind, deposits its eggs in the leaves and other tender parts of the tree. Around each puncture an excrescence is presently formed, within which the egg is hatched, and the insect passes through all the stages of its metamorphosis, until it becomes a perfect insect, when it eats its way out of its prison. The best oak-galls are heavy, knotted, and of a blue colour, and are obtained from Aleppo. They are nearly entirely soluble in water, with the assistance of heat. This soluble active matter consists of tannin, in combination with gallic acid; nine-tenths of the former, with one-tenth of the latter.

Oak-galls are supposed to be the strongest astringent in the vegetable kingdom. Both water and spirit take up nearly all their virtue, though the spirituous extract is the strongest preparation. The powder is, however, the best form; and the dose is from a few grains to half a drachm.

They are not much used in medicine, though they are said to be beneficial in intermittents. Dr. Cullen has cured agues, by giving half a drachm of the powder of galls every two or three hours during the intermission; and by it alone, or joined with camomile flowers, has prevented the return of the paroxysms. But the doctor states the amount of his results only to be this: that, "in many cases, the galls cured the intermittents; but that it failed also in many cases, in which the Peruvian bark afterwards proved successful." A fomentation, made by macerating half an ounce of bruised galls in a quart of boiling water for an hour, has been found useful for the piles, the prolapsus ani, and the fluor albus, applied cold. An injection, simply astringent, is made by diluting this fomentation, and used in gleet and leucorrhœa. The camphorated ointment of galls has been found also serviceable in piles, after the use of leeches; and is made by incorporating half a drachm of camphor with one ounce of hog's lard, and adding two drachms of galls in very fine powder. In fact, galls may be employed for the same purposes as oak bark, and are used under the same forms.

GALLA TURCICA. See *Quercus*.

GALL BLADDER. *Venicula fells*. An oblong membranous receptacle, situated under the liver, to which it is attached in the right hypochondrium. It is composed of three membranes: a common, fibrous, and villous. Its use is to retain the bile which regurgitates from the hepatic duct, there to become thicker, more acrid, and bitter, and to send it through the cystic duct, which proceeds from its neck into the ductus communis choledochus, to be sent on to the duodenum.

GALL-STONES. Biliary concretions. Hard concrete bodies, of which there are great varieties, formed in the gall-bladder of animal bodies. Gall-stones often lie quiet; so that, until dissection after death, some are never known to exist; but when they are prevented from passing through the gall ducts, they obstruct the passage of the bile into the intestines, and produce also many inconvenient symptoms, particularly the jaundice.

The diagnostics of this disorder are generally very obscure and uncertain; for other causes produce the same kind of symptoms as those which occur in this disease. The usual symptoms are, a loss of appetite, a sense of fullness in the stomach,

sickness and vomiting, langour, inactivity, sleepiness; and if the obstruction continues for a time, there is wasting of the flesh; yellowness of the eyes, skin, and urine; whitish stools; a pain in the pit of the stomach; whilst the pulse remains in its natural state. The pain excited by an obstruction of the gall ducts, in consequence of gall-stones passing through them, and this not affecting the pulse, is considered as the leading pathognomonic symptom. This pain, in some, is extremely acute, in others there is only a slight uneasiness felt about the region of the liver; but its particular seat is the gall duct, just where it enters the duodenum. In some patients there is no yellowness of the skin; in others it exists for several months. There is no disease more painful than this, in some instances; it is as frequent as any other affection of the liver; it admits of much relief from medicine, and is not immediately dangerous to the patient.

GALLIC ACID. A peculiar acid which is extracted from the nut-gall that grows on the oak. This acid is also found, in a greater or less quantity, in all sour or astringent vegetable substances.

GALLICTRICHIS. Corrupted from *callitrichis*, or *callitricum*.

GALLICUS MORBUS. The French disease. See *Dues venerea*.

GALLINAGO. (Diminutive of *gallus*, a cock.) 1. The woodcock.

2. An eminence within the prostate gland is called *caput gallinaginis*, from its fancied resemblance to a woodcock's head.

GALLIUM. See *Galium*.

GALVANISM. A professor of anatomy, in the university of Bologna, was one day making experiments on electricity, in his laboratory; near the machine were some frogs that had been flayed; the limbs of which became convulsed every time a spark was drawn from the apparatus. Galvani, surprised at this phenomenon, made it a subject of investigation, and discovered that metals, applied to the nerves and muscles of these animals, occasioned powerful and sudden contractions, when disposed in a certain manner. He gave the name of animal electricity to this order of new phenomena, from the analogy that he considered existing between these effects and those produced by electricity.

The name animal electricity has been superseded, notwithstanding the great analogy that exists between the effects of electricity and of Galvanism, in favour of the latter term; which is not only applicable to the generality of the phenomena, but likewise serves to perpetuate the memory of the discoverer.

In order to give rise to Galvanic effects, it is necessary to establish a communication

between two points of one series of nervous and muscular organs. In this manner a circle is formed, one arch of which consists of the animal parts, rendered the subject of experiment, while the other arch is composed of excitatory instruments, which generally consist of several pieces, some placed under the animal parts called supporters, others destined to establish a communication between the latter are called conductors. To form a complete Galvanic circle, take the thigh of a frog, deprived of its skin; detach the crural nerve, as far as the knee; put it on a piece of zinc; put the muscles of the leg on a piece of silver; then finish the excitatory arch, and complete the Galvanic circle by establishing a communication by means of the two supporters, by means of iron or copper wire, pewter, or lead. The instant that the communicators touch the two supporters, a part of the animal arch formed by the two supporters will be convulsed. Although this disposition of the animal parts, and of Galvanic instruments, be most favourable to the development of the phenomena, yet the composition of the animal and excitatory arch may be much varied. Thus contractions are obtained, by placing the two supporters under the nerve, and leaving the muscle out of the circle, which proves that nerves essentially constitute the animal arch.

It is not necessary for nerves to be entire, in order to produce contractions. They take place whether the organs be tied or cut through, provided there exists a simple contiguity between the divided ends. This proves that we cannot strictly conclude from what happens in muscular action, from that which takes place in Galvanic phenomena; since, if a nerve be tied or divided, the muscles on which the power is distributed lose the power of action.

The cuticle is an obstacle to Galvanic effects; they are always feebly manifested in parts covered by it. When it is moist, fine, and delicate, the effect is not entirely interrupted. Humboldt, after having detached the cuticle from the posterior part of the neck and back, by means of two blisters, applied plates of metal to the bare cutis, and at the moment of establishing a communication, he experienced sharp prickings, accompanied with a serosanguinous discharge.

If a plate of zinc be placed under the tongue, and a flat piece of silver on its superior surface, on making them touch each other, an acerb taste will be perceived, accompanied with a slight trembling.

The excitatory arch may be constructed with three, two, or even one metal only, with alloys, amalgams, or other metallic or mineral combinations, carbonated substances, &c. It is observed that metals

which are in general the most powerful exciters, induce contractions so much the more as they have an extent of surface. Metals are all more or less excitants; and it is observed that zinc, gold, silver, pewter, are of the highest rank; then copper, lead, nickel, antimony, &c.

Galvanic susceptibility, like muscular irritability, is exhausted by too long continued exercise, and is recruited by repose. Immersion of nerves and muscles in alcohol and opiate solutions diminishes and even destroys this susceptibility, in the same manner, doubtless, as the immoderate use of these substances in the living man blunts, and induces paralysis in muscular action. Immersion in oxygenated muriatic acid revests the fatigued parts, in being acted on by the stimulus. Animals killed by the repeated discharge of an electric battery, acquire an increase of Galvanic susceptibility; and this property subsists unchanged in animals destroyed by submersion in mercury, pure hydrogen gas, azot, and ammoniac; and finally, it is totally annihilated in animals suffocated by the vapour of charcoal.

Galvanic susceptibility is extinct in the muscles of animals of warm blood, in proportion as vital heat is dissipated; sometimes even when life is terminated in convulsions, contractility cannot be put into action, although warmth be not completely gone, as though the vital property were consumed by the convulsion, amidst which the animals had expired. In those of cold blood, on the contrary, it is more durable. The thighs of frogs, long after being separated from every thing, and even to the instant of incipient putrefaction, are influenced by Galvanic stimuli, doubtless, because irritability, in these animals, is less intimately connected with respiration, and life more divided among the different organs, which have less occasion to act on each other for the execution of its phenomena. The Galvanic chain does not produce sensible actions (that is contractions) until the moment it is completed, by establishing a communication with the parts constituting it. During the time it is complete, that is, through the whole space of time that the communication remains established, every thing remains tranquil; nevertheless, Galvanic influence is not suspended: in fact excitability is evidently increased, or diminished, in muscles that have been long continued in the Galvanic chain, according to the difference of the reciprocal situation of the connecting metals.

If silver has been applied to nerves, and zinc to muscles, the irritability of the latter increases in proportion to the time they have remained in the chain. By this method, the thighs of frogs have been revived in some degree, and afterwards be-

came sensible to stimuli, that before had ceased to act on them. By distributing the metals in an inverse manner, applying zinc to nerves and silver to muscles, an effect absolutely contrary is observed; and the muscles that possessed the most lively irritability when placed in the chain, seem 'to be rendered entirely paralytic if they remain long in this situation.

This difference evidently depends on the direction of the Galvanic fluid, determined towards the muscles or nerves, according to the manner in which these metals are disposed, and this is of some importance to be known for the application of Galvanic means to the cure of diseases.

Galvanic Pile.

M. Volta's apparatus is as follows:—

Raise a pile, by placing a plate of zinc, a flat piece of wet card, and a plate of silver, successively; then a second piece of zinc, &c. until the elevation is several feet high; for the effects are greater in proportion to its height; then touch both extremities of the pile, at the same instant, with one piece of iron wire; at the moment of contact, a spark is excited from the extremities of the pile, and luminous points are often perceived at different heights, where the zinc and silver come into mutual contact. The zinc end of this pile appears to be negatively electrified; that formed by the silver, on the contrary, indicates marks of positive electricity.

If we touch both extremities of the pile, after having dipped our hands into water, or, what is better, a saline solution, a commotion, followed by a disagreeable pricking in the fingers and elbow, is felt.

If we place in a tube filled with water, and hermetically closed by two corks, the extremities of two wires of the same metal which are in contact at the other extremity, one with the summit, the other with the base of the pile; these ends, even when separated only by the space of a few lines, experience evident changes at the instant the extremities of the pile are touched; the wire in contact with that part of the pile composed of zinc becomes covered with bullæ of hydrogen gas; that which touches the extremity formed by silver, becomes oxydated. Fourcroy attributes this phenomenon to the decomposition of water, by the Galvanic fluid, which abandons the oxygen to the iron that touches the positive extremity of the pile; then conducts the other gas invisibly to the end of the other wire, there to be disengaged.

From the numerous experiments of Mr. Davy, many new and important facts have been established, and Galvanism has been found one of the most powerful agents in chymistry; by its influence, platina wire has been melted; gold, silver, copper, and most of the metals, have easily been burnt; the fixed alkalis, and many of the earths,

have been made to appear as consisting of a metallic base, and oxygen; compound substances, which were before extremely difficult to decompose, are now, by the aid of Galvanism, easily resolved into their constituents.

GAMANDRA. See *Gambogia*.

GAMBIENSE GUMMI. See *Kino*.

Gamboge See *Gambogia*.

GAMBOGIA. (From the province of *Cambaya*, in the East Indies; called also *Cambodia* and *Cambogia*; hence it has obtained its name of *Cambadium*, *Cambegium*, *Gambogia*, *Gambogium*.) *Gamboge*. From its supposed virtues, it is called *gummi ad podagram*; *gammī gutta*; and, by corruption, *gotta*, *gutta gamba*, *gamon*, *germandra catagemu*, *gamboidea*, &c.; and, from its gold colour, *chrysopus*; and from its purgative quality, *succus laxativus*, *succus Indicus purgans*; and *scammonium orientale*.

A concrete vegetable juice, the produce of two trees, both called, by the Indians, *Caracapulli*, and by Linnaeus, *Gambogia gutta*. It is partly of a gummy and partly of a resinous nature. It is brought to us chiefly from *Gambaja*, in the East Indies, either in form of orbicular masses, or of cylindrical rolls of various sizes; and is of a dense, compact, and firm texture, and of a beautiful yellow colour.

In medicine, it is chiefly used as a drastic purge; it operates powerfully both upwards and downwards. Some condemn it as acting with too great violence, while others are of a contrary opinion. The dose is from two to four grains, as a cathartic; from four to eight grains it proves emetic and purgative. The roughness of its operation is said to be diminished, by giving it in a liquid form, sufficiently diluted. Rubbed with almonds, from its want of taste, it is a convenient laxative for children.

It has been given in dropsy, with cream of tartar, to correct its operation. It has also been recommended by some, to the extent of fifteen grains, joined with an equal quantity of vegetable alkali, to destroy the tape-worm. This dose is ordered in the morning, and if the worm is not expelled in two or three hours, it is repeated even to the third time, with safety and efficacy. It is asserted that it has been given to this extent even in delicate habits. This is said to be the remedy alluded to by Van Swieten, which was employed by Dr. Herenchwand, and with him proved so successful in the removal of the *tania lata*. It is an ingredient, and probably the active one, in most of the nostrums for expelling tæniæ.

Dr. Cullen says, that, on account of the quick passage of gamboge through the intestines, he was induced to give it in small and frequently repeated doses, as three or four grains, rubbed with a little sugar, every three hours; and thus found it ope-

rate without griping, or sickness, and, in three or four exhibitions, evacuate a great quantity of water, both by stool and urine.

GAMBOSIUM. See *Gambogia*.

GAMBOIDEA. See *Gambogia*.

GAMMA. (From the letter Γ , *gamma*, which it resembles.) A surgical instrument for cauterising a hernia.

GAMPHRE. (From $\gamma\alpha\mu\phi\epsilon\sigma$, crooked.) The cheek. The jaw.

GANGAMON. (From $\gamma\alpha\gamma\alpha\mu\sigma$, a fishing-net, which it was said to resemble.) A name of the omentum. Some call the contexture of nerves about the navel by this name.

GANGLION. ($\gamma\alpha\gamma\gamma\lambda\iota\sigma$, a knot.) A knot.

1. In anatomy it is applied to a natural knot-like enlargement, in the course of a nerve.

2. In surgery it is an encysted tumour, formed in the sheath of a tendon, and containing a fluid like the white of an egg. It most frequently occurs on the back of the hand or foot.

GANGRENE. (From $\gamma\alpha\sigma$, to feed upon.) *Gangrena*. An incipient mortification, so named from its eating away the flesh. Authors have generally distinguished mortification into two stages; the first, or incipient one, they name *gangrene*, which is attended with a sudden diminution of pain in the place affected; a livid discoloration of the part, which, from being yellowish, becomes of a greenish hue; a detachment of the cuticle, under which a turbid fluid is effused; lastly, the swelling, tension, and hardness, of the previous inflammation subside, and, on touching the part, a crepitus is perceptible, owing to the generation of air in the gangrenous parts.

Such is the state to which the term *gangrene* is applied. When the part has become quite cold, black, fibrous, incapable of moving, and destitute of all feeling, circulation, and life; this is the second stage of mortification, termed, *sphacelus*. *Gangrene*, however, is frequently used synonymously with the word mortification. See *Mortification*.

GARABA. An Arabic name for the disorder called *Ægilops*.

GARCINIA MANGOSTANA. (Named in honour of Dr. Garcin, who accurately described it.) The systematic name of the mangosteen tree. See *Mangosteen*.

GARGALE. ($\gamma\alpha\gamma\alpha\lambda\epsilon$.) *Gargalee*. *Gargalismo*. Irritation, or stimulation.

GARGAREON. (Hebrew.) The uvula, or glandulous body, which hangs down into the throat.

GARGARISM. *Gargarismus*. A wash for the mouth and throat.

GARGARISMA. (From $\gamma\alpha\gamma\gamma\alpha\gamma\iota\sigma$, to gargle.) A gargle.

GARGARISUM. A gargle or wash for the throat.

GARGATEUM. A bed on which lunatics, &c. were formerly confined.

Gargle See *Gargarisma*.

Garlic. See *Allium*.

Garou bark. See *Thymalea*.

GARON. (γάρων.) A kind of pickle prepared of fish; at first it was made from a fish which the Greeks call *Garos*; but the best was made from mackerels. Among the moderns, *garum* signifies the liquor in which fish is pickled.

GARROTILLO. (From *garottar*, to bind closely. Span.) A name of the cynanche maligna, from its sense of strangulation, as if the throat were bound with a cord.

GAROPHYLLUS. See *Caryophyllus*.

GAS. (From *gascht*, a German word which means an eruption of wind.) See *Gaz*.

GASTRIC ARTERY. *Arteria gastrica*. The right, or greater gastric artery, is a branch of the hepatic; the left or lesser, a branch of the splenic.

GASTRIC JUICE. *Succus gastricus*. A fluid separated by the capillary exhalant arteries of the stomach, which open upon its internal surface. The œsophagus also affords a small quantity, especially in the inferior part. Modern philosophers have paid great attention to this fluid, and from their several experiments it is known to possess the following properties. It is the principal agent of digestion, and changes the aliments into a kind of uniform soft paste: it acts on the stomach after the death of the animal. Its effects shew that it is a solvent, but of that peculiar nature that it dissolves animal and vegetable substances, uniformly, and without exhibiting a stronger affinity for the one than for the other. Although it is the most powerful agent of digestion in the stomach, its dissolvent power has need of assistance from the action of several secondary causes, as heat, which seems to augment and concentrate itself in the epigastric region. So long as the exertion of the stomach continues, there is a sort of intestine fermentation, which should not, in its full sense, be compared to the motion by which fermentative and putrescent substances are decomposed; there is also a moderate and peristaltic motion of the muscular fibres of the stomach, which press the aliment on all sides, and perform a slight trituration, while the gastric moisture softens and macerates the food before it is dissolved. By many it has been considered merely as a ferment, but this cannot be the case. See *Digestion*.

It is one of the most powerful antiseptics with which we are acquainted; and, from the experiments of *Spallanzani*, *Scopoli*, *Arminati*, and others, its nature appears to be essentially different in the several classes of animals, as they have proved by analysis. The gastric juice of the human subject,

when healthy, is inodorous, of a saltish taste, and limpid, like water, unless it be a little tinged with the yellow colour of some bile, that has regurgitated into the stomach. In quantity it is very considerable, as must be evident from the extent of the surface of the stomach, and its continual secretion; but it is most copious when solicited by the stimulus of food. Besides the properties of this fluid before mentioned, it has others, which have induced physicians and surgeons to exhibit it medicinally. It cures dyspepsia and intermittent fever. Applied externally, in form of fomentation or poultice, it cures putrid and scrofulous ulcers in a wonderful manner; and it is to be regretted that its utility is not more generally known.

GASTRITIS. (From γαστήρ, the stomach.) Inflammation of the stomach. A genus of disease in the class *pyrexia*, and order *phlegmasia*, of Cullen. It is known by pyrexia, anxiety, heat, and pain in the epigastrium, increased when any thing is taken into the stomach, vomiting, hiccup, pulse small and hard, and prostration of strength. There are two species: 1. *Gastritis phlegmonoidea*, with an inflammatory tumour. *Gastritis erysipelatosa*, when the inflammation is of a creeping or erysipelatous nature.

Gastritis is produced by acrid substances of various kinds, such as arsenic, corrosive sublimate, &c. taken into the stomach, as likewise by food of an improper nature; by taking large draughts of any cold liquor when the body is much heated by exercise, or dancing; and repelled exanthemata and gout. Besides these, it may arise from an inflammation of some of the neighbouring parts being communicated to the stomach.

The erysipelatous gastritis arises chiefly towards the close of other diseases, marking the certain approach to dissolution, and being unaccompanied with any marks of general inflammation, or by any burning pain in the stomach.

The symptoms of phlegmonous gastritis, as observed above, are, a violent burning pain in the stomach, with great soreness, distention, and flatulency; a severe vomiting, especially after any thing is swallowed, whether it be liquid or solid; most distressing thirst; restlessness, anxiety, and a continual tossing of the body, with great debility, constant watching, and a frequent, hard and contracted pulse. In some cases, a severe purging attends.

If the disease increases in violence, symptoms of irritation then ensue; there is a great loss of strength, with faintings; a short and interrupted respiration; cold, clammy sweats, hiccups, coldness of the extremities, an intermittent pulse, and the patient is soon cut off.

The event of gastritis is seldom favour-

able, as the person is usually either suddenly destroyed by the violence of the inflammation, or else it terminates in suppuration, ulceration or gangrene.

If the symptoms are very mild, and proper remedies have been employed at an early period of the disease, it may, however, terminate in resolution, and that in the course of the first, or, at farthest, the second week.

Its termination in suppuration may be known by the symptoms, although moderate, exceeding the continuance of this period, and a remission of pain occurring, whilst a sense of weight and anxiety still remain; and, on the formation of an abscess, cold shiverings ensue, with marked exacerbations in the evening, which are followed by night sweats, and other symptoms of hectic fever; and these at length prove fatal, unless the pus is thrown up by vomiting, and the ulcer heals.

Its tendency to gangrene may be dreaded, from the violence of its symptoms not yielding to proper remedies early in the disease; and when begun, it may be known by the sudden cessation of the pain; by the pulse continuing its frequency, but becoming weaker; and by delirium, with other marks of increasing debility ensuing.

Fatal cases of this disease shew, on dissection, a considerable redness of the inner coat of the stomach, having a layer of coagulable lymph lining its surface. They likewise shew a partial thickening of the substance of the organ, at the inflamed part, the inflammation seldom extending over the whole of it. Where ulceration has taken place, the ulcers sometimes are found to penetrate through all its coats, and sometimes only through one or two of them.

GASTROCELE. (From *γαστήρ*, the stomach, and *κλῆν*, a tumour.) A hernia of the stomach, occasioned by a protusion of that viscus through the abdominal parietes.

GASTROCNEMIUS. (From *γαστήρ*, the stomach, and *κνήμης*, the leg.) The muscles of the foot, which form the calf or belly of the leg.

GASTROCNEMIUS EXTERNUS. *Gemellus*. This muscle, which is situated immediately under the integuments at the back part of the leg, is sometimes called *gemellus*; this latter name is adopted by Albinus. Winslow describes it as two muscles, which he calls *gastrocnemii*; and Douglas considers this and the following as a *quadriceps*, or muscle with four heads, to which he gives the name of *extensor tarsi suavis*. It is called *bi femoro calcaneus* by Dumas. The *gastrocnemius externus* arises by two distinct heads. The first, which is the thickest and longest of the two, springs by a strong thick tendon from the upper and back part of the inner condyle of the os femoris, adhering strongly to the capsular

ligament of the joint, between which and the tendon is a considerable *bursa mucosa*. The second head arises by a thinner and shorter tendon from the back part of the outer condyle of the os femoris. A little below the joint, their fleshy bellies unite in a middle tendon, and below the middle of the tibia they cease to be fleshy, and terminate in a broad tendon, which, a little above the lower extremity of the tibia, unites with that of the *gastrocnemius internus*, to form one great round tendon, sometimes called *chorda magna*, but more commonly, *tendo Achillis*.

GASTROCNEMIUS INTERNUS. *Tibio peronei calcaneus* of Dumas. This, which is situated immediately under the last described muscle, is sometimes named *soleus*, on account of its shape, which resembles that of the sole-fish. It arises by two heads. The first springs by tendinous and fleshy fibres from the posterior part of the head of the fibula, and for some way below it. The second arises from an oblique ridge at the upper and posterior part of the tibia, which affords origin to the inferior edge of the popliteus, continuing to receive fleshy fibres from the inner edge of the tibia for some way down. This muscle, which is narrow at its origin, spreads wider as it descends, as far as its middle; after which it becomes narrower again, and begins to grow tendinous, but its fleshy fibres do not entirely disappear till it has almost reached the extremity of the tibia, a little above which it unites with the last described muscle, to form the *tendo Achillis*. This thick round chord is inserted into the lower and posterior part of the os calcis, after sliding over a cartilaginous surface on that bone, to which it is connected by a tendinous sheath that is furnished with a large *bursa mucosa*.

Both the *gastrocnemii* have the same use, viz. that of extending the foot, by drawing it backwards and downwards.

GASTROCOLICA. (From *γαστήρ*, the stomach, and *κόλον*, the colon.) A term applied to a vein which proceeds from the stomach to the colon.

GASTRODYNIA. (From *γαστήρ*, the stomach, and *δύνη*, pain.) Pain in the stomach.

GASTRO-EPIPLOIC ARTERY. *Arteria gastrico-epiploica*. The branch of the greater gastric artery that runs to the epiploon.

GASTROGRAPHY. (*Gastrographie*; from *γαστήρ*, the stomach, and *γραφή*, a suture.) The sewing of wounds of the abdomen.

GASTROTOMIA. (From *γαστήρ*, the belly, and *τομή*, to cut.) The operation of cutting open the belly and uterus, as in the Cæsarian operation.

GAULE. See *Myrtus Brabantica*.

GAZ. (From *Gascht*, German, an eruption of wind.) *Gas*. Elastic fluid.

Aëriform fluid. Elastic vapour. By the word *gaz*, we distinguish a permanently elastic aëriform fluid, or substance which has the appearance of air; that is to say, it is transparent, elastic, ponderable, invisible, (oxygenated muriatic acid gas, and nitrous acid gas, are the only exceptions to this rule,) and not condensible into a liquid or solid state by any degree of cold hitherto known.

Some of the gases exist in nature, without the aid of art, and may therefore be collected; others, on the contrary, are only producible by artificial means.

All gases are combinations of certain substances, reduced to the gaseous form by the addition of caloric and light. It is, therefore, necessary to distinguish, in every gas, the matter of heat which acted the part of a solvent, and the substance which forms the basis of the gas.

Gases are not contained in those substances from which we obtain them in the state of gas, but owe their formation to the expansive property of caloric.

The bases of some gases are known to us, and may be exhibited in an uncombined state; others again are not producible by art.

Formation of Gases.

The different forms under which bodies appear depend upon a certain quantity of caloric, chymically combined with them. The very formation of gases corroborates this truth. Their production totally depends upon the combination of the particular substances with caloric; and those we call permanently elastic are only so, because we cannot so far reduce their temperature, as to dispose them to part with it; otherwise they would undoubtedly become fluid or solid.

Water, for instance, is a solid substance in all degrees below 32° of Fahrenheit's scale; above this temperature it combines with caloric, and becomes a fluid. It retains its fluidity under the ordinary pressure of the atmosphere, till its temperature is augmented to 212° . It then combines with a larger portion of caloric, and is converted, *apparently*, into gas, or at least into elastic vapour; in which state it would continue, if the temperature of our atmosphere was above 212° . Gases are therefore solid substances, between the particles of which a repulsion is established by the quantity of caloric.

But as in the gaseous water, or steam, the caloric is retained with but little force, on account of its quitting the water when the vapour is merely exposed to a lower temperature, we do not admit steam amongst the class of gases, or permanent aëriform elastic fluids. In gases, caloric is united by a very forcible affinity, and no diminution of temperature, or pressure,

that has ever yet been effected, can separate it from them. Thus the air of our atmosphere, in the most intense cold, still remains in the aëriform state; and hence is derived the essential characters of gases, namely; *that they shall remain aëriform, under all variations of pressure and temperatures.*

In the modern nomenclature, the name of every substance existing in the aëriform state is derived from its solid base; and the term gas is used to denote its existence in this state.

In order to illustrate the formation of gases, or to shew in what manner caloric is combined with them, the following experiment may serve. Put into a retort, capable of holding half a pint of water, two ounces of muriate of soda, (common salt; pour on it half its weight of sulphuric acid, and apply the heat of a lamp; a great quantity of gas is produced, which might be collected and retained over mercury. But to serve the purpose of this experiment, let it pass through a glass receiver, having two openings, into one of which the neck of the retort passes, whilst, from the other, a bent tube proceeds, which ends in a vessel of water. Before closing the apparatus, let a thermometer be included in the receiver, to shew the temperature of the gas. It will be found that the mercury in the thermometer will rise only a few degrees; whereas the water in the vessel which receives the bent tube will soon become boiling hot.

Explanation.—Common salt consists of muriatic acid, united to soda; on presenting sulphuric acid to this union, a decomposition takes place. The sulphuric acid unites by virtue of its greater affinity to the soda, and forms sulphate of soda, or Glauber's salt; the muriatic acid becomes therefore disengaged, and takes the gaseous form, in which it is capable of existing in our temperature. To trace the caloric during this experiment, as was our object, we must remark that it first flows from the lamp to the disengaged muriatic acid, and converts it into gas; but the heat thus expended is chymically united, and therefore not appreciable by the thermometer. The caloric, however, is again evolved, when the muriatic acid gas is condensed by the water, with which it forms liquid muriatic acid.

In this experiment we therefore trace caloric in a chymical combination producing gas; and from this union we again trace it in fire, or sensible heat.

Such, in general, is the cause of the formation and fixation of gases. It may be further observed, that each of these fluids loses or suffers the disengagement of different quantities of heat, as it becomes more or less solid in its new combination,

or as that combination is capable of retaining more or less specific heat.

The discovery of æriform gaseous fluids has occasioned the necessity of some peculiar instruments, by means of which those substances may be conveniently collected, and submitted to examination. The principal ones for that purpose are styled the *pneumatic apparatus*.

The Pneumatic Reservoir, or Cistern,

Is made either of wood or strong sheet-iron, tinned, japanned, or painted. A trough of about two feet long, sixteen inches wide, and fifteen high, has been found to be sufficient for most experiments. Two or three inches below its brim, a horizontal shelf is fastened, in dimension about half, or one-third part, of the width of the trough. In this shelf are several holes: these holes must be made in the centre of a small excavation, shaped like a funnel, which is formed in the lower part of the shelf.

This trough is filled with water sufficient to cover the shelf to the height of an inch.

The use of this shelf is to support receivers, jars, or bell-glasses, which, being previously filled with water, are placed invertedly, their open end turned down upon the above-mentioned holes, through which the gases conveyed there, and directed by means of the funnel-shaped excavations, rise in the form of air-bubbles into the receiver.

When the gaseous fluids are capable of being absorbed by water, as is the case with some of them, the trough must be filled with mercury. The price and gravity of this fluid make it an object of convenience and economy, that the trough should be smaller than when water is used.

A mercurial trough is best cut in marble, free-stone, or a solid block of wood. A trough about twelve inches long, three inches wide, and four deep, is sufficient for all private experiments.

Method of collecting Gases, and transferring them from one vessel to another.

If we are desirous of transmitting air from one vessel to another, it is necessary that the vessel destined to receive it be full of water, or some fluid heavier than air. For that purpose take a wide mouthed bell-glass, or receiver; plunge it under the water in the trough, in order to fill it; then raise it, with the mouth downwards, and place it on the shelf of the trough, so as to cover one or more of the holes in it.

It will now be full of water, and continue so as long as the mouth remains below the surface of the fluid in the cistern, for in this case, the water is sustained in the vessel by the pressure of the atmosphere, in the same manner as the mercury is sustained in the barometer. It may without difficulty be imagined, that if common air (or any

other fluid resembling common air in lightness and elasticity) be suffered to enter the inverted vessel filled with water, it will rise to the upper part, on account of its levity, and the surface of the water will sub-side. To exemplify this, take a glass, or any other vessel, in that state which is usually called *empty*, and plunge it into the water with its mouth downwards; scarce any of it will enter the glass, because its entrance is opposed by the elasticity of the included air; but if the vessel be turned with its mouth upwards, it immediately fills, and the air rises in bubbles to the surface. Suppose this operation be performed under one of the jars, or receivers, which are filled with water, and placed upon the perforated shelf, the air will ascend in bubbles as before, but, instead of escaping, it will be caught in the upper part of the jar, and expelled part of the water it contains.

In this manner we see that air may be emptied out of one vessel into another by a kind of inverted pouring, by which means it is made to ascend from the lower to the upper vessel. When the receiving vessel has a narrow neck, the air may be poured in a similar manner through an inverted funnel, inserted in its mouth.

If the air is to be transferred from a vessel that is stopped like a bottle, the bottle must be unstopped, with its orifice downwards in the water: and then inclined in such a manner that its neck may come under the perforated excavation of the shelf. The gas will escape from the bottle, and, passing into the vessel adapted to receive it, will ascend in it in the form of bubbles.

In whatever manner this operation is performed, the necessity of the excavation in the lower part of the shelf may be readily conceived. It is, as mentioned before, destined to collect the gas which escapes from the vessel, and direct it in its passage towards the vessel adapted to receive it. Without this excavation, the gas, instead of proceeding to the place of its destination, would be dispersed and lost.

The vessels, or receivers, for collecting the disengaged gases, should be glass cylinders, jars, or bell-glasses, of various sizes; some of them should be open at both ends, others should be fitted with necks at the top, ground perfectly level, in order that they may be stopped by ground flat pieces of metal, glass, slate, &c. others should be furnished with ground stoppers. Some should be graduated into cubic inches, and sub-divided into decimal or other equi-distant parts. Besides these, common glass bottles, tumblers, &c. may be used.

Classification of Gases.

All the elastic æriform fluids with which we are hitherto acquainted are generally divided, by systematic writers, into two

classes; namely, those that are *respirable*, and *capable of maintaining combustion* and those that are *not respirable*, and *incapable of maintaining combustion*. This division, indeed, has its advantage; but the term *respirable*, in its physico-logical application, has been very differently employed by different writers. Sometimes by the respirability of a gas has been meant its power of supporting life, when repeatedly applied to the blood in the lungs. At other times all gases have been considered *respirable*, which were capable of introduction into the lungs by voluntary efforts, without any relation to their vitality. In the last case, the word *respirable* seems to us most properly employed, and in this sense it is here used.

Non-respirable gases are those which, when applied to the external organs of respiration, stimulate the muscles of the epiglottis in such a manner as to keep it perfectly close on the glottis; thus preventing the smallest particle of gas from entering into the bronchia, in spite of voluntary exertions.

Of respirable gases, or those which are capable of being taken into the lungs by voluntary efforts, according to their conditions, only one has the power of uniformly supporting life, namely, atmospheric air; other gases, when respired, sooner or later impair the health of the human constitution, or perhaps occasion death; but in different modes.

Some gases effect *no positive* change in the blood; animals immersed in it die of a disease produced by the privation of atmospheric air, analogous to that occasioned by their submersion in water.

Others again produce *some positive* change in the blood, as appears from the experiments of Dr. Beddoes and Professor Davy. They seem to render it incapable of supplying the nervous and muscular fibres with principles essential to sensibility and irritability. These gases, therefore, destroy animal life on a different principle.

It is obvious, therefore, that the above classification does not hold good in all respects, but is capable of misleading the student.

Gas, azotic. See *Nitrogen*.

Gas, carbonic acid. This may be obtained by pouring any acid upon calcareous earth, which thereby becomes decomposed; the effused acid combines with the lime, and forms a new neutral salt, and the carbonic acid is disengaged and escapes in the form of a colourless gas, viz. carbonic acid gas. See *Carbonic acid*.

Gas-hepatic. See *Sulphurated hydrogen gas*.

Gas hydrogen. *Inflammable air.* See *Hydrogen*.

Gas, light carbonated hydrogen. See *Carbonated hydrogen gas, light*.

Gas, heavy carbonated hydrogen. See *Carbonated hydrogen gas, heavy*.

Gazeous oxyd of carbon. See *Carbon, gazeous oxyd of*.

GEISOMA. (From *γειον*, the eaves of the house.) *Geison*. The prominent parts of the eye-brows, which hang over the eyes like the eaves of a house.

GEISON. See *Geisoma*.

GELASINOS. (From *γελω*, to laugh.) An epithet for the four middle fore-teeth, because they are shewn in laughter.

GELASMUS. (From *γελω*, to laugh.) The Sardonian laugh.

GELATINE. Gelly, or jelly. An animal substance soluble in water, but not in alcohol; capable of assuming a well-known elastic or tremulous consistence, by cooling, when the water is not too abundant, and liquifiable again, by increasing its temperature. This last property remarkably distinguishes it from albumen, which becomes consistent by heat. It is precipitated in an insoluble form by tannin, and it is this action of tannin on gelatine that is the foundation of the art of tanning leather.

Jellies are very common in our kitchens; they may be extracted from all the parts of animals, by boiling them in water. Hot water dissolves a large quantity of this substance. Acids likewise dissolve them, as do likewise more particularly the alkalis. Jelly, which has been extracted without long decoction, possesses most of the characters of vegetable mucilage; but it is seldom obtained without a mixture of albumen.

Jellies, in a pure state, have scarcely any smell or remarkable taste. By distillation, they afford an insipid and inodorous phlegm, which easily putrefies. A stronger heat causes them to swell up, become black, and emit a fetid odour, accompanied with white acrid fumes. An impure volatile alkali, together with empyreumatic oil, then passes over, leaving a spongy coal, not easily burned, and containing common salt and phosphat of lime.

The jelly of various animal substances is prepared for the use of sea-faring persons under the name of portable soup. The whole art of performing this operation consists in boiling the meat, and taking the scum off, as usual, until the soup possesses the requisite flavour. It is then suffered to cool, in order that the fat may be separated. In the next place, it is mixed with five or six whites of eggs, and slightly boiled. This operation serves to clarify the liquid, by the removal of opaque particles, which unite with the white of egg at the time it becomes solid by the heat,

and are consequently removed along with it. The liquor is then to be strained through flannel, and evaporated, on the water-bath, to the consistence of a very thick paste; after which it is spread, rather thin, upon a smooth stone, then cut into cakes, and, lastly, dried in a stove, until it becomes brittle. These cakes may be kept four or five years, if defended from moisture. When intended to be used, nothing more is required to be done, than to dissolve a sufficient quantity in boiling water, which by that means becomes converted into soup.

GELATIO. (From *gelo*, to freeze.) Freezing; or that rigidity of the body which happens in a catalepsy, as if the person were frozen.

GEMELLUS. (From *gemimus* double, having a fellow.) See *Gastrocnemius* and *Gemini*.

GEMINI. *Gemelli* of Winslow. Part of the *marcipialis* of Cowper. *Ischio spini trochanterien* of Dumas. This muscle has been a subject of dispute among anatomists since the days of Vesalius. Some describe it as two distinct muscles, and hence the name it has gotten of *gemini*. Others contend that it ought to be considered as a single muscle. The truth is, that it consists of two portions, which are united together by a tendinous and fleshy membrane, and afford a passage between them to the tendon of the obturator internus, which they inclose as it were in a purse. These two portions are placed under the glutæus maximus, between the ischium and the great trochanter.

The superior portion, which is the shortest and thickest of the two, arises fleshy from the external surface of the spine of the ischium; and the inferior, from the tuberosity of that bone, and likewise from the posterior sacro-ischiatic ligament. They are inserted, tendinous and fleshy, into the cavity at the root of the great trochanter. Between the two portions of this muscle, and the termination of the obturator internus, there is a small *bursa mucosa*, connected to both, and to that part of the capsular of the joint which lies under the *gemini*.

This muscle assists in rolling the os femoris outwards, and prevents the tendon of the obturator internus from slipping out of its place while that muscle is in action.

GEMURSA. (From *gemo*, to groan; so called from the pain it was said to occasion in walking.) The name of an excrescence between the toes.

GENIAS. (From *γενει*, the cheek.) The downy hairs which first cover the cheek; also the name of a bandage mentioned by Galen, which covers the cheek, and comes under the chin.

GENERATION. Many ingenious hy-

potheses have been instituted by physiologists to explain the mystery of generation, but the whole of our knowledge concerning it appears to be built upon the phenomena it affords; and may be seen in the works of Haller, Buffon, Cruickshanks, and Haighton. It is a sexual action, performed in different ways in most animals; many of them have different sexes, and require conjunction: such are the human species, quadrupeds, and others. The females of quadrupeds have a matrix, separated into two cavities, *uterus bicornis*, and a considerable number of tests; they have no menstrual flux; most of them bear several young at a time, and the period of their gestation is generally short. The generation of birds is very different. The males have a strong genital organ, which is often double. The vulva in females is placed behind the anus; the ovaries have no matrices, and there is a duct for the purpose of conveying the egg from the ovarium into the intestines: this passage is called the oviduct. The eggs of pullets have exhibited unexpected facts to physiologists, who examined the phenomena of incubation. The most important discoveries are those of the immortal Haller, who found the chicken perfectly formed in eggs which were not fecundated. There is no determinate conjunction between fishes; the female deposits her eggs on the sands, over which the male passes, and emits its seminal fluid, doubtless for the purpose of fecundating them; these eggs are hatched after a certain time. The males of several oviparous quadrupeds have a double or forked organ. Insects exhibit all the varieties which are observed in other animals: there are some, indeed the greater number, which have the sexes in two separate individuals; among others, the reproduction is made either with or without conjunction, as in the vine fretter; one of these insects, confined alone beneath a glass, produces a great number of others. The organ of the male, in insects, is usually armed with two hooks, to seize the female: the place of these organs is greatly varied; with some it is at the upper part of the belly, near the chest, as in the female dragonfly; in others it is at the extremity of the antenna, as in the male spider. Most worms are hermaphrodite; each individual has both sexes. Polypi, with respect to generation, are singular animals: they are reproduced by buds, or offsets: a bud is separated from each vigorous polypus, which is fixed to some neighbouring body, and grows: polypi are likewise found on their surface, in the same manner as branches issue from plants. These are the principal modes of generation in animals. In the human species, which engages our attention more particularly, the phenomena

are as follow: the mode of congress of the man with the woman requires no description; but generation does not consist in that alone; there are certain states or conditions requisite for conception to take place. The ovum must have arrived at a state of maturity. There must be such a determination of blood to the uterus, that, together with the venereal stimulus, shall induce an action in the Fallopian tubes, by which the fimbriæ grasp the ovum that is to be impregnated. During this state of the parts, the semen virile must be propelled into the uterus, in order that its subtle and vivifying portion shall pass along the tube to the ovum. Fecundation having thus taken place, a motion is induced in the vivified ovum, which ruptures the tender vesicle that contains it; the fimbriæ of the Fallopian tube then grasp and convey it into the tube, which, by its peristaltic motion, conducts it into the cavity of the uterus, there to be evolved and brought to maturity, and, at the expiration of nine months, to be sent into the world.

GENERATION, FEMALE ORGANS OF. The parts subservient to generation, in a woman, are divided into external and internal.

The external parts are, the *mons veneris*, the *labia*, the *perineum*, the *clitoris*, and the *nymphae*. To these may be added the *meatus urinarius*, or orifice of the urethra. The *hymen* may be esteemed the barrier between the external and internal parts. The internal parts of generation are, the *vagina* and *uterus*, and its appendages.

GENERATION, MALE ORGANS OF. The parts which constitute the organs of generation in men are the *penis*, *testicles*, and *vesicula seminales*.

GENIO. (From *γενω*, the chin.) Names compounded of this word belong to muscles which are attached to the chin.

GENIO-HYO-GLOSSUS. (*Musculus genio-hyo-glossus*; from *γενω*, the chin, and *γλωσσα*, the tongue, so called from its origin in the chin, and insertion in the tongue.) *Genio glossus* of some authors. This muscle forms the fourth layer between the lower jaw and os hyoides. It arises from a rough protuberance in the inside of the middle of the lower jaw; its fibres run like a fan, forwards, upwards, and backwards, and are inserted into the top, middle, and root of the tongue, and base of the os hyoides near its corner. Its use is to draw the tip of the tongue backwards into the mouth, the middle downwards, and to render its back concave. It also draws its root and the os hyoides forwards, and thrusts the tongue out of the mouth.

GENIO-HYOIDEUS. (*Musculus genio-hyoideus*; from *γενω*, the chin, and *ωιδης*, the os hyoides; so called from its origin in the chin, and its insertion in the

os hyoides.) This muscle constitutes the third layer between the lower jaw and os hyoides. It is a long, thin, and fleshy muscle, arising tendinous from a rough protuberance at the inside of the chin, and growing somewhat broader and thicker as it descends backward, to be inserted by very short tendinous fibres into both the edges of the base of the os hyoides. It draws the os hyoides forwards to the chin.

GENIOPHARYNGÆUS. The constrictor pharyngis superior.

GENIPI ALBUM. The plant which bears this name, in the pharmacopœias, is the *Artemisia rupestris* of Linnæus:—*foliis pinnatis, caulibus adscendentibus; floribus globosis, cernuis; receptaculo papposo*. It has a grateful smell, and is used in some countries in the cure of intermittents and obstructed catamenia.

GENIPI VERUM. The plant directed for medicinal purposes, under this title, is the *Achillea; foliis pinnatis, pinnis simplicibus, glabris, punctatis*, of Haller. It has a very grateful smell, and a very bitter taste, and is exhibited in Switzerland in epilepsy, diarrhœa, and debility of the stomach.

GENISTA. (From *genu*, a knee; so called from the inflection and angularity of its twigs.) 1. The name of a genus of plants in the Linnæan system. Class, *Dia-delpnia*. Order, *Decandria*.

2. The pharmacopœial name of the common broom. The tops and leaves of this indigenous plant, *Spartium scopartum* of Linnæus:—*foliis ternatis solitariisque, ramis inermibus angulatis*, are the parts that are employed medicinally; they have a bitter taste, and are recommended for their purgative and diuretic qualities, in hydroptic cases.

GENISTA CANARIENSIS. The systematic name of the tree the wood of which is called rhodium. See *Rhodium lignum*.

GENITALE. (From *gigno*, to beget.) The privy member.

GENITALIUM. (From *genitale*, the membrum virile.) A disease of the genital parts.

GENITURA. (From *gigno*.) The male seed. Also the membrum virile.

GENOX. (From *γενω*, the knee.) A moveable articulation like that of the knee.

Gening. See *Ginseng*.

GENTIANA. (From *Gentius*, king of Illyria, who first used it.)

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*. *Gentian*.

2. The pharmacopœial name of what is also called *Gentiana rubra*. *Gentian*. Felwort. The gentian that is met with in the shops is the root of the *Gentiana lutea* of Linnæus:—*corollis subquinq;fidis rotatis verticillatis, calycibus spatulatis*; and is

imported from Switzerland and Germany. It is the only medicinal part of the plant, has little or no smell, but to the taste manifests great bitterness, on which account it is in general use as a tonic, stomachic, anthelmintic, antiseptic, emmenagogue, and febrifuge. The official preparations of this root are, the *infusum gentiane compositum*, and *tinctura gentiane composita*, of the London Pharmacopœia, and the *infusum amarum*, *vinum amarum*, *tinctura amara*, of the Edinburgh Pharmacopœia; and the *extractum gentiane* is ordered by both.

GENTIANA ALBA. The root of this plant, *Laserpitum latifolium*; *foliis cordatis, inciso-serratis*, of Linnæus, possesses stomachic, corroborant, and deobstruent virtues. It is seldom used.

GENTIANA CENTAURIUM. Lesser centaury was lately so called in the Linnæan system, but now *chironia centaurium*. See *Centaurium*.

GENTIANA LUTEA. The systematic name of the official gentian. See *Gentiana*.

GENTIANA RUBRA. See *Gentiana*.

GENU. The knee.

GENUGRA. (From *γόνυ*, the knee, and *γενε*, a seizure.) A name in Paracelsus for the gout in the knee.

GEOFFRÆA. (Named in honour of Dr. Geoffrey.) 1. The name of a genus of plants in the Linnæan system. Class, *Dialyptha*. Order, *Decandria*.

2. The pharmacopœial name of the bark of the *Geoffroya inermis foliolis lanceolatis* of Swartz. The plant is a native of Jamaica, where it is distinguished by the name of cabbage-bark tree, or worm bark tree. It has a mucilaginous and sweetish taste, and a disagreeable smell. According to Dr. Wright of Jamaica, it is powerfully medicinal as an anthelmintic.

GEOFFROYA JAMAICENSIS. The systematic name of the bastard cabbage-tree. See *Cortex Geoffroye Jamaicensis*.

GEOFFROYA SURINAMENSIS. The systematic name of the tree, the bark of which is esteemed as an anthelmintic.

GERANIS. (From *γέρων*, a crane; so called from its supposed resemblance to an extended crane.) A bandage for a fractured clavicle.

GERANIUM. (From *γέρων*, a crane; so called because its pistil is long, like the bill of a crane.) Class, *Monadelphæa*. Order, *Decandria*. The name of a genus of plants in the Linnæan system. *Geranium*, or cranes-bill.

GERANIUM BATRACHIOIDES. Crowfoot cranes-bill. This is the *Geranium pratense* of Linnæus; it possesses adstringent virtues, but in a slight degree.

GERANIUM COLUMBINUM. Doves foot. *Geranium rotundifolium* of Linnæus. This

plant possesses slightly adstringent virtues.

GERANIUM MOSCHATUM. The adstringent property of this plant has induced practitioners to exhibit it in cases of debility and profluvia.

GERANIUM PRATENSE. The systematic name of the crowfoot cranes-bill. See *Geranium batrachioides*.

GERANIUM ROBERTIANUM. Stinking cranes-bill. Herb robert. This common plant has been much esteemed as an external application in erysipelatos inflammations, cancer, mastodynia, and old ulcers, but is now deservedly fallen into disuse.

GERANIUM ROTUNDFOLIUM. The systematic name of the doves foot. See *Geranium columbinum*.

GERANIUM SANGUINARIUM. Bloody cranes bill. *Geranium sanguineum* of Linnæus. The adstringent virtues ascribed to this plant do not appear to be considerable.

GERANIUM SANGUINEUM. The systematic name of the *Geranium sanguinarium*. *Germander*. See *Chamaedris*.

Germander, water. See *Scordium*.

GEROCOMIA. (From *γέρων*, an aged person, and *κομω*, to be concerned about.) That part of medicine which regards the regimen and treatment of old age.

GERONTOPOGON. (From *γέρων*, an old man, and *πογων*, a beard; so called because its downy seed, while enclosed in the calyx, resembles the beard of an aged man.) The herb old man's beard. Purple flowered tragopogon.

GERONTOXON. (From *γέρων*, an old person, and *τοξον*, a dart.) A small ulcer, like the head of a dart, appearing sometimes in the cornea of old persons. The socket of a tooth.

GEROPOGON. See *Gerontopogon*.

GETON. Quicksilver.

GEUM. 1. The name of a genus of plants in the Linnæan system. Class, *Icosandria*. Order, *Polygymia*.

2. The pharmacopœial name of the *Geum rivale* of Linnæus, the root of which is the part directed for medicinal uses. It is inodorous, and imparts an austere taste. In America it is in high estimation in the cure of intermittents, and is said to be more efficacious than the Peruvian bark. Diarrhoeas and hæmorrhages are also stopped by its exhibition.

GEUM URBANUM. The systematic name of the herb bennet, or avens. See *Caryophyllata*.

Giddiness. See *Vertigo*.

Gilead, balsam. See *Balsamum Gileadense*.

Gill-go-by-ground. See *Hederacea*.

Gillyflower. See *Caryophyllus ruber*.

GIN. Geneva. Hollands. The names of a spirit distilled from malt or rye, which afterwards undergoes the same process,

a second time, with juniper berries. This is the original and most wholesome state of the spirit; but it is now prepared without juniper-berries; and is distilled from turpentine, which gives it something of a similar flavour. The consumption of this article, especially in the metropolis, is immense, and the consequences are, as Dr. Willan observes, pernicious to the health of the inhabitants.

Ginger. See *Zinziber*.

GINGIBER. See *Zingiber*.

GINGIBRACHIUM. (From *gingiva*, the gums, and *brachium*, the arm.) A name for the scurvy, because the gums, arms, and legs, are affected with it.

GINGIDIUM. A species of *Daucus*.

GINGIHAL. See *Zingiber*.

GINGIPEDIUM. (From *gingiva*, the gums, and *pes*, the foot.) A name for the scurvy, because the arms and legs are affected.

GINGIVÆ. (From *gigno*, to beget, because the teeth are, as it were, born in them.) The gums. See *Gums*.

GINGLYMUS. (γγγλυμο, a hinge.) The hinge-like joint. A species of diarthrosis or moveable connection of bones, which admits of flexion and extension, as the knee-joint, &c.

GINSENG. (*Ginseng*, Indian.) The name of the root of the *Panax quinquefolium* of Linnaeus:—*foliis ternis quinatis*. It is imported into this country scarcely the thickness of the little finger, about three or four inches long, frequently forked, transversely wrinkled, of a horay texture, and both internally and externally of a yellowish white colour. To the taste it discovers a mucilaginous sweetness, approaching to that of liquorice, accompanied with some degree of bitterness, and a slight aromatic warmth. The Chinese ascribe extraordinary virtues to the root of ginseng, and have no confidence in any medicine unless in combination with it. In Europe, however, it is very seldom employed.

Ginseng root. See *Ginseng*.

GRA. Quick lime.

GRIER. Tartar.

GIZZARD. The gizzards, or stomach of poultry, with white flesh, have long been considered, in France, as medicinal. They have been recommended in obstructions of the urinary passages, complaints of the bladder, and nephritic pains; but particularly as a febrifuge. Bouillon Lagrange considers its principal substance as oxygenated gelatine, with a small quantity of extractive matter.

GLABELLA. (From *glaber*, smooth; because it is without hair.) The space between the eye-brows.

GLADIOLUS. (Dim. of *gladius*, a sword; so named from the sword-like

shape of its leaf.) The name of a genus of plants in the Linnaean system. Class, *Triandria*. Order, *Monogynia*.

GLADIOLUS LUTEUS. See *Fris palustris*.

GLAMA. (γλαμα.) The sordes of the eye.

GLAND. (*Glandula*; diminutive of *glans*, a gland.) A gland is an organic part of the body, composed of blood-vessels, nerves, and absorbents, and destined for the secretion or alteration of some peculiar fluid. The glands of the human body are divided, by anatomists, into different classes, either according to their structure, or the fluid they contain. According to their fabric, they are distinguished into four classes.

1. Simple glands.
2. Compounds of simple glands.
3. Conglobate glands.
4. Conglomerate glands.

According to their fluid contents, they are more properly divided into, 1. Mucous glands. 2. Sebaceous glands. 3. Lymphatic glands. 4. Salival glands. 5. Lachrymal glands.

Simple glands are small hollow follicles, covered with a peculiar membrane, and having a proper excretory duct, through which they evacuate the liquor contained in their cavity. Such are the mucous glands of the nose, tongue, fauces, trachea, stomach, intestines, and urinary bladder, the sebaceous glands about the anus, and those of the ear. These simple glands are either dispersed here and there, or are contiguous to one another, forming a heap, in such a manner that they are not covered by a common membrane, but each hath its own excretory duct, which is never joined to the excretory duct of another gland. The former are termed solitary simple glands, the latter aggregate or congregate simple glands.

The compound glands consist of many simple glands, the excretory ducts of which are joined in one common excretory duct; as the sebaceous glands of the face, lips, palate, and various parts of the skin, especially about the pubes.

Conglobate, or, as they are also called, lymphatic glands, are those into which lymphatic vessels enter, and from which they go out again: as the mesenteric, lumbar, &c. They are composed of a texture of lymphatic vessels, connected together by cellular membranes—have no excretory duct—they are largest in the fetus.

Conglomerate glands are composed of a congeries of many simple glands, the excretory ducts of which open into one common trunk: as the parotid gland, thyroid gland, pancreas, and all the salival glands. Conglomerate glands differ but little from the compound glands, yet they are composed of more simple glands than the compound.

The excretory duct of a gland is the

duct through which the fluid of the gland is excreted. The vessels and nerves of glands always come from the neighbouring parts, and the arteries appear to possess a higher degree of irritability. The use of the glands is to separate a peculiar liquor, or to change it. The use of the conglobate glands is unknown.

GLANDULA LACHRYMALIS. See *Lachrymal gland*.

GLANDULÆ MYRTIFORMES. See *Caruncule myrtiformes*.

GLANDULÆ PACCHIONIÆ. (*Pacchioni*, the name of the discoverer.) A number of small, oval, fatty substances, not yet ascertained to be glandular, situated under the dura mater, about the sides of the longitudinal sinus. Their use is not known.

GLANDULOCARNEUS. An epithet given by Ruysch to some excrescences which he observed in the bladder.

GLANS. A gland, or nut.

GLANS PENIS. The very vascular body that forms the apex of the penis. The posterior circle is termed the *corona glanica*. See *Corpus spongiosum urethrae*.

GLANS UNGUENTARIA. See *Ben nux*.

GLASS. This substance is sometimes employed by surgeons, when roughly powdered, to destroy opacities of the cornea.

Glass of antimony. A vitreous oxyd of antimony.

Glass wort, mail-seeded. See *Kali*.

GLANTUM. (*Quasi callastum*; from *Callia*, who first used it.) The herb wood.

Glauber's salt. See *Sulphas sodæ*.

GLAUCIUM. (From *γλαυκος*, blue, or yellow; so called from its colour.) The yellow-horned poppy.

GLAUCOMA. (From *γλαυκος*, blue; because of the eye becoming of a blue or sea-green colour.) *Glaucosis*. An opacity of the vitreous humour. It is difficult to ascertain, and is only to be known by a very attentive examination of the eye.

Glaucosis. See *Glaucoma*.

GLELOMA. (From *γλαῦον*, the name of a plant in Dioscorides.) Class, *Didynamia*. Order, *Gymnospermia*. The name of a genus of plants in the Linnæan system. Ground ivy.

GLECOMA HEDERACEA. The systematic name of the ground ivy. See *Hedera terrestris*.

GLECHON. (From *γλαῦον*.) Penny-royal.

GLECHONITES. (From *γλαῦον*, penny-royal.) Wine impregnated with penny-royal.

GLEET. In consequence of the repeated attacks of gonorrhœa, and the debility of the part occasioned thereby, it not unfrequently happens that a gleet, or constant small discharge, takes place, or remains behind, after all danger of infection is removed. Mr. Hunter remarks,

that it differs from gonorrhœa in being *infectious*, and in the discharge consisting of globular bodies, contained in a slimy mucus, instead of serum. It is unattended with pain, scalding in making of water, &c.

GLENE. (*γληνη*.) Strictly signifies the cavity or socket of the eye; but by some anatomists is also used for that cavity of a bone which receives another within it.

GLENOID. (*Glénoides*; from *γληνη*, a cavity, and *ωδωρ*, resemblance.) The name of some articulate cavities of bones.

GLEUCIUM. (From *γλαυκος*, must.) An ointment, in the preparation of which was must.

GLEUXIS. (From *γλυκυσ*, sweet.) A sweet wine.

GLISGERE. To increase gradually, properly as fire does: but, by physical writers is sometimes applied to the natural heat and increase of spirits; and by others to the exacerbation of fevers which return periodically.

GLISCHRASMA. (From *γλισχρσμη*, to become glutinous.) Lentor. Viscidity.

GLISCHROCHOLUS. (From *γλισχρσρ*, viscid, and *χολα*, the bile.) An epithet for bilious viscid excrements.

GLISOMARÇO. White chalk.

Globate gland. See *Gland*.

GLOBULARIA. (From *globus*, a globe; so called from the shape of its flower.) The French daisy.

GLOBULARIA ALYPHUM. The leaves of this plant are used in some parts of Spain in the cure of the venereal disease. It is said to act also as a powerful but safe cathartic.

GLOBUS HYSTERICUS. The air rising in the œsophagus, and prevented by spasm from reaching the mouth, is so called by authors, because it mostly attends hysteria, and gives the sensation of a ball ascending in the throat.

GLOMER. (A clue of thread.) Mostly applied to glands.

GLOMERATE GLAND. A gland formed of a glomer of sanguineous vessels, having no cavity, but furnished with an excretory duct; as the lachrymal and mammary glands.

GLOSSAGRA. (From *γλωσσα*, the tongue, and *αγγρα*, a seizure.) A rheumatic pain in the tongue.

GLOSSO. (From *γλωσσα*, the tongue.) Names compounded with this word belong to muscles, nerves, or vessels, from their being attached, or going to the tongue.

GLOSSO-PHARYNGEAL NERVES. The ninth pair of nerves. They arise from the processes of the cerebellum, which run to the medulla spinalis, and terminate by numerous branches in the muscles of the tongue and pharynx.

GLOSSO-PHARYNGEUS. (*Musculus glossopharyngeus*; *γλωσσοφαρυγγειος*, from *γλωσσα*, the tongue, and *φαρυγγξ*, the pharynx:

so named from its origin in the tongue, and its insertion in the pharynx.) See *Constrictor pharyngeus superior*.

GLOSSO-STAPHYLINUS. (*Musculus glossostaphylinus*; from *γλῶσσα*, the tongue, and *σταφυλινος*, the staphylinus; so named because it is fixed in the tongue, and terminates in the staphylinus. See *Constrictor iethmi faucium*.)

GLOSSOCATOCHOS. (From *γλῶσσα*, tongue, and *κατίζει*, to hold. An instrument in P. Agineta for depressing the tongue. A spatula linguæ. The antient glossocatochus was a sort of forceps, one of the blades of which served to depress the tongue, while the other was applied under the chin.

GLOSSOCELE. (From *γλῶσσα*, the tongue, and *κύμα*, a tumour.) An extrusion of the tongue.

GLOSSOCOMA. A retraction of the tongue.

GLOSSOCOMION. (From *γλῶσσα*, a tongue, and *κομῆναι*, to guard.) By this was formerly meant a case for the tongue, for a hantboy; but the old surgeons, by metaphor, use it to signify an instrument, or case, for containing a fractured limb.

GLOTTA. (*γλῶττα*, the tongue.) The tongue.

GLUCINE. For the discovery of this earth we are indebted to Vauquelin, who found it, in 1795, in the Aigue-marine or beryl, a transparent stone, of a green colour, and in the emerald of Peru. It exists combined with silice, argil, lime, and oxyd of iron in the one; and with the same earths, with oxyd of chrome, in the other. It has lately been discovered in the Gadolinite by Mr. Ekeberg.

Its name is taken from the Greek word (*γλυκύς*) which signifies sweet or saccharine, because it gives that taste to the salts it forms.

Glucine is white, light, and soft to the touch. It is insipid, and adheres to the tongue; and is infusible by itself in the fire. Its specific gravity is 2.967. It is soluble in alkalies and their carbonates, and in all the acids except the carbonic and phosphoric, and forms with them saccharine and slightly astringent salts. It is exceedingly soluble in sulphuric acid used to excess. It is fusible with borax, and forms with it a transparent glass. It absorbs one-fourth of its weight of carbonic acid. It decomposes sulphate of alumine. It is not precipitated by the hydro-sulphurets, nor by prussiate of potash, but by all the succinates. Its affinity for the acids is intermediate between magnesia and alumine.

To obtain this earth, reduce some beryl to an impalpable powder, fuse it with three times its weight of potash, and dissolve the mass in muriatic acid. Separate the silice by evaporation and filtration, and decompose the remaining fluid by adding carbo-

nate of potash; re-dissolve the deposit when washed in sulphuric acid, and by mingling this solution with sulphate of potash, sulphate of alumine will be obtained, which crystallizes.

Then mix the fluid with a solution of carbonate of ammonia, which must be used in excess; filter and boil it, and a white powder will gradually fall down, which is glucine.

GLOTTIS. (From *γλῶττα*, the tongue.) The superior opening of the larynx at the bottom of the tongue.

GLUTEAL ARTERY. A branch of the internal iliac artery.

GLUTEN. (*Quasi gelutem*; from *gelo*, to congeal.) Glue, Lentor. See *Gluten, animal and vegetable*.

GLUTEN, ANIMAL. This substance constitutes the basis of the fibres of all the solid parts. It resembles in its properties the gluten of vegetables.

GLUTEN, VEGETABLE. If wheat flour be made into a paste, and washed in a large quantity of water, it is separated into three distinct substances; a *muclaginous saccharine matter*, which is readily dissolved in the liquor, and may be separated from it by evaporation; *starch*, which is suspended in the fluid, and subsides to the bottom by repose; and *gluten*, which remains in the hand, and is tenacious, very ductile, somewhat elastic, and of a brown gray colour. This glutinous substance is obtained from several vegetables in great abundance, and, when dried, becomes a horny mass. It is soluble both in water and spirit of wine, and, if boiled with the former, it coagulates like the white of an egg. It burns like horn, and affords the same products by distillation in the dry way. It readily putrefies when kept in a cold and moist place.

GLUTEUS MAXIMUS. (From *γλῦς*, the buttocks.) *Gluteus magnus* of Albinus, *Gluteus major* of Cowper, and *Liesacro femoral* of Dumas. This broad radiated muscle, which is divided into a number of strong fasciculi, is covered by a pretty thick aponeurosis derived from the *fascia lata*, and is situated immediately under the integuments. It arises fleshy from the outer lip of somewhat more than the posterior half of the spine of the ilium, from the ligaments that cover the two posterior spinous processes; from the posterior sacro-sciatic ligament; and from the outer sides of the os sacrum and os coccygis. From these origins the fibres of the muscle run towards the great trochanter of the os femoris, where they form a broad and thick tendon, between which and the trochanter there is a considerable *bursa mucosa*. This tendon is inserted into the upper part of the *linea aspera*, for the

space of two or three inches downwards; and sends off fibres to the fascia lata, and to the upper extremity of the vastus externus. This muscle serves to extend the thigh, by pulling it directly backwards; at the same time it draws it a little outwards, and thus assists in its rotatory motion. Its origin from the coccyx seems to prevent that bone from being forced too far backwards.

GLUTEUS MEDIUS. *Ilio trochanterien* of Dumas. The posterior half of this muscle is covered by the gluteus maximus, which it greatly resembles in shape; but the anterior and upper part of it is covered only by the integuments, and by a tendinous membrane which belongs to the fascia lata. It arises fleshy from the outer lip of the anterior part of the spine of the ilium, from part of the posterior surface of that bone, and likewise from the fascia that covers it. From these origins its fibres run towards the great trochanter, into the outer and posterior part of which it is inserted by a broad tendon. Between this tendon and the trochanter there is a small thin *bursa mucosa*. The uses of this muscle are nearly the same as those of the gluteus maximus; but it is not confined like that muscle, to rolling the os femoris outwards, its anterior portion being capable of turning that bone a little inwards. As it has no origin from the coccyx, it can have no effect on that bone.

GLUTEUS MINIMUS. *Glutæus minor* of Albinus and Cowper; and *Ilio ischii trochanterien* of Dumas. This, which is likewise a radiated muscle, is situated under the gluteus medius. In adults, and especially in old subjects, its outer surface is usually tendinous. It arises fleshy between the two semicircular ridges we observe on the outer surface of the ilium, and likewise from the edge of its great niche. Its fibres run, in different directions, towards a thick flat tendon, which adheres to a capsular ligament of the joint, and is inserted into the fore and upper part of the great trochanter. A small *bursa mucosa* may be observed between the tendon of this muscle and the trochanter. This muscle assists the two former in drawing the thigh backwards and outwards, and in rolling it. It may likewise serve to prevent the capsular ligament from being pinched in the motions of the joint.

GLUTIA. (From γλῦτος, the buttocks.) The two small protuberances in the brain, called *Nates*.

GLUTUPATERS. (From *gluttus*, the throat, and *patos*, to extend.) An epithet for the stomach, which is an extension of the throat.

GLUTUS. (γλῦτος, from γλῦμι, filthy.) The buttocks.

GRUCEAMA. (From γλῦμι, sweet.) A sweet medicated wine.

GLYCYFICUS. (From γλῦμι, sweet, and *πικρὸς*, bitter; so called from its bitterish sweet taste.) The woody nightshade. See *Dulcamara*.

GLYCYRRHIZA. (From γλῦμι, sweet, and *ρίζα*, a root.) 1. The name of a genus of plants in the Linnæan system. Class, *Diadelphia*. Order, *Decandria*.

2. The pharmacopœial name of liquorice. The sweet root of the *Glycyrrhiza glabra* of Linnæus:—*leguminibus glabris, stipulis nullis, foliolo impari petiolato*. A native of the south of Europe, but cultivated in Britain. The root contains a great quantity of saccharine matter, joined with some proportion of mucilage, and hence it has a viscid sweet taste. It is in common use as a pectoral or emollient, in catarrhal defluxions on the breast, coughs, hoarsenesses, &c. Infusions, or the extract made from it, which is called *Spanish liquorice*, afford likewise very commodious vehicles for the exhibition of other medicines; the liquorice taste concealing that of unpalatable drugs more effectually than syrups or any of the sweets of the saccharine kind.

GLYCYRRHIZA ECHINATA. This species of liquorice is substituted in some places for the root of the *glabra*.

GLYCYRRHIZA GLABRA. The systematic name of the official liquorice. See *Glycyrrhiza*.

GLYCYRANCON. (From γλῦμι, sweet, and *αγκων*, the elbow; so called from its sweetish taste, and its inflexions, or elbows, at the joints.) A species of southernwood.

Glysons capsule. See *Capsule of Glyson*.

GNAPHALIUM. (From γναφαλον, cotton; so named from its soft downy surface.)

1. The name of a genus of plants in the Linnæan system. Class, *Syngnæda*. Order, *Polygamia superflua*.

2. The pharmacopœial name of the herb cotton-weed. The flores gnaphalii of the pharmacopœias, called also *flores hispidulæ*; seu *pedes casti* are the produce of the *Gnaphalium dioicum* of Linnæus. They are now quite obsolete, but were formerly used as astringents, and recommended in the cure of whooping-cough, phthisis, pulmonary, and hæmoptysis.

GNAPHALIUM ARNARIUM. The flowers of this plant, as well as those of the *gnaphalium stochas*, called in the pharmacopœias *flores elichrysi*. See *Elchrysium*.

GNAPHALIUM DIOICUM. The systematic name of the *pes casti*. See *Gnaphalium*.

GNAPHALIUM STOCHAS. The systematic name of Goldilocks. See *Elchrysium*.

GNATHUS. (From γναθία, to bend; so

called from their curvature.) The jaw, or jaw-bones. Also the cheek.

GONITIA. A term applied by Hippocrates, and others since, to some medicinal precepts wrote in the island of Gnidos.

Goat's rue. See *Galiga*.

GOTTRE. See *Bronchocole*.

GOLD. *Aurum.* A metal found in nature only in a metallic state; most commonly in grains, ramifications, leaves, or rhomboidal, octahedral, or pyramidal crystals. Its matrix is generally quartz, sand-stone, siliceous schistus, &c. It is found also in the sands of many rivers, particularly in Africa, Hungary, and France, in minute irregular grains, called *gold-dust*. Native gold, found in compact masses, is never completely pure; it is alloyed with silver, or copper, and sometimes with iron and tellurium. The largest piece of native gold that has been hitherto discovered in Europe, was found in the county of Wicklow, in Ireland. Its weight was said to be twenty-two ounces, and the quantity of alloy it contained was very small. Several other pieces, exceeding one ounce, have also been discovered at the same place, in sand, covered with turf, and adjacent to a rivulet.

Gold is also met with in a particular sort of argenteriferous copper pyrites, called in Hungary *Gelf*. This ore is found either massive, or crystallized in rhomboids, or other irregular quadrangular or polygonal masses. It exists likewise in the sulphurated ores of Nagaya in Transylvania. These all contain the metal called tellurium. Berthollet, and other French chymists, have obtained gold out of the ashes of vegetables.

Gold-cup. A vulgar name for many species of *Ranunculi*.

Golden-rod. See *Virga aurea*.

Goldlocks. See *Elychrysum*.

GOMPHIASIS. (From *γομφος*, a nail.) *Gomphiasmus*. A disease of the teeth, when they are loosened from the the sockets, like nails drawn out of the wood.

GOMPHIASMUS. See *Gomphiasts*.

GOMPHI. (From *γομφος*, a nail; so called, because they are as nails driven into their sockets.) The dentes molares, or grinding teeth.

GOMPHOMA. See *Gomphosis*.

GOMPHOSIS. (From *γομφος*, to drive in a nail.) *Gomphoma*. A species of immovable connection of bones, in which one bone is fixed in another, like a nail in a board, as the teeth in the alveoli of the jaws.

GONAGRA. (From *γονυ*, the knee, and *αγγω, capio*, to take.) The gout in the knee.

GONZ. (*γονυ*.) The seed. But in Hippocrates it is the uterus.

ΓΟΝΕΒΟΝΑ. (From *γονος*, a hard knot.)

1. The cramp.

2. A round tubercle in the trunk of a tree.

3. A hard round tumour of the nervous parts; but particularly a bronchocele, or other hard tumour of the neck.

ΓΟΝΕΒΛΙΟΝ. (From *γονυ*, round.) A pill.

ΓΟΝΟΙΔΕΣ. (From *γονυ*, seed, and *ωμοι*, form.) Resembling seed. Hippocrates often uses it as an epithet for the excrements of the belly, and for the contents of the urine, when there is something in them which resembles the seminal matter.

GONORRHEA. (From *γονυ*, the semen, and *ρρ*, to flow; from a supposition of the ancients, that it was a seminal flux.) A genus of disease in the class *locales*, and order *apoceneses*, of Dr. Cullen's arrangement, who defines it a preternatural flux of fluid from the urethra in males, without any libidinous desires, and from the vagina in females. He makes four species, viz.

1. *Gonorrhœa pura, or benigna*; a mucous discharge from the urethra, without dysuria, or lascivious inclination.

2. *Gonorrhœa impura, maligna, syphilitica, virulenta*; a discharge resembling pus, from the urethra, with heat of urine, &c. after impure coition, to which often succeeds a discharge of mucus from the urethra, with little or no dysuria, called a gleet. *Fluor albus malignus.* *Blennorrhagi* of Swediaur. In English, a *clap*, from the old French word *clapisses*, which were public shops, kept and inhabited by single prostitutes, and generally confined to a particular quarter of the town, as is even now the case in several of the great towns of Italy. In Germany, the disorder is named *tripper*, from dripping; and in French, *chaudpisse*, from the heat and scalding in making water.

No certain rule can be laid down with regard to the time that a clap will take before it makes its appearance, after infection has been conveyed. With some persons it will shew itself in the course of three or four days, whilst, with others, there will not be the least appearance of it before the expiration of some weeks. It most usually is perceptible, however, in the space of from six to fourteen days, and in a male, begins with an uneasiness about the parts of generation, such as an itching in the glans penis, and a soreness and tingling sensation along the whole course of the urethra; soon after which, the person perceives an appearance of whitish matter, at its orifice, and also some degree of pungency upon making water.

In the course of a few days, the discharge of matter will increase considera-

bly ; will assume, most probably, a greenish or yellowish hue, and will become thinner, and lose its adhesiveness ; the parts will also be occupied with some degree of redness and inflammation, in consequence of which the glans will put on the appearance of a ripe cherry ; the stream of urine will be smaller than usual, owing to the canal being made narrower by the inflamed state of its internal membrane, and a considerable degree of pain, and scalding heat, will be experienced on every attempt to make water.

Where the inflammation prevails in a very high degree, it prevents the extension of the urethra, on the taking place of any erection, so that the penis is, at that time, carried downwards, with great pain, which is much increased, if attempted to be raised towards the belly, and the stimulus occasions it often to be erected, particularly when the patient is warm in bed, and so deprives him of sleep, producing, in some cases, an involuntary emission of semen.

In consequence of the inflammation, it sometimes happens that, at the time of making water, owing to the rupture of some small blood-vessel, a slight hæmorrhage ensues, and a small quantity of blood is voided. In consequence of inflammation, the prepuce likewise becomes often so swelled at the end, that it cannot be drawn back ; which symptom is called a phymosis ; or that, being drawn behind the glans, it cannot be returned, which is known by the name of paraphymosis. Now and then, from the same cause, little hard swellings arise on the lower surface of the penis, along the course of the urethra, and these perhaps suppurate and form into fistulous sores.

The adjacent parts sympathizing with those already affected, the bladder becomes irritable, and incapable of retaining the urine for any length of time, which gives the patient a frequent inclination to make water, and he feels an uneasiness about the scrotum, perinæum, and fundament. Moreover the glands of the groins grow indurated and enlarged, or perhaps the testicle becomes swelled and inflamed, in consequence of which he experiences excruciating pains, extending from the seat of the complaint up into the small of the back ; he gets hot and restless, and a small symptomatic fever arises.

Where the parts are not occupied by much inflammation, few or none of the last-mentioned symptoms will arise, and only a discharge, with a slight heat or scalding in making water, will prevail.

If a gonorrhœa is neither irritated by any irregularity of the patient, nor prolonged by the want of timely and proper assistance, then, in the course of about a fortnight or

three weeks, the discharge, from having been thin and discoloured at first, will become thick, white, and of a ropy consistence ; and from having gradually began to diminish in quantity, will at last cease entirely, together with every inflammatory symptom whatever ; whereas, on the contrary, if the patient has led a life of intemperance and sensuality, has partaken freely of the bottle and high-seasoned meats, and has, at the same time, neglected to pursue the necessary means, it may then continue for many weeks, or months ; and, on going off, may leave a weakness or gleet behind it, besides being accompanied with the risk of giving rise, at some distant period, to a constitutional affection, especially if there has been a neglect of proper cleanliness ; for where venereal matter has been suffered to lodge between the prepuce and glans penis for any time, so as to have occasioned either excoriation or ulceration, there will always be danger of its having been absorbed.

Another risk, arising from the long continuance of a gonorrhœa, especially if it has been attended with inflammatory symptoms, or has been of frequent recurrence, is the taking place of one or more strictures in the urethra. These are sure to occasion a considerable degree of difficulty, as well as pain, in making water, and, instead of its being discharged in a free and uninterrupted stream, it splits into two, or perhaps is voided drop by drop. Such affections become, from neglect, of a most serious and dangerous nature, as they not unfrequently block up the urethra, so as to induce a total suppression of urine.

Where the gonorrhœa has been of long standing, warty excrescences are likewise apt to arise about the parts of generation, owing to the matter falling and lodging thereon ; and they not unfrequently prove both numerous and troublesome.

Having noticed every symptom which usually attends on gonorrhœa, in the male sex, it will only be necessary to observe, the same heat and soreness in making water, and the same discharge of discoloured mucus, together with a slight pain in walking, and an uneasiness in sitting, take place in females as in the former ; but as the parts in women, which are most apt to be affected by the venereal poison, are less complex in their nature, and fewer in number, than in men, so of course the former are not liable to many of the symptoms which the latter are ; and from the urinary canal being much shorter, and of a more simple form, in them than in men, they are seldom, if ever, incommoded by the taking place of strictures.

With women, it indeed often happens, that all the symptoms of a gonorrhœa are

so very slight, they experience no other inconvenience than the discharge, except perhaps immediately after menstruation, at which period, it is no uncommon occurrence for them to perceive some degree of aggravation in the symptoms.

Women of a relaxed habit, and such as have had frequent miscarriages, are apt to be afflicted with a disease known by the name of fluor albus, which is often difficult to distinguish from gonorrhœa virulenta, as the matter discharged in both is, in many cases, of the same colour and consistence. The surest way of forming a just conclusion, in instances of this nature, will be to draw it from an accurate investigation, both of the symptoms which are present and those which have preceded the discharge; as likewise, from the concurring circumstances, such as the character and mode of life of the person, and the probability there may be of her having had venereal infection conveyed to her by any connection in which she may be engaged.

Not long ago, it was generally supposed that gonorrhœa depended always upon ulcers in the urethra, producing a discharge of purulent matter; and such ulcers do, indeed, occur in consequence of a high degree of inflammation and suppuration; but many dissections of persons, who have died whilst labouring under a gonorrhœa, have clearly shewn that the disease may, and often does, exist without any ulceration in the urethra, so that the discharge which appears is usually of a vitiated mucus, thrown out from the mucous follicles of the urethra. On opening this canal, in recent cases, it usually appears red and inflamed; its mucous glands are somewhat enlarged, and its cavity is filled with matter to within a small distance from its extremity. Where the disease has been of long continuance, its surface all along, even to the bladder, is generally found pale and relaxed, without any erosion.

3. *Gonorrhœa laxorum libidinosa*; a pelucid discharge from the urethra, without erection of the penis, but with venereal thoughts while awake.

Gonorrhœa dormientium, oncirosmos. When, during sleep, but dreaming of venereal engagements, there is an erection of the penis and a seminal discharge.

GONORRHOEA BENIGNA. See *Gonorrhœa pura*.

GONORRHOEA CHORDATA. A gonorrhœa, accompanied with painful tension of the penis, called chordee.

GONORRHOEA DORMIENTIUM. Involuntary nocturnal emission.

GONORRHOEA IMPURA. The venereal gonorrhœa.

GONORRHOEA LAXORUM. Involuntary emission from debility.

GONORRHOEA LIBIDINOSA. Involuntary emission from lust.

GONORRHOEA MALIGNA. A venereal or malignant gonorrhœa.

GONORRHOEA MUcosa. A discharge a mucus from the urethra, or gleet.

GONORRHOEA ONEIROSMOS. An involuntary nocturnal emission.

GONORRHOEA PURA. A common gleet.

GONORRHOEA SPURIA. A species of gonorrhœa affecting the glans.

GONORRHOEA SYPHILITICA. The venereal gonorrhœa.

GONORRHOEA BALANI. A species of gonorrhœa affecting the glans penis only.

GONALGIA. (From *genu*, the knee, and *algos*, pain.) *Gonyalgia*. Gout in the knee.

GONRUS. A species of vermes, peculiar to hot climates.

Goose foot, stinking. See *Atriplex fastida*.

Goose-grass. See *Aparine*.

GOSSYPIUM. (From *gotia*, whence *gotipium*, Egyptian.) 1. The name of a genus of plants in the Linnæan system. Class, *Monadelphia*. Order, *Polyandria*.

2. The pharmacopœial name of the cotton tree. See *Bombax*.

GOSSYPIUM HERBACEUM. The systematic name of the cotton-plant. See *Gossypium*.

GOULARD'S EXTRACT. A saturated solution of acetat of lead. See *Acetatis liquor plumbi*.

Gourd. See *Cucurbita*.

Gourd, bitter. See *Colocynthis*.

Gout. See *Arthritis*.

GRACILIS. (*Gracilis*, from its smallness.) *Rectus interior femoris sive gracilis interior* of Winalow. *Sous pubis creti tibialis* of Dumas. This long, straight, and slender muscle is situated immediately under the integuments, at the inner part of the thigh. It arises by a broad and thin tendon, from the anterior part of the ischium and pubis, and soon becoming fleshy, descends nearly in a straight direction along the inside of the thigh. A little above the knee, it terminates in a slender and roundish tendon, which afterwards becomes flatter, and is inserted into the middle of the tibia, behind and under the sartorius. Under the tendons of this and the rectus, there is a considerable *bursa mucosa*, which on one side adheres to them and to the tendon of the semitendinosus, and on the other to the capsular ligament of the knee. This muscle assists in bending the thigh and leg inwards.

GRAMEN CANINUM. *Gramen dioscoridis*. *Gramen repens*. *Loliaceum radice repente*. Dog's grass. Couch grass. *Triticum repens* of Linnaeus. The roots are agreeably sweet, and possess aperient properties. The expressed juice is recommended to be given largely.

GRAMEN CRUCIS CYPERIoidis. *Gramen Egyptiacum*. Egyptian cock's-foot grass, or grass of the cross. The roots and plants

possess the same virtues as the dog's grass, and are serviceable in the earlier stages of dropsy. They are supposed to correct the bad smell of the breath, and to relieve nephritic disorders, colics, &c. although now neglected.

GRANIA. The sordes of the eyes.

GRANNE. (From *γραμμή*, a line; so called from its linear appearance.) The iris of the eye.

GRANA CRIDIA. See *Coccognidia*.

GRANA INSECTORIA. Kermes berries.

GRANA KERMES. Kermes berries.

GRANA PARADISI. *Cardamomum majus*. *Meleguetta mariguetta*. *Cardamomum piperatum*. Grains of Paradise. The seeds of the *Amomum grana Paradisi* of Linnæus. They are angular reddish brown seeds, smaller than pepper, and resembling very much the seeds of the cardamom minus. They are extremely hot, and similar in virtue to pepper.

GRANA TIGLIA. See *Tiglia grana*.

GRANA TINCTORIA. See *Kermes*.

GRANADILLA. (Dim. of *granado*, a pomegranate, Spanish: so called because at the top of the flower there are points, like the grains of a pomegranate.) The passion-flower, the fruit of which is said to possess refrigerating qualities.

GRANATRISTUM. A boil, or carbuncle.

GRANATUM. (From *granum*, a grain, because it is full of seed.) The pomegranate. The fruit of the *Punica granatum* of Linnæus:—*foliis lanceolatis, caule arboreo*. The rind of the fruit, and the flowers called *Balaustine flowers*, are the parts directed for medicinal use. In their smell there is nothing remarkable, but to the taste they are very adstringent, and have successfully been employed as such, in diseases both internal and external.

GRANDEBALÆ. (*Quod in grandioribus etate nascantur*, because they appear in those who are advanced in years.) The hairs under the arm-pits.

GRANDINES. Small tumours on the eyelids. See *Grando*.

GRANDINOSUM OS. The cuboid bone of the foot.

GRANDO. (*Quod similitudinem granorum habeat*, because it is in shape and size like a grain of seed.) Hail. A moveable tumour on the margin of the eyelid is so called, from its likeness to a hail stone.

GRANULATION. (*Granulatio*; from *granum*, a grain.) The little grain-like fleshy bodies, which form on the surfaces of ulcers and suppurating wounds, and serve both for filling up the cavities, and bringing nearer together and uniting their sides.

Nature, in bringing parts as nearly as possible to their original state, whose disposition, action, and structure, have been altered by accident, or disease; and after having, in her operations for this purpose, formed pus, she immediately sets about forming

a new matter upon surfaces, in which there has been a breach of continuity. This process is called *granulating* or *incarnation*; and the substance formed is called *granulations*. The colour of healthy granulations is a deep florid red. When livid, they are unhealthy, and have only a languid circulation. Healthy granulations, on an exposed or flat surface, rise nearly even with the surface of the surrounding skin, and often a little higher; but when they exceed this, and take on a growing disposition, they are unhealthy, become soft, spongy, and without any disposition to form skin. Healthy granulations are always prone to unite to each other, so as to be the means of uniting parts.

GRANUM MOSCHI. See *Abelmoschus*.

GRANUM REGIUM. The castor-oil seed.

GRAPHIOIDES. (From *γραφειν*, a pencil, and *ως*, a form.) The processus styloformis of the os temporis and ulna is so called. The musculus biventer vel digastricus was formerly so called, from its supposed origin from the process of the temple-bone, so called.

GRASSA. Borax.

GRATIOLO. (Dim. of *gratia*, so named from its supposed admirable qualities.)

1. The name of a genus of plants in the Linnæan system. Class, *Diandria*. Order, *Monogynia*.

2. The pharmacopœial name of the hedge-hyssop, called also *digitalis minima*, *gratia dei*, *gratiola centaurioides*. This exotic plant, the *Gratiola officinalis* of Linnæus:—*foliis lanceolatis serratis, floribus pedunculatis*, is a native of the south of Europe; but is raised in our gardens. The leaves have a nauseous bitter taste, but no remarkable smell: they purge and vomit briskly in the dose of half a drachm of the dry herb, and of a drachm infused in wine or water. This plant, in small doses, has been commonly employed as a cathartic and diuretic in hepatic diseases, and instances of its good effects in ascites and anasarca are recorded by many respectable practitioners. Gesner and Bergius found a scruple of the powder a sufficient dose, as in this quantity it frequently excited nausea or vomiting; others have given it to half a drachm, two scruples, a drachm, and even more.

An extract of the root of this plant is said to be more efficacious than the plant itself, and exhibited in the dose of half a drachm, or a drachm; in dysenteries, produces the best effects. We are also told by Kostrzewski that, in the hospitals at Vienna, three maniacal patients were perfectly recovered by its use; and in the most confirmed cases of lues venerea, it effected a complete cure; it usually acted by increasing the urinary cutaneous, or salivary discharges.

GRATIOLO OFFICINALIS. The syste-

matic name of the hedge-hyssop. See *Crucifolia*.

GRAVEDO. (From *gravis*, heavy.) A catarrh, or cold, with a sense of heaviness in the head.

Gravel. See *Calculus*.

Green sickness. See *Chlorosis*.

GRESSURA. (From *gredior*, to proceed.) The perinæum which goes from the pudendum to the anus.

GRIELUM. A name formerly applied to parsley; smallage.

GRIPHOMENOS. (From *γρηφορ*, a net, because it surrounds the body as with a net.) Applied to pains which surround the body at the loins.

Gromwell, common. See *Lithospermum*.

GROSSULARIA. (Dim. of *grossus*, an unripe fig; so named because its fruit resembles an unripe fig.) The gooseberry or gooseberry-bush.

GROTTO DEL CANT. A grotto near Naples, in which dogs are suffocated. The carbonic acid air rises about eighteen inches. A man therefore is not affected, but a dog forcibly held in, or who cannot rise above it, is soon killed, unless taken out. He is recovered by plunging him in an adjoining lake.

Ground liverwort. See *Lichen cinereus terrestris*.

GROUND IVY. A common plant used in the form of infusion or tea in pectoral complaints. See *Hedera*.

Groundpine. See *Chamaepitys*.

Groundnut. See *Pignut*.

Groundsel. See *Erigerum*.

GRUTUM. *Milium*. A hard white tubercle of the skin, resembling in size and appearance a millet-seed.

GRYPHOSIS. (From *γρηφορ*, to incurvate.) A disease of the nails, which turn inwards, and irritate the soft parts below.

GUAIACUM. (From the Spanish *Guaya-can*, which is formed from the Indian *Houx-acan*.) *Guaiacum Americanum*, *Lignum vite*. *Lignum sanctum*. *Lignum benedictum*. *Palus sanctus*.

1. The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Monogynia*.

2. The pharmacopœial name of the officinal guaiacum, *Guaiacum officinale* of Linnæus:—*foliis bijugis obtusis*. This tree is a native of the West Indian islands. The wood, gum, bark, fruit, and even the flowers, have been found to possess medicinal qualities. The wood is brought principally from Jamaica, in large pieces of four or five hundred weight each, and from its hardness and beauty is used for various articles of turnery ware. It scarcely discovers any smell, unless heated, or while rasping, in which circumstances it yields a light aromatic one: chewed, it impresses a slight acrimony, biting the palate and fauces. The gum, or rather resin, is ob-

tained by wounding the bark in different parts of the body of the tree, or by what has been called jagging. It exudes copiously from the wounds, though gradually; and when a quantity is found accumulated upon the several wounded trees, hardened by exposure to the sun, it is gathered and packed up in small kegs for exportation: it is of a friable texture, of a deep greenish colour, and sometimes of a reddish hue; it has a pungent acrid taste, but little or no smell, unless heated. The bark contains less resinous matter than the wood, and is consequently a less powerful medicine, though in a recent state it is strongly cathartic. "The fruit," says a late author, "is purgative, and, for medicinal use, far excels the bark. A decoction of it has been known to cure the venereal disease, and even the yaws in its advanced stage, without the use of mercury." The flowers or blossoms, are laxative, and in Jamaica are commonly given to the children in the form of syrup. It is only the wood and resin of guaiacum which are now in general medicinal use in Europe; and as the efficacy of the former is supposed to be derived merely from the quantity of resinous matter which it contains, they may be considered indiscriminately as the same medicine. Guaiacum was first introduced into the materia medica soon after the discovery of America; and previous to the use of mercury in the lues venerea, it was the principal remedy employed in the cure of that disease; its great success brought it into such repute, that it is said to have been sold for seven gold crowns a pound: but notwithstanding the very numerous testimonies in its favour, it often failed in curing the patient, and was at length entirely superseded by mer. ury; and though it be still occasionally employed in syphilis, it is rather with a view to correct other diseases in the habit, than for its effects as an anti-venereal. It is now more generally employed for its virtues in curing gouty and rheumatic pains, and some cutaneous diseases. Dr. Woodville and others frequently conjoined it with mercury and soap, and in some cases with bark or steel, and found it eminently useful as an alternative. In the pharmacopœias it is directed in the form of tincture and elixir: the latter is ordered by the Edinburgh College to be prepared in two ways, *viz.* with rectified spirit, and the virous spirit of sal ammoniac. Of these compounds, the dose may be from two scruples to two drachms; the gum is generally given from 6 grains to 20 or even more, for a dose, either in pills or in a fluid form, by means of mucilage or the yolk of an egg. The decoctum lignorum (Pharm. Eding.) of which guaiacum is the chief ingredient, is commonly taken in the quantity of a pint a day.

As many writers of the sixteenth century

contended that guaiacum was a true specific for the venereal disease, and the celebrated Boerhaave maintained the same opinion, the following observations are inserted: Mr. Pearson mentions, that when he was first intrusted with the care of the Lock Hospital, in 1781, Mr. Bromfield and Mr. Williams were in the habit of reposing great confidence in the efficacy of a decoction of guaiacum wood. This was administered to such patients as had already employed the usual quantity of mercury; but who complained of nocturnal pains, or had gummata, nodes, ozæna, and other effects of the venereal virus, connected with secondary symptoms, as did not yield to a course of mercurial frictions. The diet consisted of raisins, and hard biscuit; from 2 to 4 pints of the decoction were taken every day; the hot bath was used twice a week; and a dose of antimonial wine and laudanum, or Dover's powder, was commonly taken every evening. Constant confinement to bed was not deemed necessary; neither was exposure to the vapour of burning spirit, with a view of exciting perspiration, often practised; as only a moist state of the skin was desired. This treatment was sometimes of singular advantage to those whose health had sustained injury from the disease, long confinement, and mercury. The strength increased; bad ulcers healed; exfoliations were completed; and these anomalous symptoms, which would have been exasperated by mercury, soon yielded to guaiacum.

Besides such cases, in which the good effects of guaiacum made it be erroneously regarded as a specific for the lues venerea, the medicine was also formerly given, by some, on the first attack of the venereal disease. The disorder being thus benefited, a radical cure was considered to be accomplished; and though frequent relapses followed, yet, as these partly yielded to the same remedy, its reputation was still kept up. Many diseases also, which got well, were probably not venereal cases. Mr. Pearson seems to allow, that in syphilitic affections, it may indeed operate like a true antidote, suspending, for a time the progress of certain venereal symptoms, and removing other appearances altogether; but he observes, that experience has evinced, that the unsubdued virus yet remains active in the constitution.

Mr. Pearson has found guaiacum of little use in pains of the bones, except when it proved sudorific; but that it was then inferior to antimony or volatile alkali. When the constitution has been impaired by mercury and long confinement, and there is a thickened state of the ligaments, or periosteum, or foul ulcers still remaining, Mr. Pearson says, these effects will often subside during the exhibition of the decoction; and

it will often suspend for a short time the progress of certain secondary symptoms of the lues venerea: for instance, ulcers of the tonsils, venereal eruptions, and even nodes. Mr. Pearson, however, never knew one instance, in which guaiacum eradicated the virus; and he contends, that its being conjoined with mercury neither increases the virtue of this mineral, lessens its bad effects, nor diminishes the necessity of giving a certain quantity of it. Mr. Pearson remarks that he has seen guaiacum produce good effects in many patients, having cutaneous diseases, the ozæna, and scrophulous affections of the membranes and ligaments.

Guinea-worm. The *Gordius medinensis*.

GUM. The mucilage of vegetables. It is usually transparent, more or less brittle when dry, though difficultly pulverable; of an insipid, or slightly saccharine taste; soluble in, or capable of combining with, water in all proportions, to which it gives a gluey adhesive consistence, in proportion as its quantity is greater. It is separable, or coagulates by the action of weak acids; it is insoluble in alkohol, and in oil; and capable of the acid fermentation, when diluted with water. The destructive action of fire causes it to emit much carbonic acid, and converts it into coal without exhibiting any flame. Distillation affords water, acid, a small quantity of oil, a small quantity of ammonia, and much coal.

These are the leading properties of gums, rightly so called; but the inaccurate custom of former times applied the term gum to all concrete vegetable juices so that in common we hear of gum copal, gum sandarach, and other gums, which are either pure resins, or mixtures of resins with the vegetable mucilage.

The principal gums are, 1. The common gums, obtained from the plum, the peach, the cherry-tree, &c. 2. Gum arabic, which flows naturally from the acacia in Egypt, Arabia, and elsewhere. This forms a clear transparent mucilage with water. 3. Gum Seneca or Senegal. It does not greatly differ from gum arabic; the pieces are larger and clearer; and it seems to communicate a higher degree of the adhesive quality to water. It is much used by calico-printers and others. The first sort of gums are frequently sold by this name, but may be known by their darker colour. 4. Gum adragant or tragacanth. It is obtained from a small plant of the same name, growing in Syria, and other eastern parts. It comes to us in small white contorted pieces resembling worms. It is usually dearer than other gums, and forms a thicker jelly with water.

Mr. Willis has found, that the root of the common blue bell, *hyacinthus non scriptus*, dried and powdered, affords a mucilage possessing all the qualities of that

from gum arabic. The roots of the vernal squill, white lily and orchid, equally yield mucilage. Lord Dundonald has extracted a mucilage also from lichens.

Gums treated with nitric acid afford the acid of sugar.

Gum-boil. See *Parulis*.

GUMMA. A strumous tumour on the periosteum of a bone.

GUMMI ARABICUM. See *Acacia gummi*.

GUMMI CARANNE. See *Caranna*.

GUMMI CERASORUM. The juices which exude from the bark of cherry-trees. It is very similar to gum arabic, for which it may be substituted.

GUMMI CHIBOU. A spurious kind of gum elmi, but little used.

GUMMI COURBAIL. An epithet sometimes applied to the juice of the *Hymenaea courba*. See *Anime*.

GUMMI EUPHORBIL. See *Euphorbium*.

GUMMI GALDA. See *Galda*.

GUMMI GAMBIESE. See *Kino*.

GUMMI GUTTA. See *Gambogia*.

GUMMI HEDERE. Ivy gum. The resinous juice of the *Hedera helix* of Linnæus, or ivy. It is imported from the East Indies, though it may be collected from trees in this country. It is brought over in hard compact masses, externally of a reddish brown colour, internally of a bright brownish yellow, with reddish specks or veins. It has a strong, resinous, agreeable smell, and an adstringent taste. Though never used in the practice of the present day, it possesses corroborant, adstringent, and antispasmodic virtues.

GUMMI JUNIPERINUM. See *Sandarack*.

GUMMI KIKEKUNEMALO. See *Kikekumalo*.

GUMMI KINO. See *Kino*.

GUMMI LACCA. See *Lacca*.

GUMMI MYRRA. See *Myrrha*.

GUMMI RESINA LUTEA. An astringent gum from New Holland.

GUMMI RUBRUM ASTRINGENS GAMBIESE. See *Kino*.

GUMMI SAGAPENUM. See *Sagapenum*.

GUMMI SENEGALENSE. This is a true gum, brought from the country through which the river Senegal runs, in loose or single drops, much larger than gum arabic. It exudes from the *Mimosa Senegal* of Linnæus. And is similar in virtue and quality to the gum arabic, and the gum which exudes in this climate from the cherry-trees.

GUMMI TRASCANTHE. See *Tragacantha*.

GUM-RESIN. *Gum-resina.* Gum-resins are the juices of plants that are mixed with resin, and an extractive matter, which has been taken for a gummy substance.

They seldom flow naturally from plants, but are mostly extracted by incision, in the form of white, yellow, or red fluids, which dry more or less quickly. Water, spirit of wine, wine or vinegar, dissolve them only

in part, according to the proportion they contain of resin or extract. Gum-resins may also be formed by art, by digesting the parts of vegetables containing the gum-resin in diluted alcohol, and then evaporating it. For this reason most tinctures contain gum-resin. The principal gum-resins employed medicinally are aloes, ammoniacum, asafoetida, galbanum, gambogia, guaiacum, myrrha, olibanum, opoponax, sagapenum, sarcocolla, scammonium, and styrax.

GUMS. *Gingiva.* The very vascular and elastic substance that covers the alveolar arches of the upper and under jaws, and embraces the necks of the teeth.

GUTTA. *Atunsel.* A drop. Drops are uncertain forms of administering medicines. The shape of the bottle, or of its mouth, from whence the drops fall, as well as the consistence of the fluid, occasions a considerable difference in the quantity administered.

Gutta is also a name of the apoplexy, from a supposition that its cause was a drop of blood falling from the brain upon the heart.

GUTTA GAMBIA. See *Gambogia*.

GUTTA OPACA. A name for the cataract.

GUTTA SERENA. (So called by the Arabians.) *Amaurosis.* This is a disease of the eye attended with a diminution, or total loss of sight, and arising from a paralytic affection of the retina and optic nerve.

The symptoms of gutta serena are noted for being very irregular. In many cases, the pupil is very much dilated, immoveable and of its natural black colour. Sometimes, however, in the most complete and incurable cases, the pupil is of its natural size, and the iris capable of free motion. In some cases, the pupil has a dull, glassy or horny appearance. Sometimes its colour is greenish, occasionally whitish and opaque, so as to be liable to be mistaken for an incipient cataract. Richter mentions a degree of strabismus, as the only symptom, except the loss of sight, as invariably attendant on amaurosis.

The blindness produced by the gutta serena, is generally preceded by an imaginary appearance of numerous insects, or substances, like cobwebs, interposing themselves between objects and the eye. The origin of a cataract on the other hand, is usually attended with a simple cloudiness of vision.

GUTTA NIGRA. The black drops, occasionally called the Lancashire or the Cheshire drops. A secret preparation of opium, said to be more active than the common tincture and supposed to be less injurious, as seldom followed by headache.

GUTTA ROSACEA. Red spots upon the face and nose.

GUTTURAL ARTERY. The superior thyroideal artery. The first branch of the external carotid.

GYMNASTICA. (From *γυμνος*, naked, performed by naked men in the public games.) A method of curing diseases by exercise, or that part of physic which treats of the rules that are to be observed in all sorts of exercises, for the preservation of health. This is said to have been invented by one Herodicus, born at Salymbra, a city of Thrace; or, as some say, at Leutini in Sicily. He was first master of an academy, where young gentlemen came to learn warlike and manly exercises; and observing them to be very healthful on that account, he made exercise become an art in reference to the recovering of men out of diseases, as well as preserving them from them; and called it *Gymnastic*, which he made a great part of his practice of physic. But Hippocrates, who was his scholar, blames him sometimes for his excesses

with this view. And Plato exclaims against him with some warmth, for enjoining his patients to walk from Athens to Megara, which is about 25 miles, and to come home on foot as they went, as soon as ever they had but touched the walls of the city.

GYNECIA. (From *γυν*, woman.) It signifies the menses, and sometimes the lochia.

GYNECIUM. (From *γυν*, a woman.) A seraglio; the pudendum muliebre; also a name for *antimony*.

GYNECOMANIA. (From *γυν*, a woman, and *μανια*, madness.) That species of insanity that arises from love.

GYNECOMASTON. (From *γυν*, a woman, and *μαστορ*, a breast.) An enormous increase of the breasts of women.

GYNECOMYSTAX. (From *γυν*, a woman, and *μυσταξ*, a beard.) The hairs on the female pudenda.

H.

HABENA. A bridle. A bandage for keeping the lips of wounds together, made in the form of a bridle.

HACUB. A species of *carduus*. *Gundelia tournefortii* of Linnæus:—the young shoots of which are eaten by the Indians, but the roots are emetic.

HÆMAGOGA. (From *αἷμα*, blood, and *αγω*, to bring off.) Medicines which promote the menstrual and hæmorrhoidal discharges.

HÆMALOPIA. (From *αἷμα*, blood, and *ἰσλασις*, to see.) A disease of the eyes, in which all things appear of a red colour. A variety of the pseudoblepsia imaginaria.

HÆMALOPS. (From *αἷμα*, blood, and *ωψ*, the face.) A red or livid mark in the face or eye. A blood-shot eye.

HÆMANTHUS. (From the *αἷμα*, blood, and *ανθος*, a flower, so called from its colour.) The blood-flower.

HÆMATEMESIS. (From *αἷμα*, blood, and *εμεσις*, to vomit.) *Vomitus cruentus*.

A vomiting of blood. A vomiting of blood is readily to be distinguished from one which proceeds from the lungs, by its being usually preceded by a sense of weight, pain, or anxiety in the region of the stomach; by its being unaccompanied by any cough; by its being discharged in a very considerable quantity; by its being of a dark colour, and somewhat grumous; and by its being mixed with the other contents of the stomach.

The disease may be occasioned by any thing received into the stomach, which stu-

mulates it violently or wounds it; or may proceed from blows, bruises, or any other cause capable of exciting inflammation in this organ, or of determining too great a flow of blood to it; but it arises more usually as a symptom of some other disease (such as a suppression of the menstrual, hæmorrhoidal flux, or obstructions in the liver, spleen, and other viscera) than as a primary affection.

Hæmatemesis is seldom so profuse as to destroy the patient suddenly, and the principal danger seems to arise, either from the great debility which repeated attacks of the complaint induce, or from the lodgment of blood in the intestines, which by becoming putrid might occasion some other disagreeable disorder.

The appearances to be observed on dissection, where it proves fatal, will depend on the disease of which it has proved symptomatic.

HÆMATITES. (From *αἷμα*, blood, so named from its property of stopping blood, or from its colour.) *Lapis hæmatites*. Bloodstone. An elegant iron ore called bloodstone. Finely levigated, and freed from the grosser parts by frequent washings with water, it has been long recommended in hæmorrhages, fluxes, uterine obstructions, &c. in doses of from one scruple to three or four.

HÆMATINUS. (From *αἷμα*, blood, and *λίθος*, a stone.) An epithet of a collyrium, in which was the bloodstone.

HÆMATOCELE. (From *αἷμα*, blood,

and *κλυσ*, a tumour.) A swelling of the scrotum, or spermatic chord, proceeding from or caused by blood.

The distinction of the different kinds of hæmatocele, though not usually made, is absolutely necessary towards rightly understanding the disease; the general idea, or conception of which, appears to Mr. Pott to be somewhat erroneous, and to have produced a prognostic which is ill founded and hasty. According to this eminent surgeon, the disease properly called hæmatocele is of four kinds: two of which have their seat within the tunica vaginalis testis; one within the albuginea; and the fourth in the tunica communis, or common cellular membrane, investing the spermatic vessels.

In the passing an instrument, in order to let out the water from an hydrocele of the vaginal coat, a vessel is sometimes wounded, which is of such size, as to tinge the fluid pretty deeply at the time of its running out: the orifice becoming close, when the water is all discharged, and a plaster being applied, the blood ceases to flow from thence, but insinuates itself partly into the cavity of the vaginal coat, and partly into the cells of the dartos; making sometimes, in the space of a few hours, a tumour nearly equal in size to the original hydrocele. This is one species.

It sometimes happens in tapping an hydrocele, that although the fluid discharged by that operation be perfectly clear and limpid, yet in a very short space of time (sometimes in a few hours) the scrotum becomes as large as it was before, and palpably as full of a fluid. If a new puncture be now made, the discharge instead of being limpid (as before) is now either pure blood or very bloody. This is another species; but, like the preceding, confined to the tunica vaginalis.

The whole vascular compages of the testicle is sometimes very much enlarged, and at the same time rendered so lax and loose, that the tumour produced thereby has, to the fingers of an examiner, very much the appearance of a swelling composed of a mere fluid, supposed to be somewhat thick, or viscid. This is in some measure a deception; but not totally so: the greater part of the tumefaction is caused by the loosened texture of the testes; but there is very frequently a quantity of extravasated blood also.

If this be supposed to be an hydrocele, and pierced, the discharge will be mere blood. This is a third kind of hæmatocele; and very different, in all its circumstances, from the two preceding: the fluid is shed from the vessels of the glandular part of the testicles, and contained within the tunica albuginea.

The fourth consists of a rupture of, and an effusion of blood, from a branch of the spermatic vein, in its passage from the

groin to the testicle: In which case, the extravasation is made into the tunica communis, or cellular membrane, investing the spermatic vessels.

Each of these species, Mr. Potts says, he has seen so distinctly, and perfectly, that he has not the smallest doubt concerning their existence, and of their difference from each other.

HÆMATOCHYSIS. (From *αἷμα*, blood, and *χύνω*, to pour out.) A hæmorrhage or flux of blood.

HÆMATODES. (From *αἷμα*, blood; so called from the red colour of its flowers.) 1. An old name of the bloody cranesbill. See *Geranium sanguineum*.

2. Now applied to a fungus or fleshy excrescence, which has somewhat the appearance of blood.

HÆMATOLOGY. (From *αἷμα*, blood, and *λογία*, a discourse.) *Hæmatologia*. The doctrine of the blood.

HÆMATOMPHALOCLE. (From *αἷμα*, blood, *μφαλόν*, the navel, and *κλυσ*, a tumour.) A species of ecchymosis. A tumour about the navel from an extravasation of blood. It is mostly absorbed, but, if too considerable, a puncture may be made to evacuate the blood, as in ecchymosis. See *Ecchymosis*.

HÆMATOPEDESIS. (From *αἷμα*, blood, and *πίπτω*, to leap.) The leaping of the blood from a wounded artery.

HÆMOTOSIS. (From *αἷμα*, blood.) An hæmorrhage or flux of blood.

HÆMATOXYLON. (From *αἷμα*, blood, and *ξύλον*, wood; so called from the red colour of its wood.) The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Monogynia*.

HÆMATOXYLUM. (From *αἷμα*, blood, or *αἱματώδες*, bloody, and *ξύλον*, wood.) The name in some pharmacopœias for the logwood; which is so called from its red colour. See *Lignum campechense*.

HÆMATOXYLON CAMPECHIANUM. The systematic name of the logwood-tree. See *Lignum campechianum*.

HÆMATURIA. (From *αἷμα*, blood, and *ουρά*, urine.) The voiding of blood with urine. This disease is sometimes occasioned either by falls, blows, bruises, or some violent exertion, such as hard riding and jumping; but it more usually arises from a small stone lodged either in the kidney or ureter, which, by its size or irregularity, wounds the inner surface of the part it comes in contact with; in which case the blood discharged is most usually somewhat coagulated, and deposits a sediment of a dark-brown colour, resembling the grounds of coffee.

A discharge of blood by urine, when proceeding from the kidney or ureter, is commonly attended with an acute pain in the back, and some difficulty of making water, the urine, which comes away first, being muddy and high coloured, but to-

wards the close of its flowing, becoming transparent and of a natural appearance. When the blood proceeds immediately from the bladder, it is usually accompanied with a sense of heat and pain at the bottom of the belly.

The voiding of bloody urine is always attended with some danger, particularly when mixed with purulent matter. When it arises in the course of any malignant disease, it shews a highly putrid state of the blood, and always indicates a fatal termination.

The appearances to be observed on dissection will accord with those usually met with in the disease which has given rise to the complaint.

HÆMODIA. (From *αἷμα*, to stupefy.) A painful stupor of the teeth, caused by acrid substances touching them.

HÆMOTOE. (From *αἷμα*, blood, and *τοῦ*, to spit up.) The spitting of blood. See *Hæmoptysis*.

HÆMLOPTYSIS. (From *αἷμα*, blood, *πύσις*, to spit.) *Hæmoptoe*. A spitting of blood. A genus of disease arranged by Cullen in the class of *pyrexia*, and order *hæmorrhagia*. It is characterized by coughing up florid or frothy blood, preceded usually by heat or pain in the chest, irritation in the larynx, and a saltish taste in the mouth. There are five species of this disease: 1. *Hæmoptysis plethorica*, from fulness of the vessels. 2. *Hæmoptysis violenta*, from some external violence. 3. *Hæmoptysis phthisica*, from ulcers corroding the small vessels. 4. *Hæmoptysis calculosa*, from calculous matter in the lungs. 5. *Hæmoptysis vacaria*, from the suppression of some customary evacuation.

It is readily to be distinguished from hæmatemesis, as in this last the blood is usually thrown out in considerable quantities; and is, moreover, of a darker colour, more grumous, and mixed with the other contents of the stomach; whereas blood proceeding from the lungs is usually in small quantity, of a florid colour, and mixed with a little frothy mucus only.

A spitting of blood arises most usually between the ages of 16 and 25, and may be occasioned by any violent exertion either in running, jumping, wrestling, singing loud, or blowing wind-instruments; as likewise by wounds, plethora, weak vessels, hectic fever, coughs, irregular living, excessive drinking, or the suppression of some accustomed discharge, such as the menstrual or hæmorrhoidal. It may likewise be occasioned by breathing air which is too much rarefied to be able properly to expand the lungs.

Persons in whom there is a faulty proportion either in the vessels of the lungs, or in the capacity of the chest, being distinguished by a narrow thorax and prominent shoulders, or who are of a delicate make and sanguine temperament, seem much

disposed to this hæmorrhage; but in these, the complaint is often brought on by the concurrence of the various occasional and exciting causes before mentioned.

A spitting of blood is not, however, always to be considered as a primary disease. It is often only a symptom, and in some disorders, such as pleuritis, peripneumonies, and many fevers, often arises, and is the presage of a favourable termination.

Sometimes it is preceded (as has already been observed) by a sense of weight and oppression at the chest, a dry tickling cough, and some slight difficulty of breathing. Sometimes it is ushered in with shiverings, coldness at the extremities, pains in the back and loins, flatulency, costiveness, and lassitude. The blood which is spit up is sometimes thin, and of a florid red colour; and at other times it is thick, and of a dark or blackish cast; nothing, however, can be inferred from this circumstance, but that the blood has lain a longer or shorter time in the breast, before it was discharged.

An hæmoptoe is not attended with danger, where no symptoms of phthisis pulmonalis have preceded, or accompanied the hæmorrhage, or where it leaves behind no cough, dyspnea, or other affection of the lungs; nor is it dangerous in a strong healthy person, of a sound constitution; but when it attacks persons of a weak lax fibre, and delicate habit, it may be difficult to remove it.

It seldom takes place to such a degree as to prove fatal at once; but when it does, the effusion is from some large vessel. The danger, therefore, will be in proportion as the discharge of blood comes from a large vessel, or a small one.

When the disease proves fatal, in consequence of the rupture of some large vessels, there is found, on dissection, a considerable quantity of clotted blood betwixt the lungs and pleura, and there is usually more or less of an inflammatory appearance at the ruptured part. Where the disease terminates in pulmonary consumption, the same morbid appearances are to be met with as described under that particular head.

HÆMORRHAGIA. (From *αἷμα*, blood, and *ρρῆσις*, to break out.) A hæmorrhage; or flow of blood.

HÆMORRHAGIÆ. Hæmorrhages, or affluxes of blood. An order in the class *pyrexia* of Cullen's nosology is so called. It is characterized by pyrexia with a discharge of blood, without any external injury; the blood on venæsection exhibiting the buffy coat. The order *hæmorrhagia* contains the following genera of diseases, viz. epistaxis, hæmoptysis, phthisis, hæmorrhoids and menorrhagia.

HÆMORRHOIDAL ARTERIES. *Arterie hæmorrhoidales*. The arteries of the rectum are so called: they are sometimes two, and at other times three, in number.

1. The upper hæmorrhoidal artery, which is the great branch of the lower mesenteric continued into the pelvis. 2. The middle hæmorrhoidal, which sometimes comes off from the hypogastric artery, and very often from the pudical artery. It is sometimes wanting. 3. The lower or external hæmorrhoidal is almost always a branch of the pudical artery, or that artery which goes to the penis.

HÆMORRHOIDAL VEINS. *Vene Hæmorrhoidales.* These are two. 1. The external, which evacuates itself into the vena iliaca interna. 2. The internal, which conveys its blood into the vena portæ.

HÆMORRHOIS. (From *aqua*, blood, and *gao*, to flow.) *Æmorrhoids.* The piles. A genus of disease in the class *pyrexia* and order *hæmorrhagie* of Cullen. They are certain exorecences or tumours arising about the verge of the anus, or the inferior part of the intestinum rectum; when they discharge blood, particularly upon the patient's going to stool, the disease is known by the name of *bleeding piles*: but when there is no discharge, it is called *blind piles*. The rectum, as well as the colon, is composed of several muscular membranes, connected to each other by an intervening cellular substance; and as the muscular fibres of this intestine always tend, by their contraction, to lessen its cavity, the internal membrane, which is very lax, forms itself into several rugæ or folds. In this construction nature respects the use of the part, which occasionally gives passage to or allows the retention of the excrements, the hardness and bulk of which might produce considerable lacerations, if this intestine were not capable of dilatation. The arteries and veins subservient to this part are called hæmorrhoidal, and the blood that returns from hence is carried to the meseraic veins. The intestinum rectum is particularly subject to the hæmorrhoids, from its situation, structure, and use; for whilst the course of the blood is assisted in almost all the other veins of the body, by the distension of the adjacent muscles, and the pressure of the neighbouring parts, the blood in the hæmorrhoidal veins, which is to ascend against the natural tendency of its own weight, is not only destitute of these assistances, but is impeded in its passage: for, first, the large excrements which lodge in this intestine dilate its sides, and the different resistances which they form there are so many impediments obstructing the return of the blood; not in the large veins, for they are placed along the external surface of the intestine, but in all the capillaries which enter into its composition. Secondly, as often as these large excrements, protruded by others, approach near the anus, their successive pressure upon the internal coats of the intestine, which they dilate, drives back the blood into the veins,

and for so long suspends its course; the necessary consequence of which is, a distension of the veins in proportion to the quantity of blood that fills them. Thirdly, in every effort we make, either in going to stool, or upon any other occasion, the contraction of the abdominal muscles, and the diaphragm pressing the contents of the abdomen downwards, and these pressing upon the parts contained in the pelvis, another obstruction is thereby opposed to the return of the blood, not only in the large veins, but also in the capillaries, which, being of too weak a texture to resist the impulse of the blood that always tends to dilate them, may thereby become varicose.

The dilatation of all these vessels is the *primary cause* of the hæmorrhoids; for the internal coat of the intestine, and the cellular membrane which connects that to the muscular coat, are enlarged in proportion to the distension of the vessels of which they are composed. This distension, not being equal in every part, produces separate tumours in the gut, or at the verge of the anus, which increases according as the venal blood is obstructed in them, or circulates there more slowly.

Whatever, then, is capable of retarding the course of the blood in the hæmorrhoidal veins, may occasion this disease. Thus, persons that are generally constive, who are accustomed to sit long at stool, and strain hard; pregnant women, or such as have had difficult labours; and likewise persons who have an obstruction in their liver; are for the most part afflicted with the piles; yet every one has not the hæmorrhoids, the different causes which are mentioned above being not common to all, or at least not having in all the same effects. When the hæmorrhoids are once formed, they seldom disappear entirely, and we may judge of those within the rectum by those, which, being at the verge of the anus, are plainly to be seen. A small pile, that has been painful for some days, may cease to be so, and dry up; but the skin does not afterwards retain its former firmness, being more lax and wrinkled, like the empty skin of a grape. If this external pile swells and sinks again several times, we may perceive, after each return, the remains of each pile, though shrivelled and decayed, yet still left larger than before. The case is the same with those that are situated within the rectum; they may happen indeed never to return again, if the cause that produced them is removed; but it is probable that the excrements in passing out occasion a return of the swelling, to which the external ones are less liable: for the internal piles make a sort of knots or tumours in the intestine, which straightening the passage, the excrements, in passing out, occasion irritations there, that are more or less painful in proportion to the efforts which the person

makes in going to stool; and it is thus these tumours become gradually larger. The hæmorrhoids are subject to many variations; they may become inflamed from the above irritations to which they are exposed, and this inflammation cannot always be removed by art. In some, the inflammation terminates in an abscess, which arises in the middle of the tumour, and degenerates into a fistula. These piles are very painful till the abscess is formed. In others, the inflammation terminates by induration of the hæmorrhoid, which remains in a manner schirrous. These never lessen, but must necessarily grow larger. This schirrous sometimes ulcerates, and continually discharges a sanies, which the patient perceives by stains on his shirt, and by its occasioning a very troublesome itching about the verge of the anus. These kinds of hæmorrhoids sometimes turn cancerous. There are some hæmorrhoids, and those of different sizes, which are covered with so fine a skin as frequently to admit blood to pass through. This fine skin is only the internal coat of the rectum, greatly attenuated by the varicose distension of its vessels. The hæmorrhage may proceed from two causes; namely, either from an excoriation produced by the hardness of the excrements, or from the rupture of the tumefied vessels, which break by their too great distension. In some of these, the patient voids blood almost every time he goes to stool; in others not so constantly. We sometimes meet with men who have a periodical bleeding by the piles, not unlike the menses in women; and as this evacuation, if moderate, does not weaken the constitution, we may infer that it supplies some other evacuation, which nature either ceases to carry on, or does not furnish in due quantity; and hence also we may explain why the suppression of this discharge, to which nature had been accustomed, is frequently attended with dangerous diseases. The hæmorrhoids are sometimes distended to that degree as to fill the rectum, so that if the excrements are at all hard they cannot pass. In this case the excrements force the hæmorrhoids out of the anus, to procure a free passage, consequently the internal coat of the rectum, to which they are connected, yields to extension, and upon examining these patients immediately after having been at stool, a part of the internal coat of that gut is perceived forming a sort of ligature or stricture round the hæmorrhoids. A difficulty will occur in the return of these, in proportion to their size, and as the verge of the anus is more or less contracted. If the bleeding piles come out in the same manner upon going to stool, it is then they void most blood, because the verge of the anus forms a kind of ligature above them.

HÆMOSTASIA. (From *αἷμα*, blood, and *ἵσθαι*, to stand.) A stagnation of blood.

HÆMOSTATICA. (From *αἷμα*, blood, and *σταῖν*, to stop.) Styptics. Medicines which stop hæmorrhages.

HAGIOSPERMUM. (From *ἅγιος*, holy, and *σπέρμα*, seed, so called from its reputed virtues.) Wormseed.

HAGIOXYLUM. (From *ἅγιος*, holy, and *ξύλον*, wood, so named because of its medicinal virtues.) Guaiacum.

HAIR. *Pili. Capilli.* The hairs of the human body are thin, elastic, dry filaments, arising from the skin. They consist of the *bulb*, situated under the skin, which is a vascular and nervous vesicle; and a *trunk*, which perforates the skin and cuticle, and is covered with a peculiar vagina. The colour of hair varies; its seat, however, is in the medullary juice. The hair, according to its situation, is differently named; thus, on the head it is called *capilli*; over the eyes, *supercilia*; *cilia*, on the margin of the eyelids; *vibrissæ*, in the foramina of the nostrils; *pili auriculares*, in the external auditory passage; *mustax*, on the upper lip; and *barba*, on the lower jaw.

HALATUM. (From *αἷς*, salt,) A clyster composed chiefly of salt.

HALCHEMIA. (From *αἷς*, salt, and *χεῖν*, to pour out.) The art of fusing salts.

HALELEUM. (From *αἷς*, salt, and *λαλεῖν*, oil.) A medicine composed of salt and oil.

HALICABAUM. (From *αἷς*, the sea, and *καλαβός*, night-shade; so called because it grows upon the banks of the sea.) See *Alkekengi*.

HALINUS. (From *αἷς*, the sea, because of its saltish taste.) Sea-purslane, said to be antispasmodic.

HALINITRUM. (From *αἷς*, the sea, and *νίτρον*, nitre.) Nitre, or rather rock salt.

HALITUS. (From *halito*, to breathe out.) A vapour or gaz.

HALLUCINATIO. (From *hallucino*, to err.) A depraved or erroneous imagination.

HALO. (From *αἷος*, an area or circle.) The red circle surrounding the nipple, which becomes somewhat brown in old people, and is beset with many sebaceous glands.

HALMYRONES. (From *αἷμυρος*, salted.) A term applied to the humours; it means acrimonious. It is also applied to fevers which communicate such an itching sensation as is perceived from handling salt substances.

HAMALGAMA. See *Amalgam*.

HAMPSTEAD WATER. A good chalybeate mineral water in the neighbourhood of London.

HAMULUS. (Dim. of *hamus*, a hook.) *Hamus*. A term in anatomy, applied to any hook-like process, as the hamulus

of the pterygoid process of the sphænoid bone.

HAMUL. See *Hamulus*.

HAND. *Mamus*. The hand is composed of the carpus or wrist, metacarpus, and fingers. The arteries of the hand are the *palmary arch* and the *digital arteries*. The veins are the *digital*, the *cephalic of the thumb*, and the *salvarella*. The nerves are the *cutaneous externus* and *internus*.

HARDESIA. See *Lapis Hibernicus*.

HARE LIP. *Lagocheilus*. *Labia leporina*. A fissure or longitudinal division of one or both lips. Children are frequently born with this kind of malformation, particularly of the upper lip. Sometimes the portions of the lip, which ought to be united, have a considerable space between them; in other instances they are not much apart. The cleft is occasionally double, there being a little lobe, or small portion of the lip, situated between the two fissures. Every species of the deformity has the same appellation of hare-lip, in consequence of the imagined resemblance which the part has to the upper lip of a hare.

The fissure commonly affects only the lip itself. In many cases, however, it extends along the bones of the palate, even as far as the uvula. Sometimes these bones are totally wanting; sometimes they are only divided by a fissure.

Such a malformation is always peculiarly afflicting. In its least degree, it constantly occasions considerable deformity; and when it is more marked, it frequently hinders infants from sucking, and makes it indispensable to nourish them by other means. When the lower lip alone is affected, which is not most frequently the case, the child can neither retain its saliva, nor learn to speak, except with the greatest impediment. But when the fissure pervades the palate, the patient not only never articulates but very imperfectly, but cannot masticate nor swallow, except with great difficulty, on account of the food readily getting up into the nose.

HARMONIA. (From *apo*, to sit together.) Harmony. A species of synarthrosis, or immoveable connection of bones, in which bones are connected together by means of rough margins, not dentiform; in this manner most of the bones of the face are connected together.

HARROGATE WATER. The villages of High and Low Harrogate are situate in the centre of the county of York, adjoining the town of Knaresborough. The whole of Harrogate, in particular, has long enjoyed considerable reputation, by possessing two kinds of very valuable springs; and, some years ago, the chalybeate was the only one that was used internally, whilst the sulphureous water was confined to external use. At present, however, the

latter is employed largely as an internal medicine.

The sulphureous springs of Harrogate are four in number, of the same quality, though different in the degree of their powers. This water, when first taken up, appears perfectly clear and transparent, and sends forth a few air bubbles, but not in any quantity. It possesses a very strong sulphureous and fetid smell, precisely like that of a damp rusty gun-barrel, or bilgewater. To the taste it is bitter, nauseous, and strongly saline, which is soon borne without any disgust. In a few hours of exposure this water loses its transparency, and becomes somewhat pearly, and rather greenish to the eye; its sulphureous smell abates, and at last the sulphur is deposited in the form of a thin film, on the bottom and sides of the vessel in which it is kept. The volatile productions of this water shew carbonic acid, sulphurated hydrogen, and azotic gas.

The sensible effects which this water excites, are often a head-ach and giddiness on being first drank, followed by a purgative operation, which is speedy and mild, without any attendant gripes; and this is the only apparent effect the exhibition of this water displays.

The diseases in which this water is used are numerous, particularly of the alimentary canal, and irregularity of the bilious secretion. Under this water the health, appetite, and spirits improve; and, from its opening effects, it cannot fail to be useful in the costive habit of hypochondriasis. But the highest recommendation of this water has been in cutaneous diseases, and for this purpose it is universally employed, both as an internal medicine, and an external application: in this united form, it is of particular service in the most obstinate and complicated forms of cutaneous affections; nor is it less so in states and symptoms supposed connected with worms, especially with the round worm and ascarides, when taken in such a dose as to prove a brisk purgative; and in the latter case also, when used as a clyster, the ascarides being chiefly confined to the rectum, and therefore within the reach of this form of medicine. From the union of the sulphureous and saline ingredients, the benefit of its use has been long established in hæmorrhoidal affections.

A course of Harrogate waters should be conducted so as to produce sensible effects on the bowels; half a pint taken in the morning, and repeated three or four times, will produce it, and its nauseating taste may be corrected by taking a dry biscuit, or a bit of coarse bread, after it. The course must be continued, in obstinate cases, a period of some months, before a cure can be expected.

HARTFELL WATER. This mineral

water is near Moffat, in Scotland. it is a chalybeate dissolved by the sulphuric acid, and is much celebrated in scrophulous affections, and cutaneous diseases. It is used no less as an external application, than drank internally. The effects of this water, at first, are some degree of drowsiness, vertigo, and pain in the head, which soon go off, and may be hastened by a slight purge. It produces generally a flow of urine, and an increase of appetite. It has acquired much reputation also in old and languid ulcers, where the texture of the diseased part is very lax, and the discharge profuse and ill conditioned.

The dose of this water is more limited than that of most of the mineral springs which are used medicinally. It is of importance in all cases, and especially in delicate and irritable habits, to begin with a very small quantity, for an over dose is apt to be very soon rejected by the stomach, or to occasion griping and disturbance in the intestinal canal; and it is never as a direct purgative that this water is intended to be employed. Few patients will bear more than an English pint in the course of the day; but this quantity may be long continued in. It is often advisable to warm the water, for delicate stomachs, and this may be done without occasioning any material change in its properties.

Hartshorn. See *Cornu cervi*.

Hartshorn shavings. See *Cornu cervi*.

Hart's tongue. See *Scolopendrium*.

Hart-wort. See *Seneli*.

Hart-wort of Marseilles. See *Seneli Marseillense*.

Hay, came's. See *guncus odoratus*.

HEAD. *Caput.* The superior part of the body placed upon the neck, containing the cerebrum, cerebellum, and medulla oblongata. It is divided into the *face* and *hairy* part. On the latter is observed the *vertex*, or crown of the head; the *sinciput*, or fore part; the *occiput*, or hinder part. For the former, see *Face*. For the bones and muscles of the head, see *Cranium*. The common integuments of the head are called the scalp.

HEARING. *Auditus.* This sense is placed, by physiologists, among the animal actions. It is a sensation by which we hear the sound of sonorous bodies. The organ of hearing is the soft portion of the auditory nerve which is distributed on the vestibule, semicircular canals, and cochlea.

HEART. *Cor.* A hollow muscular viscus, situated in the cavity of the pericardium for the circulation of the blood. It is divided externally into a *base*, or its broad part; a *superior* and an *inferior surface*, and an *anterior* and *posterior* margin. Internally, it is divided into a *right* and *left ventricle*. The situation of the heart is oblique, not transverse; its base being

placed on the right of the bodies of the vertebrae, and its apex obliquely to the sixth rib on the left side; so that the left ventricle is almost posterior, and the right anterior. Its inferior surface lies upon the diaphragm. There are two cavities adhering to the base of the heart, from their resemblance called *auricles*. The right auricle is a muscular sac, in which are four *apertures*, two of the *venae cavae*, an opening into the right ventricle, and the opening of the coronary vein. The left is a similar sac, in which there are five *apertures*, viz. those of the four pulmonary veins, and an opening into the left ventricle. The cavities in the base of the heart are called *ventricles*: these are divided by a fleshy septum, called *septum cordis*, into a right and left. Each ventricle has two *orifices*; the one auricular, through which the blood enters, the other arterious, through which the blood passes out. These four orifices are supplied with *valves*, which are named from their resemblance; those at the arterious orifices are called the *semilunar*; those at the orifice of the right auricle, *mitral*; and those at the orifice of the left auricle, *tricuspid*. The *valve of Eustachius* is situated at the termination of the *vena cava inferior*, just within the auricle. The substance of the heart is muscular, its exterior fibres are longitudinal, its middle transverse, and its anterior oblique. The internal superficies of the ventricles and auricles of the heart are invested with a strong and smooth membrane, which is extremely irritable. The vessels of the heart are divided into *common* and *proper*. The *common* are, 1. The *aorta*, which arises from the left ventricle. 2. The *pulmonary artery*, which originates from the right ventricle. 3. The four pulmonary veins, which terminate in the left auricle. 4. The *vena cavae*, which evacuate themselves into the right auricle. The *proper vessels* are, 1. The *coronary arteries*, which arise from the aorta, and are distributed on the heart. 2. The *coronary veins*, which return the blood into the right auricle. The *nerves* of the heart are branches of the eighth and great intercostal pairs. The heart of the fœtus differs from that of the adult, in having a *foramen ovale*, through which the blood passes from the right auricle to the left.

Heart's ease. See *Viola tricolor*.

Heat. See *Caloric*.

HEAT, ABSOLUTE. This term is applied to the whole quantity of caloric existing in a body in chymical union.

Heat, animal. See *Animal heat*.

HEAT, FREE. If the heat which exists in any substance be from any cause forced in some degree to quit that substance, and to combine with those that surround it, then such heat is said to be free, or sensible, until the equilibrium is restored.

HEAT, LATENT. When any body is in

equilibrium with the bodies which surround it with respect to its heat, that quantity which it contains is not perceptible by any external sign or organ of sense, and is termed combined caloric, or latent heat.

Heat, sensible. See *Heat, free.*

Heavy carbonated hydrogen gas. See *Carbonated hydrogen gas, heavy.*

Ectic fever. (From *εκτε*, habit.) See *Febris hectica.*

HEDERA. (From *hæreo*, to stick, because it attaches itself to trees and old walls.) The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*. The ivy.

HEDERA ARBorea. The ivy. The leaves of this tree, *Hedera helix* of Linnæus, have little or no smell, but a very nauseous taste. Haller informs us, that they are recommended in Germany against the atrophy of children. By the common people of this country they are sometimes applied to running sores, and to keep issues open. The berries were supposed by the ancients to have a purgative and emetic quality; and an extract was made from them by water, called by Quercetanus *extractum purgans*. Later writers have recommended them in small doses as alexipharmic and sudorific: it is said, that in the plague at London, the powder of them was given in vinegar, or white wine, with good success. It is from the stalk of this tree that a resinous juice exudes very plentifully in warm climates. See *Gummi hederæ*.

HEDERA HELIX. The systematic name of the ivy-tree. See *Hedera arborea*.

HEDERA TERRESTRIS. Ground-ivy, or gill. *Glechoma hederacea* of Linnæus:—*foliis reniformibus crenatis*. This indigenous plant has a peculiar strong smell, and a bitterish somewhat aromatic taste. It is one of those plants which was formerly much esteemed for possessing virtues that, in the present age, cannot be detected. In obstinate coughs it is a favourite remedy with the poor.

Hedge hyssop. See *Gratiola*.

Hedge mustard. See *Erysimum*.

Hedge mustard, stinking. See *Alliaria*.

HEDRA. The anus. Excrement. A fracture.

HEDYOSMOS. Mint.

HELICOMA. Ulceration.

HELCONIA. (From *ελκος*, an ulcer.) An ulcer in the external or internal superficies of the cornea, known by an excavation and oozing of purulent matter from the cornea.

HELICYDRION. (From *ελκος*, an ulcer, and *ιδρω*, water.) *Helcydrium*. A moist ulcerous pustule.

HELICTER. (From *ελκε*, to draw.) An instrument for extracting the fæcus.

HELENIUM. (From *Helène*, the island where they grew.) See *Enula campana*.

HELICALIS MAJOR. See *Helicis major*.

HELICALIS MINOR. See *Helicis minor*.

HELICIS MAJOR. A proper muscle of the ear, which depresses the part of the cartilage of the ear into which it is inserted; it lies upon the upper or sharp point of the helix, or outward ring, arising from the upper and acute part of the helix anteriorly, and passing to be inserted into its cartilage a little above the tragus.

HELICIS MINOR. A proper muscle of the ear, which contracts the fissure of the ear: it is situated below the helicis major, upon part of the helix. It arises from the inferior and anterior part of the helix, and is inserted into the crus of the helix, near the fissure in the cartilage opposite to the concha.

HELIOTROPII SUCCUS. See *Bezzetia cerulea*.

HELIX. (Ελξ, from *ελα*, to turn about.) The external circle or border of the outer ear, that curls inwards.

HELLEBORASTER. (From *ελληβορ*, hellebore.) Fetid hellebore, or bear's foot. *Helleborus fetidus* of Linnæus: *caule multifloro folioso; foliis pedatis*. The leaves of this indigenous plant are recommended by many as possessing extraordinary anthelmintic powers. The smell of the recent plant is extremely fetid, and the taste is bitter and remarkably acrid, inasmuch that, when chewed, it excoriates the mouth and fauces. It commonly operates as a cathartic, sometimes as an emetic, and, in large doses, proves highly deleterious.

Hellebore, black. See *Helleborus niger*.

Hellebore, white. See *Helleborus albus*.

HELLEBORUS. (Ελληβορος: *παρα το τριβον* ελκεν, because it destroys, if eaten.) The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Polygynia*. Hellebore.

HELLEBORUS ALBUS. *Veratrum album*. *Elleborum album*. White hellebore, or veratrum. *Veratrum album* of Linnæus:—*racemo supradecomposito, corollis erectis*. This plant is a native of Italy, Switzerland, Austria, and Russia. Every part of the plant is extremely acrid and poisonous. The dried root has no particular smell, but a durable, nauseous, and bitter taste, burning the mouth and fauces: when powdered, and applied to issues, or ulcers, it produces griping and purging; if snuffed up the nose, it proves a violent sternutatory. Gesner made an infusion of half an ounce of this root with two ounces of water; of this he took two drachms, which produced great heat about the scapulæ and in the face and head, as well as the tongue and throat, followed by singultus, which continued till vomiting was excited. Bergius also experienced very distressing symptoms, upon tasting this infusion. The root, taken in large doses, discovers such acrimony, and operates by the stomach and rectum with such violence, that blood

is usually discharged; it likewise acts very powerfully upon the nervous system, producing great anxiety, tremors, vertigo, syncope, aphonia, interrupted respiration, sinking of the pulse, convulsions, spasms, and death. Upon opening those who have died of the effects of this poison, the stomach discovered marks of inflammation, with corrosions of its internal coat. The ancients exhibited this active medicine in maniacal cases, and it is said with success. The experience of Greding is somewhat similar: out of twenty-eight cases, in which he exhibited the bark of the root collected in the spring, five were cured. In almost every case that he relates, the medicine acted more or less upon all the excretions; vomiting and purging were very generally produced, and the matter thrown off the stomach was constantly mixed with bile; a florid redness frequently appeared on the face, and various cutaneous efflorescences upon the body; and, in some, pleuritic symptoms, with fever, supervened, so as to require bleeding; nor were the more alarming affections of spasms and convulsions unfrequent. Critical evacuations were also very evident; many sweating profusely, in some the urine was considerably increased, in others the saliva and mucous discharges: and uterine obstructions, of long duration, were often removed by its use. Veratrum has likewise been found useful in epilepsy, and other convulsive complaints: but the diseases in which its efficacy seems least equivocal are those of the skin, as itch, and different prurient eruptions, herpes, morbus pediculosus, lepra, scrofula, &c.; and in many of these it has been successfully employed, both internally and externally. As a powerful stimulant and irritating medicine, its use has been resorted to in desperate cases only; and even then it ought first to be exhibited in very small doses, as a grain, and in a diluted state, and to be gradually increased, according to the effects, which are generally of an alarming nature.

HELLEBORUS FORTIUS. The systematic name of the fetid hellebore. See *Helleboraster*.

HELLEBORUS NIGER. *Melampodium.* Black hellebore, or Christmas rose. *Helleborus niger* of Linnæus:—*scapo subnudo; foliis pedatis.* The root of this exotic plant is the part employed medicinally: its taste, when fresh, is bitterish, and somewhat acrid: it also emits a nauseous acrid smell; but, being long kept, both its sensible qualities and medicinal activity suffer very considerable diminution. The ancients esteemed it as a powerful remedy in maniacal cases. At present it is exhibited principally as an alterative, or, when given in a large dose, as a purgative. It often proves a very powerful emmenagogue in plethoric habits,

where steel is ineffectual, or improper. It is also recommended in dropsies, and some cutaneous diseases.

Helmet-flower, yellow. See *Anthora*.

HELMINTHAGOGUES. (*Helminthagoza*; from *αλμντ*, a worm, and *αγω*, to drive out.) Medicines which destroy and expel worms. See *Anthelmintics*.

HELMINTHIASIS. (*Ελμινθιασμος*, from *αλμνθις*, which signifies any species of worm.) A disease in which worms, or the larvæ of worms, are bred under the skin, or some external part of the body. It is endemial to Martinique, Westphalia, Transylvania, and some other places.

HELMINTHOCHORTON. See *Carolina corsicana*.

HELODES. (From *ελος*, a marsh.) A term applied to fevers generated from marsh miasma.

HELOSIS. (From *ελαω*, to turn.) An aversion to turning up of the eyelids.

HELIXE. (From *ελαω*, to draw; so called because it sticks to whatever it touches.) Pellitory of the wall.

HEMALOPIA. Corruptly written for *hemerolopia*.

HEMATURIA. See *Hematuria*.

HEMERALOPS. (From *ημερα*, the day, and *οφθαλμος*, eye.) One who can see but in the day-time.

HEMEROLOPIA. (From *ημερα*, a day, and *οφθαλμος*, to see.) A defect in the sight, which consists in being able to see in the day-time, but not in the evening. The following is Scarpa's description of this curious disorder. Hemeralopia, or *nocturnal blindness*, is properly nothing but a kind of imperfect periodical amaurosis, most commonly sympathetic with the stomach. Its paroxysms come on towards the evening, and disappear in the morning. The disease is endemic in some countries, and epidemic, at certain seasons of the year, in others. At sun-set, objects appear to persons affected with this complaint as if covered with an ash-coloured veil, which gradually changes into a dense cloud, which intervenes between the eyes and surrounding objects. Patients with hemerolopia have the pupil, both in the day and night time, more dilated, and less moveable, than it usually is in healthy eyes. The majority of them, however, have the pupil more or less moveable in the day-time, and always expanded and motionless at night. When brought into a room faintly lighted by a candle, where all the bystanders can see tolerably well, they cannot discern at all, or in a very feeble manner, scarcely any one object; or they only find themselves able to distinguish light from darkness, and at moon-light their sight is still worse. At day-break they recover their sight, which continues perfect all the rest of the day till sun-set.

HEMICRANIA. (From *ἡμικρανία*, half, and *κεφαλή*, the head.) A pain that affects only one side of the head.

HEMICERAUNIOS. (From *ἡμικρανία*, half, and *κομίζω*, to cut; so called because it was cut half way down. A bandage for the back and breast.

HEMICRANIA CLAVUS. Pain resembling the driving a nail into the head.

HEMIOPSIA. (From *ἡμιόψια*, half, and *ὄψις*, an eye.) A defect of vision, in which the person sees the half, but not the whole of an object.

HEMIPAGIA. (From *ἡμικρανία*, half, and *παῖς*, fixed.) A fixed pain on one side of the head.

HEMIPLEGIA. (From *ἡμικρανία*, half, and *πλῆξις*, to strike.) A paralytic affection of one side of the body. See *Paralysis*.

Hemlock. See *Cicuta*.

Hemlock, dropwort. See *Oenanthe*.

Hemlock, water. See *Cicuta aquatica*.

Hemorrhage from the lungs. See *Hæmoptysis*.

Hemorrhage from the nose. See *Epistaxis*.

Hemorrhage from the penis. See *Hæmaturia*.

Hemorrhage from the stomach. See *Hæmatemesis*.

Hemp. See *Cannabis*.

Hemp-agrimony. See *Eupatorium*.

Hemp, water. See *Eupatorium*.

Hembane. See *Hyosciamus*.

HEPAR. (*ἥπαρ*, the liver.) See *Liver*.

HEPAR SULPHURIS. Liver of sulphur. This is a *sulphuret* made either with potash or soda. It has a disagreeable fetid smell, but it is in high esteem as a medicine to decompose corrosive sublimate, when taken into the stomach. See *Sulphuretum potassæ*.

HEPAR UTERINUM. The placenta.

HEPATALGIA. (From *ἥπαρ*, the liver, and *ἄλγος*, pain.) Pain in the liver.

HEPATIC. Any thing belonging to the liver.

HEPATIO AIR. Hepatic airs consist of inflammable air combined with sulphur, which exists in very different proportions. Where each ingredient is combined merely to saturation, it is called simply *sulphurated hydrogen*; but where the sulphur is in excess, it is termed *super-sulphurated hydrogen*. Sulphurated hydrogen, combined with any base, forms a *hydrosulphuret*, and may be also called an *hepatule*, to distinguish it from an *hepar*, which is the union of sulphur singly with a basis.

HEPATIC ARTERY. *Arteria hepatica.* The artery which nourishes the substance of the liver. It arises from the cæliac, where it almost touches the point of the *lobulus Spigelii*. Its root is covered by the pancreas; it then turns a little forwards, and passes under the pylorus to the porta

of the liver, and runs betwixt the biliary ducts and the vena portæ, where it divides into two large branches, one of which enters the right, and the other the left lobe of the liver. In this place it is enclosed along with all the other vessels in the capsule of Glisson.

HEPATIC DUCT. *Ductus hepaticus.* The trunk of the biliary pores. It runs from the sinus of the liver towards the duodenum, and is joined by the cystic duct, to form the ductus communis choledochus, See *Biliary ducts*.

HEPATIC VEINS. See *Cavæ hepaticæ*, and *Vena portæ*.

HEPATICÆ. (From *ἥπαρ*, the liver; so called because it was thought to be useful in diseases of the liver.) The herb liverwort. See *Hepatica terrestris*.

HEPATICÆ NORILIS. *Herba trinitatis.* Hepatica, or herb trinity. This plant *Anemone hepatica* of Linnaeus, possesses mildly adstringent and corroborant virtues, with which intentions infusions of it have been drank as tea, or the powder of the dry leaves given, to the quantity of half a spoonful at a time.

HEPATICÆ TERRESTRIS. *Jecoraria.* Liverwort. This is the *Marchantia polymorpha*, which is very common in this country. It has a penetrating though mild pungency, and bitter taste, sinking, as it were, into the tongue. It is recommended as an aperient, resolvent, and antiscorbutic, and, though seldom used in this country, appears to be a plant of no inconsiderable virtue.

HEPATIRRHÆA. (From *ἥπαρ*, the liver, and *ῥέω*, to flow.) A diarrhœa, in which portions of flesh, like liver, are voided.

HEPATITIS. (From *ἥπαρ*, the liver.) *Inflammatio hepatis.* An inflammation of the liver. A genus of disease in the class *pyrexia*, and order *phlegmasia*, of Cullen, who defines it, "febrile affection, attended with tension and pain of the right hypochondrium, often pungent, like that of a pleurisy, but more frequently dull, or obtuse; a pain at the clavicle and at the top of the shoulder of the right side; much uneasiness in lying down on the left side; difficulty of breathing; a dry cough, vomiting, and hiccough."

Hepatitis has generally been considered of two kinds; one the *acute*, the other *chronic*.

Besides the causes producing other inflammations, such as the application of cold, external injuries from contusions, blows, &c. this disease may be occasioned by certain passions of the mind, by violent exercise, by intense summer heats, by long continued intermittent and remittent fevers, and by various solid concretions in the substance of the liver. In warm climates,

this viscus is more apt to be affected with inflammation than perhaps any other part of the body, probably from the increased secretion of bile which takes place when the blood is thrown on the internal parts, by an exposure to cold; or from the bile becoming acrid, and thereby exciting an irritation in the part.

The *acute* species of hepatitis comes on with a pain in the right hypochondrium, extending up the clavicle and shoulder, which is much increased by pressing upon the part, and is accompanied with a cough, oppression of breathing, and difficulty of laying on the side affected; together with nausea and sickness, and often with a vomiting of bilious matter. The urine is of a deep saffron colour, and small in quantity; there is loss of appetite, great thirst, and costiveness, with a strong, hard, and frequent pulse; and when the disease has continued for some days, the skin and eyes become tinged of deep yellow.

The *chronic* species is usually accompanied with a morbid complexion, loss of appetite and flesh, costiveness, indigestion, flatulency, pains in the stomach, a yellow tinge of the skin and eyes, clay-coloured stools, high-coloured urine, depositing a red sediment and ropy mucus; an obtuse pain in the region of the liver, extending to the shoulder, and not unfrequently with a considerable degree of asthma.

These symptoms are, however, often so mild and insignificant as to pass almost unnoticed; as large abscesses have been found in the liver, upon dissection, which in the person's life-time had created little or no inconvenience, and which we may presume to have been occasioned by some previous inflammation.

Hepatitis, like other inflammations, may end in resolution, suppuration, gangrene, or scirrhus; but its termination in gangrene is a rare occurrence.

The disease is seldom attended with fatal consequences of an immediate nature, and is often carried off by hæmorrhage from the nose, or hæmorrhoidal vessels, and likewise by sweating, by a diarrhoea, or by an evacuation of urine, depositing a copious sediment. In a few instances, it has been observed to cease on the appearance of erysipelas in some external part.

When suppuration takes place, as it generally does before this forms an adhesion with some neighbouring part, it is usually discharged by the different outlets with which this part is connected, as by coughing, vomiting, or by an abscess breaking outwardly; but in some instances, the pus has been discharged into the cavity of the abdomen, where no such adhesion has been formed.

On dissection, the liver is often found much enlarged, and hard to the touch; its

colour is more of a deep purple than what is natural, and its membranes are more or less affected by inflammation. Dissections likewise show that adhesions to the neighbouring parts often take place, and that large abscesses, containing a considerable quantity of pus, are often found in its substance.

HEPATITIS CYSTICA. Inflammation of the gall-bladder.

HEPATITIS PARENCHYMATOSA. Inflammation in the inner substance of the liver.

HEPATITIS PERITONÆALIS. Inflammation in the peritonæum covering the liver.

HEPATOCÉLE. (From *πάς*, the liver, and *κύλη*, a tumour.) An hernia, in which a portion of the liver protrudes through the abdominal parietes.

HEPATORIUM. The same as *Eupatorium*.

Hepatile. See *Sulphurated hydrogen gas*.

HEPHÆSTIAS. (From *Ἡφαιστος*, Vulcan, or fire.) A drying plaster of burnt tiles.

HEPIALUS. (From *ἥπιος*, gentle.) A mild quotidian fever.

HEPTAPHARMACUM. (From *ἑπτά*, seven, and *φάρμακον*, medicine.) A medicine composed of seven ingredients, the principal of which were ceruss, litharge, wax, &c.

HEPTAPHYLLOM. (From *ἑπτά*, seven, and *φύλλον*, a leaf; so named because it consists of seven leaves.) See *Tormentilla*.

HEPTAPLURUM. (From *ἑπτά*, seven, and *πλευρά*, a rib: so named from its having seven ribs upon the leaf.) The herb plantain.

HERACLEA. (From *Heraclea*, the city near which it grows.) Water horehound.

HERBA BRITANNICA. See *Hydrolapathum*.

HERBA SACRA. See *Verbena*.

Herb-bennet. See *Caryophyllata*.

Herb-mastich. See *Marum vulgare*.

Herb-of-grace. See *Gratiola*.

Herb-trinity. See *Hepatica nobilis*.

Hercules's allheal. See *Panax*.

HERCULES BOVI. Gold and mercury dissolved in a distillation of copperas, nitre, and sea-salt. Violently cathartic.

HEREDITARY DISEASE. (From *heres*, an heir.) A disease which is continued from parents to their children.

HERMAPHRODITE. (From *Ἑρμης*, Mercury, and *Ἀφροδίτη*, Venus, i. e. partaking of both sexes.) The true hermaphrodite of the ancients was, the man with male organs of generation, and the female stature of body, that is, narrow chest and large pelvis; or the woman with female organs of generation, and the male stature of body, that is, broad chest and narrow pelvis. The term is now, however, used to express any *lusus nature*, wherein the

parts of generation appear to be a mixture of both sexes.

ΗΕΡΜΕΤΙΣ. (From *Ερμης*, Mercury.) In the language of the ancient chymists, Hermes was the father of chymistry, and the Hermetic seal was the closing the end of a glass vessel while in a state of fusion, according to the usage of chymists.

Hermodactyl. See *Hermodactylus*.

HERMODACTYLUS. (*Ερμωδακτύλος*. Etymologists have always derived this word from *Ερμης*, Mercury, and *δακτύλος*, a finger. It is probably named from *Hermus*, a river in Asia, upon whose banks it grows, and *δακτύλος*, a date, which it is like.) The root of a species of colchicum, not yet ascertained, but supposed to be the *Colchicum illyricum* of Linnæus, of the shape of a heart, flattened on one side, with a furrow on the other, of a white colour, compact and solid, yet easy to cut or powder. This root, which has a viscous, sweetish, farinaceous taste, and no remarkable smell, is imported from Turkey. Its use is totally laid aside in the practice of the present day. Formerly, the roots were esteemed as cathartics, which power is wanting in those that reach this country.

HERNIA. (From *ἕρως*, a branch; from its protruding forward.) A rupture. Surgeons understand, by the term *hernia*, a tumour formed by the protusion of some of the viscera of the abdomen out of that cavity into a kind of sac, composed of the portion of peritoneum, which is pushed before them. However, there are certainly some cases which will not be comprehended in this definition; either because the parts are not protruded at all, or have no hernial sac, as the readers will learn in the course of this article.

The places in which these swellings most frequently make their appearance, are the groin, the navel, the labia pudendi, and the upper and fore-part of the thigh; they do also occur at every point of the anterior part of the abdomen; and there are several less common instances, in which hernial tumours present themselves at the foramen ovale, in the perineum, in the vagina, at the ischiatic notch, &c.

The parts which, by being thrust forth from the cavity, in which they ought naturally to remain, mostly produce herniæ, are either a portion of the omentum, or a part of the intestinal, canal, or both together. But the stomach, the liver, the spleen, uterus, ovaries, bladder, &c. have been known to form the contents of some hernial tumours.

From these two circumstances of situation and contents, are derived all the different appellations by which herniæ are distinguished. If a portion of intestine only forms the contents of the tumour, it is called *enterocele*; if a piece of omentum

only, *epiplocele*; and if both intestine and omentum contribute mutually to the formation of a tumour, it is called *entero-epiplocele*. When the contents of a hernia are protruded at the abdominal ring, but only pass as low as the groin, or labium pudendi, the case receives the name of *bubonocoele*, or *inguinal hernia*; when the parts descend into the scrotum, it is called an *oscheocoele*, or *scrotal hernia*. The *crural* and *femoral hernia*, is the name given to that which takes place below Poupart's ligament. When the bowels protrude at the navel, the case is named an *exomphalos*, or *umbilical hernia*; and *ventral* is the epithet given to the swelling, when it occurs at any other promiscuous part of the front of the abdomen. The *congenital rupture*, is a very particular case, in which the protruded viscera are not covered with a common hernial sac of peritoneum, but are lodged in the cavity of the tunica vaginalis, in contact with the testicle; and, as must be obvious, it is not named, like herniæ in general, from its situation, or contents, but from the circumstances of its existing from the time of birth.

When the hernial contents lie quietly in the sac, and admit of being readily put back into the abdomen, it is termed a *reducible hernia*; and when they suffer no constriction, yet cannot be put back, owing to adhesions, or their large size in relation to the aperture, through which they have to pass, the hernia is termed *irreducible*. An *incarcerated*, or *strangulated hernia*, signifies one which not only cannot be reduced, but suffers constriction: so that, if a piece of intestine be protruded, the pressure to which it is subjected stops the passage of its contents onward towards the anus, makes the bowel inflame, and brings on a train of most alarming and often fatal consequences.

The general symptoms of a hernia, which is reducible and free from strangulation, are—an indolent tumour at some point of the parietes of the abdomen; most frequently descending out of the abdominal ring, or from just below Poupart's ligament, or else out of the navel; but occasionally from various other situations.

The swelling mostly originates suddenly, except in the circumstances above related; and it is subject to a change of size, being smaller when the patient lies down upon his back, and larger when he stands up, or holds his breath. The tumour frequently diminishes when pressed, and grows large again when the pressure is removed. Its size and tension often increase after a meal, or when the patient is flatulent. Patients with hernia, are apt to be troubled with cholic, constipation, and vomiting, in consequence of the unnatural situation of the bowels. Very often, however, the functions of the viscera seem to suffer little or no interruption.

If the case be an *enterocoele*, and the portion of the intestine be small, the tumour is small in proportion; but though small, yet, if the gut be distended with wind, inflamed, or have any degree of stricture made on it, it will be tense, resist the impression of the finger, and give pain upon being handled. On the contrary, if there be no stricture, and the intestine suffers no degree of inflammation, let the prolapsed piece be of what length it may, and the tumour of whatever size, yet the tension will be little, and no pain will attend the handling it; upon the patient's coughing, it will feel as if it was blown into; and, in general, it will be found very easily returnable. A gurgling noise is often made when the bowel is ascending.

If the hernia be an *epitrocele*, or one of the omental kind, the tumour has a more flabby and a more unequal feel; it is in general perfectly indolent, is more compressible, and (if in the scrotum) is more oblong and less round than the swelling occasioned in the same situation by an intestinal hernia; and, if the quantity be large, and the patient an adult, it is, in some measure, distinguishable by its greater weight.

If the case be an *entero-epitrocele*, that is, one consisting of both intestine and omentum, the characteristic marks will be less clear in either of the simple cases; but the disease may easily be distinguished from every other one, by any body in the habit of making the examination.

HERNIA CEREBRI. *Fungus cerebri.* This name is given to a tumour which every now and then rises from the brain, through an ulcerated opening in the dura matter, and protrudes through a perforation in the cranium, made by the previous application of the trephine.

HERNIA CONGENITA. (So called because it is, as it were, born with the person.) This species of hernia consists in the adhesion of a protruded portion of intestine or omentum to the testicle, after its descent into the scrotum. This adhesion takes place while the testicle is yet in the abdomen. Upon its leaving the abdomen, it draws the adhering intestine, or omentum, along with it into the scrotum, where it forms the hernia congenita.

From the term *congenital*, we might suppose that this hernia always existed at the time of birth. The protrusion, however, seldom occurs till after this period, on the operation of the usual exciting causes of hernia in general. The congenital hernia does not usually happen till some months after birth; in some instances not till a late period. Mr. Hey relates a case, in which a hernia congenita was first formed in a young man, aged sixteen, whose right testis had, a little while before the attack of the disease, descended into the scrotum.

It seems probable that, in cases of hernia congenita, which actually take place when the testicle descends into the scrotum before birth, the event may commonly be referred, as observed above, to the testicle having contracted an adhesion to a piece of intestine, or of the omentum, in its passage to the ring. Whrissberg found one testicle, which had not passed the ring, adhering, by means of a few slender filaments, to the omentum, just above this aperture, in an infant that died a few days after birth.

Excepting the impossibility of feeling the testicle in hernia congenita, as we can in most cases of bubonocoele, (which criterion Mr. Samuel Cooper, in his *Surgical Dictionary*, observes Mr. Pott should have mentioned,) the following account is very excellent. "The appearance of a hernia, in very early infancy, will always make it probable that it is of this kind; but in an adult, there is no reason for supposing his rupture to be of this sort, but his having been afflicted with it from his infancy; there is no external mark, or character, whereby it can be certainly distinguished from the one contained in a common hernial sac; neither would it be of any material use in practice, if there was."

HERNIA CRURALIS. Femoral hernia. The parts composing this kind of hernia are always protruded under Poupart's ligament, and the swelling is situated towards the inner part of the bend of the thigh. The rupture descends on the side of the femoral artery and vein, between these vessels and the os pubis. Females are particularly subject to this kind of rupture, in consequence of the great breadth of their pelvis, while in them the inguinal hernia is rare. It has been computed, that nineteen out of twenty married women, afflicted with hernia, have this kind; but that not one out of an hundred unmarried females, or out of the same number of men, have this form of the disease. The situation of the tumour makes it liable to be mistaken for an enlarged inguinal gland; and many fatal events are recorded to have happened from the surgeon's ignorance of the existence of the disease. A gland can only become enlarged by the gradual effects of inflammation; the swelling of a crural hernia comes on in a momentary and sudden manner; and, when strangulated, occasions the train of symptoms described in the account of the inguinal hernia, which symptoms an enlarged gland could never occasion. Such circumstances seem to be sufficiently discriminative; though the feel of the two kinds of swelling is often not in itself enough to make the surgeon decided in his opinion. A femoral hernia may be mistaken for a bubonocoele, when the expanded part of the swelling lies over Poupart's ligament. As the taxis and operation for the first case ought to be done

differently from those for the latter, the error may lead to very bad consequences. The femoral hernia, however, may always be discriminated, by the neck of the tumour having Poupart's ligament above it. In the bubonocoele, the angle of the pubes is behind and below this part of the sac; but in the femoral hernia, it is on the same horizontal level, a little on the inside of it.

Until very lately, the stricture, in cases of femoral hernia, was always supposed to be produced by the lower border of the external oblique muscle, or, as it is termed, Poupart's ligament. A total change of surgical opinion on this subject has, however, latterly taken place, in consequence of the accurate observations first made in 1768, by Gimbernat, surgeon to the king of Spain. In the crural hernia, (says he,) the aperture through which the parts issue is not formed by two bands, (as in the inguinal hernia,) but it is a foramen, almost round, proceeding from the internal margin of the crural arch, (Poupart's ligament,) near its insertion into the branch of the os pubis, between the bone and the iliac vein; so that, in this hernia, the branch of the os pubis is situated more internally than the intestine, and a little behind; the vein externally, and behind; and the internal border of the arch before. Now it is this border which always forms the strangulation.

HERNIA FLATULENTA. A swelling of the side, caused by air that has escaped through the pleura.

HERNIA GUTTURIS. Bronchocele, or tumour of the bronchial glands.

HERNIA HUMORALIS. *Inflammatio testis.* *Orchitis.* Swelled testicle. A very common symptom attending a gonorrhoea, is a swelling of the testicle, which is only sympathetic, and not venereal, because the same symptoms follow every kind of irritation on the urethra, whether produced by strictures, injections, or bougies. Such symptoms are not similar to the actions arising from the application of venereal matter, for suppuration seldom occurs, and, when it does, the matter is not venereal. The swelling and inflammation appear suddenly, and as suddenly disappear, or go from one testicle to the other. The epididymis remains swelled, however, even for a considerable time afterwards.

The first appearance of swelling is generally a soft pulpy fullness of the body of the testicle, which is tender to the touch; this increases to a hard swelling, accompanied with considerable pain. The epididymis, towards the lower end of the testicle, is generally the hardest part. The hardness and swelling, however, often pervade the whole of the epididymis. The spermatic cord, and especially the *vas deferens*, are often thickened, and sore to the touch.

The spermatic veins sometimes become varicose. A pain in the loins, and sense of weakness there, and in the pelvis, are other casual symptoms. Cholicky pains; uneasiness in the stomach and bowels; flatulency; sickness; and even vomiting; are not unfrequent. The whole testicle is swelled, and not merely the epididymis, as has been asserted.

The inflammation of the part most probably arises from its sympathizing with the urethra. The swelling of the testicle coming on, either removes the pain in making water, and suspends the discharge, which do not return till such swelling begins to subside, or else the irritation in the urethra, first ceasing, produces a swelling of the testicle, which continues till the pain and discharge return; thus rendering it doubtful which is the cause and which the effect. Occasionally, however, the discharge has become more violent, though the testicle has swelled; and such swelling has even been known to occur after the discharge has ceased; yet the latter has returned with violence, and remained as long as the hernia humoralis.

Hernia humoralis, with stoppage of the discharge, is apt to be attended with strangury. A very singular thing is, that the inflammation more frequently comes on when the irritation in the urethra is going off, than when at its height.

The enlargements of the testicle, from cancer and scrophula, are generally slow in their progress: that of an hernia humoralis very quick.

HERNIA INCARCERATA. Incarcerated hernia. Strangulated hernia, or a hernia with stricture. The symptoms are a swelling in the groin, or scrotum, resisting the impression of the fingers. If the hernia be of the intestinal kind, it is generally painful to the touch, and the pain is increased by coughing, sneezing, or standing upright. These are the very first symptoms; and, if they are not relieved, are soon followed by others, viz. a sickness at the stomach, a frequent reaching, or inclination to vomit, a stoppage of all discharge per anum, attended with frequent hard pulse, and some degree of fever. These are the first symptoms; and if they are not appeased by the return of the intestine, that is, if the attempts made for this purpose do not succeed, the sickness becomes more troublesome, the vomiting more frequent, the pain more intense, the tension of the belly greater, the fever higher, and a general restlessness comes on, which is very terrible to bear. When this is the state of the patient, no time is to be lost; a very little delay is now of the utmost consequence; and if the one single remedy which the disease is now capable of be not administered immediately, it will generally baffle every other attempt.

This remedy is the operation whereby the parts engaged in the structure may be set free. If this be not now performed, the vomiting is soon exchanged for a convulsive hiccough, and a frequent gulping up of bilious matter; the tension of the belly, the restlessness, and fever, having been considerably increased for a few hours, the patient suddenly becomes perfectly easy, the belly subsides, the pulse, from having been hard, full, and frequent, becomes low, languid, and generally interrupted; and the skin, especially that of the limbs, cold and moist; the eyes have now a languor and a glassiness, a lack lustre, not easy to be described; the tumour of the part disappears, and the skin covering it sometimes changes its natural colour for a livid hue; but whether it keeps or loses its colour, it has an emphysematous feel, a crepitus to the touch, which will easily be conceived by all who have attended to it, but is not so easy to convey an idea of by words. This crepitus is the too sure indicator of gangrenous mischief within. In this state, the gut either goes up spontaneously, or is returned with the smallest degree of pressure; a discharge is made by stool, and the patient is generally much pleased at the ease he finds; but this pleasure is of short duration, for the hiccough and the cold sweats continuing and increasing, with the addition of spasmodic rigors and subsultus tendinum, the tragedy soon finishes.

HERNIA INGUINALIS. *Bubonocoe.* Inguinal hernia. *The hernia inguinalis* is so called, because it appears in both sexes at the groin. It is one of the divisions of hernia, and includes all those hernia in which the parts displaced pass out of the abdomen through the ring, that is, the arch formed by the aponeurosis of the musculus obliquus externus in the groin, for the passage of the spermatic vessels in men, and the round ligament in women. The parts displaced that form the hernia, the part into which they fall, the manner of the hernia being produced, and the time it has continued, occasion great differences in this disorder. There are three different parts that may produce a hernia in the groin, viz. one or more of the intestines, the epiploon, and the bladder. That which is formed by one or more of the intestines was called, by the ancients, *enterocoele*. The intestine which most frequently produces the hernia is the *ilium*: because, being placed in the iliac region, it is nearer the groin than the rest; but notwithstanding the situation of the other intestines, which seems not to allow of their coming near the groin, we often find the jejunum, and frequently also a portion of the colon and cæcum, included in the hernia. It must be remembered, that the mesentery and mesocolon are membranous substances,

capable of extension, which, by little and little, are sometimes so far stretched by the weight of the intestines, as to escape with the ilium, in this species of hernia. The hernia made by the epiploon is called *epiplocele*; as that caused by the epiploon and one of the intestines together is called *entero-epiplocele*. The hernia of the bladder is called *cystocoele*. Hernia of the bladder is uncommon, and has seldom been known to happen but in conjunction with some of the viscera. When the parts, having passed through the abdominal rings, descend no lower than the groin, it is called an incomplete hernia; when they fall into the scrotum in men, or into the *labia pudendi* in women, it is then termed complete.

The marks of discrimination between some other diseases and inguinal hernia are these:—

The disorders in which a mistake may possibly be made are, the *circocoele*, *bubo*, *hydrocele*, and *hernia humoralis*, or inflamed testicle.

For an account of the manner of distinguishing *circocoele* from a *bubonocoele*, see *Circocoele*.

The circumscribed incompressible hardness, the situation of the tumour, and its being free from all connection with the spermatic process, will sufficiently point out its being *bubo*, at least while it is in a recent state; and when it is in any degree suppurated, he must have a very small share of the *tactus eruditus*, who cannot feel the difference between matter, and either a piece of intestine or omentum.

The perfect equality of the whole tumour, the freedom and smallness of the spermatic process above it, the power of feeling the spermatic vessels, and the *vas deferens* in that process; its being void of pain upon being handled, the fluctuation of the water, the gradual formation of the swelling, its having begun below and proceeded upwards, its not being affected by any posture or action of the patient, nor increased by his coughing or sneezing, together with the absolute impossibility of feeling the testicle at the bottom of the scrotum, will always, to an intelligent person, prove the disease to be *hydrocele*.

Mr. Pott, however, allows that there are some exceptions, in which the testicle cannot be felt at the bottom of the scrotum, in cases of hernia. In recent *bubonocoeles*, while the hernial sac is thin, has not been long or very much distended, and the scrotum still preserves a regularity of figure, the testicle may almost always be easily felt at the inferior and posterior part of the tumour. But in old ruptures, which have been long down, in which the quantity of contents is large, the sac considerably thickened, and the scrotum of an irregular

figure, the testicle frequently cannot be felt; neither is it in general easily felt in the congenital hernia, for obvious reasons.

In the *hernia humoralis*, the pain in the testicle, its enlargement, the hardened state of the epididymis, and the exemption of the spermatic cord from all unnatural fullness, are such marks as cannot easily be mistaken; not to mention the generally preceding gonorrhœa. But if any doubt still remains of the true nature of the disease, the progress of it from above downwards, its different state and size in different postures, particularly lying and standing, together with its descent and ascent, will, if duly attended to, put it out of all doubt that the tumour is a true hernia.

When an inguinal hernia does not descend through the abdominal ring, but only into the canal for the spermatic chord, it is covered by the aponeurosis of the external oblique muscle, and the swelling is small and undefined.

Now and then, the testicle does not descend into the scrotum till a late period. The first appearance of this body at the ring, in order to get into its natural situation, might be mistaken for that of a hernia, were the surgeon not to pay attention to the absence of the testicle from the scrotum, and the peculiar sensation occasioned by pressing the swelling.

HERNIA ISCHIATICA. A rupture at the ischiatic notch. This is very rare. A case, however, which was strangulated, and undiscovered till after death, is related in Mr A. Cooper's second part of his work on hernia. The disease happened in a young man aged 27. On opening the abdomen, the ilium was found to have descended on the right side of the rectum into the pelvis; and a fold of it was protruded into a small sac, which passed out of the pelvis at the ischiatic notch. The intestine was adherent to the sac at two points: the strangulated part, and about three inches on each side, were very black. The intestines towards the stomach were very much distended with air, and here and there had a livid spot on them. A dark spot was even found on the stomach itself, just above the pylorus. The colon was exceedingly contracted, as far as its sigmoid flexure. A small orifice was found in the side of the pelvis, in front of, but a little above, the sciatic nerve, and on the fore part of the pyriformis muscle. The sac lay under the glutæus maximus muscle, and its orifice was before the internal iliac artery, below the obturator artery, but above the vein.

HERNIA INTESTINALIS. See *Hernia scrotalis*.

HERNIA LACHRYMALIS. When the tears pass through the puncta lachrymalia,

but stagnate in the sacculus lachrymalis, the tumour is styled *hernia lachrymalis*, with little propriety or precision. It is with equal impropriety called, by Anel, a *dropsey of the lachrymal sac*.

If the inner angle of the eye is pressed, and an aqueous humour flows out, the disease is the *fistula lachrymalis*.

HERNIA MESENTERICA. Mesenteric hernia. If one of the layers of the mesentery be torn by a blow, while the other remains in its natural state, the intestines may insinuate themselves into the aperture, and form a kind of hernia. The same consequences may result from a natural deficiency in one of these layers. Mr. A. Cooper relates a case, in which all the small intestines, except the duodenum, were thus circumstanced. The symptoms during life were unknown.

HERNIA MESOCOLICA. Mesocolic hernia. So named by Mr. A. Cooper, when the bowels glide between the layers and the mesocolon. Every surgeon should be aware that the intestines may be strangulated from the following causes: 1. Apertures in the omentum, mesentery, or mesocolon, through which the intestine protrudes. 2. Adhesions, leaving an aperture, in which a piece of intestine becomes confined. 3. Membranous bands at the mouths of hernial sacs, which becoming elongated by the frequent protrusion and return of the viscera, surround the intestine, so as to strangle them within the abdomen when returned from the sac.

HERNIA OMENTALIS. *Epiplocele*. A rupture of the omentum; or a protrusion of the omentum through apertures in the integuments of the belly. Sometimes, according to Mr. Sharpe, so large a quantity of the omentum hath fallen into the scrotum, that its weight, drawing the stomach and bowels downwards, hath excited vomiting, inflammation, and symptoms similar to those of the bubonocœle.

HERNIA PERINEALIS. Perineal hernia. In men, the parts protrude between the bladder and rectum; in women, between the rectum and vagina. The hernia does not project so as to form an external tumour; and, in men, its existence can only be distinguished by examining in the rectum. In women, it may be detected both from this part and the vagina.

HERNIA PHRENICA. Phrenic hernia. The abdominal viscera are occasionally protruded through the diaphragm, either through some of the natural apertures in this muscle, or deficiencies, or wounds and lacerations in it. The second kind of case is the most frequent. Morgagni furnishes an instance of the first. Two cases related by Dr. Macauley, and two others published by Mr. A. Cooper, are instances of the second sort. And another case has

been lately recorded by the latter gentleman, affording an example of the third kind. Hildanus, Paré, Petit, Schenck, &c. also mention cases of phrenic hernia.

HERNIA PUDENDALIS. Pudendal hernial. This is the name assigned, by Mr. A. Cooper, to that which descends between the vagina and ramus ischii, and forms an oblong tumour in the labium, traceable, within the pelvis, as far as the os uteri. Mr. C. thinks this case has some times been mistaken for a hernia of the foramen ovale.

HERNIA SCROTALIS. *Hernia oschealis.* *Hernia enteroschocele, oschiocele.* Paracelsus calls it *crepatura*. When the omentum, the intestine, or both, descend into the scrotum, it has these appellations; when the omentum only, it is called *epiploschiocele*. It is styled a perfect rupture, in contradistinction to a bubonocoele, which is the same disorder; but the descent is not so great. The hernia scrotalis is distinguished into the true and false; in the former, the omentum or intestine, or both, fall into the scrotum; in the latter, an inflammation, or a fluid, causes a tumour in this part, as in hernia humoralis, or hydrocele. Sometimes sebaceous matter is collected in the scrotum; and this hernia is called *steatocele*.

HERNIA THYROIDALIS. *Hernia ommis ovalis.* Thyroideal hernia. In the anterior and upper part of the obturator ligament there is an opening, through which the obturator artery, vein, and nerve proceed, and through which occasionally a piece of omentum or intestine is protruded, covered with a part of the peritoneum, which constitutes the hernial sac.

HERNIA UMBILICALIS. *Epiplocephalos.* *Omphalocele.* *Exomphalos.* *Omphalos.* and, when owing to flatulency, *Pneumatomphalos.* The exomphalos, or umbilical rupture, is so called from its situation, and has (like other hernia) for its general contents, a portion of intestine, or omentum, or both. In old umbilical ruptures, the quantity of omentum is sometimes very great. Mr. Ranby says, that he found two eels and a half of intestine in one of these, with about a third part of the stomach, all adhering together.

Mr. Gay and Mr. Nourse, found the liver in the sac of an umbilical hernia; and Bochnius says that he did also.

But whatever are the contents, they are originally contained in the sac formed by the protrusion of the peritoneum.

In recent and small ruptures, this sac is very visible; but in old and large ones, it is broken through at the knot of the navel, by the pressure and weight of the contents, and is not always to be distinguished: which is the reason why it has by some been

doubted whether this kind of rupture has a hernial sac or not.

Infants are very subject to this disease, in a small degree from the separation of the funiculus; but in general they either get rid of it as they gather strength, or are easily cured by wearing a proper bandage. It is of still more consequence to get this disorder cured in females, even than in males; that its return, when they are become adult and pregnant, may be prevented as much as possible; for at this time it often happens, from the too great distention of the belly, or from unguarded motion, when the parts are upon the stretch.

Dr. Hamilton has met with about two cases annually, for the space of seventeen years, of umbilical hernia, which strictly deserve the name of *congenital* umbilical hernia. The funis ends in a sort of bag, containing some of the viscera, which pass out of the abdomen through an aperture in the situation of the navel. The swelling is not covered with skin, so that the contents of the hernia can be seen through the then distended covering of the cord. The disease is owing to a preternatural deficiency in the abdominal muscles, and the hope of cure must be regulated by the size of the malformation and quantity of viscera protruded.

HERNIA UTERI. *Hysterocele.* Instances have occurred of the uterus being thrust through the rings of the muscles; but this is scarcely to be discovered, unless in a pregnant state, when the strugglings of a child would discover the nature of the disease. In that state, however, it could scarcely ever occur. It is the *cerexis* of Hippocrates.

HERNIA VAGINALIS. *Elytrocele.* Vaginal hernia. A tumour occurs within the os externum of the vagina. It is elastic, but not painful. When compressed, it readily recedes, but is reproduced by coughing, or even without this, when the pressure is removed. The inconveniences produced are an inability to undergo much exercise, or exertion; for every effort of this sort brings on a sense of bearing down. The vaginal hernia protrudes in the space left between the uterus and rectum. This space is bounded below by the peritoneum, which membrane is forced downwards, towards the perinaeum; but being unable to protrude further in that direction, is pushed towards the back part of the vagina. These cases probably are always intestinal. Some herniae protrude at the anterior part of the vagina.

HERNIA VARICOSA. See *Circocoele*.

HERNIA VENTOSA. See *Pneumatocoele*.

HERNIA VENTRALIS. *Hypogastrocele.* The ventral hernia may appear at almost any point of the anterior part of the belly, but is most frequently found between

the recti muscles. The portion of intestine, &c. &c. is always contained in a sac made by the protrusion of the peritonæum. Mr. A. Cooper imputes its causes to the dilatation of the natural foramina, for the transmission of vessels, to congenital deficiencies, lacerations, and wounds of the abdominal muscles, or their tendons. In small ventral herniæ, a second fascia is found beneath the superficial one; but in large ones the latter is the only one covering the sac.

HERNIA VESICALIS. *Hernia cystica.* *Cystocele.* The urinary bladder is liable to thrust forth from its proper situation, either through the opening in the oblique muscle, like the inguinal hernia, or under Poupart's ligament, in the same manner as the femoral.

This is not a very frequent species of hernia, but does happen, and has as plain and determined a character as any other.

HERNIARIA. (From *hernia*, a rupture; so called from its supposed efficacy in curing ruptures.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*.

2. Rupture-wort. This plant, though formerly esteemed as efficacious in the cure of herniæ, appears to be destitute not only of such virtues, but of any other. It is the *Herniaria glabra* of Linnæus; has no smell, nor taste.

HERNIARIA GLABRA. The systematic name of the rupture-wort. See *Herniaria*.

HERNIOTOMY. (*Herniotomia*; from *hernia*, and *tomæ*, to cut.) The operation to remove the strangulated part in cases of incarcerated herniæ.

HERPES. (From *ἔρπειν*, to creep; because it creeps and spreads about the skin.) Tetter. A genus of disease in the class *localæ*, and order *dialysis*, of Cullen, distinguished by an assemblage of numerous little creeping ulcers, in clusters, itching very much, and difficult to heal, but terminating in furfuraceous scales.

Mr. Bell, in his treatise on ulcers, arranges the herpes amongst the cutaneous ulcers, and says, that all the varieties of importance may be comprehended in the four following species: 1. *Herpes forficatus*, or what may be termed the *dry tetter*, is the most simple of all the species: it appears indiscriminately in different parts of the body; but most commonly on the face, neck, arms, and wrists, in pretty broad spots and small pimples; these are generally very itchy, though not otherwise troublesome; and, after continuing a certain time, they at last fall off in the form of a white powder, similar to fine bran, leaving the skin below perfectly sound; and again returning in the form of a red efflorescence, they fall off, and are renewed as before.

2. *Herpes pustulosus.* This species ap-

pears in the form of pustules, which originally are separate and distinct, but which afterwards run together in clusters. At first, they seem to contain nothing but a thin watery serum, which afterwards turns yellow, and, exuding over the whole surface of the part affected, it at last dries into a thick crust, or scab; when this falls off, the skin below frequently appears entire, with only a slight degree of redness on its surface; but on some occasions, when the matter has probably been more acrid, upon the scab falling off, the skin is found slightly excoriated. Eruptions of this kind appear most frequently on the face, behind the ears, and on other parts of the head; and they occur most commonly in children.

3. *Herpes miliaris.* The miliary tetter. This breaks out indiscriminately over the whole body: but more frequently about the loins, breast, perinæum, scrotum, and inguina, than in other parts. It generally appears in clusters, though sometimes in distinct rings, or circles, of very minute pimples, which, from their resemblance to the millet seed, has given rise to the denomination of the species. The pimples are at first, though small, perfectly separate, and contain nothing but a clear lymph, which, in the course of this disease, is excreted upon the surface, and there forms into small distinct scales; these, at last, fall off, and leave a considerable degree of inflammation below, that still continues to exude fresh matter, which likewise forms into cakes, and so falls off as before. The itching, in this species of complaint, is always very troublesome; and the matter discharged from the pimples is so tough and viscid, that every thing applied to the part adheres, so as to occasion much trouble and uneasiness on its being removed.

4. *Herpes exedens*, the eating and corroding tetter; so called from its destroying or corroding the parts which it attacks, appears commonly, at first, in the form of several small painful ulcerations, all collected into larger spots, of different sizes and of various figures, with always more or less of an erysipelatous-like inflammation. These ulcers discharge large quantities of a thin, sharp, serous matter; which sometimes forms into small crusts, that in a short time fall off; but most frequently the discharge is so thin and acrid as to spread along the neighbouring parts, where it soon produces the same kind of sores. Though these ulcers do not, in general, proceed farther than the cutis vera, yet sometimes the discharge is so very penetrating and corrosive as to destroy the skin, cellular substance, and, on some occasions, even the muscles themselves. It is this species that should be termed the *depascent*, or *phagedenic* ulcer, from the great destruction of parts which it frequently occasions. See *Phagedæna*.

HERPES AMBULATIVA. A species of erysipelas which moves from one part to another.

HERPES COLLARIS. Tetters about the neck.

HERPES DEPASCENS. The same as herpes exedens. See *Herpes*.

HERPES ESTHIOMENOS. Herpes destroying the skin by ulceration.

HERPES FACIEI. Red pimples common in the faces of adults.

HERPES FARINOSUS. See *Herpes*.

HERPES FERRUS. Common erysipelas.

HERPES INDICA. A fiery, itchy herpes, peculiar to India.

HERPES MILIARIS. See *Herpes*.

HERPES PENISCELIS. That species of erysipelas known by the name of shingles. See *Erysipelas*.

HERPES PUSTULOSUS. See *Herpes*.

HERPES RAPIENS. Venereal ulceration in the head.

HERPES SERPIGO. A name given to the cutaneous affection popularly called a ring-worm. See *Psoiriasis*. Till the recent observations of Dr. Willan, this disease has not been well discriminated by any author, though it is one with which few practitioners are unacquainted.

HERPES SICCUS. The dry, mealy, tetter round the knees.

HERPES SYPHILITICUS. *Herpes venereus*. An herpetic venereal eruption on the skin.

HERPES ZOSTER. Shingles encircling the body. See *Erysipelas*.

Herpetic eruptions. See *Herpes*.

HERPETON. (From *ερεω*, to creep.) A creeping pustule, or ulcer.

HEXAPHARMACUM. (From *εξ*, six, and *φαρμακον*, a medicine.) Any medicine, in the composition of which are six ingredients.

HIBERNICUS LAPIS. See *Lapis hibernicus*.

HIBISCUS. (From *ιβις*, a stork, who is said to chew it, and inject it as a clyster.) The marsh-mallow.

HIBISCUS ABELMOSCHUS. The systematic name of the plant whose seeds are called musk-seed. See *Abelmoschus*.

HICCUGH. Hiccup. A spasmodic affection of the diaphragm, generally arising from irritation produced by acidity in the stomach, error of diet, &c.

HIDROA. (From *ιδρω*, sweat.) A pustular disease, produced by sweating in hot weather.

HIDROCRISIS. (From *ιδρω*, sweat, and *κρινω*, to judge.) A judgment formed from the sweat of the patient.

HIDRONOSOS. (From *ιδρω*, sweat, and *νοσος*, a disease.) The sweating sickness.

HIDROPHRETUS. (From *ιδρω*, sweat, and *φρεσις*, a fever.) The sweating fever, or sickness. See *Sudor Anglicanus*.

HIDROTICA. (From *ιδρω*, sweat.) Medicines which cause perspiration.

HIDROTOPOIETICA. (From *ιδρω*, sweat, and *ποινω*, to make.)

HIERA PICRA. (From *ιερ*, holy, and *πικρος*, bitter.) Holy bitter. Pulvis aloeticus, formerly *hiera logadii*, made in the form of an electuary with honey. It is now kept in the form of dry powder, prepared by mixing socotorine aloes, one pound, with three ounces of white canella.

HIERABOTANE. (From *ιερ*, holy, and *βόταν*, an herb; so called from its supposed virtues. A species of verberna.

HIERACANTHA. (From *αεαξ*, a hawk, and *κνθος*, a flower; so named because it seizes passengers as a hawk does its prey.) A sort of thistle.

HIERACIUM. (From *αεαξ*, a hawk; so called because hawks feed upon it, or because it was said that hawks applied the juice of it to cleanse their eyes.) The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygonia aequalis*. Hawk-weed.

HIERACIUM PLOCELLA. The systematic name of the auricula muris. See *Pilocella*.

HIERACULUM. The hieracium, or hawk-weed.

HIERANOSOS. (From *ιερ*, holy, and *νοσος*, a disease; so called because it was supposed to be that disorder which our Saviour cured, in those who were said to be possessed of devils.) The epilepsy.

HIERATICUM. (From *ιερ*, holy.) A poultice for the stomach, so named from its supposed divine virtues.

Higmore's antrum. See *Antrum of Higmore*.

HIGUERO. The calabash-tree. Fruit said to be febrifuge.

HIMANTOSIS. (From *ιμας*, a thong of leather.) A relaxation of the uvula, when it hangs down like a thong.

HIMAS. A relaxation of the uvula.

HIN. *Hindisch.* *Hing.* Assafœtida.

HIPPOCASTANUM. (From *ιππος*, a horse, and *κασταν*, a chestnut; so called from its size.) *Castanea equestris*, *pavina*. Common horse-chestnut. *Æsculus hippocastanum*; *foliolis septemnis* of Linnæus. The fruit, when dried and powdered, is recommended as an errhine. The bark is highly esteemed on the Continent as a febrifuge; and is, by some, considered as being superior in quality to the Peruvian bark. The bark intended for medical use is to be taken from those branches which are neither very young nor very old, and to be exhibited under similar forms and doses as directed with respect to the cortex peruvianus. It rarely disagrees with the stomach; but its astringent effects generally require the occasional administration of a laxative.

During the late scarcity of grain, some attempts were made to obtain starch from the horse-chestnut, and not without success.

HIPPOCRATICUS. See *Facies hippocratica*.

HIPPOLAPATHUM. (From *ἵππος*, a horse, and *λαπάθω*, the lapathum; a species of lapathum; so named from its size.) See *Rhubarbarum monachorum*.

HIPPOMACHTHEUM. (From *ἵππος*, a horse, and *μαχθῆν*, fennel; so named from its size.) See *Saxifraga vulgaris*.

HIPPOSELINUM. (From *ἵππος*, a horse, and *σῆλον*, purslane; so named because it resembles a large kind of purslane.) *Smyrniacum. Macerona. Herba Alexandrina. Griseum agriceolinum.* Common Alexanders. This plant, *Smyrniacum olusatrum* of Linnæus, was formerly cultivated in our gardens, for culinary use, but is now superseded by celery. These seeds are bitter and aromatic, and the roots are more powerfully bitter. They stand recommended as resolvents, diuretics, and emmenagogues, though seldom used in medical prescription.

HIPS. The ripe fruit of the dog-rose. They are chiefly used as a sweet-meat, or in a preserved state, or to make up medicines. See *Confectio Rose canina*.

HIPPURIS. (From *ἵππος*, a horse, and *οὐρα*, a tail.) 1. Some herbs are thus named, because they resemble a horse's tail.

2. The name of a genus of plants in the Linnæan system. Class, *Monandria*. Order, *Monogynia*. Marc's tail.

HIPPURIS VULGARIS. The systematic name of the horse's tail. See *Equisetum*.

HIPPUS. (From *ἵππος*, a horse; because those who labour under this affection are continually twinkling and trembling, as is usual with those who ride on horse-back.) A repeated dilatation and alternate contraction of the pupil, arising from spasm, or convulsion of the iris.

HIR. (From *χῆρ*, the hand.) The palm of the hand.

HIRA. (From *hir*, the palm of the hand; because it is usually found empty.) The intestine jejunum.

HIRCUS BRUANTICUS. (*Quasi hirtus*; from his shaggy hair.) The goat which affords the oriental bezoar.

HIRCUS. (From *ἥρως*, a hedge; because it is hedged in by the eye-lash.) The angle of the eye.

HIRUDO. (*Quasi haurudo*; from *hauris*, to draw out; so named from its greediness to suck blood.) The leech.

HIRUNDINARIA. (From *hirundo*, the swallow; so called from the resemblance of its pods to a swallow.) Swallow-wort, or asclepias. See *Nummularia* and *Vincetoxicum*.

HIRUNDO. (*Ab herendo*; from its sticking its nest to the eaves of houses.) The swallow. The cavity in the bend of the arm.

HISPIDULA. (From *hispidus*, rough; so named from the rough, woolly surface of its stalks.) See *Gnaphalium*.

Hog's fennel. See *Peucedanum*.

HOLCIMUS. (From *ὥλμι*, to draw.) It sometimes means a tumour of the liver.

HOLCUS. 1. The Indian millet-seed, which is said to be nutritive.

2. The name of a genus of plants in the Linnæan system. Class, *Polygamia*. Order, *Monocia*.

Holly-knee. See *Ruscus*.

Holly, sea. See *Eryngium*.

HOLMISCUS. (Dira. of *ὄλμος*, a mortar.) A small mortar. Also the cavity of the large teeth, because they pound the food as in a mortar.

HOLHELICTIDES. (From *ὅλος*, whole, and *φακίς*, a pustule.) Little pimples all over the body.

HOLOSTES. See *Holosteus*.

HOLOSTEUS. (From *ὅλος*, whole, and *ὀστρον*, a bone.) *Holotes. Holosteum.* Glue-bone, or osteocolla.

HOLOSTEUM. See *Holosteus*.

HOLOTONICUS. (From *ὅλος*, whole, and *τῆναι*, to stretch.) Applied to diseases accompanied with universal convulsion, or rigour.

Holy thistle. See *Carduus benedictus*.

HOLY WEL WATER. A mineral water, arranged under the class of simple cold waters, remarkable for its purity. It possesses similar virtues to that of Malvern, See *Malvern water*.

HOMA. An anasarcaous swelling.

Homburg's sedative salt. See *Boracio acidi*.

HOMOGENEOUS. (*Homogeneous*; from *ὁμος*, like, and *γενος*, a kind.) Uniform, of a like kind, or species. A term used in contra-distinction to *heterogeneous*, when the parts of the body are of different qualities.

HOMOPLATHEOS. (*ὁμοπλαθεος*; from *ὁμος*, the shoulder, and *πλάς*, the blade.) See *Stipula*.

HONEY. *Mel.* A substance collected by bees from the nectary of flowers, resembling sugar in its elementary properties. It has a white or yellowish colour, a soft and grained consistence, and a saccharine and aromatic smell. Honey is an excellent food, and a softening and slightly aperient remedy: mixed with vinegar, it forms *oxymel*, and is exhibited in various forms, in medicine and pharmacy. It is particularly recommended to the asthmatic, and those subject to gravel complaints, from its detergent nature. Founded upon the popular opinion of honey, as a pectoral remedy, Dr. Hill's balsam of honey, a quack medicine, was once in demand; but this, besides honey, contained balsam of Tolu, or gum Benjamin, in solution.

Honey-suckle. See *Lonicera periclymenum*.

Hooping-cough. See *Pertussis*.

HOPLOCRISMA. (From *ὅπλον*, a weapon, and *χρῆμα*, a slave.) A slave, which was ridiculously said to cure wounds by

consent; that is, by anointing the instrument with which the wound was made.

Hops. See *Lupulus*.

HOR RUS The young sprouts of the hop plant are so called; plucked when only a foot above the ground, and boiled, they are eaten with butter as a delicacy, and are very wholesome.

HORDEOLUM. (Diminutive of *hordeum*, barley.) A little tumour on the eye-lids, resembling a barley corn. A stye. Scarpa remarks, the stye is strictly only a little boil, which projects from the edge of the eye-lids, mostly near the great angle of the eye. This little tumour, like the furunculus, is of a dark red colour, much inflamed, and a great deal more painful than might be expected, considering its small size. The latter circumstance is partly owing to the vehemence of the inflammation producing the stye, and partly to the exquisite sensibility and tension of the skin which covers the edge of the eye-lids. On this account, the hordeolum very often excites fever and restlessness, in delicate irritable constitutions; it suppurates slowly and imperfectly; and, when suppurated, has no tendency to burst.

The stye, like other furunculous inflammations, forms an exception to the general rule, that the best mode in which inflammatory swellings can end is, resolution; for, whenever a furunculous inflammation extends so deeply as to destroy any of the cellular substance, the little tumour can never be resolved, or only imperfectly so. This event, indeed, would rather be hurtful, since there would still remain behind a greater or smaller portion of dead cellular membrane; which, sooner or later, might bring on a renewal of the stye, in the same place as before, or else become converted into a hard indolent body, detourning the edge of the eye-lid.

HORDEUM. (*Ab horrore ariste*; from the unpleasantness of its beard to the touch.) 1. The name of a genus of plants in the Linnæan system. Class, *Triandria*. Order, *Digynia*. Barley.

2. The pharmacopœial name of the common barley. The seed called barley is obtained from several species of *hordeum*, but principally from the *vulgare*, or common or Scotch barley, and *distichon*, or *hordeum gallicum vel mundatum*, or French barley, of Linnæus. It is extremely nutritious and mucilaginous, and in common use as a drink, when boiled, in all inflammatory diseases and affections of the chest, especially when there is cough or irritation about the fauces. A decoction of barley, with gum, is considered a useful diluent and demulcent in dysuria and strangury: the gum mixing with the urine, sheaths the urinary canal from the acrimony of the urine. Amongst the ancients, decoctions of barley, *Kasha*, were

the principal medicine, as well as aliment, in acute diseases. Barley is freed from its shells in mills, and in this state called Scotch and French barley. In Holland, they rub barley into small round grains, somewhat like pearls, which is therefore called *pearl barley*, or *hordeum perlatum*. See *Prisan*.

HORDEUM CAUSTICUM. See *Cevadilla*.

HORDEUM DISTICHON. This plant affords the barley in common use. See *Hordeum*.

HORDEUM PERLATUM. See *Hordeum*.

HORDEUM VULGARE. The systematic name of one of the plants which afford the barley. See *Hordeum*.

Horchound. See *Manubium*.

HORMINUM. (From *agere*, to incite; named from its supposed qualities of provoking to venery. Garden clary. The *Salvia sclarea* of Linnæus.

Horn, hart. See *Cornu*.

HORRIPILATIO. A sense of creeping in different parts of the body. A symptom of the approach of fever.

Horse-chestnut. See *Hippocastanum*.

Horse-radish. See *Raphanus rusticianus*.

Horse-tail. See *Equisetum*.

HORTUS. (From *orior*, to rise, as being the place where vegetables grow up.) 1. A garden.

2. The genitals or womb of a woman, which is the repository of the human semen.

Hound's-tongue. See *Cynoglossum*.

House-leek. See *Sedum majus*.

Human body. See *Man*.

HUMECTANTIA. (From *humecto*, to make moist.) Medicines which soften and make moist the solids of the body.

HUMERAL ARTERY. *Arteria humeralis*. Brachial artery. The axillary artery, having passed the tendon of the great pectoral muscle, changes its name to the brachial or humeral artery, which name it retains in its course down the arm to the bend, where it divides into the radial and ulnar artery. In this course it gives off several muscular branches, three of which only deserve attention: 1. *The arteria profunda superior*, which goes round the back of the arm to the exterior muscle, and is often named the upper muscular artery. 2. Another, like it, called *arteria profunda inferior*, or the lower muscular artery. 3. *Ramus anastomaticus major*, which anastomoses round the elbow with the branches of the ulnar artery.

HUMERALIS MUSCULUS. See *Deltoides*.

HUMERALIS NERVUS. The cervical nerve.

HUMERI OS. (*Humerus*; from *aqueo*, the shoulder.) *Os humeri*. *Os brachii*. A long cylindrical bone, situated between the scapula and fore-arm. Its upper

extremity is formed somewhat laterally and internally, into a large, round, and smooth head, which is admitted into the glenoid cavity of the scapula. Around the basis of this head is observed a circular fossa, deepest anteriorly and externally, which forms what is called the neck of the bone, and from the edge of which arises the capsular ligament, which is farther strengthened by a strong membranous expansion, extending to the upper edge of the glenoid cavity, and to the coracoid process of the scapula; and likewise by the tendinous expansions of the muscles, inserted into the head of the humerus. This capsular ligament is sometimes torn in luxation, and becomes an obstacle to the easy reduction of the bone. The articulating surface of the head is covered by a cartilage, which is thick in its middle part, and thin towards its edges; by which means it is more convex in the recent subject than in the skeleton. This upper extremity, besides the round smooth head, affords two other smaller protuberances. One of these, which is the largest of the two, is of an irregular oblong shape, and is placed at the back of the head of the bone, from which it is separated by a kind of groove, that makes a part of the neck. This tuberosity is divided, at its upper part, into three surfaces; the first of these, which is the smallest and uppermost, serves for the insertion of the supraspinatus muscle; the second, or middlemost, for the insertion of the infraspinatus; and the third, which is the lowest and hindmost, for the insertion of the teres minor. The other smaller tuberosity is situated anteriorly, between the larger one and the head of the humerus, and serves for the insertion of the subscapularis muscle. Between these two tuberosities there is a deep groove, for lodging the tendinous head of the biceps brachii; the capsular ligament of the joint affording here a prolongation, thinner than the capsule itself, which covers and accompanies this muscle to its fleshy portion, where it gradually disappears in the adjacent cellular membrane. Immediately below its neck, the os humeri begins to assume a cylindrical shape, so that here the body of the bone may be said to commence. At its upper part is observed a continuation of the groove for the biceps, which extends downwards, about the fourth part of the length of the bone, in an oblique direction. The edges of this groove are continuations of the greater and lesser tuberosities, and serve for the attachment of the pectoralis, latissimus dorsi, and teres major muscles. The groove itself is lined with a glistening substance like cartilage, but which seems to be nothing more than the remains of tendinous fibres. A little lower down, towards the external and anterior side of the middle

of the bone, it is seen rising into a rough ridge, for the insertion of the deltoid muscle. On each side of this ridge the bone is smooth and flat, for the lodgment of the brachialis internus muscle; and behind the middle part of the outermost side of the ridge is a channel, for the transmission of vessels into the substance of the bone. A little lower down, and near the inner side of the ridge, there is sometimes seen such another channel, which is intended for the same purpose. The os humeri, at its lower extremity, becomes gradually broader and flatter, so as to have this end nearly of a triangular shape. The bone, thus expanded, affords two surfaces, of which the anterior one is the broadest, and somewhat convex; and the posterior one narrower and smoother. The bone terminates in four large processes, the two outermost of which are called *condyles*, though not designed for the articulation of the bone. These condyles, which are placed at some distance from each other, on each side of the bone, are rough and irregular protuberances, formed for the insertion of muscles and ligaments, and differ from each other in size and shape. The external condyle, when the arm is in the most natural position, is found to be placed somewhat forwarder than the other. The internal condyle is longer, and more protuberant, than the external. From each of these processes, a ridge is continued upwards, at the sides of the bone. In the interval between the two condyles are placed the two articulating processes, contiguous to each other, and covered with cartilage. One of these, which is the smallest, is formed into a small, obtuse, smooth head, on which the radius plays. This little head is placed near the external condyle, as a part of which it has been sometimes described. The other, and larger, process is composed of two lateral protuberances and a middle cavity, all of which are smooth, and covered with cartilage. From the manner in which the ulnar moves upon this process, it has gotten the name of *trochlea*, or pulley. The sides of this pulley are unequal; that which is towards the little head, is the highest of the two; the other, which is contiguous to the external condyle, is more slanting, being situated obliquely from within outwards, so that when the forearm is full extended it does not form a straight line with the os humeri, and, for the same reason, when we bend the elbow, the hand comes not to the shoulder, as it might be expected to do, but to the forepart of the breast. There is a cavity at the root of these processes, on each of the two surfaces of the bone. The cavity on the anterior surface is divided, by a ridge into two, the external of which receives the end of the radius, and the internal one lodges

the coronoid process of the ulnar in the flexions of the fore-arm. The cavity on the posterior surface, at the basis of the pulley, is much larger, and lodges the olecranon when the arm is extended. The internal structure of the os humeri is similar to that of other long-bones. In new-born infants, both the ends of the bone are cartilaginous, and the large head, with the two tubercles above, and condyles, with the two articulating processes below, become epiphyses before they are entirely united to the rest of the bone.

HUMERUS. (From *humer.*) *Adjutorium*. The shoulder, or joint which connects the arm to the body. In Hippocrates it is called brachium.

HUMILIS. (From *humi*, on the ground; so named because it turns the eye downwards, and is expressive of humility.) See *Rectus inferior oculi*.

HUMOR. (Ab. *humor*, from the ground; because moisture springs from the earth.) A general name of any fluid of the body.

HUMOR VITREUS. The vitreous humour of the eye, which takes its name from the resemblance to melted glass, is less dense than the chrystalline, but more than the aqueous humour: it is very considerable in the human eye, and seems to be formed by the small arteries that are distributed in cells of the hyaloid membrane; it is heavier than common water, slightly albuminous and saline.

Humour, Aqueous. See *Aqueous humour of the eye*.

Humour, Vitreous. See *Humor vitreus*.

HUMOURS OF THE EYE. They are three in number: the aqueous humour, chrystalline lens, and vitreous humour. See *Eye*.

HUMULUS. (From *humilis*, the ground; so named, because without factitious support it creeps along the ground.) The name of a genus of plants in the Linnæan system. Class, *Dioecia*. Order, *Pentandria*. The hop.

HUMULUS LUPULUS. The systematic name of the hop-plant. See *Lupulus*.

HURTICLE. The bluebottle, or *cyaneus*, is so called, because it is troublesome to cut down, and sometimes notches the sickle.

HYALOIDES. (*Membrana hyaloides*; from *yalos*, glass, and *idos*, likeness.) *Membrana arachnoidea*. Capsule of the vitreous humour. The transparent membrane enclosing the vitreous humour of the eye.

HYDARTHROS. (From *hyar*, water, and *arthros*, a joint.) *Hydathron*. *Hydathros*. *Spina ventosa* of the Arabian writers, Rhazes and Avicenna. White swelling. The white swelling, in this country, is a peculiarly common and exceedingly terrible disease. The varieties of white swelling are very numerous, and might usefully receive particular appellations. Systematic writers have generally been content with a

distinction into two kinds, viz. *rheumatic* and *scrophulous*. The last species of the disease they also distinguish into such tumours as primarily affect the bones, and then the ligaments and soft parts; and into other cases, in which the ligaments and soft parts become diseased, before there is any morbid affection of the bones.

These divisions, Mr. Samuel Cooper, in his treatise on the diseases of the joints, proves to be not sufficiently comprehensive; and the propriety of using the term *rheumatic* he thinks to be very questionable.

The knee, ankle, wrist, and elbow, are the joints most subject to white swellings. As the name of the disease implies, the skin is not at all altered in colour. In some instances, the swelling yields, in a certain degree, to pressure; but it never pits, and is almost always sufficiently firm to make an uninformed examiner believe, that the bones contribute to the tumour. The pain is sometimes vehement from the very first; in other instances, there is hardly the least pain in the beginning of the disease. In the majority of scrophulous white swellings, let the pain be trivial or violent, it is particularly situated in one part of the joint, viz. either the centre of the articulation, or the head of the tibia. Sometimes the pain continues without interruption; sometimes there are intermissions; and in other instances the pain recurs at regular times, so as to have been called, by some writers, periodical. Almost all authors describe the patient as suffering more uneasiness in the diseased part when he is warm, and particularly when he is in this condition in bed.

At the commencement of the disease, in the majority of instances, the swelling is very inconsiderable, or there is even no visible enlargement whatever. In the little depressions, naturally situated on each side of the patella, a fulness first shews itself, and gradually spreads all over the affected joint.

The patient, unable to bear the weight of his body on the disordered joint, in consequence of the great increase of pain thus created, gets into the habit of only touching the ground with his toes; and the knee, being generally kept a little bent in this manner, soon loses the capacity of becoming extended again. When white swellings have lasted awhile, the knee, is almost always found in a permanent state of flexion. In scrophulous cases of this kind, pain constantly precedes any appearance of swelling; but the interval between the two symptoms differs very much in different subjects.

The morbid joint, in the course of time, acquires a vast magnitude. Still the integuments retain their natural colour, and remain unaffected. The enlargement of the

articulation, however, always seems greater than it really is, in consequence of the emaciation of the limb both above and below the disease.

An appearance of blue distended veins, and a shining smoothness, are the only alterations to be noticed in the skin covering the enlarged joint. The shining smoothness seems attributable to the distension, which obliterates the natural furrows and wrinkles of the cutis. When the joint is thus swollen, the integuments cannot be pinched up into a fold, as they could in the state of health, and even in the beginning of the disease.

As the distemper of the articulation advances, collections of matter form about the part, and at length burst. The ulcerated openings sometimes heal up; but such abscesses are generally followed by other collections, which pursue the same course. In some cases, these abscesses form a few months after the first affection of the joint; on other occasions, several years elapse, and no supuration of this kind makes its appearance.

Such terrible local mischief must necessarily produce constitutional disturbance. The patient's health becomes gradually impaired; he loses both his appetite and natural rest and sleep; his pulse is small and frequent; and obstinate debilitating diarrhoea and profuse nocturnal sweats ensue. Such complaints are sooner or later followed by dissolution, unless the constitution be relieved in time, either by the amendment or removal of the diseased part. In different patients, however, the course of the disease, and its effects upon the system, vary very much in relation to the rapidity with which they occur.

Rheumatic white swellings are very distinct diseases from the *scrophulous distemper* of large joints. In the first, the pain is said never to occur without being distended with swelling. Scrophulous white swellings, on the other hand, are always preceded by a pain, which is particularly confined to one point of the articulation. In rheumatic cases, the pain is more general, and diffused over the whole joint.

With respect to the particular causes of all such white swellings as come within the class of rheumatic ones, little is known. External irritation, either by exposure to damp or cold, or by the application of violence, is often concerned in bringing on the disease; but very frequently no cause of this kind can be assigned for the complaint. As for scrophulous white swellings, there can be no doubt that they are under the influence of a particular kind of constitution, termed a *scrophulous* or *strumous* habit. In this sort of temperament, every cause capable of exciting inflammation, or any morbid and irritable state of a large joint, may bring such disorder as may end

in the severe disease of which we are now speaking.

In a man of a sound constitution, an irritation of the kind alluded to might only induce common healthy inflammation of the affected joint.

In scrophulous habits, it also seems probable that the irritation of a joint is much more easily produced than in the other constitutions; and no one can doubt that, when once excited in scrophulous habits, it is much more dangerous and difficult of removal than in other patients.

HYDATID. (*Hydatid*; from *uag*, water.)

A very singular animal, formed like a bladder, and distended with an aqueous fluid. These animals are sometimes formed in the natural cavities of the body, as the abdomen and ventricles of the brain, but more frequently in the liver, kidney, and lungs, where they produce diseased actions of those viscera. Cullen arranges these affections in the class *locales*, and order *tumores*. If the virus naturæ medicatrices are not sufficient to effect a cure, the patient mostly falls a sacrifice to their ravages. Dr. Baillie gives the following interesting account of the hydatids, as they are sometimes found in the liver:—There is no gland in the human body in which hydatids are so frequently found as the liver, except the kidneys, where they are still more common. Hydatids of the liver are usually found in the cyst, which is frequently of considerable size, and is formed of very firm materials, so as to give to the touch almost the feeling of cartilage. This cyst, when cut into, is obviously laminated, and is much thicker in one liver than another. In some livers it is not thicker than a shilling, and in others it is near a quarter of an inch in thickness. The laminae which compose it are formed of a white matter, and on the inside there is a lining of a pulpy substance, like the coagulable lymph. The cavity of the cyst I have seen, in one instance, subdivided by a partition of this pulpy substance. In a cyst may be found one hydatid, or a greater number of them. They lie loose in the cavity, swimming in a fluid; or some of them are attached to the side of the cyst. They consist of a round bag, which is composed of a white, semi-opaque, pulpy matter, and contain a fluid capable of coagulation. Although the common colour of hydatids be white, yet I have occasionally seen some of a light amber colour. The bag of the hydatid consists of two laminae, and possesses a good deal of contractile power. In one hydatid this coat, or bag, is much thicker and more opaque than in another; and even in the same hydatid, different parts of it will often differ in its thickness. On the inside of an hydatid, smaller ones are sometimes found, which are commonly not larger than the heads of pins, but sometimes they are even

larger in their size than a gooseberry. These are attached to the larger hydatid, either at scattered irregular distances, or so as to form small clusters; and they are also found floating loose in the liquor of the larger hydatids. Hydatids of the liver are often found unconnected with each other; but sometimes they have been said to enclose each other in a series, like pill-boxes. The most common situation of hydatids of the liver is in its substance, and enclosed in a cyst; but they are occasionally attached to the outer surface of the liver, hanging from it, and occupying more or less of the general cavity of the abdomen. The origin and real nature of these hydatids are not fully ascertained; it is extremely probable, however, that they are a sort of imperfect animalcules. There is no doubt at all, that the hydatids in the livers of sheep are animalcules; they have been often seen to move, when taken out of the liver and put into warm water; and they retain this power of motion for a good many hours after a sheep has been killed. The analogy is great between hydatids in the liver of a sheep, and those of the human subject. In both, they are contained in strong cysts; and in both they consist of the same white pulpy matter. There is undoubtedly some difference between them in simplicity of organization: the hydatid in the human liver being a simple uniform bag; and the hydatid in that of a sheep having a neck and mouth appending to the bag. This difference need be no considerable objection to the opinion above stated. Life may be conceived to be attached to the most simple form of organization. In proof of this, hydatids have been found in the brains of sheep, resembling almost exactly those in the human liver, and which have been seen to move, and therefore are certainly known to be animalcules. The hydatids of the human liver, indeed, have not, as far as I know, been found to move, when taken out of the body and put into warm water; were this to have happened, no uncertainty would remain. It is not difficult to see a good reason why there will hardly occur any proper opportunity of making this experiment. Hydatids are not very often found in the liver, because it is not a very frequent disease there; and the body is allowed to remain for so long a time after death before it is examined, that the hydatids must have lost their living principle, even if they were animalcules, however, it is very strong: and it appears even more difficult to account for their production, according to the common theory of generation, than for that of intestinal worms. We do not get rid of the difficulty by asserting, that hydatids in the human liver are not living animals, because in sheep they are certainly such, where the

difficulty of accounting for their production is precisely the same.

HYDRAGOGUES. (*Medicamenta hydragoga*; from *uag*, water, and *ago*, to drive out.) Medicines are so termed, which possess the property of increasing the secretions or excretions of the body, so as to cadse the removal of water from any of its cavities, such as tonics, diuretics, cathartics, &c.

HYDRARGYRI NITRICO-OXYDUM. See *Nitrico-oxydum hydrargyri*.

HYDRARGYRI OXYDUM CINEREUM.

See *Oxydum hydrargyri cinereum*.

HYDRARGYRI OXYDUM RUBRUM.

See *Oxydum hydrargyri rubrum*.

HYDRARGYRI OXYMURIAS. See *Oxymurias hydrargyri*.

HYDRARGYRI SUBMURIAS. See *Submurias hydrargyri*.

HYDRARGYRI SULPHURETUM RUBRUM. See *Sulphuretum hydrargyri rubrum*.

HYDRARGYRUS. *uagargyros*: from *uag*, water, and *argyros*, silver; so named from its having the resemblance to fluid silver.) See *Mercury*.

HYDRARGYRUS ACETATUS. *Mercurius acetatus*. *Pihle Keyseri*. By this preparation of mercury, the celebrated Keyser acquired an immense fortune in curing the venereal disease. It is an acetate of quicksilver, and therefore termed *acetis hydrargyri* in the new chemical nomenclature. The dose is from three to five grains. Notwithstanding the encomium given to it by some, it does not appear to be so efficacious as some other preparations of mercury.

HYDRARGYRUS CUM CRETA. Mercury with chalk, *Mercurius alkalisatus*. "Take of purified mercury, by weight, three ounces; prepared chalk, five ounces." Rub them together, until the metallic globules disappear. This preparation is milder than any other mercurial, and does not so easily act upon the bowels; it is, therefore, used largely by many practitioners, and possesses alterative properties in cutaneous and venereal complaints, in obstructions of the viscera, or of the prostate gland, given in the dose of ℥ss to ʒss, two or three times a day.

HYDRARGYRUS CUM SULPHURE. *Æthiops mineral*. This is a black sulphuret of mercury, and therefore called *sulphuretum hydrargyri nigrum* in the new chemical nomenclature. The mercury and sulphur are triturated together; the blended mass thus obtained consists of sulphur and an imperfect oxyd of mercury. The mercury, by this admixture of the sulphur, is deprived of its salivating power, and may be administered with safety to all ages and constitutions, as an anthelmintic and alterative.

HYDRARGYRUS PRECIPITATUS

ALBUS. White precipitated mercury. *Calx hydrargyri alba.* "Take of oxy muriate of mercury, muriate of ammonia, of each half a pound; solution of subcarbonate of potash, half a pint; distilled water, four pints." First dissolve the muriate of ammonia, then the oxy muriate of mercury, in the distilled water, and add thereto the solution of subcarbonate of potash. Wash the precipitated powder until it becomes tasteless; then dry it. It is only used externally, in the form of ointment, as an application in some cutaneous affections.

HYDRARGYRUS PRECIPITATUS CINEREUS. This preparation, ordered in the Edinburgh pharmacopœia, is an oxyd of mercury, and nearly the same with the *hydrargyri oxydum cinereum* of the London pharmacopœia; it is used as an alterative in cases of pains arising from an admixture of rheumatism with syphilis. It may be substituted for the hydrargyrus-sulphuratus ruber, in fumigating ozena, and venereal ulcerated sore throat, on account of its not yielding any vapour offensive to the patient.

HYDRARGYRUS PURIFICATUS. Purified mercury. *Argentum vivum purificatum.* "Take of mercury, by weight, six pounds; iron filings, a pound." Rub them together, and distil the mercury from an iron retort, by the application of heat to it. Purified quicksilver is sometimes administered in its metallic state, in doses of an ounce or more, in constipation of the bowels.

HYDRARGYRUS PHOSPHORATUS. This remedy has been observed to heal inveterate venereal ulcers in a very short time, nay, in the course of a very few days, particularly those about the pudenda. In venereal inflammations of the eyes, chancres, rheumatisms, and chronic eruptions, it has proved of eminent service. Upon the whole, if used with necessary precaution, and in the hands of a judicious practitioner, it is a medicine mild and gentle in its operation. The cases in which it deserves the preference over other mercurial preparations are these: in an inveterate stage of syphilis, particularly in persons of torpid insensible fibres; in cases of exostosis, as well as obstructions in the lymphatic system; in chronic complaints of the skin.

The following is the formula. \mathfrak{z} Hydrargyri phosphorati, gr. iv. Corticis cinnamomi in pulverem triti, gr. xiv. Sacchari purif. \mathfrak{z} ss. Misce. The whole to be divided into eight equal parts, one of which is to be taken every morning and evening, unless salivation takes place, when it ought to be discontinued. Some patients, however, will bear from one to two grains of the phosphat of quicksilver, without inconvenience.

HYDRARGYRUS VITRIOLATUS.—

Terpethum minerale. Mercurius emeticus flavus. Sulphas hydrargyri. Formerly this medicine was in more general use than in the present day. It is a very powerful and active alterative when given in small doses. Two grains act on the stomach so as to produce violent vomitings. It is recommended as an emetic in cases of amaurosis. In combination with antimony, it acts powerfully on the skin.

HYDRELXUM. (From *uſag*, water, and *elaxo*, oil.) A mixture of oil and water.

HYDRENTROCKLE. (From *uſag*, water, *entron*, an intestine, and *κκλ*, a tumour.) A dropsy of the scrotum, attended with rupture.

HYDROA. (From *uſag*, water.) A watery pustule.

HYDROCARDIA. (From *uſag*, water, and *καρδι*, the heart.) *Hydrocardia. Hydrops pericardii.* Dropsy of the heart. Dropsy of the pericardium. A collection of fluid in the pericardium, which may be either coagulable lymph, serum, or a puriform fluid. It produces symptoms similar to those of hydrothorax, with violent palpitation of the heart, and mostly an intermitting pulse. It is incurable.

HYDROCELE. (From *uſag*, water, and *κκλ*, a tumour.) The term *hydrocele*, if used in a literal sense, means any tumour produced by water; but surgeons have always confined it to those which possess either the membranes of the scrotum, or the coats of the testicle and its vessels. The first of these, viz. that which has its seat in the membranes of the scrotum, *anasarca integumentorum*, is common to the whole bag, and to all the cellular substance which loosely envelopes both the testes. It is, strictly speaking, only a symptom of a disease, in which the whole habit is most frequently more or less concerned, and very seldom affects the part only. The latter, or those which occupy the coats immediately investing the testicle and its vessels, *hydrocele tunice vaginalis*, are absolutely local, very seldom affect the common membrane of the scrotum, generally attack one side only, and are frequently found in persons who are perfectly free from all other complaints.

The *anasarca integumentorum* retains the impression of the finger. The vaginal *hydrocele* is elastic.

The *hydrocele* of the tunica vaginalis testis is a morbid accumulation of the water separated on the internal surface of the tunica vaginalis, to moisten or lubricate the testicle.

From its first appearance, it seldom disappears, or diminishes, but generally continues to increase, sometimes rapidly, at others more slowly. In some, it grows to a painful degree of distention in a few months; in others, it continues many years with little disturbance. As it enlarges, it

becomes more tense, and is sometimes transparent; so that if a candle is held on the opposite side a degree of light is perceived through the whole tumour; but the only certain distinction is the fluctuation, which is not found when the disease is an hernia of the omentum, or intestines, or an inflammatory or a schirrhous tumour of the testicle.

HYDROCELE CYSTATA. Encysted hydrocele of the spermatic cord resembles the common hydrocele; but the tumour does not extend to the testicle, which may be felt below or behind it, while, in the hydrocele of the vaginal coat, when large, the testicle cannot be discovered. In this disease, also, the penis is not buried in the tumour. Sometimes the fluid is contained in two distinct cells; and this is discovered by little contractions in it. It is distinguished from the anasarcaous hydrocele by a sensible fluctuation, and the want of the inelastic pitting; from hernia, by its beginning below, from its not receding in an horizontal position, and not enlarging by coughing and sneezing.

HYDROCELE FUNICULI SPERMATICI, or hydrocele of the spermatic cord. Anasarcaous hydrocele of the spermatic cord sometimes accompanies ascites, and at other times it is found to be confined to the cellular substance, in or about the spermatic cord. The causes of this disease may be, obstructions in the lymphatics leading from the part, in consequence of schirrous affections of the abdominal viscera, or the pressure of a truss applied for the cure of hernia.

When the affection is connected with anasarca in other parts, it is then so evident as to require no particular description. When it is local, it is attended with a colourless tumour in the course of the spermatic cord, soft and inelastic to the touch, and unaccompanied with fluctuation. In an erect position of the body, it is of an oblong figure; but when the body is recumbent, it is flatter, and somewhat round. Generally, it is no longer than the part of the cord which lies in the groin; though sometimes it extends as far as the testicle, and even stretches the scrotum to an uncommon size. By pressure, a great part of the swelling can always be made to recede into the abdomen. It instantly, however, returns to its former situation, on the pressure being withdrawn.

HYDROCELE PERITONÆI. Ascites, or common dropsy of the belly.

HYDROCELE SPINALIS. A watery swelling on the vertebra.

HYDROKLODES ISCHURIA. (From *υδωρ* water, and *κλυστρον*, attended with tumour.) Applied to a suppression of urine, from a rupture of the urethra.

HYDROCEPHALUS. (From *υδωρ*, water, and *κεφαλη*, the head.) *Hydrocephalus*

lum. Dropsy of the brain. Dropsy of the head. A genus of disease arranged by Cullen, in the class *cachexie*, and order *intumescentie*. It is distinguished by authors into external and internal: 1. *Hydrocephalus externus*, is a collection of water between the membranes of the brain. 2. *Hydrocephalus internus*, is when a fluid is collected in the ventricles of the brain, producing dilatation of the pupils, apoplexy, &c. See *Apoplexia*. It is sometimes of a chronic nature, when the water has been known to increase to an enormous quantity, effecting a diastasis of the bones of the head, and an absorption of the substance of the brain.

Pain in the head, particularly across the brow, stupor, dilatation of the pupils, nausea, vomiting, preternatural slowness of the pulse, and convulsions, are the pathognomic symptoms of this disease, which have been laid down by the generality of writers.

Hydrocephalus is almost peculiar to children, being rarely known to extend beyond the age of twelve or fourteen; and it seems more frequently to arise in those of a scrophulous and rickety habit than in others. It is an affection which has been observed to pervade families, affecting all or the greater part of the children at a certain period of their life; which seems to shew that, in many cases, it depends more on the general habit than on any local affection, or accidental cause.

The disease has generally been supposed to arise in consequence either of injuries done to the brain itself, by blows, falls, &c. from scirrhous tumours or excrescences within the skull, from original laxity or weakness in the brain, or from general debility and an impoverished state of the blood.

With respect to its proximate cause, very opposite opinions are still entertained by medical writers, which, in conjunction with the equivocal nature of its symptoms, prove a source of considerable embarrassment to the young practitioner.

Dr. Beddoes says, he believes it to belong to inflammations, and that, at an early period, he should be inclined to bleed as largely as in pneumonia.

Dr. Withering observes that, in a great many cases, if not in all, congestion, or slight inflammation, are the precursors to the aqueous accumulation.

Dr. Rush thinks that, instead of its being considered an idiopathic dropsy, it should be considered only as an effect of a primary inflammation, or congestion of blood in the brain. It appears (says he) that the disease, in its first stage, is the effect of causes which produce a less degree of that inflammation which constitutes phrenitis; and that its second stage is a less degree of that effusion which produces

serous apoplexy in adults. The former partakes of the nature of the chronic inflammation of Dr. Cullen, and the asthenic inflammation of Dr. Brown. There are others again, who view the subject in a very different light. Dr. Darwin supposes inactivity, or torpor of the absorbent vessels of the brain, to be the cause of hydrocephalus internus; but he confesses, in another part of his work, that the torpor of the absorbent vessels may often exist as a secondary effect.

Dr. Whytt, who has published an ingenious treatise on the disease, observes, the immediate cause of every kind of dropsy is the same, viz. such a state of the parts as makes the exhalant arteries throw out a greater quantity of fluids than the absorbents can take up. From what he afterwards mentions, he evidently considers this state as consisting in debility.

As many cases are accompanied with an increased or inflammatory action of the vessels of the brain, and others again are observed to prevail along with general anasarca, it seems rational to allow that hydrocephalus is, in some instances, the consequence of congestion, or slight inflammation in the brain; and that, in others, it arises either from general debility or topical laxity. In admitting these as incontrovertible facts, Dr. Thomas is at the same time induced to suppose that the cases of it occurring from mere debility are by no means frequent.

The great analogy subsisting between the symptoms which are characteristic of inflammation, and those which form the first stage of the acute species of hydrocephalus, (for the disease, as already observed, has been divided into the chronic and acute by some writers,) together with the good effects often consequent on blood-letting, and the inflammatory appearance which the blood frequently exhibits, seem to point out strong proof of the disease being, in most instances, an active inflammation, and that it rarely occurs from mere debility, as a primary cause.

The progress of the disorder has, by some, been divided into three stages.

When it is accompanied by an increased or inflammatory action of the brain, as not uncommonly happens, its first stage is marked with many of the symptoms of pyrexia, such as languor, inactivity, loss of appetite, nausea, vomiting, parched tongue, hot, dry skin, flushing of the face, head-ach, throbbing of the temporal arteries, and quickened pulse; which symptoms always suffer an exacerbation in the evening, but towards morning become milder.

When it is unaccompanied by any inflammatory action of the brain, many of these appearances are not to be observed. In these cases it is marked by a dejection

of countenance, loss of appetite, pains over the eyes, soreness of the integuments of the cranium to the touch, propensity to the bed, aversion to being moved, nausea, and costiveness. The disease, at length, makes a remarkable transition, which denotes the commencement of its second stage. The child screams out, without being able to assign any cause; its sleep is much disturbed; there is a considerable dilatation of the pupils of the eyes, without any contraction on their being exposed to light; lethargic torpor, with strabismus, or perhaps double vision, ensues, and the pulse becomes slow and unequal.

In the third stage, the pulse returns again to the febrile state, becoming uncommonly quick and variable; and coma, with convulsions, ensue. When the accumulation of water is very great, and the child young, the sutures recede a considerable way from each other, and the head, towards the end, becomes much enlarged.

When recoveries have actually taken place in hydrocephalus, we ought probably to attribute more to the efforts of nature than to the interference of art. In every instance it is to be regarded as of difficult cure.

An accumulation of water in the ventricles of the brain is one of the most common appearances to be observed on dissection. In different cases, this is accumulated in greater or less quantities. It sometimes amounts only to a few ounces, and occasionally to some pints. When the quantity of water is considerable, the foramen is raised at its anterior extremity, in consequence of its accumulation, and an immediate opening of communication is thereby formed between the lateral ventricles. The water is of a purer colour, and more limpid than what it is found in the dropsy of the thorax, or abdomen. It appears, however, to be generally of the same nature with the water that is accumulated in these cavities. In some instances the water in hydrocephalus contains a very small proportion of coagulable matter, and in others it is entirely free from it.

When the water is accumulated to a very large quantity in the ventricles, the substance of the brain appears to be a sort of pulpy bag, containing a fluid. The skull, upon such occasions, is very much enlarged in size, and altered in its shape; and it appears exceedingly large in proportion to the face. On removing the scalp, the bones are found to be very thin, and there are frequently broad spots of membrane in the bone. These appearances are, however, only to be observed where the disease has been of some years continuance.

In some cases, where the quantity of water collected is not great, the substance

of the brain has appeared to be indurated, and in others softened. At times, the organ has been found gorged with blood; collections also of a viscid tenacious matter have been discovered in cysts, upon its external surface, and tumours have been found attached to its substance.

HYDROCEPHALUS ACUTUS. See *Hydrocephalus*.

HYDROCEPHALUS EXTERNUS. Water between the brain and its membranes.

HYDROCEPHALUS INTERNUS. Water in the ventricles of the brain.

HYDROCOTYLE. (From *ὕδωρ*, water, and *κότυλη*, the cotula.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digymia*.

2. The name, in some pharmacopœias, for the common marsh or water cotula, or penny-wort, which is said to possess acrid qualities.

HYDROCRISTIS. (From *ὕδωρ*, water, and *κρίσις*, a vesicle.) An encysted dropsy.

HYDROGEN. (*Hydrogenium*; from *ὕδωρ*, water, and *γενναίω*, to become, or *γεννάω*, to produce, because with oxygen it produces wat. r.) Base of *inflammable air*.

Hydrogen is a substance not perceptible to our sensations in a separate state; but its existence is not at all the less certain. Though we cannot exhibit it experimentally uncombined, we can pursue it while it passes out of one combination into another; we cannot, indeed, arrest it on its passage, but we never fail to discover it, at least if we use the proper chymical means, when it presents itself to our notice in a new compound.

Hydrogen, as its name expresses, is one of the constituent elements of water. Its existence was unknown till lately. It is plentifully distributed in nature, and acts a very considerable part in the processes of the animal and vegetable economy. It is one of the ingredients in the mixtures of bitumen, of oils, fats, ardent spirit, ether, and, in fact, of all the proximate component parts of all animal and vegetable bodies. It forms a constituent part of all animal and vegetable acids. It is one of the bases of ammonia and of various other compound gases.

It possesses so great an affinity with caloric, that it can only exist in the state of gas; it is consequently impossible to procure it in the concrete or liquid state, independent of combination.

Solid hydrogen, therefore, united to caloric and light, forms **HYDROGEN GAS**.

Properties of Hydrogen Gas.

This gas, which was formerly called inflammable air, was discovered by Mr. Cavendish, in the year 1768, though it had been noticed long before by the ancients. The famous philosophical candle attests the antiquity of this discovery.

Hydrogen gas, like oxygen gas, is a triple compound, consisting of the ponderable base of hydrogen, caloric, and light. It possesses all the mechanical properties of atmospheric air. It is the lightest substance whose weight we are able to estimate: when in its purest state, and free from moisture, it is about thirteen times lighter than atmospheric air. It is not fitted for respiration; animals, when obliged to breathe in it, die almost instantaneously. It is decomposed by living vegetables, and its basis becomes one of the constituents of oil, resin, &c. It is inflammable, and burns rapidly when kindled, in contact with atmospheric air or oxygen gas, by means of the electric spark, or by an inflamed body; and burns, when pure, with a blue lambent flame; but all burning substances are immediately extinguished when immersed in it. It is, therefore, incapable of supporting combustion. It is not injurious to growing vegetables. It is unabsorbable by most substances; but water absorbs about one-thirteenth of its bulk. It is capable of dissolving carbon, sulphur, phosphorus, arsenic, and many other bodies. When its basis combines with that of oxygen gas, water is formed; with nitrogen it forms ammonia. It does not act on earthy substances. It is decomposable by a great variety of bodies.

Method of obtaining Hydrogen Gas.—A ready method of obtaining hydrogen gas consists in subjecting water to the action of a substance which is capable of decomposing this fluid.

1. For this purpose, let sulphuric acid, previously diluted with four or five times its weight of water, be poured on iron filings, or bits of zinc, in a small retort, or gas-bottle, called a pneumatic flask, or proof; as soon as the diluted acid comes in contact with the metal, a violent effervescence takes place, and hydrogen gas escapes without external heat being applied. It may be collected in the usual manner over water, taking care to let a certain portion escape, on account of the atmospheric air contained in the disengaging vessels.

The production of hydrogen gas in the above way is owing to the decomposition of water. The iron, or zinc, when in contact with this fluid, in conjunction with sulphuric acid, has a greater affinity to oxygen than the hydrogen has; the oxygen, therefore, unites to it, and forms an oxid of that metal, which is instantly attacked and dissolved by the acid; the other constituent part of the water, the hydrogen, is set free, which, by uniting with caloric, assumes the form of hydrogen gas. The oxygen is therefore the bond of union between the metal and the acid.

The hissing noise, or effervescence, ob-

servable during the process, is owing to the rapid motion excited in the mixture by means of the great number of air-bubbles quickly disengaged and breaking at the surface of the fluid.

We see also in this case, that *two* substances exert an attraction, and are even capable of decomposing, jointly, a *third*, which neither of them is able to do singly, viz. if we present sulphuric acid alone, or iron or zinc alone, to water, they cannot detach the oxygen from the hydrogen of that fluid; but if both are applied, a decomposition is instantly effected. This experiment, therefore, proves that the agency of chymical affinity between two or more bodies may lie dormant, until it is called into action by the interposition of another body, which frequently exerts no energy upon any of them in a separate state. Instances of this kind were formerly called *predisposing affinities*.

2. Iron, in a red heat, has also the property of decomposing water, by dislodging the oxygen from its combination with hydrogen, in the following manner.

Let a gun-barrel, having its touch-hole screwed up, pass through a furnace, or large crucible perforated for that purpose, taking care to incline the barrel at the narrowest part; adjust to its upper extremity a retort charged with water, and let the other extremity terminate in a tube, introduced under a receiver in the pneumatic trough. When the apparatus is thus disposed, and well luted, bring the gun-barrel to a red heat, and when thoroughly red-hot, make the water in the retort boil; the vapour, when passing through the red-hot tube, will yield hydrogen gas abundantly. In this experiment, the oxygen of the water combines with the iron at a red heat, so as to convert it into an oxyd, and the caloric applied combines with the hydrogen of the water, and forms hydrogen gas. It is, therefore, the result of a double affinity, that of the oxygen of the water with the metal, and that of its hydrogen with caloric.

The more caloric is employed in the experiment of decomposing water by means of iron, &c. the sooner is the water decomposed.

Hydrogen gas is frequently found in great abundance in mines and coal-pits, where it is sometimes generated suddenly, and becomes mixed with the atmospheric air of these subterraneous cavities. If a lighted candle be brought in, this mixture often explodes, and produces the most dreadful effects. It is called, by miners, *fire-damp*. It generally forms a cloud in the upper part of the mine, on account of its levity, but does not mix there with atmospheric air, unless some agitation takes place. The miners frequently set fire to

it with a candle, laying at the same time flat on their faces, to escape the violence of the shock. An easier and more safe method of clearing the mine is, by leading a long tube through the shaft of it, to the ash-pit of a furnace; by this means the gas will be conducted to feed the fire.

Hydrogen gas, in whatever manner produced, *always* originates from water, either in consequence of a preceding decomposition, in which it had been combined, in the state of solid or fixed hydrogen, with one of the substances employed, or from a decomposition of water actually taking place during the experiment.

There are instances recorded of a vapour issuing from the stomach of dead persons, which took fire on the approach of a candle. We even find accounts, in several works, of the combustion of living human beings, which appears to be spontaneous. Dr. Swediaur related some instances of porters at Warsaw, who, having drank abundantly of spirit, fell down in the street, with the smoke issuing out of their mouths; and people came to their assistance, saying they would take fire; to prevent which, they made them drink a great quantity of milk, or used a more singular expedient, by causing them to swallow the urine of the by-standers, immediately on its evacuation.

However difficult it may be to give credit to such narratives, it is equally difficult to reject them entirely, without refusing to admit the numerous testimonies of men, who were, for the most part, worthy of credit. *Citizen Lair* has collected all the circumstances of this nature which he found dispersed in different books, and has rejected those which did not appear to be supported by respectable testimony, to which he has added some others, related by persons still living. These narratives are nine in number; they were communicated to the Philomatic Society, at Paris, and inserted in the bulletin, Thermidor, An. 5, No. 29. The cause of this phenomenon has been attributed to a development of hydrogen gas taking place in the stomachs of these individuals.

Citizen Lair believes that the bodies of these people were not burned perfectly spontaneously, but it appeared to be owing to some very slight external cause, such as the fire of a candle, taper, or pipe.

HYDROGEN GAS, SULPHURATED. Sulphurated hydrogen gas possesses the properties of an acid; for when absorbed by water, its solution reddens vegetable blues; it combines also with alkalis, earths, and with several metallic oxyds. Sulphurated hydrogen gas possesses an extremely offensive odour, resembling that of putrid eggs. It kills animals, and extinguishes burning bodies. When mixed

with oxygen gas; or atmospheric air, it is inflammable. Mingled with nitrous gas, it burns with a yellowish green flame. It is decomposed by ammonia, by oxygenated muriatic acid gas, and by sulphureous acid gas. It has a strong action on the greater number of metallic oxyds. Its specific gravity to common air is as 1106 to 1000. It is composed, according to Thenart, of 70.857 sulphur, and 29.143 hydrogen. It has the property of dissolving a small quantity of phosphorus.

Sulphurated hydrogen gas may be obtained in several ways:—

1. Take dry sulphuret of potash, put it into a tubulated retort, lodged in a sand-bath, or supported over a lamp; direct the neck of the retort under a receiver placed in the pneumatic trough; then pour gradually upon the sulphuret diluted sulphuric or muriatic acid; a violent effervescence will take place, and sulphurated hydrogen gas will be liberated. When no more gas is produced spontaneously, urge the mixture with heat, by degrees, till it boils, and gas will again be liberated abundantly.

The water made use of for receiving it should be heated to about 80 or 90; at this temperature it dissolves little of the gas; whereas, if cold water be made use of, a vast quantity of it is absorbed.

Explanation.—Though sulphur makes no alteration on water, which proves that sulphur has less attraction for oxygen than hydrogen has, yet, if sulphur be united to an alkali, this combination decomposes water whenever it comes in contact with it, though the alkali itself has no attraction either for oxygen or hydrogen.

The formation of this gas explains this truth. On adding the sulphuret of potash to the water, this fluid becomes decomposed, part of the sulphur robs it of its oxygen, and forms with it sulphuric acid; this generated acid unites to part of the alkali, and forms sulphate of potash. The liberated hydrogen dissolves another part of the sulphur, and forms with it sulphurated hydrogen, the basis of this gas, which is retained by the separated portion of the alkali. The sulphuric acid added now extricates it from the alkali, and makes it fly off in the form of gas.

The mixture in the retort therefore contains an alkaline sulphure, properly so called, or the simple combination of sulphur with an alkali; sulphate of potash, and, lastly, sulphurated hydrogen gas, dissolved in the alkaline sulphure, or an hydrogenated sulphure.

Diluted muriatic acid seems best adapted for the production of sulphurated hydrogen gas from alkaline sulphurets. If nitric acid be made use of it, must be much diluted. Sulphuric acid yields little gas, unless assisted by heat. When

the proportion of sulphur in the sulphurets exceeds that of the alkali, the dense sulphuric acid poured upon it emits sulphuric acid gas, much loaded with sulphur. All the rest of the acids may be made use of for decomposing the sulphurets.

2. When iron and sulphur are united together, they afford a large quantity of sulphurated hydrogen gas, on submitting them to the action of heat, in contact with water.

Melt together, in a crucible, equal parts of iron filings and sulphur; the product is a black brittle mass, called sulphuret of iron. Reduce this to powder, and put it, with a little water, into a tubulated retort; add diluted muriatic acid, and apply a gentle heat, till no more gas is disengaged. The philosophy of this experiment is analogous to the former. Part of the oxygen of the water unites to part of the sulphur, and forms sulphuric acid; another part oxydates the iron, which, dissolved by the acid, forms sulphate of iron: the hydrogen of the water unites to another part of the sulphur, and forms sulphurated hydrogen, which becomes gaseous by the addition of caloric.

3. Sulphurated hydrogen gas may also be obtained by heating an alkaline sulphuret, with the addition of water, without the effusion of an acid. In this case, the water is also decomposed; its hydrogen unites with part of the sulphur, and forms sulphurated hydrogen; the oxygen of the water unites with another part of the sulphur, and produces sulphuric acid, which joins to the alkali and forms a sulphate. The sulphurated hydrogen becomes disengaged by heat in the gaseous form.

4. Sulphurated hydrogen gas may be obtained by passing hydrogen gas through sulphur, in a state of fusion.

For this purpose, put sulphur into a gun-barrel, or Wedgwood's tube, and place it across a furnace; fit to the lower extremity a bent glass tube, which goes under a receiver placed in the pneumatic trough, and adapt to the upper extremity a tubulated retort, or other apparatus proper for producing hydrogen gas. The sulphur must then be heated, and, when melted, and the hydrogen gas evolved, must be made to pass over it, which, in this manner, will dissolve part of the sulphur, and become converted into sulphurated hydrogen gas.

5. It may likewise be procured in a direct manner; for that purpose, let a small quantity of sulphur be enclosed in a jar full of hydrogen gas, and melt it by means of a burning-glass. This method does not succeed, except the hydrogen gas be as dry as possible, for its affinity to sulphur is weakened in proportion to its moisture.

HYDROGEN GAS, PHOSPHORATED. Phosphorated hydrogen gas consists of phosphorus dissolved in hydrogen gas.

Properties.—It is the most combustible substance in nature, and it is particularly distinguished from all other gases, by the property of taking fire immediately when brought in contact with atmospheric air. When mixed with oxygen gas, or with oxygenated muriatic acid gas, it burns with great vehemence. When bubbles of it are suffered to pass through water, they explode in succession as they reach the surface of this fluid. It has an insupportable odour, similar to that of putrid fish. It is partly absorbable by distilled water, freed from atmospheric air at low temperatures. Distilled water absorbs about one-fourth, and gives it out again without alteration by heat. Water containing atmospheric air decomposes it, when suffered to stand with it for some time. When exposed to vivid light, it deposits phosphorus in a crystalline form.

Methods of obtaining it.

1. Take a small retort; put into it one part of phosphorus and ten of a concentrated solution of potash, or soda; make the mixture boil, and receive the liberated gas over mercury; or, if it be intended for immediate use, it may be collected over water. In this experiment, a decomposition of the water takes place. Its oxygen unites to part of the phosphorus, and forms phosphoric acid, which joins to the potash, and forms phosphate of potash. The liberated hydrogen dissolves another part of the phosphorus, and becomes converted into phosphorated hydrogen gas.

In thus preparing this gas, the body of the retort should be filled as nearly as possible with the mixture, otherwise the first portion of gas which is produced inflames in the retort; a vacuum is formed, and the water forced up into the retort, and endangers the bursting of it.

2 Phosphorated hydrogen gas is also obtained, if, by a direct exposure to a strong heat, we effect a combination of phosphorus and lime, and then throw this compound into water, a great quantity of phosphorated hydrogen gas will soon be formed, and may be collected in the usual manner, over water or mercury.

The production of phosphorated hydrogen gas in this manner, is analogous to the first, with the only difference that here the decomposition of the water takes place at common temperatures.

3. Phosphorated hydrogen gas may also be obtained, according to Davy, in the following manner:

Let water be decomposed in the usual manner, by means of zinc and sulphuric acid, and add to the mixture a quantity of

phosphorus. The hydrogen evolved will dissolve part of the phosphorus; phosphorated hydrogen gas will be produced, and take fire at the surface of the fluid, so long as the decomposition of the water is made, with considerable rapidity. But the gas produced in this process burns with a more lambent flame than that obtained in the usual manner, probably on account of containing a larger quantity of hydrogen. The experiment is nevertheless brilliant; for the gas is disengaged in small bubbles, which cover the whole surface of the fluid; they disengage themselves rapidly, new ones are produced, and the whole fluid resembles a well of fire.

For the success of this experiment, it is essential that the water, during the action of its decomposition, be considerably heated, which may be effected by a copious addition of sulphuric acid, and that the phosphorus be present in a considerable quantity. Half a part of phosphorus cut into small pieces, one of granulated zinc, three of concentrated sulphuric acid, and five of water, answer this purpose exceedingly well.

Phosphorated hydrogen gas is also produced by nature. The air which burns at the surface of certain springs, and forms what is called *burning-springs*, and the *igneis fatui* (Jack o'lanterns), which glide along burying-grounds, or places where animal matter is putrefying, consists of hydrogen gas, holding phosphorus in solution.

Hydrogen gas, light, carbonated. See *Carbonated hydrogen gas*.

Hydrogen gas, heavy, carbonated. See *Carbonated hydrogen gas*.

HYDROLAPATHUM. (From *hydro*, water, and *lappa*, the dock.) *Herba Britannica.* *Lapathum aquaticum.* The water-dock. *Rumex hydrolapathum*, of Linnaeus:—*floribus hermaphroditis, valvulis integris graniferis, foliis lanceolatis.* The leaves of this plant manifest considerable acidity, and are said to possess a laxative quality. The root is strongly astringent, and has been much employed, both externally and internally, for the cure of some diseases of the skin, as scurvy, lepra, lichen, &c. The root powdered is said to be an excellent dentrifice.

HYDROMEL. (From *hydro*, water, and *melis*, honey.) *Mulnum.* *Aqua mulea.* *Meliceratum.* Braggat. *Hydromel.* Water impregnated with honey. After it is fermented, it is called vinous hydromel, or mead.

HYDROMETRA. (From *hydro*, water, and *metra*, the womb.) *Hydrops uteri.* Dropsy of the womb. A genus of disease in the class *cachexie*, and order *intumescentie*, of Cullen. It produces a swelling of the hypogastric region, slowly and gradually increasing, resembling the figure of the

uterus, yielding to, or fluctuating on, pressure; without ischury or pregnancy. Sauvage enumerates seven species. It must be considered as a very rare disease, and one that can with difficulty be ascertained.

HYDROMPHALUX. (From *υδω*, water, and *ομφαλος*, the navel.) A tumour of the navel containing water.

HYDROXOSIS. (From *υδω*, water, and *κρυσις*, a disease.) The sweating-sickness, called Ephidrosis and Sudor anglicus.

HYDROPEDESIS. (From *υδω*, water, and *παιδα*, to break out.) A breaking out into a violent sweat.

HYDROPHOBIA. (From *υδω*, water, and *φοβια*, to fear,) *Rabies canina*. *Cynanthropia*. *Cynolepsia*. Canine madness. This disease arises in consequence of the bite of a rabid animal, as a dog or cat, and sometimes spontaneously. It is termed hydrophobia, because persons that are thus bitten dread the sight or the falling of water when first seized. Cullen has arranged it under the class *neuroses*, and order *spasmi*, and defines it a loathing and great dread of drinking any liquids, from their creating a painful convulsion of the pharynx, occasioned most commonly by the bite of a mad animal.

There are two species of hydrophobia:

1. *Hydrophobia rabiosa*, when there is a desire of biting.
2. *Hydrophobia simplex*, when there is not a desire of biting.

Dr. James observes, that this peculiar affection properly belongs to the canine genus, viz. dogs, foxes, and wolves; in which animals only it seems to be innate and natural, scarcely ever appearing in any others, except when communicated from these. When a dog is affected with madness, he becomes dull, solitary, and endeavours to hide himself, seldom barking, but making a murmuring noise, and refusing all kinds of meat and drink. He flies at strangers; but, in this stage, he remembers and respects his master; his head and tail hang down; he walks as if over-powered by sleep; and a bite, at this period, though dangerous, is not so apt to bring on the disease in the animal bitten as one afflicted at a later period. The dog at length begins to pant; he breathes quickly and heavily; his tongue hangs out; his mouth is continually open, and discharges a large quantity of froth. Sometimes he walks slowly, as if half asleep, and then runs suddenly, but not always directly, forward. At last he forgets his master; his eyes have a dull, watery, red appearance; he grows thin and weak, often falls down, gets up, and attempts to fly at every thing, becoming very soon quite furious. The animal seldom lives in this latter state longer than thirty hours; and it is said, that his bites, towards the end of his ex-

istence, are the most dangerous. The throat of a person suffering hydrophobia is always much affected; and, it is asserted, the nearer the bite to this part the more perilous.

Hydrophobia may be communicated to the human subject from the bites of cats, cows, and other animals, not of the canine species, to which the affection has been previously communicated. However, it is from the bites of those domestic ones, the dog and cat, that most cases of hydrophobia originate. It does not appear that the bite of a person affected can communicate the disease to another; at least the records of medicine furnish no proof of this circumstance.

In the human species, the general symptoms attendant upon the bite of a mad dog, or other rabid animal, are—the part bitten, at some indefinite period, and occasionally long after the bitten part seems quite well, a slight pain begins to be felt in it, now and then attended with itching, but generally resembling a rheumatic pain. Then come on wandering pains, with an uneasiness and heaviness, disturbed sleep, and frightful dreams, accompanied with great restlessness, sudden startings, and spasms, sighing, anxiety, and a love for solitude. These symptoms continuing to increase daily, pains begin to shoot from the place which was wounded, all along up to the throat, with a straitness and sensation of choaking, and a horror and dread at the sight of water, and other liquids, together with a loss of appetite and tremor. The person is, however, capable of swallowing any solid substance with tolerable ease; but the moment that any thing in a fluid form is brought in contact with his lips, it occasions him to start back with much dread and horror, although he labours perhaps under great thirst at the time.

A vomiting of bilious matter soon comes on, in the course of the disease, and an intense hot fever ensues, attended with continual watching, great thirst, dryness and roughness of the tongue, hoarseness of the voice, and the discharge of a viscid saliva from the mouth, which the patient is constantly spitting out; together with spasms of the genital and urinary organs, in consequence of which the evacuations are forcibly thrown out. His respiration is laborious and uneasy, but his judgment is unaffected, and, as long as he retains the power of speech, his answers are distinct.

In some few instances, a severe delirium arises, and closes the tragic scene; but it more frequently happens, that the pulse becomes tremulous and irregular, that convulsions arise, and that nature, being at length exhausted, sinks under the pressure of misery.

The appearances to be observed, on dis-

section, in hydrophobia, are unusual aridity of the viscera and other parts; marks of inflammation in the fauces, gula, and larynx; inflammatory appearances in the stomach, and an accumulation or effusion of blood in the lungs. Some marks of inflammation are likewise to be observed in the brain, consisting in a serous effusion on its surface, or in a redness of the pia mater; which appearances have also presented themselves in the dog.

In some cases of dissection, not the least morbid appearance has been observed, either in the fauces, diaphragm, stomach, or intestines. The poison has therefore been conceived by some physicians to act upon the nervous system, and to be so wholly confined to it, as to make it a matter of doubt whether the qualities of the blood are altered or not.

HYDROPTHALMIA. (From *ὕδωρ*, water, and *ὀφθαλμος*, the eye. *Hydrophthalmium*. There are two diseases, different in their nature and consequences, thus termed. The one is a mere anasarca or oedematous swelling of the eyelid. The other, the true hydrophthalmia, is a swelling of the bulb of the eye, from too great a collection of the vitreous or aqueous humours.

HYDROPTHALMIUM. (From *ὕδωρ*, water, and *ὀφθαλμος*, the eye.) See *Hydrophthalmia*.

HYDROPTISOCELE. (From *ὕδωρ*, water, *φύση*, flatulence, and *κύμα*, a tumour.) Hernia, combined with hydrocele.

HYDROPIKA. (From *ὕδωρ*, the dropsy.) Medicines which relieve or cure dropsy.

HYDROPIPER. (From *ὕδωρ*, water, and *πικρὸν*, pepper; so called from its biting the tongue like pepper, and being a native of marshy places.) Biting arsmart. Lake-weed. Water-pepper. *Polygonum hydro Piper* of Linnaeus. This plant is very common in our ditches; the leaves have an acrid burning taste, and seem to be nearly of the same nature with those of the arum. They have been recommended as possessing antiseptic, aperient, diuretic virtues, and given in scurvy and cachexies, asthmas, hypochondriacal and nephritic complaints, and wandering gout. The fresh leaves have been applied externally, as a stimulating cataplasm.

HYDROPEUMOSARCA. (From *ὕδωρ*, water, *πνεῦμα*, wind, and *σὰρξ*, flesh.) A tumour of air, water, and solid substances.

HYDROPOIDES. (From *ὕδωρ*, a dropsy, and *ὅμοιος*, likeness.) A term formerly applied to liquid and watery excrements.

HYDROPS. (From *ὕδωρ*, water.) Dropsy. A preternatural collection of serous or watery fluid in the cellular substance, or different cavities of the body. It receives

different appellations, according to the particular situation in which it is lodged.

When it is diffused through the cellular membrane, either generally or partially, it is called *anasarca*. When it is deposited in the cavity of the cranium, it is called *hydrocephalus*; when in the chest, *hydrothorax*, or *hydrops pectoris*. When in the abdomen, *ascites*. In the uterus, *hydrometra*; and within the scrotum, *hydrocele*.

The causes of these diseases are, a family disposition thereto, frequent salivations, excessive and long continued evacuations, a free use of spirituous liquors, (which never fail to destroy the digestive powers,) schirrosities of the liver, spleen, pancreas, mesentery, and other abdominal viscera; preceding diseases, as the jaundice, diarrhoea, dysentery, phthisis, asthma, gout, intermittents of long duration, scarlet fever, and some of the exanthemata; a suppression of accustomed evacuations, the sudden striking in of eruptive humours, ossification of the valves of the heart, polypi in the right ventricle, aneurism in the arteries, tumours making a considerable pressure on the neighbouring parts, permanent obstruction in the lungs, rupture of the thoracic duct, exposure for a length of time to a moist atmosphere, laxity of the exhalents, defect in the absorbents, topical weakness, and general debility.

HYDROPS AD MATULAM. Diabetes.

HYDROPS ARTICULI. A white swelling of a joint is sometimes so called.

HYDROPS CYSTICUS. Any dropsy enclosed in bags or cysts.

HYDROPS GENU. An accumulation of synovia, under the capsular ligament of the knee.

HYDROPS MEDULLÆ SPINALIS. See *Hydro-rachitis* and *Spina bifida*.

HYDROPS OVARII. A dropsy of the ovarium. A species of a-cites.

HYDROPS PECTORIS. See *Hydrothorax*.

HYDROPS PERICARDII. See *Hydrocardia*.

HYDROPS PULMONUM. Water in the cellular interstices of the lungs.

HYDROPS SCROTI. See *Hydrocele*.

HYDROPS UTERI. *Hydrometra*.

HYDROPHRETUS. (From *ὕδωρ*, water, and *φύσος*, fever.) The sweating fever or sickness. See *Sudor Anglicus*.

HYDRO-RACHITIS. (From *ὕδωρ*, water, and *ῥαχίς*, the spine.) A fluctuating tumour, mostly situated on the lumbar vertebrae of new-born children. It is a genus of disease in the class *cachexia*, and order *intumescencia* of Cuilen, and is always incurable. See *Spina bifida*.

HYDROSACCHARUM. (From *ὕδωρ*, water, and *σάκχαρος*, sugar.) A drink made of sugar and water.

HYDROSARCA. (From *ὕδωρ*, water, and

sağ, the flesh.) Water in the cellular membrane. See *Anasarca*.

HYDROSARCOELE. (From *uag*, water, *sağ*, the flesh, and *sarx*, a tumour.) Sarcocoele, with an infusion of water into the cellular membrane.

HYDRASATUM. (From *uag*, water, and *gaw*, a rose.) A drink made of water, honey, and the juice of roses.

HYDRO-ELINUM. (From *uag*, water, and *ewar*, purslane.) A species of purslane, growing in marshy places.

HYDROSULPHURETUM STIBII LUTEUM. See *Sulphur antimonii precipitatum*.

HYDROSULPHURETUM STIBII RUBRUM. *Kermes mineralis*. A sulphuret of antimony formerly in high estimation as an expectorant, sudorific and antispasmodic, in difficult respiration, rheumatism, diseases of the skin and glands.

HYDROTHORAX. (From *uag*, water, and *bagaz*, the chest.) *Hydrops thoracis*. *Hydrops pectoris*. Dropsy of the chest. A genus of disease in the class *cachexia*, and order *intumescens*, of Cullen. Difficulty of breathing, particularly when in a horizontal posture; sudden startings from sleep, with anxiety, and palpitations of the heart; cough, paleness of the visage, anasarca swellings of the lower extremities, thirst, and a scarcity of urine, are the characteristic symptoms of hydrothorax; but the one which is more decisive than all the rest is, a fluctuation of water being perceived in the chest, either by the patient himself or his medical attendant, on certain motions of the body.

The causes which give rise to the disease are pretty much the same with those which are productive of the other species of dropsy. In some cases, it exists without any other kind of dropical affection being present; but it prevails very often as a part of more universal dropsy.

It frequently takes place to a considerable degree before it becomes very perceptible; and its presence is not readily known, the symptoms, like those of hydrocephalus, not being always very distinct. In some instances, the water is collected in both sacs of the pleura; but at other times, it is only in one. Sometimes it is lodged in the pericardium alone; but, for the most part, it only appears there, when, at the same time, a collection is present in one or both cavities of the thorax. Sometimes the water is effused in the cellular texture of the lungs, without any being deposited in the cavity of the thorax. In a few cases, the water that is collected is enveloped in small cysts, of a membranous nature, known by the name of hydatids, which seem to float in the cavity; but more frequently they are connected with, and attached to, particular parts of the internal surface of the pleura.

Hydrothorax often comes on with a sense of uneasiness at the lower end of the sternum, accompanied by a difficulty of breathing, which is much increased by any exertion, and which is always most considerable during night, when the body is in an horizontal posture. Along with these symptoms there is a cough, that is at first dry, but which, after a time, is attended with an expectoration of thin mucus. There is likewise a paleness of the complexion, and an anasarca swelling of the feet and legs, together with a considerable degree of thirst, and a diminished flow of urine. Under these appearances, we have just grounds to suspect that there is a collection of water in the chest; but if the fluctuation can be perceived, there can then remain no doubt as to the reality of its presence.

During the progress of the disease, it is no uncommon thing for the patient to feel a numbness, or degree of palsy, in one or both arms, and to be more than ordinarily sensible to cold. With regard to the pulse, it is usually quick at first, but, towards the end, becomes irregular and intermitting.

Our prognostic in hydrothorax must, in general, be unfavorable, as it has seldom been cured, and, in many cases, will hardly admit even of alleviation, the difficulty of breathing continuing to increase, until the action of the lungs is at last entirely impeded by the quantity of water deposited in the chest. In some cases, the event is suddenly fatal, but in others it is preceded, for a few days previous to death, by a spitting of blood.

Dissections of this disease shew that, in some cases, the water is either collected in one side of the thorax, or that there are hydatids formed in some particular part of it; but they more frequently discover water in both sides of the chest, accompanied by a collection in the cellular texture and principal cavities of the body. The fluid is usually of a yellowish colour; possesses properties similar to serum, and, with respect to its quantity, varies very much, being from a few ounces to several quarts. According to the quantity, so are the lungs compressed by it; and, where it is very considerable, they are usually found much reduced in size. When universal anasarca has preceded the collection in the chest, it is no uncommon occurrence to find some of the abdominal viscera in a sclirous state.

HYGIENE. (From *uaganu*, to be well.) *Hygieia*. *Hygeia*. Modern physicians have applied this term to that division of *therapia* which treats of the diet of the sick and the non-natural.

HYGIESTIS. See *Hygiene*.

HYGMA. (From *uag*, humid.) Liquid plasters.

HYDREPLASTRUM. (From *υδρ*, moist, and *πλαστρον*, a plaster.) A liquid plaster.

HYDROBLEPHARICUS. (From *υδρ*, humid, and *βλεφαρον*, the eye-lid.) Applied to the emunctory ducts in the extreme edge, or inner pari, of the eye-lid.

HYDROCIRCOELE. (From *υδρ*, moist, *αισος*, a varix, and *κυλη*, a tumour.) Dilated spermatic veins, with dropsy of the scrotum.

HYDROCOLLYRIUM. (From *υδρ*, liquid, and *κολλυριον*, a collyrium.) A collyrium composed of liquids.

HYGROLOGY. (*Hygrolagia*; from *υδρ*, a humour or fluid, and *λογος*, a discourse.) The doctrine of the fluids.

HYGROMA. (*υγρομα*: from *υδρ*, a liquid.) An encysted tumour, whose contents are either serum or a fluid like lymph. It sometimes happens that these tumours are filled with hyalids. Hygromatous tumours require the removal of the cyst, or the destruction of its secreted surface.

HYGROMETER. (*Hygrometrum*; from *υδρ*, moist, and *μετρον*, a measure.) Hygrometer. An instrument to measure the degrees of moisture in the atmosphere. It also means an infirm part of the body, affected by moisture of the atmosphere.

HYGROSCOPICS. Substances which have the property of absorbing moisture from the atmosphere. See *Atmosphere*.

HYGROMTRUM. (From *υδρ*, moist, and *μυρον*, a liquid ointment.) A liquid ointment.

HYDROPHOBIA. The same as hydrophobia.

HYLE. (*υλη*, matter.) The materia medica, or matter of any kind which comes under the cognizance of a medical person.

HYMEN. (From *Hymen*, the god of marriage, because this membrane is supposed to be entire before marriage, or copulation.) The hymen is a thin membrane, of a semilunar or circular form, placed at the entrance of the vagina, which it partly closes. It has a very different appearance in different women, but is generally, if not always, found in virgins, and is very properly esteemed the test of virginity, being ruptured in the first act of coition. The remnants of the hymen are called the *carunculae myrtiformes*. The hymen is also peculiar to the human species. There are two circumstances relating to the hymen which require medical assistance. It is sometimes of such a strong ligamentous texture, that it cannot be ruptured, and prevents the connection between the sexes. It is also sometimes imperforated, wholly closing the entrance into the vagina, and preventing any discharge from the uterus; but both these cases are extremely rare. If the hymen be of an unnaturally firm texture, but perforated, though perhaps with a very small opening, the inconve-

nienies thence arising will not be discovered before the time of marriage, when they may be removed by a crucial incision made through it, taking care not to injure the adjoining parts.

The imperforation of the hymen will produce its inconveniencies when the person begins to menstruate. For the menstrual blood, being secreted from the uterus at each period, and not evacuated the patient suffers much pain from the distention of the parts, many strange symptoms and appearances are occasioned, and suspicious injurious to her reputation are often entertained. In a case of this kind, for which Dr. Denman was consulted, the young woman, who was twenty-two years of age, having many uterine complaints, with the abdomen enlarged, was suspected to be pregnant, though she persevered in asserting the contrary, and had never menstruated. When she was prevailed upon to submit to an examination, the circumscribed tumour of the uterus was found to reach as high as the navel, and the external parts were stretched by a round soft substance at the entrance of the vagina, in such a manner as to resemble that appearance which they have when the head of a child is passing through them; but there was no entrance into the vagina. On the following morning, an incision was carefully made through the hymen, which had a fleshy appearance, and was thickened in proportion to its distention. Not less than four pounds of blood, of the colour and consistency of tar, were discharged; and the tumefaction of the abdomen was immediately removed. Several stellated incisions were afterwards made through the divided edges, which is a very necessary part of the operation; and care was taken to prevent a re-union of the hymen till the next period of menstruation, after which she suffered no inconvenience. The blood discharged was not putrid or coagulated, and seemed to have undergone no other change, after its secretion, but what was occasioned by the absorption of its more fluid parts. Some caution is required, when the hymen is closed in those who are in advanced age, unless the membrane be distended by the confined menses; as Dr. Denman once saw an instance of inflammation of the peritonæum being immediately produced after the operation, of which the patient died as in the true puerperal fever, and no other reason could be assigned for the disease.

The *carunculae myrtiformes*, by their elongation and enlargement, sometimes become very painful and troublesome.

HYMENEA COURBARIL. (*Hymenea*, corrupted *anime*, or *animea*.) The systematic name of the tree which affords the resin anime. See *Anime*.

HYO. Names compounded of this word

belong to muscles which originate from, or are inserted into, or connected with, the os hyoides; as, *Hyo-glossus*, *Hyo-pharyngeus*, *Genio-hyo-glossus*, &c.

HYO GLOSSUS. *Cerato-glossus* of Douglas and Cowper. *Basio-cerato-chondro-glossus* of Albinus. *Hyo-chondro-glossus* of Dumas. A muscle situated at the sides, between the os hyoides and the tongue. It arises from the basin, but chiefly from the corner of the os hyoides, running laterally and forwards to the tongue, which it pulls inwards and downwards.

HYOIDES OS. (*os hyoideum*: from the Greek letter υ , and *os*, likeness; so named from its resemblance.) This bone, which is situated between the root of the tongue and the larynx, derives its name from its supposed resemblance to the Greek letter υ , and is, by some writers, described along with the parts contained in the mouth. Ruysch has seen the ligaments of the bone so completely ossified, that the os hyoides was joined to the temporal bones by ankylosis. In describing this bone, it may be distinguished into its body, horns, and appendices. The body is the middle and broadest part of the bone, so placed that it may be easily felt with the finger in the fore part of the throat. Its fore-part, which is placed towards the tongue, is irregularly convex, and its inner surface, which is turned towards the larynx, is unequally concave. The *cornua*, or horns, which are flat, and a little bent, are considerably longer than the body of the bone, and may be said to form the sides of the υ . These horns are thickest near the body of the bone. At the extremity of each is observed a round tubercle, from which a ligament passes to the thyroid cartilage. The appendices, or lesser horns, *cornua minora*, as they are called by some writers, are two small processes, which in their size and shape are somewhat like a grain of wheat. They rise up from the articulations of the *cornua*, with the body of the bone, and are sometimes connected with the styloid process on each side, by means of a ligament. It is not unusual to find small portions of bone in these ligaments; and Ruysch, as we have already observed, has seen them completely ossified. In the fœtus, almost the whole of the bone is in a cartilaginous state, excepting a small point of bone in the middle of its body, and in each of its horns. The appendices do not begin to appear till after birth, and usually remain cartilaginous many years. The os hyoides serves to support the tongue, and affords attachment to a variety of muscles, some of which perform the motions of the tongue, while others act on the larynx and fauces.

HYOPHARYNGEUS. (From *os hyoideum*,

the hyoid bone, and *φαρυγξ*, the pharynx.) A muscle so called from its origin in the os hyoides, and its insertion in the pharynx.

HYOPHTHALMUS. (From *os*, a swine, and *οφθαλμος*, an eye; so named from the supposed resemblance of its flower to a hog's-eye.) Golden starwort; hog's-eye plant.

HYOSCIAMUS. (From *os*, a swine, and *σκιμος*, a bean; so named because hogs eat it as a medicine, or it may be because the plant is hairy and bristly, like a swine.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogymia*.

2. The pharmacopœial name of the henbane, called also *Faba ruilla*. *Apollinaria altercum*. *Agone*. *Altercagenou*. Common or black henbane. *Hyosciamus niger*; *foliis amplexicaulis sinuatis, floribus sessilibus* of Linnæus.

The leaves of this plant, when recent, have a slightly fetid smell, and a mucilaginous taste; when dried, they lose both taste and smell, and part also of their narcotic power. The root possesses the same qualities as the leaves, and even in a more eminent degree.

Henbane resembles opium in its action, more than any other narcotic does. In a moderate dose, it increases at first the strength of the pulse, and occasions some sense of heat, which are followed by diminished sensibility and motion; in some cases by thirst, sickness, stupor, and dimness of vision. In a large quantity, it occasions profound sleep, hard pulse, and sometimes fierce delirium, ending in coma, or convulsions, with a remarkable dilatation of the pupil, distortion of the countenance, a weak tremulous pulse, and eruption of petechiæ. On dissection, gangrenous spots have been found on the internal surface of the stomach. Its baneful effects are best counteracted by a powerful emetic, and by drinking largely of the vegetable acids.

Henbane has been used in various spasmodic and painful diseases, as in epilepsy, hysteria, palpitation, headache, paralysis, mania, and scirrhus. It is given in the form of the inspissated juice of the fresh leaves, the dose of which is from one to two grains; which requires to be gradually increased. It is sometimes employed as a substitute for opium, where the latter, from idiosyncrasy, occasions any disagreeable symptom. The henbane also is free from the constipating quality of the opium.

HYOSCIAMUS ALBUS. This plant, a native of the south of Europe, possesses similar virtues to the *hyosciamus niger*. See *Hyosciamus*.

HYOSCIAMUS LUTEUS. A species of tobacco.

HYOSCIAMUS NIGER. The systematic name of henbane. See *Hyosciamus*.

HYOTRICHOIDES. (From *os hyoideum*, the hy-

oid bone, and θυροειδής, the thyroid cartilage.) A muscle named from its origin in the hyoid bone and insertion in the thyroid cartilage.

HYPACTICA. (From ὑπαγω, to subdue.) Medicines which evacuate the faeces.

HYPALEIPTUM. (From ὑπαλείφα, to spread upon.) A spatula for spreading ointments with.

HYPELATA. (From ὑπέραιω, to move.) Cathartics

HYPERÆSTHESIS. (From ὑπεραίσθησις, to feel excess.) Error of appetite, whether by excess or deficiency. It is synonymous with Dr Cullen's order of *Dysorexia*.

HYPERCATHARSIS. (From ὑπερκαθάρσις, *supra*, over or above, and καθαίρω, to purge.) *Hyperineis Hyperinos.* An excessive purging from medicines.

HYPERCYPHOSIS. (From ὑπερκύψω, above, and κρυφή, the vertex.) A prominence, or protuberance. Hippocrates calls the lobes of the liver and lungs *Hypercypophones*.

HYPERCRISIS. (From ὑπερκρίσις: from ὑπερ, over or above, and κρίνω, to separate.) A critical excretion above measure; as when a fever terminates in a looseness, the humours may flow off faster than the strength can bear, and therefore it is to be checked.

HYPERDROSIS. (From ὑπερδρῖσις, in excess, and ὕδωρ, water.) A great distention of any part from water collected in it.

HYPEREMESIS. (From ὑπερῆσις, in excess, and ἔμεω, to vomit. An excessive evacuation by vomiting.

HYPEREPHRIDIOSIS. (From ὑπερῆφριδις, excess, and ἥδρω, sweat.) Immoderate sweating.

HYPERICUM. (From ὑπερ, over, and εἰμα, an image, or specter; so named because it was thought to have power over and to drive away evil spirits.) 1 The name of a genus of plants in the Linnæan system. Class. *Polyadelphia*. Order, *Polyandria*. St. John's wort.

2. The pharmacopœial name of the perforated or common St. John's wort, called also *fuga demonum*, and *androsemum*. *Hypericum perforatum* of Linnæus:—*floribus trigynis, caule ancipiti, foliis obtusis pellucido punctatis*. This indigenous plant was greatly esteemed by the ancients, internally in a great variety of diseases, and externally as an anodyne and discutient, but is now very rarely used. The flowers were formerly used in our pharmacopœia, on account of the great proportion of resinous oily matter, in which the medical efficacy of that plant is supposed to reside, but are now omitted.

HYPERICUM PERFORATUM. The systematic name of the St. John's wort. See *Hypericum*.

HYPERICUM SAXATILE. *Hypericoides. Coris lutea. Coris legitima cretica.* Bastard St. John's wort. The seeds are said to be diuretic, emmenagogue, and powerfully antispasmodic.

HYPERINA. (From ὑπερ, in excess, and ἵνω, to evacuate.) Medicines which purge excessively.

HYPERINESIS. See *Hypercatharsis*.

HYPERINOS. See *Hypercatharsis*.

HYPEROA. (From ὑπερ, above, and ὄστρον, The palate.

HYPEROPHARYNGÆUS. (From ὑπερ, above and φαρυγξ, the pharynx.) A muscle named from its situation above the pharynx.

HYPEROSTOSIS. (From ὑπερ, upon, and ὄστρον, a bone. See *Exostosis*.

HYPEROUM. (From ὑπερ, above, and ὤστρον, the roof, or palate.) A foramen in the upper part of the palate.

HYPERSARCOMA. (From ὑπερ, in excess, and σαρκώξ, flesh.) *Hypersarcosis.* A polypus in the nose. A fleshy excrescence. A polypus.

HYPERSARCOSIS. See *Hypersarcoma*.

HYPERODOS. (ὑπερὸς: from ὑπο, under, and ἔξω, passing out.) A flux of the belly.

HYPNOBATES. (From ὑπνός, sleep, and βαίνω, to go.) *Hypnobotasis.* One who walks in his sleep. See *Oneirodynia*.

HYPNOLOGIA. (From ὑπνός, sleep, and λόγος.) A dissertation, or directions for the due regulation of sleeping and waking.

HYPNOPOIETICA. (From ὑπνός, sleep, and ποιῶ, to cause.) Medicines which procure sleep. See *Anodynes*.

HYPNOTICS (*Hypnotica*, sc. *medicamenta*, ὑπνωτικά; from ὑπνός, to sleep.) See *Alnodynæ*.

HYPOÆMA. (From ὑπο, under, and αἷμα, blood; because the blood is under the cornea.) An effusion of red blood into the chambers of the eye.

HYPOCARODES. (From ὑπο and καρός, a carus.) *Hypocarothis.* One who labours under a low degree of carus.

HYPOCATHARSIS. (From ὑπο, under, and καθαίρω, to purge.) It is when a medicine does not work so much as expected, or but very little. Or a slight purging, when it is a disorder.

HYPOCASTRUM. (From ὑπο, sub, under, and καίω, to burn.) A stove, or hot-house, or any such-like contrivance; or place to sweat in, or to preserve plants from cold air.

HYPOCERCHALEON. (From ὑπο and κερχάριον, an aperity of the fauces. A stridulous kind of asperity of the fauces.

HYPOCHLOMENOS. (From ὑπο, under, and χέω, to pour.) One who labours under a catarrh.

HYPOCHLOROSIS. (From ὑπο, and χλωρόσις, the green sickness.) A light degree of chlorosis.

HYPOCHONDRIAC REGIONS. (*Regiones hypochondriacæ*; from ὑπο under, and χονδρός, a cartilage.) *Hypochondria.* The spaces in the abdomen that are under

the cartilages of the spurious ribs on each side of the epigastrium.

HYPOCHONDRIASIS. From *υπο*, *υπο* is hipped.) *Hypochondriacus morbus.* *Affectio hypochondriaca.* *Passio hypochondriaca.* The hypochondriac affection. Vapours, spleen, &c. A genus of disease in the class *neuroses*, and order *adynamiae*, of Cullen, characterized by dyspepsia; languor, and want of energy; sadness and fear, from uncertain causes; with a melancholic temperament.

The state of mind peculiar to hypochondriacs is thus described by Cullen:—"A languor, listlessness, or want of resolution and activity, with respect to all undertakings; a disposition to seriousness, sadness, and timidity, as to all future events, and apprehension of the worst or most unhappy state of them; and, therefore, often upon slight grounds, an apprehension of great evil. Such persons are particularly attentive to the state of their own health, to every the smallest change of feeling in their bodies; and from any unusual sensation, perhaps of the slightest kind, they apprehend great danger, and even death itself. In respect to these feelings and fears, there is commonly the most obstinate belief and persuasion." He adds, that it is only when the state of mind just described is joined with indigestion, in either sex, somewhat advanced in years, of a melancholic temperament, and a firm and rigid habit, that the disease takes the name of *Hypochondriacism*.

The seat of the hypochondriac passion is in the stomach and bowels; for first these parts are disordered, then the others suffer from the connection.

The causes are, sorrow, fear, or excess of any of the passions; too long continued watching; irregular diet. Those habitually disposed to it, (and these causes have little effect in other constitutions,) have generally a sallow or brown complexion, and a down-cast look; a rigidity of the solids, and torpor of the nervous system. Whatever may occasion nervous disorders in general, may be the cause of this in particular.

The signs of this complaint are so various, that to describe them is to describe almost every other disease; but, in general, there is an insurmountable indolence, dejected spirits, dread of death, costiveness, a slow and somewhat difficult inspiration, flatulencies in the primæ viæ, and various spasmodic affections. It is seldom fatal; but if neglected, or improperly treated, may bring on incurable melancholy, jaundice, madness, or vertigo, palsy, and apoplexy.

On dissections of hypochondriac persons, some of the abdominal viscera (particularly the liver and spleen) are usually found considerably enlarged. In some few instances, effusion and a turgescence of

the vessels have been observed in the brain.

HYPOCHONDRIUM. (From *υπο* under, and *χονδρος*, a cartilage.) That part of the body which lies under the cartilages of the spurious ribs.

HYPOCHYMA. (From *υπο*, and *χυς*, to pour; because the ancients thought that the opacity proceeded from something running under the crystalline humour.) *Hypochyma.* A cataract.

HYPOCISTIS. (From *υπο*, under, and *κυστις*, the cistis.) A plant called by Linnaeus *Asarum hypocistis*, a parasitical plant, growing in warm climates, from the roots of the cistis. The juice, *succus hypocistidis* is a mild astringent, of no particular smell nor flavour. It is seldom used.

HYPOCLEFTICUM. (From *υπο*, under, and *κλεψω*, to steal.) A chymical vessel for separating liquors, particularly the essential oil of any vegetable from the water; and named because it steals, as it were, the water from the oil.

HYPOCORION. (From *υπο*, under, and *κορυον*, the cavity above the upper eye-lid.) The cavity under the lower eye-lid.

HYPOCOPHOSIS. *Cophosis*, but in a less degree.

HYPOCRANIUM. (From *υπο*, under, and *κρανιον*, the skull.) A kind of abscess, so called because situated under the cranium, between it and the dura mater.

HYPOCLITHIS. In Rufus Ephesius, it is the extremity of the fore-part of the neck.

HYPODERMIS. (From *υπο*, under, and *δερμα*, the skin.) The cuticle under the clitoris, which covers it like a prepuce. The clitoris.

HYPODERMA. (From *υπο*, under, and *δω*, to bind.) An underswathe, or bandage.

HYPODERMUM. A bandage like the former.

HYPOGALA. (From *υπο*, under, and *γαλα*, milk; because it is a milk-like effusion, under the cornea.) A collection of white humour, like milk, in the chambers of the eye. There are two species of this disease: the one takes place, it is said, from a deposition of the milk, as is sometimes observed in women who suckle; the other from a depression of the milky cataract.

HYPOGASTRIC ARTERIES. See *Ilia ac arteries*.

HYPOGASTRIC REGION. (*Regio hypogastrica*; from *υπο*, under, and *γαστρο*, the stomach.) The region of the abdomen that reaches from above the pubis to within three fingers' breadth of the navel.

HYPOGASTRIUM. (From *υπο*, under, and *γαστρο*, the stomach.) The lower region of the part of the belly.

HYPOGASTROCELE. (From *υπογαστρον*, the hypogastrium, and *κελε*, a tumour.)

A tumour, or hernia, in the hypogastric region.

HYPOGLOSSIA. (From *υπο*, under, and *γλῶσσα*, the tongue.) The under part of the tongue, which adheres to the lower jaw.

HYPOGLOSSUS. (From *υπο*, under, and *γλῶσσα*, the tongue.) A nerve which goes to the under part of the tongue.

HYPOGLOTTIDES. (From *υπο*, under, and *γλῶττις*, the tongue.) They are a kind of medicine to be held under the tongue until they are dissolved.

HYPOGLUTIS. (From *υπο*, under, and *γλῦτος*, the nates.) It is the fleshy part under the nates towards the thigh. Some say it is the flexure of the coxa, under the nates.

HYPOMIA. (From *υπο*, under, and *αμος*, shoulder. In Galen's *Exegesis* it is the part subjacent to the shoulder.

HYPONOMOS. (From *υπονομος*, a phagedenic ulcer.) A subterraneous place. A deep phagedenic ulcer.

HYPOPIDIUM. (From *υπο*, under, and *πυς*, the foot.) A cataplasm for the sole of the foot.

HYPOPHORA. (From *υποφορας*, to be carried or conveyed underneath.) A deep fistulous ulcer.

HYPOPTHALMION. (From *υπο*, under, and *οφθαλμος*, the eye.) The part under the eye which is subject to swell in a cachexy, or dropsy.

HYPOPHYSIS. (From *υπο*, under, and *φωσ*, to produce.) A disease of the eye-lids, when the hairs grow so much as to irritate and offend the pupil.

HYPOPIUM. (From *υπο*, under, and *πυς*, pus; because the pus is under the cornea.) *Hypopyon*. *Pycsis*. *Abcessus oculi*. An accumulation of a glutinous yellow fluid, like pus, which takes place in the anterior chamber of the aqueous humour, and frequently also in the posterior one, in consequence of severe, acute ophthalmia, particularly the internal species.

This viscid matter of the hypopium is commonly called pus; but Scarpa contends that it is only coagulating lymph. The symptoms portending an extravasation of coagulable lymph in the eye, or an hypopium, are the same as those which occur in the highest stage of violent acute ophthalmia, viz. prodigious tumefaction of the eye-lids; the same swelling and redness as in chemosis; burning heat and pain in the eye; pains in the eye-brow and nape of the neck; fever, restlessness, aversion to the faintest light, and a contracted state of the pupil.

HYPOPIUM. (From *υπο*, under, and *πυς*, the nose.) A name for the parts of the upper lip below the nostrils.

HYPOBARA. (From *υπο*, under, and *σαρξ*, flesh.) *Hypobaroides*. An anasarea.

In Dr. Cullen's *Nosology*, it is synonymous with *Physconia*.

HYPOSPADIMOS. (From *υπο*, under, and *σπαις*, to dr.w.) The urethra terminating under the glans.

HYPOSPATHISMUS. (From *υπο*, under, and *σπαθις*, a spatula.) The name of an operation formerly used in surgery, for removing defluxions in the eyes. It was thus named from the instrument with which it was performed.

HYPOSPHAGMA. (From *υπο*, under, and *σφαζω*, to kill.) *Hyposphagma*. An extravasation of blood in the tunica adnata of the eye, from external injury.

HYPOSPLENIA. (From *υπο*, under, and *σπλην*, the spleen.) A small tumour of the spleen.

HYPOSTAPHYLE. (From *υπο*, and *σφαυλη*, the uvula.) Relaxation of the uvula.

HYPOSTASIS. (From *υποστημι*, to subside.) A sediment, as the sediment in urine.

HYPOTHENAR. (From *υπο*, under, and *θεναξ*, the palm of the hand.) A muscle which runs on the inside of the hand. Also that part of the hand which is opposite to the palm.

HYPOTHESIS. A system of general rules, founded partly on fact and partly on conjecture. A theory explains every fact: an hypothesis explains only a certain number of facts, leaving some unaccounted for, and others in opposition to it.

HYPOTHETON. (From *υπο*, under, and *τιθεμι*, to put.) A suppository, or medicine introduced into the rectum, to procure stools.

HYPOXYLON. (From *υπο*, and *ξύλον*, wood.) A species of *clavaria*, which grows under old wood.

HYPOZOMA. (From *υπο*, and *ζωμη*, to bind round.) The diaphragm.

HYPSILOID. (From *υψιλοειδης*, the hypsiloid bone, and *γλῶσσα*, the tongue.) A muscle named from its origin in the os hyoides, and its insertion in the tongue.

HYPSILOIDES. A name of the *Os Hyoides*; also of the *Basiglossus muscle*. See *Hyoglossus*.

HYPTIAMOS. (From *υπτιάζω*, to lie with the face upwards.) A supine decubiture, or a nausea, with inclination to vomit.

HYPULOS. (From *υπο*, under, and *υλος*, a cicatrix.) An ulcer which lies under a cicatrix.

Hyssop. See *Hyssopus*.

Hyssop, hedge. See *Gratiola*.

HYSSOFITES. (From *υσσωπης*, hyssop.) Wine impregnated with hyssop.

HYSSOPUS. (*υσσωπης*: from *Αζωβ*, Heb.)

1. The name of a genus of plants in the Linnæan system. Class, *Didynamia*. Order, *Gymnospermia*. Hyssop.

2 The pharmacopœial name of the common hyssop. *Hyssopus officinalis* of Linneus — *apicis secundis, foliis lanceolatis*.

This exotic plant is esteemed as an aromatic and stimulant, but is chiefly employed as a pectoral, and has long been thought useful in humoral asthmas, coughs, and catarrhal affections; for this purpose, an infusion of the leaves, sweetened with honey, or sugar, is recommended to be drank as tea.

HYSSOPUS CAPITATA. Wild thyme.

HYSSOPUS OFFICINALIS. The systematic name of the common hyssop. See *Hyssopus*.

HYSTERA. (From *ὕστερ*, behind; so called because it is placed behind the other parts.) The uterus, or womb.

HYSTERALGIA. (From *ὕστερ*, the womb, and *ἀλγος*, pain.) A pain in the womb.

HYSTERIA. (From *ὕστερ*, the womb, from which the disease was supposed to arise.) *Passio hysterica.* Hysterica. Dr. Cullen places this disease in the class *neuroses*, and order *spasmi*. There are four species:

1. *Hysteria chlorotica*, from a retention of the menses.
2. *Hysteria à leucorrhœa*, from a fluor albus.
3. *Hysteria à menorrhagia*, from an immoderate flow of the menses.
4. *Hysteria libidinosa*, from sensual desires.

The complaint appears under such various shapes, imitates so many other diseases, and is attended with such a variety of symptoms, which denote the animal and vital functions to be considerably disordered, that it is difficult to give a just character or definition of it; and it is only by taking an assemblage of all its appearances, that we can convey a proper idea of it to others.

The disease attacks in paroxysms, or fits. These are sometimes preceded by dejection of spirits, anxiety of mind, effusion of tears, difficulty of breathing, sickness at the stomach, and palpitations at the heart; but it more usually happens, that a pain is felt on the left side, about the flexure of the colon, with a sense of distention advancing upwards, till it gets into the stomach, and removing from thence into the throat, it occasions, by its pressure, a sensation as if a ball was lodged there, which by authors has been called *globus hystericus*. The disease having arrived at this height, the patient appears to be threatened with suffocation, becomes faint, and is affected with stupor and insensibility; whilst, at the same time, the trunk of the body is turned to and fro, the limbs are variously agitated, wild and irregular actions take place in alternate fits of laughter, crying, and screaming; incoherent expressions are uttered, a temporary delirium prevails, and a frothy saliva is discharged from the mouth. The spasms at length abating, a quantity of wind is evacuated upwards, with frequent sighing

and sobbing, and the woman recovers the exercise of sense and motion without any recollection of what has taken place during the fit, feeling, however, a severe pain in her head, and a soreness over her whole body.

In some cases, there is little or no convulsive motion, and the person lays seemingly in a state of profound sleep, without either sense or motion.

Hiccup is a symptom which likewise attends in some instances, on hysteria; and now and then it happens, that a fit of hysteria consists of this alone. In some cases of this nature, it has been known to continue for two or three days, during which, it frequently seems as if it would suffocate the patient, and proceeds, gradually weakening her, till it either goes off, or else occasions death by suffocation; but this last is extremely rare. Besides hiccup, other slight spasmodic affections sometimes wholly form a fit of hysteria, which perhaps continue for a day or two, and then either go off of themselves, or are removed by the aid of medicine.

In some cases, the patient is attacked with violent pains in the back, which extend from the spine to the sternum, and at length become fixed upon the region of the stomach, being evidently of a spasmodic nature, and often prevailing in so high a degree as to cause clammy sweats, a pale cadaverous look, coldness of the extremities, and a pulse hardly perceptible.

Hysteric affections occur more frequently in the single state of life than in the married; and that most usually between the age of puberty and that of thirty-five years; and they make their attack oftener about the period of menstruation than at any other.

They are readily excited in those who are subject to them, by passions of the mind, and by every considerable emotion, especially when brought on by surprise; hence sudden joy, grief, fear, &c. are very apt to occasion them. They have also been known to arise from imitation and sympathy.

Women of a delicate habit, and whose nervous system is extremely sensible, are those who are most subject to hysteric affections; and the habit which predisposes to their attacks is acquired by inactivity and a sedentary life, grief, anxiety of mind, a suppression or obstruction of the menstrual flux, excessive evacuations, and a constant use of a low diet, or of crude unwholesome food.

Hysteria differs from hypochondriasis in the following particulars, and, by paying attention to them, may always readily be distinguished from it. Hysteria attacks the sanguine and plethoric; comes on soon after the age of puberty; makes its onset suddenly and violently, so as to deprive the

patient of all sense and voluntary motion; is accompanied with the sensation of a ball rising upwards in the throat, so as to threaten suffocation; is attended usually with much spasmodic affection; is more apt to terminate in epilepsy than in any other disease; and, on dissection, its morbid appearances are confined principally to the uterus and ovaria.

The reverse happens in hypochondriasis. It attacks the melancholic; seldom occurs till after the age of thirty-five; comes on gradually; is a tedious disease, and difficult to cure; exerts its pernicious effects on the membranous canal of the intestines, as well by spasms as wind; is more apt to terminate in melancholy, or a low fever, than in any other disease; and, on dissection, exhibits its morbid effects principally on the liver, spleen, and pancreas, which are often found in a hard, scirrhusous, or corrupted state.

Another very material difference might be pointed out betwixt these two diseases, which is, that hysteria is much relieved by advancing in age, whereas hypochondriasis usually becomes aggravated.

The two diseases have often been confounded together; but, from considering the foregoing circumstances, it appears that a proper line of distinction should be drawn between them.

The hysteric passion likewise differs from a syncope, as in this there is an entire cessation of the pulse, a contracted face, and a ghastly countenance; whereas, in the uterine disorder, there is often something of a colour, and the face is more expanded; there is likewise a pulse, though languid; and this state may continue two or three days, which never happens in a syncope.

It also differs from apoplexy, in which the abolition of sense and voluntary motion is attended with a sort of snoring, great difficulty of breathing, and a quick pulse; which do not take place in hysteric cases.

It differs from epilepsy, in that this is supposed to arise in consequence of a distention of the vessels of the brain; whereas, in hysteria, the spasmodic and convulsive motions arise from a turgescence of blood in the uterus, or in other parts of the genital system.

However dreadful and alarming an hysteric fit may appear, still it is seldom accompanied with danger, and the disease never terminates fatally, unless it changes into epilepsy, or that the patient is in a very weak reduced state.

HYSTERIA CHLOROTICA. Hysterics from obstructed menses. See *Hysteria*.

HYSTERIA FEBRILIS. A tertian fever, with spasms and convulsions.

HYSTERIA A LEUCORRHEA. Hysterics from fluor albus. See *Hysteria*.

HYSTERIA LIBIDINOSA. Nymphomania, or female libidinous propensity. See *Hysteria*.

HYSTERIA A MENORRHAGIA. Hysterics from profuse menses. See *Hysteria*.

HYSTERICALGIA. (From *ὤστρος*, the womb, and *αλγος*, pain.) An epithet for any thing that excites pain in the uterus. Hippocrates applies this word to vinegar; and others signify by it the pains which resemble labour-pains, generally called *false pains*.

HYSTERITIS (From *ὤστρος*, the womb.) *Metritis*. Inflammation of the womb. A genus of disease in the class *pyrexia*, and order *phlegmasia*, of Cullen; characterized by pyrexia, heat, tension, tumour, and pain in the region of the womb; pain in the os uteri when touched, and vomiting.

In natural labours, as well as those of a laborious sort, many causes of injury to the uterus, and the peritonæum which covers it, will be applied. The long continued action of the uterus on the body of the child, and the great pressure made by its head on the soft parts, will farther add to the chance of injury. Besides these, an improper application of instruments, or an officiousness of the midwife in hurrying the labour, may have contributed to the violence. To these causes may be added exposure to cold, by taking the woman too early out of bed after delivery, and thereby throwing the circulating fluids upon the internal parts, putting a stop to the secretion of milk, or occasioning a suppression of the *lactia*.

An inflammation of the womb is sometimes perfectly distinct, but is more frequently communicated to the peritonæum, Fallopian tubes, and ovaria; and having once begun, the natural functions of the organ become much disturbed, which greatly adds to the disease.

It is oftener met with in women of a robust and plethoric habit than in those of lax fibres and a delicate constitution, particularly where they have indulged freely in food of a heating nature, and in a use of spirituous liquors. It never prevails as an epidemic, like puerperal fever, for which it has probably often been mistaken; and to this we may, with some reason, ascribe the difference in the mode of treating the disease, which has taken place among physicians.

An inflammation of the uterus shews itself usually about the second or third day after delivery, with a painful sensation at the bottom of the belly, which gradually increases in violence, without any kind of intermission. On examining externally, the uterus appears much increased in size, is hard to the feel, and, on making a pressure upon it, the patient experiences great soreness and pain.

Soon afterwards there ensues an increase

in heat over the whole of the body, with pains in the head and back, extending into the groins, rigors, considerable thirst, nausea, and vomiting. The tongue is white and dry, the secretion of milk is usually much interrupted, the lochial discharge is greatly diminished, the urine is high-coloured and scanty, and if the inflammation is extended to the bladder, is then so totally obstructed as to render the use of a catheter necessary; the body is costive, and the pulse is hard, full, and frequent.

These are the symptoms which usually present themselves, when the inflammation does not run very high, and is perfectly distinct; but when it is so extensive as to affect the peritonæum, those of irritation then generally succeed, and soon destroy the patient.

Uterine inflammation is always attended with much danger, particularly where the symptoms have ran high, and the proper means for removing them have not been timely adopted. In such cases, it may terminate either in suppuration, scirrhus, or gangrene.

Frequent rigors, succeeded by flushings of the face, quickness and weakness of the pulse, great depression of strength, delirium, and the sudden cessation of pain and soreness in the region of the abdomen, denote a fatal termination: on the contrary, the ensuing of a gentle diarrhœa, the lochial discharge returning in due quantity and quality, the secretion of milk recommencing, and the uterus becoming gradually softer and less tender to the touch, with an abatement of heat and thirst, prognosticate a favourable issue.

When shiverings attack the patient, after several days continuance of the symptoms, but little relief can be afforded by medicine, the event being generally fatal. In this case, the woman emaciates and loses her strength, becomes hectic, and sinks under colliquative sweating, or purging.

Upon opening the bodies of women who have died of this disease, and where it existed in a simple state, little or no extravasated fluid is usually to be met with in the cavity of the abdomen. In some instances, the peritoneal surfaces have been discovered free from the disease; whilst

in others, that portion which covers the uterus and posterior part of the bladder has been found partially inflamed. The inflammation has been observed, in some cases, to extend to the ovaria and Fallopian tubes, which, when cut open, are often loaded with blood. The uterus itself usually appears of a firm substance, but is larger than in its natural state, and, when cut into, a quantity of pus is often found. Gangrene is seldom, if ever, to be met with.

HYSTEROCELE, (From *uterus*, the womb, and *cele*, a tumour.) An hernia of the womb. This is occasioned by violent muscular efforts; by blows on the abdomen at the time of gestation, and also by wounds and abscesses of the abdomen which permit the uterus to dilate the part. Ruysch relates the case of a woman, who, becoming pregnant after an ulcer had been healed in the lower part of the abdomen, the tumid uterus descended into a dilated sac of the peritonæum in that weakened part, till it hung, with the included fœtus, at her knees. Yet, when her full time was come, the midwife reduced this wonderful hernia, and in a natural way she was safely delivered of a son.

HYSTEROCECISTICUS. (From *uterus*, the womb, and *cystis*, the bladder.) Applied to a suppression of urine from the pressure of the uterus against the neck of the bladder.

HYSTEROX. (From *uterus*, afterwards; so named because it comes immediately after the fœtus.) The placenta.

HYSTEROPTISA. (From *uterus*, the womb, and *ptisis*, flatus.) The womb distended with air.

HYSTEROPTOSIS. (From *uterus*, the womb, and *ptosis*, to fall.) A bearing down of the womb.

HYSTEROTOMY. (From *uterus*, the womb, and *tomia* to cut.) See *Cæsarian operation*.

HYSTRICIASIS. (From *hystrix*, a hedgehog, or porcupine.) A disease of the hairs, in which they stand erect, like porcupine quills. An account of this rare disease is to be seen in the *Philosophical Transactions*, No. 424.

HYSTRICIS LAPIS *S. Bezoar porcinum*.

HYSTRITIS. See *Hysteritis*.

I

IBERIS. (So named from Iberia, the place of its natural growth.) 1. The name of a genus of plants in the Linnæan system. Class, *Tetradynamia*. Order, *Siliculosa*.

2. The pharmacopœial name of the plant which is also called *Cardimantica*. The *Lepidium iberis* of Linnæus. Sciatica cresses. It possesses a warm, penetrating, pungent taste, like unto other cresses, and is recommended as an antiscorbutic, antiseptic, and stomachic.

IBIRAC. See *Guaiacum*.

IBIRÆUM. A wild species of liquorice found in Brasil.

IBIRA PATANGA. Logwood.

IBIS. *Ibis* was a bird much like our kingfisher, taken notice of by the Egyptians, because, when it was sick, it used to inject with its long bill the water of the Nile into its fundament, whence Langius, lib. ii. ep. ii. says they learned the use of clysters.

IBISCU. (From *ibis*, the stork, who was said to chew it, and inject it as a clyster.) Marshmallow.

IBIXUMA. (From *ibis*, the mallow, and *ibis*, glue; so named from its having a glutinous leaf, like the mallow.) The soap-tree, or *Saponaria Arbor*.

ICE. *Glacis*. Water made solid by the application of cold. It is frequently applied by surgeons to resolve external inflammatory diseases.

ICHOR. (*Ichor*.) A thin, aqueous, and acrid discharge.

ICTRYA. (*Ichthya*, a fish-hook; from *ichthys*, a fish.) The skin of the *Squatina*, or monkfish: also the name of an instrument like a fish-hook, for extracting the fœtus.

ICHTHYOCOLLA. (From *ichthys*, a fish, and *colla*, glue.) *Colla piscium*. Isinglass. Fish-glue. A substance, partly gelatinous, and partly lymphatic, which is prepared by rolling up the air-bladder of the *Accipenser sturio* of Linnæus, and several other fishes, and drying it in the air, after it has been twisted into the form of a short cord, as we receive it. It affords a viscid jelly by ebullition in water, which is used in medicine as an emollient in disorders of the throat, intestines, &c.

ICHTHYOSIS. (From *ichthys*, the scale of a fish; from the resemblance of the scales to those of a fish.) A genus of disease of the second order of Dr. Willan's diseases of the skin. The characteristic of ichthyosis is a permanently harsh, dry, scaly, and, in some cases, almost horny texture of the integuments of the body, unconnected with internal disorder. Psoriasis and Lepra differ from this affection, in

being but partially diffused, and in having deciduous scales. The arrangement and distribution of the scales in ichthyosis are peculiar. Above and below the olecranon on the arm, says Dr. Willan, and in a similar situation with respect to the petella on the thigh and leg they are small, rounded, prominent, or papillary, and of a black colour; some of the scaly papillæ have a short narrow neck, and broad irregular tops. On some part of the extremities, and on the trunk of the body, the scales are flat and large, often placed like tiling, or in the same order as scales on the back of a fish; but, in a few cases, they have appeared separate, being intersected by whitish furrows. There is usually in this complaint a dryness and roughness of the soles of the feet; sometimes a thickened and brittle state of the skin in the palms of the hands, with large painful fissures, and, on the face, an appearance of the scurf rather than of scales. The inner part of the wrists, the hams, the inside of the elbow, the furrow along the spine, the inner and upper part of the thigh, are perhaps the only portions of the skin always exempt from the scalliness. Patients affected with ichthyosis are occasionally much harassed with inflamed pustules, or with large painful boils on different parts of the body: it is also remarkable, that they never seem to have the least perspiration or moisture of the skin. This disease did not, in any case, appear to Dr. Willan to have been transmitted hereditarily; nor was more than one child from the same parents affected with it. Dr. Willan never met with an instance of the horny rigidity of the integuments, *Ichthyosis cornea*, impeding the motion of the muscles or joints. It is, however, mentioned by authors as affecting the lips, prepuce, toes, fingers, &c. and sometimes as extending over nearly the whole body.

ICTERITIA. (From *icterus*, the jaundice.) An eruption of yellowish spots. Also a yellow discoloration of the skin, without fever.

ICTERUS. (Named from its likeness to the plumage of the golden thrush, of which Pliny relates, that if a jaundiced person looks on one, the bird dies, and the patient recovers.) *Morbis arcuatus*, or *argutus*. *Aurigo*. *Morbis regius*. *Morbis leucoli*. The jaundice. A genus of disease in the class *cachexia*, and order *imperfecta*, of Cullen; characterized by yellowness of the skin and eyes; faces white, and urine of a high colour. There are six species:—

1. *Icterus calculosus*, acute pain in the epigastric region, increasing after eating; gall-stones pass by stools.

2. *Icterus spasmodicus*, without pain, after spasmodic diseases and passions of the mind.

3. *Icterus mucosus*, without either pain, gall-stones, or spasm, and relieved by the discharge of tough phlegm by stool.

4. *Icterus hepaticus*, from an induration in the liver.

5. *Icterus gravidarum*, from pregnancy and disappearing after delivery.

6. *Icterus infantum*, of infants.

It takes place most usually in consequence of an interrupted excretion of bile, from an obstruction in the ductus communis choledochus, which occasions its absorption into the blood-vessels. In some cases it may, however, be owing to a redundant secretion of the bile.

The causes producing the first of these are, the pressure of biliary calculi in the gall-bladder and its ducts; spasmodic constriction of the ducts themselves; and, lastly, the pressure made by tumours situated in adjacent parts; hence jaundice is often an attendant symptom on a scirrhus of the liver, pancreas, &c. and frequently likewise on pregnancy.

Chronic bilious affections are frequently brought on by drinking freely, but more particularly by spirituous liquors; hence they are often to be observed in the debauchee and the drinker of drams. They are likewise frequently met with in those who lead a sedentary life, and who indulge much in anxious thoughts.

A slight degree of jaundice often proceeds from the redundant secretion of the bile, and a bilious habit is therefore constitutional to some people, but more particularly to those who reside long in a warm climate.

By attending to the various circumstances and symptoms which present themselves, we shall in general be able to ascertain, with much certainty, the real nature of the cause which has given rise to disease.

We may be assured, by the long continuance of the complaint, and by feeling the liver and other parts externally, whether or not it arises from any tumour in this viscus, or the pancreas, mesentery, or omentum.

Where passions of the mind induce the disease, without any hardness or enlargement of the liver, or adjacent parts, and without any appearance of calculi in the feces, or on dissection after death, we are naturally induced to conclude that the disorder was owing to a spasmodic affection of the biliary ducts.

Where gall-stones are lodged in the ducts, acute lancinating pains will be felt

in the region of the parts, which will cease for a time, and then return again; great irritation at the stomach and frequent vomiting will attend, and the patient will experience an aggravation of the pain after eating. Such calculi are of various sizes, from a pea to that of a walnut; and, in some cases, are voided in a considerable number, being, like the gall, of a yellowish, brownish, or green colour.

The jaundice comes on with languor, inactivity, loathing of food, flatulency, acidities in the stomach and bowels, and costiveness. As it advances in its progress, the skin and eyes become tinged of a deep yellow; there is a bitter taste in the mouth, with frequent nausea and vomiting; the urine is very high coloured; the stools are of a grey or clayey appearance, and a dull obtuse pain is felt in the right hypochondrium, which is much aggravated by pressure with the fingers.

Where the pain is very acute, the pulse is apt to become hard and full, and other febrile symptoms to attend.

The disease, when of long continuance, and proceeding from a chronic affection of the liver, or other neighbouring viscera, is often attended with anasarca swellings, and sometimes with ascites.

Where jaundice is recent, and is occasioned by concretions obstructing the biliary ducts, it is probable that, by using proper means, we may be able to effect a cure; but where it is brought on by tumours of the neighbouring parts, or has arisen in consequence of other diseases, attended with symptoms of obstructed viscera, our endeavours will most likely not be crowned with success. Arising during a state of pregnancy, it is of little consequence, as it will cease on parturition.

On opening the bodies of those who die of jaundice, the yellow tinge appears to pervade even the most interior part of the body; it is diffused throughout the whole of the cellular membrane, in the cartilages and bones, and even the substance of the brain is coloured with it. A diseased state of the liver, gall-bladder, or adjacent viscera, is usually to be met with.

The *Icterus infantum*, or yellow gum, is a species of jaundice, which, for the most part, affects all children at, or soon after, their birth, and which usually continues for some days.

It has generally been supposed to arise from the meconium impacted in the intestines, preventing the flow of bile into them.

The effects produced by it are, languor, indolence, a yellow tinge of the skin, and a tendency to sleep, which is sometimes fatal, where the child is prevented from sucking.

ICTERUS ALBUS. The white jaundice, The chlorosis, or green-sickness, is sometimes thus called.

ICTUS. A stroke, or blow. Hence *ictus solis* means a stroke of the sun, or that affection which takes place from too great an influence of the sun's heat. It signifies also the pulsation of an artery, and the sting of a bee, or other insect.

IDÆUS. (From *idē*, a mountain in Phrygia, their native place.) A name of the pæony, and blackberry.

IDIOCRASIA. See *Idiosyncrasia*.

IDIOPATHIC. (*Idiopathicus*; from *idios*, peculiar, and *pathos*, an affection.) A disease which does not depend on any other disease, in which respect it is opposed to a symptomatic disease, which is dependant on another.

IDIOSYNCRASY. (*Idiosyncrasia*; from *idios*, peculiar *syn*, with, and *krasis*, a temperament.) A peculiarity of constitution, in which a person is affected by certain agents, which, if applied to a hundred other persons, would produce no effect. thus some people cannot see a finger bleed without fainting; and thus violent inflammation is induced on the skin of some persons by substances that are perfectly innocent to others.

IDIOTROPIA. (From *idios*, peculiar, and *tropeo*, to turn.) The same as *Idiosyncrasia*.

IGNATIA AMARA. The systematic name of the plant which affords St. Ignatius's bean. See *Faba indica*.

IGNATHI FABÆ. See *Faba indica*.

Ignatius's bean. See *Faba indica*.

IGNIS CALIDUS. A hot fire: so some call a gangrene: also a violent inflammation, just about to degenerate into a gangrene.

IGNIS FRIGIDUS. A cold fire. A sphaecelus hath been thus called, because the parts that are so affected become cold as the surrounding air.

IGNIS PERSICUS. A name of the erysipelas, also of the tumour called a carbuncle.

IGNIS ROTÆ. Fire for fusion. It is when a vessel which contains some matter for fusion is surrounded with live, i. e. red hot coals.

IGNIS SACER. A name of erysipelas, and of a species of *Herpes*.

IGNIS SAPIENTUM. Heat of horse-dung.

IGNIS SILVATICUS. A name of the *Impetigo*.

Ignis volagrius. A name of the *Impetigo*.

IGNIS VOLATICUS. See *Erysipelas*.

IKAN RADIX. A somewhat oval, oblong, compressed root, brought from China. It is extremely rare, and would appear to be the root of the orchis tribe.

ILAPRIS. A name in Myrepsus for the burdock.

ILECH. By this word, Paracelsus seems to mean a first principle.

ILENOS. In the Spagyric language it is the elementary air.

ILEOS CAUENTUM. Hippocrates describes it in lib. De Intern. Affect. In this disease, as well as in the scurvy, the breath is fetid, the gums recede from the teeth, hæmorrhages of the nose happen, and sometimes there are ulcers in the legs, but the patient can move about his business very well.

ILEUM. (From *ileo*, to turn about; from its convolutions.) *Ileum intestinum*. The last portion of the small intestines, about fifteen hands' breadth in length, which terminates at the valve of the cæcum. See *Intestines*.

ILEX. The name of a genus of plants in the Linnæan system. Class, *Tetrandria*. Order, *Tetragynia*.

ILEX AQUIFOLIUM. The systematic name of the common holly. See *Aquifolium*.

ILIA. (The plural of *Ile*, αλλ.) The flanks, or that part in which is enclosed the small intestines.

ILIAC ARTERIES. *Arterie iliace*. The arteries so called are formed by the bifurcation of the aorta, near the last lumbar vertebra. They are divided into *internal* and *external*. The *internal iliac*, also called the *hypogastric artery*, is distributed in the fœtus into six, and in the adult into five branches, which are divided about the pelvis, viz. the little iliac, the gluteal, the ischiatic, the pudical, and the obturator; and in the fœtus the umbilical. The *external iliac* proceeds out of the pelvis through Poupert's ligament, to form the femoral artery.

ILIAC PASSION. (ΕΙΛΑΟΣ, ΙΛΑΟΣ, ΜΑΛΑΟΣ, is described as a kind of nervous colic, whose seat is the ilium.) *Passio iliaca*. *Volvulus*. *Miserere mes*. *Convolutulus*. *Chordæperu*. *Tormentum*. A violent vomiting, in which the fecal portion of the food is voided by the mouth. See *Colica*.

ILIAC REGION. The side of the hypogastric region, just above the hips.

ILIACUS INTERNUS. *Iliacus* of Winslow. *Iliaco trachanten* of Dumas. A thick, broad, and radiated muscle, which is situated in the pelvis, upon the inner surface of the ilium. It arises fleshy from the inner lip of the ilium, from most of the hollow part, and likewise from the edge of that bone, between its anterior superior spinous process and the acetabulum. It joins with the psoas magnus, where it begins to become tendinous, and passing under the ligamentum Fallopii, is inserted in common with that muscle. The

tendon of this muscle has been seen distinct from that of the psoas, and, in some subjects, it has been found divided into two portions. The iliacus internus serves to assist the psoas magnus in bending the thigh, and in bringing it directly forwards.

ILIADUM. *Liadus*. It is the first matter of all things, consisting of mercury, salt, and sulphur. These are Paracelsus's three principles. His *iliadus* is also a mineral spirit, which is contained in every element, and is the supposed cause of diseases.

ILIASTER. Paracelsus says it is the occult virtue of nature, whence all things have their increase.

ILINEOS. (*ἰλινεος*: from *ἰλινξ*, a vortex.) A giddiness, in which all things appear to turn round, and the eyes grow dim.

ILISCUA. Avicenna says it is madness caused by love.

ILIUM OS. (From *ilia*, the small intestines: so named because it supports the *ilia*.) The haunch-bone. The superior portion of the os innominatum, which, in the fœtus, is a distinct bone. See *Innominatum os*.

ILLECEBRA. (From *ἄλλο*, to turn; because its leaves resemble worms.) *Vermicularis*. *Piper murale*. *Sedum minus*. Wall-pepper. Stone-crop. The plant thus called is the *Sedum acre* of Linnaeus; in its recent state it is extremely acrid, like the hydropiper; hence, if taken in large doses, it acts powerfully on the primæ viæ, proving both emetic and cathartic; applied to the skin as a cataplasm, it frequently produces vesications and erosions. Boerhave therefore imagines that its internal employment must be unsafe; but experience has discovered, that a decoction of this plant is not only safe, but of great efficacy in scorbutic complaints. For which purpose, a handful of the herb is directed, by Below, to be boiled in eight pints of beer, till they are reduced to four, of which three or four ounces are to be taken every, or every other, morning. Milk has been found to answer this purpose better than beer. Not only ulcers simply scorbutic, but those of a scrophulous, or even cancerous tendency have been cured by the use of this plant; of which Marquet relates several instances. He likewise found it useful as an external application in destroying fungous flesh, and in promoting a discharge in gangrenes and carbuncles. Another effect, for which this plant is esteemed, is that of stopping intermittent fevers.

ILICIUM. The name of a genus of plants in the Linnaean system.

ILICIUM ANISATUM. The systematic name of the plant, the seeds of which are called the star aniseed. See *Anisum stellatum*.

ILLOSTI. (From *ἄλλο*, the eye.) A distortion of the eyes.

ILLUTAMENTUM. An ancient form of an external medicine, like the *Ceroma*, with which the limbs of wrestlers, and others delighting in like exercises, were rubbed, especially after bathing; an account of which may be met with in Baccius de Thermiss.

ILLUTATIO. (From *in*, upon, and *lutum*, mud.) Illutation. A besmearing any part of the body with mud, and renewing it as it grows dry, with a view of heating, drying and discussing. It was chiefly done with the mud found at the bottom of mineral springs.

ILLYS. (From *ἄλλο*, the eye.) A person who squints, or with distorted eyes.

ILRS. (From *ἰλρς*, mud.) The fæces of wine. Also an epithet for sediment in stools which resemble fæces of wine; also the sediment in urine, when it resembles the same.

IMBRICILLITAS OCULORUM. Celsus speaks of the *Nyctalopia* by this name.

IMBIBITIO. (From *imbebo*, to receive into.) In chymistry it is a kind of cohobation, when the liquor ascends and descends upon a solid substance, till it is fix therewith.

IMMEBUS. A term given by Bartholine, and some other anatomists, to the *Subscapularis* muscle, because it was hidden, or, as it were, sunk.

IMPATIENS. (From *in*, not, and *patior*, to suffer; because its leaves recede from the hand with a crackling noise, as impatient of the touch.) A species of persicaria.

IMPERATORIA. (From *impero*, to overcome; so named because, its leaves extend and overwhelm the lesser herbs which grow near it.) 1. The name of a genus of plants in the Linnaean system. Class, *Pentandria*. Order, *Monogynia*.

2. The pharmacopoeial name of the *Imperatoria ostruthium* of Linnaeus. *Magistrantia*. Master-wort. The roots of this plant are imported from the Alps and Pyrenees, notwithstanding it is indigenous to this island: they have a fragrant smell, and a bitterish pungent taste. The plant, as its name imports, was formerly thought to be of singular efficacy; and its great success, it is said, caused it to be distinguished by the name of *divinum remedium*. At present it is considered merely as an aromatic, and consequently is superseded by many of that class which possess superior qualities.

IMPERATORIA OSTRUTHIUM. The systematic name of the master-wort. See *Imperatoria*.

IMPETIGINES. (The plural of *impetigo*; from *impeto*, to infect.) An order in the class *cachexie* of Cullen, the genera of

which are characterized by cachexia, deforming the external parts of the body with tumours, eruptions, &c.

IMPETIGO. This affection, as described by authors, is a disease in which several red, hard, dry, prurient spots arise in the face and neck, and sometimes all over the body, and disappear by furfuraceous or tender scales.

IMPIA HERBA. (From *in*, not, and *pius*, good; because it grows only on barren ground.) A name given to cudweed.

IMPLICATED. Celsus, Scribonius, and some others, call those parts of physic so, which have a necessary dependence on one another: but the term has been more significantly applied, by Bellini, to such fevers, where two at a time afflict a person, either of the same kind, as a double tertian; or of different kinds, as an intermittent tertian, and a quotidian, called a *Semiteritian*.

IMPLUVIUM. (From *impluo*, to shower upon.) The shower-bath. An embrocation.

IMPREGNATION. See *Conception* and *Generation*.

INANITIO. (From *inanio*, to empty.) Inanition. Applied to the body, it means evacuation; applied to the mind, it means a defect of its powers.

INCANTATION. *Incantatio. Incantamentum.* A way of curing diseases by charms, defended by Paracelsus, Helmont, and some other chymical enthusiasts.

INCENDIUM. (From *incendo*, to burn.) A burning fever, or sometimes any burning heat.

INCENSIO. The same as *Incendium*. Also a hot inflammatory tumour.

INCERNICULUM. (From *incerno*, to sift.) A strainer, or sieve. In anatomy, it is a name for the pelvis of the kidney, from its office as a strainer.

INCIDE. (From *incido*, to cut.) Medicines were formerly so called which consist of pointed and sharp particles, as acids, and most salts, which are said to incide or cut the phlegm, when they break it so as to occasion its discharge.

INCIDENTIA. (From *incido*, to cut.) *Alterantia.* Medicines supposed to cut viscid humours.

INCINEBATIO. (From *incinero*, to reduce to ashes.) Incineration. The reducing of any thing to ashes by fire.

INCISIVUS INFERIOR. See *Levator labii inferioris*.

INCISIVUS LATERALIS. See *Levator labii superioris alaeque nasi*.

INCISIVUS MEDIUS. See *Depressor labii superioris alaeque nasi*.

INCISORIUM. (From *incido*, to cut.) A table whereon a patient is laid for an operation.

INCISORIUM FORAMEN. A name of the foramen, which lies behind the dentes incisores of the upper jaw.

INCISORS. (*Dentes incisores*; from *incido*, to cut, from their use in cutting the food.) The four front teeth of both jaws are so called, because they cut the food. See *Teeth*.

INCONTINENTIA. (From *in* and *contineo*, to contain.) Inability to retain the natural evacuations.

INCRASSANTIA. (From *incrasso*, to make thick.) Medicines thickening the fluids.

INCUBUS. (From *incubo*, to lie upon; because the patient fancies that something lies upon his chest.) See *Night-mare*, and *Oncirodynia*.

INCUS. (A smith's anvil, from *incudo*, to smite upon; so named from its likeness in shape to an anvil.) The largest and strongest of the bones of the ear in the tympanum. It is divided into a body and two crura. Its body is situated anteriorly, is rather broad and thick, and has two eminences and two depressions, both covered with cartilage, and intended for the reception of the head of the malleus. Its shorter crus extends no farther than the cells of the mastoid apophysis. Its longer crus, together with the manubrium of the malleus, to which it is connected by a ligament, is of the same extent as the shorter, but its extremity is curved inwards, to receive the os orbiculare, by the intervention of which it is united with the stapes.

INDEX. (From *indico*, to point out; because it is generally used for such purposes.) The fore-finger.

Indian arrow-root. See *Maranta*.

Indian cress. See *Nasturtium indicum*.

INDIAN DATE-PLUM. The fruit of the *Diospyrus lotus* of Linnæus. When ripe it has an agreeable taste, and is very nutritious.

Indian leaf. See *Cassia lignea*.

Indian pink. See *Spigelia*.

INDIAN RUBBER. The substance known by the names Indian rubber, Elastic gum, Cayenne resin, Cautchuc, and by the French Caoutchouc, is prepared from the juice of the *Siphonia elastica*, — *foliis ternatis ellipticis integerrimis subtus canis longe petiolatis*. Suppl. plant. The manner of obtaining this juice is by making incisions through the bark of the lower part of the trunk of the tree, from which the fluid resin issues in great abundance, appearing of a milky whiteness as it flows into the vessel placed to receive it, and into which it is conducted by means of a tube or leaf fixed in the incision, and supported with clay. On exposure to the air, this milky juice gradually inspissates into a soft, reddish, elastic resin. It is formed by the Indians in South America into various figures, but is commonly brought to Europe in that of spear-shaped bottles, which are said to be formed by spreading

the juice of the *Siphonia* over a proper mould of clay; as soon as one layer is dry, another is added, until the bottle be of the thickness desired. It is then exposed to a thick dense smoke, or to a fire, until it becomes so dry as not to stick to the fingers, when, by means of certain instruments of iron, or wood, it is ornamented on the outside with various figures. This being done, it remains only to pick out the mould, which is easily effected by softening it with water. Indian rubber may be subjected to the action of some of the most powerful menstrua, without suffering the least change, while its pliability and elasticity are eminently peculiar to itself. Its proper menstruum is known to some persons in England, who keep it a profound secret, and prepare the gum into beautiful catheters, bougies, syringes, pessaries, &c.

Indian wheat. See *Zea mays*.

INDIANA RADIX. *Ipecacuanha*.

INDICA CAMOTEA. Potatoes.

INDICANS. Indicant. The proximate cause of a disease, or that from which the indication is drawn.

INDICATING DAYS, are the same as critical days.

INDICATION. (*Indicatio*; from *indico*, to shew.) An indication is that which demonstrates in a disease what ought to be done. It is three-fold; preservation, which preserves health; curative, which expels a present disease; and vital, which respects the powers and reasons of diet. The scope from which indications are taken, or determined, is comprehended in this distich:

—*Ars, etas, regio, complexio, virtus,
Mors et symptoma, repletio, tempus et
unus.*

INDICATOR. (From *indico*, to point; so named from its office of extending the index, or fore-finger) *Extensor indicis* of Cowper. *Extensor secundi intermedii indicis*, *proprius vulgo indicator* of Douglas, and *Cubito-rus phalangettien de l'index* of Dumas. An extensor muscle of the fore-finger, situated chiefly on the lower and posterior part of the fore-arm. It arises, by an acute fleshy beginning, from the middle of the posterior part of the ulna; its tendon passes under the same ligament with the extensor digitorum communis, with part of which it is inserted into the posterior part of the fore-finger.

INDICUM LIGNUM. Logwood.

INDICUS. Sweet and bitter costus.

INDICUS MORBUS. The venereal disease.

INDIGENOUS. (*Indigenus*; from *indu*, within, and *gigno*, to beget.) Applied to diseases which are local, or peculiar to any country.

INDURANTIA. (From *induro*, to harden.) Medicines which harden.

INDUSIUM. (From *induo*, to put on.)

A shirt. Also the name of the amnios, from its covering the foetus like a shirt.

INESIS. (From *nam*, to evacuate.) *Inethus.* An evacuation of the humours.

INFECTION. See *Contagion*.

INFERNAL. A name given to a caustic, *lapis infernalis*, from its strong burning property.

INFIBULATIO. (From *infibulo*, to button together.) An impediment to the retraction of the prepuce.

INFLAMMABLE. Chymists distinguish by this term such bodies of the mineral kingdom only as burn with facility, and flame in an increased temperature.

INFLAMMATION. (*Inflammati*; from *inflammo*, to burn.) *Phlogosis*. *Phlegmasia*. A genus of disease in the class *pyrexia*, and order *phlegmasia*, of Cullen.

This disease is characterized by heat, pain, redness, attended with more or less of tumefaction and fever. Inflammation is divided into two species, viz. phlegmonous and erysipelatous.

Besides this division, inflammation is either acute or chronic, local or general, simple, or complicated with diseases.

Phlegmonous inflammation is known by its bright red colour, tension, heat, and a circumscribed, throbbing, painful tumefaction of the part, tending to suppuration. Phlegmon is generally used to denote an inflammatory tumour, situated in the skin, or cellular membrane. When the same disease affects the viscera, it is usually called phlegmonous inflammation.

Erysipelatous inflammation is considered as an inflammation of a dull red colour, vanishing upon pressure, spreading unequally, with a burning pain and tumour scarcely perceptible, ending in vesicles, or desquamation. This species of inflammation admits of a division into erythema, when there is merely an affection of the skin alone, with very little of the whole system, and erysipelas, when there is general affection of the system.

The fever attending erysipelatous inflammation is generally synochus, or typhus, excepting when it affects very vigorous habits, and then it may be synocha.

The fever attending phlegmonous inflammation is almost always synocha. Persons in the prime of life, and in full vigour, with a plethoric habit of body, are most liable to the attacks of phlegmonous inflammation; whereas those advanced in years, and those of a weak habit of body, irritable, and lean, are most apt to be attacked with erysipelatous inflammation.

Phlegmonous inflammation terminates in resolution, suppuration, gangrene, and scirrhus, or induration.

Resolution is known to be about to take place when the symptoms gradually abate suppuration, when the inflammation does not readily yield to proper remedies; the

throbbing increases, the tumour points and is external, and rigors come on. Gangrene is about to take place when the pain abates, the pulse sinks, and cold perspirations come on. Scurhus, or induration, is known by the inflammation continuing a longer time than usual; the tumefaction continues, and a considerable hardness remains. This kind of tumour gives little or no pain, and, when it takes place, it is usually the sequel of inflammation, affecting glandular parts. It sometimes, however, is accompanied with lancinating pains, ulcerates, and becomes cancerous.

Erythematous inflammation terminates in resolution, suppuration, or gangrene. The symptoms of inflammation are accounted for in the following way.

The redness arises from the dilatation of the small vessels, which become sufficiently large to admit the red globules in large quantities; it appears also to occur, in some cases, from the generation of new vessels. The swelling is caused by the dilatation of the vessels, the phlethoric state of the arteries and veins, the exudation of coagulable lymph into the intestices of the cellular membrane, and the interruption of absorption.

In regard to the augmentation of heat, as the thermometer denotes, very little increase of temperature; it appears to be accounted for from the increased sensibility of the nerves, which convey false impressions to the sensorium. The pain is occasioned by a deviation from the natural state of the parts, and the unusual condition into which the nerves are thrown. The throbbing depends on the increased action of the arteries.

Blood taken from a person labouring under active inflammation, exhibits a yellowish white crust on the surface; this is denominated the buffy coriaceous, or inflammatory coat. This consists of a layer of coagulable lymph, almost destitute of red globules. Blood, in this state, is often termed *azy*. The colouring part of the blood is its heaviest constituent: and, as the blood of a person labouring under inflammation is longer coagulating than healthy blood, it is supposed that the red globules have an opportunity to descend to a considerable depth from the surface before they become entangled. The buffy coat of blood is generally the best criterion of inflammation; there are a few anomalous constitutions in which this state of blood is always found, but these are rare.

The occasional and exciting causes of inflammation are very numerous; they, however, may generally be classed under external violence, produced either by mechanical or chemical irritation, changes of temperature and stimulating foods. Fever often seems to be a remote cause; the inflammation thus produced is generally considered as critical. Spontaneous inflam-

mation sometimes occurs when no perceptible cause can be assigned for its production. Scrofula and syphilis may be considered as exciting causes of inflammation.

With regard to the proximate cause, it has been the subject of much dispute. Galen considered phlegmon to be produced by a superabundance of the humor sanguineus. Boerhaave referred the proximate cause to an obstruction in the small vessels, occasioned by a viscosity or lenton of the blood. Cullen and others attributed it rather to an affection of the vessels than a change of the fluids.

The proximate cause, at the present period, is generally considered to be a morbid dilatation, and increased action of such arteries as lead and are distributed to the inflamed part.

Inflammation of the brain. See *Phrenitis*.

Inflammation of the bladder. See *Cystitis*.

Inflammation of the eyes. See *Ophthalmia*.

Inflammation of the intestines. See *Enteritis*.

Inflammation of the kidneys. See *Nephritis*.

Inflammation of the liver. See *Hepatitis*.

Inflammation of the lungs. See *Peripneumonia*.

Inflammation of the peritoneum. See *Peritonitis*.

Inflammation of the pleura. See *Pleuritis*.

Inflammation of the stomach. See *Gastritis*.

Inflammation of the testicle. See *hernia humeralis*.

Inflammation of the uterus. See *Hysteritis*.

INFLATIO. (From *inflo*, to puff up.) A windy tumour, or swelling. See *Emphysema*.

INFLATIVA. (From *inflo*, to puff up with wind.) Medicines, or food, which cause flatulence.

INFLUENZA. (The Italian word for influence. The disease is so named because it was supposed to be produced by a peculiar influence of the stars.) See *Catarrhus à contagione*.

INFRASCAPULARIS. (From *infra*, beneath, and *scapula*, the shoulder-blade.) A muscle named from its position beneath the scapula.

INFRASPINATUS. (From *infra*, beneath, and *spina*, the spine.)

INFUNDIBULUM. (From *infundo*, to pour in.) 1. A canal that proceeds from the vulva of the brain to the pituitary gland in the sella turcica.

2. The beginnings of the excretory duct of the kidney, or cavities into which the urine is first received, are called *infundibula*.

INFUSION. (*Infusum*; from *infundo*, to pour in.) *Infusio*. A process that consists in pouring water of any required degree of temperature on such substances as

have a loose texture, as thin bark, wood in shavings, or small pieces, leaves, flowers, &c. and suffering it to stand a certain time. The liquor obtained by the above process is called an *infusion*. The following are among the most approved infusions.

INFUSUM ANTHEMIDIS. Infusion of chamomile. "Take of chamomile flowers, half an ounce; boiling water, a pint." Macerate for ten minutes, in a covered vessel, and strain. For its virtues see *Chamemelum*.

INFUSUM ARMORACIÆ COMPOSITUM. Compound infusion of horse-radish. "Take of fresh horse-radish-root, sliced, mustard-seeds bruised, of each one ounce; boiling water, a pint." Macerate for two hours, in a covered vessel, and strain; then add compound spirit of horse-radish, a fluid ounce. See *Raphanus rusticanus*.

INFUSUM AURANTII COMPOSITUM. Compound infusion of orange-peel. "Take of orange peel, dried, two drachms; lemon-peel, fresh, a drachm; cloves, bruised, half a drachm; boiling water, half a pint." Macerate for a quarter of an hour, in a covered vessel, and strain. See *Aurantium*.

INFUSUM CALUMBÆ. Infusion of calumba. "Take of calumba-root, sliced, a drachm; boiling water, half a pint." Macerate for two hours, in a covered vessel, and strain. See *Columba*.

INFUSUM CARYOPHYLLORUM. Infusion of cloves. "Take of cloves, bruised, a drachm and a half; boiling water, half a pint." Macerate for two hours, in a covered vessel, and strain. See *Caryophyllum*.

INFUSUM CASCARILLÆ. Infusion of cascarilla. "Take of cascarilla-bark, bruised, half an ounce; boiling water, half a pint." Macerate for two hours, in a covered vessel, and strain. See *Cascarilla*.

INFUSUM CATECHU. Infusion of catechu. "Take of extract of catechu, two drachms and a half; cinnamon bark, bruised, half a drachm; boiling water, half a pint." Macerate for an hour, in a covered vessel, and strain. See *Catechu*.

INFUSUM CINCHONÆ. Infusion of cinchona. "Take of lance-leaved cinchona bark, half an ounce; boiling water, half a pint." Macerate for two hours, in a covered vessel, and strain. See *Cinchona*.

INFUSUM CUSPARIÆ. Infusion of cusparia. "Take of cusparia bark, bruised, two drachms; boiling water, half a pint." Macerate for two hours, in a covered vessel, and strain. See *Augustura*.

INFUSUM DIGITALIS. Infusion of fox-glove. "Take of purple fox-glove leaves, dried and powdered, a drachm; boiling water, half a drachm." Macerate for two hours, in a covered vessel, and strain; then add spirit of cinnamon, half a fluid ounce. See *Digitalis*.

INFUSUM GENTIANÆ COMPOSITUM. Compound infusion of gentian. "Take of

gentian-root, sliced, orange-peel, dried, of each a drachm; lemon-peel, fresh, two drachms; boiling water, twelve ounces." Macerate for an hour, in a covered vessel, and strain. See *Gentiana*.

INFUSUM LINI. Infusion of linseed. "Take of linseed, an ounce; liquorice-root, sliced, half an ounce; boiling water, two pints." Macerate for two hours, near the fire, in a covered vessel, and strain. See *Linum*.

INFUSUM QUASSIÆ. Infusion of quassia. "Take of quassia wood, a scruple; boiling water, half a pint." Macerate for two hours, and strain. See *Quassia*.

INFUSUM RHÆI. Infusion of rhubarb. "Take of rhubarb root, sliced, a drachm; boiling water, half a pint." Macerate for two hours, and strain. See *Rhabarbarum*.

INFUSUM ROSÆ. "Take of the petals of red rose, dried, half an ounce; boiling water, two pints and a half; dilute sulphuric acid, three fluid drachms; double-refined sugar, an ounce and a half." Pour the water upon the petals of the rose in a covered glass vessel; then add the acid, and macerate for half an hour. Lastly, strain the infusion and add the sugar to it. See *Rosa*.

INFUSUM SENNÆ. Infusion of senna. "Take of senna-leaves, an ounce and half; ginger-root, sliced, a drachm; boiling water, a pint." Macerate for an hour, in a covered vessel, and strain the liquor. See *Senna*.

INFUSUM SIMAROUBÆ. Infusion of simarouba. "Take of simarouba-bark, bruised, half a drachm; boiling water, half a pint." Macerate for two hours, in a covered vessel, and strain. See *Simarouba*.

INFUSUM TABACI. Infusion of tobacco. "Take of tobacco-leaves, a drachm; boiling water, a pint." Macerate for an hour, in a covered vessel, and strain. See *Nicotiana*.

INGUVIES. The claw, crop, or gorge of a bird. Also gluttony.

INGRAVIDATION. (From *ingravidor*, to be great with child.) The same as impregnation, or going with child.

INGUEN. The groin. The lower and lateral part of the abdomen, above the thigh.

Inguinal ligament. See *Poupart's ligament*.

Inguinal hernia. See *Hernia*.

INUMATION. (From *inhume*, to put into the ground.) The burying a patient in warm or medicated earth. Some chymists have fancied thus to call that kind of digestion, which is performed by burying the materials in dung, or in the earth.

INION. (From *ι*, a nerve; as being the place where nerves originate.) The occiput. Blancard says it is the beginning of the spinal marrow: others say it is the back part of the neck.

INJACULATIO. (From *injaculo*, to

shoot into.) So Helmont calls a disorder which consists of a violent spasmodic pain in the stomach, and an immobility of the body.

INJECTION. (From *injicio*, to cast into.) A medicated liquor, to throw into a natural or preternatural cavity of the body by means of a syringe.

INNOMINATA ARTERIA. The first branch given off by the arch of the aorta. It soon divides into the right carotid and right subclavian arteries.

INNOMINATUM OS. (*Innominatus*; from *in*, priv. and *nomen*, a name; so called because the three bones of which it originally was formed grew together, and formed one complete bone, which was then left nameless.) A large irregular bone, situated at the side of the pelvis. It is divided into three portions, viz. the iliac, ischiatic, and pubic, which are usually described as three distinct bones.

The os ilium, or haunch bone, is of a very irregular shape. The lower part of it is thick and narrow; its superior portion is broad and thin, terminating in a ridge, called the *spine* of the ilium, and more commonly known by the name of the *haunch*. This spine rises up like an arch, being turned somewhat outward, and from this appearance, the upper part of the pelvis, when viewed together, has not been improperly compared to the wings of a phaeton. This spine, in the recent subject, appears as if tipped with cartilage; but this appearance is nothing more than the tendinous fibres of the muscles that are inserted into it. Externally, this bone is unequally prominent, and hollowed for the attachment of muscles; and internally, at its broadest fore-part, it is smooth and concave. At its lower part, there is a considerable ridge on its inner surface. This ridge, which extends from the os sacrum, and corresponds with a similar prominence, both on that bone and the ischium, forms, with the inner part of the ossa pubis, what is called the brim of the pelvis. The whole of the internal surface, behind this ridge, is very unequal. The os ilium has likewise a smaller surface posteriorly, by which it is articulated to the sides of the os sacrum. This surface has, by some, been compared to the human ear, and by others, to the head of a bird; but neither of these comparisons seem to convey any just idea of its form or appearance. Its upper part is rough and porous; lower down it is more solid. It is firmly united to the os sacrum by a cartilaginous substance, and likewise by very strong ligamentous fibres, which are extended to that bone from the whole circumference of this irregular surface. The spine of this bone, which is originally an epiphysis, has two considerable tuberosities, one anteriorly, and the other posteriorly, which is the

largest of the two. The ends of this spine too, from their projecting more than the parts of the bone below them, are called spinal processes. Before the anterior spinal process the spine is hollowed, where part of the Sartorius muscle is placed; and below the posterior spinal process there is a very large niche in the bone, which, in the recent subject, has a strong ligament stretched over its lower part, from the os sacrum to the sharp-pointed process of the ischium; so that a great hole is formed, through which pass the great sciatic nerve and the posterior crural vessels under the pyriform muscle, part of which is likewise lodged in this hole. The lowest, thickest, and narrowest part of the ilium, in conjunction with the other two portions of each os innominatum, helps to form the acetabulum for the os femoris.

The os ischium, or hip-bone, which is the lowest part of the three portions of each os innominatum, is of a very irregular figure, and usually divided into its body, tuberosity, and ramus. The body, externally, forms the inferior portion of the acetabulum, and sends a sharp-pointed process backwards, called the spine of the ischium. This is the process to which the ligament is attached, which was just now described as forming a great foramen for the passage of the sciatic nerve. The tuberosity is large and irregular, and is placed at the inferior part of the bone, giving origin to several muscles. In the recent subject it seems covered with a cartilaginous crust; but this appearance, as in the spine of the ilium, is nothing more than the tendinous fibres of the muscles that are inserted into it. This tuberosity, which is the lowest portion of the trunk, supports us when we sit. Between the spine and the tuberosity is observed a sinuosity, covered with a cartilaginous crust, which serves as a pulley, on which the obturator muscle plays. From the tuberosity, the bone, becoming narrower and thinner, forms the ramus, or branch, which, passing forwards and upwards, makes, with the ramus of the os pubis, a large hole of an oval shape, the *foramen magnum ischii*, which affords, through its whole circumference, attachment to muscles. This foramen is more particularly noticed in describing the os pubis.

The os pubis, or share-bone, which is the smallest of the three portions of the os innominatum, is placed at the upper and fore part of the pelvis, where the two ossa pubis meet, and are united to each other by means of a very strong cartilage, which constitutes what is called the *symphysis pubis*. Each os pubis may be divided into its body angle, and ramus. The body, which is the outer part, is joined to the os ilium. The angle comes forward to form the symphysis, and the ramus is a thin

apophysis, which, uniting with the ramus of the ischium, forms the *foramen magnum ischii*, or *thyroideum*, as it has been sometimes called, from its resemblance to a door, or shield. This foramen is somewhat wider above than below, and its greatest diameter is, from above downwards, and obliquely from within outwards. In the recent subject it is almost completely closed by a strong fibrous membrane, called the *obturator ligament*. Upwards and outwards, where we observe a niche in the bone, the fibres of this ligament are separated, to allow a passage to the posterior crural nerve, an artery, and vein. The great uses of this foramen seem to be to lighten the bones of the pelvis, and to afford a convenient lodgment to the obturator muscles. The three bones now described as constituting the *os innominatum* on each side, all concur to form the great acetabulum, or cotyloid cavity, which receives the head of the thigh-bone; the *os ilium* and *os ischium* making each about two-fifths, and the *os pubis* one-fifth of the cavity. This acetabulum, which is of considerable depth, is of a spherical shape. Its brims are high, and, in the recent subject, is tipped with cartilage. These brims, however, are higher above and externally than they are internally and below, where we observe a niche in the bone (which is the ischium), across which is stretched a ligament, forming a hole for the transmission of blood-vessels and nerves to the cavity of the joint. The cartilage, which lines the acetabulum, is thickest at its circumference, and thinner within, where a little hole is to be observed, in which are placed the apparatus that serves to lubricate the joint, and facilitate its motions. We are likewise able to discover the impression made by the internal ligament of the *os femoris*, which, by being attached both to this cavity and to the head of the *os femoris*, helps to secure the latter in the acetabulum. The bones of the pelvis serve to support the spine and upper parts of the body, to lodge the intestines, urinary bladder, and other viscera; and likewise to unite the trunk to the lower extremities. But besides these uses, they are destined, in the female subject, for other important purposes; and the accoucheur finds, in the study of these bones, the foundation of all midwifery knowledge. Several eminent writers are of opinion that in difficult parturition, all the bones of the pelvis undergo a certain degree of separation. It has been observed likewise, that the cartilage uniting the *ossa pubis* is thicker, and of a more spongy texture, in women than in men, and therefore more likely to swell and enlarge during pregnancy. That many instances of a partial separation of these bones, during labour, have happened, there can be no

doubt; such a separation, however, ought by no means to be considered as an uniform and salutary work of nature, as some writers seem to think, but as the effect of disease. But there is another circumstance, in regard to this part of Osteology, which is well worthy of attention; and this is, the different capacities of the pelvis in the male and female subject. It has already been observed, that the *os sacrum* is shorter and broader in women than in men; the *ossa ilia* are also found more expanded; whence it happens, that in women the centre of gravity does not fall so directly on the upper part of the thigh as in men, and this seems to be the reason why, in general, they step with less firmness, and move their hips forwards in walking. From these circumstances also, the brim of the female pelvis is nearly of an oval shape, being considerably wider from side to side, than from the symphysis pubis to the *os sacrum*; whereas, in man it is rounder, and every where of less diameter. The inferior opening of the pelvis is likewise proportionably larger in the female subject, the *ossa ischia* being more separated from each other, and the *foramen ischii* larger, so that, where the *os ischium* and *os pubis* are united together, they form a greater circle; the *os sacrum* is also more hollowed, though shorter, and the *os coccygis* more loosely connected, and therefore capable of a greater degree of motion than in men.

INNOMINATI NERVI. A name of the fifth pair of nerves.

INOCULATION. The insertion of a poison into any part of the body. It is mostly practised with that of the small-pox, because we have learnt, from experience, that by so doing we shall generally procure fewer pustules, and a much milder disease, than when the small-pox is taken in a natural way. Although the advantages are evident, yet objections have been raised against inoculation, on the notion that it exposes the person to some risk, when he might have passed through life without ever taking the disease naturally; but it is well known that he will be exposed to much greater danger, from the intercourse which he must have with his fellow-creatures, by taking the disorder in a natural way. It has also been adduced, that a person is liable to take the small-pox a second time, when produced at first by artificial means; but such instances are very rare, besides not being sufficiently authentic. We may conjecture that, in most of those cases, the matter used was not variolous, but that of some other eruptive disorder, such as the chicken-pox, which has often been mistaken for the small-pox.

To illustrate the benefits arising from inoculation, it has been calculated that a third of the adults die who take the disease

in a natural way, and about one-seventh of the children; whereas, of those who are inoculated, and are properly treated afterwards, the proportion is probably not greater than one in five or six hundred.

Inoculation is generally thought to have been introduced into Britain from Turkey, by Lady Mary Wortley Montague, about the year 1721, whose son had been inoculated at Constantinople, during her residence there, and whose infant daughter was the first that underwent the operation in this country. It appears to have been well known before this period, both in the South of Wales and Highlands of Scotland. Mr. Mungo Park, in his travels into the interior of Africa, found that inoculation had been long practised by the negroes on the Guinea coast; and nearly in the same manner, and at the same time of life, as in Europe.

It is not clearly ascertained where inoculation really originated. It has been ascribed to the Circassians, who employed it as a mean to preserve the beauty of their women. It appears more probable that accident first suggested the expedient among the different nations, to whom the small-pox had long been known, independent of any intercourse with each other; and what adds to the probability of this conjecture is, that in most places where inoculation can be traced back, for a considerable length of time, it seems to have been practised chiefly by old women, before it was adopted by regular practitioners.

Many physicians held inoculation in the greatest contempt at first, from its supposed origin; others again discredited the fact, while others, on the testimony of the success in distant countries, believed in the advantages it afforded, but still did not think themselves warranted to recommend it to the families they attended; and it was not until the experiment of it had been made on six criminals (all of whom recovered from the disease, and regained their liberty,) that it was practised, in the year 1726, on the royal family, and afterwards adopted as a general thing.

To insure success from inoculation, the following precautions should strictly be attended to.

1. That the person should be of a good habit of body, and free from any disease, apparent or latent, in order that he may not have the disease and a bad constitution, or perhaps another disorder, to struggle with at the same time.

2. To enjoin a temperate diet and proper regimen; and, where the body is plethoric, or gross, to make use of gentle purges, together with mercurial and antimonial medicines.

3. That the age of the person be as little advanced as possible, but not younger, if it can be avoided, than four months.

4. To choose a cool season of the year, and to avoid external heat, either by exposure to the sun, sitting by fires, or in warm chambers, or by going too warmly clothed, or being too much in bed.

5. To take the matter from a young subject, who has the small-pox in a favourable way, and who is otherwise healthy, and free from disease; and, when fresh matter can be procured, to give it the preference.

Where matter of a benign kind cannot be procured, and the patient is evidently in danger of the casual small-pox, we should not, however, hesitate a moment to inoculate from any kind of matter that can be procured, as what has been taken in malignant kinds of small-pox has been found to produce a very mild disease.

The mildness or malignity of the disease appears, therefore, to depend little at all on the inoculating matter. Variolous matter, as well as the vaccine, by being kept for a length of time, particularly in a warm place, is apt, however, to undergo decomposition, by putrefaction; and then another kind of contagious material has been produced.

In inoculating, the operator is to make the slightest puncture or scratch imaginable in the arm of the person, rubbing that part of the lancet which is besmeared with matter repeatedly over it, by way of insuring the absorption; and in order to prevent its being wiped off, the shirt sleeve ought not to be pulled down until the part is perfectly dry.

A singular circumstance attending inoculation is, that when this fails in producing the disease, the inoculated part nevertheless sometimes inflames and suppurates, as in cases where the complaint is about to follow; and the matter produced in those cases is as fit for inoculation as that taken from a person actually labouring under the disease. The same happens very frequently in inoculation for the cow-pox.

If, on the fourth or fifth day after the operation, no redness, or inflammation, is apparent on the edge of the wound, we ought then to inoculate in the other arm, in the same manner as before; or, for greater certainty, we may do it in both.

Some constitutions are incapable of having the disease in any form. Others do not receive the disease at one time, however freely exposed to its contagion, even though repeatedly inoculated, and yet receive it afterwards by merely approaching those labouring under it.

On the coming on of the febrile symptoms, which is generally on the seventh day in the inoculated small-pox, the patient is not to be suffered to lie a-bed, but should be kept cool, and partake freely of antiseptic cooling drinks.

INOSULATION. (From *is*, and *oculum*, a little mouth.) The running of

the veins and arteries into one another, or the interunion of the extremities of arteries and veins.

INSANIA. (From *in*, not, and *sanus*, sound.) Insanity, or deranged imagination. A genus of disease in the class *neuroses*, and order *vesania*, characterized by erroneous judgment, from imaginary perceptions, or recollections, attended with agreeable emotions in persons of a sanguine temperament. See *Mania*.

INSISSUS. (From *insideo*, to sit upon.) A vapour-bath, over which the patient sits.

INSIDIANS. (From *insidiar*, to deceive.) A name for diseases which betray no previous symptoms, but are ready to break out by surprise.

INSIPIENTIA. (From *in*, and *sapientia*, wisdom.) A low degree of delirium, without fever.

INSOLATIO. (From *in*, upon, and *sol*, the sun.) A disease which arises from a too great influence of the sun's heat upon the head.

INSPIRATION. (From *in*, and *spiro*, to breathe.) The act of drawing the air into the lungs. See *Respiration*.

INTERCOSTAL ARTERIES. *Arteria intercostales.* The arteries which run between the ribs. The superior intercostal artery is a branch of the subclavian. The other intercostal arteries are given off from the aorta.

INTERCOSTAL MUSCLES. *Intercostales externi et interni.* Between the ribs, on each side, are eleven double rows of muscles. These are the *intercostales externi* and *interni*. Galen has very properly observed, that they decussate each other like the strokes of the letter X. The *intercostales externi* arise from the lower edge of each superior rib, and, running obliquely downwards and forwards, are inserted into the upper edge of each inferior rib, so as to occupy the intervals of the ribs, from as far back as the spine to their cartilage; but from their cartilages to the sternum, there is only a thin aponeurosis covering the internal intercostals. The *intercostales interni* arise and are inserted in the same manner as the external. They begin at the sternum, and extend as far as the angles of the rib, their fibres running obliquely backwards. These fibres are spread over a considerable part of the inner surface of the ribs, so as to be longer than those of the external intercostals. Some of the posterior portions of the internal intercostals pass over one rib, and are inserted into the rib below. Verheyen first described these portions as separate muscles, under the name of *infra costales*. Winslow has adopted the same name. Cowper, and after him Douglas, calls them *costarum depressores proprii*. These distinctions, however, are altogether superfluous, as they are evidently nothing more than appendages of

the intercostals. The number of these portions varies in different subjects. Most commonly there is only four, the first of which runs from the second rib to the fourth, the second from the third rib to the fifth, the third from the fourth rib to the sixth, and the fourth from the fifth rib to the seventh. The internal intercostals of the two inferior false ribs are frequently so thin, as to be with difficulty separated from the external; and, in some subjects, one or both of them seem to be altogether wanting. It was the opinion of the ancients, that the external intercostals serve to elevate, and the internal to depress the ribs. They were probably led to this opinion, by observing the different direction of their fibres; but it is now well known, that both have the same use, which is that of raising the ribs equally during inspiration. Fallopius was one of the first who ventured to call in question the opinion of Galen on this subject, by contending that both layers of the intercostals serve to elevate the ribs. In this opinion he was followed by Hieronymus Fabricius, our countryman Mayow and Borelli. But, towards the close of the last century, Bayle, a writer of some eminence, and professor at Toulouse, revived the opinion of the ancients by the following arguments. He observed, that the oblique direction of the fibres of the internal intercostals is such, that, in each inferior rib, these fibres are nearer to the vertebrae than they are at their superior extremities, or in the rib immediately above; and that, of course, they must serve to draw the rib downwards, as towards the most fixed point. This plausible doctrine was adopted by several eminent writers, and, amongst others, by Nicholls, Hoadley, and Shreiber; but, above all, by Hamberger, who went so far as to assert, that not only the ribs, but even the sternum, are pulled downwards by these muscles, and constructed a particular instrument to illustrate this doctrine. He pretended, likewise, that the intervals of the ribs are increased by their elevation, and diminished by their depression; but he allowed that, while those parts of the internal intercostals that are placed between the bony part of the ribs pull them downwards, the anterior portions of the muscle, which are situated between the cartilages, concur with the external intercostals in raising them upwards. These opinions gave rise to a warm and interesting controversy, in which Hamberger and Haller were the principal disputants. The former argued chiefly from theory, and the latter from experiments on living animals, which demonstrate the fallacy of Hamberger's arguments, and prove, beyond a doubt, that the internal intercostals perform the same functions as the external.

INTERCOSTAL NERVE. *Nervus in-*

carotidis. Great intercostal nerve. Sympathetic nerve. The great intercostal nerve arises in the cavity of the cranium, from a branch of the sixth and one of the fifth pair, uniting into one trunk, which passes out of the cranium through the carotid canal, and descends by the sides of the bodies of the vertebrae of the neck, thorax, loins, and os sacrum: in its course it receives the small accessory branches from all the thirty pair of spinal nerves. In the neck, it gives off three cervical ganglions, the upper, middle, and lower; from which the cardiac and pulmonary nerves arise. In the thorax, it gives off the splanchnic or anterior intercostal, which perforates the diaphragm, and forms the semilunar ganglions, from which nerves pass to all the abdominal viscera. They also form in the abdomen ten peculiar plexuses, distinguished by the name of the viscus to which they belong, as the celiac, splenic, hepatic, superior, middle, and lower, mesenteric, two renal, and two spermatic plexuses. The posterior intercostal nerve gives accessory branches about the pelvis and ischiatic nerve, and at length terminates.

INTERCOSTAL VEINS. The intercostal veins empty their blood into the *vena azygos*.

INTERCURRENT FEVERS. Those which happen in certain seasons only, are called stationary; but others are called, by Sydenham, *intercurrents*.

INTEROUS. (From *inter*, between, and *cutis*, the skin.) A dropsey between the skin and the flesh. See *Anasarca*.

INTERDENTIUM. (From *inter*, between, and *dens*, a tooth.) The intervals between teeth of the same order.

INTERDITIUM. (From *inter*, between, and *digitus*, a toe, or finger.) A corn betwixt the toes, or wart betwixt the fingers.

INTERFEMINEUM. (From *inter*, between, and *femur*, the thigh.) The perineum, or space between the anus and pudendum.

INTERLUNUS MORBUS. (From *inter*, between, and *luna*, the moon; because it was supposed to affect those who were born in the wane of the moon. The epilepsy.

Intermittent fever. See *Febris intermit-tens*.

INTERNUNTII DIES. (From *internuncio*, to go between.) Applied to critical days, or such as stand between the increase of a disorder and its decrease.

INTEROSSEI MANUS. (*Interosseus musculus*; from *inter*, between, and *os*, the bone.) There are small muscles situated between the metacarpal bone, and extending from the bones of the carpus to the fingers. They are divided into *internal* and *external*; the former are to be seen only on the palm of the hand, but the latter are conspicuous both on the palm and back

of the hand.—The *interossei interni* are three in number. The first, which Albinus names *posterior indicis*, arises tendinous and fleshy from the basis and inner part of the metacarpal bone of the fore-finger, and likewise from the upper part of that which supports the middle-finger. Its tendon passes over the articulation of this part of these bones with the fore-finger, and, uniting with the tendinous expansion that is sent off from the extensor digitorum communis, is inserted into the posterior convex surface of the first phalanx of that finger. The second and third, to which Albinus gives the names of *prior annularis*, and *interosseus annularis*, arise, in the same manner, from the basis of the outsidess of the metacarpal bones that sustain the ring-finger and the little-finger, and are inserted into the outside of the tendinous expansion of the extensor digitorum communis that covers each of those fingers. These three muscles draw the fingers into which they are inserted, towards the thumb. The *interossei externi* are four in number, for among these is included the small muscle that is situated on the outside of the metacarpal bone that supports the fore-finger. Douglas calls it *extensor tertii internodii indicis*, and Winslow *semi interosseus indicis*. Albinus, who describes it among the *interossei*, gives it the name of *prior indicis*. This first interosseus externus arises by two tendinous and fleshy portions. One of these springs from the upper half of the inner side of the first bone of the thumb, and the other from the ligaments that unite the os trapezoides to the metacarpal bone of the fore-finger, and likewise from all the outside of this latter bone. These two portions unite as they descend, and terminate in a tendon, which is inserted into the outside of that part of the tendinous expansion from the extensor digitorum communis that is spread over the posterior convex surface of the fore-finger. The second, to which Albinus gives the name of *prior medii*, is not quite so thick as the last-described muscle. It arises by two heads, one of which springs from the inner side of the metacarpal bone of the fore-finger, chiefly towards its convex surface, and the other arises from the adjacent ligaments, and from the whole outer side of the metacarpal bone that sustains the middle-finger. These two portions unite as they descend, and terminate in a tendon, which is inserted, in the same manner as the preceding muscle, into the outside of the tendinous expansion that covers the posterior part of the middle finger. The third belongs likewise to the middle-finger, and is therefore named *posterior medii* by Albinus. It arises, like the last-described muscle, by two origins, which spring from the roots of the metacarpal bones of the ring and middle fingers,

and from the adjacent ligaments, and is inserted into the inside of the same tendinous expansion as the preceding muscle. The fourth, to which Albinus gives the name of *posterior annularis*, differs from the two last only in its situation, which is between the metacarpal bones of the ring and little fingers. It is inserted into the inside of the tendinous expansion of the extensor digitorum communis, that covers the posterior part of the ring-finger. All these four muscles serve to extend the fingers into which they are inserted, and likewise to draw them inwards, towards the thumb, except the third, or *posterior medii*, which, from its situation and insertion, is calculated to pull the middle finger outwards.

INTEROSSEI PEDIS. These small muscles, in their situation between the metatarsal bones, resemble the interossei of the hand, and, like them, are divided into *internal* and *external*. The *interossei pedis interni* are three in number. They arise tendinous and fleshy, from the basis and inside of the metatarsal bones of the middle, the third, and the little toes, in the same manner as those of the hand, and they each terminate in a tendon that runs to the inside of the first joint of these toes, and from thence to their upper surface, where it loses itself in the tendinous expansion that is sent off from the extensors. Each of these three muscles serves to draw the toe into which it is inserted towards the great-toe. The *interossei externi* are four in number. The first arises tendinous and fleshy from the outside of the root of the metatarsal bone of the great-toe, from the os cuneiforme internum, and from the root of the inside of the metatarsal bone of the fore-toe. Its tendon is inserted into the inside of the tendinous expansion that covers the back part of the toes. The second is placed in a similar manner between the metatarsal bones of the fore and middle toes, and is inserted into the outside of the tendinous expansion on the back part of the fore-toe. The third and fourth are placed between the two next metatarsal bones, and are inserted into the outside of the middle and third toes. The first of these muscles draws the fore-toe inwards towards the great-toe. The three others pull the toes, into which they are inserted, outwards. They all assist in extending the toes.

INTERPELLATUS MORBUS. (From *inpello*, to interrupt.) In Paracelsus it is a disease attended with irregular or uncertain paroxysms.

INTERPOLATUS DIES. (From *interpolo*, to renew.) In Paracelsus, these are the days interpolated betwixt two paroxysms.

INTERSCAPULIUM. (From *inter*, be-

tween, and *scapula*, the shoulder-blade.) That part of the spine which lies between the shoulders.

INTERSEPTUM. (From *inter*, between, and *septum*, an inclosure.) The uvula and the septum narium.

INTERSPINALES COLLI. (*Interspinales musculi*; from *inter*, between, and *spina*, the spine.) The fleshy portions between the spinous processes of the neck, that draw these processes nearer to each other.

INTERSPINALES DORSI ET LUMBORUM. These are rather small tendons than muscles, that connect the spinal and transverse processes.

INTERTRANSVERSALES LUMBORUM. Four distinct small bundles of flesh, which fill up the spaces between the transverse processes of the vertebrae of the loins, and serve to draw them towards each other.

INTERTRIGO. (From *inter*, between, and *tero*, to rub.) An excoriation about the anus, groins, axilla, or other parts of the body, attended with inflammation and moisture. It is most commonly produced by the irritation of the urine, from riding, or some acrimony in children.

INTESTINES. (*Intestina*; from *intus*, within.) The convoluted membranous tube that extends from the stomach to the anus; receives the ingested food; retains it a certain time; mixes with it the bile and pancreatic juices; propels the chyle into the lacteals, and covers the faces with mucus; is so called. The intestines are situated in the cavity of the abdomen, and are divided into the small and large, which have, besides their size, other circumstances of distinction.

The *small* intestines are supplied internally with folds, called *valvulae conniventes*, and have no bands on their external surface. The *large* intestines have no folds internally, and are supplied externally with three strong muscular bands, which run parallel upon the surface, and give the intestines a saccated appearance; and they have also small fatty appendages, called *appendiculae epiploicae*.

The first portion of the intestinal tube, for about the extent of twelve fingers' breadth, is called the *duodenum*; it lies in the epigastric region; makes three turnings, and between the first and second flexure receives, by a common opening, the pancreatic duct, and the ductus communis choledochus. It is in this portion of the intestines that chylification is chiefly performed. The remaining portion of the small intestines is distinguished by an imaginary division into the jejunum and ileum.

The *jejunum*, which commences where the duodenum ends, is situated in the umbilical region, and is mostly found empty;

hence its name; it is every where covered with red vessels, and, about an hour and an half after a meal, with lacteals.

The *Ileum* occupies the hypogastric region and the pelvis; is of a more pallid colour than the former, and terminates by a transverse opening into the large intestines, which is called the *valve of the ileum*, *valve of the cæcum*, or the *valve of Tulpus*.

The beginning of the large intestines is firmly tied down in the right iliac region, and for the extent of about four fingers' breadth is called the *cæcum*, having adhering to it a worm-like process, called the *processus cæci vermiformis*, or *appendicula cæci vermiformis*. The great intestine then commences *colon*, ascends towards the liver, passes across the abdomen, under the stomach, to the left side, where it is contorted like the letter S, and descends to the pelvis: hence it is divided in this course into the *ascending portion*, the *transverse arch*, and the *sigmoid flexure*. When it has reached the pelvis, it is called the *rectum*, from whence it proceeds in a straight line to the anus.

The intestinal canal is composed of three membranes, or coats; a common one from the peritoneum, a *muscular coat*, and a *villosa coat*, the villi being formed of the fine terminations of arteries and nerves, and the origins of lacteals and lymphatics. The intestines are connected to the body by the mesentery; the duodenum has also a peculiar connecting cellular substance, as has likewise the colon and rectum, by whose means the former is firmly accreted to the back, the colon to the kidneys, and the latter to the os coccygis, and in women to the vagina. The remaining portion of the tube is loose in the cavity of the abdomen. The arteries of this canal are branches of the *superior and inferior mesenteric*, and the *duodenal*. The veins evacuate their blood into the *vena portæ*. The nerves are branches of the eighth pair and intercostals. The *lacteal vessels*, which originate principally from the jejunum, proceed to the glands in the mesentery.

INTRICATUS (From *intrico*, to entangle; so called from its intricate folds.) A muscle of the ear.

INTUSSECI (From *intra*, within, and *secus*, towards.) Painful disorders of the internal parts.

INTROCESSIO (From *introcedo*, to go in.) *Depressio*. A depression or sinking of any part inward's.

INTUS-SUSCEPTION. (*Intus-susceptio* and *intro-susceptio*; from *intus*, within, and *susceptio*, to receive.) A disease of the intestinal tube, and most frequently of the small intestines; it consists in a portion of gut passing for some length within another portion.

INTYBUS. (From *in*, and *tuba*, a hollow

instrument, so named from the hollowness of its stalk.) See *Endytia*.

INULA. (Contracted or corrupted from *helenium*, *ἡλενιον*, fabled to have sprung from the tears of Helen.) 1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia superflua*.

2. The herb elecampane.

Inula, common. See *Enula campana*.

INULA DYSENTERICA. The systematic name of the lesser inula. See *Conyza media*.

INULA HELENIUM. The systematic name of the elecampane. See *Enula campana*.

INUSTION. (From *in*, and *uro*, to burn.) Is sometimes used for hot and dry seasons; but most commonly by surgeons for the operation of the cautery.

INVERECUNDUM OS. (From *in*, not, and *verecundus*, modest.) A name of the os frontis, from its being regarded as the seat of impudence.

Inversion of the uterus. See *Uterus, inversion of*.

INVOLUCRUM. (From *in*, and *volvo*, to wrap up; because parts are enclosed by it.) A name of the pericardium; also a name of the membrane which covers some of the viscera.

IODES. (From *ios*, brass.) Verdigrise. Green matter thrown off by vomiting.

IOXIS. (From *iox*, a violet.) A carbuncle of a violet colour.

IOXTHUS. (From *iox*, a violet, and *anthos*, a flower.) A hard pimple in the face, of a violet colour.

LOTACISMUS. (From *iota*, the Greek letter i.) A defect in the tongue, or organs of speech, which renders a person incapable of pronouncing his letters.

LOUI. A restorative alimentary liquor, prepared in Japan. It is made from the gravy of half-roasted beef; but, as to the rest, it is kept a secret.

IPECACUANHA. (Indian.) Ipecacuan. The plant from which this valuable root is obtained was long unknown; it was said by some writers to be the *Psychotria emetica*; class *Pentandria*; order *Monogynia*; by others, the *Viola ipecacuanha*, a syngenesious plant of the order *Monogynia*. It is now ascertained to be neither, but a small plant called *Callicocca ipecacuanha*. There are three sorts of ipecacuan to be met with in our shops, viz. the ash-coloured or grey, the brown, and the white.

The ash-coloured is brought from Peru, and is a small wrinkled root, bent and contorted into a great variety of figures, brought over in short pieces, full of wrinkled and deep circular fissures, down to a small white woody fibre that runs in the middle of each piece: the cortical part is compact, brittle, looks smooth, and re-

sinous upon breaking: it has very little smell; the taste is bitterish and subacid, covering the tongue, as it were, with a kind of mucilage.

The brown is small, somewhat more wrinkled than the foregoing; of a brown or blackish colour without, and white within: this is brought from Brazil.

The white sort is woody, and has no wrinkles, nor any perceptible bitterness in taste. The first, the ash-coloured or grey ipecacuan, is that usually preferred for medicinal use. The brown has been sometimes observed, even in a small dose, to produce violent effects. The white, though taken in a large one, has scarce any effect at all. Experience has proved that this medicine is the safest emetic with which we are acquainted, having this peculiar advantage—that, if it does not operate by vomit, it readily passes off by the other emunctories. Ipecacuan was first introduced as an infallible remedy against dysenteries and other inveterate fluxes, as diarrhoea, menorrhagia, leucorrhoea, &c. and also in disorders proceeding from obstructions of long standing; nor has it lost much of its reputation by time: its utility in these cases is thought to depend upon its restoring perspiration. It has also been successfully employed in spasmodic asthma, catarrhal and consumptive cases. Nevertheless, its chief use is as a vomit, and in small doses, joined with opium, as a diaphoretic. The official preparations are the *pulvis ipecacuanhe compositus*, and the *vinum ipecacuanhe*.

IQUETATA. The inhabitants of the Brazil give this name to the *Scrophularia aquatica*, which is there celebrated as a corrector of the ill flavour of senna.

IRACUNDUS. (From *ira*, anger; so called because it forms the angry look.) A muscle of the eye.

IRIS. (A rainbow; so called because of the variety of its colours.) 1. The anterior portion of the choroid membrane of the eye, which is perforated in the middle by the pupil. It is of various colours. The posterior surface of the iris is termed the *uvula*.

2. The *flower-de-luce* is also called iris, from the resemblance of its flowers to the rainbow.

3. The name of a genus of plants in the Linnæan system. Class, *Triandria*. Order, *Monogynia*.

IRIS FLORENTINA. Florentine orris, or iris. The root of this plant, *Iris florentina* of Linnæus:—*corollis barbatis, caule foliis altiore subbiflora, floribus sessilibus*; which is indigenous to Italy, in its recent state is extremely acrid, and, when chewed, excites a pungent heat in the mouth, that continues several hours; on being dried, this acrimony is almost wholly dissipated;

the taste is slightly bitter, and the smell agreeable, and approaching to that of violets. The fresh root is cathartic, and for this purpose has been employed in dropsies. It is now chiefly used in its dried state, and ranked as a pectoral and expectorant, and hence has a place in the *trechisci amyli* of the pharmacopœias.

Iris, Florentina. See *Iris Florentina*.

IRIS GERMANICA. The systematic name of the flower-de-luce. See *Iris nostras*.

IRIS NOSTRAS. Common iris, or orris. Flower-de-luce. This plant is the *Iris germanica* of Linnæus:—*corollis barbatis, caule foliis altiori multiflora, floribus inferioribus pedunculatis*. The fresh roots have a strong disagreeable smell, and an acrid nauseous taste. They are powerfully cathartic, and are given in dropsical diseases, where such remedies are indicated.

IRIS PALUSTRIS. *Gladiolus luteus*; *Acorus vulgaris*. Yellow water-flag. This indigenous plant, *Iris pseudacorus*:—*imbricis, foliis ensiformibus, petalis alternis, stigmatibus minoribus*, is common in marshes, and on the banks of rivers. It formerly had a place in the London Pharmacopœia, under the name of *gladiolus luteus*. The root is without smell, but has an acrid styptic taste, and its juice, on being snuffed up the nostrils, produces a burning heat in the nose and mouth, accompanied by a copious discharge from these organs; hence it is recommended both as an errhine and sialagogue. Given internally, when perfectly dry, its astringent qualities are such as to cure diarrhoeas. The expressed juice is likewise said to be an useful application to seriginous eruptions and scrofulous tumours.

IRIS PSEUDACORUS. The systematic name of the yellow water-flag. See *Iris Palustris*.

Irish slate. See *Lapis Hybernicus*.

IRON. *Ferrum*. Of all the metals, there is none which is so copiously and so variously dispersed through nature as iron. In animals, in vegetables, and in all parts of the mineral kingdom, we detect its presence. Mineralogists are not agreed with respect to the existence of native iron, though immense masses of it have been discovered, which could not have been the products of art; but there is much in favour of the notion that these specimens have been extracted by subterraneous fire. A mass of native iron, of 1600 pounds weight, was found by Pallas, on the river Denisei, in Siberia; and another mass of 300 pounds was found in Paraguay, of which specimens have been distributed every where. A piece of native iron, of two pounds weight, has been also met with at Kamadorf, in the territories of Neustadt, which is still preserved there. These masses evidently did not

originate in the places where they were found.

There are a vast variety of iron ores; they may, however, be all arranged under the following genera; namely, sulphurets, carburets, oxyds, and salts of iron. The sulphurets of iron from the ores, called *pyrites*, of which there are many varieties. Their colour is, in general, a straw-yellow, with a metallic lustre. They are often amorphous, and often also crystallized. Iron ores of this kind are known by the name of *mundiok*. Iron, in the state of a carburet, forms the *graphite* of Werner, (*plumbago*.) This mineral occurs in kidney-form lumps of various sizes. Its colour is a dark iron-grey, or brownish black; when cut, blueish grey. It has a metallic lustre. Its texture is fine grained. It is very brittle. The combination of iron with oxygen is very abundant. The common magnetic iron stone, *magnetical pyrites*, or *load-stone*, belongs to this class: as does *specular iron ore*, and all the different ores called *hematites*, or *blood-stone*. Iron, combined with silex, constitutes *emery*. Iron, united to carbonic acid, exists in the *sparry iron ore*. Joined to arsenic acid it exists in the ores called *arseniate of iron*, and *arseniate of iron and copper*.

Properties of iron.—Iron is distinguished from every other metal by its magnetical properties. It is attracted by the magnet, and acquires, under various conditions, the property of magnetism. Pure iron is of a whitish grey, or rather blueish colour, very slightly livid; but when polished, it has a great deal of brilliancy. Its texture is either fibrous, fine grained, or in dense plates. Its specific gravity varies from 7.6 to 7.8. It is the hardest and most elastic of all the metals. It is extremely ductile, and may therefore be drawn into wire as fine as a human hair; it is also more tenacious than any other metal, and consequently yields with equal facility to pressure. It is extremely infusible, and, when not in contact with the fuel, it cannot be melted by the heat which any furnace can excite; it is, however softened by heat, still preserving its ductility; this constitutes the valuable property of *welding*. It is very dilatable by heat. It is the only metal which takes fire by the collision of flint. Heated by the contact of air it becomes oxydated. If intensely and briskly heated, it takes fire with scintillation, and becomes a black oxyd. It combines with carbon, and forms what is called steel. It combines with phosphorus in a direct and in an indirect manner, and unites with sulphur readily, by mixture in the cold with water, and by fusion. It decomposes water in the cold slowly, but rapidly when ignited. It decomposes most of the metallic oxyds. All acids act upon iron. Very concentrated sulphuric acid

has little or no effect upon it, but when diluted it oxydates it rapidly. The nitric acid oxydates it with great vehemence. Murate of ammonia is decomposed by it. Nitrate of potash detonates very vigorously with it. Iron is likewise dissolved by alkaline sulphurets. It is capable of combining with a number of metals. It does not unite with lead or bismuth, and very feebly with mercury. It detonates by percussion with the oxygenated muriates.

Method of obtaining iron.—The general process, by which iron is extracted from its ores, is first to roast them by a strong heat, to expel the sulphur, carbonic acid, and other mineralizers, which can be separated by heat. The remaining ore, being reduced to small pieces, is mixed with charcoal, or coke; and is then exposed to an intense heat, in a close furnace, excited by bellows; the oxygen then combines with the carbon, forming carbonic acid gas during the process, and the oxyd is reduced to its metallic state. There are likewise some fluxes necessary, in order to facilitate the separation of the melted metal. The matrix of the iron ore is generally either argillaceous or calcareous, or sometimes a portion of silicious earth; but whichever of these earths is present, the addition of one or both of the others makes a proper flux. These are therefore added in due proportion, according to the nature of the ores; and this mixture, in contact with the fuel, is exposed to a heat sufficient to reduce the oxyd to its metallic state.

The metal thus obtained, and called smelted, pig, or cast, iron, is far from being pure, always retaining a considerable quantity of carbon and oxygen, as well as several heterogeneous ingredients. According as one or other of these predominates, the property of the metal differs. Where the oxygen is present in a large proportion, the colour of the iron is whitish grey, it is extremely brittle, and its fracture exhibits an appearance of crystallization: where the carbon exceeds, it is of a dark grey, inclining to blue, or black, and is less brittle. The former is the *white*, the latter the *black*, *crude iron of commerce*. The *grey* is intermediate to both. In many of these states, the iron is much more fusible than when pure; hence it can be fused and cast into any form; and when suffered to cool slowly, it crystallizes in octahedra; it is also much more brittle, and cannot therefore be either flattened under the hammer, or by the laminating rollers.

To obtain the iron more pure, or to free it from the carbon with which it is combined in this state, it must be refined, by subjecting it to the operations of melting and forging. By the former, in which the metal is kept in fusion for some time, and constantly kneaded and stirred, the quan-

tity of carbon and oxygen it contains are combined, and the produced carbonic acid gas is expelled: the metal at length becomes viscid and stiff; it is then subjected to the action of a very large hammer, or to the more equal, but less forcible, pressure of large rollers, by which the remaining oxyd of iron, and other impurities, not consumed by the fusion, are pressed out. The iron is now no longer granular nor crystallized in its texture; it is fibrous, soft, ductile, malleable, and totally infusible. It is termed forged, wrought, or bar, iron, and is the metal in a purer state, though far from being absolutely pure.

The general medicinal virtues of iron, and the several preparations of it, are to constringe the fibres, to quicken the circulation, to promote the different secretions in the remoter parts, and at the same time to repress inordinate discharges into the intestinal tube. By the use of chalybeates, the pulse is very sensibly raised; the colour of the face, though before pale, changes to a florid red; the alvine, urinary, and cuticular excretions, are increased. Fetid eructations, and black coloured faces, are marks of their taking due effect.

When given improperly, or to excess, iron produces head-ache, anxiety, heats the body, and often causes hæmorrhages, or even vomiting, pains in the stomach, spasms, and pains of the bowels.

Iron is given in most cases of debility and relaxation. In passive hæmorrhages. In dyspepsia, hysteria, and chlorosis. In most of the cachexiæ; and it has lately been recommended as a specific in cancer. In general debility, produced by disease, or excessive hæmorrhages. Where either a preternatural discharge, or suppression of natural secretions, proceeds from a langour, or sluggishness of the fluids, and weakness of the solids, this metal, by increasing the motion of the former and the strength of the latter, will suppress the flux, or remove the suppression; but where the circulation is already too quick, the solids too tense and rigid, where there is any stricture, or spasmodic contraction of the vessels, iron, and all the preparations of it, will aggravate both diseases. Iron probably has no action on the body when taken into the stomach, unless it be oxydized. But during its oxydization, hydrogen gas is evolved, and accordingly we find that fetid eructations are considered as a proof of the medicine having taken effect. It can only be exhibited internally in the state of filings, which may be given in doses from five to twenty grains. Iron wire is to be preferred for pharmaceutical preparations, both because it is the most convenient form, and because it is the purest iron.

The medicinal preparations of iron now in use are;—

1. Carbonas ferri. See *Ferri carbonas*.
2. Sulphas ferri. See *Ferri sulphas*.
3. Ferrum tartarizatum. See *Ferrum tartarizatum*.
4. Liquor ferri alkalina. See *Liquor ferri alkalini*.
5. Tinctura acetatis ferri. See *Tinctura ferri acetatis*.
6. Tinctura muriatis ferri. See *Tinctura ferri muriatis*.
7. Tinctura ferri ammoniati. See *Tinctura ferri ammoniati*.
8. Vinum ferri. See *Vinum ferri*.
9. Ferrum ammoniatum. See *Ferrum ammoniatum*.
10. Oxidum ferri rubrum. See *Oxidum ferri rubrum*.
11. Oxidum ferri nigrum. See *Oxidum ferri nigrum*.

IRRITABILITY. (From *irrito*, to provoke.) *Vis insita* of Haller. *Vis vitalis* of Goeter. Oscillation of Boerhaave. Tonic power of Stahl. Muscular power of Bell. Inherent power of Cullen. The contractility of muscular fibres, or a property peculiar to muscles, by which they contract upon the application of certain stimuli, without a consciousness of action. This power may be seen in the tremulous contraction of muscles when lacerated, or when entirely separated from the body in operations. Even when the body is dead to all appearance, and the nervous power is gone, this contractile power remains till the organization yields, and begins to be dissolved. It is by this inherent power that a cut muscle contracts, and leaves a gap; that a cut artery shrinks and grows stuff after death. This irritability of muscles is so far independent of nerves, and so little connected with feeling, which is the province of the nerves, that, upon stimulating any muscle by touching it with caustic, or irritating it with a sharp point, or driving the electric spark through it, or exciting with the metallic conductors, as those of silver, or zinc, the muscle instantly contracts, although the nerve of that muscle be tied; although the nerve be cut so as to separate the muscle entirely from all connection with the system; although the muscle be separated from the body; although the creature upon which the experiment is performed may have lost all sense of feeling, and have been long apparently dead. Thus a muscle, cut from the limb, trembles and palpitates a long time after; the heart, separated from the body, contracts when irritated; the bowels, when torn from the body, continue their peristaltic motion, so as to roll upon the table, ceasing to answer to stimuli only when they become stiff and cold; and too often, in the human body, the *vis insita* loses the exciting power of the nerves, and then palsy ensues; or, losing all governance of the nerves, the *vis insita*, acting without the

regulating power, falls into partial or general convulsions. Even in vegetables, as in the sensitive plant, this contractile power lives. Thence comes the distinction between the irritability of muscles and the sensibility of nerves; for the *irritability* of muscles survives the animal, as when it is active after death; survives the life of the part of the feelings of the whole system, as in universal palsy, where the vital motions continue entire and perfect, and where the muscles, though not obedient to the will, are subject to irregular and violent actions; and it survives the connection with the rest of the system, as when animals, very tenacious of life, are cut into parts: but *sensibility*, the property of the nerves, gives the various modifications of sense, as vision, hearing, and the rest; gives also the general sense of pleasure or pain, and makes the system, according to its various conditions, feel vigorous and healthy, or weary and low. And thus the eye feels and the skin feels: but their appointed stimuli produce no motions in these parts; they are sensible, but not irritable. The heart, the intestines, the urinary bladder, and all the muscles of voluntary motion, answer to stimuli with a quick and forcible contraction; and yet they hardly feel the stimuli by which these contractions are produced, or, at least, they do not convey that feeling to the brain. There is no consciousness of present stimulus in those parts which are called into action by the impulse of the nerves, and at the command of the will: so that muscular parts have all the irritability of the system, with but little feeling, and that little owing to the nerves which enter into their substance; while nerves have all the sensibility of the system, but no motion.

The discovery of this singular property belongs to our countrymen Glysson; but Baron Haller must be considered as the first who clearly pointed out its existence, and proved it to be the cause of muscular motion.

The laws of irritability, according to Dr. Crichton, are, 1. After every action in an irritable part, a state of rest, or cessation from motion, must take place before the irritable part can be again incited to action. If, by an act of volition, we throw any of our muscles into action, that action can only be continued for a certain space of time; the muscle becomes relaxed, notwithstanding all our endeavours to the contrary, and remains a certain time in that relaxed state, before it can be again thrown into action. 2. Each irritable part has a certain portion or quantity of the principle of irritability which is natural to it, part of which it loses during action, or from the application of stimuli. 3. By a process wholly unknown to us, it regains

this lost quantity, during its repose, or state of rest. In order to express the different quantities of irritability in any part, we say that it is either more or less redundant, or more or less defective. It becomes redundant in a part when the stimuli which are calculated to act on that part are withdrawn or withheld for a certain length of time, because then no action can take place; while, on the other hand, the application of stimuli causes it to be exhausted, or to be deficient, not only by exciting action, but by some secret influence, the nature of which has not yet been detected; for it is a circumstance extremely deserving of attention, that an irritable part, or body, may be suddenly deprived of its irritability by powerful stimuli, and yet no apparent cause of muscular or vascular action takes place at the time. A certain quantity of spirits, taken at once into the stomach, kills almost as instantaneously as lightning does: the same thing may be observed of some poisons, as opium, distilled laurel-water, the juice of the cerbera alova, &c. 4. Each irritable part has stimuli which are peculiar to it; and which are intended to support its natural action: thus, blood, which is the stimulus proper to the heart and arteries, if, by any accident, it gets into the stomach, produces sickness, or vomiting. If the gall, which is the natural stimulus to the gall-bladder, duct and duodenum, is by any accident effused into the cavity of the peritoneum, it excites too great action of the vessels of that part, and induces inflammation. The urine does not irritate the tender fabric of the kidneys, ureters, or bladder, except in such a degree as to preserve their healthy action; but if it be effused into the cellular membrane, it brings on such a violent action of the vessels of these parts, as to produce gangrene. Such stimuli are called *habitual* stimuli of parts. 5. Each irritable part differs from the rest in regard to the quantity of irritability which it possesses. This law explains to us the reason of the great diversity which we observe in the action of various irritable parts; thus the muscles of voluntary motion can remain a long time in a state of action, and if it be continued as long as possible, another considerable portion of time is required before they regain the irritability they lost; but the heart and arteries have a more short and sudden action, and their state of rest is equally so. The circular muscles of the intestines have also a quick action and short rest. The urinary bladder does not fully regain the irritability it loses during its contraction for a considerable space of time; the vessels which separate and throw out the menstrual discharge act, in general, for three or four days, and do not regain the irrita-

bility they lose for a lunar month. 6. All stimuli produce action in proportion to their irritating powers. As a person approaches his hand to the fire, the action of all the vessels in the skin is increased, and it glows with heat; if the hand be approached still nearer, the action is increased to such an unusual degree as to occasion redness and pain; and if it be continued too long, real inflammation takes place; but if this heat be continued, the part at last loses its irritability, and a sphacelus or gangrene ensues. 7. The action of every stimulus is in an inverse ratio to the frequency of its application. A small quantity of spirits taken into the stomach, increases the action of its muscular coat, and also of its various vessels, so that digestion is thereby facilitated. If the same quantity, however, be taken frequently, it loses its effect. In order to produce the same effect as at first, a larger quantity is necessary: and hence the origin of dram-drinking. 8. The more the irritability of a part is accumulated, the more that part is disposed to be acted upon. It is on this account that the activity of all animals, while in perfect health, is much livelier in the morning than at any other part of the day; for, during the night, the irritability of the whole frame, and especially that of the muscles destined for labour, viz. the muscles for voluntary action, is re-accumulated. The same law explains why digestion goes on more rapidly the first hour after food is swallowed, than at any other time; and it also accounts for the great danger that accrues to a famished person upon first taking in food. 9. If the stimuli which keep up the action of any irritable body be withdrawn for too great a length of time, that process on which the formation of the principle depends is gradually diminished, and at last entirely destroyed. When the irritability of the system is too quickly exhausted by heat, as is the case in certain warm climates, the application of cold invigorates the frame, because cold is a mere diminution of the overplus of that stimulus which was causing the rapid consumption of the principle. Under such or similar circumstances, therefore, cold is a tonic remedy; but if, in a climate naturally cold, a person were to go into a cold bath, and not soon return into a warmer atmosphere, it would destroy life just in the same manner as many poor people who have no comfortable dwellings are often destroyed from being too long exposed to the cold in winter. Upon the first application of cold the irritability is accumulated, and the vascular system therefore is exposed to great action; but, after a certain time, all action is so much diminished, that the process, whatever it be, on which the formation of

the irritable principle depends, is entirely lost. For further information on this interesting subject, see Dr. Crichton on Mental Derangement.

IRRITATION. *Irritatio.* The action produced by any stimulus.

ISCA. A sort of fungous excrescence of the oak, or of the hazel, &c. The ancients used it as the moderns used moxa.

ISCHAMON. (From *σχῆν*, to restrain, and *αἷμα*, blood.) A name for any medicine which restrains or stops bleeding.

ISCHAMUM. A species of *Andropogon*.

ISCHIAS. (*ἰσχίος*: from *ἰσχῶν*, the hip.) *Sciatica.* A rheumatic affection of the hip-joint. See *Rheumatismus*.

ISCHIATOCELE. (From *ἰσχῶν*, the hip, and *κύλη*, a rupture.) *Ischiocele.* An intestinal rupture, through the sciatic ligaments.

ISCHIO-CAVERNOSUS. See *Erector penis*.

ISCHIOCELE. See *Ischiocele*.

ISCHIUM. (From *ἰσχῆ*, the loin; so named because it is near the loin.) A bone of the pelvis of the foetus, and a part of the os innominatum of the adult. See *Innominatum os*.

ISCHNOPHONIA. (From *ἰσχνός*, slender, and *φωνή*, the voice.) A shrillness of the voice; but more frequently an hesitation of speech, or stammering; it is the *psellismus hesitans* of Cullen.

ISCHURICA. (From *ἰσχυρῆς*, a suppression of the urine.) Medicines which relieve a suppression of the urine.

ISCHURIA. (From *ἰσχεῖν*, to restrain, and *ὑπό*, the urine.) A suppression of urine. A genus of disease in the class *locales* and order *epischereses* of Cullen. There are four species of ischuria.

1. *Ischuria renalis*, coming after a disease of the kidneys, with a troublesome sense of weight in that part.

2. *Ischuria ureterica*, a disease of the kidneys, a sense of pain or uneasiness in the course of the ureters.

3. *Ischuria vesicalis*, a frequent desire to make water, with a swelling of the hypogastrium, and pain at the neck of the bladder.

4. *Ischuria urethralis*, a frequent desire to make water, with a swelling of the hypogastrium, and pain of some part of the urethra.

When there is a frequent desire of making water, attended with much difficulty in voiding it, the complaint is called a dysuria, or stranguary; and when there is a total suppression of urine, it is known by the name of an ischuria. Both ischuria and dysuria are distinguished into acute, when arising in consequence of inflammation, and chronic, when proceeding from any other cause, such as calculus, &c.

The causes which give rise to these diseases are, an inflammation of the urethra,

occasionally either by venereal sores, or by a use of acrid injections, tumour or ulcer of the prostate gland, inflammation of the bladder or kidneys, considerable enlargement of the hæmorrhoidal veins, a lodgement of indurated faeces in the rectum, spasm at the neck of the bladder, the absorption of cantharides applied externally, or taken internally, and excess in drinking either spirituous or vinous liquors; but particles of gravel sticking at the neck of the bladder, or lodging in the urethra, and thereby producing irritation, prove the most frequent cause. Gouty matter falling on the neck of the bladder will sometimes occasion these complaints.

In dysury there is a frequent inclination to make water, attended with a smarting pain, heat, and difficulty in voiding it, together with a sense of fulness in the region of the bladder. The symptoms often vary, however, according to the cause which has given rise to it. If it proceeds from a calculus in the kidney, or ureter, besides the affections mentioned, it will be accompanied with nausea, vomiting, and acute pains in the loins and region of the ureter and kidney of the side affected. When a stone in the bladder, or gravel in the urethra, is the cause, an acute pain will be felt at the end of the penis, particularly on voiding the last drops of urine, and a stream of water will either be divided into two, or be discharged in a twisted manner, not unlike a cork-screw. If a scirrhous of the prostate gland has occasioned the suppression or difficulty of urine, a hard indolent tumour, unattended with any acute pain, may readily be felt in the perinæum, or by introducing the finger in ano.

Dysury is seldom attended with much danger, unless, by neglect, it should terminate in a total obstruction. Ischury may always be regarded as a dangerous complaint, when it continues for any length of time, from the great distention and often consequent inflammation which ensue. In those cases, where neither a bougie nor a catheter can be introduced, the event, in all probability, will be fatal, as few

patients will submit to the only other means of drawing off the urine before a considerable degree of inflammation and tendency to gangrene have taken place.

Isinglass. See *Ichthyocolla*.

ISLANDICUS MUSCUS. See *Lichen islandicus*.

ISOCRONOS. (From *ισος*, equal, and *χρονος*, time.) Preserving an equal distance of time between the beats; applied to the pulse.

ISOCRATES. (From *ισος*, equal, and *κρατεις*, wine.) Wine mixed with an equal quantity of water.

ISOBROMUS. (From *ισος*, equal, and *βρομος*, to run.) The same as *Isochronos*.

ISOPYRUM. (From *ισος*, equal, and *πυρ*, fire; so named from its flame-coloured flower.) The herb aquilegia.

ISOTONUS. (From *ισος*, equal, and *τωνος*, extension.) Applied to fevers which are of equal strength during the whole of the paroxysm.

ISSUE. *Fonticulus.* An artificial ulcer, intended as a remedy on certain morbid affections, by producing a discharge of purulent matter from different parts of the body.

ISTHMIUM. (From *ισθμος*, a narrow piece of land between two seas.) The narrow passage between the mouth and gullet: the fauces.

ISTHMUS VENTRICULI. The ridge surrounding the oval fossa, or remains of the foramen ovale, in the right auricle of the human heart.

ITHMOIDES. Falsely, for *Ethmoides*.

ITINERARIUM. (From *itinerio*, to travel.) The catheter; also a staff used in cutting for the stone; it is thus named by Hil-danus.

IVA PECANGA. See *Sarsaparilla*.

Ixy. See *Hedera arborea*.

Ixy, ground. See *Hedera terrestris*.

Ixy-gum. See *Gummi hederae*.

IXIA. (From *ἰξος*, glue.) A name of the carlina, from its viscous juice. Also a preternatural distention of the veins, from *ἰξμα*, to proceed from.

IXINE. See *Carlina gummisifera*.

J.

JACEA. (*Quia prodest hominibus tristitia facientibus*; because it resists sorrow; or from *jacens*, to heal.) The herb pansy, or heart's-ease.

JACINTRUS. See *Hyacinthus*.

Jack-by-the-hedge. See *Alliaria*.

JACOBÆA. (Named because it was dedicated to St. James, or because it was directed to be gathered about the feast of

St. James.) St. James's wort. Ragwort. *Senecio jacobea* of Linnæus. The leaves of this common plant have a roughish, bitter, sub-acrid taste, extremely nauseous. A decoction is said to have been of infinite service in the cure of epidemic camp dysentery.

Jalap. See *Jalapium*.

JALAPA. See *Jalapium*.

JALAPIUM. (From *Chalapa*, or *Xala-*

pa, in New Spain, whence it is brought.) *Mechoacanna nigra*. Jalap. The plant from which this root is obtained is the *Cerovolvulus jalapa*:—caule volubili; foliis ovatis, subcordatis, obtusis, obsolete repandis, subtus vilosis; pedunculis unifloris: a native of South America. In the shops, the root is found both cut into slices and whole, of an oval shape, solid, ponderous, blackish on the outside, but grey within, and marked with several dark veins, by the number of which, and by its hardness, heaviness, and dark colour, the goodness of the root is to be estimated. It has scarcely any smell, and very little taste, but to the tongue, and to the throat, manifests a slight degree of pungency. The medicinal activity of jalap resides principally, if not wholly, in the resin, which, though given in small doses, occasions violent tormina. The root, powdered, is a very common, efficacious, and safe purgative, as daily experience evinces; but, according as it contains more or less resin, its effects must of course vary. In large doses, or when joined with calomel, it is recommended as an anthelmintic and hydrogogue. In the pharmacopœias, this root is ordered in the form of tincture and extract; and the Edinburgh College directs it also in powder, with twice its weight of crystals of tartar.

JALAPA ALBA. See *Mechoacanna*.

Jamaica bark. See *Chinchini caribæa*.

Jamaica pepper. See *Pimenta*.

JAMBELICI SALES. A preparation with sal-ammoniac, some aromatic ingredients, &c. so called from Jamblichus, the inventor of it.

JANITOR. (From *janua*, a gate.) The pylorus, so called from its being, as it were, the door or entrance of the intestines.

Japan earth. See *Catechu*.

JAPONICA TERRA. See *Catechu*.

JARGON. *Terra circona*. *Terra zercomia*. A primitive earth, lately found in the precious stone called jargon, or hyacinth of the island of Ceylon. When calcined, it has a white colour, is exceedingly heavy, and rough to the touch, has no taste, and is insoluble in water.

JASMINUM. *Jasminum*; *ιασμινος*: from *jasmen*, Arab. or from *isr*, a violet, and *isram*, odour, on account of the fine odour of the flowers.) 1. The name of a genus of plants in the Linnæan system. Class, *Dicandria*. Order, *Monogynia*.

2. The pharmacopœial name of the jessamine. The flowers of this beautiful plant, the *Jasminum officinale* of Linnæus, have a very fragrant smell, and a bitter taste. They afford, by distillation, an essential oil, which is much esteemed in Italy to rub paralytic limbs, and in the cure of rheumatic pains.

JASMINUM OFFICINALE. The systematic name of the jessamine-tree. See *Jasminum*.

JATRALIPTES. (From *ιατρος*, a physician, and *λαψα*, to anoint.) One who undertakes to cure distempers by external unction and friction: Galen makes mention of such in his time, particularly one Diotas; and Pliny informs us, that this was first introduced by Prodicus of Selymbria, who was a disciple of Æsculapius.

JATROCHYMICUS. (From *ιατρος*, a physician, and *χυμum*, chymistry.) *Chymiastr*. A chymical physician, who cures by means of chymical medicines.

JATROLIPTICE. (From *ιατρος*, a physician, and *λαψα*, to anoint.) The method of curing diseases by unction and friction.

JATROPHA. The name of a genus of plants in the Linnæan system.

JATROPHA OUCAS. The systematic name of a plant whose seeds resemble the castor-oil seeds. See *Ricinus major*.

JATROPHYSICUS. (From *ιατρος*, a physician, and *φυσικη*, nature.) An epithet bestowed on some writings, which treat of physical subjects with relation to medicines.

JECORARIA. (From *jecur*, the liver; so named from its supposed efficacy in diseases of the liver.) See *Hepatica terrestris*. Also a name given to a vein in the right hand, because it was usually opened in diseases of the liver.

JECORARIA VENA. The hepatic vein.

JECUR. The liver.

JECUR UTERINUM. The *Placenta* is, by some, thus called, from the supposed similitude of its office with that of the liver.

JEJUNUM. (From *jejunus*, empty.) *Jejunum intestinum*. The second portion of the small intestines, so called because it is mostly found empty. See *Intestines*.

JELLY. Modern chymists have given this name to the mucilaginous substance, very soluble in water, and not at all in spirits of wine, that is obtained from all the soft and white parts of animals, such as the membranes, tendons, sponuroses, cartilages, ligaments, and skin, by boiling them in water. If the decoction or jelly be strongly evaporated, it affords a dry, brittle, transparent substance, known by the name of glue.

Jerusalem consilips. See *Pulmonaria maculata*.

Jerusalem oak. See *Botrys vulgaris*.

Jerusalem sage. See *Pulmonaria maculata*.

Jessamine. See *Jasminum*.

JESUITANUS CORTEX. (*Jesuitanus*; from *jessita*, a jesuit.) A name of the Peruvian bark, because it was first introduced into Europe by Father de Lugo, a jesuit. See *Cinchona*.

JESUITICUS CORTEX. See *Cinchona*.

Jesuits bark. See *Cinchona*.

JESUITANUS. (From *jessita*, a jesuit.)

A trivial name of the Peruvian bark, because it was first introduced in Europe by Father de Lugo, a jesuit.

JET. A black bitumen, hard and compact, like certain stones, found in great abundance in various parts of France, Sweden, Germany, and Ireland. It is brilliant and vitreous in its fracture, and capable of taking a good polish by friction: it attracts light substances, and appears to be electric, like amber; hence it has been called *black amber*; it has no smell, but when heated it acquires one like bitumen judaicum.

Jews pitch. See *Bitumen judaicum*.

John's wort. See *Hypericum*.

JONTHI. See *Ionthus*.

JUDICATORIUS. (From *judico*, to discern.) An obsolete term applied to a synocha of four days, because its termination may certainly be foreseen.

JUGALE OS. (*Jugalis*; from *jugum*, a yoke; from its resemblance, or because it is articulated to the bone of the upper jaw, like a yoke.) *Os male.* *Os zygomaticum.* The ossa malarum are the prominent square bones which form the upper part of the cheeks. They are situated close under the eyes, and make part of the orbits. Each of these bones has three surfaces to be considered. One of these is exterior and somewhat convex. The second is superior and concave, serving to form the lower and lateral parts of the orbit. The third, which is posterior, is very unequal and concave, for the lodgment of the lower part of the temporal muscle. Each of these bones may be described as having four processes, formed by their four angles. Two of these may be called *orbital processes*. The superior one is connected with the orbital process of the *os frontis*; and the inferior one with the malar process of the maxillary bone. The third is connected with the temporal process of the sphenoid bone; and the fourth forms a bony arch, by its connection with the zygomatic process of the temporal bone. In infants, these bones are entire and completely ossified.

JUGLANS. (*Quasi Jovis glans*, the royal fruit, from its magnitude.) 1. The name of a genus of plants in the Linnæan system. Class, *Monocotyledon*. Order, *Polyandria*. The walnut-tree.

2. The pharmacopœial name of the walnut. The tree which bears this fruit is the *Juglans regia* of Linnæus:—*foliis ovatis glabris suberratis subaequalibus*. It is a native of Persia, but cultivated in this country. The unripe fruit, which has an astringent bitterish taste, and has been long employed as a pickle, is the part directed for medicinal use by the London College, on account of its anthelmintic virtues. An extract of the green fruit is the most convenient preparation, as it may be kept for a

sufficient length of time, and made agreeable to the stomach of the patient, by mixing it with cinnamon water.

JUGLANS REGIA. The systematic name of the walnut tree. See *Juglans*.

JUGULAR VEINS. (*Vene jugularis*; from *jugulum*, the throat.) These veins run from the head down the sides of the neck, and are divided, from their situation, into external and internal. The *external*, or *superficial jugular vein* receives the blood from the frontal, angular, temporal, auricular, sublingual or ranine, and the occipital veins. The *internal*, or *deep-seated jugular vein*, receives the blood from the lateral sinusses of the dura mater, the laryngeal and pharyngeal veins. Both jugulars unite, and form, with the subclavian vein, the superior vena cava, which terminates in the superior part of the right auricle of the heart.

JUGULUM. (From *jugum*, a yoke; because the yoke is fastened to this part.) The throat, or anterior part of the neck.

JUJUBES. See *Jujuba*.

JUJUBÆ. (*Arab.*) Jujubes. A half-dried fruit of the plumb kind, about the size and shape of an olive, the produce of the *Rhamnus zizyphus* of Linnæus. Jujubes, when in perfection, have an agreeable sweet taste, and in the southern parts of Europe, where they are common, they make an article of food in their recent state, and of medicine when half dried.

July flowers. See *Caryophyllus ruber*.

JUNCUS ODORATUS. *Fenum camelerum.* *Juncus aromaticus.* Camel-hay. Sweet rush. This dried plant, *Andropogon schenanthus* of Linnæus, is imported into this country from Turkey and Arabia. It has an agreeable smell, and a warm, bitterish, not unpleasant taste. It was formerly employed as a stomachic and deobstruent.

Juniper. See *Juniperus*.

Juniper gum. See *Sundarack*.

JUNIPERUS. (*Juniperus*; from *juvenis*, young, and *pario*, to bring forth; so called because it produces its young berries while the old ones are ripening.) 1. The name of a genus of plants in the Linnæan system. Class, *Dioecia*. Order, *Monadelphica*.

2. The pharmacopœial name of the common juniper, *Juniperus communis* of Linnæus:—*foliis ternis patentibus mucronatis, bacis longioribus*. Both the tops and berries of this indigenous plant are directed in our pharmacopœias, but the latter are usually preferred, and are brought chiefly from Holland and Italy. Of their efficacy as a stomachic, carminative, diaphoretic, and diuretic, there are several relations by physicians of great authority; and medical writers have also spoken of the utility of the juniper in nephritic cases, uterine obstructions, scorbutic affections, and some cutaneous diseases. Our pharmacopœias

direct the essential oil, and a spirituous distillation of the berries, to be kept in the shops.

JUNIPERUS COMMUNIS. The systematic name of the juniper-tree. See *Juniperus*.

JUNIPERUS LICIA. The systematic name of the plant which affords the frankincence. See *Olibanum*.

JUNIPERUS SABINA. The systematic name of the savine-tree. See *Sabina*.

JUPITER. The ancient chymical name of tin, because supposed under the government of that planet.

JUVANTIA. (From *juvo*, to assist.) Medicines, or assistances of any kind, which relieve a distemper.

JUXTANGINA. (From *juxta*, near, and *angina*, a quinsy.) A disease resembling a quinsy.

K.

KAMFERIA ROTUNDA. The systematic name of the plant which affords the officinal zedary. *Zedoria*.

KAJEPUT OLEUM. See *Cajeput oil*.

KALI. (Arab.) The *Kali* of the pharmacopœias is the vegetable alkali or potash. See *Potassa*.

KALI ACETATUM. See *Acetas potasse*.

KALI ARSATUM. See *Carbonas potasse*.

KALI ARSENICATUM. A preparation of arsenic, composed of the vegetable alkali and the oxyd of arsenic. It is the active ingredient in Fowler's mineral solution. See *Liquor arsenicalis*.

KALI CITRATUM. *Alkali volatile, succo citri saturatum.* The neutral saline liquor, a citrat of potash, is made by saturated prepared kali with lemon juice. It is the base of the saline draught; it possesses nervine and sudorific properties; and is exhibited in rheumatism, catarrh, and most febrile diseases.

KALI PREPARATUM. See *Subcarbonas potasse*.

KALI PURUM. See *Potassa fusa*.

KALI SULPHURATUM. See *Sulphuretum potasse*.

KALI TARTARIZATUM. See *Turtras potasse*.

KALI VITRIOLATUM. See *Sulphas potasse*.

KEBIRI. See *Chieri*.

KELP. The mineral alkali which is obtained in this country by burning marine plants.

KERATO PHARYNGEUS. (From *keras*, a horn, and *pharynx*, the pharynx.) A muscle so named from its shape, and insertion in the pharynx.

KERMES. (*Chermah*, Arab.) *Gratum tinctorium.* *Coccus baphica.* Round reddish grains, about the size of peas, found in Spain, Italy, and the south of France, adhering to the branches of the scarlet oak. They are the nidus of a minute red animalcule, called *Coccus quercus ilicis* of Linnæus. The *confectio alkermes*, now obsolete, was prepared with these, which possess corroborant and adstringent virtues.

KERMES MINERALIS. A preparation of antimony, so termed from its resemblance to the insect of that name. It is now disused in medicine, and gives place to the other preparations of antimony. See *Hydrosulphuretum stibii rubrum*.

Kernel wort. *Scrophularia vulgaris*.

KERYA. *Kervah*, Arab.) The *ricinus*.

KETCHUP. The prepared liquor of the mushroom.

KEYSER'S PILLS. A once celebrated mercurial medicine, the method of preparing which was purchased by the French government, and has since been published by M. Richard. The hydrargyrus acetatus is considered as an adequate substitute for the more elaborate form of Keyser. M. Richard concludes his account of Keyser's pills with observing, that he considers it to be, without exception, the most effectual remedy for the venereal disease hitherto discovered. But further trials of this remedy do not justify the sanguine accounts of its properties; though it may succeed when some of the other mercurial preparations have failed.

KIDES. A name for chilblains.

KIDNEYS. *Renes.* Two abdominal viscera, shaped like a kidney-bean, that secrete the urine. They are situated on each lumbar region, near the first lumbar vertebra, behind the peritoneum, and are composed of three substances, a cortical, which is the external, and very vascular; a tubulose, which consists of small tubes; and a papillous substance, which is the innermost. The kidneys are generally surrounded with more or less adipose membrane, and they have also a proper membrane, *membrana propria*, which is closely accreted to the cortical substance. The renal arteries, called also emulgents, proceed from the aorta. The veins evacuate their blood into the ascending cava. The absorbents accompany the blood-vessels, and terminate in the thoracic duct. The nerves of the kidneys are branches of the eighth pair and great intercostals. The

excretory duct of this viscus is called the *ureter*. At the middle or pelvis of the kidney, where the blood-vessels enter it, is a large membranous bag, which diminishes like a funnel, and forms a long canal, or ureter, that conveys the urine from the kidney to the bladder, which it perforates obliquely,

KIKEKUNEMALO. A pure resin, very similar to copal, but of a more beautiful whiteness and transparency. It is brought from America, where it is said to be used medicinally in the cure of hystericæ, tetanus, &c. It forms the most beautiful of all varnishes.

KIKI. (*Kike*, Arabian.) See *Ricinus*.

KINA KINA. See *Cinchona*.

KINKINA. See *Cinchona*.

KINO. (Indian.) *Gummi gambiense*.

Gummi rubrum adstringens gambiense. The tree from which this resin is obtained, though not botanically ascertained, is known to grow on the banks of the river Gambia, in Africa. On wounding its bark, the fluid kino immediately issues drop by drop, and, by the heat of the sun, is formed into hard masses. It is very like the resin called *Sanguis draconis*; is much redder, more firm, resinous, and adstringent, than catechu. It is now in common use, and is the most efficacious vegetable adstringent, or styptic, in the materia medica. Its dose is from twenty to thirty grains.

Knee-holly. See *Ruscus*.

Knee pan. See *Patella*.

KOLTO. (Polonese.) The plica polonica, or plaited hair.

KYNACHE. See *Cynanche*.

L.

LABDANUM. See *Ladanum*.

LABIA LEPORINA. (*Leporina*; from *lepus*, a hare, resembling a hare.) The hare-lip.

LABORATORIUM. (From *laboro*, to labour.) A place properly fitted up for the performance of chymical operations.

LABYRINTH. That part of the internal ear behind the cavity of the tympanum; it is constituted by the cochlea, vestibulum, and semicircular canals.

LAC. See *Lacca*.

LAC AMMONIACI. See *Mistura ammoniaci*.

LAC AMYGDALÆ. A very pleasant, cooling, demulcent drink, calculated to alleviate ardor urinæ, and relieve strangury. It forms a pleasant ptisan in coughs, hoarsenesses, and catarrhs. See *Mistura amygdalæ*.

LAC ASSAFŒTIDA. See *Mistura assafœtida*.

LAC SULPHURIS. See *Sulphur præcipitatum*.

LACCA. (From *lakah*, Arab.) *Gummi Lacca*. Stick-lac. Gum-lac. Seed-lac. Shell lac. The improper name of gum-lac is given to a concrete brittle substance, of a dark red colour, brought from the East Indies, incrustated on the twigs of the *Croton lacciferum* of Linnaeus:—*foliis ovatis tomentosis serrulatis petiolatis, corymbis tomentosis*, where it is deposited by a small insect, at present not scientifically known. It is found in very great quantities on the uncultivated mountains on both sides the Ganges; and is of great use to the natives in various works of art, as varnish, painting, dyeing, &c. When the resinous matter is broken off the wood into small pieces

of grains, it is termed *seed-lac*, and when melted and formed into flat plates, *shell-lac*. This substance is chiefly employed for making sealing-wax. A tincture of it is recommended as an antiscorbutic to wash the gums.

LACHRYMA ABIEGNÆ. See *Terebinthina argentatrotensis*.

LACHRYMÆ. The tears. A limpid fluid secreted by the lachrymal gland, and flowing on the surface of the eye.

LACHRYMAL BONE. See *Unguis os*.

LACHRYMAL DUCTS. *Ductus lachrymales*. The excretory ducts of the lachrymal gland, which open upon the internal surface of the upper eye-lid.

LACHRYMAL GLAND. *Glandula lachrymalis*. A glomerate gland, situated above the external angle of the orbit, in a peculiar depression of the frontal bone. It secretes the tears, and conveys them to the eye by its excretory ducts, which are six or eight in number.

LACHRYMAL NERVE. The fifth pair of nerves from the head is divided into several branches, the first of which is called the orbitary branch; this is divided into three more, the third of which is called the *lachrymal branch*; it goes off chiefly to the lachrymal gland.

LACONICUM. (Because they were much used by the people of Laconia.) A stove, or sweating-room.

LACTATION. (From *lacteo*, to suckle.) The giving suck.

LACTATS. (*Lactas*.) Salts formed by the union of the acid of sour whey, or lactic acid, with different bases; thus *aluminos lactat*, *ammoniacal lactat*, &c.

LACTEALS. *Vasa lactea.* The absorbents of the mesentery, which originate in the small intestines, and convey the chyle from thence to the thoracic duct. They are very tender and transparent vessels, possessed of an infinite number of valves, which, when distended with chyle, a milky or lacteal fluid, give them a knotty appearance. They arise from the internal surface of the villous coat of the small intestines, perforate the other coats, and form a kind of net-work, whilst the greater number unite one with another between the muscular and external coats. From thence they proceed between the laminae of the mesentery to the conglobate glands. In their course they constitute the greater part of the gland through which they pass, being distributed through them several times, and curled in various directions. The lacteals, having passed these glands, go to others, and at length seek those nearest the mesentery. From these glands, which are only four or five, or perhaps more, the lacteals pass out and ascend with the mesenteric artery, and unite with the lymphatics of the lower extremities, and those of the abdominal viscera, and then form a common trunk, the *thoracic duct*, which, in some subjects, is dilated at its origin, forming the *receptaculum chyli*. See *Nutrition*.

LACTIC ACID. (From *lac*, milk.) The acid of sour milk.

LACTICA. The Arabian name for that species of fever which the Greeks call *Typhos*, or *Typhodes*.

LACTIFUGA (From *lac*, milk, and *fugo*, to drive away.) Medicines which dispel milk.

LACTUCA. (From *lac*, milk; named from the milky juice which exudes upon its being wounded.) 1. The name of a genus of plants in the Linnaean system. Class, *Syn-genesia*. Order, *Polygamia equalis*.

2. The pharmacopoeial name of the garden-lettuce, the *Lactuca sativa* cultivated; it is esteemed as an wholesome aperient bitter anodyne, easy of digestion, but affording no nutriment. Lettuces appear to agree better with hot, bilious, melancholic temperaments, than the plegmatic. The seeds possess a quantity of oily substance, which, triturated with water, forms an emulsion esteemed by some in aridur, and some diseases of the urinary passage. Lettuce was famous for the cure of the Emperor Augustus, and formed the opiate of Galen, in his old age; a proof that, in the warmer climates, it must acquire an exaltation of its virtues above what is met with in this country.

LACTUCA GRAVEOLENS. Opium scented lettuce. Strong-scented lettuce. *Lactuca sylvestris*. *Lactuca virosa* of Linnaeus; —*foliis horizontalibus carino aculeatis lentatis*. A common plant in our hedges and

ditches. It has a strong ungrateful smell, resembling that of opium, and a bitterish acid taste: it abounds with a milky juice, in which its sensible qualities seem to reside, and which appears to have been noticed by Dioscorides, who describes the odour and taste of the juice as nearly agreeing with that of the white poppy. Its effects are also said, according to Haller, to be powerfully narcotic. Dr. Collin, at Vienna, first brought the *lactuca virosa* into medical repute, and its character has lately induced the College of Physicians at Edinburgh to insert it in the catalogue of the *Materia Medica*. More than twenty-four cases of dropsy are said, by Collin, to have been successfully treated, by employing an extract prepared from the expressed juice of this plant, which is stated not only to be powerfully diuretic, but, by attenuating the viscid humours, to promote all the secretions, and to remove visceral obstructions. In the more simple cases, proceeding from debility, the extract, in doses of eighteen to thirty grains a-day, proved sufficient to accomplish a cure: but when the disease was inveterate, and accompanied with visceral obstructions, the quantity of extract was increased to three drachms; nor did larger doses, though they excited nausea, ever produce any other bad effect; and the patients continued so strong under the use of this remedy, that it was seldom necessary to employ any tonic medicines. Though Dr. Collin began his experiments with the *lactuca* at the Pazman hospital, at the time he was trying the *arnica*, 1771, yet very few physicians, even at Vienna, have since adopted the use of this plant. Plen-ciz, indeed, has published a solitary instance of its efficacy, while Quarin informs us that he never experienced any good effect from its use; alleging, that those who were desirous of supporting its character, mixed with it a quantity of *extractum scillae*. Under these circumstances we shall only say, that the recommendation of this medicine by Dr. Collin, will be scarcely thought sufficient to establish its use in England.

LACTUCA SATIVA. The systematic name of the lettuce. See *Lactuca*.

LACTUCA SYLVESTRIS. *Scariola*. The official name of the *Lactuca scariola* of Linnaeus, which possesses a stronger degree of bitterness than the *Lactuca sativa*, and is said to be more aperient and laxative. It is nearly similar, in virtue as in taste, to endive unblanched.

LACTUCA VIROSA. The systematic name of the opium-scented lettuce. See *Lactuca graveolens*.

LACTUCELLA. (Diminutive of *lactuca*, the lettuce; so named from its milky juice.) The sow-thistle.

LACTUCIMINA. (From *lacteo*, to suckle;

so called because they happen chiefly to children while at the breast.) Aphthæ, or little ulcers, or crusty scabs, on the skin.

LACTUMEN. (From *lac*, milk; so named because it is covered with a white crust.) The achor, or scald-head; also a little crusty scab on the skin, affecting chiefly children at the breast.

LACUNE. (*Lacuna*; from *lacus*, a channel.) The mouths or openings of the excretory ducts of muciparous glands in the urethra, and other parts.

LADANUM. (From *ladon*, Arab.) *Ladanium.* This resinous juice exudes upon the leaves of the *Cistus creticus* of Linnaeus:—*aborescens extipulatus, foliis spatulato-ovatis petiolatis nervosis scabris, calycinis lanceolatis*; in Candia, where inhabitants collect it by lightly rubbing the leaves with leather, and afterwards scraping it off, and forming it into irregular masses for exportation. Three sorts of ladanum have been described by authors, but only two are to be met with in the shops. The best, which is very rare, is in dark-coloured masses, of the consistence of a soft plaster, and growing still softer on being handled; the other is in long rolls, coiled up much harder than the preceding, and not so dark. The first has commonly a small, and the last a large, admixture of fine sand, without which they cannot be collected pure, independently of designed abuses: the dust blown on the plant by winds, from the loose sands among which it grows, being retained by the tenacious juice. The soft kind has an agreeable smell, and a lightly pungent bitterish taste: the hard is much weaker. Ladanum was formerly much employed internally as a pectoral and adstringent in catarrhal affections, dysenteries, and several other diseases; at present, however, it is wholly confined to external use, and is an ingredient in the stomachic plaster, *emplastrum ladanii*.

Ladies bedstraw. See *Galium luteum*.

Ladies mantle. See *Alchemilla*.

Ladies smock. See *Cardamine*.

LETIFICANTIA. (From *letifico*, to make glad.) This term hath been applied to many compositions under the intention of cordials; but both the medicine and distinction are now quite disused.

LAGAROS. (*Λαγρος*, lax; so named from its comparative laxity.) The right ventricle of the heart.

LAGOPHTHALMIA. (From *λαγος*, a hare, and *οφθαλμος*, an eye; because it is believed that hares sleep with their eyes open.) *Logophthalmos.* The hare's eye. A disease in which the eye cannot be shut. The following complaints may arise from it: a constant weeping of the organ, in consequence of the interruption of the alternate closure and opening of the eyelids which motions so materially contri-

tribute to propelling the tears into the nose; blindness in a strong light, in consequence of the inability to moderate the rays which fall on the eye; on the same account, the sight becomes gradually very much weakened; incapacity to sleep where there is any light; irritation, pain, and redness of the eye, from this organ being exposed to the extraneous substances in the atmosphere, without the eyelids having the power of washing them away in the natural manner.

An enlargement or protrusion of the whole eye, or a staphyloma, may obviously produce lagophthalmos. But affections of the upper eyelids are the common causes. Heister says he has seen the complaint originate from a disease of the lower one. Now and then lagophthalmos depends on paralysis of the orbicularis muscle. A cicatrix, after a wound, ulcer, or burn, is the most frequent cause.

LAGOPODIUM. (From *λαγος*, a hare, and *πους*, a foot; so called because it has narrow hairy leaves, like the foot of a hare.) The herb hare's foot trefoil.

LOSOSTOMA. (From *λαγος*, a hare, and *σoma*, the mouth; so called because the upper lip is divided in the middle like that of a hare.) The hare lip.

Lakerweed. See *Hydrophyllum*.

LAMAC. Gum-arabic.

LAMBDA CIMUS. A defect in speech, which consists in an inability to pronounce certain consonants, or is that stammering, or difficulty of speech, called *Psellismus Lallans*, that is, when the letter L is pronounced too liquid, and often in the place of R.

LAMBDOIDAL SUTURE. (*Sutura Lambdoidalis*; from Λ , and *ωδω*, resemblance; because it is shaped like the letter Λ .) Occipital suture. The suture that unites the occipital bone to the two parietal bones.

LAMBITIVUM. (From *lambo*, to lick up.) A linctus or medicine to be licked up.

LAMELLA. (Dim. of *lamina*, a plate of metal.) The thin plates, or gills, of a mushroom.

LAMINA. (From *laeo*, to beat off.) A bone, or membrane, or any substance resembling a thin plate of metal. The lap of the ear.

LANIUM ALBUM. (From *Lanium*, a mountain of Ionia, where it grew, or from *lama*, a ditch, because it usually grows about ditches and neglected places.) *Urtica mortua.* *Galeopsis.* *Archangelica.* Dead nettle. White archangel nettle. Uterine hemorrhages and fluor albus are said to be relieved by infusions of this plant, from whose sensible qualities very little benefit can be expected.

LAMPANA. See *Lapsana*.

LANCETTA. (Dim. of *lancea*, a spear.) A lancet. An instrument used in phlebotomy.

LAONICA CORATIO. A method of curing the gout, by evaporating the morbid matter y topical applications.

LAPACTICA. (From *λαπαζω*, to evacuate.) Purgative medicines.

LAPARA. From (*λαπαζω*, to empty; so named from its concave and empty appearance.) The flank.

LAPAROCELE. (From *λαπαζω*, the flank, and *κελω*, a rupture.) A rupture through the side of the belly.

LAPATHUM. (From *λαπαζω*, to evacuate; so named because it purges gently.) The dock.

LAPATHUM ACETOSUM. See *Acetosa*.

LAPATHUM ACUTUM. See *Oxylapathum*.

LAPATHUM AQUATICUM. See *Hydrolapathum*.

LAPIDES CANCROBUM. See *Cancer*.

LAPIDELLUM. *Lapidellus*. (From *lapis*, a stone.) The name of a kind of spoon, formerly used to take out small stones and fragments from the bladder.

LAPILLI CANCROBUM. Crab's stones, commonly called crab's eyes. See *Cancer*.

LAPIS BEZOAR. See *Bezoar*.

LAPIS CÆRULEUS. See *Lapis lazuli*.

LAPIS CALAMINARIS. See *Calamine*.

LAPIS CALCAREUS. Hard carbonate of lime.

LAPIS CYANUS. See *Lapis lazuli*.

LAPIS HÆMATITES. See *Hæmatites*.

LAPIS HIBERNICUS. *Tegula hybernica*. *Ardesia hibernica*. *Hardesia*. Irish slate. A kind of slate, or very hard stone, found in different parts of Ireland, in a mass of a blueish black colour, which stains the hands. When dried and powdered, it is pale, or of a whitish blue, and by keeping grows black. In the fire it yields a sulphureous gas, and acquires a pale red colour, with additional hardness. It is occasionally powdered by the common people, and taken in spruce beer, against inward bruises.

LAPIS HYSTRICIS. See *Bezoar porcinum*.

LAPIS INFERNALIS. An old name for the caustic potash. See *Potassa fusa*.

LAPIS LAZULI. *Lapis cyanus*. Azure stone. A combination of silice, the blue fluat of lime and sulphat of lime, and iron. This singular mixture forms a stone, of a beautiful opaque blue, which it preserves in a strong heat, and does not suffer any alteration by the contact of air. It was formerly exhibited as a purgative and vomit, and given in epilepsy.

LAPIS MALACENSIS. See *Bezoar porcinum*.

LAPIS PORCINUS. See *Bezoar porcinum*.

LAPIS SIMIÆ. See *Bezoar simiæ*.

LAPPA MAJOR. See *Bardana*.

LAPSANA. (*Λαψαν*, from *Lampsacus*, the town near which it flourished; or from *λαπαζω*, to evacuate; because it was said

to relax the bowels.) *Lampsana*. *Napium*. *Papillaris herba*. Dock-cresses. Nipple-wort. This plant, *Lapsana communis* of Linnæus, is a lactescent bitter, and nearly similar in virtues to the cichory, dandelion, and endive. It has been employed chiefly for external purposes, against wounds and ulcerations, whence the name of nipple-wort and papillaris.

LAQUEUS GUTTURIS. A malignant inflammation of the tonsils, in which the patient appears as if he were suffocated with a noose.

LARBASON. Antimony.

Larch-tree. See *Terebinthina veneta*.

LARD. The English name of hog's fat, when melted down. Hog's lard, *adepe suelle*, forms the base of many unguents, and is often eaten by the poor instead of butter.

LARYNGOTOMY. (From *λαρυγξ*, the larynx, and *τομή*, to cut.) See *Bronchotomy*.

LARYNX. (*Λαρυγξ*, a Greek primitive.) A cartilaginous cavity, situated behind the tongue, in the anterior part of the fauces, and lined with an exquisitely sensible membrane. It is composed of the annular or cricoid cartilage, the scutiform or thyroid, the epiglottis, and two arytenoid cartilages. The superior opening of the larynx is called the *glottis*. The *laryngeal arteries* are branches of the external carotids. The *laryngeal veins* evacuate their blood into the external jugulars. The nerves of the larynx are from the eighth pair. The use of the larynx is to constitute the organ of voice, and to serve also for respiration.

LASCIVUS. (From *lacio*, to ensnare; upon account of its irregular motions.) An epithet used by Paracelsus for the chorea Sancti viti.

LASER. (A term used by the Cyrenians.) The herb laser-wort, or assafœtida.

LASERPITUM LATIFOLIUM. (From *laser*, perhaps from *lazar*, Arab.) The systematic name of the white gentian. See *Gentiana alba*.

LASERPITUM SILER. The systematic name of the heart-wort. See *Scabi*.

LATERAL OPERATION. One mode of cutting for the stone is so called.

LATERAL SINUSSES. The bifurcation and continuation of the longitudinal sinus of the dura mater. They commence about the middle of the tentorium, one passing along each horizontal crucial spine within the tentorium, and round to the foramen lacerum in basi cranii, where the internal jugular vein begins. Their use is to carry the blood from the vein into the internal jugulars, which return it to the heart.

LATEX. (*Latez*, quod in vena terre lateat.) Water, or juice. A term sometimes applied to the blood, as being the spring or source of all the humours.

LATERITIOUS SEDIMENT. (*Lo-*

teritus; from *later*, a brick.) A term applied to the brick-like sediment occasionally deposited in the urine of people afflicted with fever.

LATHYRIS. (From *lathō*, to forget; because it was thought to affect the memory.) Spurge.

LATHYRUS. (From *lathō*, to lie hid; so called from its diminutive size.) The vetch.

LATIBULUM. (From *lateo*; to lie hid.) The fomes, or hidden matter of infectious diseases.

LATISSIMUS COLLI. See *Platyema myoides*.

LATISSIMUS DORSI. (*Latissimus*, sc. *musculus*.; *Antiscaptor* of Cowper. *Dorsi-lumbo sacro humeral* of Dumas. A muscle of the humerus, situated on the posterior part of the trunk. It is a very broad, thin, and for the most part fleshy muscle, which is placed immediately under the skin, except where it is covered by the lower extremity of the trapezius. It arises tendinous from the posterior half of the upper edge of the spine of the os ilium, from the spinous processes of the os sacrum and lumbar vertebrae, and from five or six, and sometimes from seven, and even eight, of the lowermost ones of the back; also tendinous and fleshy from the upper edges and external surface of the four inferior false ribs near their cartilages, by as many distinct slips. From these different origins the fibres of the muscle run in different directions; those from the ilium and false ribs run almost perpendicularly upwards; those from the sacrum and lumbar vertebrae, obliquely upwards and forwards; and those from the vertebrae of the back, transversely outwards and forwards, over the inferior angle of the scapula, where they receive a small thin bundle of fleshy fibres, which arise tendinous from that angle, and are inserted with the rest of the muscle, by a strong, flat, and thin tendon, of about two inches in length, into the fore-part of the posterior edge of the groove observed between the two tuberosities of the os humeri, for lodging the tendon of the long head of the biceps. In dissection, therefore, this muscle ought not to be followed to its insertion, till some of the other muscles of the os humeri have been first raised. Its use is to pull the os humeri downwards and backwards, and to turn it upon its axis. Riolanus, from its use on certain occasions, gave it the name of *antertoror*. When we raise ourselves upon our hands, as in rising from off an arm-chair, we may easily perceive the contraction of this muscle. A *bursa mucoea* is found between the tendon of this muscle and the os humeri, into which it is inserted.

LAUCANIA. (From *lauro*, to receive; so called because it receives and conveys food.) The oesophagus of the throat.

Laudanum. (From *laus*, praise; so

named from its valuable properties.) See *Tinctura opii*.

Laurel, cherry. See *Laurocerasus*.

Laurel, spurge. See *Laureola*.

LAUREOLA. (Dim. of *laurus*, the laurel, named from it resembling to the laurel. Spurge laurel. The bark of this plant, *Daphne laureola* of Linnæus, is recommended to excite a discharge from the skin, in the same way as that of the *thymelea*.

LAURO CERASUS. (From *laurus*, the laurel, and *cerasus*, the cherry-tree; so called because it has leaves like the laurel.) Common or cherry laurel. *Prunus lauro-cerasus* of Linnæus:—*floribus racemosis foliis sempervirentibus dorso biglandulosis*. The leaves of the lauro-cerasus have a bitter styptic taste, accompanied with a flavour resembling that of bitter-almonds, or other kernels of the drupaceous fruits; the flowers also manifest a similar flavour. The powdered leaves, applied to the nostrils, excite sneezing, though not so strongly as tobacco. The kernel-like flavour which these leaves impart, being generally esteemed grateful, has sometimes caused them to be employed for culinary purposes, and especially in custards, puddings, blanchmange, &c.; and as the proportion of this sapid matter of the leaf to the quantity of the milk is commonly inconsiderable, bad effects have seldom ensued. But as the poisonous quality of this laurel is now indubitably proved, the public ought to be cautioned against its internal use.

The following communication to the Royal Society, by Dr. Madden, of Dublin, contains the first and principal proofs of the deleterious effects of this vegetable upon mankind:—"A very extraordinary accident that fell out here some months ago, has discovered to us a most dangerous poison, which was never before known to be so, though it has been in frequent use among us. The thing I mean is a simple water, distilled from the leaves of the lauro-cerasus; the water is at first milky, but the oil which comes over being, in a good measure, separated from the phlegm, by passing it through a flannel bag, becomes as clear as common water. It has the smell of bitter almonds, or peach-kernel, and has been for many years in frequent use among our housewives and cooks, to give that agreeable flavour to their creams and puddings. It has also been much in use among our drinkers of drams; and the proportions they generally use it in has been one part of laurel water to four of brandy. Nor has this practice, however frequent, ever been attended with any apparent ill consequences, till some time in the month of September 1728, when it happened that one Martha Boyse, a servant, who lived with a person who sold great quantities of

this water, got a bottle of it from her mistress, and gave it to her mother. Ann Boyse made a present of it to Frances Eaton, her sister, who was a shopkeeper in town, and who she thought might oblige her customers with it. Accordingly, in a few days, she gave about two ounces to a woman called Mary Whaley, who drank about two-thirds of what was filled out, and went away. Frances Eaton drank the rest. In a quarter of an hour after Mary Whaley had drank the water, (as I am informed,) she complained of a violent disorder in her stomach, soon after lost her speech, and died in about an hour, without vomiting, or purging, or any convulsion. The shopkeeper, F. Eaton, sent word to her sister, Ann Boyse, of what had happened, who came to her upon the message, and affirmed that it was not possible the cordial (as she called it) could have occasioned the death of the woman; and, to convince her of it, she filled out about three ounces and drank it. She continued talking with F. Eaton about two minutes longer, and was so earnest to persuade her of the liquor's being inoffensive, that she drank about two spoonfuls more, but was hardly well seated in her chair when she died without the least groan, or convulsion. Frances Eaton, who, as before observed, had drank somewhat more than a spoonful, found no disorder in her stomach, or elsewhere; but, to prevent any ill consequences, she took a vomit immediately, and has been well ever since."—Dr. Madden mentions another case, of a gentleman at Kilkeny who mistook a bottle of laurel-water for a bottle of pisan. What quantity he drank is uncertain, but he died in a few minutes, complaining of a violent disorder in the stomach. In addition to this, we may refer to the unfortunate case of Sir Theodosius Boughton, whose death, in 1780, an English jury declared to be occasioned by this poison. In this case, the active principle of the lauro-cerasus was concentrated by repeated distillations, and given to the quantity of one ounce; the suddenly fatal effects of which must be still in the recollection of the public. To brute animals this poison is almost instantaneously mortal, as amply appears by the experiments of Madden, Mortimer, Nicholls, Fontana, Langrish, Vater, and others. The experiments conducted by these gentlemen shew, that the laurel-water is destructive to animal life, not only when taken into the stomach, but also on being injected into the intestines, or applied externally to different organs of the body. It is remarked, by Abbé Fontana, that this poison, even "when applied in a very small quantity to the eyes, or to the inner part of the mouth, without touching the œsophagus, or being carried into the stomach, is capable of killing an animal in a few

minutes: whilst, applied in a much greater quantity to wounds, it has so fluid activity, that the weakest animals, such as pigeons, resist its action."

The most volatile is the most active part of the lauro-cerasus; and if we judge from its sensible qualities, an analogous principle seems to pervade many other vegetable substances, especially the kernels of drupaceous fruits; and in various species of the amygdalus, this sapid principle extends to the flowers and leaves. It is of importance to notice, that this is much less powerful in its action upon human subjects than upon dogs, rabbits, pigeons, and reptiles. To poison man, the essential oil of the lauro-cerasus must be separated by distillation, as in the spirituous or common laurel-water; and unless this is strongly embued with the oil, or given in a large dose, it proves innocent. Dr. Cullen observes, that the sedative power of the lauro-cerasus acts upon the nervous system in a different manner from opium and other narcotic substances, whose primary action is upon the animal functions; for the lauro-cerasus does not occasion sleep, nor does it produce local inflammation, but seems to act directly upon the vital powers. Abbé Fontana supposes that this poison destroys animal life, by exerting its effects upon the blood; but the experiments and observations from which he draws this opinion are evidently inconclusive. It may also be remarked, that many of the Abbé's experiments contradict each other. Thus it appears, from the citation given above, that the poison of this vegetable, when applied to wounds, does not prove fatal; but future experiments led the Abbé to assert, that the oil of the lauro-cerasus, "whether given internally, or applied to the wounds of animals, is one of the most terrible and deadly poisons known." Though this vegetable seems to have escaped the notice of Stœrck, yet it is not without advocates for its medical use. Linnæus informs us, that in Switzerland it is commonly and successfully used in pulmonary complaints. Langrish mentions its efficacy in agues; and as Bergius found bitter almonds to have this effect, we may, by analogy, conclude that this power of the lauro-cerasus, is well established. Baylies found, that it possessed a remarkable power of diluting the blood, and, from experience, recommended it in all cases of disease supposed to proceed from too dense a state of that fluid; adducing particular instances of its efficacy in rheumatism, asthma, and in schirrous affections. Nor does this author seem to have been much afraid of the deleterious quality of lauro-cerasus, as he directs a pound of its leaves to be macerated in a pint of water, of which he gives from thirty to sixty drops three or four times a-day.

LAURUS. The spodium of silver; so

called from Mount Laurus, where there were silver mines.

LAURUS. (From *laus*, praise; because it was usual to crown the heads of eminent men with branches of it.) 1. The name of a genus of plants in the Linnaean system. Class, *Enneundria*. Order, *Monogymia*. The laurel.

2. The pharmacopœial name of the sweet-bay. *Laurus nobilis* of Linnæus:—*foliis venosis lanceolatis perennantibus, floribus quadrifidis*. This tree is a native of Italy, but cultivated in our gardens and shrubberies as a handsome evergreen. The leaves and berries possess the same medicinal qualities, both having a sweet fragrant smell, and an aromatic adstringent taste. The laurus of honorary memory, the distinguished favourite of Apollo, may be naturally supposed to have had no inconsiderable fame as a medicine; but its pharmaceutical uses are so limited in the practice of the present day, that this dignified plant is now rarely employed, except in the way of enema, or as an external application; thus the leaves are directed in the *decoctum pro fomento*, and the berries in the *emplastrum cumini*.

LAURUS BENZOIN. The systematic name of the benjamin-tree. See *Benzoinum*.

LAURUS CAMPHORA. The systematic name of the camphire-tree. See *Camphora*.

LAURUS CINNAMOMUM. The systematic name of the cinnamon-tree. See *Cinnamomum*.

LAURUS CULILAWAN. The systematic name of the plant whose bark is called *cortex culilawan* in the shops.

LAURUS NOBILIS. The systematic name of the sweet-bay tree. See *Laurus*.

LAURUS SASSAFRAS. The systematic name of the sassafras-tree. See *Sassafras*.

Lavender, French. See *Stæchas*.

LAVENDULA. (From *lavo*, to wash; so called, because, on account of its fragrance, it was used in baths.) 1. The name of a genus of plants in the Linnaean system. Class, *Didynamia*. Order, *Gymnospermia*.

2. The pharmacopœial name of the common lavender. *Lavendula spica* of Linnæus:—*foliis sessilibus lanceolatis linearibus margine revolutis, spica interrupta nuda*. A native of the southern parts of Europe, but cultivated in our gardens on account of the fragrance of its flowers. Their taste is bitter, warm, and somewhat pungent; the leaves are weaker and less grateful. The essential oil, obtained by distillation, is of a bright yellow colour, of a very pungent taste, and possesses, if carefully distilled, the fragrance of the lavender in perfection. Lavender has been long recommended in nervous debilities, and various affections proceeding from a want of energy in the

animal functions. The College directs an essential oil, a simple spirit, and a compound tincture, to be kept in the shops.

LAVENDULA SPICA. The systematic name of the common lavender. See *Lavendula*.

LAVENDULA STÆCHAS. The systematic name of the French lavender. See *Stæchas*.

LAVER. (From *lavo*, to wash; so named because it is found in brooks, where it is constantly washed by the stream.) 1. The brook-lime.

2. The English name of a species of fucus which is eaten as a delicacy.

LAVIPEDIUM. (From *lavo*, to wash, and *pes*, the foot.) A bath for the feet.

LAWSONIA INERMIS. The systematic name of the true alkanna. See *Alkanna vera*.

LAXATIVA. (From *laxo*, to loosen.) Gentle purgatives.

LAXATOR TYMPANI. (From *laxo*, to loosen; so called from its office to relax the drum of the ear.) *Externus mallei* of Albinus. *Anterior mallet* of Winslow. *Obliquus auris* of Douglas. *Externus auris vel laxator internus* of Cowper, and *Spheni salpingo mallei* of Dumas. A muscle of the internal ear, that draws the malleus obliquely forwards towards its origin; consequently, the membrana tympani is made less concave, or is relaxed.

LAZULUS. (From *azul*, Arabian.) A precious stone, of a blue colour. The lapis lazuli.

LEAD. *Plumbum*. A metal found in considerable quantity in many parts of the earth, in different states, seldom if at all in the metallic state. It is found in that of oxyd, red lead ore, mixed with a portion of iron, clay, and other earths. The colour of this ore is aurora red, resembling red arsenic. It is found in small lumps, of an indeterminate figure, and also crystallized in four-sided rhomboidal prisms.

Combined with carbonic acid, it forms the *sparry lead ore*, so called because it has the texture and crystallization of certain spars. There are a great many varieties of this kind. It is found united with sulphuric, phosphoric, arsenic, molybdic, and chromic acids. Lastly, lead is found mineralized by sulphur, forming what is called *galena* (*sulphuret of lead*), which is by far its most abundant ore. This ore, which is very common, is found both in masses and crystals.

The primitive form of its crystals is a cube. Its colour is of a blueish lead grey. It has a considerable metallic lustre, its texture is foliated. It stains the fingers, and often feels greasy. It contains in general a minute quantity of silver.

Properties of Lead.—Lead is of a blueish white colour when fresh cut. It is malle-

able. It soon tarnishes in the atmosphere. It may easily be cut with a knife, and stains the fingers bluish-grey when rubbed. It fuses at 550° Fahr. and renders other more refractory metals fusible. It becomes vitrified in a strong and continued heat, and vitrifies various other metals. It is the least elastic of all the metals. It is very laminable, but it possesses very little ductility. Its specific gravity is 11.435. It crystallizes by cooling in small octahedra. When fused, its surface first becomes yellow and then red. It unites by fusion with phosphorus and sulphur. The greater part of the acids act upon it. The sulphuric acid requires the assistance of a boiling heat. Nitric acid is decomposed by it. Muriatic acid acts very weakly on it. Acetic acid dissolves it. Fluoric acid attacks it by heat, and slightly in the cold. It combines with other metals, but few of its alloys are applied to any use. When combined with mercury it forms a crystallizable alloy, which becomes fluid when triturated with that of bismuth.

Method of obtaining Lead.—In order to obtain lead in a great way, the ore is picked from among the extraneous matter with which it was naturally mixed. It is then pulverized and washed. It is next roasted in a reverberatory furnace, in which it is to be agitated, in order to bring all its surfaces in contact with the air. When the external parts begin to soften, or assume the form of a paste, it is covered with charcoal, the mixture is stirred, and the heat increased gradually; the lead then runs on all sides, and is collected at the bottom of the furnace, which is perforated so as to permit the metal to flow into a receptacle defended by a lining of charcoal.

The scoria remaining above in the furnace still retain a considerable proportion of lead; in order to extract it, the scoria must be fused in a blast furnace. The lead is by that means separated, and cast into iron moulds, each of which contains a portion called a *pig of lead*. These pigs are sold under the name of *ore lead*.

To disengage the silver from lead thus obtained, the metal is subjected to the action of the refining furnace. The continual application of a quantity of fresh air, which is thrown by means of large bellows upon the fused lead, which is at the same time heated as intensely as possible, oxidates the lead, and converts it into the yellow scaly oxyd, known by the name of *litharge*.

This scaly oxyd being driven off from the surface of the fused metal, as it is formed, leaves the silver alone unaltered at the bottom.

The litharge is then to be fused in contact with charcoal, that it may assume the properties of metallic lead.

In order to obtain perfectly pure lead, the lead of commerce may be dissolved in pure nitric acid, and the solution be decomposed, by adding to it, gradually, a solution of sulphate of soda, so long as a precipitate ensues. This precipitate, which is sulphate of lead, must then be collected on a filter, washed repeatedly in distilled water, and then dried. In order to reduce it to its metallic state, let it be mixed with two or three times its weight of black flux, introduce the mixture into a crucible, and expose it briskly to a red heat.

Lead, when injudiciously administered, or taken accidentally into the body, causes emaciation, violent colics, paralysis, tremors, and contractions of the limbs; and as they generally come on gradually, the cause is sometimes overlooked till it be too late. Poisoning from lead is never intentional, but only accidental; either from liquors becoming impregnated with lead, by being improperly kept in vessels lined or glazed with lead, or to which lead has been criminally added, to correct its acidity; or among manufacturers who work much with lead, as painters, or plumbers, and who are not sufficiently attentive to avoid swallowing it. The presence of lead in any suspected liquor is detected by the hydro-sulphuret of potash, which forms with it a brown precipitate, not soluble in diluted muriatic acid, and still more certainly by evaporating a portion of the liquor to dryness, and exposing the extract to a heat sufficient to reduce the lead.

The preparations of lead used in medicine are:—

1. Oxidum plumbi album. See *Cerussa* and *Plumbi carbonas*.

2. Oxidum plumbi rubrum. See *Minium*.

3. Oxidum plumbi semivitrum. See *Lithargyrum* and *Oxidum plumbi semivitrum*.

4. Supracetas plumbi. See *Plumbi superacetas*.

5. Liquor plumbi acetatis. See *Liquor plumbi acetatis*.

6. Liquor plumbi acetatis dilutus. See *Liquor plumbi acetatis dilutus*.

LAZULI LAPIS. See *Lapis lazuli*.

LEDUM PALUSTRE. The systematic name of the *rosmarinus sylvestris*. See *Rosmarinus sylvestris*.

LEENA. (From *leena*, a lioness; so named from its power.) A plaster for the hip.

LEECH. *Hirudo*. A genus of insects belonging to the order of vermes intestina. The body moves either forward or backward. There are several species, principally distinguished by their colour; but that most known to medical men is the *hirudo medicinalis*, or medicinal leech, which grows to the length of two or three

inches. The body is of a blackish brown colour, marked on the back with six yellow spots, and edged with a yellow line on each side; but both the spots and lines grow faint, and almost disappear, at some seasons. The head is smaller than the tail, which fixes itself very firmly to any thing the creature pleases. It is viviparous, and produces but one young one at a time, which is in the month of July. It is an inhabitant of clear running waters, and is well known for its use in bleeding. The species most nearly approaching this, and which it is necessary to distinguish, is the *hirudo sanguisuga*, or horse-leech. This is larger than the former; its skin is smooth and glossy; the body is depressed, the back is dusky; and the belly is of a yellowish green, having a yellow lateral margin. It inhabits stagnant waters.

The leech's head is armed with a sharp instrument that makes three wounds at once. They are three sharp tubercles, strong enough to cut through the skin of a man, or even of an ox, or horse. The mouth is, as it were, the body of the pump, and the tongue, or fleshy nipple, the sucker; by the working of this piece of mechanism, the blood is made to rise up to the conduit which conveys it to the animal's stomach, which is a membranaceous skin, divided into twenty-four small cells. The blood which is sucked out is there preserved for several months, almost without coagulating, and proves a store of provision to the animal. The nutritious parts, absorbed after digestion by animals, need not in this to be disengaged from the heterogeneous substances; nor indeed is there an anus discoverable in the leech; mere transpiration seems to be all that it performs, the matter fixing on the surface of the body, and afterwards coming off in small threads. Of this, an experiment may be tried, by putting a leech into oil, where it keeps alive for several days; upon being taken out, and put into water, there appears to loosen from its body a kind of slough, shaped like the creature's body. The organ of respiration, though unascertained, seems to be situated in the mouth; for if, like an insect, it drew breath through vent-holes, it would not subsist in oil, as, by it, these would be stopped up.

The first species only is used in medicine; being applied to the skin in order to draw off blood. With this view they are employed to bleed young children, and for the purposes of topical bleeding, in cases of inflammation, fulness, or pain. They may be employed in every case where topical bleedings are thought necessary, or where venesection cannot be performed. If the leech does not fasten, a drop of sugared milk is put on the spot it is wished to fix on, or a little blood is drawn by means of a slight puncture; after which

it immediately settles. The leech, when fixed, should be watched, lest it should find its way into the anus, when used on the hæmorrhoids, or penetrate into the œsophagus, if employed to draw the gums; otherwise it might fix upon the stomach, or intestines. In such a case, the best and quickest remedy is to swallow some salt: which is the method practised to make it loose its hold, when it sucks longer than is intended. Vegetable or volatile alkali, pepper, or acids, also make it leave the part on which it was applied. Cows and horses have been known to receive leeches, when drinking, into the throat; and the usual remedy is to force down some salt, which makes them fall off. If it is intended that the leech should draw a larger quantity of blood, the end of the tail is cut off, and it then sucks continually, to make up the loss it sustains. The discharge occasioned by the puncture of a leech is usually of more service than the process itself. When too abundant, it is easily stopped with brandy, vinegar, or other styptics, or with a compress of dry linen rags, bound strongly on the bleeding orifice. They are said to be very restless before a change of weather, if confined to glasses, and to fix themselves above the water on the approach of a fine day.

As these little animals are depen-
dent on for the removal of very dangerous diseases, and as they often seem capriciously determined to resist the endeavours made to cause them to adhere, the following directions are added, by which their assistance may, with more certainty, be obtained.

The introducing a hand, to which any ill-flavoured medicine adheres, into the water in which they are kept, will be often sufficient to deprive them of life; the application of a small quantity of any saline matter to their skin immediately occasions the expulsion of the contents of their stomach; and what is most to our purpose, the least flavour of any medication that has been applied remaining on the skin, or even the accumulation of the matter of perspiration, will prevent them from fastening. The skin should therefore, previous to their application, be very carefully cleansed from any foulness, and moistened with a little milk. The method of applying them is by retaining them to the skin by a small wine-glass, or the bottom of a large pill-box, when they will, in general, in a little time fasten themselves to the skin. On their removal, the rejection of the blood they have drawn may be obtained by the application of salt externally; but it is to be remarked, that a few grains of salt are sufficient for this purpose; and that covering them with it, as is sometimes done, generally destroys them.

LEEK. *Allium porrum*. A well-known vegetable, much employed for culinary

purposes. The recent root and juice are exhibited internally in quartan fever, in dyspepsy, dropsy, asthma, and scurvy. See *Porrum*.

LENA. (From *λενω*, a fringed edge.) The extremities of the pudenda muliebrum.

LEGUMEN. (From *lego*, to gather; so called because they are usually gathered by the hand.) All kinds of pulse are so called.

LEICHEN. See *Lichen*.

LEIENTERIA. See *Lienteria*.

LEIPOPSYCHIA. (From *λεπω*, to leave, and *ψυχη*, the soul, or life.) A swoon. See *Syncope*.

LEIPOPYELA. (From *λεπω*, to leave, and *πυρ*, heat.) A kind of ardent fever, where the internal parts are scorched with heat, while the external parts are cold.

LEIPOTHYMIA. (From *λεπω*, to leave, and *θυμος*, the mind.) See *Lipothymia*.

LEME. (From *λα*, much, and *μω*, to wink.) A defect in the eyes, when they are always winking.

LEMITHOCORPON. See *Corallina Cori-cana*.

LEMMA. (From *λεω*, to decorticate.) Bark. The skin.

LEMNIUS. (From *Lemnos*, whence it is brought.) A species of bole called terra lemnia, or earth of Lemnos.

Lemon. See *Limon*.

Lemon scurvy-grass. See *Cochlearia hortensis*.

LENIENTIA. (From *lenio*, to assuage.) Medicines which abate irritation.

LENITIVA. (From *lenis*, gentle.) Medicines which gently palliate diseases. Gentle purgatives.

LENITIVE ELECTUARY. A preparation composed chiefly of senna and some aromatics, with the pulp of tamarinds. It is given in doses of a tea-spoonful, or more, frequently repeated, as a mild laxative; and, when fresh, it answers this purpose well. See *Confectio Sennæ*.

LENS. (A *lentore*; from its glutinous quality.) 1. The lentil. *φασις* of the Greeks. *Ervum lens* of Linnæus:—*pedunculis subfloris*; *seminibus compressis, convexis*. There are two varieties; the one with large, the other small seeds. They are eaten in many places as we eat peas, than which they are more flatulent, and more difficult to digest. A decoction of these seeds is used as a lotion to the ulcerations after small-pox, and, it is said, with success.

2. See also *Chryntalline lens*.

LENTICULA. (Dim. of *lens*, a lentil.) A smaller sort of lentil. Also a freckle, or small pustule, resembling the seeds of lentil.

LENTICULAR. (From *lenticulaire*, doubly convex.) A surgical instrument, employed for removing the jagged particles of bone from the edge of the perforation made in the cranium with the trephine.

LENTICULARIA. (From *lenticula*.) A species of lentil.

LENTIGO. (From *lens*, a lentil; so named from its likeness to lentil-seeds.) A freckle.

LENTIL. An annual vegetable of the pulse kind, much used for improving the flavour of soups.

LENTISCUS. (From *lenteo*, to become clammy; so called from the gumminess of its juice.) The mastich-tree.

LENTOR. (From *lentus*, clammy.) A viscosity or siziness of any fluid.

LEONINUS. (From *leo*, the lion.) An epithet of that sort of leprosy called leontiasis.

LEONTIASIS. (From *λεων*, a lion; so called because it is said lions are subject to it.) A species of leprosy resembling the elephantiasis.

LEONTODON. (From *λεων*, the lion, and *odus*, a tooth; so called from its supposed resemblance. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia equalis*. The dandelion.

LEONTOPDIUM. (From *λεων*, a lion, and *πυρ*, a foot; so named from its supposed resemblance.) The herb lion's foot.

LEONURUS. (From *λεων*, a lion, and *ουρα*, a tail; so named from its likeness.) 1. The name of a genus of plants in the Linnæan system. Class, *Dicotyledonia*. Order, *Gymnospermia*. Lion's tail.

2. The name, in some pharmacopœias, for the lion's tail.

Leopard's bane. See *Arnica*.

LEPIDIUM. (From *λεπιδ*, a scale; so named from its supposed usefulness in cleansing the skin from scales and impurities.) Pepper-wort.

LEPIDOMARCOMA. (From *λεπιδ*, a scale, and *μαρμα*, flesh.) An irregular scaly tumour.

LEPISMA. (From *λεπιζω*, to decorticate.) Decortication. A peeling off of the skin.

LEPRA. (From *λεπιδ*, a scale; named from its appearance. The leprosy. A disease in the class *cachexie*, and order *impetiginæ*, of Cullen. Dr. Willan describes this disease as characterized by scaly patches, of different sizes, but having always nearly a circular form. In this country, three varieties of the disease are observed, which he has described under the title of *Lepra vulgaris*, *Lepra alba*, *Lepra nigricans*.

1. The *Lepra vulgaris* exhibits first small distinct elevations of the cuticle, which are reddish and shining, but never contain any fluid; these patches continue to enlarge gradually, till they nearly equal the dimensions of a crown-piece. They have always an orbicular, or oval form; are covered with dry scales, and surrounded by a red border. The scales accumulate on them, so as to form a thick prominent crust,

which is quickly re-produced, whether it fall off spontaneously, or may have been forcibly detached. This species of lepra sometimes appears first at the elbow, or on the fore-arm; but more generally about the knee. In the latter case, the primary patch forms immediately below the patella, within a few weeks, several other scaly circles appear along the fore part of the leg and thigh, increasing by degrees, till they come nearly into contact. The disease is then often stationary for a considerable length of time. If it advance further, the progress is towards the hip and loins; afterwards, to the sides, back, shoulders, and, about the same time, to the arms and hands. In the greater number of cases, the hairy scalp is the part last affected: although the circles formed on it remain for some time distinct, yet they finally unite, and cover the whole surface on which the hair grows with a white scaly incrustation. This appearance is attended, more especially in hot weather, with a troublesome itching, and with a watery discharge for several hours, when any portion of the crust is detached, which takes place from very slight impressions. The pubes in adults is sometimes affected in the same manner as the head: and if the subject be a female, there is usually an internal *pruritus pudendi*. In some cases of the disorder, the nails, both of the fingers and toes, are thickened, and deeply indented longitudinally. When the lepra extends universally, it becomes highly disgusting in its appearance, and inconvenient from the stiffness and torpor occasioned by it in the limbs. The disease, however, even in this advanced stage, is seldom disposed to terminate spontaneously. It continues nearly in the same state for several years, or sometimes during the whole life of the person affected, not being apparently connected with any disorder of the constitution.

2. *Lepra alphas*. The scaly patches in the alphas are smaller than those of the lepra vulgaris, and also differ from them in having their central parts depressed or indented. This disorder usually begins about the elbow, with distinct, eminent asperities, of a dull red colour, and not much longer than papillæ. These, in a short time, dilate to nearly the size of a silver penny. Two or three days afterwards, the central part of them suffers a depression, within which small white powdery scales may be observed. The surrounding border, however, still continues to be raised, but retains the same size and the same red colour as at first. The whole of the fore-arm, and sometimes the back of the hand, is spotted with similar patches: they seldom become confluent, excepting round the elbow, which, in that case, is covered with an uniform crust. This affec-

tion appears in the same manner upon the joint of the knee, but without spreading far along the thigh or leg. Dr. Willan has seldom seen it on the trunk of the body, and never on the face. It is a disease of long duration, and not less difficult to cure than the foregoing species of lepra; even when the scaly patches have been removed by persevering in the use of suitable applications, the cuticle still remains red, tender, and brittle, very slowly recovering its usual texture. The alphas, as above described, frequently occurs in this country.

3. The *Lepra nigricans* differs little from the lepra vulgaris, as to its form and distribution. The most striking difference is in the colour of the patches, which are dark and livid. They appear first on the legs and fore-arms, extending afterwards to the thighs, loins, neck, and hands. Their central part is not depressed, as in the alphas. They are somewhat smaller in size than the patches of the lepra vulgaris, and not only is the border livid or purplish, but the livid colour of the base likewise appears through the scaly incrustation, which is seldom very thick. It is further to be observed, that the scales are more easily detached than in the other forms of lepra, and that the surface remains longer excoriated, discharging lymph, often with an intermixture of blood, till a new incrustation forms, which is usually hard, brittle, and irregular. The lepra nigricans affects persons whose occupation is attended with much fatigue, and exposes them to cold or damp, and to a precarious or improper mode of diet, as soldiers, brewers, labourers, butchers, stage-coachmen, sculler-men, &c.; some women are also liable to it, who are habituated to poor living and constant hard labour.

LEPRA VULGARIS. The lepra vulgaris, alphas, and nigricans, have all been so described.

Leprosy. See *Lepra*.

LEPTOMICA. (From λεπτος, thin.) Attenuating medicines.

LEPTYANUS. (From λεπτος, slender.) Attenuation, or the making a substance less solid.

LEROS. (From λερος, to trifle.) A slight delirium.

LETHARGY. *Lethargus*. A heavy and constant sleep, with scarce any intervals of waking; when awakened, the person answers, but ignorant or forgetful of what he said, immediately sinks into the same state of sleep. It is considered as an imperfect apoplexy, and is mostly symptomatic.

LETHEA. (From ληθη, forgetfulness; so named because it causes forgetfulness.) The name of the poppy.

Lettuce, garden. See *Lactuca*.

LEUCACANTHA. (From λευκος, white,

and *anemone*, a thorn; so named from its white blossom.) The cotton-thistle.

LEUCANTHEMUM VULGARE. (From λευκος, white, and ανθος, a flower; so called from its white floret.) See *Bellis major*.

LEUCICETRUM. (From λευκος, white, and ιαμαρον, amber.) White amber.

LEUCOLACHANUM. (From λευκος, white, and λαχανον, a herb; so named from its colour.) Wild valerian.

LEUCOMA. (From λευκος, white) Leucoma and albugo are often used synonymously, to denote a white opacity of the cornea. Both of them, according to Scarpa, are essentially different from the nebula of the cornea; for they are not the consequence of chronic ophthalmia, attended with varicose veins, and an effusion of a milky serum into the texture of the delicate continuation of the conjuncture over the cornea; but are the result of violent acute ophthalmia. In this state, a dense coagulating lymph is extravasated from the arteries, sometimes superficially, at other times deeply into the substance of the cornea. On other occasions, the disease consists of a firm callous cicatrix on this membrane, the effects of an ulcer, or wound, with loss of substance. The term *albugo* statically belongs to the first form of the disease: *leucoma* to the last, more particularly when the opacity occupies the whole, or the chief part, of the cornea.

LEUCONYMPHEA. (From λευκος, white, and νυμφα, the water-lily.) See *Nymphaea alba*.

LEUCOPHAGIUM. (From λευκος, white, and φαγω, to eat.) A medicated white food.

LEUCOPHLEGMATIC. (*Leucophlegmatica*; from λευκος, white, and φλεγμα, phlegma.) A term applied by the older medical writers to a dropsical habit of body.

LEUCOPYPER. (From λευκος, white, and πιπερικ, pepper.) See *Piper nigrum*.

LEUCORRŒA. (From λευκος, white, and ροη, to flow.) *Fluxus albus*. The whites. An increased secretion of white mucus from the vagina of women, arising from debility, and not from the venereal virus.

LEUCORRHOÏS. (From λευκος, white, and ροη, to flow.) A discharge of mucus from the intestines.

LEVATOR. (From *levo*, to lift up.) A muscle whose office is to lift up the part to which it is attached.

LEVATOR ANGULI ORIS. *Elevator labiorum communis* of Douglas. *Caninus* of Winslow, and *Sus maxilla labialis* of Dumas. A muscle situated above the mouth, which draws the corner of the mouth upwards, and makes that part of the cheek opposite to the chin prominent, as in smiling. It arises thin and fleshy from the hollow of the superior maxillary bone, between the

root of the socket of the first grinder and the foramen infra orbitarium, and is inserted into the angle of the mouth and under lip, where it joins with its antagonist.

LEVATOR ANI. *Levator magnus, seu internus* of Douglas. *Pubo coccygi annulare* of Dumas. This muscle arises from the os pubis, within the pelvis, as far up as the upper edge of the foramen thyroideum, and joining of the os pubis with the os ischium, from the thin tendinous membrane that covers the obturator internus and coccygeus muscles, from the spinous process of the ischium. From these origins all round the inside of the pelvis, its fibres run down like rays from the circumference to a centre, to be inserted into the sphincter ani, accelerators urine, and anterior part of the two last bones of the os coccygis, surrounding the extremity of the rectum, neck of the bladder, prostate gland, and part of the testiculæ seminales. Its fibres, joining with those of its fellow, form a funnel-shaped hole, that draws the rectum upwards after the evacuation of the feces, and assists in shutting it. The levatores ani also sustain the contents of the pelvis, and assist in ejecting the semen, urine, and contents of the rectum, and perhaps, by pressing upon the veins, contribute greatly to the erection of the penis.

LEVATOR LABII INFERIORIS. *Levator menti* of Albinus. *Incisivus inferior* of Winslow. *Elevator labii inferioris proprius* of Douglas. A muscle of the mouth situated below the lips; it arises from the lower jaw, at the roots of the alveoli of two incisor teeth and the cuspidatus, and is inserted into the under lip and skin of the chin.

LEVATOR LABII SUPERIORIS ALÆQUE NASI. *Elevator labii superioris proprius* of Douglas. *Incisivus lateralis et pyramidalis* of Winslow. A muscle of the mouth and lips, that raises the upper lip towards the orbit, and a little outwards; it serves also to draw the skin of the nose upwards and outwards, by which the nostril is dilated. It arises by two distinct origins; the first, broad and fleshy, from the external part of the orbital process of the superior maxillary bone, immediately above the foramen infra orbitarium; the second, from the nasal process of the superior maxillary bone, where it joins the os frontis. The first portion is inserted into the upper lip and orbicularis muscle, the second into the upper lip and outer part of the ala nasi.

LEVATOR LABII SUPERIORIS PROPRIUS. *Musculus incisivus*. It arises under the edge of the orbit, and is inserted into the middle of the lip.

LEVATOR OCULI. See *Rectus superior oculi*.

LEVATOR PALATI. *Levator palati molle* of Albinus. *Petrosalpingo-staphilius*, vel *salpingo-staphilius internus vulgo* of Winslow. *Salpingo-staphilius* of Valsalva. *Pterigo-staphilius externus vulgo* of Douglas. *Spheno-staphilius* of Cowper. A muscle situated between the lower jaw and the os hyoides laterally. It arises tendinous and fleshy from the extremity of the petrous portion of the temporal bone, where it is perforated by the Eustachian tube, and also from the membranous part of the same tube, and is inserted into the whole length of the velum pendulum palati, as far as the root of the uvula, and unites with its fellow. Its use is to draw the velum pendulum palati upwards and backwards, so as to shut the passage from the fauces into the mouth and nose.

LEVATOR PALATI MOLLIS. See *Levator palati*.

LEVATOR PALPEBRÆ SUPERIORIS. *Aperiens palpebrarum rectus. Aperitor oculi.* A proper muscle of the upper eyelid, that opens the eye, by drawing the eyelid upwards. It arises from the upper part of the foramen opticum of the sphenoid bone, above the rectus superior oculi, near the trochlearis, and is inserted by a broad thin tendon into the cartilage that supports the upper eyelid.

LEVATOR PARVUS. See *Transversus perinei*.

LEVATOR SCAPULÆ. A muscle situated on the posterior part of the neck, that pulls the scapula upwards and a little forwards. This name, which was first given to it by Riolaus, has been adopted by Albinus. Douglas calls it *elevator seu musculus patientie*; and Winslow, *angularis vulgo levator proprius*. It is a long muscle, nearly two inches in breadth, and is situated obliquely under the anterior edge of the trapezius. It arises tendinous and fleshy from the transverse processes of the four and sometimes five superior vertebræ colli, by so many distinct slips, which soon unite to form a muscle that runs obliquely downwards an outwards, and is inserted by a flat tendon into the upper angle of the scapula. Its use is to raise the scapula upwards, and a little forwards.

LEVIGATION. The reduction of hard substances, by triture, to impalpable powders.

LEVISTICUM. (From *levo*, to assuage; so called from the relief it gives in painful flatulencies.) Lovage. The odour of this plant, *Ligustrum levisticum* of Linnæus: *foliis multiplicibus, foliis superne incisiss*, is very strong, and particularly ungrateful; its taste is warm and aromatic. It abounds with a yellowish gummy resinous juice, very much resembling opoponax. Its virtues are supposed to be similar to those of angelica and master-wort, in expelling

flatulences, exciting sweat, and opening obstructions; therefore it is chiefly used in hysterical disorders and uterine obstructions. The leaves, eaten in salad, are accounted emmenagogue. The root, which is less ungrateful than the leaves, is said to possess similar virtues, and may be employed in powder.

LEXIPHARMACA. (From *λεγω*, to terminate, and *φαρμακον*, poison.) Medicines which resist or destroy the power of poison.

LEXIPHYRETA. (From *λεγω*, to make cease, and *πυρετος*, a fever.) Febrifuge medicines.

LIBADUM. (From *λεβω*, to make moist; so called because it grows in watery places.) The lesser centaury.

LIBANOTIS. (From *λεβανος*, frankincense so called from its resemblance in smell to frankincense.) Rosemary.

LIBANUS. (From *Libanon*, a mountain in Syria, where it grows.) The frankincense-tree.

LIBOR. (From *λεω*, to distil.) A rheum or defluxion from the eyes.

LIBURNUM. (From *Liburnia*, the country where it flourished.) The mealy-tree.

LICHANUS. (From *λεχω*, to lick; so called because it is commonly used in licking up any thing.) The fore-finger.

LICHEN. (*λεχη*, or *λεχη*, a tetter or ring-worm.) Lichen is, by Dr. Wallen, defined an extensive eruption or papule affecting adults, connected with internal disorders, usually terminating in scurf, recurrent, not contagious. The varieties of lichen he considers under the denominations of *Lichen simplex*, *Lichen agrius*, *Lichen pilaris*, *Lichen lividus*, and *Lichen tropticus*.

1. The *Lichen simplex* usually commences with head-ache, flushing of the face, loss of appetite, general languor, and increased quickness of the pulse. Distinct red papulæ arise first about the cheeks and chin, or on the arms; and, in the course of three or four days, the same appearance takes place on the neck, body, and lower extremities, accompanied, with an unpleasant sensation of tingling, which is somewhat aggravated during the night. In about a week the colour of the eruption fades, and the cuticle begins to separate; the whole surface is at length covered with scurfy exfoliations, which are particularly large, and continue longest in the flexures of the joints. The duration of the complaint is seldom in any two cases alike; ten, fourteen, seventeen, or sometimes twenty days intervene betwixt the eruption and the renovation of the cuticle. The febrile state, or rather the state of irritation at the beginning of this disorder, is seldom considerable enough to confine the patient to the house. After remaining five or six

days, it is generally relieved on the appearance of the eruption. This, as well as some other species of the lichen, occurs about the beginning of summer, or in autumn, more especially affecting persons of a weak and irritable habit; hence women are more liable to it than men. Lichen simplex is also a frequent sequel of acute diseases, particularly fever and catarrhal inflammation, of which it seems to produce a crisis. In these cases the eruption has been termed, by medical writers, *scabies critica*. Many instances of it are collected under that title by Sauvages, Nosol. Method. Class x. Order 5. *Impetiginosa*.

2. The *Lichen agrius* is preceded by nausea, pain in the stomach, head-ache, loss of strength, and deep-seated pains in the limbs, with fits of coldness and shivering; which symptoms continue several days, and are sometimes relieved by the papulous eruption. The papulæ are distributed in clusters, or often in large patches, chiefly on the arms, the upper part of the breast, the neck, face, back, and sides of the abdomen; they are of a vivid red-colour, and have a redness, or some degree of inflammation, diffused round them to a considerable extent, and attended with itching, heat, and a painful tingling. Dr. Willan has observed, in one or two cases where it was produced from imprudent exposure to cold, that an acute disease ensued, with great quickness of the pulse, heat, thirst, pains of the bowels, frequent vomiting, head-ach, and delirium. After these symptoms had continued ten days, or somewhat longer, the patient recovered, though the eruption did not return. The diffuse redness connecting the papulæ, and the tendency to become pustular, distinguish the lichen agrius from the lichen simplex, and the other varieties of this complaint, in which the inflammation does not extend beyond the basis of the papulæ, and which terminates in scurf, or scales.

3. *Lichen pilaris*. This is merely a modification of the first species of lichen, and, like it, often alternates with complaints of the head, or stomach, in irritable habits. The peculiarity of the irruption is, that the small tubercles or asperities appear only at the roots of the hairs of the skin, being probably occasioned by an enlargement of their bulbs, or an unusual fulness of the blood-vessels distributed to them. This affection is distinguishable from the cutis asperina, by its permanency, by its red papulæ, and by the troublesome itching or tingling which attends it. If a part thus affected be violently rubbed, some of the papulæ enlarge to the size of wheals, but the tumour soon subsides again. The eruption continues more or less vivid for about ten days, and terminates, as usual, in small

exfoliations of the cuticle, one of which surrounds the base of each hair. This complaint, as likewise the lichen agrius, frequently occurs in persons accustomed to drink largely of spirituous liquors undiluted.

4. *Lichen lividus*. The papulæ characterizing this eruption are of a dark red, or livid hue, and somewhat more permanent than in the foregoing species of lichen. They appear chiefly on the arms and legs, but sometimes extend to other parts of the body. They are finally succeeded, though at very uncertain periods, by slight exfoliations of the cuticle, after which a fresh eruption is not preceded nor attended by any febrile symptoms. It principally affects persons of a weak constitution, who live on a poor diet, and are engaged in laborious occupations. Young persons, and often children, living in confined situations, or using little exercise, are also subject to the lichen lividus; and in them, the papulæ are generally intermixed with the petechiæ, or larger purple spots, resembling vibices. This circumstance points out the affinity of the lichen lividus with the purpura, or land-scurvy, and the connection is further proved by the exciting causes, which are the same in both complaints. The same method of treatment is likewise successful in both cases. They are presently cured by nourishing food, moderate exercise in the open air, along with the use of Peruvian and vitriolic acid, or the tincture of muriated steel.

5. *Lichen tropicus*. By this term is expressed the prickly heat, a papulous eruption, almost universally affecting Europeans settled in tropical climates. The prickly heat appears without any preceding disorder of the constitution. It consists of numerous papulæ, about the size of a small pin's head, and elevated so as to produce a considerable roughness on the skin. The papulæ are of a vivid red colour, and often exhibit an irregular form, two or three of them being in many places united together; but no redness or inflammation extends to the skin in the interstices of the papulæ.

LICHEN CANINUS. The systematic name of the ash-coloured ground liver-wort. See *Lichen cinereus terrestris*.

LICHEN CINEREUS TERRESTRIS. *Muscus caninus*. This cryptogamous plant, called ash-coloured ground liver-wort, and scientifically, *Lichen caninus* by Linnæus, has a weak, faint smell, and a sharpish taste. It was for a long time highly extolled as a medicine of singular virtue, in preventing and curing that dreadful disorder which is produced by the bite of rabid animals, but now deservedly forgotten. See *Pulvis antilyticus*.

LICHEN COCCIFERUS. See *Muscus pyridatus*.

LICHEN ISLANDICUS. The medicinal qualities of the lichen islandicus have lately been so well established at Vienna, that this plant is now admitted into the materia medica of the Edinburgh pharmacopœia. It is extremely mucilaginous, and to taste is bitter, and somewhat astringent. Its bitterness, as well as the purgative quality which its manifest, in its recent state, are in a great measure dissipated on drying, or may be extracted by a slight infusion in water, so that the inhabitants of Iceland convert it into a tolerably grateful and nutritive food. An ounce of this lichen, boiled a quarter of an hour in a pint of water, yielded seven ounces of a mucilage as thick as that procured by the solution of one pint of gum-arabic in three of water.

The medical virtues of this lichen were probably first learned from the Icelanders, who employ it in its fresh state as a laxative; but when deprived of this quality, and properly prepared, we are told that it is an efficacious remedy in consumptions, coughs, dysenteries, and diarrhœas. Scopolin seems to have been the first, who, of late years, called the attention of physicians to this remedy in consumptive disorders: and further instances of its success are related by Herz, Cramer, Tromsdorff, Ebeling, Paulisky, Stoll, and others, who bear testimony of its efficacy in most of the other complaints above-mentioned. Dr. Herz says, that since he first used the lichen in dysentery, he found it so successful, that he never had occasion to employ any other remedy; it must be observed, however, that cathartics and emetics were always repeatedly administered before he had recourse to the lichen, to which he also occasionally added opium. Dr. Chrichton informs us, that during seven months residence at Vienna, he had frequent opportunities of seeing the lichen islandicus tried in phthisis pulmonalis at the general hospitals, and confesses, "that it by no means answered the expectation he had formed of it." He adds, however, "from what I have seen, I am fully convinced in my own mind that there are only two species of this disease where this sort of lichen promises a cure. The two species I hint at are the phthisis hæmoptoica, and the phthisis pituitosa, or mucosa. In several cases of these, I have seen the patients so far get the better of their complaints as to be dismissed the hospital cured, but whether they remained long so or not I cannot take upon me to say." That this lichen strengthens the digestive powers, and proves extremely nutritious, there can be no doubt; but the great medicinal efficacy attributed to it at Vienna will not readily be credited at London. It is commonly given in the form of a decoction; an ounce and a half of the lichen being boiled in a quart of

milk. Of this a tea-cupful is directed to be drank frequently in the course of the day. If milk disagree with the stomach, a simple decoction of the lichen in water is to be used. Care ought to be taken that it be boiled over a slow fire, and not longer than a quarter of an hour.

LICHEN FIXIDATUS. The systematic name of the cup-moss. See *Muscus pyxidatus*.

LICHEN FLICATUS. The systematic name of the *Muscus arboreus*. See *Muscus arboreus*.

LICHEN PULMONARIUS. The systematic name of the officinal muscus pulmonarius quercinus. See *Pulmonaria arborea*.

LICHEN ROCCELLA. The systematic name of the roccella of the shops. See *Roccella*.

LICHEN SAXATILIS. The systematic name of the muscus crani humani. See *Umea*.

LIEN. (From λυος, soft, or smooth.) The spleen.

LIEN SINARUM. The faba Ægyptia.

LIENTERIA. (From λυος, smooth, εντερον, the intestine, and ρω, to flow.) The Latins call it *levitas intestinorum*. Lyentery. Dr. Cullen makes it a species of diarrhœa. See *Diarrhœa*.

LIFE. To live, may be defined the property of acting from an intrinsic power; hence the life of an animal body appears to be three-fold. 1. *Its chymical life*, which consists in that attraction of the elements, by which the vital principle, diffused through the solids and fluids, defends all the parts of the body from putrefaction. In this sense it may be said, that every atom of our body lives *chymically*, and that life is destroyed by putrefaction alone. 2. *Its physical life*, which consists in the irritability of the parts. This physical property remains for some time after death. Thus the heart or intestines, removed from the body whilst still warm, contract themselves on the application of a stimulus. In like manner the serpent, or eel, being cut into pieces, each part moves and palpitates for a long time afterwards. Hence these parts may be said to live *physically*, as long as they continue warm and soft. 3. *Its physiological life* consists in the action of inorganic parts proper to each, as the action of the heart and vessels; so that, these actions ceasing, the body is said to be physiologically dead. The physiological life ceases first, next the physical, and finally the chymical perishes.

LIGAMENT. (From λigo, to bind.) Ligaments are elastic and strong membranes connecting the extremities of the moveable bones. They are divided into *capsular*, which surround joints like a bag, and *connecting* ligaments. The use of the capsular ligaments is to connect the extremities of

the moveable bones, and prevent the efflux of synovia; the external and internal connecting ligaments strengthen the extremities of the moveable bones.

A Table of the principal Ligaments :

Ligaments of the lower jaw. The condyles of the lower jaw are connected with the articular sinuses of the temporal bone by two ligaments, the capsular and lateral ligament.

Ligaments of the occipital bone, and vertebrae of the neck. The condyles of the occipital bone are united with the articular depressions of the first vertebra by the capsular, broad, anterior, and posterior ligaments, the ligaments of the odontoid process, and ligamentum nuchæ.

Ligaments of the vertebrae. The vertebrae are connected together by means of their bodies and oblique processes. The bodies by a soft cartilaginous substance, and the processes by ligaments, viz. the transverse ligament of the first vertebra; the anterior and posterior common; the interspinous; the intertransverse; the intervertebral ligaments; the capsular ligaments of the oblique processes; and the ligaments of the last vertebra of the loins with the os sacrum.

Ligaments of the ribs. The posterior extremity of the ribs is united with the vertebra; the anterior with the sternum. The ligaments of the posterior extremity are, the capsular ligaments of the greater and lesser heads; the internal and external ligaments of the neck of the ribs; and a ligament peculiar to the last rib. The ligaments of the anterior extremity are, the capsular ligaments of the cartilages of the true ribs, and the ligaments of the ribs *inter se*.

Ligaments of the sternum. The ligaments connecting the three portions of the sternum to the ribs are, the membrana propria of the sternum; and the ligaments of the ensiform cartilage.

Ligaments of the pelvis. The ligaments which connect the ossa innominata with the os sacrum are, three ligamenta ileo sacra; two sacro-ischiatric ligaments; two transverse ligaments of the pelvis; the ligamentum obturans of the foramen ovale, and the ligamentum Poupartii, or inguinale. See *Pelvis*.

Ligaments of the os coccygis. The basis of the os coccygis is connected to the apex of the os sacrum, by the capsular and longitudinal ligaments.

Ligaments of the clavicle. The anterior extremity is connected with the sternum and first rib; and the posterior extremity with the acromion of the scapula, by the interclavicular, the capsular ligament, the ligamentum rhomboideum, and in the posterior extremity, the capsular ligament.

Ligaments of the scapula. The proper ligaments which connect the scapula with

the posterior extremity of the clavicle are, the conoid and trapezoid ligaments.

Ligaments of the humerus. The head of the humerus is connected with the glenoid cavity of the scapula by the capsular ligament.

Ligaments of the articulation of the cubit. The elbow-joint is formed by the inferior extremity of the humerus, and superior extremities of the ulnar and radius. The ligaments connecting these bones are, the capsular, the brachio-cubital, and the brachio-radial ligaments.

Ligaments of the radius. The radius is affixed to the humerus, cubit, and carpus, by peculiar ligaments, namely, the superior, inferior, oblique, and interosseous ligaments.

Ligaments of the carpus. The ligaments which connect the eight bones of the wrist together, and with the fore-arm and metacarpus, are, the capsular ligament of the carpus; the first and second transverse ligament; the oblique ligaments, and the capsular ligament proper to the bones of the carpus.

Ligaments of the metacarpus. The bones of the metacarpus are in part connected with the second row of bones of the carpus, and in part together, by the articular and interosseous ligaments.

Ligaments of the fingers. The fingers and phalanges are connected together, and with the metacarpus; and the thumb with the carpus, by the lateral ligaments of the fingers, and garment of the thumb with the os trapezium of the carpus.

Ligaments which keep the tendons of the muscles of the hand in their proper place. The ligaments which keep the tendons of the muscles of the hand in their place, are situated partly on the palm and partly on the back of the hand. In the back of the hand are, the external transverse ligament of the carpus, the vaginal, and the transverse ligaments of the extensor tendons. In the palm of the hand are, the internal transverse ligament of the carpus, the vaginal or crucial ligaments of the flexor tendons of the phalanges, and the accessory ligaments of the flexor tendons.

Ligaments of the articulation of the femur. The head of the os femoris is strongly annexed to the acetabulum of the os innominatum, by two very strong ligaments, the capsular ligament, and ligamentum teres, or restraining ligament.

Ligaments of the articulation of the knee. The knee-joint is formed by the condyles of the os femoris, head of the tibia and the patella. The ligaments are the capsular, the posterior, the external and the internal lateral ligaments, the crucial and the alar ligaments, the ligaments of the semilunar cartilages, and ligaments of the patella.

Ligaments of the fibula. The fibula is

connected with the tibia by means of the capsular ligament of the superior extremity, the interosseous ligament, and the ligaments of the inferior extremity.

Ligaments of the articulation of the tarsus. The inferior extremity of the tibia and fibula forms the cavity into which the astragalus of the tarsus is received. This articulation is effected by the anterior, middle, and posterior ligament of the fibula, the ligamentum tibiae deltoïdes, the capsular ligament, and the ligaments proper to the bones of the tarsus.

Ligaments of the metatarsus. The bones of the metatarsus are connected in part together, and in part with the tarsus, by means of the capsular ligament, the articular ligaments, the transverse ligaments in the back and sole of the foot, and the interosseous ligaments of the metatarsus.

Ligaments of the toes. The phalanges of the toes are united partly together, and partly with the metatarsus, by the capsular and lateral ligaments.

Ligaments which retain the tendons of the muscles of the foot in their proper place. These ligaments are found partly in the back, and partly in the sole of the foot. They are the vaginal ligament of the tibia, the transverse or crucial ligaments of the tarsus, the ligaments of the tendon of the peronei muscles, the lacinated ligament, the vaginal ligament of the extensor muscle and flexor pollicis, the vaginal ligaments of the flexor tendons, the accessory ligaments of the flexor tendons, and the transverse ligaments of the extensor tendons.

LIGAMENTUM ANNULARE. The annular ligament. A strong ligament on each ankle and each wrist.

LIGAMENTUM ARTERIOSUM. The ductus arteriosus of the foetus becomes a ligament after birth, which is so called.

LIGAMENTUM CILIARE. Behind the uvea of the human eye there arise, out of the choroid membrane, from the ciliary circle, white complicated striz, covered with a black matter. The fluctuating extremities of these striz are spread abroad even to the crystalline lens, upon which they lie, but are not affixed. Taken together they are called *ligamentum ciliare*.

LIGAMENTUM DENTICULATUM. A small ligament supporting the spinal marrow.

LIGAMENTUM FALOPII. The ligamentum rotundum uteri was so called.

LIGAMENTUM INTEROSSEUM. The ligament uniting the radius and ulna, and the tibia and fibula.

LIGAMENTUM LATUM. Broad ligament of the liver and uterus. See *Liver*, and *Uterus*.

LIGAMENTUM NUCHÆ. A strong ligament of the neck, which proceeds from one spinous process to another.

LIGAMENTUM OVARIUM. The thick round portion of the broad ligament of the uterus, by which the ovary is connected with the uterus. The ancients supposed this was hollow, to convey the female semen into the uterus.

LIGAMENTUM POUPARTII. Fallopiian ligament. Poupart's ligament. A ligament extending from the anterior superior spinous process of the ilium to the crista of the os pubis.

LIGAMENTUM ROTUNDUM. The round ligament of the uterus. See *Uterus*.

LIGATURE (From *ligo*, to bind.) A thread, or silk, of various thickness, covered with white wax, for the purpose of tying arteries, or veins, or other parts. They should be round and very firm, so as to allow being tied with some force, without risk of breaking.

The immediate effect of a tight ligature on an artery is to cut through its middle and internal coats, a circumstance that tends very much to promote the adhesion of the opposite sides of the vessel to each other. Hence the form and mode of applying a ligature to an artery should be such, as are most certain of dividing the above coats of the vessel in the most favourable manner. A broad flat ligature does not promise to answer the purpose in the best manner; because it is scarcely possible to tie it smoothly round the artery, which is very likely to be thrown into folds, or to be puckered by it, and consequently to have an irregular bruised wound made in its middle and internal coats. A ligature of an irregular form is likely to cut through these coats more completely at some parts than at others; and if it does not perfectly divide them, no adhesion can take place, and secondary hæmorrhage will follow. A fear of tying the ligature too tight may often lead to the same consequences.

LIGHT. *Lux.* The nature of light has occupied much of the attention of philosophers, and numerous opinions have been entertained concerning it. It has been sometimes considered as a distinct substance, at other times as a quality; sometimes as a cause; frequently as an effect; by some it has been considered as a compound, by others as a simple substance. Philosophers of the present day are not agreed as to the independent existence of light, or the cause by which we see.

Nature of Light.

Light is that which proceeds from any body producing the sensation of vision, or perception of other bodies, by depicting an image of external objects on the retina of the eye. Hence it announces to animals the presence of the bodies which surround them, and enables them to distinguish these bodies into transparent, opaque,

and coloured. These properties are so essentially connected with the presence of light, that bodies lose them in the dark, and become undistinguishable.

Light is regarded by philosophers as a substance consisting of a vast number of exceedingly small particles, which are actually projected from luminous bodies, and which probably never return again to the body from which they were emitted.

It is universally expanded through space. It exerts peculiar actions, and is obedient to the laws of attraction, and other properties of matter.

Explanation of certain terms of Light.

In order to facilitate the doctrine of light, we shall shortly explain a few terms made use of by philosophers when treating of it; namely:

A *ray of light* is an exceedingly small portion of light as it comes from a luminous body.

A *medium* is a body which affords a passage for the rays of light.

A *beam of light* is a body of parallel rays.

A *pencil of rays* is a body of diverging and converging rays.

Converging rays are rays which tend to a common point.

Diverging rays are those which come from a point, and continually separate as they proceed.

The rays of light are *parallel*, when the lines which they describe are so.

The *radiant point* is the point from which diverging rays proceed.

The *focus* is the point to which the converging rays are directed.

Sources of Light.

Light is emitted from the sun, the fixed stars, and other luminous bodies. It is produced by percussion, during electrification, combustion, and in various other chymical processes.

Why the sun and stars are constantly emitting light is a question, which probably will for ever baffle human understanding.

The light emitted during combustion, exists previously, either combined with the combustible body, or with the substance which supports the combustion. The light liberated during chymical action formed a constituent part of the bodies which act on each other.

Chymical Properties of Light.

The chymical effects of light have much engaged the attention of philosophers. Its influence upon animal, vegetable, and other substances, is as follows:

Effects of Light on Vegetables.

Every body knows that most of the diaceous flowers follow the sun in his course; that they attend him to his evening retreat, and meet his rising lustre in the morning with the same unerring law. It is also well known that the change of position in

the leaves of plants, at different periods of the day, is entirely owing to the agency of light, and that plants which grow in windows, in the inside of houses, are, as it were, solicitous to turn their leaves towards the light. Natural philosophers have long been aware of the influence of light on vegetation. It was first observed that plants growing in the shade, or darkness, are pale and without colour. The term *etiolation* has been given to this phenomenon, and the plants, in which it takes place, are said to be *etiolated*, or *blanched*. Gardeners avail themselves of the knowledge of this fact, to furnish our tables with white and tender vegetables. When the plants have attained a certain height, they compress the leaves, by tying them together, and by these means (or by laying earth over them) deprive them of the contact of light: and thus it is that our white celery, lettuce, cabbages, endive, &c. are obtained. For the same reason, wood is white under the green bark; roots are less coloured than plants, some of them alter their taste, &c.; they even acquire a deleterious quality when suffered to grow exposed to light. Potatoes are of this kind. Herbs that grow beneath stones, or in places utterly dark, are white, soft, aqueous, and of a mild and insipid taste. The more plants are exposed to the light, the more colour they acquire. Though plants are capable of being nourished exceedingly well in perfect obscurity, and in that state they even grow much more rapidly than in the sun, (provided the air that surrounds them is fit for vegetation,) they are colourless, and unfit for use.

Professor Davy found, by experiment, that red rose-trees, carefully excluded from light, produce roses almost white. He likewise ascertained that this flower owes its colour to light entering into its composition; that pink, orange, and yellow flowers imbibe a smaller portion of light than red ones, and that white flowers contain no light.

But vegetables are not only indebted to the light for their colour: taste and odour are likewise derived from the same source.

Light contributes greatly to the maturity of fruits and seeds. This seems to be the cause, why, under the burning sun of Africa, vegetables are in general more odouriferous, of a stronger taste, and more abounding with resin. From the same cause it happens, that hot climates seem to be the native countries of perfumes, odouriferous fruits, and aromatic resins.

The action of light is so powerful on the organism of vegetables, as to cause them to pour forth torrents of pure air from the surface of their leaves into the atmosphere, while exposed to the sun; whereas, on the contrary, when in the shade, they emit an

air of a noxious quality. Take a few handful of fresh-gathered leaves of mint, cabbage, or any other plant; place them in a bell-glass, filled with fresh water, and invert it into a basin with the same fluid. If the whole be then exposed to the direct rays of the sun, small air bubbles will appear on the surface of the leaves, which will gradually grow larger, and at last detach themselves, and become collected at the surface of the water. This is oxygen gas, formerly called vital air.

All plants do not emit this air with the same facility; there are some which yield it the moment the sun acts upon them, as the jacobaea, or rag-wort, lavender, peppermint, and some other aromatic plants. The leaves afford more air when attached to the plant than when gathered; the quantity is also greater, the fresher, and sounder they are, and if full grown and collected during dry weather. Green plants afford more air than those which are of a yellowish or white colour. Green fruits afford likewise oxygen gas; but it is not so plentifully furnished by those which are ripe. Flowers in general render the air noxious. The nasturtium indicum, in the space of a few hours, gives out more air than is equal to the bulk of all its leaves.

On the contrary, if a like bell-glass, prepared in the same manner, be kept in the dark, another kind of air will be disengaged, of an opposite quality.

There is not a substance, which, in well-closed glass vessels, and exposed to the sun's light, does not experience some alteration.

Camphor kept in glass bottles, exposed to light, crystallizes, or vegetates, into the most beautiful symmetrical figures, on that side of the glass which is exposed to the light.

Yellow wax, exposed to the light, loses its colour, and becomes bleached. Gum guaiacum, reduced to powder, becomes green, on exposure to light. Vegetable colours, such as those of saffron, logwood, &c. become pale, or white, &c.

2. Effects of Light on Animals.

The human being is equally dependent on the influence of light.

Animals in general droop when deprived of light, they become unhealthy, and even sometimes die. When a man has been long confined in a dark dungeon, (though well aired,) his whole complexion becomes sallow; pustules, filled with aqueous humours, break out on his skin; and the person who has been thus deprived of light becomes languid, and frequently dropsical.

Worms, grubs, and caterpillars, which live in the earth, or in wood, are of a whitish colour; moths, and other insects of the night, are likewise distinguishable

from those which fly by day by the want of brilliancy in their colour. The difference between those insects, in northern and southern parts, is still more obvious.

The parts of fish which are exposed to light, as the back, fins, &c. are uniformly coloured, but the belly, which is deprived of light, is white in all of them.

Birds which inhabit the tropical countries have much brighter plumage than those of the north. Those parts of the birds which are not exposed to the light are uniformly pale. The feathers on the belly of a bird are generally pale, or white; the back, which is exposed to the light, is almost always coloured; the breast, which is particularly exposed to light in most birds, is brighter than the belly.

Butterflies, and various other animals of equatorial countries, are brighter coloured than those of the polar regions. Some of the northern animals are even darker in summer and paler in winter.

3 Effects of Light on other Substances.

Metallic oxyds become combustible when exposed to light. Acids are decomposed by its contact, and various other substances change their nature.

Light carbonated hydrogen. See Carbonated hydrogen, light.

LIGNUM AGALLOCHI VERI. See *Lignum aloes*.

LIGNUM ALOES. *Lignum agallochi veri*. *Lignum calambac*. *Lignum aspalathi*. *Xylo aloes*. *Agallochum*. *Calambac*. Aloes-wood. The tree whose wood bears this name is not yet scientifically known. It is imported from China in small, compact, ponderous pieces, of a yellow rusty brown colour, with black or purplish veins, and sometimes of a black colour. It has a bitterish resinous taste, and a slight aromatic smell. It is used to fumigate rooms in eastern countries.

LIGNUM ASPALATHI. See *Lignum aloes*.

LIGNUM CALAMBAC. See *Lignum aloes*.

LIGNUM CAMPECHENSE. (*Campechensis*; so called because it was brought from Campeachy, in the bay of Honduras.) *Lignum campechianum*. *Lignum campeccanum*. *Lignum indicum*. *Lignum saphan*. Logwood. The wood of the *Hæmatoxylum campechyanum* of Linnæus; it is of a solid texture and of a dark red colour. It is imported principally as a substance for dyeing, cut intounks and logs of about three feet in length: of these pieces the largest and thickest are preserved, as being of the deepest colour. Logwood has a sweetish subadstringent taste, and no remarkable smell: it gives a purplish red tincture both to watery and spirituous infusions, and tinges the stools, and sometimes the urine, of the same colour. It is employed medicinally as an adstringent and corroborant.

In diarrhoeas it has been found peculiarly efficacious, and has the recommendation of some of the first medical authorities; also in the latter stages of dysentery, when the obstructing causes are removed, to obviate the extreme laxity of the intestines, usually superinduced by the repeated dejections. In the form of decoction the proportion is two ounces to 2 lbs. of fluid, reduced by boiling to one. An extract is ordered in the pharmacopœias. The dose from ten to forty grains.

LIGNUM INDICUM. See *Guaiacum*.

LIGNUM MOLUCCENSE. See *Lignum pavana*.

LIGNUM NEPHRITICUM. Nephritic wood. The wood of the *Gulandina moringa*; *intermis foliis sub-bipinnatis, foliis inferioribus ternatis* of Linnæus, which also affords the *maz bean*. It is brought from America in large, compact, ponderous pieces, without knots, the outer part of a whitish or pale yellowish colour, the inner of a dark brown, or red. When rasped, it gives out a faint aromatic smell. It is never used medicinally in this country, but stands high in reputation abroad, against difficulties of making urine, nephritic complaints, and most disorders of the kidneys and urinary passages.

LIGNUM PAVANÆ. *Lignum pavanum*. *Lignum moluccense*. The wood of the *Croton tiglium*; *foliis ovalis glabris acuminatis serratis, caule arboræ* of Linnæus, which affords the *grana tiglii*. It is of a light spongy texture, white within, but covered with a greyish bark; and possesses a pungent, caustic taste, and a disagreeable smell. It is said to be useful as a purgative in hydropical complaints.

LIGNUM RHODIUM. See *Rhodium lignum*.

LIGNUM SANCTUM. See *Guaiacum*.

LIGNUM SANTALI RUBRI. See *Santalum rubrum*.

LIGNUM SAPPAN. See *Lignum campechense*.

LIGNUM SERPENTUM. The wood of the *Ophioxylum serpentinum* of Linnæus. It is said to be an alexipharmic.

LIGUSTICUM LEVISTICUM. The systematic name of lovage. See *Levisticum*.

LIGUSTRUM. (From *lige*, to bind; so named from its use in making bands.) 1. The name of a genus of plants in the Linnean system. Class, *Diandria*. Order, *Monogynia*.

2. The pharmacopœial name of the herb privet.

LILIASO. (Dim. of *lilium*, the lily; so named from the resemblance of its flower to that of a lily.) *Lilium*. Spiderwort; formerly said to be alexipharmic and carminative.

LILIUM. (From *lilis*, smooth, graceful; so named from the beauty of its leaf.)

The name of a genus of plants in the Linnean system. Class, *Hexandria*. Order, *Monogynia*. The lily.

LILIUM ALBUM. The white lily. The roots of the common white lily, *Lilium candidum* of Linnæus:—*foliis sparsis, corollis campanulatis, intus glabris*, are directed by the Edinburgh pharmacopœia; they are extremely mucilaginous, and chiefly used boiled in milk and water, in emollient and suppurating cataplasms, to inflammatory tumours. These lily-roots afford a good substitute, in times of scarcity, for bread. The distilled water has been sometimes used as a cosmetic.

LILIUM CANDIDUM. The systematic name of the white lily. See *Lilium album*.

LILIUM CONVALLIUM. *Convallaria*. *Mai-anthemum*. *Convallaria maialis*. Lily of the valley. May-lily. The flowers of this plant, *Convallaria majalis*, *scapo nudo* of Linnæus, have a penetrating bitter taste, and are given in nervous and catarrhal disorders. When dried and powdered they prove strongly purgative. Watery or spirituous extracts made from them, given in doses of a scruple, or drachm, act as gentle stimulating aperients and laxatives, and seem to partake of the purgative virtue as well as the bitterness of aloes. The roots, in the form of tincture, or infusion, act as a sternutatory when snuffed up the nose, and as a laxative or purgative when taken internally.

Lily, May. See *Lilium convallium*.

Lily, white. See *Lilium album*.

Lily, water. See *Nymphaea alba* and *Nymphaea lutea*.

Lily of the valley. See *Lilium convallium*.

LIMATURA FERRI. Steel filings are considered as possessing stimulating and strengthening qualities, and are exhibited in worm cases, ataxia, leucorrhœa, diarrhœa, chlorosis, &c.

LIMAX. (From *limus*, slime; so named from its sliminess.) *Cochlea terrestris*. The snail. This animal abounds with a viscid slimy juice, which is readily given out, by boiling, to milk or water, so as to render them thick and glutinous. These decoctions are apparently very nutritious and demulcent, and are recommended in consumptive cases and emuciations.

LIME. *Calx*. An earth found in great abundance in nature, though never pure, or in an uncombined state. It is always united to an acid, and very frequently to the carbonic acid, as in chalk, common lime-stone, marble, calcareous spar, &c. It is contained in the waters of the ocean; it is found in vegetables; and is the basis of the bones, shells, and other hard parts of animals. Its combination with sulphuric acid is known by the name of sulphate of lime (*gypsum*, or plaster of Paris.) Com-

bined with fluoric acid it constitutes fluatc of lime, or Derbyshire spar.

Properties.—Lime is in solid masses, of a white colour, moderately hard, but easily reducible to powder. Its taste is bitter, urinous, and burning. It changes blue cabbage juice to a green. It is unalterable by the heat of our furnaces. It splits and falls into powder in the air and loses its strong taste. It is augmented in weight and in size by slowly absorbing water from the atmosphere. Its specific gravity is 2.3. It combines with phosphorus by heat. It unites to sulphur both in the dry and humid way. It absorbs sulphurated hydrogen gas. It unites with some of the metallic oxyds. Its slaking by water is attended with heat, hissing, splitting, and swelling up, while the water is partly consolidated and partly converted into vapour; and the lime is reduced into a very voluminous dry powder, when it has been sprinkled with only a small quantity of water. It is soluble when well prepared in 300 parts of water. It unites to acids. It renders silex and alumine fusible, and more particularly these two earths together.

Method of obtaining Lime.—Since the carbonic acid may be separated from the native carbonate of lime, this becomes a means of exhibiting the lime in a state of tolerable purity. For this purpose introduce into a porcelain or earthen retort, or rather into a tube of green glass, well coated over with lute, and placed across a furnace, some powdered Carara marble, or oyster-shell powder. Adapt to its lower extremity a bent tube of glass, conveyed under a bell. If we then heat the tube, we obtain carbonic acid gas; and lime will be found remaining in the tube, or retort.

The burning of lime in the large way, depends on the disengagement of the carbonic acid by heat; and, as lime is infusible in our furnaces, there would be no danger from too violent a heat, if the native carbonate of lime were perfectly pure; but as this is seldom the case, an extreme degree of heat produces a commencement of vitrification in the mixt stone, and enables it to preserve its solidity, and it no longer retains the qualities of lime, for it is covered with a sort of crust, which prevents the absorption of the water when it is attempted to be slaked. This is called over-burnt lime.

In order to obtain lime in a state of great purity, the following method may be had recourse to.

Take Carara marble, or oyster-shells; reduce them to powder, and dissolve the powder in pure acetous acid; precipitate the solution by carbonate of ammonia. Let the precipitate subside, wash it repeatedly in distilled water, let it dry, and then expose it to a white heat for some hours.

The acetous acid, in this operation, unites to the lime, and forms acetite of lime, disengaging at the same time the carbonic acid, which flies off in the gaseous state: on adding to the acetite of lime carbonate of ammonia, acetite of ammonia and an artificial carbonate of lime are formed; from the latter the carbonic acid is again expelled, by exposure to heat, and the lime is behind, in a state of perfect purity.

Lime trees. See *Tilia*.

Lime-water. See *Liquor calcis*.

LIMES. A fruit like a small lemon, the juice of which is a very strong acid, and very much used in the making of punch. Externally, the same acid is applied in the cutaneous affections of warm climates, and also as a remedy against the pains that precede the appearance of yaws.

LIMON. (Hebrew.) *Limonia mala*.

Malus limonia acida. Citrea malus. Citrus.

The lemon. The tree which affords this fruit is the *Citrus medica* of Linnæus:—*petiolo hnearibus*: a native of the upper part of Asia, but cultivated in Spain, Portugal, and France. The juice, which is much more acid than that of the orange, possesses similar virtues. It is always preferred where a strong vegetable acid is required. Saturated with the fixed vegetable alkali, it forms the citrat of potash, which is in frequent extemporaneous use in febrile diseases, and by promoting the secretions, especially that of the skin, proves of considerable service in abating the violence of fever. This medicine is also often employed to restrain vomiting. As an antiscorbutic, the citric acid is also very generally taken on board ships destined for long voyages; but even when well depurated of its mucilaginous parts, it is found to spoil by long keeping. To preserve it in purity for a considerable length of time, it is necessary that it should be brought to a highly concentrated state, and for this purpose it has been recommended to expose the juice to a degree of cold sufficient to congeal the aqueous and mucilaginous parts. After a crust of ice is formed, the juice is poured into another vessel; and, by repeating this process several times, the remaining juice, it is said, has been concentrated to eight times its original strength, and kept, without suffering any material change, for several years. Whytt found the juice of lemons to allay hysterical palpitations of the heart, after various other medicines had been experienced ineffectual; and this juice, or that of oranges, taken to the quantity of four or six ounces in a day, has sometimes been found a remedy in the jaundice. The exterior rind of the lemon is a very grateful, aromatic bitter, not so hot as orange-peel, and yielding in distillation a

less quantity of oil, which is extremely light, almost colourless, and generally brought from the southern parts of Europe, under the name of *Essence of Lemons*. The lemon-peel, though less warm, is similar in its qualities to that of the orange, and is employed with the same intentions. The Pharmacopœias direct a syrup of the juice, *syrupus limonis*, and the peel enters into vinous and aqueous bitter infusions; it is also ordered to be candied; and the essential oil is an ingredient in the *spiritus ammoniæ compositus*, and other formulæ.

LIMONIUM. (From *limon*, a green field; so called from its colour.) Sea-lavender. Astringent.

LIMONIUM. (From *limon*, a green-field; so called from the colour of its unripe fruit.) The lemon-tree, or citrus medica.

LINAGROSTIS. (From *linon*, cotton, and *agrostis*, grass; so called from the softness of its texture.) Cotton-grass.

LINANGINA. (From *linum*, flax, and *ango*, to strangle; so called, because if it grows among flax or hemp, it twists round it, and chokes it.) The herb dodder.

LINARIA. (From *linum*, flax, named from the resemblance of its leaves to those of flax.) *Oxyria*. *Urtaria*. Common toad-flax. *Antirrhinum linaria* of Linnæus:—*foliis lanceolato linearibus confertis, caule erecto, spicis terminalibus sessilibus, floribus, imbricatis*. A perennial indigenous plant, common in barren pastures, hedgers, and the sides of roads, flowering from July to to September. The leaves have a bitterish and somewhat saline taste, and, when rubbed between the fingers, have a faint smell, resembling that of elder. They are said to be diuretic and cathartic, and in both characters to act powerfully, especially in the first; hence the name *urinalis*. They have been recommended in dropsies, and other disorders requiring powerful evacuations. The linaria has also been used as a resolvent in jaundice, and such diseases as were supposed to arise from visceral obstructions. But the plant has been chiefly valued for its effects when externally applied, especially in hæmorrhoidal affections, for which both the leaves and flowers have been employed in various forms of ointment, fomentation, and poultice. Dr. Wolph first invented an ointment of this plant for the piles. The Landgrave of Hesse, to whom he was physician, constantly interrogated him, to discover its composition; but Wolph obstinately refused, till the prince promised to give him a fat ox annually for the discovery: hence, to the following verse, which was made to distinguish the linarian from the esculet, viz.

"*Escula lactescit, sine lacte linaria crescit,*

the hereditary) Marshal of Hesse added.

"*Esula nil nobis, sed dat linaria taurum.*"
LINCTUS. (From *lingo*, to lick.) *Lohoc*. *Eclegma*. *Elezis*. *Elegma*. *Eclectes*. *Eclectes*. *Ilinctus*. Lick and linctive. A term in pharmacy, that is generally applied to a soft and somewhat oily substance, of the consistence of honey, which is licked off the spoon, it being too solid and adhesive to be taken otherwise.

LINEA ALBA. (From *linum*, a thread, and *album*, white: so called from its appearance and colour.) *Linea centralis*. An aponeurosis that extends from the scrobiculus cordis straight down to the navel, and from thence to the pubis. It is formed by the tendinous fibres of the internal oblique ascending and the external oblique descending muscles, and the transversalis, interfaced with those of the opposite side.

LINEÆ SEMILUNARES. The lines which bound the outer margin of the recti muscles, formed by the union of the abdominal tendons.

LINEÆ TRANSVERSÆ. Lines which cross the recti muscles.

LINGUA. (From *lingo*, to lick up. The tongue. See *tongue*.)

LINGUA AVIS. The seeds of the *Fragaria*, or ash, are so called, from their supposed resemblance to a bird's tongue.

LINGUA CANINA. So called from the resemblance of its leaves to a dog's tongue. See *Cynoglossum*.

LINGUA CERVINA. See *Scolopendrium*.

LINGUALIS. (*Lingualis*, sc. *musculus*; from *lingua*, a tongue.) *Basio-glossus* of Cowper. A muscle of the tongue. It arises from the root of the tongue laterally, and runs forward between the hyo-glossus and genio-glossus, to be inserted into the tip of the tongue, along with part of the stylo-glossus. Its use is to contract the substance of the tongue, and to bring it backward.

LINIMENT. (*Linimentum*; from *lino*, to anoint.) An oily substance, of a mediate consistence between an ointment and oil, but so thin as to drop.

The following are some of the most approved forms.

LINIMENTUM ERUGINIS. Liniment of verdigris, formerly called *oxymel æruginis*, *mel Ægyptiacum*, and *unguentum Ægyptiacum*. "Take of verdigris, powdered, an ounce; vinegar, seven fluid-ounces; clarified honey, fourteen ounces." Dissolve the verdigris in the vinegar, and strain it through a linen cloth; having added the honey gradually, boil it down to a proper consistence.

LINIMENTUM AMMONIÆ CARBONATIS. Linimentum of carbonat of ammoniac, formerly called *linimentum ammoniæ* and *linimentum volatile*. "Take of solution

of subcarbonate of ammonia, a fluid-ounce; olive oil, three fluid-ounces." Shake them together until they unite. A stimulating liniment, mostly used to relieve rheumatic pains, bruises, and paralytic numbness.

LINIMENTUM AMMONIÆ FORTIUS. Strong liniment of ammonia. "Take of solution of ammonia, a fluid-ounce; olive oil, two fluid-ounces." Shake them together until they unite. A more powerful stimulating application than the former, acting as a rubefacient. In pleurodynia, indolent tumours, stiffness of the joints, and arthritic pains, it is to be preferred to the milder one.

LINIMENTUM AQUÆ CALCIS. Liniment of lime-water. "Take of lime-water, olive-oil, of each eight ounces; rectified spirit of wine, one ounce." Mix. This has been long in use as an application to burns and scalds.

LINIMENTUM CAMPHORÆ. Camphor liniment. "Take of camphor, half an ounce. Olive oil, two fluid-ounces." Dissolve the camphor in the oil.

LINIMENTUM CAMPHORÆ COMPOSITUM. Compound camphor liniment. "Take of camphor, two ounces; solution of ammonia, six fluid-ounces; spirit of lavender, a pint." Mix the solution of ammonia with the spirit in a glass retort; then, by the heat of a slow fire, distil a pint. Lastly, in this distilled liquor, dissolve the camphor. An elegant and useful stimulant application in paralytic, spasmodic, and rheumatic diseases. Also for bruises, sprains, rigidities of the joints, incipient chilblains, &c. &c.

LINIMENTUM HYDRARGYRI COMPOSITUM. Compound mercurial liniment. "Take of strong mercurial ointment, prepared lard, of each four pounds; camphor, an ounce; rectified spirit, fifteen minims; solution of ammonia, four fluid-ounces." First powder the camphor, with the addition of the spirit, then rub it with the mercurial ointment and the lard; lastly, add gradually the solution of ammonia, and mix the whole together. An excellent formula for all surgical cases, in which the object is to quicken the action of the absorbents, and gently stimulate the surfaces of parts. It is a useful application for diminishing the indurated state of particular muscles, a peculiar affection every now and then met with in practice; and it is peculiarly well calculated for lessening the stiffness and chronic thickening often noticed in the joints. If it be frequently or largely applied, it affects the mouth more rapidly than the mercurial ointment.

LINIMENTUM OPIATUM. A resolvent anodyne embrocation, adapted to remove indolent tumours of the joints, and those weaknesses which remain after strains and chilblains before they break.

LINIMENTUM SAPONIS COMPOSITUM. Compound soap liniment. *Linimentum saponis.* "Take of hard soap, three ounces; camphor, an ounce; spirit of rosemary, a pint." Dissolve the camphor in the spirit, then add the soap, and macerate in the heat of a sand-bath, until it be melted. The basis of this form was first proposed by Riverius, and it is now commonly used under the name of opodeldoc. This is a more pleasant preparation, to rub parts affected with rheumatic pains, swellings of the joints, &c. than any of the foregoing, and at the same time not inferior, except where a rubefacient is required.

LINIMENTUM SAPONIS CUM OPIO. Soap liniment, with opium. "Take of compound soap liniment, six ounces; tincture of opium, two ounces." Mix. For dispersing indurations and swellings, attended with pain, but no acute inflammation.

LINIMENTUM TEREBINTHINÆ. Turpentine liniment. "Take of resin of cerate, a pound; oil of turpentine, half a pint." Add the oil of turpentine to the cerate, previously melted, and mix. This liniment is very commonly applied to burns, and was first introduced by Mr. Kentish, of Newcastle.

LINIMENTUM TEREBINTHINÆ VITRIOLICUM. Vitriolic liniment of turpentine. "Take of olive oil, ten ounces; oil of turpentine, four ounces; vitriolic acid, three drachms." Mix. This preparation is said to be efficacious in chronic affections of the joints, and in the removal of long existing effects of sprains and bruises.

Liniment of ammonia. See *Linimentum ammoniæ.*

Liniment of camphor. See *Linimentum camphoræ.*

Liniment of mercury. See *Linimentum hydrargyri.*

Liniment of verdigris. See *Linimentum æruginis.*

Liniment of turpentine. See *Linimentum terebinthinæ.*

LINNÆA. (So named in honour of Linnæus.) The name of a genus of plants in the Linnæan system. Class, *Didynamia*. Order, *Angiospermia*.

LINNÆA BOREALIS. The systematic name of the plant named in honour of the immortal Linnæus, which has a bitter substringent taste, and is used in some places in the form of fomentation to rheumatic pains, and an infusion with milk is much esteemed in Switzerland in the cure of sciatica.

LINOSPERMUM. (From *linos*, flax, and *sperma*, seed.) Linseed.

LIN-SEED. The fruit of the flax plant, or linum, is much used in medicine. Its qualities are mucilaginous and oily. It is lubricating and emollient. It is employed

in decoction, or infusion, in heat of urine, &c. in the form of clyster, in tenesmus, in cataplasm, in quincy, and other complaints. The proportion of the seeds in the decoction are, an ounce to a pound of water. See *Linum*.

Lint. See *Linteum*.

LINTEUM. *Lint.* A soft woolly substance, made by the scraping of old linen cloth, and employed in surgery as the common dressing in all cases of wounds and ulcers, either simply or covered with different unctuous substances.

LINUM. (From *λεπτός*, soft, smooth; so called from its soft, smooth, texture.)

1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Pentagynia*.

2. The pharmacopœia name of the common flax. *Linum sylvestre*. *Linum usitatissimum* of Linnæus:—*calycibus capsulæque mucronatis, petalis crenatis, foliis lanceolatis alternis, caule subulitario*. The seeds of this useful plant, called linseed, have an unctuous, mucilaginous, sweetish taste, but no remarkable smell; on expression they yield a large quantity of oil, which, when carefully drawn, without the application of heat, has no particular taste or flavour: boiled in water, they yield a large proportion of a strong flavourless mucilage, which is in use as an emollient or demulcent in coughs, hoarsenesses, and pleuritic symptoms, that frequently prevail in catarrhal affections; and it is likewise recommended in nephritic pains and stranguries. The meal of the seeds is also much used, externally, in emollient and maturating cataplasms. The expressed oil is an official preparation, and is supposed to be of a more healing and balsamic nature than the other oils of this class: it has, therefore, been very generally employed in pulmonary complaints, and in colics and constipations of the bowels. The cake which remains after the expression of the oil contains the farinaceous part of the seed, and is used in fattening cattle, under the name of oil cake.

LINUM CATHARTICUM. *Linum minimum*. *Chamaedon*. Purging flax, or milk-mountain. This small plant, *Linum catharticum*, *foliis oppositis ovato-lanceolatis, caule dichotomo corollis acutis* of Linnæus, is an effectual and safe cathartic. It has a bitterish and disagreeable taste. A handful infused in half a pint of boiling water is the dose for an adult.

LINUM USITATISSIMUM. The systematic name of the common flax. See *Linum*.

LIPARIS. (From *λεπτός*, fat; so named from its unctuous quality.) See *Pinguicula*.

LIPAROCLE. (From *λεπτός*, fat, and *κλέα*, a tumour.) That species of sarco-

cele in which the substance constituting the disease is fat.

LIPOMA. (From *λεπτός*, fat.) A solitary, soft, unequal, indolent tumour, arising from a luxuriancy of adeps in the cellular membrane. The adipose structure forming the tumour is sometimes diseased towards its centre, and more fluid than the rest. At other times it does not appear to differ in any respect from adipose membrane, except in the enlargement of the cells containing the fat. These tumours are always many years before they arrive at any size.

LIPOTECIA. (From *λεπτός*, to leave, and *ψυχή*, the soul, or life.) A swoon.

LIPOTHYMIA. (From *λεπτός*, to leave, and *θυμός*, the mind.) *Detritum animi et anime*. *Defectio animi; dissolutio*. *Exanimatio*. *Syncope*. *Asphyxia*. *Vitium lapsus*. The greatest degree is called *Apoplechia*. *Asphyxia*. *Eclamps*. Fainting. Dr. Cullen makes it a genus of disease in the class *Neuroses*, order *Adynamia*, which he defines: the motion of the heart diminished, or at rest for some time. He distinguishes it into three species. See *Syncope*.

LIPPITUDO. (From *lippus*, bleary-eyed.) *Epiphora*. *Xerophthalmia*. Blearedness. An exudation of a puriform humour from the margin of the eye-lids. The proximate cause is a deposition of acrimony on the glandulæ meibomianæ in the margin of the eye-lids. This humour in the night glues the tarsi of the eye-lids together. The margins of the eye-lids are red and tumefy are irritated, and excite pain. An ophthalmia, fistula lachrymalis, and sometimes an ecropium, are the consequences. The species of the lippitudo are,

1. *Lippitudo infantum*, which is familiar to children, particularly of an acrimonious habit. The lippitudo of infants is mostly accompanied with tinea, or some scabby eruption, which points out that the disease originates, not from a local, but general or constitutional, affection.

2. *Lippitudo adultorum* or *senilis*. This arises from various acrimones, and is likewise common to hard drinkers.

3. *Lippitudo venerea*, which arises from a suppressed gonorrhœa, or fluor albus, and is likewise observed in children born of parents with venereal complaints.

4. *Lippitudo scrophulosa*, which accompanies other scrophulous symptoms.

5. *Lippitudo scorbutica*, which affects the scorbutic.

LIPTRIA. (From *λεπτός*, to leave, and *πύρ*, heat.) A sort of fever, where the heat is drawn to the inward parts, while the extremities are cold.

LIQUIDAMBER. The name of a genus of plants in the Linnæan system.

LIQUIDAMBER STYRACIFLUA. The

systematic name of the tree which affords both the liquid-amber and liquid storax.

See *Liquidambra*.

LIQUIDAMBRA. (From *liquidus* and *amber*.) *Styrax liquida*. *Liquidamba*. Liquid amber. A resinous juice, of a yellow colour, inclining to red, at first about the consistence of turpentine, by age hardened into a solid brittle mass. It is obtained by wounding the bark of the *Liquidambar styraciflua*; *foliis palmato-angulatis*; *foliis indivisis, acutis*, of Linnæus. This juice has a moderately pungent, warm balsamic taste, and a very fragrant smell, not unlike that of the *Styrax calamita* heightened by a little ambergris. It is seldom used medicinally. The *Styrax liquida* is also obtained by boiling from this plant. There are two sorts distinguished by authors; the one, the purer part of the resinous matter, that rises to the surface in boiling, separated by a strainer, of the consistence of honey, tenacious like turpentine, of a reddish or ash-brown colour, moderately transparent, of an acrid unctuous taste, and a fragrant smell, faintly resembling that of the solid storax, but somewhat disagreeable. The other; the more impure part which remains on the strainer, untransparent, and in smell and taste much weaker than the former. Their use is chiefly as stomachics, in the form of plaster.

LIQUIBITIA. (From *liquor*, juice, or from *elikeris*, Welsh.) See *Glycyrrhiza*.

LIQUOR ÆTHEREUS VITRIOLICUS. The liquor procured from a distillation of equal parts of sulphuric acid and spirit of wine redistilled,

LIQUOR ACETATIS AMMONIÆ. Solution of acetate of ammonia; formerly called *Aqua ammoniæ acetate*. "Take of carbonate of ammonia, two ounces; acetic acid, four pints." Add the acid to the salt, until bubbles of gas shall no longer arise, and mix.

If the acid rather predominate, the solution is more grateful to the taste; and provided that acid be correctly prepared, the proportions here given will be found sufficient; where the acid cannot be depended on, it will be right to be regulated rather by the cessation of effervescence than by quantity.

This preparation was formerly known in the shops under the name of spirit of Minderus. When assisted by a warm regimen, it proves an excellent and powerful sudorific; and, as it operates without quickening the circulation, or increasing the heat of the body, it is admissible in febrile and inflammatory diseases, in which the use of stimulating sudorifics are attended with danger. Its action may likewise be determined to the kidneys, by walking about in the cool air. The common dose is half an ounce, either by itself

or along with other medicines, adapted to the same intention.

LIQUOR ACETATIS PLUMBI. Solution of acetate of lead, formerly called *aqua lithargyri acetati*. Goulard's extract. "Take of vitrified oxyd of lead, two pounds and four ounces; acetic acid, a gallon." Mix, and boil down to six pints, constantly stirring; then set it by, that the feculencies may subside, and strain.

It is principally employed by surgeons, as a resolvent against inflammatory affections.

LIQUOR ACETATIS PLUMBI DILUTUS. Diluted solution of acetate of lead. *Aqua lithargyri acetati composita*. "Take of solution of acetate of lead, a drachm; distilled water, a pint; weak spirit, a fluid drachm." Mix. The virtues of this water, the *aqua vegeto-minerals* of former pharmacopœias, applied externally, are resolvent, refrigerant, and sedative.

LIQUOR ALUMINIS COMPOSITUS. Compound solution of alum. "Take of alum, sulphate of zinc, of each half an ounce; boiling water, two pints." Dissolve at the same time the alum and sulphate of zinc in the water, and then strain the solution through paper.

This water was long known in our shops under the title of *aqua aluminosa bateana*. It is used for cleansing and healing ulcers and wounds, and for removing cutaneous eruptions, the part being bathed with it hot three or four times a-day. It is sometimes likewise employed as a collyrium; and as an injection in fluor albus and gonorrhœa, when not accompanied with virulence.

LIQUOR AMMONIÆ. Solution of ammonia. *Aqua ammoniæ pure* of the London Pharm. 1787. "Take of muriate of ammonia, lime, newly prepared, of each two pounds; water, a pint and a half." Reduce the muriate of ammonia and the lime into powder, separately; then mix them, and introduce them into a large glass retort, into which a pint of the water has been previously poured. Having placed the retort in a sand-bath, lute on a tubulated receiver, through which the ammonia may pass into a third vessel, containing half a pint of the water, and cooled. Then at first apply a gentle heat; and increase it by degrees, until the retort becomes red.

Water of ammonia is very rarely given internally, although it may be used in doses of ten or twenty drops, largely diluted, as a powerful stimulant in asphyxia and similar diseases. Externally, it is applied to the skin as a rubefacient, and in the form of gas to the nostrils, and to the eyes as a stimulant; in cases of torpor, paralysis, rheumatism, syncope, hysteria, and chronic ophthalmia.

LIQUOR CARBONATIS AMMONIÆ. Solution of carbonate of ammonia. The *aqua*
3 M

ammonia of the Pharm. Lond. 1787. "Take of carbonate of ammonia, eight ounces; distilled water, a pint." Dissolve the subcarbonate of ammonia in the water, and filter the solution through paper. This preparation possesses the properties of ammonia in its action on the human body. See *Carbonas ammoniac*.

LIQUOR AMNII. All that fluid which is contained in the membranaceous ovum surrounding the fœtus in utero is called by the general name of the waters, the water of the amnion, or ovum, or liquor amnii. The quantity, in proportion to the size of the different parts of the ovum, is greatest by far in early pregnancy. At the time of parturition, in some cases, it amounts to or exceeds four pints; and in others it is scarcely equal to as many ounces. It is usually in the largest quantity when the child has been some time dead, or is born in a weakly state. This fluid is generally transparent, often milky, and sometimes of a yellow, or light brown colour, and very different in consistence; and these alterations seem to depend upon the state of the constitution of the parent. It does not coagulate with heat, like the serum of the blood; and, chymically examined, it is found to be composed of phlegm, earthy matter, and sea salt, in different proportions in different subjects, by which the varieties in its appearance and consistence are produced. It has been supposed to be excrementitious; but it is generally thought to be secreted from the internal surface of the ovum, and to be circulatory, as in other cavities. It was formerly imagined, that the fœtus was nourished by this fluid, of which it was said to swallow some part frequently; and it was then asserted, that the qualities of the fluid were adapted for its nourishments. But there have been many examples of children born without any passage to the stomach; and a few, of children in which the head was wanting, and which have nevertheless arrived at the full size. These cases fully prove that this opinion is not just, and that there must be some other medium by which the child is nourished, besides the waters. The incontrovertible uses of this fluid are, to serve the purpose of affording a soft bed for the residence of the fœtus, to which it allows free motion, and prevents any external injury during pregnancy: and inclosed in the membranes, it procures the most gentle, yet efficacious, dilatation of the os uteri, and soft parts, at the time of parturition. Instances have been recorded, in which the waters of the ovum are said to have been voided so early as in the sixth month of pregnancy, without prejudice either to the child or parent. The truth of these reports seem to be doubtful, because, when the membranes are intentionally broken, the action of the

uterus never fails to come on, when all the water is evacuated. A few cases have occurred to me, says Dr. Denman, in practice, which might have been construed to be of this kind; for there was a daily discharge of some colourless fluid from the vagina, for several months before delivery; but there being no diminution of the size of the abdomen, and the waters being regularly discharged at the time of labour, it was judged that some lymphatic vessel near the os uteri had been ruptured, and did not close again till the patient was delivered. He also met with one case, in which, after the expulsion of the placenta, there was no sanguineous discharge, but a profusion of lymph, to the quantity of several pints in a few hours after delivery; but the patient suffered no inconvenience, except from surprise.

LIQUOR ANTIMONII TARTARIZATI. Solution of tartarized antimony. *Vinum antimonii tartarizati* of the Pharm. Lond. 1787. "Take of the tartarized antimony, one scruple; boiling distilled water, four fluid-ounces; wine six fluid-ounces." Dissolve the tartarized antimony in the boiling distilled water, then add the wine. Half an ounce of the solution contains one grain of the salt. This preparation may be given in all cases where the tartar emetic is indicated.

LIQUOR ARSENICALIS. Arsenical solution. "Take of prepared oxyd of arsenic, in very fine powder, subcarbonate of potash from tartar, of each 64 grains; distilled water a pint." Boil them together in a glass vessel, until the arsenic be entirely dissolved. When the solution is cold, add "compound spirit of lavender, four fluid-drachms." Then add as much distilled water as may exactly fill a pint measure. This preparation accords with the formula of Dr. Fowler, of Stafford, who first introduced it in imitation of a celebrated popular remedy for intermittents, sold under the name of the tasteless ague-drop. The compound spirit of lavender is only intended to give some colour and taste, without which it would resemble common water, and hence be more liable to mistakes. Where the dose is small, and the effects so powerful, the most minute attention to its proportion and preparation become necessary. Each ounce contains four grains of the oxyd, and each drachm one-eighth of a grain; but it will rarely be proper to go beyond half the latter quantity, or one-sixteenth of a grain as a dose.

LIQUOR CALCIS. Solution of lime, formerly called *aqua calcis*. Lime-water. "Take of lime, half a pound; boiling distilled water, twelve pints." Pour the water upon the lime, and stir them together; next cover the vessel immediately, and let it stand for three hours; then keep

the solution upon the remaining lime in stopped glass bottles, and pour off the clear liquor when it is wanted for use.

Lime is soluble in about 450 times its weight of water, or little more than one grain in one fluid-ounce. It is given internally, in doses of two ounces and upwards, in cardialgia, spasms, diarrhoea, &c. and in proportionate doses in convulsions of children, arising from acidity, or ulcerated intestines, intermittent fevers, &c. Externally it is applied to burns and ulcers.

LIQUOR CUPRI AMMONIATI. Solution of ammoniated copper. *Aqua cupri ammoniati* of Pharm. Lond. 1787, and formerly called *aqua sapharina*. "Take of ammoniated copper, a drachm; distilled water, a pint." Dissolve the ammoniated copper in the water, and filter the solution through paper. This preparation is employed by surgeons for cleansing foul ulcers, and disposing them to heal.

LIQUOR FERRI ALKALINI. Solution of alkaline iron, "Take of iron, two drachms and a half; nitric acid, two fluid-ounces; distilled water, six fluid-ounces; solution of carbonate of potash, six fluid-ounces."

Having mixed the acid and water, pour them upon the iron, and when the effervescence has ceased, pour off the clear acid solution; add this gradually, and at intervals, to the solution of subcarbonate of potash, occasionally shaking it, until it has assumed a deep brown red colour, and no further effervescence takes place. Lastly, set it by for six hours, and pour off the clear solution. This preparation was first described by Stahl, and called *tinctura martis alkalina*, and is now introduced in the Lond. Pharm. as affording a combination of iron distinct from any other, and often applicable to practice. The dose is from half a drachm to a drachm.

LIQUOR OXYMURIATIS HYDRARGYRI. Solution of oxymuriate of mercury. "Take of oxymuriate of mercury, eight grains; distilled water, fifteen fluid-ounces; rectified spirit, a fluid-ounce." Dissolve the oxymuriate of mercury in the water, and add the spirit.

This solution is directed in order to facilitate the administration of divisions of the grain of this active medicine. Half an ounce of it contains one-eighth of a grain of the salt. The dose is from one drachm to half an ounce.

LIQUOR MINERALIS ANODYNUS HOFFMANNI. Hoffman's anodyne liquor. A preparation of ether extolled as an anodyne and antispasmodic. See *Spiritus atheris compositi*.

LIQUOR POTASSE. Solution of potash. *Aqua kali puri*. *Lixivium saponarium*. "Take of subcarbonate of potash, lime, newly prepared, of each a pound. Boiling distilled water, a gallon" Dissolve

the potash in two pints of the water; add the remaining water to the lime. Mix the liquors while they are hot, stir them together, then set the mixture by in a covered vessel; and after it has cooled, strain the solution through a cotton bag.

If any diluted acid dropped into the solution occasion the extrication of bubbles of gas, it will be necessary to add more lime, and to strain it again. A pint of this solution ought to weigh sixteen ounces.

LIQUOR SUBCARBONATIS POTASSE. Solution of subcarbonate of potash. *Aqua kali preparati*. *Lixivium tartari*. *Oleum tartari per deliquium*. "Take of subcarbonate of potash, a pound; distilled water, twelve fluid-ounces." Dissolve the subcarbonate of potash in the water, and then strain the solution through paper.

It possesses antacid virtues, and is a good antidote against arsenic taken into the stomach. It is also given with advantage in convulsions and spasms, from acidity in the stomach of children, in calculous diseases, gouty affections, scrophula, aphthæ, &c. The carbonate of soda is milder, and perhaps a preferable remedy for general use. The dose of the liquor potassæ subcarbonatis is from half a drachm to a drachm and a half.

LIQUOR VOLATILIS CORNU CERVI. This preparation of the volatile alkali, commonly termed hartshorn, possesses the same virtues as the carbonate of ammonia. It is in common use to smell at in faintings, &c. See *Carbonas ammoniacæ*.

Liquorice. See *Glycyrrhiza*.

Liquorice, Spanish. See *Glycyrrhiza*.

LITHAGOGA. (From *lithos*, a stone, and *ago*, to bring away.) Medicines which expel the stone.

Litharge. See *Lithargyrus*.

Litharge plaster. See *Emplastrum lithargyri*.

LITHARGYRUS. (From *lithos*, a stone, and *argyros*, silver.) *Lithargyrum*. A subcarbonate of lead, in an imperfect state of vitrification. When silver is refined by cupellation with lead, this latter metal, which is scorified, and causes the scorification of the imperfect metals alloyed with the silver, is transformed into a matter composed of small semitransparent shining plates, resembling mica; which is litharge. Litharge is more or less white or red, according to the metals with which the silver is alloyed. The white is called litharge of silver; and the red has been improperly called litharge of gold. See *Lead, Liquor plumbi acetatis*, and *Liquor plumbi acetatis dilutus*.

LITHIAS. A lithiate, or salt, formed by the union of the lithic acid, or acid of the stone sometimes found in the human bladder, with different bases; thus, *lithiate of alumine*, *lithiate of ammonia*, &c.

LITHIASIS. (From *λίθος*, a stone.)

1. The formation of stone, or gravel.

2. A tumour of the eyelid, under which is a hard concretion resembling a stone.

LITHOBALAN. (From *λίθος*, a stone, and *λαμβάνω*, to seize.) An instrument for extracting the stone from the bladder.

LITHOIDES. (From *λίθος*, a stone, and *ωδός*, a likeness; so called from its hardness.) The petrous portion of the temporal bone.

LITHOLOGY. (*Lithologia*; from *λίθος*, a stone, and *λογος*, a discourse.) A discourse or treatise on stones.

LITHOMARCA. Stone marle. Fuller's earth is one of the most useful varieties of lithomarge.

LITHONTRIPTICS. (*Lithontriptica*, sc. *medicamenta*; from *λίθος*, a stone, and *θραύω*, to break.) From the strict sense and common acceptance of the word, this class of medicine should comprehend such as possess a power of dissolving calculi in the urinary passages. It is, however, doubted by many, whether there be in nature any such substances. By this term, then, is meant those substances which possess a power of removing a disposition in the body to the formation of calculi. The researches of modern chymists have proved, that these calculi consist mostly of a peculiar acid, named the lithic or uric acid. With this substance the alkalies are capable of uniting, and forming a soluble compound; and these are accordingly the sole lithontriptics. From the exhibition of alkaline remedies, the symptoms arising from stone in the bladder are very generally alleviated; and they can be given to such an extent that the urine becomes very sensibly alkaline, and is even capable of exerting a solvent power on these concretions. Their administration, however, cannot be continued to this extent for any length of time, from the irritation they produce on the stomach and urinary organs. The use, therefore, of the alkalies, as solvents, or lithontriptics, is now scarcely ever attempted; they are employed merely to prevent the increase of the concretion, and to palliate the painful symptoms, which they do apparently by preventing the generation of lithic acid, or the separation of it by the kidneys; the urine is thus rendered less irritating, and the surface of the calculus is allowed to become smooth.

When the alkalies are employed with this view, they are generally given saturated, or supersaturated, with carbonic acid. This renders them much less irritating. It at the same time, indeed, diminishes its solvent power; for the alkaline carbonates exert no action on urinary calculi; but they are still capable of correcting that acidity in the primæ viz, which is the cause of the deposition of the lithic acid from

the urine, and therefore serve equally to palliate the disease. And when their acrimony is thus diminished, their use can be continued for any length of time.

It appears from the experiments of Fourcroy, and others, that the other ingredients of calculi, as well as the lithic acid, are dissolved by the caustic alkali, and various experiments have shewn that most calculi yield to its power. Lime-water has also been found a solvent of urinary calculi, out of the body. It is obvious, however, that what is taken by the mouth is subject to many changes in the alimentary canal, and also the lymphatic and vascular systems; and in this way, it must be exceedingly difficult to get such substances (even were they not liable to alterations) in sufficient quantity into the bladder. Indeed there are very few authenticated cases of the urine being so changed as to become a menstruum for the stone. Excepting the case of Dr. Newcombe, recorded by Dr. Whytt, the instance of Mr. Home is almost the only one. Though lithontriptics, however, may not in general dissolve the stone in the bladder, yet it is an incontrovertible fact that they frequently mitigate the pain; and, to lessen such torture as that of the stone in the bladder, is surely an object of no little importance. Lime was long ago known as a solvent of urinary calculi, and different methods were employed to administer it. One of these plans fell into the hands of a Mrs. Steevens, and her success caused great anxiety for the discovery of the secret. At last, Parliament bought the secret for the sum of 3000*l.* In many instances, stones which had been unquestionably felt were no longer to be discovered; and as the same persons were examined by the greatest skill and eminence, both before and after the exhibition of her medicines, it was no wonder that the conclusion was drawn, that the stones really were dissolved. From the cessation of such success, and from its now being known that the stones are occasionally protruded between the fasciculi of the muscular fibres of the bladder, so to be lodged in a kind of cyst on the outside of the muscular coat, and cause no longer any grievances, surgeons of the present day are inclined to suspect that this must have happened in Mrs. Steevens's cases. This was certainly what happened in one of the cases on whom the medicine had been tried. It is evident that a stone so situated would not any longer produce irritation, but would also be quite undiscoverable by the sound, for, in fact, it is no longer in the cavity of the bladder.

As soap was, with reason, supposed to increase the virtues of the lime, it led to the use of caustic alkali, taken in mucilage, or veal broth. Take of kali, prepared,

℥viij; of quick lime ℥iv; of distilled water, ℔ij. Mix them well together in a large bottle, and let them stand for twenty four hours. Then pour off the ley, filter it through paper, and keep it in well-stopped vials for use. Of this, the dose is from thirty drops to ℥ij, which is to be repeated two or three times a-day, in a pint of veal broth, early in the morning, at noon, and in the evening. Continue this plan for three or four months, living, during the course, on such things as least counteract the effect of the medicine.

The common fixed alkalis, or carbonated alkali, and the acidulous soda-water, have of late been used as lithontriptics. Honey has also been given, and Mr. Home, surgeon at the Savoy, has recorded its utility in his own and his father's cases. Bitters have likewise been tried.

Dismissing all theories, lime-water, soap, acidulous soda-water, caustic alkali, and bitters, are useful in cases of stone. Of the soap, as much may be taken as the stomach will bear, or as much as will prove gently laxative; but of the lime-water, few can take more than a pint daily.

The acidulous soda-water may be taken in larger quantities, as it is more agreeable. The acidulous salt is now prepared so as to produce the water extemporaneously. It must be swallowed, however, while the salt is dissolving, as the carbonic acid very rapidly escapes.

There is a remedy celebrated in Holland, under the name of liquor lithontriptica loosii, which contains, according to an accurate analysis, calx muriata. This, professor Hufeland recommends in the following form:

℞ Calcis muriatæ ℥j.

Aquæ distillatæ, ℥ij. ft. solutio.

Thirty drops are to be taken four times a-day, which may be increased as far as the stomach will bear.

For curing stone patients, little reliance can be placed in any lithontriptics hitherto discovered, though they may rationally be given with a confident hope of procuring an alleviation of the fits of pains attending the presence of stone in the bladder. After all, the only certain method of getting rid of the calculus is the operation. See *Lithotomy*.

LITHOSPERMUM. (From *λίθος*, a stone, and *σπέρμα*, seed; named from the hardness of its seed) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

2. The pharmacopœial name of *Milium solis*. Common gromwell. The seeds of this official plant, *Lithospermum officinale*; *seminibus levibus, corollis vix calycem superantibus, foliis lanceolatis*, of Linnæus, were formerly supposed, from their stony hardness, to be efficacious in calculous and gra-

velly disorders. Little credit is given to their lithontriptic character, yet they are occasionally used as diuretic for clearing the urinary passages, and for obviating stranguary, in the form of emulsion.

LITHOSPERMUM OFFICINALE. The systematic name of the official gromwell. See *Lithospermum*.

LITHOTOMY. (*Lithotomia*; from *λίθος*, a stone, and *τομή*, to cut.) *Cystotomia*. The operation of cutting into the bladder, in order to extract a stone. Several methods have been recommended for performing this operation, but there are only two which can be practised with any propriety. One is, where the operation is to be performed immediately above the pubes, in that part of the bladder which is not covered with peritoneum, called the *high operation*. The other, where it is done in the perineum, by laying open the neck and lateral part of the bladder, so as to allow of the extraction of the stone, called the *lateral operation*, from the prostatic gland of the neck of the bladder being laterally cut.

LITRON. (Corruptly written for *λίτρον*.) Nitre.

LITRUS. (From *λίνο*, to anoint.) A liniment.

LIVER. *πλάξ. Hepar.* A large viscus, of a deep red colour, of great size and weight, situated under the diaphragm, in the right hypochondrium, its smaller portion occupying part of the epigastric region. In the human body, the liver is divided into two principal lobes, the right of which is by far the largest. They are divided on the upper side by a broad ligament, and on the other side by a considerable depression, or fossa. Between and below these two lobes is a smaller lobe, called *lobulus Spigelii*. In describing this viscus, it is necessary to attend to seven principal circumstances:—its ligaments; its surfaces; its margins; its tubercles; its fissure; its sinus; and the pori biliari.

The *ligaments* of the liver are five in number, all arising from the peritoneum. 1. *The right lateral ligament*, which connects the thick right lobe with the posterior part of the diaphragm. 2. *The left lateral ligament*, which connects the convex surface and margin of the left lobe with the diaphragm, and, in those of whom the liver is very large, with the œsophagus and spleen. 3. *The broad or middle suspensory ligament*, which passes from the diaphragm into the convex surface, and separates the right lobe of the liver from the left. It descends from above through the large fissure to the concave surface, and is then distributed over the whole liver. 4. *The round ligament*, which in adults consists of the umbilical vein, indurated into a ligament. 5. *The coronary ligament*.

The liver has two *surfaces*, one superior,

which is convex and smooth, and one inferior, which is concave and has holes and depressions, to receive, not only the contiguous viscera, but the vessels running into the liver.

The *margins* of the liver are also two in number; the one, which is posterior and superior, is obtuse; the other, situated anteriorly and inferiorly, is acute.

The *tubercles* of the liver are likewise two in number, viz. *lobulus anonymus*, and *lobulus caudatus*, and are found near the vena portæ.

Upon looking on the concave surface of this viscus, a considerable fissure is obvious, known by the name of the *fissure of the liver*.

In order to expose the *sinus*, it is necessary to remove the gall-bladder, when a considerable sinus, before occupied by the gall-bladder, will be apparent.

The *blood-vessels* of the liver are the hepatic artery, the vena portæ, and the cavæ hepaticæ, which are described under their proper names. The *absorbents* of the liver are very numerous. The liver has *nerves* from the great intercostal and eighth pair, which arise from the hepatic plexus, and proceed along with the hepatic artery and vena portæ into the substance of the liver. With regard to the substance of the liver, various opinions have been entertained. It is, however, now pretty well ascertained to be a large gland, composed of lesser glands connected together by cellular structure. The small glands which thus compose the substance of the liver are termed *penicilli*, from the arrangement of the arterial ramifications of the vena portæ composing each gland, resembling that of the hairs of a pencil. The chief use of this large viscus is to supply a fluid, named bile, to the intestines, which is of the utmost importance in chylickation. The small *penicilli* perform this function by a specific action on the blood they contain, by which they secrete in their very minute ends the fluid termed *hepatic bile*; but whether they pour it into what is called a follicle, or not, is yet undecided, and is the cause of the difference of opinion respecting the substance of the liver. If it be secreted into a follicle, the substance is truly glandular, according to the notion of the older anatomists; but if it be secreted merely into a small vessel, called a biliary pore (whose existence can be demonstrated, corresponding to the end of each *penicilli*, without any intervening follicle, its substance is then, in their opinion, vascular. According to our notions in the present day, in either case, the liver is said to be glandular; for we connect to our senses the idea of a gland, when any arrangement of vessels performs the office of separating from the blood a fluid or sub-

stance different in its nature from the blood. The small vessels which receive the bile secreted by the *penicilli* are called *pori biliarii*; these converge together throughout the substance of the liver towards its under surface, and, at length, form one trunk, called *ductus hepaticus*, which conveys the bile into either the *ductus communis choledochus*, or *ductus cysticus*. See Gall-bladder.

Liver, acute inflammation of the. See *Hepatitis*.

Liver of sulphur. See *Sulphurets*.

Liver wort. See *Hypatica terrestris*.

Liver-wort, ash-coloured. See *Lichen cinereus terrestris*.

Liver wort, ground. See *Lichen cinereus terrestris*.

Liver-wort, Iceland. See *Lichen islandicus*.

Liver-wort, noble. See *Hepatica terrestris*.

LIVOR. (From *liveo*, to be black and blue.) A blackish mark on the body, from a blow. A dark circle under the eye.

LIX. (From *lux*, light.) Pot-ash. Wood ash.

LIXIVIA VITRIOLATA SULPHUREA. A sulphat of potash.

LIXIVIAL. Those salts are called *lixivial* which have been extracted by *lixivation*, and these chiefly are fixed alkalis; which are therefore called *lixivial salts*.

LIXIVIATION. *Lessive.* The process employed by chymists of dissolving, by means of warm water, the saline and soluble particles of cinders, the residues of distillation and combustion, coals and neutral earths, in order to obtain those particles which are termed *lixivial salts*.

LIXIVIUM. (From *lix*, wood-ash.) The liquor in which saline and soluble particles of the residues of distillation and combustion are dissolved.

LIXIVIUM SAPONARIUM. See *Liquor potassæ*.

LIXIVIUM TARTARI. See *Liquor carbonatis potassæ*.

LOBELIA. Named in honour of Lobel, a botanist. 1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia* Order, *Monogamia*.

2. The pharmacopœial name of the blue lobelia, or cardinal flower. The root of this plant, *Lobelia syphilitica*, is the part directed by the Edinburgh Pharmacopœia for medicinal use; in taste it resembles tobacco, and is apt to excite vomiting. It derived the name of *syphilitica* from its efficacy in the cure of syphilis, as experienced by the North American Indians, who considered it as a specific in that disease, and with whom it was long an important secret, which was purchased by Sir William Johnson, and since published by

different authors. The method of employing this medicine is stated as follows: A decoction is made of a handful of the roots in three measures of water. Of this half a measure is taken in the morning fasting, and repeated in the evening; and the dose is gradually increased, till its purgative effects become too violent, when the decoction is to be intermitted for a day or two, and then renewed, until a perfect cure is effected. During the use of this medicine, a proper regimen is to be enjoined, and the ulcers are also to be frequently washed with the decoction, or, if deep and foul, to be sprinkled with the powder of the inner bark of the New-Jersey tea-tree, *Ceanothus Americanus*. Although the plant thus used is said to cure the disease in a very short time, yet it is not found that the anti-syphilitic powers of the lobelia have been confirmed in any instance of European practice.

LOBELIA STYHLITICA. The systematic name of the blue lobelia of the pharmacopœias. See *Lobelia*.

LOBULUS. (Dim. of *lobus*, a lobe.) A small lobe.

LOBULUS ACCESSORIUS. See *Lobulus anonymus*.

LOBULUS ANONYMUS. *Lobulus accessorius anterior-quadratus.* The anterior point of the right lobe of the liver. Others define it to be that space of the great lobe betwixt the fossa of the umbilical vein and gall-bladder, and extending forward from the fossa for the lodgment of the porta, to the anterior margin of the liver.

LOBULUS CAUDATUS. *Processus caudatus.* A tail-like process of the liver, stretching downward from the middle of the great right lobe to the lobulus spigelii. It is behind the gall-bladder and betwixt the fossa venæ portarum, and the fissure for the lodgment of the vena cava.

LOBULUS SPIGELII. *Lobulus posterior. L. posticus-papellatus.* The lobulus spigelii is betwixt the two greater lobes, but rather belonging to the right great lobe. From its situation deep behind, and from its having a perpendicular papilla-like projection, it is called lobulus posterior, or papillatus. To the left side, it has the fissure for the lodgment of the ductus venosus; on the right, the fissure for the vena cava; and above, it has the great transverse fissure of the liver, for the lodgment of the cylinder of the porta; obliquely to the right, and upwards, it has a connection with the lower concave surface of the great lobe, by the processus caudatus, which Winslow calls one of the roots of the lobulus spigelii. It is received into the bosom of the lesser curve of the stomach.

LOCALES. The fourth class of Cullen's Nosology, which comprehends morbid affections that are partial, and includes

eight genera, viz. *dysæsthesiæ*, *dysorexiæ*, *dyscinesia*, *apocenosæ*, *epischæsæ*, *tumores*, *ecopia*, and *dialyses*.

Localis membrana. The pia mater.

LOCHIA. (From *λοχια*, to bring forth.) The cleansings. The serous, and for the most part green-coloured, discharge that takes place from the uterus and vagina of women, during the first four days after delivery.

LOCHIORRHŒA. (From *λοχια*, and *ρρœa*, to flow.) An excessive discharge of the lochia.

Locked jaw. See *Tetanus*.

Logwood. See *Lignum campechense*.

LOXCHITIS. (From *λοχχι*, a lance; so named because the leaves resemble the head of a lance.) The herb spleenwort.

LONGANUM. (From *longus*, long; so named from its length.) The intestinum rectum.

LONGINE. A disease peculiar to the female, and only daring pregnancy, and those states in which the uterine discharge is suppressed.

LONGISSIMUS DORSI. *Lumbo dorso trachelii* of Dumas. This muscle, which is somewhat thicker than the sacrolumbalis, greatly resembles it, however, in its shape and extent, and arises, in common with that muscle, between it and the spine. It ascends upwards along the spine, and is inserted by small double tendons into the posterior and inferior part of all the transverse processes of the vertebrae of the back, and sometimes of the last vertebra of the neck. From its outside it sends off several bundles of fleshy fibres, interspersed with a few tendinous filaments, which are usually inserted into the lower edge of the ten uppermost ribs, not far from their tubercles. In some subjects, however, they are found inserted into a less number, and in others, though more rarely, into every one of the ribs. Towards the upper part of this muscle is observed a broad and thin portion of fleshy fibres, which cross and intimately adhere to the fibres of the longissimus dorsi. This portion arises from the upper and posterior part of the transverse processes of the five or six uppermost vertebrae of the back, by as many tendinous origins, and is usually inserted, by six tendinous and fleshy slips, into the transverse processes of the six inferior vertebrae of the neck. This portion is described by Winslow and Albinus as a distinct muscle; by the former under the name of *transversalis major colli*, and by the latter under that of *transversalis cervicis*. But its fibres are so intimately connected with those of the longissimus dorsi, that it may very properly be considered as an appendage to the latter. The use of this muscle is to extend the vertebrae of the back, and to keep the trunk of the body erect; by

means of its appendage, it likewise serves to turn the neck obliquely backwards, and a little to one side.

LONGISSIMUS MANUS. The flexor tertii internodii pollicis.

LONGISSIMUS OCULI. The obliquus major oculi.

LONGITUDINAL SINUS. Longitudinal sinus of the dura mater. A triangular canal, proceeding in the falxiform process of the dura mater, immediately under the bones of the skull, from the crista galli to the tentorium, where it branches into the lateral sinuses. The longitudinal sinus has a number of trabeculae or fibres crossing it. Its use is to receive the blood from the veins of the pia mater, and convey it into the lateral sinuses, to be carried through the internal jugulars to the heart.

LONGUS COLLI. *Pre dorso cervical* of Dumas. This is a pretty considerable muscle, situated close to the anterior and lateral part of the vertebrae of the neck. Its outer edge is in part covered by the rectus internus major. It arises tendinous and fleshy within the thorax, from the bodies of the three superior vertebrae of the back, laterally; from the bottom and fore-part of the transverse processes of the first and second vertebrae of the back, and of the last vertebra of the neck; and likewise from the upper and anterior points of the transverse processes of the sixth, fifth, fourth, and third vertebrae of the neck, by as many small distinct tendons; and is inserted tendinous into the fore-part of the second vertebra of the neck, near its fellow. This muscle, when it acts singly, moves the neck to one side; but, when both act, the neck is brought directly forwards.

LONICERA. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

LONICERA DIERVILLA. The systematic name of a species of honey-suckle. See *Diervilla*.

LONICERA PRÆCLOMINUM. Honey-suckle. This beautiful and common plant was formerly used in the cure of asthma, for cleansing fordid ulcers, and removing diseases of the skin, virtues it does not now appear to possess.

Looseness. See *Diarrhæa*.

LOPEX RADIX. *Rudix lopesiana.* *Radix indica lopesiana.* The root of an unknown tree growing, according to some, at Goa. It is met with in pieces of different thickness, some at least of two inches diameter. The woody part is whitish, and very light; softer, more spongy, and whiter next the bark, including a denser, somewhat reddish, medullary part. The bark is rough, wrinkled, brown, soft, and, as it were, woolly, pretty thick, covered with a thin paler cuticle. Neither the woody nor corticle part has any remarkable smell

or taste, nor any appearance of resinous matter. It appears that this medicine has been remarkably effectual in stopping colliquative diarrhoeas, which had resisted the usual remedies. Those attending the last stage of consumptions were particularly relieved by its use. It seemed to act, not by an astringent power, but by a faculty of restraining and appeasing spasmodic and inordinate motions of the intestines. Dr. Gaubius, who gives this account, compares its action to that of *simarouba*, but thinks it more efficacious than this medicine.

Lopez-root. See *Lopez radix*.

LOPEZIANA RADIX. See *Lopez radix*.

LOPHADIA. (From *λοφος*, the hinder part of the neck.) *Lophia*. The first vertebra of the neck.

LORDOSIS. (From *λωσος*, curved, bent.) An inflexion of the spine, in which it is bent inwards.

LORICA. (From *lorico*, to crust over.) A kind of lute, with which vessels are coated before they are put into the fire.

LORICATION. Coating. Nicholson recommends the following composition for the coating of glass vessels, to prevent their breaking when exposed to heat. Take of sand and clay, equal parts; make them into a thin paste with fresh blood, prevented from coagulating by agitation, till it is cold, and diluted with water; add to this some hair, and powdered glass; with a brush, dipped in this mixture, besmear the glass; and when this layer is dry, let the same operation be repeated twice, or oftener, till the coat applied is about one-third part of an inch in thickness.

LORIND MATRICES. An epilepsy, or a convulsive disorder, proceeding from the uterus.

Loss of Appetite. See *Anorexia*.

LOTION. (*Lotio*, from *lavo*, to wash.) An external fluid application. Lotions are usually applied by wetting linen in them, and keeping it on the part affected.

LOTUS. (From *λω*, to desire.) 1. A tree whose fruit was said to be so delicious as to make those who tasted it to forsake all other desires: hence the proverb *Λωτον σπαρον, lotum gustavi*: I have tasted lotus.

2. The name of a genus of plants in the Linnæan system. Class, *Diadelphæa*. Order, *Decandria*.

Lousy disease. A general corruption of the humours, in consequence of which these insects are bred in ulcers, and cover the whole body.

LOVE-APPLE. The fruit of the *Solanum lycopersicum* of Linnæus. It is so much esteemed by the Portuguese and the Spaniards, that it is an ingredient in almost all their soups and sauces, and is deemed cooling and nutritive.

Loveage. See *Levisticum*.

LOXARTHEROS. (From *λοξος*, oblique, and *αρθρον*, a joint.) An obliquity of the joint, without spasm or luxation.

LUDUS HELMONTII. (The) waxen vein,

called also *ludus paracelsi*. A stony matter said to be serviceable in calculus.

LUES BRITICA. One of the pompous names for epilepsy.

LUES NEURODES CONVULSIVA. A mild typhus fever.

LUES VENEREA. (From *lues*, to dissolve, because it produces dissolution; and *veneræ*, from *Venus*, because it is propagated by acts of venery.) The plague of Venus, or the venereal disease. Dr. Cullen calls it *syphilis*. It has also been called the venereal pestilence, or pox. *Aphrodisius morbus*. *Morbus gallicus*. *Indicus morbus*. *Neapolitanus morbus*. *Paturus*. See *Syphilis* and *Gonorrhæa*.

LUJULA. (Corrupted or contracted from *Allelujah*, *Praise the Lord*; so called from its many virtues.) *Acetosella*. Wood-sorrel. *Oxalis acetosella* of Linnaeus. This delicate indigenous plant is totally inodorous, but has a grateful acid taste, which is more agreeable than the common sorrel, and approaches near to that of the juice of lemons, or the acid of tartar, with which it also corresponds in a great measure in its medical effects, being esteemed refrigerant, antiscorbutic, and diuretic. Its principal use, however, is to allay inordinate heat, and to quench thirst; for this purpose a pleasant whey may be formed by boiling the plant in milk. An essential salt is prepared from this plant, known by the name of Essential Salt of Lemons, and commonly used for taking ink-stains out of linen.

LUMBAGO. (From *lumbus*, the loin.) A rheumatic affection of the muscles about the loins.

LUMBAGO PSOADICA. *Lumbago apostematosa*. *Lumbago ab arthrocace*. Pains in the loins from abscess.

LUMBARES ARTERIÆ. The lumbal arteries.

LUMBARES NERVI. The lumbal nerves.

LUMBARES VENÆ. The lumbal nerves.

LUMBARIS EXTERNUS. See *Quadratus lumborum*.

LUMBARIS INTERNUS. See *Psoas magnus*.

LUMBAR ABSCESS. *Psoas abscess*. A species of *arthropneusis*, that receives its name from the situation in which the matter is found, namely, upon the side of the psoas muscle, or betwixt that and the iliacus internus. Between these muscles, there lies a quantity of loose cellular membrane, in which an inflammation often takes place, either spontaneously or from mechanical injuries. This terminates in an abscess that can procure no outlet but by a circuitous course, in which it generally produces irreparable mischief, without any violent symptoms occurring to alarm the patient. The abscess sometimes forms a swelling above Poupert's ligament; sometimes below it; and frequently the

matter glides under the fascia of the thigh. Occasionally, it makes its way through the sacro-ischiatic foramen, and assumes rather the appearance of a fistula in ano. The uneasiness in the loins, and the impulse communicated to the tumour by coughing, evince that the disease arises in the lumbar region; but it must be confessed, that we can hardly ever know the existence of the disorder, before the tumour, by presenting itself externally, leads us to such information. The lumbar abscess is sometimes connected with diseased vertebrae, which may either be a cause or effect of the collection of matter. The disease, however, is frequently unattended with this complication.

The situation of the symptoms of lumbar abscess renders this affection liable to be mistaken for some others, viz. lumbago and nephritic pains, and, towards its termination, for crural or femoral hernia. The first, however, is not attended with the shivering that occurs here; and nephritic complaints are generally discoverable by attention to the state of the urine. The distinction from crural hernia is more difficult. In both, a soft inelastic swelling is felt in the same situation; but in hernia, it is attended with obstructed feces, vomiting, &c. and its appearance is always sudden, while the lumbar tumour is preceded by various complaints before its appearance in the thigh. In a horizontal posture, the abscess also totally disappears, while the hernia does not.

Lumbar region. The loins.

LUMBRICALES MANUS. (*Lumbricales*, sc. *musculus*; from their resemblance to the *lumbricus*, or earth-worm.) *Fidicinales*. *Flexor primi internodii digitorum manus vel perforatus lumbricales* of Cowper. *Anuli tendino phalangiens* of Dumas. The four small fingers of the flexors of the fingers, which assist the bending of the fingers when the long flexors are in full action. They arise thin and fleshy from the outside of the tendons of the flexor profundus, a little above the lower edge of the carpal ligaments, and are inserted by long slender tendons into the outer sides of the broad tendons of the interosseal muscles, about the middle of the first joint of the fingers.

LUMBRICALES PEDIS. *Plantæ tendino phalangiens* of Dumas. Four muscles like the former, that increase the flexion of the toes, and draw them inward.

LUMBRICUS. (*à lubricitate*, from its slipperiness.) *Ascaris lumbricoides*. *Lumbricus terre*. The long round worm. A species of worm which inhabits occasionally the human intestines. It has three nipples at its head, and a triangular mouth in its middle. Its length is from four to twelve inches, and its thickness, when twelve inches long, about that of a goose-quill. They are some-

times solitary, at other times very numerous.

LUMBRICUS TERRESTRIS. See *Earthworm*.

LUNBUS VENERIS. See *Millesfolium*.

LUNA. (So named from its resemblance in brightness to silver.) The old alchemical name of silver.

Lunar caustic. See *Nitras argenti*.

LUNARE OS. One of the bones of the wrist.

LUNATICA ISCHURIA. (From *luna*, the moon.) A suppression of urine which returns monthly. It is noticed by Sauvages.

LUNG. *Pulmo.* The lungs are two viscera situated in the chest, by means of which we breathe. The lung, in the right cavity of the chest, is divided into three lobes, that in the left cavity into two. They hang in the chest, attached at their superior part to the neck, by means of the trachea, and are separated by the mediastinum. They are also attached to the heart, by means of the pulmonary vessels. The substance of the lungs is of four kinds, viz, vesicular, vascular, bronchial, and a parenchymatous substance. The vesicular substance is composed of the air-cells. The vascular invests those cells like a net-work. The bronchial is throughout the lungs, having the air cells at their extremities: and the spongy substance that connects the spaces between these parts is termed the *parenchyma*. The lungs are covered with a fine membrane, a reflexion of the pleura, called *pleura pulmonalis*. The internal surface of the air-cells is covered with a very fine, delicate, and sensible membrane, which is continued from the larynx through the trachea and bronchia. The arteries of the lungs are the pulmonary, which circulate the blood through the air-cells to undergo a certain change, and the bronchial artery, a branch of the aorta, which carries blood to the lungs for their nourishment. The pulmonary veins return the blood that has undergone this change, by four trunks, into the left auricle of the heart. The bronchial veins terminate in the vena azygos. The nerves of the lungs are from the eighth pair and great intercostal. The absorbents are of two orders; the superficial and deep-seated: the former are more readily detected than the latter. The glands of these viscera are called bronchial. They are muciparous, and situated about the bronchia. See *Respiration*.

Lung-worm, spotted. See *Pulmonaria maculata*.

LUPIA. (From *lupus*, to molest.) A genus of disease, including encysted tumours, whose contents are very thick, and sometimes solid, as *meliceria*, *atherama*, *steatoma*, and *ganglion*.

LUPULUS. (From *lupus*, grief, or dislike;

so called from its extreme bitterness.) Under this term the white lupin is directed in some pharmacopœias. The seed, the ordinary food of mankind in the days of Galen and Pliny, is now forgotten. Its farinaceous and bitter meal is occasionally exhibited to remove worms from the intestines, and made into poultices to resolve indolent tumours.

LUPINUS ALBUS. The systematic name of the white lupin. See *Lupinus*.

LUPULUS. (From *lupus*, dislike; so named from its bitterness.) *Convolutus perennis*. The hop. It is the floral leaf or bractea of this plant, *Humulus lupulus* of Linnæus, that is dried and used in various kinds of strong beer. Hops have a bitter taste, less ungrateful than most of the other strong bitters, accompanied with some degree of warmth and aromatic bitter, and are highly intoxicating. The hop-flower also exhales a considerable quantity of its narcotic power in drying; hence those who sleep in the hop-houses are with difficulty roused from their slumber. A pillar stuffed with these flowers is said to have laid our present monarch to sleep when other remedies had failed.

LUPUS. The wolf, so named from its rapacity. The cancer is also so called, because it eats away the flesh like a wolf.

LUSTRAGO. (From *lustrum*, to expiate; so called because it was used in the ancient purifications.) Flat or base vervain.

Lute. See *Lutum*.

LUTEA CORPORA. See *Corpora lutea*.

LUTEOLA. (From *lutum*, mud, because it grows in muddy places, or is the colour of mud.) *Struthium*. Dyer's weed. Dioscorides recommends it as useful in jaundice, but it is now neglected.

LUTUM. (From *lutos*, soluble.) *Cementum*. Mud. **Lute.** A composition with which chymical vessels are covered, to preserve them from the violence of the fire, and to close exactly their joinings to each other, to retain the substances which they contain when they are volatile and reduced to vapour.

LUXATION. (*Luxatura*; from *luxo*, to put out of joint.) A dislocation of a bone from its proper cavity.

LYCHARGE. (From *lykos*, a wolf, and *argos*, to strangle.) A species of quincy, in which the patient makes a noise like the howling of a wolf.

LYCHANTHROPIA. (From *lykos*, a wolf, and *anthropos*, a man.) A species of insanity, in which the patients leave their houses in the night, and wander about like wolves, in unfrequented places.

LYCHNIS. (From *lychnis*, a torch; because the ancients used its leaves rolled up for torches.) A name of several vegetable productions.

LYCOTRICHUM. (From *lykos*, a wolf, and *trichum*, to slay; so called because it was

the custom of hunters to secrete it in raw flesh, for the purpose of destroying wolves.)

LYCOPERDON. (From *λυκος*, a wolf, and *σπινθηρ*, to break wind; so named because it was supposed to spring from the dung of wolves.) 1. The name of a genus of plants in the Linnæan system. Class, *Cryptogamia*. Order, *Fungi*.

2. The pharmacopœial name of the puff-ball. *Crepitus lupi*. A round or egg shaped fungus, the *Lycoperdon bovista* of Linnæus, when fresh, of a white colour, with a very short, or scarcely any pedicle, growing in dry pasture grounds. When young, it is sometimes covered with tubercles on the outside, and is pulpy within. By age it becomes smooth externally, and dries internally, into a very fine, light, brownish dust, which is used by the common people to stop hæmorrhages.

LYCOPERDON BOVISTA. The systematic name of the puff-ball. See *Lycoperdon*.

LYCOPERDON TUBER. The systematic name of the truffle. See *Truffle*.

LYCOPERSICON. (From *λυκος*, a wolf, and *πικρον*, a peach; so called from its exciting a violent degree of lust.) Wolf's peach. Poisonous.

LYCOPODIUM. (From *λυκος*, a wolf, and *πους*, a foot; so called from its supposed resemblance.) 1. The name of a genus of plants in the Linnæan system. Class, *Cryptogamia*. Order, *Musci*.

2. The pharmacopœial name of the *Muscus clavatus*. Club moss. Wolf's claw. This plant, *Lycopodium clavatum* of Linnæus, affords a great quantity of pollen, which is much esteemed in some places to sprinkle on young children, to prevent, and in the curing parts which are fretting. A decoction of the herb is said to be a specific in the cure of the plica polonica.

LYCOPODIUM CLAVATUM. The systematic name of the club-moss. See *Lycopodium*.

LYCOPODIUM SELAGO. The systematic name of the upright club-moss. See *Muscus erectus*.

LYCOPSIS. (From *λυκος*, a wolf, and *οψη*, an aspect; so called from its being of the colour of a wolf.) *Echium Ægyptiacum*, or wall bugloss.

LYCOPUS. (From *λυκος*, a wolf, and *πους*, a foot; so named from its likeness.) The name of a genus of plants in the Linnæan system. Class, *Diandria*. Order, *Monogynia*. Wolf's claw, or water horehound. The *Lycopus europæus* is astringent.

LYOISMUS. (From *λυω*, to distort.) A dislocation.

LYEUS. (From *λυω*, to bend; so called from its flexibility.) The agnus castus.

LYMPH. *Lympha*. The liquid contained in the lymphatic vessels. It has a stitious smell, no taste, and is of a crystalline colour. Its specific gravity is greater than water; in consistence, it is thin and

somewhat plastic. The quantity in the human body appears to be very great, as the system of the lymphatic vessels forms no small part of it. Its constituent principles appear to be gelatinous albuminous water. The lymphatic vessels absorb this water from the tela cellulosa of the whole body, from all the viscera and the cavities of the viscera; and convey it to the thoracic duct, where it is mixed with the chyle.

The use of the lymph is to turn the superfluous nutritious jelly from every part, and to mix it with the chyle in the thoracic duct, there to be further converted into the nature of the animal; and lastly, it has mixed with it the superfluous aqueous vapour, which is expired into the cavity of the cranium, thorax, and abdomen.

LYMPHATIC GLANDS. *Glandule lymphaticæ.* See *Conglobate glands*.

LYMPHATICS. Absorbent vessels, that carry a transparent fluid, or lymph. They are small and transparent, and originate in every part of the body. With the lacteal vessels of the intestines, they form what is termed the *absorbent system*. Their termination is in the thoracic duct. See *Absorbent vessels*, *Lacteals*, and *Thoracic duct*.

Lymphatics of the head and neck.—Absorbents are found on the scalp and about the viscera of the neck, which unite into a considerable branch, that accompanies the jugular vein. Absorbents have not been detected in the human brain; yet there can be no doubt of there being such vessels: it is probable that they pass out of the cranium through the canalis caroticus and foramen lacerum in basi cranii, on each side, and join the above *jugular branch*, which passes through some glands as it proceeds into the chest to the angle of the subclavian and jugular veins.

The absorbents, from the right side of the head and neck, and from the right arm, do not run across the neck, to unite with the great trunk of the system; they have an equal opportunity of dropping their contents into the angle betwixt the right subclavian and the jugular vein. These vessels then uniting, form a trunk, which is little more than an inch, nay, sometimes not a quarter of an inch, in length, but which has nearly as great a diameter as the proper trunk of the left side.

This vessel lies upon the right subclavian vein, and receives a very considerable number of lymphatic vessels; not only does it receive the lymphatics from the right side of the head, thyroid gland, neck, &c. and the lymphatics of the arm, but it receives also those from the right side of the thorax and diaphragm, from the lungs of this side, and from the parts supplied by the mammary artery. Both in this and in the great trunk there are many valves.

Lymphatics of the upper extremities.—

The absorbents of the upper extremities are divided into superficial and deep-seated. The *superficial absorbents* ascend under the skin in every direction to the wrist, from whence a *branch* proceeds upon the posterior surface of the fore-arm to the head of the radius, over the internal condyle of the humerus, up to the axilla, receiving several branches as it proceeds. Another *branch* proceeds from the wrist along the anterior part of the fore-arm, and forms a *net-work*, with a *branch* coming over the ulna from the posterior part, and ascends on the inside of the humerus to the glands of the axilla. The *deep-seated absorbents* accompany the larger blood-vessels, and pass through two glands about the middle of the humerus, and ascend to the glands of the axilla. The superficial and deep-seated absorbents, having passed through the axillary glands, form *two trunks*, which unite into *one*, to be inserted with the jugular absorbents into the thoracic duct; at the angle formed by the union of the subclavian with the jugular vein.

Lymphatics of the inferior extremities.—

These are also superficial and deep-seated. The *superficial ones* lie between the skin and muscles. Those of the toes and foot form a *branch*, which ascends upon the back of the foot, over the tendon of the *cruzus anticus*, forms with other branches a *plexus* above the ankles, then proceeds along the tibia over the knee, sometimes passes through a gland, and proceeds up the inside of the thigh, to the subinguinal glands. The *deep-seated absorbents* follow the course of the arteries, and accompany the femoral artery, in which course they pass through some glands in the leg and above the knee, and then proceed to some deep-seated subinguinal glands. The absorbents from about the external parts of the pubis, as the penis, perineum, and from the external parts of the pelvis, in general proceed to the inguinal glands. The subinguinal and inguinal glands send forth several branches, which pass through the abdominal ring into the cavity of the abdomen.

Lymphatics of the abdominal and thoracic viscera.—

The absorbents of the lower extremities accompany the external iliac artery, where they are joined by many branches from the *uterus*, *urinary bladder*, *spermatic chord*, and some branches accompanying the internal iliac artery; they then ascend to the sacrum, where they form a *plexus*, which proceeds over the *psaos muscles*, and meeting with the lacteals of the mesentery, form the thoracic duct, or trunk of the absorbents, which is of a serpentine form, about the size of a crow-quill, and runs up the dorsal vertebrae, through the posterior opening of the diaphragm, between the aorta and vena azygos, to the angle formed by the union of

the subclavian and jugular veins. In this course it receives:—the *absorbents of the kidneys*, which are superficial and deep-seated, and unite as they proceed towards the thoracic duct: and the *absorbents of the spleen*, which are upon its peritoneal coat, and unite with those of the pancreas:—a *branch* from a plexus of vessels passing above and below the duodenum, and formed by the absorbents of the *stomach*, which come from the lesser and greater curvature, and are united about the pylorus with those of the pancreas and liver, which converge from the external surface and internal parts towards the porta of the liver, and also by several branches from the *gall-bladder*.

Use of Lymphatics.—The office of these vessels is to take up substances which are applied to their mouths; thus the vapour of circumscribed cavities, and of the cells of the cellular membrane, are removed by the lymphatics of those parts; and thus mercury and other substances are taken into the system when rubbed on the skin.

The principle by which this absorption takes place, is a power inherent in the mouths of absorbing vessels, a vis insita, dependent on the high degree of irritability of their internal membrane, by which the vessels contract and propel the fluid forwards. Hence the use of this function appears to be of the utmost importance, viz. to supply the blood with chyle; to remove the superfluous vapour of circumscribed cavities, otherwise dropsies, as hydrocephalus, hydrothorax, hydrocordis, ascites, hydrocele, &c. would constantly be taking place: to remove the superfluous vapour from the cells of the cellular membrane dispersed throughout every part of the body, that anasarca may not take place: to remove the hard and soft parts of the body, and to convey into the system medicines which are applied to the surface of the body.

LYPOMA. See *Lipoma*.

LYRA. (From *lyra*, a lyre, or musical instrument.) *Psalterium*. The triangular medullary space between the posterior crura of the fornix of the cerebrum, which is marked with prominent medullary fibres that give the appearance of a lyre.

LYRAUS. (From *lyra*, the lyre; so called because its leaves are divided like the strings of a lyre.) The *dornicum Germanicum*, or German leopard's-bane.

LYSISTIA. (From *lyse*, to loosen, and *stia*, a member.) The relaxation of limbs.

LYSIMACHIA. (From *Lysimachus*, who first discovered it.) The name of a genus of plants in the Linnean system. Class, *Pentandria*. Order, *Monogynia*.

LYSIMACHIA NUMMULARIA. The systematic name of the money-wort. See *Nummularia*.

LYSIMACHIA PURPUREA. The herb,

hoot, and flowers of this plant, *Lythrum salicaria* of Linnaeus, possess a considerable degree of astringency, and are used medicinally in the cure of diarrhoeas and dysenteries, fluor albus, and hæmoptysis.

LYSSOMECTUS. (From *lyssa*, canine madness, and *decupio*, to bite.) One who is mad in consequence of having been bitten by a mad animal.

LYTHRUM. (From *lythron*, blood; so

called from its resemblance in colour, and *Salicaria*, from *salix*, a willow; from the resemblance of its leaves to those of a willow.) The name of a genus of plants in the Linnaean system. Class, *Dodecandria*. Order, *Digynia*.

LYTHRUM SALICARIA. The systematic name of the common or purple willow-herb. See *Lythmachia purpurea*.

Mary E. Grace
went to Heaven M. Sept. 10th 1842

M. This letter has two significations: when herbs, flowers, chips, or such-like substances, are ordered in a prescription, and M. follows them, it signifies *manipulus*, a handful; and when any quantity of other ingredients is directed, it is a contraction of *misce*; thus, *m. f. mist.* signifies mix and make a mixture.

MACANDON. (Indian.) A tree growing in Malabar, whose fruit is roasted and eaten as a cure for dysenteries and relief in cholera morbus, and other complaints.

MACAPATI. Sarsaparilla.

MACAXOCOTLIVERA. The name of a tree in the West Indies, whose fruit is sweet and laxative. A decoction of the bark of this tree cures the itch, and the powder thereof heals ulcers.

Macedonian parsley. See *Petroselinum Macedonicum*.

MACEDONISUM SEMEN. The seeds of the *Smyrniolum olusatrum*, said to possess bitterish, aromatic, and carminative virtues.

MACER. (From *masa*, Heb.) Grecian macer, or mace. The root which is imported from Barbary by this name is supposed to be the simarouba, and is said to be antidyenteric.

MACÉRATION. (From *macero*, to soften by water.) In a pharmaceutical sense, this term implies an infusion either with or without heat, wherein the ingredients are intended to be almost wholly dissolved, in order to extract their virtues.

MACRONA. The *Smyrniolum olusatrum* or herb Alexander.

MACHERIA. The persicaria, or peach-kernels.

MACHÆRION. *Macheris*. The amputating-knife.

MACHAON. The proper name of an ancient physician, said to be one of the sons of Æsculapius; whence some authors have fancied to dignify their own inventions with his name, as particularly a collyrium, described by Scribonius, intitled, *Asclepias Machaonis*; and hence, also, medicine in general is by some called *Are Machaonia*.

MACHINAMENTUM ARISTIONIS. A machine for reducing dislocation.

MACHINULE. A word sometimes used by physical writers to express those little compositions, which are parts of stone. It is a sort of rocky marle.

MACIA. The anagallis.

MACIES. A wasting of the body. See *Atrophy* and *Tubes*.

MACIS. Mace. The middle bark of the nutmeg. A thick, tough, reticulated unctuous membrane, of a lively, reddish, yellow colour, approaching to that of saffron, which envelopes the shell of the nutmeg. See *Nux moschata*. The mace, when fresh, is of a blood-red colour, and acquires its yellow hue in drying. It is dried in the sun, upon hurdles fixed above one another, and then, it is said, sprinkled with seawater, to prevent its crumbling in carrying. It has a pleasant aromatic smell, and a warm, bitterish, moderately pungent taste. It is in common use as a grateful spice, and appears to be in its general qualities nearly similar to the nutmeg. The principal difference consists in the mace being much warmer, more bitter, less unctuous, and sitting easier on weak stomachs.

Mace possesses qualities similar to those of nutmeg, but is less astringent, and its oil is supposed to be more volatile and acrid.

MACRE. The macer.

MACROPHYSOCEPHALUS. (From *μακρος*, long, *φύσις*, nature, and *κεφαλή*, the head.) One who has a head unnaturally long and large. This word, according to Turton, is only used by Ambrose Parey.

MACROPIPER. (From *μακρος*, long, and *πιπέρων*, pepper.) See *Piper longum*.

MACROPNÆA. (From *μακρος*, long, and *πνέω*, to breathe.) A difficulty of breathing, where the inspirations are at long intervals.

MACULA. A spot. A permanent discolouration of some portion of the skin, often with a change of its texture, but not

connected with any disorder of the constitution.

MACULE ALBÆ. White specks on the eye.

MACULE HEPATICÆ. Hepatic, or liver-coloured spots on the skin.

MACULE LATÆ. Shingles, or erysipelas.

MACULE OCULORUM. Cataracts; white specks on the eye.

MACULE PESTILENTES. Petechial, or purple spots.

MACULE VENEREÆ. The venereal eruption.

MACULE VOLATICÆ. Any transitory eruption.

MAD-APPLES. The oblong egg-shaped fruit of the *Solanum melongena* of Linnæus. They are often boiled, in their native places, in soups and sauces, the same as the love-apple; are accounted very nutritive, and are much sought after by the votaries of Venus.

MADAROSIS. (From *μαδω*, bald, without hair.) A defect or loss of eye-brows, or eye-lashes, causing a disagreeable deformity, and painful sensation of the eyes in a strong light.

Madder. See *Rubia*.

Madness. See *Melancholia*, *Insania*, and *Mania*.

Madness, Canine. See *Hydrophobia*.

MADOR. Moisture. A sweating. See *Ephidrosis*.

MAGDALEON. (From *μασσω*, to knead.) A mass of plaster, or other composition, reduced to a cylindrical form.

MAGELLANICUS CORTEX. The *Winteranus cortex*, nearly allied in its properties to *canella alba*.

MAGISTERY. (From *magister*, a master.) The ancient chymists used this word to signify a peculiar and secret method of preparing any medicine, as it were, by a masterly process. A subtle preparation, as a precipitate or solution by menstruum.

MAGISTRALIA. (From *magister*, a master.) Applied, by way of eminence, to such medicines as are extemporaneous, or in common use.

MAGISTRANTIA. (From *magistro*, to rule; so called by way of eminence, as exceeding all others in virtue.) See *Imperatoria*.

MAGMA. (From *μασσω*, to blend together.) *Epilema*. A thick ointment. The faces of an ointment after the thinner parts are strained off. A confection.

MAGNES. (From *Magnes*, its inventor.) The magnet, or load-stone. A muddy iron ore, in which the iron is modified in such a manner as to afford a passage to a fluid called the magnetic fluid. The magnet exhibits certain phenomena; it is known by its property of attracting steel filings, and is found in Auvergne, in Biscay, in Spain, in Sweden, and Siberia.

MAGNES ARSENICALIS. Arsenical magnet. It is a composition of equal parts of antimony, sulphur, and arsenic, mixed and melted together, so as to become a glassy body.

MAGNES EPILEPSIÆ. The native cinabar.

MAGNESIA. 1. The ancient chymists gave this name to such substances as they conceived to have the power of attracting any principle from the air. Thus an earth which, on being exposed to the air, increased in weight, and yielded vitriol, they called *magnesia vitriolata*; and later chymists observing, in their process, that a nitrous acid was separated, and an earth left behind, supposing it had attracted the acid, called it *magnesia nitri*, which, from its colour, soon obtained the name of *magnesia alba*.

2. An earth not found pure in nature, but obtained by art from some of its combinations. It gives a peculiar character to the substances of which it forms a part. The stones which contain magnesia in a considerable quantity have generally a smooth and unctuous feel, a greenish cast, a fibrous or striated texture, and a silky lustre. Among them we may mention *talc*, *steatite*, *serpentine*, *chlorite*, *asbestos*, *actynolite*, *jaule*, or *nephritic stone*, *baikalite*, *boracite*, &c. It is likewise found neutralized with various acids. It has been discovered by Vauquelin in several sea-plants.

Properties.—Pure magnesia does not form with water an adhesive ductile mass. It is in the form of a very white spongy powder, soft to the touch, and perfectly tasteless. It is very slightly soluble in water. It absorbs carbonic acid gradually from the atmosphere. It changes very delicate blue vegetable colours to green. Its attraction to the acids is weaker than those of the alkalis. Its salts are partially decomposed by ammonia, one part of the magnesia being precipitated, and the other forming a triple compound. Its specific gravity is about 2.3. It is infusible even by the most intense heat; but when mixed with some of the other earths it becomes fusible. It combines with sulphur. It does not unite to phosphorus or carbon. It is not dissolved by alkalis in the humid way. When heated strongly, it becomes phosphorescent. With the dense acids it becomes ignited. With all the acids it forms salts of a bitter taste, mostly very soluble.

Method of obtaining Magnesia. The usual method of procuring magnesia is, to precipitate it from sulphate of magnesia by means of an alkali.

To effect this, dissolve any quantity of sulphate of magnesia in a large quantity of distilled water, and add to it gradually a solution of perfectly pure carbonate of pot-

tab, or soda, till no more precipitate ensues. Then collect the precipitate on a cloth, and boil it repeatedly in a large quantity of distilled water, till this fluid passes perfectly tasteless. It is then to be dried, and exposed in a crucible to a red heat, till a sample of it, when cold, does not occasion the least effervescence with acids.

In this process, a double decomposition takes place, the sulphuric acid of the sulphate of magnesia combines with the alkali, and forms sulphate of potash; and the carbonic acid of the alkali joins to the disengaged magnesia, and forms carbonate of magnesia; the latter is precipitated, and the sulphate of potash remains in solution. On exposing the carbonate of magnesia to heat, the acid is expelled, and the magnesia is left behind in a pure state. The magnesia of the present London Pharmacopoeia was formerly called *Magnesia calcinata; usta; pura*. It is directed to be made thus:—"Take of carbonate of magnesia, four ounces; burn it in a very strong fire, for two hours, or until acetic acid being dropped in extricates no bubbles of gas." It is given as an absorbent antacid, and ecceprotic, in cardialgia, spasms, convulsions, and tormina of the bowels of infants; pyrosis, flatulencies, and other diseases of the primæ viæ; obstipation, leucorrhœa, rickets, scrofula, crusta lactea, and podagra. The dose is from half a drachm to a drachm.

MAGNESIA CALCINATA. See *Magnesia*.

MAGNESIA OPALINA. In making the hepar antimonii, some add, to the antimony and nitre, decrepitated sal-ammoniac, and thus make the opalin. It is a weaker emetic than the liver of antimony.

MAGNESIA VITRIOLATA. See *Sulphas magnesiæ*.

MAGNESIA USTA. See *Magnesia*.

MAGNESIÆ CARBONAS. The London College direct it to be made as follows: Take of the sulphate of magnesia, subcarbonate of magnesia, of each a pound; water, three gallons. Dissolve the subcarbonate of potash in three pints of water, and strain. Dissolve also the sulphate of magnesia, separately, in five pints of the water, and strain; then add the rest of the water to the rest of the solution, apply heat, and when it boils pour in the former solution, stirring them well together; next, strain through a linen cloth; lastly, wash the powder repeatedly with boiling water, and dry it upon bibulous paper, in a heat of 200°. It is in form of very fine powder, considerably resembling flour in its appearance and feel; it has no sensible taste on the tongue; it gives a faint greenish colour to the tincture of violets, and converts turnsole to a blue. It is employed medi-

cinally as an absorbent, antacid, and purgative, in doses from half a drachm to two drachms. See *Carbonas magnesiæ*.

MAGNESIÆ SULPHAS. See *Sulphas magnesiæ*.

MAGNET. See *Magnes*.

MAGNETISM. The property which iron possesses of attracting or repelling other iron according to circumstances.

MAGNETISM, ANIMAL. A sympathy lately supposed, by some persons, to exist between the magnet and the human body; by means of which the former became capable of curing many diseases in an unknown way, somewhat resembling the performances of the old magicians. Animal magnetism is now entirely exploded.

MAGNUM DEI DONUM. So Dr. Mead calls the Peruvian bark.

MAGNUM OS. The third bone of the lower row of bones of the carpus, reckoning from the thumb towards the little finger.

MAGNUS MORBUS. The great disease. So Hippocrates calls the epilepsy.

MAGYDARIS. The root of the herb laserwort.

MAHAGONI. The systematic name of the tree the wood of which bears this name, and is so well known, is the *Swietenia mahagoni* of Linnæus. Its bark, when dried, has an adstringent bitter taste, similar to that of Peruvian bark, but stronger, for which it appears it may be substituted in the cure of fevers, and other diseases.

MARALEB. A species of *Prunus*.

MAHMOUDI. *Scammonium*.

Maidenhair. See *Adiantum*.

MAIDENHAIR, CANADA. *Adiantum Canadense*. This is the *Adiantum pedatum* of Linnæus. It is in common use in France, for the same purposes as the common adiantum is in this country, and appears to be far superior to it.

Maidenhair, English. See *Adiantum*.

MAIDENHAIR-TREE. *Ginkgo. Ginan Itsis*. In China and Japan, where this tree grows, the fruit acquires the size of a damask plumb, and contains a kernel resembling that of our apricot. These kernels always make part of the desert at all public feasts and entertainments. They are said to promote digestion, and to cleanse the stomach and bowels.

MAJANTHEMUM. See *Lilium convallium*.

MAIL-ANSCH. A species of *rhamnus* growing in Malabar. A decoction of its root is recommended against the gout; and a decoction of its leaves against the jaundice.

MAIL-CLOU. A Malabar tree, from whose bruised leaves and bark is prepared an apozem against the after pains of women in childbed, and for promoting the lochia.

MAIL-CLOU-KATOV. This is larger than

the above species. It is evergreen and astringent.

MAJORANA. (*Quod mense Maio flor.*, because it flowers in May.) Sweet marjoram. *Origanum majorana* of Linnaeus:—*foliis ovatis serratis, spicis subrotundis compactis pubescentibus*. This plant has been long cultivated in our gardens, and is in frequent use for culinary purposes. The leaves and tops have a pleasant smell, and a moderately warm, aromatic, bitterish taste. They yield their virtues to aqueous and spirituous liquors, by infusion, and to water in distillation, affording a considerable quantity of essential oil. The medicinal qualities of the plant are similar to those of the wild plant (see *Origanum*;) but being much more fragrant, it is thought to be more cephalic, and better adapted to those complaints known by the name of nervous; and may therefore be employed with the same intentions as lavender. It was directed in the *pulvis stomatatorius*, by both pharmacopœias, with a view to the agreeable odour which it diffuses to the asarabacca, rather than to its errhine power, which is very inconsiderable; but is now wholly omitted in the Pharm. Lond. In its recent state, it is said to have been successfully applied to schirrous tumours of the breast.

MAJORANA SYRIACA. See *Marum syriacum*.

MALA. (From *malus*, an apple; so called from its roundness.) A prominent part of the cheek.

MALA AESTIVA. The citron.

MALA ÆTHIOPICA. A species of *Licopersicon*.

MALA AURANTIA. See *Aurantium hispalense*.

MALA COTONEA. The quince.

MALA INSANA NIGRA. The fruit of the black fruited night-shade. See *Melengena*.

MALABAR PLUM. This fruit, which is the produce of the *Eugenia jambos*, smells, when ripe, like roses. On the coast of Malabar, where the trees grow plentifully, these plums are in great esteem. They are not only eaten fresh off the trees, but are preserved in sugar, in order to have them eatable all the year. Of the flowers, a conserve is prepared, which is used medicinally, as a mild astringent.

MALABATHRI OLEUM. Oil of cassia lignea.

MALABATHRUM. (*Malabathrum*: from *Malabar*, in India, whence it was brought, and *bethra*, a leaf, Ind.) The leaf of the tree whose bark is called cassia. See *Cassia lignea*.

MALABATHRUM. (From *malabathrum*, malabathrum.) Ointment of malabathrum. It is compounded of myrrh, spikenard, malabathrum, and many other aromatic ingredients.

Malaca bean. See *Anacardium orientale*.

MALACA RARY. The root of the *argathia alexipharmaca*.

MALACHE. (From *malachos*, soft; so called from the softness of its leaf.) The mallow.

MALACHITE. (From *malachy*, the mallow; from its resemblance in colour to the mallow.) A species of copper ore found in Siberia.

MALACIA. (From *malachos*, a ravenous fish.) *Pica*, or depraved appetite, when such things are coveted as are not proper for food.

MALACOSTEON. (From *malachos*, soft, and *ostion*, a bone.) *Mollities costarum*, or softness of the bones.

MALASTRA. (From *malassum*, to soften.) Emollient medicines.

MALÆ OS. (From *malus*, so called from its roundness.) The cheek-bone. See *Jugale os*.

MALASUETTA. *Malaguetta*. Grains of Paradise.

MALASMA. (From *malassum*, to soften.) *Basis*. It is synonymous with *Cataplasma*, from the frequency of making cataplasms to soften; but formerly *malagmar* were made of many other ingredients.

MALAXIRIS. A species of *Piper*.

MALARUM OSSA. The cheek-bones. See *Jugale os*.

MALATS. Salts formed by the union of the malic acid, or acid of apples, with different bases; thus *malat of copper*, *malat of lead*, &c.

MALE. The arm-pit.

Male fern. See *Filiix*.

Male orchis. See *Satyrion*.

Male speedwell. See *Veronica*.

MALIC ACID. *Acidum malicum*. This acid is obtained by saturating the juice of apples with alkali, and pouring in the acetous solution of lead, until it occasions no more precipitate. The precipitate is then to be edulcorated, and sulphuric acid poured on it, until the liquor has acquired a fresh acid taste, without any mixture of sweetness. The whole is then to be filtered, to separate the sulphate of lead. The filtered liquor is the malic acid, which is very pure, remains always in a fluid state, and cannot be rendered concrete. The union of this acid with different bases constitutes what are called *malats*.

MALIGNANT. *Malignus*. A term which may be applied to any disease, whose symptoms are so aggravated as to threaten destruction of the patient. It is frequently used to signify a dangerous epidemic.

Malignant fever. See *Typhus putrida*.

Malignant sore throat. See *Cynanche maligna*.

MALIS. A disease of the skin, produced by an insect lodging underneath. It is very common in Persia, where the disease is produced by the worm called *Gordius medinensis*, or *Dracunculus persicus*; in Ame-

rica, by the *Palex*; and it is sometimes produced in Europe by the *Pediculus*.

MALLAN-TODDALL. The name of a tree in Malabar, the root, bark, leaves, and fruit of which are esteemed, as a specific, in the epilepsy.

MALLEABILITY. (*Malleabilitas*; from *malleus*, a hammer.) The property which several metals possess of being extended under the hammer into thin plates, without cracking. The thin leaves of silver and gold are the best examples of malleability.

MALLANOTHE. *Pavette. Pavate. Erysi-pelas curans arbor.* A shrub which grows in Malabar. The leaves, boiled in palm oil, cure the impetigo; the root, powdered, and mixed with ginger, is diuretic.

MALLEI ANTERIOR. See *Laxator tympani*.

MALLEI EXTERNUS. See *Laxator tympani*.

MALLEI INTERNUS. See *Tensor tympani*.

MALLEOLUS. (Dim. of *malleus*, a mallet; so called from its supposed resemblance to a mallet.) The ankle, distinguished into external and internal, or *malleolus externus* and *internus*.

MALLEUS. (*Quasi molleus*; from *molle*, to soften; so called from its likeness to a little hammer.) A bone of the internal ear is so termed. It is distinguished into a head, neck, and manubrium. The head is round, and encrusted with a thin cartilage, and annexed to another bone of the ear, the incus, by ginglymus. Its neck is narrow, and situated between the head and manubrium, or handle; from which a long slender process arises, adheres to a furrow in the auditory canal, and is continued as far as the fissure in the articular cavity of the temporal bone. The manubrium is terminated by an enlarged extremity, and connected to the membrana tympani by a short conoid process.

Mallow, common. See *Malva*.

Mallow, round-leaved. See *Malva rotundifolia*.

Mallow, vervain. See *Malva alcea*.

MALOGRAIATUM. (From *malum*, an apple, and *granum*, a grain; so named from its grain-like seeds.) The pomegranate.

MALPHIGIA GLABRA. (So named in honour of Malpighius.) The systematic name of a tree which affords an esculent cherry.

MALTHE. (From *μαλασσω*, to soften.) *Malthacodes.* A medicine softened and tempered with wax.

MALTRACTICA (From *μαλθακίζω*, to soften.) Emollient medicines.

MALTHAURUM. Common salt.

MALUM. 1. A disease. 2. An apple.

3. In a strict sense, it is the disease called

Procidencia oculi; it is when the eyes exceed the bounds of the eye-lids.

MALUM MORTUUM. A disease that appears in the form of a pustule, which soon forms a dry, brown, hard, and broad crust. It is seldom attended with pain, and remains fixed for a long time before it can be detached. It is mostly observed on the tibia and on coccygis, and sometimes the face.

MALVA. (*Malva, quasi molva*; from *molle*, soft; named from the softness of its leaves.) 1. The name of a genus of plants in the Linnæan system. Class, *Monadelphia*. Order, *Polyandria*.

2. The pharmacopœical name of the common mallow. *Malva vulgaris Malva sylvestris* of Linæus:—*caule erecto herbaceo, foliis septemlobatis acutis, pedunculis petiolisque pilosis*. This indigenous plant has a strong affinity to the althæa, both in a botanical and a medical respect. See *Althæa*. The leaves and flowers are principally used in fomentations, cataplasms, and emollient enemas. The internal use of the leaves seems to be wholly superseded by the radix althææ.

MALVA ALCEA. The vervain mallow. The flowers of this plant are used medicinally in some countries.

MALVA ARBOREA. The mallow-tree. This beautiful tree is the *alcea rosacea* of Linæus. The flowers are said to possess adstringent and mucilaginous virtues. They are seldom used medicinally.

MALVA ROTUNDIFOLIA. Round-leaved mallow. The whole herb and root possess similar virtues to, and may be substituted for, the common mallow. See *Malva*.

MALVA SYLVESTRIS. The systematic name of the common mallow. See *Malva*.

MALVA VERBENACEA. *Alcea. Alcea vulgaris major.* Vervain mallow. This plant is distinguished from the common mallow, by its leaves being jagged, or cut in about the edges. It agrees in virtues with the other mallows, but it is least mucilaginous of any.

MALVAVISCUS. (From *malva*, the mallow, and *viscus*, glue; so named from its viscosity.) The marsh-mallow. See *Althæa officinalis*.

MALVA VULGARIS. See *Malva*.

MALVERN WATER. The village of Great Malvern has, for many years, been celebrated for a spring of remarkable purity, which has acquired the name of the holy well, from the reputed sanctity of its waters, and the real and extensive benefit long derived in various cases from its use.

The holy-well water, when first drawn, appears quite clear and pellucid, and does not become sensibly turbid on standing. It possesses somewhat of an agreeable

pungency to the taste; but this is not considerable. In other respects it does not differ in taste from pure good water.

The contents of Malvern holy-well are:—some carbonic acid, which is, in an uncombined state, capable of acting upon iron, and of giving a little taste to the water; but the exact quantity of which has not been ascertained:—a very small portion of earth, either lime or magnesia, united with the carbonic and marine acids:—perhaps a little neutral alkaline salt, and a very large proportion of water:—for we may add, that, the carbonic acid perhaps excepted, the foreign matter is less than that of any spring-water which we use. No iron, or metal of any kind, is found in it, though there are chalybeates in the neighbourhood.

It is singular, that, notwithstanding its apparent purity, this water is said not to keep well, and soon acquires a foetid smell, by standing in open vessels.

Malvern water, like many others, was at first only employed as an external application; and this, indeed, is still its principal use, though it is extended, with some advantage, to a few internal diseases. It has been found highly efficacious in painful and deep ulcerations, the consequence of a scrophulous habit of body, and which are always attended with much local irritation, and often general fever. Applied to the sore, it moderates the profuseness of the discharge, corrects the fetor, which so peculiarly marks a caries of the bone, promotes the granulating process, and a salutary exfoliation of the carious part; and by a long perseverance in this course, very dangerous and obstinate cases have at last been cured. Inflammation of the eye, especially the ophthalmia, which is so troublesome in scrophulous habits, often yields to this simple application, and we find that, for a great number of years, persons afflicted with sore eyes have been in the habit of resorting to Malvern holy-well. Another order of external diseases, for which this water is greatly celebrated, is cutaneous eruptions; even those obstinate cases of dry desquamations, that frequently follow a sudden application of cold in irritable habits, are often cured by this remedy. Where the skin is hot and dry, it remarkably relieves the intolerable itching of herpetic disorders, and renders the surface of the body more cool and perspirable. It appears, however, from a nice observation of Dr. Wall, that this method of treatment is not so successful in the cutaneous eruptions of very lax leucophlegmatic habits, where the extremities are cold and the circulation languid; but that it succeeds best where there is unusual irritation of the skin, and where it is apt to break in painful fissures, that ooze out a watery acrid lymph. On the first applica-

tion of this water to an inflamed surface, it will often for a time increase the pain and irritation, but these effects go off in a few days.

The great benefit arising from using Malvern waters, as an external remedy, in diseases of the skin, and surface of the body, have led to its employment in some internal disorders, and often with considerable advantage. Of these, the most important are, painful affections of the kidneys and bladder, attended with the discharge of bloody, purulent, or foetid urine; the hectic fever, produced by scrophulous ulceration of the lungs, or very extensive and irritating sores on the surface of the body; and also fistulas of long standing, that have been neglected, and have become constant and troublesome sores.

The Malvern water is in general a perfectly safe application, and may be used with the utmost freedom, both as an external dressing for sores and as a common drink.

The internal use of Malvern waters is sometimes attended at first with a slight nausea, and, not unfrequently, for the first day or two, it occasions some degree of drowsiness, vertigo, or slight pain of the head, which comes on a few minutes after drinking it. These symptoms go off spontaneously, after a few days, or may readily be removed by a mild purgative. The effects of this water on the bowels are not at all constant; frequently it purges briskly for a few days, but it is not uncommon for the body to be rendered costive by its use, especially, as Dr. Wall observes, with those who are accustomed to malt liquors. In all cases it decidedly increases the flow of urine, and the general health of the patient. The duration of a course of Malvern waters must vary very considerably, on account of the different kinds of disease for which this spring is resorted to.

MAMEI. The *mammoe*, *momin*, or toddy tree. This tree is found in different parts of the West Indies, but those on the Island of Hispaniola are the best. From incisions made in the branches, a copious discharge of pellucid liquor is obtained, which is called *momin*, or toddy wine; it must be drank very sparingly, because of its very diuretic quality. It is esteemed as an effectual preservative from the stone, as also a solvent of it when generated. There are two species.

MAMILLÆ. (Dim. *mamma*, the breast.) The breasts of men are so termed. It is likewise applied sometimes to the nipple.

MAMIRA. It is said, by Paulus Ægineta, to be the root of a plant which is of a detergent quality. Some think it is the root of the *doronicum*; but what it really is cannot be ascertained.

MAMMA. See *Breast*.

MAMMARY ARTERIES. *Ateriae mamillares.* The internal mammary artery is a branch of the subclavian, and gives off the mediastinal, thymal, and pericardiac arteries. The external mammary is a branch of the axillary artery.

MAMMARY VEINS. *Vene mamillares.* These vessels accompany the arteries, and evacuate their blood into the subclavian vein.

MAMMEE AMERICANA. The systematic name of the tree on which the mammee fruit grows. See *Mammee*.

MAMMEE. A delicious fruit, the produce of the *Mammea Americana* of Linnaeus. They have a very grateful flavour when ripe, and are much cultivated in Jamaica, where they are generally sold in the markets for one of the best fruits of the island.

MAN. *Homo.* Man is compounded of solids, fluids, a vital principle, and, what distinguishes him from every other animal, a soul.

I. The solids are divided into hard and soft, which analysis demonstrates to be formed of earthy particles, connected together by an intermediate gluten. The hard parts are the bones and cartilages. The soft parts, muscles, nerves, the viscera, and every other part except the fluids. See *Bones, Cartilages, Ligaments, Muscles, Arteries, Veins, Nerves, Lymphatics, Viscera, &c.*

II. The fluids are very various. See *Fluids*.

Anatomy demonstrates the structure of the various parts of which the human body consists. Chymistry has, of late, made great progress towards ascertaining its principles and elements, which are as follows :

The constituent principles of man are, 1. *The water*, which constitutes the greatest part of the humours, and is the vehicle of the other principles. 2. *The animal gas*, which consists of carbonated hydrogen, and is found, not only in the blood, but in all the other fluids. 3. *The inflammable gas*, emitted from the large intestines, *in flatu*. 4. *The animal gluten*, which consists of carbon and azote, and forms the fibres of the solid parts; the caseous portion of the milk; and the crur of the blood. 5. *The albumen*, present in the serum of the blood. 6. *The jelly*, found in the serum of the blood; lymph of the lymphatic vessels, and other fluids; and the interstices of all the fibres. 7. *The crur*, which is the animal gluten impregnated with oxydated iron. 8. *The mucus*, which lubricates the primæ viæ; the aerial surfaces of the lungs; the parts of generation, and the urinary passages. 9. *The animal oil*, which fills the cells of the adipose membrane. 10. *The resin*, found in

the bile. 11. *The sebatic acid*, which is present in animal oil. 12. *The phosphoric acid*, which enters into the composition of the animal earth of the bones, and the phosphorated salts of the urine. 13. *The lactic acid*, in the sugar of the serum of the milk. 14. *The sugar*, latent in the serum of the milk. 15. *The animal earth*, which is a phosphorated calx, and not only forms the greatest part of the bones, but also is found in the fibres of the soft parts, and in all the fluids. 16. *Phosphorated volatile alkali*; and 17. *Phosphorated soda*, both of which are detected in the urine. 18. *Culinary salt*, obtained from the urine, gastric juice, semen, and other humours.

The elementary principles of our body, hitherto known, are, 1. *Azot*, an element which, combined with hydrogen, constitutes volatile alkali; with the matter of heat, azotic air; with carbon, the gluten of animal fibres. *Azot* is the primary element of the animal body, for it may be extracted from almost every part of the animal, by means of the nitrous acid, this having a greater affinity with the elements than the azot itself. The mucus, jelly, membranes, tendons, ligaments, and cartilages, afford it in a less degree by means of the nitrous acid. The lymph, serum of the blood, the water of hydropic patients, the liquor amnii, and cheese give out more. The greatest quantity of azot is obtained from the coagulable lymph of the blood, and from muscle. The flesh of young animals contains less than that of old; and it is in greater quantity in sarco-phagous, than in the flesh of phytophagous animals and fish. It is not probable that the azot is produced by the decomposition of the acid of nitre; for, after having performed the separation, it is capable of saturating the same quantity of alkali as before. 2. *The matter of heat*, which enters into the composition of both solids and fluids, and which, in a separate form, constitutes the animal heat. 3. *The matter of light*, which in its free state produces vision, and, when compounded, enters as an element into the composition of oil and all other inflammable parts. The eyes of animals, which shine in the night time, owe this property to the matter of light. 4. *The electric matter*, which enters into all bodies, and affords the phenomena of animal electricity. 5. *Oxygen*, which, in combination with the matter of heat, constitutes vital air; with hydrogen, forms water; with accecent bases, the acid salts of our fluids. 6. *Hydrogen*, which, combined with oxygen, forms water; with azot, volatile alkali; with the matter of heat, inflammable air, which is emitted from the large intestines; and with carbon, animal gas; and lastly, combined with carbon and the sebatic acid, constitutes the oil of the adipose mem-

brane. 7. *Carbon*, which, in combination with hydrogen and the sebatic acid, constitutes the oil of the adipose membrane; with hydrogen alone, animal gaz; with azot, animal gluten. 8. *Sulphur*, which, combined with inflammable air, constitutes the hepatic air that exhales from muscular fibres, hair, incubated eggs, animal gluten, and, according to Lavoisier, human excrement. 9. *Phosphorus*, which, with oxygen, forms the phosphoric acid; and, with inflammable air, phosphoric air. The lucid sweat of some men, the phosphorescence, or light, given out by the putrefying bodies of some animals, and the phosphorus obtained from cheese and human bones, sufficiently shew that phosphorus constitutes an element of our body. 10. *Soda*, or the fixed mineral alkali. 11. *Potash*, or the fixed vegetable alkali. Each of these is found in several of the fluids of the human body. 12. *An earthy element*. Of the earths, no kind is so frequently detected as the *calcareous*, which is found in the bones and other parts. 13. *A metallic element*. Of so great a number of metals, iron and manganese alone are found in an organized body, whether animal or vegetable. Iron is in greater quantity in the flesh than in the bones; but in the greatest proportion in the crur or red part of the blood. 14. *An odorous principle*, perceptible in all the animal fluids; but of a peculiar kind in the human urine and excrements. 15. *The nervous fluid*, or principle contained in the nerve, and which appears to be an element *sui generis*, distinct from all known fluids, and not to be collected by art.

III. *The vital principle*. In all solid and fluid parts of a living body, there exists an element, with properties peculiar to itself, which constitutes life; hence it is justly called vital. This principle induces a mode of union in the other elements, widely differing from that which arises from the common laws of chemical affinity. By the aid of this principle, nature produces the animal fluids, as blood, bile, semen, and the rest, which can never be produced by the art of chymistry. But if, in consequence of death, the laws of vital attraction, or affinity, cease to operate, then the elements, recovering their former properties, become again obedient to the common laws of chymical affinity, and enter into new combinations, from which new principles, or the production of putrefaction, are produced. Thus the hydrogen, combining itself with the azot, forms volatile alkali; and the carbonated hydrogen, with the azot, putrid air, into which the whole body is converted. It also appears from hence, why organized bodies alone, namely, animal and vegetable, are subject to putridity, to which inorganic or mineral substances are in no degree liable, the latter

not being compounded according to the laws of vital affinity, but only according to those of chymical affinity. For the taticence, or resolution of the pyrites, or ferum sulphuratum, in the atmospheric air, is not putrefaction, but only the oxygen, furnished by the air, combining with the sulphur, and forming sulphuric acid.

Fire, as well as putridity, separates the constituent principles of animal bodies into their elements; but these, by a peculiar law, under the action of fire, again combine in a different manner, and form peculiar constituent principles, called the products of fire. Thus the hydrogen, combining with azot, is changed into volatile alkali; but with a large proportion of carbon, it forms empyreumatic oil. From what has hitherto been said, it will also appear, that the true constituent principles of the animal body cannot be detected, either by putrefaction or the action of fire; for by these means we only discover the elements of those principles. Thus, whenever volatile alkali is found to be generated, azot and hydrogen may be supposed to have been present in the natural state of the animal substance; and when empyreumatic oil is obtained, it may be concluded it is furnished by the hydrogen and carbon of the animal part.

MANACA. A Brazil shrub, whose root is powerfully emetic and cathartic.

MANCORON. According to Oribasius, a kind of sugar, which is found in a sort of cane.

MANCURANA. The *origanum vulgare*.

MANDIBULA. (From *mando*, to chew.) The lower jaw. See *Maxilla inferior*.

MANDRAGORA. (From *mandra*, a den, and *agora*, to collect; because it grows about caves and dens of beasts; or from the German *man dragen*, bearing man.) Mandrake. *Atropa mandragora* of Linnæus. The boiled root is employed in the form of poultice, to discuss indolent tumours.

MANDRAGORITES. (From *mandragora*, the mandrake.) Wine, in which the roots of the male mandrake are infused.

Mandrake. See *Mandragora*.

MANDUCATOR. (From *manduco*, to chew.) The muscles which perform the action of chewing.

MANGA. (Indian.) The mango-tree.

MANGANESE. This metallic substance seems, after iron, to be the most frequently diffused metal through the earth; its ores are very common. As a peculiar metal, it was first noticed by Galin and Scheele, in the years 1774 and 1777. It is always found in the state of an oxyd, varying in the degree of oxydation. La Peyrouse affirmed that he had found manganese in a metallic state; but there was probably some mistake in his observation

They are distinguished into *grey oxyd of manganese*, *black oxyd of manganese*, *reddish white oxyd of manganese*, and *carbonate of manganese*. All these combinations have an earthy texture; they are very ponderous; they occur both amorphous and crystallized; and generally contain a large quantity of iron. Their colour is black, blackish brown, or grey, seldom white. They soil the fingers like soot. They are sometimes crystallized in prisms, tetrahedral, rhomboidal, or striated.

Properties.—Manganese is of a whitish grey colour. Its fracture is granulated, irregular, and uneven. It is of a metallic brilliancy, which it, however, soon loses in the air. Its specific gravity is about 6.850. It is very hard, and extremely brittle. It is one of the most refractory metals, and most difficult to fuse, requiring at least 163° of Wedgwood's pyrometer. Its oxydability is so rapid, that exposure to the air is sufficient to render it red, brown, black, and friable, in a very short time; it can, therefore, only be kept under water, oil, or ardent spirit. It is the most combustible of all the metals. It decomposes water, by heat very rapidly, as well as the greater part of the metallic oxyds. It decomposes sulphuric acid. It is soluble in nitric acid. It is fusible with earthen, and colours them brown, violet, or red, according to its state of oxydation. It discolours glasses tinged by iron. It does not appear to unite with sulphur. It combines with phosphorus. It unites with gold, silver, and copper, and renders them brittle. It unites to arsenic in close vessels, but does not enter into union with mercury. It forms three differently coloured oxyds, by combining with different portions of oxygen.

Method of obtaining Manganese.—This metal is obtained by mixing the black oxyd, finely powdered, with pitch; making it into a ball, and putting this into a crucible, with powdered charcoal, one-tenth of an inch thick at the sides, and one-fourth of an inch deep at the bottom. The empty space is then to be filled with powdered charcoal; a cover is to be luted on; and the crucible exposed, for an hour, to the strongest heat that can be raised. Or, digest the black oxyd of manganese repeatedly, with the addition of one-sixteenth of sugar, in nitric acid; dilute the mixture with three times its bulk of water; filter it, and decompose it by the addition of potash; collect the precipitate, form it into a paste with oil, and put it into a crucible, well lined with charcoal. Expose the crucible for at least two hours to the strongest heat of a forge.

Manganese may also be obtained in the following manner:

Prepare a saturated solution of sulphate

of manganese, bring it to a boiling heat, and add to it, gradually, a solution of tartrate of potash, until no further precipitate ensues; then filter the solution, and wash the precipitate in water, and when dry make it into a paste with oil, and proceed as before.

In this process, the sulphuric acid unites to the potash, and forms sulphate of potash, and the tartareous acid joins to the manganese, and forms a tartrate of manganese, which is decomposable by heat.

MANGEL WUNSEL. The root of scarcity. A plant of great importance, as a substitute for bread in periods of famine. It has not, however, succeeded so well in this country as in Germany. It is properly a species of beet.

MANGIFERA INDICA. The systematic name of the mango-tree. See *Mango*.

Mango. The fruit of the *Mangifera Indica* of Linnaeus, which is cultivated all over Asia. When ripe, they are juicy, of a good flavour, and so fragrant as to perfume the air to a considerable distance. They are eaten either raw or preserved with sugar. Their taste is so luscious, that they soon pall the appetite. The unripe fruits are pickled in the milk of the cocoa-nut that has stood until sour, with salt, capsicum, and garlic. From the expressed juice is prepared a wine; and the remainder of the kernel can be reduced to an excellent flour for the making of bread.

MANGOSTANA. See *Mangosteen*.

MANGOSTEEN. A fruit about the size of an orange, which grows in great abundance on the tree called *Garcinia Mangostana* by Linnaeus, in Java and the Molucca islands. According to the concurring testimonies of all travellers, it is the most exquisitely flavoured, and the most salubrious of all fruits, it being such a delicious mixture of the tart and sweet. The flesh is juicy, white, almost transparent, and of a more delicate and agreeable flavour than the richest grape. It is eaten in almost every disorder, and the dried bark is used medicinally in dysenteries and tenesmus, and a strong decoction of it is much esteemed as a gargle in ulcerated sore throats.

Mangosteen bark. See *Mangosteen*.

MANIA. (From *μανία*, to rage.) Raving or furious madness. A genus of disease in the class *neuroses*, and order *vesania* of Cullen. The definition of mania is delirium, unaccompanied with fever; but this does not seem altogether correct, as a delirium may prevail without any frequency of pulse, or fever; as happens sometimes with women in the hysteric disease. In mania, the mind is not perfectly master of all its functions; it receives impressions from the senses, which are very different from those produced in health;

the judgment and memory are both lost, and the irritability of the body is much diminished, being incapable, it is supposed, of resisting the usual morbid effects of cold, hunger, and watching, and being likewise less susceptible of other diseases than before.

Mania may be said to be a false perception of things, marked by an incoherence, or raving, and in a resistance of the passions to the command of the will, accompanied, for the most part, with a violence of action, and furious resentment at restraint.

There are two species of madness, viz. the melancholic and furious.

Madness is occasioned by affections of the mind, such as anxiety, grief, love, religion, terror, or enthusiasm; the frequent and uncurbed indulgence in any passion, or emotion, and by abstruse study. It short, it may be produced by any thing that affects the mind so forcibly as to take off its attention from all other affairs. Violent exercise, frequent intoxication, a sedentary life, the suppression of periodical and occasional discharges and secretions, excessive evacuations, and paralytic seizures, are likewise enumerated as remote causes. Certain diseases of the febrile kind have been found to occasion madness, where their action has been very violent. In some cases it proceeds from an hereditary predisposition. Two constitutions are particularly the victims of madness; the sanguine and melancholic; by the difference of which its appearance is somewhat modified. Each species of mania is accompanied with particular symptoms. Those which attend on the melancholic are, sadness, dejection of spirits, and its attendants. Those which accompany an attack of furious madness are, severe pains in the head, redness of the face, noise in the ears, wildness of the countenance, rolling and glistening of the eyes, grinding of the teeth, loud roaring, violent exertion of strength, absurd, incoherent discourse, unaccountable malice to certain persons, particularly to the nearest relatives and friends, a dislike to such places and scenes as formerly afforded particular pleasure, a diminution of the irritability of the body, with respect to the morbid effects of cold, hunger and watching, together with a full, quick pulse.

Mania comes on at different periods of life; but, in the greater number of cases, it makes its attack between thirty and forty years of age. Females appear to be more subject to mania than males.

Dissections of maniacal cases, Dr. Thomas observes, most generally shew an effusion of water into the cavities of the brain; but, in some cases, we are able to discover evident marks of previous in-

flammation, such as thickening and opacity of the tunica arachnoides and pia mater. In a few instances, a preternatural hardness of the substance of the brain.

From Dr. Greding's observations, it appears that the skulls of the greater number of such persons are commonly very thick. Some he found of a most extraordinary degree of thickness; but it appears that the greater number of insane people die of atrophy and hydrothorax.

MANIHOT. The *Jatropha manihot*.

MANIPULUS. (*Quod manum, implet, because it fills the hand.*) A handful.

MANJAFUMERAM. A common tree in the West Indies, the flowers of which are distilled, and the water used against inflammations of the eyes.

MANNA. (From *mano*, a gift, Syr. it being the food given by God to the children of Israel in the wilderness; or from *madna*, what is it? an exclamation occasioned by their wonder at its appearance.) *Manna calabrina.* *Ros calabrinus.* *Acromeli.* *Alisar.* *Dryasomeli.* That species which is of a rosy colour, is called *maba.* *Mel aertum.* from the supposition that it descended from heaven. The condensed juice of the flowering ash, or, *Fraxinus grnus*:—*foliis ovato-oblongis serratis petiolatis, floribus corollatis.* Hort. Kew, which is a native of the southern parts of Europe, particularly Sicily and Calabria. Many other trees and shrubs have likewise been observed to emit a sweet juice, which concretes upon exposure to the air, and may be considered of the manna kind, especially the *Fraxinus rotundifolia* and *excoelior*. In Sicily, these three species of *fraxinus* are regularly cultivated for the purpose of procuring manna, and with this view are planted on the declivity of a hill with an eastern aspect. After ten years growth, the trees first begin to yield the manna, but they require to be much older, before they afford it in any considerable quantity. Although the manna exudes spontaneously upon the trees, yet, in order to obtain it more copiously, incisions are made through the bark, by means of a sharp crooked instrument; and the season thought to be most favourable for instituting this process is a little before the dog-days commence, when the weather is dry and serene. Manna is generally distinguished into different kinds, viz. the manna in tear, the canulated and flaky manna, and the common brown or fat manna. All these varieties seem rather to depend upon their respective purity, and the circumstance in which they are obtained from the plant, than upon any essential difference of the drug. The best manna is in oblong pieces, or flakes, moderately dry, friable, very light, of a whitish or pale yellow colour, and in some degree transparent: the inferior kinds are

moist, unctous, and brown. Manna is well known as a gentle purgative, so mild in its operation, that it may be given with safety to children and pregnant women, to the delicacy of whose frames and situation it is particularly adapted. It is esteemed a good and pleasant auxiliary to the purgative neutral salts. It sheathes acrimony, and is useful in coughs, disorders of the breast, and such as are attended with fever and inflammation, as in pleuritis, &c. Its is particularly efficacious in bilious complaints, and helps the discharge of mineral waters, when they are not of themselves sufficiently active. It is apt to create flatulencies and gripes; both which are prevented by a small addition of some warm carminative. It purges in doses of from $\mathfrak{z}\text{ij}$ to $\mathfrak{z}\text{ij}$; but its purgative quality is much increased, and its flatulent effects prevented, by a small addition of cassia. The dose for children is from one scruple to three. It is best dissolved in whey.

MANNA BRIGANTICA. A species of manna brought from Brianconois in Dauphiny.

MANNA CALABRINA. Calabrian manna.

MANNA CANULATA. Flaky manna, or manna concentered on straw, or chips.

MANNA SECUNDA. An inferior or second sort of manna.

MANNA THURIS. A coarse powder of olibanum.

MANNIFERA ARBOR. (From *manna*, and *fero*, to bear.) The *fraxinus ornaus*.

MASSORIUS. (From *mando*, to chew.) The masseter muscle.

MANTILE. The name of a bandage.

MANUS DEI. A name of a resolvent plaster, which is described by Lemery. Also for opium.

Mapple. See *Saccharum canadense*.

MARANDA. A species of myrtle, growing in the island of Zeylon, a decoction of the leaves of which are said to be excellent against the venereal disease.

MARANTA. 1. The name of a genus of plants in the Linnæan system. Class, *Monandria*. Order, *Monogymia*.

2. The name of the Indian arrow-root.

There are three species of Maranta, the *Arundinacea*, *Galanga*, and *Comesa*, all of them herbaceous, perennial exotics of the Indies, kept here in hot-houses for curiosity; they have thick, knotty, creeping roots, crowned with long, broad, arundinaceous leaves, ending in points, and upright stalks half a yard high, terminated by bunches of monopetalous, ringent, five-parted flowers. They are propagated by parting the roots in spring, and planting them in pots of light rich earth, and then plunging them in the bark-bed. The root of the *Maranta Galanga*, commonly called arrow-root, is used by the Indians to extract the virus communicated by their

poisoned arrows, from whence it has obtained its name. It is cultivated in gardens and provision-grounds in the West Indies; and the starch is obtained from it by the following process: The roots, when a year old, are dug up, well washed in water, and then beaten in a large deep wooden mortar to a pulp; this is thrown into a large tub of clean water: the whole is then well stirred, and the fibrous part wrung out by the hands, and thrown away. The milky liquor being passed through a hair sieve, or coarse cloth, is suffered to settle, and the clear water drained off. At the bottom of the vessel is a white mass, which is again mixed with clean water, and drained: lastly, the mass is dried on sheets in the sun, and is pure starch.

Arrow-root contains, in small bulk, a greater proportion of nourishment than any other yet known. The powder, boiled in water, forms a very pleasant transparent jelly, very superior to that of sago or tapioca; and is much recommended as a nutritious diet for children and invalids. The jelly is made in the following manner: to a desert spoonful of powder, add as much cold water as will make it into a paste; then pour on half a pint of boiling water: stir it briskly, and boil it a few minutes, when it will become a clear smooth jelly; a little sugar and sherry wine may be added for debilitated patients; but for infants, a drop or two of essence of caraway-seeds, or cinnamon, is preferable, wine being very liable to become acrescent in the stomachs of infants, and thus disagree with the bowels. Fresh milk, either alone or diluted with water, may be substituted for the water. For very debilitated frames, and especially for ricketty children, this jelly, blended with an animal jelly, as that of the stag's horn (*vasura cornu cervi*), affords a more nutritious diet than arrow-root alone, which may be done in the following manner: Boil half an ounce of stag's horn shavings, in a pint of water, for fifteen minutes; then strain, and add two desert spoonfuls of arrow-root powder, previously well mixed with a tea-cupful of water; stir them briskly together, and boil them for a few minutes. If the child should be much troubled with flatulency, two or three drops of essence of caraway-seeds, or a little grated nutmeg, may be added; but for adults, port wine, or brandy, will answer best.

MARASMUS. (From *marasmo*, to grow lean.) *Atrophæa*. Emaciation. A wasting away of the flesh, without fever or apparent disease. See *Atrophy*.

MARATHRITES. (From *μαραθρον*, fennel.) A vinous infusion of fennel; or wine impregnated with fennel.

MARATHROPHYLLUM. (From *μαραθρον*, fennel, and *φυλλον*, a leaf; so named

because its leaves resemble those of the common fennel.) See *Peucedanum officinale*.

MARATHRUM. (From *μαραν*, to wither; so called because its stalk and flowers wither in the autumn.) See *Faniculum*.

MARATHRUM SYLVESTR. See *Peucedanum*.

MARBLE. Powdered marble, which is a carbonate of lime, is used in pneumatic medicine, to give out carbonic acid gas.

Marcasite. See *Bismuth*.

MARCASSITA. (From *marcasite*, Germ.) See *Bismuth*.

MARCHANTIA POLYMORPHA. The systematic name of the liver-wort. See *Hepatica terrestris*.

MARCORES. (From *marces*, to become lean.) Universal emaciation. The first order in the class *cachexia* of Cullen's nology.

Muretail. See *Equisetum*.

MARGARITA. (From *margarith*, Rab.)

1. The pearl, *Perla*. *Uma*. A small calcareous concretion, of a bright transparent whiteness, found on the inside of the shell *Concha margaritifera* of Linnaeus, or mother-of-pearl fish. Pearls were formerly exhibited as antacids.

2. A tumour upon the eye resembling a pearl.

MARIGOLD, MARSH. *Caltha palustris* of Linnaeus. The flower buds of this very common plant may be pickled as a good substitute for capers.

Marine acid. See *Muriatic acid*.

Marine salt. See *Murias sode*.

MARIPENDAM. A plant in the island of St. Domingo: its tops are distilled, and thus a water is obtained, which is held in great esteem against pains in the stomach.

MARISCA. An excrescence about the anus, or the piles in a state of tumefaction; the *Hemorrhoids tumens* of Cullen.

MARISCUM. The mercurialis f. uticosr.

Marjoram, sweet. See *Majorana*.

Marjoram, wild. See *Origanum*.

MARJORANA. See *Origanum*.

MARMALADE. The pulp of quinces, or any other fruit, boiled into a consistence with honey.

MARMAHYGÆ. (From *μαρμαγε*, to shine.) An appearance of sparks, or coruscations, flushing before the eyes,

MARMOLARIA. (From *marmor*, marble; so named because it is spotted like marble.) Bear's-breech. See *Acanthus mollis*.

MARMORATA AURIUM. (From *marmor*.) Ear-wax.

MARMORIGE. A variety of the *Pseudoplepis Imaginaria*, in which sparks and flashes of fire are supposed to present themselves.

MARMOREUS TARTARUS. The hardest species of human calculus.

MAROCOSTINUM. A purgative extract made of the marum and costus; originally made by Minlereus.

MARROW. *Medulla.* The fat substance secreted by the small arteries of its proper membrane, and contained in the medullary cavities of the long cylindrical bones. See *Bone*.

Marrow, spinal. See *Medulla spinalis*.

MARRUBIASTRUM. Ballota, or stinking horehound.

MARRUBIUM. ((From *marrob*, a bitter juice, Heb.) *Marrub-marson*. 1. The name of a genus of plants in the Linnæan system. Class, *Didynamia*. Order, *Gymnospermia*.

2. The pharmacopœial name of the common white horehound. *Marrubium album*. *Marrubium vulgare* of Linnaeus:—*dentibus calycinis setaceis uncinatis*. The leaves of this indigenous plant have a moderately strong smell of the aromatic kind, but not agreeable; which, by drying, is improved; and in keeping for some months is in great part dissipated; their taste is very bitter, penetrating, diffusive, and durable in the mouth. That horehound possesses some share of medicinal power, may be inferred from its sensible qualities, but its virtues do not appear to be clearly ascertained. It is a favourite remedy with the common people in coughs and asthmas. The usual dose is from half an ounce to an ounce, in infusion, two or three times a day. The dose of the extract is from gr. x to 3ss.

MARRUBIUM ALBUM. See *Marrubium*.

MARRUBIUM AQUATICUM. Water horehound; opening, corroborant.

MARRUBIUM HISPANICUM. Mad-wort, or Spanish horehound.

MARRUBIUM NIGRUM FOETIDUM. The black stinking horehound, or ballota.

MARRUBIUM VERTICILLATUM. *Marrubium hispanicum*. The base horehound. Galen's madwort.

MARRUBIUM VULGARE. The systematic name of the common horehound. See *Marrubium*.

MARS. The alchemists gave this name to iron.

MARS ALKALIZATUS SOLUBILIS. Iron and fixed alkali.

MARS SACCHARATUS. Iron mixed with starch and melted sugar.

MARS SOLUBILIS. *Ferrum tartarizatum*.

MARS SULPHURATUS. Iron filings, and sulphur dephlegmated.

Marseilles hart-wort. See *Scellæ massiliense*.

Marsh mallow. See *Althea*.

Marsh trefoil. See *Trefoilium paludosum*.

MARSUPIALIS. (*Marsupialis*, &c *musculus*; from *marsupium*, a purse; so named from its resemblance.) See *Obturator internus*.

MARTAGON LILY. *Lilium martagon* of Linnaeus, who informs us the root makes part of the daily food of the Siberians.

MARTIAL. Sometimes used to express preparations of iron, or such as are impregnated therewith; as the *Martial Regulus* of antimony, &c.

MARTIATUM UNGUENTUM. Soldiers' ointment. Ointment of laurel, rue, marjoram, &c.

MARTIS ESSENTIA. A solution of lead in acid.

MARTIS LIMATURA PRÆPARATA. Purified filings of iron.

MARUM CRETICUM. See *Marum Syriacum*.

MARUM SYRIACUM. (From *mar*, bitter, Heb.) *Marum creticum*. *Majorana syriaca*. *Marum verum*. *Marum Coortus*. *Chamedrys incana maritima*. Marum germander, or Syrian herb mastich. This shrub is the *Teucrium marum* of Linnaeus: — *foliis integerrimis ovatis acutis petiolatis, subtus tomentosis, floribus sacmiosis secundis*. It grows plentifully in Greece, Egypt, Crete, and Syria. The leaves and younger branches, when recent, on being rubbed betwixt the fingers, emit a volatile aromatic smell, which readily excites sneezing; to the taste they are bitterish, accompanied with a sensation of heat and acrimony. Judging from these sensible qualities of the plant, it may be supposed to possess very active powers. It is recommended as a stimulant, aromatic, and deobstruent; and Linnaeus, Rosenstein, and Bergius, speak highly of its utility. Dose, ten grains to half a drachm of the powdered leaves, given in wine. At present, however, marum is chiefly used as an errhine.

MARUM VERUM. See *Marum syriacum*.

MARUM VULGARE. Common herb mastich. *Sampuchus*. *Clinopodium mastichina gallerum*. *Thymra hispanica*. *Jacquinia indica*. *Thymus mastichina* of Linnaeus. A low shrubby plant, a native of Spain, which is employed as an errhine. It has a strong agreeable smell, like mastich. Its virtues are similar to those of *marum syriacum*, but less powerful.

MARVISUM. Malmsey wine.

MASCHALE. (Μασχάλη.) The arm-pit.

MASCHALISTER. (From *μασχαλιστε*.) The second vertebra of the back.

MASLACH. A medicine of the opiate kind, in use amongst the Turks.

MASFETAM. *Maspetum*. The leaves of the *asafoetida* plant.

MASSA. (From *μασσα*, to blend together.) A mass. A term generally applied to the compound out of which pills are to be formed.

MASSA CARNEA JACOBI SYLVII. See *Flexor longus digitorum pedis*.

MASSALIS. A name for mercury.

MASSETER. (From *μασσαι*, to chew; because it assists in chewing.) *Zigomato-maxillaire* of Dumas. A muscle of the lower jaw, situate on the side of the face. It is a short thick muscle, which arises, by fleshy and tendinous fibres, from the lower edge of the malar process of the maxillary bone, the lower horizontal edge of the os malæ, and the lower edge of the zygomatic process of the temporal bone, as far backwards as the eminence belonging to the articulation of the lower jaw. From some little interruption in the fibres of this muscle, at their origin, some writers describe it as arising by two, and others by three, distinct portions, or heads. The two layers of fibres of which it seems to be composed cross each other as they descend, the external layer extending backwards, and the internal one slanting forwards. It is inserted into the basis of the coronoid process, and into all that part of the lower jaw which supports the coronoid and condyloid processes. Its use is to raise the lower jaw, and, by means of the above-mentioned decussation, to move it a little forwards and backwards in the act of chewing.

Massicot. Calcined white lead.

MASSOY CORTEX. See *Cortex massoy*.

Masterwort. See *Imperatoria*.

MASTICATION. (*Masticatio*; from *mastic*, to chew.) Chewing. A natural function. The mixing together and dividing of the particles of the food in the mouth, by the action of the jaws, tongue, lips, and cheeks. By means of this function, the food is lacerated and mixed with the saliva and the mucus of the mouth and fauces, and thus made into a bolus of such a consistence as to be formed into a convenient size to be swallowed. See *Deglutition*.

MASTICATORIES. (*Masticatoria*, sc. *medicamenta*; from *Mastico*, to chew.) Such medicines as are intended for chewing.

MASTICHE. (From *μασσα*, to express.) *Mastix*. Mastich. The tree which affords this resin is the *Pistachia lentiscus*; *foliis abrupte pinnatis, foliis lanceolatis* of Linnaeus. A native of the south of Europe. In the island of Chio the official mastich is obtained most abundantly, and, according to Tournefort, by making transverse incisions in the bark of the tree, from whence the mastich exudes in drops, which are suffered to run down to the ground, when, after sufficient time is allowed for their concretion, they are collected for use. Mastich is brought to us in small, yellowish, transparent, brittle tears, or grains; it has a light agreeable smell, especially when rubbed, or heated; on being chewed, it first crumbles, soon after sticks together, and becomes soft

and white, like wax, without impressing any considerable taste. It is considered to be a mild corroborant and astringent; and, as possessing a balsamic power, it has been recommended in hæmoptysis, proceeding from ulceration, leucorrhœa, debility of the stomach, and in diarrhœas and internal ulcerations. Chewing this drug has likewise been said to have been of use in pains of the teeth and gums, and in some catarrhal complaints; it is, however, in the present day, seldom used either externally or internally. The wood abounds with the resinous principle, and a tincture may be obtained from it, which is esteemed in some countries in the cure of hæmorrhages, dysenteries, and gout.

Mastich tree. See *Mastiche*.

Mastich herb, common. See *Marum vulgare*.

Mastich herb, Syrian. See *Marum Syriacum*.

Mastich wood. See *Mastiche*.

MASTICHELÆUM. (From *μαστιχ*, mastich, and *λαιον*, oil.) Oil of mastich.

MASTICHIA. (From *μαστιχ*, mastich; so called because it smells like mastich.) *Nux virginiana*, or mastich of ligon.

MASTICHINA. (Dim. of *mastiche*.) *Mastichina gallorum marum*, or mastich thyme. See *Marum vulgare*.

Masticot. See *Masticot*.

MASTIX. See *Mastiche*.

MASTODYNIA. (From *μασος*, a breast, and *δυν*, pain.) Phlegmon of the breast of women. This disease may take place at any period of life, but it most commonly affects those who give suck. It is characterized by tumefaction, tension, heat, redness, and pain; and comes sometimes in both breasts, but most commonly in one. Pyrexia generally attends the disease. It is sometimes very quickly formed, and in general without any thing preceding to shew it; but now and then a slight shivering is the forerunner. This disease terminates either in resolution, in suppuration, or schirrus. If the disease is left to itself, it generally terminates in suppuration.

The causes which give rise to this disease are those which give rise to most of the phlegmasie, as cold, violent blows, &c. In women who are lying in, or giving suck, it mostly arises either from a suppression of the lochia or a retention of milk. Mastodynia is often of long continuance; it is a very painful disease, but it is seldom fatal, unless when absolutely neglected, when it may run into scirrhous, and finally cancer. The termination of the disease by gangrene is never to be apprehended; at least few, if any, have seen the disease terminate in this way.

MASTOID. (*Mastoides*; from *μασος*, a breast, and *ωδωρ*, resemblance.) 1. Those processes of bones are so termed that are shaped like the nipple of the breast.

2. The name of a muscle which is inserted into the mastoid process.

MASTOIDEUM FORAMEN. A hole in the temporal bone of the skull.

MASTOIDEUS LATERALIS. A name for the complexus.

MASTOIDEUS. (From *μασος*, the mastoid process.) See *Sterno-cleido-mastoides*.

MATALISTA RADIX. A root said to be imported from America, where it is given as a purgative, its action being rather milder than that of jalap.

MATER. (*Matr*, a mother.) Two membranes of the brain had this epithet given them by the Arabians, who thought they gave origin to all other membranes of the body. See *Dura mater* and *Pia mater*. Also a name of the herb mugwort, because of its virtue in disorders of the womb.

MATER DURA. See *Dura mater*.

MATER HERBARUM. *Artemisia*, or common mugwort.

MATER METALLORUM. Quicksilver.

MATER PERLORUM. Mother of pearl. The concha margaritifera.

MATER PIA. See *Pia mater*.

MATERIA PERLATA. Jf, instead of crystallizing the salts contained in the liquor separated from diaphoretic antimony, an acid be poured into it, a white precipitate is formed, which is nothing else but a very refractory calx of antimony.


MATERIATURA. Castellus explains *morbi materiatura* to be diseases of intemperance.

MATERIA MEDICA. By this term is understood a general class of substances, both natural and artificial, which are used in the cure of diseases.

Cartheuser, Newman, Lewis, Gleditsch, Linnæus, Vogel, Alston, and other writers on the *Materia Medica*, have been at much labour to contrive arrangements of these articles. Some have arranged them according to their natural resemblances; others according to their real or supposed virtues; others according to their active constituent principles.

These arrangements may have their peculiar advantages. The first may be preferred by the natural historian, the second by the physiologist, and the last by the chymist. The pharmacopœias published by the Colleges of Physicians of London, Dublin, and Edinburgh, have the articles of the *Materia Medica* arranged in alphabetical order; this plan is also adopted by almost all the continental pharmacopœias.

Dr. Cullen has arranged the Materia Medica as follows :—



The following is the list of articles which come under the preceding classes :—

- | | | |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>I. NUTRIMENTS.</p> <p>1. FRUITA.</p> <p>a. <i>Fresh, sweet, acidulous,</i>
as
Prunes
Oranges
Lemons
Raspberries
Red and black currants
Mulberries
Grapes</p> <p>b. <i>Dried, sweet, acidulous,</i>
as
Raisins
Currants
Figs.</p> <p>2. OLERACIOUS HERBS.</p> <p>Water-cresses
Dandelion
Parsley
Artichoke.</p> <p>3. ROOTS.</p> <p>Carrot
Garlick
Satyrion.</p> <p>4. SEEDS AND NUTS.</p> <p>Almonds, sweet and bitter
Walnuts</p> | <p>II. MEDICINES.</p> <p>1. ASTRINGENTS.</p> <p>Olives
Red rose
Cinquefoil
Tormentil
Madder
Sorrel
Water-dock
Bisturt
Fern
Granate
Oak-bark
Galls
Logwood
Quince
Mulberry
Sloe
Gum-arabic
Catechu
Dragon's blood
Alkanet
Balustine flower
St. John's wort
Millefoil
Plantain
Convallaria
Bear's berry.</p> | <p>2. TONICS.</p> <p>Gentian
Lesser centaury
Quassia
Simarouba
Marsh trefoil
Fumatory
Camomile
Tansy
Wormwood
Southernwood
Sea-wormwood
Water-germander
Virginian snakeroot
Leopard's bane
Peruvian bark.</p> <p>3. EMOLLIENTS.</p> <p><i>Columniferous,</i>
Marsh mallow
Mallow.
<i>Farinaceous,</i>
Quince-seeds
Fenn greek-seed
Linseed,
<i>Various emollients,</i>
Pellitory
Verbascum
White lily</p> |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

4. CORROSIVES.

5. STIMULANTS.

A. *Vericillated*,

Lavender

Balm

Marjoram

Sweet marjoram

Syrian herb

d Rosemary

o Hyssop

in Ivy

e Mint

e Peppermint

w Pennyroyal

m Thyme

in Mother of thyme

rh Sage.

Umbellated,

Fennel

g Archangel

anise

a Caraway

Coriander

Cumin

ti Dill

Sassafras.

sc *Siliquous*,

N Horse radish

Water-cress

A Mustard

t Scurvy-grass.

Aliacious,

Garlick.

Coniferous,

Fir

Juniper.

Balanics,

Venice turpentine.

Common turpentine

Canada balsam

Copaiba balsam

Tolu balsam

Balm of Gilead.

Resinous,

Guaiacum

Ladanum

Storax

Benzoin.

Aromatics

Cinnamon

Nutmeg

Mace

Clove

Allspice

Canella

Cascarilla

Black pepper

Long pepper

Indian pepper

Ginger

Lesser Cardamom

Zedoary

Virginian snake-root

Ginseng

Aromatic reed

Acrid,

Wake robin

Pellitory

Stavesacre

Sedatives.

6. NARCOTICS.

Rheadaceous,

White poppy

Red poppy

Umbellated,

Hemlock

Water hemlock.

Solinaceous,

Belladonna

Henbane

Tobacco

Bitter-sweet

Stramonium.

Varia,

Laurel

Camphire

Saffron

Wine.

7. REFRIGERANTS.

Fruits of plants

Acidulous herbs and fruits.

8. ANTISPASMODICS.

Fetid herbs,

Worm-wood

Fetid goosefoot

Cumin

Pennyroyal

Rue

Savine.

Fetid gums,

Asafoetida

Galbanum

Opopanax

Valerian.

9. DILUENTS.

10. ATTENUANTS.

Alkalis

Sugar

Liquorice

Dried fruits.

11. INSPISSANTS.

Acids

Farinaceous and mucila-

ginous demulcents.

12. DEMULCENTS.

Mucilaginous,

Gum arabic

—tragacanth.

Farinaceous,

Bland oils.

13. ANTACIDS.

14. ANTALKALINES.

15. ANTISEPTICS.

Acid parts of plants

Acescent herbs

Sugar

Siliquous plants

Alliacious plants

Astringents

Bitters

Aromatics

Essential oils

Camphor

Gum resins.

Saffron

Contrainerva

Valerian

Opium

Wine

16. EMBRINES.

Asarabacca

White hellebore

Water iris

Pellitory.

17. SIALOOGUES.

Archangel

Cloves

Imperatory

Tobacco

Pepper

Pellitory

18. EXPECTORANTS.

Ivy

Horebound

Penny-royal

Elecampane

Florentine oris-root

Tobacco

Squill

Coltsfoot

Benzoin

Storax

Canada balsam

Tolu balsam.

19. EMETICS.

Asarabacca

Ipecacuan

Tobacco

Squill

Mustard

Horse-radish

Bitters.

20. CATRACTICS.

Milder,

Mild acid fruits

Cassia pulp

Tamarind

Dulcious

Sugar

Manna

Sweet roots

Bland oils

Damask rose

Violet.

Polypody

Mustard

Bitters

Balsamics.

Acrid,

Rhubarb

Seneka

Broom

Elder

Castor-oil

Senna

Black hellebore

Jalap

Scammony

Buckthorn

Tobacco

White hellebore

Coloquint
Elatarium.
21. Diuretics.

Parsley
Carrot
Fennel
Pimpinell
Eryngo
Madder
Varia
Burdock
Bitter-sweet
Wake-robin
Asarabacca
Foxglove

Tobacco
Rue
Savine
Snake-root
Squill
Bitters
Balsamics
Siliquosæ
Alliacæ.

22. DIAPHORETICS.

Saffron
Bitter-sweet
Opium
Camphor
Contra-yerva

Serpentary
Sage
Water germander
Guaiacum
Sassafras
Seneca
Vegetable acids
Essential oil
Wine
Diluents.

23. EMMENAGOGUES.

Aloe
Fetid gums
Fetid plants
Saffron.

The following is the arrangement of the Materia Medica, according to J. Murray, in his Elements of Materia Medica and Pharmacy.

A. General stimulants.

a. Diffusible } Narcotics.
Antispasmodics
b. Permanent } Tonic
Astringent.

. Local stimulants.

Emetics
Carthartics
Emmenagogues
Diuretics
Diaphoretics
Expectorants
Sialogogues
Errhines
Epispastics.
c. Chymical remedies. Refrigerents
Antacids
Lithontriptics.
Escharotics.
d. Mechanical remedies. Anthelmintics.
Demulcents
Diluents
Emollients.

Under the head of NARCOTICS are included—

Alkohol. Ether. Camphor, Papaver somniferum. Hyoscyamus niger. Atropa belladonna. Aconitum napellus. Conium maculatum. Digitalis purpurea. Nicotiana tabacum. Lactuca virosa. Datura stramonium. Rhododendrum chrysanthum. Rhus toxicodendron. Arnica montana. Strychnos nux vomica. Prunus lauro cerassus.

Under the second class, ANTISPASMODICS, are included—Moschus. Castoreum. Oleum animale empyreumaticum. Petroleum Ammonia. Ferula asafoetida. Sagapenum. Bubon galbanum. Valeriana officinalis. Crocus sativus. Melaleuca leucadendron.

Narcotics used as Antispasmodics—

Ether. Camphor. Opium.

Tonics used as Antispasmodics—

Cuprum. Zincum. Hydrargyrum. Cinchona.

The head of TONICS embrace—

1. From the mineral kingdom, Hydrargyrum. Ferrum. Zincum. Cuprum.

Arsenicum. Barytes. Calx. Acidum. Nitricum. Oxy-murias potassæ.

2. From the vegetable kingdom,

Cinchona officinalis. Cinchona caribæa. Cinchona floribunda. Angustura. Aristolochia serpentaria. Dortenia contrayerva. Croton cleutheria. Calomba. Quassia excelsa. Quassia Simarouba. Swietenia febrifuga. Swietenia mahagoni. Gentiana leutea. Anthemis nobilis. Artemisia absinthium. Chironia centaurium. Marrubium vulgare. Menyanthes trifoliata. Centaurea benedicta. Citrus aurantium. Citrus medica. Laurus cinnamomum. Laurus cassia. Canella alba. Acorus calamus. Amomum zinziber. Kamferia rotunda. Santalum album. Pterocarpus santalinus. Myristica moschata. Caryophyllus aromaticus. Capsicum annuum. Piper nigrum. Piper longum. Piper cubeba. Myrtus pimenta. Amomum repens. Carum carui. Coriandrum sativum. Pimpinella anisum. Anethum fœniculum. Anethum graveolens. Cuminum cyminum. Angelica archangelica. Mentha piperita. Mentha viridis. Mentha pulegium. Hyssopus officinalis.

The class of ASTRINGENTS comprehends the following:—

1. From the vegetable kingdom.

Quercus rubor. Quercus cerris. Tormentilla erecta. Polygonum bistorta. Anchusa tinctoria. Hæmatoxylon campechianum. Rosa gallica. Arbutus uva ursi. Mimosa catechu. Kino. Pterocarpus draco. Ficus indica. Pistachia lentiscus.

2. From the mineral kingdom.

Acidum sulphuricum. Argilla. Super sulphas argillæ et potassæ. Calx. Carbonas calcis. Plumbum. Zincum. Ferrum. Cuprum.

The articles which come under the head of EMMETICS are,

1. From the vegetable kingdom.

Callicocca ipecacuanha. Scilla maritima. Anthenus nobilis. Sinapis alba. Asarum Europæum. Nicotiana tabacum.

2. From the mineral kingdom.

Antimonium. Sulphas zinci. Sulphas cupri. Subacetes cupri. Ammonia. Hydro-sulphuretum ammoniz.

CATHARTICS include

Laxatives. Manna. Cassia fistula. Tamarindus Indica. Ricinus communis. Sulphur. Magnesia.

Purgatives. Cassia senna. Rheum palmatum. Convolvulus jalapa. Helleborus niger. Brionia alba. Cucumis colocynthis. Momordica elaterium. Rhamnus catharticus. Aloe perfoliata. Convolvulus scammonia. Gambojia gutta. Submurias hydrargyri. Sulphas magnesiz, Sulphas sodæ. Sulphas potassæ. Supertartris potassæ. Tartris potassæ et sodæ. Murias sodæ. Terebinthina veneta. Nicotiana tabacum.

The medicines arranged under **EMMENOGOGUES** are,

1. From the class of Antispasmodics, Castoreum. Ferula asafoetida. Bubon galbanum.

2. From the class of Tonics, Ferrum. Hydrargyrus. Cinchona officinalis.

3. From the class of Cathartics, Aloe. Helleborus niger. Sinapis alba. Rosmarinus officinalis. Rubia tinctorum. Ruta graveolens. Juniperus sabinæ.

The class of **DIURETICS** includes,

1. Saline diuretics.

Supertartris potassæ. Nitras potassæ. Murias ammoniz. Acetis potassæ. Potassa.

a. From the *vegetable* kingdom,

Scilla maritima. Digitalis purpurea. Nicotiana tabacum. Solanum dulcamara. Laticia virosa. Colchicum autumnale. Gratiola officinalis. Spartium scoparium. Juniperus communis. Copaifera officinalis. Pinus balsamea. Pinus larix.

b. From the *animal* kingdom.

Meloe vesicatorius.

Under the class **DIAPHORETICS** are,

Ammonia. Murias ammoniz. Acetis ammoniz. Citras ammoniz. Submurias hydrargyri. Antimonium. Opium. Camphor. Guaiacum officinale. Daphne mezereum. Smilax sarsaparilla. Laurus sassafras. Cochlearia armoracia. Salvia officinalis.

The class **EXPECTORANTS** comprehends, Antimonium. Ipecacuanha. Nicotiana tabacum. Digitalis purpurea. Scilla maritima. Allium sativum. Polygala senega. Ammoniacum. Myrrha. Styra

benzoin. Styra officinalis. Toluferia balsamum. Myroxylon peruiferum. Amyris gileadensis.

The articles of the class **SIALAGOGUES** are, Hydrargyrus. Anthemis pyrethrum. Arum maculatum. Amomum zinziber. Daphne mezereum. Nicotiana tabacum.

The class of **EMMENES** are, Iris florentina. Esculus hippocastanum. Origanum majorana. Lavendula spica. Asarum Europæum. Veratrum album. Nicotiana tabacum. Euphorbia officinalis.

In the class **ERISPASTICS** and **RUBEFACIENTS** are, Meloe vesicatorius. Ammonia. Pix Burgundica. Sinapis alba. Allium sativum.

REFRIGERANTS are constituted by the following articles. Citrus aurantium. Citrus medica. Tamarindus Indica. Acidum acetosum. Supertartris potassæ. Nitras potassæ. Boras sodæ.

The list of articles that come under the class **ANTACIDS** are, Potassa. Soda. Ammonia. Calx. Carbonas calcis. Magnesia.

In the class **LITHONTRIPTICS** are, Potassa. Carbonas potassæ. Soda. Carbonas sodæ. Sapo albus. Calx.

In the class **ESCHAROTICS** are, Acida mineralia. Potassa. Nitras argenti. Murias antimonii. Sulphas cupri. Acetis cupri. Murias hydrargyri. Subnitras hydrargyri. Oxydum arsenici album. Juniperus sabinæ.

In the class **ANTHELMINTICS** are, Dolichos pruriens. Ferri limatura. Stannum pulveratum. Olea Europæa. Artemesia santonica. Spigelia marilandica. Polypodium filix mas. Tanacetum vulgare. Geoffrea inermis. Gambojia gutta. Submurias hydrargyri.

DEMULCENTS are, Mimosa nilotica. Astragalus tragacantha. Linum usitatissimum. Althæa officinalis. Malva sylvestris. Glycyrrhiza glabra. Cynas circinalis. Orchis mascula. Maranta arundinacea. Triticum hybernium. Icthyocolla. Olea Europæa. Amygdalus communis. Sevum ceti. Cera.

Water is the principal article of the class **DILUENTS**; and as for the last class, **EMOLLIENTS**, heat conjoined with moisture is the principal, though all unctuous applications may be included.

The New London Pharmacopœia presents us with the following list for the **Materia**

Medica :—

Abietis resina
Absinthium
Acacia gummi
Acetosæ folia
Acetosella
Acetum
Acidum sulphuricum
Aconiti folia
Adeps
Ærugo

Allii radix
Aloes spicatæ
Aloes vulgaris
Althææ folia
Alumen
Ammoniz murias
Ammoniacum
Amygdala amara et dulcis
Amylum
Anethi semina

Anisi semina
Anthemidis flores
Antimonii sulphuretum
Argentum
Armoraciz radix
Arsenici oxydum
Asara folia
Asafoetida gummi
Avenæ semina
Aurantii baccæ

Aurantii cortex	Gentianæ radix	Quercus cortex
Balsamum Peruvianum	Glycyrrhizæ radix	Resina flava
Balsamum Tolutanum	Granati cortex	Rhamni baccæ
Belladonæ folia	Guaiaci resina et lignum	Rhei radix
Benzoinum	Hæmatoxyli lignum	Rhœados petala
Bistorta	Hellebori foetidi folia	Ricini semina et oleum
Cajuputi oleum	Hellebori nigri radix	Rosæ caninæ pulpa
Calamina	Hordei semina	Rosæ centifoliæ petala
Calami radix	Humuli strobili	Rosæ gallicæ petala
Calumbæ radix	Hydrargyrum	Rosmarini cacumina
Campbora	Hyoscyami folia et semina	Rubiz radix
Canellæ cortex	Jalapæ radix	Ruta folia
Capsicum baccæ	Ipecacuanhæ radix	Sabinæ folia
Carbo ligni	Juniper baccæ et semina	Saccharum
Cardamines flores	Kino	Salicis cortex
Cardamomi semina	Lapis calcareus	Sambuci flores
Caricæ fructus	Lavendulæ flores	Sapo durus et mollis
Carici semina	Lauri baccæ et folia	Sarsaparillæ radix
Caryophylli	Lichen	Sassafras lignum et radix
Caryophylli oleum	Limones	Scammonæ gummi resina
Cascarillæ cortex	Limonum cortex	Scillæ radix
Cassiz pulpa	Linum catharticum	Senegæ radix
Castoreum	Lytta	Sennæ folia
Catechu	Magnesiz sulphas	Serpentariæ radix
Centaurii cacumina	Malva	Sevum
Cera alba	Manna	Simaroubæ cortex
Cera flava	Marrubium	Sinapis semina
Cerevisiæ frumentum	Mastiche	Sodæ murias
Cetaceum	Mel	Sodæ boras
Cinchonæ lancifoliæ cordi-	Mentha piperita	Sodæ sulphas
foliæ et oblongifoliæ cor-	Mentha viridis	Soda impura
tex	Menyanthes	Spartii cacumina
Cinnamomi cortex	Mezerei cortex	Spigeliæ radix
Cinnamomi oleum	Mori baccæ	Spiritus rectificatus et tenuis
Coccus	Moschus	Spongia
Colchici radix	Myristicæ nucléi	Stannum
Colocynthis pulpa	Myrrha	Staphisagriæ semina
Conii folia	Olibanum	Styracis balsamum
Contrayervæ radix	Olivæ oleum	Succinum
Copaiba radix	Opium	Sulphur
Coriandri semina	Opopanax	Tabaci folia
Cornua	Origanum	Tamarindi pulpa
Creta	Ovum	Taraxaci radix
Croci stigmata	Papaveris capsulæ	Terebinthina Canadensi et
Cumini semina	Petroleum	Chia
Cupri sulphas	Pimentæ baccæ	Testæ
Cuspariæ cortex	Piperis longi fructus	Terebinthina vulgaris
Cydoniæ semina	Piper nigrum	Terebinthinæ oleum
Dauci radix	Pix arida	Tormentillæ radix
Dauci semina	Pix liquida	Toxicodendri folia
Digitalis folia	Plumbum	Tragacantha
Dolichi pubes	Plumbi carbonas	Tussilago
Dulcamaræ caulis	Plumbi oxydum semivitreum	Valerianæ radix
Elaterii poma	Porri radix	Veratri radix
Elemi	Potassæ nitras	Violæ flores
Euphorbiæ gummi resina	Potassæ supertartras	Vinum
Farina	Potassa impura	Ulmæ cortex
Ferrum	Pruna	Uvæ passæ
Filicis radix	Pterocarpi lignum	Uvæ ursi folia
Fucus	Pulegium	Zincum
Galbani gummi resina	Pyrethri radix	Zingiberis radix
Gallæ	Quassiz lignum	

MATLOCK. A village in Derbyshire affords a mineral water of the acidulous class; its abundant springs issue from a lime-stone rock, near the banks of the Der-

went. Several of these springs possess a temperature of 66°. Matlock water scarcely differs from common good spring water, in sensible properties. It is extremely

transparent, and exhales no vapour, except in cold weather. It holds little or no mass of aerial particles; it curdles soap, on first taken up, but it loses this effect on long keeping, perhaps from the deposition of its calcareous salts: it appears differ very little from good spring water, on tasted; and its effects seem referable to its temperature. It is from this latter circumstance that it forms a proper acid bath for the nervous and irritable, and those of a debilitated constitution; and it is usually recommended after the use of Bath and Buxton waters, and as preparatory to sea-bathing.

MATRICALIA. (From *matrix*, the womb.) Medicines appropriated to disorders of the uterus.

MATRICARIA. (From *matrix*, the womb; called from its uses in disorders of the womb. *Parthenium. Febrifuga.* Common feverfew, or febrifuge, and often, but very improperly, featherfew. Mother's wort. The leaves and flowers of this plant *Matricaria parthenium*—*foliis compositis, platis; foliis ovatis, incis; pedunculis ramosis*, have a strong, not agreeable, smell, and a moderately bitter taste, both which they communicate, by warm infusion, to water and rectified spirit. The watery infusions, pressed, leave an extract of considerable bitterness, and which discovers also a saline matter, both to the taste, and in a more sensible manner by throwing up to the surface all crystalline efflorescences in keeping. The peculiar flavour of the matricaria exists, in the evaporation, and impregnates the distilled water, on which also a quantity of essential oil is found floating. The quantity of spirituous extract, according to Reaumur's experiments, is only about one-sixth the weight of the dry leaves, whereas the watery extract amounts to one-half. This plant is evidently the *parthenium* of Dioscorides, since whose time it has been very generally employed for medical purposes. In natural affinity, it ranks with camomile and tansy, and its sensible qualities shew it to be nearly allied to them in its medicinal character. Reaumur states its virtues to be tonic, stochic, resolvent, and emmenagogue. It has been given successfully as a vermifuge, and for the cure of intermittents; but its most celebrated use is in female disorders, especially in hysteria; and hence it is supposed to have derived the name matricaria. Its smell, taste, and analysis, shew it to be a medicine of considerable activity; we may therefore say, with Gray—*Rarius hodie prescribitur, quam antea*.

MATRICARIA CAMOMILLA. The systematic name of the common feverfew. See *anemum vulgare*.

MATRICARIA PARTHENIUM. The systematic name of the feverfew. See *Matricaria*.

MATRIX. (*Matr.*) The womb. See *Uterus*.

MATRONALIS. (From *matrona*, a matron; so called because its smell is grateful to women.) The violet.

MATURANTIA. (From *mature*, to ripen.) Medicines which promote the suppuration of tumours.

MATURATION. (*Maturatio*; from *mature*, to make ripe.) A term in surgery, signifying that process which succeeds inflammation, by which pus is collected in an abscess.

Maudlin. See *Ageratum*.

MAURO-MARSON. See *Marrubium*.

MAXILLA. (From *mastrum*, to chew.) The jaw.

MAXILLA INFERIOR. *Os maxillare inferius. Mandibula.* The maxilla inferior, or lower jaw, which, in its figure, may be compared to a horse-shoe, is at first composed of two distinct bones; but these, soon after birth, unite together at the middle of the chin, so as to form only one bone. The superior edge of this bone has, like the upper jaw, a process, called the *alveolar* process. This, as well as that of the upper jaw, to which it is in other respects a good deal similar, is likewise furnished with cavities for the reception of the teeth. The posterior part of the bone, on each side, rises perpendicularly into two processes, one of which is called the *coronoid*, and the other the *condyloid*, process. The first of these is the highest; it is thin and pointed; and the temporal muscle, which is attached to it, serves to elevate the jaw. The condyloid process is narrower, thicker, and shorter than the other, terminating in an oblong rounded head, which is formed for a moveable articulation with the cranium, and is received into the fore part of the fossa described in the temporal bone. In this joint there is a moveable cartilage, which, being more closely connected to the condyle than to the cavity, may be considered as belonging to the former. This moveable cartilage is connected with both the articulating surface of the temporal bone and the condyle of the jaw, by distinct ligaments arising from its edges all round. These attachments of the cartilage are strengthened, and the whole articulation secured, by an external ligament, which is common to both, and which is fixed to the temporal bone, and to the neck of the condyle. On the inner surface of the ligament, which attaches the cartilage to the temporal bone, and backwards in the cavity, is placed what is commonly called the gland of the joint; at least the ligament is there found to be much more vascular than at any other part. At the bottom of each coronoid process, on its inner part, is a foramen, or canal, which extends under the roots of all the teeth, and terminates at the outer surface of the bone near the

shin. Each of these foramina affords a passage to an artery, vein, and nerve, which send off branches to the several teeth.

This bone is capable of a great many motions. The condyles, by sliding from the cavity towards the eminences on each side, bring the jaw horizontally forwards, as in the action of biting; or the condyles only may be brought forwards, while the rest of the jaw is tilted backwards, as is the case when the mouth is open. The condyles may also slide alternately backwards and forwards from the cavity to the eminence, and *vice versa*; so that while one condyle advances, the other moves backwards, turning the body of the jaw from side to side, as in grinding the teeth. The great use of the cartilages seems to be that of securing the articulation, by adapting themselves to the different inequalities in these several motions of the jaw, and to prevent any injuries from friction. This last circumstance is of great importance where there is so much motion, and accordingly this cartilage is found in the different tribes of carnivorous animals, where there is no eminence and cavity, nor other apparatus for grinding.

The alveolar processes are formed of an external and internal plate, united together by thin bony partitions, which divide the processes at the fore part of the jaw into as many sockets as there are teeth. But, at the posterior part, where the teeth have more than one root, each root has a distinct cell. These processes, in both jaws, begin to be formed with the teeth, accompany them in their growth, and disappear when the teeth fall. So that the loss of the one seems constantly to be attended with the loss of the other.

MAXILLA SUPERIOR. *Oss maxillare superius.* The superior maxillary bones constitute the most considerable portion of the upper jaw, are two in number, and generally remain distinct through life. Their figure is exceedingly irregular, and not easily to be described. On each of these bones is observed several eminences. One of these is at the upper and fore-part of the bone, and, from its making part of the nose, is called the *nasal process*. Internally, in the inferior portion of this process, is a fossa, which, with the *os unguis*, forms a passage for the lachrymal duct. Into this nasal process likewise is inserted the short round tendon of the *musculus orbicularis palpebrarum*. Backwards and outwards, from the root of the nasal process, the bone helps to form the lower side of the orbit, and this part is therefore called the *orbital process*. Behind this orbital process, the bone forms a considerable tuberosity, and at the upper part of this tuberosity is a channel, which is almost a complete hole. In this channel

passes a branch of the fifth pair of nerves, which, together with a small artery, is transmitted to the face through the external orbital foramen, which opens immediately under the orbit. Where the bone on each side is joined to the *os maxilæ*, and helps to form the cheeks, is observed what is called the *malar process*. The lower and anterior parts of the bone make a kind of circular sweep, in which are the *alveoli*, or sockets for the teeth; this is called the *alveolar process*. This alveolar process has posteriorly a considerable tuberosity on its internal surface. Above this alveolar process, and just behind the fore-teeth, is an irregular hole, called the *foramen incisicum*, which separating into two, and sometimes more holes, serves to transmit small arteries and veins, and a minute branch of the fifth pair of nerves, to the nostrils. There are two horizontal lamellæ behind the alveolar process, which, uniting together, form part of the roof of the mouth, and divide it from the nose. This partition, being seated somewhat higher than the lower edge of the alveolar process, gives the roof of the mouth a considerable hollowness. Where the *ossa maxillaria* are united to each other, they project some, what forwards, leaving between them a furrow, which receives the inferior portion of the *septum nasi*. Each of these bones is hollow, and forms a considerable sinus under its orbital part. This sinus, which is usually, though improperly, called *antrum Highmorianum*, is lined with the pituitary membrane. It answers the same purposes as the other sinuses of the nose, and communicates with the nostrils by an opening, which appears to be a large one in the skeleton, but which, in the recent subject, is much smaller. In the fœtus, instead of these sinuses, an oblong depression only is observed at each side of the nostrils, nor is the tuberosity of the alveolar process then formed. On the side of the palate, in young subjects, a kind of fissure may be noticed, which seems to separate the portion of the bone which contains the *dentes incisores* from that which contains the *dentes canini*. The fissure is sometimes apparent till the sixth year, but after that period it in general wholly disappears.

The *ossa maxillaria* not only serve to form the cheeks, but likewise the palate, nose, and orbits; and, besides their union with each other, they are connected with the greatest part of the bones of the face and cranium, viz. with the *ossa nasi*, *ossa malarum*, *ossa unguis*, *ossa palati*, *os frontis*, *os sphenoides*, and *os ethmoides*.

MAXILLARY ARTERIES. *Arteria maxillaria.* These are the branches of the external carotid. The *external maxillary* is the fourth branch of the carotid; it proceeds anteriorly, and gives off the facial

or mental, the coronary of the lips, and the angular artery. The *internal maxillary* is the next branch of the carotid; it gives off the sphæno maxillar, the inferior alveolar, and the spinous artery.

MAXILLARY GLANDS. *Glandule maxillariæ.* The glands so called are conglomerate, and are situated under the angles of the lower jaw. The excretory ducts of these glands are called Warthonian, after their discoverer.

MAXILLARY NERVES. *Nervi maxillares.* The superior and inferior maxillary nerves are branches of the fifth pair, or trigemini. The former is divided into the sphæno-palatine, posterior alveolar, and the infra orbital nerve. The latter is divided into two branches, the internal lingual, and one, more properly, called the inferior maxillary.

May-bily. See *Lilium convallium*.

May-weed. See *Cotula fetida*.

Mays, Indian. See *Zea mays*.

Maw worm. See *Ascaris vermicularis*.

Meadow crowfoot. See *Ranunculus pratensis*.

Meadow, queen of the. See *Ulmaria*.

Meadow saffron. See *Colchicum*.

Meadow saxifrage. See *Saxifraga vulgaris*.

Meadow sweet. See *Ulmari*.

MEADOW THISTLE, ROUND LEAVED. The leaves of this plant, *Cnicus oleraceus* of Linnaeus, are boiled, in the northern parts of Europe, and eaten as we do cabbage.

Measles. See *Rubeola*.

MEATUS AUDITORIUS EXTERNUS. See *Ear*.

MEATUS AUDITORIUS INTERNUS. The internal auditory passage is a small bony canal, beginning internally by a longitudinal orifice at the posterior surface of the petrous portion of the temporal bone, running towards the vestibulum and cochlea, and there being divided into two less cavities by an eminence. The superior and smaller of these is the orifice of the aqueduct of Fallopius, which receives the portio dura of the auditory nerve: the other inferior and larger cavity is perforated by many small holes, through which the portio mollis of the auditory nerve passes into the labyrinth.

MEATUS CÆCUS. A passage in the throat to the ear, called Eustachian's tube.

MEATUS CUTICULARES. The pores of the skin.

MEATUS CYSTICUS. The gall-duct.

MEATUS URINARIUS. In women, this is situated in the vagina, immediately below the symphysis of the pubis, and behind the nymphæ. In men, it is at the end of the glans penis.

Mecca balsam. See *Balsam of Gilead*.

Mechoacan. See *Mechoacanna*.

MECHOACANNA. (From *Mechoacan*, a

province in Mexico, whence it is brought.) *Jalappa alba*. *Rhabarbarum album*. *Mechoacam*. The root of a species of convolvulus, *Convolvulus Americanus*, or *Bryonia alba Peruviana*, brought from Mexico. It possesses aperient properties, and was long used as the common purge of this country, but is now wholly superseded by jalap.

MECHOACANNA NIGRA. See *Jalapium*.

MEXON. (From *μυρον*, bulk; so named from the largeness of its head.) The papaver, or poppy.

MEXONIS. (From *μυρον*, the poppy; so called because its juice is soporiferous, like the poppy.) The lettuce.

MEXOSTRUM. (From *μυρον*, the poppy.) 1. The inspissated juice of the poppy. Opium. 2. The green excrementitious substance that is found in the large intestines of the fœtus.

MEDIAN NERVE. The second branch of the brachial plexus.

MEDIAN VEINS. The situation of the veins of the arms is extremely different in most individuals. When a branch proceeds near the bend of the arm, inwardly from the basilic vein, it is termed the *basilic medium*; and when a vein is given off from the cephalic in the like manner, it is termed the *cephalic median*. When these two veins are present, they mostly unite just below the bend of the arm, and the common trunk proceeds to the cephalic vein.

MEDIANUM. The *Mediastinum*.

MEDIASTINUM. (*Quasi in media statur.*) The membranous septum, formed by the duplicature of the pleura, that divides the cavity of the chest into two parts. It is divided into an anterior and posterior portion.

MEDIASTINUM CEREBRI. The falciform process of the dura mater.

MEDICAGO. (From *medica*.) The herb trefoil.

MEDICAMENTARIA. Pharmacy, or the art of making and preparing medicines.

MEDICAMENTUM. (From *medico*, to heal.) A medicine.

MEDICASTER. A false pretender to the knowledge of medicine; the same as Quack.

MEDICINA DIÆTETICA. The regulation of regimen, or the non-naturals.

MEDICINA DIASOSTICA. That part of medicine which relates to preserving health.

MEDICINA GYMNASTICA. That part of medicine which relates to exercise.

MEDICINA HERMETICA. The application of chemical remedies.

MEDICINA PROPHYLACTICA. That part of medicine which relates to preserving health.

MEDICINA TRISTITILE. Common salfron.

MEDICINE. Any substance that is exhibited with a view to cure or allay the

violence of the disease. It is also very frequently made use of to express the healing art, when it comprehends anatomy, physiology, and pathology.

MEDICINAL DAYS. Such are so called by some writers, wherein the crisis or change is expected, so as to forbid the use of medicines, in order to wait nature's effort, and therefore require all assistance of art to help forward, or prepare the humours for such a crisis: but it is most properly used for those days wherein purging, or any other evacuation, is most conveniently complied with.

MEDICINAL HOURS. Are those wherein it is supposed that medicines may be taken to the greatest advantage, commonly reckoned in the morning fasting, about an hour before dinner, about four hours after dinner, and at going to bed; but in acute cases, the times are to be governed by the symptoms and aggravation of the distemper.

MEDICINALIS. (From *medicina*.) Medicinal; having a power to restore health, or remove disease.

MEDINA. In Paracelsus it is a species of ulcer.

MEDINENSIS VENA. (*Medinensis*; so called because it is frequent at Medina.) *Dracunculus*. *Gordius medinensis* of Linnaeus. The muscular hair-worm. A very singular animal, which, in some countries, inhabits the cellular membrane between the skin and muscles. See *Dracunculus*.

MEDITULLIUM. (From *medius*, the middle. See *Diploë*.)

MEDIUS VENTER. The middle venter is the thorax, or chest.

Medlar. See *Mespilus*.

MEDULLA. (*Quasi in medis ossis*.) 1. The marrow. See *Marrow*.

2. The pith or pulp of vegetables.

3. The white substance of the brain is called medulla, or the medullary part, to distinguish it from the cortical.

4. The continuation of the brain, and in the spine, is called medullary spinalis.

MEDULLA CASSIAE. The pulp of the cassia fistularis.

MEDULLA OBLONGATA. The medullary substance, of the same use as the cerebrum, that lies within the cranium, upon the basillary process of the occipital bone. It is formed by the connection of the crura cerebri and crura cerebelli, and terminates in the spinal marrow. It has several eminences, viz. pons varioli, corpora pyramidalia, and corpora olivaria.

MEDULLA SPINALIS. *Cerebrum elongatum*. *Een*. The spinal marrow. A continuation of the medulla oblongata, which descends into the specus vertebralis from the foramen magnum occipitale, to the third vertebra of the loins, where it termi-

nates in a number of nerves; which, from their resemblance, are called *cauda equina*. The spinal marrow is composed, like the brain, of a cortical and medullary substance: the former is placed internally. It is covered by a continuation of the dura mater, pia mater, and tunica arachnoidea. The use of the spinal marrow is to give off, through the lateral or intervertebral foramina, thirty pairs of nerves, called cervical, dorsal, lumbar, and sacral nerves.

MEDULLARY. (*Medullaris*; from *medulla*, marrow.) Like unto marrow.

MEDULLARY SUBSTANCE. The white or internal substance of the brain is so called.

MEGALOSPLANCHNUS. (From *μεγας*, great, and *σπλANCHNOS*, a bowel. Having some of the viscera enlarged, from schirrus or other cause.

MIGRAINE. A species of cephalalgia, or a pain generally affecting one side of the head, towards the eye, or temple, and arising from the state of the stomach.

MEIBOMIUS'S GLANDS. *Meibomii glandulae*. The small glands which are situated between the conjunctive membrane of the eye and the cartilage of the eye-lid, first described by Meibomius.

MEL. See *Honey*.

MEL ACETATUM. *Oxymel*. "Take of clarified honey, two pounds; acetic acid, two pints. Boil them down to a proper consistence, in a glass vessel, over a slow fire." This preparation of honey and vinegar possesses aperient and expectorating virtues, and is given, with these intentions, in the cure of humoral asthma, and other diseases of the chest, in doses of one or two drachms. It is also employed in the form of gargle, when diluted with water.

MEL BORACIS. Honey of borax. "Take of borax, powdered, a drachm; clarified honey, an ounce. Mix." This preparation is found very useful in aphthous affections of the fauces.

MEL DESPUMATUM. Clarified honey; See *Honey*.

MEL ROSE. Rose honey. "Take of red-rose petals, dried, four ounces; boiling water, three pints; clarified honey, five pounds." Macerate the rose petals in the water, for six hours, and strain; then add the honey to the strained liquor, and, by means of a water-bath, boil it down to a proper consistence. An admirable preparation for the base of various gargles and collutories. It may also be employed with advantage, mixed with extract of bark, or other medicines, to children, who have a natural disgust to medicines.

MEL SCILLAE. "Take of clarified honey, three pounds; vinegar of squills, two pints." Boil them in a glass vessel, with a

slow fire, to the thickness of a syrup. Aperient, expectorant, and detergent virtues, are attributed to the honey of squills. It is given in doses of two or three drachms, along with some aromatic water, as that of cinnamon, to prevent the great nausea which it would otherwise be apt to excite. In large doses it proves emetic.

MELA. (From *μαλα*, to search.) A probe.

MELÆNA. (From *μαλας*, black.) The black vomit. The black disease. Hippocrates applies this name to two diseases. In the first, the patient vomits black bile, which is sometimes bloody and sour; sometimes he throws up a thin saliva; and at others a green bile, &c. In the second, the patient is as described in the article *Morbus Niger*.

MELAINA NOSOS. See *Melena*.

MELALEUCA LEUCADENDRON. The systematic name of the plant which affords the cajeput oil. See *Cajepu* *al*.

MELAMPHYLLUM. (From *μαλας*, black, and *φυλλον*, a leaf; so named from the blackness of its leaf.) Bear's-breech. See *Acanthus mollis*.

MELAMPOBIUM. (From *Melampus*, the shepherd who first used it.) Black hellebore. See *Helleborus niger*.

MELANAGOGA. (From *μαλας*, black, and *αγω*, to expel.) Medicines which purge off black bile.

MELANCHLORUS. (*Μαλαχχλος*.) A livid colour of the skin. The black jaundice.

MELANCHOLIA. (From *μαλας*, black, and *χολα*, bile; because the ancients supposed that it proceeded from a redundancy of black bile.) Melancholy madness. A disease in the class *neuroses*, and order *Vesante*, of Cullen, characterised by erroneous judgment, but not merely respecting health, from imaginary perceptions or recollection, influencing the conduct, and depressing the mind with ill-grounded fears; not combined with either pyrexia or comatose affections; often appearing without dyspepsia, yet attended with costiveness, chiefly in persons of rigid fibres and torpid insensibility.

MELANOPYPER. (From *μαλας*, black, and *πυπερ*, pepper.) Black pepper. See *Piper nigrum*.

MELANORRHIZON. (From *μαλας*, black, and *ριζα*, a root.) A species of hellebore with black roots.

MELANTERIA. (From *μαλας*, black; so called because it is used for blacking leather.) Green vitriol.

MELANTHELEUM. (From *μαλας*, black, and *αιων*, oil.) Oil expressed from the black seeds of fennel flowers.

MELANTHIUM. (From *μαλας*, black; so named from its black seed.) The herb fennel flower.

MELAS. (From *μαλας*, black.) *Vitiligo*

nigra. **Morphea nigra.** *Lepa maculosa nigra.* A disease that appears upon the skin in black or brown spots, which very frequently penetrate deep, even to the bone, and do not give any pain or uneasiness. It is a disease very frequent in, and endemial to, Arabia, where it is supposed to be produced by a peculiar miasma.

MELASMA. (From *μαλας*, black.) *Melasma.* A disease that appears not unfrequently upon the tibia of aged persons, in form of a livid black spot, which, in a day or two, degenerates into a very foul ulcer.

MELASPERMUM. (From *μαλας*, black, and *σπερμα*, seed.) See *Melanthium*.

MELCA. (From *αμυγα*, to milk.) Milk. A food made of acidulated milk.

MELE. (From *μαλα*, to search.) A probe.

MELEAGRIS. (From *Meleager*, whose sisters were fabled to have been turned into this bird.) The guinea-fowl. Also a species of fritillaria, so called because its flowers are spotted like the guinea-fowl.

MELEGETA. *Meleguetta.* Grains of Paradise.

MELKIOS. (From *Melos*, the island where it is made.) A species of alum.

MELMELUM. (From *μαλας*, honey, and *μηλον*, an apple; so named from its sweetness.) Paradise apple.

MELI. (*Μελ*.) Honey.

MELIGERIA. See *Meliceris*.

MELICERIS. (From *μαλας*, honey, and *κερος*, wax.) An encysted tumour, whose contents resemble honey in consistence and appearance.

MELICRATON. (From *μαλας*, honey, and *κρατυμι*, to mix.) Hydromel. Mead. Wine impregnated with honey.

MELIGION. (From *μαλας*, honey.) A fetid humour, discharged from ulcers attended with a caries of the bone, of the consistence of honey.

Melilot. See *Melilotus*.

MELILOTUS. (From *μαλας*, honey, and *λωτος*, the lotus; so called from its smell, being like that of honey.) *Lotus sylvestris.* *Sertula campana.* *Trifolium cabalinum.* *Coroda regia.* *Trifolium odoratum.* Common melilot. This plant, *Trifolium melilotus officinale* of Linnaeus, has been said to be resolvent, emollient, anodyne, and to participate of the virtues of chamomile. Its taste is unpleasant, subacid, subsaline, but not bitter; when fresh it has scarcely any smell; in drying it acquires a pretty strong one, of the aromatic kind, but not agreeable. The principal use of melilot has been in clysters, fomentations, and other external applications.

MELIUM. (From *μαλας*, an apple.) Off made from the flowers of the apple-tree.

MELIPHYLLUM. (From *μαλας*, honey,

and *φύλλον*, a leaf; so called from the sweet smell of its leaf; or because bees gather honey from it.) Baum, or balm. See *Melissa*.

MELISSA. (From *μίσσα*, a bee, because bees gather honey from it.) *Citrargo. Citraria. Melinum. Melifolium. Melittis. Cedronella. Aspiastrum. Melissa citrina. Eriostem. Balm. Melissa officinalis* of Linnaeus. A native of the southern parts of Europe, but very common in our gardens. In its recent state, it has a roughish aromatic taste, and a pleasant smell of the lemon kind. It was formerly much esteemed in nervous diseases, and very generally recommended in melancholic and hypochondriacal affections; but, in modern practice, it is only employed when prepared as tea, as a grateful diluent drink in fevers, &c.

MELISSA CALAMINTHA. The systematic name of the field catmint. See *Calamintha*.

MELISSA CITRENA. See *Melissa*.

MELISSA GRANDIFLORA. The systematic name of the mountain calaminth. See *Calamintha magno flore*.

MELISSA OFFICINALIS. The systematic name of balm: See *Melissa*.

MELISSA TURCICA. See *Moldavica*.

MELISSOPHYLLUM. (From *μίσσα*, balm, and *φύλλον*, a leaf.) *Melissa*, or balm. Also a species of horehound, with leaves resembling balm.

MELITIS MELISSOPHYLLUM. The systematic name of the mountain balm. See *Melissophyllum*.

MELITISMUS. (From *μέλι*, honey.) A linctus, prepared with honey.

MELITTONA. (From *μέλι*, honey.) A confection made with honey. Honey-dew.

MELIZOMUM. (From *μέλι*, honey, and *ζυμος*, broth.) Mead. A drink prepared with honey.

MELLAGO. (From *mel*, honey.) Any medicine which has the consistence and sweetness of honey.

MELILOTUS. See *Melilotus*.

MELLINA. (From *mel*, honey.) Mead. A sweet drink prepared with honey.

MELLITA. (From *mel*, honey.) Preparations of honey.

MELO. The common melon. Musk melon. *Cucumis melo* of Linnaeus. This fruit, when ripe, has a delicious refrigerating taste, but must be eaten moderately, with pepper, or some aromatic, as all this class of fruits are obnoxious to the stomach, producing spasms and colic. The seeds possess mucilaginous qualities.

MELOCARPUS. (From *μίσσα*, an apple, and *καρπός*, fruit; from its resemblance to an apple, or its root.) The fruit of the aristolochia.

MELOE VESICATORIUS. The Spanish fly, now called *lytta*.

MELON. (*Μέλον*.) A disorder of the eye, in which it protuberates out of the socket.

Melon, common. See *Melo*.

Melon, musk. See *Melo*.

Melon, water. See *Citrullus*.

MELONGENA. Improperly called *mela insana. Solanum pomiferum.* Mad apple. The Spaniards and Italians eat it in sauce and in sweet-meats. The taste somewhat resembles citron.

MELOSIS. *Μελωσις.* A term which frequently occurs in Hippocrates, Do Capit. *Vulner.* for that search into wounds which is made by surgeons with the specillum, or probe.

MELOTIS. *Μελωτική.* Used for the lesser specillum, and often for that particular instrument contrived to search or cleanse the ear with, more commonly called *Auriscalpium*.

MELOTHRIA PENDULA. The systematic name of the small creeping cucumber plant. The inhabitants of the West Indies pickle the berries of this plant, and use them as we do capers.

MELYSOPHYLLUM. (From *μίσσα*, balm, and *φύλλον*, a leaf.) Mountain balm, or nettle. This elegant plant, *Melitis melissophyllum* of Linnaeus, is seldom used in the present day; it is said to be of service in uterine obstructions and calculous diseases.

MEMBRANA. See *Membrane*.

MEMBRANA HYALOIDEA. *Membrana arachnoidea.* The transparent membrane which includes the vitreous humour of the eye.

MEMBRANA PUPILLARIS. A very delicate membrane, of a thin and vascular texture, and an ash colour, arising from the internal margin of the iris, and totally covering the pupil, in the fetus, before the six months.

MEMBRANA RUYSCHIANA. The celebrated anatomist Ruysch discovered that the choroid membrane of the eye was composed of two laminae. He gave the name of *membrana ruyschiana* to the internal lamina, leaving the old name of choroides to the external.

MEMBRANA TYMPANA. The membrane covering the cavity of the drum of the ear, and separating it from the meatus auditorius externus. It is of an oval form, convex below the middle, towards the hollow of the tympanum, and concave towards the meatus auditorius, and convex above the middle towards the meatus, and concave towards the hollow of the tympanum. According to the observations of anatomists, it consists of six laminae; the first, and most external, is a production of the epidermis, the second is a production of the skin lining the auditory passage; the third is cellular membrane, in which

the vessels form an elegant net-work; the fourth is shining, thin, and transparent, arising from the periosteum of the meatus; the fifth is cellular membrane, with a plexus of vessels like the third; and the sixth lamina, which is the innermost, comes from the periosteum of the cavity of the tympanum. This membrane, thus composed of several laminae, has lately been discovered to possess muscular fibres.

MEMBRANALOGIA. (From *membrana*, a membrane, and *λογος*, a discourse.) Membranology. The doctrine of the common integuments and membranes.

MEMBRANE. *Membrana.* A thin expanded substance, composed of cellular texture, whose elastic fibres are so arranged and woven together, as to allow of great pliability. The membranes of the body are various, as the skin, peritoneum, pleura, dura mater, &c. &c.

MEMBRANOSUS. See *Tensor vaginæ femoris*.

MEMBRANUS. See *Tensor vaginæ femoris*.

MENBRILE OS. See *Occipital bone*.

Menagogues. See *Emmenagogues*.

MENDOSUS. (From *mendax*, counterfeit.) This term is used, by some, in the same sense as *spurius*, or *illegitimus*; *Mendosa Costæ*, false or spurious ribs; *Mendosa Sutura*, the squamous suture, in the skull, or bastard suture.

MENINGOPHYLAX. (From *μνινξ*, a membrane, and *φυλακς*, to guard.) An instrument to guard the membranes of the brain, while the bone is cut, or rasped, after the operation of the trepan.

MENINX. (From *μνιν*, to remain.) The Greek term for the membranes enveloping the brain. See *Dura mater* and *Pia mater*.

MENINX DURA. See *Dura mater*.

MENINX PIA. See *Pia mater*.

MENISPERMUM COCCULUS. The systematic name of the plant the berries of which are well known by the name of *Cocculus indicus*. Indian berries, or Indian cockles. Whilst green, they are used by the Indians to catch fish, which they have the power of intoxicating, and killing. In the same manner they catch birds, making the berry into a paste, forming it into small seeds, and putting these in places where they frequent.

MENORRHAGIA. (From *μνιν*, the menses, and *ρρηναι*, to break out.) *Hæmorrhagia uterina*. Flooding. An immoderate flow of the menses. A genus of disease in the class *pyrexia*, and order *hemorrhagia*, of Cullen, characterized by pains in the back, loins, belly; similar to those of labour, attended with a preternatural flux of blood from the vagina, more copious than natural. He distinguishes six species:—

1. *Menorrhagia rubra*, proper; from

women neither with child nor in child-birth.

2. *Menorrhagia alba*, serous; the fluor albus. See *Leucorrhœa*.

3. *Menorrhagia vitiorum*, from some local disease.

4. *Menorrhagia lochialis*, from women after delivery. See *Lochia*.

5. *Menorrhagia abortus*. See *Abortion*.

6. *Menorrhagia nabothi*, when there is a serous discharge from the vagina in pregnant women.

This disease seldom occurs before the age of puberty, and is often an attendant on pregnancy. It is in general a very dangerous disease, more particularly if it occur at the latter period, as it is then often so rapid and violent as to destroy the female in a very short time, where proper means are not soon adopted. Abortions often give rise to floodings, and at any period of pregnancy, but usually before the fifth month than at any other time. Moles, in consequence of an imperfect conception, at length become detached, and sometimes give rise to a considerable degree of hæmorrhage.

The causes which most frequently give rise to floodings are, violent exertions of strength, sudden surprises and frights, violent fits and passion, great uneasiness of mind, uncommon longings during pregnancy, over fullness of blood, profuse evacuations, general weakness of the system, external injuries, as blows and bruises, and the death of the child, in consequence of which the placenta becomes partially or wholly detached from the uterus, leaving the mouths of the vessels of the latter, which anastomosed with those of the latter, perfectly open. It is necessary to distinguish between an approaching miscarriage and a common flooding, which may be readily done by inquiring whether or not the hæmorrhage has proceeded from any evident cause, and whether it flows gently, or is accompanied with unusual pains. The former usually arises from some fright, surprise, or accident, and does not flow gently and regularly; but bursts out of a sudden, and again stops all at once, and also is attended with severe pains in the back and the bottom of the belly; whereas the latter is marked with no such occurrence. The further a woman is advanced in pregnancy, the greater will be the danger if floodings take place, as the mouths of the vessels are much enlarged during the last stage of pregnancy, and of course a vast quantity will be discharged in a short space of time.

MENSA. The second lobe of the liver was so called by the ancients.

MENSES. (From *mensis*, a month.)

MENSES CESSANTES. The menses departing.

MENSES DEFECTIVÆ. Menses defective or suppressed. The amenorrhœa of Cullen.

MENSES IMMODOICÆ. The menorrhagia rubra of Cullen.

Menses, immoderate flow of the. See *Menorrhagia*.

Menses, interruption of. See *Amenorrhœa*.

Menses, retention of. See *Catamenia*.

MENSES PHILOSOPHICÆ. A philosophical, or chymical month. According to some, it is three days and nights; others say it is ten; and there are who reckon it to be thirty or forty days.

MENSTRUATION. From the uterus of every healthy woman who is not pregnant, or who does not give suck, there is a discharge of blood at certain periods, from the time of puberty to the approach of old age; and, from the periods or returns of this discharge, it is called *Menstruation*. There are several exceptions to this definition. It is said that some women never menstruate; some menstruate while they continue to give suck; and others are said to menstruate during pregnancy; some are said to menstruate in early infancy, and others in old age; but such discharges, Dr. Denman is of opinion, may, with more propriety, be called morbid, or symptomatic: yet the definition is generally true.

At whatever time of life this discharge comes on, a woman is said to be at puberty: though of this state it is a consequence, and not a cause. The early or late appearance of the menses may depend upon the climate, the constitution, the delicacy or hardness of living, and upon the manners of those with whom young women converse. In Greece, and other hot countries, girls begin to menstruate at eight, nine, and ten years of age; but, advancing to the northern climes, there is a gradual protraction of the time till we come to Lapland, where women do not menstruate till they arrive at maturer age, and then in small quantities, at long intervals, and sometimes only in the summer. But, if they do not menstruate according to the genius of the country, it is said they suffer equal inconveniences as in warmer climates, where the quantity discharged is much greater, and the periods shorter. In this country, girls begin to menstruate from the fourteenth to the eighteenth year of their age, and sometimes at a later period, without any signs of the disease; but if they are luxuriously educated, sleeping upon down beds, and sitting in hot rooms, menstruation usually commences at a more early period.

Many changes in the constitution and appearance of women are produced at the time of their first beginning to menstruate,

Their complexion is improved, their countenance is more expressive and animated, their attitudes graceful, and their conversation more intelligent and agreeable; the tone of their voice becomes more harmonious, their whole frame, but particularly their breasts, are expanded and enlarged, and their minds are no longer engaged in childish pursuits and amusements.

Some girls begin to menstruate without any preceding indisposition; but there are generally appearances or symptoms which indicate the change that is about to take place. These are usually more severe at the first than in the succeeding periods: and they are similar to those produced by uterine irritation from other causes, as pains in the back and inferior extremities, complaints of the viscera, with various hysteric and nervous affections. These commence with the first disposition to menstruate, and continue till the discharge comes on, when they abate, or disappear, returning, however, with considerable violence in some women, at every period during life. The quantity of blood discharged at each evacuation depends upon the climate, constitution, and manner of living; but it varies in different women in the same climate, or in the same woman at different periods; in this country it amounts to about five or six ounces.

There is also a great difference in the time required for the completion of each period of menstruation. In some women the discharge returns precisely to a day, or an hour, and in others there is a variation of several days, without inconvenience. In some it is finished in a few hours, and in others it continues from one to ten days; but the intermediate time, from three to six days, is most usual.

There has been an opinion, probably derived from the Jewish legislator, afterwards adopted by the Arabian physicians, and credited in other countries, that the menstruous blood possessed some peculiar malignant properties. The severe regulations which have been made in some countries for the conduct of women, at the time of menstruation:—the expression used, *Isaiah*, chap. xxx. and in *Ezekiel*:—the disposal of the blood discharged, or of any thing contaminated with it:—the complaints of women attributed to its retention;—and the effects enumerated by grave writers, indicate the most dreadful apprehensions of its baneful influence. Under peculiar circumstances of health, or states of the uterus, or in hot climates, if the evacuation be slowly made, the menstruous blood may become more acrimonious or offensive than the common mass, or any other secretion from it; but in this country and age no malignity is suspected, the menstruous woman mixes in society as at all other

times, and there is no reason for thinking otherwise than that this discharge is of the most inoffensive nature.

At the approach of old age, women cease to menstruate; but the time of cessation is commonly regulated by the original early or late appearances of the menses. With those who began to menstruate at ten or twelve years of age, the discharge will often cease before they arrive at forty; but if the first appearance was protracted to sixteen or eighteen years of age, independently of disease, such women may continue to menstruate till they have passed the fiftieth, or even approach the sixtieth year of their age. But the most frequent time of the cessation of the menses, in this country, is between the forty-fourth and forty-eighth year; after which women never bear children. By this constitutional regulation of the menses, the propagation of the species is in every country confined to the most vigorous part of life: and had it been otherwise, children might have become parents, and old women might have had children, when they were unable to supply them with proper or sufficient nourishment. See *Catamenia*.

MENSTRUUM. Solvent. All liquors are so called, which are used as dissolvents, or to extract the virtues or ingredients by infusion, decoction, &c. The principal *menstrua*, made use of in *Pharmacy*, are water, vinous spirits, oils, acid, and alkaline liquors. Water is the *menstruum* of all salts, of vegetable gums, and of animal jellies. Of the first it dissolves only a determinate quantity, though of one kind of salt more than of another; and being thus saturated, leaves any additional quantity of the same salt untouched. It is never saturated with the two latter, but unites readily with any proportion of them, forming, with different quantities, liquors of different consistences. It takes up likewise, when assisted by trituration, the vegetable gummy resins, as ammoniacum and myrrh; the solutions of which, though imperfect, that is, not transparent, but turbid and of a milky hue, are nevertheless applicable to valuable purposes in medicine. Rectified spirit of wine is the *menstruum* of the essential oils and resins of vegetables; of the pure distilled oils of animals, and of soaps, though it does not act upon the expressed oil and fixed alkaline salt, of which soap is composed. Hence, if soap contains any superfluous quantity of either the oil or salt, it may, by means of this *menstruum*, be excellently purified therefrom. It dissolves, by the assistance of heat, volatile alkaline salts; and more readily the neutral ones, composed either of fixed alkali and the acetic acid, as the sal diureticus, or of volatile

alkali and the nitrous acid. Oils dissolve vegetable resins and balsams, wax, animal fats, mineral bitumens, sulphur, and certain metallic substances, particularly lead. The expressed oils are, for most of these bodies, more powerful *menstrua* than those obtained by distillation; as the former are more capable of sustaining, without injury, a strong heat, which is in most cases necessary to enable them to act. All acids dissolve alkaline salts, alkaline earths, and metallic substances. The different acids differ greatly in their action upon these last: one dissolving some particular metals, and another others. The vegetable acids dissolve a considerable quantity of zinc, iron, copper, and tin; and extract so much from the metallic part of the antimony as to become powerfully emetic; they likewise dissolve lead, if previously calcined by fire; but more copiously if corroded by their steam. The marine acid dissolves zinc, iron, and copper; and though it scarce acts on any other metallic substance in the common way of making solutions, may nevertheless be artfully combined with them all, except gold. The corrosive sublimate, and ammoniacal caustic of the shops, are combinations of it with mercury and the metallic part of antimony, effected by applying the acid, in the form of fume, to the subjects, at the same time strongly heated. The nitrous acid is the common *menstruum* of all metallic substances, except gold and the antimonial semi-metal, which are soluble only in a mixture of the nitrous and marine. The vitriolic acid easily dissolves zinc, iron, and copper; and may be made to corrode, or imperfectly dissolve, most of the other metals. Alkaline lixivia dissolve oils, resinous substances, and sulphur. Their power is greatly promoted by the addition of quicklime, instances of which occur in the preparation of soap and in the common caustic. Thus assisted, they reduce the flesh, bones, and other solid parts of animals, into a gelatinous matter. Solutions made in water and spirit of wine possess the virtue of the body dissolved; whilst oils generally sheathe its activity, and acids and alkalies vary its quality. Hence watery and spiritous liquors are the proper *menstrua* of the native virtues of vegetable and animal matters. Most of the foregoing solutions are easily effected, by pouring the *menstruum* on the body to be dissolved, and suffering them to stand together for some time, exposed to a suitable warmth. A strong heat is generally requisite to enable oils and alkaline liquors to perform their office; nor will acids act on some metallic bodies without its assistance. The action of watery and spiritous *menstrua* is likewise expedited by a moderate heat, though the quantity which

they afterwards keep dissolved, is not, as some suppose, by this means increased. All that heat occasions these to take up, more than they would do in a longer time in the cold, will, when the heat ceases, subside again. The action of acids on the bodies which they dissolve, is generally accompanied with heat, effervescence, and a copious discharge of fumes. The fumes which arise during the dissolution of some metals, in the vitriolic acid, prove inflammable; hence, in the preparation of the artificial vitriols of iron and zinc, the operator ought to be careful, especially where the solution is made in a narrow-mouthed vessel, lest, by the imprudent approach of a candle, the exhaling vapour be set on fire. There is another species of solution in which the moisture of air is the *menstruum*. Fixed alkaline salts, and those of the neutral kind, composed of alkaline salts and the vegetable acids, or of alkaline earths, and any acid except the vitriolic, and some metallic salts, on being exposed for some time to a moist air, gradually attract its humidity, and at length become liquid. Some substances, not dissoluble in water in its grosser form, as the butter of antimony, are easily liquified by this slow action of the aerial moisture. This process is termed *Deliquation*. The cause of solution assigned by some naturalists, namely, the admission of the fine particles of one body into the pores of another, whose figure fits them for their reception, is not just, or adequate, but hypothetical and ill-presumed; since it is found that some bodies will dissolve their own quantity of others, as water does of Epsom salt, alcohol of essential oils, mercury of metals, one metal of another, &c. whereas the sum of the pores or vacuities of every body must be necessarily less than the body itself, and consequently those pores cannot receive a quantity of matter equal to the body wherein they reside.

How a *menstruum* can suspend bodies much heavier than itself, which very often happens, may be conceived, by considering, that the parts of no fluids can be so easily separated, but they will a little resist or retard the descent of any heavy bodies through them: and that this resistance is, *ceteris paribus*, still proportional to the surface of the descending bodies. But the surface of bodies do by no means increase or decrease in the same proportion as their solidities do: for the solidity increases as the cube, but the surface only as the square of the diameter; wherefore it is plain, very small bodies will have much larger surfaces, in proportion to their solid contents, than larger bodies will, and consequently, when grown exceeding small, may easily be buoyed up in the liquor.

MENTABRA. (From *mentum*, the chin,

and *αγχα*, a prey.) *Impetigo*. An eruption about the chin, forming a tenacious crust, like that on scald heads.

MENTASTRUM. (Dim. of *mentha*, mint.) The red water-mint. See *Mentha aquatica*.

MENTHA. (From *Minthe*, the harlot who was changed into this herb.) *Hedysmum*. The name of a genus of plants in the Linnæan system. Class, *Didynamia*. Order, *Gymnospermia*. Mint.

MENTHA AQUATICA. *Mentastrium*. *Silybrium menthastrum*. *Mentha rotundifolia palustris*. Water-mint. This plant is frequent in moist meadows, marshes, and on the banks of rivers. It is less agreeable than the spear-mint, and in taste bitterer and more pungent. It may be used with the same intentions as the spear-mint, to which, however, it is much inferior.

MENTHA CATARIA. *Mentha felina*. *Herba felis*. *Calamintha*. *Nepetella*. Nep, or cat-mint. See *Nepeta*.

MENTHA CERVINA. The systematic name of the hart's penny-royal. See *Pulegium cervinum*.

MENTHA CRISPA. *Colymbifera minor*. The achilliza ageratum. This species of mentha has a strong and fragrant smell, its taste is warm, aromatic, and slightly bitter. In flatulences of the primæ viæ, hypochondriacal, and hysterical affections, it is given with advantage.

MENTHA PIPERITA. The systematic and pharmacopœial name of peppermint. *Mentha piperitis*. *Mentha piperita* of Linnaeus:—*floribus capitatis, foliis ovatis petiolatis, staminibus corollæ brevioribus*. The spontaneous growth of this plant is said to be peculiar to Britain. It has a more penetrating smell than any of the other mints; a strong pungent taste, glowing like pepper, sinking, as it were, into the tongue, and followed by a sense of coolness. The stomachic, antispasmodic, and carminative properties of peppermint, render it useful in flatulent cholics, hysterical affections, retchings, and other dispeptic symptoms, acting as a cordial, and often producing an immediate relief. Its officinal preparations are an essential oil, a simple water, and a spirit.

MENTHA PIPERITIS. See *Mentha piperita*.

MENTHA PULEGIUM. The systematic name of the pennyroyal. See *Pulegium*.

MENTHA SARACENICA. See *Balsamita mar.*

MENTHA SATIVA. See *Mentha viridis*.

MENTHA SPICATA. Hart-mint, and common spear-mint. *Mentha vulgaris*. See *Mentha viridis*.

MENTHA VIRIDIS. Spear-mint. Called also *Mentha vulgaris*. *Mentha spicata*. *Mentha viridis* of Linnaeus:—*spicis oblongis, foliis lanceolatis nudis serratis ses-*

silibus, staminibus corolla longioribus. The plant grows wild in many parts of England. It is not so warm to the taste as peppermint, but has a more agreeable flavour, and is therefore preferred for culinary purposes. Its medicinal qualities are similar to those of peppermint; but the different preparations of the former, though more pleasant, are, perhaps, less efficacious. The official preparations of spearmint are, an essential oil, a conserve, a simple water, and a spirit.

MENTI LEVATOR. See *Levator labii inferioris*.

MENTULA. (From *matah*, a staff, Heb.) The penis.

MENTULAGRA. (From *mentula*, the penis, and *agra*, a prey.) A disorder of the penis, induced by a contraction of the *erectores musculi*, and causing impotence.

MENYANTHES. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

MENYANTHES TRIFOLIATA. The systematic name of the buckbean. See *Trifolium paludosum*.

Mephetic air. See *Nitrogen*.

MEPHITIS. (From *mephuith*, a blast. Syr.) A poisonous exhalation. See *Contagion*.

MERCURIALIS. (From *Mercurius*, its inventor.) 1. The name of a genus of plants in the Linnæan system. Class, *Dioecia*. Order, *Emeandria*.

2. The pharmacopœial name of the French mercury. *Mercurialis annua* of Linnæus. The leaves of this plant have no remarkable smell, and very little taste. It is ranked among the emollient oleraceous herbs, and is said to be gently aperient. Their principal use has been in clysters.

MERCURIALIS ANNUA. The systematic name of the French mercury. See *Mercurialis*.

MERCURIALIS MONTANA. The *Mercurialis perennis* of Linnæus. See *Cynocrambe*.

MERCURIALIS PERENNIS. The systematic name of dog's mercury. See *Cynocrambe*.

MERCURIALIS SYLVESTRIS. The *Mercurialis perennis* of Linnæus. See *Cynocrambe*.

MERCURIUS ACETATUS. See *Hydrargyrum acetatus*.

MERCURIUS ALKALIZATUS. See *Hydrargyrum cum creta*.

MERCURIUS CALCINATUS. See *Oxydum hydrargyri rubrum*.

MERCURIUS CHEMICORUM. Quicksilver.

MERCURIUS CINNABARINUS. Cinnabar factitia, now called sulphuretum hydrargyri.

MERCURIUS CORROSIVUS. See *Oxymurias hydrargyri*.

MERCURIUS CORROSIVUS RUBER. See *Nitrico oxydum hydrargyri*.

MERCURIUS CORROSIVUS SUBLIMATUS. See *Oxymurias hydrargyri*.

MERCURIUS DULCIS SUBLIMATUS. Duplicated mercurial sublimate, now called submurias hydrargyri; formerly called calomelas.

MERCURIUS ENETICUS FLAVUS. See *Hydrargyrum vitriolatus*.

MERCURIUS MORTIS. See *Mercurius vite*.

MERCURIUS PRÆCIPITATUS ALBUS. See *Hydrargyrum præcipitatus albus*.

MERCURIUS PRÆCIPITATUS DULCIS. See *Submurias hydrargyri*.

MERCURIUS PRÆCIPITATUS RUBER. See *Nitrico oxydum hydrargyri*.

MERCURIUS VITÆ. The mercury of life. *Pulvis angelicus.* *Mercurius mortis.* *Algarothi pulvis.* An oxyd of antimony, formerly preferred to the vitrum antimonii, for making of emetic tartar.

MERCURY. *Hydrargyrum.* *Mercurius.* A metal found in five different states in nature. 1. Native, (*native mercury*;) adhering in small globules to the surface of cinnabar ores, or scattered through the crevices, or over the surfaces of different kinds of stones. 2. It is found united to silver, in the ore called *amalgam of silver*, or *native amalgam*. This ore exhibits thin plates, or grains; it sometimes crystallizes in cubes, parallelepipeds, or pyramids. Its colour is of a silver white, or grey; its lustre is considerably metallic. Combined with sulphur it constitutes *native cinnabar*, or sulphuret of mercury. This ore is the most common. It is frequently found in veins, and sometimes crystallized in tetrahedra, or three-sided pyramids. Its colour is red. Its streak metallic. 3. 4. Mercury, united either to muriatic or sulphuric acid, forms the ore called *horn quicksilver*, or *corneous mercury*. These ores are, in general, semi-transparent, of a grey or white colour, sometimes crystallized, but more frequently in grains. 5. United to oxygen it constitutes the ore called *native oxyd of mercury*. Mercurial ores particularly abound in Spain, Hungary, China, and South America.

Properties.—Mercury, or quicksilver, is the only one of the metals that remains fluid at the ordinary temperature of the atmosphere, but when its temperature is reduced to -40° Fahr. it assumes a solid form. This is a degree of cold, however, that only occurs in high northern latitudes, and, in our climate, mercury cannot be exhibited in a solid state, but by means of artificial cold. When rendered solid, it possesses both ductility and malleability. It crystallizes in octahedra, and contracts

strongly during congelation. It is divisible in very small globules. It presents a convex appearance in vessels to which it has little attraction, but is concave in those to which it does adhere. It becomes electric and phosphorescent by rubbing upon glass, and by agitation in a vacuum. It is a very good conductor of caloric, of electricity, and of Galvanism. The specific gravity of mercury is 13.563. Although fluid, its opacity is equal to that of any other metal, and its surface, when clean, has considerable lustre. Its colour is white, similar to silver. Exposed to the temperature of 600° Fahr. it is volatilized. When agitated in the air, especially in contact with viscous fluids, it becomes converted into a black oxyd. At a temperature nearly the same as that at which it boils, it absorbs about 14 or 16 per cent. of oxygen, and then becomes changed into a red crystallizable oxyd, which is spontaneously reducible by caloric and light at a higher temperature. The greater number of the acids act upon mercury, or are at least capable of combining with its oxyds. It combines with sulphur by trituration, but more intimately by heat. It is acted on by the alkaline sulphurets. It combines with many of the metals; these compounds are brittle, or soft, when the mercury is in large proportion. There is a slight union between mercury and phosphorus. It does not unite with carbon, or the earths.

Method of obtaining Mercury.—Mercury may be obtained pure by decomposing cinnabar, by means of iron filings. For that purpose, take two parts of red sulphuret of mercury (cinnabar), reduce it to powder, and mix it with one of iron filings, put the mixture into a stone retort, direct the neck of it into a bottle, or receiver, filled with water, and apply heat. The mercury will then be obtained in a state of purity.

In this process, the sulphuret of mercury, which consists of sulphur and mercury, is heated in contact with iron, the sulphur quits the mercury and unites to the iron, and the mercury becomes disengaged; the residue in the retort is a sulphuret of iron.

Mercury is a very useful article both in the cure of diseases and the arts. There is scarcely a disease against which some of its preparations are not exhibited; and over the venereal disease it possesses a specific power. It is considered to have first gained repute in curing this disease, from the good effects it produced in eruptive diseases. In the times immediately following the venereal disease, practitioners only attempted to employ this remedy with timorous caution, so that, of several of their formulæ, mercury scarcely composed a fourth part, and few cures were effected. On the other hand, empirics, who noticed the little efficacy of these small doses, ran into the opposite extreme,

and exhibited mercury in such large quantities, and with such little care, that most of their patients became suddenly attacked with the most violent salivations, attended with dangerous consequences. From these two very opposite modes of practice, there originated such uncertainty respecting what could be expected from mercury, and such fears of the consequences which might result from its employment, that every plan was eagerly adopted, which offered the least chance of cure without having recourse to this mineral. A medicine, however, so powerful, and whose salutary effects were seen, by attentive practitioners, amid all its inconveniences, could not sink into oblivion. After efforts had been made to discover a substitute for it, and it was seen how little confidence those means deserved on which the highest praises had been lavished, the attempts to discover its utility were renewed. A medium was pursued, between the too timid methods of those physicians who had first administered it, and the inconsiderate boldness of the empirics. Thus the causes from which both parties failed were avoided; the character of the medicine was revived in a more durable way, and from this period its reputation has always been maintained.

It was about this epoch that mercury began to be internally given; hitherto it had only been externally employed, which was done in three manners. The first was in the form of liniment, or ointment; the second as a plaster; and the third as a fumigation. Of the three methods just described, only the first is at present much in use, and even this is very much altered. Mercurial plasters are now only used as topical discutient applications to tumours and indurations. Fumigations, as anciently managed, were liable to many objections, particularly from its not being possible to regulate the quantity of mercury to be used, and from the effect of the vapour on the organs of respiration frequently occasioning trembling, palsies, &c. Frictions with ointment have always been regarded as the most efficacious mode of administering mercury.

Mercury is carried into the constitution in the same way as other substances, either by being absorbed from the surface of the body, or that of the alimentary canal. It cannot, however, in all cases, be taken into the constitution in both ways, for sometimes the absorbents of the skin will not readily receive it; at least no effect is produced, either on the disease or constitution, from this mode of application. On the other hand, the internal absorbents will, sometimes, not take up the medicine, or, at least, no effect is produced either on the disease or constitution. In many persons, the bowels can hardly bear mercury at all; and it should then be given in the

mildest form possible, conjoined with such medicines as will lessen or correct its violent effects, although not its specific ones, on the constitution. When mercury can be thrown into the constitution with propriety, by the external method, it is preferable to the internal plan; because the skin is not nearly so essential to life as the stomach, and is therefore in itself capable of bearing much more than the stomach. The constitution is also less injured. Many courses of mercury would kill the patient if the medicine were only given internally, because it proves hurtful to the stomach and intestines, when given in any form, or joined with the greatest correctors.

Mercury has two effects; one as a stimulus on the constitution and particular parts, the other as a specific on a diseased action on the whole body, or of parts. The latter action can only be computed by the disease disappearing.

In giving mercury in the venereal disease, the first attention should be to the quantity, and its visible effects in a given time; which, when brought to a proper pitch, are only to be kept up, and the decline of the disease to be watched; for by this we judge of the invisible or specific effects of the medicine, and know what variation in the quantity may be necessary. The visible effects of mercury affect either the whole constitution, or some parts capable of secretion. In the first, it produces universal irritability, making it more susceptible of all impressions. It quickens the pulse, increases its hardness, and occasions a kind of temporary fever. In some constitutions, it operates like a poison. In some it produces a kind of hectic fever; but such effects commonly diminish on the patient becoming accustomed to the medicine.

Mercury often produces pains, like those of rheumatism, and nodes of a scrophulous nature. The quantity of mercury to be thrown in for the cure of any venereal complaint, must be proportioned to the violence of the disease. A small quantity, used quickly, will have equal effects to those of a large one employed slowly; but if these effects are merely local, that is, upon the glands of the mouth, the constitution at large not being equally stimulated, the effects upon the diseased parts must be less, which may be known by the local disease not giving way in proportion to the effects of mercury on some particular part. If it be given in very small quantities, and increased gradually, so as to steal insensibly on the constitution, a vast quantity at a time may at length be thrown in without any visible effects at all.

The constitution, or parts, are more susceptible of mercury at first than afterwards.

Mercury occasionally attacks the bowels, and causes violent purging, even of blood.

This effect is remedied by intermitting the use of the medicine, and exhibiting opium. At other times, it is suddenly determined to the mouth, and produces inflammation, ulceration, and an excessive flow of saliva. To obtain relief in this circumstance, purgatives, nitre, sulphur, gum-arabic, lime-water, camphor, bark, kali-sulphuratum, blisters, &c. have been advised. Mr. Pearson, however, does not place much confidence in the efficacy of such means; and, the mercury being discontinued for a time, he recommends the patient to be freely exposed to a cold air, with the occasional use of cathartics, mineral acids, Peruvian bark, and the assiduous application of astringent gargles. The most material objection (says Mr. Pearson,) which I foresee against the method of treatment I have recommended, is the hazard to which the patient will be exposed of having the saliva suddenly checked, and of suffering some other disease in consequence of it.

The hasty suppression of a ptyalism may be followed by serious inconveniences, as violent pains, vomiting, and general convulsions.

Cold liquids taken into the stomach, or exposure of the body to the cold air, must be guarded against during a course of mercury. Should a suppression of the ptyalism take place, from any act of indiscretion, a quick introduction of mercury should be had recourse to, with the occasional use of the warm bath.

Mercury, when it falls on the mouth, sometimes produces inflammation, which now and then terminates in mortification. The ordinary operation of mercury does not permanently injure the constitution; but, occasionally, the impairment is very material; mercury may even produce local diseases, and retard the cure of chancres, buboes, and certain effects of the lues venerea, after the poison has been destroyed. Occasionally, mercury acts on the system as a poison, quite unconnected with its agency as a remedy, and neither proportionate to the inflammation of the mouth nor actual quantity of the mineral absorbed. Mr. Pearson has termed this morbid state of the system *erethismus*; it is characterised by great depression of strength, a sense of anxiety about the precordia, irregular action of the heart, frequent sighing, trembling, a small, quick, and sometimes intermitting pulse, occasional vomiting, a pale contracted countenance, a sense of coldness; but the tongue is seldom furred, and neither the natural or vital functions are much disturbed. When this effect of mercury takes place, the use of mercury should be discontinued, whatever may be the stage, extent, or violence of the venereal disease. The patient should be exposed to a dry and cool air, in such a way as not to give

fatigue; in this way the patient will often recover in ten or fourteen days. In the early stage, the erethismus may often be averted by leaving off the mercury and giving camphor mixture with volatile alkali. Occasionally, the use of mercury brings on a peculiar eruption, which has received the names of mercurial rash, eczema mercuriale, lepra mercurialis, mercurial disease, and erythema mercuriale.

In order that mercury should act on the human body, it is necessary that it should be oxydised, or combined with an acid. The mercury contained in the unguentum hydrargyri is an oxyde. The metal, however, in the mercurial ointment, is the most simple and least combined form of all its preparations, and hence, (says Mr. S. Cooper,) it not only operates with more mildness on the system, but with more specific effect on the disease. Various salts of mercury operate more quickly when given internally than mercurial frictions; but few practitioners of the present day confide in the internal use of mercury alone; particularly when the venereal virus has produced effects in consequence of absorption. Rubbing in mercurial ointment is the mode of affecting the system with mercury in the present day; and, as a substitute for this mode of applying mercury, Mr. Abernethy recommends the mercurial fumigation, where the patient has not strength to rub in ointment, and whose bowels will not bear the internal exhibition of it.

The preparations of mercury now in use are,

1. Nitrico-oxydum hydrargyri.
2. Oxydum hydrargyri cinereum.
3. Oxydum hydrargyri rubrum.
4. Oxy-murias hydrargyri.
5. Submurias hydrargyri.
6. Sulphuretum hydrargyri rubrum.
7. Hydrargyrus cum creta.
8. Hydrargyrus precipitatus albus.
9. Hydrargyrus purificatus.

Mercury, dog's. See *Cynocrambe*.

Mercury, English. See *Bonus henricus*.

Mercury, French. See *Mercurialis*.

MEROBALNEUM. (From *μερος*, a part, and *βαλανος*, a bath.) A partial bath.

MEROCELE. (From *μερος*, the thigh, and *κελη*, a tumour.) A femoral hernia. See *Hernia*.

MERON. (*Μερος*) The thigh.

MERUS. Applied to several things in the same sense as genuine, or unadulterated, as *merum vinum*, neat wine.

MESAREUM. (From *μερος*, the middle, and *αγα*, the belly.) The mesentery.

MESEMBRYANTHEUM CRYSTALLINUM. The juice of this plant, in a dose of four spoonfuls every two hours, it is asserted, has removed an obstinate spasmodic affection of the neck of the bladder, which would not yield to other remedies.

MESENTERIC. Meseraic. Belonging to the mesentery. See *Mesentery*.

MESENTERIC ARTERIES. *Arterie mesenterice.* Two branches of the aorta in the abdomen are so called. The superior mesenteric is the second branch; it is distributed upon the mesentery, and gives off the superior or right colic artery. The inferior mesenteric is the fifth branch of the aorta; it sends off the internal hæmorrhoidal.

MESENTERIC GLANDS. *Glandule mesenterice.* These are conglobate, and are situated here and there in the cellular membrane of the mesentery. The chyle from the intestines passes through these glands to the thoracic duct.

MESENTERIC PLEXUS OF NERVES. *Nervorum plexus mesentericus.* The superior, middle, and lower mesenteric plexuses of nerves are formed by the branches of the great intercostal nerves.

MESENTERIC VEINS. *Vena mesenterica.* They all run into one trunk, that evacuates its blood into the vena portæ. See *Vena portæ*.

MESENTERITIS. (From *μεσεντερ*, the mesentery.) An inflammation of the mesentery. A species of peritonitis of Cullen.

MESENTERY. (*Mesenterium*; from *μερος*, the middle, and *εντερον*, an intestine.) A membrane in the cavity of the abdomen, attached to the vertebrae of the loins, and to which the intestines adhere. It is formed of a duplicature of the peritonæum, and contains within it, adipose membrane, lacteals, lymphatics, lacteal glands, mesenteric arteries, veins, and nerves. Its use is to sustain the intestines in such a manner that they possess both mobility and firmness; to support and conduct with safety the blood-vessels, lacteals, and nerves; to fix the glands, and give an external coat to the intestines.

It consists of three parts: one uniting the small intestines, which receives the proper name of mesentery; another connecting the great intestines, termed mesocolon; and a third attached to the rectum, termed mesorectum.

MESERAIC. The same as mesenteric.

MESERION. See *Meserium*.

MESIRE. A disorder of the liver, mentioned by Avicenna, accompanied with a sense of heaviness, tumour, inflammation, pungent pain, and blackness of the tongue.

MESOCOLON. (From *μερος*, the middle, and *κολον*, the colon.) The portion of the mesentery to which the colon is attached. The mesentery and mesocolon are the most important of all the productions of the peritonæum. In the pelvis, the peritonæum spreads itself shortly before the rectum. But where that intestine becomes loose, and forms the semilunar curve, the peritonæum there rises consi-

derably from the middle iliac vessels, and region of the psoas muscle, double, and with a figure adapted for receiving the hollow colon. But above, on the left side, the colon is connected with almost no intermediate loose production to the peritonæum, spread upon the psoas muscle, as high as the spleen, where this part of the peritonæum, which gave a coat to the colon, being extended under the spleen, receives and sustains that viscus in a hollow superior recess.

Afterwards the peritonæum, from the left kidney, from the interval between the kidneys, from the large vessels, and from the right kidney, emerges forwards under the pancreas, and forms a broad and sufficiently long continuous production, called the transverse mesocolon, which, like a partition, divides the upper part of the abdomen, containing the stomach, liver, spleen, and pancreas, from the lower part. The lower plate of this transverse production is continued singly from the right mesocolon to the left, and serves as an external coat to a pretty large portion of the liver, and descending part of the duodenum. But the upper plate, less simple in the course, departs from the lumbar peritonæum at the kidney, and region of the vena cava, farther to the right than the duodenum, to which it gives an external membrane, not quite to the valve of the pylorus; and beyond this intestine, and beyond the colon, it is joined with the lower plate, so that a large part of the duodenum lies within the cavity of the mesocolon. Afterwards, in the region of the liver, the mesocolon is inflected, and descending over the kidney of the same side much shorter, it includes the right of the colon, as far as the intestinum cæcum, which rests upon the iliac muscle and the appendix, which is provided with a peculiar long curved mesentery. There the mesocolon terminates, almost at the bifurcation of the aorta.

The whole of the mesocolon and of the mesentery is hollow, so that the air may be forced in between its two laminae, in such a manner as to expand them into a bag. At the place where it sustains the colon, and also from part of the intestinum rectum, the mesocolon, continuous with the outer membrane of the intestine, forms itself into small slender bags, resembling the omentum, for the most part in pairs, with their loose extremities thicker and bifid, and capable of admitting air blown in between the plates of the mesocolon.

MESOCRANIUM. (From *mesos*, the middle, and *kranion*, the skull.) The crown of the head, or vertex.

MESOGASTRIUM. (From *mesos*, the middle, and *gaster*, the stomach.) The substance on the concave part of the stomach, which attaches itself to the adjacent parts.

MESOGLOSSUM. (From *mesos*, the middle, and *glossa*, the tongue.) A muscle inserted in the middle of the tongue.

MESOMERIA. (From *mesos*, the middle, and *meros*, the thigh.) The parts between the thighs.

MESOMPHALUM. (From *mesos*, the middle, and *omphalos*, the navel.) The middle of the navel.

MESOPHYTUM. (From *mesos*, the middle, and *ophrys*, the eyebrows.) The part between the eyebrows.

MESOPLEURUM. (From *mesos*, the middle, and *pleura*, a rib.) The space or muscles between the ribs.

MESORECTUM. (From *mesos*, the middle, and *rectum*, the straight gut.) The portion of peritonæum which connects the rectum to the pelvis.

MESOTHENAR. (From *mesos*, the middle, and *thana*, the palm of the hand.) A muscle situated in the middle of the palm of the hand.

MESPILUS. (*Oti n mesos pilos*, because it has a cap or crown in the middle of it.) The name of a genus of plants in the Linnæan system. Class, *Icosandria*. Order, *Pentagynia*.

2. The pharmacopœial name of the medlar. This fruit, and also its seeds, have been used medicinally. The tree which bears them is the *Mespilus germanica* of Linnæus. The immature fruit is serviceable in checking diarrhœas; and the seeds were formerly esteemed in allaying the pain attendant on nephritic diseases.

MESPILUS GERMANICA. The systematic name of the medlar-tree. See *Mespilus*.

METABASIS. (From *metabasis*, to digress.) *Metabole*. A change of remedy, of practice, or disease; or any change from one thing to another, either in the curative indications, or the symptoms of a distemper.

METABOL. See *Metabasis*.

METACARPAL BONES. The five longitudinal bones that are situated between the wrist and the fingers: they are distinguished into the metacarpal bone of the thumb, fore-finger, &c.

METACARPIUS. (From *metra*, after, and *carpos*, the wrist.) *Metacarpium*. That part of the hand which is between the wrist and the fingers.

METACARPUS. A muscle of the carpus. See *Adductor metacarpi minimi digiti manus*.

METACERASMA. (From *metra*, after, and *cerasma*, to mix.) *Cerasma*. A mixture tempered with any additional substance.

METACHEIRIKIA. (From *metachairō*, to perform by the hand.) Surgery. Any manual operation.

METACHORESIS. (From *metachōreō*, to digress.) The translation of a disease from one part to another.

METACINEMA. (From *metra* and *cinema*,

to remove.) Diseased pupil of the eye, or distortion of it.

METACARPUS. (From *meta*, after, and *carpus*, a knuckle.) The last joint of a finger, or that which contains the nail.

METALLAGE. (From *metallaxis*, to change.) A change in the state or treatment of a disease.

METALLURGIA. (From *metallon*, a metal, and *ergon*, work, labour.) That part of chymistry which concerns the operations of metals.

METALS. We are at present acquainted with twenty-two metals, essentially differing from each other, namely,

Arsenic. Tungsten. Molybdena. Chrome. Columbium. Titanium. Uranium. Cobalt. Nickel. Manganese. Bismuth. Antimony. Tellurium. Mercury. Zinc. Tin. Lead. Iron. Copper. Silver. Gold. Platina.

The ancient division of these substances into noble or perfect, and semi or imperfect metals, is now abandoned, and we class them into different genera; proceeding in a gradation from those which possess, in a certain sense, the *least*, to those which possess the *most* metallic properties.

Classification of Metals.

1. *Very brittle and acidifiable Metals.*—Arsenic. Tungsten. Molybdena. Chrome. Columbium.

2. *Less brittle and simply oxydable Metals.*—Titanium. Uranium. Cobalt. Nickel. Manganese. Bismuth. Antimony. Tellurium.

3. *Partly ductile and oxydable Metals.*—Mercury. Zinc.

4. *Very ductile and easily oxydable Metals.*—Tin. Lead. Iron. Copper.

5. *Exceedingly ductile and difficultly oxydable Metals.*—Silver. Gold. Platina.

All the metals are found in the bowels of the earth, though sometimes they are on the surface. They are met with in different combinations with other matters, such as sulphur, oxygen, and acids; particularly with the carbonic, muriatic, sulphuric, and phosphoric acids. They are also found combined with each other, and sometimes, though rarely, in a pure metallic state, distinguishable by the naked eye.

In their different states of combination, they are said to be mineralized, and are called **ORES**. The ores of metals are, for the most part, found in nature in mountainous districts; and always in such as form a continued chain. There are mountains which consist entirely of iron ore, but, in general, the metallic part of a mountain bears a very inconsiderable proportion to its bulk. Ores are also met with in the cavities or crevices of rocks, forming what are termed **VEXES**, which are more easily discovered in these situations than when they lie level in plains.

The metallic matter of ores is very generally incrustated, and intermingled with some

earthy substance, different from the rock in which the vein is situated; which is termed its **MATRIX**. This, however, must not be confounded with the mineralizing substance with which the metal is combined, such as sulphur, &c.

General physical Properties of Metals.

All metals are combustible bodies. They possess a certain brilliancy, in consequence of the complete reflection of the light that falls upon them, which is termed *metallic lustre*. They are the most dense and heavy substances in nature; the heaviest fossil, not metallic, having a specific gravity much below that of the lightest metal. They are the most opaque of all bodies. A stone of the greatest opacity, when divided into thin plates, has more or less transparency; whereas gold is the only metal which admits of being reduced to such a degree of thinness as to admit the smallest perceptible transmission of light. Gold leaf, which is about 1-280,000 part of an inch in thickness, transmits light of a lively green colour; but silver, copper, and all the rest of the metals, are perfectly opaque. Another property, which belongs exclusively to metals, (though they do not all possess it,) is *malleability*; by which is meant a capacity of having their surface increased, either in length or breadth, without being liable to fracture. This capacity is not precisely the same in those metals which do possess it, for some, which admit of extension when struck with a hammer, cannot be drawn into wire, which property is termed *ductility*: this property depends, in some measure, on another peculiar quality of metals, namely *tenacity*; by which is meant the power which a metallic wire, of a given diameter, has of resisting the action of a weight suspended from its extremity. All metals are fusible, though the degree of temperature at which this can be effected differs very much. Mercury is always fluid at the ordinary temperature of our atmosphere, while platina can scarcely be melted by the most intense heat of our furnaces. Metals are perfectly opaque when in a state of fusion; and are crystallizable when suffered to cool slowly and undisturbed. The tetrahedron and the cube are their primitive figures, though they very often take the octahedral form. They can likewise be volatilized at very high temperatures. They are the best conductors of caloric and electricity. Their susceptibility of combination is very great: they unite with carbon, sulphur, and phosphorus. They do not combine with earths by fusion; but their oxyds readily unite to acids, alkalies, and earths. They decompose water and several acids. Some effect this at common temperatures: some require a red heat, and others the interposition of another body. Water does not dissolve any of the metals, though it is a solvent of some of their oxyds. They are

insoluble in ardent spirit, ether, or oils. They are all capable of combining with oxygen, though many of them require very high temperatures to effect this union, and others cannot be united to it but in an indirect manner. Most of the metals can be combined with each other; they then form alloys, many of which are of the greatest utility in the arts.

METAMORPHOPSIA. (From *μεταμορψωσις*, a change, and *ὄψις*, sight.) *Visus defiguratus*. Disfigured vision. It is a defect in vision, by which persons perceive objects changed in their figures. The species are, 1. *Metamorphopsia acuta*, when objects appear much larger than their size. 2. *Metamorphopsia diminuta*, when objects appear diminished in size, arising from the same causes as the former. 3. *Metamorphopsia mutans*, objects seem to be in motion; to the vertiginous and intoxicated persons, every thing seems to stagger. 4. *Metamorphopsia tortuosa seu flexuosa*, when objects appear tortuous, or bending. 5. *Metamorphopsia inversa*, when all objects appear inverted. 6. *Metamorphopsia imaginaria*, is the vision of a thing not present, as may be observed in the delirious, and in maniacs. 7. *Metamorphopsia from a remaining impression*: it happens to those who very attentively examine objects, particularly in a great light, some time after to perceive the impression.

METAPEDIUM. (From *μετα*, after, and *πῦς*, the foot.) The metatarsus.

METAPHRENUM. (From *μετα*, after, and *φρεν*, the diaphragm.) That part of the back which is behind the diaphragm.

METAPOROPOISIS. (From *μετα*, *πορος*, a duct, and *ποιω*, to make.) A change in the pores of the body.

METAPTOSIS. (From *μεταπτεω*, to digress.) A change from one disease to another.

METASTASIS. (From *μεταστημι*, to change, to translate.) The translation of a disease from one place to another.

METASTYCRISIS. (From *μεταστυκρηνω*, to transmute.) Any change of constitution.

METATARSAL BONES. The five longitudinal bones between the tarsus and the toes; they are distinguished into the metatarsal bone of the great toe, fore toe, &c.

METATARSUS. (From *μετα*, after, and *ταρσος*, the tarsus.) That part of the foot between the tarsus and toes.

METELLA NUX. See *Nux vomica*.

METTORIUM. (From *μεττωρος*, a vapour.) 1. A flatulent dropsy.

2. A tympanic state of the abdomen, taking place suddenly in acute diseases.

METTOROS. (*Mettoros*: from *μετα*, and *ωρος*, to elevate.) Elevated, suspended, erect, sublime, tumid. Galen expounds pains of this sort, as being those which affect the peritonæum, or other more su-

perficial parts of the body: these are exposed to the more deep-seated ones.

METHUELIN. A drink prepared from honey by fermentation.

METHEMBRIUM. (From *μετα*, and *εμβριω*, a day.) A quotidian fever.

METHODICA MEDICINA. Signifies that practice which was conducted by rules, such as are taught by Galen and his followers, in opposition to the empirical practice.

METHODUS. (From *μετα*, and *ὁδος*, a way.) The method, or ratio, by which any operation or cure is conducted.

METOPION. *Metopion*. American sumach, a species of *Rhus*. It is a name of the bitter almond, also of an oil, or an ointment, made by Dioscorides, which was thus called because it had galbanum in it, which was collected from a plant called *Metopium*.

METOPION. METOPION. An ointment made of galbanum.

METOPUM. (From *μετα*, after, and *ὀφθαλμος*, the eye.) The forehead.

METOSIS. A kind of amaurosis, from an excess of short sightedness.

METRA. (From *μητρ*, a mother.) The womb.

METRECHYTA. (From *μετρα*, the womb, and *εχυνω*, to pour into.) Injections into the womb.

METRECHYTES. (From *μετρα*, the womb, and *εχυνω*, to pour in.) A syringe to inject fluids into the womb.)

METRITIS. (From *μετρα*, the womb.) Inflammation of the uterus. See *Hysteritis*.

METROCELIDES. (From *μητρ*, a mother, and *κελε*, a spot, or mole.) A mole, or mark, impressed upon the child by the mother's imagination.

METROCELLIS. (From *μητρ*, a mother, and *κελε*, a blemish.) A mole, or mark, impressed upon the child by the mother's imagination.

METRO-MANIA. A rage for reciting verses. In the *Acta Societatis Medicæ Havniensis*, published 1779, is an account of a tertian attended with remarkable symptoms; one of which was the *metro-mania*, which the patient spoke extempore, having never before had the least taste for poetry; when the fit was off, the patient became stupid, and remained so till the return of the paroxysm, when the poetical powers returned again.

METROPTOSIS. (From *μετρα*, the uterus, and *πτωω*, to fall down.) *Prolapsus uteri*. The descent of the uterus through the vagina.

METRORHAGIA. (From *μετρα*, the womb, and *ρηνω*, to break out.) An excessive discharge from the womb.

MEU. See *Meum athamanticum*.

MEUM ATHAMANTICUM. (From *μινω*, less; so called, according to Minshew, from its diminutive size.) *Meu*. Spiguel.

Baldmoney. The root of this plant, *Æthusa meum*, is recommended as a carminative, stomachic, and for attenuating viscid humours, and appears to be nearly of the same nature as lovage, differing in its smell being rather more agreeable, somewhat like that of parsnips, but stronger, and being in its taste less sweet, and more warm, or acrid.

MEXICANUM. (From *Mexico*, whence it is brought.) A name of the balsam of Peru.

Mexico seed. See *Ricinus*.

Mexico tea. See *Botrys mexicana*.

Mezereon. See *Mezereum*.

MEZEREUM. (A word of some barbarous dialect.) *Mezereon*. Spurge-olive. Widow-wail. This plant, *Daphne mezereum* of Linnaeus:—*floribus sessilibus ternis caulinis, foliis lanceolatis deciduis*, is extremely acrid, especially when fresh, and, if retained in the mouth, excites great and long continued heat and inflammation, particularly of the mouth and fauces; the berries also have the same effects, and, when swallowed, prove a powerful corrosive poison, not only to man, but to dogs, wolves, and foxes. The bark of the root is the part employed medicinally in the decoction *sarsaparilla compositum*, to assist mercury in resolving nodes and other obstinate symptoms of syphilis. The antisyphilitic virtues of mezereum, however, have been by many writers very justly doubted. The result of my own experience (says Mr. Pearson, of the Lock Hospital) by no means accords with the representation given of this root by former writers. From all that I have been able to collect, in the course of many years observation, I feel myself authorised to assert, unequivocally, that the mezereum has not the power of curing the venereal disease in any one stage, or under any one form. If a decoction of this root should ever reduce a venereal node, where no mercury has been previously given, yet the patient will by no means be exempted from the necessity of employing mercury for as long a space of time, and in as large a quantity, as if no mezereum had been taken. With respect to the power it is said to possess, of alleviating the pain, and diminishing the bulk of membranous nodes, nothing peculiar and appropriate can be ascribed to the mezereum on these accounts, since we obtain the same good effects from sarsaparilla, guaiacum, volatile alkali, blistering plasters, &c. Nevertheless, venereal nodes, which have subsided under the use of any of these articles of the materia medica, will appear again, and often with additional symptoms, if a full and efficacious course of mercury be not submitted to. It has, indeed, been alleged, that mezereum always alleviates the pain occasioned by a venereal node, and generally reduces it, where the periosteum only is affected; and that it sel-

dom fails of removing those enlargements of the periosteum which have not yielded during the administration of mercury.

That some instances of success, in cases like these, may have fallen to the share of those who made the assertion, it would not become me to deny; but I have met with few such agreeable evidences of the efficacy of this medicine. I have given the mezereum in the form of a simple decoction, and also as an ingredient in compound decoctions of the woods, in many cases, where no mercury had been previously employed, but never with advantage to a single patient. I have also tried it, in numerous instances, after the completion of a course of mercury; yet, with the exception of two cases, where the thickened state of the periosteum was removed during the exhibition of it, I never saw the least benefit derived from taking this medicine. In a few cases of anomalous pains, which I supposed were derived from irregularities during a mercurial course, the mezereum was of service, after I had tried the common decoction of the woods without success; but even in this description of cases, I have always found it a very uncertain remedy. I have made trial of this vegetable in a great number of scrofulous cases, where the membranes covering the bones were in a diseased state, and I am not sure that one single patient obtained any evident and material benefit from it.

The late Dr. Cullen, whose reports may justly claim attention from all medical men, when treating of the mezereum, in his *Materia Medica*, says, "I have frequently employed it in several cutaneous affections, and sometimes with success." It were to have been wished, that the professor of medicine had specified what those diseases of the skin were, in which the mezereum was sometimes employed with success; for, if I except an instance or two of lepra, in which the decoction of this plant conferred a temporary benefit, I have very seldom found it possessed of medicinal virtue, either in syphilis, or in the sequelæ of that disease, in scrofula, or in cutaneous affections. Indeed the mezereum is of so acrimonious a nature, often producing heat and other disagreeable sensations in the fauces, and, on many occasions, disordering the primæ viæ, that I do not often subject my patients to the certain inconveniences which are connected with the primary effects of this medicine, as they are rarely compensated by any other important and useful qualities.

MEZEREUM ACETATUM. Very thin slices of the bark of the fresh mezereon root are to be steeped for twenty-four hours in common vinegar. The late Dr. Morris usually directed the application of this remedy to issues, when a discharge from them could not be encouraged by the

common means. It generally answers this purpose very effectually in the course of one night, the pea being removed, and a small portion of the bark applied over the opening.

MIASMA. (From *μῆμα*, to infect.) See *Contagion*.

MICROCOSMIC BEZOAR. See *Calculus*.

MICROLEUCONTYPHÆA. (From *μικρός*, small, *λευκός*, white, and *νύμφα*, the water-lily.) The small white water lily.

MICRONTYPHÆA. (From *μικρός*, small, and *νύμφα*, the water-lily.) The smaller water-lily.

MICROCHIS. (From *μικρός*, small, and *ὄρχις*, a testicle.) One whose testicles are unusually small.

MICROSPHYXIA. (From *μικρός*, small, and *σπύξ*, the pulse.) A debility and smallness of the pulse.

Midriff. See *Diaphragm*.

MISMA. (From *μῆμα*, to mix.) A confection, or ointment.

MIGRANA. A corruption of *hæmicrania*.

Miffoil, common. See *Millefolium*.

MILIARIA. (From *milium*, millet; so called because the small pustules, or vesicles, upon the skin resemble millet-seed.) Miliary fever. A genus of disease in the class *pyrexie*, and order *exanthemata* of Cullen; characterized by synchus; cold stage considerable; hot stage attended with anxiety and frequent sighing; perspiration of a strong and peculiar smell; eruption, preceded by a sense of pricking, first on the neck and breast, of small red pimples, which in two days become white pustules, desquamate, and are succeeded by fresh pimples. Miliary fever has been observed to affect both sexes, and persons of all ages and constitutions; but females of a delicate habit are most liable to it, particularly in child-bed. Moist variable weather is most favourable to its appearance, and it occurs most usually in the spring and autumn. It is by some said to be a contagious disease, and has been known to prevail epidemically.

Very violent symptoms, such as coma, delirium, and convulsion-fits, now and then attend miliary fever, in which case it is apt to prove fatal. A numerous eruption indicates more danger than a scanty one. The eruption being steady is to be considered as more favourable than its frequently disappearing and coming out again, and it is more favourable when the places covered with the eruption appear swelled and stretched than when they remain flaccid. According to the severity of the symptoms, and depression of spirits, is the danger greater. See also *Sudamina*.

MILLOLUM. (Dim. of *milium*, millet.) A small tumour on the eyelids, resembling in size a millet-seed.

MILTARIS. (From *miles*, a soldier; so called from its efficacy in curing fresh wounds.) See *Millefolium*.

MILTARIS HERBA. See *Millefolium*.

MILUM. (From *milium*, a millet-seed.) *Grutum.* A very white and hard tubercle, in size and colour resembling a millet-seed. Its heat is immediately under the cuticle, so that, when pressed, the contents escape, appearing of an atheromatous nature.

MILUM SOLIS. See *Lithospermum*.

MILK. *Lac.* A fluid secreted by peculiar glands, and designed to nourish young animals in the early part of their life. It is of an opaque white colour, a mild saccharine taste, and a slightly aromatic smell. It is separated immediately from the blood, in the breasts or udders of female animals. Man, quadrupeds, and cetaceous animals, are the only creatures which afford milk. All other animals are destitute of the organs which secrete this fluid. Milk differs greatly in the several animals.

The following are the *Properties* of animal and human milk:

Milk separates spontaneously into *cream, cheese, and serum of milk*; and that sooner in a warm situation than in a cold one. In a greater temperature than that of the air, it acceces and coagulates, but more easily and quicker by the addition of acid salts, or coagulating plants. *Lime-water* coagulates milk imperfectly. It is not coagulated by *alkali* pure, for it dissolves its caseous part. With carbonated *alkali* the caseous and cremoraceous parts of milk are changed into a liquid soap, which separates in the form of white flakes: such milk, by boiling, is changed into a yellow and then into a brown colour. Milk, distilled to dryness, gives out an insipid water, and leaves a whitish brown extract, called the *extract of milk*; which, dissolved in water, makes a milk of less value. Milk, fresh drawn, and often agitated in a warm place, by degrees goes into the vinous fermentation, so that alcohol may be drawn over by distillation, which is called *spirit of milk*. It succeeds quicker, if yeast be added to the milk. Mare's milk, as it contains the greatest quantity of the sugar of milk, is best calculated for vinous fermentation.

The *Principles* of milk, or its integral parts, are, 1. The *Aroma*, or odoriferous volatile principle, which flies off from fresh-drawn milk in the form of visible vapour. 2. *Water*, which constitutes the greatest part of milk. From one pound, eleven ounces of water may be extracted by distillation. This water, with the sugar of milk, forms the *serum of the milk*. 3. *Bland oil*, which, from its lightness, swims on the surface of milk after standing, and forms the *cream of milk*. 4. *Cheese*, separated by coagulating milk, falls to the bottom of the vessel, and is the animal gluten. 5. *Sugar*, obtained from the serum of milk by evaporation. It unites the caseous and butyraceous part with the water of the milk. 6. *Some new*.

tral salts, as the *muriate* of potash and *muriate* of lime, which are accidental, not being found at all times, nor in every milk. These principles of milk differ widely in respect to quantity and quality, according to the diversity of the animals.

The *aroma* of the milk is of so different an odour, that persons accustomed to the smell, and those whose olfactory nerves are very sensible, can easily distinguish whether milk be that of the cow, goat, mare, ass, or human. The same may be said of the serum of the milk, which is properly the seat of the aroma. The *serum* of milk is thicker and more copious in the milk of the sheep and goat, than in that of the ass, mare, or human milk. The *butter* of goats and cows is easily separated from the milk, and will not again unite itself with the butter-milk. Sheep's butter is soft, and not of the consistence of that obtained from the cow and goat. Asses', mares', and human butter, can only be separated in the form of cream; which cream, by the assistance of heat, is with ease again united to the milk from which it is separated. The *cheese* of cows' and goats' milk is solid and elastic, that from asses and mares soft, and that from sheep's milk almost as soft as gluten. It is never separated spontaneously from the milk of a woman, but only by art, and is wholly fluid. The *serum* abounds most in human, asses', and mares' milk. The milk of the cow and goat contains less, and that of the sheep least of all. The *sugar* of milk is in the greatest quantity in the mares and asses, and somewhat less in the human milk.

MILK, ASSES'. Asses' milk has a very strong resemblance to human milk in colour, smell, and consistence. When left at rest for a sufficient time, a cream forms upon its surface, but by no means in such abundance as a woman's milk. Asses' milk differs from cows' milk, in its cream being less abundant and more insipid; in its containing less curd; and in its possessing a greater proportion of sugar.

MILK, COWS'. The milk of women, mares, and asses, nearly agree in their qualities; that of cows, goats, and sheep, possess properties rather different. Of these, cows' milk approaches nearest to that yielded by the female breast, but differs very much in respect to the aroma; it contains a larger proportion of cream and cheese, and less serum than human milk; also less sugar than mares' and asses' milk.

Cows' milk forms a very essential part of human sustenance, being adapted to every state and age of the body; but particularly to infants, after being weaned.

MILK, EWES'. This resembles almost precisely that of the cow; its cream, however, is more abundant, and yields a butter not so consistent as cow's milk butter. It makes excellent cheese.

MILK, GOATS'. It resembles cows', except in its greater consistence; like that milk, it throws up abundance of cream, from which butter is easily obtained.

MILK, HUMAN. The white, sweetish fluid, secreted by the glandular fabric of the breasts of women. The *secretory organ* is constituted by the great conglomerate glands, situated in the fat of both breasts, above the *musculus pectoralis major*. From each acinus composing a mammary gland, there arises a radicle of a *lactiferous* or *galactiferous* duct. All these canals, gradually converging, are terminated without anastomosis, in the papillæ of the breasts, by many orifices, which, upon pressure, pour forth milk. The smell of fresh-drawn milk is peculiar, animal, fatuous, and not disagreeable. Its taste sweetish, soft, bland, agreeable. The specific gravity is greater than water, but lighter than blood; hence it swims on it. Its colour is white and opaque. In consistence it is oily and aqueous. A drop put on the nail flows slowly down, if the milk be good.

Time of Secretion. The milk most frequently begins to be secreted in the last months of pregnancy; but, on the third day after delivery, a serous milk, called *Colostrum*, is separated; and at length pure milk is secreted very copiously into the breasts, that, from its abundance, often spontaneously drops from the nipples.

If the secretion of milk be daily promoted by suckling an infant, it often continues many years, unless a fresh pregnancy supervene. The quantity usually secreted within twenty-four hours, by nurses, is various, according as the nourishment may be more or less chylous. It appears that not more than two pounds of milk are obtained from five or six pounds of meat. But there have been known nurses, who have given from their breasts two, or even more than three pounds, in addition to that which their child has sucked. That the origin of the milk is derived from chyle carried with the blood of the mammary arteries into the glandular fabric of the breasts is evident, from its more copious secretion a little after meals; its diminished secretion from fasting; from the smell and taste of food or medicines in the secreted milk; and lastly, from its spontaneous *acrescence*; for humours perfectly animal become putrid.

The *milk of a woman* differs: 1. in respect to *food*. The milk of a woman who suckles, living upon *vegeto-animal food*, never *acresces* nor coagulates spontaneously, although exposed for many weeks to the heat of a furnace. But it evaporates gradually in an open vessel, and the last drop continues thin, sweet, and bland. The reason appears to be, that the caseous and *eremorrhaceous* parts cohere together, by means of the sugar, more intimately

than in the milk of animals, and do not so easily separate; hence its acescence is prevented. It does *accesce*, if mixed or boiled with vinegar, juice of lemons, cream or tartar, dilute sulphuric acid, or with the human stomach. It is *coagulated* with the acid of salt, or nitre, and by an acid gastric juice of the infant; for infants often vomit up the coagulated milk of the nurse. The milk of a sucking woman, who lives upon vegetable food only, like cows' milk, easily and of its own accord *acesces*, and is acted upon by all coagulating substances like the milk of animals. 2. In respect of *the time of digestion*. During the first hours of digestion the chyle is crude, and the milk less subacted; but towards the twelfth hour after eating, the chyle is changed into blood, and then the milk becomes yellowish and nauseous, and is spit out by the infant. Hence the best time for giving suck is about the fourth or fifth hour after meals. 3. In respect of *the time after delivery*. The milk secreted immediately after delivery is serous, purges the bowels of the infant, and is called *colostrum*. But in the following days it becomes thicker and more pure, and the longer a nurse suckles, the thicker the milk is secreted; thus, new-born infants cannot retain the milk of a nurse who has given suck for a twelve-month, on account of its spissitude. 4. In respect of *food and medicines*. Thus, if a nurse eat garlic, the milk becomes highly impregnated with its odour, and is disagreeable. If she indulge too freely in the use of wine or beer, the infant becomes ill. From giving a purging medicine to a nurse, the child also is purged; and, lastly, children affected with tormina of the bowels, arising from acids, are often cured by giving the nurse animal food. 5. In respect of *the affections of the mind*. There are frequent examples of infants being seized with convulsions from sucking mothers irritated by anger. An infant of one year old, while he sucked milk from his enraged mother, on a sudden was seized with a fatal hæmorrhage, and died. Infants at the breast in a short time pine away, if the nurse be afflicted with grievous care; and there are also infants who, after every coition of the mother, or even if she menstruate, are taken ill.

The use of the mother's milk is, 1. It affords the native *aliment* to the new-born infant, in which respect milk differs little from chyle. Those children are the strongest, who are nourished the longest by the mother's milk. 2. The *colostrum* should not be rejected; for it relaxes the bowels, which in new-born infants ought to be open, to clear their intestines of the *meconium*. 3. *Lactation* defends the mother from a dangerous reflux of the milk into the blood, whence lacteal metastasis, and leucorrhæa, are so frequent in lying-in women, who do not give suck. The mo-

tion of the milk also being hastened through the breast by the sucking of the child, prevents the very common induration of the breast, which arises in consequence of the milk being stagnated. 4. *Men* may live upon milk, unless they have been accustomed to the drinking of wine. For all nations, the Japanese alone excepted, use milk, and many live upon it alone.

MILK, MARES'. This is thinner than that of the cow, but scarcely so thin as human milk. Its cream cannot be converted into butter by agitation. The whey contains sugar.

MILK BLOTCHES. An eruption of white vesicles, which assume a dark colour, resembling the blackening of the small-pox, and are succeeded by scabs producing an ichthy matter, attended with considerable itchiness. It generally appears on the forehead and scalp, extending half over the face, and at times even proceeding farther. The period of its attack is the time of teething; and it is probably the same disease as the *crustæa lactea*.

Milk fever. See *Puerperal fever*.

Milk-teeth. See *Teeth*.

MILK-THISTLE. The leaves of this plant, when young, surpasses, when boiled, the finest cabbage, and possesses diuretic qualities. See also *Carduus marie*.

Milk-veitch. See *Tragacantha*.

Milk-wort. See *Polygala*.

Milk-wort, rattle-snake root. See *Seneca*.

MILLEFOLIUM. (From *mille*, a thousand, and *folium*, a leaf; named from its numerous leaves.) *Achillea Myriophylon*. *Chiliphullon*. *Lumbus veneris*. *Militaris herba*. *Stratiotes*. *Carpentaria*. *Speculum veneris*. Common yarrow, or milkfoil. The leaves and flowers of this indigenous plant, *Achillea millefolium* of Linnaeus:—*foliis bipinnatis nudis; laciniis linearibus dentatis; caulibus superne sulcatis*, have an agreeable, weak, aromatic smell, and a bitterish, rough, and somewhat pungent taste. They are both directed for medicinal use, in the Edinburgh Pharmacopœia; in the present practice, however, they are almost wholly neglected.

MILLEMORBIA. (From *mille*, a thousand, and *morbus*, a disease; so called from its use in many diseases.) See *Scrophularia vulgaris*.

MILLEPEDE. See *Millepedes*.

MILLEPEDES. (From *mille*, a thousand, and *pes*, a foot, named from their numerous feet.) *Milipeda*. Wood-lice. The *Oniscus asellus* of Linnaeus. These insects, though they obtain a place in the pharmacopœias, are very seldom used medicinally in this country; they appear to act as stimulants and slight diuretics, and for this purpose they ought to be administered in a much greater dose than is usually prescribed. The expressed juice, or forty or fifty living millepedes, given in a mild

drink, is said to cure very obstinate jaundices.

MILLET-SEED. The fruit of the *Panicum miliaceum* of Linnæus. They are esteemed as a nutritious article of diet, and are often made into puddings in this country.

MILLET-SEED, INDIAN. The fruit of the *Panicum italicum* of Linnæus. It is much esteemed in Italy, being a constant ingredient in soups, and made into a variety of forms for the table.

MILLIUM. (From *mille*, a thousand; so called from the multitude of its seed.) *Milium*. The millet.

MIL-MOUNTAIN. See *Linum Catharticum*.

MILPHOSIS. *Μιλφωσις*. A baldness of the eyebrows.

MILTOR. *Μιλτορ*. Minium, or red-lead.

Miltwaste. See *Ceterach*.

MILZADELLA. (From *milza*, the spleen, Span.; so called from its supposed virtues in diseases of the spleen.) The herb archangel.

MIMOSA CATECHU. The systematic name of the tree which affords the terra japonica. See *Catechu*.

MIMOSA NILOTICA. Supposed to be the tree which afforded the gum-arabic, but now considered to be the acacia vera. See *Arabic gum*.

MIMOSA SENEGAL. The systematic name of the tree from which the gum senegal exudes.

Mindererius spirit. See *Liquor ammoniac acetatis*.

MINERALIA. See *Minerale*.

MINERALOGY. That part of natural history which relates to minerals.

Mineral poisons. See *Poisons*.

MINERAL WATERS. *Aqua minerala*, *Aqua medicinales*. Waters holding minerals in solution are called *mineral waters*. But as all water, in a mineral state, is impregnated, either more or less, with some mineral substances, the name *mineral waters* should be confined to such waters as are sufficiently impregnated with mineral matters to produce some sensible effects on the animal economy, and either to cure or prevent some of the diseases to which the human body is liable. On this account, these waters might be with much more propriety, be called *medicinal waters*, were not the name by which they are commonly known too firmly established by long use.

The mineral waters which are the most esteemed, and consequently the most resorted to for the cure of diseases, are those of

- | | |
|-------------|----------------|
| 1. Aix. | 5. Buxton. |
| 2. Berege. | 6. Borset. |
| 3. Bath. | 7. Cheltenham. |
| 4. Bristol, | 8. Carlsbad, |

- | | |
|-----------------|-----------------------|
| 9. Epsom. | 17. Scarborough. |
| 10. Harrowgate. | 18. Spa. |
| 11. Hartfell. | 19. Sedlitz. |
| 12. Holywell. | 20. Sea-water. |
| 13. Malvern. | 21. Seltzer. |
| 14. Matlock. | 22. Tunbridge. |
| 15. Moffat. | 23. Vichy, and others |
| 16. Pyrmont. | of less note. |

For the properties and virtues of these consult their respective heads.

Fourcroy divides all mineral and medicinal waters into nine orders, viz.

1. Cold alcidulous waters.
2. Hot or thermal acidulous waters.
3. Sulphuric saline waters.
4. Muriatic saline waters,
5. Simple sulphureous waters.
6. Sulphurated gaseous waters.
7. Simple ferruginous waters.
8. Ferruginous and acidulous waters.
9. Sulphuric ferruginous waters.

Dr. Saunders arranges mineral waters into the following classes:

1. Simple cold.
2. ——— thermal.
3. ——— saline.
4. Highly carbonated alkaline.
5. Simple carbonated chalybeate.
6. Hot carbonated chalybeate.
7. Highly carbonated chalybeate.
8. Saline carbonated chalybeate.
9. Hot saline highly carbonated chalybeate.
10. Vitriolated chalybeate.
11. Cold, sulphureous.
12. Hot, alkaline, sulphureous.

In order to present the reader, under one point of view, with the most conspicuous features in the composition of the mineral waters of this and some other countries, the following Synoptical Table is subjoined, from Dr. Saunders' works on mineral waters.

The reader will please to observe, that under the head of *Neutral Purging Salts* are included the sulphats of soda and magnesia, and the muriats of lime, soda, and magnesia. The power which the earthy muriats may possess of acting on the intestinal canal is not quite ascertained, but, from their great solubility, and from analogy with salts, with similar component parts, we may conclude that this forms a principal part of their operation.

The reader will likewise observe, that where the spaces are left blank, it signifies that we are ignorant whether any of the substance at the head of the column is contained in the water; that the word *none* implies a certainty of the absence of that substance; and the term *uncertain* means that the substance is contained, but, that the quantity is not known.

A SYNOPTICAL TABLE, shewing the Composition of MINERAL WATERS.

CLASS.	NAME.	Highest Temperature. Fahrenheit.	Contained in an English Wine Pint of 28.875 Cubic Inches.									
			Azotic Gas. Cubic Inches	Carbonic Acid Gas. Cubic Inches	Sulphurated Hydrogen. Cubic Inches	Carbonated Soda. Grains.	Neutral Purging Salts. Grains.	Selenite and Earthy Carbonates. Grains.	Oxyd of Iron. Grains.			
Simpler cold	Malvern			uncertain	none	none	uncertain	uncertain	none			
	Holywell				none	none	uncertain	uncertain	none			
Simpler thermal	Bristol	74°	uncertain	3.75	none	none	2.81	3.16	none			
	Matlock	66°		uncertain	none	none	uncertain	uncertain	none			
Simple saline	Buxton	82°	0.474	uncertain	none	none	0.25	1.625	none			
	Sedlitz			1.	none	none	185.6	8.68	none			
	Epsom				none	none	40.7	8.7	none			
	Sea				none	none	237.5	6.	none			
Highly carbonated alkaline	Seltzer			17.	none	4.	17.5	8.	none			
Simple carbonated chalybeate	Tunbridge		0.675	1.325	none	none	0.344	0.156	0.125			
Hot carbonated chalybeate	Bath	116°	1.7	1.7	none	none	10.7	10.7	uncertain			
Highly carbonated chalybeate	Spa			12.79	none	1.47	4.632	1.47	0.58			
	Pyrmont			26.	none	none	7.13	23.075	0.56.			
Saline, carbonated chalybeate	Cheltenham		uncertain	5.687	uncertain	none	62.125	6.85	0.625			
	Scarborough			uncertain	none	none	20.	10.	uncertain			
Hot, saline, highly carbonated chalybeate	Vichy	120°?		uncertain	none	uncertain		uncertain	uncertain			
	Carlsbad	165°		uncertain	none	uncertain		uncertain	uncertain			
Vitriolated chalybeate	Hartfell			uncertain	none	uncertain		uncertain	uncertain			
	Harrowgate		0.875	1.	none	none	none	4.15	4.815*			
Cold sulphureous	Vichy		0.5	0.635	2.375	none	91.25	3.	none			
	Aix	143°		uncertain	1.25	none	4.5	none	none			
Hot, alkaline, sulphureous	Bonnet	132°		uncertain	uncertain	12.	5.	4.75	none			
	Berger	120°		uncertain	uncertain	uncertain	uncertain	uncertain	none			

* That is, 2.94 contained in the sulphat of iron, (this salt, when crystallized, containing 28 per cent of oxyd of iron, according to Kirwan,) and 1.875 additional of oxyd of iron.

Mr. Henry, in his epitome of chymistry, gives the following concise and accurate account for the analysis of mineral waters:

Water is never presented by nature in a state of complete purity. Even when collected as it descends in the form of rain, chymical tests detect in it foreign ingredients. And when it has been absorbed by the earth, has traversed its different strata, and is returned to us by springs, it is found to have acquired various impregnations. The readiest method of judging of the contents of natural waters is by applying what are termed tests, or reagents, *i. e.* substances which, on being added to a water, exhibits, by the phenomena they produce, the nature of the saline and other ingredients. For example, if, on adding an infusion of litmus to any water, its colour is changed to red, we infer that the water contains an uncombined acid; if this change ensue even after the water has been boiled, we judge that the acid is a fixed and not a volatile one: and if, on adding the muriate of barytes, a precipitate falls down, we safely conclude that the peculiar acid present in the water is either entirely or in part the sulphuric acid. Mr. Henry first enumerates the tests generally employed in examining mineral waters, and describes their application, and afterwards indicates by what particular tests the substances generally found in waters may be detected.

A. Infusion of Litmus. Syrup of Violets, &c.—As the infusion of litmus is apt to spoil by keeping, some solid litmus should be kept. The infusion is prepared by steeping this substance, first bruised in a mortar, and tied up in a thin rag, in distilled water, which extracts its blue colour. If the colour of the infusion tends too much to purple, it may be amended by a drop or two of pure ammonia; but of this no more should be added than what is barely sufficient, lest the delicacy of the test should be impaired. The syrup of violets is not easily obtained pure. The genuine syrup may be distinguished from the spurious by a solution of corrosive sublimate, which changes the former to green, while it reddens the latter. When it can be procured genuine, it is an excellent test of acids, and may be employed in the same manner as the infusion of litmus. Paper stained with the juice of the marsh violet, or with that of radishes, answers a similar purpose. In staining paper for the purpose of a test, it must be used unsized; or, if sized, it must previously be washed with warm water; because the alum which enters into the composition of the size will otherwise change the vegetable colour to a red.

Infusion of litmus is a test of most uncombined acids.

If the infusion reddens the unboiled but not the boiled water under examination, or if the red colour occasioned by adding the infusion to a recent water return to blue on boiling, we may infer that the acid is a volatile one, and most probably the carbonic acid. Sulphuretted hydrogen gas, dissolved in water, also reddens litmus, but not after boiling. To ascertain whether the change be produced by carbonic acid, or sulphuretted hydrogen, when experiment shews that the reddening cause is volatile, add a little lime-water. This, if carbonic acid be present, will occasion a precipitate, which will dissolve with effervescence, on adding a little muriatic acid. Sulphuretted hydrogen may also be contained in the same water, which will be ascertained by the tests hereafter to be described.

Paper tinged with litmus is also reddened by the presence of carbonic acid, but regains its blue colour by drying. The mineral and fixed acids redden it permanently. That these acids, however, may produce their effect, it is necessary that they should be present in a sufficient proportion.

Infusion of litmus reddened by vinegar—Spirituuous tincture of Brazil wood—Tincture of turmeric, and paper stained with each of these three substances—Syrup of violets. All these different tests have one and the same object.

1. Infusion of litmus reddened by vinegar, or litmus paper reddened by vinegar, has its blue colour restored by alkalis and pure earths, and by carbonated alkalis and earths.

2. Turmeric paper and tincture are changed to a reddish brown by alkalis, whether pure or carbonated, and by pure earths; but not by carbonated earths.

3. The red infusion of Brazil wood, and paper stained with it, become blue by alkalis and earths, and even by the latter, when dissolved by an excess of carbonic acid. In the last mentioned case, however, the change will either cease to appear, or be much less remarkable, when the water has been boiled.

4. Syrup of violets, when pure, is by the same causes turned green, as also paper stained with the juices of violets, or radishes.

B. Tincture of Galls.

Tincture of galls is the test generally employed for discovering iron, with all the combinations of which it produces a black tinge, more or less intense, according to the quantity of iron. The iron, however, in order to be detected by this test, must be in the state of red oxyd, or, if oxydated in

a less degree, its effects will not be apparent, unless after standing some time in contact with air. By applying this test before and after evaporation, or boiling, we may know whether the iron be held in solution by carbonic acid or a fixed acid ; for,

1. If it produce its effects before the application of heat, and not afterwards, carbonic acid is the solvent.

2. If after, as well as before, a mineral acid is the solvent.

3. If, by the boiling, a yellowish powder be precipitated, and yet galls continue to strike the water black afterwards, the iron, as often happens, is dissolved both by carbonic acid and a fixed acid. A neat mode of applying the gall test was used by M. Klaproth, in his analysis of the Carlsbad water. A slice of the gall-nut was suspended by a silken thread, in a large bottle of the recent water; and so small was the quantity of iron, that it could only be discovered in water fresh from the spring.

C. Sulphuric Acid.

1. Sulphuric acid discovers, by a slight effervescence, the presence of carbonic acid, whether uncombined or united with alkalis, or earths.

2. If lime be present, whether pure or uncombined, the addition of sulphuric acid occasions, after a few days, a white precipitate.

3. Barytes is precipitated instantly in the form of a white powder.

4. Nitrous and muriatic salts, on adding sulphuric acid and applying heat, are decomposed; and if a stopper, moistened with pure ammonia, be held over the vessel, white clouds appear. For distinguishing whether nitric or muriatic acid be present, rules will be given hereafter.

Nitric and Nitrous Acids.

These acids, if they occasion effervescence, give the same indications as the sulphuric. The nitrous acid has been recommended, as a test distinguishing between hepatic waters that contain sulphuretted hydrogen gas. In the former case, a precipitate ensues on adding nitrous acid, and a very fetid smell arises; in the latter, a slight cloudiness only appears, and the smell of the water becomes less disagreeable.

D. Oxalic Acid and Oxalates.

This acid is a most delicate test of lime, which it separates from all its combinations.

1. If a water, which is precipitated by oxalic acid, becomes milky on adding a watery solution of carbonic acid gas, or by blowing air through it by means of a quill, or glass tube, we may infer that pure lime

(or barytes, which has never yet been found pure in water) is present.

2. If the oxalic acid occasion a precipitate before, but not after boiling, the lime is dissolved by an excess of carbonic acid.

3. If, after boiling by a fixed acid, a considerable excess of any of the mineral acids, however, prevents the oxalic acid from occasioning a precipitate, even though lime be present; because some acids decompose the oxalic, and others, dissolving the oxalate of lime, prevent it from appearing.

The oxalate of ammonia, or of potash, (which may easily be formed by saturating their respective carbonates with a solution of oxalic acid) are not liable to the above objections; and are preferable, as reagents, to the uncombined acids. Yet even these oxalates fail to detect lime when supersaturated with muriatic or nitric acids; and if such an excess be present, it must be saturated before adding the test with pure ammonia. Fluat of ammonia is the best test of lime. It is made by adding carbonate of ammonia to diluted fluoric acid.

E. Pure Alkalis and Carbonated Alkalis.

1. The pure fixed alkalis precipitate all earths and metals, whether dissolved by volatile or fixed menstrua, but only in certain states of dilution: for example, sulphate of alumina may be present in water, in the proportion of 4 grains to 500, without being discovered by pure fixed alkalis. As the alkalis precipitate so many substances, it is evident they cannot afford any precise information when employed as reagents. From the colour of the precipitate, as it approaches to pure white, or recedes from it, an experienced eye will judge that the precipitated earth contains less or more of the metallic admixture.

2. Pure fixed alkalis decompose all salts with basis of ammonia, which becomes evident by its smell, and also by the white fumes it exhibits when a stopper is brought near it, moistened with muriatic acid.

3. Carbonates of potash and soda have similar effects.

4. Pure ammonia precipitates all earthy and metallic salts. Besides this property, it also imparts a deep blue colour to any liquid that contains copper in a state of solution.

Carbonate of ammonia has the same properties, except that it does not precipitate magnesia from its combinations. Hence, to ascertain whether this earth be present in any solution, add the carbonate of ammonia till no further precipitation ensues, filter the liquor, and then add pure ammonia. If any precipitation now occurs, we may infer the presence of magnesia.

F. Lime-Water.

1. Lime-water is applied for the purposes of a test, chiefly for detecting carbonic acid. Let any liquor, supposed to contain this acid, be mixed with an equal bulk of this acid. If carbonic acid be present, either free or combined, a precipitate will immediately appear, which, on adding a few drops of muriatic acid, will immediately dissolve with effervescence.

2. Lime-water will immediately shew the presence of corrosive sublimate, by a brick-dust coloured sediment. If arsenic be present in any liquid, lime-water, when added, will occasion a precipitate, consisting of lime and arsenic, which is very difficultly soluble in water. This precipitate, when mixed up with oil, and laid on the hot coals, yields the well-known garlic smell of arsenic.

G. Pure Barytes, and its Solution in Water.

1. A solution of pure barytes is even more effectual than lime-water, in detecting the presence of carbonic acid, and is much more portable and convenient; since from the crystals of this earth, the solution may at any time be prepared. In discovering fixed air, the solution of barytes is used similarly to lime-water; and, if this acid be present, gives, in like manner, a precipitate soluble with effervescence in muriatic acid.

Pure strontites has similar virtues as a test.

H. Metals.

1. Of the metals, silver and mercury are tests of the presence of sulphurets, and of sulphuretted hydrogen gas. If a little quicksilver be put into a bottle, containing water impregnated with either of these substances, its surface soon acquires a black film, and, on shaking, a blackish powder separates from it. Silver is immediately tarnished from the same cause.

2. The metals also may be used as tests of each other, on the principle of elective affinity. Thus, for example, a polished iron plate, immersed in a solution of sulphate of copper, soon acquires a coat of this metal, and the same in other similar examples.

I. Sulphate of Iron.

This is the only one of the sulphates, except that of silver, applicable to the purposes of a test. When used in this view, it is generally employed to ascertain the presence of oxygenous gas, of which a natural water may contain a small quantity.

A water suspected to contain this gas, may be mixed with a little recently dissolved sulphate of iron, and kept corked up. If an oxyd of iron be precipitated in the course of a few days, the water may be inferred to contain oxygenous gas.

Sulphate, Nitrate, and Acetate of Silver.

These solutions are, in some measure, applicable to the same purpose.

1. They are peculiarly adapted to the discovery of muriatic acid and muriates. For the silver, quitting the nitric acid, combines with the muriatic, and forms a flaky precipitate, which at first is white, but, on exposure to the sun's light, acquires a bluish colour. This precipitate, Dr. Black states to contain, in 1000 parts, as much muriatic acid as would form 425 parts and a half of chrystallised muriate of soda, which estimate scarcely differs at all from that of Klaproth. A precipitation, however, may arise from other causes, which it may be proper to state.

2. The solutions of silver in acids are precipitated by carbonated alkalis and earths. The agency of these may be prevented by previously adding a few drops of the same acid in which the silver is dissolved.

3. The nitrate and acetate of silver are decomposed by the sulphuric and sulphurous acids; but this may be prevented by adding previously a few drops of nitrate or acetate of barytes, and after allowing the precipitate to subside, the clear liquor may be decanted, and the solution of silver added. Should a precipitation now take place, the presence of muriatic acid, or some one of its combinations, may be suspected. To obviate uncertainty, whether a precipitation be owing to sulphuric or muriatic acid, a solution of sulphate of silver may be employed, which is effected only by the latter acid.

4. The solutions of silver are precipitated by extractive matters; but in this case also the precipitate is discoloured, and is soluble in nitrous acid.

K. Nitrate and Acetate of Lead.

1. Acetate of lead, the most eligible of these two tests, is precipitated by sulphuric and muriatic acids; but as, of both these, we have much better indicators, it is not necessary to enlarge on its application to this purpose.

2. The acetate is also a test of sulphuretted hydrogen and of sulphurets of alkalis, which occasion a black precipitate; and if a paper on which characters are traced with a solution of acetate of lead, be held over a portion of water containing sulphuretted hydrogen, they are soon rendered visible.

3. The acetate of lead is employed in the discovery of uncombined boracic acid, a very rare ingredient of waters. To ascertain whether this be present, some cautions are necessary. The uncombined alkalis and earths (if any be suspected) must be saturated with acetic or acetous acid. The sulphates must be decomposed by acetate or nitrate of barytes, and the

muriates by acetite or nitrate of silver. The filtered liquor, if boric acid be contained in it, will give a precipitate soluble in nitric acid of the specific gravity of 1.3.

L. Nitrate of Mercury prepared with and without Heat.

This solution, differently prepared, is sometimes employed as a test. But, since other tests answer the same purposes more effectually, it is not absolutely necessary to have these tests.

M. Muriate, Nitrate, and Acetite of Barytes.

1. These solutions are all most delicate tests of sulphuric acid and of its combinations, with which they give a white precipitate, insoluble in dilute muriatic acid. They are decomposed, however, by carbonates of alkali; but the precipitate occasioned by these is soluble in dilute muriatic and nitric acid with effervescence, and may even be prevented by adding previously a few drops of the acid contained in the barytic salt.

One hundred grains of dry sulphate of barytes (according to Klaproth, p. 168,) contains about 45 one-fifth of sulphuric acid, of the specific gravity 1850, according to Clayfield, 33 of acid of a. g. 2240, according to Thenard, after calcination about 25. These estimates differ very considerably. From Klaproth's experiments it appears that 1000 grains of sulphate of barytes indicate 595 of desiccated sulphate of soda, or 1416 of the crystallized salt. The same chymist has shewn that 100 grains of sulphate of barytes are produced by the precipitation of 71 grains of sulphate of lime.

2. Phosphoric salts also occasion a precipitate which is soluble in muriatic acid without effervescence.

N. Prussiate of Potash and Lime.

Of these two, the prussiat of potash is the most eligible. When pure, it does not speedily assume a blue colour on the addition of acid, nor does it immediately precipitate muriated barytes. Prussiat of potash is a very sensible test of iron, with the solutions of which in acids it produces a Prussian blue precipitate, in consequence of a double elective affinity. To render its effect more certain, however, it may be proper to add, previously to any water suspected to contain iron, a little muriatic acid, with a view to the saturation of uncombined alkalis, or earths, which, if present, prevent the detection of any minute portions of iron.

1. If a water, after boiling and filtration, does not afford a blue precipitate on the addition of prussiat of potash, the solvent of the iron may be inferred to be a volatile one, and probably the carbonic acid.

2. Should the precipitation ensue in the boiled water, the solvent is a fixed acid,

the nature of which must be ascertained by other tests.

O. Solution of Soap in Alcohol.

This solution may be used to ascertain the comparative hardness of waters. With distilled water it may be mixed without producing any change; but, if added to a hard water, it produces a milkiness, more or less considerable as the water is less pure; and from the degree of milkiness an experienced eye will judge of its quality. The acids, alkalis, and all earthy and metallic salts, decompose soap, and occasion that property in water termed hardness.

Alcohol.

Alcohol, when mixed with any water in the proportion of about an equal bulk, precipitates all the salts which it is capable of dissolving.

P. Hydro-sulphuret of Ammonia.

This and other sulphurets, as well as water saturated with sulphurated hydrogen, may be employed in detecting lead and arsenic, with the former of which they give a black, and with the latter a yellowish precipitate. As lead and arsenic, however, are never found in natural waters, these tests are not required.

MINERALS. (*Mineralia*; from *mina*, a mine of metal.) All substances which do not possess organization, or are not produced by an organized body, belong to the class called minerals. Among this varied class of materials, which require the attention of the chymist and manufacturer, many are compounded of such principles and formed under such circumstances and situations in the earth, that it is difficult to distinguish them without having recourse to the test of experiment; several are formed with considerable regularity as to the proportion of their principles, their fracture, their colour, specific gravity, and figure of crystallization.

Mineral bodies which enter into the composition of the globe, are classed by mineralogists under four heads:—1. Earths. 2. Salts. 3. Inflammable fossils; and 4. Metals and their ores. Under the term earths are arranged stones and earths, which have no taste, and do not burn when heated with contact of air.

Under the second, salts, or those saline substances which melt in water and do not burn, they require, according to Mr. Kirwan, less than two hundred times their weight of water to dissolve them.

By inflammable fossils are to be understood all those minerals not soluble in water, and exhibiting a flame more or less evident when exposed to fire in contact with air.

The fourth class, or ores, are compound bodies. Nature has bestowed their proper metallic appearance on some sub-

stances, and when this is the case, or they are alloyed with other metals, or semi-metals, they are called native metals. But such as are distinguished, as they commonly are, in mines, in combination with some other unmetallic substances, are said to be mineralized. The substance that sets them in that state, is called the mineralizer, and the compound of both, an ore. For example, in the common ore of copper, this metal is found oxydated, and the oxyd combined with sulphur. The copper may be considered as mineralized with oxygen and sulphur, and the compound of the three bodies forms an ore of copper.

Mineral salts. See Salts.

MINIMUM. A minim. The sixtieth part of a drachm. An important change has been adopted in the last London Pharmacopœia, for the mensuration of liquids, and the division of the wine pint, to insure accuracy in the measurement of qualities of liquids below one drachm. The number of drops contained in one drachm has been assumed to be sixty; and taking water as a standard, this number, though by no means accurate, would still be sufficient for ordinary purposes; but when other liquids of less specific gravity are used, a much larger number is required to fill the same measure, as of proof spirit, 140 drops are required to equal the bulk of 60 of water, dropped from the same vessel. If, therefore, in the composition of medicines, measures suited to the standard of water were used occasionally only, and it was generally assumed that sixty drops were equal to one fluid-drachm, and one fluid-drachm was substituted for sixty drops prescribed, twice the dose intended would be given. There are further objections to the use of drops; that their bulk is influenced by the quantity of liquid contained, in the bottle from which they fall, by the thickness of the lip, and even by the inequalities on the surface of the lip of the same bottle; that volatile liquids, to which this mode is most commonly applied, are thus exposed with extensive surfaces, and their evaporation promoted, and on all these accounts the adoption of some decisive convenient and uniform substitute became necessary. The subdivision of the wine pint has therefore been extended to the sixtieth part of the fluid-drachm, which is termed minim; and glass measures, expressive of such subdivision have been adapted by the college.

MINUM. Red lead. *See Lead.*

MINUM GRACORUM. Native cinnabar.

Mint, common. *See Mentha sativa.*

Mint, pepper. *See Mentha piperita.*

Mint, water. *See Menta aquatica.*

Miscarriage. *See Abortion.*

MISERERE MI. (*Hic compassio* on

me; so called from its unhappy torments.) The liliac passion.

Mislaw. *See Musa paradisica.*

MISOCYRACUS. Thus some were called who professed themselves enemies to the chymists, and their enthusiastic conceits.

MISFICKLE. A white, brilliant, granulated iron ore, composed of iron in combination with arsenic.

Misletoc. *See Viscum.*

MISTURA. A mixture. A fluid composed of two or more ingredients. It is mostly contracted in prescriptions thus, *mist*, e. g.—*f. mist.* which means, let it be made into a mixture.

MISTURA CAMPHORÆ. Camphire mixture. "Take of camphor, half a drachm; rectified spirit, ten minims; water, a pint. First rub the camphor with the spirit, then with the water gradually added, and strain the liquor." A very elegant preparation of camphire, for delicate stomachs, and those who cannot bear it in substance, as an antispasmodic and nervine. There is a great loss of camphire in making it as directed by the pharmacopœia. Water can only take up a certain quantity. For its virtues, *see Camphora.*

MISTURA CORNU USTI. "Take of hartshorn, burnt and prepared, two ounces; acacia gum, an ounce; water, three pints." Boil down to two pints, constantly stirring, and strain. For its virtues, *see Cornu.*

MISTURA CRETÆ. Chalk mixture. "Take of prepared chalk, half an ounce; refined sugar, three drachms; gum arabic, powdered, half an ounce." Mix. A very useful and pleasant form of administering chalk as an adstringent and antacid. It is particularly calculated for children, in whom it allays the many deranged actions of the primæ viæ, which are produced by acidities. Dose, one ounce to three, frequently. *See Creta* and *Carbonas calcis.*

MISTURA FERRI COMPOSITA. "Take of myrrh, powdered, a drachm; subcarbonate of potash, twenty-five grains; rose-water, seven ounces and a half; sulphate of iron, powdered, a scruple; spirit of nutmeg, an ounce; refined sugar, a drachm. Rub together the myrrh, the subcarbonate of potash and sugar; and, during the trituration, add gradually, first, the rose water and spirit of nutmegs, and last, the sulphate of iron. Pour the mixture immediately into a proper glass bottle, and stop it close." This preparation is the celebrated mixture of Dr. Griffiths. A chymical decomposition is effected in forming this mixture, a subcarbonate of iron is formed, and a sulphate of potash.

MISTURA GUAIACI. "Take of guaiacum gum resin, a drachm and a half; refined sugar, two drachms; mucilage of acacia gum, two fluid ounces; cinnamon water,

eight fluid ounces. Rub the guaiacum with the sugar, then with the mucilage; and, when they are mixed, pour on the cinnamon water gradually." For its virtues see *Guaiacum*.

MISTURA MOSCHI. "Take of musk, acacia gum, powdered, refined sugar, of each a drachm; rose-water, six fluid ounces." Rub the musk first with the sugar, then with the gum, and add the rose-water by degrees. An excellent diaphoretic and antispasmodic. It is by far the best way of administering musk; when boluses cannot be swallowed. Dose, one ounce to three, frequently.

Mithridate mustard. See *Thlaspi campestre*.

MITHRIDATIUM. The electuary called *Mithridate*, from Mithridates, king of Pontus and Bythinia, who experienced the virtues of the simples separately, afterwards combined them; but then the composition consisted of but few ingredients, viz. twenty leaves of rue, two walnuts, two figs, and a little salt: of this he took a dose every morning, to guard himself against the effects of poison.

MITRAL VALVES. *Valvule mitrales.* The valves of the left ventricle of the heart are so called from their resemblance to a mitre.

MIVA. An antient term for the form of a medicine, not unlike a thick syrup, now called *Marmalade*.

MIXTURE. 1. See *Mistura*.

2. Chymical mixture should be distinguished from the chymical solution; in the former, the aggregate particles can again be separated by mechanical means, and the proportion of the different particles determined; but, in solution, no mechanical power whatsoever can separate them.

MOCHLIA. (From *μοχλος*, a lever.) A reduction of the bones from an unnatural to a natural situation.

MOCA LICA. (From *μοχλω*, to move.) Violent purges.

MODIOLUS. (Dim. of *modus*, a measure.) The nucleus, as it were, of the cochlea of the ear is so termed. It ascends from the basis of the cochlea to the apex.

Mofette. See *Nitrogen*.

MOFFAT WATER. A cold sulphureous water, of a very simple composition. Moffat, a village situated about fifty-six miles south-west of Edinburgh, affords this mineral water; when first drawn, it appears rather milky and bluish; the smell is exactly similar to that of Harrogate, the smell is sulphureous and saline, without any thing bitter. It sparkles somewhat on being poured from one glass to another.

According to Dr. Garnett's analysis, a wine gallon of Moffat water contains thirty-

six grains of muriate of soda, five cubic inches of carbonic acid gas, four of azotic gas, and ten of sulphurated hydrogen, making altogether nineteen cubic inches of gas. Moffat water is, therefore, very simple in its composition, and hence it produces effects somewhat similar to those of Harrogate. It is, perhaps, on this account also that it so soon loses the hepatic gas, on which depends the greater part of its medicinal power. The only sensible effect of this water is that of increasing the flow of urine; when it purges, it appears rather to take place from the excessive dose than from its mineral ingredients. This water appears to be useful chiefly in cutaneous eruptions, and as an external application in an increased temperature, scrofula in its early stage appears to be alleviated, it is also used as an external application to irritable ulcers, and is recommended in dyspepsia, and where there is inaction of the alimentary canal.

MOGILALIA. (From *μωγ*, difficulty, and *αλω*, to speak. A difficulty of speech.

MOLA. (Heb.) 1. The knee-pan; so named because it is shaped like a mill-stone. 2. A mole, or shapeless mass of flesh in the uterus. See *Mole*.

MOLAR GLANDS. *Glandulae molares.* Two salival glands situated on each side of the mouth, between the masseter and buccinator muscles, the excretory ducts of which open near the last dens molaris.

MOLARIS. (From *molaris*, a grind-stone; because they grind the food.) A double-tooth. See *Teeth*.

Molasses. See *Treacle*.

MOLDAVICA. *Melissa Turca.* Turkey balsam. Canary balsam. Balsam of Gilead. This plant, *Dracocephalum moldavica*; *floribus verticillatis, bracteis lanceolatis, serratis capillatis* of Linnæus, affords a fragrant essential oil, by distillation, known in Germany by the name of *oleum syria*. The whole herb abounds with an aromatic smell, and an agreeable taste, joined with an aromatic flavour; it is recommended to give tone to the stomach and nervous system.

MOLE. *Mola.* By this term authors have intended to describe different productions of, or excretions from the uterus.

By some it has been used to signify every kind of fleshy substance, particularly those which are properly called polypi; by others, those only which are the consequence of imperfect conception, or when the ovum is in a morbid or decayed state; and by many, which is the most popular opinion, every coagulum of blood which continues long enough in the uterus to assume a form, and to have only the fibrous part, as it has been called, remaining; is denominated a mole.

There is surely much impropriety, says Dr. Denman, in including, under one general name, appearances so contrary, and substances so different.

For an account of the first kind, see *Polypus*.

Of the second kind, which has been defined as an *ovum deformis*, as it is the consequence of conception, it might more justly be arranged under the class of monsters; for though it has the appearance of a shapeless mass of flesh, if examined carefully with a knife, various parts of a child may be discovered, lying together in apparent confusion, but in actual regularity. The pedicle also by which it is connected to the uterus, is not of a fleshy texture, like that of the polypus, but has a regular series of vessels like the umbilical cord, and there is likewise a placenta and membranes containing water. The symptoms attending the formation, growth, and expulsion of this apparently confused mass from the uterus, corresponding with those of a well-formed child.

With respect to the third opinion of a mole, an incision into its substance will discover its true nature; for, although the external surface appears at the first view to be organized flesh, the internal part is composed merely of coagulated blood. As substances of this kind, which mostly occur after delivery, would always be expelled from the action of the uterus, there seems to be no reason for a particular inquiry, if popular opinion had not annexed the idea of mischief to them, and attributed their formation or continuance in the uterus to the negligence or misconduct of the practitioner. Hence the persuasion arose of the necessity of extracting all the coagula of blood out of the uterus, immediately after the expulsion of the placenta, or of giving medicines to force them away; but abundant experience hath proved, that the retention of such coagula is not, under any circumstances, productive of danger, and that they are most safely expelled by the action of the uterus, though at very different periods after their formation.

MOLLE. Indian mastich.

MOLLITIES OSSIUM. (*Mollities*, from *mollis*, soft.) A disease of the bones, wherein they can be bent without fracturing them, in consequence either of the inordinate absorption of the phosphate of lime, from their natural solidity is derived, or else of this matter not being duly secreted and deposited in their fabric. In rickets, the bones only yield and become distorted by slow degrees, and retain their natural inflexibility; but in the present disease they may be at once bent in any direction. The *mollities ossium* is rare, and its causes not well understood. All the cases of *mollities ossium* yet on record have proved

fatal, and no means of cure are yet known. On dissection of those who have died, all the bones, except the teeth, have been found unusually soft, so that scarce any of them could resist the knife, the periosteum has been found thicker than usual, and the bones have been found to contain a great quantity of oily matter and little earth.

MOLLITIES UNGUINUM. A preternatural softness of the nails: it often accompanies chlorosis.

MOLLIFICATION. A barbarous term of a palsy of the muscles in any particular part.

MOLUCCENSE LIGNUM. See *Lignum puvane*.

MOLYBDAT. *Molybda*. A salt formed by the union of the molybdic acid with different bases; thus, *molybdat of alumina*, *molybdat of antimony*, &c.

MOLYBDENA. (From *μολυβδης*, lead.) *Molybdis*. A metal which exists mineralized by sulphur in the ore called *sulphuret of Molybdena*. This ore, which is very scarce, is so similar in several of its properties to plumbago that they were long considered as varieties of the same substance. It is of a light lead-grey colour, its surface is smooth, and feels unctuous, its texture is lamellated, it soils the fingers, and marks paper bluish-black, or silver-grey. It may be cut with a knife. It is generally found in compact masses; seldom in particles, or crystallized. It is met with in Sweden, Spain, Saxony, Siberia, and Iceland. Scheele shewed that a peculiar metallic acid might be obtained from it; and later chymists have succeeded in reducing this acid to the metallic state. We are indebted to Mr. Hatchett, for a full and accurate analysis of this ore.

The native *sulphuret of molybdena* is the only ore hitherto known which contains this metal.

Properties of Molybdena.—*Molybdena* is either in an agglutinated blackish friable mass, having little metallic brilliancy, or in a black powder. The mass slightly united, shews, by a magnifying glass, small round brilliant grains. Its weight is from 6600 to 7500. It is one of the most infusible of the metals. It is capable of combining with a number of metals by fusion. It forms with sulphur an artificial sulphuret of molybdena analogous to its ore. It unites also to phosphorous. The affinity of molybdena for oxygen is very feeble, according to Mr. Hatchett. The alkalis have no action on molybdena in the moist way, but it enters readily into fusion with potash and soda. It is oxydable by boiling sulphuric acid, and acidifiable by the nitric acid. Muriatic acid does not act upon it. It is capable of existing in not less than four different degrees of oxygenation.

Method of obtaining Molybdena.—To ob-

tain molybdena is a task of the utmost difficulty. Few chymists have succeeded in producing this metal, on account of its great infusibility. The method recommended in general is the following:—Molabdic acid is to be formed into a paste with oil, dried at the fire, and then exposed to a violent heat in a crucible lined with charcoal. By this means the oxyd becomes decomposed; a black agglutinated substance is obtained, very brittle under the finger, and having a metallic brilliancy. This is the metal called molybdena.

MOLYBDITIS. See *Molybdena*.

MOLYBDOΣ. (*Ὀρί μολυβδός*, from its gravity.) Lead.

MOLYEA. (Dim. of *μολύ*, moly.) Garlic, whose head, like moly, is not divided into cloves.

MOMISCS. (From *μυμός*, a blemish.) That part of the teeth which is next the gums, and which is usually covered with a foul tartareous crust.

MOMORDICA. The name of a genus of plants in the Linnæan system. Class, *Monocotyledon*. Order, *Syngenesia*.

MOMORDICA ELLATERIUM. (*Momordica*; from *mordeo*, to bite; from its sharp taste.) The systematic name of the squirting cucumber. See *Elatarium*.

MONARDA PISTULOSA. The systematic name of the purple monarda. The leaves of this plant have a fragrant smell, and an aromatic and somewhat bitter taste, possessing nervine, stomachic, and deobstruent virtues. An infusion is recommended in the cure of intermittent fevers.

MONELLI. A species of *Anagallis*.

Moneywort. See *Nummularia*.

Monk's rhubarb. See *Rhabarbarum monachorum*.

Monkshead. See *Anthora*.

MONOCULUM. (From *μῦος*, single, and *oculus*, the eye.) A name given to the cæcum, or blind gut, by Paracelsus, because it is perforated only at one end.

MONOCULUS. (From *μῦος*, one, and *oculus*, an eye.) *Monopia*. A very uncommon species of monstrosity, in which there is but one eye, and that mostly above the root of the nose.

MONOHEMERA. (From *μῦος*, single, and *ἡμέρα*, a day.) A disease of one day's continuance.

MONOMACHON. The *intestinum cæcum*.

MONOPEDIA. (From *μῦος*, single, and *πῦσις*, to compress. A pain in only one side of the head.

ΜΟΝΟΦΙΑ. (From *μῦος*, single, and *ὤψ*, the eye.) See *Monoculus*.

ΜΟΝΟΧΗΝΙΣ. (From *μῦος*, one, and *ὄρχη*, a testicle.) An epithet for a person that has but one testicle.

MONS. A mount, or hill.

MONS VENERIS. The triangular

eminence immediately over the os pubis of women, that is covered with hair.

MONSTER. *Luus nature.* Dr. Denman divides monsters into, 1st, Monsters from redundancy or multiplicity of parts; 2d, Monsters from deficiency or want of parts; 3d, Monsters from confusion of parts.

To these might perhaps be added, without impropriety, another kind, in which there is neither redundancy, nor deficiency, nor confusion of parts, but an error of place, as in transposition of the viscera. But children born with diseases, as the hydrocephalus, or their effects, as in some cases of blindness, from previous inflammation, cannot be properly considered as monsters, though they are often so denominated.

Of the first order there may be two kinds, redundancy or multiplicity of natural parts, as of two heads and one body, of one head and two bodies, an increased number of limbs, as legs, arms, fingers, and toes; or excrescences or additions to parts of no certain form, as those upon the head and other parts of the body, not surprising that we should be ignorant also of the manner in which monsters or irregular births are generated or produced; though it is probable that the laws by which these are governed are as regular, both as to cause and effect, as in common or natural productions. Formerly, and indeed till within these few years, it was a generally-received opinion, that monsters were not primordial or aboriginal, but that they were caused subsequently, by the power of the imagination of the mother, transferring the imperfection of some external object, or the mark of something for which she longed, with which she was not indulged, to the child of which she was pregnant; or by some accident which happened to her during her pregnancy. Such opinions, it is reasonable to think, were permitted to pass current, in order to protect pregnant women from all hazardous and disagreeable occupations, to screen them from severe labour, and to procure for them a greater share of indulgence and tenderness than could be granted to them in the common occurrences of life. The laws and customs of every civilized nation have, in some degree, established a persuasion that there was something sacred in the person of a pregnant woman: and this may be right in several points of view; but these go a little way towards justifying the opinion of monsters being caused by the imagination of the mother. The opinion has been disproved by common observation, and by philosophy, not perhaps by positive proofs, but by many strong negative facts; as the improbability of any child being born perfect, had such a power

existed; the freedom of children from any blemish, their mothers being in situations most exposed to objects likely to produce them; the ignorance of the mother of any thing being wrong in the child, till, from information of the fact, she begins to recollect every accident which happened during her pregnancy, and assigns the worst, or the most plausible, as the cause; the organization and colour of these adventitious substances; the frequent occurrence of monsters in the brute creation, in which the power of the imagination cannot be great; and the analogous appearances in the vegetable system, where it does not exist in any degree. Judging, however, from appearances, accidents may perhaps be allowed to have considerable influence in the production of monsters of some kinds, either by actual injury upon part, or by suppressing or deranging the principle of growth, because, when an arm, for instance, is wanting, the rudiments of the deficient parts may generally be discovered.

MORBILLI. (Dim. of *morbus*, a disease.) See *Rubeola*.

MORBUS ARQUATUS. The jaundice.

MORBUS ATTONITUS. The epilepsy.

MORBUS COXARIUS. See *Arthroponia*.

MORBUS GALLICUS. The venereal disease.

MORBUS HERCULEUS. The epilepsy.

MORBUS INFANTILIS. The epilepsy.

MORBUS INDICUS. The Indian disease, the venereal disease.

MORBUS MAGNUS. The epilepsy.

MORBUS NIGER. The black disease. So Hippocrates named it, and thus described it. This disorder is known by vomiting a concrete blood of a blackish red colour, and mixed with a large quantity of insipid, acid, or viscid phlegm. This evacuation is generally preceded by a pungent tensive pain, in both the hypochondria; and the appearance of the disease is attended with anxiety, a compressive pain in the præcordia, and fainting, which last is more frequent and violent, when the blood which is evacuated is fetid and corrupt. The stomach and the spleen are the principal, if not the proper seat of this disease.

MORBUS REGIUS. The jaundice.

MORBUS SACER. The epilepsy.

MOREL. *Phallus esculentus* of Linnæus. It grows on moist banks and wet pasture, and springs up in May. It is used in the same manner as the truffle, for gravies and stewed dishes, but gives an inferior flavour.

MORETUS. (From *morum*, the mulberry.) A decoction of mulberries.

MORIA. (From *pauci*, foolish.) Idiocy. Fatuity.

MORO. (From *morom*, a mulberry.) A small abscess resembling a mulberry.

MOROSIS. (From *μωρος*, folly.) See *Amentia*.

MORPHEA ALBA. (From *μωρη*, form.) A species of cutaneous leprosy. See *Alphus*.

MORSELLUS. *Morsellus*. A lozenge.

MORSELLI. An ancient name for those forms of medicines which were to be chewed in the mouth, as a lozenge, the word signifying a little mouthful.

MORTUS DIABOLI. The frimbrix of the Fallopian tubes.

MORTA. See *Pemphigus*.

MORTARIOLUM. (Dim. of *mortarium*, a mortar.) In chymistry, it is a sort of mould for making cupels with, also a little mortar. In anatomy, it is the sockets of the teeth.

MORTIFICATION. (From *mors*, death, and *facio*, to become.) *Mortificatio.* *Gangrena.* *Sphacelus*. The loss of vitality of a part of the body. Surgeons divide mortification into two species, the one preceded by inflammation, the other without it. In inflammations that are to terminate in mortification, there is a diminution of power joined to an increased action; this becomes a cause of mortification, by destroying the balance of power and action, which ought to exist in every part. There are, however, cases of mortification that do not arise wholly from that as a cause; of this kind are the carbuncle and the slough, formed in the small-pox pustule. Healthy phlegmonous inflammation seldom ends in mortification, though it does so when very vehement and extensive. Erysipelatous inflammation is observed most frequently to terminate in gangrene; and whenever phlegmon is in any degree conjoined with an erysipelatous affection, which it not unfrequently is, it seems thereby to acquire the same tendency, being more difficult to bring to resolution, or suppuration, than the true phlegmon, and more apt to run into a mortified state.

Causes which impede the circulation of the part affected, will occasion mortification, as is exemplified in strangulated hernia, tied polypi, or a limb being deprived of circulation from a dislocated joint.

Preventing the entrance of arterial blood into a limb, is also another cause. Paralysis, conjoined with pressure, old age, and ossification of the arteries, may produce mortification; also cold, with the sudden application of warmth, and likewise excessive heat applied to a part.

The symptoms of mortification that take place after inflammation are various, but generally as follows:—the pain and sympathetic fever suddenly diminish, the part affected becomes soft, and of a livid colour, losing at the same time more or less of its sensibility.

When any part of the body loses all motion, sensibility, and natural heat, and becomes of a brown livid or black colour, it is said to be affected with sphacelus. When the part becomes a cold, black, fibrous, senseless substance, it is termed a slough. As long as any sensibility, motion, and warmth continue, the state of the disorder is said to be gangrene. This last term is synonymous with mortification.

When gangrene takes place, the patient is usually troubled with a kind of hiccough: the constitution always suffers an immediate dejection, the countenance assumes a wild cadaverous look, the pulse becomes small, rapid, and sometimes irregular; cold perspirations come on, and the patient is often affected with diarrhoea and delirium.

MORUM. The mulberry. The tree that affords this fruit is the *Morus nigra*; *foliis cordatis scabris* of Linnæus. Mulberries abound with a deep violet coloured juice, which, in its general qualities, agrees with that of the fruits called *acido dulces*, allaying thirst, partly by refrigerating, and partly by exciting an excretion of mucus from the mouth and fauces, a similar effect is also produced in the stomach, where, by correcting putrescency, a powerful cause of thirst is removed. The London College directs a *sympus mori*, which is an agreeable vehicle for various medicines. The bark of the root of this tree is said, by André, to be useful in cases of tænia.

MORUS. The name of a genus of plants in the Linnæan system. Class, *Monœcia*. Order, *Tetandria*. The mulberry-tree.

MORUS NIGRA. The systematic name of the mulberry-tree. See *Morum*.

MOSCHATA NUX. See *Nux-moschata*.

MOSCHUS. (*Mosch.*, Arab.) Musk. An unctuous substance, contained in excretory follicles about the navel of the male animal called *moschus moschiferus* by Linnæus, a ruminating quadruped, resembling the antelope, from which it does not differ sufficiently to form a particular genus; the strong and permanent smell of which is peculiar to it. It is contained in a bag placed near the umbilical region. The best musk is brought from Tonquin, in China; an inferior sort from Agria and Bengal, and a still worse from Russia.

It is slightly unctuous, of a black colour, having a strong durable smell and a bitter taste. It yields part of its active matter to water, by infusion; by distillation the water is impregnated with its flavour, alcohol dissolves it, its impurities excepted. Chewed, and rubbed with a knife on paper, it looks bright, yellowish, smooth, and free from grittiness. Laid on a red-hot iron, it catches flame and burns almost

entirely away, leaving only an exceeding small quantity of light greyish ashes. If any earthy substances have been mixed with the musk the impurities will discover them. The medicinal and chymical properties of musk and castor are very similar: the virtues of the former are generally believed to be more powerful, and hence musk is preferred in cases of imminent danger. It is prescribed as a powerful antispasmodic, in doses of three grains or upwards, even to half a drachm, in the greater number of spasmodic diseases, especially in hysteria and singultus, and also in diseases of debility. In typhus, it is employed to remove subulturn tendinum, and other symptoms of a spasmodic nature. In cholera it frequently stops vomiting, and, combined with ammonia, it is given to arrest the progress of gangrene. It is best given in the form of bolus. To children it is given in the form of enema, and is an efficacious remedy in the convulsions arising from dentition. It is also given in hydrophobia, and in some forms of mania.

MOSCHUS MOSCHIFERUS. The systematic name of the musk animal. See *Moschus*.

MOSQUITA. (From *mosquita*, a gnat, Span.) An itching eruption of the skin, produced in hot climates by the bite of gnats.

MOSSYLLUM. (*Μοσυλλον*.) The best cinnamon.

Mother of thyme. See *Serpyllum*.

Motherwort. See *Cardiaca*.

Motion, muscular. See *Muscular motion*.

Motion, peristaltic. See *Peristaltic motion*.

MOTORII OCULORUM. (*Nervi Motorii*; so called from their office.) The third pair of nerves of the brain. They arise from the crura cerebri, and are distributed on the muscles of the bulb of the eye.

Mould. See *Fontanella*.

Mountain parsley, black. See *Oreoselinum*.

Mouse-ear. See *Pilosella*.

MOUTH. *Os.* The cavity of the mouth is well known. The parts which constitute it are the common integuments, the lips, the muscles of the upper and under jaw, the palate, two alveolar arches, the gums, the tongue, the cheeks, and salivary glands. The bones of the mouth are the two superior maxillary, two palatine, the lower jaw, and thirty-two teeth. The arteries of the external parts of the mouth are branches of the infra-orbital, inferior alveolar, and fascial arteries. The veins empty themselves into the external jugulars. The nerves are branches from the fifth and seventh pair. The use of the mouth is for

mastication, speech, respiration, deglutition, suction, and taste.

MOXA JAPONICA. (Japanese.) *Artemisia Chinensis*. *Mugia patrae*. *Moxa*. Mugwort of China. A soft lanuginous substance, prepared in Japan, from the young leaves of a species of mugwort, by beating them when thoroughly dried, and rubbing them betwixt the hands, till only the fine fibres are left. Moxa is celebrated in the eastern countries for preventing and curing many disorders, by being burnt on the skin; a little cone of it laid upon the part, previously moistened, and set on fire on the top, burns down with a temperate and glowing heat, and produces a dark-coloured spot, the ulceration of which is promoted by putting a little garlic, and the ulcer is either healed up when the eschar separates, or kept running for a length of time, as different circumstances may require.

MUCILAGE. *Mucilago*. A solution of gum. See *Gum*.

MUCILAGINOUS EXTRACTS. Extracts that readily dissolve in water, scarcely at all in spirits of wine, and undergo spirituous fermentation.

MUCILAGO ACACIÆ. Mucilage of acacia. *Mucilago gummi arabici*. "Take of acacia gum, powdered, four ounces; boiling water, half a pint." Rub the gum with the water, until it incorporates into a mucilage. A demulcent preparation, more frequently used to combine medicines, than in any other form.

MUCILAGO AMYLÆ. Starch mucilage. "Take of starch, three drachms; water, a pint." Rub the starch, gradually adding the water to it; then boil until it incorporates into a mucilage. This preparation is mostly exhibited with opium, in the form of clysters, in diarrhœas and dysenteries, where the tenesmus arises from an abrasion of the mucus of the rectum.

MUCILAGO ARABICI GUMMI. See *Mucilago acaciæ*.

MUCILAGO SEMINIS CYDONIÆ. See *Decoctum cydoniæ*.

MUCILAGO TRAGACANTHÆ. Mucilage of tragacanth joined with syrup of mulberries, this forms a pleasant demulcent, and may be exhibited to children, who are fond of it. These two last mucilages are omitted in the last London Pharmacopœia, as possessing no superiority over the mucilage of acacia.

MUCOCARNEUS. In M. A. Severinus, it is an epithet for a tumour, and an abscess, which is partly fleshy and partly mucous.

MUCOUS GLANDS. *Glandule mucosæ*. Muciparous glands. Glands that secrete mucus, such as the glands of the Schneiderian membrane of the nose, the

glands of the fauces, œsophagus, stomach, intestines, bladder, urethra, &c.

MUCUS, ANIMAL. Animal mucus differs from that obtained from the vegetable kingdom, in not being soluble in water, swimming on its surface; nor capable of mixing oil with water, and being soluble in mineral acids, which vegetable mucus is not. The use of this substance is to lubricate and defend the parts upon which it is secreted, as the nose, œsophagus, stomach, intestines, urethra, vagina, &c. Mr. Everard Home, in his dissertation on the properties of pus, informs us of a curious and apparently decisive mode of distinguishing between pus and animal mucus. The property, he observes, which characterizes pus, and distinguishes it from most other substances, is, its being composed of globules, which are visible when viewed through a microscope; whereas animal mucus, and all chymical combinations of animal substances, appear in the microscope to be made up of flakes. This property was first noticed by the late Mr. J. Hunter.

MUCUS, VEGETABLE. See *Gum*.

Mugwort. See *Artemisia vulgaris*.

MULÆ. Pustules contracted either by heat or cold.

Mulberry. See *Morum*.

Mullein. See *Verbascum*.

MULSUM. *Musum*. *Mulæ*. *Hydromel*. Honey-water; though sometimes it signifies wine sweetened with honey.

MULTIFIDUS SPINÆ. (From *multus*, many, and *findo*, to divide.) *Transversæ spinalis lumborum*. *Musculus sacer*. *Semi-spinalis internus, sive transverso-spinalis dorsii*. *Semi-spinalis, sive transverso-spinalis collæ, pars interna* of Winslow. *Transversalis lumborum vulgo sacer*. *Transversalis dorsii*. *Transversalis collæ* of Douglas. *Lumbo dorsii spinal* of Dumas. The generality of anatomical writrs have unnecessarily multiplied the muscles of the spine, and hence their descriptions of these parts are confused, and difficult to be understood. Under the name of *multifidus spinæ*, Albinus has therefore very properly included those portions of muscular flesh, intermixed with tendinous fibres, which lie close to the posterior part of the spine, and which Douglas and Winslow have described as three distinct muscles, under the names of *transversales*, or *transverso-spinales*, of the loins, back, and neck. The multifidus spinæ arises tendinous and fleshy from the upper convex surface of the os sacrum, from the posterior adjoining part of the ilium, from the oblique and transverse processes of all the lumbar vertebrae, from the transverse processes of all the dorsal vertebrae, and from those of the cervical vertebrae, excepting the three first. From all these

origins the fibres of the muscles run in an oblique direction, and are inserted, by distinct tendons, into the spinous processes of all the vertebræ of the loins and back, and likewise into those of the six inferior vertebræ of the neck. When this muscle acts singly, it extends the back obliquely, or moves it to one side; when both muscles act, they extend the vertebræ backwards.

MULTIFORME OS. See *Ethmoid bone*.

MULTIPES. (From *multus*, many, and *pes*, a foot.) 1. The wood-louse. 2. The polypus. 3. Any animal having more than four feet.

Mumps. See *Cynanche*.

MUNDIFICATIVA. (From *mundus*, to cleanse.)

Mundificatoria. Medicines which purify and clean away foulness.

MUNDIFICANTIA. See *Mundificativa*.

MUNGOOS. *Radix serpentum.* This bitter root of the plant *Ophiorrhiza mungos* of Linnaeus is much esteemed in Java, Sumatra, &c. as preventing the effects which usually follow the bite of the *naja*, a venomous serpent, with which view it is eaten by them. It is also said to be exhibited medicinally in the cure of intestinal worms.

MURALIS. (From *murus*, a wall; so called because it grows upon walls.) Pellytory. See *Parietaria*.

MURARIA. (From *murus*, a wall; because it grows about walls.) A species of maiden hair.

MURIAS. A muriate, or salt, formed by the union of the muriatic acid with certain bases, as *muriate of ammonia*, &c.

MURIAS AMMONIACÆ. See *Ammonia muriata*, and *Sal ammoniac*.

MURIAS BARYTÆ. *Terra ponderosa soluta.* The muriate of barytes, or heavy earth, is a very acrid and poisonous preparation. In small doses it proves sudorific, diuretic, deobstruent, and alterative; in an over-dose, emetic, and violently purgative. The late Dr. Crawford found it very serviceable in all diseases connected with scrophula; and the Germans have employed it with great success in some diseases of the skin and viscera, and obstinate ulcers. The dose of the saturated solution in distilled water, is from five to fifteen drops for children, and from fifteen to twenty for adults.

MURIAS CALCIS. *Calx soluta.* *Sal ammoniacus fixus.* This preparation is exhibited with the same views as the muriate of barytes. It possesses deobstruent, diuretic, and cathartic virtues, and is much used by the celebrated Fourcroy against scrophula, and scrophulous diseases. Six, twelve, and twenty grains, are given to children three times a day, and a drachm to adults.

MURIAS FERRI. *Ferrum salitum.*

Olum martis per deliquitum. This preparation of iron is styptic and tonic, and may be given in chlorosis, intermittents, rachitis, &c.

MURIAS FERRI AMMONIACALIS. See *Ferrum ammoniatum*.

MURIAS HYDRARGYRI. There are two simple muriates of mercury. See *Submuriat hydrargyri*, and *Oxymuriat hydrargyri*.

MURIAS HYDRARGYRI AMMONIACALIS. See *Hydrargyrus precipitatus albus*.

MURIAS HYDRARGYRI OXYGENATUS. See *oxymuriat Hydrargyri*.

MURIAS HYPEROXYGENATUS POTASSÆ. The oxygenated muriate of potash has lately been extolled in the cure of the venereal disease. It is exhibited in doses of from fifteen to forty grains in the course of a day. It increases the action of the heart and arteries, oxygenates the blood, and proves of great service in scorbutus, asthenia, and cachectic diseases.

MURIAS POTASSÆ. *Alkali vegetabile salitum.* *Sal digestivum.* *Sal febrifugum Sylovi.* This salt is exhibited with the same intention as the muriate of soda, and was formerly in high estimation in the cure of intermittents, &c.

MURIAS SODÆ. Muriate of soda. *Alkali minerale salitum.* *Sal communis.* *Sal culinaris.* *Sal fontium.* *Sal gemme.* *Sal marinus.* *Natron muriatum.* *Soda muriata.* Common culinary salt. This salt is more abundant in nature than any other. It is found in prodigious masses in the internal part of the earth, in Calabria, in Hungary, in Muscovy, and more especially Weilicksa, in Poland, near Mount Capax, where the mines are very large, and afford immense quantities of salt. It is also obtained by several artificial means from sea-water. It possesses antiseptic, diuretic, and resolvent qualities, and is frequently employed in form of clyster, fomentation, lotion, pediluvium, and bath, in obstipation, against worms, gangrene, scrophulous, tumours, herpetic eruptions, arthritis, &c.

MURIAS STIBII HYPEROXYGENATUS. See *Antimonium muriatum*.

MURIATIC ACID GAS. The basis of this gas is still unknown. The presence of oxygen has not been even demonstrated in it, and it is only by analogy that we may venture to suppose it instrumental in this acid gas.

Properties.—It has a very pungent and suffocating odour, which excites coughing. It is readily absorbed by water, by ardent spirit, ether, fat and essential oils, melted wax, phosphorus, and many other bodies. It is a true acid. It suffocates animals, and is so very caustic as to excoriate the skin. It extinguishes a lighted taper, the flame of which becomes green, or rather light

blue, at the upper part of its disk. Light has no effect upon it. Caloric rarifies it. It is heavier than common air. The specific gravity of the former is to that of the latter as 1.750 to 1.000. When brought into contact with atmospheric air, or oxygen gas, it forms a white cloud. Ice is melted by it as speedily as if thrown into the fire. It unites to alkaline and terrene substances, and forms with them new compounds. It has no action on siliceous earths. It combines with alumine and magnesia. It absorbs oxygen, when in the state of gas, feebly, though there are methods of uniting them readily. If ammoniacal gas be mixed with it, and heat applied, both gases lose their gaseous form in a moment, and are transformed to a concrete salt. Carbonic acid gas, nitrogen gas, gaseous oxyd of nitrogen, sulphurated hydrogen gas, and carbonated hydrogen gas, have no action upon it. It has never been found in a disengaged state in nature. When electric explosions are made to pass through it, its bulk is diminished and hydrogen gas is evolved. These changes are owing to a quantity of water contained in the gas, and cease when it is deprived of moisture, as has been proved by Mr. Henry.

Method of obtaining Muriatic Acid Gas.—1. By decomposing muriate of soda, by means of sulphuric acid. For this purpose, put into a tubulated retort two parts of very dry muriate of soda, and pour on it gradually one part of concentrated sulphuric acid. A violent action takes place and muriatic acid gas becomes liberated, which must be collected over mercury in the usual manner. The sulphuric acid has a greater affinity for the soda than the muriatic acid has, it therefore unites to it and forms sulphate of soda. The muriatic acid, being liberated, takes the gaseous form, and appears as muriatic acid gas, and as the decomposition takes place very rapidly, it is not necessary to apply heat, until the disengagement of the gas begins to slacken, after which the further extrication may be assisted by the heat of a lamp.

2. Muriatic acid gas may likewise be obtained by expelling it from its combination with water.

For this purpose put concentrated muriatic acid into a retort, immerse the beak of it under a receiver placed in a mercurial pneumatic trough, and filled with that metal. On exposing the acid to a gentle heat, muriatic acid gas will be obtained. If the process be very carefully managed, nothing but water remains in the retort.

3. Muriatic acid gas is likewise produced by putting any quantity of liquid muriatic acid into a long glass tube, and adding to it about one-third or one-fourth by measure of concentrated sulphuric acid. A violent

effervescence takes place, and the whole tube becomes filled with dense white vapours, which are muriatic acid gas, condensed again by means of the moisture of the atmosphere. The sulphuric acid, added to the muriatic acid, deprives the latter of part of its water, a combination and penetration of the two liquids take place, and caloric is evolved, contributing to render the gas æriform, which is thus forced to escape.

This experiment proves that the affinity of sulphuric acid for water is greater than that of muriatic acid.

Water impregnated with this gas forms

MURIATIC ACID.

Properties.—Liquid muriatic acid, or water impregnated with muriatic acid gas, is a colourless, very odorous, and pungent fluid. It emits copious white fumes in contact with moist atmospheric air; these fumes are muriatic acid gas that escapes from it, and condenses again by combining with the humidity of the air. If a wide-mouthed bottle, containing strong muriatic acid, be opened, and the hand brought near its orifice, a sensible warmth is perceived, which arises from the combination of the acid gas with the water of the atmosphere. Liquid muriatic acid is unalterable by any known combustible body. It disengages the carbonic, phosphoric, and sulphureous acids from all their combinations, but it is constantly expelled by the action of the sulphuric acid.

Method of obtaining Muriatic Acid.—Muriatic acid is best obtained by decomposing muriate of soda, or common salt, by means of sulphuric acid, in the following manner:

Put into a tubulated retort, lodged in a sand-heat, or supported over a lamp, and connected with Pepys's distillatory vessel, or Woulf's bottles, every one containing a small quantity of distilled water, three parts of muriate of soda, and pour on it one of sulphuric acid very gradually, or rather let it be suffered to drop into the retort, by means of a funnel fastened to its tubulure, and whose inner opening may at pleasure be closed, wholly or in part, by means of a ground-glass rod. Muriatic acid gas will be plentifully disengaged, which passes through the neck of the retort, and becomes absorbed by the water. When the water in the first bottle is fully saturated, it absorbs no more, and becomes cold, but the gas continues to pass into the next bottles, and heats the water they contain. The water, thus impregnated with muriatic acid gas, is muriatic acid.

Remark.—If sulphuric acid, diluted with an equal quantity, by weight, of water, be made use of in this process, the apparatus of Pepys or Woulf may be dispensed with,

and a common receiver may be used with safety.

The salts formed by the combination of muriatic acid with different bases, are called **MURIATES**.

This acid possesses active tonic powers, in typhus, or nervous fevers, although employed on the Continent with success, and used in very liberal quantities in this country, is apt to determine to the bowels. In the fevers of children, the oxymuriatic acid is said frequently to act as a specific. Externally, the muriatic acid has been applied in the form of a bath, to the feet, in gout. In a late publication, there are accounts of its successful application as a lithontriptic.

Muriatic acid, oxygenated. See *Oxygenated muriatic acid*.

MUSADI. *Sal ammoniac.*

MUSA PARADISIACA. *Musa. Palma humilis. Ficus Indica. Bala. Platanus.* The plantain-tree. It grows spontaneously in many parts of India, but has been immemorially cultivated by the Indians in every part of the continent of South-America. It is an herbaceous tree, growing to the height of fifteen or twenty feet. The fruit are nearly of the size and shape of ordinary cucumbers, and, when ripe, of a pale yellow colour, of a mealy substance, a little clammy, a sweetish taste, and will dissolve in the mouth without chewing. The whole spike of fruit often weighs forty or fifty pounds. When they are brought to table by way of desert, they are either raw, fried, or roasted; but, if intended for bread, they are cut before they are ripe, and are then either roasted or boiled. The trees being tall and slender, the Indians cut them down to get at the fruit; and in doing this they suffer no loss, for the stems are only one year's growth, and would die if not cut; but the roots continue, and new stems soon spring up which in a year produce ripe fruit also. From the ripe plantains they make a liquor called *mistaw*. When they make this, they roast the fruit in their husks, and, after totally beating them to a mash, they pour water upon them, and, as the liquor is wanted, it is drawn off. But the nature of this fruit is such, that they will not keep long without running into a state of putrefaction; and therefore, in order to reap the advantage of them at all times, they make cakes of the pulp, and dry them over a slow fire, and, as they stand in need of *mistaw*, they mash the cakes in water, and they answer all the purposes of fresh fruit. The cakes are exceedingly convenient to make this liquor in their journeys, and they never fail to carry them for that purpose. The leaves of the tree being large and spacious, serve the Indians for table-cloths and napkins.

MUSA SAPIENTUM. The systematic name of the banana-tree. See *Banana*.

MUSCIPALA. (From *mus*, a mouse, and *capio*, to take, so called from its viscosity.) by which flies are caught as with bird-lime.) A species of lychnis.

MUSCLE. *Musculus.* The parts that are usually included under this name consist of distinct portions of flesh, susceptible of contraction and relaxation; the motions of which, in a natural and healthy state, are subject to the will, and for this reason they are called *voluntary* muscles. Besides these, there are other parts of the body that owe their power of contraction to their muscular fibres: thus the heart is a muscular texture, forming what is called a hollow muscle; and the urinary bladder, stomach, intestines, &c. are enabled to act upon their contents, merely because they are provided with muscular fibres; these are called *involuntary* muscles, because their motions are not dependent on the will. The muscles of respiration being in some measure influenced by the will, are said to have a *mixed* motion. The names by which the voluntary muscles are distinguished are founded on their size, figure, situation, use, or the arrangement of their fibres, or their origin and insertion; but, besides these particular distinctions, there are certain general ones that require to be noticed. Thus, if the fibres of a muscle are placed parallel to each other, in a straight direction, they form what anatomists term a *rectilinear* muscle; if the fibres cross and intersect each other, they constitute a *compound* muscle; when the fibres are disposed in the manner of rays, a *radiated* muscle; when they are placed obliquely with respect to the tendon, like the plume of a pen, a *penniform* muscle. Muscles that act in opposition to each other are called *antagonists*; thus every extensor has a flexor for its antagonist, and *vice versa*. Muscles that concur in the same action are termed *congeneres*. The muscles being attached to the bones, the latter may be considered as levers, that are moved in different directions by the contraction of those organs. That end of the muscle which adheres to the most fixed part is usually called the *origin*; and that which adheres to the more moveable part the *insertion* of the muscle. In almost every muscle two kinds of fibres are distinguished; the one soft, of a red colour, sensible, and irritable, called *fleshy* fibres, see *Muscular Fibres*; the other of a firmer texture, of a white glistening colour, insensible, without irritability or the power of contracting, and named *tendinous* fibres. They are occasionally intermixed, but the fleshy fibres generally prevail in the belly, or middle part of the muscle, and the tendinous ones in the extremities. If these tendinous fibres are formed into a round slender cord, they form what is called the

tendon of the muscle; on the other hand, if they are spread into a broad flat surface, it is termed an *aponeurosis*.

Each muscle is surrounded by a very thin and delicate covering of cellular membrane, which encloses it as it were like a sheath, and, dipping down into its substance, surrounds the most minute fibres we are able to trace, connecting them to each other, lubricating them by means of the fat which its cells contain in more or less quantity in different subjects, and serving as a support to the blood-vessels, lymphatics, and nerves, which are so plentifully distributed through the muscles.—This cellular membrane, which in no respect differs from what is found investing and connecting the other parts of the body, has been sometimes mistaken for a membrane peculiar to the muscles; and hence we often find writers giving it the name of *membrana propria musciosa*. The muscles owe the red colour, which so particularly distinguishes their belly part, to an infinite number of arteries, which are every where dispersed through the whole of their reticular substance; for their fibres, after having been macerated in water, are (like all other parts of the body divested of their blood) found to be of a white colour. These arteries usually enter the muscles by several considerable branches, and ramify so minutely through their substance, that we are unable, even with the best microscopes, to trace their ultimate branches. Ruysch fancied that the muscular fibre was hollow, and a production of a capillary artery; but this was merely conjectural. The veins, for the most part, accompany the arteries, but are found to be larger and more numerous. The lymphatics, likewise, are numerous, as might be expected from the great proportion of reticular substance, which is every where found investigating the muscular fibres. The nerves are distributed in such abundance to every muscle, that the muscles of the thumb alone are supplied with a greater proportion of nervous influence than the largest viscera, as the liver for instance. They enter the generality of muscles by several trunks, the branches of which, like those of the blood-vessels, are so minutely dispersed through the cellular substance, that their number and minuteness soon elude the eye and the knife of the anatomist. This has given rise to a conjecture, as groundless as all the other conjectures on this subject, that the muscular fibre is ultimately nervous.

A Table of the Muscles.

The generality of anatomical writers have arranged muscles according to their several uses; but this method is evidently defective, as the same muscle may very

often have different and opposite uses. The method here adopted is that more usually followed; they are enumerated in the order in which they are situated, beginning with those that are placed nearest the integuments, and proceeding from these to the muscles that are more deeply seated.

[The reader will be pleased to observe, that all the muscles are in pairs, except those marked thus*].

Muscles of the integuments of the cranium:

Occipito frontalis*. Corrugator supercilii.

Muscles of the eye-lids:

Orbicularis palpebrarum. Levator palpebræ superioris.

Muscles of the eye-ball.

Rectus superior. Rectus inferior. Rectus internus. Rectus externus. Obliquus superior. Obliquus inferior.

Muscles of the nose and mouth:

Levator palpebræ superioris alæque nasi. Levator labii superioris proprius. Levator anguli oris. Zygomaticus major. Zygomaticus minor. Buccinator. Depressor anguli oris. Depressor labii inferioris. Orbicularis oris*. Depressor labii superioris alæque nasi. Constrictor nasi. Levator menti vel labii inferioris.

Muscles of the external ear.

Superior auris. Anterior auris. Posterior auris. Helicis major. Helicis minor. Tragicus. Antitragicus. Transversus auris.

Muscles of the internal ear.

Laxator tympani. Membrana tympani. Tensor tympani. Stapedius.

Muscles of the lower jaw.

Temporalis. Masseter. Pterygoideus externus. Pterygoideus internus.

Muscles about the anterior part of the neck:

Platysma myoides. Sterno-cleidomastoideus.

Muscles between the lower jaw and os hyoides:

Digastricus. Mylo-hyoideus. Genio-hyoideus. Genio-glossus. Hyo-glossus. Lingualis.

Muscles situated between the os hyoides and trunk:

Sterno-hyoideus. Crico-hyoideus. Sterno-thyroideus. Thyreo-hyoideus. Cricothyroideus.

Muscles between the lower jaw and os hyoides laterally:

Stylo-glossus. Stylo-hyoideus. Stylo-pharyngeus. Circumflexus. Levator palati molli.

Muscles about the entry of the fauces.

Constrictor isthmi faucium. Palatopharyngeus. Azygos uvula*.

Muscles situated on the posterior part of the pharynx:

Constrictor pharyngis superior. Constrictor

tor pharyngis medius. Constrictor pharyngis inferior.

Muscles situated about the glottis :

Crico-arytænoideus posticus. Crico-arytænoideus lateralis. Thyreo-arytænoideus. Arytænoideus Obliquus*. Arytænoideus transversus*. Thyreo-epiglottideus. Arytæno-epiglottideus.

Muscles situated about the anterior part of the abdomen :

Obliquus descendens externus. Obliquus ascendens internus. Transversalis abdominis. Rectus abdominis. Pyramidalis.

Muscles about the male organs of generation :

Dartos*. Cremaster. Erector penis. Accelerator urinæ. Transversus perinei.

Muscles of the anus :

Sphincter ani*. Levator ani*.

Muscles of the female organs of generation :

Erector clitoridis. Sphincter vaginæ*.

Muscles situated within the pelvis :

Obturator internus. Coccygeus.

Muscles situated within the cavity of the abdomen :

Diaphragma*. Quadratus lumborum. Psoas parvus. Psoas magnus. Iliacus internus.

Muscles situated on the anterior part of the thorax :

Pectoralis major. Subclavius. Pectoralis minor. Serratus major anticus.

Muscles situated between the ribs, and within the thorax :

Intercostales externi. Intercostales interni. Triangularis.

Muscles situated on the anterior part of the neck, close to the vertebrae :

Longus colli. Rectus internus capitis major. Rectus capitis internus minor. Rectus capitis lateralis.

Muscles situated on the posterior part of the trunk :

Trapezius. Latissimus dorsi. Serratus posticus inferior. Rhomboideus. Splenius. Serratus superior posticus. Spinalis dorsi. Levatores costarum. Sacro-lumbalis. Longissimus dorsi. Complexus. Trachelo-mastoideus. Levator scapulae. Semi-spinalis dorsi. Multifidus spinæ. Semi-spinalis colli. Transversalis colli. Rectus capitis posticus minor. Obliquus capitis superior. Obliquus capitis inferior. Scalenus. Inter-spinales. Intertransversales.

Muscles of the superior extremities :

Supra-spinatus. Infra-spinatus. Teres minor. Teres major. Deltoides. Coracobrachialis. Subscapularis.

Muscles situated on the os humeri :

Biceps flexor cubiti. Brachialis internus. Biceps extensor cubiti. Anconeus.

Muscles situated on the fore arm :

Supinator radii longus. Extensor carpi radialis longior. Extensor carpi radialis brevior. Extensor digitorum communis. Ex-

tensor minimi digiti. Extensor carpi ulnaris. Flexor carpi ulnaris. Palmaris longus. Flexor carpi radialis. Pronator radii teres. Supinator radii brevis. Extensor ossis metacarpi pollicis manus. Extensor primi internodii. Extensor secundi internodii. Indicator. Flexor digitorum sublimis. Flexor digitorum profundus. Flexor longus pollicis. Pronator radii quadratus.

Muscles situated chiefly on the hand :

Lumbricales. Flexor brevis pollicis manus. Opponens pollicis. Abductor pollicis manus. Adductor pollicis manus. Abductor indicis manus. Palmaris brevis. Abductor minimi digiti manus. Adductor minimi digiti. Flexor parvus minimi digiti. Interossei interni. Interossei externi.

Muscles of the inferior extremities :

Pectinalis. Triceps adductor femoris. Obturator externus. Gluteus maximus. Gluteus minimus. Gluteus medius. Pyriformis. Gemini. Quadratus femoris.

Muscles situated on the thigh :

Tensor vaginæ femoris. Sartorius. Rectus femoris. Vastus externus. Vastus internus. Cruralis. Semi-tendinosus. Semimembranosus. Biceps flexor cruris. Popliteus.

Muscles situated on the leg :

Gastrocnemius externus. Gastrocnemius internus. Plantaris. Tibialis anticus. Tibialis posticus. Peroneus longus. Peroneus brevis. Extensor longus digitorum pedis. Extensor proprius pollicis pedis. Flexor longus digitorum pedis. Flexor longus pollicis pedis.

Muscles chiefly situated on the foot :

Extensor brevis digitorum pedis. Flexor brevis digitorum pedis. Lumbricalis pedis. Flexor brevis pollicis pedis. Abductor pollicis pedis. Adductor pollicis pedis. Abductor minimi digiti pedis. Flexor brevis minimi digiti pedis. Transversales pedis. Interossei pedis externi. Interossei pedis interni.

MUSCULAR FIBRE. The fibres that compose the body of a muscle are disposed in fasciculi, or bundles, which are easily distinguishable by the naked eye; but these fasciculi are divisible into still smaller ones; and these again are probably subdivisible *ad infinitum*. The most minute fibre we are able to trace seems to be somewhat plaited; these plaits disappearing when the fibre is put upon the stretch, seem evidently to be the effect of contraction, and have probably induced some writers to assert, that the muscular fibre is twisted or spiral. Various have been the opinions concerning the structure of these fibres; they are all of them founded only on conjecture, and therefore we shall mention only the principal ones, and this with a view rather to gratify the curiosity of the reader, than to afford him information. Borelli supposes them to be so many hol-

low cylinders, filled with a spongy medullary substance, which he compares to the pith of elder, *spongiosa ad instar sambuci*. These cylinders, he contends, are intersected by circular fibres, which form a chain of very minute bladders. This hypothesis has since been adopted by a great number of writers, with certain variations. Thus, for instance, Borelli supposes the vesicles to be of a rhomboidal shape; whereas Bernouilli contends that they are oval. Cowper went so far as to persuade himself that he had filled these cells with mercury; a mistake, no doubt, which arose from its insinuating itself into some of the lymphatics. It is observable, however, that Leeuwenhoek says nothing of any such vesicles. Here, as well as in many other of her works, Nature seems to have drawn a boundary to our inquiries, beyond which no human penetration will probably ever extend. It is surely more commendable, however, to acknowledge our ignorance, than to indulge ourselves in chimeras.

MUSCULAR MOTION. Muscular motions are of three kinds; namely voluntary, involuntary, and mixed. The *voluntary motions* of muscles are such as proceed from an immediate exertion of the active powers of the will: thus, the mind directs the arm to be raised or depressed, the knee to be bent, the tongue to move, &c. The *involuntary motions* of muscles are those which are performed by organs, seemingly of their own accord, without any attention of the mind, or consciousness of its active power: as the contraction and dilatation of the heart, arteries, veins, absorbents, stomach, intestines, &c. The *mixed motions* are those which are in part under the controul of the will, but which ordinarily act without our being conscious of their acting: as is perceived in the muscles of respiration, the intercostals, the abdominal muscles, and the diaphragm.

When a muscle acts, it becomes shorter and thicker; both its origin and insertion are drawn towards its middle. The sphincter muscles are always in action: and so likewise are antagonist muscles, even when they seem at rest. When two antagonist muscles move with equal force, the part which they are designed to move remains at rest; but if one of the antagonist muscles remains at rest, while the other acts, the part is moved towards the centre of motion.

All the muscles of living animals are constantly endeavouring to shorten themselves.

When a muscle is divided it contracts. If a muscle be stretched to a certain extent, it contracts, and endeavours to acquire its former dimensions as soon as the stretching cause is removed: this takes place in the dead body; in muscles cut out

of the body; and also in parts not muscular, and is called by the immortal Haller *vis mortua*, and by some *vis elastica*. It is greater in living than in dead bodies, and is called the *tone* of the muscles.

When a muscle is wounded, touched; or otherwise irritated, it contracts, independent of the will: this power is called *irritability*, and by Haller *vis insita*; it is a property peculiar to and inherent in the muscles. The parts of our body which possess this property are called irritable, as the heart, arteries, muscles, &c. to distinguish them from those parts which have no muscular fibres. With regard to the degree of this property peculiar to various parts, the heart is the most irritable, then the stomach and intestines; the diaphragm, the arteries, veins, absorbents, and at length the various muscles follow; but the degree of irritability depends upon the age, sex, temperament, mode of living, climate, state of health, idiosyncrasy, and likewise upon the nature of the stimulus.

When a muscle is stimulated, either through the medium of the will or any foreign body, it contracts, and its contraction is greater or less in proportion as the stimulus applied is greater or less. The contraction of muscles is different according to the purpose to be served by their contraction: thus, the heart contracts with a jerk; the urinary bladder, slowly and uniformly; puncture a muscle, and its fibres vibrate; and the abdominal muscles act slowly in expelling the contents of the rectum. Relaxation generally succeeds the contraction of muscles, and alternates with it.

The use of this property is very considerable; for upon it depends all muscular motion, and the function of every viscus, except that of the nerves.

Muscular Power. See *Irritability*.

MUSCULUS, (a diminutive of *mus*, a mouse, from its resemblance to a field mouse.) See *Muscle*.

MUSCULUS CUTANEUS. See *Platysma myoides*.

MUSCULUS FASCIE LATÆ. See *Tensor vaginæ femoris*.

MUSCULUS PATIENTIÆ. See *Levator scapule*.

MUSCULUS STAPEDIUS. See *Stapedius*.

MUSCULUS SUPERCILIJ. See *Corrugator supercilij*.

MUSCULUS TUBÆ NOYÆ. See *Circumflexus*.

MUSCUS. (From *μυσχος*, tender, so called from its delicate and tender consistence.) Moss.

MUSCUS ARBORÆUS. This plant, *Lichen plicatus* of Linnæus, we are informed by that great botanist, is applied by the Laplanders to parts which are excoriated by a long journey. It is slightly adstringent

and is applied with that intention to bleeding-vessels.

MUSCUS CANINUS. See *Lichen cinereus terrestris*.

MUSCUS CLAVATUS. See *Lycopodium*.

MUSCUS CRASSI HUMANI. See *Urina*.

MUSCUS CUMATILIA. This cryptogamous plant, *Lichen aphorus*, is said to act powerfully on the intestines, though never used in the practice of the present day.

MUSCUS ERECTUS. Upright club moss. The pharmacopœial name of the *Lycopodium selago* of Linnæus. The decoction of this plant acts violently as a vomit and a purgative, and was formerly on that account employed to produce abortions.

MUSCUS ISLANDICUS. See *Lichen islandicus*.

MUSCUS MARITIMUS. See *Corallinus*.

MUSCUS PULMONARIUS QUERCINUS. See *Pulmonaria arborea*.

MUSCUS PYXIDATUS. Cup-moss. *Musculus pyxidei terrestris*. *Lichen pyxidatus major*. These very common little plants, *Lichen cocciferus* and *pyxidatus* of Linnæus, for both are used indifferently, are employed by the common people in this country in the cure of hooping-cough, in the form of decoction.

MUSCUS SQUAMOSUS TERRESTRIS. See *Lycopodium*.

MUSLA PATTE. A name for moxa.

MUSHROOM. There are several species of the *agaricus*, which go by the term mushroom; as the *agaricus charterellus*, *deliciosus*, *violaceus*, &c. but that which is eaten in this country is the *agaricus campestris* of Linnæus. Similar to it in quality is the champignon, or *agaricus pratensis*. Broiled with salt and pepper, or stewed with cream and some aromatic, they are extremely delicious, and, if not eaten to excess, salubrious. Great care should be taken to ascertain that they are the true fungus, and not those of a poisonous nature. *Catchup* is made by throwing salt on mushrooms, which causes them to part with their juice.

Musk. See *Moschus*.

Musk-cranesbill. See *Geranium moschatum*.

Musk-melon. See *Melo*.

Musk-seed. See *Abelmoschus*.

MUSQUITTO. A species of gnat in the West Indies, which produce small tumours on whatever part they settle, attended with so high a degree of itching and inflammation, that the person cannot refrain from scratching, by a frequent repetition of which he not uncommonly occasions them to ulcerate, particularly if the subject is of a robust and full habit.

Mustard, black. See *Sinapi*.

Mustard, hedge. See *Erysimum*.

Mustard, treacle. See *Thlaspi*.

Mustard, mishridate. See *Thlaspi*.

Mustard, yellow. See *Sinapi*.

MUTITAS (From *mutus*, dumb) Dumbness. A genus of disease in the class *locales* and order *dyscinesie* of Cullen, which he defines an inability of articulation. He distinguishes three species, viz.

1. *Mutitas organica*, when the tongue is removed or injured.

2. *Mutitas autemica*, arising from an affection of the nerves of the organ.

3. *Mutitas surdorum*, depending upon being born deaf, or becoming so in their infantile years.

MUZA. See *Musa*.

MYACANTHA. (From *mys*, a mouse, and *akanthos*, a thorn, so called, because its prickly leaves are used to cover whatever is intended to be preserved from mice.) See *Ruscus*.

MYAGRO. See *Myagrum*.

MYAGRUM. (From *mys*, a fly, and *agrus*, to seize, because flies are caught by its viscosity.) A species of wild mustard.

MYCE. (From *mys*, to wink, shut up, or obstruct.) 1. It is a winking, closing, or obstruction. It is applied to the eyes, to ulcers, and to the viscera, especially the spleen, where it imports obstructions. 2. In surgery, it is a fungus, such as arises in ulcers and wounds. 3. Some writers speak of a yellow vitriol, which is called *Myce*.

MYCHTHISMOS. (From *mys*, to mutter or groan.) In Hippocrates, it is a sort of sighing or groaning during respiration, whilst the air is forced out of the lungs.

MYCONOIDES. (From *myxos*, a noise, and *oides*, a likeness,) applied to an ulcer full of mucus, and which upon pressure emits a wheezing sound.

MYCTER. The nose.

MYCTERES. *Muxræge*. The nostrils.

MYDRESIS. (From *mys*, to abound with moisture.) It imports, in general, a corruption of any part from a redundant moisture. But Galen applies it particularly to the eye-lids.

MYDOR. (From *mys*, to grow putrid.) Fungus or putrid flesh in a fistulous ulcer.

MYDRIASIS. (From *mys*, to abound in moisture; so named, because it was thought to originate in redundant moisture.) A disease of the iris. Too great a dilatation of the pupil of the eye, with or without a defect of vision. It is known by the pupil always appearing of the same latitude or size in the light. The species of mydriasis are, 1. *Mydriasis amaurotica*, which, for the most part, but not always, accompanies an amaurosis. 2. *Mydriasis hydrocephalica*, which owes its origin to an hydrocephalus internus, or internal dropsy of the ventricles of the cerebrum. It is not uncommon amongst children, and is the most certain diagnostic of the disease. 3. *Mydriasis verminosa*, or a dilatation of

the pupil from saburra and worms in the stomach or small intestines. 4. *Mydriasis a synechia*, or a dilatation of the pupil, with a concretion of the uvea with the capsule of the crystalline lens. 5. *Mydriasis paralytica*, or a dilated pupil, from a paralysis of the orbicular fibres of the iris: it is observed in paralytic disorders, and from the application of narcotics to the eye. 6. *Mydriasis spasmodica*, from a spasm of the rectilinear fibres of the iris, as often happens in hysteric and spasmodic diseases. 7. *Mydriasis*, from atony of the iris, the most frequent cause of which is a large cataract distending the pupil in its passing when extracted. It vanishes in a few days after the operation, in general; a pupil, however long dilated, may remain so from the over and long-continued distention.

MYLACRIS. (From *μύλη*, a grind-stone, so called from its shape.) The patella, or knee-pan.

MYLE. *Μύλη*. The knee-pan, or a mole in the uterus.

MYLON. See *Staphyloma*.

MYLO. Names compounded with this word belong to muscles, which are attached near the grinders; from *μύλη*, a grinder-tooth: such as,

MYLO-GLOSSI. Small muscles of the tongue.

MYLO-HYOIDEUS. *Mylo-hyoidien* of Dumas. This muscle, which was first described by Fallopius, is so called from its origin near the *dentes molares*, and its insertion into the os hyoides. It is a thin, flat muscle, situated between the lower jaw and the os hyoides, and is covered by the anterior portion of the digastricus. It arises fleshy, and a little tendinous, from all the inner surface of the lower jaw, as far back as the insertion of the pterygoideus internus, or, in other words, from between the last dens molaris and the middle of the chin, where it joins its fellow, to form one belly, with an intermediate tendinous streak, or *linea alba*, which extends from the chin to the os hyoides, where both muscles are inserted into the lower edge of the basis of that bone. This has induced Riolanus, Winslow, Albinus, and others, to consider it as a single penniform muscle. Its use is to pull the os hyoides upwards; forwards, and to either side.

MYLO-PHARYNGEUS. (*Musculus mylo-pharyngeus*, *μυλοφαρυγγιαίος*; from *μύλη*, the grinding-tooth, and *φαρυγξ*, the pharynx.) See *Constrictor pharyngeus superior*.

MYOCEPHALUM. (From *μύς*, a fly, and *κεφαλή*, a head, from its resemblance to the head of a fly.) A tumour in the uvea of the eye.

MYOCOLITIS. (From *μύς*, a muscle, and *κοιλία*, a belly. So Vogel names inflammation of the muscles of the belly.

MYODESOPSIA. (From *μύς*, a fly, and *ὄψις*, vision.) A disease of the eyes, in which the person sees black spots, an appearance of flies, cobwebs, or black wool, before his eyes.

MYOLOGY. (*Myologia*, from *μύς*, a muscle, and *λογος*, a discourse.) The doctrine of the muscles.

MYOPIA. (From *μύω*, to wink, and *ὄψις*, the eye.) Near-sighted, purblind. The myopes are considered those persons who cannot see distinctly above twenty inches. The myopia is likewise adjudged to all those who cannot see at three, six, or nine inches. The proximate cause is the adunation of the rays of light in a focus before the retina. The species are, 1. *Myopia*, from too great a convexity of the cornea. The cause of this convexity is either from nativity, or a greater secretion of the aqueous humour: hence on one day there shall be a greater myopia than on another. An incipient hydrophthalmia is the origin of this myopia. 2. *Myopia*, from too great a longitude of the bulb. This length of the bulb is native, or acquired from a congestion of the humours in the eye; hence artificers occupied in minute objects, as the engravers of seals, and persons reading much, frequently after puberty become myopes. 3. *Myopia*, from too great a convexity of the anterior superficies in the crystalline lens. This is likewise from birth. The parallel rays which fall into the cornea, by so much they fall more obliquely, so much the more convex is the cornea, or crystalline lens or vitreous humour in the anterior superficies. But the angle of refraction is equal to the angle of incidence: therefore the angle of refraction so much sooner will be formed as the cornea or lens is more convex. This perfectly accounts for short-sightedness; but an anterior too great convexity of the cornea is the most common cause. 4. *Myopia*, from too great a density of the cornea, or humours of the eye. Optics teach us by so much sooner the rays of light are forced into a focus, by so much the diaphanous body is denser. 5. *Myopia*, from a mydriasis or too dilated a pupil; for so much the wider the aperture of the diaphragma is in an optical instrument, so much the nearer is the focus. 6. *Myopia infantilis*. Infants, from the great convexity of the cornea, are often myopes; but by degrees, as they advance in years, they perceive objects more remotely, by the cornea becoming less convex.

MYOPS. (From *μύω*, to wink, and *ὄψις*, the eye.) One who is near-sighted.

MYOSIS. *Μύωσις*. A contraction or too small perforation of the pupil: it is known by viewing the diameter of the pupil, which is smaller than usual, and remains so in an obscure place, where, *αἴτιον*

rally, if not diseased, it dilates. It occasions weak sight, or a vision that remains only a certain number of hours in the day ; but, if wholly closed, total blindness. The species of this disorder are, 1. *Myosis spasmodica*, which is observed in the hysterical, hypochondriac, and in other spasmodic and nervous affections ; it arises from a spasm of the orbicular fibres of the iris. 2. *Myosis paralytica* arises in paralytic disorders. 3. *Myosis inflammatoria*, which arises from an inflammation of the iris or uvea, as in the internal ophthalmia, hypopium, or wounded eye. 4. *Myosis*, from an accustomed contraction of the pupil. This frequently is experienced by those who contemplate very minute objects ; by persons who write ; by the workers of fine needle-work ; and by frequent attention to microscopical inquiries. 5. *Myosis*, from a defect of the aqueous humour, as after extraction. 6. *Myosis nativa*, with which infants are born. 7. *Myosis naturalis*, is a coarctation of the pupil by light, or from an intense examination of minutest objects. These coarctations of the pupil are temporary, and spontaneously vanish.

MYOSITIS. (From *μῦς*, a muscle.) Inflammation of a muscle. It is the term given by Sagar to acute rheumatism.

MYOSOTIS. (*Mos*, a muscle, and *ωτ*, ear, an ear ; so called because its leaves are hairy, and grow longitudinally, like the ear of a mouse.) See *Pilosella*.

MYOTOMY. (From *μῦς*, a muscle, and *τομή*, to cut.) The dissection of the muscles.

MYRICA GALE. The systematic name of the Dutch myrtle. See *Myrtus brabantica*.

MYRIOPHYLLON. (From *μῦρος*, infinite, and *φύλλον*, a leaf, named from the number of its leaves.) See *Millefolium*.

MYRISTICA. The name of a genus of plants in the Linnæan system. Class, *Dicæcia*. Order, *Monadelphia*.

MYRISTICA AROMATICA. Swartz's name of the nutmeg-tree.

MYRISTICA MOSCHATA. The systematic name of the tree which produces the nutmeg. See *Nux moschata*.

MYRISTICA NUX. See *Nux moschata*.

MYRMECIA. (From *μῦρμηξ*, a pismire,) A small painful wart, of the size and shape of a pismire. See *Myrmecium*.

MYRMECIUM. A moist soft wart about the size of a lupine, with a broad base, deeply rooted, and very painful. It grows on the palms of the hands and soles of the feet.

MYROCOFUM. (From *μῦρος*, an ointment, and *κοπος*, labour.) An unguent to remove lassitude.

MYROBALANUS. (From *μῦρος*, an unguent, and *καλάνος*, a nut, so called be-

cause it was formerly used in ointments.) A myrobalan. A dried fruit, of the plum kind, brought from the East Indies. All the myrobalanus have an unpleasant, bitterish, very austere taste, and strike an inky blackness with a solution of steel. They are said to have a gently purgative as well as an astringent and corroborating virtue. In this country they have been long expunged from the pharmacopœias. Of this fruit there are several species.

MYROBALANUS BELLIRICA. The Belliric myrobalan. This fruit is of a yellowish grey colour, and an irregular roundish or oblong figure, about an inch in length, and three quarters of an inch thick.

MYROBALANUS CHEBULA. The chebule myrobalan. This resembles the yellow in figure and ridges, but is larger, of a darker colour, inclining to brown or blackish, and has a thicker pulp.

MYROBALANUS CITRINA. Yellow myrobalan. This fruit is somewhat longer than the Belliric, with generally five large longitudinal ridges, and as many smaller between them, somewhat pointed at both ends.

MYROBALANUS EMBLICA. The emblic myrobalan is of a dark blackish grey colour, roundish, about half an inch thick, with six hexagonal faces, opening from one another.

MYROBALANUS INDICA. The Indian or black myrobalan, of a deep black colour, oblong, octangular, differing from all the others in having no striae, or only the rudiments of one, from which circumstance they are supposed to have been gathered before maturity.

MYROBALANS. See *Myrobalanus*.

MYRON. (From *μῦρος*, to flow.) An ointment, medicated oil, or unguent.

MYROPHYLLUM. *Millefolium aquaticum*. Water-fennel. It is said to be vulnerary.

MYROXYLON. (From *μῦρος*, an ointment, and *ξύλον*, wood.) The name of a genus of plants in the Linnæan system. Class, *Diandria*. Order, *Monogynia*.

MYROXYLON PERUVIENUM. The systematic name of the tree which gives out the Peruvian balsam. See *Balsamum Peruvianum*.

MYRRHA. (*Myrrha*, Heb.) Also called *stacte*, and the worst sort *ergasma*. A botanical specimen of the tree which affords this gum-resin has not yet been obtained ; but from the account of Mr. Bruce, who says it very much resembles the *cacia vera* of Linnæus, there can be little doubt in referring it to that genus, especially as it corresponds with the description of the tree given of it by Dioscorides. The tree that affords the myrrh, which is obtained by incision, grows on the eastern coast of Arabia Felix, and in that part of Abyssinia which is situ-

ated near the Red Sea, and is called by Mr. Bruce troglodite. Good myrrh is of a turbid black red colour, solid and heavy, of a peculiar smell, and bitter taste. Its medicinal effects are warm, corroborant, and antiseptic; it has been given as an emmenagogue in a dose from 5 to 20 grains; it is also given in cachexias, and applied externally as an antiseptic and vulnerary. In doses of half a drachm, Dr. Cullen remarks that it heats the stomach, produced sweat, and agreed with the balsams in affecting the urinary passages. It has lately come more into use as a tonic in hectic cases, and is said to prove less heating than most other medicines of that class. Myrrh dissolves almost totally in boiling water, but, as the liquor cools, the resinous matter subsides. Rectified spirit dissolves less of this concrete than water; but extracts more perfectly that part in which its bitterness, virtues and flavour reside; the resinous matter which water leaves undissolved is very bitter, but the gummy matter which spirit leaves undissolved is insipid, the spirituous solution containing all the active part of the myrrh: it is applied to ulcers, and other external affections of a putrid tendency; and also as a wash, when diluted, for the teeth and gums. There are several preparations of this drug in the London and Edinburgh pharmacopœias.

MYRRHINE. (From *myrra*, myrrh; so called because it smells like myrrh.)

MYRRIS. (From *myrra*, myrrh; so named from its myrr-like smell.) Sweet cicely; anti-scorbutic.

MYRSINUM. (From *myrron*, the myrtle, and *lauro*, oil.) Oil of myrtle.

MYRTACANTHA. (From *myrtos*, a myrtle, and *anarba*, a thorn; so called from its likeness to myrtle, and from its prickly leaves.) Butcher's broom. See *Ruscus*.

MYRTIDANUM. (From *myrtos*, the myrtle.) An excrescence growing on the trunk of the myrtle, and used as an astringent.

MYRTILLUM. The berries which are directed in pharmacopœias by the name of *bacca myrtillorum*, are the fruit of the *Vaccinium myrtillus* of Linnæus. Prepared with vinegar they are esteemed as antiscorbutics, and when dry possess astringent virtues.

Myrtiform caruncles. See *Glandule myrtiformes*.

Myrtiform glands. See *Glandule myrtiformes*.

Myrtle, common. See *Myrtus*.

Myrtle, Dutch. See *Myrtus brabantica*.

MYRTO CHLIDIBES. (From *myrtos*, the clitoris, and *χλωος*, a lip.) The nymphæ of the female pudenda.

MYRTON. The clitoris.

MYRTUM. (From *myrtos*, a myrtle.) A little prominence in the pudenda of women, resembling a myrtle-berry. It also means the clitoris.

MYRTUS. (From *myrta*, myrrh, because of its smell, or from *myrrha*, a virgin who was fabled to have been turned into this tree.)

1. The name of a genus of plants in the Linnæan system. Class, *Scosandria*. Order, *Monogynia*. 2. The pharmacopœial name of the *Myrrhine*. The myrtle. *Myrtus communis* of Linnæus. The berries of this plant are recommended in alvine and uterine fluxes, and other disorders from relaxation and debility. They have a roughish, and not unpleasant taste, and appear to be moderately astringent and corroborant, partaking also of aromatic qualities.

MYRTUS BRABANTICA. *Myrtus Anglica*. *Myrtifolia belgica*. Gale. *Gagel*. *Rhus sylvestris*. *Acorn*. *Eleagnus*. *Eleagnus cordo*. *Chamaeleagnus dodonæo*.

The leaves, flowers, and seeds of this plant, *Myrica-gale* of Linnæus, sweet willow, or Dutch myrtle, have a strong fragrant smell, and a bitter taste. They are said to be used amongst the common people for destroying moths and cutaneous insects, and the infusion is given internally as a stomachic and vermifuge.

MYRTUS CARYOPHYLLATA. The systematic name of the tree which affords the cassia bark. See *Cassia caryophyllata*.

MYRTUS COMMUNIS. *Myrtus communis italica*. The systematic name of the common myrtle. See *Myrtus*.

MYRTUS FYMENTA. The systematic name of the tree which bears the Jamaica pepper. See *Pimento*.

MYSTAX. The hair which forms the beard in man, on each side the upper lip.

MYXUS. An epithet for a sort of sinking pulse, when the second stroke is less than the first, the third than the second, &c. Of this there are two kinds: the first in when the pulse so sinks as not to rise again; the other, when it returns again, and rises in some degree. Both are esteemed bad presages.

MYXOBACOMA. (From *myxa*, mucus, and *bagē*, flesh.) *Mucocarcinus*. A tumour which is partly fleshy and partly mucous.

MYXTER. (From *myxa*, the mucus of the nose.) The nose or nostril.

N.

N. In prescriptions this letter is a contraction for *numero*, in number.

NACTA. An abscess of the female breast

NADUCEM. A uterine coagulum, or mole.

NEVI MATERNI. *Macule matricies.* *Stigmata.* *Metrocildes.* Mother's marks. Marks on the skin of children, which are born with them, and which are produced by the longing of the mother for particular things, or her aversion to them; hence they resemble either mulberries, strawberries, grapes, pines, bacon, &c.

NAI CORONA. A name of the cowage.

NAIL. *Unguis.* A horny lamina, situated on the extremity of the fingers and toes.

NAKIR. According to Schenkus this means wandering pains of the limbs.

NAPELLUS. (A diminutive of *napus*, a kind of turnip; so called because it has a bulbous root like that of the *napus*.) See *Aconitum*.

NAPHE FLORES. Orange flowers are sometimes so called. See *Aurantium*.

NAPHTHA. *Naphtha.* A very fluid species of petroleum, of a lighter colour than petroleum, more or less transparent, perfectly thin and liquid, light, so as to float on water, odoriferous, volatile, and inflammable. It is found separated by nature from petroleum and bitumen, but its separation is more readily effected by art. This fluid has been used for an external application for removing old pains, nervous disorders, such as cramps, contractions of the limbs, paralytic affections, &c.

NAPIFOLIA. Bare cole.

NAPIUM. The name of the nipple wort. A species of *lapsana*. It is one of the bitter lactescent plants, similar in virtues to endive. See *Lapsana*.

NAFUS. (From *napus*, Rabb.) *Napus sylvestris.* *Bunias.* Wild navew, or rape. The *Brassica napus* of Linnæus. The seeds yield upon expression a large quantity of oil called rape oil, which is sometimes ordered in stimulating liniments.

NAPUS DULCIS. See *Rapus*.

NAPUS SYLVESTRIS. See *Rapus*.

NARCISSUS. The daffodil. A genus of plants in the Linnæan system. Class, *Hexandria*. Order, *Monogynia*.

NARCOSIS. (From *narcoo*, to stupefy.) Stupefaction, stupor, numbness.

NARCOTICS. (*Narcotica*, sc. *Medicamenta*. From *narcoo*, to stupefy.) Those

medicines which have the power of procuring sleep. See *Anodynes*.

Nard, Celtic. See *Nardus Celtica*.

Nard, Indian. See *Nardus Indica*.

NARDOSTACHYS. (From *nardos*, spike-nard, and *stachys*, sage.) A species of wild sage resembling spikenard in its leaves and smell.

NARDUS. (From *nard*, Sir.) Spike-nard.

NARDUS CELTICA. *Spica Celtic.* *Dioscoridis.* Celtic nard. *Valeriana Celtica* of Linnæus. The root of this plant, a native of the Alps, has been recommended as a stomachic, carminative, and diuretic. At present it is only used in this country in the theriaca and mithridate, though its sensible qualities promise some considerable medicinal powers. It has a moderately strong smell, and a warm, bitterish, sub-acrid taste.

NARDUS INDICA. *Spica nardi.* *Spica Indica.* Indian nard or spikenard. The root of this plant, *Andropogon nardus* of Linnæus, is an ingredient in the mithridate and theriaca; it is moderately warm and pungent, accompanied with a flavour not disagreeable. It is said to be used by the Orientals as a spice.

NARDUS ITALICA. The *lavendula latifolia*.

NARDUS MONTANA. An old name of the *asarabacca*.

NARDUS RUSTICA. An old name of the *asarabacca*. See *Asarum*.

NARES. (Pl. of *nasus*.) *Mycteres.* The nostrils. The cavity of the nostrils is of a pyramidal figure, and is situated under the anterior part of the cranium, in the middle of the face. It is composed of fourteen bones, viz. the frontal, two maxillary, two nasal, two lachrymal, two inferior spongy, the sphæmoid, the vomer, the ethmoid, and two palatine bones, which form several eminences and cavities. The eminences are the septum narium, the cavernous substance of the ethmoid bone, called the superior conchæ, and the inferior spongy bones. The cavities are three pair of pituitary sinuses, namely, the frontal, sphæmoid and maxillary; the anterior and posterior foramina of the nostrils; the ductus nasalis, the sphæno-palatine foramina, and anterior palatine foramina. All these parts are covered with periosteum, and a pituitary membrane which secretes the mucus of the nostrils. The arteries of this cavity are branches of the internal maxillary. The veins empty themselves

into the internal jugulars. The nerves are branches of the olfactory, ophthalmic, and superior maxillary. The use of the nostrils is for smelling, respiration, and speech.

NARISUSORIA. (From *nares*, the nostrils, and *fundo*, to pour.) Medicines dropped into the nostrils.

NARIS COMPRESSOR. See *Compressor naris*.

NARTA. (*Narta*, ex *nardi* odore, from its smell.) A plant used in ointments.

NARTHECIA. (From *Nartheis*, the island where it flourished.) *Narthez*. A kind of fennel.

NASALIA. (From *nasus*, the nose.) Errhines.

NASALIS LABIS SUPERIORI. See *orbicularis oris*.

NASARIUM. (From *nasus*, the nose.) The mucus of the nose.

NASCALE. (From *nasus*, the nose.) A wood of cotton pessary for the nose.

NASCAPTHUM. See *Narcapthum*.

NASI, DEPRESSOR. See *Depressor labii superioris alicque nasi*.

NASI OSSA. (*nasus*, the nose.) The two small bones of the nose that are so termed from the bridge of the nose. In figure they are quadrangular and oblong.

NASTURTIIUM. (*Quod nanum torquent*, because the seed, when bruising, irritates the nose.) The name of a genus of plants in the Linnæan system. Class *Tetradynamia*. Order *Siliquosa*.

NASTURTIIUM AQUATICUM. *Laver odoratum*. *Cratæva etum*. *Cressis*. *Cardaminea*. Water-cress. This indigenous plant *Sisymbrium nasturtium* of Linnæus; *siliquis declinatis, foliis pinnatis, foliolis subcordatis*, grows plentifully in brooks and stagnant waters. The leaves have a moderate pungent taste, emit a quick penetrating smell, like that of mustard seed, but much weaker. Water-cresses obtain a place in the materia medica, for their antiscorbutic qualities, which have been long very generally acknowledged by physicians. The most pleasant way of administering them is in form of a salad.

NASTURTIIUM HORTENSE. Dittander. This plant is the *Lepidium sativum* of Linnæus; *floribus tetradynamiis; foliis oblongis, multifidis*; it possesses warm, nervine, and stimulating qualities, and is given as an antiscorbutic, antiseptic, and stomachic, especially by the lower order.

NASTURTIIUM INDICUM. *Arctostaphylos*. *Flos sanguineus monardi*. *Nasturtium Peruvianum*. *Cardaminum minus*. Greater Indian cress or nasturtium. *Trapezium majus* of Linnæus. This plant is a native of Peru; it was first brought to France in 1684, and there called *La grande capucine*. In its recent state this plant, and more especially its flowers, have a smell and taste resembling those of water-cress; and the

leaves, on being bruised in a mortar, emit a pungent odour, somewhat like that of horse-radish. By distillation with water, they impregnate the fluid in a considerable degree with the smell and flavour of the plant. Hence the antiscorbutic character of the nasturtium seems to be well founded, at least as far as we are able to judge from its sensible qualities: therefore in all those cases, where the warm and antiscorbutic vegetables are recommended, this plant may be occasionally adopted as a pleasant and effectual variety. Patients, to whom the nauseous taste of scurvy-grass is intolerable, may find a grateful substitute in the nasturtium. The flowers are frequently used in salads, and the capsules are by many highly esteemed as a pickle. The flowers, in the warm summer months, about the time of sunset, have been observed to emit sparks like those of the electrical kind.

NATA. Natta. A species of wen with slender pendent neck. Linnæus speaks of it as rooted in a muscle.

NATES. (From *nato*, to flow: because the excrements are discharged from them.) 1. The buttocks, or the fleshy parts upon which we sit. 2. Two of the eminences of the tubercle quadrigemina of the brain are so named, from their resemblance.

NATES CEREBRI. See *Tubercula quadrigemina*.

NATRŒN. (So called from *Natron*, a lake in Judea, where it was produced.) *Natrum*. 1. The name formerly given by the college of physicians to the alkali now called soda.

2. A native salt, which is found crystallized in Egypt, in the lake called Natron, and in other hot countries, in sands surrounding lakes of salt water. It is an impure carbonate of soda.

3. The name of an impure carbonate of soda, obtained by burning *Salsoi kali* of Linnæus. Not only this, but various other plants, on being burned, are found to afford this alkali, and some in a greater proportion than this: these are

The *Salsoia sativa*, Lin. *Salsoia sonda*, Lofling. *Kali hispanicum rupestre annuum sedi-folius brevibus*. *Kali d'Alicante*. This grows abundantly on that part of the Spanish coast which is washed by the Mediterranean sea. This plant is deservedly first enumerated by Professor Murray, as it supplies all the best soda consumed in Europe, which by us is called Spanish or Alicante soda, and by the Spanish merchants Barilla de Alicante. See *Barilla*.

Salsoia soda, Lin. *Kali majus cochleato semine*. *Le salicor*. This species, which grows on the French Mediterranean coast, is much used in Languedoc for the preparation of this salt, which is usually exported to Sicily and Italy.

Salsola tragus, Lin. affords an ordinary kind of soda, with which the French frequently mix that made in Languedoc. This adulteration is also practised by the Sicilians, who distinguish the plant by the term *salvaggia*.

Salicornia herbacea, Lin. is common in salt marshes, and on the sea-shore, all over Europe. Linnæus prefers the soda obtained from this plant to that of all the others; but though the quantity of alkali which it yields is very considerable, as a great portion of it is united with muriatic acid, it is mixed with much common salt.

Salicornia arabica, Lin. *Mesembryanthemum nodiflorum*, Lin. *Plantago squarrosa*, Lin. All these, according to Alpinus, afford this alkali. It has also been procured from several of the fuci, especially *F. vesiculosus*, and distinguished here by the name kelp. Various other marine plants might also be noticed as yielding an impure soda by combustion; but the principal are confined to the genus *salsola*, and that of *salicornia*. The *salsola kali*, on the authority of Rauwolf, is the species from which the salt is usually obtained in eastern countries.

It is to be regretted that the different kinds of soda, which are brought to European markets, have not been sufficiently analysed to enable us to ascertain with tolerable certainty the respective value of each; and indeed, while the practice of adulterating this salt continues, any attempts of this kind are likely to prove fruitless. The best information on this subject is to be had from Jessica, Mascarelle, Cadet, Borlare, and Seutini. In those places where the preparation of soda forms a considerable branch of commerce, as on the coast of the Mediterranean, seeds of the *salsola* are regularly sown in a proper situation near the sea, which usually shoot above ground in the course of a fortnight. About the time the seeds become ripe, the plants are pulled up by the roots, and exposed in a suitable place to dry, where their seeds are collected; this being done, the plants are tied up in bundles, and burned in an oven constructed for the purpose, where the ashes are then, while hot, continually stirred with long poles. The saline matter, on becoming cold, forms a hard solid mass, which is afterwards broken in pieces of a convenient size for exportation.

According to chymical analysis, the impure sodas of commerce generally contain a portion of vegetable alkali, and neutral salts, as muriate of soda and sulphate of potash, and not unfrequently some portion of iron is contained in the mass; they are therefore to be considered as more or less a compound, and their goodness to be estimated accordingly. The Spanish soda, of

the best sort, is in dark coloured masses, of a bluish tinge, very ponderous, sonorous, dry to the touch, and externally abounding with small cavities, without any offensive smell, and very salt to the taste: if long exposed to the air, it undergoes a degree of spontaneous calcination. The best French soda is also dry, sonorous, brittle, and of a deep blue colour, approaching to black. The soda which is mixed with small stones, which gives out a fetid smell on solution, and is white, soft, and deliquescent, is of the worst kind. The method of purifying the impure carbonates of soda of commerce is directed in the London Pharmacopœia, under the article of Sodæ subcarbonas, and in the Edinburgh Pharmacopœia, under that of Sal alkalinus fixus fossilis purificatus. The pure crystals, thus formed of Alicant barilla, are colourless, transparent, lamellated, of a rhomboidal figure; and one hundred parts are found to contain twenty of alkali, sixteen of aerial acid, and sixty-four of water; but upon keeping the crystals for a length of time, if the air be not excluded, the water evaporates, and they assume the form of a white powder. According to Inslin, one ounce of water, at the temperature 62° of Fahr. dissolves five drachms and fifteen grains of the crystals.

NATRON MURIATUM. See *Murias sodæ*.

NATRON PREPARATUM. See *Subcarbonas sodæ*.

NATRON TARTARISATUM. See *Tartarizata sodæ*.

NATRON VITRIOLATUM. See *Sulphas sodæ*.

NATULÆ. (Dim. of *nates*, the buttocks; so called from their resemblance.) Two prominences of the brain. See *Tubercula quadrigemina*.

NATURAL ACTIONS. Those actions by which the body is preserved; as hunger, thirst, &c. See *Function*.

NATURALIA. (From *natura*, nature.) The parts of generation.

NAUSEA. (*Nauus*: from *navis*, a ship; because it is a sensation similar to that which people experience upon sailing in a ship.) *Nauisiois*. *Nautia*. An inclination to vomit without effecting it; also a disgust of food, approaching to vomiting. It is an attendant on cardialgia, and a variety of other disorders, pregnancy, &c. occasioning an aversion for food, an increase of saliva, disgusted ideas at the sight of various objects, loss of appetite, debility, &c.

NAUSIOSIS. See *Nausea*.

NAUTIA. See *Nausea*.

NAUTICUS. (*Nauticus*, a sailor; so called from the use which sailors make of it, in climbing ropes.) A muscle of the leg, exerted in climbing up.

Navew, garden. See *Rapus*.

Naveu, sweet. See *Rapus*.

Naveu, wild. See *Rapus*.

NAVICULARE OS. (From *navicula*, a little boat.) *Naviformis*. *Navicularis*. *Os scaphoides*. A bone of the carpus and tarsus is so called, from its supposed resemblance to a boat. See *Carpus* and *Tarsus*.

NAVICULARIS. See *Naviculare os*.

NAVIFORMIS. A name of the *os naviculare*.

NEAPOLITANUS MORBUS. (From *Neapolis*, or *Naples*, because it was said to have been first discovered at Naples, when the French were in possession of it.) The venereal disease.

NEBULA. (From νεφελή.) 1. A cloudy spot in the cornea of the eye.

2. The cloud-like appearance in the urine, after it has been a little time at rest.

NECK. *Collum*. The parts which form the neck are divided into external and internal. The external parts are the common integuments, several muscles, eight pair of cervical nerves, the eighth pair of nerves of the cerebrum, and the great intercostal nerve; the two carotid arteries, the two external jugular veins, and the two internal; the glands of the neck, viz. the jugular, submaxillary, cervical, and thyroid. The internal parts are, the fauces, pharynx, œsophagus, larynx, and trachea. The bones of the neck are the seven cervical vertebræ.

NECROSIS. (From νεκρός, to destroy.) This word, the strict meaning of which is only mortification, is, by the general consent of surgeons, confined to this affection of the bones. The death of parts of bones was not distinguished from caries, by the ancients. However, necrosis and caries are essentially different; for in the first, the part affected of the bone is deprived of the vital principle; but this is not the case when it is simply carious. Caries is very analogous to ulceration, while necrosis is exactly similar to mortification of the soft parts.

NECROSIS OSTILAGINEA. A painful convulsive contraction of the limbs.

NECTAR. *Nectar*. A wine made of honey.

NEDYIA. (From νῆδυς, the belly.) *Nedys*. The intestines.

NEDYS. See *Nedyia*.

NEGRO CACHEXY. *Cachexia Africana*. Mal d'estomach of the French. A propensity for eating earth, peculiar to males as well as females, in the West Indies and Africa.

NELERA. (From νεύω, to furthermost.) The lower part of the belly.

NEMOROSA. (From nemus, a grove; so called because it grows in woods.) A species of anemone.

Nep. See *Nepeta*.

NEPA THEOPHRASTI. The spartium scoparium.

NEPENTHES. (From νη neg. and πένθος, grief; so called from their exhilarating qualities.) A preparation of opium, and a kind of bugloss.

NEPETA. (From *nepte*, Germ.) *Herba felis*. *Nep.* or catmint; so called because cats are very fond of it. The leaves of this plant, *Nepeta cataria* of Linnæus;—*floribus spicatis; verticillis subpedicellatis; foliis petiolatis, cordatis, dentata-serratis*, have a moderately pungent aromatic taste, and a strong smell, like an admixture of spearmint and penny-royal. The herb is recommended in uterine disorders, dyspepsia, and flatulency.

NEPETA CATARIA. The systematic name of the catmint. See *Nepeta*.

NEPETELLA. (Dim. of *nepeta*.) The lesser catmint.

NEPHELA. (Dim. of νεφός, a cloud.) A cloud-like spot on the cornea of the eye.

NEPHELOIDES. (From νεφός, a cloud, and εἶδος, a likeness.) Cloudy. Applied to the urine.

NEPHRALGIA. (From νεφός, the kidney, and ἄλγος, pain.) Pain in the kidney.

NEPHRALGIA CALCULOSA. Pain from stone in the kidneys.

NEPHRALGIA RHEUMATICA. The lumbago, or pain in the loins.

NEPHRELMINTHICA ISCHURIA. (From νεφός, the kidney, and ἔμυς, a worm.) Applied to a suppression of urine from worms.

Nephritic wood. See *Lignum nephriticum*.

NEPHRITICA AQUA. Spirituous distillation of nutmeg and hawthorn flowers.

NEPHRITICS. (*Nephritica*, sc. *medicamenta*; from νεφός, the kidney.) Medicines are so termed that are employed in the cure of diseases of the kidneys.

NEPHRITICUM LIGNUM. See *Lignum nephriticum*.

NEPHRITIS. (From νεφός, a kidney.) Inflammation of the kidney. A genus of disease in the class *pyrexia*, and order *phlegmasia*, of Cullen; known by prexia, pain in the region of the kidneys, and shooting along the course of the ureter; drawing up of the testicles; numbness of the thigh; vomiting; urine high coloured, and frequently discharged; costiveness, and colic pains. Nephritis is symptomatic of calculus, gout, &c.

This inflammation may be distinguished from the colic, by the pain being seated very far back, and by the difficulty of passing urine, which constantly attends it; and it may be distinguished from rheumatism, as in this the pain is but little influenced, or increased, either by motion or pressure.

Nephritis is to be distinguished from a

calculus in the kidney, or ureter, by the symptoms of fever accompanying, or immediately following the attack of pain, and these continuing without any remarkable intermission; whereas, in a calculus of the kidney, or ureter, they do not occur until a considerable time after violent pain has been felt. In the latter case, too, a numbness of the thigh, and a retraction of the testicle on the affected side, usually takes place.

The causes which give rise to nephritis are, external contusions, strains of the back, acrids conveyed to the kidneys in the course of the circulation, violent and severe exercise, either in riding or walking, calculous concretions lodged in the kidneys, or ureters, and exposures to cold. In some habits, there is an evident predisposition to this complaint, particularly the gouty, and in these there are often translations of the matter to the kidneys, which very much imitate nephritis.

An inflammation of the kidney is attended with a sharp pain on the affected side, extending along the course of the ureter; and there is a frequent desire to make water, with much difficulty in making it; the body is costive, the skin is dry and hot, the patient feels great uneasiness when he endeavours to walk, or sit upright; he lies with most ease on the affected side, and is generally troubled with nausea and frequent vomiting.

When the disease is protracted beyond the seventh or eighth day, and the patient feels an obtuse pain in the part, has frequent returns of chilliness and shiverings, there is reason to apprehend that matter is forming in the kidney, and that a suppuration will ensue.

Dissections of nephritis shew the usual effects of inflammation on the kidney; and they likewise often discover the formation of abscesses, which have destroyed its whole substance. In a few instances, the kidney has been found in a scirrhous state.

NEPHROLITICA ISCHURIA. (From *νεφρος*, a kidney, and *λιθος*, a stone.) Applied to an ischury, from a stone in the kidneys.

NEPHROMETRE. The prostatic muscles.

NEPHROPLETHORICUS. (From *νεφρος*, a kidney, and *πληθωρα*, a plethora.) Applied to a suppression of urine from a plethora.

NEPHROPHLEGMA. (From *νεφρος*, a kidney, and *φλεγμα*, phlegm.) Applied to a suppression of urine from pituitous or mucous matter in the kidneys.

NEPHROPLEGICUS. (From *νεφρος*, a kidney, and *πληγμα*, a stroke.) A suppression of urine from a paralysis of the kidney is called ischuria nephroplegica.

NEPHROPTICUS. (From *νεφρος*, a kidney, and *πτωσις*, pus.) Applied to a sup-

pression of urine from purulent matter in the kidneys.

NEPHROS. (From *νεω*, to flow, and *φωρα*, to bear; as conveying the urinary fluid.) A kidney.

NEPHROSPASTICUS. (From *νεφρος*, a kidney, and *σπασμος*, to contract.) Applied to a suppression of urine from a spasm of the kidneys.

NEPHROTHRYMBOIDES. (From *νεφρος*, a kidney, and *θρομβος*, a grumous concretion, and *ωμοις*, a likeness.) Applied to a suppression of urine from grumous blood in the kidneys.

NEPHROTOMY. *Nephrotomia*; from *νεφρος*, a kidney, and *τομή*, to cut.) The operation of extracting a stone from the kidney. A proceeding which, perhaps, has never been actually put in practice. The cutting into the kidney, the deep situation of this viscus, and the want of symptoms by which the lodgment of a stone in it can be certainly discovered, will always be strong objections to the practice.

NERIUM. (From *νερος*, humid; so called because it grows in moist places.) The name of a genus of plants in the Linnean system. Class, *Pentandria*. Order, *Monogynia*.

NERIUM ANTIDYSENTERIUM. The systematic name of the tree which affords the Cadoga pala bark. See *Coniis cortex*.

NEROLI OLEUM. Essential oil of orange flowers. See *Aurantium*.

NERVALIA OSSA. (From *nervus*, a nerve.) The bones through which the nerves pass.

NERVE. *Nervus*. Formerly it meant a sinew. This accounts for the opposite meanings of the word *nervous*; which sometimes means strong, sinewy; and sometimes weak, and irritable. Nerves are long, white, medullary cords, that serve for sensation. They originate from the brain and spinal marrow; hence they are distinguished into cerebral and spinal nerves, and distributed upon the organs of sense, the viscera, vessels, muscles, and every part that is endowed with sensibility. The cerebral nerves are the olfactory, optic, motores oculorum, pathetici, or trochleatores, trigimini, or divisi, abducent, auditory, or acoustic, par vagum and lingual. Heister has drawn up the uses of these nerves in the two following verses:

*Olfacients, cernens, oculosque movens patiensque,
Gustans, abducens, audiensque, vagansque, loquensque.*

The spinal nerves are thirty pair, and are divided into eight pair of cervical, twelve pair of dorsal, five pair of lumbar, and five of sacral nerves. In the course of the nerves there are a number of knots: these are called *ganglions*; they are commonly of an oblong shape, and of a grayish colour.

somewhat inclining to red, which is, perhaps, owing to their being extremely vascular. Some writers have considered these little ganglions as so many little brains. Lancisi fancied he had discovered muscular fibres in them, but they certainly are not of an irritable nature. A late writer (Dr. Johnson) imagines they are intended to deprive us of the power of the will over certain parts, as the heart, for instance; but if this hypothesis were well founded, they should be met with only in nerves leading to voluntary muscles; whereas it is certain that the voluntary muscles receive nerves through ganglions. Dr. Munro, from observing the accurate intermixture of the minute nerves which compose them, considers them as new sources of nervous energy. The nerves, like the blood-vessels, in their course through the body, communicate with each other, and each of these communications constitute what is called a *plexus*, from whence branches are again detached to different parts of the body. The use of the nerves is to convey the principles of motion and sensibility to the brain, from all parts of the system, and from the brain to every part of the system. The manner in which this operation is effected is not yet determined. The inquiry has been a constant source of hypothesis in all ages, and has produced some ingenious ideas, and many erroneous positions, but without having hitherto afforded much satisfactory information. Some physiologists have considered a trunk of nerves as a solid cord, capable of being divided into an infinite number of filaments, by means of which the impressions of feeling are conveyed to the common sensorium. Others have supposed each fibril to be a canal, carrying a volatile fluid, which they term the *nervous fluid*. Those who contend for their being solid bodies are of opinion that feeling is occasioned by vibration; so that, for instance, according to this hypothesis, by pricking the finger, a vibration would be occasioned in the nerve distributed through its substance; and the effects of this vibration, when extended to the sensorium, would be an excitation of pain; but the inelasticity, the softness, the connection, and the situation of the nerves, are so many proofs that vibration has no share in the cause of feeling.

A Table of the Nerves.

CEREBRAL NERVES.

1. The first pair, called *olfactory*.
2. The second pair, or *optic nerves*.
3. The third pair, or *oculorum motorii*.
4. The fourth pair, or *pathetici*.
5. The fifth pair, or *trigemini*, which gives off
 - a. The *ophthalmic*, or *orbital nerve*, which sends
 1. A branch to unite with one from the

sixth pair, and form the great intercostal nerve.

2. The *frontal nerve*.
3. The *lachrymal*.
4. The *nasal*.
- b. The *superior maxillary*, which divides into
 1. The *spheno palatine nerve*.
 2. The *posterior alveolar*.
 3. The *infra orbital*.
- c. The *inferior maxillary nerve*, from which arise
 1. The *internal lingual*.
 2. The *inferior maxillary*, properly so called
6. The sixth pair, or *abducentes*, which send off
 1. A branch to unite with one from the fifth, and form the great intercostal.
 2. The seventh pair, or *auditory nerves*, these arise by two separate beginnings, viz.
 - The *portio dura*, a nerve going to the face.
 - The *portio mollis*, which is distributed on the ear.
 - The *portio dura*, or *facial nerve*, gives off the *chorda tympani*, and then proceeds to the face.
8. The eighth pair, or *par vagum*, arise from the medulla oblongata, and join with the accessory of Willis. The *par vagum* gives off
 1. The *right and left recurrent nerves*.
 2. Several branches in the chest, to form the *cardiac plexus*.
 3. Several branches to form the *pulmonic plexus*.
 4. Several branches to form the *esophageal plexus*.
 5. It then forms in the abdomen the *stomachic plexus*.
 6. The *hepatic plexus*.
 7. The *splenic plexus*.
 8. The *renal plexus*, receiving several branches from the great intercostal, which assists in their formation.
9. The ninth pair, or *lingual nerves*, which go from the medulla oblongata to the tongue.

SPINAL NERVES.

Those nerves are called *spinal*, which pass out through the lateral or intervertebral foramina of the spine.

They are divided into *cervical*, *dorsal*, *lumbar*, and *sacral nerves*.

CERVICAL NERVES.

The *cervical nerves* are eight pairs.

The first are called the *occipital*; they arise from the beginning of the spinal marrow, pass out between the margin of the occipital foramen and atlas, form a ganglion on its transverse process, and are distributed about the occiput and neck.

The second pair of cervical nerves send a

branch to the accessory nerve of Willis, and proceed to the parotid gland and external ear.

The *third* cervical pair supply the integuments of the scapula, cucullaris, and triangularis muscles, and send a branch to form with others the diaphragmatic nerve.

The *fourth, fifth, sixth, seventh, and eighth* pair all converge to form the *brachial plexus*, from which arise the six following

NERVES OF THE UPPER EXTREMITIES.

1. The *axillary* nerve, which sometimes arises from the radial nerve: It runs backwards and outwards around the neck of the humerus, and ramifies in the muscles of the scapula.

2. The *external cutaneous*, which perforates the coraco-brachialis muscle, to the bend of the arm, where it accompanies the median vein as far as the thumb, and is lost in its integuments.

3. The *internal cutaneous*, which descends on the inside of the arm, where it bifurcates. From the bend of the arm the anterior branch accompanies the basilic vein, to be inserted into the skin of the palm of the hand; the posterior branch runs down the internal part of the fore-arm, to vanish in the skin of the little finger.

4. The *median* nerve, which accompanies the brachial artery to the cubit, then passes between the brachialis internus, pronator rotundus, and the perforatus and perforans, under the ligament of the wrist to the palm of the hand, where it sends off branches in every direction to the muscles of the hand, and then supplies the digital nerves, which go to the extremities of the thumb, fore and middle fingers.

5. The *ulnar* nerve, which descends between the brachial artery and basilic vein, between the internal condyle of the humerus and the olecranon, and divides in the fore-arm into an *internal* and *external* branch. The former passes over the ligament of the wrist and sesamoid bone, to the hand, where it divides into three branches, two of which go to the ring and little finger, and the third forms an arch towards the thumb, in the palm of the hand, and is lost in the contiguous muscles. The latter passes over the tendon of the extensor carpi ulnaris and back of the hand, to supply also the two last fingers.

6. The *radial* nerve, which sometimes gives off the axillary nerve. It passes backwards, about the os humeri, descends on the outside of the arm, between the brachialis externus and internus muscles to the cubit; then proceeds between the supinator longus and brevis, to the superior extremity of the radius, giving off various branches to adjacent muscles. At this place it divides into two branches; *one* goes along the radius, between the supinator longus and radialis internus to the back of

the hand, and terminates in the interosseous muscles, the thumb and three first fingers; the *other* passes between the supinator brevis and head of the radius, and is lost in the muscles of the fore-arm.

DORSAL NERVES.

The *dorsal* nerves are twelve pairs in number. The first pair gives off a branch to the brachial plexus. All the dorsal nerves are distributed to the muscles of the back; intercostals, serrati, pectoral, abdominal muscles, and diaphragm. The five inferior pairs go to the cartilages of the ribs, and are called *costal*.

LUMBAR NERVES.

The five pair of lumbar nerves are bestowed about the loins and muscles, skin of the abdomen and loins, scrotum, ovaria, and diaphragm. The second, third, and fifth pair unite and form the *obturator nerve*, which descends over the psoas muscle into the pelvis, and passes through the foramen thyroideum to the obturator muscle, triceps, pectineus, &c.

The third and fourth, with some branches of the second pair, form the *crural nerve*, which passes under Poupart's ligament with the femoral artery, sends off branches to the adjacent parts, and descends in the direction of the sartorius muscle to the internal condyle of the femur, from whence it accompanies the saphena vein to the internal ankle, to be lost in the skin of the great toe.

The fifth pair are joined to the first pair of the sacral nerves.

SACRAL NERVES.

There are five pair of *sacral* nerves, all of which arise from the *cauda equina*, or termination of the medulla spinalis, so called from the nerves resembling the tail of a horse. The four first pair give off branches to the pelvic viscera, and are afterwards united to the last lumbar, to form a large *plexus*, which gives off

The *ischiatric nerve*, the largest in the body. The ischiatic nerve, immediately at its origin, sends off branches to the bladder, rectum, and parts of generation; proceeds from the cavity of the pelvis through the ischiatic notch, between the tuberosity of the ischium and great trochanter, to the ham, where it is called the *popliteal nerve*. In the ham it divides into two branches.

1. The *peroneal*, which descends on the fibula, and distributes many branches to the muscles of the leg and back of the foot.

2. The *tibial*, which penetrates the gastrocnemii muscles to the internal ankle, passes through a notch in the os calcis to the sole of the foot, where it divides into an *internal* and *external plantar* nerve, which supply the muscles and aponeurosis of the foot and the toes.

Physiology of the Nervous System.

The nervous system, as the organ of sense and motion, is connected with so many functions of the animal economy, that the study of it must be of the utmost importance, and a fundamental part of the study of the whole economy. The nervous system consists of the medullary substance of the brain, cerebellum, medulla oblongata, and spinalis; and of the same substance continued into the nerves, by which it is distributed to many different parts of the body. The whole of this system seems to be properly distinguished into these four parts.

1. The medullary substance contained in the cranium and vertebral cavity; the whole of which seems to consist of distinct fibres, but without the smaller fibres being separated from each other by any evident enveloping membranes.

2. Connected with one part or other of this substance are, the *nerves*, in which the same medullary substance is continued; but here more evidently divided into fibres, each of which is separated from the others by an enveloping membrane, derived from the pia mater.

3. Parts of the extremities of certain nerves, in which the medullary substance is divested of the enveloping membranes from the pia mater, and so situated as to be exposed to the action of certain external bodies, and perhaps so framed as to be affected by the action of certain bodies only; these are named the *sentient extremities* of the nerves.

4. Certain extremities of the nerves, so framed as to be capable of a peculiar contractility; and, in consequence of their situation and attachments, to be, by their contraction, capable of moving most of the solid and fluid parts of the body. These are named the *moving extremities* of the nerves: they are commonly named *moving*, or *muscular fibres*.

These several parts of the nervous system are every where the same continuous medullary substance which is supposed to be the vital solid of animals, so constituted in living animals, and in living systems only, as to admit of motions being readily propagated from any one part to every other part of the nervous system, so long as the continuity and natural living state of the medullary substance remains. In the living man, there is an immaterial thinking substance, or *mind*, constantly present, and every phenomenon of thinking is to be considered as an affection or faculty of the mind alone. But this immaterial and thinking part of man is so connected with the material and corporeal part of him, and particularly with the nervous system, that motions excited in this give occasion to thought, and thought, however occasioned, gives occasion to new motions in the nervous system. This mu-

tual communication, or influence, is assumed with confidence as a fact: but the mode of it we do not understand, nor pretend to explain; and therefore are not bound to obviate the difficulties that attend any of the suppositions which have been made concerning it. The phenomena of the nervous system occur commonly in the following order. The impulse of external bodies acts upon the sentient extremities of the nerves; and this gives occasion to perception, or thought, which, as first arising in the mind, is termed *sensation*. This sensation, according to its various modification, gives occasion to *volition*, or the willing of certain ends to be obtained by the motion of certain parts of the body; and this volition gives occasion to the contraction of muscular fibres, by which the motion of the part required is produced. As the impulse of bodies on the sentient extremities of a nerve does not occasion any sensation, unless the nerve between the sentient extremity and the brain be free; and as, in like manner, volition does not produce any contraction of muscles, unless the nerve between the brain and muscle be also free, it is concluded, from both these facts, that sensation and volition, so far as they are connected with corporeal motions, are functions of the brain alone; and it is presumed, that sensation arises only in consequence of external impulse producing motion in the sentient extremities of the nerves, and of that motion's being thence propagated along the nerves to the brain; and, in like manner, that the will operating in the brain only, by a motion begun there, and propagated along the nerves, produces the contraction of muscles. From what is now said, we perceive more distinctly the different functions of the several parts of the nervous system, as distinguished in, 1. The sentient extremities seem to be particularly fitted to receive the impressions of external bodies: and, according to the difference of these impressions, and of the condition of the sentient extremity itself, to propagate along the nerves motions of a determined kind, which, communicated to the brain, give occasion to sensation. 2. The brain seems to be a part fitted for, and susceptible of, those motions, with which sensation, and the whole consequent operations of thought, are connected: and thereby is fitted to form a communication between the motions excited in the sentient, and those in consequence arising in the moving extremities of the nerves, which are often remote and distant from each other. 3. The moving extremities are so framed as to be capable of contraction, and of having this contraction excited by motion propagated from the brain, and communicated to the contractile fibre. 4. The nerves, more strictly so called, are to

be considered as a collection of medullary fibres, each enveloped in its proper membrane, and thereby so separated from every other, as hardly to admit of any communication of motion from any one to the others, and to admit only of motion along the continuous medullary substance of the same fibre, from its origin to the extremities, or contrariwise. From this view of the parts of the nervous system; of their several functions and communication with each other, it appears, that the beginning of motion in the animal economy is generally connected with sensation: and that the ultimate effects of such motion are chiefly actions depending immediately upon the contraction of moving fibres, between which and the sentient extremities the communication is by means of the brain.

NERVÆ SPONGIOSÆ. The cavernous part of the penis.

NERVI INTERCOSTALES INNOMINATI. The fifth pair of nerves.

NERVINES. (*Nervina*, sc. *medicamenta*, from *nervus*.) Neurotics. Medicines that relieve disorders of the nerves. They are all the antispasmodics, and the various preparations of bark and iron.

NERVORUM RESOLUTIO. A species of apoplexy or palsy.

NERVOSUM OS. The occipital bone.

Nervous consumption. See *Atrophia*.

Nervous diseases. See *Neuroses*.

Nervous Fever. See *Febris nervosa*.

Nervous head ache. See *Cephalalgia*.

NERVOUS FLUID. Nervous principle. The vascularity of the cortical part of the brain, and of the nerves themselves, their softness, pulpiness, and natural humid appearance, give reason to believe that between the medullary particles of which they are principally composed, a fine fluid is constantly secreted, which may be fitted to receive and transmit, even more readily than other fluids do, all impressions which are made on it. It appears to exhale from the extremities of the nerves. The lassitude and debility of muscles from too great exercise, and the dulness of the sensorial organs from excessive use, would seem to prove this. It has no smell nor taste; for the cerebrine medulla is insipid and inodorous. Nor has it any colour, for the cerebrum and nerves are white. It is of so subtle a consistence, as never to have been detected. Its mobility is stupendous; for in less than a moment, with the consent of the mind, it is conveyed from the cerebrum to the muscles, like the electric matter. Whether the nervous fluid be carried from the organ of sense in the sensorial nerves to the cerebrum, and from thence in the motory nerves to the muscles, cannot be positively affirmed, but may be proved. The constituent principles of this liquid are perfectly unknown, as they cannot be rendered visible by art, or proved by experiment. Upon

making a ligature upon a nerve, the motion of the fluid is interrupted, which proves that something corporeal flows through it. It is therefore a weak argument to deny its existence because we cannot see it; for who has seen the matter of heat, oxygen, azot, and other elementary bodies, the existence of which no physician in the present day doubts? The electric matter, whose action on the nerves is very great, does not appear to constitute the nervous fluid; for nerves exhibit no signs of spontaneous electricity; nor can it be the magnetic matter, as the experiment of Gavius with the magnet demonstrates; nor is it oxygen, nor hydrogen, nor azot; for the first very much irritates the nerves, and the other two suspend their action. The nervous fluid, therefore, is an element sui generis, which exists and is produced in the nerves only; hence, like other elements, it is a thing unknown, and only to be known by its effects. The pulposity of some nerves, and their lax situation, does not allow them and the brain to act on the body and soul only by oscillation. Lastly, a tense chord, although tied, oscillates. The use of the nervous fluid is, 1. It appears to be an intermediate substance between the body and the soul, by means of which the latter thinks, perceives, and moves the muscles subservient to the will. Hence the body acts upon the soul, and the soul upon the body. 2. It appears to differ from the vital principle; for parts live and are irritable which want nerves, as bones, tendons, plants, and insects.

NERVIS. (From *νν*, neg. and *ννς*, to eat; so called because it is generally found empty.) The jejunum.

Nettle, common. See *Urtica*.

Nettle, dead. See *Lanum album*.

Nettle-rash. See *Urticaria*.

NEUROCHONDROBES. (From *ννς*, a sinew, and *χνδρς*, a cartilage.) A hard substance between a sinew and a cartilage.

NEUROLOGY. (From *ννς*, a nerve, and *λνγς*, a discourse.) The doctrine of the nerves.

NEURONECTORES. (From *ννς*, a nerve, and *ννς*, a matrix.) The psoas muscles are so called by Fallopius, as being the repository of so many small nerves.

NEUROSES. (From *ννς*, a nerve.) Nervous diseases. The second class of Cullen's nosology is so called; it comprehends affections of sense and motion, disturbed; without either idiopathic pyrexia, or topical diseases.

NEUROTICA. (From *ννς*, a nerve.) Nervous medicines.

NEUROTOMY. (*Neurotomia*, from *ννς*, a nerve, and *ννς*, to cut.) A dissection of the nerves. Also a puncture of a nerve.

NEUTRAL SALTS. Secondary salts. Under the name of neutral or secondary

salts are comprehended such matters as are composed of two primitive saline substances combined together. They are called neutral, because they do not possess the characters of acid nor alkaline salts, which are primitive salts; such are Epsom salts, alum, nitre, &c.

NIXUS. (From *nexus*, wind.) A complication of substances in one part, as the membrane which involves the foetus.

NICKEL. It is to Cronstedt that we are indebted for the discovery of this metal; though the substance from which he extracted it was known in the year 1694. Cronstedt proved it to be a peculiar metal in the year 1751. Nickel is found in nature generally in the metallic state, more rarely in that of an oxyd. Its ores have a coppery red colour, generally covered more or less with a greenish grey efflorescence. The most abundant ore is that termed *sulphuret of nickel*, or *kupfernickel*, which is a compound of nickel, arsenic, sulphuret of iron, and sometimes cobalt and copper. This ore occurs either massive, or disseminated, but never crystallized; it is of a copper colour, sometimes yellowish, white, or grey. It exists also combined with oxygen, and a little carbonic acid, in what is called *native oxyd of nickel*, (*nickel ochre*;) it then has an earthy appearance, and is very friable; it is found coating *kupfernickel*, and seems to originate from the decomposition of this ore. It is found contaminated with iron in the mineral substance called *martial nickel*; this native combination, when fresh broken, has a lamellated texture; when exposed to the air, it soon turns black, and sometimes exhibits thin rhomboidal plates placed irregularly over each other. It is also found united to arsenic, cobalt, and alumine in the ore, called *arseniate of nickel*.

Properties.—Nickel, when free from heterogeneous substances, is of a pale flesh colour. When fresh broken it has a strong lustre. It has a fine-grained compact texture, and can be a little flattened by hammering, similar to cast iron. It leaves a trace when rubbed upon the polished surface of a hard stone. Its specific gravity is 7.380. It requires a very intense heat for fusion. When exposed for a long time to a humid atmosphere, its surface becomes gradually covered with an oxyd of a greenish hue; this takes place likewise and more rapidly, when heated in contact with air. When fused with borax it produces a glass of a hyacinth colour. It unites with phosphorus by fusion, and forms with it a phosphuret which is very fusible, white, and in brilliant needles. With sulphur it forms, by fusion, a hard yellow mass, with small brilliant facets. Sulphuric acid, assisted by heat, dissolves it. Nitric acid acts on it more readily. Muriatic acid, when heated on it, likewise

dissolves part of it. Boracic and phosphoric acids seem to have little or no action on nickel. It readily unites with gold, and renders that metal white and brittle. It likewise fuses with platinum, silver, and bismuth. It does not alloy with mercury. It is easily oxydated by the nitrate and the super-oxygenated muriate of potash. It is *not magnetic*, nor has it the smallest effect upon the magnetic needle. This has been proved by Mr. Chevenix, who has also shown that the magnetism of common nickel is owing to the iron which so obstinately adheres to it. For a portion of iron, so small as not to be detected by the best chemical tests, when combined with nickel, is capable of communicating magnetic properties to the whole mass as strong as if the whole were composed of steel.

Method of obtaining Nickel.—To obtain nickel, the ore is first roasted, in order to free it from sulphur and arsenic; it is then changed into a greenish oxyd. This oxyd is mixed with two or three parts of black flux. The mixture is put into a crucible, and, being covered with decripitated muriate of soda, it is brought to the state of fusion, by the strongest heat of a smith's forge.

When the crucible is broken, there is found at the bottom, under brown, blackish, and sometimes blue soot, a button of a yellowish white colour, equal in weight to a tenth, a fifth, and even a half, of the ore employed. This metal, however, is still far from being pure.

In order to purify it, the button obtained is again broken into small pieces, strongly heated, and then digested with its own weight of concentrated sulphuric acid, and distilled to dryness. The dry mass is dissolved in water, and filtered. This solution, in general, deposits crystals of arsenic, and finally affords dark green crystals of sulphate of nickel. This sulphate is re-dissolved in water, and decomposed by carbonate of potash. The precipitate is dissolved in liquid ammonia; the blue solution leaves a residuum, which is filtered off, and the filtered solution saturated with nitric acid. The nickel is then precipitated in the form of a greyish green powder, by carbonate of potash. From this oxyd the metallic nickel is obtainable by exposing it to heat, when made into a mass with oil and a little charcoal powder.

The nickel obtained in this manner was, until lately, considered as perfectly pure. It possesses, however, magnetic properties. It is therefore contaminated with iron.

In order to obtain this metal in a state of absolute purity, the following method of Chevenix must be had recourse to:

Take the native sulphuret of nickel; reduce it to powder, and roast it in contact with charcoal powder over a gentle fire. When no more fumes arise, pour then nitric acid over it, and dissolve it by heat in

a Florence flask. Decant the solution, filter it through bibulous paper, and evaporate it to dryness in a glass basin. Dissolve the nitrate of nickel in a sufficient quantity of distilled water, and decompose it by the addition of the strongest liquid ammonia, taking care to add it in excess. The oxyd of nickel and cobalt will thus be re-dissolved; then let the solution stand undisturbed till a precipitate again ensues. The solution must then be evaporated; it becomes blue during this process, by the precipitation of the cobalt, which should be separated, and the evaporation be then continued to dryness; the residue will be pure oxyd of nickel.

In order to reduce this oxyd to the metallic state, let it be made into a paste with oil, mix it with about three parts of black flux, and put it into a crucible, covering it with borax and muriate of soda, and heat the crucible violently for an hour and a half in a forge, a button will then be obtained, which is pure nickel.

NICOPHORUS: (From *νικω*, victory, and *φορος*, to bear, so called because victors were crowned with it.)

NICOTIANA. (From Mr. Nicot, who first brought it into Europe.) 1. The name of a genus of plants in the Linnæan system. Class *Pentandria*. Order *Monogynia*. 2. The pharmacopœial name of the official tobacco. *Petum*, by the Indians, *Tabacum*. *Hyosciamus Peruvianus*. *Piscit*. The Virginia tobacco. *Nicotiana tabacum* of Linnæus:—*foliis lanceolato-ovatis sessilibus decurrentibus floribus acutis*, is the plant employed medicinally. It is a very active narcotic and sternutatory. A decoction of the leaves is much esteemed in some diseases of the skin, and is by some said to be a specific against the itch. The fumes and the decoction are employed in obstinate constipations of the bowels, and very frequently with success; it is necessary, however, to caution the practitioner against an effect mostly produced by its exhibition, namely, syncope; with cold sweats; and in some instances death.

NICOTIANA AMERICANA. American or Virginian tobacco. See *Nicotiana*.

NICOTIANA MINOR. Tobacco Anglicum. *Priapeia*. *Hyosciamus luteus*. English tobacco. This plant, *Nicotiana rustica* of Linnæus, is much weaker than the Virginian tobacco, the leaves are chiefly used to smoke vermin, though they promise, from their more gentle operation, to be a safer remedy in some cases than the former.

NICOTIANA RUSTICA. The systematic name of the English tobacco. See *Nicotiana minor*.

NICOTIANA TABACUM. The systematic name of the tobacco-plant. See *Nicotiana*.

NIGELLAS. (Quasi *nigrella*, from

niger, black, so named from its black seed.) 1. The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Pentagynia*. 2. The pharmacopœial name of the *Nigella sativa* of Linnæus. Devil in a bush, or fennel-flower. It was formerly employed medicinally as an expectorant and deobstruent, but is now deservedly fallen into disuse.

NIGELLA SATIVA. The systematic name of the devil in a bush. See *Nigella*.

NIGELLASTRUM. (From *Nigella*, fennel-flower.) *Pseudomelanthium*. *Lychnis segetum major*. *Gothago*. *Nigella officinarum*. *Lychnoides segetum*. Cockle, a herb resembling the nigella.

Night-blindness. See *Nyctalopia*.

NIGHT-MARE. *Incubus*. *Onirodynia gravans*. The nervous or indisposed persons are oppressed during sleep with a heavy pressing sensation on the chest, by which respiration is impeded, or the circulation of blood intercepted, to such a degree as to threaten suffocation. Frightful ideas are recollected on waking, which occupied the dreaming mind. Frequent attempts are made to cry out, but often without effect, and the horrors and agitations felt by the patient are inexpressibly frightful. The sensations generally originate in a large quantity of wind; or indigestible matter in the stomach of *supper-eaters*, which, pressing the stomach against the diaphragm, impede respiration, or render it short and convulsed. Inflated intestines may likewise produce similar effects, or mental perturbations.

There is another species of night-mare mentioned by authors, which has a more dangerous tendency; and this arises from an impeded circulation of blood in the lungs, when lying down, or too great relaxation of the heart and its impelling powers. Epilepsy, apoplexy, or sudden death, are sometimes amongst the consequences of this species of disturbed sleep. Polypi in the large vessels, aneurisms, water in the thorax, pericardium, or lungs, empyema, &c. are amongst the most dangerous causes. See *Onirodynia*.

Nightshade, American. See *Phytolacca decandria*.

Nightshade, deadly. See *Belladonna*.

Nightshade, garden. See *Solanum*.

Nightshade, Palestine. See *Solanum sanctum*.

Nightshade, woody. See *Dulcamara*.

NIGRITIES. (From *niger*, black.) A caries is called *nigrities ossium*, a blackness of the bone.

NIHILUM ALBUM. A name formerly given to the flowers, or oxyd of zinc.

NINSI RADIX *Ninjin*. *Nindsin*. This root was long supposed to be the same as ginseng. It now appears, however, to be the produce of a different plant, the stem

mineral of Linnæus, *foliis serratis, palmatis; ramis ternatis*, possessing similar though weaker properties than ginseng.

NINZIN. See *Ninzi radix*.

NIPPLE. The small projecting portion in the middle of the breasts of men and women. It is much larger in the latter, and has an opening in it, the excretory ducts of the lacteal glands.

Nipple-mort. See *Lampæana*.

NITRAS. (From *nitrum*, nitre.) A nitrate; a salt formed by the union of the nitric acid and a different bases, as the nitrate of potash, soda, silver, &c.

NITRAS AMMONIUM. *Alkali volatile nitratum.* *Sal ammoniacus nitrosus.* *Ammonia nitrosa.* A salt composed of the acid of nitre and ammonia, the virtues of which are irritating, diuretic, and deobstruent; externally, it is resolvent and sialagogue.

NITRAS ARGENTI. *Causticum lunare.* *Argentum nitratum.* "Take of silver an ounce, nitric acid, a fluid-ounce and a half, distilled water, two fluid-ounces." Mix the nitric acid and water, and dissolve the silver therein on a sand bath; then increase the heat gradually, that the nitrate of silver may be dried. Melt the salt in a crucible over a slow fire until, the water being evaporated, it shall cease to boil; then pour it quickly into moulds of convenient shape. Its virtues are corrosive and astringent. Internally it is exhibited in very small quantities, in epilepsy; and externally it is employed to destroy fungous excrescences, callous ulcers, fistulas, &c. In the latter disease it is employed as an injection: from two grains to three being dissolved in an ounce of distilled water.

NITRAS POTASSÆ. See *Nitre*.

NITRAS POTASSÆ FUSUS. *Sal prunella.* *Nitrum tabulatum.* This salt, besides the nitric acid and potash, contains a little sulphuric acid.

NITRAS SODÆ. *Alkali minerale nitratum.* *Nitrum cubicum.* Its virtues are similar to those of nitrate of potash, for which it may be safely substituted.

Nitrate of potash. See *Nitre*.

Nitrate of silver. See *Nitras argenti*.

NITRE. *vilgæ.* *Nitrum.* *Potasse nitras.* *Salpetre.* *Aldurat.* *Alkali.* *Atac.* *Baurack.* *Acuto.* *Hakimistum.* Salt petre. A perfect neutral salt, formed by the union of the nitrous acid with the vegetable alkali. Its taste is cooling, and it does not alter the colour of the syrup of violets. Nitre exists in large quantities in the earth, and is continually formed in inhabited places; it is found in great quantities upon walls which are sheltered from the rain. It is of great use in the arts; it is the principal ingredient in gunpowder; and, burned with different proportions of tartar, forms the substances called fluxes. It is of considerable importance in medicine, as a febrifuge, diuretic, and antiphlogistic remedy.

NITRIC ACID. *Acidum nitricum.* The London College direct this acid to be made by distilling equal parts, by weight, of dried nitrate of potash and sulphuric acid, two pounds of each. Mix them in a glass retort, then distil the nitric acid in a sand bath, until a red vapour arises; lastly, having added to the acid first distilled an ounce more of dry nitrate of potash, distil the nitric acid again in a similar manner.

The specific gravity of nitric acid is to that of water, as 1.600 to 1.000. An ounce diluted with water ought to dissolve of a lump of lime-stone immersed therein seven drachms.

This acid undiluted is a powerful caustic, and is sometimes employed as such by surgeons to destroy fungous excrescences. Very much diluted it is exhibited internally as a tonic and antiseptic, in the cure of typhoid fevers, scurvy, syphiloid diseases, and other cachexias.

Nitric-oxyd of mercury. See *Nitric-oxydum hydrargyri*.

NITRICO-OXYDUM HYDRARGYRI. *Hydrargirus nitratu ruber.* *Mercurius corrosivus ruber.* *Mercurius precipitatus corrosivus.* Nitric-oxyd of mercury. Red precipitate. "Take of purified mercury, by weight, three pounds:—of nitric acid, by weight, a pound and a half:—of distilled water, two pints." Mix in a glass vessel, and boil the mixture in a sand bath, until the mercury being dissolved, the water also evaporated, and a white mass remain. Rub this into powder, and put it into another shallow vessel, then apply a moderate heat, and raise the fire gradually until the red vapour shall cease to rise." This preparation is very extensively employed by surgeons as a stimulant and escharotic, but its extraordinary activity does not allow of its being given internally. Finely levigated and mixed with common cerates, it is an excellent application to indolent ulcers, especially those which remain after burns and scalds, and those in which the granulations are indolent and flabby. It is also an excellent caustic application to chancrea.

NITROGEN. (From *vilgæ*, nitre, and *γεννωμι*, to generate; so called because it is the generator of nitre.) Azote. Alkaligen. A simple body, very abundant in nature, though not producible alone, or in an insaluble state. It is not distinctly perceptible to the human senses, however aided by instruments. We know it only in its combination. But the reality of its existence is unquestionable; since we can mark its passage out of one combination into another; since we know the laws of chemical attraction to which it is subject; since we discern the precise character of those simple substances with which it is combinable, and can distinguish the nature of the new compounds which the combination pro-

duces. The separate existence and peculiar nature of this substance were first discovered by Dr. Rutherford. It is the radical principle of our atmospheric air, and other gaseous substances, and forms a constituent part of animal and vegetable substances. It is a component part of the nitric acid, and of ammonia. It probably enters into the formation of alkalies, and it may be considered as a real alkali- gen or alkalinizing principle, in opposition to oxygen, which, as we have noticed before, is the principle of acidity. One of the most remarkable combinations, into which nitrogen is known to enter, is that which takes place between it and light and caloric. The compound thus produced is called

NITROGEN GAS.

Phlogisticated air. Azotic gas. Mephitic air. Mofette.

Properties.—Nitrogen gas, or azotic gas, as it is also called, is not possessed of any remarkable property capable of characterizing it; but is principally distinguished by certain negative qualities, namely, it is extremely hurtful to respiration, and quickly kills animals. Plants thrive and even flourish in it. It has no sensible taste. It neither reddens blue vegetable colours, nor precipitates lime, or barytic water. Its weight is to common air, as .985 to 1.000. No combustible substance burns in nitrogen gas; but it is capable of combustion in combination with oxygen gas, when exposed to the action of the electric spark. It is not absorbable by water. It is capable of dissolving sulphur, phosphorus and charcoal, in minute quantities. It unites to hydrogen under certain conditions, and constitutes with it ammonia. When united to oxygen in different proportions, it produces atmospheric air, gaseous oxyd of azote or nitrogen, nitrous gas, nitrous acid and nitric acid. It is a component part of all animal substances, and communicates to them their most distinctive characters. It was discovered by Dr. Rutherford of Edinburgh.

Nitrogen gas has been found by Priestley in the Bath waters, and by Dr. Pearson in the Buxton waters.

Methods of obtaining Nitrogen Gas.—

Nitrogen gas may be obtained by various means. For instance, it has been long since ascertained that air, which has served the purposes of combustion and respiration, is no longer proper for these uses. Chymists have availed themselves of this circumstance, in order to obtain nitrogen gas in the following manner.

1 Make a quantity of sulphuret of potash or sulphuret of iron into a paste with water, and place the mixture in a saucer or plate over water, on a stand raised above the fluid; then invert over it a jar or bell-

glass, and allow this to stand for a day or two. The air contained in the bell-glass will gradually diminish, as will appear from the ascent of the water, until only about three-fourths of its original bulk remain.

When no further diminution takes place, the vessel containing the sulphuret must be removed, and the remaining air will be found to be nitrogen gas.

In this experiment the moistened sulphuret of potash has a great affinity to oxygen, it attracts and separates it from the atmospheric air, and the nitrogen gas is left behind; the sulphur is, during the experiment, converted into sulphuric acid, which unites to the alkali and forms sulphate of potash. The water with which the sulphuret is moistened likewise undergoes a decomposition, as shall be more fully proved in another place.

2. Nitrogen gas may likewise be obtained from fresh animal substances. For this purpose cut a piece of lean muscular flesh into small pieces, introduce them into a retort, and pour over them weak nitric acid. If the heat of a lamp not exceeding 80° Fahr. be then gently applied, the gas will be speedily obtained; for all animal substances are composed of nitrogen, carbon, hydrogen, and oxygen; and on adding nitric acid in this way, the equilibrium of the respective affinities is destroyed; the nitrogen gas becoming separated.

That the nitric acid does not furnish the nitrogen gas is obvious; from its saturating after its action as large a quantity of alkali as before, consequently it could not have suffered any alteration.

The fibrous part of animal matter is that which affords the most nitrogen gas; next to this all the concrete parts, such as the clot of blood; next to that, albuminous matter, such as the serum and the white of eggs; gelatinous substances afford the least.

3. Nitrogen gas may likewise be obtained, by causing oxygenated muriatic acid gas to be received in a vessel containing liquid ammonia, for ammonia consists of hydrogen and nitrogen. The hydrogen of the ammonia unites to the oxygen of the oxygenated muriatic acid, and forms water, heat is evolved, the nitrogen becomes free, and the oxygenated muriatic acid becomes converted into simple muriatic acid.

Nitrogen gaz. See *Azet*.

NITROGEN, GAZEUS OXYD OF.

This combination of nitrogen and oxygen was formerly called the dephlogisticated nitrous gaz, but now gaseous oxyd of nitrogen, or nitrous oxyd. It was first discovered by Priestley. Its nature and properties have since been investigated (though not very accurately) by a society of Dutch chymists.

Professor Davy has examined with uncommon accuracy the formation and pro-

erties of all the substances concerned in its production. He has detected the sources of error in the experiments of Priestley and the Dutch chemists, and to him we are indebted for a thorough knowledge of this gas. We shall therefore exhibit the philosophy of this gaseous fluid as we find it in his researches concerning the nitrous oxid.

Properties—It exists in the form of permanent gas. A candle burns with a brilliant flame and crackling noise in it; before its extinction the white inner flame becomes surrounded with a blue one. Phosphorus introduced into it, in a state of actual inflammation, burns with increased splendour, as in oxygen gas. Sulphur introduced into it, when burning with a feeble blue flame is instantly extinguished; but when in a state of vivid inflammation, it burns with a rose-coloured flame. Ignited charcoal burns in it more brilliantly than in atmospheric air. Iron wire, with a small piece of wood affixed to it, when inflamed, and introduced into a vessel filled with this gas, burns vehemently, and throws out bright scintillating sparks. No combustible body however burns in it, unless it be previously brought to a state of vivid inflammation. Hence sulphur may be melted, and even sublimed in it, phosphorus may be liquified in it without undergoing combustion. Nitrous oxid is pretty rapidly absorbed by water that has been boiled; a quantity of gas equal to rather more than half the bulk of the water may be thus made to disappear, the water acquires a sweetish taste, but its other properties do not differ perceptibly from common water. The whole of the gas may be expelled again by heat. It does not change blue vegetable colours. It has a distinctly sweet taste, and a faint but agreeable odour. It undergoes no diminution when mingled with oxygen or nitrous gas. Most of the liquid inflammable bodies, such as ether, alcohol, volatile and fat oils, absorb it rapidly and in great quantity. Acids exert but little action on it. The affinity of the neutro-saline solutions for gaseous oxid of nitrogen is very feeble. Green muriate and green sulphate of iron, whether holding nitrous gas in solution, or not, do not act upon it. None of the gases when mingled with it, suffer any perceptible change at common temperatures; the muriatic and sulphurous acid gases excepted, which undergo a slight expansion. Alkalies freed from carbonic acid, exposed in the dry or solid form, have no action upon it; they may however be made to combine with it in the nascent state, and then constitute saline compounds of a peculiar nature. These combinations deflagrate when heated with charcoal, and are decomposed by acids; the gaseous oxid of nitrogen being disengaged. It undergoes

no change whatever from the simple effect of light. The action of the electric spark, for a long while continued, converts it into a gas, analogous to atmospheric air and nitrous acid; the same is the case when it is made to pass through an ignited earthen tube. It explodes with hydrogen in a variety of proportions, at very high temperatures; for instance, when electric sparks are made to pass through the mixture. Sulphureted, heavy and light carbonated hydrogen gases, and gaseous oxid of carbon likewise burn with it when a strong red heat is applied. 100 parts by weight of nitrous oxid, contain 36.7 of oxygen and 63.3 of nitrogen; 100 cubic inches weigh 50 grains at 55° temperature and 30 atmospheric pressure. Animals, when wholly confined in gaseous oxid of nitrogen, give no signs of uneasiness for some moments, but they soon become restless and then die. When gaseous oxid of nitrogen is mingled with atmospheric air, and then received into the lungs, it generates highly pleasurable sensations; the effects it produces on the animal system are eminently distinguished from every other chemical agent. It excites every fibre to action, and rouses the faculties of the mind, inducing a state of great exhilaration, an irresistible propensity to laughter, a rapid flow of vivid ideas, and unusual vigour and fitness for muscular exertions, in some respects resembling those attendant on the pleasantest period of intoxication, without any subsequent languor, depression of the nervous energy, or disagreeable feelings; but more generally followed by vigour, and a pleasurable disposition to exertion, which gradually subsides.

Such are the properties that characterize the nitrous oxid.

The Dutch chemists and some French and German philosophers assert that it cannot be respired; that burning phosphorus, sulphur, and charcoal are extinguished in it, &c. It is probable they did not examine it in a state of purity, for it is otherwise difficult to account for these and many other erroneous opinions.

Methods of obtaining gaseous oxid of nitrogen.—Gaseous oxid of nitrogen is produced, when substances, having a strong affinity with oxygen are brought into contact with nitric acid, or with nitrous gas. It may therefore be obtained by various processes, in which nitrous gas or nitric acid is decomposed by substances capable of attracting the greater part of their oxygen. The most commodious and expeditious as well as cheapest mode of obtaining it, is by decomposing nitrate of ammonia, at a certain temperature, in the following manner:—

1. Introduce into a glass retort some pure nitrate of ammonia, and apply the heat of an Argand's lamp, the salt will

soon liquefy, and, when it begins to boil, gas will be evolved. Increase the heat gradually till the body and neck of the retort become filled with a semi-transparent milky-white vapour. In this state the temperature of the fused nitrate is between 340° and 480° . After the decomposition has proceeded for a few minutes, so that the gas evolved quickly enlarges the flame of a taper held near the orifice of the retort, it may be collected over water, care being taken during the whole process, never to suffer the temperature of the fused nitrate to rise above 500° Fahr. which may easily be judged of, from the density of the vapours in the retort, and from the quiet ebullition of the fused nitrate; for if the heat be increased beyond this point, the vapours in the retort acquire a reddish and more transparent appearance: and the fused nitrate begins to rise, and occupy twice the bulk it did before. The nitrous oxid, after its generation, is allowed to stand over water, for at least six hours, and is then fit for respiration or other experiments.


Explanation.—Nitrate of ammonia consists of nitric acid and ammonia; nitric

acid is composed of nitrous gas and oxygen; and ammonia consists of hydrogen and nitrogen. At a temperature of about 480° the attractions of hydrogen for nitrogen in ammonia, and that of nitrous gas for oxygen in nitric acid, are *diminished*; while, on the contrary, the attractions of the hydrogen of ammonia for the oxygen of the nitric acid, and that of the remaining nitrogen of the ammonia for the nitrous gas of the nitric acid, are *increased*; hence all the former affinities are broken, and new ones produced, namely, the hydrogen of the ammonia attracts the oxygen of the nitric acid, the result of which is *water*; the nitrogen of the ammonia combines with the liberated nitrous gas, and forms *nitrous oxid*. The water and nitrous oxid produced, probably exist in binary combination in the æriform state, at the temperature of the decomposition.

Such is the philosophy of the production of gaseous oxid of nitrogen, by decomposing nitrate of ammonia at that temperature, given by Davy.

To illustrate this complicated play of affinity more fully, the following sketch may not be deemed superfluous.

A Diagram exhibiting the production of Gaseous Oxid of Nitrogen by decomposing Nitrate of Ammonia, at 480° Fahr.



Professor Davy has likewise pointed out, that, when the heat employed for decomposing nitrate of ammonia is raised above the before-stated temperature, another play of affinities takes place, the attractions of nitrogen and hydrogen for each other, and of oxygen for nitrous gas, are still more diminished, whilst that of ni-

trogen for nitrous gas is totally destroyed, and that of hydrogen for oxygen increased to a greater extent. A new attraction likewise takes place, namely, that of nitrous gas for nitric acid to form *nitrous vapour*, and a new arrangement of principles is rapidly produced: the nitrogen of the ammonia having no affinity for any of the

single principles at this temperature, enters into no binary compound; the oxygen of the nitric acid forms water with the hydrogen, and the nitrous gas combines with the nitric acid to form *nitrous vapour*.

All these substances most probably exist in combination, at the temperature of their production; and at a lower temperature assume the form of *nitrous acid*, *nitrous gas*, *nitrogen*, and *water*; and hence we see the necessity of not heating the nitrate of ammonia above the before-stated temperature.

On account of the rapid absorption of gaseous oxid of nitrogen by water, it is economical to preserve the fluid which has been used to confine this gas, and to make use of it for collecting other quantities of it. In order to hasten its production, the nitrate of ammonia may be previously freed from its water of crystallization by gently fusing it in a glass or Wedgewood's basin for a few minutes, and then keeping it for use in a well-stopped bottle.

2. Nitrous oxid may likewise be obtained by exposing common nitrous gas to alkaline sulphates, particularly to sulphite of potash containing its full quantity of water of crystallization. The nitrous oxid produced from nitrous gas by sulphite of potash has all the properties of that generated from the decomposition of nitrate of ammonia.

The conversion of nitrous gas into nitrous oxid by these bodies, depends on the abstraction of a portion of its oxygen by the greater affinity of the sulphite presented to it. The nitrogen and remaining oxygen assume a more condensed state of existence, and constitute nitrous oxid.

3. Nitrous oxid may also be obtained by mingling together nitrous gas and sulphurated hydrogen gas. The volume of gases in this case is diminished, sulphur deposited, ammonia, water, and nitrous oxid are formed.

The change of principles which takes place in this experiment depends upon the combination of the hydrogen of the sulphurated hydrogen gas, with different portions of the oxygen and nitrogen of the nitrous gas, to form water and ammonia, and to deposit sulphur. The remaining oxygen and nitrogen being left in due proportion constitute nitrous oxid.

Remark.—This singular exertion of attraction by a simple body appears highly improbable *a priori*; but the formation of ammonia, and the non-oxygenation of the sulphur, elucidate the fact. In performing this experiment care should be taken that the gases should be rendered as dry as possible; for the presence of water considerably retards the decomposition.

4. Nitrous oxid may also be produced by presenting alkaline sulphurets to nitrous

gas. Davy observed that a solution of sulphuret of strontia, or barytes, answers this purpose best.

This decomposition of nitrous gas is not solely produced by the abstraction of oxygen from the nitrous gas, to form sulphuric acid. It depends equally on the decomposition of the sulphurated hydrogen dissolved in the solution or liberated from it. In this process sulphur is deposited and sulphuric acid formed.

5. Nitrous oxid is obtained in many circumstances similar to these in which nitrous gas is produced. Dr. Priestley found that nitrous oxid was evolved, together with nitrous gas, during the solution of iron, tin, and zinc in nitrous acid.

It is difficult to ascertain the exact rationale of these processes, for very complicated agencies of affinities take place. Either the nascent hydrogen arising from the decomposition of the water by the metallic substance may combine with portions of the oxygen and nitrogen of the nitrous gas, and thus by forming water and ammonia, convert into nitrous oxid; or the metallic substance may attract at the same time oxygen from the water and nitrous gas, whilst the nascent hydrogen of the water seizes upon a portion of the nitrogen of the nitrous gas, to form ammonia. The analogy between this process and the decomposition of nitrous gas by sulphurated hydrogen, render the first opinion most probable.

Such are the principal methods of obtaining nitrous oxid. There are no reasons, Davy thinks, for supposing that nitrous oxid is formed in any of the processes of nature, and the nice equilibrium of affinity by which it is constituted forbids us to hope for the power of composing it from its simple principles. We must be content to produce it artificially.

NITRO-MURIATIC ACID. The compound acid formed by uniting the nitrous and marine acids. It is commonly known by the name of aqua regia. See *Oxygenated muriatic acid*.

Nitrous acid. See *Acidum nitrosum*.

NITROUS GAS. The name of nitrous gas is given to an æriform fluid, consisting of a certain quantity of nitrogen gas and oxygen. It is an elastic, colourless fluid, having no sensible taste; it is neither acid nor alkaline; it is exceedingly hurtful to animals, producing instant suffocation, whenever they attempt to breathe it. The greater number of combustible bodies refuse to burn it. It is nevertheless capable of supporting the combustion of some of these bodies. Phosphorus burns in nitrous gas when introduced into it in a state of inflammation; pyrophorus takes fire in it spontaneously.

It is not decomposable by water, though 100 cubic inches of this fluid, when freed from air, absorb about 11.8 cubic inches

of gas. This solution is void of taste; it does not redden blue vegetable colours; the gas is expelled again when the water is made to boil or suffered to freeze. It has no action on nitrogen gas even when assisted by heat. It is decomposed by several metals at high temperatures.

Its specific gravity, when perfectly pure, is to that of atmospheric air as 1.093 to 1.

Ardent spirit, saccharine matters, hydrocarbonates, sulphurous acid and phosphorus have no action on it. It is not sensibly changed by the action of light. Heat dilates it. It rapidly combines with oxygen gas at common temperatures, and converts it into nitrous acid. Atmospheric air produces the same effect but with less intensity. It is absorbable by green sulphate, muriate and nitrate of iron, and decomposable by alkaline, terrene, and metallic sulphurets, and other bodies that have a strong affinity for oxygen; but it is not capable of combining with them chemically, so as to form saline compounds. From the greatest number of bodies which absorb it, it may be again expelled by the application of heat.

It communicates to flame a greenish colour before extinguishing it; when mixed with hydrogen gas it acquires the property of burning with a green flame. It is absorbable by nitric acid and renders it fuming.

When exposed to the action of caloric in an ignited porcelain tube, it experiences no alteration, but when electric sparks are made to pass through it, it is decomposed and converted into nitrous acid, and nitrogen gas. Phosphorus does not shine in it. It is composed of 56 parts of oxygen and 44 nitrogen.

Methods of obtaining nitrous gas.—

1. Put into a small proof, or retort, some copper wire or pieces of the same metal, and pour on it nitric acid of commerce diluted with water, an effervescence takes place and nitrous gas will be produced. After having suffered the gas to escape for a few minutes on account of the atmospheric air contained in the retort, collect the gas in the water-apparatus as usual. In order to obtain the gas in a pure state, it must then be shook for some time in contact with water. The water in this instance suffers no alteration, on the contrary, the acid undergoes a partial decomposition; the metal robs the nitric acid of the greatest part of its oxygen and becomes oxidated; the remainder of the acid having lost so much of its oxygen, becomes thereby so altered, that at the usual temperature it can exist no longer in the liquid state, but instantly expands and assumes the form of gas; ceasing at the same time to act as an acid, and exhibiting different properties.

Instead of presenting copper to nitric acid, iron, zinc, mercury, or silver may be made use of. The metals best suited for the production of nitrous gas are silver, mercury, and copper.

2. Nitrous gas may likewise be obtained by synthesis. This method of obtaining it we owe to Dr. Milner of Cambridge.

Into the middle of an earthen tube about 20 inches long and three-fourth wide, open at both ends, put as much coarsely-powdered manganese as is sufficient nearly to fill it. Let this tube traverse a furnace having two openings opposite to each other. To one end of the tube lute a retort containing water strongly impregnated with ammonia, and to the other adapt a bent glass tube which passes into the pneumatic trough. Let a fire be kindled in the furnace, and when the manganese may be supposed to be red-hot, apply a gentle heat to the retort and drive over it the vapour of the ammonia; the consequence will be that nitrous gas will be delivered at one end of the tube, at the time that the ammonia enters the other end; and this effect does not take place without the presence of the alkali.

Explanation.—Ammonia consists of hydrogen and nitrogen, its hydrogen combines with the oxygen which is given out by the ignited manganese, and forms water, its nitrogen unites at the same time to another portion of the oxygen, and constitutes the nitrous gas.

There is a cause of deception in this experiment, against which the operator ought to be on his guard, lest he should conclude no nitrous gas is formed, when in reality there is a considerable quantity. The ammonia, notwithstanding every precaution, will frequently pass over undecomposed. If the receiver in the pneumatic trough is filled with water, great part of this will indeed be presently absorbed; but still some portion of it will mix with the nitrous gas formed in the process. Upon admitting the atmospheric air, the nitrous gas will become decomposed, and the red nitrous fumes instantly unite with the alkali. The receiver is presently filled with white clouds of nitrate of ammonia: and in this manner a wrong conclusion may easily be drawn from the want of the orange colour of the nitrous fumes. A considerable quantity of nitrous gas may have been formed, and yet no orange colour appear, owing to this circumstance; and therefore it is easy to understand how small a quantity of nitrous gas may be most effectually disguised by the same cause.

Dr. Milner also obtained nitrous gas, by passing ammoniacal gas over sulphate of iron deprived of its water of crystallization,

Water impregnated with this gas forms
NITROUS ACID.

The common mode of obtaining nitrous acid is to decompose nitrate of potash by means of sulphuric acid with the assistance of heat. The nitric acid suffers a partial decomposition during the process, and hence it is the nitrous acid which is obtained in the first process of distillation.

It seems to be true that nitrous acid of a much darker orange red colour is obtained by decomposing nitrate of potash by means of sulphate of iron, than when the same salt is decomposed by sulphuric acid. The following is the process made use of by our manufacturers.

Take a quantity of sulphate of iron, deprived of its water of crystallization by heat, and mix it with an equal weight of dry nitrate of potash; put the mixture into a glass retort, to which a very spacious receiver has been luted, containing a little water, and begin the distillation with a very slow fire. As soon as the red vapours cease to come over, let the fire be slackened, and when the vessels are cooled, the receiver may be cautiously withdrawn, and its contents quickly transferred through a glass funnel into a bottle, furnished with a ground stopper.

NITROUS OXID. See *Nitrogen, gaseous acid of*.

NITRUM. See *Natron and nitre*.

NITRUM PURIFICATUM. See *Nitre*.

NITRUM VITRIOLATUM. Sulphuric acid and soda. See *Sulphate of soda*.

NOBILIS. (*Quæ nobilis*, from *nosco*, to know.) The heart, by way of eminence, is called *nobilis valvula*, the noble valve.

NOBLE METALS. A name formerly bestowed on the perfect metals, gold, silver, and platina.

NOCTAMBULATIO. (From *nax*, night, and *ambulo*, to walk.) *Noctisurgium*. Walking in the night, when asleep. See *Oneirodynia*.

NOCTISURGIUM. See *Noctambulatio*.

Nocturnal emissions. See *Genorrhœa dormientium*.

NODDING ONICUS. The systematic name of this plant is *Cnicus cernuus* of Linnaeus. In Siberia the tender stalks are first peeled and then boiled and eaten by the inhabitants.

NODE. *Nodus*. A hard circumscribed tumour, proceeding from a bone, and caused by a swelling of the periosteum; they appear on every part of the body, but are more common on such as are thinly covered with muscles, as the os frontis, forepart of the tibia, radius and ulna. As they increase in size they become more painful

from the distention they occasion in the periosteum. When they continue long the bone becomes completely carous.

NOVUS. (From *Anad*, to tie, Heb.) A node or swelling upon a bone. See *Node*.

NOLI ME TANGERE. A species of herpes affecting the skin and cartilages of the nose, very difficult to cure, because it is exasperated by most applications. The disease generally commences with small, superficial spreading ulcerations on the alæ of the nose, which become more or less concealed beneath furfuraceous scabs. The whole nose is frequently destroyed by the progressive ravages of this peculiar disorder, which sometimes cannot be stopped or retarded by any treatment, external or internal.

NOXA. (From *noxi*, to eat.) *Noma*. Ulcers that sometimes attack the cheek or vulva of young girls. They appear in the form of red and somewhat livid spots; are not attended with pyrexia, pain, or tumour, and in a few days become gangrenous.

NON-NATURALS. Under this term, ancient physicians comprehend air, meat and drink, sleep and watching, motion and rest, the retentions and excretions, and the affections of the mind; or, in other words, those principal matters which do not enter into the composition of the body, but at the same time are necessary to its existence.

NOXUS. (*Quasi novenus*, from *novem*, nine.) *Humeri musculus placentini*. The ninth or coracoid muscle of the shoulder.

NOFAL. *Nopalnochetzth*. The plant that feeds the cochineal insect.

NORLANDICUS BACCÆ. The fruit of the *rubus arcticus* of Linnaeus, which this illustrious character found very grateful and refreshing in his tour through the northern part of Sweden. In putrid fever, exanthematous diseases, and scurvy, they promise to be, like other summer fruits, very serviceable.

NOSE. *Nasus*. See *Nares*.

Nose, bleeding of. See *Epistaxis*.

NOSOCOMIUM. (From *nosos*, a disease, and *κομειν*, to take care of.) *Nosodychium*. An hospital.

Nosodochium. See *Nosocomium*.

NOSOLOGY. (*Nosologia*, from *nosos*, a disease, and *λογος*, a discourse.) The doctrine of the names of diseases. Modern physicians understand by nosology the arrangement of diseases in classes, genera, species, &c. The following are the approved arrangements of the several nosologists. That of Dr. Cullen is generally adopted in this country, and next to it the arrangement of Sauvage.

Synoptical View of the Classes, Orders, and Genera, according to the CULLENTIAN system.

ORDER I. FEBRES.**§ 1. Intermittentes.**

1. Tertianæ
2. Quartanæ
3. Quotidianæ.

§ 2. Continuas.

4. Synocha
5. Typhus
6. Synochus.

ORDER H.**PHLEGMASIE.**

7. Phlogosis
8. Ophthalmia
9. Phrenitis
10. Cynanche
11. Pneumonia
12. Carditis.

ORDER I. COMATA.

41. Apoplexia
42. Paralysis.

ORDER II.**ADYNAMIE.**

43. Syncope
44. Dyspepsia
45. Hypochondriasis
46. Chlorosis.

ORDER III. SPASMI.

47. Tetanus

ORDER I.**MARCORES.**

68. Tabes
69. Atrophia

ORDER II.**INTUMESCENTIÆ.****§ 1. Adiposa.**

70. Polysarcia.

§ 2. Flutuosa.

71. Pneumatosis,
72. Tympanites

ORDER I.**DYÆSTHESIE.**

91. Caligo
92. Amaurosis
93. Dysopia
94. Pseudoblepsis
95. Dysecocæ
96. Paracusis
97. Anosmia
98. Agheusia
99. Anæsthesia.

ORDER II.**DYSOREXIE.****§ 1. Appetitus erronei.**

100. Bulimia
101. Polydipsia
102. Pica
103. Satyriasis
104. Nymphomania
105. Nostalgia

CLASS I.—PYREXIE.

13. Peritonitis
14. Gastritis
15. Enteritis
16. Hepatitis
17. Splenitis
18. Nephritis
19. Cystitis
20. Hysteritis
21. Rheumatismus
22. Odontalgia
23. Podagra
24. Arthropæosis.

ORDER III.**EXANTHEMATA.**

25. Variola
26. Varicella

CLASS II.—NEUROSES.

48. Trismus
49. Convulsio
50. Chorea
51. Raphania
52. Epilepsia
53. Palpitatio
54. Asthma
55. Dyspnœa
56. Pertussis
57. Pyrosis
58. Cholica

CLASS III.—CACHEXIE.

73. Physometra
- § 3. Aquosa.**
74. Anasarca
75. Hydrocephalus
76. Hydrorachitis
77. Hydrothorax
78. Ascites
79. Hydrometra
80. Hydrocele.
- § 4. Solide.**
81. Physconia

CLASS IV.—LOCALES.

- § 2. Appetitus deficientes.**
106. Anorexia
107. Adipsia
108. Anaphrodisia

ORDER III.**DYSCINESIE.**

109. Aphonia
110. Mutitas
111. Paraphonia
112. Pællismus
113. Strabismus
114. Dysphagia
115. Contractura

ORDER IV.**APOCENOSES.**

116. Profusio
117. Ephidrosis
118. Epiphora
119. Ptyalismus
120. Eneuresis

27. Rubeola
28. Scarlatina
29. Pestis
30. Erysipelas
31. Miliaria
32. Urticaria
33. Pemphigus
34. Aphtha.

ORDER IV.**HÆMORRHAGIE.**

35. Epistaxis
36. Hæmoptysis
37. Hæmorrhoids
38. Menorrhagia
39. Catarrhus
40. Dysenteria.

ORDER IV.**VESANIÆ.**

64. Amentia
65. Melancholia
66. Mania
67. Oneirodynia.

82. Rachitis.

ORDER III.**IMPETIGINES.**

83. Scrophula
84. Syphilis
85. Scorbutus
86. Elephantiasis
87. Lepra
88. Frambæsia
89. Tricoma
90. Icterus.

121. Gonorrhœa.

ORDER V.**EPISCHESES.**

122. Obstipatio
123. Ischuria
124. Dysuria
125. Dyspermatismus
126. Amenorrhœa.

ORDER VI.**TUMORES.**

127. Aneurisma
128. Varix
129. Ecchymoma
130. Schirrus
131. Cancer
132. Bubo
133. Sarcoma
134. Veruca
135. Clavus
136. Lupia

137. Ganglion
138. Hydatid
139. Hydatidus
140. Exostosis.
ORDER VII.
ECTOPIÆ.
141. Hernia

142. Prolapsus
143. Luxatio
ORDER VIII.
DIALYSES.
144. Vulnus
145. Ulcus

146. Herpes
147. Tinea
148. Psora
149. Fractura
150. Caries

Synoptical View of the System of SAUVAGES.

CLASS I.—VITIA.

ORDER I. MACULÆ.

- Genus 1. Leucoma
2. Vitiligo
3. Ephelis
4. Gutta rosea
5. Nevus
6. Ecchymoma.

ORDER II.

EFFLORESCENTIÆ.

7. Herpes
8. Epinictis
9. Psudracia
10. Hydroa.

ORDER III. PHYMATA.

11. Erythema
12. Oedema
13. Emphysema
14. Schirrus
15. Phlegmone
16. Bubo
17. Parotis
18. Furunculus
19. Anthrax
20. Cancer
21. Paronichia
22. Phymosis.

ORDER IV.

EXCRESCENTIÆ.

23. Sarcoma
24. Condyloma
25. Verruca

26. Pterygium
27. Hordeolum
28. Bronchocele
29. Exostosis
30. Gibbositas
31. Lordosis.

ORDER V.

CYSTIDES.

32. Aneurisma
33. Varix
34. Hydatid
35. Mariaca
36. Staphyloma
37. Lupia

38. Hydrarthrus
39. Apostema
40. Exomphalus
41. Oscheocele

ORDER VI.

ECTOPIÆ.

42. Exophthalmia
43. Blepharoptosis
44. Hypostophyle
45. Paraglossa.

ORDER VI.

ECTOPIÆ.

46. Proptoma
47. Exania
48. Foxocyste
49. Hysteroptosis
50. Euterocele

51. Epiplocele
52. Gasterocele
53. Hepatocele
54. Splenocele
55. Hysterocele
56. Cystocele
57. Encephalocele
58. Hysteroloxia
59. Parochydium
60. Exarthrema
61. Diastasis
62. Laxarthrus.

ORDER VII.

PLAGÆ.

63. Vulnus
64. Punctura
65. Excoriatio
66. Contusio
67. Fractura
68. Fissura
69. Ruptura
70. Amputatura
71. Ulcus
72. Exulceratio
73. Sinus
74. Fistula
75. Rhagus
76. Eschara
77. Caries
78. Arthrocace.

CLASS II.—FEBRES.

83. Hectica

ORDER II.

REMITTENTES.

84. Amphimerina
85. Tritæophya
86. Tetartophya.

ORDER III.

INTERMITTENTES.

87. Quotidiana
88. Tertiana
89. Quartana
90. Erratica.

ORDER I.
CONTINUÆ.

79. Ephemera
80. Synocha
81. Synochus
82. Typhus

ORDER I.

EXAMTHEMATICÆ.

91. Pestis
92. Variola
93. Pemphigus
94. Rubecula
95. Miliaris
96. Purpura
97. Erysipelas
98. Scarlatina
99. Eosera

CLASS III.—PHLEGMASÆ.

100. Aphtha.

ORDER II.

MEMBRANACEÆ.

101. Phrenitis
102. Paraphrenesis
103. Pleuritis
104. Gastritis
105. Enteritis
106. Epiploitis
107. Metritis
108. Cystitis.

ORDER III.

PARENCHYMATOSÆ.

109. Cephalitis
110. Cynanche
111. Carditis
112. Peripneumonia
113. Hepatitis
114. Splenitis
115. Nephritis.

ORDER I.
TONICI PARTIALES.

- 116. Strabismus
- 117. Trismus
- 118. Obstipitas
- 119. Contractura
- 120. Crampus
- 121. Priapismus

ORDER II.
TONICI GENERALES.

- 122. Tetanus

ORDER I.
SPASMODICÆ.

- 138. Ephialtes
- 139. Sternutatio
- 140. Oscedo
- 141. Singultus

ORDER I.
DYSÆSTHESIÆ.

- 152. Cataracta
- 153. Caligo
- 154. Amblyopia
- 155. Amaurosis
- 156. Anosmia
- 157. Agheusia
- 158. Dysecœa
- 159. Paracusis
- 160. Cophosis
- 161. Anæsthesia

ORDER II.
ANEPITHYMIÆ.

- 162. Anorexia

ORDER I. VAGI.

- 183. Arthritis
- 184. Ostocopus
- 185. Rhumatismus
- 186. Catarrhus
- 187. Anxietas
- 188. Lassitudo
- 189. Stupor
- 190. Pruritus
- 191. Algor
- 192. Ardor

ORDER II. CAPITIS.

- 193. Cephalalgia
- 194. Cephalæa

ORDER I.
HALLUCINATIONES.

- 216. Vertigo
- 217. Suffusio
- 218. Diplopia
- 219. Syrosmas
- 220. Hypochondriasis
- 221. Somaambulismus.

ORDER II
MOROSITATES.

- 222. Pica

CLASS IV.—SPASMI.

- 123. Catochus.

ORDER III.
CLONICI PARTIALES.

- 124. Nystagmus
- 125. Carphologia
- 126. Pandiculatio
- 127. Apomyttosis
- 128. Convulsio
- 129. Tremor
- 130. Palpitatio

- 131. Claudicatio.

ORDER IV.
CLONICI GENERALES.

- 132. Rigor
- 133. Eclampsia
- 134. Epilepsia
- 135. Hysteria
- 136. Scelotyrbe
- 137. Beriberia.

CLASS V.—ANHELATIONES

- 142. Tussis.

ORDER II.
OPPRESSIVÆ.

- 143. Stertor
- 144. Dyspnœa
- 145. Asthma

- 146. Orthopnœa

- 147. Angina

- 148. Pleurodynæ

- 149. Rhuma

- 150. Hydrothorax

- 151. Empyema.

CLASS VI.—DEBILITATES.

- 163. Adipsia

- 164. Anaphrodisia.

ORDER III.
DYSCINESIÆ.

- 165. Mutitas
- 166. Aphonia
- 167. Psellismus
- 168. Paraphonia
- 169. Paralysis
- 170. Hemiplegia
- 171. Paraplexia.

ORDER IV.
LEIPOPSYCHIE.

- 172. Asthenia.

- 173. Leipothymia

- 174. Syncope

- 175. Asphyxia

ORDER V.
COMATA.

- 176. Catalepsia

- 177. Ecstasis

- 178. Typhomania

- 179. Lethargus

- 180. Cataphora

- 181. Carus

- 182. Apoplexia.

CLASS VII.—DOLORES.

- 195. Hemicrania

- 196. Ophthalmia

- 197. Otalgia

- 198. Odontalgia

ORDER III.
PECTORIS.

- 199. Dysphagia

- 200. Pyrosis

- 201. Cardiognus.

ORDER IV.
ABDOMINALES IN-
TERNI.

- 202. Cardialgia

- 203. Gastrodynia

- 204. Colica

- 205. Hepthalgia

- 206. Splenalgia

- 207. Nephralgia

- 208. Dystocia

- 209. Hysteralgia

ORDER V.

EXTERNI ET ARTUUM.

- 210. Mastodynus.

- 211. Rachialgia

- 212. Lumbago

- 213. Ischias

- 214. Proctalgia

- 215. Pudendagra

CLASS VIII.—VESANIAE.

- 223. Bulimia

- 224. Polydipsia

- 225. Antipathia

- 226. Nostalgia

- 227. Panophobia

- 228. Satyriasis

- 229. Nymphomania

- 230. Tarantismus

- 231. Hydrophobia

ORDER III. DELIRIA.

- 232. Paraphresyne

- 233. Amentia

- 234. Melancholia

- 235. Mania

- 236. Demonomania.

ORDER IV.

VESANIAE ANOMA-
LÆ.

- 237. Amnesia

- 238. Agrypnia.

CLASS IX.—FLUXUS.

ORDER I.
SANGUIFLUXUS.

- 239 Hæmorrhægia
- 240 Hæmoptysis
- 241. Stomacace
- 242. Hæmatemesis
- 243. Hæmaturia
- 244 Menorrhægia
- 245. Abortus.

ORDER II.
ALVIFLUXUS.

- 246. Hepatorrhœa
- 247 Hæmorrhoids
- 248. Dysenteria
- 249. Melæna

ORDER I.
MACIES.

- 275. Tabes
- 276 Phthisis
- 277. Atrophia
- 278. Aridura.

ORDER II.
INTUMESCENTIÆ.

- 279. Polysarcia
- 280. Pneumatosis
- 281. Anasarca
- 282. Phlegmatia
- 283. Physconia
- 284. Graviditas.

ORDER III.
HYDROPE PARTIALES.

- 285. Hydrocephalus
- 286. Physocephalus

- 250. Nausea
- 251. Vomitus
- 252. Ileus
- 253. Cholera
- 254. Diarrhœa
- 255. Cæliaca
- 256. Lienteria
- 257. Tenesmus.

ORDER III.
SERIFLUXUS.

- 258. Ephidrosis
- 259 Epiphora
- 260 Coryza
- 261. Ptyalismus
- 262. Avacatharsis

CLASS X.—CACHEXIÆ.

- 287. Hydrorachitis
- 288. Ascites
- 289. Hydrometra
- 290. Physometra
- 291. Tympanites
- 292. Metrorismus
- 293. Ischuria.

ORDER IV.
TUBERA.

- 294. Rachitis
- 295. Scrophula
- 296. Carcinoma
- 297. Leontiasis
- 298. Malis
- 299. Framboesia.

ORDER V.
IMPETIGINES.

- 300. Syphilis
- 301. Scorbutus

- 263. Diabetes
- 264. Eneuresis
- 265. Dysuria
- 266. Pyuria
- 267. Leucorrhœa
- 268. Gonorrhœa
- 269. Dyspermatismus
- 270. Galactorrhœa
- 271. Octorrhœa.

ORDER IV.
ABRIFLUXUS.

- 272. Flatulenta
- 273. Aëdopsophia
- 274. Dysodia.

- 302. Elephantiasis
- 303. Lepra
- 304. Scabies
- 305. Tinea.

ORDER VI.
ICTERITIAÆ.

- 306. Aurigo
- 307. Melasicterus
- 308. Phænigmus
- 309. Chlorosis.

ORDER VII.
CACHEXIÆ ANOMALÆ.

- 310. Phthiriasis
- 311. Trichoma
- 312. Alopecia
- 313. Eleosis
- 314. Gangræna
- 315. Necrosis

Synoptical View of the System of LINNÆUS.

CLASS I.—EXANTHEMATICI

ORDER I.
CONTAGIOSI.

- 1. Morta.
- 2. Pestis
- 3. Variola
- 4. Rubeola

- 5. Petechia
- 6. Syphilis.

ORDER II.
SPORADICI

- 7. Miliaria

- 8. Uredo
- 9. Aphthæ.

ORDER III.
SOLITARIÆ.

- 10. Erysipelas.

CLASS II.—CRITICI.

ORDER II.
INTERMITTENTES.

- 15. Quotidiana
- 16. Tertiana
- 17. Quartana
- 18. Duplicana
- 19. Errana.

ORDER III.
EXACERBANTES.

- 20. Amphimerina
- 21. Tritæus
- 22. Tetartophia
- 23. Hemitritæa
- 24. Hectica

ORDER I.
CONTINENTES.

- 11. Diaria
- 12. Synocha
- 13. Synochus
- 14. Lenta.

CLASS III.—PHLOGISTICI.

- 30. Proctitis
- 31. Cystitis.

ORDER II.
PARENCHYMATICI.

- 32. Sphacelismus
- 33. Cynanche
- 34. Peripneumonia

- 35. Hepatitis
- 36. Splenitis
- 37. Nephritis
- 38. Hyæteritis.

ORDER III.
MUSCULOSI.

- 32. Phlegmone

ORDER I.
INTRINSECA.

40. Cephalalgia
41. Hemicrania
42. Gravedo
43. Ophthalmia
44. Otalgia
45. Otonalgia
46. Angina
47. Soda

ORDER I.
IDEALES.

65. Delirium
66. Paraphrosyne
67. Amentia
68. Mania
69. Dæmonia
70. Vesania
71. Melancholia

ORDER II.
IMAGINARIÆ.

72. Syringmos

ORDER I.
DEFECTIVI.

90. Lassitudo
91. Languor
92. Asthenia
93. Lipothymia
94. Syncope
95. Aphyxia

ORDER II.
SOPOROSI.

96. Somnolentia
97. Typhomania
98. Lethargus

ORDER I.
SPASTICI.

121. Spasmus
122. Briapismus
123. Borborygmus
124. Trismus
125. Sardiasis
126. Hysteria
127. Tetanus
128. Catochus
129. Catalepsia

ORDER I.
SUFFOCATORII.

146. Ruucedo
147. Vociferatio
148. Risus
149. Fletus
150. Suspirium
151. Oscitatio
152. Pandiculatio
153. Singultus

ORDER I.
CAPITIS.

172. Otorrhœa

CLASS IV.—DOLOROSI.

48. Cardialgia
49. Gastrica
50. Colica
51. Hepatica
52. Splenica
53. Pleuritica
54. Pneumonica
55. Hysteralgia
56. Nephritica
57. Dysuria

58. Pudendagra
59. Præctica

ORDER II.
EXTRINSECI.

60. Arthritis
61. Ostocopus
62. Rheumatismus
63. Volatica
64. Pruritus

CLASS V.—MENTALES.

73. Phantasma
74. Vertigo
75. Phanophobia
76. Hypochondriasis
77. Somnambulismus

ORDER III.
PATHETICI.

78. Citta
79. Buhmia
80. Polydipsia

81. Satyriasis
82. Erotomania
83. Nostalgia
84. Tarantismus
85. Rabies
86. Hydrophobia
87. Cacofitia
88. Antipathia
89. Anxietas

CLASS VI.—QUICTALES.

99. Cataphora
100. Carus
101. Apoplexia
102. Paraplegia
103. Hemiplegia
104. Paralysis
105. Stupor

ORDER III.
PRIVATIVI.

106. Morosis
107. Oblivio
108. Amblyopia

109. Cataracta
110. Amaurosis
111. Scotomia
112. Cophosis
113. Anosmia
114. Ageusia
115. Aphonia
116. Anorexia
117. Adipsia
118. Anæsthesia
119. Atecnia
120. Atonia

CLASS VII.—MOTORII.

130. Agrypnia

ORDER II.
AGITATORII.

131. Tremor
132. Palpitatio
133. Orgasmus
134. Subsultus
135. Carpologia
136. Stridor
137. Hippos

138. Psellismus
139. Chorea
140. Beriberi

ORDER II.
AGITATORII.

141. Rigor
142. Convulsio
143. Epilepsia
144. Hieranosus
145. Raphania

CLASS VIII.—SUPPRESSORII.

154. Sternutatio
155. Tussis
156. Stertor
157. Anhelatio
158. Suffocatio
159. Empyema
160. Dyspnœa
161. Asthma
162. Orthopnœa
163. Ephialtes

ORDER II.
CONSTRICTORII.

164. Aglutitio
165. Flatulentia
166. Obstipatio
167. Ischuria
168. Dysmenorrhœa
169. Dyslochia
170. Aglactatio
171. Sterilitas.

CLASS IX.—EVACUATORII.

173. Epiphora
174. Hæmorrhagia
175. Coryza

176. Stomacace
177. Ptyalismus.

**ORDER II.
THORACIS.**

- 178. Stratus
- 179. Expectoration
- 180. Hæmoptysis
- 181. Vomica

**ORDER III.
ABDOMINIS.**

- 182. Ructus
- 183. Nausea
- 184. Vomica
- 185. Hæmatemesis
- 186. Iliaca

**ORDER I.
EMACIANTES.**

- 209. Phthisis
- 210. Tabes
- 211. Atrophia
- 212. Marasmus
- 213. Rachitis

**ORDER I.
HUMORALIA.**

- 227. Aridura
- 228. Dignitum
- 229. Emphysema
- 230. Oedema
- 231. Sugillatio
- 232. Inflammatio
- 233. Abscessus
- 234. Gangrena
- 235. Sphacelus

**ORDER II.
DIALYTICA.**

- 236. Fractura
- 237. Luxatura
- 238. Ruptura
- 239. Contusura
- 240. Profusio
- 241. Vunus
- 242. Amputatura
- 243. Laceratura
- 244. Punctura
- 245. Morura
- 246. Combustura
- 247. Excoriatura
- 248. Intertrigo
- 249. Rhagias

**ORDER III.
EXULCERATIONES.**

- 250. Ulcus
- 251. Cacoethes
- 252. Noma
- 253. Carcinoma
- 254. Ozena
- 255. Fistula
- 256. Caries
- 257. Arthrocace
- 258. Coccyta
- 259. Paronychia
- 260. Perino

- 187. Cholera
- 188. Diarrhoea
- 189. Lienteria
- 190. Coeliaca
- 191. Cholirica
- 192. Dysenteria
- 193. Hæmorrhoids
- 194. Tenesmus
- 195. Crepitus

**ORDER IV.
GENITALIUM.**

- 196. Enuresis
- 197. Stranguria

- 198. Diabetes
- 199. Hæmaturia
- 200. Glus
- 201. Gonorrhoea
- 202. Leucorrhoea
- 203. Menorrhagia
- 204. Parturitus
- 205. Abortus
- 206. Mola

**ORDER V.
CORPORIS EXTERNI.**

- 207. Galactia
- 208. Sudor.

CLASS X.—DEFORMES.

**ORDER II.
TUMIDOSI.**

- 214. Polysarcia
- 215. Leucophlegmatia
- 216. Anasarca
- 217. Hydrocephalus
- 218. Ascites
- 219. Hyposarca
- 220. Tympanites

- 221. Graviditas.

**ORDER III.
DECOLORES.**

- 222. Cachexia
- 223. Chlorosis
- 224. Scorbutus
- 225. Icterus
- 226. Plethora.

CLASS XI—VITIA.

- 261. Pressura
- 262. Arctura

**ORDER IV.
SCABIES.**

- 263. Lepra
- 264. Tinea
- 265. Achor
- 266. Psora
- 267. Lippitudo
- 268. Serpigo
- 269. Herpes
- 270. Varus
- 271. Bacchia
- 272. Bubo
- 273. Anthrax
- 274. Phlyctæna
- 275. Pustula
- 276. Papula
- 277. Hordeolum
- 278. Verruca
- 279. Clavus
- 280. Myrmecium
- 281. Eschara.

**ORDER V.
TUMORES PROTUBERANTES.**

- 282. Aneurisma
- 283. Varix
- 284. Schirrus
- 285. Struma
- 286. Atheroma
- 287. Anchylosus
- 288. Ganglion
- 289. Natta
- 290. Spinola
- 291. Exostosis.

**ORDER VI.
PROCIDENTIAE.**

- 292. Hernia

- 293. Prolapsus
- 294. Condyloma
- 295. Sarcoma
- 296. Pterygium
- 297. Ectropium
- 298. Phimosis
- 299. Clitorismus.

**ORDER VII.
DEFORMATIONES.**

- 300. Contractura
- 301. Gibber
- 302. Lordosis
- 303. Distortio
- 304. Tortura
- 305. Strabismus
- 306. Lagophthalmia
- 307. Nyctalopia
- 308. Presbytia
- 309. Myopia
- 310. Libarium
- 311. Lagostoma
- 312. Apella
- 313. Atreta
- 314. Plica
- 315. Hæmuties
- 316. Alopecia
- 317. Trichiasis.

**ORDER VIII.
MACULAE.**

- 318. Cicatrix
- 319. Nævus
- 320. Morphæa
- 321. Vix
- 322. Sudamen
- 323. Melasma
- 324. Hepatizon
- 325. Lentigo
- 326. Ephelis.

*Synoptical View of the System of VOZEL.*ORDER I.
INTERMITTENTES.

1. Quotidiana
2. Tertiana
3. Quartana
4. Quintana
5. Sextana
6. Septana
7. Octana
8. Nonana
9. Decimana
10. Viginti-dies
11. Menstrua
12. Tertiana duplex
13. Quartana duplex
14. Quartana triplex

ORDER II.
CONTINUAE.
§ 1. *Simplices.*

15. Quotidiana
16. Synochus
17. Amatoria
18. Phrenitis
19. Epixia
20. Causos
21. Elodes
22. Lethargus
23. Typhomania
24. Leipyria
25. Phricodes

ORDER I.
HAEMORRHAGIAE.

81. Haemorrhagia
82. Epistaxis
83. Haemoptoe
84. Haemoptysis
85. Stomacace
86. Odontorrhœa
87. Otorrhœa
88. Ophthalmorrhagia
89. Hæmatemesia
90. Hepatirrhœa
91. Catarrhexis
92. Hæmaturia
93. Cystirrhagia
94. Stymatosis
95. Hæmatopedesis

126. Gravedo
127. Flatulentia
128. Obstipatio

134. Anxietas
135. Bistrismus
136. Pruritus
137. Catapxyxis
138. Rheumatismus
139. Arthritis
140. Cephalalgia
141. Cephalœa
142. Clavus

CLASS I.—FEBRES.

26. Langodes
27. Assodes
28. Cholera
29. Synchopalis
30. Hydrophobia
31. Oscitans
32. Ictericoodes
33. Pestilentialis
34. Siriasis

§ 2 *Compositæ.*¶ 1. *Exanthematicæ.*

35. Variolosa
36. Morbillosa
37. Miliaris
38. Pustulialis
39. Scarlatina
40. Urtica
41. Bullosa
42. Varicella
43. Pemphigodes
44. Aphthosæ

¶ 2. *Inflammatoriæ.*

45. Phrenismus
46. Chemosia
47. Ophthalmites
48. Otites
49. Angina
50. Pleuritis
51. Peripneumonia
52. Mediastina

53. Pericarditis
54. Carditis
55. Paraphrenitis
56. Gastritis
57. Enteritis
58. Hepatitis
59. Splenitis
60. Mesenteritis
61. Omentitis
62. Peritonitis
63. Myocolitis
64. Pancreatica
65. Nephritis
66. Cistitis
67. Hysteritis
68. Erysipelacea
69. Podagrica
70. Panaritium
71. Cyssotis.

¶ 3. *Symptomaticæ.*

72. Apoplectica
73. Catarrhalis
74. Rheumatica
75. Hæmorrhoidalis
76. Lactea
77. Vulneraria
78. Suppuratoria
79. Lenia
80. Hectica.

CLASS II.—PROFLUVIA.

96. Menorrhagia
97. Abortio

ORDER II.

APOCENOSES.

98. Catarrhus
99. Epiphora
100. Coryza
101. Otopuosis
102. Otoplatos
103. Ptyalismus
104. Vomica
105. Diarrhœa
106. Puorrhœa
107. Dysenteria
108. Lienteria
109. Cœliaca

110. Cholera
111. Pituitaria
112. Leucorrhœis
113. Eneuresis
114. Diuresis
115. Diabetes
116. Puoturia
117. Chylaria
118. Gonorrhœa
119. Leucorrhœa
120. Exoneirosis
121. Hydropedesia
122. Galactia
123. Hypercatharsis
124. Erphyse
125. Dysodia

CLASS III.—EPISCHESES.

129. Ischuria
130. Amorrhœa
131. Dyslochia

132. Deuteria
133. Agalaxis

CLASS IV.—DOLORES.

143. Hemicrania
144. Carebaria
145. Odontalgia
146. Hæmodia
147. Odaxismus
148. Otalgia
149. Acataposis
150. Cionia
151. Humantosis

152. Cardiognius
153. Mastodynia
154. Soda
155. Periadynia
156. Pneumatosis
157. Cardialgia
158. Encausia
159. Nausea
160. Colica

161. Filema
162. Ilcus
163. Stranguria
164. Dysuria
165. Luthuriasis
166. Tenesmus
167. Clunescia

168. Cedma
169. Hysteralgia
170. Dysmenorrhœa
171. Dystochia
172. Atocia
173. Priapismus

174. Psoriasis
175. Padogra
176. Osteocopus
177. Psophos
178. Volatica
179. Epiphlogisma.

CLASS V.—SPASMI.

180. T. tanus
181. Opisthotonus
182. Episthotonus
183. Catochus
184. Tremor
185. Frigus
186. Horror
187. Rigor
188. Epilepsia
189. Heclampsia
190. Hicranosus
191. Convulsio
192. R. phania
193. Chorea

194. Crampus
195. Scelerotyrbæ
196. Angone
197. Glossocele
198. Glo-socoma
199. Hippos
200. Illosis
201. Cinclesis
202. Cataclasis
203. Cilliosis
204. Sternutatio
205. Tussis
206. Clamor
207. Trismus

208. Capistrum
209. Sardiæsis
210. Gelasmus
211. Incubus
212. Singultus
213. Palpitatio
214. Vomitus
215. Ructus
216. Ruminatio
217. Oesophagismus
218. Hypochondriasis
219. Hysteria
220. Phlogosis
221. Digitum.

CLASS VI.—ADYNAMIAE.

212. Lassitudo
223. Asthenia
224. Torpor
225. Adynamia
226. Paralysis
227. Paraplegia
228. Hemiplegia
229. Apoplexia
230. Catalepsis
231. Carus
232. Coma
233. Somnolentia
234. Hyphophasis
235. Ptosis
236. Amblyopia
237. Mydriasis
238. Amaurosis
239. Cataracta
240. Synzezis
241. Glaucoma
242. Achlys

243. Nyctalopia
244. Hermeralopia
245. Hemalopia
246. Dysicoia
247. Surditas
248. Anosmia
249. Apogeusis
250. Asaphia
251. Clangor
252. Raucitas
253. Aphonia
254. Leptophonia
255. Oxyphonia
256. Rhenophonia
257. Mutitas
258. Traulotia
259. Psallotia
260. Ischophonia
261. Battarismus
262. Suspirium
263. Oscitatio

264. Pandiculatio
265. Apnœa
266. Macropnœa
267. Dyspnœa
268. Asthma
269. Orthopnœa
270. Pnignia
271. Renchus
272. Rhoehmos
273. Lipothymia
274. Syncope
275. Asphyxia
276. Apepsia
277. Dyspepsia
278. Diaphthora
279. Anorexia
280. Anatrope
281. Adypsia
282. Acyisis
283. Agnesia
284. Anodynia.

CLASS VII.—HYPAERESTHESES.

285. Antipathia
286. Agrypnia
287. Phantasma
288. Caligo
289. Hamalopia
290. Marmaryge
291. Dysopia

292. Susurrus
293. Vertigo
294. Apogeusia
295. Polydipsia
296. Bulimus
297. Addephagia

298. Cynorexia
299. Allotriophagia
300. Malacia
301. Pica
302. Bombus
303. Celæa

CLASS VIII.—CACHEXIÆ.

304. Cachexia
305. Chlorosis
306. Icterus
307. Melanchlorus
308. A. rophia
309. Tabes
310. Phthisis
311. Hydrothorax
312. Rachitis

313. Anasarca
314. Ascites
315. Hydrocystis
316. Tympanites
317. Hysterophyse
318. Scorbutus
319. Syphilis
320. Lepra

321. Elephantiasis
322. Elephantia
323. Plica
324. Phthiriasis
325. Physconia
326. Paracystis
327. Gangrena
328. Sphacelus.

CLASS IX.—PARANOIÆ.

329. Athymia
330. Delirium

331. Mania
332. Melancholia

333. Ectasis
334. Ecplexis

335. Enthusiasmos
336. Stupiditas

337. Amentia
338. Oblivio

339. Somnium
340. Hypnobotasis.

CLASS X.—VITIA.

ORDER I.
INFLAMMATIONES.

341. Ophthalmia
342. Blepharotis
343. Erysipelas
344. Hieropyr
345. Paronychia
346. Onychia
347. Encausis
348. Phimosi
349. Paraphimosis
350. Pernio

ORDER II.
TUMORES.

351. Plegmone
352. Furunculus
353. Anthrax
354. Abcessus
355. Onyx
356. Hippopoyon
357. Phygethlon
358. Empyema
359. Phyma
360. Ectymata
361. Urticaria
362. Parulis
363. Epulis
364. Anchylopa
365. Paraglossa
366. Chilon
367. Scrophula
368. Bubon
369. Bronchocele
370. Parotis
371. Gongrona
372. Sparganosia
373. Coillima
374. Scirrhus
375. Cancer
376. Sarcoma
377. Polypus
378. Condyloma
379. Ganglion
380. Ramula
381. Terminus
382. Oedema
383. Encephalocoele
384. Hydrocephalum
385. Hydrophthalmia
386. Spina bifida
387. Hydromphalus
388. Hydrocele
389. Hydrops scroti
390. Steatitis
391. Pneumatosis
392. Emphysema
393. Hysteroptosis
394. Cystoptosis
395. Archoptoma
396. Bubonocoele
397. Oscheocoele
398. Omphalocoele

399. Merocele
400. Enterocoele ovarialis
401. Ichiatocele
402. Ilytrocele
403. Hypogastrocele
404. Cystocoele
405. Cyrtoma
406. Hydrenterocoele
407. Varix
408. Eneurisma
409. Cirrocele
410. Gastrocele
411. Hepatocele
412. Splenocoele
413. Hysterocele
414. Hygrocirsocele
415. Sarcocoele
416. Physocoele
417. Exostosis
418. Hyperostosis
419. Pædithroceae
420. Encytis
421. Staphyloma
422. Staphylosis
423. Fungus
424. Tofus
425. Flemen

ORDER III.
EXTUBERANTIAE.

426. Verruca
427. Porrus
428. Clavus
429. Callus
430. Encanthia
431. Pladarotis
432. Pinnula
433. Pterygium
434. Hordeolum
435. Grando
436. Varus
437. Gutta rosacea
438. Ephelis
439. Esoche
440. Exoche.

ORDER IV.
PUSTULÆ and PAPULÆ.

441. Epinyctis
442. Phlyctæna
443. Herpes
444. Scabies
445. Aquula
446. Hydroa
447. Variola
448. Varicella
449. Purpura
450. Encauma

ORDER V.
MACULÆ.

451. Ecchymoma
452. Petechiæ
453. Morbilli

454. Scarlatæ
455. Lentigo
456. Urticaria
457. Stigma
458. Vibex
459. Vitiligo
460. Leuce
461. Cyasma
462. Lichen
463. Selina
464. Nebula

ORDER VI.
DISSOLUTIONES.

465. Vulnus
466. Ruptura
467. Rhagas
468. Fractura
469. Fissura
470. Plicatio
471. Thlasis
472. Luxatio
473. Subluxatio
474. Diachalasis
475. Altitus
476. Porrigio
477. Aposyrma
478. Anapleusis
479. Spasma
480. Contusio
481. Diabrosis
482. Agomphiasis
483. Eschara
484. Piptonychia
485. Cacoethes
486. Therioma
487. Carcinoma
488. Phagedæna
489. Noma
490. Sycosis
491. Fistula
492. Sinus
493. Caries
494. Achores
495. Crusta lactea
496. Favus
497. Tinea
498. Argemon
499. Ægilops
500. Ozæna
501. Aphthæ
502. Intertrigo
503. Rhacosis.

ORDER VII.
CONCRETIONES.

504. Ancyloblepharon
505. Zynizesis
506. Diacrymoma
507. Ancyloglossum
508. Ancylois
509. Cicatrix
510. Dasytalon.

CLASS XI. DEFORMITATES.

- | | | |
|---------------------|-------------------|---------------------|
| 511. Phoxos | 528. Phalacroctis | 545. Leiopodes |
| 512. Gibber | 529. Alopecia | 546. Apella |
| 513. Caput obatipum | 530. Miliarosis | 547. Hypospadias |
| 514. Strabismus | 531. Ptilosis | 548. Urorrhœas |
| 515. Myopiasis | 532. Rodatio | 549. Atreta |
| 516. Lagophthalmus | 533. Phalangosis | 550. Sanioles |
| 517. Trichiasis | 534. Coloboma | 551. Cripiorchis |
| 518. Ectropium | 535. Cercosis | 552. Hermaphrodites |
| 519. Entropium | 536. Cholosia | 553. Dionysiscus |
| 520. Rhœas | 537. Gryposis | 554. Artetiscus |
| 521. Rhysemata | 538. Nævus | 555. Nefrendis |
| 522. Lagocheules | 539. Munstronitas | 556. Spanopogon |
| 523. Melachosteon | 540. Polysarcia | 557. Hyperstetiscus |
| 524. Hirsuties | 541. Ischnosis | 558. G. liancon |
| 525. Canities | 542. Rhucnosis | 559. Galbulus |
| 526. Distrix | 543. Varus | 560. Mola. |
| 527. Xurasia | 544. Valgus | |

A Synoptical View of the System of SAGAR.

CLASS I.—VITIA.

ORDER I.
MACULÆ.

1. Leucoma
2. Vitiligo
3. Ephelis
4. Nævus
5. Ecchymoma.

ORDER II.
EFFLORESCENTIÆ.

6. Pustula
7. Papula
8. Phlyctæna
9. Barchia
10. Varus
11. Herpes
12. Epinyctis
13. Hemeropathos
14. Psyrdracia
15. Hydroa.

ORDER III.
PHYMATA.

16. Erythema
17. Oedema
18. Emphysema
19. Scirrbus
20. Inflammatio
21. Bubo
22. Pirotis
23. Farunculus
24. Anthrax
25. Cancer

ORDER I.
SOLUTIONES.
recentes, cruentæ.

78. Vulnus
79. Punctura
80. Sclopetoplaga
81. Morsus
82. Excoriatio
83. Contusio
84. Ruptura

26. Paronychia
27. Phimosis.

ORDER IV.
EXCRESCENTIÆ.

28. Sarcoma
29. Condyloma
30. Verucca
31. Pterygium
32. Hordeolum
33. Trachelophyma
34. Exostosis.

ORDER V.
CYSTITIDES.

35. Aneurysma
36. Varix
37. Mariaca
38. Hydatis.
39. Staphyloma
40. Lupia
41. Hydarthrus
42. Apostema
43. Exomphalus
44. Oscheophyma.

ORDER VI.
ECTOPLÆ.

45. Exophthalmia
46. Blepharoptosis
47. Hypostaphyle
48. Paraglossa
49. Proptoma

CLASS II.—PLAGÆ.

ORDER II.
SOLUTIONES,

recentes, cruentæ, artificiales.

85. Operatio
86. Amputatio
87. Sutura
88. Paracentesis.

ORDER III.
SOLUTIONES,
incruentæ.

89. Ulcus
90. Exulceratio

50. Exania
51. Exocystis
52. Hysterptosis
53. Colpoptosis
54. Gastrocele
55. Ophthalmocoele
56. Hepatocele
57. Merocele
58. Bubonocoele
59. Opodeocoele
60. Ischiocoele
61. Colpocoele
62. Perinæocoele
63. Peritonæorixis.
64. Encephalocoele
65. Hysteroloxia
66. Parorchidium
67. Exarthrema
68. Diastasis
69. Læxarthrus
70. Gibbositas
71. Lordosis.

ORDER VII.
DEFORMITATES.

72. Lagostoma
73. Apella
74. Polymerisma
75. Epidosis
76. Anchylomerisma
77. Hirsuties.

91. Fistula
92. Sinus
93. Eschara
94. Caries
95. Arthrocece.

ORDER IV.
SOLUTIONES,
anomale.

96. Rhagas.
97. Ambustio
98. Fractura
99. Fissura.

ORDER I. MACIES.

- 100. Tabes.
- 101. Phthisis
- 102. Atrophia
- 103. Hæmatoporia
- 104. Aridura.

ORDER II.
INTUMESCENTIÆ.

- 105. Plethora
 - 106. Polysarcia
 - 107. Pneumatosis
 - 108. Anasarca
 - 109. Phlegmatia.
 - 110. Phyeconia.
 - 111. Graviditas.
- ORDER III.
HYDROPEs *partialis*.
- 112. Hydrocephalus

ORDER I. VAGI.

- 142. Arthritis
- 143. Ostocopus
- 144. Rheumatismus
- 145. Catarrhus
- 146. Anxietas
- 147. Lassitudo
- 148. Stupor
- 149. Pruritus
- 150. Algor
- 151. Ardor.

ORDER II.
CAPITIS.

- 152. Cephalalgia

ORDER I.
SANGUIFLUXUS.

- 174. Hæmorrhagia
- 175. Hæmoptysis.
- 176. Stomacace.
- 177. Hæmatemesis
- 178. Hæmaturia
- 179. Metrorrhagia
- 180. Abortus.

ORDER II.
ALVIPLUXUS,
sanguinolenti.

- 181. Hepatirrhœa
- 182. Hæmorrhœis
- 183. Dysenteria
- 184. Melæna.

ORDER I.
EGERENDORUM.

- 210. Adiæpneustia
- 211. Sterilitas
- 212. Ischuria
- 213. Dysuria

ORDER I.
TONICI PARTIALES.

- 222. Strabismus
- 223. Trismus

CLASS III.—CACHEXIÆ.

- 113. Physocephalus
- 114. Hydrorachitis
- 115. Ascites
- 116. Hydrometra
- 117. Physometra
- 118. Tympanites
- 119. Meteorismus.

ORDER IV. TUBERA.

- 120. Rachitis
- 121. Scrophula
- 122. Carcinoma
- 123. Leontiasis
- 124. Malis
- 125. Framboesia.

ORDER V.

IMPETIGINES.

- 126. Syphilis
- 127. Scorbutus

CLASS IV.—DOLORES.

- 153. Cephalæa
- 154. Hemicrania
- 155. Ophthalmia
- 156. Otalgia
- 157. Odontalgia.

ORDER III.
PECTORIS.

- 158. Pyrosis.
- 159. Cardiognmus.

ORDER IV.
ABDOMINIS.

- 160. Cardialgia
- 161. Gastrodynïa
- 162. Colica

CLASS V.—FLUXUS.

ORDER III.
ALVIPLUXUS.
non sanguinolenti.

- 185. Nausea
- 186. Vomitus
- 187. Ileus
- 188. Cholera
- 189. Diarrhœa
- 190. Cœliaca
- 191. Lienteria
- 192. Tenesmus
- 193. Proctorrhœa.

ORDER IV.
SERIFLUXUS.

- 194. Ephidrosis
- 195. Epiphora

CLASS VI.—SUPPRESSIONES.

- 214. Aglactasio
- 215. Dyslochïa.

ORDER II.

INGERENDORUM.

- 216. Dysphagia
- 217. Angina.

CLASS VII.—SPAPMI.

- 224. Obstipitas
- 225. Contractura
- 226. Crampus
- 227. Priapismus

- 128. Elephantiasis.
- 129. Lepra
- 130. Scabies
- 131. Tinea.

ORDER VI.
ICTERICIÆ.

- 132. Aurigo
- 133. Melaficterus
- 134. Phœnigmus
- 135. Chlorosis.

ORDER VII.
ANOMALÆ

- 136. Phthiriasis
- 137. Trichoma
- 138. Alopecia
- 139. Elcosis
- 140. Gangræna
- 141. Necrosis.

- 163. Hepatalgia
- 164. Splenalgia
- 165. Nephralgia
- 166. Hysteralgia.

ORDER V.

EXTERNARUM.

- 167. Mastodynia
- 168. Rachialgia
- 169. Lumbago
- 170. Ischias
- 171. Proctalgia
- 172. Pudendagra
- 173. Digiium

- 196. Coryza
- 197. Pyralismus
- 198. Anacatharsis.
- 199. Diabetes
- 200. Enuresis
- 201. Pyuria
- 202. Leucorrhœa
- 203. Lochiorrhœa
- 204. Gonorrhœa
- 205. Galactirrhœa
- 206. Otorrhœa.

ORDER V.

AERIFLUXUS.

- 207. Flatulentia
- 208. Aëdopsophia
- 209. Dysodia.

ORDER III.

IMI VENTRIS.

- 218. Dysmenorrhœa
- 219. Dystocia
- 220. Dys hæmorrhœis
- 221. Obstipatio.

ORDER II.
TONICI GENERALES.

- 228. Tetanus
- 229. Catochus.

**ORDER III.
CLONICI PARTIALES.**

- 230. Nystagmus
- 231. Carphologia
- 232. Subsaltus
- 233. Pandiculatio
- 234. Apomystosis

- 235. Convulsio
- 236. Tremor
- 237. Palpitatio
- 238. Claudicatio.

**ORDER IV.
CLONICI GENERALES.**

- 239. Phricasmus

- 240. Eclampsia
- 241. Epilepsia
- 242. Hysteria
- 243. Scelotyrbe
- 244. Beriberia.

**ORDER I.
SPASMODICÆ.**

- 245. Ephialtes
- 246. Sternutptio
- 247. Oscedo
- 248. Singultus

**ORDER I.
DYSÆSTHESIÆ.**

- 258. Amblyopia
- 259. Caligo
- 260. Cataracta
- 261. Amaurosis
- 262. Anosmia
- 263. Agheusia
- 264. Dysecora
- 265. Paracusis
- 266. Cophosis
- 267. Anæsthesia.

**ORDER II.
ANEPITHYMIÆ.**

- 268. Anorexia

**ORDER I.
CONTAGIOSA.**

- 289. Pestis
- 290. Variola
- 291. Pemphigus
- 292. Purpura

**ORDER I.
MUSCULOSÆ.**

- 299. Phlegmone
- 300. Cynanche
- 301. Myositis
- 302. Carditis.

**ORDER II.
MEMBRANACÆ.**

- 303. Phrenitis

**ORDER I.
CONTINUÆ.**

- 316. Judicatoria
- 317. Humoraria
- 318. Frigeraria
- 319. Typhus
- 320. Hectica.

**ORDER I.
HALLUCINATIONES.**

- 328. Vertigo
- 329. Suffusio
- 330. Diplopia
- 331. Syngmos
- 332. Sympochondriasis
- 333. Somnambulismus.

CLASS VIII.—ANHELATIONES.

- 249. Tussis.

**ORDER II.
SUPPRESSIVÆ.**

- 250. Stertor
- 251. Dyspnœa
- 252. Asthma

- 253. Orthopnœa
- 254. Pleurodyne
- 255. Rheuma
- 256. Hydrothorax
- 257. Empyema.

CLASS IX.—DEBILITATES.

- 269. Adipsia
- 270. Anaphrodisia.

**ORDER III.
DYSINESIÆ.**

- 271. Mutitas
- 272. Aphonía
- 273. Psellismus
- 274. Cacophonia
- 275. Paralysis
- 276. Hemiplegia
- 277. Paraplexia.

**ORDER IV.
LEIPOPSYCHIA.**

- 278. Asthenia

- 279. Lipothymia
- 280. Syncope
- 281. Asphyxia.

**ORDER V.
COMATA.**

- 282. Catalepsia
- 283. Ecstasis
- 284. Typhomania
- 285. Lethargus
- 286. Cataphora
- 287. Carus
- 288. Apoplexia.

CLASS X.—EXANTHEMATA.

- 293. Rubecola
- 294. Scarlatina.

**ORDER II.
NON CONTAGIOSA.**

- 295. Miliaris

- 296. Erysipelas
- 297. Essera
- 298. Aphtha.

CLASS XI.—PHLEGMASIA.

- 304. Diaphragmitis
- 305. Pleuritis
- 306. Gastritis
- 307. Enteritis
- 308. Epiploitis
- 309. Cystitis

- 311. Peripneumonia
- 312. Hepatitis
- 313. Splenitis
- 314. Nephritis
- 315. Metritis.

**ORDER III.
PARENCHYMATOSÆ.**

- 310. Cephalitis

CLASS XII.—FEBRES.

**ORDER II.
REMITTENTES.**

- 321. Amphimerina
- 322. Tritæophya
- 323. Tetartophya.

**ORDER III.
INTERMITTENTES.**

- 324. Quotidiana
- 325. Tertiana
- 326. Quartana
- 327. Erratica.

CLASS XIII.—VESANIÆ.

**ORDER II.
MOROSITATES.**

- 334. Pica.
- 335. Bulimia
- 336. Polydipsia
- 337. Antipathia
- 338. Nostalgia
- 339. Panophobia

- 340. Satyriasis
- 341. Nymphomania
- 342. Tarantismus
- 343. Hydrophobia
- 344. Rabies.

- ORDER III.
DELIRIA.**
- 345. Paraphrosine

346. Amentia
347. Melancholia

348. Demonomania
349. Mania.

ORDER IV. ANOMALÆ.
350. Amnesia
351. Agrypnia.

Synoptical View of the System of Dr. MACBRIDE.

CLASS I.—UNIVERSAL DISEASES.

ORDER I. FEVERS.

1. Continued
2. Intermittent
3. Remittent
4. Eruptive
5. Hectic.

ORDER II.

INFLAMMATIONS.

6. External
7. Internal

ORDER III. FLUXES.

8. Alvine
9. Hæmorrhage
10. Humoral discharge

ORDER IV.

PAINFUL DISEASES.

11. Gout
12. Rheumatism
13. Ostocopus
14. Headach
15. Toothach
16. Earach
17. Pleurodyne
18. Pain in the stomach
19. Colic

20. Lithiasis

21. Ischuria

22. Proctalgia.

ORDER V.

SPASMODIC DISEASES.

23. Tetanus
24. Catochus
25. Locked jaw
26. Hydrophobia
27. Convulsion
28. Epilepsy
29. Ecclampsia
30. Hieranosos.

ORDER VI.

**WEAKNESSES and PRI-
VATIONS.**

31. Coma
32. Palsy
33. Fainting.

ORDER VII.

**ASTHMATIC DISOR-
DERS.**

34. Dyspnœa
35. Orthopnœa

36. Asthma

37. Hydrothorax

38. Empyema.

ORDER VIII.

MENTAL DISEASES.

39. Mania
40. Melancholia.

ORDER IX.

**CACHEXIES, or Humoral
Diseases.**

41. Corpulency
42. Dropsy
43. Jaundice
44. Emphyema
45. Tympany
46. Physconia
47. Atrophia
48. Osteosarcosis
49. Sarcostosis
50. Mortification
51. Scurvy
52. Scrophula
53. Cancer
54. Lues venerea.

CLASS II.—LOCAL DISEASES.

ORDER I.

**OF THE INTERNAL
SENSES.**

55. Loss of memory
56. Hypochondriasis
57. Loss of judgment.

ORDER II.

**OF THE EXTERNAL
SENSES.**

58. Blindness
59. Depraved sight
60. Deafness
61. Depraved hearing
62. Loss of smell
63. Depraved smell
64. Loss of taste.
65. Depraved taste
66. Loss of feeling.

ORDER III.

OF THE APPETITES.

67. Anorexia.
68. Cynorexia
69. Pica
70. Polydipsia
71. Satyriasis.
72. Nymphomania
73. Anaphrodisia.

ORDER IV.

**OF THE SECRETIONS
AND EXCRETIONS.**

74. Epiphora
75. Coryza
76. Ptyalism
77. Anacatharsis

78. Otorrhœa

79. Diarrhœa

80. Incontinence of urine

81. Pyuria

82. Dysuria

83. Constipation

84. Tenesmus

85. Dysodia

86. Flatulence

87. Œdosophia.

ORDER V.

**IMPEDING DIFFERENT
ACTIONS.**

88. Aphonia

89. Mutitas

90. Paraphonia

91. Dysphagia

92. Wryneck

93. Angone

94. Sneezing

95. Hiccup

96. Cough.

97. Vomiting

98. Palpitation of the heart.

99. Chorea

100. Trismus

101. Nystagmus

102. Cramp

103. Scolotyrbæ.

104. Contraction

105. Paralysis

106. Anchylosis

107. Gibbositas

108. Lordosis

109. Hydarthrus.

ORDER VI.

**OF THE EXTERNAL
HABIT.**

110. Tumour

111. Excrescence

112. Aneurism

113. Varix

114. Papulæ

115. Phlyctenæ

116. Pustulæ

117. Scabies, or Psora

118. Impetigo

119. Leprosy

120. Elephantiasis

121. Framboesia

122. Herpes

123. Maculæ

124. Alopecia

125. Trichoma

126. Scald head

127. Phthiriasis.

ORDER VII.

DISLOCATIONS.

128. Hernia

129. Prolapsus

130. Luxation.

ORDER VIII.

**SOLUTIONS OF CON-
TINUITY.**

131. Wound

132. Ulcer

133. Fissure

134. Fistula
135. Burn, or scald

135. Excoriation
137. Fracture

138. Caries.

CLASS III.—SEXUAL DISEASES.

ORDER I.

- GENERAL, *proper to Men.*
139. Febris testicularis
140. Tabes dorsalis.

ORDER II.

- LOCAL, *proper to Men.*
141. Dyspermatusmus
142. Gonorrhœa simplex
143. Gonorrhœa virulenta
144. Priapism
145. Phymosis
146. Pharyngismus
147. Crystalline

148. Hernia humoralis

149. Hydrocele

150. Sarcocoele

151. Cirsocele.

ORDER III. GENERAL,
proper to Women.

152. Amenorrhœa

153. Chlorosis

154. Leucorrhœa

155. Menorrhagia

156. Hysteralgia

157. Graviditas

158. Abortus

159. Dystochia

160. Febris puerperalis

163. Mastodynia

ORDER IV.

LOCAL, *proper to Women.*

162. Hydrops ovarii

163. Scirrhus ovarii

164. Hydrometra

165. Physometra

166. Prolapsus uteri

167. ——— vaginæ

178. Polypus uteri

CLASS IV.—INFANTILE DISEASES.

ORDER I. GENERAL.

169. Colica meconialis
170. ——— lactentium
171. Diarrhœa infantum
172. Aphthæ

173. Eclampsia

174. Atrophia

173. Rachitis.

ORDER II. LOCAL.

176. Imperforation

177. Anchyloglossum

178. Aurigo

179. Purpura

180. Crusta lactea.

NOSTALGIA. (*Nostalga*: from *nostos*, to return, and *algos*, pain.) A vehement desire for revisiting one's country. A genus of disease in the class *Locales*, and order *Dysorexiæ* of Cullen, known by impatience when absent from one's native home, and a vehement desire to return, attended with gloom and melancholy, loss of appetite, and want of sleep.

NOTHUS. (From *notos*, spurious.) 1. Those ribs which are not attached to the sternum are called *nothæ costæ*, the spurious ribs.

2. The term is applied to such diseases as resemble others: as peripneumonia notha, &c.

NOTIUS. (From *notos*, the back.) An epithet of the spinal marrow.

NOTIODES. (From *notus*, moisture.) Applied to a fever, attended with a vitiation of the fluids, or a colliquative wasting.

NOUFER'S MEDICINE. This famous remedy was invented by Madam Noufer, as a cure for the tape-worm. See *Filix*.

NUBULA. (Dim. of *nubes*, a cloud.) A cloud in the urine. A white speck in the eye.

NUCES CALLE. Common galls.

NUCES PURGANTES. See *Ricines*.

NUCSTA. See *Nux moschata*.

NUCHA. The hind part or nape of the neck.

NUCISTA. The nutmeg.

NUCLEUS. (*E nucæ*, from the nut.) A kernel. A fruit enclosed in a hard shell.

NUCULÆ SAPONARIÆ. *Bacca bermudensis.* Soap berries. Bermudas berries. A spherical fruit, about the size of a cherry, whose cortical part is yellow, glossy, and so transparent as to shew the sphericle black nut which rattles within, and which includes a black kernel. It is the produce of the *Sapindus saponaria* of Linnæus. The

cortical part has a bitter taste, and no smell; it raises a soapy froth with water, and has similar effects with soap in washing, and it is said to be a medicine of singular and specific virtue in chlorosis.

NUMMULARIA. (From *nummus*, money; so called because its leaves are round and of the size of the old silver two-pence.)

Hirundinaria. *Centimorbia.* Moneywort. This plant, *Lysimachia nummularia* of Linnæus, is very common in our ditches. It was formerly accounted vulnerary; it possesses antiscorbutic and restraining qualities. Boerhaave looks upon it as similar to a mixture of scurvy-grass with sorrel.

NUT, COCOA. The fruit of the *Cocos nucifera* of Linnæus. Within the nut is found a kernel, as pleasant as an almond, and also a large quantity of liquor resembling milk, which the Indians greedily drink before the fruit is ripe, it being then pleasant, but when the nut is matured, the liquor becomes sour. Some full-grown nuts will contain a pint or more of this milk, the frequent drinking of which seems to have no bad effects upon the Indians; yet Europeans should be cautious of making too free with it at first, for when Lionel Wafer was at a small island in the South Sea, where the tree grew in plenty, some of his men were so delighted with it, that at parting they were resolved to drink their fill, which they did; but their appetites had like to have cost them their lives, for though they were not drunk, yet they were so chilled and benumbed, that they could not stand, and were obliged to be carried aboard by those who had more prudence than the elves, and it was many days before they recovered. The shells of these nuts being hard, and capa-

ble of receiving a polish, they are often cut transversely, when, being mounted on stands, and having their edges silvered, or gilt, or otherwise ornamented, they serve the purpose of drinking-cups. The leaves of the tree are used for thatching, for brooms, baskets, and other utensils; and of the reticular web, growing at their base, the Indian women make cauls and aprons.

Nut, Barbadoes. See *Ricinus major*.

Nut, pistachio. See *Pistachio nus*.

Nut, purging. See *Ricinus major*.

Nutmeg. See *Nux moschata*.

NUTRITION. *Nutritio.* Nutrition may be considered the completion of the assimilating functions. The food changed by a series of decompositions, animalized and rendered similar to the being which it is designed to nourish, applies itself to those organs, the loss of which it is to supply; and this identification of nutritive matter to our organs constitutes nutrition.

The living body is continually losing its constituent parts, which a variety of causes are incessantly carrying off; several of its organs are constantly engaged in separating humours which pass off loaded with a part of its substance, consumed by the uniting action of air and caloric: internal friction, agitated by a pulsatory motion, detaches its particles.

Thus the animal machine is continually destroyed, and at distant periods of life does not contain a single particle of the same constituent parts. An experiment made with madder, (*rubia tinctorum*), which, when mixed with the food, reddens the bones of animals, proves in a very decisive manner this perpetual decomposition of living animal matter. Entirely to obliterate the diffused red colour of bones, it is only necessary to suspend for a time the use of this root. Therefore, if the most compact and solid parts be in a continual motion of decomposition and recomposition, there can be no doubt but that this motion must be more rapid in those parts, the constituent principles of which are in the smallest degree of cohesion, as in fluids.

It has been an object of consideration to determine the period of the entire renovation of the body; it has been said that an interval of seven years was necessary for the same particles to be totally obliterated, and their place supplied by others; but this change should seem to be more rapid in infancy and youth; it should also seem to be retarded in manhood, and require a very long time to be accomplished in old age, when all our parts acquire a remarkable degree of consistence and fixity, at the same time that the vital actions become more languid. There is no doubt but that sex, temperature, climate, profession, mode of living, and a variety of other causes, accelerate and retard this period,

so that it is impossible to affirm any thing certain on the precise time of its duration.

In proportion as our parts are destroyed, they are renewed by homogeneous particles, or such as are exactly similar to themselves; otherwise their nature, which is always alike, would suffer continual changes. When the nutritive matter has been animalized, or a-similated to the body which it is designed to nourish, by the organs of digestion, absorption, circulation, respiration, and secretion, the parts which it supplies retain and incorporate it with their own substance. This nutritive identification is variously effected in different parts, as the brain, muscles, bones, &c.; each of these appropriates to itself, by a true secretion, that which is found analogous to its nature, and rejects the heterogeneous particles brought by different vessels, chiefly by the arteries. A bone is a secretory organ, that becomes incrustated with phosphat of lime; the lymphatic vessels, which in the work of nutrition perform the office of excretory ducts, remove this salt after it has remained a certain time in the areolæ of its texture. It is the same in muscles with respect to the fibrin, and in the brain with albumen: each part imbibes, and renders solid in its structure, such juices as are of the same nature, in consequence of a power, of which the affinity of aggregation of the chymists gives us an idea, and perhaps furnishes us with an exact model.

A part to acquire nourishment should possess sensibility and motion; a ligature placed on its arteries and nerves, by destroying both these faculties, prevents it from being nourished, or having life. The blood flowing in the veins, and the fluid of the absorbents, contain vivifying and reparatory parts, in much smaller quantity than arterial blood; it is even generally believed that lymph and venous blood do not contain any thing directly nutritious.

The mechanism of nutrition would be explained after having precisely determined the differences of composition that exists between the aliments on which we exist and the exact substance of our organs, if we could distinguish how each function divests them of their characters, to invest them with our properties for each individual part, to co-operate in changing their nutritious principle into our own peculiar structure. To resolve this problem, let us suppose a man living entirely on vegetables, which, in fact, constitute the principal part of the subsistence of the generality of men; whatever portion of the plant he may consume, whether stalk, leaves, flowers, seeds, or roots; carbon, hydrogen, and oxygen enter their composition, which may be always, by a strict analysis, resolved into water and carbonic acid; to these three constituent principles,

sometimes a small quantity of azot, salts, and other things, is united. If we then examine the nature of the organs of this man, whose diet consists exclusively in vegetables, they will be proved of a composition very different from the kind of food; azot predominates, although the vegetable substance contain it in very small quantity, and many new products will be discovered which had not been distinguished in the aliment, but which abound in the body receiving nourishment, and seem produced by the act of nutrition.

The essential part of this function, therefore, is to cause the nutritive matter to pass into a more advanced state of composition, to deprive it of a portion of its carbon and hydrogen, to give a predominance of azot, and develop several substances which were not before distinguishable. Every living body, without exception, seems to possess a faculty of forming and decomposing substances, by the assistance of which it is supported, and of giving rise to new products. The marine-plant, the ashes of which form soda, if sown in a box filled with earth that does not contain a particle of that alkali, and moistened with distilled water, furnishes it in as great a quantity as if the plant had been growing on the borders of the sea, in a swampy soil, always inundated by brackish or salt water.

Living bodies are the proper, elaboratories in which such combinations and decompositions occur as art cannot imitate; bodies that to us appear simple, as soda and silix seem to form themselves of other parts, while some bodies, the composition of which we cannot determine, as certain metals suffer inevitable decompositions; from which we may fairly conclude, that the powers of nature in the composition and decomposition of bodies, far surpass the science of chymists. For a substance to be employed in our nourishment, it should be capable of change and fermentation, that is, susceptible of experiencing an internal and spontaneous motion, by which its elements change their combination and qualities. This condition of spontaneous mutability, excludes from the class of aliment every thing which is not organized, or constituted part of a living being: thus minerals are absolutely refractory to the action of our organs, which cannot convert them into their own peculiar substance. The common principle drawn from alimentary substances, however various they may be, called by Hippocrates the aliment, is probably a composition capable of a great degree of change and fermentation; this is also the opinion of all those who have endeavoured to discover its nature. Lorry thinks it is a mucous body; Cullen considers it saccharine;

Hallé believes it to be an hydro-carbonated oxyd, which only differs from the oxalic acid by having a smaller portion of oxygen. It is obvious that these three sentiments have the greatest resemblance, since oxygen, carbon, and hydrogen, united in different proportions, form a mucus, a saccharine body, and the oxalic base. The analysis of animal substances by nitric acid, reduces it to the latter base, by taking from it a great quantity of azot, the presence of which constitutes its most remarkable character.

Halé believes that the hydro-carbonated oxyd is combined with oxygen, in the stomach and intestinal canal, whether the latter principle be introduced with the food into the primæ viæ, or furnished by the decomposed humours; the intestinal fluids suffer their azot to be disengaged, which is carried to the alimentary base, and replaces the carbon that had been attracted by the oxygen to form the carbonic acid. This gas, when in the lungs, and again subjected to the action of atmospheric oxygen, carries off a certain portion of its carbon; and as it disengages the azot from the venous blood, it effects a new combination of this principle with the chyle; and when propelled to the skin, the atmospheric oxygen again disengages its carbon, and compleas its azotification; perhaps even the cutaneous organ answers similar purposes to the lymphatic system, as the pulmonary organ may effect to the sanguiferous system.

The animalization of alimentary substance, therefore, takes place principally by the loss of carbon, which is replaced by azot in animal fluids. These support themselves in a proper state, for, as they are continually losing the carbonic principle in the intestinal, pulmonary, and cutaneous combinations, they would be too much animalized if a newly formed chyle were not to attract the excess of azot. This theory is admitted, by its author, not to account for the formation of phosphoric salts, adeps, and abundance of other productions; but without adopting it *in toto*, we are induced to conclude, from the experiments and observations on which it is established, that the oxygen of the atmospheric air is one of the most powerful agents employed by nature to convert the aliments on which we subsist into our own peculiar substance.

NUTRITUM UNGUENTUM. A composition of litharge, vinegar, and oil.

NUX AR Matica. The nutmeg.

NUX AQUATICA. See *Tribulus aquaticus*.

NUX BARBADENSIS. See *Ricinus major*.

NUX BASILICA. The walnut.

NUX BERN. See *Ben*.

NUX CATHARTICA. The garden spurge.

NUX CATHARTICA AMERICANA. See *Ricinus major*.

NUX IBERICA. The cocoa-nut.

NUX JUGLANS. See *Juglans*.

NUX MEDICA. The maldivian nut.

NUX METELLA. The nux vomica.

NUX MOSCHATA. *Nutina*. *Nux myristica*. *Chrysobalanus Galeni*. *Unguentaria*. *Assala*. *Nux aromatica*. The nutmeg. The seed, or kernel, of the *Myristica moschata* of Linnæus:—*foliis lanceolatis, fructu glabro*. It is a spice that is well known, and has been long used both for culinary and medical purposes. Distilled with water, they yield a large quantity of essential oil, resembling in flavour the spice itself; after the distillation, an insipid sebaceous matter is found swimming on the water; the decoction, inspissated, gives an extract of an unctuous, very lightly bitterish taste, and with little or no astringency. Rectified spirit extracts the whole virtue of nutmegs, by infusion, and elevates very little of it in distillation; hence the spirituous extract possesses the flavour of the spice in an eminent degree. Nutmegs, when heated, yield to the press a considerable quantity of limpid, yellow oil. There are three kinds of unctuous substances, called oil of mace, though really expressed from the nutmeg. The best is brought from the East Indies, in stone jars; this is of a thick consistence, of the colour of mace, and has an agreeable fragrant smell; the second sort, which is paler coloured, and much inferior in quality, comes from Holland, in solid masses, generally flat, and of a square figure; the third, which is the worst of all, and usually called common oil of mace, is an artificial composition of suet, palm-oil, and the like, flavoured with a little genuine oil of nutmeg. The medicinal qualities of nutmeg are supposed to be aromatic, anodyne, stomachic and adstringent; and hence it has been much used in diarrhoeas and dysenteries. To many people, the aromatic flavour of nutmeg is very agreeable: they, however, should be cautioned not to use it in large quantities, as it is apt to affect the head, and even to manifest an hypnotic power in such a degree as to prove extremely dangerous. Bontius speaks of this as a frequent occurrence in India; and Dr. Cullen relates a remarkable instance of this soporific effect of nutmeg, which fell under his own observation; and hence concludes that, in apoplectic and paralytic cases, this spice may be very improper. The official preparations of nutmeg are a spirit and an essential oil, and the nutmeg, in substance, roasted to render it more adstringent; both the spice itself and the essential oil enter several compositions, as the *confectio aromatica*, *spiritus ammonie aromaticus*, &c.

NUX MYRISTICA. See *Nux moschata*.

NUX PERGICA. The walnut.

NUX PISTACIA. See *Pistachio nut*.

NUX PURGANS. See *Ricinus major*.

NUX SCRAPIONIS. St. Ignatius's bean.

NUX VOMICA. *Nux metella*. The nux vomica, lignum colubrium, and faba sancti ignatii, have been long known in the *Materia Medica* as narcotic poisons, brought from the East Indies, while the vegetables which produced them were unknown, or at least not botanically ascertained.

By the judicious discrimination of Linnæus, the nux vomica was found to be the fruit of the tree described and figured in the *Hortus Malabaricus*, under the name of *Caniram*, *cucurbitifera malabariensis* of Plukenet; *Vomica* of Linnæus. Now called *Strychnos nux vomica*.

To this genus also, but upon evidence less conclusive, he likewise justly referred the colubrium. But the faba sancti ignatii he merely conjectured might belong to this family, as appears by the query, *An strychni species?* which subsequent discoveries have enabled us to decide in the negative; for, in the *Suppl. Plant.* it constitutes the new genus *Ignatia*, which Loureiro has lately confirmed, changing the specific name *amarra* to that of *philippina*. The strychnos and ignatia are, however, nearly allied, and both rank under the order *Solanaceæ*.

Dr. Woodville has inquired thus far into the botanical origin of these productions, from finding that, by medical writers, they are generally treated of under the same head, and in a very confused and indiscriminate manner. The seed of the fruit, or berry of this tree, *Strychnos nux vomica*, is the official nux vomica; it is flat, round, about an inch broad, and near a quarter of an inch thick, with a prominence in the middle on both sides, of a gray colour, covered with a kind of woolly matter; and internally hard and tough, like horn. To the taste it is extremely bitter, but has no remarkable smell. It consists chiefly of a gummy matter, which is moderately bitter: the resinous part is very inconsiderable in quantity, but intensely bitter; hence rectified spirit has been considered its best menstruum.

Nux vomica is reckoned amongst the most powerful poisons of the narcotic kind, especially to brute animals; nor are instances wanting of its deleterious effects upon the human species. It proves fatal to dogs in a very short time, as appears by various authorities. Hillefeld and others found that it also poisoned hares, foxes, wolves, cats, rabbits, and even some birds, as crows and ducks; and Loureiro relates, that a horse died in four hours after taking a drachm of the seed in a half-roasted state.

The effects of this baneful drug upon different animals, and even upon those of the same species, appear to be rather uncertain, and not always in proportion to the quantity of the poison given. With

some animals it produces its effects almost instantaneously; with others, nor till after several hours, when laborious respiration, followed by torpor, trembling, coma, and convulsions, usually precede the fatal spasms, or tetanus, with which this drug commonly extinguishes life.

From four cases related of its mortal effects upon human subjects, we find the symptoms corresponded nearly with those which we have here mentioned of brutes; and these, as well as the dissections of dogs killed by this poison, not shewing any injury done to the stomach or intestines, proves that the nux vomica acts immediately upon the nervous system, and destroys life by the virulence of its narcotic influence.

The quantity of the seed necessary to produce this effect upon a strong dog, as appears by experiments, need not be more than a scruple; a rabbit was killed by five, and a cat by four, grains: and of the four persons to whom we have alluded, and who unfortunately perished by this deleterious drug, one was a girl ten years of age, to whom fifteen grains were exhibited at twice for the cure of an ague. Loss, however, tells us that he took one or two grains of it in substance, without discovering any bad effect: and that a friend of his swallowed a whole seed without injury.

In Britain, where physicians seem to observe the rule *Saltem non nocere* more strictly than in any other countries, the nux vomica has been rarely, if ever, employed as a medicine. On the continent, however, and especially in Germany, they have certainly been guided more by the axiom, "What is incapable of doing much harm, is equally unable to do much good." The truth of this remark was lately very fully exemplified by the practice of Baron Stoerck, and is farther illustrated by the medicinal character given of nux vomica, which, from the time of Gesner till that of a modern date, has been recommended by a succession of authors as an antidote to the plague, as a febrifuge, as a vermifuge, and as a remedy in mania, hypochondriasis, hysteria, rheumatism, gout, and canine madness. In Sweden, it has of late years been successfully used in dysentery; but Bergius, who tried its effects in this disease, says, that it suppressed the flux for twelve hours, which afterwards returned again. A woman, who took a scruple of this drug night and morning, two successive days, is said to have been seized with convulsions and vertigo, notwithstanding which the dysenteric symptoms returned, and the disorder was cured by other medicines; but a pain in the stomach, the effect of the nux vomica, continued afterwards for a long time.

Bergius, therefore, thinks it should only

be administered in the character of a tonic and anodyne, in small doses, (from five to ten grains,) and not till after proper laxatives have been employed. Loureiro recommends it as a valuable internal medicine in fluor albus; for which purpose he roasts it till it becomes perfectly black and friable, which renders its medicinal use safe, without impairing its efficacy. It is said to have been used successfully in the cure of agues, and has also been reckoned a specific in pyrosis, or water brush.

NYCTALOPIA. (From *νύξ*, the night, and *ὤψ*, an eye.) *Imbecilitas oculorum* of Celsus. A defect in vision, by which the patient sees little or nothing in the day, but in the evening and night sees tolerably well. The proximate cause is various:

1. From a periodical amaurosis, or gutta serena, when the blind paroxysm begins in the morning, and terminates in the evening.

2. From too great a sensibility of the retina, which cannot bear the meridian light. See *Phobotomia*.

3. From an opaque spot in the middle of the crystalline lens. When the light of the sun in the meridian contracts the pupil, there is blindness; about evening, or in more obscure places, the pupil dilates, hence the rays of light pass through the limbus of the crystalline lens.

4. From a disuse of light; thus persons who are educated in obscure prisons see nothing immediately in open meridian light; but by degrees their eyes are accustomed to distinguish objects in day-light.

5. From an immoveable mydriasis; for in this instance the pupil admits too great a quantity of light, which the immobile pupil cannot moderate; hence the patient, in a strong light, sees little or nothing.

6. From too great a contraction of the pupil. This admits a sufficiency of lucid rays in bright light, but towards night the pupil dilates more, and the patient sees better.

8 *Nyctalopia endemica*. A whole people have been nyctalops, as the Æthiopians, Africans, Americans, and Asiatics. A great flow of tears are excreted all the day from their eyes; at night they see objects.

8. From a commotion of the eye; from which a man in the night saw all objects distinctly.

NYCTOBASIS. (From *νύξ*, the night, and *βάσις*, to go.) Walking in the sleep.

NYMPHÆ. (From *νύμφα*, a water-nymph; so called because it stands in the water-course.) *Ala interna minores clitoridis. Colliculum. Collicula. Myrtoscheilides. Labia minora*. Two membranous folds, situated within the labia majora, at the sides of the entrance of the vagina uteri.

NYMPHÆA. (From *νύμφα*, a water-nymph; because it grows in watery places.)

The name of a genus of plants in the Linnaean system. Class, *Polyandria*. Order, *Monogynia*. The water-lily.

NYMPHA ALBA. *Leuconymphaea*. *Nenuphar*. *Micro-leuconymphaea*. White water-lily. This beautiful plant, *Nymphaea alba* of Linnæus, was formerly employed medicinally as a demulcent, and slightly anodyne remedy. It is now laid aside.

NYMPHA GLANDIFERA. *Nymphaea Indica*, and *Madraspatana*. Names for the faba Egyptia.

NYMPHA LUTEA. *Nymphaea major lutea* of Caspar Bauhin. Yellow water-lily. *Nymphaea lutea* of Linnæus. This beautiful plant was employed formerly with the same intentions as the white, and, like it, is now fallen into disuse. Lindesholpe informs us, that, in some parts of Sweden, the roots, which are the strongest part, were, in times of scarcity, used as food, and did not prove unwholesome.

NYMPHA LOTUS. The Egyptian lotus. An aquatic plant, a native of both Indies. The root is conical, firm, about the size of a middling pear, covered with a blackish bark, and set round with fibres. It has a sweetish taste, and, when boiled, or roasted, becomes as yellow within as the yolk of an egg. The plant grows in abundance on the banks of the Nile, and is there much sought after by the poor, who, in a short time, collect enough to supply their families with food for several days.

NYMPHOIDES. (From *νυμφα*, the water-lily, and *ωδ*, likeness.) A herb resembling the water-lily.

NYMPHOMANIA. (From *νυμφα*, nymph, and *μανια*, madness) *Furor uterinus*. Called by the Arabians, *Acras*. *Brachuna*. *Arascon*. *Arasum*. *Astromania*. A genus of disease in the class *Locales*, and order *Dysorexia*, of Cullen, characterised by excessive and violent desire for coition in women. The effects, as described by Juvenal, in his sixth satire, are most humiliating to human nature. It acknowledges the same causes as satyriasis; but as females, more especially in warm

climates, have a more irritable fibre, they are apt to suffer more severely than the males.

It is a species of madness, or a high degree of hysteria. Its immediate cause is a preternatural irritability of the uterus and pudenda of women, or an unusual acrimony of the fluids in these parts. Its presence is known by the wanton behaviour of the patient; she speaks and acts with unrestrained obscenity, and, as the disorder increases, she scolds, cries, and laughs, by turns. While reason is retained, she is silent, and seems melancholy, but her eyes discover an unusual wantonness. The symptoms are better or worse until the greatest degree of the disorder approaches, and then, by every word and action, her condition is too manifest.

NYMPHOTOMIA. (From *νυμφα*, the nymph, and *τομή*, to cut.) The operation of removing the nymph when too large.

NYSTAGMUS. (From *νυσσιν*, to sleep.) A twinkling of the eyes, such as happens when a person is very sleepy. Authors also define nystagmus to be an involuntary agitation of the ocular bulb. It is known by the instability or involuntary and constant motions of the globe of the eye, from one canthus to another, or in some other directions. Sometimes it is accompanied with an hippus, or an alternate and repeated dilatation and constriction of the pupil. The species are, 1. Nystagmus, from fear. This agitation is observed under the operation for the cataract; and it is checked by persuasion, and waiting a short space of time. 2. Nystagmus, from sand, or small gravel, falling in the eye. 3. Nystagmus, from a catarrh, which is accompanied with much inflammation. 4. Nystagmus, from saburra in the primæ viæ, as is observed in infants afflicted with worms, and is known by the signs of saburra. 5. Nystagmus symptomaticus, which happens in hysteric, epileptic, and sometimes in pregnancy, and is a common symptom accompanying St. Vitus's dance.

O.

OAK. See *Quercus*.

O. k of Jerusalem. See *Botrys vulgaris*.

Oak, acia. See *Quercus marina*.

Oak, willow-leaved. See *Quercus Phellos*.

Out. See *Avena*.

OBLÆA. (From *ελαος*, a dart, or a spit.) *Obelæa sagittalis*, an epithet for the sagittal suture of the skull.

OBLÆSION. (From *ob*, against, and *lædo*, to hurt.) An injury done to any part.

OBLISCOTHECA. (From *obeliskos*, an obelisk, and *θησαυρος*, a bag; so called from the shape of its seed-bags.) The dwarf American sun-flower.

OBLIQUUS ASCENDENS ABDOMINIS. See *Obliquus internus abdominis*.

OBLIQUUS ASCENDENS INTERNUS. See *Obliquus internus abdominis*.

OBLIQUUS AURIS. See *Laxator tympani*.

OBLIQUUS CAPITIS INFERIOR. See *Obliquus inferior capitis*.

OBLIQUUS CAPITIS SUPERIOR. See *Obliquus superior capitis*.

OBLIQUUS DESCENDENS ABDOMINIS. See *Obliquus externus abdominis*.

OBLIQUUS DESCENDENS EXTERNUS. See *Obliquus externus abdominis*.

OBLIQUUS EXTERNUS. See *Obliquus externus abdominis*.

OBLIQUUS EXTERNUS ABDOMINIS. This muscle, which is so named by Morgagni, Albinus, and Winslow, is the *Obliquus descendens* of Vesalius and Douglas, and the *Obliquus major* of Haller, and some others. By Dumas it is named *Iliopubicasto-abdominal*. It is a broad, thin muscle, fleshy posteriorly, and tendinous in its middle and lower part, and is situated immediately under the integuments, covering all the other muscles of the lower belly. It arises from the lower edges of the eight, and sometimes, though rarely, of the nine inferior ribs, not far from their cartilages, by as many distinct fleshy portions, which indigitate with corresponding parts of the serratus major anticus, and the latissimus dorsi. From these several origins, the fibres of the muscle descend obliquely forwards, and soon degenerate into a broad and thin aponeurosis, which terminates in the linea alba. About an inch and a half above the pubis, the fibres of this aponeurosis separate from each other, so as to form an aperture, which extends obliquely inwards and forwards, more than an inch in length, and is wider above than below, being nearly of an oval figure. This is what is sometimes, though erroneously, called the *ring* of the abdominal muscles, for it belongs only to the external oblique there being no such opening either in the obliquus internus, or in the transversalis, as some writers, and particularly Douglas and Cheselden, would give us to understand. This opening, or ring, serves for the passage of the spermatic vessels in men, and of the round ligament of the uterus in women, and is of a larger size in the former than in the latter. The two tendinous portions, which, by their separation, form this aperture, are called the *columns* of the ring. The anterior, superior, and inner column, which is the broadest and thickest of the two, passes over the symphysis pubis, and is fixed to the opposite os pubis; so that the anterior column of the right obliquus externus intersects that of the left, and is, as it were, interwoven with it, by which means their insertion is strengthened, and their attachment made firmer. The posterior, inferior and exterior column approaches the anterior one as it descends, and is fixed behind and be-

low it to the os pubis of the same side. The fibres of that part of the obliquus externus, which arises from the two inferior ribs, descend almost perpendicularly, and are inserted, tendinous and fleshy, into the outer edge of the anterior half of the spine of the ilium. From the anterior superior spinous process of that bone, the external oblique is stretched tendinous to the os pubis, forming what is called *Poupart's*, and sometimes *Fallopian's* ligament, Fallopian having first described it. Winslow, and many others name it, the *inguinal* ligament. But, after all, it has no claim to this name, it being nothing more than the tendon of the muscle, which is turned or folded inwards at its anterior edge. It passes over the blood vessels of the lower extremity, and is thickest near the pelvis; and in women, from the greater size of the pelvis, it is longer and looser than in men. Hence we find that women are most liable to crural hernia; whereas men, from the greater size of the ring of the external oblique, are most subject to the inguinal. From this ligament, and from that part of the tendon which forms the ring, we observe a detachment of tendinous fibres, which are lost in the *fascia lata*, of the thigh. This may, in some measure, account for the pain which, in cases of strangulated hernia, is felt when the patient stands upright, and which is constantly relieved upon bending the thigh upwards. This muscle serves to draw down the ribs in expiration; to bend the trunk forwards when both muscles act, or to bend it obliquely to one side, and, perhaps, to turn it slightly upon its axis, when it acts singly; it also raises the pelvis obliquely when the ribs are fixed; it supports and compresses the abdominal viscera, assists in the evacuation of the urine and feces, and is likewise useful in parturition.

OBLIQUUS INFERIOR. See *Obliquus inferior capitis*, and *Obliquus inferior oculi*.

OBLIQUUS INFERIOR CAPITIS. This muscle, which is the *obliquus inferior sive major* of Winslow, and the *Spina axo-ido-trachelo-altoidei* of Dumas, is larger than the obliquus superior capitis. It is very obliquely situated between the two first vertebrae of the neck. It arises tendinous and fleshy from the middle and outer side of the spinous process of the second vertebra of the neck, and is inserted tendinous and fleshy into the lower and posterior part of the transverse process of the first vertebra. Its use is to turn the first vertebra upon the second, as upon a pivot, and to draw the face towards the shoulder.

OBLIQUUS INFERIOR OCULI. *Obliquus minor oculi* of Winslow, and *Maxilla sclerotici* of Dumas. An oblique muscle of the eye, that draws the globe of the eye forwards, inwards, and downwards. It arises by a narrow beginning from the

outer edge of the orbital process of the superior maxillary bone, near its junction with the lachrymal bone, and running obliquely outwards, is inserted into the sclerotic membrane of the eye.

OBLIQUUS INFERIOR SIVE MAJOR. See *Obliquus inferior capitis*.

OBLIQUUS INTERNUS. See *obliquus internus abdominis*.

OBLIQUUS INTERNUS ABDOMINIS.

This muscle, which is the *Obliquus ascendens* of Vesalius, Douglas, and Cowper, the *obliquus minor* of Haller, the *Obliquus internus* of Winslow, the *Obliquus ascendens internus* of Innes, and the *lilo-lumbo-costi-abdominal* of Dumas, is situated immediately under the external oblique, and is broad and thin like that muscle, but somewhat less considerable in its extent. It arises from the spinous processes of the three inferior lumbar vertebrae, and from the posterior and middle part of the os sacrum, by a thin tendinous expansion, which is common to it and to the serratus posticus inferior, by short tendinous fibres, from the whole spine of the ilium, between its posterior tuberosity and its anterior and superior spinous process; and from two-thirds of the posterior surface of what is called Fallopius's ligament, at the middle of which we find the round ligament of the uterus in women, and the spermatic vessels in men, passing under the thin edge of this muscle; and in the latter, it likewise sends off some fibres, which descend upon the spermatic chord, as far as the tunica vaginalis of the testis, and constitute what is called the *cremaster* muscle, which surrounds, suspends, and compresses the testicle. From these origins, the fibres of the internal oblique run in different directions; those of the posterior portion ascend obliquely forwards, the middle ones become less and less oblique, and, at length, run in an horizontal direction, and those of the anterior portion extend obliquely downwards. The first of these are inserted, by very short tendinous fibres, into the cartilages of the fifth, fourth, and third of the false ribs; the fibres of the second, or middle portion, form a broad tendon, which, after being inserted into the lower edge of the cartilage of the second false rib, extends towards the linea alba, and separates into two layers; the anterior layer, which is the thickest of the two, joins the tendon of the obliquus externus, and runs over the two upper thirds of the rectus muscle, to be inserted into the linea alba; the posterior layer runs under the rectus, adheres to the anterior surface of the tendon of the transversalis, and is inserted into the cartilages of the first of the false, and the last of the true ribs, and likewise into the linea alba. By this structure we may perceive that the greater part of the rectus is in-

closed, as it were, in a sheath. The fibres of the anterior portion of the internal oblique, or those which arise from the spine of the ilium and the ligamentum Fallopii, likewise form a broad tendon, which, instead of separating into two layers, like that of the other part of the muscle, runs over the lower part of the rectus, and adhering to the under surface of the tendon of the external oblique, is inserted into the fore part of the pubis. This muscle serves to assist the obliquus externus; but it seems to be more evidently calculated than that muscle is to draw the ribs downwards and backwards. It likewise serves to separate the false ribs from the true ribs, and from each other.

OBLIQUUS MAJOR ABDOMINIS. See *Obliquus externus abdominis*.

OBLIQUUS MAJOR CAPITIS. See *Obliquus inferior capitis*.

OBLIQUUS MAJOR OCULI. See *Obliquus superior oculi*.

OBLIQUUS MINOR ABDOMINIS. See *Obliquus internus abdominis*.

OBLIQUUS MINOR CAPITIS. See *Obliquus superior capitis*.

OBLIQUUS MINOR OCULI. See *Obliquus inferior oculi*.

OBLIQUUS SUPERIOR CAPITIS. Riolanus, who was the first that gave particular names to the oblique muscles of the head, called this muscle *obliquus minor*, to distinguish it from the inferior, which, on account of its being much larger, he named *obliquus major*. Spigelius afterwards distinguished the two, from their situation with respect to each other, into *superior* and *inferior*; and in this he is followed by Cowper and Douglas. Winslow retains both names. Dumas calls it *Traehelo-altoideo-occipital*. That used by Albinus is here adopted. This little muscle, which is nearly of the same shape as the *recti capitis*, is situated laterally between the occiput and the first vertebra of the neck, and is covered by the complexus and the upper part of the splenius. It arises, by a short thick tendon, from the upper and posterior part of the transverse process of the first vertebra of the neck, and, ascending obliquely inwards, and backwards, becomes broader, and is inserted, by a broad flat tendon, and some few fleshy fibres, into the os occipitis, behind the back part of the mastoid process, under the insertion of the complexus and splenius, and a little above that of the rectus major. The use of this muscle is to draw the head backwards, and perhaps to assist in its rotatory motion.

OBLIQUUS SUPERIOR OCULI. *Trochlearis.* *Obliquus major* of Winslow, and *Optico-trochleiscleroticius* of Dumas. An oblique muscle of the eye, that rolls the globe of the eye, and turns the pupil downwards and outwards. It arises like

the straight muscles of the eye from the edge of the foramen opticum at the bottom of the orbit, between the rectus superior and rectus internus; from thence runs straight along the papyraceous portion of the ethmoid bone to the upper part of the orbit, where a cartilaginous trochlea is fixed to the inside of the internal angular process of the os frontis, through which its tendon passes, and runs a little downwards and outwards, enclosed in a loose membranaceous sheath, to be inserted into the sclerotic membrane.

OBLIQUUS SUPERIOR SIVE MINOR. See *Obliquus superior capitis*.

OBLIQUUS SUPERIOR SIVE TROCHLEARIS. See *Obliquus superior oculi*.

OBSIDIANUM. A species of glass, so called from its resemblance to a kind of stone, which one Obsidius discovered in Ethiopia, of a very black colour, though sometimes pellucid, and of a muddy water. Pliny says also, that *obsidianum* was a sort of colour with which vessels were glazed. Hence the name is applied, by Libavius, to glass of antimony.

OBSTETRIC. (*Obstetricus*; from *obstrix*, a nurse.) Belonging to midwifery.

OBSTIPATIO. (From *obstipō*, to stop up.) Costiveness. A genus of disease in the class *Locales*, and order *Epischoeces* of Cullen, comprehending three species:

1. *Obstipatio debilitum*, in weak and commonly dyspeptic persons.

2. *Obstipatio rigidorum*, in persons of rigid fibres, and a melancholy temperament.

3. *Obstipatio obstructorum*, from obstructions. See *Colica*.

OBSTRUENTIA. (From *obstruo*, to shut up.) Medicines which close the orifices of the ducts, or vessels.

OBSTUPIFACIENTIA. (From *obstupefacio*, to stupefy.) Narcotics.

OBSTUNDENTIA. (From *obtundo*, to make blunt.) Substances which sheath or blunt irritation, and are much the same as demulcents. They consist chiefly of bland, oily, or mucilaginous matters, which form a covering on inflamed and irritable surfaces, particularly those of the stomach, lungs, and anus.

OBTURATOR EXTERNUS. *Extra-petio pubi-trochanterien* of Dumas. This is a small flat muscle, situated obliquely at the upper and anterior part of the thigh, between the pectinalis and the fore-part of the foramen thyroideum, and covered by the adductor brevis femoris. It arises tendinous and fleshy from all the inner half of the circumference of the foramen thyroideum, and likewise from part of the obturator ligament. Its radiated fibres collect and form a strong roundish tendon, which runs outwards, and, after adhering to the capsular ligament of the joint, is inserted into a cavity at the inner and back part of

the root of the great trochanter. The chief uses of this muscle are, to turn the thigh obliquely outwards, to assist in bending the thigh, and in drawing it inwards. It likewise prevents the capsular ligament from being pinched in the motions of the joint.

OBTURATOR INTERNUS. *Marsupialis*, sen *obturator internus* of Douglas. *Marsupialis seu bursalis* of Cowper, and *Intra-petio-trochanterien* of Dumas. A considerable muscle, a great part of which is situated within the pelvis. It arises, by very short tendinous fibres, from somewhat more than the upper half of the internal circumference of the foramen thyroideum of the os innominatum. It is composed of several distinct fasciculi, which terminate in a roundish tendon that passes out of the pelvis, through the niche that is between the spine and the tuberosity of the ischium, and, after running between the two portions of the gemini in the manner just now described, is inserted into the cavity at the root of the great trochanter, after adhering to the adjacent part of the capsular ligament of the joint. This muscle rolls the os femoris obliquely outwards, by pulling it towards the ischiatric niche, upon the cartilaginous surface of which its tendon, which is surrounded by a membranous sheath, moves as upon a pulley.

OBTURATOR NERVE. A nerve of the thigh, that is lost upon its inner muscles.

OCCIPITAL BONE. *Os occipitis. Os memorie. Os nervosum. Os basilare.*

This bone, which forms the posterior and inferior part of the skull, is of an irregular figure, convex on the outside and concave internally. Its external surface, which is very irregular, serves for the attachment of several muscles. It affords several inequalities, which sometimes form two semi-circular hollows, separated by a scabrous ridge. The inferior portion of the bone is stretched forwards in form of a wedge, and hence is called the *cuneiform* process, or *basilary* process. At the base of this process, situated obliquely on each side of the foramen magnum, are two flat, oblong protuberances, named *condyles*. They are covered with cartilage, and serve for the articulation of the head with the first vertebra of the neck. In the inferior portion of this bone, at the basis of the cranium, and immediately behind the cuneiform process, we observe a considerable hole, through which the medulla oblongata passes into the spine. The *nervi accessorii*, the vertebral arteries, and sometimes the vertebral veins likewise, pass through it. Man being designed for an erect posture, this foramen magnum is found nearly in the middle of the basis of the human cranium, and at a pretty equal distance from the posterior part of the occiput, and the

anterior part of the lower jaw; whereas in quadrupeds it is nearer the back part of the occiput. Besides this hole, there are four other smaller foramina, viz. two before, and two behind the condyles. The former serve for the transmission of the ninth pair of nerves, and the two latter for the veins which pass from the external parts of the head to the lateral sinuses. On looking over the internal surface of the os occipitis, we perceive the appearance of a cross, formed by a very prominent ridge, which rises upwards from near the foramen magnum, and by two transverse sinuosities, one on each side of the ridge. This cross occasions the formation of four fossæ, two above and two below the sinuosities. In the latter are placed the lobes of the cerebellum, and in the former the posterior lobes of the brain. The two sinuosities serve to receive the lateral sinuses. In the upper part of this bone is seen a continuation of the sinuosity of the longitudinal sinus; and at the basis of the cranium we observe the inner surface of the cuneiform process made concave, for the reception of the medulla oblongata. The occipital bone is thicker and stronger than any of the other bones of the head, except the petrous part of the ossa temporum; but it is of unequal thickness. At its lateral and inferior parts, where it is thinnest, it is covered by a great number of muscles. The reason for so much thickness and strength in this bone seems to be, that it covers the cerebellum, in which the least wound is of the utmost consequence; and that it is, by its situation, more liable to be fractured by falls than any other bone of the cranium. For, if we fall forwards, the hands are naturally put out to prevent the forehead's touching the ground; and if on one side, the shoulders in a great measure protect the sides of the head; but if a person fall backwards, the hind part of the head consequently strikes against the earth, and that too with considerable violence. Nature therefore has wisely constructed this bone so as to be capable of the greatest strength at its upper part, where it is the most exposed to injury. The os occipitis is joined, by means of the cuneiform process, to the sphenoid bone, with which it often ossifies, and makes but one bone in those who are advanced in life. It is connected to the parietal bones by the lambdoidal suture, and to the temporal bones by the additamentum of the temporal suture. The head is likewise united to the trunk by means of this bone. The two condyles of the occipital bone are received into the superior oblique processes of the atlas, or first vertebra of the neck, and it is by means of this articulation that a certain degree of motion of the head backwards and forwards is performed.

But it allows only very little motion to either side; and still less of a circular motion, which the head obtains principally by the circumvolution of the atlas on the second vertebra, as is described more particularly in the account of the vertebra. In the fœtus, the os occipitis is divided by an unossified cartilaginous substance into four parts. One of these, which is the largest, constitutes all that portion of the bone which is above the foramen magnum: two others, which are much smaller, compose the inside of the foramen magnum, and include the condyloid processes; and the fourth is the cuneiform process. This last is sometimes not completely united with the rest, so as to form one bone, before the sixth or seventh year.

OCCIPITALIS. See *Occipito-frontalis*.

OCCIPITO-FRONTALIS. *Digastricus cranii*. *Epicranius* of Albinus. *Frontalis et occipitalis* of Winslow and Cowper, and *Occipito-frontal* of Dumas. A single, broad, digastric, muscle, that covers the cranium, pulls the skin of the head backwards, raises the eye-brows upwards, and, at the same time, draws up and wrinkles the skin of the forehead. It arises from the posterior part of the occiput, goes over the upper part of the os parietale and os frontalis, and is lost in the eye-brows.

OCCIPUT. The hinder part of the head. See *Caput*.

OCCULT QUALITY. A term that has been much used by writers that had not clear ideas of what they undertook to explain; and which served therefore only for a cover to their ignorance.

OCCULT DISEASES, is likewise from the same mint as the former, *occulus* signifying hidden, and, therefore, nothing can be understood, when a person speaks of a hidden disease, but that it is a disease he does not understand.

OCHREA. (From *οχηω*, to carry.) A vehicle, or thin fluid.

OCHETREMA. (From *οχετος*, a duct.) The nostril.

OCHTUS. (From *οχηω*, to convey.) A canal or duct. The urinary, or abdominal passages.

OCHUS. (From *οχηω*, to carry.) The bag of the scrotum.

OCHRA. (From *οχρος*, pale; so named because it is often of a pale colour.) 1. Ochre. *Minera ferri lutea vel rubra*, An argillaceous earth impregnated with iron, of a red or yellow colour. The Armenian bole, and other earths, are often adulterated with ochre.

2. The fore-part of the tibia.

OCHREUS. (From *οχρος*, pale; so called from the pale muddy colour of its flow-ers.) A leguminous plant, or kind of pulse.

OCHTHODES. (From *οχθος*, importing the tumid lips of ulcers, callous, tumid.)

An epithet for ulcers, whose lips are callous and tumid, and consequently difficult to heal.

OCIMEASTRUM. (Dim. of *ocimum*, basil.) Wild white campion, or basil.

OCIMUM. (From *ocum*, swift; so called from its quick growth.) *Ocimum*. The name of a genus of plants in the Linnæan system. Class, *Didymia*. Order, *Gymnospermia*. See *Basilicum*.

OCIMUM BASILICUM. The systematic name of the common or citron basil. See *Basilicum*.

OCIMUM CARYOPHYLLATUM. *Ocimum minimum* of Casper, Bauhon, and Linnæus. Small or bush basil. This plant is mildly balsamic. Infusions are drank as tea, in catarrhus and uterine disorders, and the dried leaves are made into cephalic and sternutatory powders. They are, when fresh, very juicy, of a weak aromatic and very mucilaginous taste, and of a strong and agreeable smell, improved by drying.

OCTAVA. (From *octo*, eight.) An erratic intermitting fever, which returns every eighth day.

OCTAVUS HUMERI. The *Teres Minor*.

OCTAVUS HUMERI PLACENTINI. The *Teres Minor*.

OCULARES COMMUNES. A name for the nerves called *Motores Oculorum*.

OCULARIA. (From *oculus*, the eye; so called from its uses in disorders of the eye.) See *Euphrasia*.

OCULI ADDUCTOR. See *Rectus internus oculi*.

OCULI ATTOLLENS. See *Rectus superior oculi*.

OCULI CANCROBUM. See *Cancer*.

OCULI DEPRESSOR. See *Rectus inferior oculi*.

OCULI ELEVATOR. See *Rectus superior oculi*.

OCULI LEVATOR. See *Rectus superior oculi*.

OCULI OBLIQUUS INFERIOR. See *Obliquus inferior oculi*.

OCULI OBLIQUUS MAJOR. See *Obliquus superior oculi*.

OCULI OBLIQUUS MINOR. See *Obliquus inferior oculi*.

OCULUS BOVINUS. See *Proptosis*.

OCULUS BOVIS. See *Bellis major*.

OCULUS BUBULUS. See *Proptosis*.

OCULUS CHRISTI. Austrian flea bane; a species of *Inula*.

OCULUS ELEPHANTINUS. A name given to *Proptosis*.

OCULUS GENU. The knee-pan.

OCULUS LACHRYMAN. The *Epiphora*.

OCULUS MUNDI. A species of *Opal*, generally of a yellowish colour. By laying in water it becomes of an amber colour, and also transparent.

ODAXISMOS. (From *odus*, a tooth.) A

biting sensation, pain, or itching in the gums.

ODONTASOMOS. (From *odus*, a tooth, and *ago*, to draw.) The name of an instrument to draw teeth, one of which, made of lead, Forrester relates to have been hung up in the temple of Apollo, denoting, that such an operation ought not to be made, but when the tooth was loose enough to draw with so slight a force as could be applied with that.

ODONTAGRA. (From *odus*, a tooth, and *agea*, a seizure.) 1. The gout in the teeth.

2. A tooth-drawer.

ODONTALGIA. (From *odus*, a tooth, and *algos*, pain.) The tooth-ach. This well-known disease makes its attack by a most violent pain in the teeth, most frequently in the molares, more rarely in the incisori, reaching sometimes up to the eyes, and sometimes backwards into the cavity of the ear. At the same time, there is a manifest determination to the head, and a remarkable tension and inflation of the vessels takes place, not only in the parts next to that where the pain is seated, but over the whole head.

The tooth-ach is sometimes merely a rheumatic affection, arising from cold, but more frequently from a carious tooth. It is also a symptom of pregnancy, and takes place in some nervous disorders. It may attack persons at any period of life, though it is most frequent in the young and plethoric. From the variety of causes which may produce this affection, it has been named by authors odontalgia cariosa, scorbutica, catarrhalis, arthritica, gravidarum, hysterica, stomachica, and rheumatica.

ODONTALGICA. (*Medicamenta odontalgica*; from *odontalgia*, the tooth-ach.) Medicines which relieve the tooth-ach.

Many empirical remedies have been proposed for the cure of the tooth-ach, but have not in any degree answered the purpose. When the affection is purely rheumatic, blistering behind the ear will almost always remove it; but when it proceeds from a carious tooth, the pain is much more obstinate. In this case it has been recommended to touch the pained part with a hot iron, or with oil of vitriol, in order to destroy the aching nerve; to hold spirits in the mouth; to put a drop of oil of cloves into the hollow of the tooth, or a pill made of camphor, opium, and oleum caryophylli. Others recommend gum mastich, dissolved in oleum terebinthinæ, applied to the tooth upon a little cotton. The great Boerhaave is said to have applied camphor, opium, oleum caryophylli, and alcohol, upon cotton. The caustic oil which may be collected from writing paper, rolled up tight, and set fire to at the end, will sometimes destroy the exposed nervous sub-

stance of a hollow tooth. The application of *radix pyrethri*, by its power of stimulating the salivary glands, either in substance or in tincture, has also been attended with good effects. But one of the most useful applications of this kind, is strong nitrous acid, diluted with three or four times its weight of spirit of wine, and introduced into the hollow of the tooth, either by means of an hair pencil or a little cotton. When the constitution has had some share in the disease, the Peruvian bark has been recommended, and perhaps with much justice, on account of its tonic and antiseptic powers. When the pain is not fixed to one tooth, leeches applied to the gum are of great service. But very often all the foregoing remedies will fail, and the only infallible cure is to draw the tooth.

ODONTIASIS. (From *odontos*, to pull forth the teeth.) Dentition, or cutting teeth.

ODONTICA. (From *odus*, a tooth.) Remedies for pains in the teeth.

ODONTIRHEA. (From *odus*, a tooth, and *rho*, to flow.) Bleeding from the socket of the jaw, after drawing the tooth.

ODONTIS. (From *odus*, a tooth; so called because its decoction was supposed useful in relieving the tooth-ach.) *Odontitis*. A species of *lychnis*.

ODONTITIS. See *Odontis*.

ODONTOGYPHUM. (From *odus*, a tooth, and *γυφω*, to scrape.) An instrument for scaling and scraping the teeth.

ODONTOID. (*Odontoides*; from *odus*, a tooth, and *oides*, form, because it is shaped like a tooth.) Tooth-like. A process of the second vertebra of the neck is so called. See *Dentatus*.

ODONTOLITHOS. (From *odus*, a tooth, and *lithos*, a stone.) The tartar, or stony crust upon the teeth.

ODONTOPHYIA. (From *odus*, a tooth, and *φυω*, to grow.) Dentition, or cutting teeth.

ODONTOTRIKMA. (From *odus*, a tooth, and *τριχω*, to wear away.) A dentifrice, or medicine, to clean the teeth.

ODORIFEROUS GLANDS. *Glandule odoriferae*. These glands are situated around the corona glandis of the male, and under the skin of the labia majora and nymphæ of females. They secrete a sebaceous matter, which emits a peculiar odour; hence their name.

OE. (*Oon*: from *oon*, to bear; so named from its fruitfulness.) The service tree.

ECONOMY, ANIMAL. (From *oikos*, a house, and *νομος*, a law.) *Economia animalis*. The conduct of nature in preserving animal bodies is called the animal economy.

OEDEMA. (From *oedem*, to swell.) A synonym of *anasarca*. See *Anasarca*.

OEDEMATODES. Like to an *oedema*.

OEDEMSARCA. (From *oedema*, a swelling, and *sarx*, flesh.) A species of tumour mentioned by M. A. Severinus, of a middle nature, betwixt an *oedema*, or soft tumour, and *sarcoma*, or hard tumour.

OENANTHE. (From *oen*, wine, and *anthos*, a flower; so called because its flowers smell like the vine.) 1. The botanical name of a genus of the umbelliferous plants. Class, *Pentandria*. Order, *Digynia*. 2. The pharmacopœial name of hemlock dropwort. *Oenanthe cherophylli foliis*. (*Oenanthe crocata* of Linnæus. An active poison that has too often proved fatal, by being eaten in mistake instead of water parsnep. The juice, nevertheless, cautiously exhibited, promises to be an efficacious remedy in inveterate scorbutic eruptions. The root of this plant is not unpleasant to the taste, and esteemed to be most deleterious of all the vegetables which this country produces. Mr. Howell, surgeon at Haverfordwest, relates, that "eleven French prisoners had the liberty of walking in and about the town of Pembroke. Three of them being in the fields a little before noon, dug up a large quantity of this plant, which they took to be wild celery, to eat with their bread and butter for dinner. After washing it, they all three ate, or rather tasted of the roots. As they were entering the town, without any previous notice of sickness at the stomach, or disorder in the head, one of them was seized with convulsions. The other two ran home, and sent a surgeon to him. The surgeon endeavoured first to bleed, and then to vomit him; but those endeavours were fruitless, and he died presently. Ignorant of the cause of their comrade's death, and of their own danger, they gave of these roots to the other eight prisoners, who ate of them with their dinner. A few minutes afterwards, the remaining two, who gathered the plants, were seized in the same manner as the first, of which one died; the other was bled, and a vomit, with great difficulty, forced down, on account of his jaws being, as it were, locked together. This operated, and he recovered, but was some time affected with dizziness in his head, though not sick, or the least disordered in the stomach. The other eight being bled and vomited immediately were soon well. At Clonwell, in Ireland, eight boys, mistaking this plant for water-parsnep, ate plentifully of its roots. About four or five hours after, the eldest boy became suddenly convulsed, and died; and before the next morning four of the other boys died in a similar manner. Of the other three, one was maniacal several hours, another lost his hair and nails, but the third escaped unhurt. Stalpaert Vander Wiel mentions two cases of the fatal effects of this root: these, however, were

attended with great heat in the throat and stomach, sickness, vertigo, and purging; they both died in the course of two or three hours after eating the root. Allen, in his *Synopsis Medicinæ*, also relates that four children suffered greatly by eating this poison. In these cases great agony was experienced before the convulsions supervened; vomitings likewise came on, which were encouraged by large draughts of oil and warm water, to which their recovery is ascribed. The late Sir William Watson, who refers to the instances here cited, also says, that a Dutchman was poisoned by the leaves of the plant boiled in pottage. It appears, from various authorities, that most brute animals are not less affected by this poison than man; and Mr. Lightfoot informs us, that a spoonful of the juice of this plant, given to a dog, rendered him sick and stupid; but a goat was observed to eat the plant with impunity. The great virulence of this plant has not, however, prevented it from being taken medicinally. In a letter from Dr. Poulteney to Sir William Watson, we are told that a severe and inveterate cutaneous disorder was cured by the juice of the root, though not without exciting the most alarming symptoms. Taken in the dose of a spoonful, in two hours afterwards, the head was affected in a very extraordinary manner, followed with violent sickness and vomiting, cold sweats, and rigors; but this did not deter the patient from continuing the medicine, in somewhat less doses, till it effected a cure.

ENANTHE CROCATÆ. The systematic name of the hemlock dropwort. See *Oenanthe*.

ENANTHE CHEROPHYLLI FOLIIÆ. See *Oenanthe*.

ENANTHE CUCUTÆ FACIÆ LOBELII. The *Enanthe crocata*.

ENABEA. (*Onyx*: from *onaga*, the cuttings of vines.) The ashes prepared of the twigs, &c. of vines.

ENELHEUM. (From *onox*, wine, and *elaeon*, oil.) A mixture of oil and wine.

ENOGALA. (From *onox*, wine, and *gala*, milk.) A sort of potion made of wine and milk. According to some, it is wine as warm as new milk.

ENOGARUM. (From *onox*, wine, and *garum*, garum.) A mixture of wine and garum.

ENOMELI. (From *onox*, wine, and *melas*, honey.) Mead, or wine, made of honey, or sweetened with honey.

ENOPLIA. (From *onox*, wine.) The great jubeb-tree, the juice of whose fruit is like that of the grape.

ENOSTAGMA. (From *onox*, wine, and *stagna*, to distil.) Spirit of wine.

ENOTHEKA. (From *onox*, wine; so called because its dried roots smell like wine.) A species of *lysimachia*.

GENUS ANTHINOS. (From *anthos*, a flower.) Flowery wine. Galen says it is *Oenose anthosmias*, or wine impregnated with flowers, in which sense it is an epithet for the *Cyceon*.

GENUS ANTHOSMIAS. (From *anthos*, a flower, and *osmia*, a smell.) Sweet-scented wines.

GENUS APODEXUS. Wine in which the dais, or *tæda*, are boiled.

GENUS APRESMENTUS. A wine heated to a great degree, and prescribed among other things, as garlic, salt, milk, and vinegar.

GENUS DEUTERUS. (*Δευτερος*, second.) Wines of the second pressing.

GENUS DIACHROMENUS. Wine diffused in larger vessels, cooled, and strained from the lees, to render it thinner and weaker; wines thus drawn off are called *saccus*, and *saccata*, from the bag through which they are strained.

GENUS GALACTODES. (From *γала*, milk.) Wine with milk, or wine made as warm as new milk.

GENUS MALACUS. *Ænus malthacus*. Soft wine. Sometimes it means weak and thin, opposed to strong wine; or mild, in opposition to austere.

GENUS MELICEROOS. Wine in which is honey.

GENUS OENODES. Strong wine.

GENUS STRAPHIDIOS LEUCOS. White wine made from raisins.

GENUS TETHALASMENOS. Wine mixed with sea-water.

ÆSOPHAGUS. (From *αισωφαγος*, the gullet.) The muscle forming the sphincter *æsophi*.

ÆSOPHAGISINUS. (From *αισωφαγος*, the gullet.) Difficult swallowing, from spasm.

ÆSOPHAGUS. (From *αισ*, to carry, and *φαγε*, to eat; because it carries the food into the stomach.) The membranous and muscular tube that descends in the neck, from the pharynx to the stomach. It is composed of three tonics, or membranes, viz. a common, muscular, and mucous. Its arteries are branches of the *æsoophageal*, which arises from the aorta. The veins empty themselves into the *vena azygos*. Its nerves are from the eighth pair and great intercostal; and it is every where under the internal or mucous membrane supplied with glands that separate the mucus of the *æsophagus*, in order that the masticated bolus may readily pass down into the stomach.

ÆSTROMANIA. (From *αισρος*, the pudenda of a woman, and *μανια*, to rage.) A furor uterinus.

ÆSTRUM VENEREUM. (From *æstrus*, a gad-bee; because by its bite, or sting, it agitates cattle.) The venereal orgasm, or pleasant sensation experienced during coition.

CESTR. (From *ce*, a sheep, and *gros*, borders.) *Oestropos*. *Oestypum*. *Oestypus*. It frequently is met with in the ancient Pharmacy, for a certain oily substance, boiled out of particular parts of the fleeces of wool, as what grows on the flank, neck, and parts most used to sweat.

OTTA ALBA. (From *phath*, a fragment, Heb.) Van Helmont thus calls the white coagulation which arises from a mixture of a rectified spirit of wine, and of urine; but the spirit of urine must be distilled from well-fermented urine; and that must be well dephlegmated, else it will not answer.

OFFICINAL. (*Officinalis*; from *officina*, a shop.) Any medicine, directed by the colleges of physicians to be kept in the shops, is so termed.

OPTRUCATIO. The same as *Amaurosis*.

OIL. (*Oleum*; from *olea*, the olive; this name being at first confined to the oil expressed from the olive.) Oils are defined, by modern chymists, to be proper juices of a fat or unctuous nature, either solid or fluid, indissoluble in water, combustible with flame, and volatile in different degrees. They are never formed but by organic bodies; and all the substances in the mineral kingdom, which present oily characters, have originated from the action of vegetable or animal life. Oils are distinguished into fat, and essential oils; under the former head are comprehended oil of olives, almonds, rape, ben, linseed, hemp, cocoa, &c. Essential oils differ from fat oils by the following characters: their smell is strong and aromatic; their volatility is such that they rise with the heat of boiling water, and their taste is very acrid; they are likewise much more combustible than fat oils; they are obtained by pressure, distillation, &c. from strong-smelling plants, as that of peppermint, aniseed, caraway, &c. The use of fat oils in the arts, and in medicine, is very considerable; they are medicinally prescribed as relaxing, softening, and laxative remedies; they enter into many medical compounds, such as balsams, unguents, plasters, &c. and they are often used as food on account of the mucilage they contain. See *Oliwa*. Essential oils are employed as cordial, stimulant, and antispasmodic remedies.

Oil, atherial. See *Oleum atheriale*.

Oil, almond. See *Amygdala*.

Oil of allepice. See *Oleum pimentis*.

Oil of amber. See *Oleum succini*.

Oil of caraway. See *Oleum carvi*.

Oil, castor. See *Ricinus*.

Oil of chamomile. See *Oleum anthemidis*.

Oil of juniper. See *Oleum juniperi*.

Oil of lavender. See *Oleum lavendulae*.

Oil of linseed. See *Oleum lini*.

Oil of mace. See *Oleum macis*.

Oil, olive. See *Oliwa*.

Oil of origanum. See *Oleum origani*.

Oil, palm. See *Palm oil*.

Oil of pennyroyal. See *Oleum pulegii*.

Oil of peppermint. See *Oleum menthae piperitis*.

Oil, rock. See *Petroleum*.

Oil of spearmint. See *Oleum menthae viridis*.

Oil, sulphurated. See *Oleum sulphuratum*.

Oil of turpentine. See *Oleum terebinthae*.

Ointment. See *Unguentum*.

OLEA. The name of a genus of plants in the Linnæan system. Class, *Monandria*. Order, *Monogynia*.

OLEA EUROPEA. The systematic name of the plant from which the olive oil is obtained. See *Oliwa*.

OLEAMEN. (From *oleum*, oil.) A thin liniment composed of oils.

OLEANDER. (From *olea*, the olive-tree, which it resembles.) The rose bay,

OLEASTER. (Dim. of *olea*, the olive-tree.) The wild olive.

OLECRANON. (From *olæon*, the ulna, and *cranon*, the head.) The elbow, or process of the ulna, upon which a person leans.

OLENE. (*ὀλῆν*.) The cubit, or ulna.

OLEOSACCHARUM. (From *Oleum*, oil, and *saccharum*, sugar.) An essential oil, ground up with sugar.

OLEUM. See *Oil*.

OLEUM ABSTINUM. The resinous juice which exudes spontaneously from the silver and red firs. It is supposed to be superior to that obtained by wounding the tree.

OLEUM ÆTHEREUM. Atherial oil. *Oleum vini*. After the distillation of sulphuric æther, carry on the distillation with a less degree of heat, until a black froth begins to rise; then immediately remove the retort from the fire. Add sufficient water to the liquor in the retort, that the oily part may float upon the surface. Separate this, and add to it as much lime-water as may be necessary to neutralize the adherent acid, and shake them together. Lastly, collect the atherial oil which separates. This oil is used as an ingredient in the compound spirit of æther. It is a yellow colour, less volatile than æther, soluble in alcohol, and insoluble in water.

OLEUM AMYGDALÆ. See *Amygdala*.

OLEUM ANIMALE. An empyreumatic substance, obtained by distillation from animal substances. It is sometimes exhibited as an antispasmodic and diaphoretic, in the dose of from ten to forty drops.

OLEUM ANISI. Formerly *Oleum essentielle anisi*, *oleum e semibus anisi*. Oil of anise. The essential oil of aniseed possesses all the virtues attributed to the anisum, and is often given as a stimulant and carminative, in the dose of from five to eight drops.

mixed with an appropriate vehicle. See *Anisum*.

OLEUM ANTHEMIDIS. Oil of chamomile, formerly called *oleum e floribus chamamelis*. See *Chamæmelum*.

OLEUM CAMPHORATUM. In retentions of urine, rheumatic pains, distentions of the abdomen from ascites; tension of the skin from abscess, this is an excellent application. See *Camphora*.

OLEUM CARPATHICUM. A fine essential oil, distilled from the fresh cones of the tree which affords the common turpentine. See *Terebinthina vulgaris*.

OLEUM CARUI. Formerly called *Oleum essentielle carui*. *Oleum essentielle e seminibus carui*. The oil of carraways is an admirable carminative, diluted with rectified spirit into an essence, and then mixed with any proper fluid. See *Carum*.

OLEUM CARYOPHYLLIA ROMATICI. A stimulant and aromatic preparation of the clove. See *Caryophyllum aromaticum*.

OLEUM CEDRINUM. *Essentia de cedro*. The oil of the peel of citrons, obtained in a particular manner, without distillation, in Italy.

OLEUM CINNAMOMI. A warm, stimulant, and delicious stomachic. Given in the dose of from one to three drops, rubbed down with some yolk of egg, in a little wine, it allays violent emotions of the stomach from morbid irritability, and is particularly serviceable in debility of the primæ viæ, after cholera morbus.

OLEUM CORNI CERVI. This is applied externally as a stimulant to paralytic affections of the limbs.

OLEUM GABIANUM. See *Petroleum rubrum*.

OLEUM JUNIPERI. Formerly called *Oleum essentielle juniperi baccæ*. *Oleum essentielle e baccis juniperi*. Oil of juniper. Oil of juniper berries possesses stimulant, carminative, and stomachic virtues, in the dose of from two to four drops, and in a larger dose proves highly diuretic. It is often administered in the cure of dropsical complaints, when the indication is to provoke the urinary discharge.

OLEUM LAVENDULÆ. Formerly called *Oleum essentielle lavendulæ*. *Oleum essentielle e floribus lavendulæ*. Oil of lavender. Though mostly used as a perfume, this essential oil may be exhibited internally, in the dose of from one to five drops, as a stimulant in nervous head-achs, hysteria and debility of the stomach.

OLEUM LAURI. *Oleum laurinum*. An anodyne and antispasmodic application, generally rubbed on sprains and bruises unattended with inflammation.

OLEUM LIMONIS. The essential oil of lemons possesses stimulant and stomachic powers, but is principally used externally, mixed with ointments as a perfume.

OLEUM LINI. Linseed oil is emollient and demulcent, in the dose of from half an ounce to an ounce. It is frequently given in the form of clyster in colics and obstipation. Cold-drawn linseed oil, with lime-water and extract of lead, forms, in many instances, the best application for burns and scalds. See *Linum*.

OLEUM LUCII PISCIS. See *Esox lucius*.

OLEUM MACIS. *Oleum myristice expressum*. Oil of mace. A fragrant sebaceous substance, expressed in the East Indies from the nutmeg. There are two kinds. The best is brought in stone jars, is somewhat soft, of a yellow colour, and resembles in smell the nutmeg. The other is brought from Holland, in flat square cakes. The weak smell and faint colour warrants our supposing it to be the former kind sophisticated. Their use is chiefly external, in form of plaster, unguent, or liniment.

OLEUM MALABATHI. An oil similar in flavour to that of cloves, brought from the East Indies, where it is said to be drawn from the leaves of the cinnamon tree.

OLEUM MENTHÆ PIPERITÆ. Formerly called *Oleum essentielle menthæ piperitidis*. Oil of peppermint. Oil of peppermint possesses all the active principle of the plant. It is mostly used to make the simple water; mixed with rectified spirit it forms an essence, which is put into a variety of compounds, as sugar drops and trochiscs, which are exhibited as stimulants, carminatives, and stomachics.

OLEUM MENTHÆ VIRIDIS. Formerly called *Oleum essentielle menthæ sativæ*. Oil of spearmint. This essential oil is mostly in use for making the simple water, but may be exhibited in the dose of from five to two drops as a carminative, stomachic, and stimulant.

OLEUM NEROLI. *Essentia neroli*. The essential oil of the flowers of the Seville orange tree. It is brought to us from Italy and France.

OLEUM MYRISTICÆ. The essential oil of nutmeg is an excellent stimulant and aromatic, and may be exhibited in every case where such remedies are indicated, with advantage.

OLEUM MYRISTICÆ EXPRESSUM. This is commonly called oil of mace. See *Oleum macis*.

OLEUM NITRIOLI. See *Sulphureous acid*.

OLEUM OLIVÆ. See *Olea*.

OLEUM ORIGANI. Formerly called *Oleum essentielle origani*. Oil of origanum. A very acrid and stimulating essential oil. It is employed for alleviating the pain arising from caries of the teeth, and for making the simple water of marjoram.

OLEUM PALMÆ. See *Palm oil*.

OLEUM PETRÆ. See *Petroleum*.

OLEUM PINENTÆ. Oil of allspice. A stimulant and aromatic oil.

OLEUM PULEGII. Formerly called *Oleum essentielle pulegii*. Oil of penny-royal. A stimulant and antispasmodic oil, which may be exhibited in hysterical and nervous affections.

OLEUM RICINI. See *Ricinus*.

OLEUM RORIS MARINI. Formerly called *Oleum essentielle roris marini*. Oil of rosemary. The essential oil of rosemary is an excellent stimulant, and may be given with great advantage in nervous and spasmodic affections of the stomach.

OLEUM SABINÆ. A stimulating emmenagogue; it is best administered with myrrh, in the form of bolus.

OLEUM SASSAFRÆ. An agreeable stimulating stomachic carminative and sudorific.

OLEUM SINAPEOS. This is an emollient oil, the acrid principle of the mustard remaining in the seed. See *Sinapis*.

OLEUM SUCCINI. *Oleum succini rectificatum*. Oil of amber is mostly used externally, as a stimulating application to paralytic limbs, or those affected with cramp and rheumatism. Hooping-cough, and other convulsive diseases, are said to be relieved also by rubbing the spine with this oil.

OLEUM SULPHURATUM. Formerly called *Balsamum sulphuris simplex*. Sulphurated oil. "Take of washed sulphur, four ounces; olive oil, a pint." Having heated the oil in a very large iron pot, add the sulphur gradually, and stir the mixture after each addition, until they have united. This, which was formerly called simple balsam of sulphur, is an acrid stimulating preparation, and much praised by some in the cure of coughs and other phthisical complaints.

OLEUM SYRIÆ. A fragrant essential oil, obtained by distillation from the balm of Gilead plant. See *Moldavica*.

OLEUM TEMPLINUM. *Oleum templinum verum*. A terebinthinate oil obtained from the fresh cones of the *Pinus abies* of Linnæus.

OLEUM TERRÆ. See *Petroleum*.

OLEUM TEREBINTHINÆ RECTIFICATUM. "Rectified oil of turpentine, a pint; water, four pints." Distil over the oil. Stimulant, diuretic, and sudorific virtues are attributed to this preparation, in the dose of from ten drops to twenty, which are given in rheumatic pains of the chronic kind, especially sciatica. Its chief use internally, however, is as an anthelmintic and styptic. Uterine, pulmonary, gastric, intestinal, and other hemorrhages, when passive, are more effectually relieved by its exhibition than by any other medicine. Externally it is applied, mixed with

ointments, and other applications, to bruises, sprains, rheumatic pains, indolent ulcers, burns, and scalds.

OLEUM VINI. Stimulant and anodyne in the dose of from one to four drops.

OLFACTORY NERVES. (Nervi olfactorii; from olfactus, the sense of smelling.) The first pair of nerves are so termed, because they are the organs of smelling. They arise from the corpora striata, perforate the ethmoid bone, and are distributed very numerously on the pituitary membrane of the nose.

OLIBANUM. (From lebana, Chald.) Thus. Frankincense. This drug has received different appellations, according to its different appearances; the single tears are called simply olibanum, or thus; when two are joined together, thus masculum; and when two are very large, thus femininum; if several adhere to the bark, thus corticosum; the fine powder, which rubs off from the tears, mica thuris; and the coarser, manna thuris. The gum-resin that is so called, is the juice of the *Juniperus lycias*—*foliis ternis undique imbricatis ovatis obtusis*, and is brought from Turkey and the East Indies; but that which comes from India is less esteemed. It is said to ooze spontaneously from the bark of the tree, appearing in drops, or tears, of a pale yellowish, and sometimes of a reddish colour. Olibanum has a moderately strong and not very agreeable smell, and a bitterish, somewhat pungent taste; in chewing it sticks to the teeth, becomes white, and renders the saliva milky. Laid on a red-hot iron, it readily catches flame, and burns with a strong diffusive and not unpleasant smell. On trituration with water, the greatest part of it dissolves into a milky liquor, which, on standing, deposits a portion of resinous matter. The gummy and resinous parts are nearly in equal proportions; and though rectified spirit dissolves less of the olibanum than water, it extracts nearly all its active matter. In antient times, olibanum seems to have been in great repute in affections of the head and breast, coughs, hæmoptysis, and in various fluxes, both uterine and intestinal; it was also much employed externally. Recourse is seldom had to this medicine, which is now superseded by myrrh, and other articles of the resinous kind. It is, however, esteemed by many as an adstringent, and though not in general use, is considered as a valuable medicine in fluor albus, and debilities of the stomach and intestines; applied externally in the form of plaster, it is said to be corroborant, &c. and with this intention it forms the basis of the emplastrum thuris.

OLISOTROPHIA. (From *olôgê*, small, and *trôphê*, to nourish.) Deficient nourishment.

OLISTHENA. (From *olisthênai*, to fall out.) A luxation.

OLIVA. The olive. *Olea sativa*. The fruit of the *Olea Europea* of Linnaeus: *foliis lanceolatis integerrimis, racemosis, axillaribus coarctatis*. The olive-tree, in all ages has been greatly celebrated, and held in peculiar estimation, as the bounteous gift of heaven; it was formerly exhibited in the religious ceremonies of the Jews, and is still considered as emblematic of peace and plenty. The varieties of this tree are numerous, distinguished not only by the form of the leaves but also by the shape, size, and colour of the fruit; as the large Spanish olive, the small oblong Provence olive, &c. &c. These, when pickled, are well known to us by the names of Spanish and French olives, which are extremely grateful to many stomachs, and said to excite appetite and promote digestion; they are prepared from the green unripe fruit, which is repeatedly steeped in water, to which some quicklime or alkaline salt is added, in order to shorten the operation: after this they are washed and preserved in a pickle of common salt and water, to which an aromatic is sometimes added. The principal consupution, however, of this fruit is in the preparation of the common salad oil, or *oleum olivæ* of the pharmacopœias, which is obtained by grinding and pressing them when thoroughly ripe: the finer and purer oil issues first by gentle pressure, and the inferior sorts on heating what is left, and pressing it more strongly. The best olive oil is of a bright pale amber colour, bland to the taste, and without any smell: it becomes rancid by age, and sooner if kept in a warm situation. With regard to its utility, oil, in some shape, forms a considerable part of our food, both animal and vegetable, and affords much nourishment. With some, however, oily substances do not unite with the contents of the stomach, and are frequently brought up by eructation; this happens more especially to those whose stomachs abound with acid.—Oil, considered as a medicine, is supposed to correct acrimony, and to lubricate and relax the fibres; and therefore has been recommended internally, to obviate the effects of various stimuli, which produce irritation, and consequent inflammation: on this ground it has been generally prescribed in coughs, catarrhal affections, and erosions. The oil of olives is successfully used in Switzerland against the *tenia oculi superficialibus*, and it is in very high estimation in this and other countries against nephritic pains, spasms, cholic, constipation of the bowels, &c. Externally it has been found a useful application to bites and stings of various poisonous animals, as the mad dog, several serpents, &c. also to burns, tumours, and other affections, both by itself or mixed in liniments or poultices. Oil rubbed over the body is said to

be of great service in dropæles, particularly ascites. Olive oil enters several official compositions, and when united with water, by the intervention of alkali, is usually given in coughs and hoarseness.

Olive-tree. See *Olida*.

Olives. See *Oliva*.

Olive, spurge. See *Mezereum*.

OLIVARIUS. (From *oliva*, the olive.) *Oliviformis*. Resembling the olive; applied to two eminences on the lower part of the medulla oblongata, called *corpora olivaria*.

OLOPHYCTIS. (From *ολος*, whole, and *φύκτις*, a pustule.) A small hot eruption, covering the whole body; when partial, it is called *phlyctæna*.

OLUSATRUM. (*Id est, olus atrum*, the black herb, from its black leaves) Lovage.

OMAGRA. (From *ομας*, the shoulder, and *αγος*, a seizure.) The gout in the shoulder.

OMENTITIS. (*Omentitis*; from *omentum*, the caul.) Inflammation of the omentum, a species of peritonitis.

OMENTUM. (From *omen*, a guess; so called because the soothsayers prophesied from an inspection of this part) *Epiploon*. The caul. An adipose membranous viscus of the abdomen, that is attached to the stomach, and lies on the anterior surface of the intestines. It is thin and easily torn, being formed of a duplicature of the peritoneum, with more or less of fat interposed. It is distinguished into the great omentum and the little omentum.

The *omentum majus*, which is also termed *omentum gastrocolicum*, arises from the whole of the great curvature of the stomach, and even as far as the spleen, from whence it descends loosely behind the abdominal parietes, and over the intestines to the navel, and sometimes into the pelvis. Having descended thus far, its inferior margin turns inwards and ascends again, and is fastened to the colon and the spleen, where its vessels enter.

The *omentum minus*, or *omentum hepato-gastricum*, arises posteriorly from the transverse fissure of the liver. It is composed of a duplicature of peritoneum, passes over the duodenum, and small lobe of the liver; it also passes by the lobulus spigelii and pancreas, proceeds into the colon and small curvature of the stomach, and is implanted ligamentous into the œsophagus. It is in this omentum that Winslow discovered a natural opening, which goes by his name. If air be blown in at the foramen of Winslow, which is always found behind the lobulus spigelii, between the right side of the liver and hepatic vessels, the vena portarum and duodenum, the cavity of the omentum, and all its sacs may be distended.

The omentum is always double, and between its lamellæ closely connected by very tender cellular substance, the vessels

are distributed and the fat collected. Where the top of the right kidney, and the lobulus spigelii of the liver, with the subjacent large vessels, form an angle with the duodenum, there the external membrane of the colon, which comes from the peritoneum joining with the membrane of the duodenum, which also arises immediately from the peritoneum lying upon the kidney, enters the back into the transverse fissure of the liver, for a considerable space, is continuous with its external coat, contains the gall-bladder supports the hepatic vessels, and is very yellow and slippery. Behind this membranous production, betwixt the right lobe of the liver, hepatic vessels, vena portarum, biliary ducts, aorta, and adjacent duodenum, there is the natural opening just mentioned, by which air may be blown extensively into all the cavity of the omentum. From thence, in a course continuous with this membrane from the pylorus and the smaller curvature of the stomach, the external membrane of the liver joins in such a manner with that of the stomach, that the thin membrane of the liver is continued out of the fossa of the venal duct, across the little lobe into the stomach, stretched before the lobe and before the pancreas. This little omentum, or *omentum hepatico-gastricum*, when inflated, resembles a cone, and gradually becoming harder and emaciated, it changes into a true ligament, by which the œsophagus is connected to the diaphragm. But the larger omentum, the *omentum gastro-colicum*, is of a much greater extent. It begins at the first accession of the right gastro-epiploic artery to the stomach, being continued there from the upper plate of the transverse mesocolon; and then from the whole great curve of the stomach, as far as the spleen, and also from the right convex end of the stomach towards the spleen, until it also terminates in a ligament, that ties the upper and back part of the spleen to the stomach: this is the anterior lamina. Being continued downward, sometimes to the navel, sometimes to the pelvis, it hangs before the intestines, and behind the muscles of the abdomen, until its lower edge being reflected upon itself, ascends, leaving an intermediate vacuity between it and the anterior lamina, and is continued to a very great extent, into the internal membrane of the transverse colon, and lastly, into the sinus of the spleen, by which the large blood-vessels are received, and it ends finally on the œsophagus, under the diaphragm. Behind the stomach, and before the pancreas, its cavity is continuous with that of the smaller omentum. To this the *omentum colicum* is connected, which arises farther to the right than the first origin of the omentum gastrocolicum from the mesocolon, with the cavity of which it is continuous, but produced solely from the

colon and its external membrane, which departs double from the intestine; it is prolonged, and terminates by a conical extremity, sometimes of longer, sometimes of shorter extent, above the intestinum cæcum. For all the blood which returns from the omentum and mesocolon, goes into the vena portarum, and by that into the liver itself. The omentum gastrocolicum is furnished with blood from each of the gastro-epiploic arteries, by many descending articulated branches, of which the most lateral are the longest, and the lowest anastomose by minute twigs with those of the colon. It also has branches from the splenic, duodenal, and adipose arteries. The omentum colicum has its arteries from the colon, as also the smaller appendices, and also from the duodenal and right epiploic. The arteries of the small omentum come from the hepatics, and from the right and left coronaries. The omentum being fat and indolent, has very small nerves. They arise from the nerves of the eighth pair, both in the greater and lesser curvatures of the stomach. The arteries of the mesentery are in general the same with those which go to the intestine, and of which the smaller branches remain in the glands and fat of the mesentery. Various small accessory arteries go to both mesocolons, from the intercostals, spermatics, lumbar, and capsular, to the transverse portion from the splenic artery, and pancreato-duodenalis, and to the left mesocolon, from the branches of the aorta going to the lumbar glands. The veins of the omentum in general accompany the arteries, and unite into similar trunks; those of the left part of the gastrocolic omentum into the splenic, and also those of the hepato-gastric, which likewise sends its blood to the trunk of the vena portarum; those from the larger and right part of the gastrocolic omentum, from the omentum colicum, and from the appendices epiploides, into the mesenteric trunk. All the veins of the mesentery meet together in one wick, in the true trunk of the large vena portarum, being collected first into two large branches, of which the one, the mesenteric, receives the gastro-epiploic vein, the colicæ mediz, the ilio-colicæ, and all those of the small intestines, as far as the duodenum; the other, which going transversely, inserts itself into the former, above the origin of the duodenum, carries back the blood of the left gastric veins, and those of the rectum, except the lowermost, which belongs partly to those of the bladder and partly to the hypogastric branches of the pelvis. The vein which is called hæmorrhoidalis interna is sometimes inserted rather into the splenic than into the mesenteric vein. Has the omentum also lymphatic vessels? Certainly there are conglobate glands, both in the little omentum and in

the gastro-colicum; and ancient anatomists have observed pellucid vessels in the omentum; and a modern has described them for lacteals of the stomach.

OMENTUM COLICUM. See *Omentum*.

OMENTUM GASTRO-COLICUM. See *Omentum*.

OMENTUM HEPATICO-GASTRICUM. See *Omentum*.

OMO. Names compounded with this word belong to muscles which are attached to the scapula; from *omos*, the shoulder, As,

OMOCOTYLE. (From *omos*, the shoulder, and *cotyla*, a cavity.) The cavity in the extremity of the neck of the scapula, in which the head of the humerus is articulated.

OMO-HYOIDEUS. *Caroco-hyoideus* of Alpinus and Douglas, *scapulo-hyoiden* of Dumas. A muscle situated between the os hyoides and shoulder, that pulls the os hyoides obliquely downwards. It arises broad, thin, and fleshy, from the superior costa of the scapula, near the semilunar notch, and from the ligament that runs across it; thence ascending obliquely, it becomes tendinous below the sternocleidomastoideus; and growing fleshy again, is inserted into the base of the os hyoides.

OMOPLATA. (From *omos*, the shoulder, and *πλατος*, broad.) See *Scapula*.

OMOPLATO-HYOIDEUS. The same, as *Omohyoideus*.

OMOTOCOS. (From *omos*, crude, and *τοκο*, to bring forth.) A miscarriage.

OMOTRIBES. (From *omos*, crude, and *τριβω*, to bruise.) Oil expressed from unripe olives.

OMPHACINUM. (From *ομφαλινον*, the juice of unripe grapes.) Oil expressed from unripe olives.

OMPHACION. *Omphacium*. (From *ομφακος*, an unripe grape.) The juice of unripe grapes; and by some applied to that of wild apples, or crabs, commonly called *Verrucosa*.

OMPHACITIS. (From *ομφακος*, an unripe grape, because it resembles an unripe grape in its sour astringent state.) A small kind of gall; an excrescence from the oak.

OMPHACOMELI. (From *ομφακος*, an unripe grape, and *μελι*, honey.) A sort of oxymel made of the juice of unripe grapes and honey.

OMPHALOCARPUS. (From *ομφαλος*, the navel, and *καρπος*, fruit; so called because its fruit resembles a navel.) Cleavers; hayriff.

OMPHALOCELE. (From *ομφαλος*, the navel, and *κηλη*, a tumour.) An umbilical hernia. See *Hernia*.

OMPHALODES. (From *ομφαλος*, a navel; so named because the calyx is excavated in the middle like the human navel.) A plant resembling borage.

OMPHALOMANTIA. (From *ομφαλος*, the navel, and *μαντις*, to prophesy.) The foolish vaticination of midwives, who pretend to foretell the number of the future

offspring from the number of knots in the navel.

OMPHALOS. (From *ομφαλινον*, to roll up.) The navel.

OMPHALOTOMIA. (From *ομφαλος*, the navel, and *τομη*, to cut.) The separation of the navel-string.

ONASRA. (From *οναγρος*, the wild ass, so called because it is said to tame wild beasts.) Also a name for the rheumatism in the elbow.

ONEIRODYNIA. (From *ονειρον*, a dream, and *δυνω*, anxiety.) Disturbed imagination during sleep. A genus of disease in the class *Neuroses*, and order *Vesaniae* of Cullen, containing two species,

1. *Oneirodynia activa*, walking in the sleep.

2. *Oneirodynia gravans*, the incubus, or night-mare. See *Night-mare*.

ONEIROSMOS. (From *ονειρω*, to dream.) Venereal dreams.

ONEIROSONOS. (From *ονειρον*, a dream, and *σπον*, the seed.) So the Greeks call an occasional emission of the semen in sleep, when it only happens rarely.

Onion. See *Cepa*.

Onion, sea. See *Scilla*.

ONIA. (From *ονος*, an ass.) the dung of an ass. It was in repute with Hippocrates.

ONISCUS. (From *ονος*, an ass; so called because like the ass it requires much beating before it is useful.) The stock-fish. Also the slow-worm.

ONISCUS ABELLUS. The systematic name of the woodlouse. See *Millepedes*.

ONITIS. (From *ονος*, an ass, because asses covet it.) The organum plant.

ONOBRYCHIS. (From *ονος*, an ass, and *βρυχη*, to bray; so called, according to Blanchard, because the smell or taste makes asses bray.) Holy hay: saintfoin; cockshew vetch.

ONONIS. (From *ονος*, an ass, because it interrupts asses when at plough.) 1. The name of a genus of plants in the Linnæan system. Class, *Diadelphia*. Order, *Decandrio*. 2. The pharmacopœial name of the *Resta bovis*. *Arresta bovis*. *Remora aratri*. Rest harrow. The roots of this plant, *Ononis spinosa*, vel *arvensis* of Linnæus, have a faint unpleasant smell, and a sweetish, bitterish, somewhat nauseous taste. Their active matter is confined to the cortical part, which has been sometimes given in powder, or other forms, as an aperient and diuretic.

ONONIS ARVENSIS. The systematic name of the rest harrow. See *Ononis*.

ONONIS SPINOSA. The systematic name of the rest harrow. See *Ononis*.

ONOPORDIUM ACANTHIUM. (*Onopordion*; from *ονος*, an ass, and *πυλω*, to break wind; so named from its being much coveted by asses, and from the noise it makes upon pressure.) The systematic name of the cotton-thistle. See *Cardus tormentosus*.

OPUSMA ECHINIS. The systematic name of the plant whose root is called *echinus luteus* in some pharmacopœias. It is supposed to possess menagogue virtues.

OPUSULA. (From *ovv*, the nail.) A whitlow at the side of the finger nail.

ONYX. *Ovξ.* *Unguis.* An abscess, or collection of pus between the lamellæ of the cornea; so called from its resemblance to the stone called onyx. The diagnostic signs are, a white spot or speck, prominent, soft, and fluctuating. The species are:

1. *Abscessus superficialis*, arising from inflammation, not dangerous, for it vanishes when the inflammation is resolved by the use of astringent collyria.

2. *Abscessus profundus*, or a deep abscess, which is deeper seated between the lamellæ of the cornea, sometimes breaking internally, and forming an hypopium: when it opens externally it leaves a fistula upon the cornea; whenever the pus is evacuated, there remains a leucoma.

OOLIBIA. (From *ovv*, an egg, and *libe*, a likeness.) An epithet for the aqueous humour of the eye.

OPHIOGLOSSIDES. (From *ophioglosson*, ophioglossum, and *libe*, a likeness.) A fungus resembling the adder's tongue.

OPHIOGLOSSUM. (From *ovv*, a serpent, and *glossa*, a tongue; so called from the resemblance of its fruit.) Adder's tongue.

OPHIORRIZA MUNGOS. The systematic name of the plant whose root is called *radix serpentum* in the pharmacopœias. See *Mungos radix*.

OPHIOCORODON. (From *ovv*, a serpent, and *korodon*, garlic, so named because it is spotted like a serpent.) Broad-leaved garlic.

OPHIOTAPHYLUM. (From *ovv*, a serpent, and *taphala*, a berry, so called because serpents feed upon its berries.) White briary. See *Bryonia*.

OPHIOXYLUM SERPENTINUM. The systematic name of the tree whose wood is termed *lignum serpentinum*. See *Serpentinum lignum*.

OPERTS. (*Ovov*. The lowest part of the forehead, where the eye-brows grow. Also an herb so called because its juice was used to make the hair of the eye-brows black.

OPHTHALMIA. (From *ophthalmos*, the eye.) *Ophthalmia*. An inflammation of the membranes of the eye, or of the whole bulb of the eye. The symptoms which characterize this disease are a preternatural redness of the tunica conjunctiva, owing to a turgescence of its blood-vessel; pain and heat over the whole surface of the eye, often attended with a sensation of some extraneous body between the eye and eyelid and a plentiful effusion of tears. All these symptoms are commonly increased by motion of the eye, or its coverings, and likewise by exposure to light. We judge

of the depth of the inflammation by the degree of pain produced by light thrown upon the eye. When the pain produced by light is considerable, we have much reason to imagine that the parts at the bottom of the eye, and especially the retina, are chiefly affected, and, *vice versa*, when the pain is not much increased by this exposure, we conclude with great probability that the inflammation is confined perhaps entirely to the external covering of the eye. In superficial affections of this kind too the symptoms are in general local; but, whenever the inflammation is deep seated, it is attended with severe shooting pains through the head, and fever to a greater or less degree commonly takes place. During the whole course of the disease there is for the most part a very plentiful flow of tears, which frequently become so hot and acrid as to excoriate the neighbouring parts, but it often happens after the disease has been of some duration, that together with the tears a considerable quantity of a yellow purulent-like matter is discharged, and when the inflammation has either spread to the eyelids, or has been seated there from the beginning, as soon as the tarsi becomes affected a discharge takes place of a viscid glutinous kind of matter, which greatly adds to the patient's distress, as it tends to increase the inflammation by cementing the eye-lids so firmly together as to render it extremely difficult to separate them.

Ophthalmia is divided into external, when the inflammation is superficial, and internal, when the inflammation is deep seated, and the globe of the eye is much affected.

In severe ophthalmia two distinct stages are commonly observable; the first is attended with a great deal of heat and pain in the eye, and considerable febrile disorder; the second is comparatively a chronic affection without pain and fever. The eye is merely weakened, moister than in the healthy state, and more or less red.

Ophthalmia may be induced by a variety of exciting causes as operate in producing inflammation in other situations. A severe cold in which the eyes are affected at the same time with the pituitary cavities, fauces, and trachea; change of weather; sudden transition from heat to cold; the prevalence of cold winds, residence in damp or sandy countries, in the hot season; exposure of the eyes to the vivid rays of the sun; are causes usually enumerated, and considering which, it does not seem extraordinary that ophthalmia should often make its appearance as an epidemic, and afflict persons of every age and sex. Besides these exciting causes, writers also generally mention the suppression of some habitual discharge, as of the menses, bleedings from the nose, from hæmorrhoids, &c. Besides which, inflammation of the eyes may be

occasioned by the venereal and scrophulous virus.

OPHTHALMIC GANGLION. *Ganglion ophthalmicum.* Lenticular ganglion. This ganglion is formed in the orbit, by the union of a branch of the third or fourth pair with the first branch of the fifth pair of nerves.

OPHTHALMIC NERVE. *Nervus ophthalmicus.* Orbital nerve. The first branch of the ganglion or expansion of the fifth pair of nerves. It is from this nerve that a branch is given off, to form, with a branch of the sixth, the great intercostal nerve.

OPHTHALMIC EXTENT. See *Meteores oculorum.*

OPHTHALMODYNIA. (From *οφθαλμος*, an eye, and *δυναμις*, pain.) A vehement pain in the eye, without, or with very little redness. The sensation of pain is various, as itching, burning, or as if gravel were between the globe of the eye and lids. The species are :

1. *Ophthalmodynia rheumatica*, which is a pain in the muscular expansions of the globe of the eye, without redness in the albuginea. The rheumatic inflammation is serous, and rarely produces redness.

2. *Ophthalmodynia periodica*, is a periodical pain in the eye, without redness.

3. *Ophthalmodynia spasmodica*, is a pressing pain in the bulb of the eye, arising from spasmodic contractions of the muscles of the eye, in nervous, hysteric, and hypochondriac persons. It is observed to terminate by a flow of tears.

4. *Ophthalmodynia from an internal inflammation of the eye.* In this disorder, there is a pain and sensation as if the globe was pressed out of the orbit.

5. *Ophthalmodynia hydropthalmica.* After a great pain in the inferior part of the os frontis, the sight is obscured, the pupil is dilated, and the bulb of the eye appears larger, pressing on the lid. This species is likewise perceived from an incipient hydropthalmia of the vitreous humour.

6. *Ophthalmodynia arenoea*, is an itching and sensation of pain in the eye, as if sand or gravel were lodged between the globe and lid.

7. *Ophthalmodynia symptomatica*, which is a symptom of some other eye-disease, and is to be cured by removing the exciting cause.

8. *Ophthalmodynia cancerosa*, which arises from cancerous acrimony deposited in the eye, and is rarely curable.

OPHTHALMOPONIA. (From *οφθαλμος*, the eye, and *πονια*, to labour.) An intense pain in the eye, whence the light is intolerable.

OPHTHALMOPYOSIS. (From *οφθαλμος*, an eye, and *πυωσις*, a fall.) A falling down of the globe of the eye on the cheek, canthus, or upwards, the globe itself being

scarcely altered in magnitude. The cause is a relaxation of the muscles, and ligamentous expansions of the globe of the eye. The species are :

1. *Ophthalmoptosis violenta*, which is generated by a violent contusion or strong stroke, as happens sometimes in boxing. The eye falls out of the socket on the cheek or canthus of the eye, and from the elongation and extension of the optic nerve occasions immediate blindness.

2. *Ophthalmoptosis*, from a tumour within the orbit. An exostosis, tophus, abscess, encysted tumours, as, atheroma, hygroma, or schirrhous, forming within the orbit, induration of the orbital adeps, may throw the bulb of the eye out of the socket upwards, downwards, or towards either canthus.

3. *Ophthalmoptosis paralytica*, or the paralytic ophthalmoptosis, which arises from a paralysis or palsy of the recti muscles, from hence a stronger power in the oblique muscles of the bulb.

4. *Ophthalmoptosis staphylomatica*, when the staphyloma depresses the inferior eyelid and extends on the cheek.

OPIATE. (*Medicamentum opiatum*; from the effects being like that of opium.) A medicine that procures sleep, &c. See *Anodynes*.

ΟΠΙΟΝ. (*Opium*.) Opium.

ΟΠΙΣΜΟΣ. (From *οπισθεν*, opium.) An opiate confection.

ΟΠΙΣΤΗΝΑΡ. (From *οπισθεν*, backwards and *ωαγ*, the palm.) The back part of the palm.

ΟΠΙΣΤΗΟΚΡΑΝΙΟΝ. (From *οπισθεν*, backward, and *κρανιον*, the head.) The occiput, or hinder part of the head.

ΟΠΙΣΤΗΟΚΡΩΣΙΣ. (From *οπισθεν*, backward, and *κρωσις*, a gibbosity.) A curved spine.

ΟΠΙΣΤΗΟΤΟΝΟΣ. (From *οπισθεν*, backwards, and *τυνα*, to draw.) A fixed spasm of several muscles, so as to keep the body in a fixed position, and bent backwards. Cullen considers it as a variety of tetanus. See *Tetanus*.

OPIUM. (Probably from *οπιο*, juice; or from *οπι*, Arab.) *Opium Thebaicum*, from being anciently prepared chiefly at Thebes. *Opion* and *manus dei*, so called from its extensive medical properties. Called also by the Arabians *afion*. Afium, Opium is the concreted milky juice which exudes from the unripe heads or capsules of poppies, when incisions are made in them; this juice is gradually exsiccated to a proper consistence.

The species of poppy is the *papaver somniferum*, of Linnæus — *calycibus capsulaeque glabris foliis amplexicaulibus incisatis*. It is brought from Turkey, Egypt, the East Indies, and other parts of Asia, where poppies are cultivated for this use in fields, as corn among us. The manner in which it is collected has been described long ago by

Kämpfer and others; but the most circumstantial detail of the culture of the poppy, and the method of procuring the opium, is that given by Mr. Kerr, as practised in the province of Bahar: he says, "The field being well prepared by the plough and harrow, and reduced to an exact level superficies, it is then divided into quadrangular areas of seven feet long, and five feet in breadth, leaving two feet of interval, which is raised five or six inches, and excavated into an aqueduct for conveying water to every area, for which purpose they have a well in every cultivated field. The seeds are sown in October or November. The plants are allowed to grow six or eight inches distant from each other, and are plentifully supplied with water; when the young plants are six or eight inches high, they are watered more sparingly. But the cultivator spreads all over the areas a nutrient compost of ashes, human excrements, cow-dung, and a large portion of nitrous earth, scraped from the highways and old mud walls. When the plants are nigh-flowering, they are watered profusely, to increase the juice.

When the capsules are half grown, no more water is given, and they begin to collect the opium.

At sunset they make two longitudinal double incisions upon each half ripe capsule, passing from below upwards, and taking care not to penetrate the internal cavity of the capsule. The incisions are repeated every evening until each capsule has received six or eight wounds; then are they allowed to ripen their seeds. The ripe capsules afford little or no juice. If the wound was made in the heat of the day, a cicatrix would be too soon formed. The night dews, by their moisture, favour the exsillation of the juice.

Early in the morning, old women, boys, and girls, collect the juice by scraping it off the wounds with a small iron scoop, and deposit the whole in an earthen pot, where it is worked by the hand in the open sunshine, until it becomes of a considerable spissitude. It is then formed into cakes of a globular shape, and about four pounds in weight, and laid into little earthen basins to be further exsiccated. These cakes are covered over with the poppy or tobacco leaves, and dried until they are fit for sale. Opium is frequently adulterated with cow-dung, the extract of the poppy plant procured by boiling and various other substances which they keep in secrecy."

This process, however, is now but rarely practised, the consumption of this drug being too great to be supplied by that method of collection.

The best sort of the *official opium* is the expressed juice of the heads, or of the heads and the upper part of the stalks in-

spissated by a gentle heat; this was formerly called meconium, in distinction from the true opium, or juice which issues spontaneously.

The inferior sorts (for there are considerable differences in the quality of this drug,) are said to be prepared by boiling the plant in water and evaporating the strained decoction; but as no kind of our opium will totally dissolve in water, the juice is most probably extracted by expression. Newman was informed by some Turks at Genoa and Leghorn, that in some places the heads, stalks, and leaves are committed to the press together, and that this juice inspissated affords a very good opium.

On this head Dr. Lewis remarks, that the point has not yet been fully determined. It is commonly supposed, that whatever preparations the Turks may make from the poppy for their own use, the opium brought to us is really the milky juice collected from incisions made in the heads, as described by Kämpfer. It is certain that an extract made by boiling the heads, or the heads and stalks in water, is much weaker than opium; but it appears also, that the pure milky tears are considerably stronger.

The principles separable from opium are, a resin, gum, a minute proportion of saline matter, water and earth, which are intimately combined together, insomuch that all the three dissolve almost equally in water and in spirit. It is probably to the saline principle Nicholson observes in this and other vegetables that the intimacy of union is in great measure to be ascribed.

Four ounces of opium, treated with alcohol, yielded three ounces and four scruples of resinous extract; five drachms and a scruple of insoluble impurities remaining. On taking four ounces more, and applying water at first, Newman obtained two ounces five drachms and one scruple of gummy extract; the insoluble part amounting here to seven drachms and a scruple. In distillation, alcohol brought over little or nothing; but the distilled water was considerably impregnated with the peculiar ill smell of opium.

From this analysis may be estimated the effects of different solvents upon it. Alcohol and proof spirit dissolving its resin, afford tinctures possessing all its virtues. Water dissolves its gummy part, which is much less active, but a part of the resin is at the same time taken up by the medium of the gum. Wines also afford solutions possessing the virtues of opium. Vinegar dissolves its active matter, but greatly impairs its powers.

The use of this celebrated medicine, though not unknown to Hippocrates, can be clearly traced to Diagoras, who was nearly his cotemporary, and its importance has ever since been gradually advanced by

succeeding physicians of different nations. Its extensive practical utility, however, has not been long well understood; and in this country perhaps may be dated from the time of Sydenham. Opium is the chief narcotic now employed; it acts directly upon the nervous power, diminishing the sensibility, irritability, and mobility of the system; and, according to Cullen, in a certain manner suspending the motion of the nervous fluid to and from the brain, and thereby inducing sleep, one of its principal effects. From this sedative power of opium, by which it allays pain, inordinate action, and restlessness, it naturally follows that it may be employed with advantage in a great variety of diseases. Indeed, there is scarcely any disorder, in which, under some circumstances, its use is not found proper; and though in many cases it fails of producing sleep, yet, if taken in a full dose, it occasions a pleasant tranquillity of mind, and a drowsiness, which approaches to sleep, and which always refreshes the patient. Besides the sedative power of opium, it is known to act more or less as a stimulant, when given in a larger dose, exciting the motion of the blood. By a certain conjoined effort of this sedative and stimulant effect, opium has been thought to produce intoxication, a quality for which it is much used in eastern countries.

The principal indications which opium is capable of fulfilling are, supporting the actions of the system, allaying pain and irritation, relieving spasmodic action, inducing sleep, and checking morbidly increased secretions. It is differently administered, as it is designed to fulfil one or other of these indications.

Where opium is given as a stimulus, it ought to be administered in small doses, frequently repeated, and slowly increased, as by this mode the excitement it produces is best kept up. But when the design is to mitigate pain or irritation, or the symptoms arising from these, it ought to be given in a full dose, and at distant intervals, by which the state of diminished power and sensibility is most completely induced.

One other general rule, with respect to the administration of opium, is, that it ought not to be given in any pure inflammatory affection, at least until evacuations have been used, or unless means are employed to determine it to the surface, and produce a diaphoresis.

In continued fevers, not of the pure inflammatory kind, opium is administered sometimes as a general stimulus, and at other times to allay irritation. The great practical rule in such cases is, that it ought to be given in such quantities only, that the pulse becomes slower and fuller from its operation. Its exhibition is impro-

per, where local inflammation, especially of the brain, or of its membranes, exists.

In intermittent fever, the exhibition of an opiate renders the paroxysms milder, and facilitates the cure. Dr. Cullen recommends the union of opium with bark, which enables the stomach to bear the latter in larger doses, and adds considerably to its efficacy.

In the greater number of the profluvia catarrh cholera, opium is employed to lessen the discharge, and is frequently the principal remedy in effecting the cure. In passive hæmorrhagy, it proves useful by its stimulant power. In retrocedent gout it is used as a powerful stimulant.

In convulsive and spasmodic diseases it is advantageously administered, with the view of relieving symptoms, or even of effecting a permanent cure, and in several of them it requires to be given to a very great extent.

In lues venerea it promotes the action of mercury, and relieves the irritation arising either from that remedy, or from the disease.

In the year 1779, opium was introduced into practice as a specific against the lues venerea. It was employed in several of the military hospitals, where it acquired the reputation of a most efficacious remedy; and Dr. Michaelis, physician of the Hessian forces, published an account of a great number of successful experiments made with it, in the first volume of the Medical Communications in the year 1784. Opium was afterwards given as an antivenereal remedy in some foreign hospitals. Many trials were also made of its virtues in several of the London hospitals, and in the Royal Infirmary at Edinburgh. Very favourable reports of its efficacy in removing venereal complaints were published by different practitioners; but, at the same time, so many deductions were to be made, and so many exceptions are to be admitted, that it required little sagacity to discover, that most of the advocates for this medicine reposed by a slender and fluctuating confidence in its antivenereal powers. Mr. Pearson made several experiments on the virtues of opium in lues venerea, at the Lock Hospital, in the years 1784 and 1785; and published a narrative of its effects, in the second volume of the Medical Communications. "The result of my experiments," says he, "was very unfavourable to the credit of this new remedy: and I believe that no surgeon in this country relies on opium as a specific against the venereal virus. I have been long accustomed to administer opium with great freedom during the venereal course; and the experience of nearly twenty years has taught me, that, when it is combined with mercury, the proper efficacy of the latter is not in any measure increased; that it would not be safe to rely

upon a smaller quantity of the mineral specific, nor to contract the mercurial course within a shorter limit than where no opium has been employed. This representation will not, I presume, admit of controversy; yet we frequently hear people expressing themselves upon this head, as if opium manifested some peculiar qualities in venereal complaints, of a distinct nature from its well known narcotic properties, and thus afforded an important aid to mercury in the removal of lues venerea." Perhaps it may not be unuseful to disentangle this subject from the perplexity in which such indefinite language necessarily involves it. Opium, when given in conjunction with mercury, by diminishing the sensibility of the stomach and bowels, prevents many of those inconveniences which this mineral is apt to excite in the primæ viæ; and thus its admission into the general system is facilitated. Mercury will likewise often produce a morbid irritability, accompanied with restlessness and insomnolence, and it sometimes renders venereal sores painful, and disposed to spread. These accidental evils, not necessarily connected with the venereal disease, may be commonly alleviated, and often entirely removed, by a judicious administration of opium; and the patient will consequently be enabled to persist in using the mineral specific. It however, must be perfectly obvious, that opium, in conferring this sort of relief, communicates no additional virtues to mercury, and that, in reality, it assists the constitution of the patient, not the operation of the medicine with which it is combined. The salutary effects of mercury, as an antidote, may be diminished or lost by the superintention of vomiting, dysentery, &c. Opium will often correct these morbid appearances, and so will spices, wine, an appropriate diet, &c. yet it would be a strange use of words to urge, wherever these articles of food were beneficial to a venereal patient, that they concurred in augmenting the medicinal virtues of mercury. It may be supposed that the majority of medical men would understand, by the terms "to assist a medicine in curing a contagious disease," that the drug conjoined with the specific actually increased its medicinal efficacy; whereas, in the instances before us, it is the human body only, which has been aided to resist the operation of certain noxious powers, which would render a preference in the antidote prejudicial or impossible. The soothing qualities of this admirable medicine can scarcely be estimated too highly. Yet we must beware of ascribing effects to them which have no existence; since a confidence in the antivenereal virtue of opium would be a source of greater mischief, than

its most valuable properties would be able to compensate.

Opium is employed with laxatives in colic, and often prevents ileus and inflammation, by relieving the spasm.

It is often given to promote healthy suppuration, and is a principal remedy in arresting the progress of gangrene.

The sudorific property of opium is justly considered of considerable power, more especially in combination with ipecacuan or antimony. The combined powder of ipecacuan, consisting of one part of ipecacuan, one part of opium, and eight of sulphat of potash, is a very powerful sudorific, given in a dose from 15 to 25 grains. The combination of opium with antimony is generally made by adding 30 to 40 drops of antimonial wine to 25 or 30 drops of tincture of opium, and forming them into a draught.

Opium, taken into the stomach in immoderate doses, proves a narcotic poison, producing vertigo, tremors, convulsions, delirium, stupor, stertor, and, finally, fatal apoplexy.

Where opium has been taken so as to produce these dangerous consequences, the contents of the stomach are first to be evacuated by a powerful emetic, as a solution of the sulphat of zinc. Large draughts of vinegar, or any of the native vegetable acids, are then to be swallowed. Moderate doses of brandy, or a strong infusion of coffee, have also been found useful.

Respecting the external application of opium, authors seem not sufficiently agreed. Some allege, that when applied to the skin it allays pain and spasm, procures sleep, and produces all the salutary or dangerous effects which result from its internal use; while others say, that thus applied it has little or no effect whatever. It has also been asserted, that when mixed with caustic it diminishes the pain which would otherwise ensue; and if this be true, it is probably by decreasing the sensibility of the part. Injected by the rectum, it has all the effect of opium taken into the stomach; but to answer this purpose, double the quantity is to be employed. Applied to the naked nerves of animals, it produces immediate torpor and loss of power in all the muscles with which the nerves communicate.

The requisite dose of opium varies in different persons, and in different states of the same person. A quarter of a grain will in one adult produce effects which ten times the quantity will not do in another; and a dose that might prove fatal in cholera or colic, would not be perceptible in many cases of tetanus, or mania. The lowest fatal dose, to those unaccustomed to take it, seems to be about four grains; but a dangerous dose is so apt to produce

vomiting, that it has seldom time to occasion death. When given in too small a dose, it often produces disturbed sleep, and other disagreeable consequences; and in some cases it seems impossible to be made to agree in any dose or form. Often, on the other hand, from a small dose, sound sleep and alleviation of pain will be produced; while a larger one occasions vertigo and delirium. Some prefer the repetition of small doses; others the giving a full dose at once: its operation is supposed to last about eight hours; this however must depend upon circumstances. The usual dose is one grain. The official preparations of this drug are numerous. The following are among the principal: *Opium purificatum*, *pitula ex opio*, *pulvis opiatu*, *tinctura opii*, *tinctura opii camphorata*, and *confectio opii*: it is also an ingredient in the *pulvis ipecacuanha compositus*, *electuarium japonicum*, *pulvis e cretacompositus*, &c.

OPOBALSAMUM. (From *οπος*, juice, and *βαλσαμον*, balsam.) See *Balsamum Gileadense*.

OPOCALPASON. (From *οπος*, juice, and *καλπασσα*, a tree of that name.) *Opocarpison*. The juice of a tree called *Calpast*. It resembles myrrh, but is poisonous.

OPODOCELE. A rupture through the foramen ischii, or into the labia pudendi.

OPOIELDOC. A term of no meaning, frequently mentioned by Paracelsus. Formerly it signified a plaster for all external injuries, but now is confined to a camphorated soap liniment.

OPOPANAX. (From *οπος*, juice, and *παναξ*, the panacea.) The plant from whence the gum is produced is known by the names of *opoponacum*, *panax heracleum*, *panax costinum*, *panax pastinacea*, *kyna*, *Hercules all heal*, and *opoponaxwort*. *Pastinaca opopanax* of Linnaeus; *foliis pinnatis, foliolis basi antica excisis*. *Opopanax* is the gummi-resinous juice obtained by means of incisions made at the bottom of the stalk of the plant, from which it gradually exudes, and by undergoing spontaneous concretion, assumes the appearance under which we have it imported from Turkey, and the East Indies, viz. sometimes in little drops or tears, more commonly in irregular lumps, of a reddish yellow colour, on the outside, with specks of white; internally of a paler colour, and frequently variegated with large white pieces. *Opopanax* has a strong disagreeable smell, and a bitter, acrid, somewhat nauseous taste. It is only employed in the present practice as an antispasmodic, in combination with opium, although it was formerly mentioned as an attenuant, deobedient. Its antispasmodic power is more powerful than galbanum,

and more so than ammoniacum. It has no place in the Edinburgh Pharmacopœia, but is directed by the London College in the *pitula galbani composita*.

OPOPIA. (From *οπτομαι*, to see.) The bones of the eyes.

OPORICE. (From *οπορα*, autumnal fruits.) A conserve made of ripe fruits.

OPPIATIO. (From *oppilo*, to shut up.) *Oppilation* is a close kind of obstruction; for, according to Rhodius, it signifies, not only to shut out, but also to fill.

OPPILATIVA. (From *oppilo*, to shut up.) Medicines or substances which shut up the pores.

OPPONENS POLLICIS. See *Flexor ossis metacarpi pollicis*.

OPPRESSIO. The catalepsy, or any pressure upon the brain.

OPRIGOROS. (From *οψη*, late, and *γενεσθαι*, to be born.) A dens sapientiz, or late cut tooth.

OPTIC NERVES. (*Nervi optici*, from *οπτομαι*, to see: because they are the organs of sight.) The second pair of nerves of the brain, they arise from the thalami, nervorum opticozum, perforate the bulb of the eye, and in it form the retina.

OPUNTIA. (*Ab opunte*, from the city *Opus*, near which it flourished.) The prickly leaves of this plant, *Cactus opuntia* of Linnaeus, abound with a mucilaginous matter, which is esteemed in its native countries an emollient, in the form of poultice.

Orache, stinking. See *Atriplex fatida*.

Orange. See *Aurantium*.

Orange Seville. See *Aurantium*.

Orange, shaddock. See *Shaddock*.

ORBICULARE OS. (*Orbicularis*, shaped like a ring, from *orbiculus*, a little ring.) *Os pisiforme*. A name of a bone of the carpus. Also a very small round bone, not larger than a pin-head, that belongs to the internal ear.

ORBICULARIS ORIS. (*Musculus orbicularis oris*, from *orbiculus*, a little ring; so called from its shape.) *Sphincter laborum* of Douglass, *semi orbicularis* of Winslow, *constrictor oris* of Cowper, and *labial* of Dumas. A muscle of the mouth, formed in a great measure by those of the lips; the fibres of the superior descending, those of the inferior ascending, and decussating each other about the corner of the mouth; they run along the lip to join those of the opposite side, so that the fleshy fibres appear to surround the mouth like a sphincter. Its use is to shut the mouth, by contracting and drawing both lips together, and to counteract all the muscles that assist in forming it.

ORBICULARIS PALPEBRARUM. *Orbicularis*, scil. *musculus*. *Orbicularis palpebrarum ciliaris* of authors, and *maxillo palpebral* of Dumas. A muscle common to both the eyelids. It arises by a number

of fleshy fibres from the outer edge of the orbital process of the superior maxillary bone, and from a tendon near the inner angle of the eye; these fibres run a little downward and outwards, over the upper part of the cheek, below the orbit, covering the under eyelid, and surround the external angle, being closely connected only to the skin and fat; they then run over the superciliary ridge of the os frontis, towards the inner canthus, where they mix with the fibre of the os occipito-frontalis and corrugator supercilii: then covering the upper eyelid, they descend to the inner angle, opposite to their inferior origin, and firmly adhere to the internal angular process of the os frontis, and to the short round tendon which serves to fix the palpebræ and muscular fibres arising from it. It is inserted into the nasal process of the superior maxillary bone by a short round tendon, covering the anterior and upper part of the lachrymal sac, which tendon can be easily felt at the inner canthus of the eye. The use of this muscle is to shut the eye, by drawing both lids together, the fibres contracting from the outer angle towards the inner, press the eyeball, squeeze the lachrymal gland, and convey the tears towards the puncta lachrymalia.

ORBICULARIS PALPEBRARUM CILIARIS. See *Orbicularis palpebrarum*.

ORBITS. *Orbita*. The two cavities under the forehead, in which the eyes are situated, are so termed. The angles of the orbits are called *canthi*. Each orbit is composed of seven bones, viz. the frontal, maxillary, jugal, lachrymal, ethmoid, palatine, and sphenoid. The use of this bony socket is to maintain and defend the organ of sight, and its adjacent parts.

ORCHEA (From *ορχη*, a testicle.) Galen says it is the *scrotum*.

ORCHIS. (From *ορχιδας*, to desire.)

1. A testicle.

2. The name of a genus of plants in the Linnæan system. Class, *Gyandria*. Order, *Dianthia*.

ORCHIS BIFOLIA. The systematic name of the butterfly orchis. See *Satyrion*.

ORCHIS MASCUULA. The systematic name of the male orchis. See *Satyrion*.

ORCHIS MORIO. The systematic name of the orchis from whose root the salep is made. See *Salep*.

ORCHITIS. (From *ορχη*, a testicle.) See *Hernia humoralis*.

ORCHOS. (From *ορχος*, a plantation or orchard; so called from the regularity with which the hairs are inserted.) The extremities of the eye-lids, where the eyelashes grow.

ORCHOTOMY. (From *ορχη*, a testicle, and *τομή*, to cut.) Castration. The operation of extracting a testicle.

OROSSELINUM. (From *ορος*, a mountain, and *οσσην*, parsley, so named because it grows wild upon mountains.) Black mountain parsley. The root and seed of this plant, *Athaminta oroselinum* of Linnæus, *foliolis divaricatis*, as well as the whole herb, were formerly used medicinally. Though formerly in so high estimation as to obtain the epithet of *polychæta*, this plant is seldom used in the practice of the present day. An extract and tincture prepared from the root were said to be attenuant aperient, deobstruent, and lithontriptic. The oil obtained by distillation from the seed was esteemed to allay the toothach; and the whole was recommended as an antiscorbutic and corroborant.

ORESTION. (From *ορος*, a mountain.) In Dioscorides it is the *Helenium*, or a kind of elecampane growing upon mountains.

OREXIS. (From *ορεσμαι*, to desire.) *Orexia*. The appetite.

ORICIA. (From *Oricus*, a city of Epirus, near which it grows.) A species of fir or turpentine tree.

ORIENTALIA FOLIA. The leaves of senna.

ORIGANUM. (From *ορος*, a mountain, and *γαιω*, to rejoice, so called because it grows upon the side of mountains.)

1. The name of a genus of plants in the Linnæan system. Class, *Dydinamia*. Order, *Gymnospermia*.

2. The pharmacopœial name of *Majorana mancarana*. *Originanum heracleoticum*, from *Heraclea*, where the best was said to be produced. *Zazarhendi herba*. Wild-majorum. *Origanum vulgare* of Linnæus: *spicis subrotundis paniculatis conglomeratis, bracteis calyce longioribus ovatis*. This plant grows wild in many parts of Britain. It has an agreeable aromatic smell, approaching to that of marjoram, and a pungent taste, much resembling thyme, to which it is likewise thought to be more readily allied in its medicinal qualities, and therefore deemed to be emmenagogue, tonic, stomachic, &c. The dried leaves, used instead of tea, are said to be exceedingly grateful. They are employed in medicated baths and fomentations.

ORIGANUM CRETICUM. See *Dictamnus creticus*.

ORIGANUM DICTAMNUS. The systematic name of the dittany of Crete. See *Dictamnus creticus*.

ORIGANUM MAJORANA. The systematic name of sweet marjoram. See *Marjorana*.

ORIGANUM SYRIACUM. The systematic name of the Syrian herb mastich. See *Marum*.

ORIGANUM VULGARE. The systematic

name of the wild marjoram. See *Origanum*.

ORIS CONSTRICTOR. See *Orbicularis oris*.

ORLEANA TERRA. (*Orleana*, so named from the place where it grows.) The substance so called is a ceraceous mass, obtained from the seeds of the *Bixa orleana* of Linnaeus. In Jamaica and warm climates it is considered as a useful remedy in dysentery, possessing adstringent and stomachic qualities.

ORNITHOGALUM MARITIMUM. (From *ornis*, a bird, and *gala*, milk, so called from the colour of its flowers, which are like the milk found in eggs.) A kind of wild onion. See *Scilla*.

ORNITHOGLOSSUM. (From *ornis*, a bird, and *glossa*, a tongue, so called from its shape.) Bird's tongue. The seeds of the ash-tree, as sometimes so called.

ORNITHOLOGY. (From *ornis*, a bird, and *logos*, a discourse.) That part of natural history which treats of birds.

ORNITHOPODIUM. (From *ornis*, a bird, and *pus*, a foot; so called from the likeness of its pods to a bird's claw.) Bird's foot; scorpion wort.

ORNUS. (From *orn*, Heb.) The ash-tree which affords manna.

OROBANCHE. (From *oros*, the wild pea, and *αρχα*, to suffocate; so called because it twines round the orobus, and destroys it.) The great tooth wort or hypocystis.

OROBRYCHIS. (From *oros*, the wood-pea, and *βροχο*, to eat.) The same as orobus.

OROBUS. (From *oros*, to eat.) 1. The name of a genus of plants in the Linnaean system. Class, *Diadelphia*. Order, *Decandria*.

2. The pharmacopoeial name of the eruvum. See *Eruum*.

OROBUS TUBEROSUS. The heath pea. The root of this plant is said to be nutritious. The Scotch Highlanders hold them in great esteem, and chew them like tobacco.

OROSELINUM. See *Oreoselinum*.

ORPIMENT. *Orpimentum*. Native orpiment is found in yellow, brilliant, and, as it were, tacky masses, often mixed with realgar, and sometimes of a greenish colour. See *Arsenic*.

Orpine. See *Faba crassa*.

ORRHOPHYGIUM. (From *oros*, the extremity, and *φυγι*, the buttocks.) The extremity of the spine, which is terminated by the os coccygis.

ORRHOS. (From *oros*, to flow.) Serum, whey. The raphe, and the extremity of the sacrum.

Orris, common. See *Iris nostras*.

Orris, Florentine. See *Iris florentina*.

ORTHOCOLON. (From *orthos*, straight, and *κολον*, a limb.) It is a species of stiff joint,

and is when it cannot be bended, but remains straight.

ORTHOPNOEA. (From *orthos*, erect, and *πνοη*, breathing.) A very quick and laborious breathing, during which the person is obliged to be in an erect posture.

ORVALA. (*Orvale*, French.) A species of clary or horminum.

ORVIETANUM is used for a medicine that resists poisons, from a mountebank of Orvieto in Italy, who first made himself famous by taking such things upon the stage, after doses of pretended poisons. Though some say its inventor was one H. F. Orvietanus, and that it is named after him.

ORYZA. (From *arez*, Arab.) 1. The name of a genus of plants in the Linnaean system. Class, *Triandria*. Order, *Digymia*. 2. The name for rice, or the seeds of the *Oryza sativa* of Linnaeus. Rice is the principal food of the inhabitants in all parts of the East, where it is boiled and eaten, either alone or with their meat. Large quantities of it are annually sent into Europe, and it meets with a general esteem for family purposes. The people of Java have a method of making puddings of rice, which seems to be unknown here, but it is not difficult to put in practice, if it should merit attention. They take a conical earthen pot, which is open at the large end, and perforated all over: this they fill about half full with rice, and putting it into a larger earthen pot of the same shape, filled with boiling water, the rice in the first pot soon swells, and stops the perforations so as to keep out the water; by this method the rice is brought to a firm consistence, and forms a pudding, which is generally eaten with butter, oil, sugar, vinegar, and spices. The Indians eat stewed rice with good success against the bloody flux; and in most inflammatory disorders they cure themselves with only a decoction of it. The spirituous liquor called arrack is made from this grain. Rice grows naturally in moist places; and will not come to perfection, when cultivated, unless the ground be sometimes overflowed, or plentifully watered. The grain is of a grey colour when first reaped; but the growers have a method of whitening it before it is sent to market. The manner of performing this, and beating it out in Egypt, is thus described by Hasselquint: They have hollow iron cylindrical pestles, about an inch diameter, lifted by a wheel-worked with oxen. A person sits between the pestles, and, as they rise, pushes forward the rice, whilst another winnows and supplies fresh parcels. Thus they continue working until it is entirely free from chaff. Having in this manner cleaned it, they add one-thirtieth part of salt, and rub them both together, by which the grain acquires a whiteness; then it is

passed through a sieve, to separate the salt again from it. In the island of Ceylon they have a much more expeditious method of getting out the rice; for, in the field where it is reaped, they dig a round hole, with a level bottom, about a foot deep, and eight yards diameter, and fill it with bundles of corn. Having laid it properly, the women drive about half a dozen oxen continually round the pit; and thus they will tread out forty or fifty bushels a day. This is a very ancient method of treading out corn, and is still practised in Africa upon other sorts of grain.

ORYZA SATIVA. The systematic name of the rice-plant. See *Oryza*.

OS. See *Bone*.

OS EXTERNUM. The entrance into the vagina. It is so named in opposition to the mouth of the womb, which is called the *os internum*, or *os tincæ*.

OS INTERNUM. *Os tinca*, and *amphideon*, or *amphideum*. Galen calls it *oscheon*. The orifice or mouth of the womb.

OS LEONIS. The antirrhium linaria.

OS TINCÆ. See *Os internum*.

OSCHEOCELE. (From *oscheon*, the scrotum, and *cele*, a tumour) This term is sometimes given to a tumour of the scrotum, from an accumulation of water, (see *Hydrocele*); and sometimes to a scrotal hernia, (see *Hernia*.)

OSCHEON. *Οσχεον*. The scrotum. Galen gives the name to the *os uteri*.

OSCHEOPHYMA. (From *oscheon*, the scrotum, and *phyma*, a tumour. (A swelling of the scrotum.

Oscillation of Boerhaave. See *Irritability*.

OSCITANS. (From *oscito*, to gape.) The yawning fever.

OSCITATIO. (From *oscito*, to gape.) *Chasme*. *Oscedo*. Yawning. Gaping.

OSULATORIUS. (From *osculo*, to kiss; so called because the action of kissing is performed by it.) The sphincter muscle of the lips.

OSCULUM. Dim. of *os*, a mouth.) A little mouth.

Osmund royal. See *Osmunda regalis*.

OSMUNDA. (From *Osmund*, who first used it) The name of a genus of plants in the Linnæan system. Class, *Cryptogamia*. Order, *Filices*.

OSMUNDA REGALIS. The systematic name of the osmund royal. Its root possesses adstringent and styptic virtues.

OSPHYS. *Οσφυς*. The loins.

OSSA SPONGIOSA. The spongy bones are two in number, and are called *ossa spongiosa inferiora*. The ethmoid bone has two turbinated portions, which are sometimes called the superior spongy bones. These bones, which, from their shape, are sometimes called *ossa turbinata*, have, by some anatomists, been described

as belonging to the ethmoid bone; and by others, as portions of the *ossa palati*. In young subjects, however, they are evidently distinct bones. They consist of a spongy lamella in each nostril. The convex surface of this lamella is turned towards the septum narium, and its concave part towards the maxillary bone, covering the opening of the lachrymal duct into the nose. From their upper edge arise two processes: the posterior of these, which is the broadest, hangs as it were upon the edge of the antrum highmorianum; the anterior one joins the *os unguis*, and forms a part of the lachrymal duct. These bones are complete in the fœtus. They are lined with the pituitary membrane; and, besides their connection with the ethmoid bone, are joined to the *ossa maxillaria superiora*, *ossa palati*, and *ossa unguis*. Besides these *ossa spongiosa inferiora*, there are sometimes two others, situated lower down, one in each nostril. These are very properly considered as a production of the sides of the maxillary sinus turned downwards. In many subjects, likewise, we find other smaller bones, standing out into the nostrils, which, from their shape, might also deserve the name of *turbinata*, but they are uncertain in their size, situation, and number.

OSSICULA AUDITUS. The small bones of the internal ear are four in number, viz. the malleus, incus, stapes, and *os orbiculare*; and are situated in the cavity of the tympanum. See *Malleus*, *Incus*, *Stapes*, and *Orbiculare os*.

OSSIFICATION. (From *os*, a bone, and *facio*, to make.) See *Bone*.

OSSIFRAGA. (From *os*, a bone, and *frango*, to break.) A petrified root, called the bone-binder, from its supposed virtues in uniting fractured bones.

OSSIFRAGUS. See *Osteocolla*.

OSSIVORUS. (From *os*, a bone, and *voro*, to devour.) Applied to a species of tumour or ulcer, which destroys the bone.

OSTANGA. (From *ostium*, a bone, and *αγγα*, a laying hold of.) A forceps to take out bones with.

OSTARIUS. (A porter, from *ostium*, a door; so called as being the passage into the bowels.) The lower orifice of the stomach.

OSTEITES. (From *ostium*, a bone.) The bone-binder. See *Osteocolla*.

OSTEOCOLLA. (From *ostium*, a bone, and *κόλλα*, to glue.) *Ossifraga*. *Holosteus*. *Osteites*. *Amosteus*. *Osteolithes*. *Stelochites*, glue bone, stone, or Bone-binder. A particular carbonate of lime, found in some parts of Germany, particularly in the Marché of Brandenburg, and in other countries. It is met with in loose sandy grounds, spreading from near the surface to a considerable depth, into a number of

ramifications, like the roots of a tree; it is of whitish colour, soft whilst under the earth, & able when dry, rough on the surface, for the most part either hollow within, or filled with a solid wood, or with a powdery white matter. It was formerly celebrated for promoting the coalition of fractured bones, and the formation of callus; which virtues are not attributed to it in the present day.

OSTEOCOPUS. (From *ὀστος*, a bone, and *κόπος*, uneasiness.) A very violent fixed pain in any part of the bone.

OSTEOGENICA. (From *ὀστος*, a bone, and *γενναω*, to beget.) Medicines which promote the generation of a callus.

OSTEOGENY. (*Osteogenia*, from *ὀστος*, a bone, and *γενναω*, generation.) The growth of bones. Bones are either formed between membranes or in the substance of cartilages, and the bony deposition is effected by a determined action of arteries. The secretion of bone takes place in cartilage in the long bones, as those of the arm, leg, &c. and betwixt two layers of membrane, like the bones of the skull, where true cartilage is never seen. Often the bony matter is formed in distinct bags, and there it grows into form, as in the teeth; for each tooth is formed in its little bag, which by injection can be filled and covered with vessels. Any artery of the body can assume this action, and deposit bone, which is formed also where it should not be, in the tendons, and in the joints, in the great arteries, and in the valves, in the flesh of the heart itself, or even in the soft and pulpy substance of the brain.

All the bones in the fœtus are merely cartilage before the time of birth; this cartilage is never hardened into bone, but from the first it is an organized mass. It has its vessels, which are at first transparent, but which soon dilate; and whenever the red colour or the blood begins to appear in them, ossification very quickly succeeds, the arteries being so far enlarged as to carry the coarser parts of the blood. The first mark of ossification is an artery which is seen running into the centre of the jelly which is formed. Other arteries soon appear, and a net work of vessels is formed, and then a centre of ossification begins, stretching its rays according to the length of the bone, and then the cartilage begins to grow opaque, yellow, brittle; it will no longer bend, and a bony centre may easily be discovered. Other points of ossification are successively formed, preceded by the appearance of arteries. The ossification follows the vessels, and buries and hides those vessels by which it is formed. The vessels advance towards the ends of the bone, the whole body of the bone becomes opaque, and

there is left a small vascular circle only at either end; the heads are separated from the body of the bone by a thin cartilage, and the vessels of the centre, extending still towards the extremities of the bone, perforate the cartilage, pass into the head of the bone, and then its ossification also begins, and a small nucleus of ossification is formed in its centre. Thus the heads and the body are at first distinct bones, formed apart, joined by a cartilage, and not united till the age of fifteen or twenty years. Then the deposition of bone begins, and while the bone is laid by the arteries, the cartilage is conveyed away by the absorbing vessels; and while they convey away the superfluous cartilage, they model the bone into its due form, shape out its cavities, cancelli and holes, remove the thinner parts of the cartilage, and harden it into due consistence. The earth which constitutes the hardness of bone, and all its useful properties, is dead, inorganicized, and lies in the interstices of bone, where it is made up of gelatinous matter, to give it consistence and strength, furnished with absorbents to keep it in health, and carry off its wasted parts; and pervaded by vessels to supply it with new matter. During all the process of ossification, the absorbents proportion their action to the stimulus which is applied to them; they carry away the serous fluid, when jelly is to take its place; they remove the jelly as the bone is laid; they continue removing the bony particles also, which (as in a circle,) the arteries continually renew; this renovation and change of parts goes on even in the hardest bones, so that after a bone is perfectly formed, its older particles are continually being removed, and new ones are deposited in their place. The bony particles are so deposited in the flat bones of the skull as to present a radiated structure, and the vacancies between the fibres which occasion this appearance are found, by injection, to be chiefly passages for blood-vessels. As the fœtus increases in size, the osseous fibres increase in number, till a lamina is produced; and as the bone continues to grow, more lamina are added, till the more solid part of a bone is formed. The ossification which begins in cartilage is considerably later than that which has its origin between membranes. The generality of bones are incomplete until the age of puberty, or between the fifteenth and twentieth year, and in some few instances not until a later period: the small bones of the ear however are completely formed at birth.

OSTEOGRAPHY. (From *ὀστος*, a bone, and *γραφω*, to describe.) The description of the bones. See *Bone*.

OSTEOLITHOS. (From *ὀστος*, a bone, and *λίθος*, a stone.) See *Osteocolla*.

OSTEOLOGY. (From *ostion*, a bone, and *λογος*, a discourse.) The doctrine of the bones. See *Bone*.

OSTIOLA. (Dim. of *ostium*, a door.) The valves or gates of the heart.

OSTREUM. (From *οστρεον*, a shell.) The oyster. The shell of this fish is occasionally used medicinally; its virtues are similar to those of the carbonate of lime. See *Creta*.

OSTRITUM. (Blanchard calls it a corruption from *laserpitium*.) *Imperatoria*, or *masterwort*.

OSTRUTHIUM. *Laserpitium*. See *Imperatoria*.

OSYRIS. *Cassia poetica* *Lobellii*. *Cassia latinorum*. *Cassia lignea monspeliensis*. *Cassia monspeliensis*. Poet's rosemary. The whole shrub is astringent. It grows in the southern parts of Europe.

OSTALGIA. (From *ος*, the ear, and *αλγος*, pain.) The ear-ache.

OTENCHYTES. (From *otos*, the genitive of *ous*, an ear, and *εγχυω*, to pour in.) A syringe for the ears.

OTHONNA. (From *οθεν*, lint; so called from the softness of its leaves.) A species of celandine.

OTICA. (From *ος*, the ear.) Medicines against diseases of the ear.

OTITES. (From *ος*, the ear.) An epithet of the little finger, because it is commonly made use of in scratching the ear.

OTITIS. (From *ος*, the ear.) Inflammation of the internal ear. It is known by pyrexia, and an excruciating and throbbing pain in the internal ear, that is sometimes attended with delirium.

OTOPLATOS. (From *ος*, the ear.) A stinking ulcer behind the ears.

OTOPOYOSIS. (From *ος*, the ear, and *πυος*, pus.) A purulent discharge from the ear.

OTORRHEA. (From *ος*, the ear, and *ρεια*, to flow.) A discharge of blood or matter from the ear.

OVALE FORAMEN. (See *Foramen ovale*.)

OVARUM. (Dim. of *ovum*, an egg.) The ovaria are two flat oval bodies, about one inch in length, and rather more than half in breadth and thickness, suspended in the broad ligaments, about the distance of one inch from the uterus behind, and a little below, the Fallopian tubes. To the ovaria, according to the idea of their structure entertained by different anatomists, various uses have been assigned, or the purpose they answer has been differently explained. Some have supposed that their texture was glandular, and that they secreted a fluid equivalent to, and similar to, the male semen; but others, who have examined them with more care, assert that they are ovaria in the literal acceptation of the term, and include a number of vesicles,

or ova, to the amount of twenty-two, of different sizes, joined to the internal surface of the ovaria by cellular threads or pedicles; and that they contain a fluid which has the appearance of thin lymph. These vesicles are, in fact, to be seen in the healthy ovaria of every young woman. They differ very much in their number in different ovaria, but are very seldom so numerous as has just been stated. All have agreed, that the ovaria prepare whatever the female supplies towards the formation of the fœtus; and this is proved by the operation of spaying, which consists in the extirpation of the ovaria, after which the animal not only loses the power of conceiving, but desire is for ever extinguished. The outer coat of the ovaria, together with that of the uterus, is given by the peritonæum; and whenever an ovum is passed into the Fallopian tube, a fissure is observed at the part through which it is supposed to have been transferred. These fissures healing, leave small longitudinal cicatrices on the surface, which are said to enable us to determine, whenever the ovarium is examined, the number of times a woman has conceived. The corpora lutea are oblong glandular bodies of a yellowish colour, found in the ovaria of all animals when pregnant, and, according to some, when they are salacious. They are said to be calyces, from which the impregnated ovum has dropped; and their number is always in proportion to the number of conceptions found in the uterus. They are largest and most conspicuous in the early state of pregnancy, and remain for some time after delivery, when they gradually fade and wither till they disappear. The corpora lutea are very vascular, except at their centre, which is whitish; and in the middle of the white part is a small cavity, from which the impregnated ovum is thought to have immediately proceeded. The ovaria are the seat of a particular kind of dropsy, which most commonly happens to women at the time of the final cessation of the menses, though not unfrequently at a more early period of life. It is of the encysted kind, the fluid being sometimes limpid and thin, and at others discoloured and gelatinous. In some cases it has been found to contain one cyst, often in several, and in others the whole tumefaction has been composed of hydatids not larger than grapes. The ovaria are also subject, especially a short time after delivery, to inflammation, terminating in suppuration, and to scirrhus and cancerous diseases, with considerable enlargement. In the former state, they generally adhere to some adjoining part, as the uterus, rectum, the bladder, or the external integuments, and the matter is discharged from

the vagina by stool, by urine, or by any external abscess of the integuments of the abdomen.

OVIDUCT. (*Oviductus*, from *ovum*, an egg, and *ductus*, a canal.) The Fallopian tube, or canal, which runs from the ovary to the bottom of the womb.

OVIPAROUS. (From *ovum*, an egg, and *pario*, to bring forth.) Animals which exclude their young in the egg, which are afterwards hatched.

OVORUM TESTA. Egg-shells. A testaceous absorbent.

OVUM. See *Egg*.

OVUM PHILOSOPHICUM. *Ovum chymicum*. A glass body, round like an egg.

OXALATS. *Oxalae*. Salts formed by the combination of the oxalic acid with different bases; thus, *oxalat of ammonia*, &c.

OXALIC ACID. *Acidum oxalicum*. Salt of sorrel. Acid of sugar. This acid is obtained by evaporating the fresh juice of sorrel almost to the consistence of honey, when it is to be poured into a glass vessel with a narrow neck, and covered with a stratum of the oil of olives. After some weeks the sides of the bottle are invested with a crust, which is the salt of sorrel, or *oxalis potassa acidulus*. The salt of sorrel is then to be dissolved in boiling water, and a small quantity of the nitrate of barytes added to it, when the barytes will unite with the oxalic acid, and the potash with the nitric acid. The oxalat of barytes, which is precipitated, is then to be decomposed by digestion with sulphuric acid, by which means the oxalic acid is let loose. Formerly this acid was considered as different from that of sugar, but it is now proved by experiments to be the same in all its properties.

OXALIS. (From *oxus*, sharp; so called from the sharpness of its juice.) The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Pentagynia*. Wood-sorrel.

OXALIS ACETOCELLA. (Dim. of *acetosa*.) The systematic name of the wood-sorrel. See *Lujula*.

OXALWE. (From *oxus*, sharp, and *ale*, salt.) A mixture of vinegar and salt.

Ox-eye-daisy. See *Helix major*

Ox's tongue. See *Picris echioides*.

OXYCANTHA GADENI. (From *oxus*, sharp, and *acanthus*, a thorn; so called from the acidity of its fruit.) The barberry. See *Berberis*.

OXYCEDRUS. (From *oxus*, acutely, and *cedrus*, a cedar; so called from the sharp termination of its leaves.) A kind of cedar. Spanish juniper, a species of *juniperus*.

OXYCOCOOS. (From *oxus*, acid, and *coccus*, a berry, so named from its acidity) *Vaccinia palustris*. *Vitis idæa palustris*. Moor-berry. The cranberry. The berries

of the *Vaccinium oxycoccus* of Linnæus are so termed in some pharmacopœias. They are about the size of our haws, and are pleasantly acid, with which intention they are used medicinally in Sweden. In this country they are mostly preserved and made into tarts.

OXYCRATUM. (From *oxus*, acid, and *cratus*, to mix.) Oxycrates. Vinegar mixed with such a portion of water as is required, and rendered still milder by the addition of a little honey.

OXYCROCUM EMPLASTUM. (From *oxus*, acid, and *spices*, *crocus*, saffron.) A plaster in which there is much saffron, but no vinegar necessary, unless in dissolving some gums.

OXYD. *Oxyd. Oxide. Oxyde. Oxydum*. A substance formed by the union of oxygen with a basis: thus, *oxyd of iron*, *oxyd of copper*, &c.

Oxyd of carbon, gaseous. See *Carbon, gaseous oxyd of*.

OXYDATION. The operation by which a substance is made to combine with oxygen.

OXYDERICA. (From *oxus*, acute, and *deris*, to see.) Medicines which sharpen the sight.

OXYDUM. (So called from oxygen, which enters into its composition.) See *Oxyd*.

OXYDUM ANTIMONII. Oxyd of antimony. This is the *calx antimonii*, the *crocus antimonii lotus*, and the *antimonium diaphoreticum*, of old pharmacopœias. It is made thus: "Take of sulphuret of antimony, powdered, two ounces, muriatic acid, eleven fluid-ounces, nitric acid, one fluid-ounce. The acids being mixed together in a glass vessel, add the antimony gradually thereto, and digest them in a boiling heat for an hour, then strain the solution and pour it into a gallon of water, in which two ounces of the subcarbonate of potash have been previously dissolved; wash the precipitated powder by repeated effusions of water until all the acid is washed away, then dry it upon bibulous paper." This preparation possesses diaphoretic virtues, and is given in the dose of from three grains to ten.

OXYDUM ARSENICI ALBUM. See *Arsenious acid*.

OXYDUM CUPRI VIRIDE ACETATUM. See *Verdigris*.

OXYDUM FERRI LUTEUM. See *Ferri boracæ*.

OXYMURIAS HYDRARGYRI. *Hydrargyris muriatus*. Oxymuriat of mercury; Take of purified mercury by weight two pounds, sulphuric acid by weight thirty ounces, dried muriate of soda four pounds. Boil the mercury with the sulphuric acid in a glass vessel until the sulphate of mercury shall be left dry. Rub this, when it

is cold, with the muriate of soda in an earthen-ware mortar; then sublime it in a glass cucurbit, increasing the heat gradually.

An extremely acrid and violently poisonous preparation.

Given internally in small doses properly diluted, and never in the form of pill, it possesses oxygenating, antisymphilitical, and alterative virtues. Externally applied, in form of lotion, it facilitates the healing of venereal sores, and cures the itch. In gargles for venereal ulcers in the throat the oxymeriat of mercury gr. iii. or iv., barley decoction ℥j., honey of roses ℥ij., proves very serviceable; also in cases of tetters, from gr. v. to gr. x. to water ℥j.; and for films and ulcerations of the cornea, gr. i. to water ℥iv.

Mr. Pearson remarks that when the sublimate is given to cure the primary symptoms of syphilis, it will sometimes succeed; more especially, when it produces a considerable degree of soreness of the gums, and the common specific effects of mercury in the animal system. But it will often fail of removing even a recent chancre; and where that symptom has vanished during the administration of corrosive sublimate, I have known, says he, a three months' course of that medicine fail of securing the patient from a constitutional affection. The result of my observations is, that simple mercury, calomel or calcined mercury, are preparations more to be confided in for the cure of primary symptoms, than corrosive sublimate. The latter will often check the progress of secondary symptoms very conveniently, and I think it is peculiarly efficacious in relieving venereal pains, in healing ulcers of the throat, and in promoting the desquamation of eruptions. Yet even in these cases it never confers permanent benefit; for new symptoms will appear during the use of it; and on many occasions it will fail of affording the least advantage to the patient from first to last. I do, sometimes, indeed, employ this preparation in venereal cases; but it is either at the beginning of a mercurial course, to bring the constitution under the influence of mercury at an early period, or during a course of inunction, with the intention of increasing the action of simple mercury. I sometimes also prescribe it after the conclusion of a course of friction, to support the mercurial influence in the habit, in order to guard against the danger of a relapse. But on no occasion whatever do I think it safe to confide in this preparation singly and uncombined, for the cure of any truly venereal symptom.

OXYDUM HYDRARGYRI CINEREUM. Grey oxyd of mercury. "Take of submuriate of mercury, an ounce; lime-water, a

gallon." Boil the submuriate of mercury in the lime-water, constantly stirring, until a grey oxyd of mercury is separated. Wash this with distilled water, and then dry it. The dose from gr. ii. to x.

OXYDUM HYDRARGYRI NIGRUM. The black oxyd of mercury has received several names. *Ethops per se. Pulvis mercurialis cinereus. Mercurius cinereus. Turpethum nigrum. Mercurius precipitatus niger.* There are four preparations of it in high estimation:

One made by rubbing mercury with mucilage of gum-arabic. Plenck, of Vienna, has written a treatise on the superior efficacy of this medicine. It is very troublesome to make; and does not appear to possess more virtues than some other mercurial preparations.

Another made by triturating equal parts of sugar and mercury together.

The third, composed of honey or liquorice and hydrargyrus purificatus.

The fourth is the blue mercurial ointment.

All these preparations possess anthelmintic, antisymphilitic, alterative, sialagogue, and deobstruent virtues, and are exhibited in the cure of worms, syphilis, amenorrhoea, diseases of the skin, chronic diseases, obstructions of the viscera, &c.

OXYDUM HYDRARGYRI RUBRUM. *Hydrargyrus calcinatus.* Red oxyd of mercury. "Take of purified mercury by weight a pound." Pour the mercury into a glass matrass, with a very narrow mouth and broad bottom. Apply a heat of 600° to this vessel, without stopping it, until the mercury has changed into red scales; then reduce these to a very fine powder.

The whole process may probably require an exposure of six weeks.

This preparation of mercury is given with great advantage in the cure of syphilis. Its action, however, is such, when given alone, on the bowels, as to require the addition of opium, which totally prevents it. It is also given in conjunction with opium and camphire, as a diaphoretic, in chronic pains and diseases of long continuance.

It is given as an alterative and diaphoretic from gr. ss. ad. ii. every night, joined with camphor and opium, each gr. one-fourth or one-half. It is violently emetic and cathartic in gr. iv. to gr. v.

OXYDUM PLUMBI ALBUM. See *Subcarbonas plumbi*.

OXYDUM PLUMBI RUBRUM. See *Lead*.

OXYDUM PLUMBI SEMIVITREUM. See *Lithargyrus*.

OXYDUM STIBII ALBUM. See *Oxydum antimonii*.

OXYDUM STIBII SEMIVITREUM. A vitreous oxyd of antimony. It was formerly called *Vitrum antimonii*, and consists of an

oxyd of antimony with a little sulphur; it is employed to make antimonial wine.

OXYDUM STIBII SULPHURATUM. This is an oxyd of antimony with sulphur, and was formerly called *Hepar antimonis*. *Crocus metallorum*. *Crocus antimonii*. It was formerly exhibited in the cure of fevers and atonic diseases of the lungs. Its principal use now is in preparing other medicines.

OXYDUM ZINCI SUPPLINATUM. See *Zinci oxydum*.

OXYDUM ZINCI. See *Zinci Oxydum*.

OXYGARUM. (From *oxy*, acid, and *garum*, garum.) A composition of garum and vinegar.

OXYGEN. (*Oxygenium*; from *oxy*, acid, and *γενος*, to generate; because it is the generator of acidity.) This substance, although existing sometimes in a solid and sometimes in an æriform state, is never distinctly perceptible to the human senses, but in combination.

We know it only in its combination, by its effects. Nature never presents it solitary; chymists do not know how to insulate it. It is a principle which was long unknown. It is absorbable by combustible bodies, and converts them into acids. It is an indispensable condition of combustion, uniting itself always to bodies which burn, augmenting their weight, and changing their properties. It may be disengaged in the state of oxygen gas, from burnt bodies, by a joint accumulation of caloric and light. It is highly necessary for the respiration of animals. It exists universally dispersed through nature, and is a constituent part of atmospheric air, of water, of acids, and of all bodies of the animal and vegetable kingdoms.

One of the most remarkable combinations into which it is capable of entering, is that which it forms with light and caloric. The nature of that mysterious union has not been ascertained, but it is certain that, in that state, it constitutes the gaseous fluid called **OXYGEN GAS**.

Properties of Oxygen Gas.—Oxygen gas is an elastic invisible fluid, like common air, capable of indefinite expansion and compression. It has neither taste or odour, nor does it show any traces of an acid. Its specific gravity, as determined by Kirwan, is 0.00135, that of water being 1.0000; it is, therefore, 740 times lighter than the same bulk of water. Its weight is to atmospheric air as 1103 to 1000. One hundred and sixteen cubic inches of oxygen gas weigh 39.38 grains. It is not absorbed by water, but entirely absorbable by combustible bodies, which, at the same time, disengage its caloric and light, producing in consequence a strong heat and flame. It rekindles almost extinct combustible bodies. It is indispensable to respiration,

and is the cause of animal heat. It hastens germination. It combines with every combustible body, with all the metals, and with the greater number of vegetable and animal substances. It is considered as the cause of acidity; and from this last property is derived the name *oxygen*, a word denoting the origin of acidity.

The act of its combining with bodies is called *oxydation*, or *oxygenation*; and the bodies with which it is combined are called *oxyds*.

Oxygen gas is the chief basis of the pneumatic doctrine of chymistry.

Methods of obtaining Oxygen Gas.—We are at present acquainted with a great number of bodies, from which we may, by art, produce oxygen gas. It is most amply obtained from the oxyds of manganese, or mercury; from nitrate of potash; from the green leaves of vegetables, and from oxygenated muriate of potash, or soda. Besides these, there are a great many other substances, from which oxygen gas may be procured.

1. In order to procure oxygen gas in a state of great purity, pure oxygenated muriate of potash, or soda, must be made use of. With this view, put some of the salt into a small earthen or glass retort, the neck of which is placed under the shelf of the pneumatic trough, filled with water; and heat the retort by means of a lamp. The salt will begin to melt, and oxygen gas will be obtained in abundance, and of great purity, which may be collected and preserved over water.

Explanation.—Oxygenated muriate of potash consists of oxygenated muriatic acid and potash; at an elevated temperature, a decomposition of the oxygenated muriatic acid takes place; its oxygen unites to the caloric, and forms oxygen gas. The oxygenated acid becomes the more converted into simple muriatic acid, which remains in the retort united to the potash, in the form of muriate of potash.

2. Oxygen gas may likewise be obtained from the green leaves of vegetables.

For this purpose fill a bell-glass with water, introduce fresh-gathered green leaves under it, and place the bell, or receiver, inverted, in a vessel containing the same fluid; expose the apparatus to the rays of the sun, and very pure oxygen gas will be liberated.

The emission of oxygen gas is proportioned to the vigour of the plant and the vivacity of the light; the quantity differs in different plants and under different conditions.

Explanation.—It is an established fact, that plants decompose water, and probably carbonic acid, which serve for their nourishment; they absorb the hydrogen and carbon of these fluids, disengaging a part of

the oxygen in a state of purity. Light, however, favours this decomposition greatly; it seems to serve for melting the oxygen, and thus forms it into oxygen gas; in proportion as it becomes disengaged, the hydrogen becomes fixed in the vegetable, and combines partly with the carbon and partly with the oxygen, to form the oil, &c. of the vegetable.

3. Nitrate of potash is another substance frequently made use of for obtaining oxygen gas, in the following manner:

Take any quantity of this salt, introduce it into a coated earthen or glass retort, and fit to it a tube, which must be plunged into the pneumatic trough, under the receiver filled with water. When the apparatus has been properly adjusted, heat the retort gradually, till it becomes red-hot; the oxygen gas will then be disengaged rapidly.

The gas obtained in this way is also very pure, especially if the last portion be kept separate.

Explanation.—Nitrate of potash consists of nitric acid and potash. Nitric acid consists again of oxygen and nitrogen. On exposing the salt to ignition, a partial decomposition of the acid takes place; the greatest part of the oxygen of the nitric acid unites to caloric, and appears under the form of oxygen gas. The other part remains attached to the potash in the state of nitrous acid. The residue in the retort is, therefore, nitrite of potash, if the process has been carefully conducted.

Remark.—If too much heat be applied, particularly towards the end of the process, a total decomposition of the nitric acid takes place; the oxygen gas, in that case, will therefore be mingled with nitrogen gas. The weight of the two gases, when collected, will be found to correspond very exactly with the weight of the acid which had been decomposed. The residue then left in the retort is potash.

4. Black oxyd of manganese, however, is generally made use of for obtaining oxygen gas, on account of its cheapness. This native oxyd is reduced to a coarse powder; a stone, or rather an iron retort, is then charged with it and heated. As soon as the retort becomes ignited, oxygen gas is obtained plentifully.

Explanation.—Black oxyd of manganese is the metal called manganese fully saturated with oxygen, together with many earthy impurities; on applying heat, part of the solid oxygen quits the metal and unites to caloric, in order to form oxygen gas, the remainder of the oxygen remains united to the metal with a forcible affinity; the metal, therefore, re-approaches to the metallic state, or is found in the state of a grey oxyd of manganese.

One pound of the best manganese yields upwards of 1400 cubic inches of oxygen

gas, nearly pure. If sulphuric acid be previously added to the manganese, the gas is produced by a less heat, and in a larger quantity; a glass retort may then be used, and the heat of a lamp is sufficient.

5. Red oxyd of mercury yields oxygen gas in a manner similar to that of manganese.

Explanation.—This oxyd consists likewise of solid oxygen and mercury, the combination of which takes place on exposing mercury to a heat of about 610° Fahr. At this degree it attracts oxygen, and becomes converted into an oxyd; but if the temperature be increased to about 1000°, the attraction of oxygen is changed. The oxygen then attracts caloric stronger than it did the mercury; it therefore abandons it and forms oxygen gas. The mercury then re-appears in its metallic state.

6. Red oxyd of lead yields oxygen gas on the same principle.

OXYGENATED MURIATIC ACID GAS. This gas possesses an uncommonly pungent and suffocating odour. It is absolutely and in every respect non-respirable; animals immersed in it die instantly. It is absorbable by water, and forms with it what is called liquid oxygenated muriatic acid. When water is saturated with it, the compound crystallizes at low temperatures. Oxygenated muriatic acid gas is not invisible, but has a *yellow-greenish* colour. It is capable of maintaining and exciting combustion in many cases. Phosphorus, charcoal, red sulphuret of mercury, sulphuret of antimony, bismuth, iron, zinc, copper, gold, arsenic, cobalt, tin, lead, and several other combustible bodies, take fire spontaneously when introduced into it. It is heavier than atmospheric air. It weakens and reddens the flame of a taper, but does not extinguish it. It decomposes ammoniac. It thickens fat oils. It detonates with hydrogen gas. Nitrous gas immediately produces a cloud of reddish vapour with it. It is likewise decomposed by sulphurated, phosphorated, and carbonated hydrogen gases. It is not altered by exposure to light, and passes unaltered through an ignited porcelain tube. It discolours stuffs, and totally destroys most of the vegetable colours, rendering them white. It also bleaches yellow wax, &c.

This gas may be obtained in several ways.

1. Take one part of the native oxyd of manganese, one of red precipitate of mercury, or red lead, put it into a glass retort, and add four parts of concentrated muriatic acid. This, on distillation, affords a quantity of yellow æriform fluid, which is oxygenated muriatic acid gas, and by agitating it with water, it combines and forms oxygenated muriatic acid.

It may also be obtained without the red mercurial precipitate, or red lead, thus :

2. Put into a retort one part of powdered black oxyd of manganese, three or four of concentrated muriatic acid, connect the retort with the pneumatic trough, and receive the gas over water in the usual manner. When no more gas is liberated, apply the heat of a lamp, and gas will be produced abundantly, which may be kept in bottles with ground glass stoppers.

The oxyd of manganese yields up in this process part of its oxygen to part of the muriatic acid, which becomes converted into oxygenated muriatic acid gas; the oxyd of manganese being thus partly de-oxydated, is dissolved in the remaining quantity of the muriatic acid, which remains behind in the retort as muriate of manganese.

The retort containing the mixture should not be filled above one-third, for the mixture, on the application of heat, swells and is otherwise very apt to be forced over into the neck of the retort.

3. Oxygenated muriatic acid gas may likewise be obtained in an indirect manner, by decomposing muriate of soda in contact with black oxyd of manganese. For that purpose mix eight parts, by weight, of muriate of soda with three of powdered oxyd of manganese, put the mixture into a tubulated retort, and pour upon it gradually four parts of sulphuric acid, diluted previously with three of water, and which has been suffered to cool after dilution. On applying a gentle heat, gas will be produced as before.

In this operation the sulphuric acid acts on the muriate of soda, to the base of which it unites; the muriatic acid formed by this union attacks the oxyd of manganese; one part of it combines with the oxygen, and another with the oxyd brought nearer to the metallic state, and the result is sulphat of soda and muriate of manganese, which remain in the distilling vessel; and oxygenated muriatic acid, which passes in the state of gas at common temperatures.

In preparing this gas great care should be taken that it does not escape into the apartment in any considerable quantity; as it acts violently on the pituitous membrane, occasions a defluxion of the brain, blunts the senses of smell and taste, produces head-ach, and proves extremely injurious to health.

Pelletier fell a sacrifice in attempting to breathe it; a consumption was the consequence, which proved fatal.

Liquid ammonia is the remedy best calculated to check its effects, when accidentally set at liberty in places where it is prepared.

The water which adheres to the inner

side of the vessel filled with oxygenated muriatic acid gas crystallizes in the form of yellow spangles, if the temperature is near the freezing point. If a considerable quantity of gas be thus condensed, care must be taken to keep it at a low temperature; for as soon as the temperature is raised, it expands, and endangers the bursting of the vessel. When absorbed by water it forms liquid

OXYGENATED MURIATIC ACID.—

This acid is of a greenish yellow colour. It has a styptic bitter taste, and a very suffocating odour. Instead of reddening blue vegetable colours, it has the remarkable property of rendering them white. In high temperatures, when light is excluded, phosphorus remains unaltered in liquid oxygenated muriatic acid; but if light be admitted, the colour of the acid gradually disappears, and the phosphorus is converted into phosphoric acid. It thickens oils and animal fats, and renders them less disposed to combine with alkalies. Its action upon metals presents phenomena extremely curious and important; the oxygen of the acid unites with the metal, and the produced oxyd is afterwards dissolved by the de-oxydated acid.

Method of obtaining Oxygenated Muriatic Acid.—

Put into a tubulated retort, supported over a lamp, one part of black oxyd of manganese reduced to a gross powder, and pour over it three parts of concentrated muriatic acid, recline the retort in such a manner that the fluid which rises up into its neck, may easily run back again into the body, and apply a receiver with a little distilled water in it; the receiver must be luted to the retort by a fillet of paper. When the effervescence, which instantly takes place on the affusion of the acid, ceases, apply a gentle heat. Oxygenated muriatic acid gas will be evolved, and the receiver become filled with yellow vapours, which are absorbed by the water. When the water has acquired a yellowish-green colour, the receiver may be removed, and another one applied till no more gas is extricated. The process may be more elegantly conducted, by joining the apparatus of Burkiit or Pepys to the distillatory vessel. The common muriatic acid which may arise is condensed in the first bottle, and the oxygenated muriatic acid gas unites to the water in the second, third, &c.

The union of oxygenated muriatic acid with different bases forms salts, known by the name of oxygenated muriates.

OXYLYCUM. (From *ὄξύς*, acid, and *λύσις*, sweet.) Honey mixed with vinegar. An oxymel.

OXYLAPATHUM. (From *ὄξύς*, acid, and *λαπάθος*, the dock; so named from its acidity.) *Lapathum acutum. Rumex acutus*.

of Linnaeus. Sharp-pointed deck. *Rumex floribus hermaphroditis; valvula dentatis graniferis, foliis cordato oblongis acuminatis*. The decoction of the root of this plant is used in Germany to cure the itch; and it appears to have been used in the time of Dioscorides, in the cure of leprous and impetigenous affections, both alone and boiled with vinegar.

OXYMEL. (From *ὄξύς*, acid, and *μέλι*, honey.) Apomeli. Adipson. Honey and vinegar boiled to a sirup. See *Mel acetatum*.

OXYMEL ERUGINIS. See *Linimentum eruginis*.

OXYMEL COLCHICI. Oxymel of meadow saffron is an acrid medicine, but is nevertheless employed, for its diuretic virtues, in dropsies.

OXYMEL SCILLÆ. A very useful expectorant.

OXYMYRRHINE. (From *ὄξύς*, acute, and *μύρρον*, the myrtle; so called from its resemblance to myrtle, and its pointed leaves.) *Oxymyrine*. Wild myrtle.

OXYMYRSINE. See *Oxymyrhine*.

OXYNITRUM. (From *ὄξύς*, acid, and *νίτρον*, nitre.) A plaster composed chiefly of vinegar and nitre.

OXYOPIA. (From *ὄξύς*, acute, and *ὄψις*, vision.) The faculty of seeing more acutely than usual. Thus there have been instances known of persons who could see the stars in the day-time. The proximate cause is a preternatural sensibility of the retina. It has been known to precede the gutta serena; and it has been asserted that prisoners, who have been long detained in darkness, have learned to read and write in darkened places.

OXYPHLEGMASIA (From *ὄξύς*, acute, and *φλέγω*, to burn.) An acute inflammation.

OXYPHENICON. (From *ὄξύς*, acid, and *φαινέ*, the tamarind; a native of Phœnicia.) See *Tamarindus*.

OXYPHONIA. (From *ὄξύς*, sharp, and *φωνή*, the voice.) The same as *Paraphonia Clangens*. It is a howling.

OXYREOMA. (From *ὄξύς*, acid, and *ῥέω*, to break wind.) An acid eructation.

OXYRHODINON. (From *ὄξύς*, acid, and *ῥόδον*, oil of roses.) A composition of the oil of roses and vinegar.

OXYRHODIUM. (From *ὄξύς*, acid, and *ῥαζαγόν*, sugar.) A composition of vinegar and sugar.

OXYSAL DIAPHORETICUM. A preparation of Angelus Sala. It is a fixed salt, loaded with more acid than is necessary to saturate it. The salt of Juniper is of this kind.

OXYTOCA. (From *ὄξύς*, quick, and *τατα*, to bring forth.) Medicines which promote delivery.

OXYTRIPHYLLUM. (From *ὄξύς*, acid, and *τριφυλλον*, trefoil; so named from its acidity.) Wood-sorrel.

Oyster. See *Ostreum*.

Oyster-shell. See *Ostreum*.

OZÆNA. (From *ὄζα*, a stench.) An ulcer situated in the nose, discharging a fœtid purulent matter, and sometimes accompanied with caries of the bones. Some authors have signified by the term, an ill-conditioned ulcer in the antrum. The first meaning is the original one. The disease is described as coming on with a trifling tumefaction and redness about the ala nasi, accompanied with a discharge of mucus, with which the nostril becomes obstructed. The matter gradually assumes the appearance of pus, is most copious in the morning, and is sometimes attended with sneezing, and a little bleeding. The ulceration occasionally extends round the ala nasi to the cheek, but seldom far from the nose, the ala of which also it rarely destroys. The ozæna is often connected with scrophulous and venereal complaints. In the latter cases, portions of the ossa spongiosa often come away. After the complete cure of all venereal complaints, an exfoliating dead piece of bone will often keep up symptoms similar to those of the ozæna, until it is detached. Mr. Pearson remarks, that the ozæna frequently occurs as a symptom of the cachexia syphiloidea. It may perforate the septum nasi, destroy the ossa spongiosa, and even the ossa nasi. Such mischief is now more frequently the effect of the cachexia syphiloidea, than of lues venerea. The ozæna must not be confounded with abscesses in the upper jaw-bone.

OXYTUM. (From *ὄζα*, to smell; so called from its fragrance.) Sweet basil.

P.

P. A contraction of *pugillus*, a pugil, or eighth part of a handful, and sometimes a contraction of *partes*, parts.

P. Æ. A contraction of *partes æquales*.

P. P. A contraction of *pulvis putrum*, Jesuit's powder.

PABULUM. (From *pasco*, to feed.) Food, aliment. The animal heat and animal spirits are called *pabulum vite*, the food of life.

Pacchionian glands. See *Glandule Pacchione*.

PACHYTICA. (From *παχυς*, to incrassate.) Medicines which incrassate or thicken the fluids.

PACHYS. *Παχυς*, thick. The name of a disorder described by Hippocrates, but not known by us.

PADUS. The wild cluster cherry, or bird's cherry. The *Prunus padus* of Linnæus. The bark and berries of this shrub are used medicinally. The former, when taken from the tree, has a fragrant smell, and a bitter, subastringent taste, somewhat similar to that of bitter almonds. Made into a decoction, it cures intermittents, and it has been recommended in the cure of several forms of syphilis. The latter are said to cure the dysentery.

PEDANCHONE. (From *παῖς*, a child, and *αἷον*, to strangulate.) A species of quinsy common among children.

PEDARTHOCAE. (From *παῖς*, a boy, *αἰθρῶν*, a joint, and *κακόν*, an evil.) The joint evil. Severinus calls the *Spina Ventosa* by this name, as also doth Dr Cullen. By some this name is used to express a sort of anasarca.

PEONIA. (From *Pæon*, who first applied it to medicinal purposes.) 1. The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Digynia*.

2. The pharmacopœial name of the common peony. Male and female peony. This plant, *Pæonia officinalis* of Linnæus:—*foliis oblongis*, has long been considered as a powerful medicine; and, till the late revision by the London College, it had a place in the catalogue of the *Materia Medica*; in which the two common varieties of this plant are indiscriminately directed for use; and, on the authority of G. Bauhin, improperly distinguished into male and female peony.

The roots and seeds of peony have, when fresh, a faint, unpleasant smell, somewhat of the narcotic kind, and a mucilaginous, subacid taste, with a slight degree of bitterness and adstringency. In drying, they lose their smell and part of their taste. Extracts made from them by water are almost insipid, as well as inodorous; but extracts made by rectified spirits are manifestly bitterish, and considerably adstringent. The flowers have rather more smell than any of the other parts of the plant, and a rough, sweetish taste, which they impart, together with their colour, both to water and spirit.

The roots, flowers, and seeds of peony, have been esteemed, in the character of an anodyne and corroborant, but more especially the roots; which, since the days of Galen, have been very commonly employed as a remedy for the epilepsy. For this purpose, it was usual to cut the root into thin slices, which were to be attached to a string,

and suspended about the neck as an annulet; if this failed of success, the patient was to have recourse to the internal use of this root, which Willis directs to be given in the form of a powder, and in the quantity of a drachm, two or three times a-day, by which, as we are informed, both infants and adults were cured of this disease. Other authors recommended the expressed juice to be given in wine, and sweetened, with sugar, as the most effectual way of administering this plant. Many writers, however, especially in modern times, from repeated trials of the peony in epileptic cases, have found it of no use whatever; though Professor Home, who gave the radix pzoniz to two epileptics at the Edinburgh infirmary, declares that one received a temporary advantage from its use. Of the good effects of this plant, in other disorders, we find no instances recorded.

PÆONIA OFFICINALIS. The systematic name of the common pæony. See *Pæonia*.

Paigil. See *Primula veris*.

PAIN. *Dolor.* Any unpleasant sensation, or irritation.

Painter's colic. See *Colica pictorum*.

Palate. See *Palatum*.

PALATE BONE. (*Os palati*; from *pala*, to hedge in; because it is staked in, as it were, by the teeth.) These two bones are of very irregular figure. They are placed between the ossa maxillaria superiora and the os sphenoides, at the back part of the roof of the mouth, and extend from thence to the bottom of the orbit. Each of these bones may be divided into four parts, viz. the inferior, or square portion, the pterygoid process, the nasal lamella, and orbital process. The first of these, or the square part of the bone, helps to form the palate of the mouth. The upper part of its internal edge rises into a spine, which makes part of the septum narium. The *pterygoid* process, which is smaller above than below, is so named from its being united with the pterygoid processes of the sphenoid bone, with which it helps to form the pterygoid fossæ. It is separated from the square part of the bone, and from the nasal lamella, by an oblique fossa, which, applied to such another in the os maxillare, forms a passage for a branch of the fifth pair of nerves. The *nasal lamella* is nothing more than a very thin bony plate, which arises from the upper side of the external edge of the square part of the bone. Its inner surface is concave, and furnished with a ridge, which supports the back part of the os spongiosum inferius. Externally, it is convex, and firmly united to the maxillary bone. The *orbital* process is more irregular than any other part of the bone. It has a smooth surface, where it helps to form the orbit; and, when viewed in its place, we see it

contiguous to that part of the orbit which is formed by the os maxillare, and appearing as a small triangle at the inner extremity of the orbital process of this last mentioned bone. This fourth part of the os palati likewise helps to form the zygomatic fossa on each side, and there its surface is concave. Between this orbital process and the sphenoid bone, a hole is formed, through which an artery, vein, and nerve, are transmitted to the nostrils. The ossa palati are complete in the fetus. They are joined to the ossa maxillaria superiora, os sphenoides, os ethmoides, ossa spongiosa inferiora, and vomer.

PALATI CIRCUMFLEXUS. See *Circumflexus*.

PALATI LEVATOR. See *Levator palati*.

PALATI TENSOR. See *Circumflexus*.

PALATO-PHARYNGEUS. (*Musculus palato-pharyngeus*; so called from its origin in the palate and insertion in the pharynx.) *Thyro-staphilinus* of Douglas. *Thyro-pharyngo-staphilinus* of Winslow, and *palato-pharyngien* of Dumas. A muscle situated at the side of the entry of the fauces. It arises by a broad beginning from the middle of the velum pendulum palati at the root of the uvula posteriorly, and from the tendinous expansion of the circumflexus palati. The fibres are collected within the posterior arch behind the tonsils, and run backwards to the top and lateral part of the pharynx, where the fibres are scattered and mixed with those of the stylo-pharyngeus. It is inserted into the edge of the upper and back part of the thyroid cartilage. Its use is, to draw the uvula and velum pendulum palati downwards and backwards, and at the same time to pull the thyroid cartilage and pharynx upwards, and shorten it; with the *constrictor superior pharyngis* and tongue, it assists in shutting the passage into the nostrils; and in swallowing, it thrusts the food from the fauces into the pharynx.

PALATO-SALPINGEUS. (From *palatum*, the palate, and *σαλπηξ*, a trumpet; so called from its origin in the palate, and its trumpet-like shape.) See *Circumflexus*.

PALATO-STAPHILINUS. See *Azygos uvule*.

PALATUM. (From *palo*, to hedge in, because it is staked in, as it were, by the teeth.) The palate, or roof of the mouth.

PALATUM MOLLE. The soft palate. This lies behind the bony palate; and from the middle of it the uvula hangs down.

PALEA DE MECHA. A name given by some to the *Juncus Odoratus*.

PALIMPSSA. (From *παλιν*, repetition, and *πσσα*, pitch.) Dioscorides says, that dry pitch is thus named, because it is prepared of pitch twice boiled.

PALINDOXIA. (Παλιν, again, and *δο-*

μα, a course.) Is used by Hippocrates for any recurgitation of humours to the more noble parts: and sometimes for the return of a distemper.

PALIVERT. (From *παλιν*, to move, and *νεφρ*, urine; so called from its diuretic qualities.) A species of white thorn.

PALLIATIVA. (From *pallio*, to dissemble.) Medicines given only with an intent to relieve pains in a fatal disease.

PALM OIL. *Oleum palmæ.* This oil is produced chiefly from the fruit of the *Cocos butyracea* of Linnæus:—*inermis, frondibus pennatis: foliis simplicibus*, by bruising and dissolving the kernels of the fruit in water, without the aid of heat, by which the oil is separated, and rises to the surface, and, on being washed two or three times, is rendered fit for use. When brought into this country, it is of the consistence of an ointment, and of an orange yellow colour, with little taste, and of a strong, though not disagreeable, smell. Its use is confined to external applications in pains, tumours, and sprains; but it appears to possess very little if any advantage over other bland oils.

PALMA. (From *παλιν*, to move.)

1. The palm of the hand.

2. The name of a genus of plants in the Linnæan system, so called because the leaves are extended from the top, like the fingers upon the hand.

PALMA CHRISTI. See *Ricinus*.

PALMARIS BREVIS. (*Palmaris*; from *palmæ*, the hand.) *Palmaris brevis vel caro quadrata* of Douglas, and *Palmare cutané* of Dumas. A small, thin, cutaneous, flexor muscle of the hand, situated on the fore-arm, between the wrist and the little-finger. Fallopius tells us that it was discovered by Cananus. Winslow names it *palmaris cutaneæ*. It arises from a small part of the internal annular ligament, and inner edge of the aponeurosis palmaris, and is inserted by small bundles of fleshy fibres into the os pisiforme, and into the skin and fat that cover the abductor minimi digiti. This muscle seems to assist in contracting the palm of the hand.

PALMARIS CUTANEUS. See *Palmaris brevis*.

PALMARIS LONGUS. *Ulnaris gracilis* of Winslow, and *Epitrochlo carpi palmaris* of Dumas. A flexor muscle of the arm, situated on the fore-arm, immediately under the integuments. It arises tendinous from the inner condyle of the os humeri, but soon becomes fleshy, and after continuing so about three inches, terminates in a long slender tendon, which, near the wrist, separates into two portions, one of which is inserted into the internal annular ligament, and the other loses itself in a tendinous membrane, that is nearly of a triangular shape, and extends over the

palm of the hand, from the carpal ligament to the roots of the fingers, and is called *aponeurosis palmaris*. Some of the fibres of this expansion adhere strongly to the metacarpal bones, and separate the muscles and tendons of each finger. Several anatomical writers have considered this aponeurosis as a production of the tendon of this muscle, but seemingly without reason, because we now and then find the latter wholly inserted into the carpal ligament, in which case it is perfectly distinct from the aponeurosis in question; and, in some subjects, the palmaris longus is wanting, but the aponeurosis is always to be found. Rhodius, indeed, says that the latter is now and then deficient; but there is good reason to think that he was mistaken. This muscle bends the hand, and may assist in its pronation; it likewise serves to stretch the aponeurosis palmaris.

PALMOS. (From *παλλω*, to beat.) A palpitation of the heart.

PALMULA. (Dim. of *palma*, the hand; so called from its shape.) 1. A date.

2. The broad and flat end of a rib.

PALPEBRÆ. (*A palpitando*, from its frequent motion.) The eyelids, distinguished into upper and under: at each end they unite and form the canthus.

Palpebræ superioris levator. See *Levator palpebræ superioris*.

Palpebrum aperiens rectus. See *Levator palpebræ superioris*.

PALPITATIO. 1. A palpitation or convulsive motion of a part.

2. Palpitation of the heart, which is either constant or frequently returning. A genus of disease in the class *Neurosis*, and order *Spasmi*, of Cullen.

Palsy. See *Paralysis*.

PALUDAPIUM. (From *palus*, a lake, and *apum*, smallage; so named because it grows in and about rivulets.) A species of smallage.

PALUS SANCTUS. A name of guaiacum.

PANPHILIUM. (From *πας*, all, and *φιλος*, grateful; so called from its extensive usefulness.) A plaster described by Galen.

PANPINIFORMIS. (From *pampinus*, a tendril, and *forma*, a likeness.) Resembling a tendril; applied to the spermatic chord and the thoracic duct.

PANACEA. (From *πας*, the neuter of *πας*, all, and *αἰσθωμαι*, to cure.) An epithet given by the ancients to those remedies which they conceived would cure every disease. Unfortunately for those of the present day there are no such remedies.

PANACEA DUC HOLSATIÆ. The sulphate of potash.

PANACEA DUPLICATA. Sulphate of potash.

PANACEA VEGETABILIS. Saffron.

PANADA. (Dim. of *pane*, bread, Ital.) *Panata. Panatella.* Bread boiled in water to the consistence of pap. Dry biscuits soaked are the best for this purpose.

PANALETHES. (From *πας*, all, and *αἰσθωμαι*, true.) A name of a cephalic plaster, from its universal efficacy.

PANARITIA. Corrupted from *paronychia*.

PANARIS. (Corrupted from *paronychia*.) See *Paronychia*.

PANAVA. See *Lignum pavane*.

PANAX. (From *πας*, all, and *αἰσθωμαι*, a cure.) The pharmacopoeial name of *Hercules's* alchal or wound wort. *Laserpittum chinorum* of Linnaeus. The seeds and roots of this plant are warm, and similar in flavour and quality to those of the parsnip. The roots and stalks have a much stronger smell, which resembles that of *opopanax*, and Boerhave relates, that on wounding the plant in the summer, he obtained a yellow juice, which being inspissated a little in the sun, agreed perfectly in both respects with that exotic gum resin.

PANAX QUINQUEFOLIUM. The systematic name of the plant which affords the ginseng root. See *Ginseng*.

PANCHRESTOS. (From *πας*, all, *χρηστος*, useful, so named from its general usefulness.) *Panchreston.* An epithet of a collyrium described by Galen. It is also of the same signification as *Panacea*, but little used.

PANCHYMAGOSA. (From *πας*, all, *χυμος*, succus, humour, and *αγω*, *duco*, to lead or draw.) Is ascribed to such medicines as are supposed to purge all humours equally alike: but this is a conceit now not minded.

PANGENUS. (From *πας*, all, and *γενος*, common) Epidemic; applied to popular diseases, which attack all descriptions of persons.

PANCRATIUM. (From *πας*, all, and *αγαν*, to conquer; so called from its virtues in overcoming all obstructions.) See *Scilla*.

PANCREAS. (From *πας*, all, and *γενος*, flesh; so called from its fleshy consistence.) A glandular viscus of the abdomen, of a long figure, compared to a dog's tongue, situated in the epigastric region under the stomach. It is composed of innumerable small glands, the excretory ducts of which unite and form one duct, called the pancreatic duct, which perforates the duodenum with the ductus communis choledochus, and conveys a fluid, in its nature similar to saliva, into the intestines. The pancreatic artery is a branch of the splenic. The veins evacuate themselves into the par vagum and great intercostal. The use of the pancreas is to secrete the pancreatic juice, which is to be mixed with the

chyle in the duodenum. The quantity of the fluid secreted is uncertain; but it must be very considerable, if we compare it with the weight of the saliva, the pancreas being three times larger, and seated in a warmer place. It is expelled by the force of the circulating blood, and of the incumbent viscera in the full abdomen; as the liver, stomach, spleen, mesenteric and splenic arteries, and the aorta. Its great utility appears from its constancy, being found in almost all animals; nor is it refuted by the few experiments in which a part of it was cut out from a robust animal, without occasioning death; because the whole pancreas cannot be removed without the duodenum: for even a part of the lungs may be cut out without producing death, but they are not therefore useless. It seems principally to dilute the viscid cystic bile, to mitigate its acrimony, and to mix it with the food. Hence it is poured into a place remote from the cystic duct, as often as there is no gall-bladder. Like the rest of the intestinal humours, it dilutes and resolves the mass of aliments, and performs every other office of the saliva.

Pancreatic duct. See *Ductus pancreaticus*.

Pancreatic juice. See *Pancreas*.

PANCREAS. (From *παρ*, all, and *αἴμα*, a fountain.) A name of the pancreas from its great secretion.

PANDALITUM. A whitlow.

PANDEMIC. (*Pandemicus*; from *παρ*, all, and *δῆμος*, the people.) A synonym of epidemic. See *Epidemic*.

PANDICULATIO. (From *pandicula*, to gape and stretch.) Pandiculation, or that restless stretching and gaping which accompanies the cold fit of an ague.

PANICUM. (A *paniculus*, from its many particles; the spike consisting of innumerable thick seeds, disposed in many panicles.) The name of a genus of plants in the Linnæan system. Class, *Triandria*, Order, *Digynia*.

PANICUM ITALICUM. The systematic name of the plant which affords the Indian millet-seed. See *Millet-seed*, *Indian*.

PANICUM NYLIACUM. The systematic name of the plant which affords the millet seed. See *Millet-seed*.

PANIS CUCULIS. See *Acetosella*.

PANIS PORCINUS. A species of cyclamen.

PANNICULUS. (From *pannus*, cloth.)

1. A piece of fine cloth.

2. The cellular and carnosous membranes are so called from their resemblance to a piece of fine cloth.

PANNONICA. (From *pannus*, a rag; so called because its stalk is divided into many uneven points, like the end of a piece of rag.) Hawkweed.

PANNUS. (From *πῆμα*, to labour.) 1. A piece of cloth.

2. A tent for a wound.

3. A speck in the eye, resembling a bit of rag.

4. An irregular mark upon the skin.

PANOCHE. Buboec in the groin.

PANOPHOBIA. (From *παρ*, all, and *φοβος*, fear.) That kind of melancholy which is attended with groundless fears.

Pansies. See *Viola tricolor*.

PANTASOSA. (From *παρ*, all, and *αἶμα*, to drive out.) Medicines which expel all morbid humours.

PANTOLMIUS. (From *παρ*, all, and *τελλω*, to dare; so named from its general uses.) A medicine described by Æginetus.

PANTOPHOBIA. The same as hydrophobia.

PANUS. (From *πῆμα*, to work.) 1. A weaver's roll.

2. A soft tumour shaped like a weaver's roll.

PAPAVER. (*Papaver*, from *pappa*, pap; so called because nurses used to mix this plant in children's food to relieve the colic and make them sleep.) 1. The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order *Monogynia*. The poppy.

2. The pharmacopœical name of the white poppy. *Papaver album*. *Papaver somniferum* of Linnæus:—*calycibus capsulisque glabris, foliis amplexicaulibus incisiss.* It is from heads of this plant that the opium is obtained. See *Opium*. They are also directed for medicinal use in the form of fomentation, and in the *syrupus papaveris*, a useful anodyne, which often succeeds in procuring sleep where opium fails; it is, however, more especially adapted to children. The seeds of this species of poppy contain a bland oil, and in many places are eaten as food; as a medicine, they have been usually given in the form of emulsion in catarrhs, stranguries, &c.

PAPAVER ERBATICUM. Red or corn poppy. *Papaver rhæas* of Linnæus:—*capsulis glabris globosis, caule piloso multifloro, foliis pennatifidis incisiss.* The heads of this species, like those of the *somniferum*, contain a milky juice of a narcotic quality; from which an extract is prepared, that has been successfully employed as a sedative. The flowers have somewhat of the smell of opium, and a mucilaginous taste, accompanied with a slight degree of bitterness. A syrup of these flowers is directed in the London Pharmacopœia, which has been thought useful as an anodyne and pectoral, and is prescribed in coughs and catarrhal affections. See *Syrupus rhæadæ*.

PAPAVER NIGRUM. The black poppy. The seeds and heads of the *papaver commi-*

ferum, are also called in some pharmacopœias *semina* and *capitula papaveris nigra*. See *Papaver*.

PAPAYER RHÆAS. The systematic and pharma: opœ:al name of the red corn poppy. See *Papaver erraticum*.

PAPAYER SOMNIFERUM. The systematic name of the white poppy. See *Papaver erraticum*.

PAPAW. The fruit of the *Carica papaya* of LINNÆUS, a native of both Indies, and Guinea coast of Africa. When the roundish fruit are nearly ripe, the inhabitants of India boil and eat them with their meat, as we do turnips. They have somewhat the flavour of a pompon. Previous to boiling, they soak them for some time in salt and water, to extract the corrosive juice, unless the meat they are to be boiled with should be very salt and old, and then this juice being in them will make them as tender as a chicken. But they mostly pickle the long fruit, and thus they make no bad succedaneum for mango. The buds of the female flowers are gathered, and made into a sweetmeat; and the inhabitants are such good husbands of the produce of this tree, that they boil the shells of the ripe fruit into a repast, and the insides are eaten with sugar in the manner of melons.

PAPILLA. 1. The nipple of the breast. See *Nipple*.

2. The fine terminations of nerves, &c. as the nervous papillæ of the tongue, skin, &c.

PAPILLÆ MEDULLARÆ. Small eminences on the medulla oblongata.

PAPILLARIS HERBA. See *Lapsana*.

PAPPUS. The hair on the middle of the chin. See *Capillus*.

PAPULA. (*Papula*, dim. of *pappa*, a dug or nipple.) A very small and accumulated elevation of the cuticle, with an inflamed base, not containing a fluid, nor tending to suppuration. The duration of papulæ is uncertain, but they terminate for the most part in scurf.

PAR CUCULLARE. So Casserius calls the *Musculus Crico-Arytanoideus*.

PAR VAGUM. (*Par*, a pair.) The eighth pair of nerves. They arise from the corpora olivaria of the medulla oblongata, and proceed into the neck, thorax, and abdomen. In the neck the par vagum gives off two branches, the lingual and superior laryngeal; and, in the thorax, four branches, the recurrent laryngeal, the cardiac, the pulmonary, and the œsophageal plexuses. At length the trunks of the nervi vagi, adjacent to the mediastinum, run into the stomach, and there form the stomachic plexus, which branches to the abdominal plexuses.

PARACENTESIS. (From *παράκέντιω*, to pierce through.) The operation of tap-

ping to evacuate the water in ascites, dropsy of the ovarium, uterus, &c.

PARACASTICOS. (From *παράκαζω*, to decline.) *Paracme*. The declension of any distemper; also, according to Galen, that part of life, where a person is said to grow old, and which he reckons from 35 to 49, when he is said to be old.

PARACOE. (From *παρά*, dim, and *ακουω*, to hear.) Difficult hearing; dulness of hearing.

PARACOLLETTICA. (From *παράκολλωμαι*, to glue together.) Agglutinants; substances which unite parts preternaturally separated.

PARACOPSE. (From *παράκωπτω*, to be delirious.) In Hippocrates it is a slight delirium.

PARACRISIS. (From *παράκρνω*, to deprecate.) A slight disarrangement of the faculties, where the patient is inattentive to what is said to him.

PARACUSIS. (From *παρα*, wrong, and *ακουω*, to hear.) Depraved hearing. Deafness. A genus of disease in the Class *Locales*, and Order *Dysæsthesiæ* of Cullen. There are two species, 1. *Paracusis imperfecta. Surditas*. When existing sounds are not heard as usual. 2. *Paracusis imaginaria*, called also *susurrus. Syrygmus. Syrygmus. Tinnitus aurium*. When imaginary sounds are heard, not from without, but excited within the ear.

PARACYNARCHE. From *παρά*, a dog, and *αγγω*, to strangle.) A species of *Quinty*: it being a distemper to which dogs are subject.

PARADISUS. (Heb.) A pungent seed resembling the cardamom, is named *granum paradisi*, from its virtues

Paradisi grana See *Grana paradisi*.

PARAGLOSSA. (From *παρά*, and *γλωσσα*, the tongue.) A prolapsus of the tongue, a swelled tongue.

PARAGOSE. (From *παράγω*, to adduce.) Signifies that fitness of the bones to one another, as is discernible in their articulation; and bones which are thereby easier of reduction, when dislocated, are by Hippocrates called *παράγωγοι*.

PARALAMPSIS. (From *παράλαμπω*, to shine a little.) Some writers use this word to express a cicatrix in the transparent part of the cornea of the eye.

PARALLAGMA. (From *παράλλαττω*, to change.) *Parallaxis*. The transmutation of a solid part from its proper place, as where one part of a broken bone lies over another.

PARALLAXIS. See *Parallagma*.

PARALLELA. (From *παράλληλος*, parallel.) A sort of scurf or leprosy, affecting only the palms of the hands, and running down them in parallel lines; it happens sometimes in the venereal disease.

PARALOGIA. (From *παράλογος*, to talk absurdly.) A delirium in which the patient talks wildly.

PARALOPHIA. (From *παρα*, near, and *λοφία*, the first vertebra of the back.) Thus some anatomical writers, as Keil, &c. express the lower and lateral part of the neck near the vertebrae.

PARALYSIS. (From *παράλυσις*, to loose or weaken.) *Catalysis.* *Attonitus morbus.* *Stuper.* *Tremor.* The palsy. A genus of disease in the Class *Neuroses*, and Order *Comata*, of Cullen, known by a loss or diminution of the power of voluntary motion, affecting certain parts of the body, often accompanied with drowsiness. In some instances, the disease is confined to a particular part; but it more usually happens that one entire side of the body from the head downwards is affected. The species are: 1. *Paralysis partialis*, partial, or palsy of some particular muscle. 2. *Paralysis hemiplegica*, palsy of one side, longitudinally. 3. *Paralysis paraplegica*, palsy of one half of the body, taken transversely, as both legs and thighs. 4. *Paralysis venenata*, from the sedative effects of poisons. Paralysis is also symptomatic of several diseases, as worms, scrophula, syphilis, &c.

It may arise in consequence of an attack of apoplexy. It may likewise be occasioned by any thing that prevents the flow of the nervous power from the brain into the organs of motion; hence tumours, ever distention and effusion, often give rise to it. It may also be occasioned by morbid matter to the head, by the suppression of usual evacuations, and by the pressure made on the nerves by luxations, fractures, wounds, or other external injuries. The long continued application of sedatives will likewise produce palsy, as we find those, whose occupations subject them to the constant handling of white lead, and those who are much exposed to the poisonous fumes of metals, or minerals, are very apt to be attacked with it. What ever tends to relax and enervate the system, may likewise prove an occasional cause of this disease.

Palsy usually comes on with a sudden and immediate loss of the motion and sensibility of the parts; but in a few instances, it is preceded by a numbness, coldness, and paleness, and sometimes by slight convulsive twitches. When the head is much affected, the eye and mouth are drawn on one side, the memory and judgment are much impaired, and the speech is indistinct and incoherent. If the disease affects the extremities, and has been of long duration, it not only produces a loss of motion and sensibility, but likewise a considerable flaccidity and

wasting away in the muscles of the parts affected.

When palsy attacks any vital part, such as the brain, heart, or lungs, it soon terminates fatally. When it arises as a consequence of apoplexy, it generally proves very difficult of cure. Paralytic affections of the lower extremities ensuing from any injury done to the spinal marrow, by blows and other accidents, usually prove incurable. Palsy, although a dangerous disease in every instance, particularly at an advanced period of life, is sometimes removed by the occurrence of a diarrhoea or fever.

The morbid appearances to be observed on dissections in palsy, are pretty similar to those which are to be met with in apoplexy; hence collections of blood, and of serous fluids, are often found effused on the brain, but more frequently the latter, and in some instances the substance of this organ seems to have suffered an alteration. In palsy, as well as in apoplexy, the collection of extravasated fluid is generally on the opposite side of the brain to that which is affected.

PARALYSIS HERBA. (From *παράλυσις*, to weaken; so called from its use in paralytic disorders.) The cowslip and primrose are sometimes so termed. See *Primula veris*, and *Primula vulgaris*.

PARAMERIA. (From *παρα*, near, and *μῆρος*, the thigh.) The inward parts of the thigh.

PARANESUS. (From *παρα*, near, and *μῆρος*, the middle.) The ring finger, or that which is between the middle and the little fingers.

PARANGA. (From *παρα*, dim, and *ῥα*, to understand.) Alienation of mind; defect of judgment.

PARAPECHYON. (From *παρα*, near, and *πῆχυς*, the cubit.) That part of the arm from the elbow to the wrist.

PARAPHORA. (From *παράφωρος*, to transfer.) A slight kind of delirium, or light-headedness in a fever: some use this word for a delirium in general.

PARAPHRENESIS. A delirium; also paraphrenitis.

PARAPHROSYNE. (From *παράφροσις*, to be estranged in mind.) The same as *Mania*.

PARAPHONIA. (From *παρα*, wrong, and *φωνή*, sound.) Alteration of the voice. A genus of disease in the Class *Locales*, and Order *Dyscinesia* of Cullen, comprehending six species, viz. *paraphonia puberum*, *paraphonia rauca*, *paraphonia resonans*, *paraphonia palatina*, *paraphonia clangens*, and *paraphonia comatosa*.

PARAPHORA. (From *παράφωρος*, to transfer.) A slight alienation of the mind.

PARAPHRENITIS. (From *παρα*, dim,

and *egg*, the mind.) *Diaphragmitis*. An inflammation of the diaphragm. A genus of disease in the Class *Pyrexia* and Order *Phlegmaria* of Cullen, known by delirium, with difficulty of breathing, and pain in the region of the diaphragm.

PARAPHIMOSIS. (From *παγα*, about, and *πυμω*, to bridle.) The paraphimosis is a disorder wherein the prepuce, being retracted toward the root of the penis, cannot be returned again over the glans, but makes a sort of ligature behind the corona. It is easily known; the glans is uncovered, the skin tumefied on the corona, and above it forms a circular collar or stricture, which, from the skin being unequally extended, becomes indented, and makes several rings round the part. This disease may proceed from two causes; as first from the imprudence of young people, and sometimes also of grown persons, who having the end of their prepuce too strait, cannot uncover their glans without pain, and when they have done it, neglect returning it so soon as they ought; and thus the contracted part of the prepuce forms a constriction behind the glans. Soon after, the glans and penis swell, and the prepuce being consequently very much distended, is affected in the same manner; an inflammation seizes upon both, and swellings quickly appear upon the stricture formed by the prepuce, so that the whole may be liable to a gangrene, if not speedily relieved. The second thing that may produce a paraphimosis, is a venereal virus. In adults, whose glans is uncovered, there frequently arise venereal chancres in the prepuce after impure coition, which before they digest, are generally attended with inflammation, more or less considerable. This inflammation is alone sufficient to render the prepuce too strait for the size of the penis, in consequence of which a swelling or inoculation may ensue like that before mentioned; and this is what is termed a paraphimosis.

PARAPLEGIA. (From *παρεπλησσω*, to strike inharmoniously.) Palsy of one half of the body taken transversely. A species of paralysis. See *Paralysis*.

PARAPOPLEXIA. (From *παγα* dim, and *εποπληξια*, an apoplexy.) A slight apoplexy.

PARATHRONA. (From *παγα*, and *αρθρον*, a joint.) A slight luxation; a tumour from protrusion, as an hernia.

PARATHRONATA. Plural of *parathroma*, and synonymous with *ectopie*.

PARASYTOS. (From *παγα*, and *συμτος*, number.) A pulse not suitable to the age of the person.

PARASCEPASTRA. (From *παγα*, and *επισκεψω*, to cover.) A cap or bandage to go round the whole head.

PARASCHIDE. (From *παγα*, and *σχίζω*,

to cleave.) A fragment or fissure in a broken bone.

PARASITIC. (*Parasitica*; from *πασιτικός*, a parasite or hanger on.) Animals and plants are so termed that receive their nourishment in the bodies of others, as worms, polypes, hydatids, &c.

PARASPHAGIS. (From *παγα*, near, and *σφαγν*, the throat.) The part of the neck contiguous to the clavicles.

PARASTATA. (From *πασιςτις*, to stand near.) It signifies any thing situated near another.

PARASTATON. (From *πασιςτις*, to stand near.) In Hippocrates it signifies the *Epididymis*. Herophilus and Galen called these the *Varicose Parastata*, to distinguish them from the *Glandulose Parastata*, now called *Prostate*. Rufus Ephesius called the tube Fallopiæ by the name of *Parastata Varicose*.

PARASTREMA. (From *παγαστρεψω*, to distort, or pervert.) A perversion, or convulsive distortion of the mouth, or any part of the face.

PARASTRANCHE. A species of Quinsy. See *Paracynanche*.

PARATHENAR. (From *παγα*, near, and *σναγ*, the sole of the foot.) A muscle situated near the sole of the foot.

Parathenar minor. See *Flexor brevis minimi digiti pedis*.

PARDALIUM. (From *παρδος*, the panther.) An ointment smelling like the panther.

Paragoric Elixir. See *Tinctura opti camphorata*.

PAREGORICS. (*Medicamenta paregorica*, *παεργωγικά*; from *παεργωγν*, to mitigate, to assuage.) Medicines that allay pain are so termed.

PARRIA. *Παρια*. That part of the face which is between the eyes and chin.

PARRIA BRAVA. See *Pariera brava*.

PAROEPHALIS. (From *παγα*, near, and *εγκεφαλον*, the brain.) See *Cerebellum*.

PARENCHYMA. (From *παεργωγν*, to strain through, because the ancients believed the blood was strained through it.) The spongy and cellular substance that connects parts together. It is now only, in anatomy, applied to the connecting medium of the substance of the lungs.

PARESIS. (From *παεσμι*, to relax.) An imperfect palsy.

PARIERA BRAVA. (Span.) *Parcyra. Ambutua. Butua. Overo butua.* The root of this plant, *Cissampelos pareira* of Linnaeus, *foliis pellatis cordatis emarginatis*. Class, *Dissecta*. Order, *Monodelphia*. A native of South America and the West Indies, has no remarkable smell, but to the taste it manifests a notable sweetness of the liquorice kind, together with a considerable bitterness, and a slight rough-

ness covered by the sweet matter. The facts adduced on the utility of the *radix pariera brava* in the nephritic and calculous complaints, are principally mentioned by foreigners, and no remarkable instances of its efficacy are recorded by English practitioners.

PARIETAL BONES. (*Parietalia*, from *paries*, a wall, because they defend the brain like walls.) *Ossa verticis*. *Ossa synipitidis*. *Ossa verticalia vel bregmatica*. Two arched and somewhat quadrangular bones, situated one on each side of the superior part of the cranium. Each of these bones forms an irregular square. They are thicker above than below; but are somewhat thinner, and at the same time more equal and smooth than the other bones of the cranium. The only foramen we observe in them, is a small one towards the upper and posterior part of each. It has been named the parietal foramen, and serves for the transmission of a small vein to the longitudinal sinus. In many subjects this foramen is wanting. On the inner surface of these bones are the marks of the vessels of the dura mater, and of the convoluted surface of the brain. On the inside of their upper edge we may likewise observe a considerable furrow, which corresponds with the longitudinal sinus of the dura mater; and lower down, towards their posterior and inferior angle, is a smaller one for part of the lateral sinuses. These bones are joined to each other by the sagittal suture; to the os sphenoides, and ossa temporum, by the squamous suture; to the os occipitis by the lambdoidal suture; and to the os frontis by the coronal suture. Their connexion with this latter bone is well worthy our attention. We shall find that in the middle of the suture, where the os frontis from its size and flatness is the most in danger of being injured, it rests upon the arch formed by the parietal bones; whereas at the sides, the parietal bones are found resting upon the os frontis because this same arch is there in the greatest danger from pressure. In new born infants, the ossa parietalia are separated from the middle of the divided os frontis by a portion of the cranium, then unossified. When the finger is applied to this part, the motion of the brain, and the pulsation of the arteries of the dura mater, may be easily distinguished. In general the whole of this part is completely ossified before we are seven years of age.

PARIETARIA. (From *paries* a wall; because it grows upon old walls, and among rubbish.) 1. The name of a genus of plants in the Linnæan system. Class, *Polygamia*. Order, *Monœcia*.

2. The pharmacopœical name of the wall pellitory. *Parietaria officinalis* of

Linnæus:—*foliis lanceolato-ovatis, pedunculis dichotomis, calycibus diphyllis*. This plant has no smell, and its taste is simply herbaceous. In the practice of the present day it is wholly laid aside, although it was formerly in high estimation as a diuretic.

PARIETARIA OFFICINALIS. The systematic name of the wall pellitory. See *Parietaria*.

PARIS. (So called in reference to the youth of that name, who adjudged the golden apple to Venus, this herb bearing but one seed.) 1. The name of a genus of plants in the Linnæan system. Class, *Oc-tandria*. Order, *Tetragynia*.

2. The pharmacopœical name of the herb Paris. The herb true-love. *Paris quadrifolia* of Linnæus. The colour and smell of this plant indicate its possessing narcotic powers. The leaves and berries are said to be efficacious in the cure of whooping-cough, and to act like opium. Great caution is requisite in their exhibition, as convulsions and death are caused by an overdose. The root possesses emetic qualities.

PARIS QUADRIFOLIA. The systematic name of the herb Paris. See *Paris*.

PARIETARIA. (From *paries*, and *terram*: the part of the throat near which the tonsils are.) A part of the throat near the tonsils, or disorders of the tonsils.

PARIETIMOTOMUS. (From *paries*, the tonsils, and *tema*, to cut.) An instrument with which the tonsils were formerly scarified.

PARODONTIS. (From *paries*, near, and *odus*, a tooth.) A painful tubercle upon the gums.

PARONYCHIA. (From *paries*, about and *onyx*, the nail.) *Panaris*. *Punarium*. A whitlow, or whitloe. Any collection of pus formed in the fingers is termed by authors panaris, or whitloe, and is an abscess of the same nature with those arising in other parts of the body. These abscesses are situated more or less deep, which has induced the writers upon the subject to divide them into several species: accordingly they have ranged them under four heads, agreeable to the places where they are formed. The first kind of panaris is formed under the cuticle, on one side of the nail, and sometimes all round it. The second is seated in the fat lying under the skin, between that and the sheath which involves the flexor tendons. The third is described by authors to be formed within the sheath; and they still add a fourth species, arising between the periosteum and the bone.

PAROPTIS. (From *paries*, near, and *optis*, the eye.) The external angles of the eyes.

PAROPTESIS. (From *paries*, and *optas*, to roast.) A provocation of sweat, by mak-

ing a patient approach the fire, or by placing him in a bagnio.

PARORASIS. (From *para*, dim, and *oza*, to see.) An imbecility of sight.

PARORCHIDIUM. (From *para*, and *orchis*, a testicle.) A tumour in the groin, occasioned by the testicle, which is passing into the scrotum.

PAROTID GLAND. (*Glandula parotidea*; from *para*, about, and *ot*, the ear.)

Parotis. A large conglomerate and salival gland, situated under the ear, between the mamillary process of the temple bone and the angle of the lower jaw. The excretory duct of this gland opens in the mouth, and is called, from its discoverer, the *Stenonian duct*.

PAROTIDEA. (From *parotis*, the parotid gland.) The trivial name of a species of quinsay, in which the parotid gland, neck and throat, are considerably affected. See *Cynanche*.

PAROTIS. (From *para*, near, and *ot*, the ear.) See *Parotid gland*.

PAROXYSM. (*Paroxysmus*; from *para*, to aggravate.) A periodical exacerbation or fit of a disease.

Parsley, black mountain. See *Oreoselinum*.

Parsley, common. See *Petroselinum*.

Parsley, Macedonian. See *Petroselinum Macedonicum*.

Parnip. See *Pastinaca*.

Parnip, water. See *Sium*.

PARTHENIASTRUM. (Dim. of *parthenium*, tansy.) A species of tansy, or bastard feverfew.

PARTHENIS. The same as *parthenium*.

PARTHENIUM. (From *παρθενος*, a virgin; so called because of its uses in diseases of young women.) See *Matricaria*.

Parthenium nas. See *Tanacetum*.

PARULIS. (From *para*, near, and *ulor*, the gum.) An inflammation, boil, or abscess in the gums.

PARYGROX. (From *para*, and *υγρος*, humid.) A liquid or moist preparation for allaying a topical inflammation.

PASIPHILUS. (From *pas*, all, and *φιλος*, grateful; from its general usefulness.) A name given to a vitriolic plaster.

PASSA. (From *para*, to sprinkle over.) See *Catapasma*.

PASSA. (From *pauo*, to spread.) In *Paracelsus* it is a whitloe.

Passa minor. See *Uva passa minor*.

PASSAVANTICUS. (From *pas*, all, and *ανωσ*, to dry up.) An epithet given by Schroder to a powder which dries up and evacuates morbid humours.

PASSIFLORA. The name of a genus of plants in the Linnæan system. Class, *Gynandria*. Order, *Pentandria*.

Passiflora laurifolia. Bay-leaved

passion-flower. A native of Surinam. The fruit have a delicious smell and flavour, and are excellent for quenching thirst, abating heat of the stomach, increasing the appetite, recruiting the spirits, and allaying the heat in burning fevers.

PASSIFLORA MALIFORMIS. Apple-shaped granadilla. The fruit of this species of passion flower is esteemed a delicacy in the West Indies, where it is served up at table in desserts: they are not unwholesome.

Passion, colic. See *Diarrhœa*.

Passion, hysteric. See *Hysteria*.

Passion, iliac. See *Iliac passion*.

PASSULUM MAJORE. See *Uva passa major*.

PASSULATUM. (From *passa*, a fig, or raisin.) Is a term given by Dispensatory writers to some medicines, where raisins are the chief ingredient, as the electarium *passulatum*, &c.

PASSUM. (From *passa*, a grape, or raisin.) Raisin-wine.

PASTA RECIA. (From *parso*, to sprinkle.) A lozenge, or small cake, sprinkled over with some dry powdered substance.

PASTILLUM. (Dim. of *pasta*, a lozenge.) *Pastillus.* A little lump of paste, or ball, made to take like a lozenge, a troch, or pastil.

PASTINACA. (A *pasta*; from its usefulness as a food.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digymia*. *Parnip*.

2. The pharmacopœial name of the *parnip*. The cultivated or garden *parnip* is the *Pastinaca sativa* of Linnæus:—*folioli simpliciter pinnatis. Elaphoboscum*. Its roots are sweet and nutritious, and in high esteem as an article of food. They possess an aromatic flavour, more especially those of the wild plant, and are exhibited in calculous complaints for their diuretic and sheathing qualities.

PASTINACA OPOPANAX. The systematic name of the plant which yields *opopanax*. See *Opopanax*.

PASTINACA SATIVA. The systematic name of the *parnip*. See *Pastinaca*.

PATELLA. (Dim. of *patina*, a dish; so named from its shape.) *Rotula*. The knee-pan. A small flat bone, which, in some measure, resembles the common figure of the heart, with its point downwards, and is placed at the fore part of the joint of the knee. It is thicker in its middle part than at its edge. Anteriorly it is a little convex, and rough for the insertion of muscles and ligaments; posteriorly, it is smooth, covered with cartilage, and divided by a middle longitudinal ridge, into two slightly concave surfaces, of which the external one is the largest and deepest. They are both exactly adapted to the pul-

key of the os femoris. The edges of this posterior surface are rough and prominent, where the capsular ligament is attached, and below is a roughness at the point of the bone, where the upper extremity of a strong tendinous ligament is fixed, which joins this bone to the tuberosity at the upper end of the tibia. This ligament is of considerable thickness, about an inch in breadth, and upwards of two inches in length. The rotula is composed internally of a cellular substance, covered by a thin bony plate; but its cells are so extremely minute, that the strength of the bone is, upon the whole, very considerable. In new-born children, it is entirely cartilaginous. The use of this bone seems to be, to defend the articulation of the joint of the knee from external injury. It likewise tends to increase the power of the muscles which act in the extension of the leg, by removing their direction farther from the centre of motion, in the manner of a pulley. When we consider the manner in which it is connected with the tibia, we find that it may very properly be considered as an appendix to the latter, which it follows in all its motions, so as to be to the tibia what the olecranon is to the ulna; with this difference, however, that the rotula is moveable, whereas the olecranon is a fixed process. Without this mobility, the rotatory motion of the leg would have been prevented.

PATHETICI. (*Patheticus*; from *παθος*, an affection; because they direct the eyes to express the passions of the mind.) *Trochleatores.* The fourth pair of nerves. They arise from the crura of the cerebellum laterally, and are distributed in the musculus obliquus superior seu trochlearis.

PATHOGNOMONIC. (*Pathognomonicus*: from *παθος*, a disease, and *γνωσκω*, to know.) A term given to those symptoms which are peculiar to a disease. They are also termed proper or characteristic symptoms.

PATHOLOGY. (*Pathologia*; from *παθος*, a disease, and *λογος*, a discourse.) The doctrine of diseases. It comprehends *nomenclology, etiology, symptomatology, semiotics, and therapia.*

PATIENTIA. (From *patior*, to bear, or suffer.) The name of the herb monk's rhubarb, from its gentle purging qualities. See *Rhabarbarum.*

Patience, garden. See *Rhabarbarum monachorum.*

PATOR NARIUM. (From *pateo*, to be opened.) The sinus, cavity, or chasm of the nose.

PATRUM CORTEX. (So called from the Jesuits, called fathers in the church of Rome, who first spread its use in Europe.) See *Cinchona.*

PATURIA. The venereal disease.

PAULINA, CONFECTIO. (From *Paulina*, to rest.) It is a warm opiate, such as the London College have called *Confectio Opia*, in their Dispensatory. It is the *Paulina* of Aristarchus, which is the same with the *Confectio Archigenis.*

Paul's betony. Set *Veronica.*

PAVANA. See *Lignum pavane.*

PAVOR. (From *paveo*, to fear; so called from the dread there is of approaching or touching a person affected with it.) The itch.

PEA. A species of pulse of great variety, and much in use as a nourishing article of diet.

Peach. See *Persica.*

Peagle. See *Primula veris.*

Pearl. See *Margarita.*

Pearl-ashes. See *Potash.*

Pearl-barley. See *Hordeum.*

PEAR. Of pears there are a many varieties, affording an wholesome nourishment.

PECHEDIUM. *Παχιδιον.* The perinium.

PECHURIM CORTEX. An highly aromatic bark, the produce of a species of *Laurus*. It is extremely fragrant, like unto that of cinnamon, which it greatly resembles in its properties. In Lisbon it is much esteemed in the cure of dysenteries, and for allaying obstinate vomitings.

PECHURIM FABA. See *Faba pechurim.*

PECHURIM. See *Pechurim.*

PECHYAGRA. (From *παχυς*, the cubit, and *αγγρα*, a seizure.) The gout in the elbow.

PECHYS. *Παχυς.* The cubit, or elbow.

PECHTYRABE. An epithet for the scurvy.

Pecquet's duct. See *Thoracic duct.*

PECTEX. The pubes, or share-bone.

PECTINALIS. (*Pectinalis*, sc. *musculus*; so named from it arising at the *pecten*, or pubis.) *Pectineus* of authors, and *Pubio femoral* of Dumas. This is a small flat muscle, situated obliquely between the pubis and the little trochanter, at the upper and anterior part of the thigh. It arises broad and fleshy from all the anterior edge of the os pectinis, or pubis, as it is more commonly called, as far as its spine, and, descending obliquely backwards and outwards, is inserted by a short and broad tendon, into the upper and anterior part of the *linea aspera* of the os femoris, a little below the lesser trochanter. This muscle serves to bend the thigh, by drawing it upwards and inwards, and likewise assists in rolling it outwards.

PECTINATI MUSCULI. (*Pectinati*; from *pecten*, a comb: so named from their supposed resemblance.) The fasciculated muscular fibres of the right auricle of the heart.

Pectineus. See *Pectinalis.*

Pectoralis. See *Pectoralis major*.

PECTORALIS MAJOR. (*Musculus pectoralis*; from *pectus* the breast.) *Pectoralis* of authors, and *sterno-costo-clavio-humeral* of Dumas. This is a broad, thick, fleshy, and radiated muscle, situated immediately under the integuments, and covering almost the whole anterior part of the breast. Winslow calls it *pectoralis major*, to distinguish it from the *serratus anticus*, which he has named *pectoralis minor*. It arises from the cartilaginous extremities of the fifth and sixth ribs, from the last of which its tendinous fibres descend over the upper part of the obliquus externus and rectus abdominis, helping to form a part of the sheath in which the latter is included. It likewise springs from almost the whole length of the sternum by short tendinous fibres, which evidently decussate those on the other side; and tendinous and fleshy from more than a third of the anterior part of the clavicle. From these origins the fibres run in a folding manner towards the axilla, and are inserted by a broad tendon into the os humeri, above the insertion of the deltoid muscle, and at the outer side of the groove which lodges the tendon of the long head of the biceps: some of its fibres likewise extend into that groove; and, from the lower part of this tendon, which is spread near two inches along the os humeri, we find it sending off other fibres, which help to form the fascia that covers the muscles of the arm. It often happens that that part of the pectoralis which arises from the clavicle, is separated from the inferior portion, so as to appear like a distinct muscle. This has induced Winslow to divide it into parts, one of which he calls the *clavicular*, and the other the *thoracic* portion. Sometimes these two portions are inserted by separate tendons, which cross one another at the upper and inner part of the os humeri, the tendon of the thoracic portion being inserted at the outer edge of the bicipital groove, immediately behind the other. This muscle, and the latissimus dorsi, form the cavity of the axilla, or arm-pit. The use of the pectoralis is to move the arm forwards, or to raise it obliquely towards the sternum. It likewise occasionally assists in moving the trunk upon the arm; thus, when we exert any efforts with the hand, as in raising ourselves from off an arm-chair, or in sealing a letter, the contraction of this muscle is particularly observable. To these uses Haller adds that of assisting in perspiration, by raising the sternum and ribs. He tells us he well remembers, that when this muscle was affected by rheumatism, his breathing was incommoded: and that, when troubled with difficulty of respiration, he had often found himself greatly relieved by raising and drawing back his

shoulders, keeping his arms at the same time firmly fixed. Winslow, however, has denied this use, and Albinus has omitted it, probably because it does not take place in a natural state.

PECTORALIS MINOR. *Serratus anticus* of Albinus. Douglas and Cowper call this muscle *Serratus minor anticus*, and Winslow gives it the name of *Pectoralis minor*, and Dumas calls it *Costo-coracoidien*. It is a fleshy and pretty considerable muscle, situated at the anterior and lateral part of the thorax, immediately under the pectoralis major. It arises from the upper edges of the third, fourth, and fifth ribs, near where they join with their cartilages by an equal number of tendinous and fleshy digitations, which have been compared to the teeth of a saw, whence this and some other muscles, from their having a similar origin, or insertion, have gotten the name of *serrati*. From these origins it becomes thicker and narrower as it ascends, and is inserted by a flat tendon into the upper part of the coracoid process of the scapula. The principal use of this muscle is to draw the scapula forwards and downwards; and when that is fixed, it may likewise serve to elevate the ribs.

PECTORALS. (*Medicamenta pectoralia*; from *pectus*, the breast.) Medicines that relieve disorders of the chest.

PECTUS. The breast. See *Thorax*.

PECTUSCULUM. (Dim. of *pectus*, the breast; so named from its shape.) The metatarsus.

PEDEULATIO. *Morbus Pedicularis. phlegmasis.* That disease of the body in which lice are continually bred on the skin.

PEDES HIPPOCAMPI. (*Pes*, a foot, so named from their resemblance to the feet of the *hippocampus*, or sea-horse.) Two columns at the end of the fornix of the brain, which diverge posteriorly.

PEDETHMUS. (From *πρὸς*, to leap.) The motion which is sensible in the arteries from the impulse of the blood. The pulse.

PEDIASMUS. (From *πῆδος*, a field.) An epithet of a species of wild myrrh.

PEDICULARIA. (From *pediculus*, a louse; so called from its use in destroying lice.) The herb staves-acre. See *Staphisagria*.

PEDICULUS. (Dim. of *pes*, a foot; so named from its many small feet.) 1. A louse.

2. A pedicle or foot-stalk of a flower, or leaf.

Pedicus. See *Extensor digitorum brevis*.

PEDILUVIUM. (From *pedes*, the feet, and *lavo*, to wash.) A bath for the feet.

PEDION. (From *πῆς*, the foot.) The sole of the foot.

PEDORA. (From *pes*, a foot.) The sordes of the eyes, ears, and feet.

PEGANELNUM. (From *αργαίον*, rue, and *ελαιον*, oil.) Oil of Rue.

PEGANERUM. (From *αργαίον*, rue.) A plaster composed of rue.

PEGANUM. (From *αργαίον*, to compress; so called because, by its dryness, it condenses the seed.) Rue.

PEGE. (Πηγή, a fountain.) The internal angles of the eyes are called *Pege*.

PELADA. A species of baldness, a shedding of the hair from a venereal cause.

PELAGRA. The disease called the *pelagra* does not appear to have been noticed by any of our nosologists. Indeed few accounts of it have hitherto been published, although the peculiar symptoms with which it is attended, and the fatal consequences which generally ensue from it, render it equally curious and important. In certain districts, as Milan and Padua, in Italy, where it is peculiarly prevalent, it is computed to attack five inhabitants out of every hundred. The following account of this singular disease is extracted from Dr. Jansen's treatise on the subject, who had seen the disease at Milan:

About the month of March or April, when the season invites the farmers to cultivate their fields, it often happens that a shining red spot suddenly arises on the back of the hand, resembling the common erysipelas, but without much itching or pain, or indeed any other particular inconvenience. Both men and women, girls and boys, are equally subject to it. Sometimes this spot affects both hands, without appearing on any other part of the body. Not uncommonly it arises also on the shins, sometimes on the neck, and now and then, though very rarely, on the face. It is sometimes also seen on the breasts of women, where they are not covered by the clothes, but such parts of the body as are not exposed to the air, are very seldom affected; nor has it ever been observed to attack the palm of the hand, or the sole of the foot. This red spot elevates the skin a little, producing numerous small tubercles of different colours; the skin becomes dry and cracks, and the epidermis sometimes assumes a fibrous appearance. At length it falls off in white furfuraceous scales; but the shining redness underneath still continues, and, in some instances, remains through the following winter. In the mean time, excepting this mere local affection, the health is not the least impaired, the patient performs all his rural labours as before, enjoys a good appetite, eats heartily, and digests well. The bowels are generally relaxed at the very commencement of the disease, and continue so throughout its whole course. All the other excretions are as usual; and, in females, the menses return at their accustomed periods, and in

the proper quantity. But what is most surprising is, that in the month of September, when the heat of the summer is over, in some cases sooner, in others later, the disorder generally altogether disappears, and the skin resumes its natural healthy appearance. This change has been known to take place as early as the latter end of May or June, when it has only been in its earliest stage. The patients, however, are not now to be considered as well; the disease hides itself, but is not eradicated: for, no sooner does the following spring return, but it quickly re-appears, and generally is accompanied with severer symptoms. The spot grows larger, the skin becomes more unequal and hard, with deeper cracks. The patient now begins to feel uneasiness in the head, becomes fearful, dull, less capable of labour, and much wearied with his usual exertions. He is exceedingly affected with the changes of the atmosphere, and impatient both of cold and heat. Nevertheless he generally gets through his ordinary labour, with less vigour and cheerfulness indeed than formerly, but still without being obliged to take to his bed: and, as he has no fever, his appetite continues good, and the chylopoietic viscera perform their proper functions. When the *pelagra* has even arrived at this stage, the returning winter, nevertheless, commonly restores the patient to apparent health; but the more severe the symptoms have been, and the deeper root the disease has taken, the more certainly does the return of spring produce it with additional violence. Sometimes the disease in the skin disappears, but the other symptoms remain notwithstanding. The powers both of the mind and body now become daily more enfeebled; peevishness, watchings, vertigo, and, at length, complete melancholy, supervene. Nor is there a more distressing kind of melancholy any where to be seen, than takes place in this disease. "On entering the hospital at Legnano," says Dr. Jansen, "I was astonished at the mournful spectacle I beheld, especially in the women's ward. There they all sat, indolent, languid, with downcast looks, their eyes expressing distress, weeping without cause, and scarcely returning an answer when spoken to; so that a person would suppose himself to be among fools and mad people: and, indeed, with very good reason; for gradually this melancholy increases, and at length ends in real mania.

"Many, as I had an opportunity of observing in this hospital, were covered with a peculiar and characteristic sweat, having a very offensive smell, which I know not how better to express than by comparing it to the smell of mouldy bread. A person accustomed to see the disease

would at once recognise it by this single symptom. Many complained of a burning pain at night in the soles of the feet, which often deprived them of sleep. Some with double vision; others with fatuity; others with visceral obstructions; others with additional symptoms. Nevertheless, fever still keeps off, the appetite is unimpaired, and the secretions are regularly carried on. But the disease goes on increasing, the nerves are more debilitated, the legs and thighs loose the power of motion, stupor or delirium come on, and the melancholy terminates in confirmed mania. In the hospital at Legnano, I saw both men and women in this maniacal state. Some lay quiet; others were raving, and obliged to be tied down to the bed, to prevent them from doing mischief to themselves and others. In almost all these, the pulse was small, slow, and without any character of fever. One woman appeared to have a slight degree of furor uterinus; for, at the sight of men, she became merry, smiled, offered kisses, and by her gestures desired them to come towards her. Some were occupied in constant prayers; some pleased themselves with laughter, and others with other things. But it was remarkable, that all who were in this stage of the disease, had a strong propensity to drown themselves. They now begin to grow emaciated, and the delirium is often followed by a species of tabes. A colliquative diarrhœa comes on, which no remedy can stop, as also has been observed in nostalgia. Sometimes, in the pelagra, the diarrhœa comes on before the delirium, and the delirium and stupor mutually interchange with each other. The appetite often suddenly failed, so that the sick will sometimes go for near a week without tasting food. Not uncommonly it returns as suddenly, so that they eagerly devoured whatever was offered them, and this even at times when they are horribly convulsed. The convulsions with which they are attacked, are most shocking to see, and are of almost every kind, catalepsy excepted, which has been described by writers. I saw one girl in bed, who was violently distorted by opisthotonos every time she attempted to rise. Some are seized with emprosthotonos; and others with other species of tetanus. At length, syncope and death close the tragedy, often without any symptom of fever occurring through the whole course of the disease. The first stage of the pelagra, in which the local affection only takes place, Dr. Jansen observes, continues in some instances for a great length of time; persons being occasionally met with in whom it has lasted six or eight, or even fifteen years, disappearing regularly every win-

ter, or returning again in the spring. This occasions some of the inhabitants to pay little attention to it; although, in other cases, it reaches its greatest height after the second or third attack. It appears that this disease is not infectious, and that the causes producing it are yet uncertain. It has been supposed, by some, to arise from the heat of the sun's rays; and hence it is now and then called *mal de sole*; but this does not produce any similar disease in other parts of the world, where it is in an equal or even much greater degree than at Milan; no disease, in any respect resembling it, having hitherto been noticed in such regions, except the lepra asturiensis described by Thiery, and after him by Sauvages. In this, a tremor of the head and trunk of the body takes place, which does not happen in the pelagra. This, however, is the principal difference in the two diseases.

PELARIUM. (From *πηλος*, mud; so called from its muddy consistence.) A collyrium.

Pelato salpingæus. See *Circumflexus*.

PELECANUS. (From *πελεκας*, to perforate; so named from its curvature at the end resembling the beak of the Pelican.)

1. The Pelican.

2. An instrument to draw teeth.

PELECINUM. (From *πελεκινος*, a hatchet; so called because its seeds are shaped like a two-edged hatchet.) The hatchet vetch.

PELIOMA. (From *μελος*, black.) An echymosis when liver-coloured.

Pelitory, bastard. See *Ptarmica*.

Pelitory of Spain. See *Pyrethrum*.

Pellitory, wall. See *Parietaria*.

PELMA. (From *πελαω*, to move forwards.) The sole of the foot, or a sock adapted to the sole of the foot.

PELTALIS CARTILAGO. (From *pelta*, a buckler; so called from its shape.) The scutiform cartilage of the larynx.

PELVIC LIGAMENTS. The articulation of the os sacrum with the last lumbar vertebra, and with the ossa innominata, is strengthened by means of a strong transverse ligament, which passes from the extremity and lower edge of the last lumbar vertebra, to the posterior and internal surface of the spine of the ilium. Other ligaments are extended posteriorly from the os sacrum to the ossa ilia on each side, and, from the direction of their fibres, may be called the lateral ligaments. Besides these, there are many shorter ligamentous fibres, which are seen stretched from the whole circumference of the articulating surfaces of these two bones. But the most remarkable ligaments of the pelvis are the two *sacro-ischiatric* ligaments, which are placed towards the posterior and inferior part of the pelvis. One of

these may be called the greater, and the other the lesser sacro-ischiatic ligament. The first of these is attached to the posterior edge of the os sacrum, to the tuberosity of the ilium, and to the first of the three divisions of the os coccygis. Its other extremity is inserted into the inner surface of the tuberosity of the ischium. At its upper part it is of considerable breadth, after which it becomes narrower, but expands again before its insertion into the ischium, and extending along the tuberosity of that bone to the lower branch of the os pubis, where it terminates in a point, forms a kind of falx, one end of which is loose, while the other is fixed to the bone. The lesser sacro-ischiatic ligament is somewhat thicker than the former, and is placed obliquely before it. It extends from the transverse processes of the os sacrum, and the tuberosity of the spine of the ilium, on each side, to the spine of the ischium. These two ligaments not only serve to strengthen the articulation of the ossa innominata with the os sacrum, but to support the weight of the viscera contained in the pelvis, the back and lower part of which is closed by these ligaments. The posterior and external surface of the greater ligament likewise serves for the attachment of some portions of the gluteus maximus and gemini muscles. The symphysis pubis is strengthened internally by a transverse ligament, some of the fibres of which are extended to the obturator ligament.

PELVIS. (From *πελως*, a basin, because it is shaped like a basin used in former times.) The cavity below the belly. The pelvis consists, in the child, of many pieces, but, in the adult, it is formed of four large bones, of the os sacrum behind, the ossa innominata on either side, and the os coccygis below. See *Os sacrum*, *Ossa innominata*, and *Os coccygis*. It is wide and expanded at its upper part, and contracted at its inferior aperture. The upper part of the pelvis, properly so called, is bounded by an oval ring, which parts the cavity of the pelvis from the cavity of the abdomen. This circle is denominated the brim of the pelvis; it is formed by a continued and prominent line along the upper part of the sacrum, the middle of the ilium, and the upper part, or crest, of the pubis. This circle of the brim supports the impregnated womb, keeps it up against the pressure of labour pains; and sometimes this line has been "as sharp as a paper-folder, and has cut across the segment of the womb;" and so, by separating the womb from the vagina, has rendered delivery impossible; and the child escaping into the abdomen the woman has died. The lower part of the pelvis is denominated

the outlet. It is composed by the arch of the pubis and by the sciatic ligaments; it is wide and dilatable, to permit the delivery of the child, but which being sometimes too wide, permits the child's head to press so suddenly, and with such violence upon the soft parts, that the perineum is torn.

The marks of the female skeleton have been sought for in the skull, as in the continuation of the sagittal suture; but the truest marks are those which relate to that great function by which chiefly the sexes are distinguished; for while the male pelvis is large and strong, with a small cavity, narrow openings, and bones of greater strength, the female pelvis is very shallow and wide, with a large cavity and slender bones, and with every peculiarity which may conduce to the easy passage of the child.

The office of the pelvis is to give a steady bearing to the trunk, and to connect it with the lower extremities, by a sure and firm joining, to form the centre of all the great motions of the body, to contain the internal organs of generation, the urinary bladder, the rectum, and occasionally part of the small intestines, and to give support to the gravid uterus.

PELVIS AURIUM. The cochlea in the ear.

PELVIS CEREBRI. The infundibulum in the brain.

PEMPHIGORES. (From *πνευμα*, a blast of wind.) A fever distinguished by flatulences and inflations, in which a sort of aerial vapour was said to pass through the skin.

PEMPHIGUS. (From *πνευμα*, a bubble, or vesicle.) *Febris bullosa*. *Exanthemata serosa*. *Morta*. *Pemphigus helveticus*. *Pemphigus major*. *Pemphigus minor*. The vesicular fever. A fever attended by successive eruptions of vesicles about the size of almonds, which are filled with a yellowish serum, and in three or four days subside. The fever may be either synocha or typhus. It is a genus of disease in the Class *Pyrexia* and Order *Exanthemata* of Cullen. The latest writers on this disease contend, that it is sometimes acute and sometimes a chronic affection; that the former is constantly attended with fever, the latter is constantly without; that in neither case is it an acrimonious or contagious matter thrown out by the constitution, but pure serum, secreted by the cutaneous exhalant arteries. So rare was this disease when Dr. Cullen wrote, that he never saw it but once, in a case which was shewn to him by Dr. Home. Dr. David Stuart, then physician to the hospital at Aberdeen, published an account of it in the Edinburgh Medical Commentaries.

The patient was a private soldier of the seventy-third regiment, aged eighteen, formerly a pedlar, and naturally of a healthy constitution. About twenty days before, he had been seized with the measles, when in the country; and, in marching to town, on the second day of their eruption, he was exposed to cold; upon which they suddenly disappeared. On his arrival at Aberdeen, he was quartered in a damp under-ground apartment. He then complained of sickness at stomach, great oppression about the præcordia, head-ache, lassitude and weariness on the least exertion, and stiffness and rigidity of his knees and other joints. He had been purged, but with little benefit. About ten days before, he observed, on the inside of his thighs, a number of very small, distinct, red spots, a little elevated above the surface of the skin, and much resembling the first appearance of the small-pox. This eruption gradually spread itself over his whole body, and the pustules continued every day to increase in size.

Upon being received into the hospital, he complained of head ache, sickness at stomach, oppression about the præcordia, thirst, sore throat, with difficulty of swallowing; his tongue was foul, his skin felt hot and feverish; pulse from 110 to 120, rather depressed; belly costive, eyes dull and languid, but without delirium. The whole surface of the skin was interspersed with vesicles, or phlyctenæ, of the size of an ordinary walnut; many of them were larger, especially on the arms and breast. In the interstices, between the vesicles, the appearance of the skin was natural, nor was there any redness round their base; the distance from one to another was from half an inch to a hand-breadth, or more. In some places two or three were joined together, like the pustules in the confluent small-pox. A few vesicles had burst of themselves, and formed a whitish scab, or crust. These were mostly on the neck and face; others shewed a tolerable laudable pus. However, by far the greatest number were perfectly entire, turgid, and of a bluish colour. Upon opening them, it was evident that the cuticle elevated above the cutis, and distended with a thin, yellowish, semi-pellucid serum, formed this appearance. Nor was the surface of the cutis ulcerated, or livid; but of a red florid colour, as when the cuticle is separated by a blister, or superficial burning. No other person laboured under a similar disease, either in the part of the country from which he came, or where he resided in Aberdeen.

Since the publication of this case of pemphigus, by Dr. Stuart, observations on this disease have been published by Dr. Dickson, of Dublin, by Mr. Gaitskell and

Mr. Upton, in the Mem. of the Medical Society of London. Some subsequent observations on pemphigus were published in the London Med. Journal, by Mr. Thomas Christie. From a case which Mr. Christie describes, he is disposed to agree with Dr. Dickson, in thinking that sometimes, at least, pemphigus is not contagious. He remarks, however, that the pemphigus described by some foreign writers was extremely infectious; circumstances which, he thinks, may lead to a division of the disease into two species, the pemphigus simplex and complicatus, both of which, but especially the last, seem to vary much with respect to mildness and malignity.

PEMPHIGUS MAJOR. A title under which pemphigus is spoken of by Sauvages, who defines it an eruption of phlyctenæ about the size of an hazel-nut, filled with a thin yellow serum.

PEMPHIGUS MINOR. In this species the vesicles are no larger than garden-peas.

PENPHIS. A species of *Lithrum*.

PENPTEUS. (From *πεντος*, the fifth.) An ague, the paroxysm of which returns every fifth day.

PENEA. A species of *Polygala*.

PENEA MUCRONATA. The systematic name of the plant which affords the sarcocolla. See *Sarcocolla*.

PENETRANTIA. (From *penetro*, to pierce through.) Medicines which pass through the pores and stimulate.

PENICILLIFORM. (*Penicilliformis*; from *penicillum*, a pencil, and *forma*, likeness.) Resembling a painter's pencil.

PENICILLUS. (Dim. of *peniculum*, a brush.) 1. A tent, or pledget.

2. The glandular or secreting extremities of the vena portæ. See *Liver*.

PENIDIUM. A kind of clarified sugar, with a mixture of starch, made up into small rolls. The confectioners call it barley-sugar.

PENIDIUM SACCHARATUM. See *Penidium*.

PENIS. (*A pendendo*, from its hanging down.) *Membrum virile*. The cylindrical part that hangs down, under the mons veneris, before the scrotum of males. It is divided by anatomists into the root, body, and head called the *glans penis*. It is composed of common integuments, two corpora cavernosa, and one corpus spongiosum, which surrounds a canal, the *urethra*, that proceeds from the bladder to the apex of the penis, where it opens by the *meatus urinarius*. See *Urethra*. The fold of the skin that covers the glans penis is termed the prepuce. The arteries of the penis are from the hypogastric and schiatic. The vein of the penis, *vena magna ipsius penis*, empties itself into

the hypogastric vein. The absorbents of this organ are very numerous, and run under the common integuments to the inguinal glands: absorbents also are found in great plenty in the urethra. The glands of the penis are, Cowper's glands, the prostate, muciparous, and odoriferous glands. The nerves of the penis are branches of the sacral and ischiatic.

PENIS CEREBRI. The pineal gland.

Penis, erector. See *Erector penis*.

PENIS MULIEBRIS. See *Clitoris*.

Pennyroyal. See *Pulegium*.

Pennyroyal, harts. See *Pulegium cervinum*.

PENTADACTYLON. (From *πεντε*, five, and *δακτυλος*, a finger; so called because it has five leaves upon each stalk, like the fingers upon the hand.) The herb cinquefoil; also a name for the ricinus, the fruit of which resembles a hand.

PENTANTHRUM. (From *πεντε*, five, and *ανθρον*, ointment.) An ointment composed of five ingredients.

PENTANURON. (From *πεντε*, five, and *νυρον*, a string; so called because it has five-ribbed leaves.) Ribwort.

PENTAPHARMACON. (From *πεντε*, five, and *φαρμακον*, remedium, remedy.) Any medicine consisting of five ingredients.

PENTAPHYLLOIDES. (From *πενταφυλλον*, cinquefoil, and *ειδος*, likeness; so called from its resemblance to cinquefoil.) Barren strawberry.

PENTAPHYLLUM. (From *πεντε*, five, and *φυλλον*, a leaf; so named because it has five leaves on each stalk.) Common cinquefoil, or five-leaved grass. The roots of this plant, *Potentilla reptans* of Linnæus: — *foliis quinatis, caule repente, pedunculis unifloris*, have a bitterish styptic taste. They were used by the antients in the cure of intermittents; but the medicinal quality of cinquefoil is confined, in the present day, to stop diarrhœas and other fluxes.

PENTAPLEURUM. The same as pentanuron.

PENTATOMON. (From *πεντε*, five, and *τομη*, to cut; so called because its leaves are divided into five segments.) Cinquefoil.

PENTOROBUS. (From *πεντε*, five, and *ροβος*, the wood-pea; so called because it has five seeds resembling the wood-pea.) The herb peony.

Peony, common. See *Pœonia*.

PEPANSIS. (From *πεπασθαι*, to concoct.)

Pepasmus. The maturation or concoction of humours.

PEPASNOS. The same as pepansis.

PEPASTICA. (From *πεπασθαι*, to concoct.) Digestive medicines.

PEPITA NUX. Ignatius's bean.

PEPLION. (From *πεπλος*, the herb

devil's-milk; so called from its resemblance.) *Peplos*. Wild parsley.

PERO. (From *πειρω*, to ripen.) See *Cucurbita*.

Pepper, black. See *Piper nigrum*.

Pepper, Guinea. See *Piper Indicum*.

Pepper, Jamaica. See *Pimenta*.

Pepper, long. See *Piper longum*.

Pepper, poorman's. See *Hydropiper*.

Pepper, wall. See *Illecebra*.

Pepper, water. See *Hydropiper*.

Pepper, white. See *Piper nigrum*.

Peppermint. See *Mentha piperitis*.

Peppermint. See *Lepidium*.

PARTICOS. (From *πειρω*, to ripen.) Such a thing as promotes digestion, or is digestive.

PERACUTE. Very sharp. Diseases are thus called when greatly inflamed, or aggravated beyond measure.

PARACEPIA. Paraleptici, or paraley-breakstone.

PERCOLATION. (*Percolatio*, straining through; from *per*, through, and *colo*, to strain.) It is generally applied to animal secretion, from the office of the glands resembling that of a strainer, in transmitting the liquors that pass through them.

PERDETUM. In Paracelsus it is the root of skirret.

Perennial worm-grass. See *Spigelia*.

PERETRIOR. (From *περαω*, to dig through.) The perforating part of the trepan.

PERDICIUM. (From *περδικη*, a partridge; so called because partridges were said to feed upon it.) Pellitory of the wall.

PERFOLIATA. (From *per* and *folium*, so called because the leaves surround the stem, like those of a cabbage.) Round-leaved hare's-ear, or therox wax. This plant, *Euphorbia rotundifolia* of Linnæus, was formerly celebrated for curing ruptures, mixed into a poultice with wine and oatmeal.

Perforans. See *Flexor profundus perforans*.

Perforans, seu flexor profundus. See *Flexor longus digitorum pedis profundus perforans*.

Perforans, seu flexor tertii internodii digitorum pedis. See *Flexor longus digitorum pedis profundus perforans*.

Perforans vulgo profundus. See *Flexor profundus perforans*.

PERFORATA. (From *perforo*, to pierce through; so called because its leaves are full of holes.) See *Hypericum*.

Perforatus. See *Flexor brevis digitorum pedis*, and *Flexor sublimis perforatus*.

Perforatus, seu flexor secundi internodii digitorum pedis. See *Flexor brevis digitorum pedis sublimis perforatus*.

PERIANNIA. (From *περιαννα*, to hang round.) An amulet, or charm, which was hung round the neck to prevent infection.

PERIBLEPSIS. (From *περίβλεψα*, to stare about.) That kind of wild look which is observed in delirious persons.

PERIBOLE. (From *περίβωλον*, to surround.) Sometimes it signifies the dress of a person; at others, a translation of the morbid humours to the surface of the body.

PERIBROSIS. An ulceration, or erosion, at the corners or uniting parts of the eyelids. This disorder most frequently affects the internal commissure of the eyelids. The species are 1. *Peribrosis*, from the acrimony of the tears, as may be observed in the epiphora. 2. *Peribrosis*, from an *xylops*, which sometimes extends to the commissure of the eyelids.

PERICARDITIS. (From *περικαρδίον*, the pericardium.) Inflammation of the pericardium. See *Carditis*.

PERICARDIUM. (From *περί*, about, and *καρδία*, the heart.) The membranous bag that surrounds the heart. Its use is to secrete and contain the vapour of the pericardium, which lubricates the heart, and thus preserves it from concreting with the pericardium.

PERICARPIA. (From *περί*, about, and *carpus*, the wrist.) Are medicines that are applied to the wrist.

PERICENMIA. (From *περί*, about, and *κνήμη*, the tibia.) The parts about the tibia.

PERICHONDRIUM. (From *περί*, about, and *χόνδρος*, a cartilage.) The membrane that covers a cartilage.

PERICHRISIS. (From *περί*, about, and *χρίω*, to anoint.) A liniment.

PERICHRISTA. (From *περί*, around, and *χρίω*, to anoint.) Any medicines with which the eyelids are anointed, in an ophthalmia.

PERICLASIS. (From *περί*, about, and *κλάω*, to break.) It is a term used by Galen for such a fracture of the bone as quite divides it, and forces it through the flesh into sight. Or a fracture with a great wound, wherein the bone is laid bare.

PERICLYNENUM. (From *περικλυζω*, to roll round; so called because it twists itself round whatever is near it.) The honey-suckle, or woodbine.

PERICRANIUM. (From *περί*, about, and *κράνιον*, the cranium.) The membrane that is closely connected to the bones of the head.

PERIDESMICA. (From *περί*, about, and *δεσμός*, a ligature.) Applied to an ischuria, or suppression of urine, from stricture in the urethra.

PERIDROMOS. (From *περί*, about, and *δρομος*, a course.) The extreme circumference of the hairs of the head.

PERIEGRIA. *Περίεργα*. Is any needless caution or trouble in an operation, as *περίεργος*, is one who despatches it with unnecessary circumstances: both the

terms are met with in Hippocrates, and others of the Greek writers.

PERISTROCHOS. (From *περιστρέω*, to surround, or to guard.) An epithet for diseases, signs, or symptoms, importing their being salutary, and that they prognosticate the recovery of the patient.

PERISGRAPHE. (From *περίγραφε*, to circumscribe.) An inaccurate description, or delineation. In Vesalius, *perigraphæ* signifies certain white lines and impressions, observable in the musculus rectus of the abdomen.

PERIN. (From *περα*, a bag.) A testicle. Some explain it the *Perineum*; others say it is the *Anus*.

PERINMOELLE. (From *περίνηλον*, the perineum, and *κλάω*, a rupture.) A rupture in the perineum.

PERINÆUM. (From *περίνηον*, to flow round, because that part is generally moist.) The space between the anus and organs of generation.

Perineus transversus. See *Transversus perinei*.

PERINICTIDES. (From *περί*, and *νύξ*, the night.) Little swellings like nipples; or, as others relate, pustules, or pimples, which break out in the night.

PERIOSTEUM. (From *περί*, about, and *ὀστέον*, a bone.) The membrane which invests the external surface of all the bones, except the crowns of the teeth. It is of a fibrous texture, and well supplied with arteries, veins, nerves, and absorbents. It is called *pericranium*, on the cranium; *periorbita*, on the orbits; *perichondrium*, when it covers cartilage; and *peridesmium*, when it covers ligament. Its use appears to be to distribute the vessels on the external surfaces of bones.

PERIPHIMOSIS. See *Phimosis*.

PERIPNEUMONIA. See *Pneumonia*.

PERIPNEUMONIA. (From *περί*, and *πνεύμων*, the lung.) Peripneumony, or inflammation of the lungs. See *Pneumonia*.

PERIPNEUMONIA NOTHA. Bastard or spurious peripneumony. Practitioners, it would appear, do not all affix this name to the same disease; some affirming it to be a rheumatic affection of the respiratory muscles, while others consider it as a mild peripneumony. It is characterised by difficulty of breathing, great oppression at the chest, with obscure pains, coughs, and occasionally an expectoration. Spurious peripneumony is sometimes so slight as to resemble only a violent catarrh; and, after the employment of a few proper remedies, goes off by a free and copious expectoration; but sometimes the symptoms run high, and an effusion of serum into the bronchiz takes place, which destroys the patient.

PERIPHEMA. (From *περί*, about, and *ἔμωρ*, pus.) Is a collection of matter about any part, as round a tooth in the gums:

PERIPLASIS. (From *περι*, about, *πλάσσειν*, to break.) A breaking off, or a separation round about, either of corrupted bones or of dead flesh.

PERIPLUGA. (From *περιπλύνω*, to flow about.) Is a reflux of humours from the habit of the body into any of the larger emunctories for its excretion, as in an hydropsical case, of water upon the bowels or kidneys, where it passes away by urine, or stool.

PERIPTHEMUS. (From *περι*, about, and *κνυφος*, gibbous.) An incision made across the forehead, or from one temple to another, over the upper part of the os frontis, over the coronary suture. It was formerly used when a considerable inflammation or defluxion in the eyes attended.

PERISTALTIC MOTION. (*Peristalticus*; from *περιστάλλω*, to contract.) The vermicular motion of the intestines, by which they contract and propel their contents. A similar motion takes place in the Fallopian tubes, after conception, by means of which the ovum is translated from the ovarium into the uterus.

PERISTAPHYLINUS. (From *περι*, about, and *σταφυλινος*, the staphylinus.) A muscle which is connected with the staphylinus.

PERISTERIUM. (From *περιστήρ*, a pigeon; so called because pigeons cover it.) The herb vervain. See *Verbena*.

PERISTOMA. (From *περιστήνω*, to strew about.) *Peristoma* properly signifies any covering, but is applied, by Pecquet, to the mucous or villous coat or lining of the intestines, the same which Blasius calls *Muscum Villosum*; Bartholine, *Crusta Membranosa*; and De Graaf, *Crusta Vermicularis*.

PERISTOTILE. (From *περιστάλλω*, to compress.) 1. The time between a contraction and dilatation of the heart.

2. A pause, or intermission between the systole and diastole, which is by most denied to be perceived in healthy persons, but when dying it is very sensibly felt.

PERITERION. (From *περι*, and *τρώγω*, to preserve.) The perforating part of the trepan.

PERITONORIXIS. (From *περιτοναϊος*, the peritonæum, and *γίγνομαι*, to break.) A bursting of the peritonæum, and consequent hernia.

PERITONEUM. (From *περιτύνω*, to extend round.) A strong simple membrane, by which all the viscera of the abdomen are surrounded. It has an exceedingly smooth, exhaling and moist internal surface. Outwardly, it is every where surrounded by cellular substance, which, towards the kidneys, is very loose and very fat; but is very short at the lower tendon of the transverse muscles. It begins from the diaphragm, which it completely lines; and, at the last fleshy fibres of the ribs,

and the external lumbar fibres, it completes the septum, in conjunction with the pleura, with which it is continuous through the various intervals of the diaphragm. Posteriorly, it descends before the kidneys; anteriorly, behind the abdominal muscles; it dips into the pelvis; from the bones of the pubis, passes over the bladder, and descends behind; and being again carried backwards, at the entrance of the ureters in two lunar folds, it rejoins upon the intestinum rectum, that part of itself which invests the loins, and in this situation lies before the rectum. The cellular texture which covers the peritonæum on the outside, is continued into sheaths in very many places; of which, one receives the testicle on each side, another the iliac vessels of the pelvis, viz. the obturatoria, those of the penis, bladder, and aorta, and, ascending to the breast, accompany the œsophagus and vertebræ; by means of which, there is a communication between the whole body and the peritonæum, well known in dropsical people. It has various prolongations for covering the viscera. The shorter productions of this membrane are called ligaments; and are formed by a continuous reduplication of the peritonæum, receding from its inner surface, enclosing cellular substance, and extending to some viscus, where its plates separate, and, having diverged, embrace the viscus; but the intermediate cellular substance always accompanies this membranaceous coat, and joins it with the true substance of the viscus. Of this short kind of production, three belong to the liver, one or two to the spleen, and others to the kidneys, and to the sides of the uterus and vagina. By this means, the tender substance of the viscera is defended from injury by any motion or concussion, and their whole mass is prevented from being misplaced by their own weight, and from injuring themselves, being securely connected with the firm sides of the peritonæum.

PERITONITIS. (From *περιτοναϊος*, the peritonæum.) An inflammation of the peritonæum. A genus of disease in the Class *Pyrexie* and Order *Phlegmasie* of Cullen, known by the presence of pyrexia, with pain in the abdomen, that is increased when in an erect position, but without other proper signs of abdominal inflammation. When the inflammation attacks the peritoneum of the viscera, it takes the name of the viscus; thus, *peritonitis hepatis*, *peritonitis intestinalis*, *peritonitis omentalis*, or *epiploitis*, or *omentitis*, *peritonitis mesenterii*.

All these Dr. Cullen considers under the general head of gastritis, as there are no certain signs by which they can be distinguished from each other, and the method of cure must be the same in all. He however distinguishes three species.

1. *Peritonitis propria*; when the peritoneum, strictly so called, is inflamed.

2. *Peritonitis omentalis. Omentitis. Epiplöitis*, when the omentum is affected.

3. *Peritonitis mesenterica*, when the mesentery is inflamed.

PERIZOMA. (From *περιζώνωμι*, to gird round.) Strictly signifies a girdle; but by Hildanus, and some other chyrurgical writers, it is applied to such instruments for supporting ruptures, which we commonly call trusses. Some also express by it the diaphragm.

PERLA. (Ital. and Span. *perl*, Welch, *perlen*, Germ.) See *Margarita*.

Perls. See *Margarita*.

PERNIO. A kibe or chilblain. A species of *erythema* of Cullen. Chilblains are painful inflammatory swellings, of a deep purple or leaden colour, to which the fingers, toes, heels, and other extreme parts of the body are subject on being exposed to a severe degree of cold. The pain is not constant, but rather pungent and shooting at particular times, and an insupportable itching attends. In some instances the skin remains entire, but in others it breaks and discharges a thin fluid. When the degree of cold has been very great, or the application long continued, the parts affected are apt to mortify and slough off, leaving a foul ill-conditioned ulcer behind. Children and old people are more apt to be troubled with chilblains than those of a middle age; and such as are of a scrophulous habit, are remarked to suffer severely from them.

Peroneus anticus. See *Peroneus brevis*.

PERONEUS BREVIS. (*Peroneus*, sc. *musculus*, *περωναίος*, from *perone*, the fibula.) This muscle is the *peroneus secundus* seu *anticus* of Douglas, the *peroneus medius* seu *anticus* of Winslow, the *peroneus secundus* of Cowper, and *petit-peroneo-sus-metatarsien* of Dumas. It arises by an acute, thin, and fleshy origin from the anterior and outer part of the fibula, its fibres continuing to adhere to the lower half of that bone. Its round tendon passes through the groove in the malleolus externus, along with that of the *peroneus longus*, after which it runs in a separate groove to be inserted into the upper and posterior part of the tubercle at the basis of the metatarsal bone, that supports the little toe. Its use is to assist the *peroneus longus*.

PERONEUS LONGUS. This muscle, which is the *peroneus primus* seu *posticus* of Douglas, *peroneus maximus* seu *posterior* of Winslow, *peroneus primus* of Cowper, and *tebi-peroneo-tarsien* of Dumas, is situated somewhat anteriorly along the outer side of the leg. It arises tendinous and fleshy from the external lateral part of the head of the tibia, and likewise from the upper

anterior surface and outer side of the *perone* or fibula, its fibres continuing to adhere to the outer surface of the latter to within three or four inches of the malleolus externus. It terminates in a long round tendon, which runs obliquely behind the malleolus internus, where it passes through a cartilaginous groove in common with the *peroneus brevis*, being bound down by an annular ligament. When it has reached the os calcis, it quits the tendon of the *peroneus brevis*, and runs obliquely inwards along a groove in the os cuboidei, under the muscles on the sole of the foot, to be inserted into the outside of the posterior extremity of the metatarsal bone, that supports the great toe. Near the insertion of this muscle we find a small *bursa mucosa*. This muscle draws the foot outwards, and likewise assists in extending it.

Peroneus maximus. See *Peroneus longus*.

Peroneus medius. See *Peroneus brevis*.

Peroneus posticus. See *Peroneus longus*.

Peroneus primus. See *Peroneus longus*.

Peroneus secundus. See *Peroneus brevis*.

PERONEUS TERTIUS. This is the name given by Albinus to a muscle which, by some writers, is called *nonus Vesalii*, or Vesalius's ninth muscle of the foot; but by most considered in the present day as a portion of the *extensor longus digitorum pedis*. It is situated at the anterior, inferior, and outer part of the leg, along the outer edge of the last-described muscle, to which it is intimately united. It arises fleshy from the anterior surface of the lower half of the fibula, and from the adjacent part of the interosseous ligament. Its fibres run obliquely downwards, towards a tendon which passes under the annular ligament, and then running obliquely outwards, is inserted into the root of the metatarsal bone that supports the little toe. This muscle assists in bending the foot.

PERONE. (From *πύγῃ*, to fasten; so called because it fastens together the tibia and the muscles.) The fibula.

PERSICA. (From *Persia*, its native soil.) The peach. The fruit of the *Amygdalus Persica* of Linnæus. It is known to be grateful and wholesome, seldom disagreeing with the stomach, unless this organ is not in a healthy state, or the fruit has been eaten to excess, when effects similar to those of the other dulco-acid summer fruits may be produced. The flowers, including the calyx, as well as the corolla, are the parts of the persica used for medicinal purposes. These have an agreeable but weak smell, and a bitterish taste. Boileuc observes, "that when distilled, without addition, by the heat of a water bath, they yield one-sixth their weight, or more, of a whitish liquid, which communicates to a considerable quantity of other liquids a flavour

like that of the kernels of fruits. These flowers have a cathartic effect, and, especially to children, have been successfully given in the character of a vermifuge; for this purpose, an infusion of a drachm of the flowers dried, or half an ounce in their recent state, is the requisite dose. The leaves of the persica are also found to possess an anthelmintic power, and from a great number of experiments appear to have been given with invariable success both to children and adults. However, as the leaves and flowers of the persica manifest, in some degree, the quality of those of the laurocerasis, they ought to be used with caution.

PERSICARIA. (From *persica*, the peach-tree, so called because its blossoms are like those of the peach.) *Persicaria mitis*. *Plumbago*. Arsmart. This plant, *Polygonum persicaria* of Linnaeus, is said to possess vulnerary and antiseptic properties; with which intentions it is given in wine to restrain the progress of gangrene.

PERSICARIA MITIS. See *Persicaria*.

PERSICARIA URENS. See *Hydropiper*.

PERSICUS IGNIS. A carbuncle. Avicenna says, it is that species of carbuncle which is attended with pustules and vesications.

PERSISTENS FEBRIS. A regular intermitting fever, the paroxysms of which return at constant and stated hours.

PERSONATA. (From *persona*, a disguised person, because, according to Pliny, the ancient actors used to mask themselves with the leaves of this plant.) See *Bardana*.

PERSPIRATION. *Perspiratio*. The vapour that is secreted by the extremities of the cutaneous arteries from the external surface of the body. It is distinguished into *sensible* and *insensible*. The former is separated in the form of an invisible vapour, the latter so as to be visible in the form of very little drops adhering to the epidermis. The *secretory organ* is composed of the extremities of the cutaneous arteries. The *smell* of the perspirable fluid, in an healthy man, is fatuous and animal; its *taste* manifestly salt and ammoniacal. In *consistence* it is vaporous and aqueous; and its *specific gravity* is greater than water. For the most part it is yellowish, from the passage of the subcutaneous oil, and sebaceous matter of the subcutaneous glands. Sometimes it is reddish, from the globules of the cruror passing through, especially under the axillæ. The *quantity* is sometimes so profuse, as not only conspicuously to moisten the linen, but also the thicker garments.

The *constituent principles* of the perspirable fluid appear to be, 1. *Water*, attenuated into vapour, by the matter of heat 2. *Animal gas*, or carbonated hydrogen. As the production of carbonated

air with the oxygen of the atmosphere shews. 3. *Azotic gas*. For water, in which a man has bathed, soon becomes putrid. Carbonated hydrogen, chemically combined with azot, would appear to constitute *putrid miasma*. May not this be the origin of putrid fever, in those narrow confined chambers in which there are many persons? 4. The *glandular emagma* and *subcutaneous oil*; hence linen is stained with a yellowish colour, and leanness is brought on. 5. The *serum of the blood*. This affords an immense quantity of water, and the albuminous and saline part of the sweat. It makes the linen of a viscid rigidity, and of a salt taste. Glass-blowers sometimes excrete so acid a sweat, that salt has been seen collected in crystals on their faces.

Perspiration varies in respect to, 1. The *temperature of the atmosphere*. Thus men have a more copious, viscid, and higher coloured sweat on the summer days, and in warm countries, than in colder regions. 2. *Sex*. The sweat of a man is said to smell more acrid than that of a woman. 3. *Age*. The young are more subject to sweat than the aged, who during the excessive heat of the summer scarcely sweat at all. 4. *Ingesta*. An alliacious sweat is perceived from eating garlic; a leguminous from peas; an acid from acids; a fetid from animal food only; and a rancid sweat from fat foods, as is observed in Greenland. A long abstinence from drink causes a more acrid and coloured sweat; and the drinking a great quantity of cold water in summer, a limpid and thin sweat. 5. *Medicines*. The sweat of those who have taken musk, even moderately, and assafœtida, or sulphur, smells of their respective natures. 6. *Region of the body*. The sweat of the head is greasy; on the forehead it is more aqueous; under the axillæ very unguinous; and in the interstices of the toes, it is very fetid, forming in the most healthy man blackish sordes. 7. *Diseases*. In this respect it varies very much, in regard to quantity, smell, and colour; for the sweat of gouty persons is said to turn vegetable juices to a red colour, and is of a cretaceous nature. Some men also have a lucid sweat, others a sweat tinging their linen of a cerulean colour.

The use of the insensible perspiration are, 1. To *liberate* the blood from superfluous animal gas, azot, and water. 2. To eliminate the noxious and heterogeneous excrements; hence the acid, rancid, leguminous, or putrid perspiration of some men. 3. To *moisten* the external surface of the body, lest the epidermis cutis, and its nervous papillæ be dried up by the atmospheric air. 4. To *counter-balance* the suppressed pulmonary transpiration of the lungs; for when it is suppressed, the cu-

taneous is increased; hence the nature of both appears to be the same.

The use of the sensible perspiration, or sweat, in an healthy man, is scarcely observable, unless from an error of the non-naturals. Its first effect on the body is always prejudicial, by exhausting and drying it; although it is sometimes of advantage. 1. By supplying a watery excretion: thus when the urine is deficient, the sweat is often more abundant. In this manner an aqueous diarrhoea is frequently cured by sweating. 2. By eliminating, at the same time, any morbid matter. Thus various miasmata are critically expelled, in acute and chronic diseases, with the sweat.

PERTUSSIS. (From *per*, much, and *tussis*, cough.) The hooping-cough. A genus of disease in the class *neuroses*, and order *spasmi* of Cullen, known by a convulsive strangulating cough, with hooping, returning by fits that are usually terminated by a vomiting, and being contagious.

Children are most commonly the subjects of this disease, and it seems to depend on a specific contagion, which affects them but once in their life. The disease being once produced, the fits of coughing are often repeated without any evident cause; but in many cases, the contagion may be considered as only giving the predisposition, and the frequency of the fits may depend upon various exciting causes, such as violent exercise, a full meal, the having taken food of difficult digestion, and irritation of the lungs by dust, smoke, or disagreeable odours. Emotions of the mind may likewise prove an exciting cause.

Its proximate or immediate cause seems to be a viscid matter or phlegm lodged about the bronchiz, trachea, and fauces, which sticks so close as to be expectorated with the greatest difficulty. Some have supposed it to be a morbid irritability of the stomach, with increased actions of its mucous glands; but the affection of the stomach which takes place in the disease, is clearly only of a secondary nature, so that this opinion must be erroneous.

The hooping-cough usually comes on with a difficulty of breathing, some degree of thirst, a quick pulse, and other slight febrile symptoms, which are succeeded by a hoarseness, cough, and difficulty of expectoration. These symptoms continue perhaps for a fortnight or more, at the end of which time the disease puts on its peculiar and characteristic form, and is now evident, as the cough becomes convulsive, and is attended with a peculiar sound, which has been called a hoop.

When the sonorous inspiration has happened, the conghing is again renewed, and continues in the same manner as before, till either a quantity of mucus is thrown

up from the lungs, or the contents of the stomach are evacuated by vomiting. The fit is then terminated, and the patient remains free from any other for some time, and shortly afterwards returns to the amusements he was employed in before the accession of the fit, expresses a desire for food, and when it is given to him, takes it greedily. In those cases, however, where the attack has been severe, he often seems much fatigued, makes quick inspirations, and falls into a faint.

On the first coming on of the disease, there is little or no expectoration, or if any, it consists only of thin mucus, and as long as this is the case the fits of coughing are frequent, and of considerable duration; but on the expectoration becoming free and copious, the fits of coughing are less frequent, as well as of shorter duration.

By the violence of coughing, the free transmission of blood through the lungs is somewhat interrupted, as likewise the free return of the blood from the head, which produces that turgescence and suffusion of the face which commonly attends the attack, and in some instances brings on a hæmorrhage either from the nose or ears.

The disease having arrived at its height, usually continues for some weeks longer, and at length goes off gradually. In some cases it is however protracted for several months, or even a year.

Although the hooping-cough often proves tedious, and is liable to return with violence on any fresh exposure to cold, when not entirely removed, it nevertheless is seldom fatal, except to very young children, who are always likely to suffer more from it than those of a more advanced age. The danger seems indeed always to be in proportion to the youth of the person, and the degree of fever, and difficulty of breathing, which accompany the disease, as likewise the state of debility which prevails.

It has been known in some instances to terminate in apoplexy and suffocation. If the fits are put an end to by vomiting, it may be regarded as a favourable symptom, as may likewise the taking place of a moderate and free expectoration, or the ensuing of a slight hæmorrhage from the nose or ears.

Dissections of those who die of the hooping-cough usually shew the consequence of the organs of respiration being affected, and particularly those parts which are the seat of catarrh. When the disease has been long protracted, it is apt to degenerate into pulmonary consumption, asthma, or visceral obstructions, in which last case the glands of the mesentery are found in a hard and enlarged state.

Peruvian balsam. See *Balsamum Peruvianum*.

Peruvian bark. See *Cinchona*.

PERUVIANUS CORTEX. See *Cinchona*.

PERUVIANUS CORTEX FLAVUS. See *Cinchona*.

PERUVIANUS CORTEX RUBER. See *Cinchona*.

PERVIGILIUM. (From *per*, much, and *vigilo*, to watch.) Watching, or a want of sleep. See *Vigilance*.

PERVINCA. (From *pervincio*, to tie together) So called because its stringy roots were used for binding substances together. The herb periwinkle.

PES ALEXANDRINUS. See *Pyrethrum*.

PES CAPRÆ. Goat's foot, a species of *Oxalis*; also a species of *Convulvulus*.

PES CATI. See *Gnaphalium*.

PES COLUMBINUS. See *Geranium columbinum*.

PES LEONIS. The ladies mantle is sometimes so called. See *Alchemilla*.

PES TIGRIDIS. Tiger's foot, a species of *Ipomœa*.

PESSARY. (*Pessarum*, from *press*, to soften.) An instrument that is introduced into the vagina to support the uterus.

Pestilent wort. See *Petasites*.

PESTIS. The plague. A genus of disease in the class *pyrexia* and order *exanthemata* of Cullen, characterised by typhus, which is contagious in the extreme, prostration of strength, buboes, and carbuncles, petechiz, hæmorrhage, and colliquative diarrhœa.

By some writers the disease has been divided into three species; that attended with buboes; that attended with carbuncles; and that accompanied with petechiz. This division appears wholly superfluous. Dr. Russel, in his elaborate treatise on the plague, makes mention of many varieties; but when these have arisen, they seem to have depended in a great measure on the temperament and constitution of the air at the time the disease became epidemical, as likewise on the patient's habit of body at the time of his being attacked with it.

The plague is by most writers considered as the consequence of a pestilential contagion, which is propagated from one person to another by association, or by coming near infected materials.

It has been observed that it generally appears as early as the fourth or fifth day after infection: but it has not yet been ascertained how long a person who has laboured under the disease is capable of infecting others, nor how long the contagion may lurk in an unfavourable habit without producing the disease, and may yet be communicated, and the disease excited, in habits more susceptible of the infection. It has generally been supposed, however, that a quarantine of 40 days is much longer than is necessary for persons, and probably for goods also. Experience has not yet determined how much of this term may be abated. "If I am not much

mistaken," observes Dr. Thomas, "the Board of Trade has, however, very lately, under the sanction of the College of Physicians, somewhat abridged it."

It sometimes happens that after the application of the putrid vapour, the patient experiences only a considerable degree of languor and slight head-ache for many days previous to a perfect attack of the disease: but it more usually comes to pass, that he is very soon seized with great depression of strength, anxiety, palpitations, syncope, stupor, giddiness, violent head-ache, and delirium, the pulse becoming at the same time very weak and irregular.

These symptoms are shortly succeeded by nausea, and a vomiting of a dark bilious matter, and in the further progress of the disease, carbuncles make their appearance; buboes arise in different glands, such as the parotid, maxillary, cervical, axillary, and inguinal; or petechiz hæmorrhages and a colliquative diarrhœa ensue, which denote a putrid tendency prevailing to a great degree in the mass of the blood.

Such are the characteristic symptoms of this malignant disease, but it seldom happens that they are all to be met with in the same person. Some, in the advanced state of the disease, labour under buboes, others under carbuncles, and others again are covered with petechiz.

The plague is always to be considered as attended with imminent danger, and when it prevailed in this country about 200 years ago, proved fatal to the most of those who were attacked with it. It is probable, however, that many of them died from want of care and proper nourishment, as the infected were forsaken by their nearest friends; because in Turkey and other countries, where attention is paid to the sick, a great many recover.

When the disease is unattended by buboes, it runs its course more rapidly, and is more generally fatal, than when accompanied by such inflammations. The earlier they appear, the milder usually is the disease. When they proceed kindly to suppuration they always prove critical, and ensure the patient's recovery. A gentle diaphoresis, arising spontaneously, has been known in many instances likewise to prove critical. When carbuncles shew a disposition to become gangrenous, the event will be fatal. Petechiz, hæmorrhages and colliquative diarrhœa denote the same termination.

Dissections of the plague have discovered the gall-bladder full of black bile, the liver very considerably enlarged, the heart much increased in size, and the lungs, kidneys, and intestines beset with carbuncles. They have likewise discovered all the other appearances of putrid fever.

PETALODES. (From *πέταλον*, a leaf or

thin scale.) Is by Hippocrates applied to an urine which hath in it flaky substances resembling leaves.

PETASITES. (From *πῆταςος*, a hat, so named because its leaves are shaped like a hat.) Butterbur. Pestilentwort. *Tussilago petasites* of Linnæus. The roots of this plant are recommended as aperient and alexipharmic, and promise, though now forgotten, to be of considerable activity. They have a strong smell, and a bitterish acrid taste, of the aromatic kind, but not agreeable.

PETECULA. (From the Italian *petecchio*, a flea-bite, because they resemble the bites of fleas.) A red or purple spot that mostly appears in contagious diseases, and resembles flea-bites.

PETRAPIUM. (From *petra*, a rock, and *apium*, parsley, so called because it grows in stony places.) See *Petroselinum Macedonicum*.

PETRELEUM. (From *petra*, a rock, and *ελαιον*, oil.) An oil or liquid bitumen which distils from rocks.

PETROLEUM. (From *petra*, a rock, and *oleum*, oil.) The name of petroleum is given to a liquid bituminous substance which flows between rocks, or in different places at the surface of the earth. The more fluid species are distinguished by the name of *naphtha*, and the thicker by those of *piscephaltum* and *pisceillum*. See *Naphtha*, *Mineral pitch*, &c.

PETROLEUM BARBADENSE. Barbadoes tar. This is chiefly obtained from the island of Barbadoes, and is sometimes employed externally in paralytic diseases.

PETROLEUM RUBRUM. *Oleum gabianum*. Red petroleum. A species of rock-oil of a blackish red colour, of thicker consistency, less penetrating and more disagreeable smell than the other kinds of petroleum. It abounds about the village of Gabian in Languedoc.

PETROLEUM SULPHURETUM. A stimulating balsamic remedy given in coughs, asthmas, and other affections of the chest.

Petropharyngeus. A muscle which arises in the apophysis petrosa, and is inserted into the pharynx.

Petro-salpinge staphylinus. See *Levator palati*.

PETROSELINUM. (From *petra*, a rock, and *σαλιον*, parsley.) *Petroselinum vulgare*. *Apium hortense*. Coramon parsley. *Apium petroselinum* of Linnæus:—*foliis caulibus linearibus, involucrellis minutis*. Both the roots and seeds of this plant are directed by the London College for medicinal use: the former have a sweetish taste, accompanied with a slight warmth or flavour, somewhat resembling that of carrot; the latter are in taste warmer and more aromatic than any other part of the plant, and manifest considerable bitterness. The roots are said to be aperient

and diuretic, and have been employed in nephritic pains and obstructions of urine. The seeds possess aromatic and carminative powers, but are seldom prescribed.

PETROSELINUM MACEDONICUM. *Apium petraeum*. *Petrapium*. Macedonian parsley. This plant, *Bubon Macedonicum* of Linnæus, is similar in quality to the common parsley, but weaker and less grateful. The seeds enter the celebrated compounds mithridate and theriaca.

PETROSELINUM VULGARE. See *Petroselinum*.

PETROSILEX. A species of coarse flint, of a deep blue or yellowish green colour. It is interspersed in veins through rocks; and from this circumstance derives its name.

PEUCEDANUM. (From *πυκνὴ*, the pine-tree; so called from its leaves resembling those of the pine-tree.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*. 2. The pharmacopœial name of the hog's fennel and sulphur-wort. *Marathrum sylvestre*. *Marathrophyllum*. *Pinnastellum*. *Hieracleum ptericinum*. The plant which bears these names in the pharmacopœias is the *Peucedanum officinale* of Linnæus:—*foliis quinque parvis, filiformibus linearibus*. The root is the officinal part; it has a strong fetid smell, somewhat resembling that of sulphureous solutions, and an acrid, unctuous, bitterish taste. Wounded, when fresh in the spring or autumn, particularly in the former season, in which the root is most vigorous, it yields a considerable quantity of yellow juice, which soon dries into a solid gummy resin, which retains the taste and strong smell of the root. This, as well as the root, is recommended as a nervine and anti-hysterical remedy.

PEUCEDANUM OFFICINALE. The systematic name of the hog's fennel. See *Peucedanum*.

PEUCEDANUM SILAUS. The systematic name of the meadow saxafrage. See *Saxifraga vulgaris*.

PEYERI GLANDULÆ. Peyer's glands. Brunner's glands. Small glands situated under the villous coat of the intestines.

PEZIZA AURICULA. See *Auricula jude*.

PHENOMENA. *Phenomenon*. (From *φαῖναι*, to make appear.) All those appearances in the human body which are contrary to the usual process of nature.

PHAGEDÆNA. (From *φαγεῖν*, to eat.) A species of ulcer that spreads very rapidly.

PHAGEDENICS. (*Phagedenica*, sc. *medicamenta*; from *φαγεῖν*, to eat.) Applications that destroy fungous flesh.

PHALACRUM. (From *φαλακρος*, bald.) A surgical instrument, with a blunt smooth top; as a probe.

PHALANX. See *Phalanx*.

PHALANXIS. (From φαλαγξ, a row of soldiers.) 1. An affection of the eye-lids, where there are two or more rows of hairs upon them.

2. A morbid inversion of the eyelids.

PHALANX. (From φαλαγξ, a battalion.) The small bones of the fingers and toes, which are distinguished into the first, second, and third phalanx.

PHALARIS CANARIENSIS. (From φαλος, white, shining; so named from its white, shining seed, and canariensis, from its being the principal food of the canary-birds.) Canary-grass. The seed of this plant is well known to be the common food of canary-birds. In the Canary islands, the inhabitants grind it into meal, and make a coarse sort of bread with it.

PHALLUS ESCULENTUS. The systematic name of the morel fungus. See *Morel*.

PHANTASMA. (From φανταζω, to make appear.) Imagination Depraved vision.

PHARIS. (From *Pharos*, the island from whence it was brought.) A violent kind of poison.

PHARMACEUTICA. (From φαρμακω, to exhibit medicines.) Pharmacuetica, or the doctrine of compounding and dosing medicines. See *Pharmacy*.

PHARMACODYNIA. (From φαρμακον, a medicine, and δυναμις, chemistry.) Pharmacentic chemistry, or that part of chemistry which respects the preparation of medicines.

PHARMACOPŒIA. (From φαρμακον, a medicine, and ποιω, to make.) A dispensatory, or book of directions for the composition of medicines approved of by medical practitioners, or published by authority. The following are the most noted, viz.

P. Amstelodamensis.

P. Argentoratensis.

P. Augeratanensis.

P. Bateana.

P. Brandenburgensis.

P. Brandenburgica.

P. Bruzellensis.

P. Edinburgensis.

P. Hafniensis.

P. Londinensis.

P. Norimbergensis.

P. Parisiensis.

P. Ratibonensis.

P. Regia.

PHARMACOPOLA. (From φαρμακον, a medicine, and πωλω, to sell.) An apothecary, or vender of medicines.

PHARMACOPOLIUM. (From φαρμακον, a medicine, and πωλω, to sell.) A druggist's or apothecary's shop.

PHARMACOPSIA. (From φαρμακον, a medicine, and πωσις, a potion.) A liquid medicine.

PHARMACOTHECA. (From φαρμακον, a medicine, and τιθεμι, to place.) A medicine-chest.

PHARMACY. The art of preparing remedies for the treatment of diseases.

The articles of the *Materia Medica*, being generally unfit for administration in their original state, are subjected to various operations, mechanical or chemical, by which they become adapted to this purpose. Herein consists the practice of pharmacy, which therefore requires a previous knowledge of the sensible and chemical properties of the substances operated on. The qualities of many bodies are materially changed by heat, especially in conjunction with air and other chemical agents; the virtues of others reside chiefly in certain parts, which may be separated by the action of various menstrua, particularly with the assistance of heat; and the joint operation of remedies on the human body is often very different from what would be anticipated, from that which they exert separately; hence, in the preparations and compositions of the *Pharmacopœias*, we are furnished with many powerful as well as elegant forms of medicine.

PHARYNGEUS. (From φάρυγξ, the pharynx.) Belonging to or affecting the pharynx; thus cynanche pharyngea, &c.

PHARYNGEUS. φάρυγξ. The pharynx, or fauces.

PHARYNGOSTRAPHIUS. A muscle originating in the pharynx and terminating in the septum, above the uvula.

PHARYNGOTOMIA. (From φάρυγξ, the pharynx, and τομή, to cut.) The operation of cutting the pharynx.

PHARYNX (ἄρσενος, because it conveys the food into the stomach.) The muscular bag at the back part of the mouth. It is shaped like a funnel, adheres to the fauces behind the larynx, and terminates in the œsophagus. Its use is to receive the masticated food, and to convey it into the œsophagus.

PHASEOLUS VULGARIS. (From φασολος, a little ship, or galliot, which its pods were supposed to resemble.) The systematic name of the kidney-bean. See *Bean*, kidney.

PHASANTUM. (From φασγανον, a knife; so called because its leaves are shaped like a knife, or sword.) The herb sword-grass.

PHATNUM. (From φάτην, a stall.) The socket of a tooth.

PHELLANDRIUM AQUATICUM. (*Phellandrium*; from φελλος, the cork-tree, and ὕδωρ, water; so called because it floats upon the water like cork.) The systematic name of the water-fennel. See *Penniculus aquaticum*.

PHENOS. (From φεμεν, to shut up.) A medicine against a dysentery.

PHILABEUS. (From φιλω, to love, and ὄλεος, a brother; so called because, by its roughness, it attaches itself to whatever is near it.) See *Asperine*.

PHILASTRUM. (From *φιλεω*, to love, and *αἰδωτος*, a man; so called from its uses.) 1. Medicines relieving the pain of the stone.

2. The herb goose-grass, because it sticks to the garments of those who touch it. See *Aperine*.

PHILOXUM. (From *Philo*, its inventor.) A warm opiate.

PHILTRUM. (From *φιλεω*, to love.) 1. A philtre, or imaginary medicine, to excite love.

2. The depresso on the upper lip. where lovers salute.

PHILTIA. (The name of the daughter of Chiron, who first applied it medicinally.) Mock privet.

PHIMOSIS. (From *φιμεω*, to bind up.) A constriction or straitness of the extremity of the prepuce, which, preventing the glans from being uncovered, is often the occasion of many troublesome complaints. It may arise from different causes both in children and grown persons. Children have naturally the prepuce very long; and as it exceeds the extremity of the glans, and is not liable to be distended, it is apt to contract its orifice. This often occasions a lodgment of a small quantity of urine between that and the glans, which if it grows corrosive, may irritate the parts so as to produce an inflammation. In this case, the extremity of the prepuce becomes more contracted, and consequently the urine more confined. Hence the whole inside of the prepuce excoriates and suppurates; the end of it grows thick and swells, and in some months becomes callous. At other times it does not grow thick, but becomes so strait and contracted as hardly to allow the introduction of a probe. The only way to remove this disorder is by an operation. A phimosis may affect grown persons from the same cause as little children; though there are some grown persons who cannot uncover their glans, or at least not without pain, and yet have not the extremity of the prepuce so contracted as to confine the urine from passing, we notwithstanding find them sometimes troubled with a phimosis, which might be suspected to arise from a venereal taint, but has in reality, a much more innocent cause. There are, we know, sebaceous glands situated in the prepuce, round the corona, which secrete an unctuous humour, which sometimes becomes acrimonious, irritates the skin that covers the glans, and the irritation extending to the internal membrane of the prepuce, they both become inflamed, and yield a purulent serum, which cannot be discharged, because the glans is swelled, and the orifice of the prepuce contracted. We find also some grown persons, who, though they never uncovered the glans, have been subject to phimosis from a venereal cause.

In some it is owing to gonorrhoea, where the matter lodged between the prepuce and the glans occasioned the same excoriation as the discharge before mentioned from the sebaceous glands. In others, it proceeds from venereal chancres on the prepuce, the glans, or the frænum; which producing an inflammation either on the prepuce or glans, or both, the extremity of the fore-skin contracts, and prevents the discharge of the matter. The parts, in a very little time, are greatly tumefied, and some times a gangrene comes on in less than two days.

PHLEBORRHAGIA. (From *φλεβ*, a vein, and *ρρηνμι*, to break out.) A rupture of a vein.

PHLEBOTOMY. (*Phlebotomia*; from *φλεβ*, a vein, and *τομω*, to cut.) The opening of a vein.

PHLEGM. (From *φλεγω*, to burn, or to excite.) In chemistry it means water from distillation, but, in the common acceptance of the word, it is a thick and tenacious mucus secreted in the lungs.

PHLEGMASIA. (From *φλεγμα*, phlegm, and *εγω*, to drive out.) Medicines which promote the discharge of phlegm.

PHLEGMASIA. (From *φλεγμα*, to burn.) An inflammation.

PHLEGMASIA DOLENS. A disease noticed by some of the French writers, under the name of *L'enflure des jambes et des cuisses de la femme accouchée*; whilst others have called it *dépôt du lait*, from its supposed cause. By the Germans it is called *Edema lacteum*, and by the English *the white leg*. This disease principally affects women in the puerperal state; in a few instances it has been observed to attack pregnant women; and, in one or two cases, nurses on losing their children, have been affected by it. Women of all descriptions are liable to be attacked by it during and soon after childbed; but, those whose limbs have been pained or anasarcons during pregnancy, and who do not suckle their offspring, are more especially subject to it. It has rarely occurred oftener than once to the same female. It supervenes to easy and natural, as well as to difficult and preternatural, births. It sometimes makes its appearance in twenty-four or forty-eight hours after delivery, and at other times, not till a month or six weeks after; but in general, the attack takes place from the tenth to the sixteenth day of the lying-in. It has in many instances, attacked women who were recovering from puerperal fever; and, in some cases, has supervened, or succeeded to thoracic inflammation. It not uncommonly begins with coldness and rigors; these are succeeded by heat, thirst, and other symptoms of pyrexia; and then pain, stiffness, and other symptoms of topical inflammation supervene. Sometimes

the local affection is from the first accompanied with, but is not preceded by, febrile symptoms. Upon other occasions, the topical affection is neither preceded by puerperal fever, nor rigors, &c. but soon after it has taken place, the pulse becomes more frequent, the heat of the body is increased, and the patient is affected with thirst, head-ache, &c. The pyrexia is very various in degree in different patients, and sometimes assumes an irregular remittent or intermittent type. The complaint generally takes place on one side only at first, and the part where it commences is various; but it most commonly begins in the lumbar, hypogastric, or inguinal region, on one side, or in the hip, or top of the thigh, and corresponding labium pudendi. In this case the patient first perceives a sense of pain, weight, and stiffness, in some of the above-mentioned parts, which are increased by every attempt to move the pelvis, or lower limb. If the part be carefully examined, it generally is found rather fuller or hotter than natural, and tender to the touch, but not discoloured. The pain increases, always becomes very severe, and, in some cases, is of the most excruciating kind. It extends along the thigh, and when it has subsisted for some time longer or shorter in different patients, the top of the thigh and the labium pudendi become greatly swelled, and the pain is then sometimes alleviated, but accompanied with a greater sense of distention. The pain next extends down to the knee, and is generally the most severe on the inside and back of the thigh, in the direction of the internal cutaneous and the crural nerves; when it has continued for some time, the whole of the thigh becomes swelled, and the pain is somewhat relieved. The pain then extends down the leg to the foot, and is commonly the most severe in the direction of the posterior tibial nerve; after some time, the parts last attacked begin to swell, and the pain abates in violence, but is still very considerable, especially on any attempt to move the limb. The extremity being now swelled throughout its whole extent, appears perfectly or nearly uniform, and it is not perceptibly lessened by an horizontal position, like an oedematose limb. It is of the natural colour, or even whiter; is hotter than natural; excessively tense, and exquisitely tender when touched. When pressed by the finger in different parts, it is found to be elastic, little, if any, impression remaining, and that only for a very short time. If a puncture, or incision, be made into the limb, in some instances, no fluid is discharged; in others, a small quantity only issues out, which coagulates soon after; and in others, a larger quantity of fluid escapes, which does not coagulate; but the whole of the effused matter cannot be drawn off in this way. The

swelling of the limb varies both in degree and in the space of time requisite for its full formation. In most instances, it arrives at double the natural size, and in some cases at a much greater. In lax habits, and in patients whose legs have been very much affected with anasarca during pregnancy, the swelling takes place more rapidly than in those who are differently circumstanced; it sometimes arrives, in the former class of patients, at its greatest extent in twenty-four hours, or less, from the first attack.

Instead of beginning invariably at the upper part of the limb, and descending to the lower, this complaint has been known to begin in the foot, the middle of the leg, the ham, and the knee. In whichsoever of these parts it happens to begin, it is generally soon diffused over the whole of the limb, and, when this has taken place, the limb presents the same phenomena, exactly, that have been stated above, as observable when the inguen, &c. are first affected.

After some days, generally from two to eight, the febrile symptoms diminish, and the swelling, heat, tension, weight, and tenderness of the lower extremity, begin to abate, first about the upper part of the thigh, or about the knee, and afterwards in the leg and foot. Some inequalities are found in the limb, which, at first, feel like indurated glands, but, upon being more nicely examined, their edges are not so well defined as those of conglobate glands; and they appear to be occasioned by the effused matter being of different degrees of consistence in different points. The conglobate glands of the thigh and leg are sometimes felt distinctly, and are tender to the touch, but are seldom materially enlarged; and as the swelling subsides, it has happened, that an enlargement of the lymphatic vessels, in some part of the limb, has been felt, or been supposed to be felt.

The febrile symptoms having gradually disappeared, the pain and tenderness of the limb being much relieved, and the swelling and tension being considerably diminished, the patient is debilitated and much reduced, and the limb feels stiff, heavy, benumbed and weak. When the finger is pressed strongly against it for some time, in different points, it is found to be less elastic than at first, in some places retaining the impression of the finger for a longer, in other places for a shorter time, or scarcely at all. And, if the limb be suffered to hang down, or if the patient walk much, it is found to be more swelled in the evening, and assumes more of an oedematose appearance. In this state the limb continues for a longer or shorter time, and is commonly at length reduced wholly, or nearly to the natural size.

Hitherto the disease has been described

as affecting only one of the inferior extremities, and as terminating by resolution, or the effusion of a fluid that is moved by the absorbents; but, unfortunately, it sometimes happens, that after it abates in one limb, the other is attacked in a similar way. It also happens, in some cases, that the swelling is not terminated by resolution; for sometimes a *suppuration* takes place in one or both legs, and ulcers are formed which are difficult to heal. In a few cases, a gangrene has supervened. In some instances, the patient has been destroyed by the violence of the disease, before either suppuration or gangrene have happened.

The *predisposing causes* of this disease, when it occurs during the pregnant or puerperal state, or in a short time afterwards, appear to be, 1st, *The increased irritability and disposition to inflammation which prevailed during pregnancy, and in a still higher degree for some time after parturition.* 2dly, *The over-distended, or relaxed state of the blood-vessels of the inferior part of the trunk and of the lower extremities, produced during the latter months of utero-gestation.*

Amongst the *exciting causes* of this disease may be enumerated, 1st, *Contusions*, or violent exertions of the lower portions of the abdominal and other muscles inserted in the pelvis, or thighs, or of the muscles of the inferior extremities, and contusions of the cellular texture connected with these muscles, during a tedious labour. 2dly, *The application of cold and moisture*, which are known to act very powerfully upon every system in changing the natural distribution of the circulating fluids, and, consequently, in a system predisposed by parturition, may assist in producing the disease, by occasioning the fluids to be impelled, in unusual quantity, into the weakened vessels of the lumbar, hypogastric, and inguinal regions, and of the inferior extremities. 3dly, *Suppression*, or diminution of the lochia, and of the secretion of milk, which, by inducing a plethoric state of the sanguiferous system, may occasion an inflammatory diathesis, may favour congestion, and the determination of an unusual quantity of blood to the vessels of the parts just mentioned, and thus contribute to the production of an inflammation of these parts. 4thly, *Food taken in too large quantity, and of a too stimulating quality*, especially when the patient does not give suck. This cause both favours the production of plethora, and stimulates the heart and arteries to more frequent and violent action; the effects of which may be expected to be particularly felt in the lumbar, hypogastric, or inguinal regions, and in the lower extremities, from the state of their blood-vessels. 5thly, *Standing, or walking too much*, before the

arteries and veins of the lower half of the body have recovered sufficiently from the effects of the distention which existed during the latter months of pregnancy. This must necessarily occasion too great a determination of blood to these parts, and consequently too great a congestion in them; whence they will be more stimulating than the upper parts of the body, and inflammation will sometimes be excited in them.

From an attentive consideration of the whole of the phenomena observable in this disease, and of its remote causes and cure, no-doubt remains, Dr. Hull thinks, that the *proximate cause* consists in an *inflammatory affection, producing suddenly a considerable effusion of serum and coagulating lymph from the exhalents into the cellular membrane of the limb.*

PHLEGMASIÆ. Inflammations. The second order in the class *pyrexie* of Cullen's nosological arrangement, characterised by pyrexia, with topical pain and inflammation; the blood, after venesection, exhibiting a buffy coat.

PHLEGMATORRAGIA. (From φλεγμα, mucus, and ῥαγνμι, to break out.) A discharge of thin mucous phlegm from the nose, through cold.

PHLEGMON. (From φλεγω, to burn.) *Phlegmone.* An inflammation of a bright red colour, with a throbbing and pointed tumour, tending to suppuration.

PHLOGISTON. (From φλογίζω, to burn.) The inflammable principle. Staal gave this term to a principle which he imagined was pure fire, or the matter of fire fixed in combustible bodies, in order to distinguish it from fire in action, or in a state of liberty.

Phlogisticated air. See *Nitrogen gas*.

PHLOGOSIS. (From φλογω, to inflame.) Inflammation.

PHLYCTÆNÆ. (Φλυκταιναι, small bladders.) *Phlyctis. Phlysis.* Small pellucid vesicles, that contain a serous fluid. Linnæus and Vogel use this term as synonymous with hydætes.

PHLYZACIUM. (From φλυζω, to be hot.) A pustule on the skin, excited by fire, or heat. See *Pustule*.

PHŒNIX DACTYLIFERA. (*Phanix*; from *Phanecia*, its native soil.) The systematic name of the date-tree. See *Dactylus*.

PHOSPHATS. (*Phosphas*; from *phosphorus*.) Salts formed by the union of phosphoric acids with different bases; thus, *phosphat of ammonia, phosphat of lime, &c.*

PHOSPHITES. *Phosphis.* Salts formed by the combination of phosphorus acid with different bases; thus, *aluminous phosphite, ammoniacal phosphite, &c.*

Phosphorated hydrogen gas. See *Hydrogen gas, phosphorated*.

PHOSPHORIC ACID. *Acidum phosphoricum.* This acid may be obtained from

bones, in the following manner:—Three parts of diluted sulphuric acid are to be effused upon four parts of pulverised ashes of bones, while continually stirred. By this means the sulphuric acid combines with the calcareous earth, and disengages the phosphoric acid. The mass is then to be repeatedly washed in water, and the ley slowly evaporated: the sulphate of lime, which still adheres, will be thus gradually expelled; and at last, phosphoric acid, in a dry and vitreous form, will be obtained.

PHOSPHORUS. (From *phos*, light, and *phero*, to carry.) *Autophosphorus*. Phosphorus has never been found pure in nature. It is always met with united to oxygen, or in the state of phosphoric acid. In that state it exists very plentifully, and is united to different animal, vegetable, and mineral substances.

Properties.—Phosphorus is a flesh-coloured or yellowish semi-transparent substance, of the consistence of wax, but brittle during frost. In atmospheric air, it is luminous at common temperatures without emitting heat. It has a rough disagreeable taste, and its odour resembles that of garlic. Its specific gravity is 2.033, water being 1.000. Phosphorus crystallises in laminae, in needles, or elongated octahedra. Exposed to the light, it becomes covered with a crust, which is first white, next orange, and at last red. It becomes liquid at a temperature of 99° Fahr. It takes fire spontaneously, and burns rapidly in the open air, at 122° Fahr. with a brilliant white flame, and becomes converted into phosphoric acid. It is volatilised at 554° Fahr. It is soluble in caustic alkalies, by the assistance of heat. Expressed and essential oils take up a small quantity, and are rendered luminous. Sulphuric ether, nitric ether, and ardent spirit, dissolve it sparingly in the cold. It combines with lime, strontia, barytes, sulphur, and with metals. It is soluble in hydrogen gas, and decomposes nitric acid, and metallic solutions. It acts strongly, and frequently like poison, on living animals.

Methods of obtaining Phosphorus.—For some time, phosphorus was made in very inconsiderable quantities, and by a tedious and disagreeable process, consisting in evaporating considerable quantities of urine, and decomposing them by various means.

The following processes, now employed, are more easy and expeditious.

Gibbert's Process.—According to this method, phosphorus may be obtained very economically, and without any offensive preparation. It consists in pouring a concentrated solution of nitrate of lead, by a little at a time, into a quantity of urine, until no more cloudiness is produced by a further addition of the solution. The mixture is then to be diluted with soft water,

and suffered to stand undisturbed; when the precipitate is fully subsided, the clear fluid is to be separated. The precipitate is then formed into a paste, with charcoal powder, and the mass is to be dried gradually in an earthen pan, and submitted to distillation.

In this process the phosphoric acid of the urine, unites to the lead of the nitrate of lead, and the nitric acid joins to the ammonia and soda of the urine; hence phosphate of lead, and nitrate of soda and ammonia are formed. The former, being insoluble, falls to the bottom, and the latter salt remains in the super-natant fluid. On adding charcoal to the phosphate of lead, and exposing it to a high temperature, the union is again broken; the phosphoric acid becomes decomposed, its oxygen unites to the charcoal, and forms carbonic acid gas, which flies off during the distillation; the phosphorus appears in its simple state, and the metal is left behind in the retort, together with the super-abundant quantity of charcoal.

Nicola's Process.—Take a quantity of bones of adult animals, burn them to whiteness in an open fire, and reduce them to a fine powder. Upon three pounds of this powder, after having been put into a matrass, there may be poured two pounds of concentrated sulphuric acid of commerce; four or five pounds of water must be afterwards added by degrees, to assist the action of the acid. During the whole process, the operator must place himself and the vessel, so that the fumes of the mixture may be blown from him. The whole is then to be left in a gentle-heated sand-bath, for about twelve hours, or more, taking care to supply the loss of water which happens by evaporation. The next day, a large quantity of water must be added, the clear water afterwards decanted, and the rest strained through a cloth or sieve. The residuary matter is to be edulcorated by repeated effusions of hot water, till it passes tasteless. The water which has been used to wash out the adhering acid, is mixed with the before decanted or strained liquor, and the whole fluid is gradually evaporated in a flat earthen basin, to the consistence of syrup. It is then to be mixed with an equal weight of charcoal powder, and submitted to distillation in an iron or earthen retort. Instead of applying a receiver, the neck of the retort may be immersed in a basin of water, to a small depth, and the phosphorus, as it comes over, will fall in drops to the bottom.

In this process, the sulphuric acid unites with the calcareous earth of the bones, and forms sulphate of lime; and the phosphoric acid of the bones becomes disengaged, and remains dissolved in the liquor. The charcoal, at an elevated temperature, takes the oxygen from this acid,

and carbonic acid gas is formed, and phosphorus passes over.

Method of purifying Phosphorus.—Phosphorus obtained in either manner is of a dirty blackish colour, and soiled with a certain quantity of charcoal and half-burnt phosphorus, which gives it that appearance. In order to purify it, it must be put into a piece of chamois leather, and closely tied up in it. The whole is then immersed in a vessel of boiling water, the phosphorus melts, and may be pressed through the leather, taking care to keep it under the water.

It is better, however, to purify phosphorus by a second distillation.

Dr. Higgins purified it by means of hydrogen gas.

Methods of moulding Phosphorus into Cylinders.—In order to form phosphorus into sticks, a funnel with a long neck may be used, the lower orifice of which is closed with a cork; the funnel is then to be filled with water, and phosphorus put in it, and this being plunged into boiling water, the heat communicated to the funnel melts the phosphorus, which runs into the neck and acquires that form. The funnel is then removed into a vessel of cold water, and when it is thoroughly cooled, the cork is taken out and the phosphorus thrust out of its mould with a piece of wood, and then preserved in water.

Pelletier invented another method, which is as follows:

Take a few tubes of any length, the apertures of which are of such a size that they can be exactly closed with the extremity of the finger. Melt the phosphorus in boiling water, and apply to it one of the ends of the tube, while you hold the other in your mouth; make a short inspiration, that the phosphorus may ascend a little way in the tube; stop the inspiration when the phosphorus has risen a sufficient height, and close the extremity of the tube with the fore-finger, and immerse it in a basin of cold water. The phosphorus will soon become fixed, and by a slight shake may be separated from the tube.

The earliest account we have concerning the medicinal use of phosphorus, is in the seventh volume of Haller's Collection of Theses, relating to the history and cure of diseases. The original dissertation is entitled, *De Phosphori Loco Medicamenti adsumpti virtute medica, aliquot casibus singulibus confirmata, Auctore J. Gabi Mentz*. There are three cases of singular cures performed by means of phosphorus, narrated in this thesis; the history of these cases and cures was sent to Dr. Gabi Mentz, by his father.

The first instance is of a man who laboured under a putrid fever; for whom the best alexipharmic medicines, as they are called, together with a proper regimen,

were prescribed. A diarrhœa, however, ensued, accompanied with great anxiety about the præcordia, delirium, and general prostration of the powers of life. Proper remedies were tried to stop the diarrhœa, and check the disease, but in vain. For three days the patient had been insensible and exhausted. In this extremity the physician had recourse to phosphorus; two grains of which were exhibited, together with a sufficient quantity of theriaca, to make them into a bolus. This occasioned a gentle sweat, and general quiet. The dose was twice repeated in the evening, and again towards morning, with the addition of another grain. The sweat became copious, and the memory and the use of the external senses were restored. The patient, thus revived, was afterwards completely re-established by other remedies.

The second case, is that, of a man who laboured under a bilious fever. Although various remedies appear to have been judiciously employed, yet the disease gained ground, until at last the patient was almost quite exhausted. Three grains of phosphorus were exhibited at eleven o'clock in the forenoon, which produced a little quiet; but the patient became so thirsty that he could not refrain from drinking. After this he was quiet for two hours, and a profuse sweat broke out all over his body. The physician seeing this, ordered him another dose in the evening. He slept and perspired the whole evening, and by means of proper remedies, was afterwards completely cured.

The third case is entitled a malignant catarrhal fever, with petechia. It seems to have been the common *typhus putridus* of Cullen, accompanied with cough and catarrhal symptoms in the beginning. We are informed, that on the third day of the disease, the patient was deprived of the use of his external senses; that he became delirious and exceedingly exhausted. Two grains of phosphorus were given to him at two o'clock, and two more in the evening, which restored him to his senses, and occasioned a copious sweat. Proper remedies were afterwards employed, which accomplished his recovery.

Dr. Mentz appears to have been one of the first practitioners who tried this heroic remedy internally. Dr. Morgenstern and Dr. Hatman seem both to have afterwards employed it.

The following twelve cases, relating to this interesting subject, are translated from a thesis which is very little known in this country. It is the inaugural dissertation of one Dr. Wolff, who graduated at Gottingen in the year 1791: and who states that the cases are extracted from the private diary of his father, a physician of emi-

nence in Poland, who practised physic with great success and reputation for upwards of thirty years.

Case I.—In the month of August, 1763, I was called to a woman twenty-five years old, whom I found in a state of low delirium. Her pulse was small, weak, and tremulous, almost vanishing, as it were, under the finger. Her whole chest and arms were disfigured with livid spots. Her neighbours and attendants informed me that she had been seized with a fever about eleven days before I saw her: and that she had been attended by some ignorant practitioner, who, finding that his remedies did not succeed, had deserted her the day before, declaring that God alone could cure her.

The case appeared to me one of those desperate states of disease, in which a practitioner either ought to refrain from doing any thing, or to make trial of some new, bold, and powerful remedy, which might act as an uncommon stimulus to the nerves, and rouse their suppressed energy. Such a remedy I expected to find in phosphorus, and accordingly I ordered my patient five drops of its solution, in ether, which contained three grains of phosphorus. They were exhibited in a spoonful of Rhenish wine, and the patient swallowed a few cupfuls of an infusion of the flores tilizæ after them. I visited the patient three hours afterwards, and, not finding any change, I repeated the dose. Two hours were scarce elapsed when the pulse began to rise, and the whole body to be diffused with an equal heat; immediately afterwards, the pulse became undulatory, a breathing sweat (*sudor halituosus*) broke out, and at the same time the delirium subsided.

I exhibited a third dose at the end of sixteen hours: a number of red spots then appeared on the skin, and the patient complained very much of a sense of oppression and pain at the præcordia, and in the abdomen. These symptoms I endeavoured to allay by diluents and frequent emollient clysters, which brought away a great quantity of foul fæces. The Peruvian bark completed the cure.

Case II.—A young woman, twenty-two years of age, was, for the first time, delivered of a healthy child. She recovered perfectly well for the first nine days; but being then greatly frightened by some sudden noise in the house, she was seized with a chilly fit, afterwards with flushes of heat, and soon became delirious. I was called to her on the third day, after various remedies had been tried in vain by another physician. Finding that she had a hard pulse, with great oppression in her chest, and a foul tongue, I ordered her to be bled, to take a solution of Glauber's salts, and

to receive some antiphlogistic and emollient clysters. The other physician strongly opposed this advice, contending strenuously for his heating diaphoretic plan; and I therefore returned home. He continued to administer his alexipharms and cardiac remedies to the unhappy patient. Three days elapsed before I heard any thing of her; but some of her friends then waited on me, entreating me with great earnestness to visit her, as her physician had deserted her, declaring that it was impossible for any one to save her.

I found her with a tremulous intermittent pulse, cold extremities, and wandering in her intellects. Of the solution of phosphorus I immediately gave her five drops, in a little Rhenish wine; and in about two hours after, an equal degree of heat diffused itself over her body, and her senses returned. Upon repeating the dose, a sweat broke out, which relieved her so much that I afterwards could proceed with the proper remedies for the further cure of the complaint.

Case III.—A young man, twenty-two years old, was seized with a putrid-gastric fever, which was accompanied with a white miliary eruption. On the seventh day of the disease, while under the influence of a profuse sweat, he was taken from his bed, at his own desire, and had his linen changed. An hour was scarcely elapsed when he was seized with great anxiety, the miliary eruption almost totally disappeared, and he began to grow delirious. Being called to him, I gave him the solution of phosphorus, with an infusion of elder flowers and Rhenish wine. In about an hour afterwards, the sweat and eruption returned, and he was snatched from the jaws of death.

Case IV.—A youth sixteen years old, was seized with a putrid fever, on the seventh day of which he was affected with diarrhœa of so severe a kind, that he had forty-eight stools in the course of a day. Being sent for the day after, I found him with an hippocratic countenance, and *subultus tendinum*. Having exhibited the phosphorus twice to him, a general breathing sweat broke out, and the diarrhœa subsided. Afterwards, proper remedies were exhibited, and he was restored to health in fifteen days.

Seven Cases.—During the years 1770 and 1771, while the war was carried on upon the borders of our remotest provinces between the Turks and Russians, a putrid fever took its rise in the camps, and spread itself to us. It was then that I ordered this divine remedy, with the happiest effects, to seven sick who laboured under the putrid fever, some of whom had eruptions, with great prostration of strength.

Case XII.—A young lady of quality,

twenty-five years old, endowed with such an exquisite degree of nervous sensibility, that she used frequently to be affected with spasms and convulsive motions without any apparent cause, became pregnant in the year 1774, which was the seventh of her marriage. On the third day preceding delivery, she was seized with a disease which might be justly said to be somewhat between catalepsy and coma vigil. She lay stretched out upon her back, her eyes open and fixed, her legs and arms were quite flexible, and remained in any posture in which they were put; her pulse and respiration were entirely natural, and she swallowed whatever fluid was put into her mouth.

After she had remained three days in this state, she was delivered of a dead child, which, from the marks of putrescency on its body, must have died some days before.

Evacuations carefully employed, antispasmodic and nervous remedies exhibited, both by the mouth and anus, embrocations and liniments, produced such a salutary change, that in fifteen days the patient was able to leave her bed, and the greatest hope was therefore entertained of her complete recovery. But, on the twenty-eighth day after delivery, when I went again to visit my patient, (who lived at a considerable distance from me,) I found her complaining of rigour and chilliness of the inferior extremities, her pulse was frequent and small, and her face was more flushed than usual. Frictions, clysters, the pediluvium, and antispasmodics, were all tried, but without producing any favourable event, and, towards the middle of the night, she became totally rigid; yet she had complete possession of her mind, although all the external senses, the sight excepted, were abolished. I gave to her three drops of the solution of phosphorus in a spoonful of wine. Scarcely had an hour elapsed when a warm sweat broke out, and, together with it, a white miliary eruption. Afterwards, other remedies were exhibited, which effected her recovery.

Such are the cases related by Dr. Wolff, which it is presumed will be found sufficiently interesting to awaken and arrest the attention of the practitioners of this country. A medicine of greater powers cannot be named, if the facts which have been related are correctly true. Little doubt can be entertained that many trials will soon be made with it in this country; but it is to be hoped they will be made with caution, and under as favourable circumstances for the reception of such a powerful stimulus as the nature of the diseases in which it is to be given will admit of. The following particulars concerning this subtle remedy, will serve as a salutary caution against the too free exhibition of it in the time of making experiments:—

The dangerous consequences which are

likely to follow the injudicious administration of phosphorus, cannot be impressed on the mind more strongly than by the relation of a few cases and experiments which are mentioned by Wackard, in the fourth part of his miscellaneous writings, (*Vermischte Medicinische Schriften*, von M. A. Wackard.) These cases and experiments I literally translated from the original German; a work from which the medical practitioner may reap much information, being replete with practical remarks made by a very attentive and accurate observer.

A Jew, of a phlegmatic habit, was struck with apoplexy. He was speechless and lame, and could not void his feces except he was assisted by art; his appetite, however, was good. Mineral baths, and many remedies, were tried to restore him, but in vain. I was at last tempted to make a trial of phosphorus. At first, from two to three grains were given to him, and it was my intention to have increased the dose to five or six. The first dose was given in a conserve, but the following day in honey. His excrements were luminous. Suddenly, about the middle of the third night, he was seized with violent vomiting, during which he was very ill; but, after the vomiting had ceased, he found himself better. The pulse was small and quick. I did not venture to give any more phosphorus, particularly as his relations were afraid of it. A blister was applied, and cooling and demulcent drinks were exhibited. He looked very ill, seemed quite exhausted, and appeared to suffer pain in the abdomen, which was a little tumefied. He took some nourishment every day; but what he ate was very small in quantity. The vomiting ceased entirely. He lived four days longer, and died. I did not see him the day when that event happened; but he had all the symptoms of having died of gangrene. There were even externally, according to the report of the surgeon, many large gangrenous spots. This case caused me a great deal of uneasiness. The patient could not speak for some time before his death, and could not therefore describe his sufferings.

Almost about the same time that Professor Zessler exhibited the phosphorus to his two patients, the intelligence was communicated to a man who had a tendency to apoplexy and to palsy. He had read in *Mellin's Materia Medica* all that was said in praise of phosphorus, and he was, in consequence, desirous of making trial of it; but, before he began to take it, he was seized with a kind of apoplectic fit, in which he lay as if deprived of all power. He caused three grains of phosphorus to be rubbed with a little oil, and to be given to him; he took his dose morning and evening. It seemed to revive him, for he got up and walked about the house. He

continued to take this medicine several days, upon which he became feverish, and was affected with nausea and loss of appetite. A gentle emetic was given him. Two days after this I saw him for the first time; he told me that he had been violently affected with pain in the abdomen, particularly towards the evening. He was very dry, and his eyes were yellow. When I pressed my hand on the abdomen it did not give him pain. I ordered him whey, nitre, acid drinks, cream of tartar, and clysters. The third day after this he grew worse. I happened to be out of town, and another physician was sent for, who gave him musk, and various other remedies. I visited him in the evening and found him gangrenous. His hands were applied to his head, as if he suffered acute pain there; his screams were loud and unremitting; he was delirious and spoke unintelligibly, and there was not any pulse at the wrist. He died about midnight. Upon inspecting the body, the liver appeared obstructed, and the stomach was gangrenous near the cardia. Is it not to be feared that the phosphorus had occasioned this? or was this the effect of a gouty or rheumatic acrimony thrown upon that part? It was afterwards remarked, that a number of people were seized at this time with vomiting and violent pains in the bowels, and also with the iliac passion. Two months were scarcely elapsed when I met with a case in which all the circumstances of the complaint were similar to those narrated in the above-mentioned case, but where the patient had not swallowed a single grain of phosphorus.

The following are some experiments which Dr. Weickard made with phosphorus on dogs:—

"It is now," he observes, "upwards of a year since six grains of phosphorus were offered to a hungry dog. The phosphorus was enclosed in a piece of meat; the dog smelt it, and refused to take it; it was therefore forced down his throat. He immediately began to run about the room, exhibiting much anxiety, and seemed desirous to go out. He grew sick, and vomited the bolus, which seemed to be on fire; upon this, the dog again appeared impatient to get out. The smell of the phosphorus, however, ceased to be emitted by the vomited matter. The dog smelt it again, and ate it. He immediately became extremely lively, springing quickly from one table to another. The experiments ought to have stopped here; but some one brought another piece of meat, in which there were eight grains of phosphorus enclosed. This was also forced down the dog's throat, and he again vomited it on fire; he did not, however, exhibit the same anxiety to leave the room which he had done the first time, but waited patiently, until the disagreeable savor which

arose and filled the apartment, had escaped from his morsel, after which he ate it greedily. His inclination to leap and run about was now uncommonly great; nothing was too high or too distant for him; but as we were afraid he might do some injury by this kind of exercise, he was dismissed the apartment. He again vomited once or twice; but since that time has remained perfectly well."

Such are the principal facts related by Weickard. We now come to the latest publication on the subject of phosphorus. It is a letter on its medicinal virtues by M. Alphonso Leroi, and is printed in the only volume of *Memoirs* which the Society of Emulation at Paris has yet published. One of the first experiments which M. Leroi made was upon himself; he swallowed three grains of phosphorus with some theriaca. It is a wonder, he says, that he was not killed by it; for phosphorus does not require more air than is commonly contained in the stomach, in order to take fire, and burn in such a manner as to have eaten through that organ. "For two hours," the Professor observes, "I found myself extraordinarily incommoded; I drank frequently little draughts of cold water. After some hours had elapsed the uneasiness ceased. My urine was high-coloured; but the following day my muscular force was doubled, and I felt an insupportable venereal irritation. I afterwards gave this remedy to the young man whose case is related in the *Gazette de Santé*, for the 29th of August, 1779. It was truly owing to phosphorus that his life was saved in a most malignant fever, in which the prostration of strength was so great as not to leave many other resources but this remedy."

M. Leroi then proceeds to inform us, that he has frequently employed phosphorus since that time. He at first prescribed it in the form of a lobeoch. It is always difficult to pulverize it; but this difficulty he overcame in the following manner: "He puts the phosphorus into warm water, and agitates it violently; it divides itself like oil, into a great number of little globules, and, if cold water be now added, it is precipitated to the bottom in the form of a powder. Of this powder one or two grains are to be taken, and rubbed with a little sugar, a drop or two of oil, and some yolk of an egg. A quarter of a grain every day is quite sufficient to produce great effects."

At other times, he says, he has given it in a mixture of oil, syrup, and some aromatic distilled water. He asserts, that Kunkle gave phosphorus internally in England in the form of pills, which were luminous; and he informs us that he himself has found out the manner of composing them; but they require so nice a manipulation, that

he has intrusted the secret only to Messrs. Pelletier, the brothers, being warned by his own experience of the mischief which arises from giving it improperly prepared. Each pill contains the eighth of a grain of phosphorus. They are endowed with a soporific and calming property. Professor Leroi has frequently employed them in cases of rheumatismus, in a great number of nervous diseases, pituitous diseases, and in many acute and chronic complaints. He believes that this remedy is capable of prolonging life beyond the natural period; and after having spoken of its great restorative power, he related the following case in confirmation of this opinion:—

“I was one day called to an old man, aged eighty-seven, the uncle of Madame de Fourqueux, in whom life seemed to be almost totally extinguished. I composed for him a mixture of six ounces of different distilled aromatic waters, one ounce of oil containing three grains of phosphorus, and two ounces of syrup. Of this he took three table-spoonfuls every day; and besides this, he took eight drops of volatile alkali, in a glass of sweet aromatic water, twice a-day, before his meals. By these means I illuminated the dying embers of life, and he survived seven years after an attack of weakness in which it must have appeared like madness to attempt to interfere.

“I may safely affirm, that I have been as much occupied in seeking out the cases in which this medicine proves hurtful, as in detecting those in which it is serviceable; and I can assert, not only from my own experience, but also from that of the late M. Lecoindre, my pupil, who was physician to the hospital at Nambouillet, that we have not found it hurtful in any one case: that it has only been useful in some, and that only in such cases in which life was already extinct is some parts of such patients as had not a sufficient portion of living principle in them to reanimate the whole frame.”

Mr. Leroi affirms, that the divisibility of phosphorus is almost infinite; and, in proof of this assertion, he states, that “the body of a woman who died of a putrid fever, and who had taken one grain of phosphorus, was entirely luminous within. The hands of the late Rielle, the anatomist, who opened the body, were luminous some hours after they had been washed!”

Upon maturely considering all the facts which have been brought forward concerning it, little room for doubt will be left in the mind, as to its uncommon powers; but, although this be granted, experience it still wanting to point out the diseases, and various stages of disease, in which it may be employed with equal safety and utility, as also to determine the circumstances under which it would

be dangerous even to try it. Thus much may doubtless be asserted concerning it, that it is one of the highest stimuli which we have in the catalogue of the materia medica; and that, although it is affirmed by M. Leroi, and others, to be “calming and sedative,” it is only so in such cases as wine, æther, Hoffman’s anodyne liquor, and opium, are also found to be calming and sedative, that is, in cases where the arterial action of the whole frame is nearly exhausted, although still quick. Every practitioner should be cautioned against exhibiting it in any inflammatory disease, where much strength exists; and in all cases, very small doses should be first exhibited, and those with the utmost circumspection.

Phosphuret. See Phosphuretum.

PHOSPHURETUM. (*Phosphuretum*, from *phosphorus*.) A combination of phosphorus not oxygenated, with different bases; as, *phosphuret of copper*, *phosphuret of iron*, &c.

PHOTOPHOBIA. (From *φως*, light, and *φοβία*, to dread.) Such an intolerance of light, that the eye, or rather the retina, can scarcely bear its irritating rays. Such patients generally wink, or close their eyes in light, which they cannot bear without exquisite pain, or confused vision. The proximate cause is too great a sensibility in the retina. The species are,

1. *Photophobia inflammatoria*, or dread of light from an inflammatory cause, which is a particular symptom of the internal ophthalmia.

2. *Photophobia*, from the disuse of light, which happens to persons long confined in dark places or prisons; on the coming out of which into light the pupil contracts, and the persons cannot bear light. The depression of the cataract occasions this symptom, which appears as though fire and lightening entered the eye, not being able to bear these strong rays of light.

3. *Photophobia nervæ*, or a nervous photophobia, which arises from an increased sensibility of the nervous expansion and optic nerve. It is a symptom of the hydrophobia, and many disorders, both acute and nervous.

4. *Photophobia*, from too great light, as looking at the sun, or at the strong light of modern lamps.

PHOTOPSIA. (From *φως*, light, and *οψις*, vision.) Lucid vision. An affection of the eye in which the patient perceives luminous rays, ignited lines, or coruscations.

PHRAGMUS. (From *φρασσα*, to enclose or fence; so called from their being set round like a fence of stakes.) The rows of teeth.

PHRENES. (*Phren*. from *φην*, the mind, because the ancients imagined it was the seat of the mind.) The diaphragm.

PHRENESIS. See *Phrenitis*.

PHRENIC NERVE. (*Nervus phrenicus*, from *φνις*, the diaphragm). Diaphragmatic nerve. It arises from a union of the branches of the third, fourth, and fifth cervical pairs, on each side, passes between the clavicle and subclavian artery, and descends from thence by the pericardium to the diaphragm.

PHRENICÆ ARTERIÆ. The arteries going to the diaphragm.

PHRENICÆ VENÆ. The veins coming from the diaphragm.

PHRENITIS. (*φνιτις*, from *φνις*, the mind). *Phrenetiasis*. *Phrenismus*. *Cephalitis*. *Sphacelismus*. *Cephalalgia inflammatoria*. By the Arabians *karabitus*. Phrensy or inflammation of the brain. A genus of disease in the class *pyrexia*, and order *phlegmasia* of Cullen; characterised by strong fever, violent headache, redness of the face and eyes, impatience of light and noise, watchfulness, and furious delirium. It is symptomatic of several diseases, as worms, hydrophobia, &c. Phrenitis often makes its attacks with a sense of fulness in the head, flushing of the countenance, and redness of the eyes, the pulse being full, but in other respects natural. As these symptoms increase, the patient becomes restless, his sleep is disturbed, or wholly forsakes him. It sometimes comes on, as in the epidemic, of which Saalman gives an account, with pain, or a peculiar sense of uneasiness of the head, back, loins, and joints; in some cases, with tremor of the limbs, and intolerable pains of the hands, feet, and legs. It now and then attacks with stupor, and rigidity of the whole body, sometimes with anxiety and a sense of tension referred to the breast, often accompanied with palpitation of the heart. Sometimes nausea and a painful sense of weight in the stomach, are among the earliest symptoms. In other cases, the patient is attacked with vomiting, or complaints of the heart-burn, and griping pains in the bowels. When the intimate connexion which subsists between the brain and every part of the system is considered, the variety of the symptoms attending the commencement of phrenitis is not so surprising, nor that the stomach in particular should suffer, which so remarkably sympathises with the brain. These symptoms assist in forming the diagnosis between phrenitis, and synocha. The pain of the head soon becomes more considerable, and sometimes very acute. 'If the meninges,' says Dr. Fordyce, 'are affected, the pain is acute; if the substance only, obtuse, and sometimes but just sensible.' And Dr. Cullen remarks, 'I am here, as in other analogous cases, of opinion, that the symptoms above mentioned of an acute inflammation, always

mark inflammations of membranous parts, and that an inflammation of parenchyma, or substance of viscera, exhibits, at least commonly, a more chronic inflammation.

The seat of the pain is various: sometimes it seems to occupy the whole head; sometimes, although more circumscribed, it is deep-seated, and ill defined. In other cases it is felt principally in the forehead or occiput. The redness of the face and eyes generally increases with the pain, and there is often a sense of heat and throbbing in the head, the countenance acquiring a peculiar fierceness. These symptoms for the most part, do not last long before the patient begins to talk incoherently, and to shew other marks of delirium. Sometimes, however, Saalman observes, delirium did not come on till the fifth, sixth, or seventh day. The delirium gradually increases, till it often arrives at a state of phrensy. The face becomes turgid, the eyes stare, and seem as if bursting from their sockets, tears, and sometimes even blood, flowing from them; the patient in many cases resembling a furious maniac, from whom it is often impossible to distinguish him, except by the shorter duration of his complaint. The delirium assists in distinguishing phrenitis and synocha, as it is not a common symptom in the latter. When delirium does attend synocha, however, it is of the same kind as in phrenitis.

We should, *a priori*, expect in phrenitis considerable derangement in the different organs of sense, which so immediately depend on the state of the brain. The eyes are incapable of bearing the light, and false vision, particularly that termed *muce volitantes*, and flashes of light seeming to dart before the eyes, are frequent symptoms. The hearing is often so acute, that the least noise is intolerable; sometimes, on the other hand, the patient becomes deaf; and the deafness, Saalman observes, and morbid acuteness of hearing, sometimes alternate. Affections of the smell, taste, and touch, are less observable.

As the organs of sense are not frequently deranged in synocha, the foregoing symptoms farther assist the diagnosis between this complaint and phrenitis.

The pulse is not always so much disturbed at an early period, as we should expect from the violence of the other symptoms, compared with what we observe in idiopathic fevers. When this circumstance is distinctly marked, it forms, perhaps, the best diagnosis between phrenitis and synocha, and gives to phrenitis more of the appearance of mania. In many cases, however, the fever runs as high as the delirium; then the case often almost exactly resembles a case of violent synocha, from which it is the more difficult to distinguish it if the pulse be full and strong. In ge-

neral, however, the hardness is more remarkable than in synocha, and in many cases the pulse is small and hard, which may be regarded as one of the best diagnostics between the two complaints, the pulse in synocha being always strong and full. In phrenitis it is sometimes, though rarely, intermitting. The respiration is generally deep and slow, sometimes difficult, now and then interrupted with hicough, seldom hurried and frequent; a very unfavourable symptom. In many of the cases mentioned by Saalman, pneumonia supervened.

The deglutition is often difficult, sometimes convulsive. The stomach is frequently oppressed with bile, which is an unfavourable symptom; and complete jaundice, the skin and urine being tinged yellow, sometimes supervenes. Worms in the stomach and bowels are also frequent attendants on phrenitis, and, there is reason to believe, may have a share in producing it. The hydrocephalus internus, which is more allied to phrenitis than dropsy of the brain, properly so called, seems often, in part at least, to arise from derangement of the primæ viz, particularly from worms. We cannot otherwise account for the frequent occurrence of these complaints.

Instead of a superabundance of bile in the primæ viz, there is sometimes a deficiency, which seems to afford even a worse prognosis. The alvine faces being of a white colour, and a black cloud in the urine, are regarded by Lobb as fatal symptoms. The black cloud in the urine is owing to an admixture of blood: when unmixed with blood, the urine is generally pale.

There is often a remarkable tendency to the worst species of hæmorrhages, towards the fatal termination of phrenitis. Hæmorrhagy from the eyes has already been mentioned. Hæmorrhagy from the intestines also, tinging the stools with a black colour, is not uncommon. These hæmorrhages are never favourable; but the hæmorrhages characteristic of synocha, particularly that from the nose, sometimes occur at an earlier period, and, if copious, generally bring relief. More frequently, however, blood drops slowly from the nose, demonstrating the violence of the disease, without relieving it. In other cases there is a discharge of thin mucus from the nose.

Tremors of the joints, convulsions of the muscles of the face, grinding of the teeth, the face from being florid suddenly becomes pale, involuntary tears, a mucus from the nose, the urine being of a dark red or yellow colour, or black, or covered with a pellicle, the faces being either vilous or white, and very fetid, profuse sweat of the head, neck, and shoulders,

paralysis of the tongue, general convulsions, much derangement of the internal functions, and the symptoms of other visceral inflammations, particularly of the pneumonia, supervening, are enumerated by Saalman as affording the most unfavourable prognosis. The delirium changing to coma, the pulse at the same time becoming weak, and the deglutition difficult, was generally the forerunner of death. When, on the contrary, there is a copious hæmorrhagy from the hæmorrhoidal vessels, from the lungs, mouth, or even from the urinary passages, when the delirium is relieved by sleep, and the patient remembers his dreams, when the sweats are free and general, the deafness is diminished or removed, and the febrile symptoms become milder, there are hopes of recovery.

In almost all diseases, if we except those which kill suddenly, as the fatal termination approaches, nearly the same train of symptoms supervenes, viz. those denoting extreme debility of all the functions. Saalman remarks that the blood did not always shew the buffy coat.

Phrenitis, like most other complaints, has sometimes assumed an intermitting form, the fits coming on daily, sometimes every second day. When phrenitis terminates favourably, the typhus, which succeeds the increased excitement, is generally less in proportion to that excitement, than in idiopathic fevers; a circumstance which assists in distinguishing phrenitis from synocha.

The imperfect diagnosis between these complaints is further assisted by the effects of the remedies employed. For in phrenitis in removing the delirium and other local symptoms, the febrile symptoms in general soon abate. Whereas in synocha, although the delirium and head-ache be removed, yet the pulse continues frequent, and other marks of indisposition remain for a much longer time.

It will be of use to present at one view the circumstances which form the diagnosis between phrenitis and synocha.

Synocha generally makes its attack in the same manner; its symptoms are few and little varied. The symptoms at the commencement of phrenitis are often more complicated, and differ considerably in different cases. Derangement of the internal functions are comparatively rare in synocha. In phrenitis it almost constantly attends, and often appears very early. The same observation applies to the derangement of the organs of sense. In synocha, the pulse from the commencement, is frequent, slow, and rapid. In phrenitis, symptoms denoting the local affection often became considerable before the pulse is much disturbed. In phrenitis we have seen that the pulse sometimes very suddenly loses its strength, the worst

species of hæmorrhages, and other symptoms denoting extreme debility, shewing themselves; such symptoms are generally the forerunner of death. But that when the termination is favourable, the degree of typhus which succeeds it is less in proportion to the preceding excitement in synocha. Lastly, if we succeed in removing the delirium and other symptoms affecting the head, the state of the fever is found to partake of this favourable change more immediately and completely than in synocha, where, although we succeed in relieving the head-ache or delirium, the fever often suffers little abatement.

With regard to the duration of phrenitis, Eller observes, that when it proves fatal, the patient generally dies within six or seven days. In many fatal cases, however, it is protracted for a longer time, especially where the remissions have been considerable. Upon the whole, however, the longer it is protracted, providing the symptoms do not become worse, the better is the prognosis.

PHRENITIASIS. See *Phrenitis*.

Phreny. See *Phrenitis*.

PHTHEIRIASIS. (From *phug*, a louse.)

See *Phthiriasis*.

PHTHEIROCTONUM. See *Phthetrectonum*.

PHTHIRIASIS. (From *phug*, a louse.)

Morbus Pediculorus. Pediculatio. Phthiriasis. A disease in which several parts of the body generate lice, which often puncture the skin, and produce little sordid ulcers.

PHTHEIROCTONUM. (From *phug*, a louse, and *ctono*, to kill, because it destroys lice.) *Phthirium.* The herb *staphis-agra*, or *Staves-acre*.

PHTHISIS. (From *phim*, to consume.) *Tubes Pulmonalis.* Pulmonary consumption. A genus of disease in the class *pyrexia*, and order *hæmorrhagie* of Cullen; known by emaciation, debility, cough, hectic fever, and purulent expectoration.

Species: 1. *Phthisis incipiens*, incipient, without any expectoration of pus.

2. *Phthisis humida*, with an expectoration of pus.

3. *Phthisis scrophulosa*, from scrophulous tubercles in the lungs, &c.

4. *Phthisis hæmoptica*, from hæmoptysis.

5. *Phthisis exanthematica*, from exanthemata.

6. *Phthisis chlorotica*, from chlorosis.

7. *Phthisis syphilitica*, from a venereal ulcer in the lungs.

The causes which predispose to this disease are very numerous. The following are, however, the most general: hereditary disposition; particular formation of body, and obvious by a long neck, prominent shoulders, and narrow chest; scrophulous diathesis, indicated by a fine clear skin, fair hair, delicate

rosy complexion, large veins, thick upper lip, a weak voice, and great sensibility; certain diseases, such as syphilis, scrophula, the small-pox, and measles; particular employments exposing artificers to dust, such as needle pointers, stone-cutters, millers, &c. or to the fumes of metals or minerals under a confined and unwholesome air; violent passions, exertions or affections of the mind, as grief, disappointment, anxiety, or close application to study, without using proper exercise; frequent and excessive debaucheries, late watching, and drinking freely of strong liquors: great evacuations, as diarrhæa, diabetes, excessive venery, fluor albus, immoderate discharge of the menstrual flux, and the continuing to suckle too long under a debilitated state; and, lastly, the application of cold, either by too sudden a change of apparel, keeping on wet clothes, lying in damp beds, or exposing the body too suddenly to cool air, when heated by exercise; in short, by any thing that gives a considerable check to the perspiration. The more immediate or occasional causes of phthisis are, hæmoptysis, pneumonic inflammation proceeding to suppuration, catarrh, asthma, and tubercles, the last of which is by far the most general. The incipient symptoms usually vary with the cause of the disease; but when it arises from tubercles, it is usually thus marked: It begins with a short dry cough, that at length becomes habitual, but from which nothing is spit up for some time, except a frothy mucus that seems to proceed from the fauces. The breathing is at the same time somewhat impeded, and upon the least bodily motion is much hurried: a sense of straitness, with oppression at the chest, is experienced; the body becomes gradually leaner, and great languor, with indolence, dejection of spirits, and loss of appetite, prevail. In this state the patient frequently continues a considerable length of time, during which he is, however, more readily affected than usual by slight colds, and upon one or other of the occasions the cough becomes more troublesome and severe, particularly by night, and it is at length attended with an expectoration, which towards morning is more free and copious. By degrees the matter which is expectorated becomes more viscid and opaque, and now assumes a greenish colour and purulent appearance, being on many occasions streaked with blood. In some cases, a more severe degree of hæmoptysis attends, and the patient spits up a considerable quantity of florid, frothy blood. The breathing at length becomes more difficult, and the emaciation and weakness goes on increasing. With these, the person begins to be sensible of pain in some part of the thorax, which, however, is usually

felt at first under the sternum, particularly on coughing. At a more advanced period of the disease a pain is sometimes felt on one side, and at times prevails in so high a degree, as to prevent the person from lying easily on that side : but it more frequently happens, that it is felt only on making a full inspiration, or coughing. Even where no pain is felt it often happens, that those who labour under phthisis cannot lie easily on one or other of their sides, without a fit of coughing being excited, or the difficulty of breathing being much increased. At the first commencement of the disease, the pulse is often natural, or perhaps is soft, small, and a little quicker than usual : but when the symptoms which have been enumerated have subsisted for any length of time, it then becomes full, hard, and frequent. At the same time the face flushes, particularly after eating, the palms of the hands and soles of the feet are affected with burning heat ; the respiration is difficult and laborious ; evening exacerbations become obvious, and by degrees, the fever assumes the hectic form. This species of fever is evidently of the remittent kind, and has exacerbations twice every day. The first occurs usually about noon, and a slight remission ensues about five in the afternoon. This last is, however, soon succeeded by another exacerbation, which increases gradually until after midnight ; but about two o'clock in the morning a remission takes place, and this becomes more apparent as the morning advances. During the exacerbations the patient is very sensible to any coolness of the air, and often complains of a sense of cold when his skin is, at the same time, preternaturally warm. Of these exacerbations, that of the evening is by far the most considerable. From the first appearance of the hectic symptoms, the urine is high coloured, and deposits a copious branny red sediment. The appetite, however, is not greatly impaired, the tongue appears clean, the mouth is usually moist, and the thirst is inconsiderable. As the disease advances, the fauces put on rather an inflamed appearance, and are beset with aphthæ, and the red vessels of the tunica adnata become of a pearly white. During the exacerbations, a florid circumscribed redness appears on each cheek ; but at other times the face is pale, and the countenance somewhat dejected. At the commencement of hectic fever, the belly is usually costive ; but in the more advanced stages of it, a diarrhœa often comes on, and this continues to recur frequently during the remainder of the disease ; colliquative sweats likewise break out, and these alternate with each other, and induce vast debility. In the last stage of the disease the emaciation is so great, that the patient

has the appearance of a walking skeleton ; his countenance is altered, his cheeks are prominent, his eyes look hollow and languid, his hair falls off, his nails are of a livid colour, and much incurvated, and his feet are affected with œdematous swellings. To the end of the disease the senses remain entire, and the mind is confident and full of hope. It is, indeed, a happy circumstance attendant on phthisis, that those who labour under it are seldom apprehensive or aware of any danger ; and it is no uncommon occurrence to meet with persons labouring under its most advanced stage, flattering themselves with a speedy recovery, and forming distant projects under that vain hope. Some days before death the extremities become cold. In some cases a delirium precedes that event, and continues until life is extinguished.

As an expectoration of mucus from the lungs may possibly be mistaken for purulent matter, and may thereby give us reason to suspect that the patient labours under a confirmed phthisis, it may not be amiss to point out a sure criterion, by which we shall always be able to distinguish the one from the other. The medical world are indebted to the late Mr. Charles Darwin for the discovery, who has directed the experiment to be made in the following manner :

Let the expectorated matter be dissolved in vitriolic acid, and in caustic luvium, and add pure water to both solutions. If there is a fair precipitation in each, it is a certain sign of the presence of pus ; but if there is not a precipitate in either, it is certainly mucus.

Mr. Everard Home, in his dissertation on the properties of pus, informs us of a curious, and apparently a decisive mode of distinguishing accurately between pus and animal mucus. The property, he observes, which characterises pus, and distinguishes it from most other substances, is, its being composed of globules, which are visible when viewed through a microscope ; whereas animal mucus, and all chemical combinations of animal substances appear in the microscope to be made up of flakes. This property was first noticed by the late Mr. John Hunter.

Pulmonary consumption, is in every case to be considered as attended with much danger ; but it is more so when it proceeds from tubercles, than when it arises in consequence either of hæmoptysis, or pneumonic suppuration. In the last instance the risk will be greater where the abscess breaks inwardly, and gives rise to empyema, than when its contents are discharged by the mouth. Even cases of this nature have, however, been known to terminate in immediate death. The impending danger is generally to be judged of, however, by the hectic symptoms ; but more parti-

cularly by the fetor of the expectoration, the degree of emaciation and debility, the coilliquative sweats, and the diarrhoea. The disease has, in many cases, been found to be considerably retarded in its process by pregnancy; and in a few has been alleviated by an attack of mania.

The morbid appearance most frequently to be met with on the dissections of those who die of phthisis, is the existence of tubercles in the cellular substance of the lungs. These are small tumours which have the appearance of indurated glands, are of different sizes, and are often found in clusters. Their firmness is usually in proportion to their size, and when laid open in this state they are of a white colour, and of a consistence nearly approaching to cartilage. Although indolent at first, they at length become inflamed, and lastly form little abscesses or vomicae, which breaking, and pouring their contents into the bronchiz, give rise to a purulent expectoration, and thus lay the foundation of phthisis. Such tubercles or vomicae are most usually situated at the upper and back part of the lungs; but in some instances they occupy the outer part, and then adhesions to the pleura are often formed.

When the disease is partial, only about a fourth of the upper and posterior part of the lungs is usually found diseased; but in some cases life has been protracted till not one twentieth part of them appeared, on dissection, fit for performing their function. A singular observation, confirmed by the morbid collections of anatomists, is, that the left lobe is much oftener affected than the right.

PHTHISIS ISCHIADICA. A wasting of the thigh and leg from an abscess or other cause in the hip.

PHTHISIS SUPILLI. An amaurosis.

PHTHORIA. (From *phtho*, an abortion.) Medicines which promote abortion.

PHU. (*ph*, or *phw*, from *phua*, Arab.) See *Valeriana major*.

PHYETHLON. (From *phu*, to grow.) A red and painful tubercle in the arm-pits, neck and groins.

PHYLACTERIUM. (From *phylaxo*, to preserve.) An amulet or preservative against infection.

PHYLLANTHUS EXBOLICA. The systematic name of the Indian tree from which the emblic mirobalan is obtained.

PHYLLITIS. (From *phyllos*, a leaf; so called because the leaves only appear.) See *Scolopendrium*.

PHYMA. (From *phu*, to produce.) A small tubercle on any external part of the body.

PHYSALIS. (From *phu*, to inflate; so called because its seed is contained in a kind of bladder, and *alkekengi*, alkakangi, Arab. the haliacabus or winter cherry.)

The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

PHYSALIS ALKEKENGI. The systematic name of the winter cherry. See *Alkekengi*.

PHYSCONIA. (From *phu*, a big-bellied fellow.) *Hypocæca*. *Hypersarcidæ*. Enlargement of the abdomen. A genus of disease in the class *cachexia*, and order *intumescencia* of Cullen; known by a tumour occupying chiefly one part of the abdomen, increasing slowly, and neither sonorous nor fluctuating. Species: 1. *Hepatica*. 2. *Splenica*. 3. *Renalis*. 4. *Uterina*. 5. *Ab ovario*. 6. *Mesenterica*. 7. *Omentalis*. 8. *Visceralis*.

PHYSEMA. (From *phu*, to inflate.) *Phyæsis*. A windy tumour.

PHYSTER MACROCEPHALUS. (*Physter*, from *phu*, to inflate; so named from its action of blowing and discharging water from its nostrils, and *macrocephalus*, from *μακρο*, long, and *κεφαλη*, the head; from the length of its head.) See *Ambergria*.

PHYSIOGNOMY. (*Physiognomia*, from *phu*, nature, and *gno*, to know.) The art of knowing the disposition of a person from the countenance.

PHYSIOLOGY. (*Physiologia*, from *phu*, nature, and *λογος*, a discourse.) The science which treats of the actions and powers of an animated body.

PHYSCOLE. (From *phu*, wind, and *αλλε*, a tumour.) A species of hernia, whose contents are distended with wind.

PHYSCOPHALUS. (From *phu*, wind, and *κεφαλη*, the head.) Emphysema of the head. See *Pneumotisis*.

PHYSOMETRA. (From *phu*, to inflate, and *μετρα*, the womb.) *Hysterophyæ*. A windy swelling of the uterus. A tympany of the womb. A genus of disease in the class *cachexia* and order *intumescencia* of Cullen; characterised by a permanent elastic swelling of the hypogastrium, from flatulent distention of the womb. It is a rare disease, and seldom admits of a cure.

PHYTRUMA. (From *phu*, to generate; so called from its great increase and growth.) The herb rocket.

PHYTOLACCA DECANDRIA. (*Phytolacca*. From *phu*, a plant, and *λακκα*, gum lac; so called because it is of the colour of lacca, and *decandria*, from its Linnæan arrangement.) *Solanum magnnm virginianum rubrum* Pork-physic. Pork-weed. Poke-weed. Red weed of Virginia. Red nightshade. The systematic name of the American nightshade. *Solanum racemosum Americanum*. In Virginia and other parts of America, the inhabitants boil the leaves, and eat them in the manner of spinach. They are said to have an anodyne quality, and the juice of the root is violently cathartic. The Portuguese

had formerly a trick of mixing the juice of the berries with their red wines, in order to give them a deeper colour; but it was found to debase the flavour. This was represented to his Portuguese majesty, who ordered all the stems to be cut down yearly before they produced flowers, thereby to prevent any farther adulteration. This plant has been used as a cure for cancers, but to no purpose.

PHYTOLOGY. (*Phytologia*. From *φυτον*, an herb, and *λογος*, a discourse.) That part of the science of natural history which treats on plants.

PHYTONIMESALIA. (From *φυτον*, a plant, and *mineralis*, a mineral.) Substances of a vegetable and mineral nature; as amber and coral.

PIA MATER. (*Pia mater*, the natural mother; so called because it embraces the brain, as a good mother folds her child.) *Localis membrana. Meningis tenuis.* A thin membrane, almost wholly vascular, that is firmly accreted to the convolutions of the cerebrum, cerebellum, medulla oblongata, and medulla spinalis. Its use appears to be, to distribute the vessels to, and contain the substance of, the cerebrum.

PICA. (*Pica*, the magpie; so named because it is said the magpie is subject to this affection.) *Picano. Malacia. Alitriphagia. Citta. Cissa.* Longing. Depraved appetite, with strong desire for unnatural food. It is very common to pregnant women and chlorotic girls, and by some it is said in men who labour under suppressed hemorrhoids.

PICUA. (*Picus*, pitch.) The common or red fir or pitch tree is so termed. The cones, branches, and every part of the tree, affords the common resin called frankincense. See *Terebinthina vulgaris*.

PICURUM. See *Pechurum*.

PICUS ECHIOIDES. (From *πιςος*, bitter, and *echioides*, from *εχis*, a viper, and *ωσις*, resemblance.) The systematic name of the common ox-tongue; the leaves are frequently used as a pot-herb by the country people, who esteem it good to relax the bowels.

PICTORIUS. (From the *Picts*, who were subject to this disease.) Applied to a species of colic. It should be rather called colica pictorum, the painter's colic, because from their use of lead they are much afflicted with it.

PIESTRUM. (From *πιεω*, to press.) An instrument to compress the head of a dead fœtus, for its more easy extraction from the womb.

PIE-NUT. Earth-nut. Ground-nut. The bulbous root of the *Hunium bulbocastanum* of Linnaeus. *Pig-nuts*, so called because that animal is very fond of them, and will dig with its snout to some depth for them, are of a dirty brown colour, a little bigger than a hazel-nut, and flavoured like the chestnut. They are said to possess a styptic

quality, and are deemed serviceable in laxity of the kidneys.

PIGMENTUM. (From *pingo*, to paint.) Pigment. This name is given by anatomists to a mucous substance found in the eye, which is of two kinds. *The pigment of the iris*, is that which covers the anterior and posterior surface of the iris, and gives the beautiful variety of colour in the eyes. *The pigment of the choroid membrane*, is a black or brownish mucus which covers the anterior surface of the choroid membrane, contiguous to the retina and the interior surface of the ciliary processes.

PILA HYSTRICIS. The bezoar hystricis.

PILA MARINA. A species of alcyonium found on sea-coasts amongst wrack. It is said to kill worms, and when calcined to be useful in scrophula.

Piles. See *Hemorrhoids*.

Pile-wort. See *Chaledonium minus*.

PILI COXEI. The hair of the head, eyebrows, and eyelids, are so termed, because they grow *in utero*.

PILI POSTEI. The hair which grows from the surface of the body after birth is so termed, in contradistinction to that which appears before birth; as the hair of the head, eyebrows, and eyelids.

PILLOLLA. (From *pilus*, hair, because its leaves are hairy.) *Auricula muris. Myosotis.* Mouse-ear. This common plant, *Hieracium pilocella* of Linnaeus, contains a bitter lactescent juice, which has a slight degree of astringency. The roots are more powerful than the leaves. They are very seldom used in this country.

Pills, aloetic, with myrrh. See *Pilule aloeticæ cum myrrha*.

Pills, compound aloetic. See *Pilule aloes composite*.

Pills, compound gamboge. See *Pilule gambogice composite*.

Pills, compound galbanum. See *Pilule galbanice composite*.

Pills, compound squill. See *Pilula scilla composite*.

Pills of iron with myrrh. See *Pilula ferri cum myrrha*.

Pills, soap, with opium. See *Pilule saponis cum opio*.

Pills, mercury. See *Pilule hydrargyri*.

Pills, submuriate of mercury. See *Pilula hydrargyri submuriatæ*.

PILULA. A pill. A small round form of medicine the size of a pea. The consistence of pills is best preserved by keeping the mass in bladders, and occasionally moistening it. In the direction of masses to be thus divided, the proper consistence is to be looked for at first, as well as its preservation afterwards, for if the mass then become hard and dry it is unfit for that division for which it was originally intended; and this is in many instances such an objection to the form that it is doubtful whether for the purposes of the pharmacopœia the

greater number of articles had not better be kept in powder, and their application to the formation of pills, left to extemporaneous direction.

PILULÆ ALOES COMPOSITÆ. Compound aloetic pills. "Take of Spike-aloe, powdered, an ounce; extract of gentian, half an ounce; oil of caraway, forty minims; syrup, as much as is sufficient." Beat them together, until they form an uniform mass. From fifteen to twenty-five grains prove moderately purgative and stomachic.

PILULÆ ALOES CUM MYRRHÆ. Aloetic pills with myrrh. "Take of spike-aloe, two ounces; saffron, myrrh, of each an ounce; syrup, as much as is sufficient." Powder the aloe and myrrh separately; then beat them all together until they form an uniform mass. From ten grains to a scruple of this pill, substituted for the *pilula Ruff*, prove stomachic and laxative, and are calculated for delicate females, especially where there is uterine obstruction.

PILULÆ CAMBOGIÆ COMPOSITÆ. Compound gamboge pills. "Take of gamboge, powdered, extract of spike aloe, powdered, compound cinnamon powder, of each a drachm; soap, two drachms." Mix the powders together; then having added the soap, beat the whole together until they are thoroughly incorporated. These pills are now first introduced in the London Pharmacopœia, as forming a more active purgative pill than the *pil. aloes cum myrrha*, and in this way supplying an article very commonly necessary in practice. The dose is from ten grains to a scruple.

PILULÆ CUPRI. An excellent tonic and diaphoretic pill, which may be given with advantage in dropsical diseases, where tonics and diuretics are indicated.

PILULÆ FERRI CUM MYRRHÆ. Pills of iron and myrrh. "Take of myrrh, powdered, two drachms; subcarbonate of soda, sulphate of iron, sugar, of each a drachm." Rub the myrrh with the subcarbonate of soda; add the sulphate of iron, and rub them again; then beat the whole together until they are thoroughly incorporated. These pills answer the same purpose as the *mistura ferri composita*. The dose is from ten grains to one scruple.

PILULÆ GALBANI COMPOSITÆ. Compound galbanum pills. Formerly called *pilule gummosæ*. "Take of galbanum resin, an ounce; myrrh, sagapenum, of each an ounce and half; assafœtida gum resin, half an ounce; syrup, as much as is sufficient." Beat them together until they form an uniform mass. A stimulating antispasmodic and emmenagogue. From a scruple to half a drachm may be given three times a day in nervous disorders of the stomach and intestines, in hysterical affections and hypochondriasis.

PILULÆ HYDRARGYRI. Mercurial pills. Often from its colour called the blue pill. "Take of purified mercury, two drachms. Confection of red roses, three drachms; liquorice-root, powdered, a drachm." Rub the mercury with the confection, until the globules disappear; then add the liquorice-root, and beat the whole together, until they are thoroughly incorporated.

An alterative and antivenereal pill, which mostly acts on the bowels if given in sufficient quantity to attempt the removal of the venereal disease, and therefore requires the addition of opium. The dose is from five grains to a scruple. Three grains of the mass contain one of mercury. Joined with the squill pill, it forms an excellent expectorant and alterative, calculated to assist the removal of dropsical diseases of the chest, and asthmas attended with visceral obstruction.

PILULÆ HYDRARGYRI SUBMURIATÆ. Pills of submuriate of mercury. "Take of submuriate of mercury, precipitated sulphuret of antimony, of each a drachm; guaiacum gum resin, powdered, two drachms." Rub the submuriate of mercury first with the precipitated sulphuret of antimony, then with the guaiacum gum resin, and add as much copaiba as may be requisite to give the mass a proper consistence. This is intended as a substitute for the famed Plummer's pill. It is exhibited as an alterative in a variety of diseases, especially cutaneous eruptions, pains of the venereal or rheumatic kind, cancerous and schirrous affections, and chronic ophthalmia. The dose is from five to ten grains. In about five grains of the mass there is one grain of the submuriate of mercury.

PILULÆ SAPONIS CUM OPIO. Pills of soap and opium. Formerly called *pilule saponacæ*. "Take of hard opium, powdered, half an ounce; hard soap, two ounces." Beat them together until they are thoroughly incorporated. The dose is from three to ten grains. Five grains of the mass contain one of opium.

PILULÆ SCILLÆ COMPOSITÆ. Compound squill pills. "Take of squill root fresh, dried and powdered, a drachm; ginger-root, powdered, hard soap, of each three drachms; ammoniacum, powdered, two drachms." Mix the powders together; then beat them with the soap, adding as much syrup as may be sufficient to give a proper consistence.

An attenuant, expectorant, and diuretic pill, mostly administered in the cure of asthma and dropsy. The dose is from ten grains to a scruple.

PILUS. (Πίλος, wool carded.) The short hair which is found all over the body. See *Capillus*.

PIMENTA. (From *Ramienta*. Sp. pepper.) *Piper caryophyllatum*. Cocculi

*indi aromatici. Piper chape. Amomum pimenta. Caryophyllus aromaticus. Caryophyllus Americanus. Piper odoratum Jamaicense. Jamaica pepper, or allspice. The berries of the Myrtus pimenta of Linnæus:—**foribus trichotomo-paniculatis, foliis oblongo-lanceolatis.* This spice, which was first brought over for dietetic uses, has been long employed in the shops as a succedaneum to the more costly oriental aromatics: it is moderately warm, of an agreeable flavour, somewhat resembling that of a mixture of cloves, cinnamon, and nutmegs. Both pharmacopœias direct an aqueous and spirituous distillation to be made from these berries; and the Edinburgh College orders the *oleum essentielle piperis Jamaicensis.*

PIMENTO. See *Pimenta.*

Pimpinel. The *Anagallis arvensis* is sometimes so called. See *Anagallis.*

Pimpinel, water. See *Becabunga.*

PIMPINELLA. (*Quasi bipinnella* or *bipennula*, from the double penate order of its leaves.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria.* Order, *Digynia.* *Pimpinella.*

2 The systematic name of the *Pimpinella alba.* *Pimpinella nostras.* Several species of *pimpinella* were formerly used officinally; but the roots which obtain a place in the *materia medica* of the Edinburgh Pharmacopœia, are those of the Burnet saxifrage, the *Pimpinella saxifraga* of Linnæus.—*foliis pinnatis, foliolis radicalibus subrotundis, summis linearibus.* They have an unpleasant smell; and a hot, pungent, bitterish taste: they are recommended by several writers as a stomachic: in the way of gargle, they have been employed for dissolving viscid mucus, and to stimulate the tongue when that organ becomes paralytic.

PIMPINELLA ALBA. A variety of the *pimpinella magna*, whose root is indifferently used with that of the greater *pimpinell*, called *radix pimpinelle nigra.* See also *Pimpinella.*

PIMPINELLA ANISUM. The systematic name of the anise plant. See *Anisum.*

PIMPINELLA ITALICA. The root which bears this name in some pharmacopœias is that of the *Sanguisorba officinalis* of Linnæus. It is now fallen into disuse.

PIMPINELLA MAGNA. The systematic name of the greater *pimpinella.* See *Pimpinella nigra.*

PIMPINELLA NIGRA. The root of this plant, *Pimpinella magna* of Linnæus, has been lately extolled in the cure of erysipelatous ulcerations, tinea capitis, rheumatism, and other diseases.

PIMPINELLA NOSTRAS. See *Pimpinella.*

PIMPINELLA SAXIFRAGA. The systematic name of the Burnet saxifrage. See *Pimpinella.*

PINASTELLUM. (From *pinus*, the pine-tree; so called because its leaves resemble those of the pine-tree.) Hog's fennel. See *Peucedanum.*

PINEA. The stone-pine. The young and fresh fruit of this plant, *Pinus pined* of Linnæus, are eaten in some countries in the same manner as almonds here, either alone, or mixed with sugar. They are nutritive, demulcent, and laxative.

PINEAL GLAND. (Called *pineal* from *pineæ*, a pine-apple, from its supposed resemblance to that fruit.) *Glandula pinealis.* *Conarium.* A small heart-like substance, about the size of a pea, situated immediately over the corpora quadrigemina, and hanging from the *thalami nervorum opticorum* by two crura or peduncles. Its use is not known. It was formerly supposed to be the seat of the soul.

Pins-apple. See *Ananas.*

Pine-thistle. See *Carlina gummiifera.*

PINEUS FURSANS. See *Ricinus major.*

PINGUEDO. (From *pinguis*, fat.) Fat. See *Fat.*

PINGUICULA. (From *pinguis*, fat, so called because its leaves are fat to the touch.) *Sanicula montana.* *Sanicula eboracensis.* *Viola palustris.* *Liparis. Cucullata.* *Dodecatheon Plinii.* Butter-wort. Yorkshire fanicle. The remarkable unctuousity of this plant, *Pinguicula vulgaris* of Linnæus, has caused it to be applied to chaps, and as a pomatum to the hair. Decoctions of the leaves in broths are used by the common people in Wales as a cathartic.

PINHONES INDICI. See *Ricinus major.*

Pink, Indian. See *Spigelia.*

PINNA. (*Pinna*, a wing.) The name of the lateral and inferior part of the nose, and the broad part of the ear.

PINNACULUM. (Dim. of *pinna*, a wing.) A pinnacle. A name of the uvula from its shape.

PINUS. The name of a genus of plants in the Linnæan system. Class, *Monoecia.* Order, *Monadelphica.* The pine-tree.

PINUS ABIES. The Norway spruce fir, which affords the *pix Burgundica.* See *Burgundy pitch.*

PINUS BALSAMEA. The systematic name of the tree which affords the Canada balsam. See *Balsamum Canadense.*

PINUS LARYX. The systematic name of the tree which gives us the agaric and Venice turpentine. See *Agaricus albus*, and *Terebinthina veneta.*

PINUS PICCA. The systematic name of the silver fir. See *Terebinthina communis.*

PINUS PINA. The systematic name of the stone-pine tree. See *Pinea.*

PINUS STYLVESTIS. The systematic name of the Scotch fir. See *Pix liquida.*

PIPER. (From *πιπτε*, to concoct,

because by its heat it assists digestion.) Pepper. The name of a genus of plants in the Linnæan system. Class, *Diandria*. Order, *Trigynia*.

PIPER ALBUM. *Leucopiper*. See *Piper nigrum*.

PIPER BRASILIANUM. *Guinea pepper*.

PIPER CALEUTICUM. *Guinea pepper*.

PIPER CARYOPHYLLATUM. *Jamaica pepper*.

PIPER CAUDATUM. See *Cubeba*.

PIPER CUBEDA. The plant whose berries are called cubeba. See *Cubeba*.

PIPER DECORTICATUM. White pepper.

PIPER FAVASCI. The clove-berry tree.

PIPER GUINEENSE. See *Piper indicum*.

PIPER HISPANTICUM. See *Piper indicum*.

PIPER INDICUM. *Capaicum*. *Lada chilli*. *Cape melago*. *Solanum arene*. *Silvastrum Plinii*. *Piper Brazilianum*. *Piper Guineense*. *Piper Caleuticum*. *Piper Hispanicum*. *Piper Lusitanicum*. Cayenne pepper. Guinea pepper. This species of pepper is obtained from the *Capaicum anuum* of Linnæus:—*caule herbaceo, pedunculis solitariis*. As an aromatic of the stimulant kind, it is efficacious in some paralytic and gouty cases, or to promote excitement where the bodily organs are languid and torpid.

Piper indicum. Guinea or Indian pepper.

PIPER JAMAICENSE. See *Pimenta*.

PIPER LONGUM. *Macropiper*. *Acapathii*. *Cutu tripali*. *Pimphilum*. Long pepper. *Piper longum* of Linnæus:—*foliis cordatis petiolatis sessilibusque*. The berries or grains of this plant are gathered while green, and dried in the heat of the sun, when they change to a blackish or dark grey colour. They possess precisely the same qualities as the *piper indicum*, only in a weaker degree.

PIPER LUSITANICUM. See *Piper indicum*.

PIPER MURALE. See *Illecebra*.

PIPER NIGRUM. *Melanopiper*. *Molagocodi*. *Lada*. *Piper aromaticum*. Black pepper. This species of pepper is obtained in the East Indies, from the *Piper nigrum* of Linnæus:—*foliis ovatis septemnerviis glabris, petiolis simplicissimis*. Its virtues are similar to those of the other peppers. The black and white pepper are both obtained from the same tree, the difference depending on their preparation and degrees of maturity.

PIPERITIS. (From *piper*, pepper, so called because its leaves and roots are biting like pepper to the taste.) The herb dittany or lepidium.

PIQUETI RECEPTACULUM. The thoracic duct.

PIRAMIDALIA CORPORA. See *Corpora pyramidalia*.

PISIFORME OS. The fourth bone of the first row of the carpus.

Pismire. See *Formica*.

Piso-a-bed. See *Taraxacum*.

PISSASPHALTUS. (From *πῖσσα*, pitch, and *ασφαλτος*, bitumen.) The thicker kind of rock oil.

PISTACIA. (*Πιστία*, supposed to be a Syrian word.) The name of a genus of plants in the Linnæan system. Class, *Diccia*. Order, *Pentandria*.

PISTACIA LENTISCUS. The systematic name of the tree which affords the mastick. See *Mastiche*.

PISTACIA NUX. *Pistachio-nut*. An oblong pointed nut, about the size and shape of a filbert, including a kernel of a pale greenish colour, covered with a yellow or greenish skin. It is the produce of a large tree, the *Pistacia vera* of Linnæus:—*foliis imparipennatis, foliis subovatis recurvis*. *Pistachio-nuts* have a sweetish unctuous taste, resembling that of sweet almonds, and, like the latter, afford an oil, and may be formed into an emulsion.

PISTACIA TEREBINTHUS. The systematic name of the tree which gives out the Cyprus turpentine. See *Chio turpentine*.

PISTACIA VARA. The systematic name of the tree which affords the *nucis pistacie*. See *Pistacia nux*.

Pistachio-nut. See *Pistacia nux*.

PISTOLOCHIA. (From *πιστος*, faithful, and *λοχια*, parturition, so called because it was thought to promote delivery.) Birth wort.

PITCH. *Pix*. *Pix sicca*. The juice of a species of fir, extracted by incisions made in the bark of the tree. It is sometimes used as a detergent by surgeons.

Pitch, Burgundy. See *Pix arida*.

Pitch, Jews. See *Bitumen judaicum*.

Pitch-tree. See *Picea*.

PITTACIUM. (From *πῖττα*, pitch.) A pitch plaster.

PITTOTA. (From *πῖττα*, pitch.) Medicines in which pitch is the principal ingredient.

PITUITA. Phlegm, or viscid and glutinous mucus.

PITUITARY GLAND. *Glandula pituitaria*. A gland situated within the cranium, between a duplicature of the dura mater, in the sella turcica of the sphenoid bone.

PITUITARY MEMBRANE. *Membrana pituitaria*. Schneiderian membrane. The mucus membrane that lines the nostrils and sinuses, communicating with the nose, is so called, because it secretes the mucus of those parts, to which the ancients have assigned the name of *pituita*.

PITYRIASIS. (From *πυρρος*, bran, so named from its branny-like appearance.) A genus in the second order, or scaly diseases, of Dr. Willan's cutaneous diseases. The pityriasis consists of irregular patches of small thin scales, which repeatedly form and separate, but never collect into crusts, nor are attended with redness

or inflammation, as in the lepra and scaly tetter. Dr. Willan distinguishes pityriasis from the porrigo of the Latins, which has a more extensive signification, and comprehends a disease of the scalp, terminating in ulceration; whereas the former is, by the best Greek authors, represented as always dry and scaly. Thus, according to Alexander and Paulus, pityriasis is characterised by "the separation of slight furfureous substances from the surface of the head, or other parts of the body, without ulceration." Their account of this appearance is conformable to experience; and the two varieties of it which they have pointed out may be denominated, *Pityriasis capitis*, and *Pityriasis versicolor*.

Pityriasis capitis, when it affects very young infants, is termed by nurses the dandriff. It appears at the upper edge of the forehead and temples, as a slight whitish scurf set in the form of a horse-shoe; on other parts of the head there are large scales, at a distance from each other, flat and semipellucid. Sometimes, however, they nearly cover the whole of the hairy scalp, being close together, and imbricated. A similar appearance may take place in adults; but it is usually the effects of lepra, scaly tetter, or some general disease of the skin.

Elderly persons have the *pityriasis capitis* in nearly the same form as infants; the only difference is, that this complaint in old people occasions larger exfoliations of the cuticle.

The *pityriasis versicolor* chiefly affects the arms, breast, and abdomen. It is diffused very irregularly; and being of a different colour from the usual skin colour, it exhibits a singular chequered appearance. These irregular patches, which are at first small, and of a brown or yellow hue, appear at the scrobiculus cordis, about the mammaræ, clavicles, &c. Enlarging gradually, they assume a tessellated form; in other cases they are branched, so as to resemble the foliaceous lichens growing on the bark of trees; and sometimes, when the discolouration is not continuous, they suggest the idea of a map being distributed on the skin like islands, continents, peninsulas, &c. All the discoloured parts are slightly rough, with minute scales, which soon fall off, but are constantly replaced by others. This scurf, or scalliness, is most conspicuous on the sides and epigastric region. The cuticular lines are somewhat deeper in the patches than on the contiguous parts; but there is no elevated border, or distinguishing boundary between the discoloured part of the skin, and that which retains its natural colour. The discolouration rarely extends over the whole body. It is strongest and fullest round the umbilicus, on the breasts,

and sides; it seldom appears in the skin over the sternum, or along the spine of the back. Interstices of proper skin colour are more numerous, and largest at the lower part of the abdomen and back, where the scales are often small, distinct, and a little depressed. The face, nates, and lower extremities are least affected; the patches are found upon the arms, but mostly on the inside, where they are distinct and of different sizes.

The *pityriasis versicolor* is not a cuticular disease; for when the cuticle is abraded from any of the patches, the sallow colour remains as before in the skin or rete mucosum. This singular appearance is not attended with any internal disorder, nor with any troublesome symptom, except a little itching or irritation felt on getting into bed, and after strong exercise, or drinking warm liquors. There is in some cases a slight exanthema, partially distributed among the discoloured patches; and sometimes an appearance like the lichen pileus; but eruptions of this kind are not permanent, neither do they produce any change in the original form of the complaint. The duration of the *pityriasis versicolor* is always considerable. Dr. Willan has observed its continuance in some persons for four, five, or six years. It is not limited to any age or sex. Its causes are not pointed out with certainty. Several patients have referred it to fruit taken in too great quantities; some have thought it was produced by eating mushrooms; others by exposure to sudden alterations of cold and heat. In some individuals, who had an irritable skin, and occasionally used violent exercise, the complaint has been produced, or at least much aggravated, by wearing flannel next to the skin. It is likewise often observed in persons who had resided for a length of time in a tropical climate.

PIX ARIDA. Formerly called *Pix Burgundica*. The prepared resin of the *Pinus abies* of Linnaeus:—*foliis solitariis subtragonis acutiusculis distichis, ramis infra nudis coni scylindraceis*. It is of a solid consistence, yet somewhat soft, of a reddish brown colour, and not disagreeable smell. It is used externally as a stimulant in form of plaster in catarrh, pertussis, and dyspnoea.

PIX BURGUNDICA. *Burgundy pitch*. See *Pix arida*.

PIX LIQUIDA. *Tar*. Tar is produced from the *Pinus sylvestris* of Linnaeus:—*foliis geminis rigidis, conis ovato conicis longitudine, foliorum subgeminis basi rotundatis*, by cutting it into pieces, which are enclosed in a large oven constructed for the purpose. It is well known for its economical uses. Tar-water, or water impregnated with the more soluble parts of tar, was sometime ago a very fashion-

able remedy in a variety of complaints, but is in the present practice fallen into disuse.

PLACEDO. I will please : an epithet given to any medicine adapted more to please than benefit the patient.

PLACENTA. (From *πλάκωσις*, a cake, so called from its resemblance to a cake.) The placenta is a circular, flat, vascular, and apparently fleshy substance, different in its diameter in different subjects, but usually extending about six inches, or upwards, over about one-fourth part of the outside of the ovum in pregnant women. It is more than one inch in thickness in the middle, and becomes gradually thinner towards the circumference from which the membranes are continued. The placenta is the principal medium by which the communication between the parent and child is preserved ; but though all have allowed the importance of the office which it performs, there has been a variety of opinions on the nature of that office, and of the manner in which it is executed.

That surface of the placenta which is attached to the uterus by the intervention of the connecting membrane, is lobulated and convex ; but the other, which is covered with the amnion and chorion, is concave and smooth, except the little eminences made by the blood-vessels. It is seldom found attached to the same part of the uterus in two successive births ; and, though it most frequently adheres to the anterior part, it is occasionally fixed to any other, even to the os uteri, in which state it becomes a cause of a dangerous hæmorrhage at the time of parturition. The placenta is composed of arteries and veins, with a mixture of pulpy or cellular substance. Of these there are two orders, very curiously interwoven with each other. The first is a continuation of those from the funis, which ramify on the internal surface of the placenta, the arteries running over the veins, which is a circumstance peculiar to the placenta ; and then, sinking into its substance, anastomose and divide into innumerable small branches. The second order proceeds from the uterus ; and these ramify in a similar manner with those from the funis, as appears when a placenta is injected from those of the parent. The veins, in their ramifications, accompany the arteries as in other parts. There have been many different opinions with respect to the manner in which the blood circulates between the parent and child, during its continuance in the uterus. For a long time it was believed that the intercourse between them was uninterrupted, and that the blood propelled, by the powers of the parent pervaded, by a continuance of the same force, the vascular system of the fœtus ; but repeated attempts having been made

without success to inject the whole placenta, funis and fœtus, from the vessels of the parent, or any part of the uterus, from the vessels of the funis, and it is now generally allowed, that the two systems of vessels in the placenta, one of which may be called maternal, the other fetal, are distinct. It is also admitted, that the blood of the fœtus is, with regard to its formation, increase, and circulation, unconnected with, and totally independent of the parent ; except that the matter by which the blood of the fœtus is formed must be derived from the parent. It is thought that which has probably undergone some preparatory changes in its passage through the uterus, is conducted by the uterine or maternal arteries of the placenta to some cells or small cavities, in which it is deposited ; and that some part of it, or something secreted from it, is absorbed by the fetal veins of the placenta, and by them conveyed to the fœtus for its nutriment. When the blood which circulates in the fœtus requires any alteration in its qualities, or when it has gone through the course of the circulation, it is carried by the arteries of the funis to the placenta, in the cells of which it is deposited, and then absorbed by the maternal veins of the placenta, and conducted to the uterus, whence it may enter the common circulation of the parent. Thus it appears, according to the opinion of Harvey, that the placenta performs the office of a gland, conveying air, or secreting the nutritious juices from the blood brought from the parent by the arteries of the uterus, and carried to the fœtus by the veins of the funis, in a manner probably not unlike to that in which milk is secreted and absorbed from the breasts. The veins in the placenta are mentioned as the absorbents, because no lymphatic vessels have yet been found in the placenta or funis ; nor are there any nerves in these parts ; so that the only communications hitherto discovered between the parent and child, is by the sanguineous system. The proofs of the manner in which the blood circulates between the parent and child are chiefly drawn from observations made upon the funis. When it was supposed that the child was supplied with blood in a direct stream from the parent, it was asserted that, on the division of the funis, if that part next to the placenta was not secured by a ligature, the parent would be brought into extreme danger by the hæmorrhage which must necessarily follow. But this opinion, which laid the foundation of several peculiarities in the management of the funis and placenta, is proved not to be true : for, if the funis be compressed immediately after the birth of the child, and whilst the circulation in it is going on, the arteries between the part

compressed and the child throb violently, but those between the compression and the placenta have no pulsation; but the vein between the part compressed and the placenta swells, and that part next to the fœtus becomes flaccid. But if, under the same circumstances, the funis be divided, and that part next the child be not secured, the child would be in danger of losing its life by the hæmorrhage; yet the mother would suffer no inconvenience if the other part was neglected. It is moreover proved, that a woman may die of an hæmorrhage occasioned by a separation of the placenta, and the child be nevertheless born, after her death, in perfect health. But if the placenta be injured, without separation, either by the rupture of the vessels which pass upon its inner surface, or in any other way, the child being deprived of its proper blood, would perish, yet the parent might escape without injury. See also *Secundines*.

PLACENTULA. (Dim. of *placenta*.) A small placenta.

PLACAROTIS. (From *πλαστος*, moist, flaccid.) A fungous and flaccid tumour within the eye-lid.

PLANTAGO. (From *planta*, the sole of the feet; so called from the shape of its leaves, or because its leaves lie upon the ground and are trodden upon.) 1. The name of a genus of plants in the Linnæan system. Class, *Tetrandria*. Order, *Monogynia*. The plantain.

2. The pharmacopœial name of the *centinervia*. *Polyneuron*. *Plantago latifolia*. *Plantago major* of Linnæus:—*foliis ovatis glabris, scapo tereti, spica fœsculis imbricatis*. This plant was retained until very lately in the *materia medica* of the Edinburgh College, in which the leaves are mentioned as the pharmaceutical part of the plant; they have a weak herbaceous smell, an austere, bitterish, subsaline taste; and their qualities are said to be refrigerant, attenuating, subtyptic, and diuretic.

PLANTAGO LATIFOLIA. See *Plantago*.

PLANTAGO MAJOR. The systematic name of the broad-leaved plantain. See *Plantago*.

PLANTAGO PSYLLIUM. The systematic name of the branching plantain. See *Psyllium*.

Plantain. See *Plantago*.

Plantain-tree. See *Musa paradisiaca*.

PLANTARIS. (*Plantaris*, sc. *musculus*. From *planta*, the sole of the foot, to which it belongs.) *Tibialis gracilis* vulgo *plantaris* of Winslow. *Extensor tarsi minor* vulgo *plantaris* of Douglas. A muscle of the foot, situated on the leg, that assists the soleus, and pulls the capsular ligament of the knee from between the bones. It is sometimes, though seldom, found wanting on both sides. This long and slender muscle, which is situated under the *gastrocnemius externus*, arises, by a thin

fleshy origin, from the upper and back part of the outer condyle of the os femoris. It adheres to the capsular ligament of the joint, and, after running obliquely downwards and outwards, for the space of three or four inches, along the second origin of the *gastrocnemius internus*, and under the *gastrocnemius externus*, terminates in a long, thin, and slender tendon, which adheres to the inside of the tendo Achillis, and is inserted into the inside of the posterior part of the os calcis. This tendon sometimes sends off an aponeurosis that loses itself in the capsular ligament, but it does not at all contribute to form the aponeurosis that is spread over the sole of the foot, as was formerly supposed, and as its name would seem to imply. Its use is to assist the *gastrocnemii* in extending the foot. It likewise serves to prevent the capsular ligament of the knee from being pinched.

PLANUM OS. (*Planus*, soft, smooth; applied to a bone whose surface is smooth or flat.) The papyraceous or orbital portion of the ethmoid bone was formerly so called.

Plaster, ammoniacum. See *Emplastrum ammoniaci*.

Plaster, ammoniacum, with mercury. See *Emplastrum ammoniaci cum hydrargyro*.

Plaster, blistering fly. See *Emplastrum lytta*.

Plaster, compound Galbanum. See *Emplastrum Galbani compositum*.

Plaster, compound pitch. See *Emplastrum picis compositum*.

Plaster, common. See *Emplastrum cumini*.

Plaster, lead. See *Emplastrum plumbi*.

Plaster, mercurial. See *Emplastrum hydrargyri*.

Plaster of opium. See *Emplastrum opii*.

Plaster, resin. See *Emplastrum resine*.

Plaster, soap. See *Emplastrum saponis*.

Plaster, wax. See *Emplastrum ceræ*.

PLATA. (From *πλάτυς*, broad.) The shoulder-blade.

PLATIASMUS. (From *πλάτυς*, broad.) A defect in the speech in consequence of too broad a mouth.

PLATINA. A metal so called. (The name platina was given to this metal by the Spaniards from the word *plata*, which signifies silver in their language, by way of comparison with that metal, whose colour it imitates; or from the river *Plata*, near which it is found.) It exists in nature, only in a metallic state in small grains, combined with iron, copper, plumbago, &c. The largest mass of which we have heard, is one of the size of a pigeon's egg, in the possession of the Royal Society of Bergara. It is found in the parishes of Novita and Citaria, north from Choco in Peru, and near Carthagera in South America. It was unknown in Europe before the year 1748. Don Antonio Ulloa then gave the first in-

formation concerning its existence in the narrative of his voyage with the French academicians to Peru.

Properties.—Platina purified from all extraneous mixture is of a white colour, intermediate between that of silver and tin. It is the hardest of all metals. Its specific gravity being from 20.6 to 23, makes it by far the heaviest body known. It is hard, malleable, ductile, laminable like gold; but to what degree is not yet ascertained. We have seen platina drawn into a wire of a smaller diameter than the two-thousandth part of an inch. It is the most infusible of all the metals. It cannot be melted (in a considerable quantity at least,) by the most violent heat of our furnaces, but may be fused by the heat of a burning lens, or by the assistance of oxygen gas. When pure, its parts may be made to combine or weld, by hammering in a white heat; a property confined to this metal and iron. It suffers no alteration from the action of air; neither water, the earths, nor the saline-terrene substances have any power of re-action upon it. Potash acts upon it at high temperature. It is not oxidated when exposed red-hot to the air, for a very long time. It may however be oxidated by the galvano-electric spark, and by the nitro-muriatic acid. This acid dissolves it and assumes first a yellow and afterwards a deep orange colour. The solution tinges animal substances with a dark colour, and may be decomposed by alkalies and by muriate and nitrate of ammonia, which have no effect on solutions of gold. No other acid exercises any action upon it. The oxid of platina which is a triple compound, consisting of ammonia, muriatic acid, and platina, may be reduced by the action of a violent heat. Platina combines with phosphorus and sulphur with considerable facility. It unites with the greater number of the metals by fusion. Of these alloys, that with copper is the most valuable, as it is ductile, susceptible of a fine polish, and does not tarnish on exposure to air; the rest of them are very little known.

Method of obtaining Platina.—The processes most commonly employed to obtain pure platina, are as follows:

1. Take equal parts of platina in grains, and acidulous tartre of potash, put the mixture into a well luted crucible, and expose it for two hours to a violent heat. The platina fuses, but it becomes brittle and whiter than platina is in common; then expose it to a very strong heat under a muffle, by which means all the arsenic combined with it will be disengaged; and the platina remain behind in a malleable state.

2. Platina may likewise be obtained pure, by decomposing the nitro-muriatic

solution of common platina, by muriate of ammonia, heating the precipitate intensely, and stamping it when of a white heat into one mass: or, by assisting the fusion with a stream of oxygen gas.

3. Jannetty's process, which is considered as the best, and which is generally used for obtaining malleable platina, is as follows:

Triturate common Platina with water, to wash off every contaminating matter that water can carry away. Mix the platina with about one-fifth part of arsenious acid and one-fifteenth part of potash; putting the whole in a proper crucible, in the following manner: having well heated the crucible and the furnace receiving it, put in one-third of the mixture, apply to this a strong heat, and add one-third more: after a renewed application of heat, throw in the last portion. After a thorough fusion of the whole, cool and break the mass. Then fuse it a second time, and, if necessary, even a third time, till it ceases to be magnetic. Break it into small pieces, and melt those pieces in separate crucibles, and in portions of a pound and a half of the platina to each crucible, with an equal quantity of arsenious acid and half a pound of potash. After cooling the contents of the different crucibles in a horizontal position, in order to have them throughout of equal thickness, heat them under a muffle to volatilize the arsenious acid, and maintain them in this state, without increase of heat, for the space of six hours. Heat them, next, in common oil, till the oil shall have evaporated to dryness. Then immerse them in nitric acid, boil them in water, heat them to redness in a crucible, and hammer them into a dense mass. They are now fit to be heated in a naked fire, and hammered into bars for the purposes of commerce.

Mr. Richter directs, in order to purify platina, and render it malleable, the following process: dissolve platina of commerce in nitro muriatic acid, and let fall into this solution potash, until a precipitate begins to appear; then add a solution of sulphate of potash, till the whole is precipitated. Wash the precipitate till the water that passes do not change its colour by adding to it prussiate of potash. Dry the precipitate, and mix with it 1.5 times its weight of soda, freed from its water of crystallization; press it into a crucible, but not so as to fill it, heat it gradually, and raise the heat till it fuses.

PLATYCORIA. (From *πλατος*, broad, and *κορη*, the pupil of the eye.) An enlarged pupil.

PLATYPHTHALMUS. (From *πλατος*, broad, and *οφθαλμος*, the eye; so called because it is used by women to enlarge the eye.) Antimony.

PLATYPHYLLUM. (From *πλατύς*, broad, and *φύλλον*, a leaf.) Broad leaved.

PLATYSMA MYOIDES. (From *πλατύς*, broad, *μύς*, a muscle, and *οἶδος*, resemblance.) *Musculus cutaneus* of Winslow. *Quadratus genæ vel Latissimus colli* of Douglas. *Latissimus colli* of Albinus. *Quadratus genæ, seu tetragonus* of Winslow, and *thoraco-maxillæ facialis* of Dumas. A thin muscle on the side of the neck, immediately under the skin, that assists in drawing the skin of the cheek downwards; and when the mouth is shut, it draws all that part of the skin to which it is connected below the lower jaw upwards.

PLECTANÆ. (From *πλεκτω*, to fold.) The horns of the uterus.

PLECTRUM. (From *πλεττα*, to strike, so named from their resemblance to a drum-stick.) A drum-stick. The styloid process of the os petrosus and the uvula.

PLEROSIS. See *Plethora*.

PLESMONE. See *Plethora*.

PLETHORA. (From *πληθω*, to fill.) *Plesmone*. *Plerosis*. An excessive fullness of vessels, or a redundancy of blood.

PNEUMONIA. See *Pulmonia*.

PLEURA. (*Πνευγ.*) A membrane which lines the internal surface of the thorax, and covers its viscera. It forms a great process, the mediastinum, which divides the thorax into two cavities. Its use is to render the surface of the thorax moist by the vapour it exhales. The cavity of the thorax is every where lined by this smooth and glistening membrane, which is in reality two distinct portions or bags, which by being applied to each other laterally, form the septum called mediastinum: this divides the cavity into two parts, and is attached posteriorly to the vertebrae of the back; and anteriorly to the sternum. But the two laminae of which this septum is formed, do not every where adhere to each other; for at the lower part of the thorax they are separated, to afford a lodgment to the heart; and at the upper part of the cavity they receive between them the thymus gland. The pleura is plentifully supplied with arteries and veins from the internal mammary, and the intercostals. Its nerves, which are very inconsiderable, are derived chiefly from the dorsal and intercostal nerves. The surface of the pleura, like that of the peritonæum and other membranes lining cavities, is constantly bedewed with a serous mixture, which prevents adhesions of the viscera. The mediastinum, by dividing the breast into two cavities, obviates many inconveniences to which we should otherwise be liable. It prevents the two lobes of the lungs from compressing each other when we lie on one side, and consequently contributes to

the freedom of respiration, which is disturbed by the least pressure on the lungs. If the point of a sword penetrates between the ribs into the cavity of the thorax, the lungs on that side cease to perform their office, because the air being admitted through the wound, prevents the dilatation of that lobe, while the other lobe, which is separated from it by the mediastinum, remains unhurt, and continues to perform its functions as usual.

PLEURITIS. (From *πλινγν*, the membrane lining the lungs.) Pleurisy, or inflammation of the pleura. A species of pneumonia of Cullen. See *Pneumonia*. In some instances the inflammation is partial, or affects one place in particular, which is commonly on the right side; but in general, a morbid affection is communicated throughout its whole extent. The disease is occasioned by exposure to cold, and by all the causes which usually give rise to all inflammatory complaints; and it attacks chiefly those of a vigorous constitution and plethoric habit. In consequence of the previous inflammation, it is apt at its departure to leave behind a thickening of the pleura, or adhesions to the ribs and intercostal muscles, which either lay the foundation of future pneumonic complaints, or render the patient more susceptible of the changes in the state of the atmosphere than before.

It comes on with an acute pain in the side, which is much increased by making a full inspiration, and is accompanied by flushing in the face, increased heat over the whole body, rigors, difficulty of laying on the side affected, together with a cough and nausea, and the pulse is hard, strong, and frequent, and vibrates under the finger when pressed upon, not unlike the tense string of a musical instrument. If blood is drawn, and allowed to stand for a short time, it will exhibit a thick sily or buffy coat on its surface. If the disease be neglected at its onset, and the inflammation proceeds with great violence and rapidity, the lungs themselves become affected, the passage of the blood through them is stopped, and the patient is suffocated; or from the combination of the two affections, the inflammation proceeds on to suppuration, and an abscess is formed. The prognostic in pleurisy must be drawn from the severity of the symptoms. If the fever and inflammation have run high, and the pain should cease suddenly, with a change of countenance and a sinking of the pulse, great danger may be apprehended; but if the heat and other febrile symptoms abate gradually, if respiration is performed with greater ease and less pain, and a free and copious expectoration ensues, a speedy recovery may be expected.

The appearances on dissection are much the same as those mentioned under the head of pneumonia, viz. an inflamed state of the pleura, connected with the lungs, having its surface covered with red vessels, and a layer of coagulated lymph lying upon it, adhesions too, of the substance of the lungs to the pleura. Besides these, the lungs themselves are often found in an inflamed state, with an extravasation either of blood or coagulated lymph in their substance. Tubercles and abscesses are likewise frequently met with.

PLEUROCOLLISIS. (From *πλευρα*, the pleura, and *κολλησις*, to adhere.) An adhesion of the pleura to the lungs or some neighbouring part.

PLEURODYNIA. (From *πλευρα*, and *δυνι*, pain.) A pain in the side, from a rheumatic affection of the pleura.

PLEURO-PNEUMONIA. (From *πλευρα*, and *πνευμονια*, an inflammation of the lungs.) An inflammation of the lungs and pleura.

PLEUROTHORAX. (From *πλευρα*, the pleura, and *θερος*, upright, and *πναι*, to breathe.) A pleurisy in which the patient cannot breathe without keeping his body upright.

PLEUROSTHONOS. (From *πλευρα*, the side, and *συνα*, to stretch.) A spasmodic disease in which the body is bent to one side.

PLEXUS. (From *plector*, to plait or knit.) A net-work of vessels. The union of two or more nerves is also called a plexus.

PLEXUS CARDIACUS. The cardiac plexus of nerves is the union of the eighth pair of nerves and great sympathetic.

PLEXUS CHOROIDES. The choroid plexus is a net-work of vessels situated in the lateral ventricles of the brain.

PLEXUS PAMPINIFORMIS. The plexus of vessels about the spermatic chord.

PLEXUS PULMONICUS. The pulmonary plexus is formed by the union of the eighth pair of nerves with the great sympathetic.

PLEXUS RETICULARIS. A net-work of vessels under the fornx of the brain.

PLICA. (From *plico*, to entangle. This disease is commonly distinguished by the adjective *Polonica*, it being peculiar to the inhabitants of Poland and Lithuania.) *Helotie*. *Kolto*. *Rhopulosis*. *Plica Polonica*. *Trichoma*. Plaited hair. A disease of the hairs, in which they become long and coarse, and matted and glued into inextricable tangles. It is peculiar to Poland and Tartary, and generally appears during the autumnal season.

PLICARIA. (From *plico*, to entangle; so called because its leaves are entangled together in one mass.) Wolf's-claw, or club moss.

PLUMBAG. PANTHOR. The fourfold bandage.

Plum, Malabar. See *Malabar plum*.

PLUMBAGO. (From *plumbum*, lead; so called because it is covered with lead-coloured spots.) 1. Lead-wort. See *Perisicaria*.

2. An ore of a shining blue black colour, a greasy feel, and tuberculated when fractured. It is by many erroneously taken for molybdena, from which it is easily distinguished by its fracture, that of the latter being always lamellated.

PLUMBAGO EUPHRA. The systematic name of the tooth-wort. See *Dentaria*.

PLUMBI CARBONAS. Subcarbonate of lead called ceruse, or white lead. It is sometimes employed medicinally in form of powder and ointment, to children whose skin is fretted. It should however be cautiously used, as there is great reason to believe that complaints of the bowels of children originate from its absorption.

Plumbi, erydum semivitreum. See *Litharge*.

PLUMBI SUPERACETAS. See *Supracetas plumbi*.

PLUMBI SUBCARBONAS. See *Plumbi carbonas*.

PLUMBUM. See *Lead*.

PLUMBUM CANDIDUM. See *Stannum*.

PLUMBUM CINEREUM. Bismuth.

PLUMBUM NIGRUM. Black lead.

PLUMBUM RUBRUM. The philosopher's stone.

PLUMBUM USTUM. Burnt lead.

PLUMBERI PILULE. Plummer's pills. A composition of calomel antimony, guaiacum and balsam of copaivi. See *Pilule hydrargyri submuriatis*.

PLUMS. Three sorts of plums are ranked amongst the articles of the materia medica; they are all met with in the gardens of this country, but the shops are supplied with them moderately dried, from abroad.

1. The *pruna brignolencia*; The Brignole plum, or prune, brought from Brignole in Provence; it is of a reddish yellow colour, and has a very grateful, sweet, subacid taste. 2. The *pruna Gullica*; the common or French prune. 3. The *pruna damascena*, or damson. See *Damson*. All these fruits possess the same general qualities with the other summer fruits. The pruneloes, in which the sweetness has a greater mixture of acidity than in the other sorts, are used as mild refrigerants in fevers and other hot indispositions. The French prunes and damsons are the most emollient and laxative; they are often taken by themselves to gently move the belly, where there is a tendency to inflammations. Decoctions of them afford a useful basis for laxative or purgative mixtures, and the pulp in substance for electuaries.

PLUNKET'S CANCER REMEDY. "Take crows' foot, which grows in low grounds, one handful; dogs' fennel, three sprigs; both well pounded; crude brimstone in powder, three middling thimbles full; white arsenic, the same quantity; incorporated all in a mortar, and made into small balls the size of a nutmeg, and dried in the sun." These balls must be powdered and mixed with the yolk of an egg, and laid over the sore or cancer upon a piece of pig's bladder, or stripping of a calf when dropped, which must be cut to the size of the sore, and smeared with the yolk of an egg. This must be applied cautiously to the lips or nose lest any part of it get down; nor is it to be laid on too broad on the face, or too near the heart; nor to exceed the breadth of half-a-crown; but elsewhere as far as the sore goes. The plaster must not be stirred until it drops off of itself, which will be in a week. Clean bandages are often to be put on.

PNEUMATIC APPARATUS. The discovery of æriform fluids has, in modern chemistry, occasioned the necessity of some peculiar instruments, by means of which those substances may, in distillations, solutions, or other operations, be caught, collected, and properly managed. The proper instruments for this are styled the pneumatic apparatus. Any kind of air is specifically lighter than any liquid; and therefore, if not decomposed by it, rises through it in bubbles. On this principle rests the essential part of the apparatus, adapted to such operations. Its principal part is the pneumatic trough, which is a kind of reservoir for the liquid, through which the gas is conveyed and caused to rise, and is filled either with water or with quicksilver. Some inches below its brim an horizontal shelf is fastened, in dimension about half or the third part of the trough, and provided on its foremost edge with a row of holes, into which, from underneath, short-necked funnels are fixed. The trough is filled with water sufficient to cover the shelf, to support the receivers, which being previously filled with water or mercury, are placed invertedly, their open end turned down upon the above-mentioned holes, through which afterwards the gases, conveyed there and directed by means of the funnels, rise in the form of air-bubbles.

In some cases the trough must be filled with quicksilver, because water decomposes some kinds of air by absorbing their basis. The price and specific gravity of that metal make it necessary to give to the quicksilver-trough smaller dimensions. It is either cut in marble, or made of wood well joined. The late Karsten has contrived an apparatus, which, to the advantage of saving room, adds that of great convenience.

To disengage gases, retorts of glass, either common or tubulated, are employed, and placed in a sand-bath, or heated by the fire of a lamp. Earthen, or coated glass retorts, are put in the naked fire. If necessary, they are joined with a metallic or glass conveying pipe. When, besides the æriform, other fluids are to be collected, the middle or intermediate bottle finds its use; and to prevent, after cooling, the rising of the water from the trough into the disengaging vessels, the tube of safety is employed. For the extrication of gases taking place in solutions, for which no external heat is required, the bottle called disengaging bottle, or proof, may be used. For receivers, to collect the disengaged airs, various cylinders of glass are used, whether graduated or not, either closed at one end, or open at both; and, in this last case, they are made air tight by a stopper fitted by grinding. Besides these, glass bells and common bottles are employed.

To combine with water, in a commodious way, some gases that are only gradually and slowly absorbed by it, the glass apparatus of Parker is serviceable.

PNEUMATOCELE. (From *πνευμα*, wind, and *cele*, a tumour.) Any species of hernia, that is distended with flatus.

PNEUMATOPHALUS. (From *πνευμα*, wind, and *ομφαλος*, the navel.) A flatulent, umbilical hernia.

PNEUMATOSIS. (From *πνευματος*, to inflate.) *Emphysema*. Windy swelling. A genus of disease in the class *cachexia*, and order *intumescentia* of Cullen, known by a collection of air in the cellular texture under the skin, rendering it tense, elastic, and crepitating. The species of pneumatosis are:

1. *Pneumatosis spontanea*, without any manifest cause.
2. *Pneumatosis traumatica*, from a wound.
3. *Pneumatosis venenata*, from poisons.
4. *Pneumatosis hysterica*, with hysteria.

PNEUMONIA. (From *πνευμων*, a lung.) *Pneumonitis*. *Peripneumonia*. *Peripneumonia vera*. Inflammation of the lungs. A genus of disease in the class *pyrexia* and order *phlegmasia* of Cullen; characterised by pyrexia, difficult respiration, cough, and a sense of weight and pain in the thorax. The species of pneumonia, according to the above nosologist, are,

1. *Peripneumonia*. The pulse not always hard, but sometimes soft: an obtuse pain in the breast: the respiration always difficult; sometimes the patient cannot breathe, unless in an upright posture; the face swelled, and of a livid colour; the cough for the most part moist, frequently bloody.

2. *Pleuritis*. The pulse hard: a pungent pain in one side, aggravated during the time of inspiration; an uneasiness

when lying on one side; a very painful cough, dry in the beginning of the disease, afterwards moist, and frequently bloody. See *Pleuritis*.

With respect to pneumonia, the most general cause of this inflammation, is the application of cold to the body, which gives a check to the perspiration, and determines a great flow of the blood to the lungs. It attacks principally those of a robust constitution and plethoric habit, and occurs most frequently in the winter season and spring of the year; but it may arise in either of the other seasons, when there are sudden vicissitudes from heat to cold.

Other causes, such as violent exertions in singing, speaking, or playing on wind instruments, by producing an increased action of the lungs, have been known to occasion peripneumony. Those who have laboured under a former attack of this complaint, are much predisposed to returns of it.

The true peripneumony comes on with an obtuse pain in the chest or side, great difficulty of breathing, (particularly in a recumbent position, or when lying on the side affected,) together with a cough, dryness of the skin, heat, anxiety, and thirst. At the first commencement of the disease the pulse is usually full, strong, hard, and frequent; but in a more advanced stage it is commonly weak, soft, and often irregular. In the beginning, the cough is frequently dry and without expectoration; but in some cases it is moist even from the first, and the matter spit up is various both in colour and consistence, and is often streaked with blood.

If relief is not afforded in time, and the inflammation proceeds with such violence as to endanger suffocation, the vessels of the neck will become turgid and swelled; the face will alter to a purple colour; an effusion of blood will take place into the cellular substance of the lungs, so as to impede the circulation through that organ, and the patient will soon be deprived of life.

If these violent symptoms do not arise, and the proper means for carrying off the inflammation have either been neglected, or have proved ineffectual, although adopted at an early period of the disease, a suppuration may ensue, which event is to be known by frequent slight shiverings, and an abatement of the pain and sense of fullness in the part, and by the patient not being able to lay on the side which was affected, without experiencing great uneasiness.

When peripneumony proves fatal, it is generally by an effusion of blood taking place into the cellular texture of the lungs, so as to occasion suffocation, which usually happens between the third and seventh

day; but it may likewise prove fatal, by terminating either in suppuration or gangrene.

When it goes off by resolution, some very evident evacuation always attends it; such as a great flow of urine, with a copious sediment, diarrhoea, a sweat diffused over the whole body, or a hæmorrhage from the nose; but the evacuation which most frequently terminates the complaint, and which does it with the greatest effect, is a free and copious expectoration of thick white or yellow matter, slightly streaked with blood, and by this the disease is carried of generally in the course of ten or twelve days.

Our opinion as to the event, is to be drawn from the symptoms which are present. A high degree of fever, attended with delirium, great difficulty of breathing, acute pain, and dry cough, denote great danger: on the contrary an abatement of the febrile symptoms, and of the difficulty of breathing, and pain, taking place on the coming on of a free expectoration, or the happening of any other critical evacuation, promises fair for the recovery of the patient. A termination of the inflammation in suppuration, is always to be considered as dangerous.

On dissection, the lungs usually appear inflamed, and there is often found an extravasation, either of blood, or of coagulable lymph in their cellular substance. The same appearances likewise present themselves in the cavity of the thorax and within the pericardium. The pleura, connected with the lungs, is also in an inflamed state, having its surface every where crowded with red vessels. Besides these, abscesses are frequently found in the substance of the lungs, as likewise tubercles and adhesions to the ribs are formed. A quantity of purulent matter is often discovered also in the bronchiae.

PNEUMOPLEURITIS. (From *πνευμα*, the lungs, and *πleuritis*, an inflammation of the pleura.) An inflammation of the lungs and pleura.

PNIGALUM. (From *πνιγω*, to suffocate.) The night-mare. A disorder in which the patient appears to be suffocated.

PNIX. (From *πνιγω*, to suffocate.) An hysterical sense of suffocation.

PODAGRA. (From *πους*, the foot, and *αλγω*, a taking, or seizure.) *Febris podagrica. Arthritis. Dolor podagricus.* The gout. A genus of disease in the class *pyrexia* and order *phlegmania* of Cullen; known by pyrexia, pain in the joints, chiefly of the great toe, and especially the hands and feet, returning at intervals: previous to the attack, the functions of the stomach are commonly disturbed. Species:

1. *Podagra regularis. Arthritis podagra.*

Arthritis rachialgica. *Arthritis estiva* of Sauvage. The regular gout.

2. *Podagra atonica.* *Arthritis melancholica, hiemalis, chlorotica, and asthmatica* of Sauvage. The atonic gout.

3. *Podagra retrograda.* The retrocedent gout.

4. *Podagra aberrans.* Misplaced or wandering gout. See *Arthritis*.

PODAGRARIA. (From *podagra*, the gout; so called because it was thought to expel the gout.) Gout-weed. A species of *agropodium*.

PODONIPTUM. (From *πους*, a foot, and *νίψω*, to wash, or bathe.) A bath for the feet.

PODOPHYLLUM. (From *πους*, a foot, and *φύλλον*, a leaf; so named from its shape.) A species of wolf's-bane.

PODOTHÉCA. (From *πους*, a foot, and *τίθημι*, to put.) A shoe, or stocking. An anatomical preparation, consisting of a kind of shoe of the scarf-skin, with the nails adhering to it, taken from a dead subject.

POISON. *Venenum.* That substance which, when applied externally, or taken into the human body, uniformly effects such a derangement in the animal economy as to produce disease, may be defined a poison. It is extremely difficult, however, to give a definition of a poison; and the above is subject to great inaccuracy. Poisons are divided, with respect to the kingdom to which they belong, into animal, vegetable, mineral, and halituous, or aerial.

Poisons are only deleterious in certain doses; for the most active, in small doses, form the most valuable medicines. There are, nevertheless, certain poisons, which are really such in the smallest quantity, and which are never administered medicinally; as the poison of hydrophobia, the plague. There are likewise substances which are innocent when taken into the stomach, but which prove deleterious when taken into the lungs, or when applied to an abraded surface; thus carbonic acid is continually swallowed with fermented liquors, and thus the poison of the viper may be taken with impunity; whilst inspiring carbonic acid kills, and the poison of the viper inserted into the flesh, often proves fatal.

Several substances also act as poisons when applied either externally or internally, as arsenic.

When a substance produces disease, not only in mankind, but in all animals, it is distinguished by the term *common poison*, as arsenic, sublimate, &c. whilst that which is poisonous to man only, or to animals, and often to one genus, is said to be a *relative poison*; thus aloes are poisonous to dogs and wolves; the phellandrium aquaticum kills horses, whilst oxen devour it greedily, and with impunity. It appears,

then, that substances act as poisonous only in regard to their dose, the part of the body they are applied to, and the subject.

Poisons enter the body in the following ways:

1. Through the œsophagus alone, or with the food.

2. Through the anus by clysters.

3. Through the nostrils.

4. Through the lungs with the air.

5. Through the absorbents of the skin, either whole, ulcerated, cut, or torn.

Poison oak. See *Rhus toxicodendron*.

POLEMONIUM. (From *Polemon*, its inventor.) Wild sage.

Poley mountain. See *Polium creticum*, and *Polium montanum*.

POLIUM. (From *πολιος*, white; so called from its white capillaments.) Poley. *Teucrium of Linnæus*; of which botanists enumerate several species.

POLIUM CRETICUM. Candy poley-mountain. *Teucrium creticum* of Linnæus. The tops and whole herb enter the antiquated compounds *mithridate* and *theriaca*. The plant is obtained from the island of Candy; has a moderately aromatic smell, and a nauseous bitter taste. It is placed amongst the aperients and corroborants.

POLIUM MONTANUM. Poley-mountain of Montpellier. This plant, *Teucrium capitatum* of Linnæus, bears the winter of our climate, and is generally substituted for the candy-species.

POLLEX. The thumb, or great-toe.

POLYCHÆRSTUS. (From *πολυς*, much, and *χρῆσις*, useful.) Many virtues, or uses. Applied to many medicines from their extensive usefulness.

POLYDIPSIA. (From *πολυς*, much, and *διψα*, thirst.) Excessive thirst. A genus of disease in the class *locales* and order *dyseorexis* of Cullen. It is mostly symptomatic of fever, dropsy, excessive discharges, or poisons.

POLYGALA. (From *πολυς*, much, and *γάλα*, milk; so named from its abundance of its milky juice.) 1. The name of a genus of plants in the Linnæan system. Class, *Diadelphia*. Order, *Octandria*.

2. The pharmacopœial name of *Polygala*. Common milk-wort. The root of this plant, *Polygala vulgaris* of Linnæus, is somewhat similar in taste to that of the seneka, but much weaker. The leaves are very bitter, and a handful of them, infused in wine, is said to be a safe and gentle purge.

POLYGALA AMARA. This is a remarkably bitter plant, and, though not used in this country, promises to be as efficacious as those in greater repute. It has been given freely in phthisis pulmonalis, and, like other remedies, failed in producing a cure: yet, as a palliative, it claims attention. Its virtues are balsamic, demulcent, and corroborant.

POLYGALE SENECAL. The systematic name of the rattlesnake milk-wort. See *Seneca*.
POLYGALE VULGARIS. The systematic name of the common milk-wort. See *Polygala*.

POLYGONATUM. (From *πολυ*, many, and *γων*, a joint; so named from its numerous joints or knots.) *Sigillum Solomonis*. Solomon's seal. The *Convallaria polygonatum* of Linnæus.

POLYGONUM. (From *πολυ*, many, and *γων*, a joint; so named from its numerous joints.) The name of a genus of plants in the Linnæan system. Class, *Octandria*. Order, *Trigynia*. Knot-grass.

POLYGONUM AVICULARE. The systematic name of the knot-grass. See *Centummodia*.

POLYGONUM BACCIFERUM. A species of equisetum, or horse-tail.

POLYGONUM BISTORTA. The systematic name of the officinal bistort. See *Bistorta*.

POLYGONUM DIVARICATUM. The systematic name of the eastern buck-wheat plant. See *Buck-wheat*, eastern.

POLYGONUM FASOPHYUM. The systematic name of the buck-wheat. See *Buck-wheat*.

POLYGONUM HYDROPIPER. The systematic name of the poor man's pepper. See *Hydropiper*.

POLYGONUM LATIFOLIUM. Common knot-grass.

POLYGONUM MAS. Common knot-grass.

POLYGONUM MINUS. Rupture-wort.

POLYGONUM PERSICARIA. The systematic name of arsmart. See *Persicaria*.

POLYGONUM SILENEOIDES. Parsley break-stone.

POLYPODIUM. (From *πολυ*, many, and *πους*, a foot; so called because it has many roots.) The name of a genus of plants in the Linnæan system. Class, *Cryptogamia*. Order, *Filices*. Fern, or polypody.

POLYPUS. (From *πολυ*, many, and *πους*, a foot; from its sending off many ramifications, like legs.) 1. The name of a genus of zoophytes.

2. In Cullen's Nosology, a synonym with *sarcoma*. A kind of tumour, which is generally narrow where it originates, and then becomes wider, somewhat like a pear; which most commonly is met with in the nose, uterus, or vagina; and which received its name from an erroneous idea, that it is usually had several roots, or feet, like polypi, or zoophytes.

Polypi vary from each other according to the different causes that produce them, and the alterations that happen in them. Sometimes a polypus of the nose is owing to a swelling of the pituitary membrane, which swelling may possess a greater or less space of the membrane, as also its cellular substance, and may affect either one or both nostrils. At other times, it arises from an ulcer produced by a caries of some of the

bones which form the internal surface of the nostrils. Polypuses are sometimes so soft, that upon the least touch they are lacerated, and bleed; at other times they are very compact, and even scirrhus. Some continue small a great while; others increase so fast as, in a short time, to push out at the nostrils, or extend backwards towards the throat. Le Dran mentions, that he has known them fill up the space behind the uvula, and, turning towards the mouth, have protruded the fleshy arch of the palate so far forwards as to make it parallel with the third *dentes molares*. There are others which, though at first free from any malignant disposition, become afterwards carcinomatous, and even highly cancerous. Of whatever nature the polypus is, it intercepts the passage of the air through the nostril, and when large, forces the *septum narium* into the other nostril, so that the patient is unable to breathe, unless through the mouth. A large polypus pressing in like manner upon the spongy bones, gradually forces them down upon the maxillary bones, and thus compresses and stops up the orifice of the *ductus lachrymalis*; nor is it impossible for the sides of the *canalis nasalis* to be pressed together. In which case the tears, having no passage through the nose, the eye is kept constantly watering, and the *sacculus lachrymalis*, not being able to discharge its contents, is sometimes so much dilated as to form what is called a flat *scrofula*. The above writer has seen instances of polypuses so much enlarged as to force down the *ossa palati*.

The polypus of the uterus is of three kinds, in respect to situation. It either grows from the fundus, the inside of the cervix, or from the lower edge of the os uteri. The first case is the most frequent, the last the most uncommon. Polypi of the uterus are always shaped like a pear, and have a thin pedicle. They are almost invariably of that species which is denominated fleshy, hardly ever being scirrhus, cancerous or ulcerated.

The coagulable substance which is found in the cavities of the heart of those who are some time in *articulo mortis*, is improperly called a polypus.

POLYSARCHIA. (From *πολυ*, much, and *σαρξ*, flesh.) *Polyomatia*. *Obesitas*. *Corpulencia*. *Steutites* of Vogel. Troublesome corpulency, or fatness. A genus of disease in the class *cachexie* and order *intumescencie* of Cullen.

POLYSONATIA. (From *πολυ*, much, and *συνα*, a body.) See *Polyarchia*.

POLYSPASTUM. (From *πολυ*, much, and *σπασ*, to draw.) A forcible instrument for reducing luxations.

POLYTRICHUM. (From *πολυ*, many, and *τριχ*, air; so called from its resemblance to a woman's hair, or because, in ancient times, women used to dye the hair

with it, to keep it from shedding.) *Polytricon*. 1. The name of a genus of plants in the Linnæan system. Class, *Cryptogamia*. Order, *Musci*.

2. The systematic name of the golden maidenhair. See *Adiantum aureum*.

POLYTRICUM COMMUNE. The systematic name of the golden maidenhair. See *Adiantum aureum*.

POLYURICA ISCHURIA. (From *πολυς*, much, and *ουρον*, urine.) Strangury, from long retention of urine.

POMACEUM. (From *pomum*, an apple.) Cider, or the fermented juice of apple.

Pomegranate. See *Granatum*.

POMPHOLYGODES. (From *πομφολυξ*, a bubble.) Urine, with bubbles on the surface.

POMPHOLYX. (From *πομφος*, a bladder.) 1. A bubble. 2. The whitish powder called tutty, a kind of zinc, which adheres to the covers of the crucibles in making brass, in the form of small bubbles. See *Tutia*.

POMPHOS. (From *πομφο*, to put forth.) A bladder, or watery pustule.

POMUM. An apple. See *Apples*.

POMUM ADAMI. (*Pomum*, an apple; so called because it was thought to have originated in consequence of a whimsical supposition that part of the forbidden apple which Adam eat, stuck in the throat, and thus became the cause.) The protuberance in the anterior part of the neck, formed by the fore-part of the thyroid cartilage.

POMUM AXORIS. A name for the solanum lycopersicum.

PONS VAROLII. *Corpus annulare*. *Processus annularis*. *Eminentia annularis*. Varolius's bridge. An eminence of the medulla oblongata, first described by Varolius. It is formed by the two exterior crura of the cerebellum becoming flattened and passing over the crura of the cerebrum.

PONTICA VINA. Acid, feculent, and tartarous wine.

PONTICUM MEL. A sort of poisonous honey.

Poorman's pepper. *Lepidium*.

Poplar. See *Populus*.

POPLES. *Ignis*. *Ignys*. The ham, or joint of the knee.

POPLITEAL ARTERY. (*Arteria poplitea*; from *poples*, the ham.) The continuation of the crural artery, through the hollow of the ham.

POPLITEUS. (*Popliteus musculus*; from *poples*, the ham.) A small triangular muscle lying across the back part of the knee-joint.

Poppy, red corn. See *Papaver erraticum*.

Poppy, white. See *Papaver album*.

POPULAGO. (From *populus*, the poplar; because its leaves resemble those of the poplar.) Marsh marigold.

POPULUS. (From *πολυς*, many; because of the multitude of its shoots.) 1. The name of a genus of plants in the Linnæan system. Class, *Diœcia*. Order, *Ociandria*.

2. The pharmacopœial name of the *Populus nigra* of Linnæus, the black poplar; called also *Egeiros*. The young buds, *scuti*, or rudiments of the leaves, which appear in the beginning of the spring; were formerly employed in an official ointment. At present they are almost entirely disregarded, though they should seem, from their sensible qualities, to be applicable to purposes of some importance. They have a yellow, unctuous, odorous, balsamic juice.

POPULUS BALSAMIFERA. See *Tucamahaca*.

POPULUS NIGRA. The systematic name of the black poplar. See *Populus*.

PORCUS. A name for the pudendum muliebree.

PORI BILIARI. The billiary pores or ducts, that receive the bile from the penicilli of the liver, and convey it to the hepatic duct. See *Liver*.

POROCLEL. (From *πορος*, a callus, and *κλελη*, a tumour.) A hard tumour of the testicle.

POROMPHALUM. (From *πορος*, a callus, and *ομφαλος*, the navel.) A hard tumour of the navel.

PORRIGO. (*A porrigendo*, from its spreading abroad.) A disease very common among children, in which the skin of the hairy part of the head becomes dry and callous, and comes off like bran upon combing the head.

PORRUM. Porret, or common leek. *Allium porrum* of Linnæus. Every part of this plant, but more particularly the root, abounds with a peculiar odour. The expressed juice possesses diuretic qualities, and is given in the cure of dropsical diseases, and calculous complaints.

PORTA. (*A portando*, because through it the blood is carried to the liver.) That part of the liver where its vessels enter.

PORTA VENA. See *Vena porta*.

PORTAIGUILLE. The acutenaculum.

PORTIO DURA. (One branch of the seventh pair of nerves is called *portio dura*, the hard portion, either from its being more firm than the other, or because it runs into the hard part of the skull; and the other the *portio mollis*, or soft portion.) Facial nerve. This nerve arises near the pons, from the crus of the brain, enters the petrous portion of the temporal bone, gives off a branch into the tympanum, which is called the chorda tympani, and then proceeds to form the *pes anserinus* on the face, from whence the integuments of the face are supplied with nerves. See *Facial nerve*.

PORTIO MOLLIS. Auditory nerve.

This nerve arises from the medulla oblongata and fourth ventricle of the brain, enters the petrous portion of the temporal bone, and is distributed on the internal ear, by innumerable branches, not only to the cochlea, but also to the membrane lining the vestibulum and semicircular canals, and is the immediate organ of hearing.

PORTLAND POWDER. A celebrated gout remedy. It consists of various bitters; principally of horehound, birthwort, of the tops and leaves of germander, ground-pine, and centaury, dried, powdered, and sifted. It is now fallen into disuse.

PORTORIUM. (From *porta*, a door; because it is, as it were, the door or entrance of the intestines.) The right orifice of the stomach.

PORTULACA. (From *porto*, to carry, and *lac*, milk; because it increases the animal milk.) 1. The name of a genus of plants in the Linnæan system. Class, *Dodecandria*. Order, *Digymia*.

2 The pharmacopœial name of the purslane. *Andrachne*. *Allium gallicum*. The plant which is so called in dietical and medical writings, is the *Portulaca oleracea* of Linnæus; it abounds with a watery and somewhat acid juice, and is often put into soups, or pickled with spices. It is said to be antiseptic and aperient.

PORTULACA OLERACEA. The systematic name of the eatable purslane. See *Portulaca*.

PORUS. A pore, or duct.

POSCA. Vinegar and water mixed.

POSSERTUM. Posset. Milk curdled with wine, treacle, or any acid.

POST BRACHIALE. (From *post*, after, and *brachium*, the arm.) The metacarpus.

POSTERIOR ANNULARIS. (*Mucellus posterior annularis*.) An external interosseal muscle of the hand, that extends and draws the ring-finger inwards.

POSTERIOR INDICIS. (*Musculus posterior indicis*.) An internal interosseal muscle of the hand, that extends the fore-finger obliquely, and draws it outwards.

POSTERIOR MEDII. An external interosseal muscle of the hand, that extends the middle finger, and draws it outwards.

POTAMOGEITON. (From *ποταμος*, a river, and *γειτον*, adjacent; so named because it grows about rivers.) The name of a genus of plants in the Linnæan system. Class, *Tetrandria*. Order, *Trygynia*.

Potash, acetate of. See *Acetas potasse*.

Potash, carbonate of. See *Carbonas potasse*.

Potash, fused. See *Potassa fusa*.

Potash, solution of. See *Liquor potasse*.

Potash, subcarbonate of. See *Potassa subcarbonas*.

Potash, subcarbonate of, solution of. See *Liquor potasse subcarbonatis*.

Potash, sulphate of. See *Potasse sulphas*.

Potash, sulphuret of. See *Potasse sulphuretum*.

Potash, super-sulphate of. See *Potasse super-sulphas*.

Potash, tartrate of. See *Potasse tartreas*.

Potash with lime. See *Potassa cum calce*.

Potash. See *Potassa*.

POTASSA. (So called from the pots, or vessels in which it was first made.) Potash. *Kali*. Vegetable alkali. This alkali may be obtained from several substances.

1. By evaporating the lixivium of the ashes of wood and some plants. This is calcined, and by this means disengaged from all the blackening principles.

The ashes are more or less rich in alkali, according to the nature of the wood which affords them. In general, hard woods contain the most. The ashes of beech afford from 11 to 13lb. per quintal, according to the experiments which have been made, in the large way; those of box afforded from 12 to 14lb. Wormwood affords a good quantity.

To extract this alkali, nothing more is necessary than to wash the ashes, and to concentrate the solution in boilers of cast iron. It is on account of the alkalies that wood-ashes are employed in the lixiviums used by laundresses, or bleachers. The use of alkali, in this case, is to combine with the fat substances, and to render them soluble in water.

Almost all the pot-ash sold in commerce for the use of our glass-houses, our soap-makers, our bleaching-grounds, &c. is fabricated in the north, where the abundance of wood admits of its being applied to this single purpose.

2. The lees of wine is almost totally converted into this alkali by combustion. This salt is called *cenures gravelles*; it has almost always a greenish colour, and considered as very pure.

3. The combustion of tartar of wine likewise affords an alkali of considerable purity. It is usually burned wrapped up in paper, in small packets, which are dipped in water, and afterwards exposed upon burning coals. In order to purify it, the residue of the combustion is dissolved in water, the solution concentrated by fire, the foreign salts separated in proportion as they precipitate, and a very pure alkali is at last obtained, which is known by the name of *salt of tartar*. To procure salt of tartar more speedily, as well as more economically, a mixture of equal parts of nitrate of pot-ash, or common nitre, and tartar, may be burned. The residue, after lixiviation affords a beautiful salt of tartar. Salt of tartar is the alkali most commonly employed for medical uses.

4. If salt-petre be fused upon charcoal, the acid is decomposed and dissipated, while the alkali remains alone and disengaged; this is called extemporaneous alkali.

When the salt has been brought to the greatest state of purity, it attracts the humidity of the air, and is resolved into a liquor. In this state it is known by the very improper name of *Oil of tartar per deliquium*.

From all these salts, which are impure and imperfect carbonates of potash, the pure alkali may be extracted.

Methods of obtaining pure Potash—It has long been a desideratum with chemists to possess a method of preparing potash in a state of absolute purity; the strong tendency which it has to combination, renders this extremely difficult. The following are the methods now made use of.

1. Bouillon la Grange's apparatus consists of several boxes of common deal. At the bottom put river-sand, which must be well washed, and over it add another stratum, but of a finer kind, and cover the whole with a cloth, besprinkled with wood-ashes. In the bottom of each box a hole is made, into which is fitted a glass tube, for the purpose of affording a passage to the liquor as it filters through the sand.

Having arranged the apparatus in this manner, take equal parts of quick-lime and pot-ash of commerce, if the lime is very caustic; but in the contrary case, it requires twenty-parts of lime to fifteen of pot-ash: put water into an iron kettle, bring it nearly to a state of ebullition, and then add lime, which, by its slaking, will bring the water to that state completely. When it is slaked, mix the pot-ash, and form of the whole a thick liquid, which must be suffered to cool a little. Then pour the mixture into the boxes, and immediately throw water over it; but, to prevent the water from making holes, when added, place over it a small board, which will rise with the water.

Care must be taken to place earthen pans, or other vessels, to receive the liquor which runs through the tubes; and, that the ley may not absorb carbonic acid from the atmosphere, the vessels must be closed with care, in such a manner as to exclude the external air. It will be necessary also to keep water always over the mixture, which must be collected till it passes tasteless from the tubes.

The liquors obtained are nearly of the same degree of strength till towards the end of the process; when they grow weak suddenly.

To evaporate the water, use should be made of cast-iron pots, beginning with the last portions, which are a little weaker; and to prevent the necessity of keeping the strongest a long time in contact with

the air when boiling, a strong ebullition is requisite. When the fluid is concentrated to a certain degree, any sulphate of potash that may be present will crystallize, and be precipitated.

To obtain dry potash, pour the concentrated liquor into a small basin, and proceed with the evaporation, till a little of it, poured on an iron plate, or a marble slab, becomes solid.

Then put the concrete potash into a jar, and pour over it very strong alcohol; the potash alone dissolves in it; the sulphate and muriate of potash, with the portions of earth and even of carbonic acid, which it obstinately retains, or which it may have acquired from the air during the evaporation, remain at the bottom of the solution. Afterwards decant the pure liquor, and distil it in a retort till it becomes colourless. It must then be evaporated in a silver basin. On cooling, it crystallizes in white laminae, which are sometimes three-tenths of an inch in length: or, instead of suffering it to crystallize, it may be evaporated to dryness.

2. Lowitz has given another method. According to this chemist, the whole of the operation for obtaining potash of the greatest purity, and without the least colour, consists in this: A lixivium of potash, freed from carbonic acid in the usual manner, is evaporated to a thick pellicle. After the cooling, the foreign salt which has crystallized is to be separated, and the evaporation of the lixivium continued in an iron pot. During this second evaporation, the pellicle of foreign salts, particularly of carbonate of potash, which continues to be formed, must be carefully taken off with an iron skimmer. When no more pellicle is formed, and the matter ceases to boil up, it is removed from the fire, and suffered to cool, continually stirred with an iron spatula. It is then to be dissolved in double the quantity of cold water, and the solution filtered, and evaporated in a glass retort, till it begins to deposit regular crystals. If the mass should consolidate ever so little by cooling, a small quantity of water is to be added, and it must be heated again to render it fluid. After the formation of a sufficient quantity of regular crystals, the fluid, which is very brown, is to be decanted, and the salt, after being suffered to drain, must be re-dissolved in the same quantity of water. The decanted fluid must be kept in a well-closed bottle, and suffered to become clear by subsiding during several days. It must then be decanted for a second evaporation and crystallization. The process must be repeated as long as the crystals afford, with the least possible quantity of water, solutions perfectly limped. These solutions are to be preserved in well-closed bottles, to defend them from the access of air.

The greatest difficulty of this process arises from the facility with which the fluid assumes a solid form. To obviate this inconvenience, a small portion of it may be concentrated to the point at which it becomes converted into a solid mass by cooling. The saturation of a lixivium considerably evaporated, may be ascertained by throwing small pieces of this mass into it during its cooling. When these are no longer dissolved, it is a proof that it is at the required point.

With regard to the foreign salts which are mixed with the potash, the greatest portion separates by crystallization, after the first evaporation. The rest is separated during the second concentration, by the continual skimming of the pellicle. The little which may remain with the potash must precipitate for want of water of solution, in a lixivium, wherein the alkali itself is no longer dissolved but by its own water of crystallization.

Remarks.—The property of alkalies, to dissolve in highly rectified alcohol, with the exclusion of every foreign salt, would afford an excellent means of obtaining potash very pure, if their mutual action did not afford a new source of impurity. For when an alkali, absolutely pure and crystallized, is dissolved in spirit of wine, even without heat, the fluid assumes a very brown colour, which becomes still deeper after being decanted from the saline mass.

The crystallization of potash is very different, accordingly as the crystals are formed with cold or heat. In the first case, the crystals obtained are octahedra in groups, which contain 0.43 water of crystallization, and excite, by their solution in water, even in the summer, a degree of cold very near the point of aqueous congelation. In the second case, very thin crystalline transparent blades of extraordinary magnitude are formed, which, by an assemblage of lines crossing each other in infinite directions, present an aggregate of cells or cavities, most commonly so perfectly closed that the vessel may be inverted without the escape of the smallest drop of lixivium, though sometimes included to the amount of an ounce or two. For this reason, it is necessary to break this fine crystallization, that the fluid may run off. The crystals present, in their regular formation, rectangular tetragonal blades, which, as they contain little water of crystallization, produce a considerable degree of heat when dissolved in water.

By exposing such alkaline crystals to a red heat, in a silver crucible, they become fused; and, after cooling, afford a mass as white as snow, extremely caustic and deliquescent.

As the crystals and the lixivium, during the length of time required to drain the salt, may frequently become charged with

a portion of carbonic acid, it is advisable, in order to avoid this inconvenience as much as possible, that the lixivium, as soon as it is brought to the requisite point of concentration, should be poured into a narrow-necked bottle, and well closed therein to crystallize. After the crystals are formed, the bottle is to be reversed without opening, and kept in a temperature rather warm until the crystals are well dried. During the winter, the liquor, after the first crystallization, continues to crystallize without being submitted to a new evaporation, provided only that it be exposed to a temperature somewhat colder than that wherein the first crystals were formed.

Properties of pure Pot-ash.—Pot-ash, in a crystallized form, consists of soft, quadrangular, compressed prisms, which are extremely caustic and deliquescent. It dissolves all soft animal matters when brought into contact with it. It liquifies by a gentle heat, and rises in fumes at high temperatures. It does not unite in a direct manner with phosphorus. It easily combines with sulphur and metallic oxids. It dissolves alumine in the humid way, and even a small quantity of silex; but it does not act on glue, or zircon, nor on magnesia, or lime. It contracts no union with barytes. It absorbs water and carbonic acid rapidly. It fuses into glass with silex, by the aid of fire. It combines with the acids, and forms salts, which in general do not yield their acid unless to barytes. It changes blue vegetable colours green, and possesses all the general properties of alkalies.

Decomposition of Potassa.

Potassa, from its analogy to ammonia, has often been conjectured to be a compound body; but nothing satisfactory had ever been proved. Its decomposition, however, has since been effected by the labours of Mr. Davy, and the discovery of its constituents will ever be regarded as the most important in chemistry. Mr. Davy was led to institute experiments, with a view to decompose the fixed alkalies by his previous discovery, that, by the powers of galvanism, the principles of bodies were separated, according to a certain law, some being attracted to the one galvanic pole, others to the other; and that the strength of these attracting forces are proportional to the energies of the opposite electricities in the galvanic circle, and to the conducting power and the degree of concentration of the substances submitted to their action.

In his first attempts, he exposed the alkali, dissolved in water, to the action of the galvanic battery, but the water alone of the solution was decomposed. Potassa, in fusion, was then placed in the galvanic circuit; a vivid light, and appearances of

the production of combustible matter were observed; on submitting the solid alkali, rendered a conductor of electricity by being very slightly moistened, to the galvanic action, these appearances were still better marked; and it was in this way Mr. Davy succeeded in effecting the complete decomposition.

"A small piece of pure potash, which had been exposed a few seconds to the atmosphere, so as to give conducting power to the surface, was placed upon an insulated disc of platina, connected with the negative side of the battery in a state of intense activity; and a platina wire, communicating with the positive side, was brought in contact with the upper surface of the alkali. Under these circumstances a vivid action was observed to take place. The potash began to fuse at both its points of electrization. There was a violent effervescence at its upper surface; at the lower, or negative surface, there was no liberation of elastic fluid; but small globules having a high metallic lustre, and being precisely similar in visible characters to quicksilver, appeared; some of which burnt, with an explosion and bright flame, as soon as they were formed, and others remained, and were merely tarnished, and finally covered with a white film, which formed on their surfaces. These globules numerous experiments soon shewed to be the substance I was in search of, and a peculiar inflammable principle the basis of potash."

The platina, Mr. Davy found, had no share in its production; as it was equally produced when even charcoal, and other metals, were used for completing the electrical circuit. The phenomena too were quite independent of the presence of air.

As, in the decomposition of compound substances by galvanism, Mr. Davy had found that when combustible bases had been developed at the negative side, oxygen is evolved at the positive side, it was reasonable to conclude that this happened in the above experiment, and that the effervescence observed at the part of the potassa in immediate contact with the positive wire, was due to the disengagement of oxygen gas. He accordingly found, performing the experiments in glass tubes, so as to collect the aerial product given out at the positive surface, that this, on the most delicate examination, proved to be pure oxygen; and, unless an excess of water was present, no gas was evolved from the negative surface, nothing being produced there but the inflammable globules.

By these analytic experiments, therefore, potassa is proved to be a compound of a peculiar substance, highly inflammable, and having a metallic lustre, with

oxygen. And this Mr. Davy soon confirmed, by synthetic experiments.

He found that the metallic lustre of this inflammable substance immediately became destroyed in the atmosphere, and a white crust formed upon it which was pure potassa. This was soon dissolved by attracting humidity from the air; a new quantity was formed on the surface, until the whole disappeared, and had formed a saturated alkaline solution. When the globules were placed in tubes containing atmospheric air, or oxygen gas, an absorption of oxygen took place, and a crust of alkali was formed. When they were strongly heated, confined in given portions of oxygen, a rapid combustion, with a brilliant white flame, was produced, and they were converted into a white and solid mass, which was found to be potassa; oxygen was absorbed, and nothing emitted which affected the purity of the residual air. From subsequent experiments, Mr. Davy endeavoured to determine the proportions of these elements. From the products of the combustion of the base, compared with the quantity of oxygen absorbed, he inferred taking the mean of two experiments, that potassa consists of 86.1 of base, and 13.9 of oxygen. From the results of the decomposition of water by the base, the proportions indicated were 84 of base and 16 of oxygen. The mean of these will be 85 and 15.

The decomposition of potassa may be effected, without difficulty, with a galvanic battery of not very high powers. One of 100 plates, of six inches square, is sufficient.

An important confirmation of the discovery is that which has been received from an experiment performed by Guy-Lussac and Thenard. It occurred to these chemists, that potassa might be decomposed by causing a substance to act on it having a strong attraction for oxygen. They put clean iron filings into a gun-barrel, bent so that the filings in the curvature could be raised to a sufficient heat, by the barrel being placed across a furnace. With one extremity of it, a tube is connected containing solid potassa. When the iron filings in the barrel are brought to a white heat, the alkali in the tube which has been previously kept cold by a freezing mixture, is melted by applying heat to it by a portable furnace; and it is allowed to run through, by a small aperture, upon the iron filings. It suffers decomposition, the iron attracts its oxygen, and the inflammable base is sublimed to the other end of the tube, which is kept cold, so as to condense it; a tube of safety, containing a little mercury, being connected with the extremity, to allow of the disengagement of any aerial matter, and at the

same time exclude the air. Hydrogen gas is disengaged during the process, which appears to be derived from the decomposition of water contained in the alkali, and the result is most successful when the alkali is in the driest state. The base of the alkali is obtained in the form of brilliant laminae, adhering to the side of the gun-barrel. There is also found in the curvature where the slings were contained, an alloy of it with iron. According to Mr. Davy, who performed the experiment with success, the base thus obtained is rather heavier than that procured by electricity, probably from containing a little iron.

To the matter discovered by these researches, as the base of potassa, Mr. Davy has given the name of

POTASSIUM;

the termination of this name being that assigned to metals, and this substance being presumed to be metallic.

Potassium, at the temperature of 60° of Fahrenheit, appears in the form of small globules, possessing the metallic lustre and opacity, and having the general appearance of quicksilver, so that by the eye it is not easy to distinguish between them. At this temperature, however, it is rather imperfectly fluid, so that it does not easily recover its globular form. At 70° it is more liquid and mobile; and at the temperature of 100° is so completely so, that different globules can be easily run into one. At 50° it becomes a soft and malleable solid, which has the lustre of polished silver, and at 32° it becomes harder and brittle, displaying, when broken, a crystallized texture. Though so fusible, it is not very volatile, but requires a temperature approaching a red-heat to convert it into vapour. It condenses unchanged. It is a perfect conductor of electricity, and is also an excellent conductor of heat.

It is the lightest substance known; it is therefore lighter than ether, or alcohol. Mr. Davy found that it did not even sink in naphtha twice distilled, the specific gravity of which was about 770. Its specific gravity he estimates, compared with that of water, is as 6 to 10, at 62° of Fahrenheit. In its solid form it is rather heavier, but still, when cooled to 40°, it swims in distilled naphtha.

Its chemical relations are not less singular than its physical properties.

It combines with oxygen slowly and without flame, at all temperatures below that of its volatilization; but, at this temperature, combustion of it takes place, the heat is intense, and the light white and vivid. It appears to combine with different proportions of oxygen; at least an oxide of it, in a lower state of oxydation than potassa, can be obtained. Potassium

inflames in other gases which can afford oxygen, as the oxy muriatic acid gas. It is so liable to oxygenation, that it is difficult to preserve it unchanged; the best method is to keep it in naphtha; for, although it does not sink in that fluid, it is enveloped by a film of it, which protects it from the action of the air.

When heated in hydrogen gas, at a degree below its point of vaporization, it diminishes in volume, and the gas explodes with the production of alkaline fumes, when it is allowed to pass into the air, a portion of the potassium appears, therefore, to have been dissolved; but, by cooling, this is in a great measure deposited, as the gas loses its property of detonating, spontaneously.

Potassium, when brought into contact with water, decomposes it with great violence, an explosion is produced, with flame, and potassa is formed. Placed on ice, it instantly burns with a bright flame, melting the ice. So strong is the action of this substance on water, that it discovers, by the decomposition it produces, the smallest quantity of water in other liquids, as in alcohol, or in ether. Potassium, thrown into solutions of the mineral acids, inflames and burns on the surface, and the compound of potassa, with the acid employed, is formed.

Potassium combines with the primary inflammables. When brought in contact with phosphorus, under exposure to air, both bodies become fluid, burn, and phosphate of potassa is formed. When the experiment is made under naphtha, so as to exclude the air, they combine, and form a compound less fusible than either of its ingredients. It has the lustre of polished lead.

When potassium is brought in contact with sulphur in fusion, under the vapour of naphtha, they combine rapidly with the evolution of heat and light, and a grey substance, in appearance like sulphuret of iron, is formed, a little sulphuretted hydrogen being evolved. When the union is effected in the atmosphere, inflammation takes place.

With the metals, potassium enters readily into combination. With mercury it produces some singular results. When one part of it is added to eight or ten parts of mercury in volume, at 60° Fahrenheit, they constantly unite and form a substance exactly like mercury in colour, but which has less coherence; for small portions of it appear like flattened spheres. When a globule is made to touch a globule about twice as large, they combine with considerable heat; the compound is fluid at the temperature of its formation; but, when cold, it appears as a solid metal, similar in colour to silver. If the quantity

of the basis of potassa is still further increased, so as to be about one-thirtieth the weight of mercury, the amalgam increases in hardness, and becomes brittle. The solid amalgam, in which the basis is in the smallest proportion, seems to consist of about one part in weight of base, and seventy parts of mercury, and is very soft and malleable.

When these compounds are exposed to air, they rapidly absorb oxygen; potassa, which deliquesces, is formed, and, in a few minutes, the mercury is found pure and unaltered. When a globule of the amalgam is thrown into water, it rapidly decomposes it, with a hissing noise; potassa is formed, pure hydrogen is disengaged, and the mercury remains free.

The fluid amalgam of mercury and potassium dissolves all the metals; and, in this state of union, mercury acts on platinum and iron.

When potassium is heated with gold, or silver, or copper, in a close vessel of pure glass, it rapidly acts upon them; and when the compounds are thrown into water, the fluid is decomposed, potassa formed, and the metals appear to be separated unaltered. It reduces the metallic oxides when heated with them, and when the potassium is in excess, it combines with the reduced metal. In consequence of this property, it decomposes flint glass and green glass, reducing the metallic oxides they contain, forming potassa, which dissolves the glass. At a red heat, it acts even on the purest glass, attracting part of the oxygen of the alkali in the glass, and forming the substance which is potassium in the first degree of oxygenation.

The preparations of this alkali that are used in medicine, are

1. Potassa fusa.
2. Liquor potassæ.
3. Potassa cum calce.
4. Subcarbonas potassæ.
5. Carbonas potassæ.
6. Sulphas potassæ.
7. Super-sulphas potassæ.
8. Tartas potassæ.
9. Acetas potassæ.
10. Citras potassæ.
11. Oxymurias potassæ.
12. Sulphuretum potassæ.

POTASSA CARBONAS. See *Carbonas potassæ*.

POTASSA CUM CALCE. Potash with lime. *Calx cum kali puro. Causticum commune fortius. Lapis infernalis sive septicus.* "Take of solution of potash, three pints; fresh lime, a pound." Boil the solution of potash down to a pint, then add the lime, previously slaked by the addition of water, and mix them together intimately. This is in common use with surgeons as a caustic, to produce ulcerations, and to open abscesses.

POTASSA FUSA. Fused potash. *Kali purum. Alkali vegetabile fixum cæsium.* "Take of solution of potash, a gallon." Evaporate the water, in a clean iron pot, over the fire, until, when the ebullition has ceased, the potash remains in a state of fusion; pour it upon a clean iron plate, into pieces of convenient form. This preparation of potash is violently caustic, destroying the living animal fibre with great energy.

POTASSÆ ACETAS. See *Acetas potassæ*.

POTASSÆ SUBCARBONAS. Subcarbonate of potash, formerly called *Kali preparatum. Sal absinthii. Sal Tartari. Sal plantarum.* "Take of impure potash, powdered, three pounds; boiling water, three pints and a half." Dissolve the potash in water, and filter; then pour the solution into a clean iron pot, and evaporate the water over a moderate fire until the liquor thickens; then let the fire be withdrawn, and stir the liquor constantly with an iron rod, until the salt concretes into granular crystals.

A purer subcarbonate of potash may be prepared in the same manner from tartar, which must first be burnt, until it becomes ash-coloured.

This preparation of potash is in general use to form the citrat of potash for the saline draughts. A scruple is generally directed to be saturated with lemon-juice. In this process, the salt which is composed of potash and carbonic acid is decomposed. The citric acid having a greater affinity for the potash than the carbonic, seizes it and forms the citrat of potash, whilst the carbonic acid flies off in the form of air. The subcarbonate of potash possesses antacid virtues, is an antidote against white arsenic, and may be exhibited with advantage in convulsions and other spasms of the intestines arising from acidity, in calculous complaints, leucorrhœa, scrophula, and aphthous affections. The dose is from ten grains to half a drachm.

POTASSÆ SULPHAS. Formerly called *Kali vitriolatum. Alkali vegetabile vitriolatum. Sal de duobus. Arcanum duplicatum. Sal polychrestus. Nitrum vitriolatum. Tartarum vitriolatum.* "Take of salt which remains after the distillation of nitric acid, two pounds; boiling water, two gallons." Mix them, that the salt may be dissolved; next add as much carbonate of potash as may be requisite for the saturation of the acid; then boil the solution, until a pellicle appears upon the surface, and, after straining, set it by, that crystals may form. Having poured away the water, dry the crystals on bibulous paper. Its virtues are cathartic, diuretic, and deobstruent; with which intentions it is administered in a great variety of diseases, as constipation, suppression of the

lochia, fevers, icterus, dropsies, milk tumours, &c. The dose is from one scruple to half an ounce.

POTASSÆ SULPHURETUM. See *Sulphuretum potassæ*.

POTASSÆ SUPERSULPHAS. Super-sulphate of potash. "Take of the salt which remains after the distillation of nitric acid, two pounds; boiling water, four pounds." Mix them together, so that the salt may be dissolved, and strain the solution; then boil it until a pellicle appear upon the surface, and set it by, that crystals may form. Having poured away the water, dry these crystals upon bibulous paper.

POTASSÆ TARTRAS. Tartrate of potash, formerly called *Kali tartariatum*. *Tartarum solubile*. *Tartaris tartariatum*, *Sal vegetabilis*. *Alkali vegetabile tartariatum*. "Take of subcarbonate of potash, a pound; supertartrate of potash, three pounds; boiling water a gallon." Dissolve the subcarbonate of potash in the water; next add the supertartrate of potash, previously reduced to powder, gradually, until bubbles of gas shall cease to arise. Strain the solution through paper, then boil it until a pellicle appear upon the surface, and set it by, the crystals may form. Having poured away the water, dry the crystals upon bibulous paper. Diuretic, deobstruent, and eccoprotic virtues are attributed to this preparation.

POTATON, COMMON. The root of the *Solanum tuberosum* of Linnæus. *Solanum esculentum*. *Kippa*. *Kelengu*. *Pupus Americus*. *Puppus Americanus*. *Convolvus Indicus*. A native of Peru. An extremely nutritious and wholesome vegetable.

POTATON, SPANISH. The root of the *Convolvulus batatas* of Linnæus. It is a native of the Indies. It is firm, and of a pale brown on the outside; white within, and very sweet, like chestnuts, and the only esculent root of the genus convolvulus.

POTENTILLA. (*A potentia*, from its efficacy.) 1. The name of a genus of plants in the Linnæan system. Class, *Icosandria* Order, *Polygynia*.

2. The pharmacopœial name of the wild tansy. *Argentina*. *Anserina*. The leaves of this plant, *Potentilla anserina* of Linnæus, possess mildly astringent and corroborant qualities; but are seldom used, except by the lower orders.

POTENTILLA ANSERINA. The systematic name of the silver-weed, or wild tansy. See *Potentilla*.

POTENTILLA REPTANS. The systematic name of the common cinquefoil. See *Pentaphyllum*.

POTERIUM. (From *potagion*, a cup; so named from the shape of its flowers.) The name of a genus of plants in the

Linnæan system. Class, *Monocotyledon*. Order, *Polyandria*.

POTERIUM SANGUISORRHA. The systematic name of the Burnet saxifrage, the leaves of which are often put into cool tankards; they have an astringent quality.

POUPARTII LIGAMENTUM. See *Poupurt's ligament*.

POUPART'S LIGAMENT. *Ligamentum Poupartii*. Fallopian ligament. Inguinal ligament. A strong ligament, or rather a tendinous expansion of the external oblique muscle, going across from the inferior and anterior spinous process of the ilium, to the crista of the os pubis. It is under this ligament the femoral vessels pass; and when the intestine or omentum passes underneath it, the disease is called a femoral hernia.

Powder, antimonial. See *Pulvis Antimoniæ*.

Powder of burnt hartshorn with opium. See *Pulvis cornu cum opio*.

Powder, compound, of aloe. See *Pulvis aloes compositus*.

Powder, compound, of chalk. See *Pulvis cretæ compositus*.

Powder, compound, of chalk with opium. See *Pulvis cretæ compositus cum opio*.

Powder, compound, of cinnamon. See *Pulvis cinnamoni compositus*.

Powder, compound, of contrayerva. See *Pulvis contrayervæ compositus*.

Powder, compound, of ipecacuanha. See *Pulvis Ipecacuanhæ compositus*.

Powder, compound, of kino. See *Pulvis kino compositus*.

Powder, compound, of scammony. See *Pulvis scammonie compositus*.

Powder, compound, of senna. See *Pulvis sennæ compositus*.

Powder, compound, of tragacanth. See *Pulvis tragacanthæ compositus*.

Power, muscular. See *Irritability and Muscular motion*.

Power, tonic. See *Irritability*.

Precipitate, red. See *Nitricæ-ærythrum hydrargyri*.

Precipitate, white. See *Calx hydrargyri alba*.

PRÆCORDIA. (From *præ*, before, and *cardia*, the heart.) The fore part or the region of the thorax.

PRÆFURNIUM. (From *præ*, before, and *furnus*, a furnace.) The mouth of a chemical furnace.

PRÆPARANTIA MEDICAMENTA. Medicines preparing the peccant fluids to pass off.

PRÆPARANTIA VASA. The spermatic vesicles of the testicles.

PRÆPUTIUM. (From *præputio*, to cut off before, because some nations used to cut it off in circumcision.) *Epagogion*, of Dioscorides. *Poshe*. The prepuce. The membranous or cutaneous fold that covers the glans penis and clitoris.

Præpuce. See *Præputium*.

PRASIUM. (From *πρᾶσις*, a square border; so called from its square stalks.) *Marrubium*, or horsehound.

PRASUM. (From *πρᾶσις*, to burn, because of their hot taste.) The leek.

PRAXIS. (From *πρᾶξις*, to perform.) The practice of medicine.

PRECIPITATION. (*Præcipitatio*; from *precipito*, to cast down.) When two bodies are united, for instance, an acid and an oxyd, and a third body is added, such as an alkali, which has a greater affinity with the acid than the metallic oxyd, the consequence is, that the alkali combines with the acid, and the oxyd, thus deserted, appears in a separate state at the bottom of the vessel in which the operation is performed. This decomposition is commonly known by the name of *precipitation*, and the substance that sinks is named a *precipitate*. The substance, by the addition of which the phenomenon is produced, is denominatèd the *precipitant*.

PREDISPOSING CAUSE. *Causa prædisponens. Causa prægumena.* That which renders the body susceptible of disease. The most frequent predisposing causes of diseases are, the temperament and habit of the body, idiosyncrasy, age, sex, and structure of the part diseased.

PREDISPOSITION. *Prædispositio.* That constitution, or state of the solids, or fluids, or of both, which disposes the body to the action of disease.

PREGNANCY. *Utero-gestation.* The particular manner in which pregnancy takes place has hitherto remained involved in obscurity, notwithstanding the laborious investigation of the most eminent philosophers of all ages.

Although pregnancy is a state which (with a few exceptions) is natural to all women, it is in general the source of many disagreeable sensations, and often the cause of diseases which might be attended with the worst consequences if not properly treated.

It is now, however, universally acknowledged, that those women who bear children enjoy, usually, more certain health, and are much less liable to dangerous diseases, than those who are unmarried, or who prove barren.

Signs of pregnancy.—The womb has a very extensive influence, by means of its nerves, on many other parts of the body; hence, the changes which are produced on it by impregnation, must be productive of changes on the state of the general system. These constitute the signs of pregnancy.

During the first fourteen or fifteen weeks, the signs of pregnancy are very ambiguous, and cannot be depended on; for, as they proceed from the irritation of the womb on other parts, they may be occasioned by

every circumstance which can alter the natural state of that organ.

The first circumstance which renders pregnancy probable, is the suppression of the periodical evacuation, which is generally accompanied with fulness in the breasts, head-ache, flushings in the face, and heat in the palms of the hands.

These symptoms are commonly the consequences of suppression, and therefore are to be regarded as signs of pregnancy, in so far only as they depend on it.

As, however, the suppression of the periodical evacuation often happens from accidental exposure to cold, or from the change of life in consequence of marriage, it can never be considered as an infallible sign.

The belly, some weeks after pregnancy, becomes flat, from the womb sinking, and hence drawing down the intestines along with it; but this cannot be looked upon as a certain sign of pregnancy, because an enlargement of the womb from any other cause will produce the same effect.

Many women, soon after they are pregnant, become very much altered in their looks, and have peculiar irritable feelings, inducing a disposition of mind which renders their temper easily ruffled, and incite an irresistible propensity to actions of which on other occasions they would be ashamed.

In such cases, the features acquire a peculiar sharpness, the eyes appear larger, and the mouth wider than usual; and the woman has a particular appearance, which cannot be described, but with which women are well acquainted.

These breeding symptoms, as they are called, originate from the irritation produced on the womb by impregnation; and as they may proceed from any other circumstance which can irritate that organ, they cannot be depended on when the woman is not young, or where there is not a continued suppression for at least three periods.

The irritations on the parts contiguous to the womb are equally ambiguous: and therefore the signs of pregnancy, in the first four months, are always to be considered as doubtful, unless every one enumerated be distinctly and unequivocally present.

From the fourth month, the signs of pregnancy are less ambiguous, especially after the womb has ascended into the cavity of the belly. In general, about the fourth month, or a short time after, the child becomes so much enlarged, that its motions begin to be felt by the mother; and hence a sign is furnished at that period called quickening. Women, very improperly consider this sign as the most unequivocal proof of pregnancy; for though, when it

occurs about the period described, preceded by the symptoms formerly enumerated, it may be looked upon as a sure indication that the woman is with child; yet, when there is an irregularity, either in the preceding symptoms or in its appearance, the situation of the woman must be doubtful.

This fact will be easily understood; for as the sensation of the motion of the child cannot be explained, or accurately described, women may readily mistake other sensations for that of quickening. Flatus has often been so pent up in the bowels, that the natural pulsation of the great arteries, of which people are conscious only in certain states of the body, has frequently been mistaken for this feeling.

After the fourth month, the womb rises gradually from the cavity of the basin, enlarges the belly, and pushes out the navel: hence the protrusion of the navel has been considered one of the most certain signs of pregnancy in the latter months. Every circumstance, however, which increases the bulk of the belly occasions this symptom; and therefore it cannot be trusted to, unless other signs concur.

The progressive increase of the belly, along with suppression, after having been formerly regular, and the consequent symptoms, together with the sensation of quickening at the proper period, afford the only true marks of pregnancy.

These signs, however, are not to be entirely depended on; for the natural desire which every woman has to be a mother, will induce her to conceal, even from herself, every symptom which may render her situation doubtful, and to magnify every circumstance which can tend to prove that she is pregnant.

Beside quickening and increase of bulk of the belly, another symptom appears in the latter months, which, when preceded by the ordinary signs, renders pregnancy certain beyond a doubt. It is the presence of milk in the breasts. When, however, there is any irregularity in the preceding symptoms, this sign is no longer to be considered of any consequence.

As every practitioner must naturally wish to distinguish pregnancy from disease, the disorders which resemble it should be thoroughly understood, and also their diagnostics. It is, however, necessary to remark, that wherever any circumstance occurs which affords the most distant reason to doubt the case, recourse ought to be had to the advice of an experienced practitioner, and every symptom should be unreservedly described to him.

ΠΡΕΣΒΥΟΠΙΑ. (From *prehendo*, to surprise; so named from its sudden seizure.) The catalepsy.

PRESBYOPIA. (From *πρεσβυς*, old, and *ὄψις*, sight, because it is frequent with old men.) That defect of the sight by which objects close are seen confusedly, but, at remoter distances, distinctly. As the myopia is common to infants, so the presbyopia is a malady common to the aged. The proximate cause is a tardy adunation into the focus, so that it falls beyond the retina. The species are,

1. *Presbyopia* from a flatness of the cornea. By so much the cornea is flatter, so much the less and more tardy it refranges the rays into the focus. This evil arises, 1st, From a want of aqueous or vitreous humour, which is common to the aged; or may arise from some disease; 2d, From a cicatrix, which diminishes the convexity of the cornea; 3d, From a natural conformation of the cornea.

2. *Presbyopia* from too flat a crystalline lens. This evil is most common to the aged, or it may happen from a wasting of the crystalline lens.

3. *Presbyopia* from too small density of the cornea or humours of the eye. By so much more these humours are thin or rarified, so much the less they refrange the rays of light. Whosoever is a myope from this cause is cured in older age; for age induces to a greater density of the cornea and lens. From this it is an observed fact, that the *presbyopes* are often cured spontaneously, and throw away their glasses, which younger persons in this disease are obliged to use.

4. *Presbyopia* from a custom of viewing continually remote objects; hence artificers who are occupied in remote objects are said to contract this malady. The reason of this phenomenon is not very clear.

5. *Presbyopia senilis.* From a multitude of causes aged persons are *presbyopes*; from a penury of humours, which render the cornea and lens flatter, and the bulb shorter. When in senile ages, from dryness, the bulb of the eye becomes flatter and shorter, and the cornea flatter, those who were short-sighted or myopes before, see now without their concave glasses.

6. *Presbyopia* from too close a proximity of objects. The focus is shorter of distant, but longer of nearer objects.

7. *Presbyopia* from a coarctated pupil. By so much smaller is the aperture of the diaphragm in an optic tube, so much remoter is the focus.

8. *Presbyopia mercurialis*, which arises from the use of mercurial preparations. The patient feels a pressing pain in the eye, which, from being touched is increased, and the bulb of the eye appears as if rigid, and with difficulty can be moved. Near objects the patient can scarce distinguish.

and distant only in a confused manner. Many have supposed this disorder an imperfect amaurosis.

PRESBYTÆ. See *Presbyopia*.

PRESBYTIA. (From *πρεσβυς*, old; because it is usual to old people.) See *Presbyopia*.

PRESURA. (From *πρεσβα*, to inflame.) Inflammation at the ends of the fingers from cold.

PRIAPEIA. See *Nicotiana minor*.

PRIAPISCUS. (From *πριαπός*, the penis.)

1. A tent made in the form of a penis. 2. A bougie.

PRIAPISMUS. (From *πριαπός*, a heathen god, whose penis is always painted erect.) Priapism. A continual erection of the penis. Cælius Aurelianus says it is a palsy of the seminal vessels, by which the disorder is produced.

Priapism See *Priapismus*.

PRIAPUS. (Πριαπός, a heathen god, remarkable for the largeness of his genitals.)

1. The penis.

2. A name of the nepenthes or wonderful plant, from the appendages at the end of the leaves resembling an erected penis.

PRIMÆ VIÆ. The first passages. The stomach and the intestinal tube are so called, and the lacteals the *secundæ viæ*.

Primary teeth. See *Teeth*.

Primrose. See *Primula vulgaris*.

PRIMULA. (From *primulus*, the beginning; so called because it flowers in the beginning of the spring.) The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

PRIMULA VERIS. (From *primulus*, the beginning, so called because it flowers in the beginning of the spring.) *Verbasculum*. The cowslip, paigil, or peagle. The flowers of this plant have a moderately strong and pleasant smell, and a somewhat roughish bitter taste. Vinous liquors impregnated with their flavour by maceration or fermentation, and strong infusions of them drank as tea, are supposed to be mildly corroborant, antispasmodic, and anodyne. An infusion of three pounds of the fresh flowers in five pints of boiling water is made in the shops into a syrup of a fine yellow colour, and agreeably impregnated with the flavour of the cowslip.

PRIMULA VULGARIS. The primrose. The leaves and root of this common plant possess sternutatory properties.

PRINCEPS ALEXIPHARMACORUM. Angelica. Which by some was formerly so much esteemed as to obtain this name.

PRINCIPLES. *Principia*. Primary substances. According to modern chemists, this term is applied to those particles which are composed of two or more elements (See *Elements*), that may again be

decomposed by the action of fire or putridity, such as water, gum, resin, &c.

PRIONONES. (From *πριον*, a saw.) Serated: applied to the sutures of the skull.

PRIOR ANNULARIS. (*Musculus prior annularis*.) Fourth interosseus of Winslow. An internal interosseous muscle of the hand. See *Interossei manus*.

PRIOR INDICIS. *Extensor tertii interodii indicis* of Douglas. *Interossei manus internus* of Albinus. *Interossei* of Winslow and Cowper, and *sous-metacarpo-lateri-phalangiens* of Dumas. An internal interosseal muscle of the hand, which draws the forefinger inwards towards the thumb, and extends it obliquely.

PRIOR MEDII. (*Musculus prior medii*.) *Interossei manus bicipites*, seu *externi* of Albinus. *Second interosseus* of Douglas, and *sous-metacarpo-lateri-phalangiens* of Dumas. An external interosseous muscle of the hand. See *Interossei manus*.

PROBANG. A flexible piece of whalebone with sponge fixed to the end.

PROBÆ. (From *probo*, to try; because surgeons try the depth and extent of wounds, &c. with it.) *Stylus*. A surgical instrument of a long and slender form.

PROBOLÆ. (From *προβέλλω*, to project.) A prominence. Anapophysis.

PROBOSCIS. (From *προ*, before, and *βροσκα*, to feed. (A snout or trunk as that of an elephant by which it feeds itself.

PROCARDIUM. (From *προ*, before, and *καρδια*, the stomach or heart.) The pit of the stomach.

PROCATARCTIC CAUSE. (*Causa procatarctica*, from *προκαταρχω*, to go before.) See *Exciting cause*.

PROCESS. (*Processus*, from *procedo*, to go before.) An eminence of a bone; as the spinous and transverse processes of the vertebrae.

PROCESSUS CAUDATUS. See *Lobulus caudatus*.

PROCESSUS CECI VERMIFORMIS. See *Intestines*.

PROCESSUS CILIARIS. See *Ciliar ligament*.

PROCESSUS MAMILLARES. A name formerly applied to the olfactory nerves.

PROCIDENTIA. (From *procido*, to fall down.) A falling down of any part; thus, *procidentia ani, uteri, vaginae*, &c.

PROCONDYLUS. (From *προ*, before, and *κονδυλος*, the middle joint of the finger.) The first joint of a finger next the metacarpus.

PROCTALGIA. (From *πρωκτος*, the fundament, and *αλγος*, pain.) A violent pain at the anus. It is mostly symptomatic of some disease, as piles, scirrhus, prurigo, cancer, &c.

PROCTITIS. (From *πρωκτος*, the anus.) *Clonedia. Cyscolitis.* Inflammation of the internal or mucous membrane of the lower part of the rectum.

PROCTOLEUCORRHEA. (From *πρωκτος*, the anus, *λευκος*, white, and *ρρεω*, to flow.) *Proctorrhœa.* A purging of white mucus with heat and itching.

PROCTORRHEA. (From *πρωκτος*, the anus, and *ρρεω*, to flow.) See *Proctoleucorrhœa*.

PROFLUVIA. (From *profluo*, to run down.) Fluxes. The fifth order in the class *pyrexia* of Cullen's nosology, characterised by pyrexia, with increased excretions.

PROFLUVII CORTEX. See *Cenosi cortex*.

PROFUNDUS. See *Flexor profundus perforans*.

PROFUSIO. A loss of blood. A genus of disease in the class *locales* and order *apo censes* of Cullen.

PROGLOSIS. (From *προ*, before, and *γλωσσα*, the tongue.) The tip of the tongue.

PROGNOSIS. (From *προ*, before, and *γνωσις*, to know.) The art of foretelling the event of diseases from particular symptoms.

PROGNOSTIC. (*Prognosticus*, from *προγνωσις*, to know before hand.) Applied to those symptoms which may be foretold before they appear.

PROLAPSUS. (From *prolabor*, to slip down.) *Procidencia. Delapsio. Exania. Proptoma. Proptosis.* A protrusion. A genus of disease in the class *locales* and order *ectopia* of Cullen; distinguished by the falling down of a part that is uncovered.

PROLEPTICUS (From *προλαμβάνω*, to anticipate.) Applied to those diseases whose paroxysms anticipate each other, or return after less and less intervals of intermission.

PRONALACTERIUM. (From *προ*, before, and *μαλασσω*, to soften.) The room where the body was softened previous to bathing it.

PRONETOPIDIUM. (From *προ*, before, and *μετωπον*, the forehead.) *Prometoposis.* The skin upon the forehead.

PRONETOPIS. See *Prometopidium*.

PRONATION. The act of turning the palm of the hand downwards. It is performed by rotating the radius upon the ulna, by means of several muscles which are termed pronators; as,

PRONATOR QUADRATUS. See *Pronator radii quadratus*.

PRONATOR RADII BREVIS. See *Pronator radii quadratus*.

PRONATOR RADII QUADRATUS. *Pronator quadratus* of Douglas and Albinus. *Pronator quadratus sive transversus* of Winslow. *Pronator radii brevis seu quadratus*

of Cowper. *Cubito-radialis* of Dumas. This, which has gotten its name from its use and its shape, is a small fleshy muscle, situated at the lower and inner part of the fore-arm, and covered by the tendons of the flexor muscles of the hand. It arises tendinous and fleshy from the lower and inner part of the ulna, and runs nearly in a transverse direction, to be inserted into that part of the radius which is opposite to its origin, its inner fibres adhering to the interosseous ligament. This muscle assists in the pronation of the hand, by turning the radius inwards.

PRONATOR RADII TERES of Innes and Cowper. *Pronator teres* of Albinus and Douglas. *Pronator teres. sive obliquus* of Winslow. *Epirochlo-radialis* of Dumas. This is a small muscle, situated at the upper and anterior part of the fore-arm. It is called *teres*, to distinguish it from the pronator quadratus. It arises tendinous and fleshy from the anterior and inferior part of the outer condyle of the os humeri; and tendinous from the coronoid process of the ulna, near the insertion of the brachialis internus. The median nerve passes between these two portions. From these origins the muscle runs obliquely downwards and outwards, and is inserted, tendinous and fleshy, into the anterior and convex edge of the radius, about the middle of that bone. This muscle, as its name indicates, serves to turn the hand inwards.

PRONERVATIO. (From *pro*, before, and *nervus*, a string.) A tendon or string like the end of a muscle.

PROPHYLACTIC. (*Prophylactica*, from *προ*, before, and *φύλασσω*, to defend.) Any means made use of to preserve health.

PROPRIETATIS ELIXIR. Elixir of aloes and tinctura aloes composita.

PROPTOMA. (From *προπτωω*, to fall down.) *Procidencia.* A relaxation such as that of the scrotum, of the under lip, of the breasts in females, of the præpuce, or of the ears.

PROPTOMA. (From *προ*, before, and *πυρ*, pus.) A premature collection of pus.

PRONA. (From *πρωγα*, the prow of a vessel.) The occiput.

PROSARTHROSIS. (From *προσ*, to, and *αρθρω*, to articulate.) That articulation which has manifest motion.

PROSPERMA. (From *προσπυρνωμι*, to fix near.) A fixing of humours in one spot.

PROSTASIS. (From *προσιστημι*, to predominate.) An abundance of morbid humours.

PROSTATE GLAND. (*Glandula prostata*, from *προ*, before, and *ιστημι*, to stand; because it is situated before the urinary bladder.) *Corpus glandulosum*

Adenoides. A very large, heart-like, firm gland, situated between the neck of the urinary bladder and bulbous part of the urethra. It secretes the lacteal fluid, which is emitted into the urethra by ten or twelve ducts that open near the verumontanum during coition. This gland is liable to inflammation and its consequences.

PROSTATE, INTERIOR. See *Transversus perinei alter*.

PROTOGALA. (From *πρωτος*, first, and *γала*, milk.) The first milk after delivery.

PROXIMATE CAUSE. *Causa proxima*. The proximate cause of a disease may be said to be in reality the disease itself. All proximate causes are either diseased actions of simple fibres, or an altered state of the fluids.

PRUNA. (Pl. of *prunum*.) Plums or prunes. See *Plums*.

Prune. See *Plums*.

PRUNELLA. (From *pruna*, a burn, because it heals burns.) *Brunella*. *Consolidula minor*. *Symphitum minus*.

1. The name of a genus of plants in the Linnæan system. Class, *Didynamia*. Order, *Gymnospermia*.

2. The pharmacopœial name of what is also called self-heal. *Prunella vulgaris* of Linnæus, *foliis omnibus ovato-oblongis, serratis, petiolatis*, it is recommended as an adstringent in hæmorrhages and fluxes, as in gargles against aphthæ and inflammation of the fauces.

PRUNELLA VULGARIS. The systematic name of the self-heal. See *Prunella*.

Prunelloe. See *Plum*.

PRUNUM GALLICUM. The common prune or plum. The plant which affords this fruit is the *Prunus domestica* of Linnæus:—*Pedunculis subsolitariis, foliis lanceolato ovatis convolutis, ramis muticis; gemme florifera aphylla*. Prunes are considered as emollient, cooling, and laxative, especially the French prunes, which are directed in the decoction of senna, and other purgatives: and the pulp is ordered in the electuarius of senna. See *Plum*.

PRUNUS SILVESTRE. The sloe or fruit of the *Prunus spinosa* of Linnæus:—*Prunus pedunculis solitariis foliis lanceolatis glabris, ramis spinosis*. It is sometimes employed in gargles, to tumefactions of the tonsils and uvula, and from its adstringent taste was formerly much used in hæmorrhages, &c.

PRUNUS. The name of a genus of plants in the Linnæan system. Class, *Icosandria*. Order, *Monogynia*.

PRUNUS AVIUM. The systematic name of the black cherry-tree. See *Cerasa nigra*.

PRUNUS CERASUS. The systematic name of the red cherry-tree. See *Cerasa rubra*,

PRUNUS DOMESTICA. The systematic name of the damson-tree. See *Damson*.

PRUNUS LAURO-CERASUS. The systematic name of the poison laurel. See *Lauro-cerasus*.

PRUNUS PADUS. The systematic name of the bird cherry-tree. See *Padus*.

PRUNUS SPINOSA. The systematic name of the sloe-tree. See *Prudus sylvestris*.

PRURIGO (From *prurio*, to itch.)

Pruritus. *Scabies*; *Psora*. *Darta*. *Libido*. *Pavoni*. The prurigo is a genus of disease in the order *papulous eruptions* of Dr. Willan's cutaneous diseases. As it arises from different causes, or at different periods of life, and exhibits some varieties in its form, he describes it under the titles of prurigo mitis, prurigo formicans, and prurigo senilis. In these the whole surface of the skin is usually affected; but there are likewise many cases of local prurigo, which will be afterwards noticed according to their respective situations.

1. The *prurigo mitis* originates without any previous indisposition, generally in spring, or the beginning of summer. It is characterised by soft and smooth elevations of the cuticle, somewhat larger than the papulæ of the lichen, from which they also differ by retaining the usual colour of the skin; for they seldom appear red, or much inflamed, except from violent friction. They are not, as in the other case, accompanied with tingling, but with a sense of itching almost incessant. This is, however, felt more particularly on undressing, and often prevents rest for some hours after getting into a bed. When the tops of the papulæ are removed by rubbing or scratching, a clear fluid oozes out from them, and gradually concretes into thin black scabs.

This species of prurigo mostly affects young persons, and its cause may, I think, says Dr. Willan, in general be referred to sordes collected on the skin, producing some degree of irritation, and also preventing the free discharge of the cutaneous exhalation; the bad consequences of which must necessarily be felt at that season of the year when perspiration is most copious. Those who have originally a delicate or irritable skin, must likewise, in the same circumstances, be the greatest sufferers.

The eruption extends to the arms, breast, back, and thighs, and often continues during two or three months of the summer, if not relieved by proper treatment. When persons affected with it neglect washing the skin, or are uncleanly in their apparel, the irruption grows more inveterate, and at length, changing its form, often terminates in the itch. Pustules arise among the papulæ, some filled with lymph, others

with pus. The *acarus scarabei* begins to breed in the furrows of the cuticle, and the disorder becomes contagious.

2. The *Prurigo formicans* is a much more obstinate and troublesome disease than the foregoing. It usually affects persons of adult age, commencing at all seasons of the year indifferently; and its duration is from four months to two or three years, with occasional short intermissions. The papulæ are sometimes larger, sometimes more obscure, than in the preceding species; but, are, under every form, attended with an incessant, almost intolerable itching. They are diffused over the whole body, except the face, feet, and palms of the hands; they appear, however, in greatest number on those parts which, from the ordinary mode of dress, are subjected to tight ligatures; as about the neck, joints, and thighs.

The itching is complicated with other sensations, which are variously described by patients. They sometimes feel as if small insects were creeping on the skin; sometimes as if stung all over by ants; sometimes as if hot needles were piercing the skin in divers places. On standing before a fire, or undressing, and more particularly on getting into bed, these sensations become most violent, and usually preclude all rest during the greater part of the night. The *prurigo formicans* is by most practitioners deemed contagious, and confounded with the itch. In endeavouring to ascertain the justness of this opinion, Dr. Willan has been led to make the following remarks: 1. The eruption is, for the most part, connected with internal disorder, and arises where no source of infection can be traced. 2. Persons affected may have constant intercourse with several others, and yet never communicate the disease to any of them. 3. Several persons of one family may have the *prurigo formicans* about the same time; but he thinks this should be referred rather to a common predisposition than to contagion, having observed that individuals of a family are often so affected, at certain seasons of the year, even when they reside at a distance from each other.

Although the *prurigo formicans* is never, like the former species, converted into the itch, yet it does occasionally terminate in a pustular disease, not contagious.

3. *Prurigo senilis*. This affection does not differ much in its symptoms and external appearances from the *prurigo formicans*; but has been thought by medical writers to merit a distinct consideration, on account of its peculiar inveteracy. The *prurigo* is perhaps aggravated, or becomes more permanent, in old age, from the dry, condensed state of the skin and cuticle, which often takes place at that

period. Those who are affected with it in a high degree have little more comfort to expect during life, being incessantly tormented with a violent and universal itching. The state of the skin in the *prurigo senilis*, is favourable to the production of an insect, the *pediculus humanus*, more especially to the variety of it usually termed body-lice.

These insects, it is well known, are bred abundantly among the inhabitants of sordid dwellings, of jails, workhouses, &c. and in such situations prey upon persons of all ages indiscriminately. But in the *prurigo senilis* they arise, notwithstanding every attention to cleanliness or regimen, and multiply so rapidly that the patient endures extreme distress, from their perpetual irritation. The nits or eggs are deposited on the small hairs of the skin, and the pediculi are only found on the skin or on the linen, not under the cuticle, as some authors have represented. In connexion with the foregoing series of complaints, Dr. Willan mentions some pruriginous affections which are merely local. He confines his observations to the most troublesome of these, seated in the podex, præputium, urethra, pubes, scrotum, and pudendum muliebre. Itching of the nostrils, eye-lids, lips, or of the external ear, being generally symptomatic of other diseases, do not require a particular consideration.

1. *Prurigo podicis*. *Ascarides* in the rectum excite a frequent itching and irritation about the sphincter ani, which ceases when the cause is removed by proper medicines. A similar complaint often arises, independently of worms, hæmorrhoidal tumours, or other obvious causes, which is mostly found to affect persons engaged in sedentary occupations; and may be referred to a morbid state of secretion in the parts, founded, perhaps, on a diminution of constitutional vigour. The itching is not always accompanied with an appearance of papulæ or tubercles; it is little troublesome during the day-time, but returns every night soon after getting into bed, and precludes rest for several hours. The complaint continues in this form during three or four-months, and has then an intermission, till it is produced again by hot weather, fatigue, watching, or some irregularity in diet. The same disease occurs at the decline of life, under a variety of circumstances.

Women after the cessation of the catamenia, are liable to be affected with this species of *prurigo*, more especially in summer or autumn. The skin between the nates is rough and papulated, sometimes scaly, and a little humour is discharged by violent friction. Along with this complaint, there is often an eruption of itching papulæ on the neck, breast, and back;

a swelling and inflammation of one or both ears, and a discharge of matter from behind them, and from the external meatus auditorius. The prurigo podicis sometimes occurs as a symptom of the lues venerea.

2. The *prurigo preputii* is owing to an altered state of secretion on the glans penis, and inner surface of the preputium. During the heat of summer there is also, in some persons, an unusual discharge of mucus, which becomes acrimonious, and produces a troublesome itching, and often an excoriation of these parts. Washing of them with water, or soap and water, employed from time to time, relieves the complaint, and should indeed be practised as an ordinary point of cleanliness, where no inconvenience is immediately felt. If the fluid be secreted in too large a quantity, that excess may be restrained, by washes made with the extract of lead, or by applying the unguentum cerusæ acetate.

3. *Prurigo urethralis*. A very troublesome itching sometimes takes place at the extremity of the urethra in females, without any manifest cause. It occurs as well in young women as in those who are of an advanced age. On examination, no stricture nor tumour has been found along the course of the urethra. Probably, however, the itching may be occasioned by a morbid state of the neck of the bladder, being in some instances connected with pain and difficulty of making water.

An itching at the extremity of the urethra in men is produced by calculi, and by some diseases of the bladder. In cases of stricture an itching is also felt, but near the place where the stricture is situated. Another cause of it is small broken hairs, which are sometimes drawn in from the pubes, between the preputium and glans, and which afterwards becoming fixed in the entrance of the urethra, occasion an itching or slight stinging, particularly on motion. Mr. J. Pearson, surgeon of the Lock Hospital, has seen five cases of this kind, and gave immediate relief by extracting the small hair from the urethra.

4. *Prurigo pubis*. Itching papule often arise on the pubes, and become extremely sore if their tops are removed by scratching. They are occasioned sometimes by neglect of cleanliness, but more commonly by a species of pediculus, which perforates the cuticle, and thus derives its nourishment, remaining fixed in the same situation. These insects are termed by Linnaeus, &c. *pediculi pubis*; they do not, however, affect the pubes only, but often adhere to the eye-brows, eye-lids, and axillæ. They are often found, also, on the breast, abdomen, thighs, and legs, in persons of the sanguine temperament; who have those parts covered with strong

hairs. It is remarkable that they seldom or never fix upon the hairy scalp. The great irritation produced by them on the skin solicits constantly scratching, by which they are torn from their attachments; and painful tubercles arise at the places where they had adhered. When the pediculi are diffused over the greater part of the surface of the body, the patient's linen often appears as if sprinkled with drops of blood.

5. *Prurigo scroti*. The scrotum is affected with a troublesome and constant itching from ascarides within the rectum, from friction by violent exercise in hot weather, and very usually from the pediculi pubis. Another and more important form of the complaint appears in old men, sometimes connected with the prurigo podicis, and referable to a morbid state of the skin, or superficial glands of the part. The scrotum, in this case, assumes a brown colour, often also becoming thick, scaly, and wrinkled. The itching extends to the skin covering the penis, more especially along the course of the urethra; and has little respite, either by day or night.

6. The *Prurigo pudendi muliebris*, is somewhat analogous to the prurigo scroti in men. It is often a symptomatic complaint in the lichen and lepra; it likewise originates from ascarides irritating the rectum, and is in some cases, connected with a discharge of the fluor albus.

A similar affection arises in consequence of the change of state in the genital organs at the time of puberty, attended with a series of most distressing sensations. Dr. Willan confines his attention to one case of the disorder, which may be considered as idiopathic, and which usually affects women soon after the cessation of the catamenia. It chiefly occurs in those who are of the phlegmatic temperament, and inclined to corpulency. Its seat is the labia pudendi, and entrance into the vagina. It is often accompanied with an appearance of tension or fulness of those parts, and sometimes with inflamed itching papule on the labia and mons veneris. The distress arising from a strong and almost perpetual itching in the above situation, may be easily imagined. In order to allay it in some degree, the sufferers have frequent recourse to friction, and to cooling applications: whence they are necessitated to forego the enjoyment of society. An excitement of venereal sensations also takes place from the constant direction of the mind to the parts affected, as well as from the means employed to procure alleviation. The complicated distress thus arising, renders existence almost insupportable, and often produces a state of mind bordering on phrensy.

Deep ulcerations of the parts seldom

take place in the prurigo pudendi; but the appearance of aphthæ on the labia and nymphæ, is by no means unusual. From intercourse with females under these circumstances, men are liable to be affected with aphthous ulcerations on the glans, and inside of the præputium, which prove troublesome for a length of time, and often excite an alarm, being mistaken for chancres.

Women, after the fourth month of their pregnancy, often suffer greatly from the prurigo pudendi, attended with aphthæ. These, in a few cases, have been succeeded by extensive ulcerations, which destroyed the nymphæ, and produced a fatal hectic: such instances are, however, extremely rare. The complaint has, in general, some intervals or remissions; and the aphthæ usually disappear soon after delivery, whether at the full time, or by a miscarriage.

PRURITUS. (From *pruria*, to itch.) See *Prurigo*.

PRUSSIATS. Salts formed by the union of the prussic acid, or colouring matter of prussian blue, with different bases; thus, *prussiat of alumine*, *prussiat of ammoniac*, &c.

PSALLOIDES. (From *ψαλλος*, a stringed instrument, and *υδος*, a likeness; because it appears as if stringed like a dulcimer.) Applied by the ancients to the inner surface of the fornix of the brain.

PSALTERIUM. (A harp; because it is marked with lines that give it the appearance of a harp.) *Lyra*. The medullary body that unites the posterior crura of the fornix of the brain.

PSAMMISXUS. (From *ψαμμος*, sand.) An application of hot sand to any part of the body.

PSAMNODES. (From *ψαμμος*, sand.) Applied to urine which deposits a sandy sediment.

PSPELLISMUS. (From *ψαλλίζω*, hesitation of speech.) *Psellotis*. Defect of speech. A genus of disease in the class *locales* and order *dyscenæria* of Cullen.

PSELLOTIS. See *Psellismus*.

PSEUDO. (*ψευδς*, false.) *Pseudus*. Spurious; prefixed to many substances which are only fictitious imitations; as *pseudomomum*, a spurious kind of amonium, &c.

PSEUDO-ACORUS. See *Iris palustris*.

PSEUDOBLEPSIS. (From *ψυδος*, false, and *ὁρασις*, sight.) *Phantasma*. *Suffusio*. Imaginary vision of objects. A genus of disease in the class *locales* and order *dysæthesia* of Cullen; characterised by depraved sight, creating objects, or representing them different from what they are.

Species:

1. *Pseudoblepsis imaginaria*, in which objects are perceived that are not present.

2. *Pseudoblepsis mutans*, in which objects that are present appear somewhat changed.

PSEUDOPHYTETERUM. See *Piarnica*.

PSIDIUM. The name of a genus of plants in the Linnæan system.

PSIDIUM POMIFERUM. The systematic name of the apple guava. This plant, and the *pyriferum*, bear fruits, the former like apples, the latter like pears. The apple kind is most cultivated in the Indies, on account of the pulp having a fine acid flavour, whereas the pear species is sweet, and therefore not so agreeable in warm climates. Of the inner pulp of either, the inhabitants make jellies; and of the outer rind they make tarts, marmalades, &c. The latter they also stew and eat with milk, and prefer them to any other stewed fruits. They have an adstringent quality, which exists also in every part of the tree, and abundantly in the leaf-buds, which are occasionally boiled with barley and liquorice, as an excellent drink against diarrhæas. A simple decoction of the leaves, used as a bath, are said to cure the itch, and most cutaneous eruptions.

PSIDIUM PYRIFERUM. The systematic name of the pear guava. See *Psidium pomiferum*.

PSILOTERA. (From *ψιλος*, to denude.) Applications to remove the hair.

PSILOTERUM. (From *ψιλος*, to depilate; so called because it was used by depilatories.) The white briony.

PSIMMYTHISM. (From *ψω*, to smooth; so called because of its use as a cosmetic.) Cerus, or white lead.

PSOÆ. (*ψοας*, the loins.) *Alopeces*. *Ne-phrometra*. *Neurometeres*. The name of two pair of muscles in the loins.

PSOAS ABCESS. See *Lumbar abcess*.

PSOAS MAGNUS. (From *ψοα*, the loin; because it is situated in the loins.) *Psoas, seu lumbaris internus* of Winslow. *Pre-lumbo-trochantin* of Dumas. This is a long, thick, and very considerable muscle, situated close to the forepart and sides of the lumbar vertebræ. It arises from the bodies of the last vertebræ of the back, and of all the lumbar vertebræ laterally, as well as from the anterior surfaces of their transverse processes, by distinct tendinous and fleshy slips, that are gradually collected into one mass, which becomes thicker as it descends, till it reaches the last of the lumbar vertebræ, where it grows narrower again, and, uniting its outer and posterior edge, (where it begins to become tendinous) with the iliacus internus, descends along with that muscle under the ligamentum fallopii, and goes to be inserted tendinous at the bottom of the trochanter minor of the os femoris, and fleshy into the bone a little below that process. Between the tendon of this mus-

cle and the ischium, we find a considerable bursa mucosa. This muscle, at its origin, has some connexion with the diaphragm, and likewise with the quadratus lumborum. It is one of the most powerful flexors of the thigh forwards, and may likewise assist in turning it outwards. When the inferior extremity is fixed, it may help to bend the body forwards, and in an erect posture, it greatly assists in preserving the equilibrium of the trunk upon the upper part of the thigh.

PSOAS PARVUS. *Pre-humbo-pubius* of Dumas. This muscle, which was first described by Riolanus, is situated upon the psoas magnus, at the anterior part of the loins. The psoas parvus arises thin and fleshy from the side of the uppermost vertebra of the loins, and sometimes also from the lower edge of the last vertebra of the back, and from the transverse processes of each of these vertebrae; it then extends over part of the psoas magnus, and terminates in a thin flat tendon, which is inserted into that part of the brim of the pelvis, where the os pubis joins the ilium. From this tendon a great number of fibres are sent off, which form a thin fascia; that covers part of the psoas magnus and iliacus internus, and gradually loses itself on the fore part of the thigh. In the human body this muscle is very often wanting; but in a dog, according to Douglas, it is never deficient. Riolanus was of opinion, that it occurs oftener in men than in women; Winslow asserts just the contrary; but the truth seems to be, that it is as often wanting in one sex as in the other. Its use seems to be to assist the psoas magnus in bending the loins forwards; and when we are lying upon our back, it may help to raise the pelvis.

Psoas sive lumbaris internus. See *Psoas magnus*.

PSORA. *Yupa. Scabies.* The itch. A genus of diseases in the class *locales* and order *dialyses* of Cullen: appearing first on the wrists and between the fingers in small pustules with watery heads. It is contagious.

PSORIASIS. (From *ψωρα*, to itch.) The disease to which Dr. Willan gives this title is characterised by a rough and scaly state of the cuticle, sometimes continuous, sometimes in separate patches, of various sizes, but of an irregular figure, and for the most part accompanied with rhagades or fissures of the skin. From the lepra it may be distinguished, not only by the distribution of the patches, but also by its cessation and recurrence at certain seasons of the year, and by the disorder of the constitution with which it is usually attended. Dr. Willan gives the following varieties:

Psoriasis guttata. This complaint appears in small, distinct, but irregular

patches of laminated scales, with little or no inflammation round them. The patches very seldom extend to the size of a sixpence. They have neither an elevated border, nor the oval or circular form by which all the varieties of lepra are distinguished; but their circumference is sometimes angular, and sometimes goes into small serpentine processes. The scale formed upon each of them is thin, and may be easily detached, leaving a red, shining base. The patches are often distributed over the greatest part of the body, but more particularly on the back part of the neck, the breasts, arms, loins, thighs, and legs. They appear also upon the face, which rarely happens in lepra. In that situation they are red and more rough than the adjoining cuticle, but not covered with scales. The psoriasis guttata often appears on children in a sudden eruption, attended with a slight disorder of the constitution, and spreads over the body within two or three days. In adults it commences with a few scaly patches on the extremities, proceeds very gradually, and has a longer duration than in children. Its first occurrence is usually in the spring season, after violent pains in the head, stomach, and limbs. During the summer it disappears spontaneously, or may be soon removed by proper applications, but it is apt to return again early in the ensuing spring, and continues so to do for several successive years. When the scales have been removed, and the disease is about to go off, the small patches have a shining appearance, and they retain a dark red, intermixed with somewhat of a bluish colour, for many days, or even weeks, before the skin is restored to its usual state. In the venereal disease there is an eruption which very much resembles the psoriasis guttata, the only difference being a slighter degree of scalliness, and a different shade of colour in the patches, approaching to a livid red, or very dark rose colour. The patches vary in their extent, from the section of a pea, to the size of a silver penny, but are not exactly circular. They rise at first very little, if at all, above the cuticle. As soon, however, as the scales appear on them, they become sensibly elevated; and sometimes the edge or circumference of the patch is higher than the little scales in its centre. This eruption is usually seen upon the forehead, breast, between the shoulders, or in the inside of the fore-arms, in the groins, about the inside of the thighs, and upon the skin covering the lower part of the abdomen. The syphilitic psoriasis guttata is attended with, or soon followed by, an ulceration of the throat. It appears about six or eight weeks after a chancre has been healed by an ineffectual course of mercury. A similar appearance takes place at nearly the same period, in

some cases where no local symptoms had been noticed. When a venereal sore is in a discharging state, this eruption, or other secondary symptoms, often appear much later than the period above mentioned. They may also be kept back three months, or even longer, by an inefficient application of mercury. If no medicines be employed, the syphilitic form of the psoriasis guttata will proceed during several months, the number of the spots increasing, and their bulk being somewhat enlarged, but without any other material alteration.

2. The *Psoriasis diffusa* spreads into large patches irregularly circumscribed, reddish, rough, and chappy, with scales interspersed. It commences, in general, with numerous minute asperities, or elevations of the cuticle, more perceptible by the touch than by sight. Upon these, small distinct scales are soon after formed, adhering by a dark central point, while their edges may be seen white and detached. In the course of two or three weeks all the intervening cuticle becomes rough and chappy, appears red, and raised, and wrinkled, the lines of the skin sinking into deep furrows. The scales which form among them are often slight, and repeatedly exfoliate. Sometimes, without any previous eruption of papulæ, a large portion of the skin becomes dry, harsh, cracked, reddish, and scaly, as above described. In other cases, the disorder commences with separate patches of an uncertain form and size, some of them being small, like those in the psoriasis guttata, some much larger. The patches gradually expand till they become confluent, and nearly cover the part or limb affected. Both the psoriasis guttata and diffusa likewise occur as a sequel of the lichen simplex. This transition takes place more certainly after frequent returns of the lichen. The parts most affected by psoriasis diffusa are the cheeks, chin, upper eyelids, and corners of the eyes, the temples, the external ear, the neck, the fleshy parts of the lower extremities, and the forearm, from the elbow to the back of the hand, along the supinator muscle of the radius. The fingers are sometimes nearly surrounded with a loose scaly incrustation; the nails crack and exfoliate superficially. The scaly patches likewise appear, though less frequently, on the forehead and scalp, on the shoulders, back, and loins, on the abdomen, and instep. This disease occasionally extends to all the parts above-mentioned at the same time; but, in general, it affects them successively, leaving one place free, and appearing in others; sometimes again returning to its first situation. The psoriasis diffusa is attended with a sensation of heat, and with a very troublesome itching, especially at night. It exhibits small, slight, distinct scales,

having less disposition than the lepra to form thick crusts. The chaps or fissures of the skin, which usually make a part of this complaint, are very sore and painful, but seldom discharge any fluid. When the scales are removed by frequent washing, or by the application of unguents, the surface, though raised and uneven, appears smooth and shining; and the deep furrows of the cuticle are lined by a slight scaliness. Should any portion of the diseased surface be forcibly excoriated, there issues out a thin lymph, mixed with some drops of blood, which slightly stains and stiffens the linen, but soon concretes into a thin, dry scab; this is again succeeded by a white scaliness, gradually increasing, and spreading in various directions. As the complaint declines, the roughness, chaps, scales, &c. disappear, and a new cuticle is formed, at first red, dry, and shrivelled, but which, in two or three weeks, acquires the proper texture. The duration of the psoriasis diffusa is from one to four months. If, in some constitutions, it does not then disappear, but becomes, to a certain degree, permanent, there is, at least, an aggravation or extension of it, about the usual periods of its return. In other cases, the disease, at the vernal returns, differs much as to its extent, and also with respect to the violence of the preceding symptoms. The eruption is, indeed, often confined to a single scaly patch, red, itching, and chapped, of a moderate size, but irregularly circumscribed. This solitary patch is sometimes situated on the temple, or upper part of the cheek, frequently on the breast, the calf of the leg, about the wrist, or within and a little below the elbow joint, but especially at the lower part of the thigh, behind. It continues in any of these situations several months, without much observable alteration. The complaint denominated with us the baker's itch, is an appearance of psoriasis diffusa on the back of the hand, commencing with one or two small, rough, scaly patches, and finally extending from the knuckles to the wrist. The rhagades, or chaps and fissures of the skin, are numerous about the knuckles and ball of the thumb, and where the back of the hand joins the wrist. They are often highly inflamed, and painful, but have no discharge of fluid from them. The back of the hand is a little raised or tumefied, and, at an advanced period of the disorder, exhibits a reddish, glossy surface, without crusts or numerous scales. However, the deep furrows of the cuticle are, for the most part, whitened by a slight scaliness. This complaint is not general among bakers; that it is only aggravated by their business, and affects these who are otherwise disposed to it, may be collected from the following circumstances: 1. It disappears

about Midsummer, and returns in the cold weather at the beginning of the year; 2. Persons constantly engaged in the business, after having been once affected with the eruption, sometimes enjoy a respite from it for two or three years; 3. When the business is discontinued, the complaint does not immediately cease. The grocers' itch has some affinity with the bakers' itch, or tetter; but, being usually a pustular disease at its commencement, it properly belongs to another genus. Washerwomen, probably from the irritation of soap, are liable to be affected with a similar scaly disease on the hands, and arms, sometimes on the face and neck, which, in particular constitutions, proves very troublesome, and of long duration.

3. The *Psoriasis gyrata* is distributed in narrow patches or stripes, variously figured; some of them are nearly longitudinal; some circular, or semicircular with vermiform appendages; some are tortuous, or serpentine; others like earth-worms or leeches: the furrows of the cuticle being deeper than usual, make the resemblance more striking, by giving to them an annulated appearance. There is a separation of slight scales from the diseased surface, but no thick incrustations are formed. The uniform disposition of these patches is singular; I have seen a large circular one situated on each breast above the papilla; and two or three others of a serpentine form, in analogous situations along the sides of the chest. The back is often variegated in like manner, with convoluted tetter, similarly arranged on each side of the spine. They likewise appear, in some cases, on the arms and thighs, intersecting each other in various directions. A lighter kind of this complaint affects delicate young women and children in small scaly circles or rings, little discoloured; they appear on the cheeks, neck, or upper part of the breast, and are mostly confounded with the herpetic, or pustular ring-worm. The *psoriasis gyrata* has its remissions and returns, like the *psoriasis diffusa*; it also exhibits, in some cases, patches of the latter disorder on the face, scalp, or extremities, while the trunk of the body is chequered with the singular figures above described.

4. *Psoriasis palmaria*. One very obstinate species of tetter is nearly confined to the palm of the hand. It commences with a small, harsh, or scaly patch, which gradually spreads over the whole palm, and sometimes appears in a slighter degree on the inside of the fingers and wrist. The surface feels rough from the detached and raised edges of the scaly lamina; its colour often changes to brown, or black, as if dirty; yet the most diligent washing produces no favourable effect. The cuticular furrows are deep, and cleft at the bottom longitudinally, in various places, so as to

bleed on stretching the fingers. A sensation of heat, pain, and stiffness in the motions of the hand, attends this complaint. It is worst in winter or spring, and occasionally disappears in autumn or summer, leaving a soft, dark-red cuticle; but many persons are troubled with it for a series of years, experiencing only very slight remissions. Every return or aggravation of it is preceded by an increase of heat and dryness, with intolerable itching. Shoemakers have the *psoriasis palmaria* locally, from the irritation of the wax they so constantly employ. In braziers, tinmen, silversmiths, &c. the complaint seems to be produced by handling cold metals. A long predisposition to it from a weak, languid, hectic state of the constitution may give effect to different occasional causes. Dr. Willan has observed it in women after lying-in; in some persons it is connected or alternates with arthritic complaints. When the palms of the hands are affected as above stated, a similar appearance often takes place on the soles of the feet; but with the exception of rhagades or fissures, which seem less liable to form there, the feet being usually kept warm and covered. Sometimes, also, the *psoriasis palmaria* is attended with a thickness of the præputium, with scaliness and painful cracks. These symptoms at last produce a phymosis, and render connubial intercourse difficult or impracticable; so great, in some cases, is the obstinacy of them; that remedies are of no avail, and the patient can only be relieved by circumcision. This affection of the præputium is not exactly similar to any venereal appearance, but rhagades or fissures, and indurated patches within the palm of the hand, take place in syphilis, and somewhat resemble the *psoriasis palmaria*. The venereal patches are, however, distinct, white, and elevated, having nearly the consistence of a soft corn. From the rhagades there is a slight discharge, very offensive to the smell. The soles of the feet are likewise, in this case, affected with the patches, not with rhagades. When the disease yields to the operation of mercury, the indurated portions of cuticle separate, and a smooth new cuticle is found formed underneath. The fingers and toes are not affected with the patches, &c. in venereal cases.

5. *Psoriasis labialis*. The *psoriasis* sometimes affects the prolabium without appearing on any other part of the body. Its characteristics are, as usual, scaliness, intermixed with chaps and fissures of the skin. The scales are of a considerable magnitude, so that their edges are often loose, while the central points are attached, a new cuticle gradually forms beneath the scales, but is not durable. In the course of a few hours it becomes dry, shrivelled, and broken; and, while it ex-

foliates, gives way to another layer of tender cuticle, which soon, in like manner, perishes. These appearances should be distinguished from the light chaps and roughness of the lips produced by very cold or frosty weather, but easily removed. The psoriasis labialis may be a little aggravated by frost or sharp winds, yet it receives no material alleviation from an opposite temperature. It is not, indeed, confined within any certain limit, or period of duration, having, in several instances, been protracted through all the seasons. The under lip is always more affected than the upper; and the disease takes place more especially in those persons whose lips are full and prominent.

6. *Psoriasis scrotalis*. The skin of the scrotum may be affected in the psoriasis diffusa like other parts of the surface of the body; but sometimes a roughness and scaldiness of the scrotum appears as an independent complaint, attended with much heat, itching, tension, and redness. The above symptoms are succeeded by a hard, thickened, brittle texture of the skin, and by painful chaps or excoriations, which are not easy to be healed. This complaint is sometimes produced under the same circumstances as the prurigo scroti, and appears to be in some cases a sequel of it. A species of the psoriasis scrotalis likewise occurs in the lues venerea, but merits no particular attention, being always combined with other secondary symptoms of the disease.

7. *Psoriasis infantilis*. Infants between the ages of two months and two years, are occasionally subject to the dry tetter. Irregular, scaly patches, of various sizes, appear on the cheeks, chin, breast, back, nates, and thighs. They are sometimes red, and a little rough, or elevated; sometimes excoriated, then again covered with a thin incrustation; and lastly intersected by chaps or fissures. The general appearances nearly coincide with those of the psoriasis diffusa; but there are several peculiarities in the tetter of infants which require a distinct consideration.

8. *The Psoriasis inveterata*, is characterised by an almost universal scaldiness, with a harsh, dry, and thickened state of the skin. It commences from a few irregular, though distinct patches on the extremities. Others appear afterwards on different parts, and, becoming confluent, spread at length over all the surface of the body, except a part of the face, or sometimes the palms of the hands, and soles of the feet. The skin is red, deeply furrowed or wrinkled, stiff and rigid, so as somewhat to impede the motion of the muscles, and of the joints. So quick, likewise, is the production and separation of scales, that large quantities of them are found in the bed on which a person affected

with the disease has slept. They fall off in the same proportion by day, and being confined within the linen, excite a troublesome and perpetual itching.

PSORICA. (From *ψαγε*, the itch.) Medicines to cure the itch.

PSOROPHTHALMIA. (From *ψαρο*, a scab, and *οφθαλμος*, an eye.) An inflammation of the eye-lids, attended with ulcerations, which itch very much. By psorophthalmia Mr. Ware means a case, in which the inflammation of the eye-lids is attended with an ulceration of their edges, upon which a glutinous matter lodges, incrusts, and becomes hard, so that in sleep, when they have been long in contact, they become so adherent, that they cannot be separated without pain. The proximate cause is an acrimony deposited in the glands of the eyelids. The species of the psorophthalmia are:

1. *Psorophthalmia crustacea*, which forms dry or humid crusts in the margins of the eyelids.

2. *Psorophthalmia herpetica*, in which small papulæ, itching extremely, and terminating in scurf, are observed.

PSYRAGOGICA. (From *ψυχε*, the mind, and *αγω*, to move.) Medicines which recover in syncope or apoplexy.

PSYCHOTRIA KNETICA. (From *ψυχος*, cold, and *κρηνω*, to nourish, because it grows in cold places.) See *Ipecacuanha*.

ΨΥΧΟΤΡΟΦΗΝ. (From *ψυχος*, cold, and *τροφη*, to nourish; so called because it grows in places exposed to the cold.) The herb betony.

ΨΥΧΟΛΟΥΤΗΚ. (From *ψυχε*, cold, and *λουω*, to wash.) A cold bath.

ΨΥΚΤΙΚΑ. (From *ψυχε*, to refrigerate.) Refrigerating medicines.

ΨΥΔΡΑCΙÆ. (From *ψυχε*, cold.) Red and somewhat elevated spots, which soon form broad and superficial vesicles, such as those produced by the stinging-nettle, the bites of insects, &c. See *Pustule*.

PSYLLIUM. (From *ψαλλος*, a flea; so called because it was thought to destroy fleas.) *Pulicaris herba*. *Crystallion* and *cynomoia* of Orisbanus. Flea-wort. The seeds of this plant, *Plantago psyllum* of Linnæus:—*caule ramoso herbaceo, foliis subdentatis, recurvatis; capitulis aphyllis*, have a nauseous mucilaginous taste, and no remarkable smell. The decoction of the seeds is recommended in hoarseness and asperity of the fauces.

PTARICA. (From *πταγω*, to sneeze; so called because it irritates the nose, and provokes sneezing.) *Pseudopyrethrum*. *Pyrethrum sylvestre*. *Draco sylvestris*. *Tarthen sylvestris*. *Sternutamentoria*. *Dracunculus pratensis*. Sneezewort. Bastard pellitory. *Achillea ptarmica* of Linnæus:—*foliis lanceolatis, acuminatis, argute serratis*. The flowers and roots of this plant

have a hot biting taste, approaching to that of pyrethrum, with which they also agree in their pharmaceutical properties. Their principal use is as a masticatory and sternutatory.

PTERIS. The name of a genus of plants in the Linnæan system. Class, *Cryptogamia*. Order, *Filices*.

PTERIS AQUILINA. (From *πτερον*, wing; so called from the likeness of its leaves to wings, and *aquilina*, from *aquila*, an eagle, from its resemblance to an eagle's wings.) The systematic name of the common brake, or female fern. See *Felix femina*.

PTEROCARPUS The name of a genus of plants in the Linnæan system.

PTEROCARPUS SANTALINUS. The systematic name of the red jaunder's tree. See *Santalum rubrum*.

PTERYGIUM. (*Πτερυξ*, a wing.) A membranous excrescence which grows upon the internal canthus of the eye chiefly, and expands itself over the albuginea and cornea towards the pupil. It appears to be an extension or prolongation of the fibres and vessels of the caruncula lachrymalis, or semilunar membrane, appearing like a wing. The species of pterygium are four:

1. *Pterygium tenue*, seu *ungula*, is a pellucid pellicle, thin, of a cineritious colour, and unpainful; growing out from the caruncula lachrymalis, or membrana semilunaris.

2. *Pterygium crassum*, seu *pannus*, differs from the ungula, by its thickness, red colour, and fullness of the red vessels on the white of the eye, and it stretches over the cornea like fasciculi of vessels.

3. *Pterygium malignum*, is a pannus of various colours, painful, various, and arising from a cancerous acrimony.

4. *Pterygium pingue*, seu *pinguicula*, is a mollicie like lard or fat, soft, without pain, and of a light yellow colour, which commonly is situated in the external angle of the eye, and rarely extends to the cornea; but often remains through life.

PTERYGO. Names compounded of this word belong to muscles which are connected with the pterygoid process of the sphenoid bone; as *pterygo-pharyngeus*, &c.

Pterygo-pharyngeus. See *Constrictor pharyngis superior*.

Pterygo-staphilinus externus. See *Levator palati*.

PTERYGOID PROCESS. (*Pterygoideus*, from *πτερυξ*, a wing, and *σῆμα*, resemblance.) A wing-like process of the sphenoid bone.

PTERYGOIDEUM OS. See *Ethmoid bone*.

PTERYGOIDEUS EXTERNUS. (*Pterygoideus*, from its belonging to the process pterygoideus.) *Pterygoideus minor* of Winslow. *Pterygo-coli-maxillaire* of Dumas. A muscle placed, as it were, hori-

zontally along the basis of the scull, between the pterygoid process and the condyle of the lower jaw. It usually arises by two distinct heads; one of which is thick, tendinous, and fleshy, from the outer wing of the pterygoid process of the os sphenoides, and from a small part of the os maxillare adjoining to it; the other is thin and fleshy, from a ridge in the temporal process of the sphenoid bone, just behind the slit that transmits the vessels to the eye. Sometimes this latter origin is wanting, and, in that case, part of the temporal muscle arises from this ridge. Now and then it affords a common origin to both these muscles. From these origins the muscle forms a strong fleshy belly, which descends almost transversely outwards and backwards, and is inserted tendinous and fleshy into a depression in the fore part of the condyloid process of the lower jaw, and into the anterior surface of the capsular ligament that surrounds the articulation of that bone. All that part of this muscle, which is not hid by the pterygoideus internus, is covered by a ligamentous expansion, which is broader than that belonging to the pterygoideus internus, and originates from the inner edge of the glenoid cavity of the lower jaw, immediately before the styloid process of the temporal bone, and extends obliquely downwards, forwards, and outwards, to the inner surface of the angle of the jaw. When these muscles act together, they bring the jaw horizontally forwards. When they act singly, the jaw is moved forwards, and to the opposite side. The fibres that are inserted into the capsular ligament, serve likewise to bring the moveable cartilage forwards.

PTERYGOIDEUS INTERNUS. *Pterygoideus major* of Winslow. *Pterygo-anguli-maxillaire* of Dumas. This muscle arises tendinous and fleshy from the whole inner surface of the external ala of the pterygoid process, filling all the space between the two wings; and from that process of the os palati that makes part of the pterygoid fossa. From thence growing larger, it descends obliquely downwards, forwards, and outwards, and is inserted, by tendinous and fleshy fibres, into the inside of the lower jaw, near its angle. This muscle covers a great part of the *pterygoideus externus*; and along its posterior edge we observe a ligamentous band, which extends from the back part of the styloid process to the bottom of the angle of the lower jaw. The use of this muscle is to raise the lower jaw, and to pull it a little to one side.

PTERYGOIDEUS MAJOR. See *Pterygoideus internus*.

PTERYGOIDEUS MINOR. See *Pterygoideus externus*.

PTILOSI. (From *πλος*, bald.) See *Madurosia*.

PTISANA. (From *πτισσιν*, to decorticate, bruise or pound.) *Ptissane*. Barley deprived of its husks, pounded and made into balls.

PTOSIS. (From *πτω*, to fall.) *Blepharoptosis*. An inability of raising the upper eye-lid. The affection may be owing to several causes, the chief of which are a redundancy of the skin on the eye-lid; a paralytic state of the levator muscle, and a spasm of the orbicularis.

PTOSIS IRIDIS. *Prolapsus iridis*. A prolapsus of the iris through a wound of the cornea. It is known by a blackish tubercle, which promines a little from the cornea in various forms. The species of the ptosis of the iris are,

1. *Ptosis recens*, or a recent ptosis from a side wound of the cornea, as that which happens, though rarely, in or after the extraction of the cataract.

2. *Ptosis inveterata*, in which the incarcerated prolapsed iris is grown or attached to the wound or ulcer, and by the air has become callous or indurated.

PTYALAGOGA. (From *πτυαλον*, spittle, and *αγω*, to excite.) *Ptyalagoga*. Medicines which promote a discharge of the saliva, or cause salivation.

PTYALISMOS. See *Ptyalismus*.

PTYALISMUS. (From *πτυαλίζω*, to spit.) A ptyalism or salivation, or increased secretion of saliva from the mouth.

PTYALUM. (From *πλυω*, to spit up.) The saliva or mucus from the bronchia.

PTYASMAGOGA. (From *πτυασμα*, sputum, and *αγω*, to expel.) See *Ptyalagoga*.

PUBES. The external part of the organs of generation of both sexes, which are covered with hair.

PUBIS OS. A bone of the foetal pelvis. See *Innomination os*.

PUDENDA. (From *pudor*, shame.) The parts of generation.

PUDENDAGRA. (From *pudenda*, the private parts, and *αγω*, a seizure.) *Cedma*. A pain in the private parts. By some it is called the venereal disease. Others define it to be pain or uneasiness in the genital parts of men or women, somewhat resembling a diarrhoea, but without a dysuria. Dr. Berdoe asserts, in his Essay on the Pudendagra, that it is distinct from the venereal disease, and also, that it is proper to women, but that a woman labouring under it, can communicate some inflammatory symptoms to the penis of a man who cohabits with her.

PUDENDUM MULIEBRE. The female parts of generation.

PUDICAL ARTERY. *Arteria pudica vel pudenda*. Pudendal artery. A branch of the internal iliac distributed on the organs of generation.

PURILIS MORBUS. The epilepsy.

PUERPERAL FEVER. Child-bed fever. Cullen considers this disease as a species of continued fever.

Puffball. See *Lycoperdon*.

PUGILLUS. (From *pugnus*, the fist.) *Dracma*. A pugil. The eighth part of a handful.

PULBGIUM. (From *pulex*, a flea; because the smell of its leaves, burnt, destroys fleas.) *Pulegium regale*. *Pulegian latifolium glechon*. Poddling-grass. Common pennyroyal. *Mentha pulegium* of Linnaeus:—*floribus verticillatis, foliis ovatis obtusis subcrenatis, caulibus subteretibus repentibus*. This plant is considered as a carminative, stomachic, and emmenagogue; and is in very common use in hysterical disorders. The official preparations of pennyroyal are, a simple water, a spirit, and an essential oil.

PULBGIUM GRÆVILLUM. Hart's pennyroyal. *Mentha cervina* of Linnaeus. This plant possesses the virtues of pennyroyal in a very great degree; but is remarkably unpleasant. It is seldom employed but by the country people, who substitute it for pennyroyal.

PULICARIA. (From *pulex*, a flea; so named because it was thought to destroy fleas if hung in a chamber.) See *Pygium*.

PULMO. (Plin. *πνευμον*. Attice *πνευμον*, unde, per metathesin *pulmo*.) Lang. See *Lung*.

PULMONARIA. (From *pulmo*, the lungs; so called because of its virtues in affections of the lungs.) The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*. Lungwort.

* **PULMONARIA ARBorea.** *Musci pulmonarius quertinus*. This subastringent, and rather acid plant, *Lichen pulmonarius* of Linnaeus, was once in high estimation in the cure of diseases of the lungs, especially coughs, asthmas, and catarrhs. Its virtues are similar, and in no way inferior to those of the lichen islandicus.

PULMONARIA MACULATA. *Symphytum maculatum*. Spotted lung-wort. Jerusalem cowslips. Jerusalem sage. *Pulmonaria officinalis* of Linnaeus. This plant is rarely found to grow wild in England; but is very commonly cultivated in gardens, where its leaves become broader, and approach more to a cordate shape. The leaves, which are the part medicinally used, have no peculiar smell; but, in their recent state, manifest a slightly astringent and mucilaginous taste; hence it seems not wholly without foundation that they have been supposed to be demulcent and pectoral. They have been recommended in hemoptoës, trickling coughs, and catarrhal defluxions upon the lungs. The name *pulmonaria*, however,

seems to have arisen rather from the speckled appearance of these leaves, resembling that of the lungs, than from any intrinsic quality which experience discovered to be useful in pulmonary complaints.

PULMONARIA OFFICINALIS. The systematic name of the spotted lung-wort. See *Pulmonaria maculata*.

Pulmonary consumption. See *Phthisis*.

PULMONARY VESSELS. The pulmonary artery, *arteria pulmonalis*, arises from the right ventricle of the heart, and soon divides into the right and left, which ramify throughout the lungs, and form a beautiful net-work on the air vesicles, where they terminate in the veins, *vene pulmonales*, whose branches at length form four trunks, which empty themselves into the left auricle of the heart.

PULMONIA. (From *pulmo*, the lungs) An inflammation of the lungs.

PULMONICA. (From *pulmo*, the lungs.) Medicines for the lungs.

PULSATILLA NIGRIFANS. (From *pulso*, to beat about; so called from its being perpetually agitated by the air.) This plant, *Anemone pratensis* of Linnaeus:—*pedunculo involucreto, petalis apice reflexis, foliis bipinnatis*, has been received into the Edinburgh Pharmacopœia upon the authority of Baron Stœrck, who recommended it as an effectual remedy for most of the chronic diseases affecting the eye, particularly amaurosis, cataract, and opacity of the cornea, proceeding from various causes. He likewise found it of great service in venereal nodes, nocturnal pains, ulcers, caries, indurated glands, suppressed menses, seriginous eruptions, melancholy, and palsy. The plant, in its recent state, has scarcely any smell; but its taste is extremely acrid, and, when chewed, it corrodes the tongue and fauces.

PULSE. *Pulsus*. The beating of the artery at the wrist is termed the pulse. It depends upon, and is synchronous with, that of the heart: hence physicians feel the pulse, to ascertain the quickness or tardiness of the blood's motion, the strength of the heart, &c.

PULSILEGIUM. (From *pulsus*, the pulse, and *lego*, to tell.) An instrument for measuring the pulse.

PULVINAR. (From *pulvis*, dust or chaff, with which they are filled.) A medicated cushion.

PULVINARIUM. See *Pulvinar*.

PULVIS. A powder. *Pulvinarium* This form of medicine is either coarse or very fine, simple or compound. In the compounded powders the intimate and complete admixture of the several ingredients, and more especially of those to which any of the more active substances, as opium, scammony, &c. are added, can-

not be too strongly recommended, and for this purpose it may be proper to pass them, after they are mixed mechanically, through a fine sieve.

PULVIS ALOES COMPOSITUS. Compound powder of aloë. Formerly called *pulvis aloes cum guaiaco*. *Pilule aromatics, pilule de diambre*. "Take of extract of spiked aloë, an ounce and half; guaiacum gum resin, an ounce; compound powder of cinnamon, half an ounce." Powder the extract of aloë and guaiacum gum resin separately: then mix them with the compound powder of cinnamon. The dose is from gr. x. to ℥j. It is a warm aperient, laxative powder, calculated for the aged, and those affected with dyspeptic gout, attended with costiveness and spasmodic complaints of the stomach and bowels.

PULVIS ALOES CUM CANELLA. A cathartic, deobstruent powder, possessing stimulating and aloetic properties omitted in the last London Pharmacopœia, as rather suited to the purpose of extemporaneous prescription.

PULVIS ALOES CUM FERRO. This possesses aperient and deobstruent virtues; and is mostly given in chlorosis and constipation. In the London Pharmacopœia this prescription is omitted for the same reason as *pulvis aloes cum canella*.

PULVIS ALOES CUM GUAIACO. See *Pulvis aloes compositus*.

PULVIS ANTIMONIALIS. Antimonial powder. "Take of sulphuret of antimony, powdered, a pound; hartshorn shavings, two pounds." Mix and throw them into a broad iron pot heated to a white heat, and stir the mixture constantly until it acquires an ash colour. Having taken it out, reduce it to powder, and put it into a coated crucible, upon which another inverted crucible, having a small hole in its bottom, is to be luted. Then raise the fire by degrees to a white heat, and keep it so for two hours. Reduce the residuary mass to a very fine powder. The dose is from five to ten grains. It is in high esteem as a febrifuge, sudorific, and antispasmodic. The diseases in which it is mostly exhibited are, most species of asthenic and exanthematous fevers, acute rheumatism, gout, diseases arising from obstructed perspiration, dysuria, nervous affections, and spasms.

This preparation was introduced into the former London Pharmacopœia as a substitute for a medicine of extensive celebrity, Dr. James's powder; to which, however, the present form more nearly assimilates in its dose, and it is more manageable in its administration, by the reduction of the proportion of antimony to one half.

PULVIS AROMATICUS. See *Pulvis cinnamomi compositus*.

PULVIS CERUÆ COMPOSITUS. This is mostly used in the form of collyrium, lotion, or injection, as a mucilaginous sedative.

PULVIS CHELARUM CANCRI COMPOSITUS. An antacid and adstringent powder, mostly given to children with diarrhœa and acidity of the primæ viæ.

PULVIS CINNAMOMI COMPOSITUS. Compound powder of cinnamon. Formerly called *pulvis aromaticus: species aromaticæ: species diambæ sine odoratis*. "Take of cinnamon bark, two ounces; cardamom-seeds, an ounce and half; ginger-root, an ounce; long pepper, half an ounce." Rub them together, so as to make a very fine powder. The dose is from five to ten grains. An elegant stimulant, carminative, and stomachic powder.

PULVIS CONTRAYERVÆ COMPOSITUS. "Take of contrayerva root, powdered, five ounces; prepared shells, a pound and half." Mix. A febrifuge diaphoretic, mostly given in the dose of from one to two scruples in slight febrile affections.

PULVIS CORNU USTI CUM OPIO. Powder of burnt hartshorn with opium. *Pulvis opiatus*. "Take of hard opium, powdered, a drachm; hartshorn, burnt and prepared, an ounce; cochineal, powdered, a drachm." Mix. This preparation affords a convenient mode of exhibiting small quantities of opium, ten grains containing one of the opium. It is absorbent and anodyne.

PULVIS CRETÆ COMPOSITUS. Compound powder of chalk. *Pulvis e bolo compositus sine opio. Species e scordio sine opio. Diacordium, 1720.* "Take of prepared chalk, half a pound; cinnamon bark, four ounces; tormentill root, acacia gum, of each three ounces; long pepper, half an ounce." Reduce them separately into a very fine powder, and then mix. The dose is from ʒss. to ʒi. An adstringent, carminative, and stomachic powder, exhibited in the cure of diarrhœa, pyrosis, and diseases arising from acidity of the bowels, inducing much pain.

PULVIS CRETÆ COMPOSITUS CUM OPIO. Compound powder of chalk with opium. *Pulvis e bolo compositus cum opio. Species e scordio cum opio.* "Take of compound powder of chalk, six ounces and a half. Hard opium, powdered, four scruples." Mix. The dose from one scruple to two. The above powder, with the addition of opium, in the proportion of one grain to two scruples.

PULVIS IPECACUANHÆ COMPOSITUS. Compound powder of ipecacuanhæ. "Take of ipecacuanhæ root, powdered, hard opium, powdered, of each a drachm; sulphate of potash, powdered, an ounce." Mix. A diaphoretic powder, similar to that of Dr. Dover, which gained such repute in the cure of rheumatisms, and

other diseases arising from obstructed perspiration and spasm. The dose is from five grains to a scruple.

PULVIS KINO COMPOSITUS. Compound powder of kino. "Take of kino, 15 drachms; cinnamon bark, half an ounce; hard opium, a drachm." Reduce them separately to a very fine powder; and then mix. The proportion of opium this adstringent contains is one part to twenty. The dose is from five grains to a scruple.

PULVIS MYRRHÆ COMPOSITUS. A stimulant, antispasmodic, and emmenagogue powder, mostly exhibited in the dose of from fifteen grains to two scruples, in uterine obstructions and hysterical affections.

PULVIS OPIATUS. See *Pulvis cornu cum opio*.

PULVIS SCAMMONIÆ COMPOSITUS. Compound powder of scammony. *Pulvis comiti Warwicensis.* "Take of scammony gum resin, hard extract of jalap, of each two ounces; ginger-root, half an ounce." Reduce them separately to a very fine powder, and then mix. From ten to fifteen grains or a scruple are exhibited as a stimulating cathartic.

PULVIS SCAMMONII CUM ALOE. A stimulating cathartic, in the dose of from ten to fifteen grains.

PULVIS SCAMMONII CUM CALOMELANÆ. A vermifugal cathartic, in the dose of from ten to fifteen grains.

PULVIS SENNÆ COMPOSITUS. Compound powder of senna. *Pulvis diacenne.* "Take of senna leaves, supertartrate of potash, of each two ounces; scammony gum resin, half an ounce; ginger-root, two drachms." Reduce the scammony gum resin separately, the rest together, to a very fine powder; and then mix. The dose is from one scruple to one drachm. A saline stimulating cathartic.

PULVIS TRAGACANTHÆ COMPOSITUS. Compound powder of tragacanth. *Species diatragacanthæ frigide.* "Take of tragacanth, powdered, acacia gum, powdered, starch, of each an ounce and half; refined sugar, three ounces." Powder the starch and sugar together; then add the tragacanth and acacia gum, and mix the whole. Tragacanth is very difficultly reduced to powder. The dose is from ten grains to a drachm. A very useful demulcent powder, which may be given in coughs, diarrhœas, stranguary, &c.

Pumpion, common. See Cucurbita.

PUNCTA LACHRYMALIA. (From *punctum*, a point.) Lachrymal points. Two small orifices, one of which is conspicuous in each eyelid, at the extremity of the tarsus, near the internal canthus.

PUNCTUM AUREUM. Formerly, when a hernia of the intestines was reduced by

an incision made through the skin and membrana adiposa, quite down to the upper part of the spermatic vessels, a golden wire was fixed and twisted, so as to prevent the descent of any thing down the tunica vaginalis.

PUNICA. The name of a genus of plants in the Linnæan system. Class, *Isosandria*. Order, *Menogynia*.

PUNICA GRANATUM. The systematic name of the pomegranate. See *Granatum*.

PUPIL (*Pupilla*, from *pupa*, a babe; because it reflects the diminished image of the person who looks upon it like a puppet.) The round opening in the middle of the iris, in which we see ourselves in the eye of another.

PUPILLA. See *Pupil*.

PUPILLARIS MEMBRANA. (From *pupilla*, the pupil.) *Pupille velum*. A fine vascular membrane, which in the fœtus of 5, 6, or 7 months has no pupil, grows across the part where the pupil is afterwards seen.

PURGAMENTUM. A purge.

PURGATIVA. *Purgentia*. *Cathartica*. *Catocharthica*. *Catoretica*. *Cutoteretica*. *Dejectoria*. *Abduca*. Purgative medicines.

Purging flax. See *Linum catharticum*.

Purging-nut. See *Ripinus major*.

PURPURA ALBA. *Purpura rubra*. Many writers term the miliary fever, when the pustules are white, *purpura alba*, and when they are red, *purpura rubra*.

PURPURA SCORBUTICA. Petechial eruptions in scurvy.

Purslane. See *Portulaca*.

PUS. Matter. A whitish, bland, cream-like fluid, heavier than water, found in phlegmonous abscesses, or on the surface of sores. It is distinguished, according to its nature, into laudable or good pus, scrophulous, serous, and ichorous pus, &c.

Pus taken from a healthy ulcer, near the source of circulation, as on the arm or breast, Mr. Home observes, readily separates from the surface of the sore, the granulations underneath being small, pointed, and of a florid red colour, and has the following properties: it is nearly of the consistence of cream; is of a white colour; has a mawkish taste; and, when cold, is inodorous; but, when warm, has a peculiar smell. Examined in a microscope, it is found to consist of two parts, of globules, and a transparent colourless fluid; the globules are probably white, at least they appear to have some degree of opacity. Its specific gravity is greater than that of water. It does not readily go into putrefaction. Exposed to heat, it evaporates to dryness; but does not coagulate. It does not unite with water in the heat of the atmosphere, but

falls to the bottom; yet, if kept in a considerable degree of heat, rises and diffuses through the water, and remains mixed with it, even after having been allowed to cool, the globules being decomposed.

Pus varies in its appearance, according to the different circumstances which affect the ulcer that forms it; such as, the degree of violence of the inflammation, also its nature, whether healthy or unhealthy; and these depend upon the state of health, and strength of the parts yielding pus. These changes arise more from indolence and irritability, than from any absolute disease; many specific diseases, in healthy constitutions, producing no change in the appearance of the matter from their specific quality. Thus, the matter from a gonorrhœa, from the small-pox pustules, the chicken pox, and from an healthy ulcer, has the same appearance, and seems to be made up of similar parts, consisting of globules floating in a transparent fluid, like common pus; the specific properties of each of these poisons being superadded to those of pus. Matter from a cancer may be considered as an exception; but a cancerous ulcer is never in a healthy state.

In indolent ulcers, whether the indolence arises from the nature of the parts, or the nature of the inflammation, the pus is made of globules and flaky particles, floating in a transparent fluid; and globules and flakes are in different proportions, according to the degree of indolence: this is particularly observable in scrophulous abscesses, preceded by a small degree of inflammation. That this flaky appearance is no part of true pus, is well illustrated by observing, that the proportion it bears to the globules is greatest where there is the least inflammation; and in those abscesses that sometimes occur, which have not been preceded by any inflammation at all, the contents are wholly made up of a curdy or flaky substance, of different degrees of consistence, which is not considered to be pus, from its not having the properties stated in the definition of that fluid.

The constitution and part must be in health to form good pus; for very slight changes in the general health are capable of producing an alteration in it, and even of preventing its being formed at all, and substituting in its place coagulating lymph.

This happens most readily in ulcers in the lower extremities, owing to the distance of the parts from the source of the circulation, rendering them weaker. And it is curious to observe the influence that distance alone has upon the appearance of pus.

Pus differs from chyle in its globules,

being larger, not coagulating by exposure to the air, nor by heat, which those of chyle do.

The pancreatic juice contains globules, but they are much smaller than those of pus.

Milk is composed of globules, nearly of the same size as those of pus, but much more numerous. Milk coagulates by runnet, which pus does not; and contains oil and sugar, which are not to be discovered in pus.

The cases in which pus is formed are, properly speaking, all reducible to one, which is, the state of parts consequent to inflammation. For, as far as we yet know, observes Mr. Home, pus has in no instance been met with unless preceded by inflammation; and although, in some cases a fluid has been formed independent of preceding inflammation, it differs from pus in many of its properties.

In considering the time required for the formation of pus, it is necessary to take notice of the periods which are found, under different circumstances, to intervene between a healthy or natural state of the parts, and the presence of that fluid after the application of some irritating substance to the skin.

In cases of wounds made into muscular parts, where blood-vessels are divided, the first process which takes place is the extravasation of red blood; the second is the exudation of coagulating lymph, which afterwards becomes vascular; and the third, the formation of matter, which last does not, in common, take place in less than two days; the precise time will, however, vary exceedingly, according to the nature of the constitution, and the state of the parts at the time.

If an irritating substance is applied to a cuticular surface, upon which it raises a blister, pus will be formed in about twenty-four hours.

PUSTULA. (Dim. of *pus*, matter.) See *Pustule*.

PUSTULE. (*Pustula*, a little pimple, from *pus*, corruption.) *Ecthyma*, *Eczema*. Dr. Willan defines a pustule to be an elevation of the cuticle, sometimes globate sometimes conoidal in its form, and containing pus, or a lymph which is in general discoloured. Pustules are various in their size, but the diameter of the largest seldom exceeds two lines. There are many different kinds of pustules, properly distinguished in medical authors, by specific appellations, as 1. *Phlyzaciun*, a small pustule containing pus, and raised on a hard, circular, inflamed base, of a vivid red colour. It is succeeded by a thick, hard, dark-coloured scab. 2. *Ptydraciun*, according to Dr. Willan, a minute pustule, irregularly circumscribed, producing but

a slight elevation of the cuticle, and terminating in a laminated scab. Many of these pustules usually appear together, and become confluent. When mature they contain pus; and, after breaking, discharge a thin watery humour.

POSTULA ORIS. The aphthæ.

PUTAMEN. (From *puto*, to cut.) The bark or paring of any vegetable. The putamen, or green rind of the walnut, has been celebrated as a powerful antivenereal remedy, for more than a century and a half; and Petrus Borellus has given directions for a decoction not unlike that which is commonly called the Lisbon diet-drink, in which the walnut, with its green bark, forms a principal ingredient. Ramazzini, whose works were published early in the present century, has likewise informed us, that in his time the green rind of the walnut was esteemed a good antivenereal remedy in England. This part of the walnut has been much used in decoctions, during the last fifty years, both in the green and dried state; it has been greatly recommended by writers on the continent, as well as by those of our own country; and is, without doubt, a very useful addition to the decoction of the woods. Mr. Pearson has employed it during many years, in those cases where pains in the limbs and indurations of the membranes have remained, after the venereal disease has been cured by mercury; and he informs us, that he has seldom directed it without manifest advantage.

Brambilla and Girtanner also contend for the antivenereal virtues of the green bark of the walnut; but the result of Mr. P.'s experience will not permit him to add his testimony to theirs. I have given it, says he, in as large doses as the stomach could retain, and for as long a time as the strength of the patients, and the nature of their complaints, would permit; but I have uniformly observed, that if they who take it be not previously cured of *lues venerea*, the peculiar symptoms will appear, and proceed in their usual course, in defiance of the powers of this medicine. The *Decoctum Lunitanicum* may be given with great advantage in many of those cutaneous diseases which are attended with aridity of the skin; and I have had some opportunities of observing, that when the putamen of the walnut has been omitted, either intentionally or by accident, the same good effects have not followed the taking of the decoction, as when it contained this ingredient.

PUTREFACTION. Putrid fermentation. Putrefactive fermentation. That process by which a substance is decomposed and dissipated in the air in the form of putrid gas. Every living body, when deprived of life, performs a retrograde process, and becomes

decomposed. This is called fermentation in vegetables, and putrefaction in animals. The same causes, the same agents, and the same circumstances, determine and favour the decomposition in vegetables and animals, and the difference of the products which are obtained, arises from the difference of the constituent parts of each. The requisites to this process are, 1. A certain degree of humidity. 2. The access of atmospheric air. 3. A certain degree of heat. See also *Fermentation*.

Putrid fever. A species of typhus. See *Typhus gravior*.

PYLORIC ARTERY. *Arteria pylorica*. A branch of the hepatic artery.

PYLORUS. (From *πύλωρ*, to guard an entrance; because it guards, as it were, the entrance of the bowels.) *Juniter*. *Portorium*. *Ostium*. The inferior aperture of the stomach, which opens into the intestines.

PROPTICA. (From *πύωρ*, pus, and *ποιω*, to make.) Suppurative medicines.

PROTHREA. (From *πύωρ*, pus, and *γίω*, to flow.) A purulent discharge from the belly.

PYOTURIA. (From *πύωρ*, pus, and *ύωρ*, urine.) *Pyuria*. A mucous or purulent urine.

PYRAMIDALIS. (*Pyramidalis*, sc. *musculus*; from *πυγάμικ*, a pyramid.) Fallopius, who is considered as the first accurate describer of this muscle, first gave it the name of *pyramidalis*, from its shape, hence it is called *pyramidalis Fallopii* by Douglas. But Vesalius seems to have been acquainted with it, and to have described it as a part of the rectus. It is called *pyramidalis vel succenturiatus* by Cowper. And *pubio-umbilical* by Dumas. It is a very small muscle, situated at the bottom of the fore part of the rectus, and is covered by the same aponeurosis that forms the anterior part of the sheath of that muscle. It arises, by short tendinous fibres, from the upper and fore part of the pubis. From this origin, which is seldom more than an inch in breadth, its fibres ascend somewhat obliquely, to be inserted into the linea alba, and inner edge of the rectus, commonly at about the distance of two inches from the pubis, and frequently at a greater or less distance, but always below the umbilicus. In some subjects the pyramidalis is wanting on one or both sides, and when this happens, the internal oblique is usually found to be of greater thickness at its lower part. Now and then, though rarely, there are two at one side, and only one at the other, and M. Sabattier has even seen two on each side. Fallopius, and many others after him, have considered it as the congener of the internal oblique; but its use seems to be to assist the lower part of the rectus.

PYRAMIDALES FACIÆ. See *Levator labii superioris alicque nasi*.

PYRENOIDES. (From *πύρην*, a kernel, and *ειδος*, likeness; so called from its kernel-like shape.) Applied to the process odontoid of the second vertebra.

PYRETERIUM. (From *πύρ*, fire, and *τήρ*, to keep.) The fire-hole of a furnace.

PYRETHRUM. (From *πύρ*, fire, because of the hot taste of its root.) *Buphthalmum creticum*. *Bellis montana putescens acris*. *Dentaria*. *Herba salivaria*. *Pes alexandrinus*. Pellitory of Spain. *Anthemis pyrethrum* of Linnæus:—*caulibus simplicibus unifloris decumbentibus, foliis pinato-multifidis*. This root, though cultivated in this country, is generally imported from Spain. Its taste is hot and acrid, its acrimony residing in a resinous principle. The ancient Romans, it is said, employed the root of this plant as a pickle. In its recent state, it is not so pungent as when dried, and yet, if applied to the skin, it produces inflammation. Its qualities are stimulant; but it is never used, except as a masticatory, for relieving tooth-aches, rheumatic affections of the face, and paralysis of the tongue, in which it affords relief by stimulating the excretory ducts of the salivary glands.

PYRETHRUM SILVESTRE. See *Ptar-mica*.

PYRETOLOGY. (*Pyretologia*; from *πύρ*, fire, or heat, and *λογος*, a discourse.) A discourse, or doctrine on fevers.

PYREXIA. (From *πύρ*, fire.) Fever.

PYREXIÆ. Febrile diseases. The first class of Cullen's nosology; characterised by frequency of pulse after a cold shivering, with increase of heat, and especially, among other impaired functions, a diminution of strength.

PYRIFORMIS. (From *pyrus*, a pear, and *forma*, a shape, shaped like a pear.) *Pyiformis*, seu *iliacus externus* of Douglas and Cowper. Spigelius was the first who gave a name to this muscle, which he called *pyiformis*, from its supposed resemblance to a pear. It is the *pyiformis sive pyramidalis* of Winslow, and *sacro-trochanterien* of Dumas. A small radiated muscle, situated under the glutæus maximus, along the inferior edge of the glutæus minimus. It arises by three and sometimes four tendinous and fleshy origins, from the anterior surface of the second, third, and fourth pieces of the os sacrum, so that this part of it is within the pelvis. From these origins the muscle grows narrower, and passing out of the pelvis, below the niche in the posterior part of the ilium, from which it receives a few fleshy fibres, is inserted by a roundish tendon of an inch in length, into the upper part of the cavity at the root of the trochanter major. The use of this muscle is to assist in moving the

thigh outwards, and moving it a little upwards.

PYRITES. (From *pyr*, fire; so called because it strikes fire with steel.) A metallic substance, formed of iron united with sulphur, from which all the sulphur of commerce is obtained.

PYRITES ARSENICALES. See *Arsenic*.

PYRMONT WATER. *Aqua pyrmontana*. A celebrated mineral spring at Pyrmont, a village in the circle of Westphalia, in Germany. It is of an agreeable though strongly acidulated taste, and emits a large portion of gas; which affects the persons who attend at the well, as well as those who drink the fluid, with a sensation somewhat resembling that produced by intoxication. A general view of the analysis of this water will shew that it stands the first in rank of the highly carbonated chalybeates, and contains such an abundance of carbonic acid, as not only to hold dissolved a number of carbonic salts, but to shew all the properties of this acid uncombined, and in its most active form. Pyrmont water is likewise a strong chalybeate, with regard to the proportion of iron; and it is besides a very hard water, containing much selenite and earthy carbonates. The diseases to which this mineral water may be advantageously applied, are the same as those for which the Spa, and others of the acidulated chalybeates, are resorted to, that is, in all cases of debility that require an active tonic that is not permanently heating; various disorders in the alimentary canal, especially bilious, vomiting, and diarrhoea, and complaints that originate from obstructed menstruation. At Pyrmont, the company generally drink this water by glassfuls, in a morning, to the quantity of two, three, or more English pints. Its common operation is by urine; but, if taken copiously, it generally proves laxative; and when it has not this effect, and that effect is wanted, they commonly mix, with the first glass drunk in the morning, from one to five or six drachms of some purging salt.

PYROLA. (From *pyrus*, a pear; so named because its leaves resemble those of the pear-tree.) 1. The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Monogynia*.

2. The pharmacopœial name of the round-leaved wintergreen. This elegant little plant, *Pyrola rotundifolia* of Linnæus, is now forgotten in the practice of medicine. It possesses gently adstringent

qualities, and has a somewhat bitter taste.

PYROLA ROTUNDIFOLIA. The systematic name of the wintergreen. See *Pyrola*.

PYRO-LIGNEOUS ACID. *Acidum pyro-ligneum*. An acid liquor of a brown colour, of a pretty strong and peculiar smell, obtained by distillation from wood, especially the beech, birch, and box. It is thought to be the acetic acid.

PYROMETER. (From *pyr*, fire, and *metron*, measure.) An instrument to measure those higher degrees of heat to which the thermometer cannot be applied. See *Caloric*.

PYRO-MUCOUS ACID. *Acidum pyro-mucosum*. Syrupous acid. The acid liquor obtained by distillation from insipid, saccharine, gummy, farinaceous mucilages. The celebrated Gren is of opinion, that it is a mixture of acetic with oxalic acid, and does not deserve to be received in the system of chemistry as a peculiar acid.

PYRO-TARTROUS ACID. *Acidum pyro-tartrosum*. See *Tartar*, spirit of.

PYROSIS. (From *pyro*, to burn.) *Pyrosis Suecica* of Sauvages. *Cardialgia sputatoria* of Linnæus. A disease called in Scotland the water-brash; in England, black-water. A genus of disease in the class *neuroses* and order *spasmi* of Cullen; known by a burning pain in the stomach, attended with copious eructation, generally of a watery insipid fluid.

PYROTECHNIA. (From *pyr*, fire, and *technia*, an art.) Chemistry, or that art by which the properties of bodies are examined by fire.

PYROTICA. (From *pyro*, to burn.) Caustics.

PYRUS. The name of a genus of plants in the Linnæan system. Class, *Icosandria*, Order, *Pentagynia*.

PYRUS CYDONIA. The systematic name of the quince tree. See *Cydonium malum*.

PYRUS MALUS. The systematic name of the apple-tree. See *Apples*.

PYULCUM. (From *pus*, pus, and *exco*, to draw.) An instrument to extract the pus from the cavity of any sinuous ulcer.

PYURIA. See *Pyoturia*.

PYXACANTHA. (From *pyxis*, a box, and *acantha*, a thorn.) The barberry, or thorny box-tree.

PYXIS. *Πύξις*. Properly a box; but, from its resemblance, the cavity of the hip-bone, or acetabulum, has been sometimes called os pyxidis.

Q

Q. P. An abbreviation of *quantum placet*, as much as you please.

Q. S. The contractions for *quantum sufficit*.

Q. V. An abbreviation of *quantum vis*, as much as you will.

QUADRATUS. See *Depressor labii inferioris*.

QUADRATUS FEMORIS. (*Quadratus*; from *quadra*, a square; so called from its supposed shape.) *Tuber-ischio-trochanterien* of Dumas. A muscle of the thigh, situated on the outside of the pelvis. It is a flat, thin, and fleshy muscle, but not of the shape its name would seem to indicate. It is situated immediately below the gemini. It arises tendinous and fleshy from the external surface and lower edge of the tuberosity of the ischium, and is inserted by short tendinous fibres into a ridge which is seen extending from the basis of the trochanter major to that of the trochanter minor. Its use is to bring the os femoris outwards.

QUADRATUS GENÆ. See *Platysma myoides*.

QUADRATUS LABII INFERIORIS. See *Depressor labii inferioris*.

QUADRATUS LUMBORUM. *Quadratus, seu Lumbaris externus* of Winslow. *Ilio-lumbi-costalis* of Dumas. A muscle situated within the cavity of the abdomen. This is a small, flat, and oblong muscle, that has gotten the name of *quadratus* from its shape, which is that of an irregular square. It is situated laterally, at the lower part of the spine. It arises tendinous and fleshy from about two inches from the posterior part of the spine of the ilium. From this broad origin it ascends obliquely inwards, and is inserted into the transverse processes of the four superior lumbar vertebrae, into the lower edge of the last rib, and, by a small tendon, that passes up under the diaphragm, into the side of the last vertebra of the back. When this muscle acts singly, it draws the loins to one side; when both muscles act they serve to support the spine, and perhaps to bend it forwards. In laborious respiration, the quadratus lumborum may assist in pulling down the ribs.

QUADRATUS MAXILLÆ INFERIORIS. See *Platysma myoides*.

QUADRATUS RADII. See *Pronator radii quadratus*.

QUADRIGA. (From *quatuor*, four, and *jugum*, a yoke.) A bandage which resembles the trappings of a four horse cart.

QUARTANA. *Febris quartana*. A fourth-day ague. Of this species of ague,

as well as the other kinds, there are several varieties noticed by authors. The most frequent of these are, 1. The double quartan, with two paroxysms, or fits, on the first day, none on the second and third, and two again on the fourth day. 2. The double quartan, with a paroxysm on the first day, another on the second, but none on the third. 3. The triple quartan, with three paroxysms every fourth day. 4. The triple quartan, with a slight paroxysm every day, every fourth paroxysm being similar. See also *Febris intermittens*.

QUARTZ. This name is given to the opaque, or irregularly figured vitrifiable stone.

QUASSIA. (From a slave of the name of *Quassi*, who first used it with uncommon success as a secret remedy in the malignant endemic fevers which frequently prevailed at Surinam.) 1. The name of a genus of plants in the Linnæan system. Class *Decandria*. Order, *Monogymia*.

2. The pharmacopœial name of the bitter quassia. The root, bark, and wood of this tree, *Quassia amara* of Linnæus — *floribus hermaphroditis, foliis impari-pinnatis, foliolis oppositis sessilibus, petiolo articulato alato, floribus racemosis*, are all comprehended in the catalogues of the *Materia Medica*. The tree is a native of South America, particularly of Surinam, and also of some of the West-India islands.

The roots are perfectly ligneous; they may be medically considered in the same light as the wood, which is now most generally employed, and seems to differ from the bark in being less intensely bitter; the latter is therefore thought to be a more powerful medicine. Quassia has no sensible odour; its taste is that of a pure bitter, more intense and durable than that of almost any other known substance; it imparts its virtues more completely to watery than to spirituous menstria, and its infusions are not blackened by the addition of martial vitriol. The watery extract is from a sixth to a ninth of the weight of the wood, the spirituous about a twenty-fourth. Quassia, as before observed, derived its name from a negro named Quassi, who employed it with uncommon success as a secret remedy in the malignant endemic fevers, which frequently prevailed at Surinam. In consequence of a valuable consideration, this secret was disclosed to Daniel Rolander, a Swede, who brought specimens of the quassia wood to Stockholm, in the year 1756; and, since then, the effects of this drug have been generally tried in Europe, and numerous tes-

timonies of its efficacy published by many respectable authors. Various experiments with quassia have likewise been made, with a view to ascertain its antiseptic powers; from which it appears to have considerable influence in retarding the tendency to putrefaction; and this, Professor Murray thinks, cannot be attributed to its sensible qualities, as it possesses no adstringency whatever; nor can it depend upon its bitterness, as gentian is much bitterer, yet less antiseptic. The medicinal virtues ascribed to quassia are those of a tonic, stomachic, antiseptic, and febrifuge. It has been found very effectual in restoring digestion, expelling flatulences, and removing habitual costiveness, produced from debility of the intestines, and common to a sedentary life. Dr. Lettsom, whose extensive practice gave him an opportunity of trying the effects of quassia in a great number of cases, says, "In debility, succeeding febrile diseases, the Peruvian bark is most generally more tonic and salutary than any other vegetable hitherto known; but in hysterical atony, to which the female sex is so prone, the quassia affords more vigour and relief to the system than the other, especially when united with the vitriolum album, and still more with the acid of some absorbent." In dyspepsia, arising from hard drinking, and also in diarrhoeas, the doctor exhibited the quassia with great success. But, with respect to the tonic and febrifuge qualities of quassia, he says, "I by no means subscribe to the Linnæan opinion, where the author declares, 'me quidem judice chinchinam longe superat.'" It is very well known, that there are certain peculiarities of the air, and idiosyncrasies of constitution, unfavourable to the exhibition of Peruvian bark, even in the most clear intermissions of fever; and writers have repeatedly noticed it. But this is comparatively very rare. About Midsummer, 1785, Dr. L. met with several instances of low remittent and nervous fevers, wherein the bark uniformly aggravated the symptoms, though given in intermissions the most favourable to its success, and wherein quassia, or snake-root, was successfully substituted. In such cases, he mostly observed, that there was great congestion in the hepatic system, and the debility at the same time discouraged copious evacuations. And in many fevers, without evident remissions to warrant the use of the bark, whilst, at the time, increasing debility began to threaten the life of the patient, the Doctor found that quassia, or snake-root, singly or combined, upheld the vital powers, and promoted a critical intermission of fever, by which an opportunity was offered for the bark to effect a cure. It may be given in infusion, or in pills made from the watery extract; the former is gene-

rally preferred, in the proportion of three or four drachms of the wood to twelve ounces of water.

QUASSIA AMARA. The systematic name of the bitter quassia-tree. See *Quassia*.

QUASSIA SIMAROUNA. The systematic name of the Simarouba quassia. See *Simarouba*.

QUASSY. See *Quassia*.

QUATRIO. (From *quatuor*, four; so called because it has four sides.) The astragalus.

Queen of the meadow. See *Ulmaria*.

QUEROUILA. (*Quercula*, dim. of *quercus*, the oak; so called because it has leaves like the oak.) An antiquated name of the germander. See *Chamedrys*.

QUERCUS. (From *quero*, to enquire; because divinations were formerly given from oaks by the Druids.) 1. The name of a genus of plants in the Linnæan system. Class, *Monocotyledon*. Order, *Polyandria*. The oak.

2. The pharmacopœial name of the oak. *Quercus robur* of Linneus — *foliis oblongis glabris sinuatis, lobis rotundatis, glandibus oblongis*. The oak. *Balanus*. This valuable tree is indigenous to Britain. Its adstringent effects were sufficiently known to the ancients, but it is the bark which is now directed for medicinal use by our pharmacopœias. Oak-bark manifests to the taste a strong adstringency, accompanied with a moderate bitterness. Like other adstringents, it has been recommended in agues, and for restraining hæmorrhages, alvine fluxes, and other immoderate evacuations. A decoction of it has likewise been advantageously employed as a gargle, and as a fomentation or lotion in *proidentia recti et uteri*. Galls, which, in the warm climate of the East, are found upon the leaves of this tree, are occasioned by a small insect with four wings, called *Cynips quercus folii*, which deposits an egg in the substance of the leaf, by making a small perforation through the under surface. The gall presently begins to grow to a considerable size. Two sorts of galls are distinguished in the shops; one said to be brought from Aleppo, the other from Turkey and the southern parts of Europe. The former are generally of a blueish colour, or of a greyish, or black, verging to blueness; unequal and warty on the surface; hard to break; and of a close compact texture; the other of a light brownish or whitish colour, smooth, round, easily broken, less compact, and of a much larger size. The two sorts differ only in size and strength, two of the blue galls being supposed equivalent in this respect to three of the others. Galls appear to be the most powerful of the vegetable adstringents. As a medicine, they are to be considered as applicable to the same indications as the oak-bark, and by possess-

ing a greater degree of adstringent and styptic power, seem to have an advantage over it, and to be better suited for external use. Reduced to fine powder, and made into an ointment, they have been found of great service in hæmorrhoidal affections.

QUERCUS CERRIS. The systematic name of the tree which affords the Turkey galls. See *Quercus*.

QUERCUS ESCULUS. The systematic name of the Italian oak, whose acorns are, in times of scarcity, said to afford a meal of which bread is made.

QUERCUS MARINA. The sea oak. Sea wrack. This sea-weed is the *Fucus vesiculosus* of Linnæus:—*fronde plana dichotoma costata integerrima, vesiculis axillaribus geminis, terminalibus tuberculatis*. It is said to be a useful assistant to sea-water, in the cure of disorders of the glands. Burnt in the open air, and reduced to a black powder, it forms the *æthiops vegetabilis*, which, as an internal medicine, is similar to burnt sponge.

QUERCUS PHellos. The systematic name of the willow-leaved oak, whose acorns are

much sweeter than chesnuts, and much eaten by the Indians. They afford, by expression, an oil little inferior to oil of almonds.

QUERCUS ROBUR. The systematic name of the oak-tree. See *Quercus*.

QUERCUS SUBER. The systematic name of the cork tree. See *Suber*.

Quick grass. See *Gramen caninum*.

Quick-lime. See *Lime*.

Quicksilver. See *Hydrargyrum*.

QUID PRO QVO. These words are applied the same as *succedaneum*, when one thing is made use of to supply the defect of another.

QUINA QUINA. The Peruvian bark.

Quince. See *Cydonium malum*.

Quince, Bengal. See *Bengal quince*.

Quincy. See *Cynanche*.

QUINQUEFOLIUM. (From *quinque*, five, and *folium*, a leaf; so called because it has five leaves on each foot-stalk.) *Pentaphyllum*. Cinquefoil or five-leaved grass. See *Pentaphyllum*.

QUINQUINA. See *Cinchona*.

Quinsy. See *Cynanche*.

QUOTIDIAN. See *Febris intermittens*.

R

R. or R̄. This letter is placed at the beginning of a prescription as a contraction of *recipe*, do thou take: thus, *R̄. Magnes. ʒj*, signifies, Take a drachm of magnesia.

RABIES CANINA. (*Rabies*; from *rabio*, to be mad, and *canis*, a dog.) See *Hydrophobia*.

RACHIALGIA. (From *gaxis*, the spine, and *algus*, pain.) A pain in the spine. It was formerly applied to several species of cholic which induced pain in the back.

RACHITIS. (From *gaxis*, the spine of the back; so called because it was supposed to originate in a fault of the spinal marrow.) *Crytonosus*. The English disease. The rickets. A species of disease in the class *cachexia*, and order *intumescentie* of Cullen; known by a large head, prominent forehead, protruded sternum, flattened ribs, big belly, and emaciated limbs, with great debility. It is usually confined in its attack between the two periods of nine months and two years of age, seldom appearing sooner than the former, or shewing itself for the first time, after the latter period. The muscles become flaccid, the head enlarges, the carotids are distended, the limbs waste away, and their epiphyses increase in bulk. The bones and spine of the back are variously distorted; disinclination to muscular exertion follows; the abdomen swells and grows hard; the

stools are frequent and loose; a slow fever succeeds, with cough and difficulty of respiration: atrophy is confirmed, and death ensues. Frequently it happens that nature restores the general health, and leaves the limbs distorted.

After death, the liver and the spleen have been found enlarged and scirrhus; the mesenteric glands indurated, and the lungs either charged with vomica, or adhering to the pleura; the bones soft, the brain flaccid, or oppressed with lymph, and the distended bowels loaded most frequently with slime, sometimes with worms.

It is remarkable, that in the kindred disease, which Hoffmann and Sauvage call the atrophy of infants, we have many of the same symptoms and the same appearances nearly after death. They who perish by this disease, says Hoffmann, have the mesenteric glands enlarged and scirrhus; the liver and spleen obstructed, and increased in size; the intestines are much inflated, and are loaded with black and foetid matters, and the muscles, more especially of the abdomen, waste away.

RACKASIRA BALSAMUM. See *Balsamum rackasira*.

RACOSIS. (From *gaxos*, a rag.) A ragged excoriation of the relaxed scrotum.

RADIAL ARTERY. *Arteria radialis*.

A branch of the humeral artery, that runs down the side of the radius.

RADIALIS EXTERNUS BREVIOR. See *Extensor carpi radialis brevis*.

RADIALIS EXTERNUS LONGIOR. See *Extensor carpi radialis longior*.

RADIALIS EXTERNUS PRIMUS. See *Extensor carpi radialis longior*.

RADIALIS INTERNUS. See *Flexor carpi radialis*.

RADIALIS SECUNDUS. See *Extensor carpi radialis brevis*.

RADICAL. That which is considered as constituting the distinguishing part of an acid, by its union with the acidifying principle, or oxygen, which is common to all acids. Thus sulphur is the radical of the sulphuric and sulphurous acids. It is sometimes called the base of the acid; but base is a term of more extensive application.

RADICAL VINEGAR. See *Acetum*.

RADICULA. (Dim. of *radix*, a root.) A little root; the fibrous part of a root. The common radish is sometimes so called. See *Raphanus hortensis*.

Radish, horse. See *Raphanus rusticanus*.

Radish, garden. See *Raphanus hortensis*.

RADIUS. (A spoke, a staff, or beam; so called from its resemblance.) This bone has gotten its name from its supposed resemblance to the spoke of a wheel, or to a weaver's beam; and sometimes, from its supporting the hand, it has been called *manubrium manus*. Like the ulna, it is of a triangular figure, but it differs from that bone, in growing larger as it descends, so that its smaller part answers to the larger part of the ulna, and vice versa. Of its two extremities, the uppermost and smallest is formed into a small rounded head, furnished with cartilage, and hollowed at its summit, for an articulation with the little head at the side of the pulley of the os humeri. The round border of this head, next the ulna, is formed for an articulation with the lesser symphyoid cavity of that bone. This little head of the radius is supported by a neck, at the bottom of which, laterally, is a considerable tuberosity, into the posterior half of which is inserted the posterior tendon of the biceps, while the anterior half is covered with cartilage, and surrounded with a capsular ligament, so as to allow this tendon to slide upon it as upon a pulley. Immediately below this tuberosity, the body of the bone may be said to begin. We find it slightly curved throughout its whole length, by which means a greater space is formed for the lodgment of muscles, and it is enabled to cross the ulna without compressing them. Of the three surfaces to be distinguished on the body of the bone, the external and internal ones are the broadest and flattest. The anterior surface is nar-

rower and more convex. Of its angles, the external and internal ones are rounded; but the posterior angle, which is turned towards the ulna, is formed into a sharp spine, which serves for the attachment of the interosseous ligament, of which mention is made in the description of the ulna. This strong ligament, which is a little interrupted above and below, serves not only to connect the bones of the fore-arm to each other, but likewise to afford a greater surface for the lodgment of muscles. On the fore part of the bone, and at about one-third of its length, from its upper end, we observe a channel for vessels, slanting obliquely upwards. Towards its lower extremity, the radius becomes broader, of an irregular shape, and somewhat flattened, affording three surfaces, of which the posterior one is the smallest; the second, which is a continuation of the internal surface of the body of the bone, is broader and flatter than the first; and the third, which is the broadest of the three, answers to the anterior and external surface of the body of the bone. On this last, we observe several sinuosities, covered with a thin layer of cartilage, upon which slide the tendons of several muscles of the wrist and fingers. The lowest part of the bone is formed into an oblong articulating cavity, divided into two by a slight transverse rising. This cavity is formed for an articulation with the bones of the wrist. Towards the anterior and convex surface of the bone, this cavity is defended by a remarkable eminence, called the *styloid* process of the radius, which is covered with a cartilage that is extended to the lower extremity of the ulna; a ligament is likewise stretched from it to the wrist. Besides this large cavity, the radius has another much smaller one, opposite its styloid process, which is lined with cartilage, and receives the rounded surface of the ulna. The articulation of the radius with the lesser symphyoid cavity of the ulna, is strengthened by a circular ligament, which is attached to the two extremities of that cavity, and from thence surrounds the head of the radius. This ligament is narrowest, but thickest at its middle part. But, besides this ligament, which connects the two bones of the fore-arm with each other, the ligaments which secure the articulation of the radius with the os humeri, are common both to it and to the ulna, and therefore cannot well be understood till both these bones are described. These ligaments are a capsular and two lateral ligaments. The capsular ligament is attached to the anterior and posterior surfaces of the lower extremity of the os humeri, to the upper edges and sides of the cavities we remarked at the bottom of the pulley and little head, and

likewise to some part of the condyles: from thence it is spread over the ulna, to the edges of the greater sygmoid cavity so as to include in it the end of the olecranon and of the coronoid process; and is likewise fixed round the neck of the radius, so as to include the head of that bone within it. The lateral ligaments may be distinguished into external and internal, or, according to Winslow, into *brachio-radialis*, and *brachio-cubitalis*. They both descend laterally from the lowest part of each condyle of the os humeri, and, from their fibres spreading wide as they descend, have been compared to a goose's foot. The internal ligament, or *brachio-cubitalis*, which is the longest and thickest of the two, is attached to the coronoid process of the ulna. The external ligament, or *brachio radialis*, terminates in the circular ligament of the radius. Both these ligaments adhere firmly to the capsular ligament, and to the tendons of some of the adjacent muscles. In considering the articulation of the fore-arm with the os humeri, we find that when both the bones are moved together upon the os humeri, the motion of the ulna upon the pulley allows only of flexion and extension; whereas, when the palm of the hand is turned downwards, or upwards, or in other words, in pronation and supination, we see the radius moving upon its axis, and in these motions its head turns upon the little head of the os humeri at the side of the pulley, while its circular edge rolls in the lesser sygmoid cavity of the ulna. At the lower end of the fore-arm the edge of the ulna is received into a superficial cavity at the side of the radius. This articulation, which is surrounded by a loose capsular ligament, concurs with the articulation above, in enabling the radius to turn with great facility upon its axis; and it is chiefly with the assistance of this bone that we are enabled to turn the palm of the hand upwards or downwards, the ulna having but a very inconsiderable share in these motions.

RADIX. A root.

RADIX ACORI. Galanga, or galangal.

RADIX BENGALE. See *Cassumunar*.

RADIX BRASILIENSIS. See *Ipecacuanha*.

RADIX CALAGUALE. See *Calaguale radix*.

RADIX CALAGUELLE. See *Calaguale radix*.

RADIX CASSUMUNJAR. See *Cassumunar*.

RADIX CHYNLEN. See *Chynlen radix*.

RADIX COLOMBO. See *Colomba*.

RADIX DULCIS. See *Glycyrrhiza*.

RADIX IKAN. See *Ikan radix*.

RADIX INDIANA. See *Ipecacuanha*.

RADIX INDICA LOPEZIANA. See *Lopez radix*.

RADIX MATAHSTA. See *Matahsta radix*.

RADIX ROSEA. See *Rhodiola*.

RADIX RUBRA. See *Rubia*.

RADIX TIMAC. See *Timac*.

RADIX URSINA. See *Meum*.

RADULA. (From *rado*, to scrape off.) A wooden spatula, or scraper.

RAGWORT. A poultice made of the fresh leaves is said to have a surprising effect in removing pains of the joints, and to remove the sciatica, or hip gout, in two or three applications, when ever so violent. The root is of an healing, adstringent nature. A decoction of it is good for wounds and bruises. See *Jacobaea*.

Raisin. See *Uva passa major*.

RAMALIS VENA. (From *ramale*, a dead bough.) Applied to the vena portæ, from its numerous ramifications, which resemble a bough stripped of its leaves.

RAMEX. (From *ramus*, a branch; from it protruding forwards, like a bud.) A rupture.

RANA ESCULENTA. The French frog. The flesh of this species of frog, very common in France, is highly nutritious and easily digested.

RANCID. Oily substances are said to have become rancid when, by keeping, they acquire a strong offensive smell, and altered taste.

RANINE ARTERY. *Arteria ranina*. Sublingual artery. The second branch of the external carotid.

RANULA. (From *rana*, a frog; so called from its resemblance to a frog, or because it makes the patient croak like a frog.) *Batrachos*. *Hypoglossus*. *Hypoglossum*. *Rana*. An inflammatory, or indolent tumour, under the tongue. These tumours are of various sizes and degrees of consistence, seated on either side of the frænum. Children, as well as adults, are sometimes affected with tumours of this kind; in the former, they impede the action of sucking; in the latter, of mastication, and even speech. The contents of them are various; in some, they resemble the saliva, in others, the glairy matter found in the cells of swelled joints. Sometimes, it is said that a fatty matter has been found in them; but from the nature and structure of the parts, we are sure that this can seldom happen; and, in by far the greatest number of cases, we find that the contents resemble the saliva itself. This, indeed, might naturally be expected, for the cause of these tumours is universally to be looked for in an obstruction of the salivary ducts. Obstructions here may arise from a cold, inflammation, violent fits of the toothache, attended with swelling in the inside of the mouth; and, in not a few cases, we find the ducts obstructed by a stony matter seemingly separated from the saliva, as the calculous matter is from the urine;

but where inflammation has been the cause, we always find matter mixed with the other contents of the tumour. As these tumours are not usually attended with much pain, they are sometimes neglected, till they burst of themselves, which they commonly do when arrived at the bulk of a large nut. As they were produced originally from an obstruction in the salivary duct, and this obstruction cannot be removed by the bursting of the tumour, it thence happens that they leave an ulcer extremely difficult to heal, nay, which cannot be healed at all till the cause is removed.

RANUNCULOIDES. (From *ranunculus*, and *uider*, resemblance; so named from its resemblance to the *ranunculus*.) The *Caltha palustris* or marsh marygold.

RANUNCULUS. (Dim. of *rana*, a frog; because it is found in fenny places, where frogs abound.) The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Polygynia*.

The great acrimony of most of the species of *ranunculus* is such, that, on being applied to the skin, they excite itching, redness, and inflammation, and even produce blisters, tumefaction, and ulceration of the part. On being chewed, they corrode the tongue; and, if taken into the stomach, bring on all the deleterious effects of an acrid poison. The corrosive acrimony which this family of plants possesses, was not unknown to the ancients, as appears from the writings of Dioscorides; but its nature and extent had never been investigated by experiments, before those instituted by C. Krapf, at Vienna, by which we learn, that the most virulent of the Linnæan species of *ranunculus*, are the *bulbosus*, *sceleratus*, *acris*, *arvensis*, *thora*, and *illyricus*.

The effects of these were tried, either upon himself or upon dogs, and shew that the acrimony of the different species is often confined to certain parts of the plant, manifesting itself either in the roots, stalks, leaves, flowers, or buds; the expressed juice, extract, decoction, and infusion of the plants, were also subjected to experiments. In addition to these species mentioned by Krapf, we may also notice the *R. Fammula*, and especially the *R. Alpestris*, which according to Haller, is the most acrid of this genus. Mr. Curtis observes, that even pulling up the *ranunculus acris*, the common meadow species, which possesses the active principle of this tribe, in a very considerable degree, throughout the whole herb, and carrying it to some little distance, excited a considerable inflammation in the palm of the hand in which it was held. It is necessary to remark, that the acrimonious quality of these plants is not of a fixed nature; for it may be completely dissipated by heat;

and the plant, on being thoroughly dried, becomes perfectly bland. Krapf attempted to counteract this venomous acrimony of the *ranunculus* by means of various other vegetables, none of which was found to answer the purpose, though he thought that the juice of sorrel, and that of unripe currants, had some effect in this way; yet these were much less availing than water; while vinegar, honey, sugar, wine, spirit, mineral acids, oil of tartar, p. d. and other sapid substances, manifestly rendered the acrimony more corrosive. It may be also noticed, that the virulency of most of the plants of this genus, depends much upon the situation in which they grow, and is greatly diminished in the cultivated plant.

RANUNCULUS ABORTIVUS. The systematic name of a species of *ranunculus*, which possesses acrid and vesicating properties.

RANUNCULUS ACRIS. The systematic name of the meadow crow-foot. See *Ranunculus pratensis*.

RANUNCULUS ALBUS. The plant which bears this name in the pharmacopœias is the *Anemone nemorosa* of Linnæus. The bruised leaves and flowers are said to cure tinea capitis applied to the part. The inhabitants of Kamtschatka, it is believed, poison their arrows with the root of this plant.

RANUNCULUS BULBOSUS. Bulbous root-ed crow-foot. The roots and leaves of this plant, *Ranunculus bulbosus* of Linnæus:—*culycibus retroflexis, pedunculis sulcatis, caule erecto multifloro, foliis compositis*, have no considerable smell, but a highly acrid and fiery taste. Taken internally, they appear to be deleterious, even when so far freed from the caustic matter by boiling in water, as to discover no ill quality to the palate. The effluvia, likewise, even when freely inspired, is said to occasion head-aches, anxieties, vomitings, &c. The leaves and roots, applied externally, inflame and ulcerate, or vesicate the parts, and are liable to affect also the adjacent parts to a considerable extent.

RANUNCULUS FIGARIA. The systematic name of the pilewort. See *Chelidonium minus*.

RANUNCULUS FLAMMULA. The systematic name of the smaller water crow-foot, or spearwort. Its virtues and qualities are similar to those of the *Ranunculus bulbosus*.

RANUNCULUS PALUSTRIS. Water crow-foot. The leaves of this species of crow-foot, *Ranunculus sceleratus* of Linnæus, are so extremely acrid, that the beggars in Switzerland are said, by rubbing their legs with them, to produce a very fetid and acrimonious ulceration.

RANUNCULUS PRATENSIS. Meadow crow-foot. *Ranunculus acris* of Linnæus. This,

and some other species of *ranunculus*, have, for medical purposes, been chiefly employed externally as a vesicatory, and are said to have the advantage of a common blistering plaster, in producing a quicker effect, and never causing strangury; but, on the other hand, it has been observed, that the *ranunculus* is less certain in its operation, and that it sometimes occasions ulcers, which prove very troublesome and difficult to heal. Therefore their use seems to be applicable only to certain fixed pains, and such complaints as require a long continued topical stimulus or discharge from the part, in the way of an issue, which, in various cases, has been found to be a powerful remedy.

RANUNCULUS SCLEBRATUS. The systematic name of the marsh crow-foot. See *Ranunculus palustris*.

RAPA. *Rapum. Rapus. Napus. Napa dulcis.* The turnip. *Brassica rapa* of Linnæus. Turnips are accounted a salubrious food, demulcent, detergent, somewhat laxative and diuretic, but liable, in weak stomachs, to produce flatulencies, and prove difficult of digestion. The liquor pressed out of them, after boiling, is sometimes taken medicinally in coughs and disorders of the breast. The seeds are occasionally taken as diuretics; they have no smell, but a mild acrid taste.

Rape. See *Rapus*.

RAPHANIA. (From *raphanus*, the radish, or sharlock; because the disease is said to be produced by eating the seeds of that plant.) *Convulsio raphania, vel ab ustilagine Eclampsia typhodes. Convulsio soloniensis. Necrosis ustilaginea.* Cripple disease. A genus of disease in the class *neuroses*, and order *spasmi*, of Cullen; characterised by a spasmodic contraction of the joints, with convulsive motions, and a most violent pain returning at various periods. It begins with cold chills and lassitude, pain in the head, anxiety about the præcordia. These symptoms are followed by spasmodic twichings in the tendons of the fingers and of the feet, discernible to the eye, heat, fever, stupor, delirium, sense of suffocation, aphonia, and horrid convulsions of the limbs. After these, vomiting and diarrhœa come on, with a discharge of worms. About the eleventh or the twentieth day, copious sweats succeed, or purple exanthemata, or tæta, or rigidity of all the joints.

RAPHANUS. (*Ραφανος, ὡραία το γάλας φαινομένη*: from its quick growth.) The horse-radish. A genus of plants in the Linnæan system. Class, *Tetradynamia*. Order, *Siliculosa*.

RAPHANUS HORTENSIS. *Radicula. Raphanus niger.* The radish. The several varieties of this plant, *Raphanus sativus* of Linnæus, are said to be employed medi-

cinally in the cure of calculeous affections. The juice, made into a syrup, is given to relieve hoarseness. Mixed with honey, or sugar, it is administered in pittingous asthma; and, as antiscorbutics, their efficacy is generally acknowledged.

RAPHANUS NIGER. See *Raphanus hortensis*.

RAPHANUS RUSTICANUS. *Armoracia. Raphanus marinus. Raphanus sylvestris.* Horse-radish. The plant which affords this root is the *Cochlearia armoracia*; *foliis radicalibus lanceolatis crenatis, caulibus incis* of Linnæus. Horse-radish has long been received into the materia medica, is also well known at our tables. "It affects the organs both of taste and smell with a quick penetrating pungency; nevertheless it contains in certain vessels a sweet juice, which sometimes exudes in little drops upon the surface. Its pungent matter is of a very volatile kind, being totally dissipated in drying, and carried off in evaporation, or distillation by water, and rectified as the pungency exhales, the sweet matter of the root becomes more sensible, though this also is, in a great measure, dissipated or destroyed. It impregnates both water and spirit, by infusion, or by distillation, very richly with its active matters. In distillation with water, it yields a small quantity of essential oil, exceedingly penetrating and pungent."

Dr. Cullen has mentioned every thing necessary to be known respecting the medicinal virtues of horse-radish, we shall therefore transcribe all that the ingenious professor has written on this subject. "The root of this only is employed; and it affords one of the most acrid substances of this order (*Siliquose*), and therefore proves a powerful stimulant, whether externally or internally employed. Externally, it readily inflames the skin, and proves a rubefacient that may be employed with advantage in palsy and rheumatism; and, if its application be long continued, it produces blisters. Taken internally, it may be so managed as to relieve hoarseness, by acting on the fauces. Received into the stomach, it stimulates this, and promotes digestion; and therefore is properly employed as a condiment with our animal food. If it be infused in water, and a portion of this infusion be taken with a large draught of warm water, it readily proves emetic, and may either be employed by itself to excite vomiting, or to assist the operation of other emetics. Infused in water, and taken into the stomach, it proves stimulant to the nervous system, and is thereby useful in palsy; and, if employed in large quantity, it proves heating to the whole body; and thereby it proves often useful in chronic rheumatism,

whether arising from scurvy or other causes. Bergius has given us a particular method of exhibiting this root, which is, by cutting it down, without bruising, into small pieces; and these, if swallowed without chewing, may be taken down in large quantities, to that of a table-spoonful. And the author alleges, that, in this way, taken in the morning for a month together, this root has been extremely useful in arthritic cases; which, however, I suppose to have been of the rheumatic kind. It would seem, in this manner employed, analogous to the use of unbruised mustard-seed; it gives out in the stomach its subtle volatile parts, that stimulate considerably without inflaming. The matter of horse-radish, like the same matter of the other siliqueous plants carried into the blood-vessels, passes readily into the kidneys, and proves a powerful diuretic, and is therefore useful in dropsy; and we need not say, that, in this manner, by promoting both urine and perspiration, it has been long known as one of the most powerful antiscorbutics."

RAPHANUS SATIVUS. The systematic name of the radish plant. See *Raphanus hortensis*.

RAPHANUS SYLVESTRIS. The poor man's pepper is sometimes so called. See *Lepidium*.

RAPHE SCROTI. (*Papa*, a suture.) The rough eminence which divides the scrotum, as it were, in two. It proceeds from the root of the penis inferiorly towards the perineum.

RAPHE CEREBRI. The longitudinal eminence of the corpus callosum of the brain is so called, because it appears somewhat like a suture.

RAPISTRUM. (From *rapa*, the turnip, because its leaves resemble those of turnip.) *Lampæna. Miagra.* Charlock, or wild mustard.

RAPUM. (*Ety.* uncertain.) See *Rapa*.

RAPUNCULUS. (Dim. of *rapa*, the turnip.) The wild turnip.

RAPUNCULUS VIRGIANUS. The name given by Morrison to the blue cardinal flower. See *Lobelia*.

RAPUS. See *Rapa*.

RASH. *Exanthema.* A rash consists of red patches on the skin, variously figured; in general confluent, and diffused irregularly over the body, leaving interstices of a natural colour. Portions of the cuticle are often elevated in a rash, but the elevations are not acuminated. The eruption is usually accompanied with a general disorder of the constitution, and terminates in a few days by cuticular exfoliations.

RASPATORIUM. (From *rado*, to scrape.) A surgeon's rasp.

Raspberry. See *Rubus idæus*.

RASURA. (From *rado*, to scrape.) 1. A rasure or scratch. 2. The raspings or shavings of any substance.

RATIFIA. A liquor prepared by imparting to ardent spirits the flavour of various kinds of fruits.

Rattle snake root. See *Seneca*.

RAUCEDO. (From *raucus*, hoarse.) *Raucitas.* Hoarseness. It is always symptomatic of some other disease.

RE-AGENTS. Tests. Those substances which are used in chemistry to detect the substance for which they are used. In the application of tests there are two circumstances to be attended to, viz. To avoid deceitful appearances, and to have good tests.

The principal tests are the following:

1. *Litmus.* The purple of litmus is changed to red by every acid; so that this is the test generally made use of to detect excess of acid in any fluid. It may be used either by dipping into the water a paper stained with litmus, or by adding a drop of the tincture to the water to be examined, and comparing its hue with that of an equal quantity of the tincture in distilled water.

Litmus already reddened by an acid will have its purple restored by an alkali; and thus it may also be used as a test for alkalis, but it is much less active than other direct alkaline tests.

2. *Red cabbage* has been found by Mr. Watt to furnish as delicate a test for acids as litmus, and to be still more sensible to alkalis. The natural colour of an infusion of this plant is blue, which is changed to red by acids, and to green by alkalis in very minute quantities.

3. *Brazil wood.* When chips of this wood are infused in warm water they yield a red liquor, which readily turns blue by alkalis, either caustic or carbonated. It is also blueed by the carbonated earths held in solution by carbonic acid, so that it is not an unequivocal test of alkalis till the earthy carbonates have been precipitated by boiling. Acids change to yellow the natural red of brazil wood, and restore the red when changed by alkalis.

4. *Violets.* The delicate blue of the common scented violet is readily changed to green by alkalis, and this affords a delicate test for these substances. Syrup of violets is generally used as it is at hand, being used in medicine. But a tincture of the flower will answer as well.

5. *Turmeric.* This is a very delicate test for alkalis, and on the whole perhaps is the best. The natural colour either in watery or spirituous infusion is yellow, which is changed to a brick or orange red by alkalis, caustic or carbonated, but not by carbonated earths, on which account it is preferable to Brazil wood.

The pure earths, such as lime and barytes, produce the same change.

6. *Rhubarb*. Infusion or tincture of rhubarb undergoes a similar change with turmeric and is equally delicate.

7. *Sulphuric acid*. A drop or two of concentrated sulphuric acid, added to water that contains carbonic acid, free or in combination, causes the latter to escape with a pretty brisk effervescence, whereby the presence of this gaseous acid may be detected.

8. *Nitric and oxymuriatic acid*. A peculiar use attends the employment of these acids in the sulphuretted waters, as the sulphuretted hydrogen is decomposed by them, its hydrogen absorbed, and the sulphur separated in its natural form.

9. *Oxalic acid and oxalat of ammonia*. These are the most delicate tests for lime and all soluble calcareous salts. Oxalat of lime, though nearly insoluble in water, dissolves in a moderate quantity in its own or any other acid, and hence in analysis oxalat of ammonia is often preferred, as no excess of this salt can re-dissolve the precipitated oxalat of lime. On the other hand the ammonia should not exceed, otherwise it might give a false indication.

10. *Gallie acid and tincture of galls*. These are tests of iron. Where the iron is in very minute quantities, and the water somewhat acidulous, these tests do not always produce a precipitate, but only a slight reddening, but its action is much heightened by previously adding a few drops of any alkaline solution.

11. *Prussiat of potash and lime*. The presence of iron in water is equally well indicated by these prussiates, and if the prussiat of potash is properly prepared, it will only be precipitated by a metallic salt, so that manganese and copper will also be detected, the former giving a white precipitate, the latter a red precipitate.

12. *Lime-water* is the common test for carbonic acid, it decomposes all the magnesian salts, and likewise the aluminous salts, it likewise produces a cloudiness with most of the sulphats owing to the formation of selenite.

13. *Ammonia*. This alkali when perfectly caustic serves as a distinction between the salts of lime and those of magnesia, as it precipitates the earth from the latter salts, but not from the former. There are two sources of error to be obviated, one is that of carbonic acid being present in the water, the other is the presence of aluminous salts.

14. *Carbonated alkalis*. These are used to precipitate all the earths, where carbonate of potash is used particular care should be taken of its purity, as it generally contains silex.

15. *Muriated alumine*. This test is pro-

posed by Mr. Kirwan to detect carbonate of magnesia, which cannot, like carbonated lime, be separated by ebullition, but remains till the whole liquid is evaporated.

16. *Barytic salts*. The nitrate, muriate, and acetite of barytes are all equally good tests of sulphuric acid in any combination.

17. *Salts of silver*. The salts of silver are the most delicate tests of muriatic acid, in any combination, producing the precipitated luna cornea. All the salts of silver likewise give a dark brown precipitate with the sulphuretted waters, which is as delicate a test as any that we possess.

18. *Salt of lead*. The nitrate and acetite of lead are the salts of this metal employed as tests. They will indicate the sulphuric, muriatic, and boracic acid, and sulphuretted hydrogen or sulphuretted kali.

19. *Soap*. A solution of soap in distilled water or in alcohol is curdled by water containing any earthy or metallic salt.

20. *Tartareous acid*. This acid is of use in distinguishing the salts with potash, (with which it forms a precipitate of cream of tartar,) from those of soda, from which it does not precipitate. The potash however must exist in some quantity to be detected by the test.

21. *Nitro-muriate of platina*. This salt is still more discriminative between potash and the other alkalis, than acid of tartar, and will produce a precipitate with a very weak solution of any salt with potash.

22. *Alcohol*. This most useful re-agent is applicable in a variety of ways in analysis. As it dissolves some substances found in fluids, and leaves others untouched, it is a means of separating them into two classes, which saves considerable trouble in the further investigation. Those salts which it does not dissolve, it precipitates from their watery solution, but more or less completely according to the salt contained, and the strength of the alcohol, and as a precipitant it also assists in many decompositions.

REALGAR. *Arlada. Arladar. Auripigmentum rubrum. Arsenicum rubrum factitium, Abessi.* A metallic substance of a red colour, more or less lively and transparent, and often crystallized in brilliant needles; formed by a combination of arsenic with sulphur. See *Arsenic*.

RECEPTACULUM CHYLI. (*Receptaculum, from recepto, to receive.*) *Receptaculum Pequeti*, because Pequet first attempted to demonstrate it. *Diversorium. Sacculus chyliiferus.* The existence of such a receptacle in the human body is doubted. In brute animals the receptacle of the chyle is situated on the dorsal vertebrae where the lacteals all meet. See *Absorbents*.

RECTIFICATION. (*Rectificatio, from*

rectifico, to make clean.) A second distillation, in which substances are purified by their more volatile parts being raised by heat carefully managed; thus, spirit of wine, ether, &c. are rectified by their separation from the less volatile and foreign matter which altered or debased their properties.

RECTOR SPIRITUS. The aromatic part of plants.

RECTUM. (So named from an erroneous opinion that it was straight.) *Rectum intestinum.* *Speuthymenes.* *Longanum,* or *longum.* *Archeus.* *Cysarces.* The last portion of the large intestines terminating in the anus. See *Intestines.*

RECTUS ABDOMINIS. *Pubis-sternal* of Dumas. This long and straight muscle is situated near its fellow, at the middle and fore part of the abdomen, parallel to the linea alba, and between the aponeuroses of the other abdominal muscles. It arises sometimes by a single broad tendon from the upper and inner part of the os pubis, but more commonly by two heads, one of which is fleshy, and originates from the upper edge of the pubis, and the other tendinous, from the inside of the symphysis pubis, behind the pyramidalis muscle. From these beginnings, the muscle runs upwards the whole length of the linea alba, and, becoming broader and thinner as it ascends, is inserted by a thin aponeurosis into the edge of the cartilago ensiformis, and into the cartilages of the fifth, sixth, and seventh ribs. This aponeurosis is placed under the pectoral muscle, and sometimes adheres to the fourth rib. The fibres of this muscle are commonly divided by three tendinous intersections, which were first noticed by Berenger, or, as he is commonly called, Carpi, an Italian anatomist, who flourished in the sixteenth century. One of these intersections is usually where the muscle runs over the cartilage of the seventh rib; another is at the umbilicus; and the third is between these two. Sometimes there is one, and even two, between the umbilicus and the pubis. When one, or both of these occur, however, they seldom extend more than half way across the muscle. As these intersections seldom penetrate through the whole substance of the muscle, they are all of them most apparent on its anterior surface, where they firmly adhere to the sheath; the adhesions of the rectus to the posterior layer of the internal oblique, are only by means of cellular membrane, and of a few vessels which pass from one to another.

Albinus and some others have seen this muscle extending as far as the upper part of the sternum.

The use of the rectus is to compress the fore part of the abdomen, but more parti-

cularly the lower part; and, according to the different positions of the body, it may likewise serve to bend the trunk forwards, or to raise the pelvis. Its situation between the two layers of the internal oblique, and its adhesions to this sheath, secure it in its place, and prevent it from rising into a prominent form when in action; and lastly, its tendinous intersections enable it to contract at any of the intermediate spaces.

RECTUS ABDUCENS OCULI. See *Rectus externus oculi.*

RECTUS ABDUCENS OCULI. See *Rectus internus oculi.*

RECTUS ANTERIOR BREVIS. See *Rectus capitis internus minor.*

RECTUS ANTERIOR LONGUS. See *Rectus capitis internus major.*

RECTUS ATTOLLENS OCULI. See *Rectus superior oculi.*

RECTUS CAPITIS ANTIQUS LONGUS. See *Rectus capitis internus major.*

RECTUS CAPITIS INTERNUS MAJOR. *Rectus internus major* of Albinus, Douglas, and Cowper. *Trachelo-basilare* of Dumas. *Rectus anterior longus* of Winslow. This muscle is situated on the anterior part of the neck, close to the vertebrae. It was known to most of the ancient anatomists, but was not distinguished by any particular name until Cowper gave it the present appellation, and which has been adopted by most writers except Winslow. It is a long muscle, thicker and broader above than below, where it is thin, and terminates in a point. It arises, by distinct and flat tendons, from the anterior points of the transverse processes of the five inferior vertebrae of the neck, and, ascending obliquely upwards, is inserted into the anterior part of the cuneiform process of the occipital bone. The use of this muscle is to bend the head forwards.

RECTUS CAPITIS INTERNUS MINOR. Cowper, who was the first accurate describer of this little muscle, gave it the name of *rectus internus minor*, which has been adopted by Douglas and Albinus. Winslow calls it *rectus anterior brevis*, and Dumas *petit-trachelo-basilare*. It is in part covered by the rectus major. It arises fleshy from the upper and fore part of the body of the first vertebra of the neck, near the origin of its transverse process, and, ascending obliquely inwards, is inserted near the root of the condyloid process of the occipital bone, under the last-described muscle. It assists in bending the head forwards.

RECTUS CAPITIS LATERALIS. *Rectus lateralis Fallopi* of Douglas. *Transversalis anticus primus* of Winslow. *Rectus lateralis* of Cowper, and *Tracheli alioide basilare* of Dumas. This muscle seems to have been first described by Fallopius.

Winslow calls it *transversalis anticus primus*. It is somewhat larger than the *rectus minor*, but resembles it in shape, and is situated immediately behind the internal jugular vein, at its coming out of the cranium. It arises fleshy from the upper and fore part of the transverse process of the vertebra of the neck, and, ascending a little obliquely upwards and outwards, is inserted into the occipital bone, opposite to the stylo-mastoid hole of the os temporis. This muscle serves to pull the head to one side.

RECTUS CAPITIS POSTICUS MAJOR. This, which is the *rectus major* of Douglas and Winslow, the *rectus capitis posticus minor* of Albinus, and the *spino-axoideo-occipital* of Dumas, is a small, short, and flat muscle, broader above than below, and is situated, not in a straight direction, as its name would insinuate, but obliquely, between the occiput and the second vertebra of the neck, immediately under the complexus. It arises, by a short thick tendon, from the upper and posterior part of the spinous process of the second vertebra of the neck; it soon becomes broader, and ascending obliquely outwards, is inserted, by a flat tendon, into the external lateral part of the lower semi-circular ridge of the os occipitis. The use of this is to extend the head, and pull it backwards.

RECTUS CAPITIS POSTICUS MINOR. This is the *rectus minor* of Douglas and Winslow, and the *tuber-altoideo-occipital* of Dumas. It is smaller than the last-described muscle, but resembles it in shape, and is placed close by its fellow, in the space between the *recti majores*. It arises, by a short thick tendon, from the upper and lateral part of a little protuberance in the middle of the back part of the first vertebra of the neck, and, becoming broader and thinner as it ascends, is inserted, by a broad flat tendon, into the occipital bone, immediately under the insertion of the last-described muscle. The use of it is to assist the *rectus major* in drawing the head backwards.

RECTUS CRURIS. See *Rectus femoris*.

RECTUS DEPRIMENS OCULI. See *Rectus inferior oculi*.

RECTUS EXTERNUS OCULI. *Abductor oculi. Indignabundus.* The outer straight muscle of the eye. It arises from the bony partition between the foramen opticum and lacerum, being the longest of the straight muscles of the eye, and is inserted into the sclerotic membrane, opposite to the outer canthus of the eye. Its use is to move the eye outwards.

RECTUS FEMORIS. *Rectus sive Gracilis anterior* of Winslow. *Rectus cruris* of Albinus, and *ilio-trochantericus* of Dumas. A straight muscle of the thigh, situated immediately at the fore part. It

arises from the os ilium by two tendons. The foremost and shortest of these springs form the outer surface of the inferior and anterior spinous process of the ilium; the posterior tendon, which is thicker and longer than the other, arises from the posterior and outer part of the edge of the cotyloid cavity, and from the adjacent capsular ligament. These two tendons soon unite, and form an aponeurosis, which spreads over the anterior surface of the upper part of the muscle; and through its whole length we observe a middle tendon, towards which its fleshy fibres run on each side in an oblique direction, so that it may be styled a penniform muscle. It is inserted tendinous into the upper edge and anterior surface of the patella, and from thence sends off a thin aponeurosis, which adheres to the superior and lateral part of the tibia. Its use is to extend the leg.

RECTUS INFERIOR OCULI. *Depressor oculi. Deprimens. Humilis.* The inferior of the straight muscles of the eye. It arises within the socket, from below the optic foramen, and passes forwards to be inserted into the sclerotic membrane of the bulb on the wider part. It pulls the eye downwards.

RECTUS INTERNUS FEMORIS. See *Gracilis*.

RECTUS INTERNUS OCULI. *Adducens oculi. Bibitorius.* The internal straight muscle of the eye. It arises from the inferior part of the foramen opticum, between the obliquus superior, and the rectus inferior, being, from its situation, the shortest muscle of the eye, and is inserted into the sclerotic membrane opposite to the inner angle. Its use is to turn the eye towards the nose.

RECTUS LATERALIS PALLOPI. See *Rectus capitis lateralis*.

RECTUS MAJOR CAPITIS. See *Rectus capitis posticus major*.

RECTUS SUPERIOR OCULI. *Attollens oculi. Levator oculi. Superbus.* The uppermost straight muscle of the eye. It arises from the upper part of the foramen opticum of the sphenoid bone below the levator palpebræ superioris, and runs forward to be inserted into the superior and fore part of the sclerotic membrane by a broad and thin tendon.

RECURRENT NERVES. Two branches of the par vagum in the cavity of the thorax. The right is given off near the subclavian artery, which it surrounds, and is reflected upwards to the thyroid gland; the left a little lower, and reflected around the aorta to the œsophagus, as far as the larynx. They are both distributed to the muscles of the larynx and pharynx.

REDDLE. A species of ochre or argillaceous earth, of a dark red colour, some-

times used medicinally as a tonic and antacid.

Red Saunders. See *Santulum rubrum*.

REFRIGERANTIA. (From *refrigere*, to cool.) Refrigerants. Medicines, which allay the heat of the body or the blood.

REFRIGERATORIUM. (From *refrigere*, to cool.) A vessel filled with water to condense vapours, or to make cool any substance which passes through it.

REGIMEN. (From *rego*, to govern.) A term employed in medicine to express the plan or regulation of the diet.

REGINA PRATI. See *Ulmaria*.

REGIS MORBUS. (From *rex*, a king.) The royal disease, a term applied to the jaundice from its golden colour, and to a preparation of nitre, because it dissolves only gold.

Regular gout. See *Arthritis*.

REGULUS. (Dim. of *rex*, a king; so called because the alchemists expected to find gold, the king of metals, collected at the bottom of the crucible after fusion.) A name given by the alchemists to metallic matters when separated from other substances by fusion; as, regulus of antimony, regulus of arsenic, &c.

Regulus of antimony. The pure metal antimony. See *Antimony*.

Regulus of arsenic. Pure arsenic. See *Arsenic*.

REMEDIUM DIVINUM. See *Imperatoria*.

Remittent fever. See *Febria continua*.

REMORA ARATRI. (From *remoror*, to hinder, and *aratrum*, a plough; so called because it hinders the plough.) See *Ononis*.

Remote cause. See *Exciting cause*.

REN. (*Ren, aro tu ren*; because through them the urine flows.) The kidney. See *Kidneys*.

RENAL ARTERIES. *Arterix renales.* See *Emulgent vessels*.

RENAL GLANDS. *Glandule renales.* Renal capsules. Supra-renal glands. The supra-renal glands are two hollow bodies, like glands in fabric, and placed one on each side upon the kidney. They are covered by a double tunic, and their cavities are filled with a liquor of a brownish red colour. Their figure is triangular; and they are larger in the fœtus than the kidneys; but in adults they are less than the kidneys. The right is affixed to the liver, the left to the spleen and pancreas, and both to the diaphragm and kidneys. They have arteries, veins, and lymphatics and nerves; their arteries arise from the diaphragmatic, the aorta, and the renal arteries. The vein of the right supra-renal gland empties itself into the vena cava; that of the left into the renal vein; their lymphatic vessels go directly to the thoracic duct; they have nerves common alike to these glands and the kidneys. They have no excretory duct, and their use is

at present unknown. It is supposed they answer one use in the fœtus, and another in the adult, but what these uses are is uncertain. Boerhaave supposed their use to consist in their furnishing lymph to dilute the blood returned after the secretion of the urine in the renal vein; but this is very improbable, since the vein of the right supra-renal gland goes to the vena cava, and the blood carried back by the renal vein wants no dilution. It has also been said, that these glands not only prepare lymph, by which the blood is fitted for the nutrition of the delicate fœtus; but that in adults they serve to restore in themselves, to the blood of the vena cava, the irritable parts which it loses by the secretion of bile and urine. Some, again, have considered them as diverticula in the fœtus, to divert the blood from the kidneys, and lessen the quantity of urine. The celebrated Morgagni believed their office to consist in conveying something to the thoracic duct. It is singular, that in children who are born without the cerebrum, these glands are extremely small, and sometimes wanting.

RENAL VEINS. See *Emulgent vessels*.

Renal vessels. See *Emulgent vessels*.

RENET. The gastric juice and contents of the stomach of calves. It is much employed in preparing cheese, and, in pharmacy, for making whey. To about a pound of milk, in a silver or earthen basin placed on hot ashes, add three or four grains of rennet, diluted with a little water; as it becomes cold the milk curdles, and the whey, or serous part, separates itself from the caseous part. When these parts appear perfectly distinct, pour the whole upon a strainer, through which the whey will pass, while the curds remain behind. This whey is always rendered somewhat whitish, by a very small and much divided portion of the caseous part; but it may be separated in such a manner, that the whey will remain limpid and colourless, and this is what is called clarifying it. Put into a basin the white of an egg, a glass of the serum of milk, and a few grains of tartareous acidulum in powder; whip the mixture with an oziertwig, and, having added the remainder of the unclarified whey, place the mixture again over the fire until it begins to boil. The tartareous acidulum completes the coagulation of the white part of the milk which remains; the white of egg, as it becomes hot, coagulates and envelops the caseous part. When the whey is clear, filter it through paper; what passes will be perfectly limpid, and have a greenish colour. This is clarified whey.

RENUERS. (From *renuo*, to nod the head back in sign of refusal; so called from its office of jerking back the head.) A muscle of the head.

REPELLENTIA. (From *repello*, to

drive back.) Repellents. Applications are sometimes so named which make diseases recede, as it were, from the surface of the body.

REPULSION. All matter possesses a power which is in constant opposition to attraction. This agency, which is equally powerful and equally obvious, acts an important part in the phenomena of nature, and is called *the power of repulsion*.

That such a force exists which opposes the approach of bodies towards each other is evident from numberless facts.

Newton has shown that when a convex lens is put upon a flat glass it remains at a distance of the one-hundred-and-thirty-seventh part of an inch, and a very considerable pressure is required to diminish this distance; nor does any force which can be applied bring them into actual mathematical contact; a force may indeed be applied sufficient to break the glasses into pieces, but it may be demonstrated that it does not diminish their distance much beyond the one-thousandth part of an inch. There is therefore a repulsive force which prevents the two glasses from touching each other.

Boscovich has shown that when an ivory billiard-ball sets another in motion by striking against it, an equal quantity of its own motion is lost, and the ball at rest begins to move while the other is still at a distance.

There exists therefore a repulsion between bodies; this repulsion takes place while they are yet at a distance from each other; and it opposes their approach towards each other.

The cause or the nature of this force is equally inscrutable with that of attraction, but its existence is undoubted; it increases as far as has been ascertained inversely as the square of the distance, consequently at the point of contact it is infinite.

The following experiments will serve to prove the energy of repulsion more fully.

Experiment.—When a glass tube is immersed in water the fluid is attracted by the glass, and drawn up into the tube; but, if we substitute mercury instead of water, we shall find a different effect. If a glass tube of any bore be immersed in this fluid it does not rise, but the surface of the mercury is considerably below the level of that which surrounds it.

In this case therefore a repulsion takes place between the glass and the mercury, which is even considerably greater than the attraction existing between the particles of the mercury, and hence the latter cannot rise in the tube, but is repelled, and becomes depressed.

Experiment.—When we present the north pole of a magnet A, to the same pole of another magnet B, suspended on a pivot, and at liberty to move, the magnet B

will recede as the other approaches; and by following it with A, at a proper distance, it may be made to turn round on its pivot with considerable velocity.

In this case there is evidently some agency which opposes the approach of the north poles of A and B, which acts as an antagonist, and causes the moveable magnet to retire before the other. There is therefore a *repulsion* between the two magnets, a repulsion which increases with the power of the magnets; which may be made so great that all the force of a strong man is insufficient to make the two north poles touch each other.—The same repulsion is equally obvious on electrical bodies, for instance.

Experiment.—If two small cork balls be suspended from a body with silk threads so as to touch one another, and if we charge the body in the usual manner with electricity, the two cork balls separate from each other, and stand at a distance proportional to the quantity of electricity with which the body is charged; the balls of course repel each other.

Experiment.—If we rub over the surface of a sheet of paper the fine dust of lycopodium or puff ball, and then let water fall on it in small quantities, the water will instantly be repelled and form itself into distinct drops which do not touch the lycopodium, but roll over it with uncommon rapidity. That the drops do not touch the lycopodium, but are actually kept at a distance above it, is obvious from the copious reflection of white light.

Experiment.—If the surface of water contained in a basin be covered over with lycopodium, a solid substance deposited at the bottom of the fluid may be taken out of it with the hand without wetting it. In this case the repulsion is so powerful as to defend the hand completely from the contact of the fluid.

RES NATURALES. (From *natura*, nature.) The naturals. According to Boerhaave, these are life, the cause of life, and its effects. These, he says, remain in some degree, however disordered a person may be.

RESEDA. (From *resedo*, to appease; so called from its virtue of allaying inflammation.) The herb wild rocket.

Resin, black. See *Resina nigra*.

Resin, elastic. See *Indian rubber*.

Resin tree, elastic. See *Indian rubber*.

Resin, white. See *Resina suva*.

Resin, yellow. See *Resina suva*.

RESINA. (*Resina*, from *res*, to flow.) Resin. The essential properties of resin are, being in the solid form, insoluble in water, perfectly soluble in alcohol, and in essential and expressed oils, and being incapable of being volatilized without decomposition.

Resins are obtained chiefly from the vegetable kingdom, either by spontaneous

exudation, or from incisions made into vegetables affording juices which contain this principle. These juices contain a portion of essential oil, which, from exposure to the air, is either volatilized or converted into resinous matter, or sometimes the oil is abstracted by distillation. In some plants the resin is deposited, in a concrete state, in the interstices of the wood, or other parts of the plant.

Resins, when concrete, are brittle, and have generally a smooth and conchoidal fracture; their lustre is peculiar, they are more or less transparent, and of a colour which is usually some shade of yellow, or brown; they are of a greater specific gravity than water; they are often odorous and sapid, easily fusible, and, on cooling, become solid.

RESINA ALBA. See *Resina flava*.

RESINA ELASTICA. See *Indian rubber*.

RESINA FLAVA. *Resina alba*. Yellow resin, what remains in the still after distilling oil of turpentine, by adding water to the common turpentine. It is of very extensive use in surgery as an active detergent, and forms the base of the *unguentum resina flavae*.

RESINA LUTEA NOVI BELGII. Botany-bay gum. All the information that has been hitherto collected respecting the history of the yellow gum is the following:

The plant that produces it is low and small, with long grassy leaves; but the fructification of it shoots out in a singular manner from the centre of the leaves, on a single straight stem, to the height of twelve or fourteen feet. Of this stem, which is strong and light, like some of the reed class, the natives usually make their spears. The resin is generally dug up out of the soil under the tree, not collected from it, and may perhaps be that which Tasman calls, 'gum lac of the ground.'

Mr. Boles, surgeon of the Lady Penrhyn, gives a somewhat different account; and as this gentleman appears to have paid considerable attention to the subject, his account may certainly be relied upon. After describing the tree in precisely the same manner as above, he observes, that at the top of the trunk of the tree, long grassy leaves grow in great abundance. The gum is found under these leaves in considerable quantities; it commonly exudes in round tears, or drops, from the size of a large pea to that of a marble, and sometimes much larger. These are by the heat of the sun frequently so much softened, that they fall on the ground, and in this soft state adhere to whatever they fall upon; hence the gum is frequently found mixed with dirt, wood, the bark of the tree, and various other substances: so that one lump has been seen composed of many small pure pieces of various sizes united

together, which weighed nearly half a hundred weight. It is produced in such abundance, that one man may collect thirty or forty pounds in the space of a few hours. The convicts have another method of collecting it: they dig round the tree, and break off pieces of the roots which always have some, and frequently considerable quantities of the gum in them. This gum appears nearly, but not entirely, the same as that which exudes from the trunk of the tree; the former is often mixed with a strong smelling resinous substance of a black nature, and is so interwoven in the wood itself, that it is with difficulty separated. The latter appears a pure unmixed resinous substance.

Several experiments have been made, principally with the view of determining what menstruum would dissolve the gum the most readily, and in the greatest quantity, from which it appears alcohol and ether dissolve the most.

The diseases in which this resin is administered, are those of the *primæ viæ*, and principally such as arise from spasm, a debility, a loss of tone, or a diminished action in the muscular fibres of the stomach and bowels, such as loss of appetite, sickness, vomiting, flatulency, heart-burn, pains in the stomach, &c. when they were really idiopathic complaints, and not dependent upon any disease in the stomach, or affections of other parts of the body communicated to the stomach. In debilities and relaxations of the bowels, and the symptoms from thence arising, such as purging and flatulency, it has been found of good effect. In certain cases of diarrhoea, however, (and it seemed those in which an unusual degree of irritability prevailed,) it did not answer so well, unless given in small doses, and combined with opiates, when the patient seemed to gain greater advantage than when opiates only were had recourse to. In case of amenorrhœa, depending on (what most of those cases do depend upon) a sluggishness, a debility, and acidity of the system, this medicine, when assisted by proper exercise and diet, has, by removing the symptoms of dyspepsia, and by restoring the tone and action of the muscular fibres, been found very serviceable. This medicine does not, in the dose of about half a drachm, appear to possess any remarkably sensible operation. It neither vomits, purges, nor binds the belly, nor does it materially increase the secretion of urine or perspiration. It has indeed sometimes been said to purge, and at others to occasion sweating, but they are not constant effects, and when they do occur, it generally depends on some accidental circumstance. It should seem to possess, in a very extensive degree, the property of allaying morbid irritability, and of restoring tone, strength,

and action to the debilitated and relaxed fibre. When the gum itself is given, it should always be the pure unmixed part; if given in the form of a draught, it should be mixed in water with mucilage of gum arabic; if made into pills, a small portion of Castile soap may be employed; it was found the *lixiv. sapon.* dissolved it entirely. It is commonly, however, made into a tincture by mixing equal parts of the gum and rectified spirit; one drachm of this tincture, (containing half a drachm of the pure gum,) made into a draught with water and syrup, by the assistance of 15 grains of gum arabic in mucilage, forms an elegant medicine, and at the same time very palatable.

RESINA NIGRA. Colophonia. What remains in the retort after distilling the balsam of turpentine from the common turpentine.

RESOLVENTS. (*Medicamenta Resolventia*, from *resolvo*, to loosen.) This term is applied by surgeons to such substances as discuss inflammatory tumours.

RESOLUTION. (*Resolutio*, from *resolvo*, to loosen.) A termination of inflammatory affections in which the diseases disappear without any abscess, mortification, &c. being occasioned. The term is also applied to the dispersion of swellings, indurations, &c.

RESPIRATION. (*Respiratio*, from *respiro*, to take breath.) Of all the changes the blood suffers in passing through our various organs, there are none more essential or remarkable than those occasioned by the influence of the air, which is alternately received into, and expelled from the lungs during the act of respiration. The blood which the veins return to the heart, and which the right ventricle sends into the pulmonary artery, is blackish and heavy; its temperature only 30 degrees of Reaumur's thermometer: if suffered to remain still, it coagulates slowly, and separates a great portion of serum. That which the pulmonary veins bring back to the left cavities of the heart, and which is conveyed into every part of the body by means of the arteries, is, on the contrary, of a red vermillion colour, frothy, lighter, and two degrees warmer; it is also more easily coagulable, and separates a smaller proportion of serum. All these differences, which are so easily perceptible, are dependent upon the modifications arising from having been in contact with the atmospheric air.

In man, and in all animals of warm blood that have an heart composed of two auricles and two ventricles, the blood which has been carried into all the organs by the arteries, and brought back by the veins to the heart, cannot be returned into the arteries without having first passed through the lungs, forming a medium which the blood must necessarily traverse

to pass from the right into the left cavities of the heart; this passage constitutes the pulmonary or smaller circulation.

Mayow has given the most accurate idea of the respiratory organ, in comparing it to a pair of bellows, in the inside of which was an empty bladder, the neck of which was adapted to the instrument, and gave entrance to a column of air when its parietes were separated: the air, in fact, does not enter the lungs but when the thorax dilates and enlarges by the separation of its parietes. To effect respiration, which may be defined the alternate entrance and egress of air in the lungs, the thorax must expand to receive the air, and contract to expel it. The dilatation is called inspiration, the contraction expiration; the latter is always shorter than the other, its causes are more mechanical, and the muscular powers have less influence.

The thorax in its usual state dilates only by depressing the diaphragm. The curved fibres of this muscle, that are made straight by contraction, causes it to descend towards the abdomen, and thus depressing its viscera, push forwards the anterior parietes of this cavity which sink down when expiration succeeds; the diaphragm is then relaxed and ascends, being pressed back by the abdominal viscera, on which the large muscles react. When we have occasion for the admission of a great quantity of air into the chest, it not only enlarges in length by the descent of the diaphragm, but its capacity is likewise increased in every direction. The intercostal muscles then contract, and approximate the ribs between which they are placed; yet the intercostal spaces become larger, particularly at the anterior part; for whenever oblique lines tend to become perpendicular to a vertical line, and to form right angles with it, the intercepted spaces augment in proportion as the lines, having been more oblique, approach the horizontal direction; besides, as the ribs present a double curvature in respect to their length, one on the front, the other on their sides, the convexity of the former is outwards, they separate from the axis of the chest, the cavity of which is enlarged transversely, while the latter curvature, agreeable to its edges, being augmented by a true rotatory motion, has its inferior edge pushed forwards. The thorax therefore increases both in its right and transverse diameter, each of which has been estimated to expand two lines; the extension of the vertical diameter dependent on the descent of the diaphragm is much more considerable.

When any cause whatever renders inspiration difficult, and prevents the diaphragm from descending towards the abdomen, or in any other manner impedes the motion of inspiration, the intercostal muscles not only evidently act to induce a dilatation

of the thorax, but also several other auxiliary muscles, as the *scaleni subcapulares*, *pectorales*, *serrati majores*, *latissimi dorsi*, in contracting raise the ribs, and increase the diameter of the thorax in several directions; the fixed point of these muscles should then be their movable part, because the cervical spine, the clavicle, scapula and humerus are fixed by other powers which it would be needless to enumerate. Inspiration is a state truly active, an effort of the contractile organs, which must cease when they fall into a state of relaxation. Expiration, which succeeds, is a passive motion in which few muscles co-operate, and chiefly depends on the re-action of the elastic parts constituting the structure of the parietes of the chest. It has been seen that the cartilages of the ribs experience a degree of rotation, carrying their upper edge backwards and downwards: when the cause that is productive of this action ceases to act, the parts return upon themselves, and carry back the sternum on the spine, towards which the ribs descend by their own gravity. The diaphragm is pushed nearer the thorax by the abdominal viscera, on which the large muscles of the abdomen react.

In every effort of expiration, as coughing and vomiting, the muscles react, not only in consequence of their own elasticity, but they still contract and approximate the spine, propelling the viscera towards the thorax. The *musculus triangularis* of the sternum, the *subcostales*, and the *serratus minor inferior*, may be ranked among the expirators, but they are seldom employed, and form too slender and weak powers to contribute much to the contraction of this cavity. When the chest enlarges, the lungs dilate, following the parietes which expand, and each time the thorax dilates in an adult man, from thirty to forty cubic inches of atmospheric air enter into the lungs, and, when in a state of purity, composed of seventy-three parts of azot, twenty-seven parts of oxygen and one-hundredth or two-hundredths of carbonic acid. After the atmospheric air has remained for some time in the pulmonary structure, it is expelled by the efforts of expiration, and its quantity is diminished to thirty-eight inches. And, if examined, the composition is not the same: there are found certainly seventy-three hundredths of azot, but the oxygen, its vital and respirable part, has suffered great diminution, its proportion is only fourteen-hundredths; carbonic acid constitutes the remaining thirteen hundredths, and sometimes one or two parts of hydrogen gas are found. It is likewise altered by the admixture of an aqueous vapour, which condenses in cold weather in passing out of the nostrils and mouth. This is known by the name of be *humour* of pulmonary exhalation.

These changes, compared with those the blood has suffered in its passage through the lungs, manifestly indicate a reciprocal action of this liquid and the oxygen of the atmosphere. The dark venous blood, slow of coagulation, and separating much serum, loaded with hydrogen and carbon, possesses only thirty degrees of heat, gives off to the oxygen of the atmosphere its hydrogen and carbon, to constitute the carbonic acid, and pulmonary vapour; and, as oxygen cannot enter these new combinations without disengaging a portion of caloric, which rarefies it into gas, the blood seizes this heat, now liberated with so much greater facility as it proportionably loses its hydrogen and carbon, and, agreeable to the ingenious experiments of Crawford, its capacity for caloric augments in the relation of 10 : 11 : 5.

The blood in its passage through the lungs, is deprived of hydrogen and carbon, and, in becoming arterial, is loaded with oxygen and caloric, which are formed into oxyds of hydrogen and carbon; these, by a fresh addition of oxygen, which is derived from the atmospheric air in the lungs, form water and carbonic acid.

Arterial blood becomes venous when any cause suspends or retards is course, as proved by the following experiment of J. Hunter. He tied the carotid of a dog in two places, at about four inches distance; the blood which came out of that portion of the artery between the ligatures, when opened several hours afterwards, was coagulated and dark like that of the veins.

The blood flowing in small veins very much resembles that of the arteries; and often in a copious bleeding the colour of the blood, at first very dark, becomes gradually paler, so that, towards the end of the operation, the blood which flows exhibits the qualities of that in the arteries, which is occasioned by rendering a more easy and direct passage of the blood from the arteries into the veins by emptying the venous system.

Richerand, in explaining the theory of respiration, thinks it probable that the oxygen absorbed in uniting with the hydrogen of every part of the body, produces the water that dilutes the venous blood, which is more fluid, and abounds more in serum than arterial blood; whilst this in uniting with carbon, forms an oxyd that gives this blood a dark colour, constituting one of its principal characteristics. The water in the lungs exhales, dissolved by the air, and forms pulmonary transpiration or exhalation; the oxyd of carbon, more completely burnt by a super-addition of oxygen, constitutes carbonic acid, which gives the air, passed by expiration, the property of precipitating lime-water.

By means of the absorption of oxygen by venous blood, we can explain how the

phenomena of respiration continues in every part of the body, and give rise to heat uniformly spread throughout all our organs. In proportion as the blood loses its caloric, for which its affinity or capacity diminishes as it becomes venous, the parts which give off their hydrogen and carbon seize it. If the lungs were the only organs in which the matter of heat could be disengaged, the temperature of these viscera should be much higher than that of other parts, and experience proves that it is not sensibly augmented.

Rest harrow. See *Oenitis*.

RESTA BOVIS. The rest harrow is so called because it hinders the plough; hence *resta bovis*. See *Oenitis*.

RESUSCITATION. The restoring of persons, apparently dead, to life. Under this head, strictly speaking, is considered the restoring of those who faint, or have breathed noxious and irrespirable air; yet it is chiefly confined to the restoring of those who are apparently dead from being immersed in a fluid, or by hanging. Dr. Curry, of Guy's hospital, has written a very valuable treatise on this subject; and such is its importance, that we have thought proper to insert the following account:

"From considering," he observes, "that a drowned person is surrounded by water instead of air, and that in this situation he makes strong and repeated efforts to breathe, we should expect that the water would enter and completely fill the lungs. This opinion, indeed, was once very general, and it still continues to prevail among the common people. Experience, however, has shewn, that unless the body lies so long in the water as to have its living principle entirely destroyed, the quantity of fluid present in the lungs is inconsiderable; and it would seem that some of this is the natural moisture of the part accumulated; for, upon drowning kittens, puppies, &c. in ink, or other coloured liquors, and afterwards examining the lungs, it is found that very little of the coloured liquor has gained admittance to them. To explain the reason why the lungs of drowned animals are so free from water, it is necessary to observe, that the muscles which form the opening into the wind-pipe are exquisitely sensible, and contract violently upon the least irritation, as we frequently experience when any part of the food or drink happens to touch them. In the efforts made by a drowning person, or animal, to draw in air, the water rushes into the mouth and throat, and is applied to these muscles, which immediately contract in such a manner as to shut up the passage into the lungs. This contracted state continues as long as the muscles retain the principle of life, upon which the power of muscular contraction depends; when that is gone, they become relaxed, and the water enters the wind-

pipe, and completely fills it. On dissecting the body of a recently drowned animal, no particular fulness of the vessels within the skull, nor any disease of the brain or its membranes, are visible. The lungs are also sound, and the branches of the wind-pipe generally contain more or less of a frothy matter, consisting chiefly of air, mixed with a small quantity of colourless fluid. The *right* cavity of the heart, and the trunks of the large internal veins which open into it, and also the trunk and larger branches of the artery which carries the blood from this cavity through the lungs, are all distended with dark coloured blood, approaching almost to blackness. The *left* cavity of the heart, on the contrary, is nearly, or entirely empty, as are likewise the large veins of the lungs which supply it with blood, and the trunk and principal branches of the great artery which conveys the blood from hence to the various parts of the body. The external blood-vessels are empty; and the fleshy parts are as pale as if the animal had been bled to death. When a body has lain in the water for some time, other appearances will also be observable; such as, the skin livid, the eyes blood shot, and the countenance bloated and swollen; but these appearances, though certainly unfavourable, do not absolutely prove that life is irrecoverably gone. It is now known, that in the case of drowning, no injury is done to any of the parts essential to life; but that the *right* cavity of the heart, together with the veins and arteries leading to and from that cavity, are turgid with blood, whilst every other part is almost drained of this fluid. The practice of holding up the bodies of drowned persons by the heels, or rolling them over a cask, is unnecessary; the lungs not being filled with any thing that can be evacuated in this way. Therefore such a practice is highly dangerous, as the violence attending it may readily burst some of those vessels which are already overcharged with blood, and thus convert what was only suspended animation, into absolute and permanent death. The operation of inflating the lungs is a perfectly safe, and much more effectual method of removing any frothy matter they may contain; and whilst it promotes the passage of the blood through them, also renders it capable of stimulating the *left* cavity of the heart, and exciting it to contraction. As soon as the body is taken out of the water, it should be stripped of any clothes it may have on, and be immediately well dried. It should then be wrapped in dry, warm blankets, or in the spare clothes taken from some of the by-standers, and be removed as quickly as possible to the nearest house that can be got convenient for the purpose. The fittest will be one that has a tolerably large apartment, in which a fire is ready,

or can be made. The body may be carried in men's arms, or laid upon a door; or, in case the house be at a distance from the place, if a cart can be procured, let the body be placed in it, on one side, upon some straw, with the head and upper parts somewhat raised; and in this position a brisk motion will do no harm. Whatever be the mode of conveyance adopted, particular care should be taken that the head be neither suffered to hang backwards, nor to bend down with the chin upon the breast. When arrived at the house, lay the body on a mattress, or a double blanket, spread upon a low table, or upon a door supported by stools; the head and chest being elevated by pillows. As the air of a room is very soon rendered impure by a number of people breathing in it, for this reason as well as to avoid the confusion and embarrassment attending a crowd, no more persons should be admitted into the apartment where the body is placed, than are necessary to assist immediately in the recovery: in general six will be found sufficient for this purpose, and these should be the most active and intelligent of the by-standers. It will be found most convenient to divide the assistants into two sets; one set being employed in restoring the heat of the body, while the other institutes an artificial breathing in the best manner they are able. Every skilful person should be provided with a flexible tube made of elastic gum, half a yard in length, to introduce into the wind-pipe, and also with a similar tube to which a syringe can be affixed, to be put into the œsophagus. Should these not be at hand, air should be thrown into the lungs in the best manner that can be suggested at the time. Should it still be found that the air does not pass readily into the lungs, immediate recourse must be had to another and more effectual method for attaining that object. As this method, however, requires address, and also some knowledge of the parts about the throat, we would recommend that when there is not a medical gentleman present, the mode already described be tried repeatedly before this be attempted. As a quantity of frothy matter occupying the branches of the wind-pipe and preventing the entrance of the air into the lungs, is generally the circumstance which renders this mode of inflation necessary, the mouth should be opened from time to time to remove this matter as it is discharged. While one set of the assistants are engaged in performing artificial respiration, the other should be employed in communicating heat to the body. The warm bath has been usually recommended for this purpose; but wrapping the body in blankets, or woollen clothes, strongly wrung out of warm water, and renewing them as they grow cool, besides being a speedier and more practicable method of imparting

heat, has this great advantage, that it admits of the operation of inflating the lung being carried on without interruption. Until a sufficient quantity of warm water can be got ready, other methods of restoring warmth may be employed; such as the application of dry, warm blankets round the body and limbs; bags of warm grains or sand, bladders or bottles of hot water, or hot bricks applied to the hands, feet, and under the arm-pits, the bottles and bricks being covered with flannel: or the body may be placed before the fire, or in the sunshine if strong at the time, and be gently rubbed by the assistants with their warm hands, or with cloths heated at the fire or by a warming-pan. The restoration of heat should always be gradual, and the warmth applied ought never to be greater than can be comfortably borne by the assistants. If the weather happen to be cold, and especially if the body has been exposed to it for some time, heat should be applied in a very low degree at first: and if the weather be under the freezing point, and the body when stripped, feels cold and nearly in the same condition with one that is frozen, it will be necessary at first to rub it well with snow, or wash it with cold water; the sudden application of heat in such cases, having been found very pernicious. In a short time, however, warmth must be gradually applied. To assist in rousing the activity of the vital principle, it has been customary to apply various stimulating matters to different parts of the body. But as some of these applications are in themselves hurtful, and the others serviceable only according to the time and manner of their employment, it will be proper to consider them particularly. The application of all such matters in cases of apparent death, is founded upon the supposition that the skin still retains sensibility enough to be affected by them. It is well known, however, that even during life, the skin loses sensibility in proportion as it is deprived of heat, and does not recover it again until the natural degree of warmth be restored. Previous to the restoration of heat, therefore, to a drowned body, all stimulating applications are useless, and so far as they interfere with the other measures, are also prejudicial. The practice of rubbing the body with salt or spirits is now justly condemned. The salt quickly frets the skin, and has in some cases produced sores, which were very painful and difficult to heal after recovery. Spirits of all kinds evaporate fast, and thereby, instead of creating warmth, as they are expected to do, carry off a great deal of heat from the body. Spirit of hartshorn, or of salt volatile, are liable to the same objection as brandy or other distilled spirits, and are besides very distressing to the eyes of the assistants. When there is reason to think

the skin has in any degree recovered its sensibility, let an assistant moisten his hand with spirit of hartshorn, or *essu de face*, and hold it closely applied to one part: in this way evaporation is prevented, and the full stimulant effect of the application obtained. A liniment composed of equal parts of spirit of hartshorn and salad oil, well shaken together, would appear to be sufficiently stimulating for the purpose, and as it evaporates very slowly, will admit of being rubbed on without producing cold. The places to which such remedies are usually applied, are, the wrists, ankles, temples, and the parts opposite the stomach and heart. The intestines, from their internal situation and peculiar constitution, retain their irritability longer than the other parts of the body, and, accordingly, various means have been proposed for increasing the action of their fibres, in order to restore activity of the whole system. Tobacco-smoke, injected by way of clyster, is what has been generally employed with this view, and the *fumigator*, or instrument for administering it, makes a part of the apparatus which is at present distributed by the different societies established for the recovery of drowned persons. Of late, however, the use of tobacco-smoke has been objected to, and upon very strong grounds; for when we consider that the same remedy is successfully employed with the very opposite intention, namely, that of lessening the power of contraction in the muscles, and occasioning the greatest relaxation consistent with life, it must be acknowledged to be a very doubtful, if not dangerous remedy, where the powers of life are already nearly exhausted. Instead of tobacco-smoke then, we would recommend a clyster, consisting of a pint or more of water, moderately warmed, with the addition of one or two table-spoonfuls of spirit of hartshorn, a heaped tea-spoonful of strong mustard, or a table-spoonful of essence of peppermint: in defect of one or other of these, half a gill or more, of rum, brandy, or gin may be added, or the warm water given alone. This step however, need not be taken, until artificial respiration has been begun; for it will answer but little purpose to stimulate the heart through the medium of the intestines, unless we, at the same time, supply the left cavity with blood fitted to act upon it; which we cannot do without first removing the collapsed state of the lungs, and promoting the passage of the blood through them by a regular inflation. As the stomach is a highly sensible part, and intimately connected with the heart and brain, the introduction of some moderately warm and stimulating liquor into it, seems well calculated to rouse the dormant powers of life. This is very conve-

niently done by means of the syringe and flexible tube. The quantity of fluid thrown in ought not to exceed half a pint, and may be either warm negus, or water with the addition of one or other of the stimulating matters recommended above, using, however, only half the quantities mentioned there. As soon as the pulse or beating of the heart can be felt, the inside of the nostrils may be occasionally touched with a feather dipped in spirit of hartshorn, or sharp mustard; it being found by experience, that any irritation given to the nose, has considerable influence in exciting the action of the muscles concerned in respiration. When the natural breathing commences, the flexible tube and canula should be withdrawn, and any farther inflation that may be necessary, performed by blowing into the nostril. Letting blood has been generally thought requisite in every case of suspended animation. The practice, however, does not appear to have been founded upon any rational principle at first, and it has been continued from the force of custom, rather than from any experience of its good effects. In the case of drowned persons there is not, as in those who suffer from hanging or apoplexy, any unusual fullness of the vessels of the brain; and the quantity of blood that can be drawn from the external veins, will not sensibly diminish the accumulation of it in those near the heart. Besides, blood-letting, which always tend to *lessen* the action of the heart and arteries in the living body, cannot be supposed to have a directly opposite effect in cases of apparent death; on the contrary, if employed here, it will hazard the entire destruction of those feeble powers which yet remain, and to increase and support which all our endeavours should be directed. When the several measures recommended above have been steadily pursued for an hour or more, without any appearance of returning life, electricity should be tried; experience having shewn it to be one of the most powerful stimuli yet known, and capable of exciting contraction in the heart and other muscles of the body, after every other stimulus had ceased to produce the least effect. Moderate shocks are found to answer best, and these should, at intervals, be passed through the chest in different directions, in order, if possible, to rouse the heart to act. Shocks may likewise be sent through the limbs, and along the spine; but we are doubtful how far it is safe or useful to pass them through the brain, as some have recommended. The body may be conveniently insulated, by placing it on a door, supported by a number of quart bottles, whose sides are previously wiped with a towel to remove any moisture they may have contracted. By experiments made

on different animals, it is found that the blood passes through the lungs most readily when they are fully distended with air; consequently, that if the lungs of a drowned person are inflated, and kept in the expanded state whilst the electric shock is passed through the chest, the blood accumulated in the *right* cavity of the heart and its vessels, will move forward without any resistance, should the heart be brought to contract upon it. As soon as the shock is given, let the lungs be emptied of the air they contain, and filled again with fresh air; then pass another shock, and repeat this until the heart is brought into action, or until it appear that all farther attempts are useless. In order more certainly to pass the shock through the heart, place the knob of one discharging rod above the collar-bone of the right side, and the knob of the other above the short ribs of the left: the position of the discharging rods, however, may be changed occasionally, so as to vary the direction of the shock. Two thick brass wires, each about eighteen inches long, passed through the two glass tubes, or wooden cases, well varnished, and having at one end a knob, and at the other a ring to fasten the brass chain to, form very convenient discharging rods; and by means of them, the shock may be administered without the risk of its being communicated to the assistants, or carried off by the skin being wet. When the patient is so far recovered as to be able to swallow, he should be put into a warm bed, with his head and shoulders somewhat raised by means of pillows. Plenty of warm wine-*whew*, *ale-posset*, or other light and moderately nourishing drink, should now be given, and gentle sweating promoted, by wrapping the feet and legs in flannels well wrung out of hot water. If the stomach and bowels feel distended and uneasy, a clyster, consisting of a pint of warm water, with a table spoonful of common salt, or an ounce or more of Glauber's or Epsom salt, dissolved in it, may be administered. The general practice, in this case, is to give an emetic; but considering that the powers of the machine are still very weak, the agitation of vomiting is certainly hazardous. The patient should on no account be left alone, until the senses are perfectly restored, and he be able to assist himself; several persons having relapsed and been lost, from want of proper attention to them, after the vital functions were, to all appearance, completely established. Either from the distention which the arteries of the lungs have suffered, or from the sudden change from great coldness to considerable warmth, it now and then happens, that the patient is attacked, soon after recovery, with inflammation of some of the parts within the chest. This occurrence is pointed out by

pain in the breast or side, increased on inspiration, and accompanied with frequent, and full or hard pulse, and sometimes with cough. Here the taking away some blood from the arm, or the application of cupping-glasses, leeches, or a blister, over the seat of the pain, will be very proper; but the necessity for these measures, as well as the time for putting them in practice, should be left to the judgment and discretion of a medical person. Dull pain in the head, lasting sometimes for two or three days, is by no means an unfrequent complaint in those who are recovered from this and from the other states of suspended animation; and here also a moderate bleeding from the neck, either with the lancet or with cupping-glasses, may prove serviceable.

In hanging the external veins of the neck are compressed by the cord, and the return of the blood from the head thereby impeded. from the moment that suspension takes place; but as the heart continues to act for a few seconds after the wind-pipe is closed, the blood which is sent to the head during this interval, is necessarily accumulated there. Hence it is, that in hanged persons the face is greatly swolen, and of a dark red or purple colour: the eyes are commonly suffused with blood, enlarged, and prominent. On dissection, the blood-vessels of the brain are found considerably distended; but in general, no further marks of disease appear within the skull. The lungs are found generally quite collapsed, and free from frothy matter. The heart and the large blood-vessels adjoining to it, exhibit the same appearances as in the bodies of drowned persons. From the great accumulation of blood in the vessels of the head, many have been of opinion, that hanging kills chiefly by inducing apoplexy; but the following experiment made at Edinburgh several years ago, by an eminent medical professor there, clearly proves, that in hanging, as well as in drowning, the exclusion of air from the lungs is the immediate cause of death. A dog was suspended by the neck with a cord, an opening having been previously made in the wind-pipe, below that place where the cord was applied, so as to admit air into the lungs. In this state he was allowed to hang for three quarters of an hour, during which time the circulation and breathing went on. He was then cut down, without appearing to have suffered much from the experiment. The cord was now shifted below the opening into the wind-pipe, so as to prevent the ingress of air to the lungs; and the animal being again suspended, he was completely dead in a few minutes. Upon the whole, then, it appears, that the same measures recommended for drowned persons, are also necessary here;

with this addition, that opening the jugular veins, or applying cupping-glasses to the neck, will tend considerably to facilitate the restoration of life, by lessening the quantity of blood contained in the vessels of the head, and thereby taking off the pressure from the brain. Except in persons who are very full of blood, the quantity taken away need seldom exceed an ordinary tea cupful, which will in general be sufficient to unload the vessels of the head, without weakening the powers of life.

RETE. A net. A congeries of vessels, or any animal substance resembling a net.

RETE MALPHIGII. A fine net-work of the extremities of the pulmonary arteries.

RETE MIRABILE. A net-work of blood-vessels in the basis of the brain of quadrupeds.

RETE MUCOSUM. *Corpus reticulare. Corpus mucosum. Mucus Mulphigii.* A mucous substance, deposited, in a net-like form, between the epidermis and cutis, which covers the sensible cutaneous papillæ, connects the epidermis with the cutis, and gives the colour to the body: in Europeans it is of a white colour, in Ethiopians black. See *Skin*.

RETICULAR. (*Reticularis*, from *rete*, a net.) Interwoven like a net.

RETIFORM. (*Retiformis*, from *rete*, a net, and *forma*, resemblance.) Net-like.

RETINA. (From *rete*, a net.) *Amphiblastroides*. The third or innermost membrane of the eye, expanded round the choroid coat, to the ciliary ligament. It is the true organ of vision, and is formed by an expansion of the pulp of the optic nerve.

RETINACULUM. (From *retinere*, to prop or restrain.) An instrument for keeping the bowels in their place.

RETORT. (*Retorta*, from *reterquere*, to bend back again; probably so called because its neck was curved and bent back again.) A chemical vessel employed for many distillations, and most frequently for those which require a degree of heat superior to that of boiling water. They differ in form and materials: when pierced with a little hole in their roof, they are called tubulated retorts. They are made of common glass, stone ware, and iron.

RETRACTOR ANGULI ORIS. See *Buccinator*.

RETRAHENTES AURIS. *Posterior auris* of Winslow. *Retrahentes auriculæ* of Albinus. *Deprimens auriculæ* of Douglas. *Retrahentes auriculam* of Cowper, and *mastoidæ conchionis* of Dumas. Two small bundles of muscular fibres which arise from the external and posterior part of the mastoid process of the temporal bone immediately above the insertion of the sterno-cleido-mastoidæus muscle. They

are inserted into that part of the back of the ear which is opposite to the septum which divides the concha and scapha. Their use is to draw the ear backwards, and stretch the concha.

Retrocédens gout. See *Arthritis*.

RETROVERSION. See *Uterus*, *Retroversion of*.

REVERBATORY FURNACE. See *Furnace*.

REVULSION. (*Revulsio*, from *revello*, to draw away.) An old term used by the humoral pathologists, signifying the drawing of humours a contrary way.

RHABBARBARUM. (From *Rha*, and *barbarus*, wild; so called because it was brought from the banks of the Rha, now called the Wolgar, in Russia.) *Rheon. Rheum. Rhaum. Barbaria. Lappathum orientale. Lappathum chinense. Rhubarbarum verum. Rhubarbarum tartaricum.* Rhubarb. It was not until the year 1732 that naturalists became acquainted with any plant which seemed to afford the *Rhabbarbarum officinale*, when some plants received from Russia by Jussieu at Paris, and Rhau at Chelsea, were said to supply this important desideratum, and as such were adopted by Linnæus, in his first edition of the species *Plantarum*, under the name of *Rheum Rhubarbarum*. This, however, was not generally received as the genuine rhubarb plant; and with a view to ascertain this matter more completely, Kaw Boerhaave procured from a Tartarian rhubarb merchant the seeds of those plants whose roots he annually sold, and which were admitted at Petersburg to be the true rhubarb. These seeds were soon propagated, and were discovered by De Gorter to produce two distinct species, viz. the *R. Rhubarbarum* of Linnæus, or as it has since been called, the *R. Undulatum*, and another species, a specimen of which was presented to Linnæus, who declared it to be a new one, and was first mentioned in the second edition of the species *Plantarum*, in 1762, by the name of *R. Palmatum*. Previous to this time De Gorter had repeatedly sent its seeds to Linnæus, but the young plants which they produced constantly perished; at length he obtained the fresh root, which succeeded very well at Upsall, and afterwards enabled the younger Linnæus to describe this plant, ann. 1767. But two years antecedent to this, Dr. Hope's account of the *Rheum palmatum*, as it grew in the botanic garden near Edinburg, had been read before the Royal Society at London; and of the great estimation in which this plant was held by him, we have the following proof: "From the perfect similarity of this root with the best foreign rhubarb, in taste, smell, colour, and purgative qualities, we cannot doubt of our being at last

possessed of the plant which produces the true rhubarb, and may reasonably entertain the agreeable expectation of its proving a very important acquisition to Britain."

But from the relation we have given, it appears that both the seeds of the *R. Palmatum*, and the *R. Undulatum*, were transmitted to Petersburg, as those of the true rhubarb; we are therefore to conclude, that the former species has an equal claim to this importance with the latter; and from further inquiries made in Russia, there is the best authority for believing that the *R. Compactum* also affords this very useful drug. The seeds of the *R. Palmatum* were first introduced into Britain in 1762, by Dr. Hounsly (who sent them from Russia,) and were supposed to be a part of that already mentioned; and since their prosperous cultivation by the late professor of botany at Edinburgh, the propagation of this plant has been gradually extended to most of our English gardens, and with a degree of success which promises in time to supersede the importation of the foreign root. Two sorts of rhubarb roots are usually imported into this country for medical use, viz. the Chinese, and the Tartary rhubarb; the first is in oblong pieces, flattish on one side, and convex on the other; compact, hard, heavy, internally of a dull red colour, variegated with yellow and white, and when recently powdered, appears yellow, but on being kept becomes gradually redder. The second is the most valuable, and is brought to us in roundish pieces, with a large hole through the middle of each; it is more soft and friable than the former sort, and exhibits, when broken, many streaks of a bright red colour. "The marks of the goodness of rhubarb are, the liveliness of its colour when cut; its being firm and solid, but not flinty or hard; its being easily pulverable, and appearing when powdered of a fine bright yellow colour; its imparting to the spittle when chewed a deep saffron tinge, and not proving slimy or mucilaginous in the mouth; its taste is subacid, bitterish, and somewhat styptic; the smell lightly aromatic."

The purgative qualities of rhubarb are extracted more perfectly by water than by rectified spirit: the root remaining after the action of water is almost if not wholly inactive; whereas after repeated digestion in spirit, it proves still very considerably purgative. The virtue of a watery infusion, on being inspissated by a gentle heat is so much diminished, that a drachm of the extract is said to have scarcely any greater effect than a scruple of the root in substance. The spirituous tincture loses less; half a drachm of this extract proving

moderately purgative. The qualities of this root, says Dr. Cullen, are that of a gentle purgative, and so gentle that it is often inconvenient on account of the bulk of the dose required, which in adults must be from ʒss to ʒj. When given in a large dose it will occasion some griping, as other purgatives do; but it is hardly ever heating to the system, or shews the other effects of the more drastic purgatives. The purgative quality is accompanied with a bitterness, which is often useful in restoring the tone of the stomach when it has been lost; and, for the most part, its bitterness makes it sit better on the stomach than many other purgatives do. Its operation joins well with neutral laxatives; and both together operate in a lesser dose than either of them would singly. Some degree of stypticity is always evident in this medicine; and as this quality acts when that of the purgative has ceased, so, in cases of diarrhoea, when any evacuation is proper, rhubarb has been considered as the most proper remedy to be employed. It must, however, be remarked here, that in many cases of diarrhoea, no further evacuation than what is occasioned by the disease, is necessary or proper. The use of rhubarb, in substance, for keeping the belly regular, for which it is frequently employed, is by no means proper, as the adstringent quality is ready to undo what the purgative has done; but it is found that the purpose mentioned may be obtained by it, if the rhubarb is chewed in the mouth, and no more is swallowed than what the saliva has dissolved. And it must be remarked, that in this way employed it is very useful to dyspeptic persons. Analogous to this, is the use of rhubarb in solution, in which it appears to me, that the adstringent quality is not so largely extracted as to operate so powerfully as when the rhubarb was employed in substance."

The official preparations of this drug are, a watery and a vinous infusion, a simple and a compound tincture. It is also an ingredient in different compositions, as, the elixir ex aloë et rheo, pilulæ stomachicæ, and some others.

RHABBARBARUM ALBUM. See *Mecher-anna*.

RHABBARBARUM ANTIQUORUM. See *Rhe-ponticum*.

RHABBARBARUM BIOSCORIDIS. See *Rhe-ponticum*.

RHABBARBARUM MONACHORUM. *Hippopathum. Patensia.* Monks rhubarb. Garden patience. This root, which is supposed to possess the virtues of rhubarb, but in an inferior degree, is obtained from the *Rumex patensia* of Linnæus, and, according to Professor Murray, from the *Rumex alpinus* of Linnæus. It is obviously

more adstringent than rhubarb, but comes very far short of its purgative virtue.

RHABBARUM SIBIRICUM. The plant directed for medicinal purposes by this name is the *Rheum undulatum*; *foliis subvillosis undulatis petiolis equalibus* of Linnæus. It possesses similar virtues to those of the palmate species, and is in common use in Russia.

RHABBARUM TARTARICUM. See *Rhabbarum*.

RHABBARUM VERUM. See *Rhabbarum*.

RHACHIALGIA. (From *ραχις*, the spine of the back, and *αλγος*, pain.) A pain in the spine of the back.

RACHIS. (*Ραχις*, the spine of the back.) See *Spine*.

RHACHISAGRA. (From *ραχις*, the spine of the back, and *αγρα*, a prey.) A species of gout fixed in the spine of the back.

RHACHITA. (From *ραχις*, the spine of the back.) A muscle belonging to the spine of the back.

RHACHITIS. See *Rachitis*.

RHACOSIS. (From *ρακος*, a rag.) A ragged excoriation of the relaxed scrotum.

RHAGADES. (From *ρηννυμι*, to break or bruise.) Chaps. Clefts. Malignant, dry, and deep cutaneous fissures.

RHAGOIDES. (From *ραγ*, a grape-berry, and *οειδης*, a likeness; so called from its likeness in colour to a grape-seed.) Applied to the retiform tunic of the eye.

RHAMNI BACCÆ. Buckthorn-berries.

RHAMNUS. (From *ραιο*, to destroy, because of its many thorns.) The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*. Buckthorn.

RHAMNUS CATHARTICUS. The systematic name of the buckthorn. See *Spina oertina*.

RHAMNUS FRANGULA. The systematic name of the black alder. See *Frangula*.

RHAMNUS SIZYBIUS. The systematic name of the tree which affords the jujubs. See *Jujube*.

RHAPHANUS. See *Raphanus*.

RHAPONIC. A term applied to rhubarb.

Rhapontic, rhubarb. See *Rhaponticum*.

RHAPONTICUM. (The Rha of Pontus, i. e. the Rha, in Russia, a river on whose banks it grew.) *Rhabbarum dioscoridis*. *Rhabbarum antiquorum*. The root of this plant, *Rheum rhaponticum* of Linnæus, appears to have been the true rhubarb of the ancients. By some it is confounded with the modern rhubarb, though considerably different from that root in appearance, as well as in quality. The rhapontic is of a dusky colour on its surface, and a

loose spongy texture; is more adstringent than rhubarb, and less purgative; in this last intention, two or three drachms are required for a dose.

RHAPONTICUM VULGARE OFFICINARUM. *Centaurium majus*. Greater centaurry. The root of this plant *Centauria centaurum* of Linnæus, was formerly used as an aperient and corroborant in alvine fluxes. It is now totally discarded from the *Materia Medica* of this country.

RHATANIA. This substance has been long known to the manufacturers of port wine; it is the production of Peru, and is probably the root of the *cinchona cordifolia*. It is described as externally resembling the root of the *rubia tinctorum*, to the taste, being aromatic, bitter and very adstringent; its infusion or decoction turns black with sulphate of iron and precipitates tannin. The principal virtues appear to reside in the cortical part of the root which is thick and resinous. An opinion prevails that the substance sold in the shops under the name of foreign extract of bark is made from the rhatania.

It is well known that the medical virtues of this root are powerfully tonic. In debility of the digestive organs, in chronic rheumatism, fluor albus, and in intermittent fevers it has been employed with good effect. While given in doses similar to cinchona, it has the advantage of being only one third the price of that substance.

RHEI RADIX. Rhubarb root.

RHEUME. (From *ρεο*, to flow.) A defluxion, a common cold or catarrh.

RHEUM. (From *Rha*, a river in Russia.) The name of a genus of plants in the Linnæan system. Class, *Enneandria*. Order, *Trigynia*. See *Rhabbarum*.

RHEUM PALMATUM. The systematic name of the official rhubarb. See *Rhabbarum*.

RHEUM RHAPONTICUM. The systematic name of the rhapontic rhubarb. See *Rhaponticum*.

RHEUM UNDULATUM. The systematic name of the official rhubarb sibiricum. See *Rhabbarum sibiricum*.

RHEUMA. (From *ρεο*, to flow.) The discharge from the nostrils or lungs arising from cold; hence the following lines of the school of Salerno:

Si fuit ad pectus, dicatur rheuma catarrhus,

Ad fauces branchus, ad nares esto coryza.

RHEUMATISMUS. (From *ρευματισμος*, to be afflicted with defluxions.) *Dolores rheumatici et arthritici* of Hoffman. *Myositis* of Sagar. This term is so called from its being formerly used in the same sense as *rheuma*; but, in the present day, the meaning of this word is applied to a genus of disease in the class *pyrexie*, and order

phlegmasia of Cullen; characterised by pyrexia, pains in the joints, increased by the action of the muscles belonging to the joint, and heat on the part. The blood, after venesection, exhibits an inflammatory crust. Rheumatism terminates in arthrodynia, lumbago, and ischias. Rheumatism is distinguished into *acute* and *chronic*. The acute is preceded by shivering, heat, thirst, and frequent pulse; after which the pain commences, and soon fixes on the joints. The chronic rheumatism is distinguished by pain in the joints, without pyrexia, and is divided into three species; *lumbago*, affecting the loins; and *ischias*, or *sciatica*, affecting the hip, and *arthrodynia*, or pains in the joints. The acute species mostly terminate in one of these species.

Rheumatism may arise at all times of the year, when there are frequent vicissitudes of the weather, from heat to cold, but the spring and autumn are the seasons in which it is most prevalent; and it attacks persons of all ages; but very young people are less subject to it than adults.

Obstructed perspiration, occasioned either by wearing wet clothes, laying in damp linen, or damp rooms, or by being exposed to cool air when the body has been much heated by exercise, is the cause which usually produces rheumatism. Those who are much afflicted with this complaint, are very apt to be sensible of the approach of wet weather, by finding wandering pains about them at that period.

Acute rheumatism usually comes on with lassitude and rigors, succeeded by heat, thirst, anxiety, restlessness, and a hard pulse; soon after which, excruciating pains are felt in different parts of the body, but more particularly in the joints of the shoulder, wrist, knees, and ankles, or perhaps in the hip; and these keep shifting from one joint to another, leaving a redness and swelling in every part they have occupied, as likewise a great tendency to the touch. Towards evening there is usually an exacerbation, or increase of fever; and, during the night, the pains become more severe, and shift from one joint to another.

Early in the course of the disease, some degree of sweating usually occurs; but it is seldom so copious as either to remove the pains or to prove critical. In the beginning, the urine is without any sediment; but as the disease advances in its progress, and the fever admits of considerable remissions, a lateritious sediment is deposited; but this by no means proves critical.

Chronic rheumatism is attended with pains in the head, shoulders, knees, and other large joints, which at times are confined to one particular part, and at others shift from one joint to another, without

occasioning any inflammation or fever; and in this manner the complaint continues often for a considerable time, and at length goes off.

No danger is attendant on chronic rheumatism; but a person having been once attacked with it, is ever afterwards more or less liable to returns of it; and an incurable anchylosis is sometimes formed, in consequence of very frequent relapses. Neither is the acute rheumatism frequently accompanied with much danger; but, in a few instances, the patient has been destroyed by general inflammation, and, now and then, by a metastasis to some vital part, such as the head and lungs. Acute rheumatism, although accompanied with a considerable degree of inflammation in particular parts, has seldom been known to terminate in suppuration; but a serous or gelatinous effusion takes place.

Rheumatism seldom proving fatal, very few opportunities have offered for dissections of the disease. In the few which have occurred, the same appearances have been observed in inflammatory fever, as effusion within the cranium, and now and then affections of some of the viscera.

RHIBESIA. (From *ribes*, a currant.) See *Ribes nigrum*, *Ribes rubrum*, and *Fruita, summer*.

RHINCEUS. (*Rhinenceus*, sc. *musculus*; from *rhin*, the nose.) See *Compressor naris*.

RHINENCHYTES. (From *rhin*, the nose, and *enchyon*, to pour in.) A syringe for the nose.

RHINOPHONIA. (From *rhin*, the nose, and *phonia*, the voice.) A nasal voice.

RHIZAGRA. (From *rhiza*, the root, and *aggrao*, to seize.) An instrument for taking out the roots or stumps of teeth.

RHODIA. (From *rhodon*, a rose; so called because its root smells like the damask rose.) The radix rhodiz of some pharmacopœias is the produce of the *Rhodiola rosea* of Linnaeus, called rose-wort. When dry, it has a very pleasant smell, resembling that of the damask rose. In this odorous matter the medical virtue of the root resides. Poultices in which this root enters as a chief ingredient are said to allay violent pains of the head.

RHODIUM LIQNUM. (From *rhodon*, a rose; a wood which smells like roses.) Rhodium, or rose-wood. The wood or root of a tree supposed to be the *Genista canariensis* of Linnaeus. It is brought from the Canary islands. An essential oil is obtained from it, which is used principally as a perfume, but possesses cordial and corroborant virtues. It is also an ingredient in the composition of powders for the destruction of rats.

RHODODENDRON. (From *rhodon*, a rose; and *dendron*, a tree; so called because its flowers resemble the rose.) 1.

The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Monogynia*.

2. The pharmacopœial name of the oleander. Rose-bay. Yellow-flowered rhododendron. *Rhododendron chrysanthemum* Linnæi:—*foliis oblongis impunctis supra scabris venosis, corolla rotata irregulari, gemmâ florifera ferrugineo-tomentosa*. This species of rhododendron has not yet been introduced in Britain; it is a native of Siberia, affecting mountainous situations, and flowering in June and July.

This plant and its medical virtues were first described in 1747, by Gmelin and Haller. Little attention, however, was paid to it, till the year 1779, when it was strongly recommended by Koelpin as an efficacious medicine, not only in rheumatism and gout, but even in venereal cases; and it is now very generally employed in chronic rheumatisms, in various parts of Europe. The leaves, which are the part directed for medicinal use, have a bitterish subadstringent taste. Taken in a large dose, they prove a narcotic poison; and, in moderate doses, they are said to occasion heat, thirst, a degree of delirium, and a peculiar sensation of the parts affected.

As a powerful and active medicine, this shrub, says Dr. Woodville, may probably be found an addition to the materia medica. Dr. Home, who tried it unsuccessfully in some cases of acute rheumatism, says, "It appears to be one of the most powerful sedatives which we have, as, in most of the trials, it made the pulse remarkably slow, and in one patient reduced it to thirty-eight beats. And in other cases, in which the rhododendron has been used at Edinburgh, it has been productive of good effects, and accordingly it is now introduced into the Edinburgh Pharmacopœia." The manner of using this plant by the Siberians, was by putting two drachms of the dried leaves in an earthen pot, with about ten ounces of boiling water, keeping it near a boiling heat for a night; and this they took in the morning, and by repeating it three or four times, generally effected a cure.

RHODODENDRON CHRYSANTHEMUM. The systematic name of the oleander, or rose-bay. See *Rhododendron*.

RHODOLA ROSEA. The systematic name of the rose-wort. See *Rhodia*.

RHODOMELI. (From *μέλι*, the rose, and *μέλι*, honey.) Honey of roses.

RHODOS PETALA. Red poppy petals.

RHŒAS. (From *ῥέω*, to flow.) The wild poppy is sometimes so called. See *Papaver erraticum*.

RHOMBOIDEUS. (From *ῥόμβος*, a geometrical figure, whose sides are equal but not right-angled, and *ειδος*, resemblance.) *Rhomboideus major* and *minor*. *Rhomboideus* of Douglas, Winslow, and Cowper; and

Cervici dorso scapulaire of Dumas. This muscle, which is so named from its shape, is situated immediately under the trapezius. We find it usually, though not always, divided into two portions, which Albinus describes as two distinct muscles. The uppermost of these, or *rhomboideus minor*, arises tendinous from the spinous process of the three inferior vertebræ of the neck, and from the ligamentum colli; the lowermost, or *rhomboideus major*, arises tendinous from the spinous processes of the back: the former is inserted into the basis of the scapula, opposite to its spine; the latter into all the basis of the scapula, below its spine. Its use is to draw the scapula obliquely upwards, and directly backwards.

RHOPALOSIS. (From *ῥοπαλόν*, a club.) A disorder in which the hair cleaves together, and hangs down in clusters resembling clubs. The plaited hair.

Rhubarb. See *Rhubarbium*.

Rhubarb, monks. See *Rhabarbarum monachorum*.

Rhubarb, rhapsodic. See *Rhaponticum*.

RHUS. (From *ῥέω*, to flow: so called because it stops fluxes.) The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Trigynia*. The sumach-tree.

RHUS BELGICA. The Dutch myrtle is sometimes so termed. See *Myrtus brabantica*.

RHUS CORIARIA. See *Sumach*.

RHUS RADICANS. A poisonous plant, *Rhus vernix* of Linnæus, the efficacy of which Dr. Fresnoi has endeavoured to prove, in the disease called paralysis, and herpetic affections. He, in order that others should not suffer by his experiments, began by taking an infusion of one of the three folioli, of which each leaf of this plant consists; and as this dose produced no sensible effect, he increased the number to twelve. His urine and perspiration were increased in quantity, and he had some pains in his belly. He relates seven cases, in which he thinks he can remove all doubt of the efficacy of this infusion, in herpetic affections. From these the following are selected:

"A country woman," says Dr. Fresnoi, "came to me in the month of July, 1780. To consult me about the herpes farinosa, with which her face had been covered for more than a year. She was ordered to make an infusion of this plant; and, in six weeks, was entirely free from the disease."

He likewise relates five cases of paralysis, which were cured by the use of this plant.

The leaves of this plant are to be cut when in the greatest vigour, about the month of June. "Those who cut this plant," says Dr. F. "wear leathern gloves, on account of its poisonous qualities." The

same gentleman observes, he saw one case in which inflammation of the eye-lids was produced by the vapour from the plant. Four pounds of the leaves being distilled with thirty-two pounds of water, gives it a slight odour, although the plant is entirely free from it. Its taste is pungent, and inflames the mouth. The decoction which remains in the still is brown, and is generally covered with a light brown pellicle. When strained and evaporated, it gives a shining black extract. The leaves inflame and swell the hands and arms of those who take them out of the still, and brings on an itching, which remains for several days. Forty-two pounds of the leaves afford twenty ounces of extract, of a proper consistence for pills.

"A girl, in Flanders," says Dr. Fresnoi, "already subject to fits, laid down some flowers in her bed-room. Next day she told me that she had undergone a great change; that she had had no fits, and slept much better. It occurred to me," says Dr. F. "that the flowers occasioned this change. Next day, the flowers being removed, and the window opened, the convulsions re-appeared; on their being again introduced, the fits disappeared; which proved plainly it was the effect of the flowers. The success of the extract in tussis convulsiva exceeded my hopes; forty-two children being cured of this disorder in Valenciennes, during the end of the year 1786. Four grains of extract dissolved in four ounces of syrup, of which one table spoonful given to the child every third hour, generally abates the cough, and mostly leaves them.

RHUS TIFINUM The systematic name of the Virginian sumach, whose seeds are said to be useful in stopping hæmorrhages.

RHUS VERNIX. The systematic name of a species of sumach which exudes a poisonous resin. See *Rhus radicans*.

RHYAS. (*pus*, a disease of the eye.) A decrease or defect of the lachrymal caruncle. The proximate cause is a native defect; or it may originate from excision, erosion, or acrimony. This disorder is commonly incurable, and it induces an incurable *epiphora*, or a continual weeping.

RHYTINOSIS. See *Rutidosis*.

RIBES. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

RIBES NIGRUM. Black currant. This indigenous plant, *Ribes nigrum* of Linnaeus:—*racemis pilosis, floribus oblongis*, affords larger berries than those of the red, which are said to be peculiarly useful in sore throats, and to possess a diuretic power in a very considerable degree. The leaves of the black currant are extremely fragrant, and have been likewise recommended for their medicinal virtue, which Bergius states to be *mundificans, pellens,*

diuretica. The officinal preparations of the berries are the *syrupus ribis nigri*, and the *succus ribis nigri inspissatus*.

RIBES RUBRUM. *Grossularia nem spinosa*. The red currant. *Ribes rubrum* of Linnaeus:—*inermis, racemis glabris pendulis, floribus planiusculis*. The white currant-tree is merely a variety of the red, the fruit of both is perfectly analogous; therefore what is said of the one applies to the other. The red currant is abundantly cultivated in gardens, and, from its grateful acidity, is universally accepted, either as nature presents it, or variously prepared by art, with the addition of sugar. Considered medicinally, it is esteemed to be moderately refrigerant, antiseptic, attenuant, and aperient. It may be used with considerable advantage to allay thirst, in most febrile complaints, to lessen an increased secretion of bile, and to correct a putrid and scorbutic state of the fluids, especially in sanguine temperaments; but, in constitutions of a contrary kind, it is apt to occasion flatulency and indigestion.

RIBS. (*Costa*, a *costodiendo*; because they guard the vital viscera.) The ribs are the long curved bones which are placed in an oblique direction at the sides of the chest. Their number is generally twelve on each side; but, in some subjects, it has been found to be thirteen, and in others, though more rarely, only eleven. They are distinguished into true and false ribs. The seven upper ribs, which are articulated to the sternum, are called *true ribs*; and the five lower ones, which are not immediately attached to that bone, are called *false ribs*. At the posterior extremity of each rib we observe a small head, divided by a middle ridge into two articulating surfaces, covered with cartilage, which are received into two cavities contiguous to each other, and formed in the upper and lower part of each dorsal vertebra, as we have already observed in our description of the spine. This articulation, which is secured by a capsular ligament, is a species of *ginglimus*, and allows only of motion upwards and downwards. The head of each rib is supported by a short neck, and immediately beyond this we find a flattened tubercle, affording an oblong and slightly convex surface, which is articulated with the transverse process of the lowest of the two dorsal vertebrae, with which its head is articulated. At some little distance from this tuberosity, the rib makes a considerable curve, which is usually called its angle. From the tubercle to the angle the ribs are of considerable thickness, and approaching to a cylindrical shape; but, from the angle to their anterior extremity, they become thinner and flatter. To this anterior extremity is fixed a long, broad, and strong cartilage, which, in each of the true ribs, reaches to the

sternum, where its articulation is secured by a capsular ligament, and by other ligamentous fibres. The cartilages of the sixth and seventh ribs being longer than the rest, are extended upwards, in order to reach the sternum, the inferior portion of which is about on a level with the fifth rib. The cartilages of these two ribs are usually united into one, so as to leave no space between them. The false ribs are supported in a different manner; their cartilages terminate in an acute point before they reach the sternum, the eighth rib being attached by its cartilage to the lower edge of the cartilage of the seventh, or last of the true ribs; the ninth in the same manner to the eighth; and the tenth to the ninth; the cartilages of each rib being shorter than that of the rib above it. The eleventh and twelfth, which are the two lowermost ribs, are not fixed at their anterior extremities like the other ribs, but hang loose, and are supported only by their ligamentous fibres, and by muscles and other soft parts.

The external surface of each rib is somewhat convex, and its internal surface slightly concave. On the inferior and interior surface of these bones we observe a long fossa, for the lodgment of the intercostal vessels and nerves. This channel, however, does not extend through the whole length of the rib, being observable neither at the posterior extremity, where the vessels have not yet reached the bone, nor at the fore end, where they are distributed to the parts between the ribs. We seldom see any marks of it in the short ribs, as in the first, second, eleventh, and twelfth.

Thus far we have given a description, which is applicable to the ribs in general; but, as we find them differing from each other in shape, length, situation, and other respects, it will be right to speak of each rib in particular.

The *first* rib, which is the shortest of any, is likewise the most curved. It is broader than the other ribs, and, instead of being placed, as they are, obliquely, and with its edges upwards and downwards, it is situated nearly in a transverse direction, one of its edges being placed inwards, or nearly so. Of these edges, the inner one is sharp, and the outer one somewhat rounded. Its inner surface is smooth, and its superior surface is sometimes slightly depressed anteriorly by the clavicle. The head of this rib, instead of being angular, is flattened, and slightly convex, being received into a cavity, which is formed wholly in the first vertebra, and not by two vertebrae, as is the case with the other ribs.

The *second* rib is longer than the first, but shorter than the ribs below it. Its angle is placed at a small distance from its tuberosity, and its head is articulated with two vertebrae, like the other ribs. The

other ten tribes, the two last only excepted, differ from the general description we have given, chiefly in the difference of their length, which goes on gradually increasing, from the first or uppermost, to the seventh or last of the true ribs, and as gradually diminishing from that to the twelfth. Their obliquity, in respect to the spine, likewise increases as they descend, as does the distance between the head and angle of each rib, from the first rib to the ninth. The two lowest ribs differ from all the rest in the following particulars:—Their heads, like that of the first rib, are rounded, and received into a cavity formed entirely in the body of one vertebra; they have no tubercle for their articulation with the transverse processes, to which they are only loosely fixed by ligaments, and, in this respect, the tenth rib is sometimes found to agree with them: they are much shorter than the rest of the false ribs, and the twelfth is still shorter than the eleventh. The length of the latter, however, is different in different subjects, and is not always found to be the same on both sides in the same skeleton. Anteriorly, as we have already observed, their cartilages are short and loose, not being attached to the cartilages of the other ribs; and this seems to be, because the most considerable motions of the trunk are not performed on the lumbar vertebrae alone, but likewise on the lower vertebrae of the back; so that if these two ribs had been confined anteriorly, like the rest, and likewise united to the bodies of two vertebrae, and to the transverse process, this disposition would have impeded the motion of the two last vertebrae of the back, and consequently would have affected the motion of the trunk in general.

The use of the ribs is to give form to the thorax, and to cover and defend the lungs; also assist in breathing; for they are joined to the vertebrae by regular hinges, which allow of short motions, and to the sternum by cartilages, which yield to the motion of the ribs, and return again when the muscles cease to act.

Ribwort. The *Plantago lanceolata* of Linnaeus, or narrow-leaved plantain, is sometimes so termed.

Rice. See *Oryza*.

RICINUS. (*Quasi ex aspe,* a dog's nose; because they stick to the noses of dogs.) 1. The name of a genus of plants in the Linnaean system. Class, *Monocotyledon*. Order, *Monadelphica*.

2. The pharmacopœial name of the plant which affords the seed from which the castor-oil is prepared. *Casaputia major*. *Bicinus vulgaris*. *Palma christi*. *Ricinus communis* of Linnaeus:—*foliis peltatis sub-palmatis serratis*. This plant appears to be the *Kari*, or *Egeron*, of Dioscorides, who observes, that the seeds are powerfully cathartic; it is also mentioned by *Etieri*,

Paulus Aegineta, and **Pliny**. The ricinus was first cultivated in England, in the time of **Turner**, and is now annually reared in many gardens in the neighbourhood of London; and in that of **Dr. Saunders**, at **Highbury**, the plant grew to a state of great perfection. An oil extracted from the seeds of this plant, and known by the name of *oleum ricini*, *palma christi*, or *castor oil*, is the drug to which the pharmacopœias refer, and which has lately come into frequent use, as a quick but gentle purgative.

The London College directs this oil to be expressed from the seeds in the same way as that of the oil of almonds, and without the assistance of heat, by which the oil would seem to be obtained in the purest state. However, we have some reason to believe that this method is seldom practised, and that the oil usually employed here is imported from the West Indies, where it is commonly prepared in the following manner:—"The seeds being freed from the husks, or perds, which are gathered upon their turning brown, and when beginning to burst open, are first bruised in a mortar, afterwards tied up in a linen bag, and then thrown into a large pot, with a sufficient quantity of water (about eight gallons to one gallon of the seeds,) and boiled till the oil is risen to the surface, when it is carefully skimmed off, strained, and kept for use. Thus prepared, the oil is entirely free from acrimony, and will stay upon the stomach when it rejects all other medicines." **Mr. Long** remarks, that the oil intended for medicinal use, is more frequently cold drawn, or extracted from the bruised seeds by means of a hand-press. But this is thought more acrimonious than that prepared by coction. **Dr. Browne** is also of this opinion, and prefers the oil prepared by coction to that by expression; he attributes its greater mildness to the action of the fire, observing, that the expressed oil, as well as the mixed juices of the seeds, are far more active and violent in their operation.

Dr. Cullen observes, that "this oil, when the stomach can be reconciled to it, is one of the most agreeable purgatives we can employ. It has this particular advantage, that it operates sooner after its exhibition than any other purgative I know of, as it commonly operates in two or three hours. It seldom gives any griping, and its operation is generally moderate, to one, two, or three stools only. It is particularly suited to cases of costiveness, and even to cases of spasmodic cholera.

In the West Indies, it is found to be one of the most certain remedies in the dry belly-ache, or colica pistonum. It is seldom found heating or irritating to the rectum; and therefore is sufficiently well suited to hemorrhoidal persons.

The only inconvenience attending the use of this medicine is, that as an oil it is nauseous to some persons; and that, when the dose is large, it occasions sickness at the stomach for some time after it is taken. To obviate these inconveniences, several means have been tried; and it is found that the most effectual means is the addition of a little ardent spirit. In the West Indies, they employ rum; but that I might not withdraw any part of the purgative, I employ the *Tinct. senna comp.* This added, in the proportion of one to three parts of the oil, and very intimately mixed, by being shaken together in a phial, both makes the oil less nauseous to the taste, and makes it sit more easy on the stomach. The common dose of this oil is a tablespoonful, or half an ounce; but many persons require a double quantity.

RICINUS COMMUNIS. The systematic name of the plant which affords the castor-oil. See *Ricinus*.

RICINUS MAJOR. Ricinoides. Pinus purgans. Pinhenes indici. Faba cathartica. Nux cathartica americana. Nux barbadensis. An oblong black seed, the produce of the *Jatropha curcas*; *foliis cordatis angulatis* of **Linnaeus**. It affords a quantity of oil, which is given, in many places, as the castor-oil is in this country, to which it is very nearly allied. The seeds of the *Jatropha multifida* are of an oval and triangular shape, of a pale brown colour, are called purging-nuts, and give out a similar oil.

RICINUS VULGARIS. See *Ricinus*.

Rickets. See *Rachitis*.

RIGOR. A coldness, attended by a shivering, more or less perfect.

RIMA. A fissure, or opening; as the *rima laryngis*, *rima vulvae*.

RIMA GLOTTIDIS. The opening of the larynx, through which the air passes in and out of the lungs.

RINCLA. (Dim. of *rima*, a fissure.) A small fissure.

RINXUS. (From *gr*, the nose.) See *Compressor naris*.

Ring-worm. A species of herpes. See *Herpes*.

RINAGON. See *Cassumunar*.

RISUS SARDONICUS. See *Sardonic laugh*.

ROASTING. A chemical process, generally performed in crucibles, by which mineral substances are divided, some of their principals being volatilized and others changed, so as to prepare them for other operations.

ROB. (*Rob*, dense, Arabian.) An old term for an inspissated juice.

ROBORANTS. (*Medicamenta roborantia*; from *robore*, to strengthen.) Strengthening medicines. See *Stimulants*.

ROCCOLLA. The principal use of this plant, *Lichan roccolla* of **Linnaeus**, is as a

blue dye. It has been employed medicinally with success in allaying the cough attendant on phthisis, and in hysterical coughs.

Rochelle salt. See *Soda tartarizata*.

ROCKAMOLE. The *Allium scorodophrastrum* of Linnaeus. The root is used for pickles and high-seasoned dishes.

Rock-oil. See *Petroleum*.

Rock samphire. See *Samphire*.

Rocket, garden. See *Eruca*.

Rocket, Roman. See *Eruca*.

Rocket, wild. See *Eruca sylvestris*.

RORELLA. See *Ros solis*.

ROS CALABRIVUS. The officinal manna is sometimes so termed.

ROS SOLIS. (*Ros, dew*.) *Rorella*. Sundew. This elegant little plant, *Drosera rotundifolia* of Linnaeus:—*capis radicata; foliis orbiculatis*, is said to be so acrid as to ulcerate the skin, and remove warts and corns, and to excite a fatal coughing and delirium in sheep who eat it. It is seldom given medicinally in this country but by the lower orders, who esteem a decoction of it as serviceable in asthmas and coughs.

ROSA. 1. The name of a genus of plants in the Linnæan system. Class, *Iceandria*. Order, *Polygynia*. The rose.

2. A name sometimes given to the erysipelas, because it begins with a redness like that of a rose.

ROSA ALBA. The white rose. The flowers of this species possess similar but inferior virtues to those of the damask. They are directed in some officinal preparations.

ROSA CANINA. *Rosa sylvestris*. The pharmacopœial and systematic name of the dog-rose. See *Cynobatus*.

ROSA CENTIFOLIA. The pharmacopœial and systematic name of the damask rose. *Rosa damascena*. *Rosa pallida*. The damask rose. The pharmacopœias direct a syrup to be prepared from the petals of this rose. *Rosa centifolia* of Linnaeus:—*germinibus ovatis pedunculisque hispidis, caule hispido aculeato, petiolis inermibus*, which is found to be a pleasant and useful laxative for children, or to obviate costiveness in adults. Most of the roses, though much cultivated in our gardens, are far from being distinctly characterised. Those denominated varieties are extremely numerous, and often permanently uniform; and the specific differences as hitherto pointed out, are in many respects so adequate to the purpose of satisfactory discrimination, that it becomes a difficult matter to distinguish which are species and which are varieties only. The London College, following Gerard and Parkinson, has still retained the name *rosa damascena*; but the damask rose is another species, widely different from the centifolia, as appears from the description given of it by Du Roi and Miller.

The petals are directed for medicinal use; they are of a pale red colour, and of a very fragrant odour, which, to most people, is extremely agreeable; and therefore this and most of the other roses are much used as nosegays. We may remark, however, that, in some instances, they have, under certain circumstances, produced alarming symptoms. The petals "impart their odorous matter to watery liquors, both by infusion and distillation. Six pounds of fresh roses impregnate, by distillation, a gallon, or more, of water, strongly with their fine flavour. On distilling large quantities, there separates from the watery fluid a small portion of a fragrant butyraceous oil, which liquifies by heat, and appears yellow, but concretes in the cold into a white mass. An hundred pounds of the flowers, according to the experiments of Tachenius and Hoffman, afforded scarcely half an ounce of oil." The smell of the oil exactly resembles that of roses, and is therefore much used as a perfume. It possesses very little pungency, and has been highly recommended for its cordial and analeptic qualities. These flowers also contain a bitterish substance, which is extracted by water along with the odorous principle, and remains entire in the decoction after the latter has been separated by distillation, or evaporation.

This fixed sapid matter of the petals manifests a purgative quality; and it is on this account that the flowers are received in the *Materia Medica*.

ROSA DAMASCENA. See *Rosa centifolia*.

ROSA GALLICA. The pharmacopœial and systematic name of the red rose. *Rosa rubra*. The flowers of this species, *Rosa gallica* of Linnaeus:—*germinibus ovatis pedunculisque hispidis, caule petiolisque hispido aculeatis*, are valued for their adstringent qualities, which are most considerable before the petals expand; and therefore in this state they are chosen for medicinal use, and ordered by the pharmacopœias in different preparations, as those of a conserve, or confection, a honey, an infusion, and a syrup. The infusion of roses is a grateful cooling subadstringent, and useful in hæmoptysis, and other hæmorrhagic complaints; its efficacy, however, depends chiefly on the acid.

ROSA PALLIDA. See *Rosa centifolia*.

ROSA RUBRA. See *Rosa gallica*.

ROSA SYLVESTRIS. The dog-rose, or *Rosa canina* of Linnaeus. See *Cynobatus*.

ROSACEA. The term *gutta rosacea* is applied to little rosy-coloured spots upon the face and nose.

Rosa. See *Rosa*.

Rose, damask. See *Rosa centifolia*.

Rose, dog. See *Cynobatus et Rosa canina*.

Rosa radix. See *Rhodia*.

Rose, red. See *Rosa gallica*.

Rose root. See *Rhodia*.

Rose, white. See *Rosa alba*.

ROSEBAY WILLOW HERB. This is the *Epilobium angustifolium* of Linnæus, common in our woods, in moist situations. The young shoots are said to be little inferior to asparagus, when boiled.

Rosemary. See *Rosmarinus*.

Rosewood. See *Rhodium lignum*.

Rosewort. See *Rhodia*.

Rosin. See *Resina*.

ROSMARINUS (*Quasi rosa, smugens*, because it smells like myrrh.) 1. The name of a genus of plants in the Linnæan system. Class, *Diandria*. Order, *Monogynia*.

2. The pharmacopœial name of the common rosemary. *Rosmarinus hortensis. Libanotis coronaria. Dendrobibanus. Rosmarinus officinalis* of Linnæus. The leaves and tops of this plant have a fragrant aromatic smell, and a bitterish pungent taste. Rosemary is reckoned one of the most powerful of those plants which stimulate and corroborate the nervous system; it has therefore been recommended in various affections supposed to proceed from debility, or defective excitement of the brain and nerves, as in certain head-aches, deafness, giddiness, and in some hysterical and dyspeptic symptoms. The official preparations of rosemary are, an essential oil from their leaves, or from the herb in flower, a conserve of the flowers, and a spirit formerly called Hungarian water, from the flowery tops. The tops are also used in the compound spirit of lavender, and soap liniment.

ROSMARINUS HORTENSIS. See *Rosmarinus*.

ROSMARINUS OFFICINALIS. The systematic name of the common rosemary. See *Rosmarinus*.

ROSMARINUS SYLVESTRIS. The plant which bears this name in the pharmacopœias, is the *Ledum palustre* of Linnæus. It has a bitter subadstringent taste, and was formerly used in Switzerland in the place of hops. Its medicinal use is confined to the continent, where it is occasionally given in the cure of hooping-cough, sore throat, dysentery, and exanthematous diseases.

ROSTRUM. (From *rodo*, to gnaw; because birds use it to tear their food with.) A beak. The piece of flesh which hangs between the division of the hare-lip is called *rostrum leporinum*.

Rotang cane. See *Sanguis draconis*.

ROTATOR. (From *roto*, to turn.) A muscle whose office is to wheel about the thigh.

ROTULA. (Dim. of *rota*, a wheel; so called from its shape.) See *Patella*.

Round-leaved sorrel. See *Rumex scutellus*.

ROUND LIGAMENTS. *Ligamenta rotunda.* A bundle of vessels and fibres

contained in a duplicature of the peritonæum, that proceed from the sides of the uterus, through the abdominal ring, and disappear in the pudenda.

RUBEDO. (From *ruber*, red.) A diffused, but not spotted, redness in any part of the skin; such as that which arises from blushing.

RUBEFACIENTS. (*Rubefaciendia*; from *rubefacio*, to make red.) Those substances which, when applied a certain time to the skin, induce a redness without blistering.

RUBEOLA. (From *ruber*, red; from *rubio*, to become red.) *Morbilli.* The measles. A genus of disease in the class *pyrexie*, and order *exanthematosa* of Cullen; known by synocha, hoarseness, dry cough, sneezing, drowsiness; about the fourth day, eruption of small red points, discernible by the touch, which, after three days, end in mealy desquamation. The blood, after venesection, exhibits an inflammatory crust. In addition to the symptoms already related, it is remarkable, that the eyes and eye-lids always shew the presence of this disease, being somewhat inflamed and suffused with tears. The synocha continues during the whole progress of the disease. In systems of nosology, several varieties of measles are mentioned, but they may all be comprehended under two heads; the one attended with more or less of the symptoms of general inflammation; the other accompanied by a paratrid diathesis.

The measles may prevail at all seasons of the year as an epidemic, but the middle of winter is the time they are usually most prevalent; and they attack persons of all ages, but children are most liable to them. They prove most unfavourable to such as are of a plethoric or scrophulous habit. Like the small-pox, they never affect persons but once in their life; their contagion appears to be of a specific nature. The eruption is usually preceded by a general uneasiness, chilliness and shivering, pain in the head, in grown persons; but in children, a heaviness and soreness in the throat, fever, sickness, and vomiting, as happen in most fevers; but the chief characteristic symptoms are, a heaviness about the eyes, with swelling, inflammation, and a defluxion of sharp tears, and great acuteness of sensation, so that they cannot bear the light without pain, together with a discharge of serous humour from the nostrils, which produces sneezing. The heat, and other febrile symptoms, increase very rapidly; to which succeeds a frequent and dry cough, a stuffing, great oppression, and oftentimes retching to vomit, with violent pains in the loins, and sometimes a looseness; at other times there is great sweating, the tongue foul and white, the thirst very great, and, in general, the fever runs

much higher than in the milder sort of the regular small-pox. The eruptions appear about the fourth or fifth day, and sometimes about the end of the third. On the third or fourth day from their first appearance, the redness diminishes, the spots, or very small pustules, dry up, the cuticle peels off, and is replaced by a new one. The symptoms do not go off on the eruption, as in the small-pox, except the vomiting; the cough and head-ache continue, with the weakness and defluxion on the eyes, and a considerable degree of fever.

On the ninth or eleventh day, no trace of redness is to be found, but the skin assumes its wonted appearance; yet, without there have been some considerable evacuations either by the skin, or by vomiting, the patient will recover strength, but the cough will continue, the fever return with new violence, and bring on great distress and danger.

In the more alarming cases, spasms of the limbs, subsultus tendinum, delirium, or what more frequently happens, coma supervene. This last symptom so frequently attends the eruptive fever of measles, that by some practitioners it is regarded as one of its diagnostics.

In measles, as in other febrile diseases, the symptoms generally suffer some remission towards the morning, returning however in the evening with increased severity.

The measles, even when violent, are not usually attended with a putrid tendency; but it sometimes happens, that such a disposition prevails both in the course of the disease and at its termination.

In such cases petechiæ are to be observed interspersed among the eruptions, and these last become livid, or assume almost a black colour. Hemorrhages break out from different parts of the body, the pulse becomes frequent, feeble, and perhaps irregular, universal debility ensues, and the patient is destroyed.

In those cases where there is much fever, with great difficulty of breathing, and other symptoms of pneumonic inflammation, or where there is great debility with a tendency to putrescency, there will always be considerable danger; but the consequences attendant on the measles are in general more to be dreaded than the immediate disease; for although a person may get through it, and appear for a time to be recovered, still hectic symptoms and pulmonary consumption shall afterwards arise, and destroy him, or an ophthalmia shall ensue.

Measles, as well as small-pox, not unfrequently call into action a disposition to scrophula, where such happens to exist in the habit.

Another bad consequence of the measles is, that the bowels are often left by them in a very weak state; a chronic diarrhœa

remaining which has sometimes proved fatal. Dropsy has also been known as a consequence of measles.

The morbid appearances to be observed on dissections of those who die of measles are pretty much confined to the lungs and intestines: the former of which always shew strong marks of inflammation, with sometimes a tendency to sphacelus.

Where the patient dies under the eruption, the trachea and larger branches of the bronchiæ, as in the small-pox, are often covered with it, which Dr. Thomas observes may account for the increase of the cough, after the appearance of the irruption.

RUBIA. (From *ruber*, red; so called from its red roots.) 1. The name of a genus of plants in the Linnæan system. Class, *Tetrandria*. Order, *Monogynia*.

2. The pharmacopœial name of what is also called *Erythrodanum*. *Rubia major*. *Radix rubra*. Dyer's madder. *Rubia tinctorum* of Linnæus:—*foliis annuis, caule aculeato*. The roots of this plant have a bitterish, somewhat austere taste, and a slight smell, not of the agreeable kind. It was formerly considered as a deobstruent, detergent, and diuretic, but it is now very seldom used.

RUBIA TINCTORUM. The systematic name of the madder-plant. See *Rubia*.

RUBICO CURRI. See *Verdigris*.

RUBICO FERRI. (*Rubigo a colore rubro*, from its reddish colour.) Rust of iron. See *Carbonas ferri*.

RUBINUS. (From *ruber*, red, so named from its colour.) A carbuncle.

RUBUS. (From *ruber*, red, so called from its red fruit.) The name of a genus of plants in the Linnæan system. Class, *Icosandria*. Order, *Polygynia*.

RUBUS ARCTICUS. The systematic name of the shrubby strawberry. See *Bacca Norlandicæ*.

RUBUS CASSIUS. The systematic name of the dewberry plant, whose fruit resembles the blackberry in appearance and qualities.

RUBUS CHAMÆMORUS. The systematic name of the clodberry-tree. See *Chamæmorus*.

RUBUS FRUCTICOSUS. The systematic name of the common bramble, which affords blackberries. See *Blackberry*.

RUBUS IDEUS. *Batinn.* *Moron.* *Rubus ideus* of Linnæus:—*foliis quinatopinnatis ternatisque, caule aculeato, petiolis canaliculatis*. The raspberry. The fruit of this plant has a pleasant sweet taste, accompanied with a peculiar grateful flavour, on account of which it is chiefly valued. Its virtues consist in allaying heat and thirst, and promoting the natural excretions. A grateful syrup prepared from the juice is directed for officinal use.

RUCTUS. An eructation.

Rue. See *Ruta*.

Rue, goats. See *Galiga*.

RUSI PILULÆ. Rufus's pills. A compound very similar to the aloetic pills with myrrh.

RUM. A spirituous liquor, well known, the produce of the sugar-cane.

RUMEX. The name of a genus of plants in the Linnæan system. Class, *Hexandria*. Order, *Trigynia*. The dock.

RUMEX ACETOSA. The systematic name of the common sorrel. See *Acetosa*.

RUMEX ACUTUS. The systematic name of the sharp-pointed wild-dock. See *Oxylapathum*.

RUMEX ALPINUS. The systematic name of the plant which affords the monks' rhubarb. See *Rhabarbarum monachorum*.

RUMEX AQUATICUS. The water-dock. See *Hydrolapathum*.

RUMEX CRISPUS. The systematic name of the crisp-leaved dock. See *Oxylapathum*.

RUMEX HYDROLAPATHUM. The systematic name of the water-dock. See *Hydrolapathum*.

RUMEX PATIENTIA. The systematic name of the garden patience. See *Rhabarbarum monachorum*.

RUMEX SANGUINEUS. The systematic name of the bloody dock, whose root has an austere and adstringent taste, and is sometimes given by the vulgar in the cure of dysentery.

RUMEX SCUTATUS. The systematic name of the French sorrel, sometimes called *acetosa rotundifolia* in the shops. Its virtues are similar to those of common sorrel. See *Acetosa*.

RUPELLENSIS SAL. (From *rupellum Rochelle*, where it was first made by Dr. Seignette.) A term applied to Rochelle salt. Now called soda tartarizata.

RUPURA. See *Hernia*.

Rupture. See *Hernia*.

Rupture wort. See *Herniaria*.

RUSCUS. (*A russo colore*, from the carnation colour of its berries.) 1. The name of a genus of plants in the Linnæan system. Class, *Dioecia*. Order, *Syngenesia*.

2. The pharmacopœial name of the butchers' broom, or knee-holly. *Bruscus. Oxymyrrhina. Oxomyrrine. Myrtacantha. Myncantha. Scopæ regia.* Wild myrtle. A small evergreen shrub, the *ruscus aculeatus* of Linnæus. *Folia supra floriferis nudis.* It grows in woods and thickets in this country. The root, which is somewhat thick, knotty, and furnished with long fibres, externally brown, internally white, and of a bitterish taste, has been recommended as an aperient and diuretic in dropsies, urinary obstructions, and nephritic cases. It is seldom used in this country.

RUSCUS ACULEATUS. The systematic name of butchers' broom. See *Ruscus*.

RUSCUS HYPOGLOSSUS. The systematic name of the uvularia. See *Uvularia*.

RUSK-RUT. The root of the *Cyperus esculentus* of Linnæus, a native of Italy, where it is collected and eaten, being more delicately and pleasantly tasted than our chesnuts.

Rush, sweet. See *Juncus odoratus*.

Russia ashes. The impure potash, as imported from *Russia*.

RUTA. (From *pus*, to preserve, because it preserves health.) 1. The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Monogynia*.

2. The pharmacopœial name of the common rue. *Ruta graveolens* of Linnæus:—*foliis decompositis, floribus lateralibus quadrifidis.* Rue has a strong and graceful smell, and a bitter, hot, penetrating taste; the leaves are so acrid, that by much handling they have been known to irritate and inflame the skin; and the plant, in its natural or uncultivated state, is said to possess these sensible qualities still more powerfully. The imaginary quality of the rue, in resisting and expelling contagion, is now laid aside. It is doubtless a powerful stimulant, and is considered, like other medicines of the fetid kind, as possessing attenuating, deobstruent, and antispasmodic powers. In the former London Pharmacopœia it was directed in the form of an extract; and was also an ingredient in the *putris e myrrha comp.*, but are now omitted. The dose of the leaves is from 15 grains to two scruples.

RUTA GRAVEOLENS. The systematic name of the common rue. See *Ruta*.

RUTA MURARIA. The plant to which this name is given in the Pharmacopœias, is the *Asplenium ruta muraria* of Linnæus. It is supposed by some to possess specific virtues in the cure of ulcers of the lungs, and is exhibited in the form of decoction.

RUTIDOSIS. A corrugation and subsiding of the cornea of the eye. The species are, 1. *Rutidosia*, from a wound or puncture penetrating the cornea. 2. *Rutidosia*, from a fistula penetrating the cornea. 3. *Rutidosia*, from a deficiency of the aqueous humour, which happens from old age, fevers, great and continued evacuations, and in extreme dryness of the air. 4. *Rutidosia* of dead persons, when the aqueous humour exhalles through the cornea, and no fresh humour is secreted; so that the cornea becomes obscure and collapsed; this is a most certain sign of death.

RUTULA. (From *ruta*, rue.) A small species of rue.

RUYSCHIANA TUNICA. The internal surface of the choroid membrane of the human eye, which this celebrated anatomist imagined was a distinct lamina from the external surface.

RYAS. See *Rhas*.

RYE, COMMON. A very common bread-

corn among the northern parts of Europe; it is less nourishing than wheat, but a sufficiently nutritive and wholesome grain. It is more than any other grain strongly

disposed to accegency; hence it is liable to ferment in the stomach, and to produce purging, which people on the first using it commonly experience.

S

S. A. The contraction of *secundum artem*.

S, or *ss.* Immediately following any quantity, imports *semis*, or half.

SABADILLA. See *Cavadilla*.

SABINA. (Named from the Sabines, whose priests used it in their religious ceremonies.) *Savina. Subina. Sterilis. Brastus.* Common or barren Savin. *Juniperus sabina* of Linnaeus:—*foliis oppositis erectis decurrentibus, oppositionibus pyxidalis.* Savin is a native of the south of Europe and the Levant; it has long been cultivated in our gardens, and from producing male and female flowers on separate plants it was formerly distinguished into the barren and berry-bearing savin. The leaves and tops of this plant have a moderately strong smell of the disagreeable kind, and a hot, bitterish, acrid taste. They give out great part of their active matter to watery liquors, and the whole to rectified spirit. Distilled with water they yield a large quantity of essential oil. Decoctions of the leaves, freed from the volatile principle by inspissation to the consistence of an extract, retain a considerable share of their pungency and warmth along with their bitterness, and have some degree of smell, but not resembling that of the plant itself. On inspissating the spirituous tincture, there remains an extract consisting of two distinct substances, of which one is yellow, unctuous, or oily, bitterish, and very pungent; the other black, resinous, less pungent, and sub-adstringent. Savin is a powerful and active medicine, and has been long reputed the most efficacious in the materia medica, for producing a determination to the uterus, and thereby proving emmenagogue; it heats and stimulates the whole system very considerably, and is said to promote the fluid secretions. The power which this plant possesses (observes Dr. Woodville) in opening uterine obstructions, is considered to be so great, that we are told it has been frequently employed, and with too much success, for purposes the most infamous and unnatural. It seems probable, however, that its effects in this way have been somewhat overrated, as it is found, very frequently, to fail as an emmenagogue, though this, in some measure, may be ascribed to the smallness of the dose in which it has been usually prescribed by physicians; for Dr. Cullen observes, "that savin is a very

acrid and heating substance, and I have been often, on account of these qualities, prevented from employing it in the quantity necessary to render it emmenagogue. I must own, however, that it shews a more powerful determination to the uterus than any other plant I have employed; but I have been frequently disappointed in this, and its heating qualities always require a great deal of caution." Dr. Home appears to have had very great success with this medicine, for in five cases of amenorrhoea which occurred at the Royal Infirmary at Edinburgh, four were cured by the sabina, which he gave in powder from a scruple to a drachm twice a day. He says it is well suited to the debile, but improper in plethoric habits, and therefore orders repeated bleedings before its exhibition. Country people give the juice from the leaves and young tops of savin mixed with milk to their children, in order to destroy the worms; it generally operates by stool, and brings them away with it. The leaves cut small, and given to horses, mixed with their corn, destroy the bots. Externally savine is recommended as an escharotic to foul ulcers, syphilitic warts, &c. A strong decoction of the plant in lard and wax forms an useful ointment to keep up a constant discharge from blisters, &c. See *Ceratium Sabina*.

SABULOUS. (*Sabulosus.* Gritty, sandy.) A term often applied to the calcareous matter in urine.

SABURRA. Dirt, sordes, filth. Foulness of the stomach, of which authors mention several kinds, as the acid, the bitter, the empyreumatic, the insipid, the putrid.

SACCATUM DROPR. Ascites saccatus. See *Ascites*.

SACCHARI ACIDUM. Acid of sugar. If one part of sugar be distilled with three parts of nitric acid, till nitrous gas ceases to be developed, and then re-distilled with three parts of the same acid, a white crystallized salt is found in the liquid residue, which is the acid of sugar.

SACCHARUM. (*Σακχαρος*, from *sacchar*, Arab.) The Arabians call it *suchar*, *succhar*, *sutter*, *suchar*, *zucuro*, and *zozar*; the Greeks called it *sackohar*, *salchalion*, and *spodium*. Sugar. The cane from which the sugar is obtained in the West and East Indies, is the *Arundo saccharifera* of Linnaeus:—*floribus paniculatis*,

foliis plamis. It is prepared from the expressed juice boiled with the addition of quick lime or common vegetable alkali. It may be extracted also from a number of plants, as the maple, birch, wheat, corn, beet-root, skirret, parsnips, and dried grapes, &c. by digesting in alcohol. The alcohol dissolves the sugar, and leaves the extractive matter untouched, which falls to the bottom. It may be taken into the stomach in very large quantities, without producing any bad consequences, although proofs are not wanting of its mischievous effects, by relaxing the stomach, and thus inducing disease. It is much used in pharmacy, as it forms the basis of syrups, lozenges, and other preparations. It is very useful as a medicine, although it cannot be considered to possess much power, to favour the solution or suspension of resins, oils, &c. in water, and is given as a purgative for infants. Dr. Cullen classes it with the attenuantia, and Bergius states it to be saponacea, edulcorans, relaxans, pectoralis, vulneraria, antiseptica, nutrienda. In catarrhal affections both sugar and honey are frequently employed: it has also been advantageously used in calculous complaints; and from its known power in preserving animal and vegetable substances from putrefaction, it has been given with a view to its antiseptic effects. The candy, by dissolving slowly in the mouth is well suited to relieve tickling coughs and hoarseness. Sugar is every where the basis of that which is called sweetness. Its presence is previously necessary in order to the taking place of vinous fermentation. Its extraction from plants which afford it in the greatest abundance, and its refinement for the common uses of life, in a pure and separate state, are among the most important of the chemical manufactures.

The following is the mode of its manufacture in the West Indies: The plants are cultivated in rows, on fields enriched by such manures as can most easily be procured, and tilled with the plough. They are annually cut. The cuttings are carried to the mill. They are cut into short pieces, and arranged in small bundles. The mill is wrought by water, wind, or cattle. The parts which act on the canes are upright cylinders. Between these the canes are inserted, compressed, squeezed till all their juice is obtained from them, and are themselves sometimes, even reduced to powder. One of these mills of the best construction, bruises canes to such a quantity as to afford, in one day, 10,000 gallons of juice, when wrought with only ten mules. The expressed juice is received into a leaden bed. It is thence conveyed into a vessel called the receiver. The juice is found to consist of eight parts of pure water, one part of sugar, one part

of oil and gummy mucilage. From the greener parts of the canes there is apt to be at times derived an acid juice, which tends to bring the whole unseasonably into a state of acid fermentation. Fragments of the ligneous part of the cane, some portions of mud or dirt which unavoidably remain on the canes, and a blackish substance called the crust, which coated the canes at the joints, are also apt to enter into contaminating mixture with the juice. From the receiver the juice is conducted along a wooden gutter lined with lead, to the boiling-house. In the boiling-house it is received into copper pans or caldrons, which have the name of clarifiers. Of these clarifiers the number and the capacity must be in proportion to the quantity of canes, and the extent of the sugar plantation on which the work is carried on. Each clarifier has a syphon or cock, by which the liquor is to be drawn off. Each hangs over a separate fire; and this fire must be so confined, that by the drawing of an iron slider fitted to the chimney, the fire may be at any time put out. In the progress of the operations, the stream of juice from the receiver fills the clarifier with fresh liquor. Lime in powder is added in order to take up the oxalic acid, and the carbonaceous matters which are mingled with the juice. The lime also in the new salts, into the composition of which it now enters, adds itself to the sugar, as a part of that which is to be obtained from the process. The lime is to be put in in the proportion of somewhat less than a pint of lime to every hundred gallons of liquor. When it is in too great quantities, however, it is apt to destroy a part of the pure saccharine matter. Some persons employ alkaline ashes, as preferable to lime, for the purpose of extracting the extraneous matter; but it is highly probable, that lime, judiciously used, might answer better than any other substance whatsoever. The liquor is now to be heated almost to ebullition. The heat dissolves the mechanical union, and thus favours the chemical changes in its different parts. When the proper heat appears, from a rising scum on the surface of the liquor, to have been produced, the fire is then extinguished by the application of the damper. In this state of the liquor, the greater part of the impurities, being different in specific gravity from the pure saccharine solution, and being also of such a nature as to yield more readily to the chemical action of heat, are brought up to the surface in a scum. After this scum has been sufficiently formed on the cooling liquor, this liquor is carefully drawn off, either by a syphon, which raises a pure stream through the scum, or by a cock drawing the liquor at the bottom from under the scum. The scum in either case sinks down unbroken, as the liquor flows;

and is now, by cooling, of such tenacity, as not to send to any intermixture with the liquor. The liquor drawn, after this purification from the boiler, is received into a gutter or channel, by which it is conveyed to the grand copper, or evaporating boiler. If made from good canes, and properly clarified, it will now appear almost transparent. In this copper, the liquor is heated to actual ebullition. The scum raised to the surface by the boiling is skimmed off as it rises. The ebullition is continued till there be a considerable diminution in the quantity of the liquor. The liquor now appears nearly of the colour of Madeira wine. It is at last transferred into a second and smaller copper. An addition of lime-water is here made, both to dilute the thickening liquor, to detach the superabundant acid, and to favour the formation of the sugar. If the liquor be now in its proper state, the scum rises in large bubbles, with very little discolouration. The skimming and the evaporation together produce a considerable diminution in the quantity of the liquor. It is then transferred into another smaller boiler. In this last boiler, the evaporation is renewed, and continued till the liquor is brought to that degree of thickness at which it appears fit to be finally cooled. In the cooler, a shallow wooden vessel of considerable length and wideness, commonly of such a size as to contain a hogshead of sugar, the sugar as it cools, granulates, or runs into an imperfect crystallization, by which it is separated from the melasses, a mixed saccharine matter too impure to be capable even of this imperfect crystallization. To determine whether the liquor be fit to be taken from the last boiler to be finally cooled, it is necessary to take out a portion from the boiler, and try separately, whether it does not separate into granulated sugar and melasses. From the cooler the sugar is removed to the curing-house. This is a spacious airy building. It is provided with a capacious cistern for the reception of melasses, and over the cistern is erected a frame of strong joist-work, unfilled and uncovered. Empty hogsheads, open at the head, bored at the bottom with a few holes, and having a stalk of plaintain leaf thrust through each of the holes, while it rises at the same time through the inside of the hogshead, are disposed upon the frames. The mass of the saccharine matter from the coolers is put into these hogsheads. The melasses drip into the cistern through the spongy plaintain stalks in the holes. Within the space of three weeks the melasses are sufficiently drained off, and the sugar remains dry. By this process it is at last brought into the state of what is called muscovado or raw sugar. This is the ge-

neral process in the British West Indies. In this state our West India sugar is imported into Britain. The formation of loaves of white sugar is a subsequent process. In the French West India Isles it has long been customary to perform the last part of this train of processes in a manner somewhat different, and which affords the sugar in a state of greater purity. This preparation, taking the sugar from the cooler, then puts it, not into hogsheads with holes in the bottom as above, but into conical pots, each of which has at its bottom a hole half an inch in diameter, that is, in the commencement of the process, stopped with a plug. After remaining some time in the pot, the sugar becomes perfectly cool and fixed. The sugar is then removed out of the hole; the pot is placed over a large jar, and the melasses are suffered to drip away from it. After as much of the melasses as will easily run off has been thus drained away, the surface of the sugar in the jar is covered with a stratum of fine clay, and water is poured upon the clay. The water oozing gently through the pores of the clay, pervades the whole mass of sugar, redissolves the melasses still remaining in it, with some parts of the sugar itself, and carrying these off by the holes in the bottom of the pot, renders that which resists the solution much purer than the muscovado sugar made in the English way. The sugar prepared in this manner is called clayed sugar. It is sold for a higher price in the European markets than the muscovado sugar; but there is a loss of sugar in the process by claying, which deters the British planters from adopting this practice so generally as do the French.

The raw sugars are still contaminated and debased by a mixture of acid carbonaceous matter, oil, and colouring resin. To free them from these is the business of the European sugar-bakers. A new solution; clarification with alkaline substances fitted to attract away the oil, acid, and other contaminating matters; slow evaporation; and a final cooling in suitable moulds; are the processes which at last produce loaves of white sugar.

The melasses being nothing else but a very impure refuse of the sugar from which they drip, are susceptible of being employed in a new ebullition, by which a second quantity of sugar may be obtained from them. The remainder of the melasses is employed to yield rum by distillation.

In rum, alcohol is mixed with oil, water, oxalic acid, and a mixture of empyreumatic matter. The French prepare, from the mixture of melasses with water, a species of wine of good quality. In its preparation, the solution is brought into

fermentation, then passed through strainers to purify it, then put in casks; after clearing itself in these, transferred into others, in which it is to be preserved for use. The ratio of these processes is extremely beautiful; they are all directed to purify the sugar from contaminating mixtures, and to reduce it into that state of dryness or crystallization, in which it is susceptible of being the most conveniently preserved for agreeable use. The heat in general acts both mechanically to effect a sufficient dissolution of the aggregation of the parts of the cane juice, and chemically to produce in it new combinations into which caloric must enter as an ingredient. The first gentle heat is intended chiefly to operate with the mechanical influence, raising to the surface impurities, which are more easily removed by skimming, than by any other means; a gentle, not a violent heat, is in this instance employed, because a violent heat would produce empyreumatic salts, the production of which is to be carefully avoided. A boiling heat is, in the continuation of the processes, made use of, because, after the first impurities have been skimmed off, contaminating empyreumatic salts are less readily formed, because a boiling heat is necessary to effect a complete development of the saccharine matter, and because the gradual concentration of the sugar is, by such a heat, to be best accomplished. Lime is employed, because it has a stronger affinity than sugar with all the contaminating matters, and particularly because it attracts into a neutral combination that excess of oxalic acid which is apt to exist in the saccharine solution. Skimming removes the new salts which the most easily assume a solid form. The dripping carries away a mixture of water, oil, earth, sugar, from the crystallized sugar: for, in all our crystallizations we can never perform the process in the great way, with such nicety as to preserve it free from an equality of proportions, that must necessarily occasion a residue. Repeated solution, clarification, evaporation, are requisite to produce pure white sugar from the brown and raw sugars; because the complete purification of this matter from acid and colouring matter, is an operation of great difficulty, and not to be finally completed without processes which are longer than can be conveniently performed at the first, upon the sugar plantation. From vegetables of European growth, sugar is not to be easily obtained, unless the process of germination be first produced in them; or unless they have been penetrated by intense frost. Germination, or thorough freezing, develops sugar into all vegetables in which its principles of hydrogen and carbon, with a

small proportion of oxygen, exist in any considerable plenty. It is not improbable, but that if penetration by a freezing cold could be commanded at pleasure, with sufficient cheapness, it would enable us to obtain saccharine matter in a large proportion, from a variety of substances, from which even germination does not yield a sufficient quantity. In the beet, and some other European vegetables, sugar is naturally formed by the functions of vegetation to perfect combination. From these the sugar is obtained by rasping down the vegetable, extracting by water its saccharine juice, evaporating the water charged with the juice to the consistency of syrup, clarifying, purifying, and crystallizing it, just in the same manner as sugar from the sugar-cane.

SACCHARUM ACERNUM. See *Saccharum canadense*.

SACCHARUM ALBUM. White or refined sugar.

SACCHARUM ALUMINIS. Alum mixed with dragon's blood and dried.

SACCHARUM CANADENSE. The sugar obtained from a species of maple-tree, the *Acer pseudo-platanus* of Linnaeus, in Canada, and imported into some parts of Europe. It is supposed to be efficacious in disorders of the breast. Every part of the plant contains a sweet saccharine juice. The trunk, root, or branches, wounded early in the spring, bleed a large quantity of clear liquor, which in its dilute state tastes somewhat sweetish, and being inspissated, yields the concrete sugar, with a syrupy matter resembling melasses. The unboiled juice has been drank as an antiscorbutic. The Canada sugar is much esteemed in France in disorders of the breast.

SACCHARUM CANDIDUM. Sugar candy.

SACCHARUM NON PURIFICATUM. Brown sugar. It is often exhibited as a laxative in clysters, and internally to children.

SACCHARUM OFFICINARUM. The systematic name in some pharmacopœias of the sugar-cane. See *Saccharum*.

SACCHARUM PURIFICATUM. Double refined or loaf sugar. See *Saccharum*.

SACCHARUM SATURNI. See *Plumbi superacetat*.

SACCHO-LACTIC ACID. The sugar of milk in combination with oxygen.

SACCHOLAT. *Saccholat*. A salt formed by the combination of the saccholactic acid with different bases, as saccholat of iron, saccholat of ammonia, &c. &c.

SACCULI ADIPOSI. The bursæ mucosæ of the joints.

SACCULUS. (D.m. of *saccus*, a bag.) A little bag.

SACCULUS CRYLIFERTUS. The receptacle of the chyle.

SACCULUS CORDIS. The pericardium or receptacle of the heart.

SACCULUS LACHRYMALIS. See *Saccus lachrymalis*.

SACCUS LACHRYMALIS. The lachrymal sac is situated in the internal canthus of the eye, behind the lachrymal caruncle, in a cavity formed by the os unguis. It receives the tears from the puncta lachrymalia, and conveys them into the ductus lachrymalis.

SACER. (From *sagur*, secret, Heb.) **Sacred.** Applied to some diseases which were supposed to be immediately inflicted from heaven, as *sacer morbus*, the epilepsy, *sacer ignis*, erysipelas, &c. A bone is called the *os sacrum*, because it was once offered in sacrifices. It also means belonging to the *os sacrum*.

SACK. A wine used by our ancestors, which some have taken to be Rhenish, and others Canary wine. Probably it was what is called dry mountain, or some Spanish wine of that sort. Howell, in his French and English Dictionary, 1650, translates sack by the words *vin d'Espagne*. *Vin sec*.

SACRA HERBA. Common vervain.

SACRA TINCTURA. Ales, canella alba and mountain wine.

SACRAL. Of or belonging to the sacrum; as sacral arteries, veins, nerves, muscles, &c.

SACRO COCCYGEUS. A muscle of the coccyx of the back.

SACRO-LUMBALIS. *Sacro-lumbaris* of authors. *Lumbo-costo-trachelien* of Dumas. A long muscle, thicker and broader below than above, and extending from the *os sacrum* to the lower part of the neck, under the serrati postici, rhomboideus, trapezius, and latissimus dorsi. It arises in common with the longissimus dorsi, tendinous without, and fleshy within, from the posterior part of the *os sacrum*; from the posterior edge of the spine of the ilium; from all the spinous processes, and from near the roots of the transverse processes of the lumbar vertebrae. At the bottom of the back it separates from the longissimus dorsi, with which it had before formed, as it were, only one muscle, and ascending obliquely outwards, gradually diminishes in thickness, and terminates above in a very narrow point. From the place where it quits the longissimus dorsi, to that of its termination, we find it fleshy at its posterior, and tendinous at its anterior edge. This tendinous side sends off as many long and thin tendons as there are ribs. The lowermost of these tendons are broader, thicker, and shorter than those above; they are inserted into the inferior edge of each rib, where it begins to be curved forwards towards the sternum, excepting only the uppermost and last tendon, which

ends in the posterior and inferior part of the transverse process of the last vertebra of the neck. From the upper part of the five, six, seven, eight, nine, ten, or eleven lower ribs, (for the number, though most commonly seven or eight, varies in different subjects,) arise as many thin bundles of fleshy fibres, which, after a very short progress, terminate in the inner side of this muscle, and have been named by Steno, *musculi ad sacro lumbalem accessorii*. Besides these, we find the muscle sending off a fleshy slip from its upper part, which is inserted into the posterior and inferior part of the transverse processes of the five inferior vertebrae of the neck, by as many distinct tendons. This is generally described as a distinct muscle. Diemerbroeck, and Douglas and Albinus after him, call it *cervicalis descendens*. Winslow names it *transversalis collateralis colli*. Morgagni considers it as an appendage to the *sacro lumbalis*. The uses of this muscle are to assist in erecting the trunk of the body, in turning it upon its axis or to one side, and in drawing the ribs downwards. By means of its upper slip, it serves to turn the neck obliquely backwards, or to one side.

SACRO-SCIATIC LIGAMENTS. The ligaments which connect the *ossa innominata* with the *os sacrum*.

SACRUM. (So called from *sacer*, sacred; because it was formerly offered in sacrifices.) *Os sacrum*. *Os basilare*. The *os sacrum* derives its name from its being offered in sacrifice by the ancients, or perhaps from its supporting the organs of generation, which they considered as sacred. In young subjects it is composed of five or six pieces, united by cartilage; but in more advanced age it becomes one bone, in which, however, we may still easily distinguish the marks of the former separation. Its shape has been sometimes compared to an irregular triangle; and sometimes, and perhaps more properly, to a pyramid, flattened before and behind, with its basis placed towards the lumbar vertebrae, and its point terminating in the coccyx. We find it convex behind, and slightly concave before, with its inferior portion bent a little forwards. Its anterior surface is smooth, and affords four, and sometimes five transverse lines, of a colour different from the rest of the bone. These are the remains of the intermediate cartilages by which its several pieces were united in infancy. Its posterior convex surface has several prominences, the most remarkable of which are its spinous processes; these are usually three in number, and gradually become shorter, so that the third is not so long as the second, nor the second as the first. This arrangement enables us to sit with ease. Its transverse processes are formed into one oblong pro-

cess, which becomes gradually smaller as it descends. At the superior part of the bone we observe two oblique processes, of a cylindrical shape, and somewhat concave, which are articulated with the last of the lumbar vertebrae. At the base of each of these oblique processes is a notch, which, with such another in the vertebra above it, forms a passage for the twenty-fourth spinal nerve. In viewing this bone, either before or behind, we observe four, and sometimes five holes on each side, situate at each extremity of the transverse lines which mark the divisions of the bone. Of these holes, the anterior ones, and of these again, the uppermost, are the largest, and afford a passage to the nerves. The posterior holes are smaller, covered with membranes, and destined for the same purpose as the former. Sometimes at the bottom of the bone there is only a notch, and sometimes there is a hole common to it and the os coccygis. The cavity between the body of this bone and its processes, for the lodgment of the spinal marrow, is triangular, and becomes smaller as it descends, till at length it terminates obliquely on each side at the lower part of the bone. Below the third division of the bone, however, the cavity is no longer completely bony, as in the rest of the spine, but is defended posteriorly only by a very strong membrane; hence a wound in this part may be attended with the most dangerous consequences. This bone is articulated above, with the last lumbar vertebra: laterally, it is firmly united, by a broad irregular surface to the ossa innominata, or hip bone: and below it is joined to the os coccygis. In women the os sacrum is usually shorter, broader, and more curved, than in men, by which means the cavity of the pelvis is more enlarged.

Safflower. See *Carthamus*.

Saffron. See *Crocus*.

Saffron, bastard. See *Carthamus*.

Saffron, meadow. See *Colchicum*.

Saffron of steel. A red oxide of iron.

SAGAPENUM. (The name is derived from some eastern dialect.) *Serapium*. It is conjectured that this concrete gummi-resinous juice is the production of an oriental umbelliferous plant. Sagapenum is brought from Persia and Alexandria in large masses, externally yellowish, internally paler, and of an horny clearness. Its taste is hot and biting, its smell of the alliaceous and fetid kind, and its virtues are similar to those which have been ascribed to assafoetida, but weaker, and consequently it is less powerful in its effects.

Sage. See *Salvia*.

Sage of Bethlehem. See *Pulmonaria*.

Sage of Jerusalem. See *Pulmonaria maculosa*.

Sage of virtue. See *Salvia hortensis minor*.

SAGITAL SUTURE. (*Sutura sagittalis*, from *sagitta*, an arrow.) *Sutura virgata, obsoleta, rhabdoides*. The suture which unites the two parietal bones. It has been named *sagittal*, from its lying between the coronal and lambdoidal sutures, as an arrow betwixt the string and the bow.

SAGITTARIA ALEXIPHARMACA. *Malacca radix. Cumna indica. Arundo indica.* Arrow-root; dartwort.

SAGITTARIA SAGITTIFOLIA. (From *sagitta*, an arrow.) The systematic name of the common arrow-head, whose roots are esculent but not very nutritious.

SAGO. *Sagu.* A dry fecula, obtained from the pith of a species of palm, the *Cycas circularis* of Linnæus, in the islands of Java, Molucca, and the Philippines. The same substance is also brought from the West-Indies, but it is inferior to that brought from the East. Sago becomes soft and transparent by boiling in water, and forms a light and agreeable liquid, much recommended in febrile, phthisical, and calculous disorders, &c. To make it palatable, it is customary to add to it, when boiled or softened with water, some lemon juice, sugar and wine.

Sagu. See *Sage*.

Saint Anthony's fire. See *Erysipelas*.

Saint Ignatius's bean. See *Faba indica*.

Saint James's wort. See *Jacobaea*.

Saint John's wort. See *Hypericum*.

Saint Vitus's dance. See *Chorea sancti Viti*.

SAL ABSINTHII. Salt of wormwood. This salt is an imperfect carbonat of potash. See *Potasse subcarbonas*.

SAL ACETOCELLÆ. The salt of wood-sorrel, usually vended for salt of lemons, is an acidulated oxyfat of potash, and called in the new chemical nomenclature *oxylus potasse acidulus*.

SAL ALKALINUS FIXUS. See *Alkali, fixed*.

SAL ALKALINUS VOLATILIS See *Ammonia*.

SAL AMMONIAC. *Murias ammonia*. A saline concrete formed by the combination of the muriatic acid with ammonia. This salt is obtained from several sources. It is found in places adjacent to volcanoes.

1. It appears in the form of an efflorescence, or groups of needles, separate or compacted together, generally of a yellow or red colour, and mixed with arsenic and orpiment; but no use is made of that which is procured in this way.

2. In Egypt it is made in great quantities from the soot of camel's dung, which is burnt at Cairo instead of wood. This

soot is put into large round bottles, a foot and a half in diameter, and terminating in a neck two inches long. The bottles are filled up with this matter to within four inches of the neck. Each bottle holds about forty pounds of soot, and affords nearly six pounds of salt. The vessels are put into a furnace in the form of an oven, so that only the necks appear above. A fire of camel's dung is kindled beneath it, and continued for three days and three nights. On the second and the third day the salt is sublimated. The bottles are then broken, and the salt is taken out in cakes. These cakes, which are sent just as they have been taken out of the bottles in Egypt, are convex, and unequal on the one side; on the middle of this side they exhibit each a tubercle, corresponding to the neck of the bottle in which it was prepared. The lower side is concave, and both are sooty.

3. In this country sal ammoniac is likewise prepared in great quantities. The volatile alkali is obtained from soot, bones, and other substances known to contain it. To this the sulphuric acid is added, and this vitrolic ammoniac is decomposed by muriate of soda or common salt in a double affinity. The liquor obtained in consequence of this decomposition contains sulphate of soda and muriate of ammonia. The first is crystallized, and the second sublimated so as to form cakes, which are then exposed to sale.

Ammoniacal muriate has a poignant, acrid, and urinous taste. Its crystals are in the form of long hexahedral pyramids, a number of them are sometimes united together in an acute angular direction, so as to exhibit the form of feathers. M. Rome de Lille thinks the crystals of ammoniacal muriate to be octohedrons bundled together. This salt is sometimes, but not frequently, found in cubic crystals in the middle of the concave hollow part of the sublimated cakes. It possesses one singular physical property, a kind of ductility or elasticity, which causes it to yield under the hammer, or even the fingers, and makes it difficult to reduce it to a powder. Muriate of ammonia is totally volatile, but a very strong fire is requisite to sublime it. It is liable to no alteration from air; it may be kept for a long time without suffering any change; it dissolves very readily in water. Six parts of cold water are sufficient to dissolve one of the salt. A considerable cold is produced as the solution takes place, and this cold is still keener when the salt is mixed with ice. This artificial cold is happily applied to produce several phenomena which could not otherwise take place, such as the congelation of water on certain occasions,

the crystallization of certain salts, the fixation and preservation of certain liquids, naturally very subject to evaporation, &c.

SAL AMMONIACUM ACETOSUM. See *Liquor ammoniac acetatis*.

SAL AMMONIACUM LIQUIDUM. See *Liquor ammoniac acetatis*.

SAL AMMONIACUM MARTIALE. See *Ferrum ammoniacum*.

SAL AMMONIACUM SECRETUM GLAUBERI. See *Sulphur ammoniac*.

SAL AMMONIACUM VEGETABILE. See *Liquor ammoniac acetate*.

SAL AMMONIACUS FIXUS. The muriate of lime was formerly so termed.

SAL AMMONIACUS NITROSUS. See *Nitras ammoniac*.

SAL ANTIMONIUM. Tartar emetic.

SAL ARGENTI. Salt of Silver. See *Nitras argenti*.

SAL CATHARTICUS AMARUS. See *Magnesia sulphas*.

SAL CATHARTICUS ANGLICANUS. See *Magnesia sulphas*.

SAL CATHARTICUS GLAUBERI. See *Soda sulphas*.

SAL COMMUNIS. See *Soda murias*.

SAL CORNU CERVI VOLATILE. See *Carbonus Ammoniac*.

SAL CULINARIUS. See *Soda murias*.

SAL DE DUOBUS. See *Potassa sulphas*.

SAL DIURETICUS. See *Potassa acetat*.

SAL DIGESTIVUS SYLVII. A natural salt, formed of muriatic acid and potash. See *Potassa murias*.

SAL EPSOMENSIS. See *Magnesia sulphas*.

SAL ESSENTIALIS TARTARI. See *Tartar, acid of*.

SAL FERRIFUGUS SYLVII. See *Potassa murias*.

SAL FONTIUM. See *Soda murias*.

SAL FOSSILIS. See *Soda murias*.

SAL GEMME. Common or rock salt. See *Soda murias*.

SAL GLAUBERI. See *Soda sulphas*.

SAL HERBARUM. See *Potassa subcarbonas*.

SAL MARINUS. See *Soda murias*.

SAL MARTIS. See *Ferri sulphas*.

SAL MARTIS MURIATICUM SUBLIMATUM. See *Ferrum ammoniacum*.

SAL MICROCOSMICUS. The compound saline matter obtained by inspissating human urine.

SAL MIRABILIS GLAUBERI. See *Soda sulphas*.

SAL MURIATICUS. See *Soda murias*.

SAL PLANTARUM. See *Potassa subcarbonas*.

SAL POLYCHRESTUS. See *Potassa sulphas*.

SAL POLYCHRESTUS GLAUBERI. See *Potassa sulphas*.

SAL POLYCHRESTUS SEIGNETTI. See *Soda tartarizata*.

SAL FRUNELLÆ. Nitrat of potash cast into flat cakes or round balls.

SAL RUFELLENSIS. See *Soda tartarizata*.

SAL SATURNI. See *Plumbi superacetat*.

SAL SEDATIVUS. See *Boracic acid*.

SAL SEDATIVUS HOMBERGII. See *Boracic acid*.

SAL SEDLICIENSIS. See *Sulphas Magnesiæ*.

SAL BRIQUETTI. See *Soda tartarizata*.

SAL SUCCINI. The succinic acid. See *Succinic acid*.

SAL TARTARI. See *Potassa carbonas* and *Potassa*.

SAL THERMARUM CAROLINARUM. A union of sulphuric acid with magnesia. See *Sulphas Magnesiæ*.

SAL VEGETABILIS. See *Potasse tartaras*.

SAL VOLATILE. See *Spiritus ammoniæ compositus* and *Ammonia*.

SAL VOLATILIS SALIS AMMONIACI. See *Carbonas Ammonia*.

SALF.P. Salap. Salab. The root of the *Orchis morio* of Linnæus:—*bulbis indivisis, nectarii labio quadrilobis crenulato, cornu obtuso, petalis dorsulibus reflexis*. This farinaceous powder is imported from Turkey. It may be obtained from several other species of the same genus of plants. It is an insipid substance, of which a small quantity, by a proper management, converts a large portion of water into a jelly, the nutritive powers of which have been greatly over-rated. Salep forms a considerable part of the diet of the inhabitants of Turkey, Persia, and Syria. The method of preparing salep is as follows: The new root is to be washed in water, and the fine brown skin which covers it is to be separated by means of a small brush or by dipping the root in warm water, and rubbing it with a coarse linen cloth. The roots thus cleaned, are to be spread on a tin plate, and placed in an oven, heated to the usual degree, where they are to remain six or ten minutes. In this time they will have lost their milky whiteness, and acquired a transparency like horn, without any diminution of bulk. Being arrived at this state, they are to be removed in order to dry and harden in the air, which will require several days to effect; or they may be dried in a few hours, by using a very gentle heat. Salep, thus prepared, contains a great quantity of vegetable aliment; as a wholesome nourishment it is much superior to rice; and has the singular property of concealing the taste of salt water. Hence, to prevent the dreadful calamity of famine at sea, it has been proposed, that the powder of it should constitute part of the provisions of every ship's company. With regard to its medicinal properties, it may be observed, that its resto-

rative, mucilaginous, and demulcent qualities, render it of considerable use in various diseases, when employed as aliment, particularly in sea-scurvy, diarrhœa, dysentery, symptomatic fever, arising from the absorption of pus, and the stone or gravel.

SALICORNIA. The name of a genus of plants in the Linnæan system. Class, *Monandria*. Order, *Monogynia*.

SALICORNIA EUROPEA. The systematic name of the jointed glass wort, which is gathered by the country people and sold for samphire. It forms a good pickle with vinegar, and is little inferior to the samphire.

SALINE SUBSTANCES. The number of saline substances is very considerable; and they possess peculiar characters, by which they are distinguished from other substances. These characters are founded on certain properties, which it must be confessed, are not accurately distinctive of their true nature. All such substances, however, as possess several of the four following properties are considered as saline: 1. A strong tendency to combination, or a very strong affinity of composition; 2. A greater or lesser degree of sapidity; 3. A greater or lesser degree of solubility in water; 4. Perfect incombustibility.

SALIVICA. See *Nardus celtica*.

SALIVA. (So called, a *saline vapour*, from its salt taste, or from *salax*, spittle.) The fluid which is secreted by the salivary glands into the cavity of the mouth. The secretory organ is composed of three pair of salivary glands. 1. The *parotid glands*, which evacuate their saliva by means of the *Stenonian duct* behind the middle dens molaris of the upper jaw. 2. The *submaxillary glands*, which pour out their saliva through the *Warthonian ducts* on each side of the frenulum of the tongue by a narrow osculum. 3. The *sublingular glands*, situated between the internal surface of the maxilla and the tongue, and pour out their saliva through numerous *Riverian ducts* at the apex of the tongue.

The saliva in the cavity of the mouth has mixed with it, 1. The *mucus of the mouth*, which exhales from the labial and genal glands. 2. A *roscid vapour*, from the whole surface of the cavity of the mouth. The saliva is continually swallowed with, or without masticated food, and some is also spit out. It has no colour nor smell; it is *tasteless*, although it contains a little salt, to which the nerves of the tongue are accustomed. Its *specific gravity* is somewhat greater than water. Its *consistence* is rather plastic and spumous, from the entangled and atmospheric air. The quantity of twelve pounds is supposed to

be secreted in twelve hours. During mastication and speaking the secretion is augmented, from the mechanical pressure of the muscles upon the salivary glands. Those who are hungry secrete a great quantity, from the sight of agreeable food. It is imperfectly dissolved by water; somewhat coagulated by alcohol of wine; and is congealed with more difficulty than water. It is inspissated by a small dose, and dissolved in a large dose, of mineral acids. It is also soluble in aërated alkali. Caustic alkali and quick lime extract volatile alkali from saliva. It corrodes copper and iron, and precipitates silver and lead in the form of corneous luna. It assists the spirituous fermentation of farinaceous substances; hence barbarous nations prepare an inebriating drink from the chewed roots of the *Jatropha Manihot* and *Piper Methisticum*. It possesses an antiseptic virtue, according to the experiments of the celebrated Pringle. It easily becomes putrid in warm air, and gives off volatile alkali.

Constituent Principles. Saliva appears to consist of water, albumen, ammoniacal salt, and animal earth. Of water, there is four-fifths given out by distillation. The albumen is detected by alcohol of wine. The ammoniacal salt is demonstrated by triturating quick lime with saliva; and the animal earth from salival calculus, and the products of fire.

The use of the saliva is, 1. It augments the taste of the food, by the evolution of sapid matter. 2. During mastication, it mixes with, dissolves, and resolves into its principles, the food; and changes it into a pulaceous mass, fit to be swallowed: hence it commences chymification. 3. It moderates thirst, by moistening the cavity of the mouth and fauces.

SALIVAL DUCTS. The excretory ducts of the salival glands. That of the parotid gland is called the *Stenonian duct*; those of the submaxillary glands the *Warthonian ducts*; and those of the sublingual, the *Reverian ducts*.

SALIVAL GLANDS. Those glands which secrete the saliva are so termed. See *Saliva*.

SALIVANTIA. (From *saliva*, spittle.) Medicines which excite salivation.

SALIVARIA. (From *saliva*, the spittle; so called because it excites a discharge of saliva.) Pellitory of Spain.

SALIVARIS HERBA. See *Pyrethrum*.

SALIVATIO. An increased secretion of saliva. See *Ptyalismus*.

SALIX. (From *sala*, Heb.) 1. The name of a genus of plants in the Linnæan system. Class, *Diœcia*. Order, *Diandria*.

2. The pharmacopœial name of *Salix alba*. Crack willow. The bark of the

branches of the *Salix fragilis* of Linnæus: —*foliis serratis glabris ovato lanceolatis, petiolis densato glandulosis*. It manifests a considerable degree of bitterness to the taste, and is very adstringent. It is recommended as a good substitute for Peruvian bark, and is said to cure intermittents and other diseases requiring tonic and adstringent remedies. Not only the bark of this species of salix, but that of several others possess similar qualities, particularly of the *Salix alba* and *S. pentandria*, both of which are recommended in the foreign pharmacopœias. But Dr Woodville is of opinion, that the bark of the *salix triandria* is more effectual than that of any other of this genus; at least its sensible qualities give a decided preference. The trials Dr. Cullen made was with the bark of the *salix pentandria*, the bark taken from its branches, the third of an inch diameter, and of four or five years growth. Nevertheless, he adds, in intermittent fevers, he has always failed with this bark.

SALIX ALBA. See *Salix*.

SALIX CAPREA. The systematic name of a species of willow, the bark of whose branches possess the same virtues with that of the *fragilis*. See *Salix*.

SALIX FRAGILIS. The systematic name of the common crack willow. See *Salix*.

SALIX PENTANDRIA. The bark of the branches of this species of willow possesses the same virtues as that of the *fragilis*. See *Salix*.

SALIX VITULINA. The bark of the branches of this species of willow may be substituted for the *fragilis*. See *Salix*.

SALPINGO-PHARYNGEUS. This muscle is composed of a few fibres of the palatopharyngeus, which it assists in dilating the mouth of the Eustachian tube.

SALPINGO-STAPHILINUS. See *Levator palati*.

SALPINGO-STAPHILINUS INTERNUS. See *Levator palati*.

Salsofy. The root of the purple goat's beard. See *Tragopogon pratense*.

SALSOLA. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*.

SALSOLA KALI. Snail-seeded glass-wort or salt-wort. *Kali spinosum cochleatum. Tragus sive Tragum Matthioli*. The systematic name of the plant which affords the mineral alkali. See *Soda* and *Barilla*.

SALSOLA SATIVA. The systematic name of a plant, which affords the mineral alkali. See *Soda* and *Barilla*.

SALSOLA SODA. The systematic name of a plant which affords mineral alkali. See *Barilla*, and *Soda*.

Salt, cathartic. See *Sulphas Magnæ, Soda sulphas*.

Salt, common. See *Murias Soda*.
Salt, Epsom. See *Sulphas Magnesia*.
Salt privé. See *Potassa nitras*.
Salt, Rochelle. See *Soda tartarizata*.
Salt, sea. See *Murias Soda*.
Salt of steel. See *Ferri sulphas*.

SALTS. See *Saline substances*. Salts, with respect to their chemical properties, are divided into two classes; into acid salts or acids, and into alkaline salts or alkalis; and from the mutual combination of these two arises a third class, viz. that of neutral salts.

SALTS, ACID. These are distinguished by their sour taste when diluted with water. See *Acia*.

SALTS, ALKALINE. These possess a urinous, burning, and caustic taste, turn the syrup of violets to a green, have a strong affinity for acids, dissolve animal substances, unite readily with water, combines with oils and fat, and render them miscible with water, dissolve sulphur, and are crystallizable. See *Alkali*.

SALTS, NEUTRAL. Secondary salts. Under the name of neutral or secondary salts are comprehended such matters as are composed of two primitive saline substances combined together. These salts are called neutral, because they do not possess the characters of primitive salts; that is to say, they are neither acid nor alkaline salts; such as Epsom salts, alum, nitre, &c.

SALTS, PRIMITIVE. Simple salts. Under this order is comprehended those salts which were formerly thought to be simple or primitive, and which are occasionally called simple salts. The accurate experiments of the moderns have proved that these are for the most part compounded; but the term is retained with greater propriety when it is observed, that these salts compose, when united, salts which are termed secondary. These salts are never met with perfectly pure in nature, but require artificial processes to render them so. This order is divided into three genera, comprehending saline terrestrial substances, alkalis, and acids.

SALTS, SECONDARY. See *Neutral salts*.

Saltwort. See *Salsola kali*.

SALVATELLA. (*Salvatella*, sc. *vena*, from *salus*, health, because the opening of it was formerly thought to be of singular use in melancholy.) This vein runs along the little finger, unites upon the back of the hand with the cephalic of the thumb, and empties its blood into the internal and external cubital veins.

SALVIA. (*A salvendo*.) 1. The name of a genus of plants in the Linnæan system. Class, *Diantheria*. Order, *Monogynia*.

2. The pharmacopœial name of the

common sage. *Eleoaphates*. *Salvia officinalis* of Linnæus:—*foliis lanceolatis-ovatis integris crenulatis, floribus spicatis, calicibus acutis*. In ancient times sage was celebrated as a remedy of great efficacy, as would appear from the following lines of the school of Salerno:

Cur moriatur homo, cui salvia crescit in herbo?

Contra vim mortis, non est medicamen in herbo.

Salvia saluatrix, natura conciliatrix.

Salvia cum ruta faciunt tibi pocula tuta.

But at present it is not considered as an article of much importance. It has a fragrant strong smell; and a warm, bitterish, aromatic taste, like other plants containing an essential oil. It has a remarkable property in resisting the putrefaction of animal substances, and is in frequent use among the Chinese as a tonic, in the form of tea, in debility of the stomach and nervous system.

SALVIA HORTENSIS MINOR. The small sage, or sage of virtue. A variety of the official sage, possessing similar virtues.

SALVIA OFFICINALIS. The systematic name of the garden sage. See *Salvia*.

SALVIA SOLAREA. The systematic name of the garden clary, called *herminum* in the pharmacopœias. The leaves and seeds are recommended as corroborants and antispasmodics, particularly in leucorrhœas and hysterical weaknesses. They have a bitterish warm taste, and a strong smell, of the aromatic kind.

SAMBUCUS. (From *sabucca*, Heb. a musical instrument formerly made of this tree.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Trigynia*.

2. The pharmacopœial name of the elder-tree. *Sambucus vulgaris*. *Sambucus arboræ*. *Acta*. *Infelix lignum*. *Sambucus nigra* of Linnæus:—*cymis quinque partitis, foliis pinnatis, caule arboræ*. This indigenous plant has an unpleasant narcotic smell, and some authors have reported its exhalations to be so noxious, as to render it unsafe to sleep under its shade. The parts of this tree that are proposed for medicinal use in the pharmacopœias are the inner bark, the flowers, and the berries. The first has scarcely any smell, and very little taste; on first chewing, it impresses a degree of sweetness, which is followed by a very slight but durable acrimony, in which its powers seem to reside. From its cathartic property it is recommended as an effectual hydragogue by Sydenham and Boerhaave; the former directs three handfuls of it to be boiled in a quart of milk and water, till only a pint remains, of which one half is to be taken night and morning, and repeated for seven

ral days; it usually operates both upwards and downwards, and upon the evacuation it produces, its utility depends. Boerhaave gave its expressed juice in doses from a drachm to half an ounce. In smaller doses it is said to be an useful aperient and deobstruent in various chronic disorders. The flowers have an agreeable flavour; and infusions of them, when fresh, are gently laxative and aperient. When dry, they are said to promote chiefly the cuticular excretion, and to be particularly serviceable in erysipelatous and eruptive disorders. Externally they are used in ointments, &c. and in the London Pharmacopœia are directed in the form of an ointment. The berries in taste are somewhat sweetish, and not unpleasant; on expression they yield a fine purple juice, which proves an useful aperient and resolvent in sundry chronic diseases, gently loosening the belly, and promoting the urine and perspiration.

SAMBUCUS EBULUS. The systematic name of the dwarf elder. See *Ebulus*.

SAMBUCUS NIGRA. The systematic name of the elder-tree. See *Sambucus*.

SAMPHERE. The *Crithum martimum* of Linnaeus. It is a low perennial plant, and grows about the sea-coast in several parts of the island. It has a spicy aromatic flavour, which induces the common people to use it as a pot-herb. Pickled with vinegar and spice it makes a wholesome and elegant condiment which is in much esteem.

SAMPSUCHUS. See *Sambucus*.

SAMPSYCHUM. (From *σας*, to preserve, and *ψυχη*, the mind;) because of its cordial qualities. See *Sambucus*.

SANATIVA. (From *sano*, to cure.) Medicines which heal diseases.

SANCTUM SEMEN. The worm-seed, or antonicum.

SANDARACHA. (From *saghad narak*, Arab.) A gummy resin; also a sort of arsenic. See *Sandrack*.

SANDARACHA ARABUM. This resinous juice appears to have been the produce of a large species of juniper-tree.

Sanders. See *Santalum rubrum*.

SANDRACK. (An Arabian word.) *Sandaracha*. Gum juniper. A concrete resin which exudes in white tears, more transparent than mastich, from the bark of the *Juniperus communis* of Linnaeus. See *Juniperus*. Sandrack is almost totally soluble in alcohol, with which it forms a white varnish that dries speedily. Reduced to powder it is called *pounce*, which prevents ink from sinking into paper from which the exterior coating of size has been scraped away.

SANDIX. (From *sani dak*, red, Arab.) Ceruss burnt till it becomes red.

SANGUIFICATION. (*Sanguificatio*,

from *sanguis*, blood.) A natural function of the body, by which the chyle is changed into blood. The uses of sanguification are the generation of blood, which serves to fill the blood-vessels, to irritate and stimulate the heart and arteries, to generate or cause heat, to secrete the humours, and to excite the vital action.

SANGUINALIS. (From *sanguis*, blood; so named from its uses in stopping bleedings.) The *Polygonum aviculare*, or knot-grass, is sometimes so called. See *Centumnodia*.

SANGUINARIA. (From *sanguis*, blood; so named from its use in stopping bleedings.) The *Polygonum aviculare*, or knot-grass is sometimes so termed. See *Centumnodia*.

Sanguineous apoplexy. See *Apoplexia*.

SANGUIFUGIUM. (From *sanguis*, blood, and *purgo*, to purge.) A gentle fever, or such a one as by its discharges is supposed to purify the blood.

SANGIUS. See *Blood*.

SANGUIS DRACONIS. *Cinnabaris græcorum. Draconthema. Asagen. Asegen.* Dragon's blood. The red resinous juice which is obtained by wounding the bark of the *Culamus rotang*;—*caudice densissime aculeato, aculeis erectis, spadice erecto.* It is chiefly obtained from the Molucca islands, Java, and other parts of the East Indies. It is generally much adulterated, and varied in goodness and purity. The best kind is of a dark red colour, which, when powdered, changes to crimson; it readily melts and catches flame; has no smell, but to the taste discovers some degree of warmth and pungency. The ancients Greeks were well acquainted with the adstringent power of this drug; in which character it has since been much employed in hæmorrhages, in alvine fluxes. At present, however, it is not used internally, being superseded by more certain and effectual remedies of this numerous class.

SANGUIS HERCULIS. A name for the crocus.

SANGUISORBA. The name of a genus of plants in the Linnæan system. Class, *Triandria*. Order, *Monogynia*.

SANGUISORBA OFFICINALIS. The systematic name of the Italian pimpinel. See *Pimpinella Italica*.

SANGUISUGA. (From *sanguis*, blood, and *sugo*, to suck.) The leech, or blood-sucker. See *Leech*.

Sanicle. See *Sanicula*.

Sanicle, Yorkshire. See *Pinguicula*.

SANICULA. (From *sano*, to heal; so called from its virtues in healing.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*.

2. The pharmacopœial name of *sanicle*.

Cucullata. Dodecatheon. Symphytum patrum Sanicula mus. Diapensia certusa. This herb, *Sanicula europea* of Linnæus, was formerly recommended as a mild adstringent, and is supposed to have received its name from its sanative power. Its sensible qualities are a bitterish and somewhat austere taste, followed by an acrimony which chiefly affects the throat. It is only in use in the present day amongst the country people.

SANICULA EBORACENSIS. Pinguicula. Sanicula montana. Viola palustris. The Yorkshire sanicle or butter-wort. See *Pinguicula*.

SANICULA EUROPEA. The systematic name of the sanicle. See *Sanicula*.

SANICULA MAS. See *Sanicula*.

SANIES. *Icher.* This term is sometimes applied to a thin, limpid, and greenish discharge; and at other times to a thick and bloody kind of pus.

SANTALUM. (From *zandal*, Arab.) The name of a genus of plants in the Linnæan system. Saunders.

SANTALUM ALBUM. The systematic name of the yellow saunders. See *Santalum citrinum*.

SANTALUM ALBUM. The white saunders of the shop is said to be the alburnum of the tree whose medullary part is called *Santalum citrinum*.

SANTALUM CITRINUM. Santalum pallidum. Yellow saunders. *Santalum album* of Linnæus. White saunders wood is of a pale white colour, often with a yellowish tinge, and, being destitute of taste or odour, it is superseded by the *santalum citrinum*, which is of a brownish yellow colour, of a bitterish aromatic taste, and of a pleasant smell, approaching to that of the rose. Both kinds are brought from the East Indies in billets, consisting of large thick pieces, which, according to Rumphius, are sometimes taken from the same, and sometimes from different trees. For though the white and yellow saunders are the wood of the same species of tree, yet the latter, which forms the central part of the tree, is not always to be found in sufficient quantity to repay the trouble and expense of procuring it especially, unless the trees be old; while the white, which is the exterior part of the wood, is always more abundant, and is consequently much cheaper.

Yellow saunders, distilled with water, yields a fragrant essential oil, which thickens in the cold into the consistence of a balsam, approaching in smell to ambergris, or a mixture of ambergris and roses; the remaining decoction, inspissated to the consistence of an extract, is bitterish and slightly pungent. Rectified spirit extracts, by digestion, considerably more than water; the colour of the tincture is

a rich yellow. The spirit distilled off is slightly impregnated with the fine flavour of the wood; the remaining brownish extract has a weak smell, and a moderate balsamic pungency.

The wood is valued highly on account of its fragrance, hence the Chinese are said to fumigate their clothes with it, and to burn it in their temples in honour of their gods. Though still retained in the *Materia Medica* of the Edinburgh Pharmacopœia, it cannot be thought to possess any considerable share of medicinal power. Hoffman considers its virtues as similar to those of ambergris; and some others have esteemed it in the character of a corroborant and restorative.

SANTALUM FALLIDUM. See *Santalum citrinum*.

SANTALUM RUBRUM. Red saunders. *Pterocarpus santalinus* of Linnæus. There is some reason to believe that several red woods, capable of communicating this colour to spirituous liquors, are sold as red saunders; but the true officinal kind appears, on the best authority, to be of this tree, which is extremely hard, of a bright garnet red colour, and bears a fine polish. It is only the inner substance of the wood that is used as a colouring matter, and the more florid red is mostly esteemed. On being cut it is said to manifest a fragrant odour, which is more especially in old trees. According to Lewis, this wood "is of a dull red, almost blackish colour on the outside, and a deep brighter red within; its fibres are now and then curled, as in knots. It has no manifest smell, and little or no taste; even of extracts made from it with water, or with spirit, the taste is not considerable."

To watery liquors it communicates only a yellowish tinge, but to rectified spirit a fine deep red. A small quantity of an extract made with this menstruum, tinged a large one of fresh spirit of the same colour; though it does not, like most other resinous bodies, dissolve in expressed oils. Cf distilled oils, there are some, as that of lavender, which receive a red tincture from the wood itself, and from its resinous extract, but the greater number do not. Red saunders has been esteemed as a medicine; but its only use attaches to its colouring property. The juice of this tree, like that of some others, affords a species of sanguis draconis.

SANTOLINA. (From *santalum*, saunders, because it smells like the saunders wood.) See *Abrotanum femina*.

SANTOLINA CHANE-CYPARISSUS. The systematic name of the lavender cotton. See *Abrotanum femina*.

SANTONICUM. (From *Santonis*, its native place.) This plant and its seeds have received several names. *Abrothium*

santonicum Alexandrinum. Semontina. Absinthium scriphtum. Egyptium. Scheba Arabum. Zedoaria sem. Xantolna. Lumbricorum semina. Cina. Semen contra. Semen sanctum. It is the Tartarean southern-wood or wormseed, *Artemisia santonica* of Linnaeus — *foliis caulibus linearibus pinnato-multifidis, ramis indivisis, apicis secundis reflexis, floribus quinquefloris.* The seeds are small, light, and oval, composed of a number of thin membranous coats of a yellowish green colour, with a cast of brown, easily friable, upon being rubbed between the fingers, into a fine chaffy kind of substance. They are brought from the Levant : have a moderately strong and not agreeable smell, somewhat of the wormwood kind, and a very bitter sub-acrid taste. Their virtues are extracted both by watery and spirituous menstrua. They are esteemed to be stomachic, emmenagogue, and anthelmintic; but it is especially for the last-mentioned powers that they are now administered, and from their efficacy in this way they have obtained the name of wormseed. To adults the dose in substance is from one to two drachms, twice a day. Lewis thinks that the spirituous extract is the most eligible preparation of the santonicum, for the purposes of an anthelmintic.

SAPHENA. (*Vena saphena* : from *saphes*, visible.) The large vein of the leg, which ascends along the little toe over the external ankle, and evacuates part of the blood from the foot into the popliteal veins.

SAPIENTIE DENTES. The four last grinders are so called, because they appear when the person is supposed to be at years of discretion. See *Teeth*.

SAPINUS SAPONARIA. The systematic name of the plant which affords soap-nuts. See *Saponaria mucula*.

SAPU. Soap. A composition of oils, or fats, with an alkali. The medicinal soap, *sapo duras, sapo amygdalinus*, is made with oil of sweet almonds, and half its weight of potash or caustic alkali. Common or soft soap, *sapo mollis*, is made of tallow; Spanish, or Castile soap, of oil of almonds, and soda, or barilla. Black soap is a composition of train oil and an alkali; and green soap of hemp, linseed, or rape oil. The *sapo ex oleo olivæ et natro confectus* of the former London Pharmacopœia, or the *sapo albus crispans* of the Edinburgh Pharmacopœia, (white Spanish soap,) being made of the finer kinds of olive oil, is the best, and therefore preferred for internal use. Soap was imperfectly known to the ancients. It is mentioned by Pliny as made of fat and ashes, and as an invention of the Gauls. Aretæus and others inform us, that the Greeks obtained their knowledge of its medical use from the Romans.

Its virtues, according to Bergius, are detergent, resolvent, and aperient, and its use recommended in jaundices, gout, calculous complaints, and in obstructions of the viscera. The efficacy of soap, in the first of these diseases, was experienced by Sylvius, and since recommended very generally by various authors who have written on this complaint; and it has also been thought of use in supplying the place of bile in the primæ viæ. The utility of this medicine, in ictical cases, was inferred chiefly from its supposed power of dissolving biliary concretions; but this medicine has lost much of its reputation in jaundice, since it is now known, that gallstones have been found in many after death, who had been daily taking soap for several months, and even years. Of its good effects in urinary calculous affections, we have the testimonies of several, especially when dissolved in lime-water, by which its efficacy is considerably increased; for it thus becomes a powerful solvent of mucus, which an ingenious modern author supposes to be the chief agent in the formation of calculi; it is, however, only in the incipient state of the disease that these remedies promise effectual benefit, though they generally abate the more violent symptoms, where they cannot remove the cause. With Boerhaave, soap was a general medicine; for as he attributed most complaints to viscosity of the fluids, he, and most of the Boerhaavian school, prescribed it, in conjunction with different resinous and other substances, in gout, rheumatism, and various visceral complaints. Soap is also externally employed as a resolvent, and gives name to several official preparations.

SAPU TEREBINTHINÆ. Starkey's soap. "*℞ Kali præparati calidi, ℥j. Olei terebinth. ℥iii.*" The hot kali præparatum is to have the oil of turpentine gradually blended with it, in a heated mortar. Indolent swellings were formerly rubbed with this application, and perhaps some chronic affections of the joints might still be benefited by it.

SAPONARIA. (From *sapo*, soap; so called because its juice, like soap, cleans clothes.) 1. The name of a genus of plants in the Linnaean system. Class, *Decandria*. Order, *Digynia*.

2. The pharmacopœial name of the soapwort, Bruise-wort, *Struthium. Lanaria. Lychnis syriacus. Ithuxum.* The root of this plant *Saponaria officinalis* of Linnaeus — *Catycibus cylindricis, foliis ovato-lanceolatis*, is employed medicinally; it has no peculiar smell; its taste is sweetish, glutinous, and somewhat bitter. On being chewed for some time, it is said to discover a degree of acrimony, which continues to affect the mouth a considerable time. According to

Neuman, two ounces of the root yielded eleven drachms of watery extract; but Cartheuser, from a like quantity, only obtained six drachms and twenty four grains. This extract manifested a sweetish taste, followed by an acrid quality. The spirituous extract is less in quality, but of a more penetrating acrid taste. Decoctions of the root, on being sufficiently agitated, produce a saponaceous froth, a similar soapy quality is observable also in the extract, and still more manifestly in the leaves, in-somuch that they have been used by the medicant monks as a substitute for soap in washing of their clothes, and Bergius, who made several experiments with the saponaria, declares that it had all the effects of soap itself.

From these peculiar qualities of the saponaria, there can be little doubt of its possessing a considerable share of medical efficacy, which Dr. Woodville says he could wish to find faithfully ascertained.

The diseases for which the saponaria is recommended, as syphilis, gout, rheumatism, and jaundice, are not, perhaps, the complaints in which its use is most availing; for a fancied resemblance of the roots of saponaria with those of sarsaparilla, seems to have led physicians to think them similar in their effects; and hence they have both been administered with the same intentions, particularly in fixed pains, and venereal affections. Bergius says, "in arthritide, cura mercuriale, &c. nullum aptiorem potum novi." However, according to several writers, the most inveterate cases of syphilis were cured by a decoction of this plant, without the use of mercury.

Haller informs us, that Boerhaave entertained an high opinion of its efficacy in jaundice, and other visceral obstructions.

SAPONARIA NUCULA. *Bacca bermudensis.* Soap-berries. A spherical fruit, about the size of a cherry, whose cortical part is yellow, glossy, and so transparent as to shew the spherical black nut which rattles within, and which includes a white kernel. It is the produce of the *Sapindus saponaria* of Linnæus, which grows in Jamaica. It is said that the cortical part of this fruit has a bitter taste, and no smell; that it raises a soapy froth with water, and has similar effects with soap in washing; that it is a medicine of singular and specific virtue in chlorosis. They are not known in the shops of this country.

SAPONARIA OFFICINALIS. The systematic name of the soap-wort. See *Saponaria*.

SAPONULES. *Saponuli.* Combinations of the volatile or essential oils with different bases; as *saponule of alumine*.

SAPONULES, ACID. Combinations of the volatile or essential oils with different acids.

SAPOTA. The oval fruited sapota, whose seeds are sometimes given in the form of emulsion in calculous complaints, is the *Acras sapota* of Linnæus. It is a native of South America, and bears a fruit like an apple; which has, when ripe, a lascivious taste, resembling that of the marmalade of quinces, whence it is called natural marmalade.

SAPPAN LIGNUM. Logwood has been so called See *Lignum campechense*.

SAPPHARINA AQUA. *Aqua cupri ammoniati.* Made by a solution of sal ammoniac in lime-water, standing in a copper vessel.

SAPPHIRE. A gem of a sky-blue colour.

Saracens consoud. See *Virga aurea*.

SARCUM. (Dim. of *sagξ*, flesh.) A carbuncle, or small fleshy excrescence.

SARCITES. (From *sagξ*, flesh.) An anasarca.

SARCOCELE. (From *sagξ*, flesh, and *κῆλη*, a tumour.) *Hernia carnea.* This is a disease of the body of the testicle, and, as the term implies, consists, in general, in such an alteration made in the structure of it, as produces a resemblance to a hard fleshy substance, instead of that fine, soft, vascular texture, of which it is, in a natural and healthy state, composed.

The ancient writers have made a great number of distinctions of the different kinds of this disease, according to its different appearances, and according to the maligness, or malignity of the symptoms with which it may chance to be attended. Thus, the *sarcocele*, the *hydro-sarcocele*, the *scirrhus*, the *cancer*, the *cara adnata ad testem*, and the *cara adnata ad vasa*, which are really little more than descriptions of different states and circumstances of the same disease, are reckoned as so many different complaints, requiring a variety of treatment, and deriving their origin from a variety of different humours.

Every species of sarcocele consists primarily in an enlargement, induration, and obstruction of the vascular part of the testicle; but this alteration is, in different people, attended with such a variety of circumstances, as to produce several different appearances, and to occasion the many distinctions which have been made.

If the body of the testicle, though enlarged, and indurated to some degree, be perfectly equal in its surface, void of pain, has no appearance of fluid in its tunica vaginalis, and produces very little uneasiness, except what is occasioned by its mere weight, it is usually called a simple sarcocele, or an indolent scirrhus; if, at the same time that the testis is enlarged and hardened, there be a palpable accumulation of fluid in the vaginal coat, the

disease has by many been named a *hydro-sarcocoele*; if the lower part of the spermatic vessels, and the epididymis were enlarged, hard, and knotty, they supposed it to be a fungous, or morbid accretion, and called it the *caro adnata ad vasa*; if the testicle itself was unequal in its surface, but at the same time not painful, they distinguish it by the title of *caro adnata ad testem*; if it was tolerably equal, not very painful, nor frequently so, but at the same time hard and large, they gave it the appellation of an occult or benign cancer; if it was ulcerated, subject to frequent acute pain, to hæmorrhage, &c. it was known by that of a malignant or confirmed cancer. These different appearances, though distinguished by different titles, are really no more than so many stages (as it were) of the same kind of disease, and depend a great deal on several accidental circumstances, such as age, habit, manner of living, &c. It is true, that many people pass several years with this disease, under its most favourable appearances, and without encountering any of its worst; but, on the other hand, there are many, who, in a very short space of time, run through all its stages. They who are most conversant with it, know how very convertible its milkiest symptoms are into its most dreadful ones, and how very short a space of time often intervenes between the one and the other.

There is hardly any disease affecting the human body which is subject to more variety than this is, both with regard to its first manner of appearance, and the changes which it may undergo.

Sometimes the first appearance is a mere simple enlargement and induration of the body of the testicle; void of pain, without inequality of surface, and producing no uneasiness, or inconvenience, except what is occasioned by its mere weight. And some people are so fortunate to have it remain in this state for a very considerable length of time without visible or material alteration. On the other hand, it sometimes happens, that very soon after its appearance in this mild manner, it suddenly becomes unequal and knotty, and is attended with very acute pains darting up to the loins and back, but still remaining entire, that is, not bursting through the integuments. Sometimes the fury of the disease brooks no restraint; but, making its way through all the membranes which envelope the testicle, it either produces a large, foul, stinking, phagedenic ulcer, with hard edges, or it thrusts forth a painful gleetng fungus, subject to frequent hæmorrhage.

Sometimes an accumulation of water is made in the tunica vaginalis, producing that mixed appearance, called the *hydro-sarcocoele*.

Sometimes there is no fluid at all in the cavity of the tunica vaginalis; but the body of the testicle itself is formed into cells, containing either a turbid kind of water, a bloody sanies, or a purulent fetid matter. Sometimes the disorder seems to be merely local, that is, confined to the testicle, not proceeding from a tainted habit, nor accompanied with diseased viscera, the patient having all the general appearances and circumstances of health, and deriving his local mischief from an external injury. At other times, a pallid, leaden countenance, indigestion, frequent nausea, choleric pains, sudden purgings, &c. sufficiently indicate a vitiated habit, and diseased viscera, which diseased viscera may also sometimes be discovered and felt.

The progress also which it makes from the testis upward, toward the process, is very uncertain; the disease occupying the testicle only, without affecting the spermatic process, in some subjects, for a great length of time; while, in others, it totally spoils the testicle very soon, and, almost as soon, seizes on the spermatic chord.

SARCOCOLLA. (From *σαρξ*, flesh, and *κόλλα*, glue; because of its supposed power of gluing together wounds) A concrete gummi-resinous juice, supposed to be the produce of the *Penae mucronata* of Linnæus. It is brought from Persia and Arabia in small grains of a pale yellow colour, having also sometimes mixed with them a few of a deep red colour. Its taste is bitter, but followed with some degree of sweetness. It has been chiefly used for external purposes, and, as its name imports, has been thought to agglutinate wounds and ulcers; but this opinion now no longer exists.

SARCOEPHLOCLE. Enlarged testicle, with rupture, containing omentum.

SARCOLOGY. (*Sarcologia*. From *σαρξ*, flesh, and *λογία*, a discourse) The doctrine of the muscles and soft parts.

SARCOMA. (From *σαρξ*, flesh.) *Sarcosis*. *Porrus*. *Sarcophya*. *Nævus*. A fleshy excrescence. A genus of disease in the class *locales*, and order *tumores* of Cullen.

SARGOMPHALUS. (From *σαρξ*, flesh, and *ομφαλος*, the navel.) A fleshy excrescence about the navel.

SARCOPHYIA. (From *σαρξ*, flesh, and *φυειν*, to grow.) A fleshy excrescence.

SARCOPTODES. (From *σαρξ*, flesh, and *πύον*, pus.) Applied to the purulent, fleshy discharge which is thrown up in some stages of consumption.

SARCOSIS. (From *σαρξ*, flesh.) A fleshy tumour. The generation of flesh.

SARCOITIA. (From *σαρξ*, flesh.) Medicines which promote the generation of flesh in wounds.

SARDIATIS. (From *sardurus*, the sardonian, or herb, which, being eaten, causes convulsive laughter.) The *Risus sardonius* or a convulsive involuntary laughter.

SARDONIA. (From *Sardonis*, its native soil.) A kind of smallage.

SARDONIC LAUGH. *Risus sardoniacus*; so called from the herb *sardonis*, which being eaten is said to cause a deadly convulsive laughter. Hence *risus sardonius*, the sardonian laughter, or spasmodic grin.

SARDONICUS RISUS. See *Sardonic laugh*.

SARSAPARILLA. (This word is of Spanish origin, signifying a red tree.) *Smilax aspera Peruviana. Sarsa. Caribellanli. Iva pecanga. Macapasti. Zarza. Zarsaparilla. Salsaparilla. Zarzaparilla.* The root of this plant, *Smilax sarsaparilla* of Linnaeus:—*caule aculeato angulato, foliis inermibus ovatis retusumucronatis trinerviis*, has a farinaceous, somewhat bitter taste, and no smell. About two centuries ago it was introduced into Spain, as an undoubted specific in syphilitic disorders; but owing to difference of climate, or other causes, it has not answered the character which it had acquired in the Spanish West Indies. It is now considered as capable of improving the general habit of body, after it has been reduced by the continued use of mercury.

To refute the opinion that sarsaparilla possesses antisymphilitic virtues, Mr. Pearson, of the Lock Hospital, divides the subject into two distinct questions. 1. Is the sarsaparilla root, when given alone, to be safely relied on in the treatment of lues venerea? The late Mr. Bloomfield, his predecessor, and during some years his colleague, at the Lock Hospital, has given a very decided answer to this question: "I solemnly declare (says he) I never saw a single instance in my life where it cured that disorder without the assistance of mercury, either at the same time with it, or when it had been previously taken before the decoction was directed." Mr. Pearson's experience, during many years, coincides entirely with the observations of Mr. Bloomfield. He has employed the sarsaparilla, in powder and in decoctions, in an almost infinite variety of cases, and feels himself fully authorised to assert, that this plant has not the power of curing any one form of the lues venerea. The sarsaparilla, indeed, like the guaiacum, is capable of alleviating symptoms derived from the venereal virus; and it sometimes manifests the power of suspending, for a time, the destructive ravages of that contagion; but where the poison has not been previously subdued by mercury, the symptoms will quickly return; and, in addition to them, we often see the most indubitable proofs that the disease is making an actual

progress, during the regular administration of the vegetable remedy.

2. When the sarsaparilla root is given in conjunction with mercury, does it render the mercurial course more certain and efficacious? In replying to this query it is necessary to observe, that the phrase, "to increase the efficacy of mercury," may imply, that a smaller quantity of this mineral antidote will confer security on an infected person, when sarsaparilla is added to it; or it may mean, that mercury would be sometimes unequal to the cure, without the aid of sarsaparilla. If a decoction of this root did indeed possess so admirable a quality, that the quantity of mercury necessary to effect a cure might be safely reduced, whenever it was given during a mercurial course, it would form a most valuable addition to our *Materia Medica*. This opinion has been, however, unfortunately falsified by the most ample experience, and whoever shall be so unwary as to act upon such a presumption, will be sure to find his own and his patient's expectations egregiously disappointed.

If the sarsaparilla root be a genuine antidote against the syphilitic virus, it ought to cure the disease when administered alone; but, if no direct proof can be adduced of its being equal to this, any arguments founded on histories where mercury has been previously given, or where both the medicines were administered at the same time, must be ambiguous and undecisive.

It appears probable, that Sir William Fordyce, and some other persons, entertained a notion, that there were certain venereal symptoms which commonly resisted the potency of mercury, and that the sarsaparilla was an appropriate remedy in these cases. This opinion, it is presumed, is not correct, for it militates against all Mr. P. has ever observed of the progress and treatment of lues venerea. Indeed those patients who have lately used a full course of mercury, often complain of nocturnal pains in their limbs; they are sometimes afflicted with painful enlargements of the elbow and knee-joints; or they have membranous nodes, cutaneous exulcerations, and certain other symptoms, resembling those which are the offspring of the venereal virus.

It may and does often happen, that appearances like these are mistaken for a true venereal affection, and, in consequence of this error, mercury is administered, which never fails to exasperate the disease. Now, if a strong decoction of sarsaparilla-root be given to persons under these circumstances, it will seldom fail of producing the most beneficial effects; hence it has been contended, that symptoms derived from the contagion

of lues venerea, which could not be cured by mercury, have finally yielded to this vegetable remedy. It must be acknowledged, that representations of this kind have a specious and imposing air; nevertheless, Mr. Pearson endeavours to prove, that they are neither exact nor conclusive. If any of the above-named symptoms should appear near the conclusion of a course of mercury, when that medicine was operating powerfully on the whole system, it would be a strange and inexplicable thing if they could possibly be derived immediately from the uncontrolled agency of the venereal virus.

This would imply something like a palpable contradiction, that the antidote should be operating with sufficient efficacy to cure the venereal symptoms, for which it was directed, while, at the same time, the venereal virus was proceeding to contaminate new parts, and to excite a new order of appearances.

One source, and a very common one, to which some of the mistakes committed upon this subject may be traced, is a persuasion that every morbid alteration which arises in an infected person is actually tainted with the venereal virus, and ought to be ascribed to it as its true cause.

Every experienced surgeon must, however, be aware, that very little of truth and reality exists in a representation of this kind. The contagious matter, and the mineral specific, may jointly produce, in certain habits of body, a new series of symptoms, which, strictly speaking, are not venereal, which cannot be cured by mercury, and which are sometimes more to be dreaded than the simple and natural effects of the venereal virus.

Some of the most formidable of these appearances may be sometimes removed by sarsaparilla, the venereal virus still remaining in the system; and, when the force of that poison has been completely subdued by mercury, the same vegetable is also capable of freeing the patient from what may be called the sequelæ of a mercurial course.

The root of the sarsaparilla is sometimes employed in rheumatic affections, scrofula, and cutaneous complaints, where an acrimony of the fluids prevail.

SARSAPARILLA GERMANICA. The root of the *Carex arenaria* of Linneus, which grows plentifully on the sea coast, is so termed, and it appears, that the *carex disticha* and *hirta* have also been collected, and their roots used indifferently instead of the true sarsaparilla. The root of the *carex arenaria* has been found serviceable in some mucal affections of the trachea, in rheumatic pains, and gouty affections.

SARTORIUS. (*Sartorius*, so. *muscu-*

lus; from *sartor*, a tailor, because tailors cross their legs with it.) *Sartorius seu longissimus femoris* of Cowper, and *Ilio cresti tibialis* of Dumas. This flat and slender muscle, which is the longest of the human body, and from an inch and a half to two inches in breadth, is situated immediately under the integuments, and extends obliquely from the upper and anterior part of the thigh, to the upper, anterior, and inner part of the tibia, being enclosed by a thin membranous sheath, which is derived from the adjacent *fascia lata*. It arises, by a tendon of about half an inch in breadth, from the outer surface and inferior edge of the anterior superior spinous process of the ilium, but soon becomes fleshy, and runs down a little way obliquely inwards, and then for some space upon the rectus, nearly in a straight direction; after which it passes obliquely over the vastus internus, and the lower part of the adductor longus, and then running down between the tendons of the adductor magnus and the gracilis, is inserted, by a thin tendon, into the inner part of the tibia, near the inferior part of its tuberosity, and for the space of an inch or two below it. This tendon sends off a thin aponeurosis, which is spread over the upper and posterior part of the leg. This muscle serves to bend the leg obliquely inwards, or to roll the thigh outwards, and at the same time to bring one leg across the other, on which account Spigelius first gave it the name of *sartorius*, or the tailor's muscle.

SASSAFRAS. (*Quasi saxifraga*; from *saxum*, a stone, and *frango*, to break; so called because a decoction of its wood was supposed good for the stone.) *Cornus mas edulota*. *Lignum purpureum*. *Anhuiba*. The wood of the sassafras-tree, *Laurus sassafras* of Linneus:—*foliis trilobis integrisque*, is imported from North America, in long straight pieces, very light, and of a spongy texture, and covered with a rough fungous bark. It has a fragrant smell, and a sweetish, aromatic, subacid taste; the root, wood, and bark, agree in their medicinal qualities, and are all mentioned in the pharmacopœias; but the bark is the most fragrant, and thought to be more efficacious than the woody part; and the branches are preferred to the large pieces. The medical character of this drug was formerly held in great estimation, and publications were professedly written on the subject. It is now, however, thought to be of little importance, and seldom used but in conjunction with other medicines, as a corrector of the fluids. It is an ingredient in the *decoctum sarsaparille compositum*, or *decoctum lignorum*; but the only official preparation of it is the essential oil, which is carminative and stimu-

lant, and which may be given in the dose of two drops to ten.

SATANUS DEVORANS. The red lions, or antimony.

SATELLITE VEINS. The veins which accompany the brachial artery as far as the bend of the cubit.

SATH. The penis.

SATURANTIA. Medicines which neutralize the acid in the stomach.

SATURATION. A term employed in pharmacy and chemistry to express the state of a body which has a power of dissolving another, to a certain extent only, in which it has effected that degree of solution: thus, nitric acid, for instance, can only dissolve a certain quantity of lime, beyond which it does not act, having lost its former affinity; this degree of solution is termed the point of saturation, and it is then said that the nitric acid is saturated with lime.

SATUREJA. (From *satyri*, the lustful satyr, because it make those who eat it lascivious. Blanch.) 1. The name of a genus of plants in the Linnæan system. Class, *Didynamia*. Order, *Gymnospermia*.

2. The pharmacopœial name of the summer savory. *Satureja sativa*. *Culina sativa* Plinii. *Thymra*. This low shrub is the *Satureja sativa* of Linnæus, cultivated in our gardens for culinary purposes. It has a warm, aromatic, penetrating taste, and smells like thyme, but milder. It is an ingredient in most of the warm stews and made dishes.

SATUREJA CAPITATA. The systematic name of the ciliated savory. See *Thymus creticus*.

SATUREJA HORTENSIS. The systematic name of the summer savory. See *Satureja*.

SATUREJA SATIVA. See *Satureja*.

SATURNUS. (From the planet, or bea-then god, of that name.) The chemical name of lead.

SATYRIASIS. (From *satyros*, a satyr, because they are said to be greatly addicted to venery.) *Satyriasmus*. *Priapismus*. *Salucitas*. *Brachuna*. *Arascon*. Excessive and violent desire for coition in men. A genus of disease in the class *locales*, and order *dysorexie* of Cullen.

SATYRIUM. (From *satyros*, an animal given to venery, so called because it was supposed to excite venery if only held in the hand.) Dog-stones. Male orchis. The root of the *Orchis mascula* of Linnæus:—*bulbis indivisis, nectaris labio quadrilobo ornulata, cornu obtuso petalis dorsalis reflexis*, which has a place in the *Materia Medica* of the Edinburgh Pharmacopœia, on account of the glutinous slimy juice which it contains. The root of the *orchis bifolia* is also collected. *Satyrium* root has a sweetish taste, a faint and somewhat unpleasant smell. Its mucilaginous or gelatinous

quality has recommended it as a demulcent. *Salep*, which is imported here from the East, is a preparation of this root, which, considered as an article of diet, is accounted extremely nutritious, as containing a great quantity of farinaceous matter in a small bulk. The supposed aphrodisiac qualities of this root, which have been noticed ever since the days of Dioscorides, seem, says Dr. Woodville, to be founded on the fanciful doctrine of signatures, thus orchis, i. e. *eggs*, testiculus, habet, radices, instar testiculorum. See *Salep*.

SATYRIUM. See *Satyrium*.

Sauce alone. See *Miliaria*.

Saunders, red. See *Santalum rubrum*.

Saunders, yellow. See *Santalum citrinum*.

SAUR KRAUT. *Souf* crou. Cabbage preserved in brine. Antiscorbutic.

Savin See *Sabina*.

Savin ointment. See *Ceratum sabine*.

SAYINA. See *Sabina*.

Savoury. See *Satureja*.

SAXIFRAGA. The name of a genus of plants in the Linnæan system. Class, *Dicandria*. Order, *Digynia*.

SAXIFRAGA ALBA. Called by Orbesius *Beste*. *Sanicula pedum*. White saxifrage. *Saxifraga granulata* of Linnæus, who describes the taste of this plant to be acrid and pungent, which we have not been able to discover: neither the tubercles of the root nor the leaves manifest to the organs of taste any quality likely to be of medicinal use, and therefore, though this species of saxifrage has been long employed as a popular remedy in nephritic and gravelly disorders, yet we do not find, either from its sensible qualities, or from any published instances of its efficacy, that it deserves a place in the *Materia Medica*. The superstitious doctrine of signatures suggested the use of the root, which is a good example of what Linnæus has termed *radix granulata*. The bulbs, or tubercles of such roots answer an important purpose in vegetation, by supplying the plants with nourishment and moisture, and thereby enabling them to resist the effects of that drought to which the dry soils they inhabit peculiarly expose them.

SAXIFRAGA ANGELICA. See *Saxifraga vulgaris*.

SAXIFRAGA CRASSIFOLIA. The roots of this species of saxifrage is extolled by Professor Pallas as an antiseptic.

SAXIFRAGA GRANULATA. The systematic name of the white saxifrage. See *Saxifraga alba*.

SAXIFRAGA RUBRA. See *Filipendula*.

SAXIFRAGA VULGARIS. *Saxifraga angelica*. *Hippomarathrum*. *Peniculum erratum*. English, or meadow saxifrage. *Pecculanum silaus* of Linnæus. The roots, leaves, and seeds of this plant have been

commended as aperients, diuretics, and carminatives; and appear, from their aromatic smell, and moderately warm, pungent, bitterish taste, to have some claim to these virtues. They are rarely used.

Saxifrage burnet. See *Pimpinella*.

Saxifrage, English. See *Saxifraga vulgaris*.

Saxifrage, meadow. See *Saxifraga vulgaris*.

Saxifrage, white. See *Saxifraga alba*.

SCAB. A hard substance covering superficial ulcerations, and formed by a concretion of the fluid discharged from them.

SCABIES. See *Psora*.

SCABIOSA. (From *scaber*, rough; so called from its rough hairy surface.) 1. The name of a genus of plants in the Linnean system. Class, *Tetrandria*. Order, *Monogynia*.

2 The pharmacopœial name of the common scabious. This herb, *Scabiosa arvensis*, *corollis quadrifidis, radiantibus; foliis pinnatifidis, incis; caule hispido* of Linneus, and its flowers are sometimes used medicinally. The whole plant possesses a bitter and subadstringent taste, and was formerly much employed in the cure of some leprous affections, whence its name, and diseases of the lungs.

SCABIOSA ARVENSIS. The systematic name of the common field scabious. See *Scabiosa*.

SCABIOSA SUCCISA. The systematic name of the devil's bit scabious. See *Morus diabolus*.

SCALA TYMPANI. The superior spiral cavity of the cochlea.

SCALA VESTIBULI. The inferior spiral cavity of the cochlea.

Scald head. See *Tinea capitis*.

SCALE. *Synonyma.* A lamina of morbid cuticle, hard, thickened, whitish, and opaque, of a very small size and irregular, often increasing into layers, denominated crusts. Both scales and crusts repeatedly fall off, and are reproduced in a short time.

SCALENUS. (*Scalenus* *sc. Musculus*; from *scalenus*, irregular, or unequal.) Anatomical writers have differed greatly in their descriptions of this muscle, which is situated at the side of the neck, between the transverse processes of the cervical vertebrae and the upper part of the thorax. The ancients, who gave it its name from its resemblance to an irregular triangle, considered it as one muscle. Vesalius and Winslow divide it into two, Fallopius and Cowper into three, Douglas into four, and Albinus into five portions, which they describe as distinct muscles. Without deviating in the least from anatomical accuracy, it may be considered as one muscle divided into three portions. The anterior portion arises commonly from the trans-

verse processes of the six inferior vertebrae of the neck, by as many short tendons, and descending obliquely outwards, is inserted tendinous and fleshy, into the upper side of the first rib, near its cartilage. The axillary artery passes through this portion, and sometimes divides it into two slips, about an inch and a half above its insertion. The middle portion arises by distinct tendons, from the transverse processes of the four last vertebrae of the neck, and descending obliquely outwards and a little backwards, is inserted tendinous into the outer and upper part of the first rib, from its root to within the distance of an inch from its cartilage. The space between this and the anterior portion, affords a passage to the nerves going to the upper extremities. It is in part covered by the third or posterior portion, which is the thinnest and longest of the three. This arises from the transverse processes of the second, third, fourth, and fifth vertebrae of the neck, by distinct tendons, and is inserted into the upper edge of the second rib, at the distance of about an inch and a half from its articulation, by a broad flat tendon. The use of the scalenus is to move the neck to one side, when it acts singly, or to bend it forwards, when both muscles act; and when the neck is fixed, it serves to elevate the ribs, and dilate the chest.

SCALENUS PRIMUS. See *Scalenus*.

SCALENUS SECUNDUS. See *Scalenus*.

SCALENUS TERTIUS. See *Scalenus*.

SCALPELLUM. A scalpel, or common dissecting knife. A raspatory.

SCALPRUM. A denticular raspatory, or rugine, used in trepanning.

SCAMMONIUM. (A corruption of the Arabian word *chamosah*.) *Convolvulus syriacus*. *Scammonium syriacum*. *Diugrydium*. *Scammony*. The concrete gummi-resinous juice of the *Convolvulus scammonia* of Linneus—*foliis sagittatis postice truncatis, pedunculis teretibus subtifloris*. This plant grows plentifully about Maraash, Antioch, Edlib, and towards Tripoli, in Syria. No part of the dried plant possesses any medicinal quality, but the root, which Dr. Russel administered in decoction, and found it to be a pleasant and mild cathartic. It is from the milky juice of the root that we obtain the official scammony, which is procured in the following manner, by the peasants, who collect it in the beginning of June: having cleared away the earth from about the root, they cut off the top in an oblique direction, about two inches below where the stalks spring from it. Under the most depending part of the slope they fix a shell, or some other convenient receptacle, into which the milky juice gradually flows. It is left there about twelve hours, which time is sufficient for draining off the whole

juice; this, however, is in small quantity, each root affording but a very few drachms. This juice from the several roots is put together, often into the leg of an old boot, for want of some more proper vessel, where, in a little time, it grows hard, and is the genuine scammony. The smell of scammony is rather unpleasant, and the taste bitterish and slightly acrid. The different proportions of gum and resin, of which it consists, have been variously stated, but as proof spirit is the best menstruum for it, these substances are supposed to be nearly in equal parts. It is brought from Aleppo and Smyrna in masses, generally of a light shining grey colour, and friable texture; of rather an unpleasant smell, and bitterish and slightly acrid taste. The scammony of Aleppo is by far the purest. That of Smyrna is ponderous, black, and mixed with extraneous matters. Scammony appears to have been well known to the Greek and Arabian physicians, and was exhibited internally as a purgative, and externally for the itch, tinea, fixed pains, &c. It is seldom given alone, but enters several compounds, which are administered as purgatives.

Scammony. See *Scammonium*.

SCANDIX. The name of a genus of plants in the Linnean system. Class, *Pentandria*. Order, *Digynia*.

SCANDIX CERSEFOLIUM. The systematic name of the official chervil. See *Cereseolum*.

SCANDIX ODORATA. The systematic name of the sweet cicely, which possesses virtues similar to the common chervil. See *Cereseolum*.

SCAPHA. (A skiff, or cock-boat; from *scavo*, to make hollow; because formerly they were made by excavating a large tree.) 1. The excavation or cavity of the auricle, or external ear, between the helix and antihelix.

2. The name of a double-headed roller.

Scaphoid. See *Scaphoides*.

SCAPHOIDES. (From *scapho*, a little vessel, or boat, and *oides*, resemblance.) Boat-like. See *Naviculare os*.

SCAPULA. (From the Hebrew *schapha*.) *Omeplata*. *Oo homoplatea*. *Scapula*. *Epinasion*. The shoulder-blade. This bone, which approaches nearly to a triangular figure, is fixed, not unlike a buckle, to the upper, posterior, and lateral part of the thorax, extending from the first to about the seventh rib. The anterior and internal surface is irregularly concave, from the impression, not of the ribs, as the generality of anatomists have supposed, but of the subscapularis muscle. Its posterior and external surface is convex, and divided into two unequal fossae by a considerable spine, which, rising small from the posterior edge of the scapula, becomes

gradually higher and broader as it approaches the anterior and superior angle of the bone, till at length it terminates in a broad and flat process, at the top of the shoulder, called the *processus acromion*. On the anterior edge of this *processus acromion*, we observe an oblong, concave, articulating surface, covered with cartilage, for the articulation of the scapula with the clavicle. At its lower part, the acromion is hollowed, to allow a passage to the supra and infra spinati muscles. The ridge of the spine affords two rough, flat surfaces, for the insertion of the trapezius and deltoid muscles. Of two the fossae into which the external surface of the bone is divided by the spine, the superior one, which is the smallest, serves to lodge the supra spinatus muscle; and the inferior fossa, which is much larger than the other, gives origin to the infra spinatus. The triangular shape of the scapula leads us to consider its angles and its sides. The upper posterior angle is neither so thick, nor has so rough a surface, as the inferior one; but the most remarkable of the three angles of this bone is the anterior one, which is of great thickness, and formed into a glenoid cavity of an oval shape, the greatest diameter of which is from below upwards. This cavity, in the recent subject, is furnished with cartilage, and receives the head of the os humeri. The cartilaginous crust, which surrounds its brims, makes it appear deeper in the fresh subject than in the skeleton. A little beyond this glenoid cavity, the bone becomes narrower, so as to give the appearance of a neck; and above this rises a considerable process, which, from being thick at its origin, becomes thinner, and, in some degree, flattened at its extremity. This process projects considerably, and is curved downwards. From its supposed resemblance to the beak of a bird, it is called the *coracoid* process. From the whole external side of this process, a strong and broad ligament is stretched to the *processus acromion*, becoming narrower as it approaches the latter process, so as to be of a somewhat triangular shape. This ligament, and the two processes with which it is connected, are evidently intended for the protection of the joint, and to prevent a luxation of the os humeri upwards. Of the three sides of the scapula, the posterior one, which is the longest, is called the *basis*. This side is turned towards the vertebrae. Its other two sides are called *costae*. The superior *costa*, which is the upper and shortest side, is likewise thinner than the other two, having a sharp edge. It is nearly horizontal, and parallel with the second rib; and is interrupted, near the basis of the coracoid process, by a semicircular niche, which is closed by a liga-

ment that extends from one end of it to the other, and affords a passage to vessels and nerves. Besides this passage, there are other niches in the scapula for the transmission of vessels; viz. one between the coracoid process and the head of the bone, and another between its neck and the processus acromion. The third side of the scapula, or the inferior costa, as it is called, is of considerable thickness, and extends obliquely from the neck of the bone to its inferior angle, reaching from about the third to the eighth rib. The scapula has but very little cellular substance, and is of unequal thickness, being very thin at its middle part, where it is covered by a great number of muscles, and having its neck, the acromion, and coracoid process, of considerable strength. In the fetus, the basis and the neck of the scapula, together with its glenoid cavity, acromion, coracoid process, and the ridge of the spine, are so many epiphyses with respect to the rest of the bone, to which they are not completely united till a considerable time after birth. The scapula is articulated to the clavicle and os humeri, to which last it serves as a fulcrum; and, by altering its position, it affords a greater scope to the bones of the arm in their different motions. It likewise affords attachment to a great number of muscles, and posteriorly serves as a defence to the thorax.

SCAPULARIA. (From *scapula*, the shoulder-bone.) A bandage for the shoulder-blade.

SCAPULARIÆ ARTERIÆ. The scapular arteries and veins are branches of the subclavian and axillary.

SCARBOROUGH WATER. A ferruginous spring at Scarborough in Yorkshire. There are two species of chalybeate water found in this spot, and they differ considerably in their composition, though they rise nearly contiguous to each other. The one is a simple carbonated chalybeate, similar to the Tunbridge water: the other, which is better known and more frequent, and more particularly distinguished as Scarborough water, has, in conjunction with the iron, a considerable admixture of a purging salt, which adds much to its value. The diseases in which it is ordered are similar to those in which Cheltenham water is prescribed, only it is necessary to increase the purgative effect of this water by adding similar salts. It is, therefore, chiefly as an alternative that this water can be employed in its natural state.

Scarborough has an advantage belonging to its situation which Cheltenham does not possess, that of affording an opportunity for sea-bathing, the use of which will, in many cases, much assist in the plan of cure for many of the disorders for which the mineral water is resorted to.

Scarf-skin. See *Cuticle* and *Skin*.

SCARIFICATION. (*Scarificatio*, from *scarifico*, to scarify.) A superficial incision made with a lancet, or a surgical instrument called a scarificator, for the purpose of taking away blood, or letting out fluids, &c.

SCARIOLA. See *Lactuca sylvestris*.

SCARIOLA GALLORUM. The lactuca scariola is sometimes so termed. See *Lactuca sylvestris*.

SCARLATINA. (From *scarlatto*, a deep red. Ital.) The scarlet fever, a genus of disease in the class *pyrexia*, and order *exanthemata* of Cullen; characterised by contagious synocha; the fourth day the face swells; a scarlet eruption appears on the skin in patches; which, after three or four days, ends in the desquamation of the cuticle, or is succeeded by anasarca. It has two species: 1. *Scarlatina simplex*, the mild. 2. *Scarlatina cynanchica*, or *anginosa*, with ulcerated sore-throat.

Some have asserted that scarlatina never attacks the same person a second time; more extensive observation has confuted this opinion.

Scarlatina attacks persons of all ages, but children and young persons are most subject to it, and it appears at all seasons of the year; but it is more frequently met with towards the end of autumn, or beginning of winter, than at any other periods, at which time it very often becomes a very prevalent epidemic. It is, beyond all doubt, a very contagious disease.

The one to which it bears the greatest resemblance is the measles; but from this it is readily to be distinguished by the absence of the cough, watery eye, running at the nose and sneezing, which are the predominant symptoms in the early stage of the measles, but which do not usually attend on the scarlatina, or at least in any high degree.

It begins like other fevers, with languor, lassitude, confusion of ideas, chills and shiverings, alternated by fits of heat. The thirst is considerable, the skin dry, and the patient is often incommoded with anxiety, nausea, and vomiting.

Soon after the appearance of these symptoms, a stiffness is perceived in the neck, accompanied with a soreness in the throat, and a difficulty and pain in swallowing. On inspecting the internal fauces, they are found very red, and more or less swelled. On the tonsils, velum pendulum palati and uvula, the parts most affected with inflammation, there generally, not always, appears a number of small whitish or greyish specks or sloughs.

About the third day, the scarlet efflorescence appears on the skin, which seldom produces however any remission of the fever. On the departure of the efflorescence, which usually continues out only for

three or four days, a gentle sweat comes on, the fever subsides, the cuticle or scarf-skin then falls off in small scales, and the patient gradually regains his former strength and health.

On the disappearance of the efflorescence in scarlatina, it is, however, no uncommon occurrence for an anasarous swelling to affect the whole body, but this is usually of a very short continuance.

Scarlatina in several instances approaches very near to the cynanche maligna, and the patient is then seized not only with a coldness and shivering, but likewise with great languor, debility and sickness, succeeded by heat, nausea, vomiting of bilious matter, soreness of the throat, inflammation, and ulceration in the tonsils, a frequent and laborious breathing, and a quick and small depressed pulse. When the efflorescence appears, which is usually on the third day, it brings no relief, on the contrary the symptoms are much aggravated, and fresh ones arise.

In the progress of the disease, one universal redness, unattended however by any pustular eruption, pervades the face, body, and limbs, which parts appear somewhat swollen. The eyes and nostrils partake likewise more or less of the redness, and in proportion as the former have an inflamed appearance, so does the tendency to delirium prevail.

When scarlatina anginosa is to terminate in health, the fiery redness abates gradually, and is succeeded by a brown colour, and the skin becomes rough, peels off in small scales, the tumefaction subsides, and health is gradually restored. On the contrary, when it is to terminate fatally, the febrile symptoms run very high from the first of its attack, the skin is intensely hot and dry, the pulse is very frequent but small, great thirst prevails, the efflorescence makes its appearance on the second day, or sooner, and about the third or fourth is probably interspersed with large livid spots, and a high degree of delirium ensuing, or hæmorrhages breaking out, the patient is cut off about the sixth or eighth day. In some cases a severe purging arises, which never fails to prove fatal. Some again, where the symptoms do not run so high, instead of recovering, as is usual, about the time the skin begins to regain its natural colour, fall into a kind of lingering way, and are carried off in the course of a few weeks.

Scarlatina of itself is not usually attended with danger, although a considerable degree of delirium sometimes prevails for a day or two; but when it partakes much of the nature of cynanche maligna, or degenerates into typhus putrida, which it is apt to do, it often proves fatal. The

same morbid appearances are to be observed on dissection, when it destroys the patient, as in these diseases.

SCARLATINA ANGINOSA. See *Scarlatina*.

SCARLATINA CYNANCHICA. See *Scarlatina*.

SCARLATINA SIMPLEX. See *Scarlatina*.

Scarlet fever. See *Scarlatina*.

Sceleton. See *Skeleton*.

SCELOTYRE. (From *σκαλοτ*, the leg, and *τυρε*, riot, intemperance.) A debility of the legs from scurvey or an intemperate way of life.

SCHEROMA. A dryness of the eye from the want of the lacrymal fluid. The effects of this lacrymal fluid being deficient, the eyes become dry, and in their motions produce a sensation as though sand, or some gritty substances, were between the eye and the eyelid; the vision is obscured, the globe of the eye appears foulish and dull, which is a bad omen in acute diseases. The species are, 1. *Scheroma febrile*, or a dryness of the eyes, which is observed in fevers, complicated with a phlogistic density of the humours. 2. *Scheroma exhaustorum*, which happens after great evacuations, and in persons dying. 3. *Scheroma inflammatorum*, which is a symptom of the ophthalmia sicca. 4. *Scheroma itinerantium*, or the dryness of the eyes, which happens in sandy places to travellers, as in hot Syria, or from dry winds, which dries up the humidity necessary for the motion of the eyes.

SCHIEDACKBON. (From *σχιδάξ*, a splinter.) A longitudinal fracture of the bone.

SCHIEFLAUM. (From *σχινος*, the mastic, and *λαυον*, oil.) Oil of mastic.

SCHNEIDER'S MEMBRANE. The very vascular pituitary membrane of the nose, called Schneiderian, from its discoverer.

SCHONWANTHUS. (From *σχονος*, a rush, and *ανθος*, a flower.) Sweet rush, or camel's hay. See *Juncus odoratus*.

SCHONOLAVRUS. (From *σχονος*, a rush, and *λαγος*, a hare, and *εξα*, a tail; so called from its resemblance to a hare's tail.) Hare's tail. Cotton-grass.

SCIATIC ARTERY. *Arteria sciatica*. Ischiatic artery. A branch of the internal iliac.

SCIATIC NERVE. *Nervus sciaticus*. Ischiatic nerve. A branch of a nerve of the lower extremity, formed by the union of the lumbar and sacral nerves. It is divided near the popliteal cavity into the tibial and peroneal, which are distributed to the leg and foot.

SCIATIC NOTCH. Ischiatic notch. See *Ischium*.

SCIATIC VEIN. *Vena sciatica*. The vein which accompanies the sciatic artery in the thigh.

SCIATICA. *Icthia.* A rheumatic affection of the hip-joint.

SCIATICA CRASSA. The ibexis or cardimantica, raised in gardens for culinary purposes.

SCILLA. (From *scilla*, to dry; so called from its properties of drying up humours.) 1. The name of a genus of plants in the Linnæan system. Class, *Hexandria*. Order, *Monogynia*.

2. The pharmacopœial name of the medicinal squill or sea-onion. *Ornithogalum muritimum.* *Squilla.* *Scilla maritima* of Linnæus:—*nudiflora, bracteis refractis.* A native of Spain, Sicily, and Syria, growing on the sea coast. The red-rooted variety has been supposed to be more efficacious than the white, and is therefore still preferred for medicinal use. The root of the squill, which appears to have been known as a medicine in the early ages of Greece, and has so well maintained its character ever since as to be deservedly in great estimation, and of very frequent use at this time, seems to manifest a poisonous quality to several animals. In proof of this, we have the testimonies of Hillefeld, Bergius, Vogel, and others. Its acrimony is so great, that even if much handled it excoriates the skin, and if given in large doses, and frequently repeated, it not only excites nausea, tormina, and violent vomiting, but it has been known to produce strangury, bloody urine, hypercatharsis, cardialgia, hæmorrhoids, convulsions, with fatal inflammation, and gangrene of the stomach and bowels. But as many of the active articles of the materia medica, by injudicious administration, become equally deleterious, these effects of the scilla do not derogate from its medicinal virtues; on the contrary, we feel ourselves fully warranted, says Dr. Woodville, in representing this drug, under proper management, and in certain cases and constitutions, to be a medicine of great practical utility, and real importance in the cure of many obstinate diseases. Its effects, as stated by Bergius, are incitens, diuretica, emetica, subpurgans, hydragoga, expectorans, emmenagoga. In hydropsical cases it has long been esteemed the most certain and effectual diuretic with which we are acquainted; and in asthmatic affections, or dyspnœa, occasioned by the lodgment of tenaceous phlegm, it has been the expectorant usually employed. The squill, especially in large doses, is apt to stimulate the stomach, and to prove emetic; and it sometimes acts on the intestines, and becomes purgative; but when these operations take place, the medicine is prevented from reaching the blood vessels and kidneys, and the patient is deprived of its diuretic effects, which are to be obtained by giving the squill in smaller doses, repeated at more distant intervals,

or by the joining of an opiate to this medicine, which was found by Dr. Cullen to answer the same purpose. The Doctor further observes, that from a continued repetition of the squill, the dose may be gradually increased, and the interval of its exhibitions shortened; and when in this way the dose comes to be tolerably large, the opiate may be most conveniently employed to direct the operation of the squill more certainly to the kidneys. "In cases of dropsy, that is, when there is an effusion of water into the cavities, and therefore that less water goes to the kidneys, we are of opinion that neutral salt, accompanying the squill, may be of use in determining this fluid more certainly to the kidneys; and whenever it can be perceived that it take this course, we are persuaded that it will be always useful, and generally safe, during the exhibition of the squills, to increase the usual quantity of drink."

The diuretic effects of squills have been supposed to be promoted by the addition of some mercurial; and the less purgative preparations of mercury, in the opinion of Dr. Cullen, are best adapted to this purpose; he therefore recommends a solution of corrosive sublimate, as being more proper than any other, because most diuretic. Where the primæ viæ abound with mucous matter, and the lungs are oppressed with viscid phlegm, this medicine is likewise in general estimation.

As an expectorant, the squill may be supposed not only to attenuate the mucous follicles to excite a more copious excretion of it from the lungs, and thereby lessen the congestion, upon which the difficulty of respiration very generally depends. Therefore in all pulmonic affections, excepting only those of actual or violent inflammation, ulcer, and spasm, the squill has been experienced to be an useful medicine. The official preparations of squills are, a conserve, dried squills, a syrup, and vinegar, an oxymel, and pills. Practitioners have not, however, confined themselves to these. When this root was intended as a diuretic, it has most commonly been used in powder, as being in this state less disposed to nauseate the stomach; and to the powder it has been the practice to add neutral salts, as nitre, or crystals of tartar, especially if the patient complained of much thirst; others recommend calomel; and with a view to render the squills less offensive to the stomach, it has been usual to conjoin an aromatic. The dose of dried squills is from two to four or six grains once a day, or half this quantity twice a day; afterwards to be regulated according to its effects. The dose of the other preparations of this drug, when fresh, should be four times this weight; for this root loses in the process of drying four

fifths of its original weight, and this loss is merely a watery exhalation.

SCILLA EXSICCATA. Dried squill.

SCILLA HISPANICA. The Spanish squill.

SCILLA MARITIMA. The systematic name of the official squill. See *Scilla*.

SCILLA ACETUM. Squills macerated in vinegar.

SCILLA CONSERVA. Squills beat up with sugar.

SCILLA MEL. Tincture of squills boiled with honey.

SCILLA OXYMEL. Vinegar of squills boiled with honey.

SCILLA PILULE. Squill pills. Dried squills, ginger, soap, and ammoniacum.

SCILLA TINCTURA. Squills digested in spirit of wine.

SCILLITES. (From *σκίλλα*, the squill.) A wine impregnated with squills.

SCINCUS. (From *sheque*, Heb.) The skink. This amphibious animal is of the lizard kind, and caught about the Nile, and thence brought dried into this country, remarkably smooth and glossy, as if varnished. The flesh of the animal, particularly of the belly, has been said to be diuretic, alexipharmic, aphrodisiac, and useful in leprous disorders.

SCIRRHOMA. (From *σκίρρομα*, to harden.) *Scirrhomia*. A hard tumour. See *Scirrhus*.

SCIRRHUS. (From *σκίρρομα*, to harden.) A genus of disease in the class *localis*, and order *tumores* of Cullen; known by a hard tumour of a glandular part, indolent, and not readily suppurating. The following observations of Mr. Pearson are deserving of attention. A scirrhus, he says, is usually defined to be a hard, and almost insensible tumour, commonly situated in a glandular part, and accompanied with little or no discolouration of the surface of the skin. This description agrees with the true or exquisite scirrhus; but when it has proceeded from the indolent to the malignant state, the tumour is then unequal in its figure, it becomes painful, the skin acquires a purple or livid hue, and the cutaneous veins are often varicose. Let us now examine whether this enumeration of symptoms be sufficiently accurate for practical purposes.

It is probable, that any gland in the living body may be the seat of a cancerous disease; but it appears more frequently as an idiopathic affection in those glands that form the several secretions than in the absorbent glands: and of the secreting organs, those which separate fluids that are to be employed in the animal economy, suffer much oftener than the glands which secrete the excrementitious parts of the blood. Indeed, it may be doubted whether an absorbent gland be ever the primary seat of a true scirrhus. Daily experience evinces, that these glands may suffer contamination from

their connexion with a cancerous part, but under such circumstances, this morbid alteration being the effect of a disease in that neighbouring part, it ought to be regarded as a secondary or consequent affection. I never yet met with an unequivocal proof of a primary scirrhus in an absorbent gland; and if a larger experience shall confirm this observation, and establish it as a general rule, it will afford material assistance in forming the diagnosis of this disease. The general term scirrhus hath been applied, with too little discrimination, to indurated tumours of lymphatic glands. When these appendages of the absorbent system enlarge in the early part of life, the disease is commonly treated as strumous; but as a similar alteration of these parts may, and often does occur at a more advanced period, there ought to be some very good reasons for ascribing malignity to one rather than the other. In old people the tumour is indeed often larger, more indurated, and less tractable than in children; but when the alteration originated in the lymphatic glands, it will very rarely be found to possess any thing cancerous in its nature.

If every other morbid alteration in a part were attended with pain and softness, then induration and defective sensibility might point out the presence of a scirrhus. But this is so far from being the case, that even encysted tumours, at their commencement, frequently excite the sensation of impenetrable hardness. All glands are contained in capsulæ, not very elastic, so that almost every species of chronic enlargement of these bodies must be hard; hence this induration is rather owing to the structure of the part, than to the peculiar nature of the disease; and as glands in their healthy state are not endowed with much sensibility, every disease that gradually produces induration, will rather diminish than increase their perceptive powers. Induration and insensibility may therefore prove that the affected part does not labour under an acute disease; but these symptoms alone can yield no certain information concerning the true nature of the morbid alteration. Those indolent affections of the glands that so frequently appear after the meridian of life, commonly manifest a hardness and want of sensation, not inferior to that which accompanies a true scirrhus; and yet these tumours will often admit of a cure by the same mode of treatment which we find to be successful in scrophula; and when they prove unobquerable by the powers of medicine, we generally see them continue stationary and innocent to the latest period of life. Writers have indeed said much about certain tumours changing their nature, and assuming a new character; but I strongly suspect that the doctrine of the mutation of

diseases into each other, stands upon a very uncertain foundation. Improper treatment may, without doubt, exasperate diseases, and render a complaint which appeared to be mild and tractable, dangerous, or destructive; but to aggravate the symptoms, and to change the form of the disease, are things that ought not to be confounded. I do not affirm, that a breast which has been the seat of a mammary abscess, or a gland that has been affected with scrophula, may not become cancerous; for they might have suffered from this disease had no previous complaint existed; but these morbid alterations generate no greater tendency to cancer than if the parts had always retained their natural condition. There is no necessary connexion between the cancer and any other disease, nor has it ever been clearly proved that one is convertible into the other.

Chirurgical writers have generally enumerated tumour as an essential symptom of the scirrhus; and it is very true, that this disease is often accompanied with an increase of bulk in the part affected. From long and careful observation, I am however induced to think, that an addition to the quantity of matter is rather an accidental than a necessary consequence of the presence of this peculiar affection.

When the breast is the seat of a scirrhus, the altered part is hard, perhaps unequal in its figure, and definite; but these symptoms are not always connected with an actual increase in the dimensions of the breast. On the contrary, the true scirrhus is frequently accompanied with a contraction and diminution of bulk, a retraction of the nipple, and a puckered state of the skin.

The irritation produced by an indurated substance lying in the breast, will very often cause a determination of blood to that organ, and a consequent enlargement of it; but I consider this as an inflammatory state of the surrounding parts, excited by the scirrhus, acting as a remote cause, and by no means essential to the original complaint. From the evicent utility of topical blood-letting under these circumstances, a notion has prevailed that the scirrhus is an inflammatory disease; but the strongly-marked dissimilarity of a phlegmon and an exquisite scirrhus, in their appearances, progress, and mode of termination, obliges me to dissent from that opinion. That one portion of the breast may be in a scirrhus state, while the other parts are in a state of inflammation, is agreeable to reason and experience; but that an inflammation, which is an acute disease, and a scirrhus, whose essential characters are almost directly the reverse of inflammation, shall be co-existent in the same part, is not a very intelligible proposition. Tumor and inflamma-

tion are commonly met with on a variety of other occasions, and in this particular instance they may be the effects of the disease, but are not essentially connected with its presence.

An incipient scirrhus is seldom accompanied with a discolouration of the skin; and a dusky redness, purple, or even livid appearance of the surface, is commonly seen when there is a malignant scirrhus. The presence or absence of colour can, however, at the best, afford us but a very precarious criterion of the true nature of the complaint. When the disease is clearly known, an altered state of the skin may assist us in judging of the progress it has made; but as the skin may suffer similar variations in a number of very dissimilar diseases, it would be improper to found an opinion upon so delusive a phenomenon.

SCLAREA. (From *σκληρος*, hard; because its stalks are hard and dry, Blanch.) The garden clary. See *Salvia sclarea*.

SCLAREA HISPANICA. Wild clary, or horn-minum sylvestre.

SCLERIASIS. (From *σκληρον*, to harden.) *Scleroma*. *Sclerosis*. A hard tumour or induration.

SCLEROPHTHALMIA. (From *σκληρος*, hard, and *ὀφθαλμος*, the eye.) A protrusion of the eye-ball. An inflammation of the eye, attended with hardness of the parts.

SCLEROSARCOMA. (From *σκληρος*, hard, and *σάρκωμα*, a fleshy tumour.) A hard fleshy excrescence on the gums.

SCLEROSIS. See *Scleriosis*.

SCLEROTIC COAT. (*Tunica sclerotica*; from *σκληρον*, to harden; so called from its hardness.) *Sclerotic*. The outermost coat of the eye, of a white colour, dense, and tenacious. Its anterior part, which is transparent, is termed the *cornea transparens*. It is into this coat of the eye that the muscles of the bulb are inserted.

SCLEROTIS. See *Sclerotic coat*.

SCLOPETARIA AQUA. (From *sclopetum*, a gun; so called from its supposed virtues in healing gun-shot wounds.) Arquebuseade. It is made of sage, mugwort, and mint, distilled in wine.

SCLOPETOPLAGA. (From *sclopetum*, a gun, and *plaga*, a wound.) A gun-shot wound.

SCOLIASIS. (From *σκολις*, to twist.) A distortion of the spine.

SCOLOPENDRIA. The spleenwort or miltwaste is sometimes so called. See *Ceterach*.

SCOLOPENDRIUM. (From *σκολοπενδρα*, the earwig; so called because its leaves resemble the earwig.) *Phyllitis*. *Lingua cervina*. Harts-tongue. This indigenous plant as *Asplenium scolopendrium* of Linnæus:—*frondibus simplicibus cordato lingularis integerrimis stilibus hirsutis*, grows on most shady banks, walls, &c. It has a slightly adstring-

gent and mucilaginous sweetish taste. When fresh and rubbed, it imparts a disagreeable smell. Harts-tongue, and the *five capillary* herbs, of which it is one, was formerly much used to strengthen the viscera, restrain hæmorrhages and alvine fluxes, and to open obstructions of the liver and spleen, and for the general purposes of demulcents and pectorals.

SCOLOPHACHÆRIUM. (From *σκολοψα*, the woodcock, and *μαχαίρα*, a knife; so called because it is bent a little at the end like a woodcock's bill.) An incision-knife.

SCOLYMUS. (From *σκολος*, a thorn; so named from its prickly leaves.) The artichoke is sometimes so called. See *Ci-nara*.

SCOPA REGIA. The butcher's broom, or knee-holly, was formerly so termed. See *Ruscus*.

SCORBUTIA. (From *scorbutus*, the scurvy.) Medicines for the scurvy.

SCORBUTUS. (From *schorboot*, Germ.) *Gingibrachium*, because the gums and arms, and *gingripedium*, because the gums and legs, are affected by it. The scurvy. A genus of disease in the class *cachexia*, and order *impetiginos* of Cullen; characterised by extreme debility; complexion pale and bloated; spongy gums; livid spots on the skin; breath offensive; oedematous swellings in the legs; hæmorrhages; foul ulcers; fetid urine; and extremely offensive stools. The scurvy is a disease of a putrid nature, much more prevalent in cold climates than in warm ones, and which chiefly affects sailors, and such as are shut up in besieged places, owing, as is supposed, to their being deprived of fresh provisions, and a due quantity of acescent food, assisted by the prevalence of cold and moisture, and by such other causes as depress the nervous energy, as indolence, confinement, want of exercise, neglect of cleanliness, much labour and fatigue, sadness, despondency, &c. These several debilitating causes, with the concurrence of a diet consisting principally of salted or putrescent food, will be sure to produce this disease. It seems, however, to depend more on a defect of nourishment, than on a vitiated state; and the reason that salted provisions are so productive of the scurvy, is, most probably, because they are drained of their nutritious juices, which are extracted and run off in brine. As the disease is apt to become pretty general amongst the crew of a ship when it has once made its appearance, it has been supposed by many to be of a contagious nature; but the conjecture seems by no means well founded.

A preternatural saline state of the blood has been assigned as its proximate cause. It has been contended by some physicians, that the primary morbid affection in this

disease is a debilitated state of the solids, arising principally from the want of aliment.

The scurvy comes on gradually, with heaviness, weariness, and unwillingness to move about, together with dejection of spirits, considerable loss of strength, and debility. As it advances in its progress, the countenance becomes sallow and bloated, respiration is hurried on the least motion, the teeth become loose, the gums are spongy, the breath is very offensive, livid spots appear on different parts of the body, old wounds which have been long healed up break out afresh, severe wandering pains are felt, particularly by night, the skin is dry, the urine small in quantity, turning blue vegetable infusions of a green colour; and the pulse is small, frequent, and, towards the last, intermitting; but the intellects are, for the most part, clear and distinct.

By an aggravation of the symptoms, the disease, in its last stage, exhibits a most wretched appearance. The joints become swelled and stiff, the tendons of the legs are rigid and contracted, general emaciation ensues, hæmorrhages break forth from different parts, fetid evacuations are discharged by stool, and a diarrhoea or dysentery arises, which soon terminates the tragic scene.

Scurvy, as usually met with on shore, or where the person has not been exposed to the influence of the remote causes before enumerated, is unattended by any violent symptoms, as slight blotches, with scaly eruptions on different parts of the body, and a sponginess of the gums, are the chief ones to be observed.

In forming our judgment as to the event of the disease, we are to be directed by the violence of the symptoms, by the situation of the patient with respect to a vegetable diet, or other proper substitutes, by his former state of health, and by his constitution not having been impaired by previous diseases.

Dissections of scurvy have always discovered the blood to be in a very dissolved state. The thorax usually contains more or less of a watery fluid, which, in many cases, possesses so high a degree of acrimony, as to excoriate the hands by coming in contact with it; the cavity of the abdomen contains the same kind of fluid; the lungs are black and putrid; and the heart itself has been found in a similar state, with its cavity filled with a corrupted fluid. In many instances, the epiphyses have been found divided from the bones, the cartilages separated from the ribs, and several of the bones themselves dissolved by caries. The brain seldom shews any marks of disease.

SCORODON. (From *σκορδον*, garlic; so called because it smells like garlic.) Fra-

cago palustris. *Chamedrys palustris allium redolens*. Water germander. *Teucrium scordium* of Linnaeus. The leaves of this plant have a smell somewhat of the garlic kind, from which circumstance it is supposed to take its name: to the taste they are bitterish and slightly pungent. The plant was formerly in high estimation, but is now justly fallen into disuse, although recommended by some in antiseptic cataplasms and fomentations.

SCORIE. (*Scoria*, from *σκαπ*, excrement.) **Dross.** The refuse or useless parts of any substance.

SCORODOPHABUM. (From *σκαποδον*, garlic, and *πρασον*, the leek.) The wild garlic or leek snailot.

SCORODUM. (*Αρω του σκαυ οζου*, from its filthy smell.) Garlic.

SCORPIACA. (From *σκαρπιος*, a scorpion.) Medicines against the bite of serpents.

SCORPIOIDES. (From *σκαρπιος*, a scorpion, and *ιδιος*, a likeness; so called because its leaves resemble the tail of a scorpion.) **Scorpioides.** Resembling the scorpion. **Scorpion-wort.** Bird's foot.

SCORPIURUS. See *Scorpioides*.

SCORZONERA. (From *escorza*, a serpent, Span.; so called because it is said to be effectual against the bite of all venomous animals.) 1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia equalis*.

2. The pharmacopœial name of the plant called also *escorzenera*. *Viperaria*. *Serpentaria hispanica*. Goats' grass. *Vipers' grass*. The roots of this plant, *Scorzonera humilis* of Linnaeus:—*caule fulvulo, unifloro; foliis lato-lanceolatis, nervosis, planis*, has been sometimes employed medicinally as alexipharmics, and in hypochondriacal disorders, and obstructions of the viscera. The *Scorzonera hispanica* mostly supplies the shops, whose root is esculent, oleraceous, and against diseases inefficacious.

SCORZONERA HISPANICA. The systematic name of the esculent vipers' grass.

SCORZONERA HUMILIS. The systematic name of the official vipers' grass. See *Scorzonera*.

SCOTODINE. (From *σκοτος*, darkness, and *δινος*, a giddiness.) *Scotodinia*. *Scotodinos*. *Scotoma*. *Scotomia*. Giddiness with impaired sight.

SCROBICULUS CORDIS. (Dim. of *scroba*, a ditch.) The pit of the stomach.

SCROFULA. (From *scrofa*, a swine; because this animal is said to be much subject to a similar disorder.) *Scrophula*. *Struma*. *Coiras*. *Charas*. *Ecruelles*, Fr. *Scrophula*. The king's evil. A genus of disease in the class *cachexia*, and order *impetigines* of Cullen. He distinguishes four species. 1. *Scrophula vulgaris*, when it is without other disorders external and permanent. 2. *Scrophula mesenterica*, when internal, with loss of appetite, pale coun-

tenance, swelling of the belly, and an unusual fecor of the excrements. 3. *Scrophula fugax*. This is of the most simple kind; it is seated only about the neck, and for the most part is caused by the resorption from sores on the head. 4. *Scrophula Americana*, when it is joined with the yaws. *Scrophula* consists in hard indolent tumours of the conglobate glands in various parts of the body; but particularly in the neck, behind the ears, and under the chin, which after a time suppurate and degenerate into ulcers, from which, instead of pus, a white curdled matter, somewhat resembling the coagulum of milk, is discharged.

The first appearance of the disease is most usually between the third and seventh year of the child's age; but it may arise at any period between these and the age of puberty; after which it seldom makes its first attack. It most commonly affects children of a lax habit, with smooth fine skins, fair hair, and rosy cheeks. It likewise is apt to attack such children as shew a disposition to rachitis, and marked by a protuberant forehead, enlarged joints, and a tumid abdomen. Like this disease, it seems to be peculiar to cold and variable climates, being rarely met with in warm ones. *Scrophula* is by no means a contagious disease; but, beyond all doubt, is of an hereditary nature, and is often entailed by parents on their children. There are, indeed, some practitioners who wholly deny that this, or any other disease, can be acquired by an hereditary right; but that a peculiar temperament of body, or predisposition in the constitution to some diseases, may extend from both father and mother to their offspring, is, observes Dr. Thomas, very clearly proved. For example, we very frequently meet with gout in young persons of both sexes, who could never have brought it on by intemperance, sensuality, or improper diet, but must have acquired the predisposition to it in this way.

Where there is any predisposition in the constitution to scrophula, and the person happens to contract a venereal taint, this frequently excites into action the causes of the former; as a venereal bubo not unfrequently becomes scrophulous, as soon as the virus is destroyed by mercury. The late Dr. Cullen supposed scrophula to depend upon a peculiar constitution of the lymphatic system. The attacks of the disease seem much affected or influenced by the periods of the seasons. They begin usually some time in the winter and spring, and often disappear, or are greatly amended, in summer and autumn. The first appearance of the disorder is commonly in that of small oval or spherical tumours under the skin, unattended by any pain or discolouration. These appear, in general,

upon the sides of the neck, below the ear, or under the chin; but, in some cases, the joints of the elbows or ankles, or those of the fingers and toes, are the parts first affected. In these instances, we do not, however, find small moveable swellings; but, on the contrary, a tumor almost uniformly surrounding the joint, and interrupting its motion.

After some length of time the tumors become larger and more fixed, the skin which covers them acquires a purple or livid colour, and, being much inflamed, they at last suppurate and break into little holes, from which, at first, a matter somewhat puriform oozes out; but this changes by degrees into a kind of viscid serous discharge, much intermixed with small pieces of a white substance, resembling the curd of milk.

The tumors subside gradually, whilst the ulcers at the same time open more, and spread unequally in various directions. After a time, some of the ulcers heal; but other tumors quickly form in different parts of the body, and proceed on, in the same slow manner as the former ones, to suppuration. In this manner the disease goes on for some years, and appearing at last to have exhausted itself, all the ulcers heal up, without being succeeded by any fresh swellings; but leaving behind them an ugly puckering of the skin, and a scar of considerable extent. This is the most mild form under which scrophula ever appears. In more virulent cases, the eyes are particularly the seat of the disease, and are affected with ophthalmia, giving rise to ulcerations in the tarsi, and inflammation of the tunica adnata, terminating not unfrequently in an opacity of the transparent cornea.

In similar cases, the joints become affected, they swell and are incommoded by excruciating deep-seated pain, which is much increased upon the slightest motion. The swelling and pain continue to increase, the muscles of the limb become at length much wasted. Matter is soon afterwards formed, and this is discharged at small openings made by the bursting of the skin. Being, however, of a peculiar acrimonious nature, it erodes the ligaments and cartilages, and produces a caries of the neighbouring bones. By an absorption of the matter into the system, hectic fever at last arises, and, in the end, often proves fatal.

When scrophula is confined to the external surface, it is by no means attended with danger, although on leaving one part, it is apt to be renewed in others; but when the ulcers are imbued with a sharp acrimony, spread, erode, and become deep, without shewing any disposition to heal; when deep-seated collections of matter form amongst the small bones of the hands and feet, or in the joints, or tubercles in the

lungs, with hectic fever, arise, the consequences will be fatal.

On opening the bodies of persons who have died of this disease, many of the viscera are usually found in a diseased state, but more particularly the glands of the mesentery, which are not only much tumefied, but often ulcerated. The lungs are frequently discovered bent, with a number of tubercles or cysts, which contain matter of various kinds. Scrophulous glands, on being examined by dissection, feel somewhat softer to the touch than in their natural state, and when laid open, they are usually found to contain a soft curdy matter, mixed with pus.

SCROPHULARIA. (From *scrofula*, the king's evil; so called from the unequal tubercles upon its roots, like scrophulous tumors.) The name of a genus of plants in the Linnæan system. Class, *Didymia*. Order, *Angiosperma*. The fig-wort.

SCROPHULARIA AQUATICA. See *Betonica aquatica*.

SCROPHULARIA MINOR. The pile-wort is sometimes so called. See *Chetidonium minus*.

SCROPHULARIA NODOSA. The systematic name of the fig-wort. See *Scrophularia vulgaris*.

SCROPHULARIA VULGARIS. *Millemorbis*. *Scrophularia*. Common fig-wort or kernel-wort. The root and leaves of this plant, *Scrophularia nodosa* of Linnæus:—*foliis cordatis, trinerviatis; caule obtusangulo*, have been celebrated both as an internal and external remedy against inflammations, the piles, scrophulous tumors, and old ulcers; but they are now only used in this country by the country people.

SCROTAL HERNIA. *Hernia scrotalis*. *Scrotocoele*. A protrusion of any part of an abdominal viscus or viscera into the scrotum. See *Hernia*.

SCROTOSKLE. (From *scrotum*, and *skala*, a tumour.) A rupture or hernia in the scrotum.

SCROTUM. (*Quasi mortuum*, a skin or hide.) *Bursa testium*. *Oscheus*. *Oscheon*. *Orchea* of Galen. The common integuments which cover the testicles.

SCRUPULUS. (Dim. of *scrupus*, a small stone.) A scruple or weight of 20 grains.

SCURF. *Furfura*. Small exfoliations of the cuticle, which take place after some eruptions on the skin, a new cuticle being formed underneath during the exfoliation.

Scurvy. See *Scorbutus*.

Scurvy-grass. See *Cochlearia hortenensis*.

Scurvy-grass, lemon. See *Cochlearia hortenensis*.

Scurvy-grass, Scotch. See *Brassica maritima*.

SCUTIFORM CARTILAGE. See *Thyroid cartilage*.

SCUTELLARIA GALERICULATA. The systematic name of the skull-cap. See *Ternaria*.

SCYBALA. *Συκάλα.* Dry hard excrements.

SCYTHICUS. (From *Scythia*, its native soil.) An epithet of the liquorice-root or any thing brought from *Scythia*.

SEA-AIR. Is prescribed in a variety of complaints, being considered as more medicinal and salubrious than that on land, it is supposed to possess in its composition a greater quantity of oxygen. This is a most powerful and valuable remedy. It is resorted to with the happiest success against most cases of debility, and particularly against scrophulous diseases affecting the external parts of the body. See *Bath, cold*.

Sea-holly. See *Eryngium*.

Sea-moss. See *Corallina*.

Sea-oak. See *Quercus marina*.

Sea-onion. See *Scilla*.

Sea-salt. See *Murias soda*.

SEA-SICKNESS. A nausea or tendency to vomit, which varies, in respect of duration, in different persons upon their first going to sea. With some it continues only for a day or two; while with others it remains throughout the voyage. The diseases in which sea-sickness is principally recommended are asthma and consumption.

SEA-WATER. This is arranged amongst the simple saline waters. Its chemical analysis gives a proportion of one of saline contents to about twenty-three one-fourth of water; but on our shores it is not greater than one of salt to about thirty of water. Sea-water on the British coast may, therefore, be calculated to contain in the wine pint, of muriated soda 186.5 grains, of muriated magnesia fifty-one, of selenite six grains; total two hundred and forty-three one-half grains, or half an ounce and three and one-half grains of saline contents. The disorders for which the internal use of sea-water has been and may be resorted to, are in general the same for which all the simple saline waters may be used. The peculiar power of sea-water and sea-salt as a discutient, employed either internally or externally in scrophulous habits, is well known, and is attended with considerable advantage when judiciously applied.

Sea-wrack. See *Quercus marina*.

Sealed earths. See *Terra sagillata*.

SEARCHING. The operation of introducing a metallic instrument through the urethra into the bladder, for the purpose of ascertaining whether the patient has stone or not.

SEBACEOUS GLANDS. (*Glandula sebacea*, from *sebum*, suet.) Glands which secrete a sebaceous or suetty humour.

SEBADIALLA. See *Cevadilla*.

SEBATE. (From *sebum*, suet.) *Sebas*. The name in the new chemistry of every compound of the acid of fat.

SEBESTEN. (An Egyptian word.) *Sebestina* *Sebesten*. The dark black fruit of the *cordia myxa*; *foliis ovatis, subra glabris; corymbis lateralibus; calycibus decemstriatis* of Linnæus. It possesses glutinous and aperient qualities, and is exhibited in form of decoction in various diseases of the chest, hoarseness, cough, difficult respiration, &c.

SECALE. 1. The name of a genus of plants in the Linnæan system. Class, *Triandria*. Order, *Digynia*. Rye.

2. The common name of the seed of the *Secale cereale* of Linnæus. It is principally used as an article of diet, and in the northern countries of Europe is employed for affording an ardent spirit.

SECALE CEREALE. The systematic name of the rye-plant. See *Secule*.

SECONDARY. This term in general denotes something that acts as second or in subordination to another. Thus in diseases, we have *secondary symptoms*.

SECONDARY FEVER. That febrile affection which arises after a crisis, or the discharge of some morbid matter, as after the declension of the small-pox or the measles.

SECRETION. The word secretion is used to express that function by which an organ separates from the blood the constituent parts of a fluid that does not exist in it with its characteristic properties.

The difference of secreted humours is visibly connected with those of the organs employed for their formation. Thus arterial exhalation, that take place throughout the whole extent of internal surfaces, and preserves their contiguity, affords nothing but an albuminous serosity, which is only the serum of the blood slightly altered by the weak action of organization very little complicated. The analysis of the water in dropsy, which is merely in the serosity that continually transudes the surface of serous membranes, as the pleura or peritonæum, has demonstrated that this fluid has the greatest resemblance to the serum of the blood, and is only distinguished from it, by the variable proportions of albumen, and the different salts it contains in solution.

This first kind of secretion, this perspiratory transudation, would seem, then, to be a simple filtration or percolation of a liquor already formed in the blood through the porous parts of arteries; yet we must here acknowledge a peculiar action of membranes, the surface of which it perpetually lubricates; without this action the serum would remain united to the other constituent of the blood. This kind of action is termed exhalation. The distin-

guishable character of this kind of secretion is the absence of any mediate structure between the *vas efferens* and the excretory duct: the minute arteries and veins that run into the structure of membranes also constitute both.

After serous transudation, requiring only a very simple organisation, follows the secretion by cryptæ, glandular follicles, and mucous lacunæ. Each of these small glands contained in the texture of membranes lining the internal surface of the digestive, arterial, and urinary passages, and which, when conglomerated, form amygdalæ, &c. may be compared to a small bottle, the bottom of which is round, and the neck short; the membranous parietes of these vesicular cryptæ are supplied with a great quantity of vessels and nerves. It is to the peculiar action of these parietes that the secretion of mucus by these glands should be attributed. These mucous liquids are less fluid and more viscid than the fluid produced by the first kind of secretion, but contain more albumen and salts; they are more different from the serum of the blood, and of a more excrementitious nature; the nature of these bottle-kind of glands is turned towards the parts to which the mucous membrane adheres, their mouth or neck opens on the surface contiguous to these membranes.

The apertures by which the mucous glands discharge themselves are easy to be perceived on the amygdalæ, mucous surface of the urethra, rectum, &c.

Secretion and excretion are facilitated by the irritation occasioned by the presence of air, aliment, or urine; by the compression induced by them, and, finally, by the peristaltic contractions of the muscular fibres to which mucous membranes adhere throughout the whole extent of the *primæ viæ*.

The fluids, which are considerably different from the blood, require for their secretion organs of a more complicated structure; these are called conglomerate glands; to distinguish them from lymphatic glands, which are named conglobate. These glands are visceral masses, formed by an assemblage of nerves and every species of vessels, disposed in packets, and united by cellular structure; a proper membrane, or an elongation of that which lines the cavity that includes them, surrounds their external surface, and separates them from the circumjacent parts.

The arteries do not form an immediate continuation with their excretory ducts, as Rysch affirmed; nor do there exist immediate glands between these vessels as Malpighi believed; it seems more probable that each gland has its cellular or parenchymatous substance in the areolæ; into which the arteries pour the materials of the fluid they prepare; in consequence of a

power peculiar to them, and which forms their distinguishing character. Lymphatics and excretory ducts arise from the sides of these little cells, and both these species of vessels absorb; one attracts the secreted liquor, carrying it into receptacles, where it accumulates, while the other receives that part which the action of the organ could not completely elaborate, or the residue of the secretion.

The nerves which always enter more or less into the structure of secretory organs, and come principally from the great sympathetics, terminate variously in their substance, and furnish each of them with a particular sensibility, by means of which they distinguish in the blood brought thither by the vessels the constituent parts or materials of the humour they are destined to prepare, and select it by a real preference. Besides they cause them to take on a peculiar mode of activity, the exercise of which causes these separate elements to undergo a certain composition, and impresses the fluid produced with specific properties, always relative to the mode of action of which they are the result. Thus the liver retains the constituent principles of bile contained in the blood of the *vena portæ*, elaborates, combines them, and forms the bile, and animal fluid, distinguishable by certain characteristic properties that are subject to variations according as the blood contains the elements which enter into its composition in a greater or less degree; according to the increased or diminished disposition of the gland to retain them, and to effect a more or less complete mixture of them. The qualities of the bile dependent on the concurrence of all these circumstances, should present so many differences as the blood contains principles, and as the hepatic organ may offer varieties relative to the composition of the former, and degree of activity of the latter. Hence arise alterations of the bile, the most inconsiderable of which being compatible with health, escape observation, while those which are more complete, and derange the natural order of the functions, become evident by diseases of which they may be sometimes considered the effect, and at other times the cause. These alterations of the bile (and what is here said of the secretion of this humour may extend to almost every other secretion of the animal economy) never extend so far as to prevent it from being distinguished; it always preserves in a greater or less degree its essential and primitive characters, it never acquires the qualities of another liquor so as to resemble serum, urine, or saliva, &c.

The action of secretory glands is not continual, most of them are subject to the alternate state of action and rest, all, as Barden observed, are asleep or awake when any irritation operates on them, or

in their vicinity, and determines their immediate or sympathetic action. Thus saliva is secreted in greater quantity during mastication, &c.

When a secretory organ enters into action, the surrounding parts, or such as are situated in its vicinity, for instance, the liver, is comprised in the sphere of action of the duodenum, since the repletion of this intestine irritates it, determines a more abundant afflux of humours, and a more copious secretion of bile.

The blood conveyed to a secretory gland, before it arrives there, suffers preparatory changes which dispose it to furnish the constituents of the liquor about to be secreted.

The celerity with which the blood arrives at an organ; the length, diameter, angles of the vessels, and the disposition of their ultimate ramifications, are all circumstances which ought to be observed in the examination of each secretion, since they have an influence on the nature of the secreted fluid, and on the mode in which secretion is effected. When a gland is irritated it becomes the centre of fluxion, and acts on the blood brought by its vessels. Secretion dependent on a peculiar action inherent in a glandular organ is assisted by the action of surrounding muscles.

The glands, after having remained for a longer or shorter time in a state of excitation, relax, become collapsed, and fluids are not conveyed to them in such abundance, they remain in a state of sleep, and during repose renew their sensibility, which is consumed by long exertion.

A remarkable circumstance in secretions is, that they mutually replace and supply each other, so that when the urine is less copious perspiration is more abundant. A sudden coldness of the skin frequently occasions diarrhoeas, the humours are immediately repelled towards the intestinal tube, and pass off by the mucous glands of the intestines, the action of which is considerably increased.

SECRUM ACRI. The systematic name of the stone-cross. See *Illecebra*.

SECTIO CESAREA. See *Cesarian operation*.

SECTIO FRONICA. *Sectio hypogastrica.* The high operation for the stone. See *Lithotomy*.

SECUNDINES. The placenta and membranes which are expanded from its edge, and which form a complete involucre of the fetus and its waters, go under the common term of after-birth, or secundines.

The membranes of the ovum have usually been mentioned as two, the amnion and the chorion; and the latter has again been divided into the true and the false.

The third membrane, which, from its appearance, has likewise been called the vilous or spongy, and from the consideration of it as the inner lamina of the uterus, is cast off like the exuviz of some animals. The decidua has been described by Harvey, not as one of the membranes of the ovum, but as a production of the uterus. The following is the order of the membranes of the ovum, at the full period of gestation: 1st, There is the outer or connecting, which is flacculent, spongy, and extremely vascular, completely investing the whole ovum, and lining the uterus. 2dly, The middle membrane, which is nearly pellucid, with a very few small blood-vessels scattered over it, and which forms a covering to the placenta and funis, but does not pass between the placenta and uterus. 3dly, The inner membrane, which is transparent, of a firmer texture than the others, and lines the whole ovum, making, like the middle membrane, a covering for the placenta and funis with the two last. The ovum is clothed when it passes from the ovarium into the uterus, where the first is provided for its reception.

These membranes, in the advanced state of pregnancy, cohere slightly to each other, though, in some ova, there is a considerable quantity of fluid collected between them, which being discharged when one of the outer membranes is broken, forms one of the circumstances which have been distinguished by the name of by or false waters.

Between the middle and inner membrane, upon or near the funis, there is a small, flat, and oblong body, which, in the early part of pregnancy, seems to be a vesicle containing milky lymph, which afterwards becomes of a firm and apparently fatty texture. This is called the *vesicula umbilicalis*; but its use is not known. See *Placenta*.

SECUNDUM ARTEM. According to art. A term frequently used in prescription, and denoted by the letters S. A. which are usually affixed, when the making up of the recipe in perfection requires some uncommon care and dexterity.

SECURIDACA. (From *securis*, an axe; so called because its leaves resemble a small axe.) *Henbane*.

SEDATIVES. (*Medicamenta sedativa.* from *sedo*, to ease or assuage.) *Sedantia*, medicines which have the power of diminishing the animal energy, without destroying life. They are divided into *sedativa soporifica*, as opium, papaver, hyosciamus, and *sedativa refrigerantia*, as neutral salts, acids, &c.

Sedative salt of Homberg. See *Boracic acid*.

SEDENTARIA OSSA. The os coccygis and ischia.

Sedge. See Iris palustris.

SEDIMENT. The heavy parts of liquids, which fall to the bottom.

Sediments latritious. See Latritious sediment.

SEDLITZ WATER. *Seydchitz water.* A simple saline mineral water. From chemical analysis it appears, that it is strongly impregnated with vitriolated magnesia or Epsom salt, and it is to this, along with probably the small quantity of muriat of magnesia, that it owes its bitter and saline taste, and its purgative properties. The diseases in which this water is recommended are, crudities of the stomach, hypochondriasis, amenorrhœa, and the anomalous complaints succeeding the cessation of the catamenia, œdematous tumours of the legs in literary men, hæmorrhoidal affections, and scorbutic eruptions.

SEDUM. (From *sede*, to assuage; so called because it allays inflammation.) The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Pentagynia*.

SEDUM LUTEUM MURALE. Navel wort.

SEDUM MAJUS. *Aizoon. Barba Jovis.* House-leek or sangreen. *Sempervivum tectorum* of Linneus. The leaves of this plant have no remarkable smell, discover to the taste a mild subacid austerité; they are frequently applied by the vulgar to bruises and old ulcers.

SEDUM MINUS. See *Illecebra*.

SEDUM TELEPHIUM. The systematic name of the *orpine*. See *Faba crassa*.

SEEING. A sensation by which we perceive bodies around us, and their sensible qualities. The organ of sight is formed of three parts perfectly distinct, which serve to protect the globe of the eye, to withdraw it suddenly from the influence of light, and to preserve it in a condition necessary for the exercise of these functions; These consist in the supercilia, palpebræ, and lachrymal passages, parts accessory to the organ. The eye-ball itself presents two portions very different from each other, one formed by almost the whole, and may be called an optic instrument; the other formed by a medullary expansion of the optic nerve is the immediate organ of sight, this is the retina, alone adapted to receive the impression of light, and to be affected by the delicate contact of this extremely subtle fluid. This impression or sensation is transmitted to the cerebral organ by the optic nerve, of which the retina is merely the expanded extremity.

The eye-brows, as being accessory organs to vision, have the effect of diminishing the effect of a too strong light by partly absorbing its rays. The supercilia answers this purpose better in proportion to the projection formed, and the darker colour

of the hair; thus we knit the brow transversely in passing from dark to a lighted place, the strong light of which has a disagreeable effect on the organ of sight. Hence arose the custom of certain southern people in whom the eye-brows are thicker and of a darker colour, to make them blacker in order the better to fulfil the intention for which they were designed. The eye-lids, as concerned in the organs of vision, shade the eyes from the continual action of light, these like all other organs have occasion for repose, which could not have been procured had the rays of light constantly excited their sensibility. A removal of the eyelids occasions loss of sleep. The cilia, or hairs, growing upon the margin are destined to prevent insects or other light bodies in the atmosphere from insinuating between the globe of the eye and its covering. The anterior part of the eye, thus defended against external injuries, is continually moistened by the tears, they also guard against the effects of friction, to which the eye is exposed. (See *Eye*.)

Luminous rays, emanating from a light object, form a cone, the apex of which corresponds to the point of a body which we are looking at, and its base is applied to the anterior part of the cornea; all those rays which touch the mirror of the eye pass through it, experience a refraction proportioned to the density of the cornea, and to the convexity of this membrane, greater than that of the atmosphere: when approaching the perpendicular they pass through the aqueous humours less dense, and meet with the iris. All those rays which fall on this membrane are reflected, and shew its colour different in different individuals. It is only the most central rays that penetrate the pupil, and serve for sight: these enter the pupil in greater or less number, according as it may be more or less dilated. The pupil becomes larger or smaller conformable to the expansion or contraction of the iris. The motions of the iris depend entirely on the mode in which light affects the retina; it is of itself insensible to the impression of luminous rays, as proved by Fontana, who always found it immoveable when he directed rays of light exclusively to it.

The rays to which the pupil give passage pass through the aqueous humour of the posterior chamber; and soon come into contact with the crystalline lens, which powerfully refracts them, on account of its density and lenticular form. When more approaching the perpendicular by this body, they proceed as far as the retina, through the vitreous humour, that is less dense, and which preserves, without augmentation, the effect of the refraction produced by the crystalline lens; the rays

assembled into one focus strike only a single point of the retina, and produce an impression that gives us an idea of certain properties of the bodies it reflects. It is generally thought that luminous pyramids which emanate from all points of the object we behold, decussate in passing through the globe of the eye, so that the object itself is figured in a reverse direction. Although the image of each object is traced at the same time in each of our eyes, we have but one sensation, because both sensations are in harmony or combined, and only serve by assisting each other to render the impression stronger and more durable. The correspondence of affection requires the direction of the optic axes on the same objects, and however little this direction be changed we really see double, which happens in strabismus, or squinting.

If the eyes possess a too energetic power of refraction either from too great a convexity of the cornea or crystalline lens, or more considerable density of the humours and excessive depth of the globe of the eye, the luminous rays being united too soon, cross each other, again diverge, fall scattered on the retina, and produce only a confused sensation. In this disease of vision called myopia, patients can only distinguish very near objects, whence rays are given off which require an instrument possessing a considerable power of refraction. In presbyopia, on the contrary, the cornea being too flat, the crystalline not very convex, or being deep seated, the humours not sufficiently abundant, cause the rays not to be yet assembled, when they fall on the retina; so that patients can only observe with distinct objects, because the rays that come from them being very convergent, have not occasion to be much refracted. The sensibility of the retina is, under certain circumstances, so much raised, that the eye hardly supports the weakest light. Persons in this situation are called nyctalopes, who distinguish objects in the midst of utter darkness, as a few rays are sufficient to affect their organ of vision. The eyes are not immoveable in the part they occupy, they are directed towards all the objects of which we wish to form a knowledge by different motions, regulated by four recti and two oblique muscles, and it is observed that there is such a correspondence of action in muscles that move both eyes, that these organs turn at the same time towards the object, so that the visual axes are exactly parallel.

SODIUM'S SALT. A neutral salt, which consists of soda, potash, and tartaric acid. It was prepared and made known by a Frenchman named Peter Seignette, towards the end of the last century. It was then employed in preference to many other medicines long known, which had

been equally serviceable; and by these means, without much trouble, he was enabled to acquire a fortune. It must, however, be allowed that he was a skilful chemist, who, by his writings, and the invention of various other medicines, had obtained considerable reputation as a physician and naturalist. He was established as an apothecary at Rochelle; published papers on various natural objects which he had observed in his neighbourhood, in the Memoirs of the Academy of Sciences at Paris, as well as in other works; and died on the 11th of March, 1719.

He recommended this salt, which enriched him, and rendered his name famous, in some small treatises, printed in particular about the year 1762. He called it sometimes alkaline salt, sometimes *sal polycrest*, and sometimes *Rochelle salt*. After his death, his son continued to prepare and to vend it with the greatest success.

Seignette discovered this salt while he was engaged in making soluble tartar, and, according to the old opinion, imagining that both the fixed alkalis were the same, used soda instead of potash. By this means he procured, not without surprise, a salt different from the common soluble tartar which he wished to prepare, and from the other well-known salt also. He was induced, therefore, to examine it. The experiments of learned chemists discovered the component parts of this salt. The mode of preparing it was then made publicly known; and, by more accurate examination, the difference, before overlooked, between vegetable and mineral alkali, was determined; by which new light was thrown upon chemistry, and an important service rendered to a variety of arts.

Among those who contributed to bring this salt into repute was Nicholas Lemery, to whom Seignette sent a large quantity of it, which he distributed at Paris, though unacquainted with its component parts. Its composition was discovered at the same time, about the year 1731, by two French chemists, Baldue and Geoffroy, the former published his observations in the Memoirs of the Academy of Sciences; and the latter communicated his to Sir Hans Sloane, who caused them to be printed in the Philosophical Transactions. Newman, therefore, was not the first who made known the composition of Seignette's salt, in his treatise on salt-petre; for Newman's salt is essentially different; and he himself confesses that he was not acquainted with the Rochelle salt. See *Sodu tartarizata*.

SELENITE. (From *σεληνη*, the moon.) A white stone, having a figure upon it resembling a moon. Sulphat of lime.

SELENIUM. (From *σεληνη*, the moon;

so called from its usefulness in lunacy.) A kind of peony.

Self-heal. See *Prunella*.

SELINX. A disease of the nails, in which white spots are occasionally seen in their substance.

SELINUM. (From *selene*, the moon; from its supposed usefulness in disorders proceeding from the influence of the moon.) Marsh smallage.

SELLA TURCICA. (*Sella*, quasi *sedda*, from *sedes*, to sit, and *turca*; from its supposed resemblance to a Turkish saddle.) *Ephippium*. A cavity in the sphenoid bone, containing the pituitary gland, surrounded by the four clinoid processes.

SELTZER WATER. A saline water, slightly alkaline, highly acidulated with carbonic acid, containing more of this volatile principle than is sufficient to saturate the alkali, and the earths which it holds in solution. It is particularly serviceable in relieving some of the symptoms that indicate a morbid affection of the lungs; in slow hectic fever, exanthematous eruptions of the skin, foulness of the stomach, bilious vomiting, acidity and heart-burn, spasmodic pains in any part of the alimentary canal, and bloody or highly offensive stools. On account of its property in relieving spasmodic pains, and from its rapid determination to the kidneys, and perhaps its alkaline contents, it has been sometimes employed with great advantage in diseases of the urinary organs, especially those that are attended with the formation of calculus. A large proportion of the Seltzer water, either genuine or artificial, that is consumed in this country, is for the relief of these disorders. Even in gonorrhoea, either simple or venereal, Hoffman asserts that advantage is to be derived from this medicine. The usual dose is from half a pint to a pint.

SEMECARPUS ANACARDIUM. The systematic name of the tree which is supposed to afford the Mollucca bean. See *Amacardium orientale*.

SEMIOSIS. (From *σημνω*, to notify.) See *Semiotica*.

SEMEN. 1. The seed, kernel, or stone of the fruit of any vegetable.

2. The seed or prolific liquor secreted in the testicles, and carried through the epididymis and vas deferens into the vesiculae seminales, to be emitted *sub coitu* into the female vagina, and there, by its aura, to penetrate and impregnate the ovulum in the ovarium.

In castrated animals, and in eunuchs, the vesiculae seminales are small, and contracted; and a little lymphatic liquor, but no semen, is found in them. The semen is detained for some time in the vesiculae seminales, and rendered thicker from the continual absorption of its very thin part,

by the oscula of the lymphatic vessels. In lascivious men, the semen is sometimes, though rarely, propelled by nocturnal pollution from the vesiculae seminales, through the ejaculatory ducts, (which arise from the vesiculae seminales, perforate the urethra transversely, and open themselves by a narrow and very nervous mouth at the sides of the caput gallinaginis,) into the urethra, and from it to some distance. But in chaste men the greatest part is again gradually absorbed from the vesiculae seminales through the lymphatic vessels, and conciliates strength to the body. The smell of semen is specific, heavy, affecting the nostrils, yet not disagreeable. The same odour is observed in the roots of the orchis, juice of chestnuts, and the anthers of many plants. The smell of the semen of quadrupeds, when at heat, is so penetrating as to render their flesh fetid and useless, unless castrated. Thus the flesh of the stag, *temper coitus*, is unfit to eat. The taste of semen is fatuous and somewhat acrid. In the testes its consistence is thin and diluted; but in the vesiculae seminales, viscid, dense, and rather pellucid; and by venery and debility it is rendered thinner.

Specific gravity. The greatest part of the semen sinks to the bottom in water, yet some part swims on its surface, which it covers like very fine threads mutually connected together in the form of a cobweb.

Colour. In the testicles it is somewhat yellow, and in the vesiculae seminales it acquires a deeper hue. That emitted by pollution or coition, becomes white from its mixture with the whitish liquor of the prostate gland during its passage through the urethra. In those people who labour under jaundice, and from the abuse of saffron, the semen has been seen yellow, and in an atrabiliary young man, black.

Quality. Semen exposed to the atmospheric air, loses its pellucidity, and becomes thick, but after a few hours it is again rendered more fluid and pellucid than it was immediately after its emission. This phenomenon cannot arise from water or oxygen attracted from the air. At length it deposits a phosphorated calx, and forms a corneous crust.

Experiments with semen prove that it turns the syrup of violets green, and dissolves earthy, mediate, and metallic salts. Fresh semen is insoluble in water, until it has undergone the above changes in atmospheric air. It is dissolved by alkaline salts. By ætherial oil it is dried into a pellucid pellicle, like the cortex of the brain. It is dissolved by all acids, except the oxygenated acid of salt, by which it is coagulated in the form of white flakes. It is also acted upon by alcohol of wine.

By *dry distillation* semen gives out a small portion of empyreumatic oil, and

volatile alkali. The remaining incinerated carbone affords soda and phosphorated calx.

The *constituent principles of semen*. Chemical analysis demonstrates that one hundred parts of semen contain, 1. *Of water*, ninety parts, 2. *Of animal gluten*, six parts, 3. *Of phosphorated calx*, one part, 4. *Of pure soda*, three parts, 5 By microscopical examination, it is asserted that an immense number of very small animalcules with round tails, called *spermatic animalcules*, may be seen, 6. *The odorous principle*, which flies off immediately from fresh semen. It appears to consist of a peculiar vital principle, and by the ancients was called *aura seminis*.

Use. 1. Emitted into the female vagina *sub coitu*, it possesses the wonderful and stupendous power of impregnating the ovulum in the female ovarium. The odorous principle, or *aura spermatica* only, appears to penetrate through the cavity of the uterus and fallopian tubes to the female ovarium, and there to impregnate the albuminous latex of the mature ovulum by its vital power. The other principles of the semen appear to be only a vehicle of the seminal *aura*. 2. In chaste men, the semen returning through the lymphatic vessels into the mass of the blood, gives strength to the body and mind; hence the bull is so fierce and brave, the castrated ox so gentle and weak; hence every animal languishes *post coitum*; and hence *tabes dorsalis* from onanism. 3. It is by the stimulus of the absorbed semen, at the age of puberty, into the mass of the humours; that the beard and hair of the pubes, but in animals the horns, are produced; and the weeping voice of the boy changed into that of a man.

SEMEN ADJOWARN. A seed imported from the East, of a pleasant smell, a grateful aromatic taste, somewhat like savory. It possesses exciting, stimulating, and carminative virtues, and is given in the East in nervous weakness, dyspepsia, flatulency, and heart-burn.

SEMEN AGAVE. An East Indian seed, exhibited there in atonic gout.

SEMEN CONTRA. See *Santonicum*.

SEMEN SANCTUM. See *Santonicum*.

SEMI. (*Semi*, from *μεν*, half.) *Semis* in composition universally signifies half, as *semicupium*, a half-bath, or bath up to the navel; *semilunaris*, in the shape of a half-moon.

SEMICIRCULAR CANALS. These canals are three in number, and take their name from their figure. They belong to the organ of hearing, and are situated in the petrous portion of the temporal bone, and open into the vestibulum.

SEMICUPIUM. *Excathisma. Incessio.* A half-bath, or such as receives only the hips, or extremities.

SEMI INTEROSSEUS INDICIS. See *Abductor indicis manus*.

SEMILUNAR VALVES. The three valves at the beginning of the pulmonary artery and aorta are so termed, from their half-moon shape.

SEMIOTICE. (From *σημειον*, a sign.) *Semeiosis*. That part of pathology which treats on the signs of diseases.

SEMIMEMBRANOSUS. *Isthio-popliti-femoral* of Dumas. This muscle arises from the outer surface of the tuberosity of the ischium, by a broad flat tendon which is three inches in length. From this tendon it has gotten the name of semi-membranosus. It then begins to grow fleshy, and runs at first under the long head of the biceps, and afterwards between that muscle and the semi tendinosus. At the lower part of the thigh it becomes narrower again, and terminates in a short tendon, which is inserted chiefly into the upper and back part of the head of the tibia, but some of its fibres are spread over the posterior surface of the capsular ligament of the knee. Between this capsular ligament and the tendon of the muscle, we find a small bursa mucosa. The tendons of this and the last described muscle form the inner ham-string. This muscle bends the leg, and seems likewise to prevent the capsular ligament from being pinched.

SEMI-NERVOSUS. See *Semitendinosus*.

SEMINIS EJACULATOR. See *Accelerator urinae*.

SEMI-ORBICULARIS ORIS. See *Orbicularis oris*.

SEMI-SPINALIS COLLI. *Semi-spinalis sive transverso-spinalis colli* of Winslow, *Spinalis cervicis* of Albinus, *Spinalis colli* of Douglas, *Transversalis colli* of Cowper, and *Transverso-spinal* of Dumas. A muscle situated on the posterior part of the neck, which turns the neck obliquely backwards, and a little to one side. It arises from the transverse processes of the uppermost six vertebrae of the back by as many distinct tendons, ascending obliquely under the complexus, and is inserted into the spinous processes of all the vertebrae of the neck, except the first and last.

SEMI SPINALIS DORSI. *Semi-spinalis externus seu transverso-spinalis dorsi* of Winslow. *Semi spinatus* of Cowper, and *Transverso-spinal* of Dumas. A muscle situated on the back, which extends the spine obliquely backwards. It arises from the transverse processes of the seventh, eighth, ninth, and tenth vertebrae of the back, by as many distinct tendons, which soon grow fleshy, and then become tendinous again, and are inserted into the spinous processes of all the vertebrae of the back above the eighth, and into the lowermost of the neck, by as many tendons.

SEMI-SPINALIS EXTERNUS. See *Semi-spinalis dorsi*.

SEMI SPINATUS. See *Semi-spinalis dorsi*.

SEMI-TENDINOSUS. This muscle, which is the *semi-nervosus* of Douglas and Winslow, and *Ischio-creti tibial* of Dumas, is situated obliquely along the back part of the thigh. It arises tendinous and fleshy from the inferior, posterior, and outer part of the tuberosity of the ischium, in common with the long head of the biceps cruris, to the posterior edge of which it continues to adhere, by a great number of oblique fibres, for the space of two or three inches. Towards the lower part of the os femoris, it terminates in a round tendon, which passes behind the inner condyle of the thigh bone, and becoming flat, is inserted into the upper and inner part of the ridge of the tibia, a little below its tuberosity. This tendon sends off an aponeurosis, which helps to form the tendinous fascia that covers the muscles of the leg. This muscle assists in bending the leg, and at the same time draws it a little inwards.

SEMPERVIVUM. (From *semper*, always, and *vivo*, to live; so called because it is always green.) 1. The name of a genus of plants in the Linnæan system. Class, *Dodecandria*. Order, *Polygymia*.

2. The pharmacopœial name of some plants. See *Sedum majus*.

SEMPERVIVUM ACRE. The stone-crop is occasionally so termed. See *Illecebra*.

SEMPERVIVUM TECTORUM. The systematic name of the house-leek. See *Sedum majus*.

SENECIO. (*Senecio*, from *senesco*, to grow old; so called because it has a greyish down upon it, like the beard of old men.) 1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*, Order, *Polygamia superflua*.

2. The pharmacopœial name also of the groundsel. See *Erigerum*.

SENECIO VULGARIS. The systematic name of groundsel. See *Erigerum*.

SENECIO JACOBÆA. The systematic name of the ragwort. See *Jacobæa*.

SENECTA ANGUIM. The cast skin of a serpent; its decoction is said to cure deafness.

SENEGA. See *Seneka*.

SENEGA GUM. See *Gummi senegalense*.

Senegaw milkwort. See *Seneka*.

SENEKA. (So called because the Seneca or Senegaw Indians use it against the bite of the rattle-snake.) The rattle-snake-root milkwort. *Polygala senega* of Linnæus:—*Floribus imberbibus spicatis, caule erecto herbaceo simplicissimo, foliis lato lanceolatis*. The root of this plant was formerly much esteemed as a specific against the poison of the rattle-snake, and as an antiphlogistic in pleurisy, pneumonia, &c. but it is now very much laid aside. Its dose is

from ten to twenty grains; but when employed, it is generally used in the form of decoction, which, when prepared according to the formula of the Edinburgh Pharmacopœia, may be given every second or third hour.

Seengreen. See *Sedum majus*.

SENNA. (From *senna*, an Arabian word, signifying acute; so called from its sharp pointed leaves.) *Senna alexandrina*. *Senna italica*. *Folium orientale*. *Senna*, or Egyptian cassia. *Cassia senna* of Linnæus:—*foliis sejugis subovatis, petiolis eglandulatis*. The leaves of senna, which are imported here from Alexandria for medicinal use, have a rather disagreeable smell, and a subacid, bitterish, nauseous taste. They are in common use as a purgative. The formulæ given of the senna by the colleges, are those of infusion, a powder, a tincture, and an electuary. See *Infusum senna*, &c.

SENNA ALEXANDRINA. See *Senna*.

SENNA ELECTUARIUM. See *Electuarium senna*.

SENNA ITALICA. See *Senna*.

SENNA PAUPHUM. Bastard senna, or milk-vetch.

SENNA SCORPIUM. The scorpion senna.

SENNA EXTRACTUM. Extract of senna.

SENNA INFUSUM. See *Infusum senna*.

SENNA INFUSUM TARTARIZATUM. *Senna*, coriander, and cream of tartar, infused in water.

SENSATION. Sensation, or feeling, is the consciousness of a change taking place in any part, from the contact of a foreign body with the extremities of our nerves. The seat of sensation is in the pulp of the nerves.

The impression produced on any organ by the action of an external body constitutes sensation. This sensation, transmitted by nerves to the brain, is perceived, that is, felt by the organ; the sensation then becomes *perception*; and this first modification implies, as must be evident, the existence of a central organ, to which impressions produced on the senses are conveyed. The cerebral fibres are acted on with greater or less force by the sensations propagated by all the senses influenced at the same time; and we could only acquire confused notions of all bodies that produce them, if one particular and stronger perception did not obliterate the others, and fix our attention. In this collective state of the mind on the same subject, the brain is weakly affected by several sensations which leave no trace behind. It is on this principle that, having read a book with great attention, we forget the different sensations produced by the paper and characters.

When a sensation is of short duration the knowledge we have of it is so weak that soon afterwards there does not re-

main any knowledge of having experienced it. In proportion as a sensation, or an idea, which is only a sensation transformed or perceived by the cerebral organ, has produced in the fibres of this organ a stronger or weaker impression, the remembrance of it becomes more or less lively and permanent. Thus we have a *remembrance* of it, that is, call to mind that we have already been affected in the same manner: a *memory*, or the act of recalling the object of the sensation with some of its attributes, as colour, volume, &c.

When the brain is easily excitable, and, at the same time, accurately preserves impressions received, it possesses the power of representing to itself ideas with all their connexions, and all the accessory circumstances by which they are accompanied, of reproducing them in a certain degree, and of recalling an entire object, while the memory only gives us an idea of its qualities. This creative faculty is called *imagination*. When two ideas are brought together, compared, and their analogy considered, we are said to form a *judgment*; several judgments connected together constitute reasoning.

Besides the sensations that are carried from the organs of sense to the brain, there are others, internal, that seem to be transmitted to it by a kind of sympathetic reaction. It is well known what uneasiness the affection of certain organs conveys to the mind, how much an habitual obstruction of the liver is connected with a certain order of ideas; these internal sensations are the origin of our moral faculties, in the same manner as impressions that are conveyed by the organs of sense are the source of intellectual faculties. We are not on that account to place the seat of the passions of the mind in the viscera; it is only necessary to remember that the appetites, whence arise the passions, reside in this organ, and are a phenomenon purely physical, while passion consists, at the same time, in the intellectual exertion. Thus an accumulation of semen in the cavities that are employed as a reservoir for it, excites the appetite for venery, very distinct from the passion of love, although it may be frequently the determinate cause of it.

The senses may be enumerated under the following heads, viz. the sense of vision, hearing, smelling, tasting, touching.

SENSIBILITY. The capability which a nerve possesses of conveying the sensation produced by the contact of another body with it. All parts possessed of a power of producing a change, so as to excite a sensation, are called *sensible*; those which are not possessed of this property, *insensible*. To the insensible parts by nature belong all our fluids, the blood, bile, saliva, &c. and much of the solids, the

hair, epidermis, nails, &c.; but the sensible parts are the skin, eyes, tongue, ear, nose, muscles, stomach, intestine, &c.

SENSORIUM. See *Cerebrum*.

SENSORIUM COMMUNE. See *Cerebrum*.

SENSUS EXTERNI. The external senses are seeing, hearing, tasting, smelling, and feeling.

SENSUS INTERNI. The internal senses are imagination, memory, judgment, attention, and the passions.

SENTIENT EXTREMITIES. The extremities of the nerves.

SEPARATORIUM. (From *separo*, to separate.) An instrument for separating the pericranium from the skull, and a chemical vessel for separating essential parts of liquids.

SEPIA OFFICINALIS. *Sepium.* *Precipians magnum.* The cuttle fish. The systematic name of the fish whose shell possesses calcareous qualities, and is often mixed into tooth-powders.

SEPIE OS. See *Sepia officinalis*.

SEPTENARY YEARS. Climacteric years. A supposed period, or succession of years in human life, at which important constitutional changes are considered to take place; and the end of this period is therefore judged critical. This period is fixed at every seventh year. The grand climacteric is fixed at 63, and, passing that time, age, it is considered, may be protracted to go to 90. So general is this belief, that the passing of 60 generally gives much anxiety to most people.

SEPTIFOLI. See *Tormentilla*.

SEPTIC. (*Septica*; from *σπνσ*, to putrefy.) Relating to putrefaction.

SEPIUM. See *Sepia officinalis*.

SEPTIFOLIA. (From *septem*, seven, and *folium*, a leaf; so named from the number of its leaves.) Coralwort, or setfoil toothwort.

SEPTINERVIA. (From *septem*, seven, and *nervus*, a string; so called from the seven strings upon its leaf.) A species of plantain.

SEPTUM CEREBELLI. A process of the dura mater, dividing the cerebellum perpendicularly into two principal parts.

SEPTUM CEREBRI. The falciform process of the dura mater is sometimes so called. See *Falciform process*.

SEPTUM CORDIS. (*Septum*; from *sepio*, to separate.) The partition between the two ventricles of the heart.

SEPTUM LUCIDUM. *Septum pellucidum.* The thin and tender portion of the brain, dividing the lateral ventricles from each other.

SEPTUM NARIUM. *Interseptum.* The partition between the nostrils.

SEPTUM PALATI. The partition of the palate.

SEPTUM PELLUCIDUM. See *Septum lucidum*.

SEPTUM THORACIS. See *Mediastinum*.

SEPTUM TRANSVERSUM. See *Diaphragm*.

SERAPIAS. (From *Serapis*, a lascivious idol; so called because it was thought to promote venery; or from the testiculated shape of its root.) The name of a genus of plants in the Linnæan system. Class, *Gynandria*. Order, *Diandria*.

SERAPINUM. The gum-resin sagapenum is sometimes so called. See *Sagapenum*.

SERIPHUM. (From *Serphus*, an island upon which it grew.) Flix-weed.

SERIS. *Agave*. Endive.

Sermountum See *Seviti*.

Serous apoplexy. See *Apoplexia*.

SERPENTARIA GALLORUM. The arum dracunculus. See *Dracunculus*.

SERPENTARIA HISPANICA. The viper's grass. See *Scorsonera*.

SERPENTARIA VIRGINIANA. (So called from the resemblance of its roots to the tail of the rattle-snake.) *Colebrina virginiana*. *Viperina virginiana*. *Aristolochia*. *Pestilochia*. *Contrayerva virginiana*. Virginian snake-root. The plant which affords this root is the *Aristolochia serpentaria* of Linnæus:—*foliis cordatis oblongis planis, caulis infirmis flexuosis teretibus, floribus solitariis*. *Caule geniculato valde nodoso*. *Flores ad radicem*. Snake-root has an aromatic smell, approaching to that of valerian, but more agreeable; and a warm, bitterish, pungent taste. It was first recommended as a medicine of extraordinary power in counteracting the poisonous effects of the bites of serpents; this, however, is now wholly disregarded; but as it possesses tonic and antiseptic virtues, and is generally admitted as a powerful stimulant and diaphoretic, it is employed, in the present day, in some fevers where these effects are required. A *tinctura serpentarie* is directed both by the London and Edinburgh Pharmacopœias.

SERPENTUM LIGNUM. The nature of this root does not appear to be yet ascertained. It is the produce of the *Ophiaryum serpentium* of Linnæus: by whom it is said to be very bitter. In the cure of the bite of venomous serpents and malignant diseases it is said to be efficacious.

SERPENTUM RADIX. See *Mungos radix*.

SERPIGO. (From *serpo*, to creep; because it creeps on the surface of the skin by degrees.) A ring-worm, or tetter. See *Herpes*.

SERPYPILLUM. (From *serpo*, to creep, or a *serpendo*, by reason of its creeping nature.) *Serpillum*. *Gilarum*. *Serpillium vulgare minus*. Wild or mother of thyme. *Thymus serpyllum* of Linnæus:—*erectus, foliis revolutis ovatis, floribus verticillatispicatis*. This plant has the same sensible

qualities as those of the garden thyme, (see *Thymus*), but has a milder and rather more grateful flavour.

SERPYPILLUM CITRATUM. Lemon thyme. A variety of the *Thymus serpyllum* of Linnæus. It is very pungent, and has a particularly grateful odour, approaching to that of lemons.

SERPYPILLUM VULGARE MINUS. See *Serpillum*.

SERRATA. (From *serra*, a saw; so called from its serrated leaves.) See *Serratula*.

SERRATULA. (From *serra*, a saw; so called from its serrated leaves.) The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia æqualis*.

SERRATULA ANARA. The systematic name of a species of saw-wort which is said to cure agues.

SERRATUS ANTICUS. See *Pectoralis minor*.

SERRATUS MAGNUS. (*Serratus*; from *serra*, a saw; so called from its saw-like appearance.) *Serratus major anticus*, of Douglas and Cowper. *Serratus major* of Winslow, and *Costo basi-scapularis* of Dumas. This muscle is so named by Winslow and Albinus. Douglas calls it *serratus major anticus*, but improperly, as it is seated at the side, and not at the anterior part of the thorax. It is a broad fleshy muscle, of a very irregular shape, and is in part covered by the subscapularis, pectoralis, and latissimus dorsi. It arises, by fleshy digitations, from the eight superior ribs, and is inserted fleshy into the whole basis of the scapula internally, between the insertion of the rhomboides, and the origin of subscapularis, being folded, as it were, about the two angles of the scapula. This muscle may easily be divided into two and even three portions. The latter division has been adopted by Winslow. The first of these portions is the thick and short part of the muscle that arises from the first and second ribs, and is inserted into the upper angle of the scapula, its fibres ascending obliquely backwards. The second portion arises from the second rib, behind the origin of the first portion, and likewise from the third and fourth ribs; this portion is thin and short, and its fibres run nearly in a horizontal direction, to be inserted into the basis of the scapula. The third, and most considerable portion, is that which arises from the fifth, sixth, seventh, and eighth ribs, and is inserted into the lower angle of the scapula. The *serratus magnus* serves to move the scapula forwards, and it is chiefly by the contraction of this muscle that the shoulder is supported, when loaded with any heavy weight. The ancients, and even many of the moderns, particularly Douglas and Cowper, supposed

its chief use to be to dilate the thorax, by elevating the ribs; but it can only do this when the scapula is forcibly raised.

SERRATUS MAJOR ANTICUS. See *Serratus magnus*.

SERRATUS MINOR ANTICUS. See *Pectoralis minor*.

SERRATUS POSTICUS INFERIOR.

Dorso-lumbo-costal of Dumas. This is a thin muscle of considerable breadth situated at the bottom of the back, under the middle part of the latissimus dorsi. It arises by a broad thin tendon, in common with that of the last-described muscle, from the spinous processes of the two, and sometimes of the three inferior dorsal vertebrae, and from three, and sometimes four of those of the lumbar vertebrae: It then becomes fleshy, and, ascending a little obliquely outwards and forwards, divides into three, and sometimes four fleshy lips, which are inserted into the lower edges of the three or four inferior ribs, at a little distance from their cartilages. Its use seems to be to pull the ribs downwards, backwards, and outwards.

SERRATUS SUPERIOR POSTICUS.

Cervico-dorso-costal of Dumas. This is a small, flat, and thin muscle, situated at the upper part of the back, immediately under the rhomboideus. It arises, by a broad thin tendon, from the lower part of the ligamentum colli, from the spinous process of the last vertebra of the neck, and the two or three uppermost of the back, and is inserted into the second, third, fourth, and sometimes fifth ribs, by as many distinct slips. Its use is to expand the thorax, by pulling the ribs upwards and outwards.

SERTULA CAMPANA. See *Melilotus*.

SERUM. (From *serus*, late; because it is the remainder of the milk, after its better parts have been taken from it.) The serum of the blood. The yellow and somewhat greenish fluid which separates from the blood when cold and at rest. See *Blood*.

SERUM ALUMINOSUM. Alum whey.

SERUM LACTIS. Whey.

SERVICE-TREE. The fruit of this tree is considered powerfully astringent, and recommended in fluxes and dysenteries. It is given in the form of rob, and it is equally useful in distilling brandy and making cider.

SESAMOID BONES. (*Ossa sesamoides*; from *σάμη*, an Indian grain, and *οἶδος*, likeness.) This term is applied to the little bones, which, from their supposed general resemblance to the seeds of the sesamum, are called *ossa sesamoides*. They are found at the articulations of the great toes, and sometimes at the joints of the thumbs; now and then we meet with them upon the condyles of the os femoris, at the lower extremity of the fibula, under the os cuboides of the tarsus, &c. They

do not exist in the fœtus, but as we advance in life, begin first to appear in a cartilaginous state, and, at length, in adult subjects, are completely ossified. Age and hard labour seem to add to the number and size of these bones, and being most commonly found wherever the tendons and ligaments are most exposed to pressure from the action of the muscles, they are now generally considered by anatomists as the ossified parts of tendons and ligaments. These bones are usually smooth and flat on the side of the bone on which they are placed; their upper surface is convex, and, in general, adheres to the tendon that covers it, and of which it may, in some measure, be considered as a part. Although their formation seems to be owing to accidental circumstances, yet, as the two at the first joint of the great-toe are much larger than the rest, and are seldom wanting in an adult, it would seem as if these bones were of some utility; perhaps by removing the tendons farther from the centre of motion, and thus increasing the power of the muscles. The *ossa sesamoides* of the great-toe and thumb seem likewise to be of use, by forming a groove for lodging the flexor tendons secure from compression.

Sesamoidal bones. See *Sesamoid bones*.

SESAMUM. (An Egyptian word.) 1. The name of a genus of plants in the Linnean system.

2. The pharmacopœial name of the *Sesamum orientale* of Linnæus. The seed and leaves are used medicinally in some countries on account of the bland oil the former contains, and for the mucilaginous nature of the latter.

SESAMUM ORIENTALE. The systematic name of the sesamum of the pharmacopœias. See *Sesamum*.

SESELI. (Παρα το σάωρα ἄλλοι; because it is salutary for young fawns.) *Siler montanum*. Hart-wort. Sermountain. The seeds and roots of this plant, *Laserpitium siler* of Linnæus, which grows in the southern parts of Europe, are directed as officinals. They have an agreeable smell, and a warm, glowing, aromatic taste; and, though neglected in this country, do not appear to be deservedly so.

SESELI CARIUM. There is great confusion amongst the species of the seseli. The plant which bears this epithet in the pharmacopœias is the *Tordylium officinale* of Linnæus. The seeds are said to be diuretic.

SESELI MASSILIENSE. Hart-wort of Marsailles. This plant is the *Seseli tortuosum* of Linnæus. The seeds are directed for medicinal use, and have a warm biting taste, and a greater degree of pungency than those of the *Laserpitium*.

SESELI TORTUOSUM. The systematic name of the hart-wort of Marsailles. See *Seseli massiliense*.

SESSUI. This word, joined with any number, weight, measure, &c. signifies one integer and an half; as *sessui gratum*, a grain and a half.

SETACEUM (From *seta*, a bristle; because horse-hairs were first used to keep open the wound) A seton. See *Seton*.

SETON. *Setaceum*. An artificial ulcer made under the skin by means of an instrument called the seton needle, which carries with it a portion of thread or silk, that is moved backwards or forwards, and thus keeps up a constant irritation.

Setterwort. See *Helleboraster*.

SEBUM CETI. See *Spermaceti*.

SEBUM OVILE. *Sebum ovillum*. Mutton suet.

SEXUAL ACTION. Sexual functions. Those functions proper to each sex, by which the species is propagated, as the excretion of semen in men; menstruation, conception, the evolution of the fetus, parturition, &c. in women.

SEXUAL SYSTEM OF PLANTS. Linnæan system. The sexual system of plants was invented by the immortal Linnæus, professor of physic and botany at Upsal in Sweden. It is founded on the parts of fructification, viz. the stamens and pistils; these having been observed with more accuracy since the discovery of the uses for which nature has assigned them, a new set of principles have been derived from them, by means of which the distribution of plants has been brought to a greater precision, and rendered more conformable to true philosophy, in this system, than in any one of those which preceded it. The author does not pretend to call it a natural system, he gives it as artificial only, and modestly owns his inability to detect the order pursued by nature in her vegetable productions; but of this he seems confident, that no natural order can ever be framed without taking in the materials out of which he has raised his own; and urges the necessity of admitting artificial systems for convenience, till one truly natural shall appear. Linnæus has given us his *Fragmenta methodi naturalis*, in which he has made a distribution of plants under various orders, putting together in each such as appear to have a natural affinity to each other; this, after a long and fruitless search after the natural method, he gives as the result of his own speculation, for the assistance of such as may engage in the same pursuit.

Not able to form a system after the natural method, Linnæus was more fully convinced of the absolute necessity of adopting an artificial one. For the student to enter into the advantages this system maintains over all others, it is necessary that he be instructed in the science of botany, which will amply repay him for his inquiry. The following is a short outline of the sexual system:

The parts of the fructification of a plant are,

1. The *calyx*, called also the empalement, or flower-cup.

2. The *corole*, or foliation, which is the gaudy part of the flower, called vulgarly the leaves of the flower.

3. The *stamens*, or threads, called also the chives; these are considered as the male parts of the flower.

4. The *pistil*, or pointal, which is the female part.

5. The *pericarp*, or seed-vessel.

6. The *seed*.

7. The *receptacle*, or base, on which these parts are seated.

The four first are properly parts of the flower, and the three last parts of the fruit. It is from the number, proportion, position, and other circumstances attending these parts of the fructification that the classes and orders, and the genera they contain, are to be characterised, according to the sexual system.

Such flowers as want the stamens, and have the pistil, are termed *female*.

Those flowers which have the stamens, and want the pistils, are called *male*.

Flowers which have both stamens and pistils are said to be *hermaphrodite*.

Neuter flowers are such as have neither stamens nor pistils.

Hermaphrodite flowers are sometimes distinguished into *male hermaphrodites* and *female hermaphrodites*. This distinction takes place when, although the flower contains the parts belonging to each sex, one of them proves abortive or ineffectual; if the defect be in the stamina, it is a female hermaphrodite, if in the pistil, a male one.

Plants, in regard to sex, take also their denominations in the following manner:

1. *Hermaphrodite plants* are such as bear flowers upon the same root that are all hermaphrodite.

2. *Androgynous plants*, are such as, upon the same root, bear both male and female flowers, distinct from each other, that is, in separate flowers.

3. *Male plants*, such as bear male flowers only upon the same root.

4. *Female plants*, such as bear female flowers only upon the same root.

5. *Polygamous plants*, such as, either on the same or on different roots, bear hermaphrodite flowers, and flowers of either or both sexes.

The first general division of the whole body of vegetables is, in the sexual system, divided into twenty-four classes; these again are subdivided into orders; the orders into genera, the genera into species; and the species into varieties, where they are worthy of note.

A Table of the Classes and Orders.

CLASSES.	ORDERS.
1. Monandria.	Monogynia. Digynia.
2. Diandria.	Monogynia. Digynia. Trigynia.
3. Triandria.	Monogynia. Digynia. Trigynia.
4. Tetrandria.	Monogynia. Digynia. Tetragynia.
5. Pentandria.	Monogynia. Digynia. Trigynia. Tetragynia. Pentagynia. Pologynia.
6. Hexandria.	Monogynia. Digynia. Trigynia. Tetragynia. Pologynia.
7. Heptandria.	Monogynia. Digynia. Tetragynia. Heptagynia.
8. Octandria.	Monogynia. Digynia. Trigynia. Tetragynia.
9. Enneandria.	Monogynia. Trigynia. Hexagynia.
10. Decandria.	Monogynia. Digynia. Trigynia. Pentagynia. Decagynia.
11. Dodecandria.	Monogynia. Digynia. Trigynia. Pentagynia. Dodecagynia.
12. Icosandria.	Monogynia. Digynia. Trigynia. Pentagynia. Pologynia.
13. Polyandria.	Monogynia. Digynia. Trigynia. Tetragynia. Pentagynia. Hexagynia. Pologynia.
14. Didynamia.	Gymnospermia. Angiospermia.
15. Tetradynamia.	Siliculosa. Siliquosa.
16. Monadelphia.	Pentandria. Decandria. Enneandria. Dodecandria. Polyandria.
17. Diadelphia.	Pentandria. Hexandria.
18. Polyadelphia.	Pentandria. Icosandria. Polyandria.
19. Syngenesia.	Polygamia æqualis. Polygamia superflua. Polygamia frustanea. Polygamia necessaria. Polygamia segregata. Monogamia.
20. Gynandria.	Diandria. Triandria. Tetrandria. Pentandria. Hexandria. Decandria. Dodecandria. Polyandria.
21. Monoecia.	Monandria. Diandria. Triandria. Tetrandria. Pentandria. Hexandria. Heptandria. Polyandria. Monadelphia. Syngenesia. Gynandria.
22. Dioecia.	Monandria. Diandria. Triandria. Tetrandria. Pentandria. Hexandria. Octandria. Enneandria. Decandria. Dodecandria. Polyandria. Monadelphia. Syngenesia. Gynandria.
23. Polygamia.	Monoecia. Dioecia. Trioecia.
24. Cryptogamia.	Filices. Musci. Algæ. Fungi.
Appendix.	Palme.

Explanation of these terms.

As these terms in the Greek language, from whence they are taken, are all expressive of the principal circumstance that obtains in the class to which they are applied, the explanation of them will give the reader a good insight into the proper characters of the several classes, and the sexual distinctions on which they are founded.

Monandria; from *μῶνος*, one, and *ἄνδρ*, a husband, that is, a stamen.

Diandria; from *δύο*, two, and *ἄνδρ*, a husband.

Triandria; from *τρεῖς*, three, and *ἄνδρ*, a husband.

Tetrandria; from *τέσσαρες*, four, and *ἄνδρ*, a husband.

Pentandria; from *πέντε*, five, and *ἄνδρ*, a husband.

Hexandria; from *ἕξ*, six, and *ἄνδρ*, a husband.

Heptandria; from *ἑπτά*, seven, and *ἄνδρ*, a husband.

Octandria; from *οκτώ*, eight, and *ἄνδρ*, a husband.

Enneandria; from *εννέα*, nine, and *ἄνδρ*, a husband.

Decandria; from *δέκα*, ten, and *ἄνδρ*, a husband.

It is necessary to observe here, that the flowers must all be hermaphrodite in these classes; for should the female part be wanting, the plant would belong to some other class, notwithstanding the number of stamina may be such as would otherwise refer it to one of these.

Dodecandria; from *δωδεκά*, twelve, and *ἄνδρ*, a husband.

Notwithstanding the term implies that the flowers have twelve husbands, the class is not confined to this number, but includes all such hermaphrodite flowers as are furnished with any number of stamina, from twelve to nineteen inclusive. No flowers have yet been discovered that have eleven stamina, which is the reason no class has been allotted to that number.

Icosandria; from *ἰκκοί*, twenty, and *ἄνδρ*, a husband.

Here, again, the title is to be understood with considerable latitude; for though it means that the flowers have twenty stamens, yet the plants belonging to this class are rarely found with less, and they frequently have a greater number, and are therefore not to be known with certainty from the next class.

Polyandria; from *πολύς*, many, and *ἄνδρ*, a husband.

This class comprehends those hermaphrodite plants whose flowers have more stamens disunited than twenty.

Didynamia; from *dis*, two, and *dynamis*, power.

This term imports the *power* or *superiority* of two, and is applied to this class, because its flowers have four stamens, of which there are two longer than the rest. This circumstance alone is sufficient to distinguish this from the fourth class, where the four stamens are equal.

Tetradynamia; from *τετρας*, four, and *dynamis*, power.

This term implies the power or superiority of four, and accordingly there are in the flowers of this class six stamens, four of which are longer than the rest, which circumstance distinguishes them from those of the sixth class where they are equal.

Monadelphia; from *μῑς*, one, and *ἀδελφός*, a brotherhood.

The word here compounded with the numerical term, signifies a brother. This relation is employed to express the union of the filaments of the stamens, which in this class do not stand separate, but join at the base, and form one substance, out of which they proceed as from a common mother, and the title, therefore, expresses a single brotherhood, meaning, that there is but one set of stamens so united, which distinguishes this class from the two following. The number of stamens, it is to be recollected, is not limited.

Diadelphia; from *dis*, two, and *ἀδελφός*, a brotherhood.

This term implies a double brotherhood, or two sets of stamens, united in the manner explained in the former class. The number of stamens is not limited.

Polyadelphia; from *πολυς*, many, and *ἀδελφός*, a brotherhood.

Many brotherhoods or sets of stamens is meant by this term.

Syngenesia; from *σύν*, together, and *γενεσις*, generation.

This term implies *congeneration*, alluding to the circumstance of the stamens; in which, though the stamens stand separate, yet their anthers or tops, which are the parts more immediately subservient to generation, are united in a cylinder, and perform their office together.

Gynandria; from *γυνή*, a wife, and *ἀνρ*, a husband.

This term alludes to the singular circumstance of this class, in the flowers of which the stamens grow upon the pistil; so that the male and female parts are united, and do not stand separate, as in other hermaphrodite flowers.

Monœcia; from *μῑς*, one, and *οἶκος*, a house.

The word here compounded with the numerical term, signifies a *house* or *habitation*.

To understand the application of this title, it must be observed, that the plants of this class are not hermaphrodite, but androgynous; the flowers that have the stamens wanting the pistil, and those that have the pistil wanting the stamens, so that *monœcia* signifying a *single house*, alludes to this circumstance, that in this class the male and female flowers are both found on the same plant or house.

Diœcia; from *dis*, two, and *οἶκος*, a house.

This term signifies two houses, and is applied to this class, the plants of which are male and female, to express the circumstance of the *male* flowers being on one plant, and the *female* on another; the contrary of which is the case of the androgynous class *Monœcia*.

Polygœnia; from *πολυς*, many, and *γᾱμος*, nuptials.

This term implies plurality of marriages. This class produces, either upon the same or different plants, *hermaphrodite* flowers, and also flowers of *one sex* only, be it male or female; or flowers of each sex; and the latter receiving impregnation from, or giving it to the hermaphrodites, as their sex happens to be the parts essential to generation in the hermaphrodite flowers, do not confine themselves to the corresponding parts within the same flower, but become of *promiscuous* use, which is the reason of giving this title.

Cryptogœnia; from *κρυπτός*, concealed, and *γᾱμος*, nuptials.

This term means a concealment of marriages; the class consists, therefore, of such plants as either bear their flowers concealed within the fruit, or have them so small as to be imperceptible.

Explanation of the titles of the orders.

Monogynia; from *μῑς*, one, and *γυνή*, a woman, that is, a pistil.

Diagynia; from *dis*, two, and *γυνή*, a woman.

Trigynia; from *τρεῖς*, three, and *γυνή*, a woman.

Tetragynia; from *τετρας*, four, and *γυνή*, a woman.

Pentagynia; from *πεντε*, five, and *γυνή*, a woman.

Hexagynia; from *ἕξ*, six, and *γυνή*, a woman.

Decagynia; from *δεκα*, ten, and *γυνή*, a woman.

Polygynia, from *πολυς*, many, and *γυνή*, a woman.

These are the titles that occur in the thirteen first classes, and the general explanation of one pistil, two pistils, &c. will be sufficient to make it appear how they are employed in the class.

The class *didynamia* contains the orders,

Gynœspermia; from *γᾱμος*, naked, and *σπέρμα*, a seed.

Angiosperma; from *αγγος*, a vessel, and *σπέρμα*, a seed; which are distinguished by the seed being either naked, or enclosed in a pericarp or seed-vessel.

The two orders in the class *Tetradynamia* are founded on a distinction in the pericarp.

Siliculosæ; means a little *siliqua*.

Siliqua; which is a particular kind of seed-vessel.

To explain the orders contained in the class *Syngenesia*, viz. *Polygamia equalis*, *Polygamia superflua*, *Polygamia frustranea*, *Polygamia necessaria*, *Polygamia segregata*, *Monogamia*, it is necessary to explain what is meant by polygamy in flowers. It has been before observed, what is meant by *polygamous plants*: but, in respect to flowers, the term is applied to a single flower only, for the flowers of this class being compound, a polygamy arises from the intercommunication of the several florets in one and the same flower. Now, the *polygamy of flowers*, in this sense of the word, affords four cases, which are the foundations of the four first orders of this class: *equal polygamy*, is when all the flowers are hermaphrodite: *superfluous polygamy*, is when some of the florets are hermaphrodite, and others female only; for, in this case, as the fructification is perfected in the hermaphrodites, the addition of the females is a superfluity: *frustraneous polygamy*, is when some of the florets are hermaphrodite, and others neuter; for, in this case, the addition of the neuters is of no assistance to the fructification: *necessary polygamy*, is when some of the florets are male, and the rest female; for, in this case, there being no hermaphrodites, the polygamy arising from the composition of the florets of different sexes, is necessary to perfect the fructification: *polygamia segregata* implies separation; the plants of this order having partial cups growing out of the common calyx which surround and divide the florets: the order *Monogamia* signifies a single marriage, and is opposed to the polygamy of the four other orders; for in this, although the anthers are united, which is the essential character of the flowers of this class, the flower is simple, and not compounded of many florets, as in the other orders.

The title of the other order to that of *Trioecia*, in the class *Polygamia*, have already been explained.

Trioecia; from *τρεῖς*, three, and *οἶκος*, a house; because the polygamy is on three distinct plants, one producing male flowers, another female, and a third hermaphrodite or androgynous.

The class *Cryptogamia* contains the orders of

Filices, or ferns;

Musci, or mosses;

Alga, or flags;

Fungi, or mushrooms.

This short explanation of the Linnæan system has been introduced, in order to convey a general idea to medical students of its nature, and also the meanings of the several terms.

The various medicinal plants will be found systematically arranged under the title *Materia Medica*.

Srydschults water. See *Sedlitz water*.

Shullos. A species of onion.

Sharp-pointed dock. See *Oxylapathum*.

Shedding teeth. The primary or milk-teeth. See *Teeth*.

Shells, prepared. See *Testa preparata*.

Sherbes. A compound liquor prepared for punch before the spirit is added.

Shingles. See *Erysipelas*.

SHRUB. A compound prepared from spirits, lemon-juice and sugar.

SIABON. *Σιαβον*. The jaw

SIABONAGRA. (From *σιαβον*, the jaw, and *αγρα*, a seizure.) The gout in the jaw.

SIALAGOGUES. (*Medicamenta Sialagoga*; from *σιαλαγωγα*, saliva, and *αγω*, to expel.) Those medicines are so called, which excite an uncommon flow of saliva: such are mercurial preparations, pyrethrum, &c. They are divided into *sialagoga topica*, as scilla, nicotiana, piper, &c. and *sialagoga interna*, as the various preparations of mercury.

Sibbens A disease resembling syphilis.

SICCANTIA. (From *sicco*, to dry.) Drying medicines.

SICCABIA. (From *συχος*, weak, weary.)

An unpleasant lassitude and debility peculiar to women with child.

SICULA. (Dim. of *sica*, a short sword; so called from its dagger-like root.) The beet.

SICYDOK. (From *σικυος*, a cucumber.)

A transverse fracture like a cucumber broken in two parts.

SICYONE. (From *σικυος*, a cucumber or gourd; so named from its resemblance to a gourd.) A cucurbit.

SIDERATIO. (From *sidus*, a planet, because it was thought to be produced by the influence of the planets.) An apoplexy; a blast: a slight erysipelas.

SIDERIUM. (From *sidus*, iron;) an herb so called from its supposed virtues in healing wounds made by iron instruments.

SIGESBECKIA ORIENTALIS. The systematic name of a plant which is said to be useful in removing strangury, and in calculous diseases, gout, and fluor albus.

SIGHT. See *Sreing*.

SIGILLATA TERRA. Sealed earth; a species of holar earth made into cakes.

SIGILLUM BEATÆ MARIE. Black briony.

SIGILLUM HERMETICUM. An hermetic seal; mad: by closing the end of a glass tube by melting it.

SIGILLUM SALOMONIS. (Dim. of *signum*, a sign. It is called *sigillum salomonis*, Solomon's seal, because it has upon its root the resemblance of an impression made by a seal.) Solomon's seal. *Convolutus polygonatum* of Linnæus:—*foliis alternis amplexicaulis, caule ancipiti, pedunculis axillaribus subumbellatis*. The roots are applied externally as adstringents, and are administered internally as corroborants.

SIGMOID. (*Sigmoides*, from the Greek letter Σ , and *ωμοι*, a likeness; resembling the Greek letter sigma.) Applied to the valves of the heart, and sometimes to the cartilages of the aspera arteria, or the semilunar apophysis of the bones.

SIGMOIDEA FLEXURA. The sigmoid flexure, or turn of the colon.

SIGMOIDES RACIOSUS. Valves of the heart.

SIGNA CRITICA. Signs of the crisis of disease.

SIGNA DIAGNOSTICA. Diagnosis or distinguishing signs.

SILEX MONTANUM. See *Silex*.

SILEX MONTANUM. Common hartwort.

SILEX. (*Silag*, Heb.) Silix, or siliceous earth, is the principal constituent part of a very great number of the compound earths and stones forming the immense mass of the solid nucleus of the globe. It is the basis of almost all the scintillating stones, such as *flint*, *rock crystal*, *quartz*, *agate*, *calcedony*, *jasper*, &c. The sand of rivers and of the seashore, chiefly consist of it. It is deposited in vegetable substances forming petrified wood, &c. It is likewise precipitated from certain springs in a stactical form. It has been discovered in several waters in a state of solution, and is found in many plants, particularly grasses and equisetums. Professor Davy has proved that it forms a part of the epidermis of these vegetables. It is never met with absolutely pure in nature.

Properties.—Silix, when perfectly pure, exists in the form of a white powder. It is insipid and inodorous. It is rough to the touch, cuts glass, and scratches or wears away metals. Its specific gravity is about 2.66. It is unalterable by the simple combustible bodies. When mixed with water it does not form a cohesive mass. Its molecules when diffused in water are precipitated with the utmost facility. It is not acted on by any acid, except the fluoric. When in a state of extreme division it is soluble in alkalis; fused with them it forms glass. It melts with the phosphoric and boracic acids. It is unchangeable in the air, and unalterable by oxygen and the rest of the gaseous fluids. It has been considered as insoluble in water, but it appears when in a state of

extreme division to be soluble in a minute quality.

Method of obtaining silix.—Silix may be obtained, tolerably pure, from flints, by the following process: procure some common gun-flints; expose them in a crucible to a red heat, and then plunge them into cold water; by this treatment they will become brittle, and easily reducible to powder. Mix them, when pulverized, with three or four times their weight of carbonate of potash, and let the mixture be fused, in a dull red heat, in a silver crucible. We shall thus obtain a compound of alkali and silix, called siliceous potash. Dissolve this compound in water, filter the solution, and add to it diluted sulphuric or muriatic acid. An immediate precipitation now ensues, and as long as this continues, add fresh portions of acid. Let the precipitate subside; pour off the fluid that floats above it; and wash the precipitate with hot water till it comes off tasteless. This powder when dry is silix.

In this process the acid added to the solution of flint unites to the potash, and forms sulphate or muriate of potash; the siliceous earth is therefore precipitated.

It is necessary to add an excess of acid, in order that all the foreign earths which are present may be separated.

If the solution of flints be diluted with a great quantity of water, as for instance, in the proportion of 24 parts to one, and in this state an acid be poured upon it, no perceptible precipitation will ensue; the silix continues suspended in the fluid, and is invisible on account of its transparency; but it may be made to appear by evaporating part of the water.

The solution of flint, on account of its affinity with the carbonic acid is also in course of time decomposed by mere contact with air.

Another method of obtaining silix exceedingly pure is to separate it from fluoric acid.

SILICA. (From *silex*.) Siliceous earth. See *Silex*.

SILIGO. *Σιλιγίς*. Fine wheat or rye.

SILIQUA. (From *silo*, a nose turned up, a hooked nose.) A pod or receptacle for seed, consisting of two valves, and in which the seeds are fixed alternately to each suture. Also some plants which bear pods.

SILIQUA DULCIS. Sweet-pod. The fruit so called is the produce of the *Ceratonia siliqua* of Linnæus. They are about four inches in length, and as thick as one's finger, compressed and unequal, and mostly bent; they contain a sweet brown pulp, which is given in form of decoction, as a pectoral in asthmatic complaints and coughs.

SILIQUA HIRSUTA. The cowage is sometimes so called. See *Dolichos*.

SILICUASTRUM. (From *siliqua*, a pod:

named from its pods.) Judas-tree. The capsicum or Guinea-pepper was so termed by Pliny. See *Piper indicum*.

SILVQUOSA INDICA. An American plant; its juice is alexipharmic.

Silk-worm, acid of. See *Bombic acid*.

SILPHIUM SUMMI. (Zalaph, Arab.) Assafetida, or the plant which affords it.

SILVER. *Argentum*. This metal is found both native and mineralized, and combined with lead, copper, mercury, cobalt, sulphur, arsenic, &c. The principal ores of this metal are the following: Native silver, antimoniated silver, sulphuret of silver, sulphurated oxid of silver and antimony, muriate of silver, native oxid of silver, &c. It is found in different parts of the earth. The mines of the Erzgebirge or the metalliferous rocks of Mexico and Potosi, Bohemia, Norway, Transylvania, &c. are the richest.

Native silver possesses all the properties of this metal, and it appears in series of octahedra inserted in one another; in small capillary flexible threads intertwined together; in plates; or in masses. The colour of native silver is white, often tarnished. Silver alloyed with gold forms the *auriferous native silver ore*. The colour of this ore is a yellowish white. It has much metallic lustre. The *antimoniated silver ore* belongs to this class. Silver combined with sulphur, forms the *sulphurated oxid of silver*, or *vitreous silver ore*. This ore occurs in masses, sometimes in threads, and sometimes crystallized in cubes or regular octahedra. Its colour is dark bluish grey, inclined to black. Its fracture is uneven, and its lustre metallic. It is soft enough to be cut with a knife. It is sometimes found alloyed with antimony (grey silver ore.) Silver united to muriatic acid forms the *corneous silver ore* (*muriate of silver*), which appears under different colours and shapes. Silver united to oxygen constitutes the *calcareous silver ore*, of which there are several varieties. The colour of these ores is a lead grey, or greyish black. They occur massive, disseminated, and crystallized.

Germany, and other countries of Europe, but more especially Peru and Mexico in South America, contain the principal silver mines. There are, however, silver mines in Ireland, Norway, France, and many other parts of the world.

Properties.—Pure silver is very brilliant, white and sonorous. It is the most splendid of all the metals. Its specific gravity is from 10.474 to 11.091, according to the state of its density. It is exceedingly ductile and tenacious. It may be beat out into leaves only one hundred and sixty thousandth part of an inch thick, and drawn into wire the thousandth part of an inch thick. It melts at 28° of Wedgwood's

pyrometer. Exposed to a temperature considerably higher it becomes volatilized. Atmospheric air has no effect upon it, except when it contains sulphureous vapours, sulphurated or phosphorated hydrogen gases. It unites to phosphorus and sulphur. It slightly unites with the brittle acidifiable metals; but it readily enters into combination with the greater number of the other metals. With gold it forms what is termed *green gold*. Copper renders it harder without much impairing its ductility. Mercury and silver combine and form a crystallizable and fusible alloy. It unites with the rest of the metals except cobalt and nickel. It is oxidated and dissolved by several of the acids. The nitric acid attacks it rapidly in the cold. The sulphuric acid requires a boiling heat. The muriatic acid does not act upon it. The acid solutions of silver are decomposable by the alkalis, earths, and by the greater number of the metals.

Method of obtaining silver.—Different methods are employed in different countries to extract silver from its ores. In Mexico, Peru, &c. the mineral is pounded, roasted, washed, and then triturated with mercury in vessels filled with water. A mill is employed to keep the whole in agitation. The silver combines by that means with the mercury. The alloy thus obtained is afterwards washed, to separate any foreign matters from it, and then strained and pressed through leather. This being done, heat is applied to drive off the mercury from the silver, which is then melted and cast into bars or ingots.

In order to extract silver from sulphurated or vitreous silver ore, the mineral is roasted, and then melted with lead and borax, or some other flux to assist the fusion. By the first operation the sulphur is volatilized, and by the second the silver is obtained, though for the most part alloyed with the other metals, from which it is separated by cupellation, or fusion with lead or bismuth.

A perfect metal, of a white colour, and of the most lively brilliancy; next to gold, the most malleable of all metals. It is sometimes found pure, but for the most part in combination with tin or lead. It has nether taste nor smell; its specific gravity is such, that it loses about the eleventh part of its weight by immersion in water; and a cubic foot of this metal weighs 370 pounds. Native silver is found in the greatest abundance in Peru and Mexico. From this metal is obtained the officinal *Argentum nitras*.

Silver-weed. See *Potentilla*.

SIMAROUBA. (A patronymic name of America.) *Simarouba Euonymus*. *Simarouba quassia*. *Quassia simarouba* of Linnæus:—*floribus monoicis, foliis abrupte pinnatis, foliolis alternis subpetiolatis petiolo*

nuda, floribus paniculatis. The bark of this tree, which is met with in the shops, is obtained from the roots; and, according to Dr. Wright of Jamaica, it is rough, scaly, and warted; the inside, when fresh, is a full yellow, but when dried paler: it has but little smell; the taste is bitter, but not disagreeable. It is esteemed, in the West Indies, in dysenteries and other fluxes, as restoring tone to the intestines, allaying their spasmodic motions, promoting the secretions by urine and perspiration, and removing looseness of spirits attending those diseases. It is said also that it soon disposes the patient to sleep; takes off the gripes and tenesmus, and changes the stools to their natural colour and consistence.

SINAROURUM INFUSUM. See *Infusum sinaroubae*.

SINNA LAPI. See *Bryozoa sinia*.

Simple substances. See *Elements*.

SIMPLEX OCVLUS. A bandage for the eye.

SINAPIE. See *Sinapis*.

SINAPHELIUM. (From *sinari*, mustard, and *saeny*, oil.) Oil of mustard.

SINAPROS SEMEN. Mustard-seed.

SINAPI NIGRUM. See *Sinapis*.

SINAPIS. (*Οτι ονυχας ονυχας*, because it hurts the eyes.) *Erucos. Nypus*. 1. The name of a genus of plants in the Linnaean system. Class, *Tetradynamia*. Order, *Siliqueosae*. Mustard.

2. The pharmacopoeial name of the black mustard. *Nypus. Eruca. Sinape. Sinapi.* Common black mustard. *Sinapis nigra* of Linnaeus:—*siliquis glabris racemo appressis*. The seeds of this species of mustard, which are directed by the London College, and those of the *Sinapis alba*, which are preferred by that of Edinburgh, manifest no remarkable difference to the taste, nor in their effects, and therefore answer equally well for medicinal and culinary purposes. They have an acrid, pungent taste, and, when bruised, this pungency shews its volatility by powerfully affecting the organs of smell. Mustard is considered as capable of promoting appetite, assisting digestion, attenuating viscid juices, and, by stimulating the fibres, it proves a general remedy in paralytic affections. Joined to its stimulant qualities, it frequently, if taken in considerable quantity, opens the body, and increases the urinary discharge, and hence it has been found useful in dropsical complaints. Externally, flower of mustard is frequently used mixed with vinegar as a stimulant or sinapism.

SINAPIS ALBA. The systematic name of the white mustard plant, which is directed for medicinal use in the Edinburgh Pharmacopoeia. It is somewhat less pungent than the black species. See *Sinapis*.

SINAPIS NIGRA. The systematic name

of the common black pepper. See *Sinapis*.

SINAPIS SEMEN ALBUM. White mustard-seed.

SINAPIS SEMEN NIGRUM. See *Sinapis*.

SINAPISMUS. *Sinapismus. Cataplasma sinapis.* A sinapism or mustard poultice. A term given to a mixture of mustard and vinegar in form of poultice, generally applied to the calves of the legs, and soles of the feet as a stimulant, and employed in low states of fevers and other diseases, and intended to supersede the use of a blister.

SINAPIUM. (From *sinari*, mustard.) An infusion or decoction of mustard-seed.

SINCIPUT. The fore part of the head.

See *Caput*.

SINE PARI. Several muscles, veins, arteries, &c. are so called which are without a fellow. See *Azygos*.

SINGULTUS. *Lygmos.* The hiccough. A convulsive motion of the diaphragm and parts adjacent.

SINUS. 1. A cavity or depression.

2. In surgery it means a long, narrow, hollow track, leading from some abscess, diseased bone, &c.

3. The veins of the dura mater are so termed. They are several in number, the principal of which are, 1. The *longitudinal sinus*, which rises anteriorly from the crista galli, ascends and passes between the two laminae of the falciiform process to where this process ends. It then opens into, 2. *Two lateral sinuses*, distinguished into right and left, which lie in the crucial apine of the os occipitis: 3. The *inferior longitudinal*, which is a small sinus situated at the acute inferior margin of the falx.

SINUS COXA. The sinus of the coccyx.

SINUS CERNI PITUITARIUS. See *Antrum of Highmore*.

SINUS LONGITUDINALIS. See *Longitudinal sinus*.

SINUS MAXILLARIS. The antrum of Highmore. A cavity in the cheek.

SINUS MUXIBERIS. Sinus pudoris. The vagina.

SINUS VENÆ PORTARUM. The entrance into the liver.

SINUSES LATERAL. See *Lateral sinuses*.

SIPHILIS. See *Syphilis*.

SIPHONIA ELASTICA. The systematic name of the elastic resin-tree. See *Indica rubber*.

SIRIASIS. (From *σῆσις*, a cavity.) An inflammation of the brain peculiar to children, and attended with a hollowness of the eyes and depression of the fontanella.

SIRIUM MYRTIFOLIUM. The systematic name of the tree which is supposed to

afford the yellow Saunders. See *Santalum album*.

SISARUM. (*Sisa*, Heb.) Siser of skirret.

SISER. See *Sisarum*.

SISON AMMI. The systematic name of the plant which affords the amomum verum of the shops. See *Amomum*.

SISYMBRIUM. (From *σισυβος*, fringe; so named from its fringed roots.) The name of a genus of plants in the Linnæan system. Class, *Tetradymania*. Order, *Siliquosa*. The water-cress.

SISYMBRIUM NASTURTIUM. The systematic name of the water-cress. See *Nasturtium aquaticum*.

SISYMBRIUM SOPHIA. The systematic name of the herb sophia. See *Sophia chirurgorum*.

SITIOLOGY. (*Sitiologia*. From *σitos*, aliment, and *λογος*, a discourse or treatise.) A doctrine or treatise on aliment.

SIUM. (From *siu*, to move, from its agitation in water.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digynia*.

2. The pharmacopœial name of the creeping water-parsnep, *Sium nodiflorum* of Linnæus. This plant is admitted into the London Pharmacopœia in the character of an antiscorbutic. It is not nauseous, and children take it readily if mixed with milk.

SIUM AROMATICUM. The amomum verum is sometimes so called. See *Amomum*.

SIUM NANSL. The systematic name of the plant whose root is called *radix nansl* in some pharmacopœias.

SIUM NODIFLORUM. The systematic name of the creeping water-parsnep. See *Sium*.

SKELETON. (*Sceletus*, from *σκελετος*, to dry.) When the bones of the body are preserved in their natural situation, and deprived of the flesh, the assemblage is called a skeleton. See *Bones*.

SKELETON, ARTIFICIAL. The assemblage of all the bones of the animal, when hung in their respective situations by means of wire. See *Bone*.

SKELETON, NATURAL. A skeleton is so termed in opposition to an artificial one, when the bones are retained in their proper places by means of their natural ligaments.

SKIN. (*Δερμς*. *Pellis*. *Cutis*.) When carefully dissected off and separated from all adventitious matter in a middle-sized man, it weighs about four pounds and a half.

The skin, though apparently a simple membrane, is in reality laminated, consisting of several subdivisions; the outermost lamina is termed with us scarf skin, or cu-

ticle; the second has no English name, is known only to anatomists, and is called *rete mucosum*; after these two are removed we come to, as is commonly thought, the surface of the skin itself.

When a blister has been applied to the skin of a Negro, if it has not been very stimulating, in twelve hours after a thin transparent greyish membrane is raised, under which we find a fluid. This membrane is the cuticle or scarf skin. When this, with the fluid, is removed, the surface under them appears black; but if the blister had been very stimulating, another membrane, in which this black colour resides, would also have been raised with the cuticle; this is *rete mucosum*, which is itself double, consisting of another grey transparent membrane, and of a black web, very much resembling the *nigrum pigmentum* of the eye. When this membrane is removed, the surface of the true skin (as has hitherto been believed) comes in view, and is white, like that of a European. The *rete mucosum* gives the colour to the skin; is black in the Negro; white, brown, or yellowish, in the European. The reason why this membrane is black in the Negro is, perhaps, that his body may be better able to defend itself against the sun's rays, and that the heat may be prevented from penetrating. The intention of a similar membrane behind the retina in the eye, appears to be not only that of absorbing the superfluous rays of light; but, like the *amalgam* behind the looking-glass, it may enable the retina to reflect the rays, in order to perfect vision. It is not very improbable that some such purpose, as enabling the cuticle to reflect the sun's rays in those warm climates, where the inhabitants originally go naked, may be the intention of nature, in giving them the black membrane. Perhaps too, the circumstance of the countenance's becoming brown, when exposed to the sun's rays in summer, in our own climate, may be a process of nature to defend herself against the access of external heat into the body.

Both cuticle and *rete mucosum* send innumerable processes into the pores of the true skin; the process of the *rete mucosum* is always within that of the cuticle, and in contact with the sides of the pore, as formed by the true skin. These processes are remarkable in the cuticle and *rete mucosum* of the elephant, some of them are almost an inch long; the cuticle, or *rete mucosum*, or a membrane very similar, having the same properties with these, appears to be also continued into the inside of the mouth, over the tongue, internal surface of the lungs, œsophagus, stomach, and intestinal tube. In most of the last-named parts, the cuticle, however, forms

sheaths for *villi*, and not processes which line pores. On viewing the surface of the skin, even with the naked eye, we find it porous; more so in some places than in others; and the pores are also larger in some parts than others. These pores are ducts of sebaceous glands, and serve not only to transmit hairs, but, it is supposed, the greatest part of the perspirable matter itself. Absorption on the skin also, in all probability, begins on the sides of these pores. They are particularly remarkable about the mouth, nose, palms of the hands, soles of the feet, on the external ear, scalp, *mons veneris*, and around the nipple in women.

The skin itself was given to man not only for feeling, in a general sense, but for perspiration, absorption, and particularly for *touch*, in which he excels all other animals, and which resides, principally in the *tips of the fingers*. He was intended for examining, reasoning, forming a judgment, and acting accordingly; he was fitted by this sense to examine accurately the properties of surrounding bodies, not capable of being examined by his other senses. This, among other reasons, was one why he was made erect, that the points of his fingers should not be made callous, or less sensible, by *walking on them*.

The skin of human bodies is always of a white colour, in the dead body, let the colour of the *rete mucosum* be what it may, it is extremely full of pores, and extremely vascular; a child in full vigour comes into the world, from this circumstance, *scarlet*; it is endowed with intense sensibility: almost all the pain, in the different operations of surgery, is past when we have divided the skin. Some parts of the skin have more feeling than others; the lips, for example, as Haller says "*ad basia destinata*." The *glans clitoridis*, and the *glans penis*, with a similar intention; there, though the nerves are not so large as in some other parts, they are longer, more numerous, and endowed with more exquisite feeling; but where the common offices of life merely are intended, the marks of superior feeling or touch, in the skin, are the projections, above the common surface, of those packets of arteries, veins, absorbents, called *villi*; the nerves are there not only also longer, but larger, as in the points of the fingers and toes.

We are not certain that the skin is muscular, but it has properties very like those of muscle, it contracts, relaxes, and even vibrates, in some places, on certain occasions. It is extremely distensible, the skin of the *perineum* has stretched in labour from a quarter of an inch to six inches. It is also extremely elastic, and instantly after labour has returned again to the ori-

ginal quarter of an inch; it is thickest on those parts intended by nature to bear weight or pressure; of course it is thickest on the back, on the soles of the feet, and palmar of the hands. It is thinner on the fore part of the body, on the insides of the arms and legs, and where its surfaces touch opposite surfaces. It is extremely thin on the lips, and allows the colour of the blood to shine through it. It is also extremely thin on the *glans penis* in men, *glans clitoridis* in women, and on the inside of the *labia pudendi*. Skin dried and dressed is extremely strong and durable, and therefore employed in making harness for horses, clothing for men, and a variety of other purposes.

Skin, scarf See *Cuticle* and *Skin*.

Shink See *Scincus*.

SKULL. The skull or *cranium* is that bony box which contains the brain: it forms the forehead, and every part of the head except the face. It consists of eight bones, namely one *os frontis*, one *os occipitis*, one *os sphenoides*, one *os ethmoidale*, two *ossa temporalia*, and two *os parietalia*.

Sliters. See *Millepedes*.

SLEEP. *Somnus.* That state of the body in which the internal and external senses and voluntary motions are not exercised. The end and design of sleep is both to renew, during the silence and darkness of the night, the vital energy which has been exhausted through the day, and to assist nutrition.

Sloe. See *Prunus sylvestris*.

Smallage. See *Asium*.

Small-pox See *Variola*.

SMELLING. The sense of smelling is performed by means of a soft, pulpy, vascular, papilous, porous membrane, which lines the whole internal cavity of the nostrils, and is thicker upon the septum, and principal cavity of the nose, but thinner in the sinuses. It is plentifully supplied with very soft nerves, the middle one of which descend from the first pair, through the holes of the *os cribosum* to the *septum narium*; but in such a manner, that it is very difficult to trace them to their extremities and into the septum. Other lateral nerves come from the second branch of the fifth pair and its branches, from that which crosses the pterygoid canal, and from another which descends through the canals of the palate; and in the maxillary sinus from the infra orbital branch, from the dental branch, and from the anterior nerve of the palate. The anterior part of the septum has a twig from the ophthalmic of the first branch of the fifth pair.

The nostrils are supplied with very numerous arteries; from the three nasal branches of the internal maxillary, above,

both from the ethmoidal branches, and the frontal and nasal branches, with lateral arteries from the smaller ophthalmic branch of the internal carotid, and from branches of the palatine artery, and in the sinuses from the infra-orbital, and from the superior dental one. These arteries have the property of exuding blood easily, and in great quantity, without any lesion of consequence. The correspondent veins form a very large plexus upon the external pterygoid muscle; then communicate with the sinuses of the dura mater; and, lastly, meet in the external branch of the internal jugular. The arteries supply nourishment, warmth, and mucus.

The air, filled with the very subtle, invisible, pungent, oily, saline, and volatile effluvia, which exhale from almost every known body, being received into the nostrils, by the action of respiration, and by a peculiar effort for drawing the air into them, carries these particles to the nerves, widely naked, and constantly soft. By these there is excited in the nerves a kind of sensation which we call smell, by which we distinguish the several kinds of oils and salts, in a manner somewhat indistinct, difficultly reducible to classes, difficultly recalled to the memory, nevertheless sufficiently for our purposes. This sense informs us of unwholesome putridity, of excessive acrimony, and of the bland and useful nature of substances. And as salt, united with oil, is an object of taste, and as oils, combined with salts, constitute odours, the affinity of the two senses, which is necessary to derive utility from either, is apparent. But volatile particles chiefly are distinguished by smell, and fixed ones by the taste; perhaps because the thick mucous cuticle, spread over the tongue, intercepts the action of the more subtle salts, which easily affect the softer and less covered nerves of the nostrils. We are ignorant of the reasons why some smells please, and others displease; perhaps custom may have some influence in this respect.

The action of smells is strong, but of short continuance; because particles in a very minute state are applied to naked nerves, in the immediate vicinity of the brain. Hence the deleterious and refreshing actions of odours, by which people are resuscitated from faintings, and even from drowning. Hence the violent sneezing, excited by acrid particles, the evacuation of the bowels, by the smell of purgatives, and the power of antipathics. Hence the pernicious effects of excessive sneezing, more especially blindness, from the great sympathy of the nerves. Amongst the various parts of the nostrils, the septum, and the ossa turbinata, and their anterior portions, especially form the organ of smell: since these parts are multiplied

in quick-scented animals, forming beautiful spires in quadrupeds; and in fish, being distributed in parallel laminae elegantly toothed.

SMILAX. (From *σμίλας*, to cut; so called from the roughness of its leaves and stalk.) The name of a genus of plants in the Linnæan system. Class, *Diœcia*. Order, *Octandria*. Rough bind-weed.

SMILAX, CHINA. The systematic name of the China root tree. See *China*.

SMILAX, CHINESE. See *China*.

SMILAX SARSAPARILLA. The systematic name of the plant which affords the sarsaparilla. See *Sarsaparilla*.

SMYERNION HORTENSE. The masterwort has been so termed. See *Imperatoria*.

SMYERNION OLIVACEUM. The systematic name of the plant called Alexanders. See *Hippocelinum*.

Snail. See *Limax*.

Snakeroot, Virginian. See *Serpentaria Virginiana*.

Snailseeded glasswort. See *Salsola kali*.

Snakeweed. See *Bistorta*.

Snakerwood. See *Colubrinum lignum*.

Sneezewort. See *Ptarmica*.

SNEEZING. A convulsive action of the muscles of the chest from irritation of the nostrils.

Snuff. See *Nicotiana*.

Soap. See *Sapo*.

Soap-berry. See *Sapœnaria nucule*.

Soupwort. See *Sapœnaria*.

Socotrine aloes. An epithet of the best aloes which are brought from Socotria. See *Aloes*.

SODA. (An Arabian word.) The name now generally given by chemists and physicians to the mineral alkali.

It is obtained from several sources, but principally from plants growing on the sea coast. See *Natron* and *Burilla*. It occurs in the mineral kingdom, united with sulphuric, muriatic, and boracic acids; it is also found in large quantities in Egypt, combined with carbonic acid. It appears to be deposited in large impure masses, under the surface of the earth, in various countries, from which it is extracted by running water. Thus it is found after the spontaneous evaporation of the water, mixed with sand in the bottom of lakes in Hungary; in the neighbourhood of Bilin in Bohemia; and in Switzerland. It occurs also in China, and near Tripoli in Syria, Egypt, Persia, and India. It frequently oozes out of walls and crystallizes on their surface. Like potash, it is procured by lixiviation from the ashes of burnt plants, but only from those which grow upon the sea shores. The variety of plants employed for this purpose is very considerable. In Spain, soda is procured from the different species of the *salsola*, *salicornia* and *batis maritima*. The *zostera maritima* is burnt in some pla-

ces on the borders of the Baltic. In this country we burn the various species of *fuci*; and in France they burn the *chenopodium maritimum*.

The alkali thus procured is more or less pure according to the nature of the particular plant from which it is obtained. The greatest part however is a subcarbonate of soda.

In order to obtain it in a state of purity, the carbonate of soda must be treated, like potash of commerce, with lime and ardent spirit.

Properties of Soda.—Soda differs particularly from potash by the following properties:—In the fire it is rather more fusible. When exposed to the contact of the air it attracts water and carbonic acid, but it does not liquify like potash, it merely acquires a pasty consistence, and at last crumbles into powder. It is not altered by light. It attracts sulphur and sulphurated hydrogen more feebly. It adheres less strongly to the acids. It fuses and dissolves alumine more easily. All its other properties, its volatilization by a very high degree of heat, its acrid causticity, its solubility, its combinations with sulphur, &c. resemble those of potash.

Mr. Davy, having discovered the composition of potassa, submitted soda likewise to the same modes of analysis, and discovered that it too contains a substance of very singular properties, analogous to the base of potassa, though still essentially different from it, and that this base combined with oxygen forms soda.

When soda was submitted to the Galvanic action, in the mode already described with regard to potassa, the results were similar, globules of a metallic appearance were produced at the negative surface, which often burnt at the moment of their formation, and sometimes exploded with violence, separating into smaller globules which darted through the air in a state of vivid combustion. When these were produced an æriform fluid was disengaged at the positive surface, which proved to be pure oxygen. The reproduction of soda from this substance was similar to that of potassa from the base of the alkali. When the base of soda was exposed to the air, a crust of alkali formed on its surface, and oxygen was absorbed. When heated, confined in a portion of oxygen gas, a rapid combustion with a brilliant white flame took place, and soda was produced in the state of a solid white mass. The theory of the decomposition of soda is the same with that of potassa. The combustible base, like other combustible substances, is repelled by positively electrified surfaces, and attracted by negatively electrified surfaces; and the oxygen follows the contrary order. Hence their separation and evolution.

From the results of the combustion of the base of soda in oxygen gas, Mr. Davy inferred that one hundred parts of soda consist of eighty of base and twenty of oxygen. From the results of its oxidation by the decomposition of water, the proportions were estimated to be seventy-seven of base and twenty-three of oxygen. The mean proportions of these two modes are 78.5, and 21.5. The base of soda therefore combines with rather a larger proportion of oxygen than the base of potassa.

To the base of soda Mr. Davy from its analogy to the metals has given the name of

SODIUM.

The following are its properties:—

Sodium is white, opaque, and, when examined under a film of naphtha, has the lustre and general appearance of silver. It is exceedingly malleable, and is much softer than any of the common metallic substances. When pressed upon by a platina blade, with a small force, it spreads into thin leaves, and a globule of the one-tenth or one-twelfth of an inch in diameter is easily spread over a surface of a quarter of an inch, and this property does not diminish when it is cooled to 32° Fahrenheit.

It conducts electricity and heat in a similar manner to the basis of potassa; and small particles of it inflame by the Galvanic spark, and burn with bright explosions.

Its specific gravity is less than that of water. It swims in oil of sassafras of 1.096 water being one, and sinks in naphtha of specific gravity 861. By mixing together oil of sassafras and naphtha, until a fluid was formed in which the globule of sodium remained at rest above or below, Mr. Davy found its specific gravity to be to that of water nearly as nine to ten, or more accurately as 9348 to 1.

Sodium has a much higher point of fusion than potassim, its parts begin to lose their cohesion at about 120° Fahrenheit, and it is perfectly fluid at about 180°, so that it readily fuses under boiling naphtha, it is also less volatile, it remains fixed in a state of ignition at the point of fusion of plate glass.

When sodium is exposed to the atmosphere it immediately tarnishes, and by degrees becomes covered with a white crust, which deliquesces much more slowly than the substance that forms on the basis of potassa. This crust is pure soda.

The basis of soda combines with oxygen slowly without any luminous appearance, at common temperatures, and, when heated, this combination becomes much more rapid, but no light is emitted until it has acquired a temperature nearly that of ignition. The flame that it produces in oxygen gas is white, and it sends forth bright sparks, occasioning a very beautiful effect;

in common air it burns with light of the colour of that produced during the combustion of charcoal, but much brighter.

When thrown upon water it produces a violent effervescence with a loud hissing noise, it combines with the oxygen of the water to form soda, which is dissolved, and its hydrogen is disengaged. In this operation there is no luminous appearance; but when it is thrown into hot water, the decomposition is more violent, and in this case a few scintillations are observed at the surface of the fluid, owing to small particles of the base, which are thrown out of the water sufficiently heated to burn in passing through the atmosphere. And when a globule is brought in contact with a small particle of water, or with moistened paper, the heat produced (there being no medium to carry it off rapidly,) is usually sufficient for the ascension of the sodium.

The base of soda, when thrown upon the strong acids, acts upon them with great energy. When nitrous acid is employed, a vivid inflammation is produced; with muriatic and sulphuric acid there is much heat generated, but no light. When plunged beneath the surface of the acids, it is rapidly oxygenated; soda is produced and combines with the acid.

Sodium, in its degrees of oxidation, has precisely similar habits with the base of potassa. When it is fused with dry soda in certain quantities, there is a division of oxygen between the alkali and the base, and a deep brown fluid is produced, which becomes a dark grey solid on cooling, and which attracts oxygen from the air, or which decomposes water, and becomes soda.

The same body is often formed in the analytic processes of decomposition, and it is generated when the basis of soda is fused in tubes of the purest plate glass.

Sodium, when heated with hydrogen gas, does not, like potassium, appear to be dissolved, and as no luminous appearance attends the oxidation of it in cold water, it does not appear to be capable of combining even with nascent hydrogen.

There is scarcely any difference in the visible phenomena of the agencies of the basis of soda, and that of potassa on sulphur, phosphorus, and the metals.

It combines with sulphur in close vessels filled with the vapour of naphtha with great vividness, with light, heat, and often with explosion from the vaporization of a portion of sulphur, and the disengagement of sulphuretted hydrogen gas. The sulphuretted base of sulphur is of a deep grey colour. The phosphuret has the appearance of lead, and forms phosphate of soda by exposure to the air, or by combustion.

Sodium combines with the metals; in the quantity of one-fortieth, it renders mercury a fixed solid of the colour of silver, and

the combination is attended with a considerable degree of heat. It makes an alloy with tin, without changing its colour, and it acts upon gold and lead when heated. In its state of alloy it is soon converted into soda by exposure to air, or by the action of water, which it decomposes with the evolution of hydrogen. The amalgam of mercury and sodium seems to form triple compounds with other metals. It likewise combines with sulphur, and forms a triple compound of a dark grey colour.

SODA ACETATA. A neutral salt formed of a combination of acetic acid with the mineral alkali. Its virtues are similar to those of the acetate of potash.

SODA BORAXATA. See *Borax*.

Soda, carbonates of. See *Soda carbonas*.

SODA HISPANICA. See *Barilla*.

SODA HISPANICA PURIFICATA. See *Soda carbonas*.

SODA IMPURA. Impure soda. See *Natron*, *Barilla* and *Soda*.

SODA MURIATA. See *Murias sode*.

SODA MURIATICA. See *Murias sode*.

SODA PHOSPHORATA. Phosphorated soda. *Alkali minerale phosphoratum* of Bergman. This preparation is a phosphate of soda, and therefore called *phosphas sode* in the new chemical nomenclature. It is cathartic in the dose of half an ounce; dissolved in gruel it is not unpleasant, and is said to be useful, in scrophula, bronchocele, rachitis, and gout, in small doses.

Soda, subcarbonate of. See *Soda subcarbonas*.

Soda, subcarbonate of, dried. See *Soda subcarbonas exsiccata*.

Soda, sulphate of. See *Soda sulphas*.

SODA TARTARIZATA. Tartarized soda, formerly known by the names of *sal rupellensis*, *sal polychrestum Seignetti*, and lately by that of *natron tartarizatum*. "Take of subcarbonate of soda twenty ounces; supertartrate of potash, powdered, two pounds; boiling water ten pints." Dissolve the supercarbonate of soda in the water, and add gradually the supertartrate of potash; filter the solution through paper, and evaporate it until a pellicle forms upon the surface; then set it by that crystals may form. Having poured away the water, dry these crystals upon bibulous paper. This salt consists of tartaric acids, soda, and potash, the soda only combining with the superabundant acid of the super salt; it is therefore a triple salt, and it has been judged by the London College more convenient to express this difference by the adjective *tartarizata*, than to introduce the three words necessary to its description. It possesses mildly cathartic, diuretic, and deobstruent virtues, and is administered in doses from one drachm to an ounce, as a cathartic, and in the dose of twenty to thirty grains in abdominal phys-

conia, and torpidity of the kidneys. See *Seignet's salt*.

Soda, tartarized. See *Soda tartarizata*.

SODÆ BORAS. Borate of soda.

See *Borax*.

SODÆ CARBONAS. Carbonate of soda. "Take of subcarbonate of soda; a pound; subcarbonate of ammonia, three ounces; distilled water, a pint." Having previously dissolved the soda in water, add the ammonia, then by means of a sand-bath apply a heat of 180° for three hours, or until the ammonia be driven off. Lastly, set the solution by to crystallize. The remaining solution may be evaporated and set by in the same manner that crystals may again form.

SODÆ MURIAS. See *Murias soda*.

SODÆ SUBCARBONAS Subcarbonate of soda, formerly called *natron preparatum* and *salsode*. "Take of impure soda, powdered, a pound; boiling distilled water, a gallon." Boil the soda in the water for half an hour, and strain the solution; let the solution evaporate to two pints, and be set by, that crystals may form. Throw away the remaining solution. This salt consists of soda saturated with carbonic acid, and is therefore called *soda subcarbonas*. It is given in doses of from ten grains to half a drachm as an attenuant and antacid; and, joined with bark and aromatics, it is highly praised by some in the cure of scrophula. It is likewise a powerful solvent of mucus, a deobstruent and diuretic, and an antidote against oxyd of arsenic and the corrosive sublimate. The other diseases in which it is administered are those arising from an abundance of mucus in the primæ viæ; calculous complaints, gout, some affections of the skin, rickets, tinea capitis, crusta lactea, and worms. Externally it is recommended by some in the form of lotion, to be applied to scrophulous ulcers.

SODÆ SUBCARBONAS EXSICCATA. Dried subcarbonate of soda. "Take of subcarbonate of soda, a pound." Apply a gentle heat to the soda in a clean iron vessel, until it becomes perfectly dry, and constantly stir it with an iron rod. Lastly, reduce it into powder. Its virtues are similar to those of the subcarbonate.

SODÆ SULPHAS. Sulphate of sodæ, commonly known by the name of *natron triolatum*, and formerly by *sul catharticus Glauberi*. "Take of the salt which remains after the distillation of muriatic acid, two pounds. Boiling distilled water, two pounds and a half." Dissolve the salt in water, then add gradually as much subcarbonate of soda as may be required to saturate the acid; boil the solution away until a pellicle forms upon the surface, and, after having strained it, set it by that crystals may form. Having poured away the water, dry these crystals upon bibulous

paper. It possesses cathartic and diuretic qualities, and is in high esteem as a mild cathartic. It is found in the mineral kingdom formed by nature, but that which is used medicinally is prepared by art. The dose is from one drachm to one ounce.

SOL. The sun. Gold was so called by the older chemists.

SOLANEN. (From *soler*, to comfort.) Anise-seed is named *solanen intestinum*, from the comfort it affords in disorders of the intestines.

SOLANOIDES. (From *solanum*, nightshade, and *oides*, likeness.) Bastard nightshade.

SOLANUM. (From *soler*, to comfort, because it gives ease by its stupifying qualities.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria* Order, *Monogynia*. Nightshade.

2. The pharmacopœial name of the *solanum nigrum* of Linnæus; its virtues are very nearly allied to those of the *belladonna*, which consult.

SOLANUM DULCINARA. The systematic name of the bitter sweet. See *Dulcamara*.

SOLANUM PERTIDUM. The thorn-apple plant is sometimes so called. See *Stramonium*.

SOLANUM LETHALE. See *Atropa belladonna*.

SOLANUM LIGUOSUM. The bitter sweet is sometimes so termed. See *Dulcamara*.

SOLANUM MELONGENA. The systematic name of the mad apple plant. See *Mad apple*.

SOLANUM NIGRUM. The systematic name of the garden nightshade. See *Solanum*.

SOLANUM SANCTUM. The systematic name of the Palestine nightshade. The fruit of this plant is globular, and in Egypt much eaten by the inhabitants.

SOLANUM TUBEROSUM. See *Potatoes*.

SOLANUM VESICARIUM. The winter cherry plant is so called by Caspar Bauhin. See *Attekengi*.

SOLDANELLA. (*A solidando*, from its uses in healing fresh wounds.) The sea convolvulus. See *Brassica marina*.

SOLIN. (Σολιν.) A tube or channel. A cradle for a broken limb.

SOLENARIUM. (Dim of *σολιν*, a tube.) A catheter.

SOLEUS. (From *solea*, a sole, from its shape being like the sole fish.) See *Gastrocnemius internus*.

SOLIDAGO. (From *solido*, to make firm; so called from its uses of consolidating wounds.) The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia superflua*. The herb comfrey.

SOLIDAGO VIRGAUREA. The systematic name of the golden rod. See *Virga aurea*.

SOLIDS, in anatomy, are the bones, ligaments, membranes, muscles, nerves, and vessels.

SOLIUM. (From *solus*, alone; so called because it infests the body singly.) The tape-worm.

SOLOXON'S SEAL. The *convallaria polygonatum*. Useful as an outward application for bruises; dried and powdered is said to be antidyenteric, and if beaten into a conserve with sugar whilst it is green, is recommended in leucorrhœa.

SOLSEQUIUM. (From *sol*, the sun, and *sequor*, to follow, so called because it turns its flowers towards the sun.) Marygold or turnsole.

SOLVENT See *Menstruum*.

SOLUTION. An intimate commixture of solid bodies with fluids, into one seemingly homogeneous liquor. The dissolving fluid is called a menstruum or solvent.

SOLUTIVA. (From *solvere*, to loosen.) Laxative medicines. Gentle purgatives.

Somnambulism. Sleep-walking. See *Oneirodynia*.

SOMNIFERA. (From *somnus*, sleep, and *fero*, to bring.) Opiates; medicines which induce sleep.

SONCHITES. (From *σῶνχις*, the sow-thistle; so named from its resemblance to the sonchus.) The herb hawkweed.

SONCHUS. (Παρα το σῶν χυμῶν, from its wholesome juice.) The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, The sow-thistle. All the species of sonchus abound with a milky juice, which is very bitter, and said to possess diuretic virtues. The *sonchus oleraceus* of Linnæus is sometimes employed with that intention. Boiled it may be eaten as a substitute for cabbage.

SONCHUS OLERACEUS. The systematic name of the sow-thistle. See *Sonchus*.

SOOT. *Fuligo*. A volatile matter, arising from coals, wood, and other fuel, along with the smoke. It is used as a material for making muriate of ammonia. At no very remote period our dispensaries contained directions for a tincture of soot; the most material ingredient of which, however, was assafoetida.

SOPHIA. (From *sopos*, wise; so named from its great virtues in stopping fluxes.) Flux-weed or flux-weed.

SOPHIA CHIRURGICUM. This plant, *Sisymbrium sophia* of Linnæus, is now almost banished from practice. It was formerly in high estimation in the cure of wounds. It has been given internally in hysterical affections and uterine hæmorrhages, and the seeds are said to be efficacious in destroying intestinal worms.

SOPHISTICATION. A term employed in pharmacy, to signify the counterfeiting or adulterating any medicine. This practice unhappily obtains with most dealers in drugs, &c.; and the cheat is carried on so artificially by many as to prevent a discovery even by persons of the most discerning faculties.

SOPHONISTÈRES. (From *σφονίζω*, to become wise; so called because they do not appear till after puberty.) The last of the grinding-teeth.

SOPHORA HEPTAPHYLLA. The systematic name of the shrub whose root and seeds are sometimes called *anticholerica*: they are both intensely bitter, and said to be useful in cholera, colic, and dysury.

SOPHENTIA. (From *sopio*, to make sleep.) Opiates.

SOPOR. Profound sleep.

SOPORIFEROUS. (*Soporifera medicamenta*. From *sopor*, sleep, and *fero*, to bear.) A term given to those medicines which induce sleep. See *Anodynes*.

SORA. (Arab.) The nettle-rash.

SORBASTRELLA. (From *sorbeo*, to suck up, because it stops hæmorrhages.) The herb burnet.

SORBUS. (From *sorbeo*, to suck up; because its fruit stops fluxes.) The name of a genus of plants in the Linnæan system. Class, *Icosandria*. Order, *Trigynia*. The service-tree.

SORBUS AUCUPARIA. The wild service-tree. The berries of this plant are adstringent, and, it is said, have been found serviceable in allaying the pain of calculous affections in the kidneys.

SORDES. When the matter discharged from ulcers is rather viscid or glutinous, it is thus named. This matter is frequently of a brownish red colour, somewhat resembling the grounds of coffee, or grumous blood mixed with water. *Sordes*, *Sanies*, and *Ichor*, are all of them much more fetid than purulent matter, and none of them are altogether free from acrimony; but that which is generally termed *Ichor* is by much the most acrid of them, being frequently so sharp and corrosive as to destroy large quantities of the neighbouring parts.

SORE-BAY. A disease which Dr. Mosely considers as a true cancer, commencing with an ulcer. It is endemic at the Bay of Honduras.

Sore-throat. See *Cynanche*.

Sorrel, common. See *Acetosa*.

Sorrel, French. See *Rumex scutatus*.

Sorrel, roundleaved. See *Rumex scutatus*.

Sorrel wood. See *Lujula*.

SOUND. An instrument which surgeons introduce through the urethra into the bladder, to discover whether there is a stone in this viscus or not.

Sour dock. See *Acetosa*.

Southernwood. See *Abrotanum*.

Sow-bread. See *Arthamita*.

Sow-breed. See *Cyclamen*.

SPA WATER. This mineral water appears to be a very strongly acidulous chalybeate, containing more iron and carbonic acid than any other mineral spring. What applies to the use of chalybeates will apply to this water.

Spain, pellicery of. See *Pyrethrum*.

Spanish fly. See *Cantharides*.

Spanish liquorice. See *Glycyrrhiza*.

SPANGIOSIS. (From *εργαζω*, to swell.)

A milk abscess

SPARTIUM. The name of a genus of plants in the Linnæan system. Class, *Dicladelia*. Order, *Decandria*.

SPARTIUM SCOPARIUM. The systematic name of the common broom. See *Genista*.

SPASM. (*Spasmus*, from *σπασσω*, to draw.)

A spasm or convulsion. An involuntary contraction of the muscular fibres, or that state of the contraction of muscles which is not spontaneously disposed to alternate with relaxation. When the contractions alternate with relaxation, which are frequently and preternaturally repeated, they are called convulsions. Spasms are distinguished by authors into clonic and tonic spasms. In *clonic* spasms, which are the true convulsions, the contractions and relaxations are alternate, as in epilepsy; but in *tonic* spasms the member remains rigid, as in locked jaw. See *Convulsion*, *Tonic spasm*, and *Tetanus*.

SPASMI. Spasmodic diseases. The third order of the class *neuroses* of Cullen; characterised by a morbid contraction or motion of muscular fibres.

Spasmodic colic. See *Colica*.

SPASMOLOGY. (*Spasmologia*, from *σπασμος*, a spasm, and *λογος*, a discourse.) A treatise on convulsions.

SPASMUS CYNICUS. The *spasmus cynicus*, or sardonic grin, is a convulsive affection of the muscles of the face and lips on both sides, which involuntarily forces the muscles of those parts into a species of grinning distortion. If one side only be affected, the disorder is nominated *tortura oria*. When the masseter, buccinator, temporal, nasal, and labial muscles, are involuntarily excited to action, or contorted by contraction or relaxation, they form a species of malignant sneer. It sometimes arises from eating hemlock, or other acrid poisons, or succeeds to an apoplectic stroke.

SPATHONELA. (From *σπάθη*, a sword, and *μύλη*, a probe.) An edged probe.

SPATULA. (Dim. of *spatha*, a broad instrument.) An instrument for spreading salve. Also a name of the herb spurge-wort, from its broad leaves.

Spear-mint. See *Mentha sativa*.

Spearwort, water. See *Flammula*.

SPECIFIC. A remedy that has an infallible efficacy in the cure of disorders. The existence of such remedies is doubted.

SPEIllum. (From *specio*, to examine.)

A probe.

SPECULUM ANI. An instrument for distending the anus whilst an operation is performed upon the parts within.

SPECULUM MATRICIS. An instru-

ment to assist in any manual operation belonging to the womb.

SPECULUM OCULI. (*Speculum*, from *specio*, to view.) An instrument used by oculists to keep the eyelids open and the eye fixed.

SPECULUM ORIS. An instrument to force open the mouth.

SPECULUM VENERIS. See *Miliefolium*.

Speech. See *Vox*.

Speedwell, female. See *Helatine*.

Speedwell, male. See *Veronica*.

Speedwell, mountain. See *Veronica*.

SPERMA-CETI. (From *σπέρμα*, seed, a *σπέρμα*, to sow, and *cete* or *cetæ*, the whale.) *Cetaceum*. An oily, concrete, crystalline, semi-transparent matter, obtained from the cavity of the cranium of several species of whales, but principally from the *Physeter macrocephalus*, or spermaceti whale. It was formerly very highly esteemed, and many virtues were attributed to it; but it is now chiefly employed in affections of the lungs, primæ viæ, kidneys, &c. as a softening remedy, mixed with mucilages. It is also employed by surgeons as an emollient in form of cerates, ointments, &c.

SPERMATICA. Belonging to the testicle and ovary, as the spermatic artery, chord and veins.

SPERMATOCELE. (From *σπέρμα*, seed, and *κύημα*, from *κύμα*, seed, and *κύμα*, a tumour.) *Epididymis distensa*. A swelling of the testicle or epididymis from an accumulation of semen. It is known by a swelling of those organs, pain extending to the loins without inflammation.

SPERMATO-ÆSTICA. (From *σπέρμα*, and *αἷμα*, to make.) Medicines which increase the generation of seed.

SPHACELISMUS. (From *σφραγιζω*, to gangrene.) A gangrene. Also a phrenitis.

SPHACELUS. (From *σφαινω*, to destroy.) A mortification of any part. See *Gangrene*.

SPHENOIDES OS. (From *σφην*, a wedge, and *οἶος*, a likeness; because it is fixed in the cranium like a wedge.) *Os cuneiforme*, or *multiforme*. *Os unguis*. *Papillare os*. *Basilare os*. *Os polymorphos*. Pterygoid bone. The os sphenoides, or cuneiforme, as it is called from its wedge-like situation amidst the other bones of the head, is of a more irregular figure than any other bone. It has been compared to a bat with its wings extended. This resemblance is but faint, but it would be difficult perhaps to find any thing it resembles more.

We distinguish in this bone its body or middle part, and its wings or sides, which are much more extensive than its body.

Each of its wings or lateral processes is divided into two parts. Of these the uppermost and most considerable portion,

helping to form the deepest part of the temporal fossa on each side, is called the *temporal process*. The other portion makes a part of the orbit, and is therefore named the *orbital process*. The back part of each wing, from its running out sharp to meet the os petrosum, has been called the *spinous process*; and the two processes, which stand out almost perpendicular to the basis of the skull, have been named *pterygoid* or *aliform* processes, though they may be said rather to resemble the legs than the wings of the bat. Each of these processes has two plates and a middle fossa facing backwards; of these plates the external one is the broadest, and the internal one the longest. The lower end of the internal plate forms a kind of hook, over which passes the round tendon of the *musculus circumflexus palati*. Besides these, we observe a sharp middle ridge, which stands out from the middle of the bone. The fore part of it, where it joins the nasal lamella of the ethmoidal bone, is thin and straight; the lower part of it is thicker, and is received into the vomer.

The cavities observable on the external surface of the bone, are where it helps to form the temporal, nasal, and orbital fossae. It has likewise two fossae in its pterygoid processes. Behind the edge, which separates these two fossae, we observe a small groove, made by a branch of the superior maxillary nerve in its passage to the temporal muscle. Besides these, it has other depressions, which serve chiefly for the origin of muscles.

Its foramina are four on each side. The three first serve for the passage of the optic, superior maxillary, and inferior maxillary nerves; the fourth transmits the largest artery of the dura mater. On each side we observe a considerable fissure, which, from its situation, may be called the superior orbital fissure. Through it pass the third and fourth pair of nerves, a branch of the fifth, and likewise the sixth pair. Lastly, at the basis of each pterygoid process, we observe a foramen which is named *pterygoidean*, and sometimes *Vidian*, from Vidian who first described it. Through it passes a branch of the external carotid, to be distributed to the nose.

The os sphenoides on its internal surface affords three fossae. Two of these are considerable ones; they are formed by the lateral processes, and make part of the lesser fossae of the basis of the skull. The third, which is smaller, is on the top of the body of the bone, and is called *sella turcica*, from its resemblance to a Turkish saddle. In this the pituitary gland is placed. At each of its four angles is a process. They are called the *clinoid* processes, and are distinguished by their situation into anterior and posterior processes.

The two latter are frequently united into one.

Within the substance of the os sphenoides, immediately under the sella turcica, we find two cavities, separated by a thin bony lamella. These are the sphenoidal sinuses. They are lined with the pituitary membrane, and, like the frontal sinuses, separate a mucus which passes into the nostrils. In some subjects there is only one cavity; in others, though more rarely, we find three.

In infants the os sphenoides is composed of three pieces, one of which forms the body of the bone and its pterygoid processes, and the other two its lateral processes. The clinoid processes may even then be perceived in a cartilaginous state, though some writers have asserted the contrary; but we observe no appearance of any sinus.

This bone is connected with all the bones of the cranium, and likewise with the ossa maxillaria, ossa malarum, ossa palati, and vomer. Its uses may be collected from the description we have given of it.

SPHENOIDAL SUTURE. *Sutura sphenoidalis*. The sphenoidal and ethmoidal sutures are those which surround the many irregular processes of these two bones, and join them to each other and to the rest.

Spheno-salpingo-staphilinus. See *Circumflexus*.

Spheno-staphilinus. See *Levator palati*.

SPHERITIS. (From *sphaigē*, a globe; so called from its round head.) *Spherocephalia elatior*. *Spherocephalus*. The globe-thistle.

SPHEROCEPHALUS. See *Spheritis*.

SPHEROMA. (From *sphaigē*, a globe.) A fleshy globular protuberance.

SPHENO-MAXILLARIS. An artery and a fissure of the orbit of the eye is so called.

SPHINCTER. (From *σφίγω*, to shut up.) The name of several muscles, whose office is to shut or close the aperture around which they are placed.

SPHINCTER ANI. *Sphincter externus* of Albinus and Douglas. *Sphincter cutaneus* of Winslow, and *coccigio-cutaneusphincter* of Dumas. A single muscle of the anus, which shuts the passage through the anus into the rectum, and pulls down the bulb of the urethra, by which it assists in ejecting the urine and semen. It arises from the skin and fat that surrounds the verge of the anus on both sides, near as far as the tuberosity of the ischium; the fibres are gradually collected into an oval form, and surround the extremity of the rectum. It is inserted by a narrow point into the perineum, acceleratores urinæ, and transversarii perinei; and behind into the extremity of the os coccygis, by an acute termination.

Sphincter ani cutaneus. See *Sphincter ani*.

Sphincter ani externus. See *Sphincter ani*.

SPHINCTER ANI INTERNUS. Albinus and Douglas call the circular fibres of the muscular coat of the rectum, which surrounds its extremity, by this name.

SPHINCTER CUTANEUS. See *Sphincter ani*.

SPHINCTER EXTERNUS. See *Sphincter ani*.

SPHINCTER GULÆ. The muscle which contracts the top of the throat.

SPHINCTER LABIORUM. See *Orbicularis oris*.

SPHINCTER ORIS. See *Orbicularis oris*.

SPHINCTER VAGINÆ. *Constrictor cunni* of Albinus. *Second muscle of the clitoris* of Douglas, and *anulo-synde-smo-clitoridien* of Dumas. This muscle arises from the sphincter ani and from the posterior side of the vagina near the perineum; from thence it runs up the side of the vagina, near its external orifice, opposite to the nymphæ, covers the corpus cavernosum, and is inserted into the crus and body or union of the crura clitoridis. Its use is to contract the mouth of the vagina.

SPRINGONTA. (From *εργω*, to bind.) Astringent medicines.

SPONDYLIIUM. (From *σπονδυλος*, vertebra; named from the shape of its root; or probably because it was used against the bite of a serpent called *σπονδυλις*.) This is supposed to be the brankursine. See *Banca urina*.

SPICA. 1. An ear of corn. 2. A bandage resembling an ear of corn.

SPICA BREVIS. Fox-tail plant.

SPICA CÆLTICA. See *Nardus celtica*.

SPICA FÆMINA. Common lavender.

SPICA INDICA. See *Nardus indica*.

SPICA INGUINALIS. A bandage for ruptures in the groin.

SPICA INGUINALIS DUPLEX. Double bandage for ruptures.

SPICA MAS. Broad-leaved lavender.

SPICA NARDI. See *Nardus indica*.

SPICA SIMPLEX. A common roller or bandage.

SPIGELIA. (From *spica*, an ear of corn; so called from its apicated top.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogymia*.

2. The name in some pharmacopæias for the *Spigelia anthelmia* of Linnæus directed as an anthelmintic; its virtues are very similar to those of the Indian pink. See *Spigelia marilandica*.

SPIGELIA ANTHELMIA. The systematic name of the spigelia of some pharmacopæias. See *Spigelia*.

SPIGELIA LONICERA. See *Spigelia marilandica*.

SPIGELIA MARILANDICA. *Spigelia lonicera*. Perennial worm-grass, or Indian pink. *Spigelia marilandica* of Linnæus;—caule tetragono, foliis oppositis. The whole of this plant, but most commonly the root, is employed as an anthelmintic by the Indians and inhabitants of America. Dr. Hope has written in favour of this plant, in continued and remitting low worm-fevers; besides its property of destroying the worms in primæ viæ, it acts as a purgative.

Spigelian lobe. See *Liver*.

Spiguel. See *Meum athamanticum*.

Spike. See *Nardus indica*.

Spikenard. See *Nardus indica*.

SPILANTHUS ACNELLA. The systematic name of the balm-leaved spilanthus, which possesses a bitter taste and a fragrant smell. The herb and seed are said to be diuretic and menagogue, and useful in dropsies, jaundice, fluor albus, and calculous complaints.

SPINA. (*Quasi speculina*, dim. of *spica*.) 1. A thorn. 2. The back-bone; so called from the thorn-like processes of the vertebrae. 3. The shin-bone.

SPINA ACIDA. See *Berberis*.

SPINA ACUTA. The hawthorn.

SPINA ÆGYPTIACA. The Egyptian thorn or sloe-tree. See *Mimosa*.

SPINA ALBA. The white-thorn tree.

SPINA ARABICA. The chardon or Arabian thistle.

SPINA BIFIDA. *Hydrops medullæ spinalis.* *Hydrocele spinalis.* *Hydro-rachitis spinosa.* A tumour upon the spine of newborn children immediately about the lower vertebrae of the loins, and upper parts of the sacrum; at first, it is of a dark blue colour; but in proportion as it increases in size, approaches nearer and nearer to the colour of the skin, becoming perfectly diaphanous.

From the surface of this tumour a pellucid watery fluid sometimes exudes, and this circumstance has been noticed by different authors. It is always attended with a weakness, or, more properly speaking, a paralysis of the lower extremities. The opening of it rashly has proved quickly fatal to the child. Talpius, therefore, strongly dissuades us from attempting this operation. Acrel mentions a case where a nurse rashly opened a tumour, which, as she described it, was a blood bag on the back of the child at the time of its birth, in bigness equal to a hen's egg, in two hours after which the child died. From the dissection it appeared that the bladder laid in the middle of the os sacrum, and consisted of a coat, and some strong membrane, which proceed from a long fissure of the bones. The extremity of the spinal marrow lay bare, and the spinal duct, in the os sacrum, was uncommonly wide, and distended by the pressure of the waters.

Upon tracing it to the head, the brain was found nearly in its natural state, but the ventricles contained so much water that the infundibulum was quite distended with it, and the passage between the third and fourth ventricle was greatly enlarged.

He likewise takes notice of another case, where a child lived about eight years labouring under this complaint, during which time it seemed to enjoy tolerable health, though pale. Nothing seemed amiss in him, but such a degree of debility as rendered him incapable to stand on his legs.

The tumour, as in the former case, was in the middle of the os sacrum, of the bigness of a man's fist, with little discolouring; and upon pressing it became less. When opened it was found full of water, and the coats were the same as in the former, but the separation of the bones was very considerable. The spinal marrow, under the tumour, was as small as a pack-thread, and rigid; but there were no morbid appearances in the brain.

SPINA BURGHI MONSPELIENSIS. Evergreen privet.

SPINA CERVINA. (So called from its thorns resembling those of the stag.) *Rhamnus catharticus*. *Rhamnus solutivus*. *Spina insectoria*. *Cervispina*. Purging buckthorn. The fruit or berries of this shrub, *Rhamnus catharticus* of Linnaeus: — *spinis terminalibus floribus quadrifidis diclis, foliis ovatis, caule erecto*, have been long received into the materia medica: they contain a pulpy deep green juice, of a faint unpleasant smell, a bitterish, acrid, nauseous taste, which operate briskly by stool, producing thirst, dryness of the mouth and fauces, and severe gripings, unless some diluting liquor be drank plentifully after it: at present it is rarely prescribed except as a drastic purge. The dose is said to be about 20 of the fresh berries in substance; twice or thrice that number in decoction; a drachm or a drachm and a half of the dried berries; an ounce of the expressed juice or half an ounce of the rob or extract, obtained by inspissating the juice.

SPINA HIRCI. The goats-thorn of France yielding gum tragacanth.

SPINA INFECTORIA. See *Spina cervina*.

SPINA PURGATRIX. The purging thorn.

SPINA SOLSTITIALLIS. The calcitrapa officinalis Barnaby's thistle.

SPINA VENTOSA. (The term of spina seems to have been applied by the Arabians to this disorder, because it occasions a prickling in the flesh like the puncture of thorns; and the epithet ventosa is added, because, upon touching the tumour, it seems to be filled with wind, though this is not the cause of the distention.) *Spina*

ventositas. *Teredo*. *Fungus articulari*. *Arthrocece*. *Sideratio ossis*. *Cancer ossis*. *Gangrena ossis*, and some French authors *exostosis*. When children are the subjects of this disease, M. Severinus calls it *Pedarthrocece*. A tumour arising from an internal caries of a bone. It most frequently occurs in the carpus and tarsus, and is known by a continual pain in the bone, and a red swelling of the skin, which has a spongy feel.

SPINACHIA. See *Spinacia*.

SPINACIA. (From *spinaria*, Spain, whence it originally came, or from its spinous seed.) *Spinachia*. Spinage. This plant *Spinacia oleracea* of Linnaeus is sometimes directed for medicinal purposes in the cure of phthisical complaints; made into a poultice, by boiling the leaves and adding some oil, it forms an excellent emollient. As an article of food it may be considered as similar to cabbage and other oleraceous plants. See *Brassica capitata*.

SPINACIA OLERACEA. The systematic name of spinage. See *Spinacia*.

SPINÆ CRATES. The spine of the back.

SPINÆ VENTOSITAS. A caries or decay of a bone.

Spinal marrow. See *Medulla spinalis*.

SPINALIS CERVICIS. This muscle, which is situated close to the vertebræ at the posterior part of the neck and upper part of the back, arises, by distinct tendons, from the transverse processes of the five or six uppermost vertebræ of the back, and, ascending obliquely under the complexus, is inserted, by small tendons, into the spinous processes of the sixth, fifth, fourth, third, and second vertebræ of the neck.

Its use is to extend the neck obliquely backwards.

SPINALIS COLLI. See *Semi-spinalis colli*.

SPINALIS DORSI. *Transversalis dorsi* of Winrow and *inter-epineux* of Dumas. This is the name given by Albinus to a tendinous and fleshy mass, which is situated along the spinous processes of the back and the inner side of the longissimus dorsi.

It arises tendinous and fleshy from the spinous processes of the uppermost vertebræ of the loins, and the lowermost ones of the back, and is inserted into the spinous processes of the nine uppermost vertebræ of the back.

Its use is to extend the vertebræ, and to assist in raising the spine.

SPINALES LUMBORUM. Muscles of the loins.

SPINÆ. (*Spina*, from *spina*, thorn; so called from the spine-like processes of the vertebræ.) *Spina dorsii*. *Columna spinalis*. *Columna vertebralis*. A bony column or pillar extending in the posterior

part of the trunk from the great occipital foramen to the sacrum. It is composed of twenty-four bones called vertebrae. See *Vertebra*.

SPIROSA. See *Spina bifida*.

SPIROSUM SYRIACUM. The Syrian broom.

SPIRÆA AFRICANA. African meadow sweet.

SPIRÆA. (From *spira*, a pillar; so named from its spiral stalk.) Meadow sweet. The name of a genus of plants in the Linnæan system. Class, *Pentagynia*. Order, *Icosandria*.

SPIRÆA FILIPENDULA. The systematic name of the officinal dropwort. See *Filipendula*.

SPIRÆA ULMARIA. The systematic name of the meadow sweet. See *Ulmaria*.

SPIRIT. *Spiritus*. This name was formerly given by chemists to all volatile substances collected by distillation. Three principal kinds were distinguished: inflammable or ardent spirits, acid spirits, and alkaline spirits. The word spirit is now almost exclusively confined to alcohol.

SPIRITUS ÆTHERIS NITRICI. *Spiritus ætheris nitrosi*. *Spiritus nitri dulcis*. A febrifuge, diaphoretic, and diuretic compound mostly administered in asthenia, nervous affections, disuria, and calculous affections.

SPIRITUS ÆTHERIS VITRIOLICI. *Spiritus vitrioli dulcis*. A diaphoretic, antispasmodic, and tonic preparation, mostly exhibited in nervous debility and weakness of the primæ viæ.

SPIRITUS ÆTHERIS VITRIOLICI AROMATICUS. An excellent stimulating and stomachic compound, which is administered in debility of the stomach and nervous affections.

SPIRITUS ÆTHERIS VITRIOLICI COMPOSITUS. A stimulating anodyne, supposed to be the celebrated *liquor nothæ anodynus* of Hoffman. It is exhibited in fevers, nervous affections, hysteria, &c.; and in most cases of fever where medicines are rejected by the stomach, this is of infinite service.

SPIRITUS AMMONIÆ. Spirit of ammonia. Formerly called *spiritus salis ammoniaci dulcis*. *Spiritus salis ammoniaci*. "Take of rectified spirit, two pints; solution of ammonia, a pint." Mix. A stimulating antispasmodic exhibited in cases of asphyxia, asthenia, and in nervous diseases, but mostly used as an external stimulant against rheumatism, sprains and bruises.

SPIRITUS AMMONIÆ AROMATICUS. Aromatic spirit of ammonia. Formerly known by the name of *spiritus ammoniac compositus*. *Spiritus volatilis aromaticus*: *Spiritus salis volatilæ oleosus*. "Take of

spirit of ammonia, two pints; oil of lemon, oil of cloves, of each two fluid-drachms." Mix. A stimulating antispasmodic and sudorific in very general use to smell at in faintings and lowness of spirits. It is exhibited internally in nervous affections, hysteria, and weakness of the stomach. The dose is from half a drachm to a drachm.

SPIRITUS AMMONIÆ FETIDUS. Fetid spirit of ammonia. Formerly called *spiritus volatilæ fetidus*. "Take of spirit of ammonia, two pints; assafoetida, two ounces." Macerate for twelve hours, then by a gentle fire distil a pint and a half into a cooled receiver. A stimulating antispasmodic, often exhibited to children against convulsions, and to gouty and asthmatic persons. The dose is from half a fluidrachm to a drachm.

SPIRITUS AMMONIÆ SUCCINATUS. Succinated spirit of ammonia. Formerly known by the names of *eau de luce*: *Spiritus salis ammoniacæ succinatus*: *Liquor cornu cervi succinatus*. "Take of mastic, three drachms; alcohol, nine fluidrachms; oil of lavender, fourteen minims; oil of amber, four minims; solution of ammonia, ten fluidounces." Macerate the mastic in the alcohol that it may dissolve, and pour off the clear tincture, to this add the remaining articles, and shake them together. This preparation is a compound succinate of ammonia. It is much esteemed as a stimulant and nervine medicine, and is employed internally and externally against spasms, hysteria, syncope, vertigo, and the stings of insects. The dose is from ten minims to half a fluidrachm.

SPIRITUS ANISI. Spirit of aniseed. Formerly called *spiritus anisi compositus*: *Aqua seminum anisi composita*. "Take of aniseed, bruised, half a pound; proof spirit, a gallon; water sufficient to prevent empyreuma." Macerate for 24 hours, and distil a gallon by a gentle fire. A stimulating carminative and stomachic calculated to relieve flatulency, borborygmi, colic, and spasmodic affections of the bowels. The dose is from half a fluidrachm to a drachm.

SPIRITUS ARMORACIÆ COMPOSITUS. Compound spirit of horse-radish, formerly called *spiritus raphani compositus*: *Aqua raphani composita*. "Take of horse-radish root, fresh and sliced, dried orange peel, of each a pound; nutmegs, bruised, half an ounce; proof spirit, a gallon; water sufficient to prevent empyreuma." Macerate for 24 hours, and distil a gallon by a gentle fire. A very warm stimulating compound given in gouty, rheumatic, and spasmodic affections of the stomach and in scorbutic disorders. The dose is from half a fluidrachm to half an ounce.

SPIRITUS CAMPHORÆ. Spirit of camphor. Formerly known by the names of *spiritus camphoratus*: *Spiritus vinosus camphoratus*: *Spiritus vini camphoratus*: "Take of camphor, four ounces; rectified spirit, two pints." Mix, that the camphor may be dissolved. A stimulating medicine, used as an external application against chilblains, rheumatism, palsy, numbness, and gangrene.

SPIRITUS CARUI. Spirit of carraway. Formerly called *aqua seminum carui*. "Take of carraway-seeds, bruised, a pound and half; proof spirit, a gallon; water sufficient to prevent empyreuma. Macerate for 24 hours, and distil a gallon by a gentle fire. The dose is from a fluidrachm to half an ounce.

SPIRITUS CINNAMOMI. Spirit of cinnamon. Formerly called *aqua cinnamomi spirituosus*. *Aqua cinnamomi fortis*. "Take of cinnamon bark, bruised, a pound; proof spirit, a gallon; water sufficient to prevent empyreuma." Macerate for 24 hours, and distil a gallon by a gentle fire. Spirit of cinnamon is mostly used in conjunction with other carminatives to give a pleasant flavour; it may be exhibited alone as a carminative and stimulant. The dose is from a fluidrachm to half an ounce.

SPIRITUS CORNU CERVI. See *Liquor carbonatis ammoniac*.

SPIRITUS JUNIPERI COMPOSITUS. Compound spirit of juniper. Formerly called *aqua juniperi composita*. "Take of juniper-berries, bruised, a pound; carraway-seeds, bruised, fennel-seeds, bruised, of each an ounce and half; water sufficient to prevent empyreuma." Macerate for 24 hours, and distil a gallon by a gentle fire.

SPIRITUS LAVENDULÆ. Spirit of lavender. Formerly called *spiritus lavendula simplex*. "Take of fresh lavender flowers, two pounds; rectified spirit, a gallon; water sufficient to prevent empyreuma." Macerate for 24 hours, and distil a gallon by a gentle fire. Though mostly used as a perfume, this spirit may be given internally as a stimulating nervine and antispasmodic. The dose is from a fluidrachm to half an ounce.

SPIRITUS LAVENDULÆ COMPOSITUS. Compound spirit of lavender. Formerly called *spiritus lavendula compositus matthiæ*. "Take of spirit of lavender, three pints; spirit of rosemary, a pint; cinnamon bark, bruised, nutmegs, bruised, of each half an ounce; red saunders wood, sliced, an ounce." Macerate for fourteen days, and strain. An elegant and useful antispasmodic, and stimulant in very general use against nervous diseases, lowness of spirits, and weakness of the stomach, taken on a lump of sugar.

SPIRITUS LUMBRICORUM. The spirit obtained by the distillation of the earth-worm is similar to hartshorn.

SPIRITUS MENTHÆ PIPERITÆ. Spirit of peppermint. Formerly called *spiritus menthe piperitidis*: *Aqua menthe piperitidis spirituosus*. "Take of peppermint, dried, a pound and half; proof spirit, a gallon; water sufficient to prevent empyreuma." Macerate for 24 hours, and distil a gallon by a gentle fire. This possesses all the properties of the peppermint with the stimulating virtues of the spirit. The dose from one fluidrachm to an ounce.

SPIRITUS MENTHÆ VIRIDIS. Spirit of spearmint. Formerly called *spiritus menthe sativæ*: *Aqua menthe vulgaris spirituosus*. "Take of spearmint, dried, a pound and half; proof spirit, a gallon; water sufficient to prevent empyreuma." Macerate for twenty-four hours, and distil a gallon. This is most commonly added to carminative or antispasmodic draughts, and seldom exhibited alone. The dose from one fluidrachm to an ounce.

SPIRITUS MILLEPEDARUM. A fluid volatile alkali, whose virtues are similar to hartshorn.

SPIRITUS MYRDERERI. See *Liquor acetatis ammoniac*.

SPIRITUS MYRISTICÆ. Spirit of nutmeg. Formerly called *aqua nucis moschate*. "Take of nutmegs, bruised, two ounces; proof spirit, a gallon; water sufficient to prevent empyreuma." Macerate for 24 hours, and distil a gallon by a gentle fire. A stimulating and agreeable spirit possessing the virtues of the nutmeg. The dose from one fluidrachm to an ounce.

SPIRITUS NITRI DULCIS. See *Spiritus ætheris nitrici*.

SPIRITUS NITRI DUPLEX. The nitrous acid. See *Acidum nitrosum* and *Nitric acid*.

SPIRITUS NITRI FUMANS. See *Acidum nitrosum* and *Nitric acid*.

SPIRITUS NITRI GLAUBERI. See *Acidum nitrosum* and *Nitric acid*.

SPIRITUS NITRI SIMPLEX. The dilute nitrous acid. See *Acidum nitrosum dilutum*.

SPIRITUS NITRI VULGARIS. This is now called *acidum nitrosum dilutum*.

SPIRITUS PIMENTÆ. Spirit of pimento. Formerly called *spiritus pimento*. "Take of allspice, bruised, two ounces; proof spirit, a gallon; water sufficient to prevent empyreuma." Macerate for 24 hours, and distil a gallon by a gentle fire. A stimulating aromatic tincture mostly employed with adstringent and carminative medicines. The dose is from half a fluidrachm to half an ounce.

SPIRITUS PULEGII. Spirit of penny-royal. Formerly called *aqua pulegii spirituosus*. "Take of pennyroyal, dried, a pound and half; proof spirit, a gallon; water sufficient to prevent empyreuma." Macerate for 24 hours, and distil a gallon by a gentle fire. This is in very general use

as an emmenagogue amongst the lower orders. It possesses nerve and carminative virtues. The dose is from half a fluid-drachm to half an ounce.

SPIRITUS RECTOR. Boerhaave and other chemists give this name to a very attenuated principle, in which the smell of odorant bodies peculiarly resides. Now called *aroma*.

SPIRITUS ROSMARINI. Spirit of rosemary. "Take of rosemary tops, fresh, a pound and half; proof spirit, a gallon; water sufficient to prevent empyreuma." Macerate for 24 hours, and distil a gallon by a gentle fire. A very fragrant spirit, mostly employed for external purposes in conjunction with other solvents.

SPIRITUS SALIS AMMONIACI AQUOSUS. See *Liquor carbonatus ammoniac*.

SPIRITUS SALIS AMMONIACI DULCIS. See *Spiritus ammoniac*.

SPIRITUS SALIS AMMONIACI SIMPLEX. See *Liquor carbonatus ammoniac*.

SPIRITUS SALIS GLAUBERI. See *Acidum muriaticum*.

SPIRITUS SALIS MARINI. See *Acidum muriaticum*.

SPIRITUS VINI RECTIFICATUS. See *Alkohol*. Rectified spirit of wine is in general use to dissolve resinous and other medicines. It is seldom exhibited internally, though it exists in the diluted state in all vinous and spirituous liquors.

SPIRITUS VINI TENUIOR. Proof spirit, which is half the strength of rectified, is much employed for preparing tinctures of genuine resinous juices, barks, roots, &c.

SPIRITUS VITRIOLI. See *Sulphuric acid*.

SPIRITUS VITRIOLI DULCIS. See *Spiritus atheris vitriolici*.

SPIRITUS VOLATILIS FETIDUS. See *Spiritus ammoniac fetidus*.

SPISSAMENTUM. (From *spissus*, to thicken.) A substance put into oils and ointments to make them thick.

Spitting of blood. See *Hæmatemesis* and *hæmoptysis*.

SPLANCHICA. (From *σπλῆγχον*, an intestine.) Remedies for diseased bowels.

SPLANCHNOLOGIA. (From *σπλῆγχον*, an entrail, and *λογος*, a discourse.) Splanchnology, or the doctrine of the viscera.

SPLANCHNIC NERVE. The great intercostal nerve. See *Intercostal nerve*.

SPLEEN *σπλῆν*. *Lien*. The spleen or milt is a spongy viscus of a livid colour, and so variable in form, situation, and magnitude, that it is hard to determine either. Nevertheless, in a healthy man it is always placed on the left side, in the left hypochondrium, between the eleventh and twelfth false ribs. Its circumference is oblong and round, resembling an oval figure. It is larger, to speak generally,

when the stomach is empty, and smaller when it is compressed, or evacuated by a full stomach.

It should particularly be remembered of this viscus, that it is convex towards the ribs, and concave internally: also, that it has an excavation, into which vessels are inserted.

It is connected with the following parts:

1. With the stomach, by a ligament and short vessels.
2. With the omentum, and the left kidney.
3. With the diaphragm, by a portion of the peritonæum.
4. With the beginning of the pancreas, by vessels.
5. With the colon, by a ligament.

In man the spleen is covered with one simple, firm membrane, arising from the peritonæum, which adheres to the spleen, very firmly, by the intervention of cellular structure.

The vessels of the spleen are, the splenic artery coming from the celiac artery, which, considering the size of the spleen, is much larger than is requisite for the mere nutrition of it. This goes by serpentine movements, out of its course, over the pancreas, and behind the stomach, and after having given off branches to the adjacent parts, it is inserted into the concave surface of the spleen. It is afterwards divided into smaller branches, which are again divided into other yet smaller, delivering their blood immediately to the veins, but emitting it no where else. The veins, at length, come together into one, called the splenic vein, and having received the large coronary vein of the stomach, besides others, it constitutes the left principal branch of the vena portæ.

The nerves of the spleen are small; they surround the arteries with their branches; they come from the particular plexus, which is formed of the posterior branches of the eighth pair, and the great intercostal nerve.

Lymphatic vessels are almost only seen creeping along the surface of the human spleen.

The use of spleen has not hitherto been determined; yet if its situation and fabric be regarded, one would imagine its use to consist chiefly in affording some assistance to the stomach during the progress of digestion.

Spleenwort. See *Ceterach*.

SPLENALGIA. (From *σπλῆν*, the spleen, and *αλγος*, pain.) A pain in the spleen or its region.

SPLENETICA. (From *σπλῆν*, the spleen.) Medicines which relieve diseases of the spleen.

SPLENITIS. (From *σπλῆν*, the spleen.) Inflammation of the spleen. A genus of disease in the class *pyrexia* and order *phlegmasia* of Cullen; characterized by pyrexia, tension, heat, tumour, and pain in the left hypochondrium, increased

by pressure. This disease, according to Juncker, comes on with a remarkable shivering, succeeded by a most intense heat, and very great thirst; a pain and tumour are perceived in the left hypochondrium, and the paroxysms for the most part assume a quartan form; when the patients expose themselves for a little to the free air, their extremities immediately grow very cold. If an hæmorrhagy happen, the blood flows out of the left nostril. The other symptoms are the same with those of the hepatitis. Like the liver, the spleen often is also subject to a chronic inflammation which often happens after agues, and is called the ague cake, though that name is also frequently given to a scirrhus tumour of the liver succeeding intermittents. The causes of this disease are in general the same with those of other inflammatory disorders; but those which determine the inflammation to that particular part more than another, are very much unknown. It attacks persons of a very plethoric and sanguine habit of body rather than others.

SPLENIUS. (From *σπλην*, the spleen; so named from its resemblance in shape to the spleen, or according to some it derives its name from *splenium*, a *ferula*, or splint, which surgeons apply to the sides of a fractured bone.) *Splenius capitis* and *splenius colli* of Albinus, and *cervico-dorsomastoidien et dorso-trachelien* of Dumas. The splenius is a flat, broad, and oblong muscle, in part covered by the upper part of the trapezius, and obliquely situated between the back of the ear, and the lower and posterior part of the neck.

It arises tendinous from the four or five superior spinous processes of the dorsal vertebra; tendinous and fleshy from the last of the neck, and tendinous from the ligamentum colli, or rather the tendons of the two splenii unite here inseparably; but about the second or third vertebra of the neck they recede from each other, so that part of the complexus may be seen.

It is inserted, by two distinct tendons, into the transverse processes of the two first vertebra of the neck, sending off some few fibres to the complexus and levator scapula; tendinous and fleshy into the upper and posterior part of the mastoid process, and into a ridge on the occipital bone, where it joins with the root of that process.

This muscle may easily be separated into two parts. Eustachius and Fallopius were aware of this; Winslow has distinguished them into the *superior* and *inferior* portions; and Albinus has described them as two distinct muscles, calling that part which is inserted into the mastoid process and os occipitis, *splenius capitis*, and that which is inserted into the vertebra of the neck, *splenius colli*. We have here follow-

ed Douglas, and the generality of writers, in describing these two portions as one muscle, especially as they are intimately united near their origin.

When this muscle acts singly, it draws the head and upper vertebra of the neck obliquely backwards; when both act, they pull the head directly backwards.

SPLENIUM. (From *σπλην*, the spleen; so called from its efficacy in disorders of the spleen.) 1. Spleen wort. 2. A compress shaped like the spleen.

SPLENIUS CAPITIS. See *Splenius*.

SPLENIUS COLLI. See *Splenius*.

SPLENOCELE. (From *σπλην*, the spleen and *αλλη*, a tumour.) A rupture of the spleen.

SPLINT. A long piece of wood, tin, or strong pasteboard employed for preventing the ends of broken bones from moving, so as to interrupt the process by which fractures unite.

SPODIUM. *Σαπιδιον*. The *spodium* of Dioscorides and of Galen are now not known in the shops. It is said to have been produced by burning cadmia alone in the furnace; for having thrown it in small pieces into the fire, near the nozzle of the bellows, they blow the most fine and subtle parts against the roof of the furnace; and what was reflected from thence was called *spodium*. It differed from the pompholyx in not being so pure, and in being more heavy. Pliny distinguishes several kinds of it, as that of copper, silver, gold, and lead.

SPODIUM ARABUM. Burnt ivory or ivory black.

SPODIUM GRÆCORUM. The white dung of dogs.

SPOLIARIUM. A private room at the baths.

SPONDILIUM. (From *σπονδυλος*, a vertebra; so named from the shape of its root, or probably because it was used against the bite of a serpent called *σπονδυλις*.) The herb all-heal. Cow-parsnep.

SPONDYLUS. *Σαπιδυλα*. Some have thought fit to call the spine, or backbone thus, from the shape and fitness of the vertebra, to move every way upon one another.

Sponge. See *Spongia*.

Sponge-tent. See *Spongia preparata*.

SPONGIA. *Σπογγίος, σπογγία*. Sponge. A sea production, the *Spongia officinalis* of Linnæus; the habitations of insects. Burnt sponge is said to cure effectually the bronchocele, and to be of infinite utility in scrophulous complaints. Sponge tents are employed by surgeons to dilate fistulous ulcers, &c.

SPONGIA OFFICINALIS. The systematic name of the sponge. See *Spongia*.

SPONGIA PRÆPARATA. Prepared sponge. *Sponge tent.* This is formed by dipping pieces of sponge in hot melted emplastrum ceræ compositum, and pressing

them between two iron plates. As soon as cold, the substance thus formed may be cut into pieces of any shape. It was formerly used for dilating small openings, for which it was well adapted, as when the wax melted, the elasticity of the sponge made it expand and distend the opening, in which it had been put. Mr. Cooper informs us that the best modern surgeons seldom employ it.

SPONGIA USTA. Burnt sponge. "Cut the sponge into pieces, and beat it that any extraneous matters may be separated; then burn it in a close vessel until it becomes black and friable; lastly rub it to a very fine powder." This preparation is exhibited with bark in the cure of scrophulous complaints, and forms the basis of a lozenge which has been known to cure the bronchocele in many instances. The dose is from a scruple to a drachm.

SPONGIOSA OSSA. *Ossa turbinata inferiora.* These bones are situated in the under part of the side of the nose, they are of a triangular form and spongy appearance, resembling the os spongiosum superius; externally they are convex; internally they are concave; the convexity is placed towards the septum nasi, and concavity outwards. The under edge of each bone is placed horizontally near the outer part of the nose, and ending in a sharp point behind. At the upper part of the bone are two processes, the anterior of which ascends and forms part of the lacrymal groove, and the posterior descends and forms a hook to make part of the maxillary sinus.

The connexion of this bone is to the os maxillare, os palati and os unguis by a distinct suture in the young subject; but in the adult, by a concretion of substance.

The ossa spongiosa afford a large surface for extending the organ of smell by allowing the membrane of the nose to be expanded, upon which the olfactory nerves are dispersed.

In the fetus, these bones are almost complete.

SPONGIOSUM OS. 1. The ethmoid bone.

2. See *Spongiosa ossa*

SPONGIOSUS. (*σπογγιδις*, from *σπογγη*, a sponge, and *μορφη*, a shape.) Is the same as *Os cribriforme*, because it is hollow and porous like a sponge or sieve.

SPORADIC. (*Sporadicus*, from *σπορειν*, to sow.) An epithet for such infectious and other diseases as seize a few persons at any time or season.

Spotted lungwort. See *Pulmonaria*.

SPRUCE. 1. A particular species of fir. 2. A fermented liquor called spruce-beer prepared from the spruce fir. From the quantity of carbonic acid it contains, it is found a useful antiscorbutic.

Spurge flax. See *Thymalea*.

Spurge laurel. See *Laureola*.

Spurge olive. See *Mentha*.

Spruin. See *Sublunus*.

SPUTANES. See *Sputum*.

SPUTUM. (From *spuo*, to spit.) *Sputamen.* Saliva. Any kind of expectoration.

SQUAMARIA. (From *squama*, a scale; so called from its scaly roots.) The great tooth wort.

SQUAMOSE SUTURE (*Sutura squamosa*; from *squama*, a scale; because the bones lie over each other like scales.) The suture which unites the squamose portion of the temporal bone with the parietal.

Squill. See *Scilla*.

SQUILLA. See *Scilla*.

SQUINANTHEUS. (From *squinanthia*, the quince; so named from its uses in the quince) *Squinanthum.* The sweet rush was once so called. See *Juncus odoratus*.

STACHYS. (*σταχυς*, a spike; so named from its spicated stalk and seed.) The wild sage. The base borehound or marrubium hispanicum.

STACHYS FETIDA. Yellow archangel. Hedge-nettle.

STACHYS PALUSTRIS. Clown's wound-wort or all-heal.

STACTE. (*στακτη*, from *σαζω*, to distil.) Signifies that kind of myrrh which distils or falls in drops from the trees. It is also used by some writers for a more liquid kind of amber than what is commonly met with in the shops; whence in Scribonius Largus, *Ægineta*, and some others, we meet with a collyrium, and several other forms, wherein this was the chief ingredient, distinguished by the name of *Stactica*.

STACTICON. Instillation. An eye-water.

STAGMA. (*σταγμα*, to distil.) Any distilled liquor. The vitriolic acid.

STALAGMUS. (From *σταλαγμα*, to distil.) Distillation.

STALTICA. (From *σταλω*, to contract.) Healing applications.

STANNI PULVIS. Tin finely filed is exhibited internally as a vermifuge.

STANNUM. See *Tin*.

STAPEDIS MUSCULUS. See *Stapedius*.

STAPEDIUS. (*Stapedius*, sc. *musculus*; from *stapes*, one of the bones of the ear.) *Musculus stapes* of Cowper, and *pyramidal-stapedien* of Dumas. A muscle of the internal ear, which draws the stapes obliquely upwards towards the cavern, by which the posterior parts of its base is moved inwards, and the anterior part outwards.

STAPES. (*In quo pes stat*.) A bone of the internal ear, so called from its resemblance to a stirrup.

STAPHILINUS. See *Azygos uvule*.

STAPHILINUS EXTERNUS. See *Circumflexus palati*.

STAPHIS. *Staquis*, is strictly a grape, or a bunch of grapes; whence from their like-

ness thereunto it is applied to many other things, especially the glandulous parts of the body, whether natural or dustempered.

STAPHISAGRIA. (*Σταφίς αψία*, wild vine; from the resemblance of its leaves to those of the vine.) *Staphys. Pedicularia.* Staves acre. *Delphinium staphisagria* of Linnæus:—*nectaris tetraphyllis petulo brevioribus, foliis palmatis, lobis obtusis.* The seeds, which are the only parts directed for medicinal use, are usually imported here from Italy; they are large, rough, of an irregular triangular figure, and of a blackish colour on the outside, but yellowish within; their smell is disagreeable, and somewhat fetid; to the taste they are very bitter, acrid, and nauseous. It was formerly employed as a masticatory, but is now confined to external use in some kinds of cutaneous eruptions, but more especially for destroying lice and other insects; hence by the vulgar it is called louse-wort.

STAPHYLE. (*Σταφύλη*. A grape or raisin; so called from its resemblance.) The uvula.

STAPHYLINUS. (*Staphilinus*, sc. *musculus*, from *σταφύλη*, the uvula.) See *Amygos uvule*.

STAPHYLINUS EXTERNUS. See *Circumflexus palati*.

STAPHYLINUS GRACORUM. *Staphylinus sylvestris.* The wild carrot.

STAPHYLOMA. (From *σταφύλη*, a grape; so named from its being thought to resemble a grape.) *Staphyloma.* A disease of the eye-ball in which the cornea loses its natural transparency, rises above the level of the eye, and successively even projects beyond the eye-lids, in the form of an elongated, whitish, or pearl-coloured tumour which is sometimes smooth, sometimes uneven, and is attended with a total loss of sight. The proximate cause is an effusion of thick humour between the lamellæ of the cornea, so that the internal and external superficies of the cornea, very much protuberates. The remote causes are, an habitual ophthalmia, great contusion, and frequently a deposition of the variolous humour in the small-pox. The species are:

1st. *Staphyloma totale*, which occupies the whole transparent cornea; this is the most frequent species. The symptoms are, the opaque cornea protuberates, and if in the form of a cone increasing in magnitude, it pushes out and inverts the lower eye-lid; and sometimes the morbid cornea is so long elongated, as to lay on the cheek, causing friction and excoriation. The bulb of the eye being exposed to the air, sordes generate, the inferior palpebra is irritated by the cilia, and very painful red and small papillæ are observable.

2nd. *Staphyloma racemosum*, is a staphy-

loma formed by carnosous tubercles, about the size of a small pin's head.

3d. *Staphyloma parziale*, which occupies some part of the cornea: it exhibits an opaque tumour prominent from the cornea, similar to a small blueish grape.

4th. *Staphyloma sclerotica*, is a bluish tumour attached to some part of the sclerotica, but arising from the tunica albuginea.

5th. *Staphyloma pellucidum*, in which the cornea is not thickened or increased, but very much extended and pellucid.

6th. *Staphyloma complicatum*, which is complicated with an ulcer, eczema, caruncles, or any other disorder of the eye.

7th. *Staphyloma iridis.* For this species see *Ptoxis iridis*.

STAR THISTLE. The roots of this plant, *Cardina acanthis* of Linnæus, are said to be diuretic, and by some recommended in gravel and jaundice.

STARCH. *Amylum.* The fecula of wheaten flour. See *Amylum*.

Starch is one of the constituent parts in all mealy farinaceous seeds, fruits, roots, and other parts of plants. Our common starch is made from wheat. It is not necessary that the grain be first bruised in mills. The entire corn, well cleansed, is soaked in cold water until the husk separates; and the grains, having become quite soft, give out by pressure a milky fluid. The grains are then taken out of the water by means of a sieve, put into a coarse linen sack, and transferred into the treading-tub; where they are trodden, after cold water has been poured upon them.

By this operation the starchy part is washed out, and mingling with the water makes it milky. The water is now drawn off, running through a sieve into the settling tub. Fresh water is again effused upon the grains, and the same operation is continued till the water in the treading-tub is no longer rendered milky. The starch here precipitates by repose from the water that held it suspended; during which, especially in a warm season, the mucilaginous saccharine matter of the flour, that was dissolved by the water, goes into the acetous fermentation. From this cause the starch grows still purer and whiter. The water is next let off from the starch, which is several times more washed with clear fresh water; the remaining part of which is suffered to drip through linen cloths supported by hurdles, upon which the wet starch is placed. When the starch has fully subsided, it is wrapt in, wrung between these cloths, or pressed, to extort still more of the remaining liquid.

It is afterwards cut into pieces, which are laid in airy places on slightly burnt bricks to be completely dried, partly by

the free currency of air, and partly by the bricks imbibing their moisture. Lastly, the outer crust is scraped off, and they are broken into smaller pieces.

In the sacks wherein the corn was trodden there remain its husks and glutinous parts; and this residuum is employed as food for cattle.

STATICE. (From *σταῖς*, to stop, so named from its supposed property of restraining hæmorrhages.) The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Pentagynia*. The herb sea-thirst.

STATICE LIMONIUM. The systematic name of the thrift or sea frisk. See *Behen rubrum*.

STATIONARIA FEBRIS. A stationary fever. So Sydenham called those fevers which happen when there are certain general constitutions of the years, which owe their origin neither to heat, cold, dryness, nor moisture, but rather depend on a certain secret and inexplicable alteration in the bowels of the earth, whence the air becomes impregnated with such kinds of effluvia as subject the body to particular distempers, so long as that kind of constitution prevails, which, after a certain course of years, declines and gives way to another.

Stavesacre. See *Staphisagria*.

STEATOCELE. (From *σῆς*, suet, and *κύημα*, a tumour.) A collection of a suetty substance in the scrotum.

STEATOMA. (From *σῆς*, suet.) An encysted tumour, whose contents are of a suetty consistence.

STEEL. *Chalybs*. The best, hardest, finest, and closest grained iron, combined with carbon by a particular process.

STELOCHITES. See *Osteocolla*.

STELLA. (From *στᾶρα*, to arise.) A star. A bandage with many crossings like a star.

STELLARIA. (From *stella*, a star; so named from the star-like disposition of its leaves.) Stitch-wort. Ladies mantle.

STENA. (From *σῆμα*, to stand.) The penis.

Stemless milkvetch. See *Astragalus ex-capus*.

STENOTHORACES. (From *στενός*, narrow, and *θώραξ*, the chest.) Those who have narrow chests are so called.

STERILITY. Barrenness, in opposition to fertility. In women this sometimes happens from a miscarriage, or violent labour injuring some of the genital parts; but one of the most frequent causes is the suppression of the menstrual flux. There are other causes, however, arising from various diseases incident to those parts; by which the uterus may be unfit to receive or retain the male seed;—from the tubes fallopianæ being too short, or having lost their erective power; in either of

which cases no conception can take place;—from universal debility and relaxation; or a local debility of the genital system; by which means, the parts having lost their tone, or contractile power, the semen is thrown off immediately *post coitum*;—from imperforation of the vagina, of the uterus, or tube; or from diseased ova, &c.

STERNO. Names compounded of this word belong to muscles which are attached to the sternum; as,

STERNO-CLEIDO HYOIDEUS. See *Sternohyoideus*.

STERNO-CLEIDO MASTOIDEUS. *Sternomastoideus* and *cleidomastoideus* of Albinus. *Mastoideus* of Douglas and Cowper, and *terno-clavio-mastoides* of Dumas. A muscle, on the anterior and lateral part of the neck, which turns the head to one side and bends it forward. It arises by two distinct origins; the anterior tendinous and fleshy, from the top of the sternum near the junction with the clavicle; the posterior fleshy, from the upper and anterior part of the clavicle; both unite a little above the anterior articulation of the clavicle, to form one muscle, which runs obliquely upwards and outwards to be inserted, by a thick strong tendon, into the mastoid process of the temporal bone, which it surrounds; and gradually becoming thinner, is inserted as far back as the lambdoidal suture.

STERNO COSTALES. Vesalius considered these as forming a single muscle on each side, of a triangular shape; hence we find the name of *triangularis* adopted by Douglas and Albinus; but Verheyen, who first taught that they ought to be described as four or five distinct muscles, gave them the name of *sterno costales*; and in this he is very properly followed by Winslow, Haller, and Lietaud.

These muscles are situated at each side of the under surface of the sternum, upon the cartilages of the third, fourth, fifth, and sixth ribs. Their number varies in different subjects; very often there are only three, sometimes five, and even six, but most usually we find only four.

The lowermost of the sterno costales, or what would be called the inferior portion of the triangularis, arises tendinous and fleshy from the edge and inner surface of the lower part of the cartilago ensiformis, where its fibres intermix with those of the diaphragm and transversalis abdominis. Its fibres run nearly in a transverse direction, and are inserted, by a broad thin tendon, into the inner surface of the cartilage of the sixth rib, and lower edge of that of the fifth.

The second and largest of the sterno costales, arises tendinous from the cartilago ensiformis and lower part of the sternum, laterally, and, running a little obliquely outwards, is inserted into the lower

edge of the cartilage of the fifth, and sometimes of the fourth rib.

The third arises tendinous from the sides of the middle part of the sternum, near the cartilages of the fourth and fifth ribs, and, ascending obliquely outwards, is inserted into the cartilage of the third rib.

The fourth and uppermost, which is the most frequently wanting, arises tendinous from the beginning of the cartilage of the third rib and the adjacent part of the sternum, and running almost perpendicularly upwards, is inserted by a thin tendon (which covers a part of the second internal intercostal), into the cartilage and beginning of the bony part of the second rib.

All these muscles are more or less intermixed with one another at their origin, and this probably occasioned them to be considered as one muscle. Fallopius informs us, that the plate Vesalius has given of them was taken from a dog, in which animal they are much larger than in man. Douglas has endeavoured to account for this difference, but his explanation is far from being satisfactory.

STERNO HYOIDEUS. As this muscle arises from the clavicle, as well as from the sternum, Winslow calls it *sternocleidohyoideus*. It is a long, flat, and thin muscle, situated obliquely between the sternum and os hyoides, behind the lower part of the mastoideus, and covering the *sterno-thyroideus* and the *hyo-thyroideus*. It arises, by very short tendinous fibres, from the cartilaginous part of the first rib, from the upper and inner part of the sternum, from the capsular ligament that connects that bone with the clavicle, and commonly from a small part of the clavicle itself; from thence, ascending along the anterior and lateral part of the neck, we see it united to its fellow, opposite to the inferior part of the larynx, by means of a thin membrane, which forms a kind of *tinea alba*. After this the two muscles separate again, and each passing over the side of the thyroid cartilage, is inserted into the basis of the os hyoides, immediately behind the insertion of the last-described muscle.

Its use is to draw the os hyoides downwards.

STERNO MASTOIDEUS. See *Sternocleido-mastoideus*.

STERNO THYROIDEUS. *Sterno-thyroideus* of Dumas. This is flat and thin, like the preceding muscle, but longer and broader. It is situated at the fore part of the neck, between the sternum and thyroid cartilage, and behind the *sterno hyoideus*. It arises broad and fleshy from the upper and inner part of the sternum, between the cartilages of the first and second ribs, from each of which it receives some few fibres, as well as from the clavicle where it joins with the sternum. From thence, growing somewhat narrower, it ascends, and, passing

over the thyroid gland and the cricoid cartilage, is inserted tendinous into the lower and posterior edge of the rough line of the thyroid cartilage, immediately under the insertion of the last-described muscle. Now and then a few of its fibres pass on to the os hyoides. Its use is to draw the thyroid cartilage, and consequently the larynx, downwards.

STERNUM. *Pectoris os*. The breast-bone. The sternum os pectoris, or breast-bone, is the oblong, flat bone, placed at the fore part of the thorax. The ossification of this bone in the fœtus beginning from many different points at the same time, we find it, in young subjects, composed of several bones united by cartilages; but as we advance in life, most of these cartilages ossify, and the sternum, in the adult state, is found to consist of three, and sometimes only of two pieces, the two lower portions being united into one; and very often, in old subjects, the whole is formed into one bone. But, even in the latter case, we may still observe the marks of its former divisions; so that, in describing the bone, we may very properly divide it into its upper, middle, and inferior portions.

The upper portion forms an irregular square, which, without much reason, has, by many writers, been compared to the figure of a heart as it is painted on cards. It is of considerable thickness, especially at its upper part. Its anterior surface is irregular, and slightly convex; posteriorly, it is somewhat concave. Its upper middle part is hollowed, to make way for the trachea arteria. On each side, superiorly, we observe an oblong articulating surface, covered with cartilage in the recent subject, for receiving the ends of the clavicles. Immediately below this, on each side, the bone becomes thinner, and we observe a rough surface for receiving the cartilage of the first rib, and, almost close to the inferior edge of this, we find the half of such another surface, which, combined with a similar surface in the middle portion of the sternum, serves for the articulation of the cartilage of the second rib.

The middle portion is much longer, narrower, and thinner than the former; but is somewhat broader and thinner below than above, where it is connected with the upper portion. The whole of its anterior surface is slightly convex, and within it is slightly concave. Its edges, on each side, affords four articulating surfaces, for the third, fourth, fifth, and sixth ribs; and parts of articulating surfaces at its upper and lower parts, for the second and seventh ribs. About the middle of this portion of the sternum we sometimes find a considerable hole, large enough in some subjects to admit the end of the little finger. Sylvius seems to have been the first

who described it. Riolaus and some others after him have, without reason, supposed it to be more frequent in women than in men. In the recent subject it is closed by a cartilaginous substance; and, as it does not seem destined for the transmission of vessels, as some writers have asserted, we may, perhaps very properly, with M. Hunauld, consider it as an accidental circumstance, occasioned by an interruption of the ossification, before the whole of this part of the bone is completely ossified.

The third and inferior portion of the sternum is separated from the former by a line, which is seldom altogether obliterated, even in the oldest subjects. It is smaller than the other parts of the bone, and descends between the ribs, so as to have been considered as an appendix to the rest of the sternum. From its shape, and its being constantly in a state of cartilage in young subjects, it has been commonly named *cartilago xiphoides, ensiformis*, or sword-like cartilage; though many of the ancients gave the name of xiphoides to the whole sternum; comparing the two first bones to the handle, and this appendix to the blade of the sword. The shape of this appendix varies in different subjects; in some it is longer and more pointed; in others shorter and more obtuse. Veslingius has seen it reaching as low as the navel, and incommoding the motion of the trunk forwards. In general it terminates obtusely, or in a single point; sometimes, however, it is bifurcated, and Eustachius and Haller have seen it trifid. Very often we find it perforated, for the transmission of branches of the mammary artery. In the adult it is usually ossified and tipped with cartilage, but it very often continues cartilaginous through life, and Haller once found it in this state in a woman who died in her hundredth year.

The substance of the sternum, internally, is of a light, spongy texture, covered externally with a thin bony plate; hence it happens that this bone is easily fractured. From the description we have given of it, its uses may be easily understood. We have seen it serving for the articulation of seven true ribs on each side, and hence we shall find it of considerable use in respiration. We likewise observed, that it is articulated with each of the clavicles. It serves for the origin and insertion of several muscles; it supports the mediastinum; and lastly, defends the heart and lungs; and it is observable, that we find a similar bone in almost all animals that have lungs, and even in such as have no ribs, of which latter we have an instance in the frog.

STERNUTAMENTORIA. See *Plarmica*.

STERTOR A noisy kind of respiration, as is observed in apoplexy. A snoring, or snorting.

STHENIA. A term employed by the followers of Dr. Brown, to denote that state of the body which disposes to inflammatory diseases in opposition to those of debility, which arise from *asthenia*.

STIBIALIA. (From *stibium*, *antimony*.)

Antimonials. Medicines whose chief ingredient is antimony.

STIBII ESSENTIA. Antimonial wine.

STIBIUM. (*Στίβιον*: from *stibis*, to shine.) An ancient name of antimony. See *Antimonium*.

STIGMA. (*Στίγμα*: from *stige*, to inflict blows.) A small red speck in the skin, occasioning no elevation of the cuticle. Stigmata are generally distinct, or apart from each other. They sometimes assume a livid colour, and are then termed *petechia*.

STILBOMA. (From *stibis*, to polish.) A cosmetic.

STILLICIDIUM. (From *stille*, to drop, and *cado*, to fall.) A strangury, or discharge of the urine drop by drop. Also the pumping upon a part.

STIMM. *Στίμι*, Antimony.

STIMULANTS. (*Stimulantia*, sc. *medicamenta*, from *stimulo*, to stir up.) Medicines are so termed which possess a power of exciting the animal energy. They are divided into, 1. *Stimulantia sensu*, as *sinapi*, *santharides*, *mercurii preparationes*. 2. *Stimulantia defensibilia*, as *alkali volatile*, *electricity*, *heat*, &c. 3. *Stimulantia corporalia*, as *cinnamomum*, *sux moschata*, *vinum*, &c.

STIMULUS. Any thing which irritates.

Stinking lettuce. See *Lactuca grossula*.

STIZOLOBIUM. The cowage is sometimes so called. See *Dolichos*.

STOECHAS. (From *στοχαστε*, the islands on which it grew.) French lavender.

STOECHAS ARABICA. French lavender. *Spica hortulana.* *Stucadore.* *Lavandula stoechas* of Linnæus. This plant is much less grateful in smell and flavour than the common lavender, to which it is allied in its properties.

STOECHAS CITRINA. See *Elichrysum*.

STOMACACE. (From *stoma*, the mouth, and *zaco*, evil.) Canker. A fætor in the mouth, with a bloody discharge from the gums. It is generally a symptom of the scurvy. It is also a name for the scurvy.

STOMACH. (The word *stomachus* properly belongs to the upper orifice of the stomach, though given to the whole viscus.) *Ventriculus.* *Anocelia.* *Gaster.* *Nodus.* A membranous receptacle, situated in the epigastric region, which receives the food from the œsophagus; its figure is somewhat oblong and round: it is largest on the left side, and gradually diminishes towards its lower orifice, where it is the least. Its superior orifice, where the œsophagus terminates, is called the *cardia*; the inferior orifice, where the intestine begins, the *pylorus*.

The anterior surface is turned towards the abdominal muscles, and the posterior opposite the lumbar vertebrae. It has two curvatures; the first is called the great curvature of the stomach, and extends downwards, from one orifice to the other, having the omentum adhering to it; the second is the small curvature, which is also between both orifices, but superiorly and posteriorly. The stomach, like the intestinal canal, is composed of three coats, or membranes: 1. The *outermost*, which is very firm, and from the peritonæum. 2. The *muscular*, which is very thick, and composed of various muscular fibres; and, 3. The *innermost*, or *villous coat*, which is covered with exhaling and inhaling vessels, and mucus. These coats are connected together by cellular membrane. The glands of the stomach which separate the mucus are situated between the villous and muscular coat, in the cellular structure. The arteries of the stomach come chiefly from the celiac artery, and are distinguished into the coronary, gastro-epiploic, and short arteries; they are accompanied by veins which have similar names, and which terminate in the vena portæ. The nerves of the stomach are very numerous, and come from the eighth pair and intercostal nerves. The lymphatic vessels are distributed throughout the whole substance, and proceed immediately to the thoracic duct. The use of the stomach is to excite hunger and partly thirst, to receive the food from the œsophagus, and to retain it, till, by the motion of the stomach, the admixture of various fluids, and many other changes, it is rendered fit to pass the right orifice of the stomach, and afford chyle to the intestines.

Stomach, inflammation of. See *Gastritis*.

STOMACHICA PASSIO. A disorder in which there is an aversion to food, even the thought of it begets a nausea, anxiety, cardialgia, an effusion of saliva, and often a vomiting. Fasting is more tolerable than eating; if obliged to eat, a pain follows that is worse than hunger itself.

STOMACHICS. (*Stomachica*, sc. *medicamenta*; from *σμάχης*, the stomach.) Medicines which excite and strengthen the action of the stomach.

STOMACHUS. See *Stomach*.

Stone. See *Calculus*.

Stonacrop. See *Illecebra*.

STORAX. *Storaç.* See *Syrax*.

Storax, liquid. See *Liquidambra*.

STORAX LIQUIDA. See *Liquidambra*.

STORAX RUBRA OFFICINALIS. *Cascarilla bark*.

Storax, white. See *Balsamum peruvianum*.

STRABISMUS. See *Strabismus*.

STRABISMUS. (From *στράβω*, to squint.) *Strabismus*. *Strabotitas*. Squint-

ing. An affection of the eye by which a person sees objects in an oblique manner, from the axis of vision being distorted. Cullen arranges this disease in the class *locales*, and order *dyscinesie*. He distinguishes three species.

1. *Strabismus habitusalis*, when from a custom of using only one eye.

2. *Strabismus commodus*, when one eye in comparison with the other, from greater weakness, or mobility, cannot accommodate itself to the other.

3. *Strabismus necessarius*, when some change takes place in the situation or figure of the eye, or a part of it.

STRABOTITAS See *Strabismus*.

STRANEN CAMELORUM. Camel's hay, or *juncus odoratus*.

STRAMONIUM. See *Stramonium*.

STRAMONIUM. (From *stramen*, straw; so called from its fibrous roots.) *Dutray. Barryo coccalon. Solanum manicum* of Dioscorides, and *Stramonium spinosum* of Gerard. *Solanum fastidium* of Bauhin. *Stramonium majus album*. Common thorn-apple. *Datura stramonium* of Linnaeus:—*pericarpus spinosus erectus ovatis, foliis ovatis glabris*. This plant has been long known as a powerful narcotic poison. In its recent state it has a bitterish taste, and a smell somewhat resembling that of poppies, especially if the leaves be rubbed between the fingers. Instances of the deleterious effects of the plant are numerous, more particularly of the seed. An extract prepared from the seeds is recommended by Baron Stoerck in maniacal, epileptic, and convulsive affections; and is said by some to succeed, while, in the hands of others, it has failed. In this country, says Dr. Woodville, we are unacquainted with any practitioners whose experience tends to throw light on the medical character of this plant. It appears to us, continues Dr. Woodville, that its effects as a medicine are to be referred to no other power than that of a narcotic. And Dr. Cullen, speaking on this subject, says, "I have no doubt that narcotics may be a remedy in certain cases of mania and epilepsy; but I have not, and I doubt if any other person has, learned to distinguish the cases to which such remedies are properly adapted. It is therefore that we find the other narcotics, as well as the stramonium, to fail in the same hands in which they had in other cases, seemed to succeed. It is this consideration that has occasioned my neglecting the use of stramonium, and therefore prevented me from speaking more precisely from my own experience on this subject."

The extract of this plant has been the preparation usually employed, and from one to ten grains and upwards a day; but the powdered leaves, after the manner of those directed of cicuta, would seem to be

more certain and convenient. Greding found the strength of the extract to vary exceedingly; that which he obtained from Ludwig was much more powerful than that which he had of Stoerk. Externally, the leaves of stramonium have been applied to inflammatory tumours and burns, and it is said with success, and, of late, the dried leaves have been smoked as a remedy in asthma; but it does not appear that they have been more efficacious in this way than tobacco.

STRAMONIUM OFFICINALE. See *Stramonium*.

STRAMONIUM SPINOSUM. See *Stramonium*.

STRANGALIS. (From *σπασμα*, to torment.) A hard painful tumour in the breast, from milk.

STRANGURY. (*Stranguria*. From *σπασμα*, a drop, and *ουρα*, urine.) A difficulty of making water, attended with pain and dripping. See *Ischuria*.

STRATIGOTES. (From *στρατος*, an army; so named from its virtues in healing fresh wounds, and its usefulness to soldiers.) See *Millefolium*.

STRATIOTICUM. See *Millefolium*.

Strawberry. See *Fragaria*.

STREATHAM WATERS. A weak purging water, drunk from one, two, or more pints in a morning.

STREMA. (*Στρεμμα*; from *σπασμα*, to turn.) A strain, or sprain, of the parts about a joint.

STRICTURE. A diminution, or contracted state of some tube, or duct, of the body; as the œsophagus, intestines, urethra, vagina, &c. They are either organical or spasmodic.

STRIDOR DENTIUM. Grinding of the teeth.

STRIGIL. *Strigilis*. An instrument to scrape off the sweat during the gymnastic exercises of the ancients, and in their baths; *strigils* were made of metals, horn, ivory, and were curved. Some were made of linen.

STRIGMENTUM. The strigment, filth, or sordes, scraped from the skin, in baths and places of exercises.

STROPHOS. (From *σπασμα*, to turn.) A twisting of the intestines.

STROPHULUS. A papulous eruption peculiar to infants, and exhibiting a variety of forms, which are described by Dr. Willan, under the titles of *interunctus*, *albidus*, *confertus*, *volaticus*, and *candidus*.

1. *Strophulus interunctus*, (from *interuncto*, a spot here and there,) usually called the *red-gum*, and, by the French, *Efflorescence benigne*. The papulæ characterizing this affection, rise sensibly above the level of the cuticle, are of a vivid red colour, and commonly distinct from each other. Their number and extent varies much in different cases. They appear most con-

stantly on the cheeks, fore-arm, and back of the hand, but are sometimes diffused over the whole body. The papulæ are, in many places, intermixed with stigmata, and often with red patches of a larger size, which do not, however, occasion any elevation of the cuticle. A child's skin thus variegated, somewhat resembles a piece of red printed linen; and hence this eruption was formerly called the *red gum*, a term which is still retained in several counties of England, and may be found in old dictionaries. Medical writers have changed the original word for one of a similar sound, but not more significant. The *strophulus interunctus* has not, in general, any tendency to become pustular, a few small pustules containing a straw-coloured watery fluid, occasionally appear on the back of the hand, but scarcely merit attention, as the fluid is always re-absorbed in a short time, without breaking the cuticle. The eruption usually terminates in scurf, or exfoliation of the cuticle; its duration, however, is very uncertain; the papulæ and spots sometimes remain for a length of time, without an obvious alteration; sometimes disappear and come out again daily; but, for the most part, one eruption of them succeeds another, at longer intervals, and with more regularity. This complaint occurs chiefly within the two first months of lactation. It is not always accompanied with, or preceded by any disorders of the constitution, but appears occasionally in the strongest and most healthy children. Some authors connect it with aphthous ulcerations common in children, supposing the latter to be a part of the same disease diffused along the internal surfaces of the mouth and intestines. The fact, however, seems to be, that the two affections alternate with each other; for those infants, who have the papulous eruption on the skin are less liable to aphthæ; and when the aphthæ take place to a considerable degree, the skin is generally pale and free from eruption. The *strophulus interunctus* is, by most writers, said to originate from an acidity, or acrimonious quality of the milk taken into a child's stomach, communicated afterwards to the blood, and stimulating the cutaneous excretories. This opinion might, without difficulty, be proved to have little foundation. The pre-disposition to the complaint may be deduced from the delicate and tender state of the skin, and from the strong determination of blood to the surface, which evidently takes place in infants. The papulous eruption is, in many cases, connected with a weak, irritable state of the alimentary canal, and consequent indigestion. For if it be by any means suddenly repelled from the surface, diarrhœa, vomiting, spasmodic affections of the bowels, and often general disturbance

of the constitution succeed; but as soon as it reappears, those internal complaints are wholly suspended. Dr. Armstrong and others have particularly noted this reciprocation, which makes the red gum, at times, a disease of some importance, though in its usual form, it is not thought to be in any respect dangerous. On their remarks a necessary caution is founded, not to expose infants to a stream of very cold air, nor to plunge them unseasonably in a cold bath. The most violent, and even fatal symptoms have often been the consequence of such imprudent conduct.

2. The *Strophulus albidus*, by some termed the *white gum*, is merely a variety of strophulus intertinctus, but deserves some notice on account of the different appearance of its papulæ. In place of those described as characterizing the red gum, there is a number of minute, whitish specks, a little elevated, and sometimes, though not constantly, surrounded by a slight redness. These papulæ, when their tops are removed, do not discharge any fluid; it is, however, probable that they are originally formed by the deposition of a fluid, which afterwards concretes under the cuticle. They appear chiefly on the face, neck, and breast, and are more permanent than the papulæ of the red gum. In other respects, they have the same nature and tendency, and require a similar plan of treatment. Although a distinctive name has been applied to this eruption, when occurring alone, yet it is proper to observe that, in a great number of cases, there are red papulæ and spots intermixed with it, which prove its connexion with the strophulus intertinctus.

3. The *Strophulus confertus*. (From *confertus*, to crowd together.) An eruption of numerous papulæ, varying in their size, appears on different parts of the body in infants, during dentition, and has thence been denominated the *tooth-rash*. It is sometimes also termed the *rank red gum*. About the fourth or fifth month after birth, an eruption of this kind usually takes place on the cheeks and sides of the nose, extending sometimes to the forehead and arms, but rarely to the trunk or body. The papulæ on the face are smaller, and set more closely together than in the red gum; their colour is not so vivid, but they are generally more permanent. They terminate at length with slight exfoliations of the cuticle, and often appear again in the same places, a short time afterwards. The papulæ which, in this complaint, occasionally appear on the back or loins, are much larger, and somewhat more distant from each other, than those on the face. They are often surrounded by an extensive circle of inflammation, and a few of them contain a semi-pellucid watery fluid, which is reabsorbed when the in-

flammation subsides. In the seventh or eighth, the strophulus confertus assumes a somewhat different form; one or two large irregular patches appear on the arms, shoulder, or neck; in which the papulæ are hard, of a considerable size, and set so close together, that the whole surface is of a high red colour. Most commonly the fore arm is the seat of this eruption, the papulæ rising first on the back of the hand, and gradually extending upwards along the arm. Sometimes, however, the eruption commences at the elbow, and proceeds a little upwards and downwards on the outside of the arm. It arrives at its height in about a fortnight, the papulæ then begin to fade, and becomes flat at the top, afterwards the cuticle exfoliates from the part affected, which remains discoloured, rough, and irregular, for a week or two longer.

An obstinate and very painful modification of this disease takes place, though not often, on the lower extremities. The papulæ spread from the calves of the legs to the thighs, nates, loins, and round the body, as high as the navel: being very numerous and close together, they produce a continuous redness over all the parts above mentioned.

The cuticle presently, however, shrivelled, cracks in various places, and finally separates from the skin in large pieces. During this process a new cuticle is formed, notwithstanding which the complaint recurs in a short time, and goes through the same course as before. In this manner successive eruptions take place, during the course of three or four months, and perhaps do not cease till the child is one year old, or somewhat more. Children necessarily suffer great uneasiness from the heat and irritation occasioned by so extensive an eruption, yet, while they are affected with it, they often remain free from any internal or febrile complaint. This appearance should be distinguished from the intertrigo of infants, which exhibits a uniform, red, smooth, shining surface, without papulæ; and which affects only the lower part of the nates and inside of the thighs, being produced by the stimulus of the urine, &c. with which the child's clothes are almost constantly wetted. The strophulus confertus, where the child is otherwise healthy, is generally ascribed to a state of indigestion, or some feverish complaint of the mother, or nurse. Dr. Willan, however, asserts that he has more frequently seen the eruption when no such cause was evident. It may, with more probability, be considered as one of the numerous symptoms of irritation arising from the inflamed and painful state of the gums in dentition: since it always occurs during that process, and disappears soon after the first teeth have cut the gums.

4. The *Strophulus voluticus*, (from *volo*, to fly,) is characterized by an appearance of small circular patches, or clusters of papulæ, arising successively on different parts of the body. The number of papulæ in each cluster is from six to twelve. Both the papulæ and their interstices are of a high red colour. These patches continue red, with a little heat, or itching, for about four days, when they turn brown, and begin to exfoliate. As one patch declines, another appears at a small distance from it; and in this manner the complaint often spreads gradually over the face, body, and limbs, not terminating in less than three or four weeks. During that time the child has sometimes a quick pulse, a white tongue, and seems uneasy and fretful. In many cases, however, the eruption takes place without any symptoms of internal disorder. The above complaint has been by some writers denominated *ignis voluticus infantum*; under this title Astruc and Lorry have described one of the forms of *crusta lactea*, in which a successive eruption of pustules takes place on the same spot generally about the mouth or eyes, in children of different ages, and sometimes in adults. The *macula volutice infantum* mentioned by Wittichius, Sennerius, and Sebizeus, agree in some respect with the *strophulus voluticus*; but they are described by other German authors as a species of *erysipelas*, or as irregular efflorescences affecting the genitals of infants, and often proving fatal. The *strophulus voluticus* is a complaint by no means frequent. In most cases which have come under Dr. Willan's observation, it appeared between the third and sixth month; in one instance, however, it occurred about ten days after birth, and continued three weeks, being gradually diffused from the cheeks and forehead to the scalp, afterwards to the trunk of the body and to the extremities; when the patches exfoliated, a red surface was left, with slight border of detached cuticle.

5. *Strophulus candidus*. (From *candeo*, to shine.) In this form of *strophulus*, the papulæ are larger than in any of the foregoing species. They have no inflammation round their base; their surface is very smooth and shining, whence they appear to be of a lighter colour than the adjoining cuticle. They are diffused, at a considerable distance from each other, over the loins, shoulders, and upper part of the arms; in any other situation they are seldom found.

This eruption affects infants about a year old, and most commonly succeeds some of the acute diseases to which they are liable. Dr. Willan has observed it on their recovery from a catarrhal fever, and after inflammations of the bowels, or lungs. The papulæ continue hard and

elevated for about a week, then gradually subside and disappear.

STRUMA. (From *struo*, to heap up.) This term is applied by some authors to scrofula, and by others to an induration of the thyroid gland, which is endemial to the Tyrolese, Swiss, and others.

STRUMEX. (From *struma*, a scrophulous tumour.) A herb so called from its uses in healing strumous tumours.

STRUTHIUM. (From *στρουθιον*, a sparrow; so named from the resemblance of its flowers to an unfledged sparrow.) The master-wort. See *Imperatoria*.

STRICHNOMANIA. (From *στροχνη*, nightshade, and *μανια*, madness.) So the ancients called the disorder produced by eating the deadly nightshade.

STRYCHNOS. (From *στροχνη*, to torment; so named from its properties of producing insanity.) The name of a genus of plants in the Linnæan system.

STRYCHNOS RUX VOMICA. The systematic name of the tree whose seed is called the poison nut. See *Nuxvomica*.

STRYCHNOS VELUTIBILIS. The systematic name of the tree which is supposed to afford the Jesuits bean. See *Faba indica*.

STUPEFACIENTIA. (From *stupefacio*, to stupefy.) Narcotica.

STUPA. (From *στυπη*, to bind.) *Stupa*, *Stuppa*. A stupe; the same as fomentation.

STUPOR. (From *stupes*, to be senseless.) Insensibility.

STUPOR DENTITUM. Commonly called tooth on-edge.

STUPPA. See *Stupha*.

Stye. See *Hordolium*.

STYRIA. (From *Styx*, a name given by the poets to one of the rivers in hell.) A water made from sublimate, and directed in old dispensatories, so called from a supposition of its poisonous qualities. The *Aqua Regia* is also thus sometimes called, from its corrosive qualities.

STYLIFORM. (*Styliformis*; from *stylus*, a bodkin, and *forma*, a likeness.) Shaped like a bodkin, or style.

STYLISCV. (From *στυλιν*, a bodkin.) A tent made in the form of a bodkin.

STYLO. Names compounded of this word belong to muscles which are attached to the styloid process of the temporal bone; as,

STYLO-CERATO-HYOIDEUS. See *Stylo-hyoideus*.

STYLO-CHONDRO-HYOIDEUS. See *Stylo-hyoideus*.

STYLO-GLOSSUS. (*Musculus stylo-glossus*.) *Stylo-glossæ* of Dumas. A muscle situated between the lower jaw and os hyoides laterally, which draws the tongue aside and backwards. It arises tendinous and fleshy from the styloid process, and from the ligament which connects that process to the angle of the lower jaw, and

is inserted into the root of the tongue, runs along its sides, and is insensibly lost near its tip.

STYLO-HYOIDEUS. (*Musculus stylo-hyoideus*.) *Stylo-hyoideus* of Dumas. A muscle situated between the lower jaw and os hyoides laterally, which pulls the os hyoides to one side and a little upwards.

It is a small, thin, fleshy muscle, situated between the styloid process and os hyoides, under the posterior belly and middle tendon of the digastricus, near the upper edge of that muscle.

It arises by a long thin tendon, from the basis and posterior edge of the styloid process, and, descending in an oblique direction, is inserted into the lateral and anterior part of the os hyoides, near its horn.

The fleshy belly of this muscle is usually perforated on one or both sides, for the passage of the middle tendon of the digastricus.

Sometimes, though not always, we find another smaller muscle placed before the stylo-hyoideus, which, from its having nearly the same origin and insertion, and the same use, is called *stylo-hyoideus-alter*. It seems to have been first known to Eustachius; so that Douglas was not aware of this circumstance when he placed it amongst the muscles discovered by himself. It arises from the apex of the styloid process, and sometimes, by a broad and thin aponeurosis, from the inner and posterior part of the angle of the lower jaw, and is inserted into the appendix, or little horn, of the os hyoides.

The use of these muscles is to pull the os hyoides to one side, and a little upwards.

STYLO-HYOIDEUS ALTER. See *Stylo-hyoideus*.

STYLO-MASTOID FORAMEN. *Foramen stylo-mastoideum*. A hole between the styloid and mastoid process of the temporal bone, through which the portio dura of the auditory nerve passes to the temples.

STYLO-PHARYNGEUS. (*Musculus stylo-pharyngeus*.) *Stylo-thyro-pharyngien* of Dumas. A muscle situated between the lower jaw and os hyoides laterally, which dilates and raises the pharynx and thyroid cartilage upwards. It arises fleshy from the root of the styloid process, and is inserted into the side of the pharynx and back part of the thyroid cartilage.

STYMATOSIS. (From *σῡς*, to have a priapism.) A violent erection of the penis, with a bloody discharge.

STYPERIA. (From *σῡς*, to bind; so called from its adstringent properties.) Alum.

STYPTICS. (*Medicamenta styptica*, from *σῡς*, to adstringe.) A term given to those substances which possess the power of stopping hæmorrhages, such as turpentine, alum, &c.

STYRACIFLUA. (From *styrax*, storax,

and *fluō*, to flow.) Liquid storax. See *Styrax*.

STYRAX. (From *σῡραξ*, a reed, in which it was used to be preserved.) 1. The name of a genus of plants in the Linnæan system. Class, *Decandria*. Order, *Monogynia*.

2. The pharmacopœial name of the *Styrax calamita*. Official storax. *Styrax officinalis*.—*foliis ovatis, subtus villosis, racemis simplicibus folio brevioribus*. There are two kinds of storax to be found in the shops; the one is usually in irregular compact masses, free from impurities, of a reddish brown appearance, and interspersed with whitish tears, somewhat like gum ammoniac, or benzoin; it is extremely fragrant, and upon the application of heat readily melts. This has been called *storax in lump*, *red storax*; and in separate tears, *storax in tears*. The other kind, which is called the *common storax*, is in large masses, very light, and bears no external resemblance whatever to the former storax, as it seems almost wholly composed of dirty saw-dust, caked together by resinous matter. Storax was formerly used in catarrhal complaints, coughs, asthmas, obstructions, &c. In the present practice it is almost totally disregarded, notwithstanding it is an efficacious remedy in nervous diseases.

STYRAX ALBA. See *Balsamum peruvianum*.

STYRAX BENZOIN. The systematic name of the tree which affords the gum benzoin. See *Benzoïn*.

STYRAX CALAMITA. Storax in the cane, because it was formerly brought to us in reeds, or canes. See *Styrax*.

STYRAX COLATA. Strained storax.

STYRAX LIQUIDA. Liquid storax. See *Liquidambar*.

STYRAX OFFICINALIS. The systematic name of the tree which affords the solid styrax. See *Styrax*.

STYRAX RUBRA. Red storax, or in the tear.

SUBACETAS CUPRI. See *Verdigris*. *Subacetate of copper*. See *Verdigris*.

SUBALARIS VENA. The vein of the axilla or arm-pit.

SUBCARBONAS POTASSÆ. See *Potasse subcarbonas*.

SUBCARBONAS FERRI. See *Carbonas ferri*.

SUBCARBONAS PLUMBI. Subcarbonate of lead. Ceruse. White lead. This article is made in the large way in white lead-manufactories, by exposing thin sheets of lead to the vapour of vinegar. The lead is curled up and put into pots of earthen ware in which the vinegar is, in such a way as to rest just above the vinegar. Hundreds of these are arranged together and surrounded with dung, the heat from which volatilizes the acetic acid, which is decomposed by the lead, and an

imperfect carbonate of lead is formed, which is of a white colour. This preparation is seldom used in medicine or surgery but for the purpose of making other preparations, as the superacetate. See *Pulvis cerusee compositus*.

SUBCARTILAGINEUM. (From *sub*, under, and *cartilago*, a cartilage.) The hypochondrium, or part of the body which lies under the cartilages of the spurious ribs.

SUBCLAVIAN ARTERY. (From *sub*, under, and *clavia*, a key, because the clavicles were supposed to resemble the key of the ancients.) The right subclavian arises from the arteria innominata, and proceeds under the clavicle to the axilla. The left subclavian arises from the arch of the aorta, and ascends under the left clavicle to the axilla. The subclavians in their course give off the internal mammary, the cervical, the vertebral, and the superior intercostal arteries.

SUBCLAVIAN VEIN. This receives the blood from the veins of the arm, and runs into the vena cava superior.

SUBCLAVIUS. (*Musculus subclavius*; from *sub*, under, and *clavicula*, the channel bone, as being situated under the clavicle, or channel bone.) *Subclaviusmus*. *Coste claviculaire* of Dumas. A muscle, situated on the anterior part of the thorax, which pulls the clavicle downwards and forwards. It arises tendinous from the cartilage which joins the first rib to the sternum, is inserted after becoming fleshy into the inferior part of the clavicle, which it occupies from within an inch of the sternum as far outwards as to its connexion, by a ligament, with the coracoid process of the scapula.

SUBCRURÆI. Two little muscular slips sometimes found under the cruræus; they are inserted into the capsular ligament which they pull up. The cruræus assists in extending the leg.

SUBCUTANEOUS GLANDS. *Glandulae subcutaneæ*. These are sebaceous glands lying under the skin, which they perforate by their excretory ducts.

SUBCUTANEUS. The platysma myoides muscle.

SUBER. The cork-tree. The fruit of this tree, *Quercus suber* of Linnaeus, are much more nutritious than our acorns, and are sweet and often eaten when roasted in some parts of Spain. The bark, called cork, when burnt, is applied as an adstringent application to bleeding piles and to allay the pain usually attendant on hæmorrhoids, when mixed with an ointment. Pessaries and other surgical instruments are also made of this useful bark.

SUBLIMAMENTUM. (From *sublimis*, to lift up.) The pendulous substance which floats in the middle of the urine.

Sublimatæ. See *Oxymurias hydrargyri*.

SUBLIMATION. (*Sublimatio*, from

sublimis, to raise or sublime.) This chemical process differs from evaporation only in being confined to solid substances. It is usually performed either for the purpose of purifying certain substances, and disengaging them from extraneous matters; or else to reduce into vapour, and combine, under that form, principles which would have united with greater difficulty if they had not been brought to that state of extreme division.

As all fluids are volatile by heat, and consequently capable of being separated, in most cases, from fixed matters, so various solid bodies are subjected to a similar treatment. Fluids are said to distil, and solids to sublime, though sometimes both are obtained in one and the same operation. If the subliming matter concretes into a solid, hard mass, it is commonly called a sublimate; if into a powdery form, flowers.

The principal subjects of this operation are, volatile alkaline salts; neutral salts, composed of volatile salts and acids, as sal ammoniac, the salt of amber, and flowers of benzoin, mercurial preparations, and sulphur. Bodies of themselves not volatile are frequently made to sublime by the mixture of volatile ones, thus iron is carried by sal ammoniac in the preparations of the flores martialis, or ferrum ammoniacale.

The fumes of solid bodies in close vessels rise but a little way, and adhere to that part of the vessel where they concrete.

SURLIMIS. See *Flexor brevis digitorum pedis*, and *Flexor sublimis perforatus*.

SUBLINGUAL GLANDS. *Glandule sublinguales* vel *Bartholinianæ* vel *Brownianæ*. The glands which are situated under the tongue, and secrete saliva. Their excretory ducts are called *Rivierian* from their discoverer.

SUBLUXATIO. A sprain.

SUBMERSION. (*Submersio*, from *sub*, under, and *mergo*, to sink.) Drowning. A variety of the apoplexia suffocata. Sauvages terms it asphyxia immersorum.

SUBMURIAS HYDRARGYRI. Submuriate of mercury. *Calomelas*. "Take of oxymuriate of mercury, a pound; purified mercury, by weight one ounce." Rub them together until the metallic globules disappear, then sublime; take out the sublimed mass, and reduce it to powder, and sublime it in the same manner twice more successively. Lastly, bring it into the state of very fine powder by the same process which has been directed for the preparation of chalk.

Submuriat, or mild muriat of mercury, is one of the most useful preparations of mercury. As an anti-venereal it is given in the dose of a grain night and morning, its usual determination to the intestines being prevented, if necessary, by opium. It is

the preparation which is perhaps most usually given in the other diseases in which mercury is employed, as in affections of the liver, or neighbouring organs, in cutaneous diseases, chronic rheumatism, tetanus, hydrophobia, hydrocephalus, and febrile affections, especially those of warm climates. It is employed as a cathartic alone, in doses from v. to xii. grains, or to promote the operation of other purgatives. Its anthelmintic power is justly celebrated; and it is perhaps superior to the other mercurials in assisting the operation of diuretics is dropsy. From its specific gravity it ought always to be given in the form of a bolus or pill.

SUBORBITARIUS. The suborbital nerve; a branch of the fifth pair.

SUBSCAPULARIS. (*Musculus subscapularis*, from *sub*, under, and *scapula*, the shoulder-blade.) *Sous-scapulo-trachien* of Dumas. The name of this muscle sufficiently indicates its situation. It is composed of many fasciculi of tendinous and fleshy fibres, the marks of which we see imprinted on the under surface of the scapula. These fasciculi, which arise from all the basis of that bone internally, and likewise from its superior, as well as from one half of its inferior costa, unite to form a considerable flat tendon which adheres to the capsular ligament, and is inserted into the upper part of the lesser tuberosity at the head of the os humeri.

The principal use of this muscle is to roll the arm inwards. It likewise serves to bring it close to the ribs; and, from its adhesion to the capsular ligament, it prevents that membrane from being pinched.

SUBSULTUS TENDINUM. (*Subsultus*, from *subsulto*, to leap.) Weak convulsive motions or twitchings of the tendons, mostly of the hands, generally observed in the extreme stages of putrid fever.

SUBUBERES. (From the two opposite prepositions *sub* and *ex*, and *abern*, the breasts.) *Eruberes*. Hath been used by some writers for those infants who yet suck, in distinction from those who are weaned, and then are called *eruberes*.

SUCCAGO. The rob or conserve of fruit.

SUCCEDANEUM. A medicine substituted for others.

SUCCENTURIATI MUSCULI. The pyramidal muscles of the belly.

SUCCENTURIATI RENES. Two glands lying above the kidneys.

SUCCINAS AMMONIACÆ. See *Spiritus ammoniac succinatus*.

SUCCINATE. *Succinas*. Salts formed by the combination of the acid of amber, or succinic acid, with different bases; as, *succinate of potash*, *succinate of copper*, &c.

SUCCINGEUS MEMBRANA. The diaphragm.

SUCCINIC ACID. *Acidum succinicum*.

Succinum. The succinic acid is drawn from amber by sublimation in a gentle heat, and rises in a concrete form into the neck of the subliming vessel. The operation must not be pushed too far, nor by too strong a fire, otherwise the oil of amber rises along with the acid.

SUCCINUM. (From *succus*, juice; because it was thought to exude from a tree.) See *Amber*.

SUCCINUM CINEREUM. The ambergris is so called by some authors. See *Ambergri-sea*.

SUCCINUM ERISUM. The ambergris is sometimes so called. See *Ambergri-sea*.

SUCCINUM OLEUM. See *Oleum succini*.

SUCCINUM PREPARATUM. Prepared amber. See *Amber*.

SUCCISA. (From *succido*, to cut; so named from its being indented and as it were cut in pieces.) Devil's bit. A species of the genus *Scabiosa*.

SUCCI SCORBUTICI. The juice of English scurvy-grass.

Succory. See *Cichorium*.

SUCCUBUS. See *Incubus*.

SUCCUS. Juice.

SUCCUS ACACIÆ VERÆ. See *Acacia vera*.

SUCCUS ACONITI SPISSATUS. See *Aconitum*.

SUCCUS BACCÆ SAMBUCI SPISSATUS. An aperient and deobstruent extract, often employed diluted with water in the cure of catarrhal affections.

SUCCUS BELLADONNÆ SPISSATUS. See *Belladonna*.

SUCCUS CICUTÆ SPISSATUS. See *Cicuta*.

SUCCUS COCHLIARIÆ COMPOSITUS. A warm aperient and diuretic, mostly exhibited in the cure of diseases of the skin arising from scurvy.

SUCCUS CYRENIIACUS. Juice of laserwort.

SUCCUS GASTRICUS. See *Gastric juice*.

SUCCUS GLYCYRRHIZÆ. Spanish liquorice.

SUCCUS HELIOTROPIL. See *Bezetta ceru-lia*.

SUCCUS HYOSCIAMI SPISSATUS. See *Hyosciamus*.

SUCCUS INDICUS PURGANS. Gamboge.

SUCCUS LACTUCÆ VIROSÆ SPISSATUS. See *Lactuca grævesolens*.

SUCCUS LIMONIS SPISSATUS. See *Limon*.

SUCCUS LIQUORITIL. Spanish liquorice.

SUCCUS PRUNORUM SYLVESTRUM. See *Asacia Germanica*.

SUCCUS SPISSATUS BACCORUM SAMBUCI. Rob of elder-berries.

SUDAMINA. (*Sudamen*, from *sudor*, sweat.) *Hydraea*. Boas. Vesicles resembling millet-seeds in form and magnitude, which appear suddenly, without fever, especially in the summer-time after much labour and sweating.

SUDATIO. (From *suder*, sweat.) A sweating. See *Ephidrosis*.

SUDATORIUM. (From *sude*, to sweat.) A stew or sweating-house.

SUDOR ANGILICUS. Called also *Hydromela*. *Hydropyretos*. Gargatio. The sweating sickness of England; an endemic fever. Dr. Cullen thinks it a species of typhus. This disorder is thus named from its first appearing in this island, and acquires the title of sudor from the patient suddenly breaking out into a profuse sweat, which forms the great character of the disease.

SUDORIFICA. (*Sudorifica*, sc. *medicamenta*, from *suder*, sweat, and *ficio*, to make) *Hydrotica*. *Hydrotopoea*. A synonym of diaphoretics. See *Diaphoretica*.

SUFFUMENTUM. (From *suffimen*, a perfume.) *Hypocapsium*. A suffumigation, a perfume.

SUFFUTUS. The same.

SUFFOCATIO HISTERICA. A convulsive affection of the throat.

SUFFOCATIO STRIDULA. The croup.

SUFFUMIGATIO. (*Suffumigatio*, from *suf*, under, and *fumigo*, to cloke.) The burning odorous substances to remove an evil smell, or destroy miasma.

SUFFUSIO. (From *suffundo*, to pour down; so called because the ancients supposed the opacity proceeded from something running under the crystalline humour.) A cataract.

SUFFUSIO AURIGNOA. A jaundice.

Sugar. See *Saccharum*.

Sugar of lead. See *Plumbi superacetas*.

Sugar of milk. A substance produced from whey, which, if not sour, contains a saline substance to which this name has been given.

SUGILLATIO. (*Sugillatio*. From *sugilla*, to stain.) A bruise. A spot or mark made by a leech or cupping-glass.

SULCUS. A groove or furrow; generally applied to the bones.

SULPHAS. (From *sulphur*, brimstone.) A sulphat or salt formed by the union of the sulphuric acid with different bases.

SULPHAS ALUMINOSUS. Alum. See *Alumina*.

SULPHAS AMMONIÆ. *Alkali volatile vitriolatum* of Bergman. *Sal ammoniacum seoretum* of Glauber. *Vitriolum ammoniacale*. This salt has been found native in the neighbourhood of some volcanoes. It is esteemed diuretic and deobstruent, and exhibited in the same diseases as the muriate of ammonia.

SULPHAS CUPRI. *Vitriolum cupri*. *Vitriolum ceruleum*. *Vitriolum Romanum*. *Cuprum vitriolatum*. The sulphat of copper possesses acrid and styptic qualities; is esteemed as a tonic, emetic, adstringent, and escharotic, and is exhibited internally in the cure of dropsies, hæmorrhages, and as a speedy emetic. Externally it is applied to stop hæmorrhages, to hæmor-

rhoids, leucorrhœa, phagedenic ulcers, proud flesh, and condylomata.

SULPHAS FERRI. See *Ferri sulphas*.

SULPHAS HYDRARGYRI. See *Hydrargyri vitriolatum*.

SULPHAS MAGNESIÆ. *Sulphus magnesia purificata*. *Magnesia vitriolata*. *Sal catharticum amarum*. *Sal catharticum amarum*. Sulphate of magnesia. Epsom salt. Bitter purging salt.

The sulphate of magnesia exists in several mineral springs, and in sea water.

It is from these saline solutions that the salt is obtained; the method generally adopted for obtaining it is evaporation, which causes the salt to crystallize in tetrahedral prisms. It has a very bitter taste, and is soluble in its own weight of water at 60°, and in three-fourths of its weight of boiling water. Sulphate of magnesia, when perfectly pure, effloresces; but that of commerce generally contains foreign salts, such as the muriate of magnesia, which renders it so deliquescent that it must be kept in a close vessel or bladder. By the action of heat it undergoes the watery fusion, and loses its water of crystallization, but does not part with its acid. One hundred parts of crystallized sulphate of magnesia consist of 29.35 parts of acid, 17 of earth, and 53.65 of water. The alkalis, strontia, baryta, and all the salts formed by these salifiable bases, excepting the alkaline muriates, decompose sulphate of magnesia. It is also decomposed by the nitrate, carbonate and muriate of lime.

Epsom salt is a mild and gentle purgative, operating with sufficient efficacy, and in general with ease and safety, rarely occasioning any gripes or the other inconveniences of resinous purgatives. Six or eight drachmas may be dissolved in a proper quantity of common water; or four, five, or more in a pint or quart of the purging mineral water. These solutions may likewise be so managed as to produce evacuation from the other emunctories; if the patient be kept warm they increase perspiration, and by moderate exercise in the cool air, the urinary discharge. Some allege that this salt has a peculiar effect in allaying pain, as in colic, even independently of evacuation.

It is, however, principally used for the preparation of the carbonate of magnesia.

SULPHAS POTASSÆ. See *Potasse sulphas*.

SULPHAS SODÆ. See *Sodæ sulphas*.

SULPHAS ZINCI. See *Zinci sulphas*.

Sulphate. See *Sulphas*.

SULPHURE. *Sulphia*. A salt formed by the combination of the sulphureous acid with different bases: as, *aluminous sulphite*, *ammoniacal sulphite*, &c.

SULPHUR. *Abric*. *Alcubrit*. *An-*

pater. Appebrioc. Aquala. Aquila. Cibur. Chybur. Cibur. Sulphur, which is also known by the name of brimstone, is the only simple combustible substance which nature offers pure and in abundance. It was the first known of all. It is found in the earth, and exists externally in depositions, in sublimed incrustations, and on the surface of certain waters, principally near burning volcanoes. It is found combined with many metals. It exists in vegetable substances, and has lately been discovered in the albumen of eggs.

Sulphur in the mineral kingdom is either in a loose powder, or compact; and then either detached or in veins. It is found in the greatest plenty in the neighbourhood of volcanoes or pseudo-volcanoes, whether modern or extinct as at *Solfatara*, &c. and is deposited as a crust on stones contiguous to them, either crystallized or amorphous. It is frequently met with in mineral waters, and in caverns adjacent to volcanoes; sometimes also in coal-mines. It is found in combination with most of the metals. When united to iron it forms the mineral called *martial pyrites*, or *sulphur pyrites*. All the ores known by the name of *pyrites*, of which there are a vast variety, are combinations of sulphur with different metals; and hence the names of copper, tin, arsenical, &c. *pyrites*. It exists likewise in combination with alumine and lime; it then constitutes different kinds of *schistus*, or *alum ores*.

Physical properties.—Sulphur is a combustible, dry, and exceedingly brittle body, of a pale lemon-yellow colour. Its specific gravity is 1.990. It is destitute of odour, except when rubbed or heated. It is of a peculiar faint taste. It frequently crystallizes in entire or truncated octahedra, or in needles. If a piece of sulphur of a considerable size, be very gently heated, as for example, by holding it in the hand and squeezing it firmly, it breaks to pieces with a crackling noise. It is a non-conductor of electricity, and hence it becomes electric by friction. When heated, it first softens before it melts, and its fusion commences at 180° Fahr.; at 289° it becomes volatilized, and takes fire at 302°. In the beginning of fusion it is very fluid, but by continuing the heat it grows tough, and its colour changes to a reddish brown. If in this condition it be poured into water, it remains as soft as wax, and yields to any impression. In time however it hardens again and recovers its former consistence.

It unites with most of the earths and with all alkalis, and becomes soluble, when thus combined, in water. It unites to most of the metals, and renders them brittle and fusible. It is soluble in oils; water takes up a minute quantity, as does ardent spirit by means of heat. It dis-

solves in hydrogen gas. It does not combine with charcoal, but unites to phosphorus by fusion.

Sulphur, like all combustible bodies burns in proportion to the quantity of oxygen which combines with it. Sulphur, heated in a close vessel, sublimates without alteration. It is not changed by exposure to air. It is attacked by the nitric acid when poured on it in its fused state.

If melted sulphur be exposed in the open air to an increase of heat, it takes fire and burns with a blue flame and suffocating vapour. The result of this combustion is sulphureous acid. According to the slow or rapid combustion, it absorbs different quantities of oxygen, and the produced acid differs in its properties.

Method of obtaining sulphur.—A prodigious quantity of sulphur is obtained from *Solfatara* in Italy. This volcanic country every where exhibits marks of the agency of subterraneous fires; almost all the ground is bare, and white; and is every where sensibly warmer than the atmosphere, in the greatest heat of summer; so that the feet of persons walking there are burnt through their shoes. It is impossible not to observe the sulphur, for a sulphureous vapour which arises through different apertures is every where perceptible, and gives reason to believe that there is a subterraneous fire underneath from which that vapour proceeds.

From *pyrites* sulphur is extracted in the large way by the following process:

Pyrites is broken into small pieces, and put into large earthen tubes which are exposed to the heat of a furnace. A square vessel of cast iron, containing water, is connected as a receiver with the tube in the furnace. The action of the fire proceeds, and the sulphur, being thus melted, is gradually accumulated on the water in the receiver. It is then removed from this receiver, and melted in large iron ladles; in consequence of which, the earthy parts with which it was contaminated are made to subside to the bottom of the ladle, leaving the purified sulphur above. It is then again melted and suffered to cool gradually, in order to free it from the rest of the impurities. It is then tolerably pure, and constitutes the sulphur we meet with in large masses or lumps in the market.

In order to form it into rolls, it is again melted and poured into cylindrical wooden moulds, in these it takes the form in which we usually see it in commerce, as roll sulphur.

Flowers of sulphur, as they are called, are formed by subliming purified sulphur with a gentle heat in close rooms, where the sublimed sulphur is collected, though the article met with in general under that name is nothing but sulphur finely powdered.

Method of purifying sulphur.—Take one part of flowers of sulphur, boil it in twenty parts of distilled water in a glass vessel for about a quarter of an hour; let the sulphur subside, decant the water, and then wash the sulphur repeatedly in distilled water: having done this, pour over it three parts of pure nitro-muriatic acid, diluted with one part of distilled water, boil it again in a glass vessel for about a quarter of an hour, decant the acid, and wash the sulphur in distilled water till the fluid passes tasteless, or till it does not change the blue colour of tincture of cabbage, or litmus. The sulphur thus carefully treated is pure sulphur fit for philosophical experiments.

Sulphur has been long an esteemed article of the *Materia Medica*; it stimulates the system, loosens the belly, and promotes the insensible perspiration. It pervades the whole habit, and manifestly transpires through the pores of the skin, as appears from the sulphureous smell of persons who have taken it, and from silver being stained in their pockets of a blackish colour. In the stomach it is probably combined with hydrogen. It is a celebrated remedy against cutaneous diseases, particularly psora, both given internally and applied externally. It has likewise been recommended in rheumatic pains, flying gout, rickets, atrophy, coughs, asthmas, and other disorders of the breast and lungs, and particularly in catarrhs of the chronic kind.

In hemorrhoidal affections it is almost specific; but in most of these cases it is advantageously combined with some cooling purgative, especially supertartrate of potash.

The preparations of sulphur directed to be used by the London and Edinburgh Colleges, are the Sulphur lotum, Sulphur præcipitatum, and Sulphur sublimatum.

SULPHUR ANTIMONII PRÆCIPITATUM. *Sulphur auratum antimonii.* This preparation of antimony appears to have rendered that called *kermis mineral* unnecessary. It is a yellow hydro-sulphuret of antimony, and therefore called *hydro-sulphuretum stibii luteum*, in the new chemical nomenclature. As an alterative and sudorific it is in high estimation, and given in diseases of the skin and glands; and, joined with calomel, it is one of the most powerful and penetrating alteratives we are in possession of.

SULPHUR AURATUM ANTIMONII. See *Sulphur antimonii præcipitatum*.

SULPHUR LOTUM. Washed sulphur. *Flores sulphuris loti.* "Take of sublimed sulphur, a pound." Pour on boiling water: so that the acid, if there be any, may be entirely washed away; then dry it. The dose is from half a drachm to two drachms.

SULPHUR PRÆCIPITATUM. *Lac sulphuris.* "Take of sublimed sulphur, a pound, fresh lime, three pounds." Boil the sulphur and lime together in water, then strain the solution through paper, and drop in as much muriatic acid as may be necessary to precipitate the sulphur; lastly, wash this by repeated affusions of water until it is tasteless. This preparation is mostly preferred to the flowers of sulphur, in consequence of its being freed from its impurities. The dose is from half a drachm to three drachms.

SULPHUR SUBLIMATUM. Sublimed sulphur. See *Sulphur*.

Sulphureum. See *Pascedanum*.

Sulphurated hydrogen gas. See *Hydrogen gas, sulphurated*.

SULPHUR. *Sulphuretum.* A combination of sulphur with different alkaline, earthy, and metallic bases.

SULPHUREOUS ACID GAS. Several philosophers have paid attention to the properties of this gas, and its combinations; but Berthollet was the first who published an accurate account of it, which Fourcroy and Vauquelin, have investigated still more completely. Their paper is inserted in the *Annales de Chimie*, II. 54, and copied into *Nicholson's Journal*, vol. I. p. 313.

When sulphur is united to oxygen it forms an acid. When the quantity of oxygen is sufficient to oxygenate a given quantity of sulphur completely, the result is a perfect acid called sulphuric acid. If a less quantity of oxygen is made use of, the result is an imperfect acid, capable of taking the gaseous state. To this gaseous fluid the name of sulphureous acid gas is given. It becomes obvious from this, that sulphureous acid gas has the same relation to sulphuric acid as phosphorous acid has to the phosphoric acid. At least this gas occupies the middle place between sulphur and sulphuric acid.

Properties.—Sulphureous acid gas is a permanently elastic æriform fluid at the ordinary pressure and temperature of our atmosphere. Its odour is strong and suffocating. It cannot maintain combustion, nor the respiration of animals. Its weight is more than double that of atmospheric air. Its specific gravity according to Bergman is 0.00246, and 0.00251, according to Lavoisier. It is not inflammable. One hundred cubic inches of it weigh nearly 63 grains. It is composed, according to Fourcroy, of 85 sulphur and 15 oxygen. It is acid; it first reddens and then destroys the greater part of the vegetable colours. It exerts little action on the metals, and has a weak attraction to alkalis and earths. It has the property of whitening silk and giving it a lustre. Priestley, Bergman, Berthollet, &c. say that at high temperatures it deposits sulphur, but Fourcroy and Vauquelin, in con-

sequence of some new experiments, deny this assertion. Phosphorus has no action upon it. Its attraction for oxygen when dry, at the common temperature, is very feeble, but, if the smallest quantity of moisture be present, the union of the two gases is much favoured. At high temperatures the combination is more speedily effected. When a mixture of sulphureous acid gas and oxygen gas is made to pass through an ignited tube, the two gases combine and sulphuric acid is formed. When passed into water cooled down to the freezing point, the union is very rapid. Water cooled to forty degrees absorbs one-third part of its weight of sulphureous acid gas. It speedily melts ice. Water saturated with it may be frozen without parting with its gas; but when water, which has been saturated with it, is exposed to heat, it is filled with a vast number of bubbles which continually increase and rise to the surface; these bubbles are the gas separating from it. It is absorbed by oil, ether, and sulphuric acid, the latter when saturated with it acquires the property of smoking when exposed to air; its colour is altered to a yellowish brown, and its odour is penetrating like that of the gas. When the acid thus saturated with the gas is exposed a few degrees below the freezing point, in congeals or freezes into a crystalline mass. The same happens when submitted to distillation. It decomposes nitric and oxygenated muriatic acid, and dissolves camphor. Sulphureous acid gas and hydrogen gas have no action upon each other at common temperatures, but if they are passed through an ignited tube, a decomposition is effected. Charcoal likewise decomposes sulphureous acid gas at a high temperature. Monge and Clouet affirm that by extreme artificial cold and a strong pressure exerted at the same time, they rendered sulphureous acid gas fluid.

Methods of obtaining sulphureous acid gas.

—1. Take one part of mercury and four of concentrated sulphuric acid; put them into a glass retort connected with the pneumatic quicksilver-trough and apply the heat of a lamp to the mixture till it boils. On continuing the heat after the mercury has been acted upon, sulphureous acid gas will be obtained in abundance.

In this case a partial decomposition of the sulphuric acid takes place. The mercury abstracts part of its oxygen and becomes converted into an oxid; the sulphuric acid in consequence of the loss of oxygen, becomes converted into sulphureous acid, which takes the gaseous form at the common temperature.

2. Sulphureous acid gas may likewise be obtained by the slow combustion of sulphur.

Put pounded sulphur into a shallow earthen dish or crucible; melt and set fire to it;

and when it is completely inflamed cover it with a large bell-glass perfectly dry; remove the whole on a dish and surround the apparatus with mercury. The sulphur will burn for some time, and the bell-glass will be filled with white vapour. When the vapour has subsided, the gas may be transferred as usual.

Explanation.—During the burning of the sulphur in a confined quantity of atmospheric air, it absorbs the oxygen contained in it; this however, not being sufficient to oxygenate the sulphur completely, the result is an imperfect aeriform acid, or sulphureous acid gas.

3. Sulphureous acid gas may also be obtained by decomposing sulphite of potash or soda, either by heat alone, or by the affusion of acids.

4. If charcoal be moistened with sulphuric acid, and then exposed to distillation, the products are carbonic acid gas and sulphureous acid gas.

During this operation the charcoal attracts part of the oxygen of the sulphuric acid, and forms carbonic acid gas. The sulphuric acid is therefore partly de-oxidated and becomes converted into sulphureous acid gas.

5. Sulphureous acid gas is likewise formed by distilling sulphur with the oxida of mercury, lead, tin, manganese, &c.

In these processes the sulphur attracts the oxygen of the metallic oxids, and becomes converted into sulphureous acid gas, while the oxids are partially restored to the metallic state.

Water impregnated with sulphureous acid gas forms

SULPHUREOUS ACID.

1. To prepare sulphureous acid, take one part of mercury and four of concentrated sulphuric acid, put them into a retort furnished with a receiver, and connected with Burkitt's or Pepys's apparatus. The sulphureous acid gas which is disengaged is absorbed by the water in the vessel, and constitutes sulphureous acid.

2. Sulphureous acid is likewise formed during the slow combustion of sulphur.

The salts formed by the combination of sulphureous acid with different bases, are called sulphites.

SULPHURETUM AMMONIÆ. *Hepar sulphuris volatile.* Boyle's or Beguine's fuming spirit. Sulphuret of ammonia is obtained in the form of a yellow fuming liquor, by the ammonia and sulphur uniting whilst in a state of gas during distillation. It excites the action of the absorbent system and diminish arterial action, and is given internally in diseases arising from the use of mercury, phthisis, diseases of the skin, and phlegmasiæ: externally it is prescribed in the form of bath in paralysis, contractura, psora, and cutaneous diseases.

SULPHURETUM CALCIS. *Hepar calcis.* Sulphuret of lime. It is principally used as a bath in various diseases of the skin.

SULPHURETUM HYDRARGYRI NIGRUM. See *Hydrargyrus cum sulphure.*

SULPHURETUM HYDRARGYRI RUBRUM. *Cinnabaris antimonii.* Red sulphuret of mercury. *Hydrargyrus sulphureus ruber.* "Take of purified mercury, by weight forty ounces; sublimed sulphur, eight ounces." Having melted the sulphur over the fire, mix in the mercury, and as soon as the mass begins to swell, remove the vessel from the fire, and cover it with considerable force to prevent inflammation; then rub the mass into powder, and sublime. This preparation is esteemed a mild mercurial alterative, and given to children in small doses. In surgery its chief and almost only use is in the administration of quicksilver by fumigation. Thus employed it has proved extremely serviceable in venereal cases. Ulcers and excrescences about the pudendum and anus in women, are particularly benefited by it; and in these cases it is most conveniently applied by placing a red hot heater at the bottom of a night stool-pan, and after sprinkling on it a few grains of the red sulphuret quicksilver, placing the patient on the stool. To fumigate ulcers in the throat, it is necessary to receive the fumes on the part affected, through the tube of a funnel. By enclosing the patient naked in a box, it has on some occasions been contrived to fumigate the whole body at once, and in this way the specific powers of the quicksilver have been very rapidly excited.

This mode of curing the lues venerea, is spoken of as true; and the subject has of late years been revived in a treatise by Sabonette, and by trials made in Bartholomew's hospital.

Mr. Pearson, from his experiments on mercurial fumigation, concludes, that where checking the progress of the disease suddenly is an object of great moment, and where the body is covered with ulcers or large and numerous eruptions, and in general to ulcers, fungi, and excrescences, the vapour of mercury is an application of great efficacy and utility; but that it is apt to induce a ptyalism rapidly, and great consequent debility, and that for the purpose of securing the constitution against a relapse, as great a quantity of mercury must be introduced into the system, by inunction, as if no fumigation had been employed.

SULPHURETUM POTASSÆ. *Kali sulphuratum. Hepar sulphuris.* Sulphuret of potash. Liver of sulphur. The virtues and uses of this preparation are similar to those of the *sulphuretum ammoniac.*

SULPHURETUM SODÆ. A combination of soda and sulphur.

SULPHURETUM STIBII NATIVUM. *Sulphuretum stibii nigrum Antimonium crudum.* Native sulphuret of antimony. It is from this ore that all our preparations of antimony are made. See *Antimony.*

SULPHURIC ACID. *Acidum sulphuricum.* Oil of vitriol. Vitriolic acid. Sulphuric acid consists of sulphur, which constitutes its basis, and of oxygen. It frequently occurs in combination with alkalis, earths, and metals; but seldom in a state of purity, and whether at all, has been doubted. For chemical, medical, and other purposes, it is obtained by the rapid combustion of sulphur and the decomposition of metallic and earthy sulphuric acid salts by fire. If melted sulphur be exposed in open air to an increased heat, it takes fire, is entirely consumed, and burns with a flame, and an acid suffocating vapour. The sulphur is by this combustion changed to an acid. If the heat by which burning sulphur is consumed be only weak, its flame is blue, and the acid then generated is imperfect, very volatile, and æriiform. This imperfect sulphuric acid is the *sulphureous acid.* But if the heat be stronger, the flame of the sulphur is white and lively, and a perfect *sulphuric acid* in the vaporous state is formed. The sulphuric acid may be obtained by different methods; that prepared in England is by the combustion of sulphur; for which purpose peculiar apartments, with their insides lined with lead, are constructed, in order to enclose the burning sulphur. But because the sulphur would in that situation be soon extinguished, and never burn with due vivacity, about the eighth part of its weight of nitre is mingled with it to supply vital air, without which no combustion can exist. Both the water at the bottom of the chamber, and also the aqueous vapours conveyed into it, imbibe the sulphuric acid as it forms. The weak or diluted acid thus obtained is collected and concentrated by evaporating the superfluous water in glass vessels. Sulphuric acid, concentrated in this manner, is sold under the name of *English oil of vitriol, oleum vitrioli*, to distinguish it from another species, called *Saxon oil of vitriol*, which is obtained from green vitriol by distillation. Every kind of concentrated sulphuric acid, formerly found in commerce, was prepared by distillation from green vitriol, or copper; and it is on this account that the sulphuric acid is generally called *vitriolic acid.* Oil of vitriol is a very strong acid, burns and corrodes the skin. When pure, it has neither colour nor smell, but is very apt to turn more or less brown, and of a sulphureous odour, by combustible, animal, and vegetable matter, as is usually the case with what is sold in the shops. If sulphuric acid be highly concentrated, it congeals at a temperature of 150°, and becomes a

crystalline mass, formerly termed glacial oil of vitriol. Mixed with water it becomes hot, and is then termed spirit of vitriol. Sulphuric acid is a powerful antiseptic and tonic: it is given, properly diluted, in the dose of from one to three drops with cinchona and other medicines in the cure of fevers and debilities, and it is often applied externally, when very much diluted, against psora and some chronic affections of the skin.

SULPHURIS FLORES. See *Sulphur sublimatum*.

SULPHUROUS ACID. See *Sulphureous acid gas*.

Sulphur, washed. See *Sulphur lotum*.

Sulphur, precipitated. See *Sulphur precipitatum*.

Sultan flower. The *Centaurea moschata* of Linnaeus.

SUMACH. (*Sumak*, from *samak*, to be red; so called from its red berry.) Elm-leaved sumach. This plant, *Rhus coriaria* of Linnaeus:—*foliis pinnatis obtusis, caule serratis corollibus subtus villosis*, is a small tree, a native of the south of Europe. It is singular that this is the only species of the genus *rhhus* which is perfectly innocent; the others being active poisons. Both the leaves and berries of this plant are used medicinally, as astringents and tonics; the former are the most powerful, and have been long in common use, where they may be easily obtained in various complaints indicating this class of remedies. The berries, which are red, and of a roundish compressed figure, contain a pulpy matter, in which is lodged a brown, hard, oval seed, manifesting a considerable degree of adstringency. The pulp, even when dry, is grateful, and has been discovered to contain an essential salt, similar to that of wood sorrel. An infusion of the dried fruit is not rendered black by a solution of iron: hence it appears to be destitute of adstringency. But its acidity is extremely grateful; therefore, like many other fruits, these berries may be advantageously taken to allay febrile heat, and to correct bilious putrescency.

Sumach, elm leaved. See *Sumach*.

SUMEN. (Arab.) The lower or fat part of the belly.

Sun-dew. See *Ros solis*.

SUPERBUS. See *Rectus superior oculi*.

SUPERCILUM See *Eye-brow*.

SUPERCILIUM VENERIS. The milfoil or yarrow was once so termed. See *Millefolium*.

SUPERFETATION. (*Superfatio*, from *super*, above or upon, and *fatus*, a foetus.) The impregnation of a woman already pregnant.

SUPERGENITALIS. (From *super*, above, and *genita*, the testicles.) The epididymis, or body above the testicles.

SUPERGENITALIS. (From *super*, above,

and *genu*, the knee.) The patella, or kneecap.

SUPERIMPREGNATIO. (From *super*, above, and *impregnatio*, a conception.) Superfétation.

SUPERIOR AURIS. See *Attollens aurem*.

SUPERLIGULA. (From *super*, above, and *ligula*, a little tongue, the glottis.) The epiglottis.

SUPERFUSATIO. (From *super*, beyond, and *purge*, to purge.) An excessive evacuation by stool.

SUPERSCAPULARIS. (From *super*, upon, and *scapula*, the shoulder-blade.) A muscle seated upon the scapula.

SUPINATION. (*Supinatio*, from *supinus*, placed upward.) The act of turning the palm of the hand upwards, by rotating the radius upon the ulna.

SUPINATOR. (From *supinus*, upwards.) A name given to those muscles which turn the hand upwards.

SUPINATOR BREVIS. See *Supinator radii brevis*.

SUPINATOR LONGUS. See *Supinator radii longus*.

SUPINATOR RADII BREVIS. *Supinator brevis*, *sive minor*, of Winslow, and *epicondyle-radial*, of Dumas. This small muscle, which is tendinous externally, is situated at the upper part of the forearm under the supinator longus, the extensor carpi radialis brevis, the extensor carpi ulnaris, the extensor digitorum communis, and the extensor minimi digiti.

It arises tendinous from the lower and anterior part of the outer condyle of the os humeri, and tendinous and fleshy from the outer edge and posterior surface of the ulna, adhering firmly to the ligament that joins the radius to that bone. From these origins its fibres descend forwards and inwards, and are inserted into the upper, inner, and anterior part of the radius around the cartilaginous surface upon which slides the tendon of the biceps, and likewise into a ridge that runs downwards and outwards below this surface.

This muscle assists in the supination of the hand by rolling the radius outwards.

SUPINATOR RADII LONGUS. *Supinator longus* of Albinus. *Supinator longus sive major* of Winslow, and *humero-radial* of Dumas. A long flat muscle, covered by a very thin tendinous fascia, and situated immediately under the integuments along the outer convex surface of the radius. It arises by very short tendinous fibres, from the anterior surface and outer ridge of the os humeri, about two or three inches above its external condyle, between the brachialis internus and the triceps brachii; and likewise from the anterior surface of the external intermuscular membrane, or ligament, as it is called. About the middle of the radius, its fleshy fibres terminate in a flat tendon, which is

inserted into the inner side of the inferior extremity of the radius, near the root of its styloid process.

This muscle not only assists in rolling the radius outwards, and turning the palm of the hand upwards, on which account Riolanus first gave it the name of *supinator*, but it likewise assists in pronation, and in bending the fore-arm.

SUPPOSITORIUM. (From *sub*, under, and *pono*, to put.) A suppository, i. e. a substance to put into the rectum, there to remain and dissolve gradually.

Suppressed menses. See *Amenorrhœa*.

SUPPURATION. (*Suppuratio*, from *suppurare*, to suppurate.) That morbid action by which pus is deposited in inflammatory tumours. See *Pus*.

SUPRA COSTALES. A portion of the intercostal muscles. See *Intercostal muscles*.

SUPRA SPINATUS. (*Musculus supra-spinatus*.) *Supra spinatus seu supercapularis* of Cowper, and *sous-spiro-scapulo trochiterien* of Dumas. This muscle, which was first so named by Riolanus, from its situation, is of considerable thickness, wider behind than before, and fills the whole of the cavity or fossa that is above the spine of the scapula. It arises fleshy from the whole of the base of the scapula that is above its spine, and likewise from the spine itself, and from the superior costa. Opposite to the basis of the coracoid process, it is found beginning to degenerate into a tendon, which is at first covered by fleshy fibres, and then passing under the acromion, adheres to the capsular ligament of the os humeri, and is inserted into the upper part of the large tuberosity at the head of the os humeri. This muscle is covered by a thin fascia, which adheres to the upper edge and superior part of the basis, as well as to the upper edge of the spine of the scapula. The principal use of the supra spinatus seems to be to assist in raising the arm upwards; at the same time, by drawing the capsular ligament upwards, it prevents it from being pinched between the head of the os humeri and that of the scapula. It may likewise serve to move the scapula upon the humerus.

SURA. (Arab.) The calf of the leg; the fibula.

SURFEIT. The consequence of excess in eating or drinking, or of something unwholesome or improper in these articles. It consists in a heavy load or oppression of the stomach, with nausea, sickness, impeded perspiration, and at times eruptions on the skin.

SURGERY. (*Chirurgia*, from *χρῆς*, the hand, and *εργον*, labour.) A branch of the science of medicine, having for its object the cure of external diseases.

SUS SCROFA. The systematic name of the hog which affords the lard.

Suspended animation. See *Resuscitation*.

SUSPENSORIUM. (From *suspendere*, to hang) A suspensor; a bag, or bandage, to suspend the scrotum.

SUSPENSORIUM HEPATIS. The broad ligament of the liver.

SUSPENSORIUS TESTIS. The cremaster muscle of the testicle.

SUSURRUS. (From *susurre*, to murmur.) An imaginary sound in the ear.

SUTURE. (*Satura*, from *suo*, to join together.) Called also *clavata commissura*. 1. In surgery this term signifies the uniting the lips of a wound by sewing. A number of different kinds of sutures have been recommended by writers on surgery, but all of them are now reduced to two: namely, the *twisted*, and the *interrupted*, called also the *knotted suture*. The twisted suture is made in the following manner: having brought the divided parts nearly into contact, a pin is to be introduced from the outside inwards, and carried out through the opposite side to the same distance from the edge that it entered at on the former side; a firm wax ligature is then to be passed around it, making the figure of 8, by which the wounded parts are drawn gently into contact. The number of pins is to be determined by the extent of the wound; half an inch, or at most three quarters, is the proper distance between two pins. The interrupted suture is practised where a number of stitches is required, and the interruption is only the distance between the stitches.

2. In anatomy the word suture is applied to the union of bones by means of dentiform margins, as in the bones of the cranium. See *Temporal*, *sphenoidal*, *zygomatic*, *transverse*, *coronal*, *lambdoidal*, and *frontal* sutures.

Swallow-wort. See *Vincetoxicum*.

Sweat. See *Perspiration*.

Sweating, *immense.* See *Ephedrosia*.

Sweet margoram. See *Marjorana*.

Sweet navel. See *Rapum*.

Sweet rush. See *Juncus odoratus*.

Sweet sultan. The *Centaurea moschata* of Linnaeus.

Sweet willow. See *Myrtus Brabantica*.

SWIETENTIA. (Named after Swieten.) The bark of the wood of this tree is of a red colour internally; has an adstringent bitter taste; yields its active matter to water. It has been prepared as a substitute for Peruvian bark, and has been used as such with advantage. Dose half a drachm.

SWIETENTIA MAHAGONI. The systematic name of the mahogany tree. See *Mahoezmi* and *Swietenia*.

Swine-pox. See *Varicella*.

SYCOMA. (From *συκκα*, a fig.) A wart or excrescence resembling a fig.

SYCOZIS. The ophthalmia trachoma of Sauvages; also a fungous ulcer; and by others the tumour on the anus called *ma-fisca*.

SYLPHIUM. *Assafetida* is so termed by some writers. See *Assafetida*.

SYMBLEPHARUM. (From *συμ*, with, and *βλεφαρον*, the eyelid.) A concretion of the eyelid to the globe of the eye; also a concretion of the palpebra with the eye. This chiefly happens in the superior, but very rarely in the inferior palpebra. The causes of this concretion are a bad conformation of the parts, or from ulcers of the cornea, the membrana conjunctiva, or internal superficies of the palpebræ, or imprudent scarifications, or burns, especially if the eye remains long closed. There are two species, the partial, or total; in the former the adhesion is partial, in the latter, the membrana conjunctiva and cornea are concreted together.

SYMBOLE. (From *συμβάλλω*, to knit together.) Is said either of the fitness of parts with one another, or of the consent between them by the intermediation of nerves, and the like.

SYMBOLOREM. (From *συμβολον*, a sign, and *λογος*, a discourse.) The doctrine of the signs and symptoms of disease.

SYMMETRY. The exact and beautiful proportion of parts to one another.

Sympathetic nerve. See *Intercostal nerve*.

SYMPATHY. (From *συμπασχω*, to suffer together, to sympathise.) *Sympathia.* All the body is sympathetically connected together, and dependent, the one part upon the rest, constituting a general sympathy. But sometimes we find particular parts more intimately dependent upon each other than upon the rest of the body, constituting a particular sympathy. Action cannot be greatly increased in any one organ, without being diminished in some other; but certain parts are more apt to be affected by the derangement of particular organs than others; and it was the observance of this fact which gave foundation to the old and well-known doctrine of sympathy, which was said to proceed, "*tum ob communionem et similitudinem generis, tum ob viciniam.*" It may be thought that this position of action being diminished in one organ, by its increase, either in the rest or in some other part, is contradicted by the existence of general diseases or actions affecting the whole system. But in them we find, in the first place, that there is always some part more affected than the rest. This local affection is sometimes the first symptom, and affects the constitution in a secondary way, either by the irritation which it produces, or by an extension of the specific action. At other times the local affection is coeval with the general disease, and is called sympathetic. It is observed in the second place, that as there is some part which is always more affected than the rest, so also is there some organ which has its action, in consequence of this, diminished lower than that of the rest of the system, and

most commonly lower than its natural standard. From the extensive sympathy of the stomach with almost every part of the body, we find that this most frequently suffers, and has its action diminished in every disease, whether general or local, provided that the diseased action arises to any considerable degree. There are also other organs which may, in like manner, suffer from their association or connexion with others which become diseased. Thus, for instance, we see, in the general disease called puerperal fever, that the action of the breasts is diminished by the increased inflammatory action of the uterus.

In consequence of this balance of action, or general connexion of the system, a sudden pain, consequent to violent action of any particular part, will so weaken the rest as to produce fainting, and occasionally death. But this dependence appears more evidently in what may be called the smaller systems of the body, or those parts which seem to be more intimately connected with each other than they are with the general system. Of this kind is the connexion of the breasts with the uterus of the female; of the urethra with the testicles of the male; of the stomach with the liver; and of the intestines with the stomach, and of this again with the brain; of the one extremity of the bone with the other; and of the body of the muscle with its insertion; of the skin with the parts below it.

These smaller systems, or circles, shall be treated regularly; but, first, it may be proper to observe, that these are not only intimately connected with themselves, but also with the general system, an universal sympathy being thus established.

That there is a very intimate connexion between the breast and uterus has been long known; but it has not been very satisfactorily explained. Fallopius, and all the older authors, declare plainly that the sympathy is produced by an anastomosis of vessels; Bartholin adding, that the child being born, the blood no longer goes to the uterus, but is directed to the breast and changed into milk. But none of all those who talk of this derivation, assign any reasonable cause which may produce it.

In pregnancy, and at the menstrual periods, the uterus is active; but, when the child is delivered, the action of the uterus subsides, whilst the breasts in their turn become active, and secrete milk.

If, at this time, we should again produce action in the uterus, we diminish that of the breasts, and destroy the secretion of milk, as is well illustrated by the case of inflammation of the uterus, which is incident to lying-in women. When the uterus at the cessation of the menses, ceases to be active, or to secrete, we often find that the breasts have an action excited in them, becoming slowly inflamed, and assuming

a cancerous disposition. The uterus and breasts seem to be a set of glands balancing each other in the system, one only being naturally active, or secreting properly at a time; and accordingly we seldom, if ever, find that when the uterus yields the menstrual discharge, the milk is secreted in perfection, during the continuance of this discharge, nor do we ever find them both inflamed at the same time.

The uterus has not only this connexion with the breasts, but it has also a very particular sympathy with the stomach, which again sympathises with the brain; and thus we see how a disorder of the uterus may induce an extensive series of affections, each dependent on the other.

The organs of generation in the male form likewise a little system, in which all the parts exhibit this sympathy with each other. They likewise give us a very good instance of the association of action, or sympathy, in the common acceptance of that word.

Sympathy is divided into, first, the sympathy of equilibrium, in which one part is weakened by the increased action of another; and, secondly, the sympathy of association, in which two parts act together at the same time.

The sympathy of association is produced suddenly, and for a short time. The sympathy of equilibrium is produced more slowly, and continues to operate for much longer time.

It is curious enough that most, or at least many, of those organs, which seem to be connected by the sympathy of equilibrium, exhibit likewise more or less of the sympathy of association, when under the circumstances in which this can take place.

The sympathy of equilibrium is seen in the effects of inflammation of the end of the urethra on the testicle; which often diminishes its action, and produces a very disagreeable sensation of dullness, or, if this inflammation be suddenly diminished, the action of the testicle is as suddenly increased, and swelling takes place. The same is seen in the connexion of the urethra with the bladder and prostate gland, as is mentioned in all the dissertations on gonorrhœa. These parts likewise affect the stomach greatly, increased action in them weakening that organ much. This is seen in the effects of swelled testicle, or excessive venery, or inflamed bladder, and in a stone; all which weaken the stomach, and produce dyspepsia. The same remark applies to the kidney; vomiting and flatulence being produced by nephritis.

The sympathy of association, or an instance of sympathy in the common acceptance of the word, is likewise seen in the connexion betwixt the glands and testicles in coition; but for this purpose, the ac-

tion in the glands must be sudden and of short duration; for, if continued long, weakness of the testicles, or diminished action, is induced. In those parts which exhibit this natural association of action, if the action of one part be suddenly and for a short time increased, the action of the sympathising part will likewise be increased; as we see in the instance already given of coition, and likewise in paroxysms of the stone, in which the glans penis, after making water, becomes very painful.

But if the action be more slowly induced, and continued for a long time, then this association is set aside, by the stronger and more general principle of the equilibrium of action, and the sympathising part is weakened. Hence, violent inflammation of the end of the urethra produces a weakness and irritability of the bladder, dullness of the testicle, &c.

There is also an evident sympathy of equilibrium betwixt the stomach and lower tract of intestines; which two portions may be said in general to balance each other in the abdomen. When the action of the intestines is increased in diarrhœa, the stomach is often weakened, and the patient tormented with nausea. This will be cured, not so easily by medicines taken into the stomach, as by anodyne clysters, which will abate the action of the intestines. When the intestines are inflamed, as in strangulated hernia, vomiting is a never-failing attendant.

When again the stomach is inflamed, the intestines are affected, and obstinate constiveness takes place; even in hysterical affections of the stomach, the intestines are often deranged. Injections of cold water frequently relieve these affections of the stomach, by their action on the intestines.

The liver and stomach are also connected with one another. When the liver is inflamed, or has its action increased, the stomach is weakened, and dyspeptic symptoms take place. When the stomach is weakened, as, for instance, by intoxication, then the action of the liver is increased, and a greater quantity than usual of bile is secreted. The same takes place in warm climates, where the stomach is much debilitated.

If the liver has its action thus frequently increased, it assumes a species of inflammation, or becomes, as it is called, scorbutous. This is exemplified, in the habitual dram-drinkers, and in those who stay long in warm countries and use freedoms with the stomach. The liver likewise sympathises with the brain; for when this organ is injured, and its action much impaired, as in compression, inflammation and suppuration have been often known to take place in the liver.

Besides this connexion of the stomach with the liver, it is also very intimately de-

pendent on the brain, being weakened when the action of the brain is increased, as we see in inflammation of that organ. The brain again is affected with pain, when the stomach is weakened by intoxication, or other causes; and this pain will be often relieved by slowly renewing the action of the stomach, by such stimuli as are natural to it, such as small quantities of soups frequently repeated. A slight increase of action in the stomach, at least if not of a morbid kind, affects the brain so as to produce sleep, diminishing its action. This we see in the effects of a full meal, and even of a draught of warm water. The stomach likewise sympathises with the throat, squeamishness and anorexia being often produced by inflammation of the tonsils. This inflammation is frequently abated by restoring or increasing the action of the stomach. Hence the throat, in slight inflammation, is frequently easier after dinner; hence, likewise, the effects of emetics in cynanche.

The extremities of bones and muscles also sympathise in the same manner. When one end of a bone is inflamed, the action of the other is lessened, and pain is produced; for a painful sensation may result both from increased and diminished action. When the tendon of a muscle is inflamed, the body of that muscle often is pained, and *vice versa*.

Lastly, the external skin sympathises with the parts below it. If it be inflamed, as in erysipelas, the parts immediately beneath are weakened, or have their natural action diminished. If this inflammation affect the face, or scalp, then the brain is injured; and head-ache, stupor, or delirium, supervene. If it attack the skin of the abdomen, then the abdominal viscera are affected, and we have vomiting and purging, or obstinate costiveness, according to circumstances. This is illustrated by the disease of children, which is called by the women the bowel-hive, in which the skin is inflamed, as they suppose, from some morbid matter within.

If the internal parts be inflamed, the action of the surface is diminished, and, by increasing this action, we can lessen or remove the disease below; as we see daily proved by the good effects of blisters. When the stomach, intestines, or kidney, have been very irritable, a sinapism has been known to act like a charm; and, in the deep seated inflammations of the breasts, bowels, or joints, no better remedy is known, after the use of the lancet, than blisters.

The utility of issues in diseases of the lungs, the liver, and the joints, is to be explained on the same principle. In these cases we find that issues do little good unless they be somewhat painful, or be in the state of healthy ulcers. An indolent flap-

by sore, however large the discharge, (which is always thin, and accompanied with little action,) does no good, but only adds to the misery of the patient. We may, however, err on the other hand by making the issues too painful, or by keeping them active too long, for, after they have removed the inflammatory disease below, they will still operate on these parts, lessening their action, and preventing the healing process from going on properly. This is seen in cases of curvature of the spine, where at first the inflammation of the vertebra is diminished by the issues; but, if they be kept long open after this is removed, they do harm. We often see the patient recover rapidly after his surgeon has healed the issue in despair, judging that it could do no farther service, but only increase the weakness of his patient.

It is a well-established fact, that when any particular action disappears suddenly from a part, it will often speedily affect that organ which sympathises most with the part that was originally diseased. This is best seen in the inflammatory action, which, as practical writers have well observed, occasionally disappears quickly from the part first affected, and then shews itself in some other.

From the united testimony of all these facts, Mr. Burns, of Glasgow, maintains the doctrine just delivered, and proposes to introduce it into pathological reasonings. In the whole of the animal economy, we discover marks of the wisdom of the Creator, but perhaps in no part of it more than in this, of the existence of the sympathy of equilibrium, for if a large part of the system were to have its action much increased, and all the other parts to continue acting in the same proportionate degree as formerly, the whole must be soon exhausted; (for increased action would require for its support an increased quantity of energy.)

But upon this principle, when action is much increased in one part, it is to a certain degree diminished in some other, the general sum or degree of action in the body is thus less than it otherwise would be, and consequently the system suffers less.

SYMPHISIS. (From *συμ*, together, and *φω*, to grow.) Mediate connexion. A genus of the connexion of bones, in which they are united by means of an intervening body. It comprehends four species, viz. synchondrosis, syssarcosis, synneurosis, and syndenmosis.

SYMPHYTUM. (From *συμφυω*, to unite; so called because it is supposed to unite and close the lips of wounds together.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*, Order, *Monogynia*.

2. The pharmacopœial name of the *consolida major*. Comfrey. This plant, *Symphytum officinale*:—*foliis ovatis lanceolatis decurrentibus*, is administered where the althiza cannot be obtained, its roots abounding with a viscid glutinous juice whose virtues are similar to that of the *althiza*.

SYMPHYTUM MACULOSUM. See *Pulmonaria*.

SYMPHYTUM MINUS. See *Prunella*.

SYMPHYTUM OFFICINALE. The systematic name of the comfrey. See *Symphytum*.

SYNACHE. See *Cynanche*.

SYNACHICA. (From *συαχην*, the quince; so called from its uses in that disease.) Quince-wort.

SYNARTHIROSIS. (From *συν*, together, and *αρθρον*, a joint.) Immoveable connexion. A genus of connexion of bones, in which they are united together by an immoveable union. It has three species, viz. suture, harmony, and gomphosis.

SYNASTOMOSIS. Is used in the same sense as *Anastomosis*.

SYNCHONDROSIS. (From *συν*, with, and *χονδρος*, a cartilage.) A species of symphysis, in which one bone is united with another by means of an intervening cartilage, as the vertebrae and the bones of the pubis.

SYNCHONDROTOMIA. (From *συνχονδρῶσις*, the symphysis of the pubis, and *τομή*, to cut.) The operation of dividing the symphysis of the pubis.

SYNCHYSIS. (From *συψυω*, to confound.) A solution of the vitreous humour into a fine attenuated aqueous fluid. In Cullen's Nosology it is a variety of his species *caligo pupillæ*.

SYNCIPITIS OSSA. See *Parietal bones*.

SYNCIPUT. The forepart of the cranium.

SYNCOPE. (From *συν*, with, and *κτείνω*, to cut, or strike down.) *Animi deliquium*. *Leipothymia*. Fainting or swooning.

A genus of disease in the class *neuroses* and order *adynamie* of Cullen, in which the respiration and action of the heart either cease, or become much weaker than usual, with paleness and coldness, arising from diminished energy of the brain, or from organic affections of the heart. Species: 1. *Syncope cardiaca*, the cardiac syncope, arising without a visible cause, and with violent palpitation of the heart, during the intervals, and depending generally on some organic affection of the heart or neighbouring vessels. 2. *Syncope occasionalis*, the existing cause being manifested.

SYNCOPE ANGINOSA. See *Angina pectoris*.

SYNDESMOLOGIA. (From *συνδεσμος*, a ligament, and *λογος*, a discourse.) The doctrine of the ligaments.

SYNDESMO-PHARYNGEUS. See *Constrictor pharyngis medius*.

SYNDESMOSIS. (From *συνδεσμος*, a ligament.) That species of symphysis or mediate connexion of bones in which they are united by ligament, as the radius with the ulna.

SYNDESMUS. (From *συνδεω*, to bind together.) A ligament.

SYNECHIA. *Συναχία*. A concretion of the iris with the cornea, or with the capsule of the crystalline lens. The proximate cause is the contact from inflammation or adhesion of these parts, the consequence of inflammation. The remote causes are, a collapse of the cornea, a prolapse of the iris, a swelling or tumified cataract, hypopium, or a natural formation. The species of this disorder are:

1. *Synechia anterior totalis*, or a concretion of the iris with the cornea. This species is known by inspecting the parts. The pupil in this species is dilated or coarctated, or it is found concreted; from whence various lesions of vision.

2. *Synechia anterior partialis*, when only some part of the iris is accreted. This concretion is observed in one or many places; from hence the pupil is variously disfigured, and an inordinate motion of the pupil is perceived.

3. *Synechia anterior composita*, when not only the whole iris, but also a prolapse of the crystalline lens, unites with the cornea.

4. *Synechia posterior totalis*, or a concretion of the whole uvea, with the ciliary processes and the capsule of the crystalline lens.

5. *Synechia posterior partialis*, when only some part of the capsule of the crystalline lens is concreted with the uvea and cornea. This accretion is simplex, duplex, triplex, or in many places it may happen.

6. *Synechia complicata*, with an amaurosis, cataract, mydriasis, myosis, or synchysis.

SYNEUROSIS. (From *συν*, with, and *νευρον*, a nerve, because the ancients included membranes, ligaments, and tendons, under the head of nerves.) A species of symphysis, in which one bone is united to another by means of an intervening membrane.

SYNZESIS. *Confidentia*. A perfect concretion and coarctation of the pupil. It is known by the absence of the pupil, and a total loss of vision. The species are:

1. *Synzesis nativa*, with which infants are sometimes born. In this case, by an error of the first conformation of the pupil, there is no perforation; it is very rarely found.

2. *Synzesis accidentalis*, a concretion of

the pupil, from an inflammation or ex-
ulceration, of the uvea or iris, or from
a defect of the aqueous or vitreous hu-
mour.

3. *Synizesis*, from a recession of the iris
or cornea. From whatever cause it may
happen, the effect is certain, for the pupil
contracts its diameter; the longitudinal
fibres, separated from the circle of the
cornea, cannot resist the orbicular fibres:
from hence the pupil is wholly or partially
contracted.

4. *Synizesis complicata*, or that which is
complicated with an amaurosis, synechia,
or other ocular disease. The amaurosis or
gutta serena is known by the total absence
of light to the retina; we can distinguish
this not only by the pupil being closed,
but likewise the eyelids, for whether the
eyelids be open or shut, all is darkness to
the patient. The other complicated cases
are known by viewing the eye, and consid-
ering the parts anatomically.

5. *Synizesis spuria*, is a closing of the
pupil by mucus, pus, or grumous blood.

SYNOCHA. (From *συναχα*, to conti-
nue.) Inflammatory fever. A species of
continued fever; characterised by increas-
ed heat; pulse frequent, strong, hard;
urine high-coloured; senses not im-
paired. This fever is so named from its
being attended with symptoms denoting
general inflammation in the system, by
which we shall always be able readily to
distinguish it from either the nervous or
putrid. It makes its attack at all seasons
of the year, but is most prevalent in the
spring; and it seizes persons of all ages
and habits, but more particularly those in
the vigour of life, with strong elastic fibres,
and of a plethoric constitution. It is a
species of fever almost peculiar to cold and
temperate climates, being rarely, if ever
met with in very warm ones, except
amongst Europeans lately arrived; and
even then, the inflammatory stage is of
very short duration, as it very soon as-
sumes either the nervous or putrid type.

The exciting causes are sudden transi-
tions from heat to cold, swallowing cold
liquors when the body is much heated by
exercise, too free a use of vinous and spi-
rituous liquors, great intemperance, violent
passions of the mind, the sudden suppres-
sion of habitual evacuations, and the sud-
den repulsion of eruptions. It may be
doubted if this fever ever originates from
personal infection; but it is impossible
for it to appear as an epidemic amongst
such as are of a robust habit, from a pe-
culiar state of the atmosphere. It comes
on with a sense of lassitude and inactivity,
succeeded by vertigo, rigors, and pains
over the whole body, but more particular-
ly in the head and back; which symptoms
are shortly followed by redness of the face
and eyes, great restlessness, intense heat,

and unquenchable thirst, oppression of
breathing, and nausea. The skin is dry
and parched; the tongue is of a scarlet
colour at the sides, and furred with white
in the centre; the urine is red and scanty;
the body is costive; and there is a quick-
ness, with a fulness and hardness in the
pulse, not much affected by any pressure
made on the artery. If the febrile symp-
toms run very high, and proper means
are not used at an early period, stupor and
delirium come on, the imagination be-
comes much disturbed and hurried, and the
patient raves violently. The disease usual-
ly goes through its course in about fourteen
days, and terminates in a crisis, either by
diaphoresis, diarrhoea, hæmorrhage from
the nose, or the deposit of a copious sedi-
ment in the urine; which crisis is usually
preceded by some variation in the pulse.

Our judgment as to the termination of
the disease, must be formed from the vio-
lence of the attack, and the nature of the
symptoms. If the fever runs high, or
continues many days with stupor or deli-
rium, the event may be doubtful; but if
to these are added, picking at the bed-
clothes, startings of the tendons, involun-
tary discharges by stool and urine, and
hiccough, it will then certainly be fatal. On
the contrary, if the febrile heat abates,
the other symptoms moderate, and there
is a tendency to a crisis, we may then ex-
pect a recovery. In a few instances, this
fever has been known to terminate in
mania.

On opening those who die of an inflam-
matory fever, an effusion is often perceived
within the cranium, and now and then, to-
pical affections of some of the viscera are
to be observed.

SYNOCHUS. (From *συναχα*, to conti-
nue.) A mixed fever. A species of
continued fever, commencing with symp-
toms of synocha and terminating in ty-
phus; so that synocha and typhus, blended
together in a slight degree, seem to consti-
tute this species of fever, the former be-
ing apt to preponderate at its commence-
ment, and the latter towards its termina-
tion.

Every thing which has a tendency to
enervate the body, may be looked upon as
a remote cause of this fever; and accord-
ingly we find it often arising from great
bodily fatigue, too great an indulgence in
sensual pleasures, violent exertion, intem-
perance in drinking, and errors in diet, and
now and then likewise from the suppres-
sion of some long accustomed discharge.
Certain passions of the mind (such as grief,
fear, anxiety, and joy,) have been enu-
merated amongst the causes of fever, and in
a few instances it is probable they may
have given rise to it; but the concurrence
of some other powers seems generally ne-
cessary to produce this effect. The most

usual and universal cause of this fever is the application of cold to the body; and its morbid effects seem to depend partly upon certain circumstances of the cold itself, and partly upon certain circumstances of the person to whom it is applied.

The circumstances which seem to give the application of cold due effect, are its degree of intensity, the length of time which it is applied, its being applied generally, or only in a current of air, its having a degree of moisture accompanying it, and its being a considerable or sudden change from heat to cold. The circumstances of persons rendering them more liable to be affected by cold, seem to be debility, induced either by great fatigue, or violent exertions, by long fasting, by the want of natural rest, by severe evacuations, by preceding disease, by errors in diet, by intemperance in drinking, by great sensuality, by too close an application to study, or giving way to grief, fear, or great anxiety, by depriving the body of a part of its accustomed clothing, by exposing any one particular part of it, whilst the rest is kept of its usual warmth, or by exposing it generally or suddenly to cold when heated much beyond its usual temperature; these we may therefore look upon as so many causes giving an effect to cold which it otherwise might not have produced. Another frequent cause of fever seems to be breathing air contaminated by the vapours arising either directly or originally from the body of a person labouring under the disease. A peculiar matter is supposed to generate in the body of a person affected with fever, and this floating in the atmosphere, and being applied to one in health, will no doubt often cause fever to take place in him, which has induced many to suppose that this infectious matter is produced in all fevers whatever, and that they are all more or less contagious.

The effluvia arising from the human body, if long confined to one place without being diffused in the atmosphere, will, it is well known, acquire a singular virulence, and will if applied to the bodies of men, become the cause of fever. Exhalations, arising from animal or vegetable substances in a state of putrefaction, have been looked upon as another general cause of fever: marshy or moist grounds, acted upon by heat for any length of time, usually send forth exhalations which prove a never-failing source of fever, but more particularly in warm climates. Various hypotheses have been maintained, with respect to the proximate cause of fever; some supposing it to be a lentor or viscosity prevailing in the mass of blood, and stagnating in the extreme vessels; others, that it is a noxious matter introduced into or generated in the body, and that the in-

creased action of the heart and arteries is the effect of nature to expel the morbid matter; others, that it consisted in an increased secretion of bile; and others again, that it is to be attributed to a spasmodic constriction of the extreme vessels on the surface of the body; which last was the doctrine taught by the late Dr. Cullen.

An attack of this fever is generally marked by the patient's being seized with a considerable degree of languor or sense of debility, together with a sluggishness in motion, and frequent yawning and stretching; the face and extremities at the same time become pale, and the skin over the whole surface of the body appears constricted; he then perceives a sensation of cold in his back, passing from thence over his whole frame, and this sense of cold continuing to increase; tremors in the limbs and rigors of the body succeed.

With these there is a loss of appetite, want of taste in the mouth, slight pains in the head, back, and loins, small and frequent respirations. The sense of cold and its effects after a little time become less violent, and are alternated with flushings, and at last, going off altogether, they are succeeded by great heat diffused generally over the whole body; the face looks flushed, the skin is dry as likewise the tongue; universal restlessness prevails, with a violent pain in the head, oppression at the chest, sickness at the stomach, and an inclination to vomit. There is likewise a great thirst and costiveness, and the pulse is full and frequent, beating perhaps 90 or 100 strokes in a minute. When the symptoms run very high, and there is a considerable determination of blood to the head, a delirium will arise. In this fever, as well as most others, there is generally an increase of symptoms towards evening.

If the disease is likely to prove fatal, either by its continuing a long time, or by the severity of its symptoms, then a starting of the tendons, picking at the bed-clothes, involuntary discharges by urine and stool, coldness of the extremities, and hiccoughs, will be observed; where no such appearances take place, the disease will go through its course.

As a fever once produced will go on, although its cause be entirely removed, and as the continued or fresh application of a cause of fever neither will increase that which is already produced, nor occasion a new one, there can be no certainty as to the duration of fever, and it is only by attending to certain appearances or changes, which usually take place on the approach of a crisis, that we can form any opinion or decision. The symptoms pointing out the approach of a crisis, are the pulse becoming soft, moderate, and near,

and ulcers of the skin, pains, swelling, and its natural speed; the tongue losing its fur and becoming clean, with an abatement of thirst; the skin being covered with a gentle moisture, and feeling soft to the touch; the secretory organs performing their several offices; and the urine depositing flaky crystals of a dirty red colour, and becoming turbid on being allowed to stand any time.

Many physicians have been of opinion, that there is something in the nature of all acute diseases, except those of a putrid kind, which usually determines them to be of a certain duration, and therefore that these terminations, when salutary, happen at certain periods of the disease rather than at others, unless disturbed in their progress by an improper mode of treatment, or the arising of some accidental circumstance. These periods are known by the spellation of critical days; and from the time of Hippocrates down to the present, have been pretty generally admitted. The truth of them, Dr. Thomas thinks can hardly be disputed, however they may be interrupted by various causes. A great number of phenomena shew us, that both in the sound state and the diseased, nature has a tendency to observe certain periods; for instance, the vicissitudes of sleeping and watching occurring with such regularity to every one; the acute periods that the menstrual flux observes, and the exact time of pregnancy in all viviparous animals, and many other such instances that might be adduced, all prove this law.

With respect to diseases, every one must have observed the definite periods which take place in regular intermitents, as well those universal as topical, in the course of true inflammation, which at the fourth, or at the farthest the seventh day, is resolved, or after this period, changes into either abscess, gangrene, or schirrus, in exanthematous eruptions, which if they are favourable and regular, appear on a certain and definite day; for example, the small-pox about the fourth day. All these appear to be founded on immutable laws, according to which the motions of the body in health and in disease are governed.

The days on which it is supposed the termination of continued fevers principally happens, are the third, fifth, seventh, ninth, eleventh, fourteenth, seventeenth, and twentieth.

A simple continued fever terminates always by a regular crisis in the manner before mentioned, or from the febrile matter falling on some particular parts, it excites inflammation, abscess, eruption, or destroys the patient.

Great anxiety, loss of strength, intense heat, stupor, delirium, irregularity in the pulse, twitchings in the fingers and hands,

picking at the bed-clothes, startings of the tendons, hiccoughs, involuntary evacuations by urine and stool, and such like symptoms, point out the certain approach of death.

On the contrary, when the senses remain clear and distinct, the febrile heat abates, the skin is soft and moist, the pulse becomes moderate and is regular, and the urine deposits flaky crystals, we may then expect a speedy and happy termination to the disease.

The usual appearances which are to be observed on dissection of those who die of this fever, are an effusion within the cranium, and topical affections, perhaps of some viscera.

SYNOVIA. (A term of no radical meaning, coined by Paracelsus.) *Hydathros. Mucilago.* An unctuous fluid secreted from certain glands in the joint in which it is contained. Its use is to lubricate the cartilaginous surfaces of the articulatory bones, and to facilitate their motions.

SYNOVIAL GLANDS. *Glandule synoviae.* The assemblage of a fatty fimbriated structure within the cavities of some joints.

SYNTENOSIS. (From *σύν*, with, and *τενών*, a tendon.) A species of articulation, where the bones are connected together by tendons.

SYNTESIS. (From *συντελέω*, to dissolve.) A marasmus or colliquative wasting of the body.

SYNTHESIS. (From *συντίθημι*, to compose.) Combination. See *Analysis*.

SYNTHETISMUS. (From *συντίθω*, to concur.) The reduction of a fracture.

SYNOLOTTICA. (From *συνέλω*, to cicatrise.) Medicines which heal wounds.

SYPHYLIS. (The name of a shepherd, who fed the flocks of king Alcibious, who, proud of their number and beauty, insulted the sun; as a punishment for which, fable relates, that this disease was sent on earth; or from *σιφλος*, filthy.) *Lues venerea. Morbus Gallicus.* A genus of disease in the class *cachexia* and order *impetiginosae* of Cullen. Towards the close of the memorable fifteenth century, about the year 1494 or 1495 the inhabitants of Europe were greatly alarmed by the sudden appearance of this disease. The novelty of its symptoms, and the wonderful rapidity with which it was propagated throughout every part of the known world, soon made it an important object of medical inquiry.

In the common language, it is said a person has syphilis or is poked, when the venereal poison has been received into, or is diffused through the system, and there produces its peculiar effects, as ulcers of the mouth, fauces, spots, tetters,

earies of the bones, &c. But as long as the effects of the poison are local and confined to or near the genitals, the disorder is not called syphilis, lues venerea, nor pox; but distinguished by some particular name according to its different seat or appearance; such as gonorrhœa venerea, chancre or bubo.

The venereal disease is always produced by a poison. Concerning the nature of this poison, we know no more than we do about that of the small-pox or any other contagion; we know only that it produces peculiar effects. The smallest particle of this poison is sufficient to bring on the most violent disorder over the whole body. It seems to spread and diffuse itself by a kind of fermentation and assimilation of matter; and, like other contagions, it requires some time after being applied to the human body, before it produces that effect. It is not known whether it has different degrees of acrimony and volatility, or whether it is always the same in its nature, varying only with regard to the particular part to which it is applied, or according to the different habit and constitution or particular idiosyncrasy of the person who receives the infection. We know that mercury possesses a certain and specific power of destroying the venereal virus; but we are quite uncertain whether it acts by a sedative, adstringent, or evacuant quality; or if not perhaps rather by a chemical elective attraction whereby both substances uniting with one another are changed in a third, which is no more hurtful, but has some new properties entirely distinct from those which any of them had before they were united. The variolous miasma, we know, produces its effects in about twenty or twenty-four days after the infection is received from the atmosphere, and eight or ten days if by inoculation, but the venereal virus seems to keep no particular period. At some times, and perhaps in particular persons, Dr. Swediaur has seen chancres arise in the space of twelve hours, nay, in a still shorter time, indeed he mentions in a few minutes, after an impure coition; whereas, in most cases, they make their appearance only in as many days. The generality of men feel the first symptoms of a clap between the second and fifth days after an impure coitus; but there are instances where they do not appear till after as many weeks or months. About ten years ago, Dr. S. was consulted by a young man, who was seized with a violent discharge from the glans along with a phymosis, but without any chancres, four weeks after coition; and during all the intervals, he felt not the least symptom of the disease. Some years ago, a gentleman went out from London in seemingly perfect health, to the East

Indies; but on his arrival in that hot climate, after a voyage of four months a violent clap broke out before he went on shore, though he could have received no infection during the voyage, as there was not a woman on board. There are instances which render it probable that the virus may lie four, five, or six weeks, and perhaps longer, on the surface of the genitals before it is absorbed; and were it not then to produce a shanker, might probably not be absorbed at all. We see daily examples, where common women communicate the infection to different men in the space of several weeks, while they themselves have not the least symptom of syphilis local or universal, the poison lying all that time in the vagina harmless, and generally without being absorbed. How long the venereal virus may lurk in the body itself, after it has been absorbed into the mass of blood, before it produces any sensible effect, is a matter of equal uncertainty. There is scarce a practitioner who has not observed instances of its remaining harmless for weeks or even months in the body. Dr. Swediaur had a case, where, after lying dormant for half a year, it broke out with unequivocal symptoms. But the following instance, if to be depended upon, is still more extraordinary.

Some years ago, says the above writer, I was consulted by a gentleman about a sore throat, which I declared to be venereal. My patient was astonished; and assured me that for nine years past he had not had the least venereal complaint, nor had he any reason to believe he had since received any infection: but that he had been in the East Indies, where he was affected with a violent clap. On his return to Europe, being to appearance in good health, he married, and continued perfectly free of any such complaint ever since. By a mercurial course, however, the complaint for which he applied to me was completely removed. With regard to its effects, the venereal poison follows no constant rule: for though, in general, it affects first the throat, where it produces ulcerations, while in others it exerts its virulence on the skin or bones. Whilst the greatest part of mankind are thus easily affected by this poison, there are some few who seem to be altogether unsusceptible of the infection with the variolous contagion, though they go into infected places, and expose themselves to inoculation or every hazard by which the disease is generally communicated.

Some persons are more liable than others to be infected who are seemingly of the same habit; nay, the very same person seems to be more liable to be infected at one time than another, and those who have been once infected seem to be more liable

to catch the infection a second time, than those who never were infected before with the disease. The climate, season, age, state of health, idiosyncrasy, are, perhaps, as in other diseases, the necessary predisposing causes. The same difference is observable in the progress made by the disease after the patient is infected. In some the progress is slow, and the disease appears scarce to gain any ground, while in others it advances with the utmost rapidity, and speedily produces the most terrible symptoms. Whether the venereal poison can be absorbed into the system, without a previous excoriation; or ulceration of the genitals, or some other parts of the surface of the body, is still a matter of doubt. Several cases, however, have occurred which render it highly probable, if not certain, that the poison really is now and then absorbed, without any previous excoriation or ulceration whatsoever, and thus produces buboes and other venereal symptoms in the body.

It has been asserted by the earliest and even by some late writers, that it may be caught by lying in the same bed or living in the same room with or after an infected person. What may have been the case at the commencement of the disease, cannot be said, but the most accurate observations and experiments which have been made upon the subject, do not confirm this to be the case in our times. Nor are nurses infected in the Lock Hospital, where they live night and day with patients in all stages of the distemper. The fact seems to be, that patients in our times are apt to impose upon themselves or upon physicians and surgeons, with regard to this matter; and the above opinion easily gains ground among the vulgar, especially in countries where people are more influenced by prejudices, superstition, servile situation in life, or other circumstances. Hence, we sometimes hear the most ridiculous accounts given in those countries by friars and common soldiers, of the manner by which they came to this disorder; such as piles, gravel, colics, contusions, fevers, little-houses, lying in suspected beds, or lying in bed with a suspected person, retention of the semen, coition with a woman in menstruation, the use of cider, bad wine or beer, &c.

Another question undecided is, whether the venereal poison ever infects any fluid of our body besides the mucous and lymphatic system. Does the venereal poison in an infected woman ever affect the milk, and consequently can the infection be conveyed to the infant by the milk alone, without any venereal ulcer on or about the nipples? It is equally a matter of uncertainty whether the venereal disease is ever conveyed from an infected father or mother, by coition, to the fœtus, provided their geni-

tals are sound; or, whether a child is ever affected with venereal symptoms in the uterus of an infected mother. Such infected infants as came under the observation of Dr. Swediaur or of his friends, whose practice afforded them frequent opportunities of seeing new-born infants, seemed rather to militate against the opinion. Neither he nor any of them, have ever been able to observe ulcerations or other symptoms of a venereal kind upon new-born children; and such as make their appearance, four, six, or eight, or more days afterwards, on the genitals, anus, lips, mouth, &c. may rather be supposed to arise by infection during the passage from ulcers in the vagina of the mother, the skin of the infant being then nearly in as tender a state as the glans penis, or the labia; and this perhaps at the time when an absorption of the venereal poison might easier take place without a previous excoriation, or ulceration of the skin. All the ways, therefore, by which we see, in our days, the venereal poison communicated from an unhealthy to an healthy person may be reduced to the following heads:

1. By the coition of an healthy person with another who is infected with venereal symptoms of the genitals.

2. By the coition of an healthy person with another apparently healthy, in whose genitals the poison lies concealed, without having yet produced any bad symptom. Thus a woman who has perhaps received the infection from a man two or three days before, may during that time infect, and often does infect the man or men who have to do with her afterwards, without having any symptoms of the disease visible upon herself; and *vice versa*, a man may infect a woman in the same manner. Such instances occur in practice every day.

3. By sucking; in this case the nipples of the wet nurse may be infected by venereal ulcers in the mouth of the child: or *vice versa*, the nipples of the nurse being infected, will occasion venereal ulcers in the child's nose, mouth, or lips. It is uncertain as mentioned above, whether the venereal poison was ever propagated by means of the milk from the breast.

4. By exposing to the contact of venereal poison any part of the surface of the body, by kissing, touching, &c. especially if the parts so exposed have been previously excoriated, wounded, or ulcerated by any cause whatever. In this manner we frequently see venereal ulcers arise in the scrotum and thighs; and there are some well attested instances where the infection took place in the fingers of midwives or surgeons. Several instances are recorded of venereal ulcers in the nostrils, eye-lids, and lips of persons who had touched their own genitals, or those of others, affected

at the time with local venereal complaints, and then rubbed their nostrils, &c with the fingers, without previously washing the hands. There was a few years ago in London, a melancholy example of a young lady, who, after having drawn a decayed tooth, and replaced it with one taken immediately from a young woman apparently in perfect health, was soon after affected with an ulcer in the mouth. The sore manifested symptoms of a venereal nature; but such was its obstinacy, that it resisted the most powerful mercurial remedies, terminating at last in a caries of the maxilla with a most shocking erosion of the mouth and face, by which the unhappy patient was destroyed. During all this, however, we are informed that not the smallest venereal symptom was perceived in the woman from whom the sound tooth was procured.

5. By wounding any part of the body with a lancet or knife infected with the venereal virus. In this instance there is a similarity between the venereal poison and that of the small-pox. There are several examples of the latter being produced by bleeding with a lancet which had been previously employed for the purpose of inoculation, or of opening variolous pustules, without being properly cleaned afterwards. In Moravia, in the year 1577, a number of persons who assembled in a house for bathing, had themselves, according to the custom of that time, scarified by the barber, were all of them infected with the venereal disease, and treated accordingly. Krato, the physician, and Jordan, who gave a description of this distemper, are both of the opinion that it was communicated by means of the scarifying instrument. And Van Swieten relates several instances where the lues was communicated by a similar carelessness in cleaning the instrument used in bleeding or scarification.

The venereal poison applied to the urethra and vagina produce a clap. See *Gonorrhœa*. Coming into contact with other parts it produces a chancre or bubo and constitutional symptoms. Chancre is the primary and immediate consequence of inoculation with true venereal matter in any of the ways which have been mentioned, and may arise in any part of the human body: but it generally shews itself in the pudenda, because the infecting medium is there first taken up in the one sex, and communicated by contact to the other. It is not, however, peculiar to these parts, for whenever the same kind of fluid is applied to a scratch on the hand, finger, lip or nipple, the same consequence will follow. There can be no doubt but that the slightest abrasion possible, or breach of the cuticle, is sufficient to give a speedy admission to this destructive poison. A chancre makes

its appearance either with a slight inflammation which afterwards ulcerates, or there arises a small pimple or pustule filled with a transparent fluid, which soon breaks and forms into a spreading ulcer. The period at which it makes its appearance after infection is very various, being most commonly in five or six days, but in some cases not till after the expiration of as many weeks. There is both a local and general predisposition to chancres: Jews and Mahomedans, from the constant exposure of the glans and loss of the prepuce, have the cuticle of the glans penis of much firmer texture than those who have not been circumcised; and they are from this circumstance, much less subject to chancres than the rest of mankind. For the same reason they who, from the shortness of the prepuce, generally keep the glans uncovered, are neither so liable to the disease as those who have long narrow preputia; for persons thus formed constantly keep the surface of the glans and prepuce moist and tender, and almost at every cohabitation are liable to abrasions and to excoriations.

There is an intermediate state of the venereal disease between a local and constitutional affection, which arises from the absorption of venereal matter from some surface to which it has been applied. The glands situated nearest the parts thus affected are apt to become swelled and inflamed, and so as to give rise to what is termed *bubo*; and the parts of generation usually coming first in contact with the matter, so the glands in the groin generally afford this particular symptom. In most cases the venereal virus is absorbed from a chancre or an ulcer in the urethra; but instances have occurred where a bubo has arose without either gonorrhœa or any kind of ulceration, and where the matter appears to have been absorbed, without any erosion of the skin or mucous membrane.

A bubo comes on with a pain in the groin, accompanied with some degree of hardness and swelling, and is at first about the size of a kidney bean, but continuing to increase, it at length becomes as large as an egg, occasions the person to experience some difficulty in walking, and is attended with a pulsation and throbbing in the tumour, and a great redness of the skin. In some cases the suppuration is quickly completed, in others it goes on very slow, and in others again the inflammatory appearances go off without any formation of pus. In a few instances the glands have been known to become scirrhus. The following are the characteristics of a venereal bubo. The swelling is usually confined to one gland, the colour of the skin where inflammation prevails is of a florid red, the pain is very acute, the progress from inflammation to

suppuration and ulceration is generally very rapid, the suppuration is large in proportion to the size of the gland, and there is only one abscess.

A bubo is never attended with danger, where the inflamed gland proceeds on regularly to suppuration, but in particular cases it acquires an indolence after coming to a certain length, arising from a scrophulous taint, or by being combined with erysipelas it terminates in gangrene and occasions a great loss of substance. This termination is, however, more frequently met with in hospitals than in private practice, and may partly be attributed to the contaminated state of the air of the wards wherein venereal patients are lodged.

A constitutional taint is the third form under which it has been mentioned, that the venereal poison is apt to shew itself, and which always arises in consequence of the matter being absorbed and carried into the circulating mass of fluids. The absorption of it may, however, take place in three ways.

1st. It may be carried into the circulation, without producing any evident local effect on the part to which it was first applied.

2ndly. It may take place in consequence of some local affection, such as either gonorrhoea, chancre, or bubo: And,

3dly. It may ensue from an application of the matter to a common sore or wound, similar to what happens in inoculating for the small-pox.

The most general way, however, in which a constitutional taint is produced, is by an absorption of the matter, either from a chancre or a bubo.

When venereal matter gets into the system, some symptoms of it may often be observed in the course of six or eight weeks, or probably sooner: but in some cases, it will continue in the circulating mass of fluids for many months before any visible signs of its effects are produced. The system being completely contaminated, it then occasions many local effects in different parts of the body, and shews itself under a variety of forms, many of which put on the appearance of a distinct disease. We may presume that this variety depends wholly on the difference of constitution, the different kind of parts affected, and the different state these parts were in at the time the matter or poison was applied.

The first symptoms usually shew themselves on the skin and in the mouth or throat. When on the skin, reddish and brownish spots appear here and there on the surface, and eruptions of a copper colour are dispersed over different parts of the body, on the top of which these soon forms a thick scurf or scale. This scurf falls off after a short time and is succeeded by another, and the same

happening several times, and at length casting off deep scabs, an ulcer is formed which discharges an acrid foetid matter. When the matter is secreted in the glands of the throat and mouth, the tongue will often be affected so as to occasion a thickness of speech, and the tonsils, palate, and uvula will become ulcerated so as to produce a soreness and difficulty of swallowing, and likewise a hoarseness in the voice. In the venereal ulcer of the tonsil a portion of it seems as if it was dug out; it is, moreover, very foul, and has a thick white matter adhering to it, which cannot be washed off. By these characteristic marks it may, in general, readily be distinguished from any other species of ulceration in these parts.

If the disease affects the eyes, obstinate inflammation, and sometimes ulceration will also attack these organs.

The matter sometimes falls on deep seated parts, such as the tendons, ligaments, and periosteum, and occasions hard, painful swellings to arise, known by the name of nodos.

When the disease is suffered to take its own course, and not counteracted by proper remedies, the patient will in the course of time, be afflicted with severe pains, but more particularly in the night time; his countenance will become sallow, his hair will fall off, he will lose his appetite, strength and flesh, his rest will be much disturbed by night, and a small fever of the hectic kind will arise. The ulcers in the mouth and throat being likewise suffered to spread, and to occasion a caries of the bones of the palate, an opening will be made from the mouth to the nose, and the cartilages and bones of the nose being at length corroded away, this will sink on a level with the face. Some constitutions will bear up for a considerable time against the disease, whilst others again will soon sink under a general weakness and irritation produced by it. If the disorder is recent, and the constitution not impaired by other diseases, a perfect cure may easily be effected; but where it is of long standing, and accompanied with the symptoms of irritation which have been mentioned, the cure will prove tedious, and in many cases uncertain, as the constitution and strength of the patient may not admit of his going through a course of medicine sufficient to destroy the poison, or his health may be in such a state, as that only a very small quantity of mercury can be administered even at considerable intervals.

The general appearances to be observed on dissection of those who die of lues, are, caries of the bones, but more particularly those of the cranium, often communicating ulceration to the brain itself, together with enlargements and indurations of the lym-

phatic glands, scirrhus of several of the organs, particularly the liver and lungs, and exostoses of many of the hardest bones.

SYPHILIS INDICA. The yaws.

SYPHILIS POLOINICA. A variety of venereal disease.

SYPHILIS VENEREA. See *Syphilis*.

SYRIUM OLEUM. A fragrant essential oil, obtained by distilling the canary balsam plant or moldavica.

Syrian herb mastich. See *Marum syriacum*.

STRIGUUS. See *Paracusis*.

STRINGA. (From *εγγυζ*, a pipe; so called because from its branches pipes were made after the removal of the pith.) The pipe-tree.

STRINGOS. See *Paracusis*.

STRINGOTOMUM. (From *εγγυζ*, a fistula, and *τομή*, to cut.) An instrument to cut fistulas.

STRINX. (Heb.) A pipe. A syringe. A fistula.

STRYMIANUS. (From *εμεμαζα*, to evacuate.) A gentle evacuation by vomit or stool.

SYRUPUS. (*Serab*, a potion, Arab.) 1. When sugar is dissolved in any vegetable liquor, to the consistence of honey, a medical preparation is formed called syrup; which, if obtained from a single plant, is called *simple*: but if from more than one, *compound*. To keep syrups without fermenting, it is necessary that their temperature should be attended to, and kept as near 55° as possible. A good cellar will answer this purpose, for there are few summers in which the temperature of such a place rises to 60°.

2. The pharmacopœial name of a simple syrup. *Syrupus simplex*. "Take of refined sugar, two pounds and a half; water, a pint." Dissolve the sugar in the water in a water-bath; then set it aside for 24 hours; take off the scum, and if there be any faculencies, pour off the clear liquor from them.

Syrup of buckthorn. See *Syrupus rhamni*.

Syrup of ginger. See *Syrupus zingiberis*.

Syrup of lemon. See *Syrupus limonis*.

Syrup of marsh-mallows. See *Syrupus altheæ*.

Syrup of mulberry. See *Syrupus mori*.

Syrup of orange. See *Syrupus aurantii*.

Syrup of poppy. See *Syrupus papaveris*.

Syrup of red poppy. See *Syrupus Rhæados*.

Syrup of roses. See *Syrupus roseæ*.

Syrup of saffron. See *Syrupus croci*.

Syrup of senna. See *Syrupus sennæ*.

Syrup of Tolu. See *Syrupus Tolubanus*.

SYRUPUS ACETI. Sugar and vinegar. A refrigerating and antiseptic syrup.

SYRUPUS ALTHEÆ. Syrup of marsh-mallow. *Syrupus ex althea*. *Syrupus de althea*. "Take of the fresh root of marsh-mallow, bruised, half a pound; refined sugar, two pounds; water, a gallon." Boil down the water with the marsh-mallow-root to half, and press out the liquor when cold. Set it by for 24 hours, that the faculencies may subside; then pour off the liquor, and having added the sugar, boil it down to a proper consistence. An emollient and demulcent; mostly given to alay tickling coughs, hoarseness, &c. in conjunction with other remedies.

SYRUPUS AURANTII. Syrup of orange. *Syrupus corticis aurantii*. *Syrupus e corticibus aurantium*. *Syrupus de cortice aurantium*. "Take of fresh orange-peel, two ounces; boiling water, a pint; refined sugar, three pounds." Macerate the orange-peel in the water for 12 hours in a covered vessel; then pour off the liquor, and add the sugar. A pleasant bitter and stomachic.

SYRUPUS CARTOPHYLLI RUBRI. A warm and stimulating syrup.

SYRUPUS COLCHICI. An acrid and diuretic compound given in dropsies.

SYRUPUS CORTICIS AURANTII. See *Syrupus aurantii*.

SYRUPUS CROCI. Syrup of saffron. "Take of saffron, an ounce; boiling-water, a pound; refined sugar, two pounds and a half." Macerate the saffron in the water for 12 hours in a covered vessel, then strain the liquor, and add the sugar. This imparts a beautiful colour to liquors, and is sometimes employed as a cordial. Amongst the vulgar, syrup of saffron is in high esteem in measles, small-pox, &c.

SYRUPUS LIMONIS. Syrup of lemon. *Syrupus succi limonis*. *Syrupus e succo limonium*. *Syrupus e succo citriorum*. "Take of lemon-juice, strained, a pint; refined sugar, two pounds." Dissolve the sugar in the lemon-juice in the manner directed for a syrup. A very pleasant, cooling, and acrid syrup, which may be exhibited with advantage, in gastritis and bilious affections.

SYRUPUS MORI. Syrup of mulberry. *Syrupus mororum*. "Take of mulberry-juice, strained, a pint; refined sugar, two pounds." Dissolve the sugar in the mulberry-juice in the manner directed for syrup. Syrup of mulberries is very grateful and aperient, and may be given with such intentions to children.

SYRUPUS PAPAVERIS. *Syrupus papaveris albi*. *Syrupus e meconio*. *Syrupus de meconio*, sive *diacodium*. "Take of capsules of white poppy, dried and bruised, the seeds being separated, 14 ounces; refined sugar, two pounds; boiling-water, two

gallens and a half." Macerate the capsules in the water for 12 hours, then boil it down by means of a water-bath to one gallon, and press out the liquor. Boil down the liquor again to two pints, and strain it while hot. Set it by for 12 hours, that the feculencies may subside: then boil down the clear liquor to a pint, and add the sugar in the manner directed for a syrup. It should be kept in stone bottles and in a cellar. A useful anodyne preparation which may be added with advantage to a vast variety of medicines against diseases of the bowels, coughs, &c.

SYRUPUS PAPAVERIS ERRATICI. See *Syrupus rheados*.

SYRUPUS RHEADOS. *Syrupus papaveris erratici.* *Syrupus de papavere errático.* Syrup of red-poppy. "Take of red-poppy petals, fresh, a pound; boiling-water, a pint and two fluidounces; refined sugar, two pounds and a half." This is a very mild anodyne, and used more for the colour than for its medical properties.

SYRUPUS RHAMNI. Syrup of buckthorn. "Take of the fresh juice of buckthorn-berries, four pints; ginger-root, sliced, allspice, powdered, of each half an ounce; refined sugar, three pounds and a half." Set by the juice for three days that the feculencies may subside, and strain. To a pint of the clear juice add the ginger, the ginger-root, and allspice; then macerate in a gentle heat four hours, and strain; boil down what remains to one pint and a half, mix the liquors, and add the sugar in the manner directed for syrup.

This preparation, in doses of three or four spoonfuls, operate as a brisk cathartic. The principal inconvenience attending it is, that it is very unpleasant and occasions a thirst and dryness of the mouth and fauces, and sometimes violent gripes; these effects may be prevented by drinking liberally of water-gruel, or other warm liquids during the operation.

SYRUPUS RIBIS NIGRI. Syrup of black currants. Aperient and diuretic qualities are attributed to this preparation.

SYRUPUS ROSÆ. Syrup of roses. *Syrupus rosarum solutivus.* *Syrupus e rosis siccis.* "Take of damask-rose petals, dried, seven ounces; refined sugar, six pounds; boiling-water, four pints." Macerate the rose-petals in the water for 12

hours, and strain: then evaporate the strained liquor by means of a water-bath to two pints and a half; then add the sugar in the manner described for a syrup. A useful laxative for children. From ʒj. to ʒss.

SYRUPUS RUBI IDÆI. Syrup of raspberry. A pleasant aperient syrup for children.

SYRUPUS SCILLITICUS. Expectorant and diuretic.

SYRUPUS SENNÆ. Syrup of senna: "Take of senna-leaves, half an ounce; fennel-seed, bruised, a drachm; manna, refined sugar, of each a pound; water, boiling, a pint" Macerate the senna-leaves and fennel-seeds in the water for 12 hours; strain the liquor, and mix with it manna and sugar. A useful purgative for children.

SYRUPUS TOLUTANUS. Syrup of Tolu. "Take of balsam of Tolu, an ounce; water, boiling, a pint; refined sugar, two pounds." Boil the balsam in the water half an hour in a covered vessel, occasionally stirring it, strain the liquor when it is cold, and then add the sugar, in the manner directed for syrup. A useful balsamic syrup, calculated to allay tickling coughs and hoarsenesses.

SYRUPUS VIOLE. A pleasant laxative for young children.

SYRUPUS ZINGIBERIS. Syrup of ginger. "Take of ginger-root, sliced, two ounces; water, boiling, a pint; refined sugar, two pounds." Macerate the ginger-root in the water for twenty-four hours, and strain, then add the sugar in the manner directed for syrup. A carminative and stomachic syrup. Dose from one to three drachms.

SYSSARCOSIS. (From *σύν*, and *σάρξ*, flesh.) A species of union of bones in which one bone is united to another by means of an intervening muscle. In this manner the os hyoides is connected with the sternum and other parts.

System, absorbent. See *Absorbents* and *Lymphatics*.

System, genital. The parts of generation.

System, nervous. See *Nerves*.

System, vascular. The arteries and veins.

SYSTOLE. (From *συστάναι*, to contract.) The contraction of the heart.

T

T BANDAGE. A bandage so named from its figure. It is principally used for supporting the dressings, after the operation for fistula in ano, in diseases of the perinæum, and those of the groins, anus, &c.

TABAGUM. (From *Tobago*, the island from whence it was first brought.) Tobacco. See *Nicotiana*.

TABELLA. (Dim. of *tabula*, a table.) A lozenge.

TABES. (From *tabesco*, to consume or pine away.) A wasting of the body. A genus of disease in the class *cachexie* and order *marcores* of Cullen; characterised by emaciation and weakness, attended with hectic fever, but without any cough or spitting, which last symptoms distinguish it from phthisis. It has three species: 1. *Tabes purulenta*, from an ulcerous discharge; 2. *Tabes scrofulosa*, from a scrofulous habit; 3. *Tabes venenata*, from poison. See *Atrophy*.

TABES COXARIA. *Phthisis ischiadica*. A wasting of the thigh and leg from an abscess, or other cause, in the hip.

TABES DORSALIS. *Lordosis*. Dr. Cullen makes it a variety of *atrophia inanitorium*. Hippocrates calls it *tabes ossis sacri*. At present by the name of *tabes dorsalis* is understood a wasting of the body, attended at first with pain in the back or loins, and afterwards also in the neck and head, caused by a too early or a too frequent use of venery.

TABES NUTRICUM. A variety of *atrophia inanitorium*.

TABES OSSIS SACRI. Hippocrates' name for *tabes dorsalis*.

TABES PULMONALIS. A name for phthisis.

TABES RENALIS. An abscess of the kidney.

TABES SYPHILITICA. A variety of the *atrophia cacoehymica*.

TACAMAHACCA. (Indian.) The name of a resinous substance which exudes both spontaneously and when incisions are made into the stem of the *Fugera octandria* of Linnæus:—*foliolis tomentosis*, and not as was formerly supposed from the *Populus balsamifera* Linneæ. Two kinds of *tacamahacca* are met with in the shops. The best, called, from its being collected in a kind of gourd shell, *tacamahacca* in shells, is somewhat unctuous and soft, of a pale yellowish or greenish colour, a bit-terish aromatic taste, and a fragrant delightful smell, approaching to that of la-

vender and ambergris. The more common sort is in semi-transparent grains, of a whitish, yellowish, brownish, or greenish colour, and of a less grateful smell than the former. *Tacamahacca* was formerly in high estimation as an ingredient in warm stimulating plaisters; and although seldom used internally, it may be given with advantage as a corroborant and adstringent balsamic.

TACTUS. See *Touch*.

TEDA. (*Δαδα*, from *δαο*, to burn.) A torch. A species of pine which burns like a torch. A medicated torch for fumigations.

TENIA. (*Tania*, a Hebrew word, signifying a fillet; so named from its resemblance to a fillet or piece of tape.) The tape-worm. A genus of intestinal worms; characterised by a long, flat, and jointed body. Species: 1. *Tenia osculis marginalibus*, the long tape-worm, and the soleum of authors, which is peculiar to this country, Russia, France, &c.: 2. *Tenia osculis superficialibus*, the broad tape-worm, which is peculiar to the inhabitants of Switzerland, &c. See *Worms*.

Talc. See *Talcum*.

TALCUM. (From *talk*, German.) *Tale*. A white, grey, yellow, or greenish substance of a soft and soapy touch, formed of transparent laminæ placed upon each other. Talc is composed of pure magnesia mixed with near twice its weight of silice and less than its weight of alumine. There are several different appearances of *talc*. The greenish foliaceous Venice *talc* was formally used medicinally, as possessing antacid and aperient qualities.

TALPA. (From *ταλαο*, blind.) *Talporia*. A mole. Also a tumour resembling a mole in eating and creeping under the skin.

TALUS. A synonym of *Astragalus*. See *Astragalus*.

TAMALAPATRA. The Indian leaf is so termed by some authors. See *Cassia lignea*.

Tamarind See *Tamarindus*.

TAMARINDUS. (From *tamar* or *tamarindi*, which is in the Arabian language a synonym of the dactylus or date.) *Oxyphenicon*. *Saliqua arabica*. *Balam pulli*. *Tamaræa*. *Zecia oxyphencia acacia indica*. The tamarind. The tree which affords this fruit is the *Tamarindus indica* of Linnæus. The pulp of the tamarind, with the seeds, connected together by numerous

tough strings or fibres, are brought to us freed from the outer shell, and commonly preserved in syrup. According to Long, tamarinds are prepared for exportation at Jamaica, in the following manner: "The ruit or pods are gathered in June, July, and August, when full ripe, which is known by their fragility or easy breaking on small pressure between the finger and thumb. The fruit taken out of the pod, and cleared from the shelly fragments, is placed in layers in a cask, and boiling syrup, just before it begins to granulate, is poured in, till the cask is filled: the syrup pervades every part quite down to the bottom, and when cool the cask is headed for sale." The tamarind is employed as a laxative, and for abating thirst or heat in various inflammatory complaints, and for correcting putrid disorders, especially of a bilious kind, in which the cathartic, antiseptic, and refrigerant qualities of the fruit have been found equally useful. When intended merely as a laxative, it may be of advantage, (Dr. Woodville observes) to join it with manna or purgatives of a sweet kind, by which its use is rendered safer and more effectual. Three drachms of the pulp are usually sufficient to open the body, but to prove moderately cathartic, one or two ounces are required. It is an ingredient in the *confectio cassie* and *confectio sennæ*.

TAMARINDUS INDICA. The systematic name of the tamarind-tree. See *Tamarindus*.

TAMARISCUS. (From *Tamarik*, abster-tion, Heb. named from its properties of cleansing and purifying the blood.) Tamarisk. The bark, wood, and leaves of this tree, *Tamarix gallica* of Linnæus, were formerly employed medicinally, though seldom used at present. The former for its aperient and corroborant virtues in obstructions of the liver; the latter in icterus, hæmoptysis, and some affections of the skin.

TAMARIX GALLICA. The systematic name of the tamarisk-tree. See *Tamariscus*.

Tame-poison. See *Vincetoxicum*.

TANACETUM. (Corrupted from *tanasia*, *athanasia*, the old name for tansy.) 1. The name of a genus of plants in the Linnæan system. Class, *Syngenesia*. Order, *Polygamia superflua*. Tansy.

2. The pharmacopœial name of the common tansy. *Tanasia*. *Athanasia*. *Parthenium mas*. *Tanacetum vulgare* of Linnæus:—*foliis bipinnatis incisive serratis*. The leaves and flowers of tansy have a strong, not very disagreeable smell, and a bitter somewhat aromatic taste. The virtues of tansy are tonic, stomachic, anthelmintic, emmenagogue, and resolvent. It has been much used as a vermifuge; and testimonies of its efficacy are given by many respectable physicians. Not only the leaves

but the seeds have been employed with this intention, and substituted for those of santonium. We are told by Dr. Clark, that in Scotland tansy was found to be of great service in various cases of gout; and Dr. Cullen, who afterwards was informed of the effect it produced upon those who had used the herb for this purpose says, "I have known several who have taken it without any advantage, and some others who reported that they had been relieved from the frequency of their gout." Tansy is also recommended in the hysteria, especially when this disease is supposed to proceed from menstrual obstructions.

This plant may be given in powder to the quantity of a drachm or more for a dose; but it has been more commonly taken in infusion, or drank in tea.

TANACETUM BALSAMITA. The systematic name of the officinal alecost. See *Balsamita mas*.

TANACETUM HORTENSE. See *Balsamita mas*.

TANACETUM VULGARE. The systematic name of the common tansy. See *Tanacetum*.

TANASIA. See *Tanacetum*.

Tansy. See *Tanacetum*.

Tansy, wild. See *Potentilla*.

Type-worm. See *Tegmæ*.

Tapioca. See *Cassava*.

Tupping. See *Parancatesis*.

TAPBUS BARBATUS. See *Verbascum*.

Tar. See *Pix liquida*.

Tar, Barbadoes. See *Petroleum barbadense*.

TAR-WATER. A once celebrated remedy, but now neglected more than it deserves. It is made by infusing tar in water, stirring it from time to time, and lastly pouring off the clear liquor now impregnated with the colour and virtues of the tar. It is drank in many chronic affections, particularly of the lungs.

TARANTISMUS (From *tarantula*, the animal whose bite is supposed to be cured only by music.) The desire of dancing which is produced by the bite of the tarantula.

TARANTULA. (From *Taranta*, a city in Naples, where they abound.) A kind of venomous spider, whose bite is said to be cured by music.

TARAXACUM. (From *ταρασσω*, to alter or change; because it alters the state of the blood.) *Dens leonis*. The dandelion, or pissabed. *Leontodon taraxacum* of Linnæus:—*caule squamis inferne reflexis, foliis runcinatis denticulatis levibus*. The young leaves of this plant in a blanched state have the taste of endive, and make an excellent addition to those plants eaten early in the spring as salads; and Murray informs us, that at Goettingen, the roots are roasted and substituted for coffee by the poorer inhabitants, who find that an

infusion prepared in this way can hardly be distinguished from that of the coffee-berry. The expressed juice of dandelion is bitter and somewhat acid; but that of the root is bitterer, and possesses more medicinal power than any other part of the plant. It has been long in repute as a detergent and aperient, and its diuretic effects may be inferred from the vulgar name it bears in most of the European languages, *quasi lecti minga et urinaria herba dicitur*; and there are various proofs of its efficacy in jaundice, dropsy, consumption, and some cutaneous disorders.

The leaves, roots, flower, stalks, and juice of dandelion, have all been separately employed for medicinal purposes, and seem to differ rather in degree of strength than in any essential property; therefore the expressed juice, or a strong decoction of the roots have most commonly been prescribed, from one ounce to four, two or three times a day. The plant should be always used fresh; even extracts prepared from it appear to lose much of their power by keeping.

TABAXIS. (From *ταβαιν*, to disturb.) A slight ophthalmia or inflammation of the eye.

TARCHON SYLVESTRIS. See *Ptarmica*.

Tare. See *Errum*.

Tarsi, extensor minor. See *Plantares*.

TARSUS. *Τάρσος*. 1. The instep or that part of the foot which is between the leg and metatarsus: it is composed of seven bones, viz. the astragalus, os calcis, os naviculare, os cuboides, and three ossa cuneiformia.

2. The thin cartilage situated at the edges of the eyelids to preserve their firmness and shape.

TARTAR. (*Tartarum*, from *ταρταρος*, infernal; because it is the sediment or dregs.) 1. The concretion which fixes to the inside of hogsheads containing wine. It is alloyed with much extractive and colouring matter, from which it is purified by decoction with argillaceous earths and subsequent crystallization. By this means it becomes perfectly white, and shoots out crystals of tartar, consisting of a peculiar acid, called acid of tartar and potash, and is a super-tartrate of that alkali which when powdered is the cream or tartar of the shops. Its virtues are eccoprotic, diuretic, and refrigerant, and it is exhibited in abdominal physconia, dropsy, inflammatory and bilious fevers, dyspepsia from rancid or fat substances, bilious diarrhoea and colic, hæmorrhoids and obstipation.

2. A name heretofore given to many official preparations, containing the acid of tartar; but in consequence of recent changes in the chemical nomenclature superseded by appellations more expressive of the respective compositions.

TARTARIC ACID. *Acidum tartari-*

cum. Sal essentialis tartari. Acidum tartari essentialis. Tartareous acid. To obtain the pure tartaric acid, take two pounds of the crystals, and dissolve them in water, into which chalk is to be thrown by degrees till the liquid is saturated. A precipitate is formed, which is a true tartrate of lime, is tasteless, and cracks between the teeth. This tartrate is put into a cucurbit, and nine ounces of sulphuric acid, with five ounces of water, are poured on it. After twelve hours digestion, with occasional stirring, the tartaric acid is set at liberty in the solution, and may be cleared off the sulphate of lime by means of cold water. The virtues of this acid are antiseptic, refrigerant, and diuretic. It is used in acute fevers, scurvy, and hæmorrhage.

Tartar, cream of. The popular name of the purified super-tartrate of potash.

Tartar, emetic. See *Antimonium tartarizatum*.

Tartar, oil of. See *Liquor carbonatis potassæ*.

Tartar, regenerated. See *Potassæ acetatæ*.

Tartar, salt of. See *Potassæ subcarbonatæ*.

Tartar, soluble. See *Potassæ tartaræ*.

Tartar, spirit of. If the crystals of tartar be distilled by a strong heat, without any additional body, they furnish an empyreumatic acid, called the pyrotartareous acid, or spirit of tartar, and a very fetid empyreumatic oil.

Tartar, vitriolated. See *Potassæ sulphatæ*.

TANTARUM EMETICUM. See *Antimonium tartarizatum*.

TANTARUM REGENERATUM. See *Potassæ acetatæ*.

TANTARUM SOLUBILE. See *Kali tartarizatum*.

TANTARUS AMMONIÆ. See *Tartarus ammoniacæ*.

TANTARUS CHALYBEATUS. See *Tartarus potassæ acidulus ferratus*.

TARTARUS AMMONIACÆ. *Alkali volatile tartarizatum* of Bergman. *Sal ammoniacum tartareum.* *Tartarus ammoniæ.* A salt composed of tartaric acid and ammonia, its virtues are diaphoretic, diuretic, and deobstruent. It is prescribed in fevers, atonic exanthemata, catarrh, arthritic and rheumatic arthrodynia, hysteric spasms, &c.

TARTARUS POTASSÆ. See *Potassæ tartaræ*.

TARTARUS POTASSÆ ACIDULUS. Cream of tartar. See *Tartar*.

TARTARUS POTASSÆ ACIDULUS FERRATUS. *Globuli martiales.* *Tartarus chalybeatus.* *Murs solubilis.* *Ferrum potabile.* Its virtues are adstringent. It is principally used externally in the form of fomentation or bath in contusions, distortions and luxations.

TARTARUS POTASSÆ ACIDULUS STIBIATUS. See *Antimonium tartarizatum*.

TARTARUS SODÆ. See *Soda tartarizata*.

TASTE. *Gustus.* The organ of taste differs but slightly from that of touch: It appears, by certain experiments, to be seated chiefly in those nervous papillæ of the tongue which are formed from the minute ends of the ninth or lingual pair of nerves; for neither does sugar, applied to any other part of the mouth, excite the least sense of taste in the mind; nor any other sapid body, unless it contain something vehemently penetrating; in which case the palate, root of the tongue, uvula, and even the œsophagus, are affected by the sapid acrimony. That sensation, which is sometimes excited in the stomach, œsophagus, and fauces, by the regurgitation of the aliments, seems also to belong to the tongue, to which the sapid vapours are applied.

Nature designed the diversity of flavours, that animals might know those things most proper for their food; for in general, there is no aliment unhealthy, that is of an agreeable taste; nor is any thing ill tasted that is fit for the food of man. We here take no notice of excess, by which the most healthy food may become prejudicial, or of minerals, which are not furnished by nature, but prepared by art. Thus nature has invited man to take the food necessary for his subsistence, both by the pain called hunger, and by the pleasure arising from taste. But animals, which do not learn from example and the instruction of others, distinguish flavours most accurately, and, admonished by that test, abstain cautiously from unhealthy food; and, therefore, herbivorous animals especially, to which a very great diversity of aliments mixed with noxious plants are offered, are furnished with such long papillæ, and so elegant a structure of the tongue, for which man has less occasion.

TAXIS. An operation, by which those parts which have quitted their natural situation are replaced by the hand without the assistance of instruments, as in reducing hernia, &c.

TEA. See *Thea*.

TEAR. *Lachryma.* The limpid fluid secreted by the lachrymal glands, and flowing on the surface of the eye.

The organ which secretes this liquid is formed by the lachrymal glands, one of which is situated in the external canthus of each orbit, and emits six or seven excretory ducts, which open on the internal surface of the upper eyelid above its tarsus, and pour forth the tears. The tears have mixed with them an arterious, rosicid vapour, which exhales from the internal surface of the eyelids, and external of the tunica conjunctiva, into the eye. Perhaps the aqueous humour also transudes through

the pores of the cornea on the surface of the eye. A certain part of this aqueous fluid is dissipated in the air; but the greatest part, after having performed its office, is propelled by the orbicular muscle, which so closely constricts the eyelid to the ball of the eye, as to leave no space between, unless in the internal angle, where the tears are collected. From this collection the tears are absorbed by the orifices of the puncta lachrymalia; from thence they are propelled through the lachrymal canals, into the lachrymal sac, and flow through the ductus nasalis into the cavity of the nostrils, under the inferior concha nasalis. The *lachrymal sac*, appears to be formed of longitudinal and transverse muscular fibres; and its three orifices furnished with small sphincters, as the spasmodic constriction of the puncta lachrymalia proves, if examined with a probe.

The tears have no smell but a saltish taste, as people who cry perceive. They are of a transparent colour and aqueous consistence.

The *quantity*, in its natural state, is just sufficient to moisten the surface of the eye and eyelids; but from sorrow, or any kind of stimulus applied to the surface of the eye, so great is the quantity of tears secreted, that the puncta lachrymalia are unable to absorb them. Thus the greatest part runs down from the internal angle of the eyelids, in the form of great and copious drops upon the cheeks. A great quantity also descends, through the lachrymal passages into the nostrils; hence those who cry have an increased discharge from the nose.

Use of the tears.—1. They continually moisten the surface of the eye and eyelids, to prevent the pellucid cornea from drying and becoming opaque, or the eye from concreting with the eyelids. 2. They prevent that pain, which would otherwise arise from the friction of the eyelids against the bulb of the eye from continually winking. 3. They wash and clean away the dust of the atmosphere, or any thing acrid that has fallen into the eye. 4. Crying unloads the head of congestions.

TEETH. (*Dens*, a tooth; *quasi edens*, from *edo*, to eat.) Small bones fixed in the alveoli of the upper and under jaw. In early infancy Nature designs us for the softest aliment, so that the gums alone are then sufficient for the purpose of mastication; but as we advance in life, and require a different food, she wisely provides us with teeth. These are the hardest and whitest of our bones, and, at full maturity, we usually find thirty-two in both jaws; viz. sixteen above, and as many below. Their number varies indeed in different subjects; but it is seldom seen to exceed thirty-two, and it will very rarely be found to be less than twenty-eight.

Each tooth may be divided into two parts; viz. its body, or that part which appears above the gums; and its fangs or root, which is fixed into the socket. The boundary between these two, close to the edge of the gum, where there is usually a small circular depression, is called the neck of the tooth. The teeth of each jaw are commonly divided into three classes; but before each of these is treated of in particular, it will be right to say something of their general structure.

Every tooth is composed of its *cortex* or *enamel*, and its internal bony substance. The enamel, or as it is sometimes called, the vitreous part of the tooth, is a very hard and compact substance, of a white colour, and peculiar to the teeth. It is found only upon the body of the tooth, covering the outside of the bony or internal substance. When broken it appears fibrous or striated; and all the striz are directed from the circumference to the centre of the tooth. This enamel is thickest on the grinding surface, and on the cutting edges or points of the teeth, becoming gradually thinner as it approaches the neck, where it terminates insensibly. Some writers have described it as being vascular, but it is certain that no injection will ever reach this substance; that it receives no tinge from madder; and that it affords no appearance of a circulation of fluids. The bony part of a tooth resembles other bones in its structure, but is much harder than the most compact part of bones in general. It composes the inner part of the body and neck, and the whole of the root of the tooth. This part of a tooth, when completely formed, does not, like the other bones, receive a tinge from madder, nor do the minutest injections penetrate into its substance, although many writers have asserted the contrary. Mr. Hunter has been therefore induced to deny its being vascular, although he is aware that the teeth, like other bones, are liable to swellings, and that they are found ankylosed with their sockets. He supposes, however, that both these may be original formations; and, as the most convincing proof of their not being vascular, he reasons from the analogy between them and other bones. He observes, for instance, that in a young animal that has been fed with madder, the parts of the teeth which were formed before it was put on madder diet will appear of their natural colour, but that such parts as were formed while the animal was taking the madder, will be of a red colour; whereas, in other bones, the hardest parts are susceptible of the dye, though more slowly than the parts which are growing. Again, he tells us, that if you leave off feeding the animal with madder a considerable time before

you kill it, you will find the above appearances still subsisting, with this addition that all the parts of the teeth which were formed after leaving off the madder will be white. This experiment proves that a tooth once tinged does not lose its colour; whereas other bones do (though very slowly) return again to their natural appearance: and, as the dye in this case must be taken into the habit by absorbents, he is led to suspect that the teeth are without absorbents as well as other vessels. These arguments are very ingenious, but they are far from being satisfactory. The facts adduced by Mr. Hunter are capable of a different explanation from that which he has given them; and when other facts are added relative to the same subject, it will appear that this bony part of a tooth has a circulation through its substance, and even lymphatics, although, from the hardness of its structure, we are unable to demonstrate its vessels. The facts which may be adduced are, 1st. We find that a tooth recently drawn and transplanted into another socket, becomes as firmly fixed after a certain time, and preserves the same colour as the rest of the set; whereas a tooth that has been long drawn before it is transplanted, will never become fixed. Mr. Hunter, indeed, is aware of this objection, and refers the success of the transplantation, in the first instance, to the living principle possessed by the tooth, and which he thinks may exist independent of a circulation. But however applicable such a doctrine may be to zoophytes, it is suspected that it will not hold good in man, and others of the more perfect animals: and there does not appear to be any doubt but that, in the case of a transplanted tooth, there is a real union by vessels. 2dly. The swellings of the fangs of a tooth, which in many instances are known to be the effects of disease, and which are analogous to the swelling of other bones, are a clear proof of a similarity of structure, especially as we find them invested with a periosteum. 3dly. It is a curious fact, though as yet perhaps not generally known, that, in cases of phthisis pulmonalis, the teeth become of a milky whiteness, and in some degree, transparent; does not this prove them to have absorbents?

Each tooth has an inner cavity, which, beginning by a small opening at the point of the fang, becomes larger, and terminates in the body of the tooth. This cavity is supplied with blood-vessels and nerves, which pass through the same hole in the root. In old people this hole sometimes closes, and the tooth becomes then insensible.

The teeth are invested with a periosteum from their fangs to a little beyond the bony sockets, where it is attached to the gums. This membrane seems to be of

mon to the tooth which it encloses, and to the sockets which it lines. The teeth are likewise secured in their sockets by a red substance called the *gums*, which every where covers the alveolar processes, and has as many perforations as there are teeth. The gums are exceedingly vascular, and have something like cartilaginous hardness and elasticity, but do not seem to have much sensibility. The gums of infants, which perform the offices of teeth, have a hard ridge extending through their whole length; but in old people, who have lost their teeth, this ridge is wanting. The three classes into which the teeth are commonly divided, are *incisores*, *canini*, and *molars* or *grinders*.

The *incisores* are the four teeth in the fore part of the jaws; they derive their name from their use in dividing and cutting the food in the manner of a wedge, and have each of them two surfaces, which meet in a sharp edge. Of these surfaces, the anterior one is convex, and the posterior one somewhat concave. In the upper jaw they are usually broader and thicker, specially the two first, than those of the under jaw, over which they generally fall obliquely being placed a little obliquely.

The *canini* or *cuspidati* are the longest of all the teeth, deriving their name from their resemblance to a dog's tusk. There is one of these teeth on each side of the incisores, so that there are two in each jaw. They are the longest of all the teeth. Their fangs differ from that of the incisores only in being much larger, and their shape may be easily described to be that of an incisor with its edge worn off, so as to end in a narrow point instead of a thin edge. The *canini* not being calculated for dividing like the incisores, or for grinding, seem to be intended for laying hold of substances.

Mr. Hunter remarks of these teeth, that we may trace in them a similarity in shape, situation, and use, from the most imperfect carnivorous animal which we believe to be the human species, to the lion, which is the most perfectly carnivorous.

The *molars*, or *grinders*, of which there are ten in each jaw, are so called, because from their size and figure they are calculated for grinding the food. The *canini* and *incisores* have only one fang, but the three last grinders in the under jaw have constantly two fangs, and the same teeth in the upper jaw three fangs. Sometimes these fangs are divided into two points near their base, and each of these points has, perhaps, been sometimes considered as a distinct fang. The grinders likewise differ from each other in their appearance. The two first on each side, which Mr. Hunter appears to have distinguished very properly by the name of *bluspides*, seem to be of a middle nature between the incisores and grinders; they

have in general only one root, and the body of the tooth terminates in two points, of which the anterior one is the highest, so that the tooth has in some measure the appearance of one of the canini. The two grinders beyond these, on each side, are much larger. Their body forms almost a square with rounded angles; and their grinding surface has commonly five points or protuberances, two of which are on the inner, and three on the outer part of the tooth. The last grinder is shorter and smaller than the rest, and, from its coming through the gums later than the rest, and sometimes not appearing till late in life, is called *dens sapientie*. The variation in the number of teeth usually depends on these *dentes sapientie*.

Having thus described the appearance of the teeth in the adult; the manner of their formation and growth in the fœtus is next to be considered. We shall find that the alveolar process, which begins to be formed at a very early period, appears about the fourth month, only as a shallow longitudinal groove, divided by slight ridges into a number of intermediate depressions, which are to be the future alveoli or sockets. These depressions are at first filled with small pulpy substances, included in a vascular membrane; and these pulpy substances are the rudiments of the teeth. As these advance in their growth, the alveolar processes become gradually more completely formed. The surface of the pulp first begins to harden; the ossification proceeding from one or more points, according to the kind of tooth that is to be formed. Thus, in the incisores and *canini*, it begins from one point; in the *bluspides*, from two points, corresponding with the future shape of those teeth; and in the *molars* from four or five points. As the ossification advances, the whole of the pulp is gradually covered with bone, excepting its under surface, and then the fang begins to be formed. Soon after the formation of this bony part, the tooth begins to be incrustated with its enamel; but in what manner this is deposited we are as yet unable to explain.—Perhaps the vascular membrane, which encloses the pulp, may serve to secrete it. It gradually crystallizes upon the surface of the bony part, and continues to increase in thickness, especially at the points and basis of the tooth, till some time before the tooth begins to pass through the gum; and when this happens, the enamel seems to be as hard as it is afterwards, so that the air does not appear to have the least effect in hardening it, as has been sometimes supposed.—While the enamel is thus forming, the lower part of the pulp is gradually lengthened out and ossified, so as to form the fang. In those teeth which are to have more than one fang, the ossification

begins from different parts of the pulp at one and the same time. In this manner are formed the incisors, the canini, and two molares on each side, making in the whole twenty teeth, in both jaws, which are sufficient for the purposes of manducation early in life. As the fangs of the teeth are formed, their upper part is gradually pushed upwards, till at length, about the seventh, eighth, or ninth month after birth, the incisors, which are the first formed, begin to pass through the gum. The first that appears is generally in the lower jaw. The canini and molares not being formed so soon as the incisors, do not appear till about the twentieth or twenty-fourth month. Sometimes one of the canini, but more frequently one of the molares appears first.

The danger to which children are exposed, during the time of dentition, arises from the pressure of the teeth in the gum, so as to irritate it, and excite pain and inflammation. The effect of this irritation is, that the gum wastes, and becomes gradually thinner at this part, till at length the tooth protrudes. In such cases therefore we may, with great propriety, assist nature by cutting the gum. These twenty teeth are called *temporary*, or *milk* teeth, because they are all shed between the age of seven and fourteen, and are supplied by others of a firmer texture, with large fangs, which remain till they become affected by disease, or fall out in old age, and are therefore called the *permanent*, or *adult* teeth. The rudiments of these adult teeth begin to be formed at different periods. The pulp of the first adult incisor, and of the first adult grinder, may be perceived in a fœtus of seven or eight months, and the ossification begins in them about six months after birth. Soon after birth the second incisor, and canine tooth on each side, begin to be formed. About the fifth or sixth year the first bicuspid and about the seventh the second bicuspid begins to ossify. These bicipides are destined to replace the temporary grinders. All these permanent teeth are formed in a distinct set of alveoli; so that it is not by the growing of one tooth under another in the same socket, that the uppermost tooth is gradually pushed out, as is commonly imagined; but the temporary teeth, and those which are to succeed them, being placed in separate alveoli, the upper sockets gradually disappear, as the under ones increase in size, till at length the teeth they contain, having no longer any support, consequently fall out. But, besides these twenty teeth, which succeed the temporary ones, there are twelve others to be added to make up the number thirty-two. These twelve are three grinders on each side in both jaws; and in order to make room for this addition, we find the jaws grow as the teeth grow, so that they appear as com-

pletely filled with twenty teeth, as they are afterwards with thirty-two. Hence, in children the face is flatter and rounder than in adults. The first adult grinder usually passes through the gum about the twelfth year; the second, which begins to be formed in the sixth or seventh year, cuts the gum about the seventeenth or eighteenth; and the third or dens sapientie, which begins to be formed about the twelfth year, passes through the gum between the age of twenty and thirty. The dentes sapientie have, in some instances, been cut at the age of forty, fifty, sixty, and even eighty years; and it sometimes happens that they do not appear at all. Sometimes likewise it happens, that a third set of teeth appear about the age of sixty or seventy. Diermebroeck tells us that he himself, at the age of fifty-six, had a fresh canine tooth in the place of one he had lost several years before; M. du Fay saw two incisors and two canini cut the gum in a man aged eighty-four; Mr. Hunter has seen two fore-teeth shoot up in the lower jaw of a very old person; and an account was lately published of a man who had a complete set at the age of sixty. Other instances of the same kind are to be met with in authors. The circumstance is curious, and from the time of life at which it takes place, and the return of the catamenia, which sometimes happens to women at the same age, it has been very ingeniously supposed, that there is some effort in nature to renew the body at that period.

The teeth are subject to a variety of accidents. Sometimes the gums become so affected as to occasion them to fall out, and the teeth themselves are frequently rendered carious by causes which have not hitherto been satisfactorily explained. The disease usually begins on that side of the tooth which is not exposed to pressure, and gradually advances till an opening is made into the cavity: as soon as the cavity is exposed, the tooth becomes liable to considerable pain, from the air coming into contact with the nerve. Besides these accidental means by which the teeth are occasionally affected, old age seldom fails to bring with it sure and natural causes for their removal. The alveoli fill up, and the teeth consequently fall out. The gums then no longer meet in the fore part of the mouth, the chin projects forwards, and the face being rendered much shorter, the whole physiognomy appears considerably altered. Having thus described the formation, structure, growth, and decay of the teeth, it remains to speak of their uses; the chief of which we know to be in mastication. And here we cannot help observing the great variety in the structure of the human teeth, which fits us for such a variety of food, and which, when compared with the teeth given to other animals, ma-

in some measure enable us to explain the nature of the aliment for which man is intended by Nature. Thus, in ruminant animals, we find incisores only in the lower jaw, for cutting the grass, and molares for grinding it; in graminivorous animals, we see molares alone; and in carnivorous animals, canine teeth for catching at their prey, and incisores and molares for cutting and dividing it. But, as man is not designed to catch and kill his prey with his teeth, we observe that our canini are shaped differently from the fangs of beasts of prey, in whom we find them either longer than the rest of the teeth, or curved. The incisores likewise are sharper in those animals than in man. Nor are the molares in the human subject similar to the molares of carnivorous animals; they are flatter in man than in these animals; and, in the latter, we likewise find them sharper at the edges, more calculated to cut and tear the food, and, by their greater strength, capable of breaking the bones of animals. From these circumstances, therefore, we may consider man as partaking of the nature of these different classes; as approaching more to the carnivorous than to the herbivorous tribe of animals; but upon the whole, formed for a mixed aliment, and fitted equally to live upon flesh and upon vegetables. Those philosophers, therefore, who would confine a man wholly to a vegetable food, do not seem to have studied nature. As the molares are the last teeth that are formed, so they are usually the first that fall out; this would seem to prove, that we require the same kind of aliment in old age as in infancy. Besides the use of the teeth in mastication, they likewise serve a secondary purpose, by assisting in the articulation of the voice.

Teething. See *Dentition* and *Teeth*.

TEGULA HIBERNICA. See *Lapis Hibernicus*.

TEGUMENTS, COMMON. Under this term anatomists comprehend the cuticle, rete mucosum, skin, and adipose membrane, as being the covering to every part of the body except the nails. See *Skin*.

TELA. A web of cloth. The cellular membrane is so called from its likeness to a fine web.

TELA CELLULOSA. See *Cellular membrane*.

TELEPHIUM. (Because it heals old ulcers, such as that of Telephus, made by Ulysses.) See *Faba crassa*.

TELLURIUM. A very scarce metal of a tin white colour, and a high metallic lustre, found in nature alloyed with gold, silver, and lead, in the aurum paradoxicum and sylvanite.

TEMPERAMENTUM. (From *tempero*, to mix together.) The peculiar constitution of the humours. Temperaments have been variously distinguished: the di-

vision most generally received is into the sanguinous, phlegmatic, choleric, and melancholic.

TEMPLE. The lateral and flat parts of the head above the ears.

TEMPORALIS ARTERIA. The temporal artery. A branch of the external carotid, which runs on the temples and give off the frontal artery.

TEMPORAL BONES. *Ossa temporalia.* *Ossa temporum.* These two bones, which are situated one on each side of the head, are of a very irregular figure. They are usually divided into two parts, one of which from the manner of its connexion with the neighbouring bones, is called *os squamosum*, and the other *os petrosum*, from its irregularity and hardness.

In both these parts there are processes and cavities to be described. Externally there are three processes; one anterior, called *zygomatic process*, which is stretched forwards to join with the *os malæ*, and thus forms the bony jugum under which the temporal muscle passes; one posterior, called the *mastoid* or *maxillary process*, from its resemblance to a nipple; and one inferior, called the *styloid process*, from its shape, which is said to resemble that of the ancient *stylus scriptorius*. In young subjects this process is united with the bone by an intermediate cartilage, which sometimes, even in adults, is not completely ossified. Three muscles have their origin from this process, and borrow half of their names from it, viz. *stylo-glossus*, *stylo-hyoidens*, and *stylo-pharyngeus*. Round the root of this process there is a particular rising of the *os petrosum*, which some writers describe as a process, and, from its appearance with the *styloid*, have named it *vaginalis*, others describe the semi-circular ridge of the *meatus auditorius externus* as a fifth process, to which they give the name of *auditory*. The depressions and cavities are, 1. A large fossa, which serves for the articulation of the lower jaw; it is situated between the *zygomatic auditory*, and *vaginal processes*, and is separated in its middle by a fissure into which the ligament that secures the articulation of the lower jaw with this bone is fixed. The fore part of this cavity, which receives the condyle of the jaw, is covered with cartilage; the back part only with the periosteum. 2. A long fossa behind the *mastoid process*, where the *digastric muscle* has its origin. 3. The *meatus auditorius externus*, the name given to a large funnel-like canal that leads to the organ of hearing. 4. The *stylo-mastoid hole*, so called from its situation between the *styloid* and *mastoid processes*. It is likewise called the *aqueduct of Fallopius*, and affords a passage to the *portio dura* of the auditory or seventh pair of nerves. 5. Below and on the fore part of the last fora-

men we observe part of the jugular fossa, a thimble-like cavity, in which the beginning of the internal jugular vein is lodged: 6. Before, and a little above this fossa is the orifice of a foramen, through which pass the internal carotid artery and two filaments of the intercostal nerve. This conduit runs first upward and then forward, forming a kind of elbow, and terminates at the end of the os petrosum. 7. At this part of the ossa temporum we observe the orifice of a canal which runs outwards and backwards in an horizontal direction, till it terminates in the cavity of the ear called tympanum. This canal, which in the recent subject is continued from the ear to the mouth, is called the *Eustachian tube*. We shall speak of it more particularly hereafter. 8 A small hole behind the mastoid process, which serves for the transmission of a vein to the lateral sinus. But this, like other foramina in the skull that serve only for the transmission of vessels, is neither uniform in its situation, nor to be met with in every subject. The internal surface of these bones may easily be divided into three parts. The first, uppermost and largest, is the squamous part, which is slightly concave from the impression of the brain. Its semicircular edges is sloping, so that the external lamella of the bone advances farther than the internal, and thus rests more securely on the parietal bones. The second and middlemost, which is the petrous part of the bone, forms a hard, craggy protuberance, nearly of a triangular shape. On its posterior side we observe a large foramen, which is the meatus auditorius internus; it receives the double nerve of the seventh pair, viz the portio dura and portio mollis of that pair. About the middle of its anterior surface is a small foramen, which opens into the aqueduct of Fallopius, and receives a twig of the portio dura of the seventh pair of nerves. This foramen having been first described by Fallopius, and by him named *hiatus*, is sometimes called *hiatus Fallopii*. Besides these, we observe other smaller holes for the transmission of blood-vessels and nerves. Below this craggy protuberance is the third part, which, from its shape and connexion with the os occipitis by means of the lambdoidal suture, may be called the lambdoidal angle of the temporal bone. It is concave from the impression of the brain; it helps to form the posterior and inferior fossae of the skull, and has a considerable furrow, in which is lodged part of the lateral sinus. The temporal bones differ a little in their structure from the other bones of the cranium. At their upper parts they are very thin, and almost without diploë, but below they have great strength and thickness. In the *foetus*, the thin upper part, and the lower craggy part, are separated by a cartilagi-

nous substance; there is no appearance either of the mastoid or styloid processes, and, instead of a long funnel-like meatus auditorius externus, there is only a smooth bony ring, within which the membrana tympani is fastened. Within the petrous part of these bones there are several cavities, processes, and bones, which belong altogether to the ear, do not enter into the formation of the cranium, and are described under the article Ear. The ossa temporum are connected by suture with the ossa parietalia, the os occipitis, the ossa malarum, and the os sphenoides, and are articulated with the lower jaw.

TEMPORALIS (*Temporalis* sc. *mass-culus*.) *Arcardi-temporo-maxillaris* of Dumas. This muscle, which Winslow has named the *crotophyte*, arises fleshy from the lower, lateral, and anterior part of the parietal bone; from all the squamous portion of the temporal bone; from the lower and lateral part of the os frontis; from the posterior surface of the os malæ; from all the temporal process of the sphenoid bone; and sometimes from a ridge at the lower part of this process. This latter portion, however, is often common to this muscle and the pterygoideus externus. It is of a semicircular shape, and its radiated fibres converge, so as to form a strong middle tendon, which passes under the jugum, and is inserted into the coronoid process of the lower jaw, to which it adheres on every side, but more particularly at its fore part, where the insertion is continued down to the body of the bone. This muscle is covered by a pretty strong fascia, which some writers have erroneously described as a part of the aponeurosis of the occipito-frontalia. This fascia adheres to the bones, round the whole circumference of the origin of the muscle, and, descending over it, is fixed below to the ridge where the zygomatic process begins, just above the meatus auditorius; to the upper edge of the zygomatic process itself, and anteriorly to the os malæ. This fascia serves as a defence to the muscle, and likewise gives origin to some of its fleshy fibres. The principal use of the temporal muscle is to draw the lower jaw upwards, as in the action of biting; and as it passes a little forwards to its insertion, it may at the same time pull the condyle a little backwards, though not so much as it would have done if its fibres had passed in a direct line from their origin to their insertion, because the posterior and lower part of the muscle passes over the root of the zygomatic process, as over a pulley.

TENDO ACHILLIS. See *Achillis tendo*.

TENDON. (*Tendo*, from *tendo*, to stretch.) The white and glistening extremity of a muscle. See *Muscle*.

TENESMUS. (*From tango*, to con-

strings; so called from the perception of a continual constriction or bound state of the part.) A continual inclination to go to stool, without a discharge.

TENSOR. (From *tendo*, to stretch.) A muscle whose office is to extend the part to which it is fixed.

TENSOR PALATI. See *Circumflexus*.

TENSOR TYMPANI. *Internus auris* of Douglas and Cowper. *Internus mallei* of Winslow, and *sulpingo-malleus* of Dumas. A muscle of the ear, which pulls the malleus and the membrane of the tympanum towards the petreous portion of the temporal bone; by which the membrana tympani is made more concave and tense.

TENSOR VAGINÆ FEMORIS. *Fascialis.* *Membranaceus* of Douglas. *Membranus vel fascia lata* of Cowper, and *llo aponeurosi-femoral* of Dumas. *Musculus aponeurosis*, vel *fascia lata* of Winslow. A muscle, situated on the outside of the thigh, which stretches the membranous fascia of the thigh, assists in the abduction of the thigh, and somewhat in its rotation inwards. It arises by a narrow, tendinous, and fleshy beginning from the external part of the anterior, superior, spinous process of the ilium, and is inserted a little below the great trochanter into the membranous fascia.

TENT. A roll of lint for dilating openings, sinuses, &c. See *Spongia preparata*.

TENTORIUM. A process of the dura mater, separating the cerebrum from the cerebellum. It extends from the internal horizontal spine of the occipital bone, directly forwards to the sella turcica of the sphenoid bone.

TEREBELLA. (Dim. of *terebra*, a piercer or gimblet.) A trepan or instrument for sawing out circular portions of the skull. A trephine.

TEREBINTHINA. (From *resiniferus*, the turpentine-tree.) Turpentine, the produce of pine-trees.

TEREBINTHINA ARGENTAROTENSIS. Strasburg turpentine. This species is generally more transparent and less tenacious than either the Venice or Chio turpentine. It is of a yellowish brown colour, and of a more agreeable smell than any of the turpentine, except the Chio. It is extracted in several parts of Germany, from the red and silver fir, by cutting out, successively, narrow strips of the bark. In some places a resinous juice is collected from under the bark called *Cuchrym abregna*, and *oleum abietinum*.

TEREBINTHINA CANADENSIS. Canada turpentine. A production of the *pinus balsamea*. See *Balsamum Canadense*.

TEREBINTHINA CHIA. Cyprus turpentine. The resin obtained from the *pisacia terebinthus*. See *Chio turpentine*.

TEREBINTHINA COMMUNIS. See *Terebinthina vulgaris*.

TEREBINTHINA CYPRICA. See *Chio turpentine*.

TEREBINTHINA VENETA. Venice turpentine; so called because we are supplied with it from the Venetians. This species of turpentine issues spontaneously through the bark of the *Pinus larix*; *foliis fasciculatis mollibus obtusiusculis bracteis extra squamam strobilorum extantibus*. Hort. Kew: It is usually thinner than any of the other sorts; of a clear whitish or pale yellowish colour; a hot, pungent, bitterish, disagreeable taste; and a strong smell, without any thing of the aromatic flavour of the chian kind. For its virtues see *Turpentine*.

TEREBINTHINA VULGARIS. Common turpentine. The liquid resin of the *pinus sylvestris*. See *Turpentine*.

TEREBINTHINÆ OLEUM. The oil distilled from the liquid resin of the *pinus sylvestris*.

TERES. Round, smooth. 1. The name of some muscles and ligaments.

2. The name of the *ascaris lumbricoides*, or round worm, which infests the intestines.

Teres ligamentum. The ligament at the bottom of the socket of the hip joint.

TERES MAJOR. (*Teres* sc. *Musculus major*. *Teres*, round, smooth.) Eriolanus, who was the first that distinguished this and the other muscles of the scapula by particular appellations, gave the name of *teres* to this and the following muscle, on account of their long and round shape. *Anguli-scapulae*. *Humeral*, of Dumas. This muscle, which is longer and thicker than the *teres minor*, is situated along the inferior costa of the scapula, and is in part covered by the *deltoides*.

It arises fleshy from the outer surface of the inferior angle of the scapula, (where it covers some part of the *infra spinatus* and *teres minor*, with both which its fibres intermix,) and likewise from the lower and posterior half of the inferior costa of the scapula. Ascending obliquely towards the *os humeri*, it passes under the long head of the *triceps brachii*, and then becomes thinner and flatter to form a thin tendon of about an inch in breadth, and somewhat more in length, which runs immediately behind that of the *latissimus dorsi*, and is inserted along with it into the ridge at the inner side of the groove that lodges the long head of the *biceps*. These two tendons are included in a common capsula, besides which the tendon of this muscle adheres to the *os humeri*, by two other capsulae which we find placed one above the other.

This muscle assists in the rotatory motion of the arm, and likewise in drawing it downwards and backwards; so that we may consider it as the congener of the *latissimus dorsi*.

TERES MINOR. *Margininus scapulo-trochiterion* of Dumas. This muscle seems to

have been first described by Fallopius. The *teres minor* is a thin fleshy muscle, situated along the inferior edge of the *infra-spinatus*, and is in part covered by the posterior part of the *deltoideus*.

It arises fleshy from all the convex edge of the inferior costa of the scapula; from thence it ascends obliquely upwards and forwards, and terminates in a flat tendon, which adheres to the lower and posterior part of the capsular ligament of the joint, and is inserted into the lower part of the great tuberosity of the *os humeri*, a little below the termination of the *infra-spinatus*.

The tendinous membrane, which is continued from the *infra-spinatus*, and spread over the *teres minor*, likewise forms a thin septum between the two muscles. In some subjects, however, they are so closely united, as to be with difficulty separated from each other. Some of the fibres of the *teres minor* are intermixed with those of the *teres major* and *subscapularis*.

The uses of this muscle are similar to those of the *infra-spinatus*.

TERRA LEMNIA. Earth of Lemnos. See *Bolo*.

TERRA LIVONICA. See *Bolo*.

TERRA MERITA. The curcuma or turmeric root is sometimes so called.

TERRA MORTUA. See *Terra diuinata*.

TERRA PONDEROSA SALITA. See *Murias baryte*.

TERRA SIGILLATA. See *Bolo*.

TERRÆ OLEUM. See *Petroleum*.

TERRÆ ABSORBENTIA. Absorbent earths, distinguishable from other earthy and stony substances by their solubility in acids, as chalk, crabs, claws, oyster-shells, egg-shells, pearl, coral, &c.

TETTERA. (From *τεττω*, a crane.) The middle and lateral parts of the neck.

Tertian ague. See *Febris intermittens*.

TERTIANA DUPLEX. A tertian fever that returns every day; but the paroxysms are unequal, every other fit being alike.

TERTIANA DUPLICATA. A tertian fever returning every other day; but there are two paroxysms in one day.

TERTIANA FEBRIS. See *Febris intermittens*.

TERTIANA TRIPLEX. A tertian fever returning every day; every other day there are two paroxysms, and but one in the intermediate one.

TERTIANARIA. (From *tertiana*, a species of intermittent fever which is said to be cured by this plant.) The plant which is thus called in some pharmacopœias is the *Scutellaria galericulata foliis cordato lanceolatis, cr-natis; floribus axillaribus* of Linnaeus, which is common in the hedges and ditches of this country. It has a bitter taste and a garlic smell, and is said to be serviceable against that species of ague which attacks the patient every third day.

TERTIUM SAL. (From *tertius*, third.)

A neutral salt as being the product of an acid, and an alkali making a third body different from either.

TERRERA. (From *terrera*, four.) A four-square bone. The cuboid bone.

TESTA PROBATHIX. (*Quasi testa*, from *terreo*, to burn.) A cupel or test. A pot for separating baser metals from gold and silver.

TESTUDO. (From *testa*, a shell; because it is covered with a shell.) A tortoise, a snail. An ulcer, which, like a snail, creeps under the skin.

TESTES CEREBRI. See *Tubercula quæ drigemina*.

TESTICLE. *Testes*. *Orchis*. Also called *dytidium*, and by some *perin*. Two little oval bodies situated within the scrotum, and covered by a strong, white and dense coat, called *tunica albuginea testis*. Each testicle is composed of small vessels, bent in a serpentine direction, arising from the spermatic artery, and convoluted into little heaps, separated from one another by cellular partitions. In each partition there is a duct receiving semen from the small vessels; and all the ducts constitute a net which is attached to the *tunica albuginea*. From this net-work twenty or more vessels arise, all of which are variously contorted, and, being reflected, ascend to the posterior margin of the testis, where they unite into one common duct, bent into serpentine windings, and forming a hard body called the *epididymis*. The spermatic arteries are branches of the aorta. The spermatic veins empty themselves into the vena cava and emulgent vein. The nerves of the testicle are branches of the lumbar and great intercostal nerve. The use of the testicle is to secrete the semen.

Testicle, swelled See *Orchitis*.

TESTICULUS. (*Testiculus*, dim. of *testis*.) A small testicle. Also the orchis plant, so named from the resemblance of its roots to a testicle.

TESTICULUS CANINUS. See *Satanion*.

TESTIS. (A witness, the testes being the witnesses of our manhood.) See *Testicle*.

TETANOMATA. (From *τετανω*, to smooth.) *Tetanothra*. Medicines which smooth the skin, and remove wrinkles.

TETANUS. (From *τενω*, to stretch.) Spasm with rigidity. *Convulsio indica*. *Holotonicos*. *Rigor nervosus*. A genus of disease in the class *neuroses* and order *spasmi* of Cullen; characterised by a spasmodic rigidity of almost the whole body. The varieties of tetanus are, 1. *Opiathotonos*, where the body is thrown back by spasmodic contractions of the muscles. 2. *Empirodotonos*, the body being bent forwards. 3. *Trismus*, the locked jaw. Tetanus is often symptomatic of syphilis and worms.

These affections arise more frequently in warm climates than in cold ones, and are very apt to occur when much rain or mois-

ture quickly succeeds excessive dry and sultry weather. They attack persons of all ages, sexes, temperaments, and complexions, but the male sex more frequently than the female, and those of a robust and vigorous constitution than those of a weak habit. An idea is entertained by many, Dr. Thomas observes, that negroes are more predisposed to attacks of tetanus than white people; they certainly are more frequently affected with it, but this circumstance does not arise from any constitutional predisposition, but from their being more exposed to punctures and wounds in the feet, by nails, splinters of wood, pieces of broken glass, &c. from usually going bare-footed.

Tetanic affections are occasioned either by exposures to cold, or by some irritation of the nerves, in consequence of local injury by puncture, incision, or laceration. Lacerated wounds of tendinous parts prove, in warm climates, a never-failing source of these complaints. In cold climates, as well as in warm ones, the locked jaw, or trismus, frequently arises in consequence of the amputation of a limb.

When the disease has arisen in consequence of a puncture, or any other external injury, the symptoms shew themselves generally about the eighth day, but when it proceeds from an exposure to cold, they generally make their appearance much sooner.

In some instances it comes on suddenly, and with great violence; but it more usually makes its attack in a gradual manner; in which case, a slight stiffness is at first perceived in the back part of the neck, which, after a short time, becomes considerably increased, and at length renders the motion of the head both difficult and painful.

With the rigidity of the head there is likewise an uneasy sensation at the root of the tongue, together with some difficulty in swallowing, and a great tightness is perceived about the chest, with a pain at the extremity of the sternum, shooting into the back. A stiffness also takes place in the jaws, which soon increases to such a height, that the teeth become so closely set together as not to admit of the smallest opening. This is what is termed the locked jaw.

In some cases, the spasmodic affection extends no farther. In others the spasm at this stage of the disease, returning with great frequency become likewise more general, and now affect not only the muscles of the neck and jaws, but likewise those of the whole spine, so as to bend the trunk of the body very forcibly backwards, and this is what is named opisthotonos. Where the body is bent forwards the disease is called emprosthotonos.

During the whole course of the disorder, the abdominal muscles are violently affected with spasm, so that the belly is strongly

retracted, and feels very hard, most obstinate costiveness prevails, and both the flexor and extensor muscles of the lower extremities are commonly affected at the same time, so as to keep the limbs rigidly extended.

The flexors of the head and trunk become at length so strongly affected, as to balance the action of the extensors, and to keep the head and trunk so rigidly extended and straight as to render it incapable of being moved in any direction. The arms which were little affected before, are now likewise rigidly extended, the tongue also becomes affected with spasm, and being convulsively darted out, is often much injured by the teeth at that moment snapping together. It is to this state of the disease that the term tetanus has been strictly applied.

The disorder continuing to advance, every organ of voluntary motion becomes affected; the eyes are rigid and immoveable in their sockets, the countenance is hideously distorted, and expresses great distress; the strength is exhausted, the pulse becomes irregular, and one universal spasm puts a period to a most miserable state of existence.

Attacks of tetanus are seldom attended with any fever, but always with violent pain, and the spasms do not continue for a constancy, but the muscles admit of some remission in their contraction, which is renewed every ten or fifteen minutes, especially if the patient makes the least attempt to speak, drink, or alter his position.

When tetanic affections arise in consequence of a wound, puncture, or laceration, in warm climates, Dr. Thomas observes, they are almost sure to prove fatal. The locked jaw, in consequence of an amputation, likewise proves usually fatal. When these affections are produced by an exposure to cold, they may in most cases be removed by a timely use of proper remedies, although a considerable space will probably elapse before the patient will be able to recover his former strength.

On dissections of this disease, slight effusions within the cranium have been observed in a few instances; but in by far the greater number, nothing particular has been discovered, either in the brain, or any other organ.

TETARTAEUS. (*Tetartaeus*, fourth.) A quartan fever.

TETRAMYTRUM. (From *τετρας*, four, and *μυτρον*, an ointment.) An ointment of four ingredients.

TETRANGURIA. (From *τετρας*, four, and *αγρος*, a cup; so called because its fruit resembles a cup divided into four parts.) The citrill.

TETRAPHARMACUM. (From *τετρας*, four, and *φαρμακον*, a drug.) A medicine composed of four ingredients.

Tettors. See *Herpes*.

TEUCRIUM. (From *Teucer*, who discovered it.) The name of a genus of plants in the Linnæan system. Class, *Didynamia*. Order, *Verticillata*. The herb speedwell.

TEUCRIUM CAPITATUM. Th. systematic name of the poley mountain of Montpellier. See *Polium montanum*.

TEUCRIUM CHAMÆDRYS. The systematic name of the common germander. See *Chamedrys*.

TEUCRIUM CHAMÆPITYS. The systematic name of the ground pine. See *Chamæpitys*.

TEUCRIUM CRITICUM. The systematic name of the poley mountain of Candy. See *Polium criticum*.

TEUCRIUM MARUM. The systematic name of the Syrian herb mastich. See *Marum Syriacum*.

TEUCRIUM MONTANUM. The systematic name of the common poley mountain. See *Polium montanum*.

TEUCRIUM POLICUM. The systematic name of the golden poley mountain. See *Polium montanum*.

TEUCRIUM SCORDIUM. The systematic name of the water germander. See *Scordium*.

TERETRUM. (From *tereo*, to pierce.) The trepan.

TERNIALIA BENZOÏN. The Benjamin gum-tree.

TERMINTHUS. (From *τερμινθος*, the turpentine-tree.) *Albatis* Black and ardent pustules, mostly attacking the legs of females; so called from their resemblance to the fruit of the turpentine-tree.

TERNARY. Consisting of the number three, which some chemical and mystical writers have made strange work with; but the most remarkable distinction of this kind, and the only one worth notice, is that of Hippocrates, who divides the parts of a human body into continentes, contenta, and impetum facientes, though the latter is resolvable into the mechanism of the two former, rather than any thing distinct in itself.

TERRA. Earth, as distinguished from minerals and metals and precious stones.

TERRA CARIOSA. Rotten bone, a species of non-effervescent chalk, of a brown colour.

TERRA CATECHU. See *Catechu*.

TERRA DAMNATA. *Terra mortua*. Condemned earth, is the remainder after some distillations, where all that will rise is drawn off; the same as *Cuput mortuum*.

TERRA FOLIATA TARTARI. The acetate of potash.

TERRA JAPONICA. *Japan earth*. *Catechu vulgo*. *Terra Japonica*. *Cachou*. *Fauvel*. *Catchu*. *Caschu*. *Catechu*. *Catchu*. *Cashew*. *Caitchu*. *Castjoe*. *Cachu*. *Cate*. *Kuath*. The natives call it *Cutt*, the English who reside there *Cutch*. It

was called *Japan earth*, because it was long supposed to be an earthy substance from Japan. It is the inspissated juice of a species of *mimosa*, which grows in great abundance in the kingdom of Bahar. It is prepared from a decoction of the inner part of the wood. From the negligent method in which it is dried in little kilns dug for that purpose, it acquires the earthy appearance it in general has, from which circumstance it takes its name. In the kingdom of Bahar, besides being much used in medicine, it is employed for many purposes in arts, particularly for painting the beams of houses, to defend them from vermin. See *Catechu*.

TEUTHION. (Τευθιον) The herb polium.

THALAMI NERVORUM OPTICORUM (Θαλαμῶν, a bed.) Two bodies, which form in part the optic nerve, placed near to each other, in appearance white, protruding at the base of the lateral ventricles, and running in their direction inwards, a little downwards, and upwards.

THALASSOMELI. (From *θαλασσα*, the sea, and *μελι*, honey.) A medicine composed of sea-water and honey.

THALICTRUM. (From *θαλλω*, to flourish.) 1. The name of a genus of plants in the Linnæan system. Class, *Polyumbria*. Order, *Polyginia*.

2. The pharmacopœial name of what is also called *Rhabarbarum pauperum*. The root of this plant, *Thalictrum flavum* of Linnæus, is said to be aperient and stomachic, and to come very near in its virtues to rhubarb. It is a common plant in this country, but seldom used medicinally.

THALICTRUM FLAVUM. The systematic name of the poor man's rhubarb. See *Thalictrum*.

THAPSIA. (From *Thapsus*, the island where it was found.) The deadly carrot. *Thapsia asclepias* of Linnæus. The root operates violently both upwards and downwards, and is not used in the present practice.

THAPSUS. (From the island *Thapsus*.) The great white mullein, or cows lung-wort.

THEA. Tea. There are two species of this tree; viz. 1. The bohea, or black tea; and 2. The viridis, or green tea; both of which are natives of China or Japan where they attain the height of five or six feet.

Great pains are taken in collecting the leaves singly, at three different times, viz. about the middle of February, in the beginning of March, and in April. Although some writers assert, that they are first exposed to the steam of boiling water, and then dried on copper-plates; yet it is now understood that such leaves are simply dried on iron plates, suspended over a fire, till they become dry and shrivelled; when

cool, they are packed in tin boxes to exclude the air, and in that state exported to Europe.

Teas are divided in Britain into three kinds of *green*, and five of *bohea*. The former class includes, 1. *Imperial* or *bloom tea*, having a large leaf, a faint smell, and being of a light green colour. 2. *Hyson*, which has small curled leaves, of a green shade inclining to blue. 3. *Single tea*, thus termed from the place where it is cultivated. The *boheas* comprehend: 1. *Souchong*, which, on infusion, imparts a yellowish green colour. 2. *Camho*, a fine tea, emitting a fragrant violet smell, and yielding a pale shade; it receives its name from the province where it is reared. 3. *Pekoe tea* is known by the small white flowers that are mixed with it. 4. *Congo* has a larger leaf than the preceding variety, and yields a deeper tint to water; and 5. *Common bohea*, the leaves of which are of an uniform green colour. There are besides other kinds of tea, sold under the names of *gunpowder tea*, &c. which differ from the preceding, only in the minuteness of their leaves, and being dried with additional care.

Much has been said and written on the medicinal properties of tea; in its natural state it is a *narcotic* plant, on which account the Chinese refrain from its use till it has been divested of this property by keeping it at least for twelve months. If, however, good tea be drunk in moderate quantities, with sufficient milk and sugar, it invigorates the system, and produces a temporary exhilaration; but when taken too copiously, it is apt to occasion weakness, tremor, palsies, and various other symptoms arising from narcotic plants, while it contributes to aggravate hysterical and hypochondriacal complaints. Tea has also been supposed to possess considerable diuretic and sudorific virtues, which, however, depend more on the quantity of warm water employed as a vehicle, than the quality of the tea itself. Lastly, as infusions of these leaves are the safest refreshment after undergoing great bodily fatigue or mental exertion; they afford an agreeable beverage to those who are exposed to cold weather; at the same time tending to support and promote perspiration, which is otherwise liable to be impeded.

THEA GERMANICA. Fluellin; male speed-well. See *Veronica*.

THEBAICA. (*A. Thebaide regions*, from *Thebe*, where it flourished.) The Egyptian poppy.

THERESII FORAMINA. The orifices of veins in the cavities of the heart.

THECA VERTEBRALIS. (*Theca*, from *τὸ θῆμα*, to place.) The vertebral canal.

THELIPTEIRIS. (From *θηλες*, female, and *πτερος*, fern.) The female fern.

THENAR. (*Thenar*, sc. *musculus*.) See *Flexor brevis pollicis manus*.

THEOBROMA CACAO. (*Theobroma*, from *theos*, the gods, and *βρομα*, food; so called from the deliciousness of its fruit.) Cacao Indian. The systematic name of the tree which affords cocoa and chocolate.

THEODORICUM. (From *theos*, the gods, and *doron*, a gift.) The pompous name of some antidotes.

THERAPEIA. (From *θεραπεω*, to heal.) *Therapia*. The art of healing diseases.

THERAPEUTICS. (*Therapeutica*, from *θεραπεω*, to cure.) *Therapia*. *Methodus medendi*. That branch of medicine which treats of the operation of the different means employed for obviating diseases, and of the application of these means.

THERIACA. (From *θηρ*, a viper or venomous wild beast.) Treacle, or molasses: also a medicine appropriated to the cure of the bites of venomous animals, or to resist poisons.

THERIACA ANDROMACHI. The Venice or Mithridate treacle; a composition of sixty-one ingredients, prepared, pulverised, and with honey formed into an electuary.

THERIACA CELESTIS. Liquid laudanum.

THERIACA COMMUNIS. Common treacle, or molasses.

THERIACA DAMOCRATIS. An old preparation usually called *Confectio Damocratis*.

THERIACA EDINENSIS. Edinburgh theriaca. The thebaic electuary.

THERIACA JERMANORUM. A rob of juniper-berries.

THERIACA LONDINENSIS. A cataplasm of cummin seed, bay-berries, germander, snake-root, cloves and honey.

THERIACA RUSTICORUM. The roots of the common garlic were so called. See *Allium*.

THERIONIA. (From *θησιον*, to rage like a wild beast.) A malignant ulcer.

THERMUM. Warm baths or springs. See *Medicinal waters*.

THERMOMETER. (*Thermometrum*, from *θερμη*, heat, and *μετρον*, to measure.) An instrument for measuring the degrees of heat. See *Caloric*.

Thigh-bone. See *Femer*.

THIRST. *Sitis*. The sensation by which we experience a desire to drink. The seat of this sensation appears to be either in the fauces or the stomach.

Thistle, carline. See *Chamelon album*.

Thistle, holy. See *Carduus benedictus*.

Thistle, pine. See *Curlina gummifera*.

THLASPI. (From *θλαω*, to break, because its seed appears as if it were broken or bruised.) 1. The name of a genus

of plants in the Linnæan system. Class, *Tetradynamia*. Order, *Siliculosæ*.

2. The pharmaceutical name of the herb penny-cress. Two species of *thlaspi* are directed in some pharmacopæias for medicinal uses;—the *Thlaspi arvense* of Linnaeus, or treacle mustard, and *Thlaspi campestre* of Linnaeus, or mithridate mustard. The seeds of both have an acrid biting taste approaching to that of common mustard, with which they agree nearly in their pharmaceutical qualities. They have also an unpleasant flavour, somewhat of the garlic or onion kind.

THLASPI ARVENSE. The systematic name of the treacle mustard. See *Thlaspi*.

THLASPI CAMPESTRE. The systematic name of the mithridate mustard. See *Thlaspi*.

THORACIC DUCT. *Ductus thoracicus*. *Ductus Pecquetii*. The trunk of the absorbents; of a serpentine form, and about the diameter of a crow-quill. It lies upon the dorsal vertebrae, between the aorta and vena azygos, and extends from the posterior opening of the diaphragm to the angle formed by the union of the subclavian and jugular veins, into which it opens and evacuates its contents. In this course the thoracic duct receives the absorbent vessels from every part of the body.

THORAX. (From *θυρεα*, to leap, because in it the heart leaps.) The chest. That part of the body situated between the neck and the abdomen. The external parts of the thorax are, the common integuments, the breasts, various muscles, and the bones of the thorax. (See *Bones* and *Respiration*.) The parts within the cavity of the thorax are, the pleura and its productions, the lungs, heart, thymus gland, œsophagus, thoracic duct, arch of the aorta, part of the vena cava, vena azygos, the eighth pair of nerves, and part of the great intercostal nerve.

Thorn, Egyptian. The *Mimosa nilotica* of Linnaeus. See *Arabic gum*.

Thorn-apple. See *Stramonium*.

Thorn, black. See *Prunus sylvestris*.

THROMBUS. (From *θρομβω*, to disturb.) A small tumour which sometimes arises after bleeding, from the blood escaping from the vein into the cellular structure surrounding it.

Thrush. See *Aphthæ*.

THRYPTICA. (From *θρυπτεω*, to break.) Lithontriptics, medicines which are said to have the power of destroying stones in the bladder.

TAURIS CORTEX. The cascarilla and eleuthera barks, were so called. See *Cascarilla*.

TAUS. (From *θυσια*, to sacrifice, so called from its great use in sacrifices.) See *Olibanum*.

THUS JUDÆORUM. See *Thymiana*.

THUS MASCULUM. See *Olibanum*.

TRUTA OCCIDENTALIS. (*Thuja*, from *θεω*, odour, so named from its fragrant smell.) The systematic name of the tree of life. See *Arbor vitæ*.

TRYLACITIS. (From *θυλαξ*, a seed-vessel, so called from its large head.) The white garden poppy.

TRYMBRA. (From *θυμ*, thyme; so named because it smells like thyme.) See *Satureja*.

TRYMBRA HISPANICA. The name given by Tournefort to the common herb mastic. See *Marum vulgare*.

Thyme, lemon. See *Sorphyllum citratum*.

Thyme, mother of. See *Sorphyllum*.

TRYHELOA. (From *θυμα*, an odour, because of its smell.) *Oncoron*. Spargelax. Flax-leaved Daphne. This plant, *Daphne genkwa*: *panicula terminali foliis linearibus lanceolatis acuminate* of Linnaeus, affords the *garcin bark*, which very much resembles that of our mezereum. Garcin bark is to be immersed in vinegar for about an hour before it is wanted, a small piece, the size of a sixpence, thus steeped, is applied to the arm or any other part, and renewed once a day in winter and twice in summer. It produces a serous exudation from the skin without irritating or blistering. It is recommended, and is in frequent use in France and Russia against some diseases of the eyes.

TRYMIAMA. (From *θυμα*, an odour, so called from its odiferous smell.) Musk-wood. *Thus judæorum*. A bark in small brownish grey pieces, intermixed with bits of leaves, seeming as if the bark and leaves had been bruised and pressed together, brought from Syria, Sicilia, &c. and supposed to be the produce of the liquid storax tree. This bark has an agreeable balsamic smell, approaching to that of liquid storax, and a sub-acrid bitterish taste accompanied with some slight adstringency.

TRYMIUM. (From *θυμ*, thyme, because it is of the colour of thyme.) A small wart upon the skin.

TRYMOXALIS. (From *θυμ*, thyme, *οξύς*, acid, and *αλς*, salt.) A composition of thyme, vinegar and salt.

THYMUS. (*ἄρο τε θυμ*, because it was used in faintings; or from *θυμα*, an odour, because of its fragrant smell.) 1. A gland of considerable size in the fœtus, situated in the anterior duplicature or space of the mediastinum, under the superior part of the sternum. An excretory duct has not yet been detected, but lymphatic vessels have been seen going from it to the thoracic duct. Its use is unknown.

2. A small indolent carnosous tubercle like a wart arising about the anus, or the pudenda, resembling the flowers of thyme, from whence it takes its name.

3. The name of a genus of plants in the Linnæan system. Class, *Didymamia*. Order, *Gymnospermia*. Thyme.

4. The pharmacopœial name of the common thyme. This herb, the *Thymus vulgaris* of Linnaeus;—*erectus foliis revolutis ovatis, floribus verticillato spicatis*, has an agreeable aromatic smell, and a warm pungent taste. Its virtues are said to be resolvent, emmenagogue, tonic, and stomachic; yet there is no disease mentioned in which its use is particularly recommended by any writer on the materia medica.

THYMUS CITRATUS. See *Serpyllum citratum*.

THYMUS CRETICUS. The plant which bears this name in some pharmacopœias is the *Satureja capitata* of Linnaeus, which possesses similar virtues to our thyme, but in a stronger degree.

THYMUS MASTICHINA. The systematic name of the common herb mastich. See *Marum vulgare*.

THYMUS SERPYLLUM. The systematic name of the mother of thyme. See *Serpyllum*.

THYMUS VULGARIS. The systematic name of the common thyme. See *Thymus*.

THYRO. Names compounded with this word belong to muscles which are attached to the thyroid cartilage; as,

THYRO-ARYTENOIDEUS. (*Musculus thyro-arytenoideus*.) A muscle situated about the glottis, which pulls the arytenoid cartilage forwards nearer to the middle of the thyroid, and consequently shortens and relaxes the ligament of the larynx.

THYRO-HYOIDEUS. *Musculus thyro-hyoideus*.) A muscle, situated between the os hyoides and trunk, which pulls the os hyoides downwards, and the thyroid cartilage upwards.

THYRO-PHARYNGÆUS. See *Constrictor pharyngis inferior*.

THYRO-PHARYNGO-STAPHILINUS. See *Palato-pharyngeus*.

THYRO-STAPHILINUS. See *Palato-pharyngeus*.

THYROID CARTILAGE. (*Cartilago thyroidea*, from *thyros*, a shield, and *adoc*, resemblance, from its supposed resemblance to a shield.) Scutiform cartilage. The cartilage which is placed perpendicular to the cricoid cartilages of the larynx, constituting the anterior, superior, and largest part of the larynx. It is harder and more prominent in men than in women, in whom it forms the *pomum adami*.

THYROID GLAND. *Glandula thyroidea*. A large gland situated upon the cricoid cartilage, trachea, and horns of the thyroid cartilage. It is uncertain whether it be conglobate or conglomerate. Its

excretory duct has never been detected, and its use is not yet known.

TIBIA. (*Tibia*, the hautboy, qu. *tubia*, from *tuba*, a tube; so called from its pipe-like shape.) *Focile majus*; *Arundo major*; *Fosilius*; and, from its resemblance to an old musical instrument, *Canna major*; *Canna-domestica cruris*. The largest bone of the leg. It is of a long, thick and triangular shape, and is situated on the internal part of the leg. Its upper extremity is large, and flattened at its summit, where we observe two articulating surfaces, a little concave, and separated from each other by an intermediate irregular protuberance. Of these two cavities, the internal one is deepest, and of an oblong shape, while the external one is rounded, and more superficial. Each of these, in the recent subject, is covered by a cartilage, which extends to the intermediate protuberance, where it terminates. These two little cavities receive the condyles of the os femoris, and the eminence between them is admitted into the cavity which is seen between the two condyles of that bone; so that this articulation affords a specimen of the complete ginglymus. Behind the intermediate protuberance, or tubercle, is a pretty deep depression, which serves for the attachment of a ligament, and likewise to separate the two cavities from each other. Under the edge of the external cavity is a circular flat surface, covered with cartilage, which serves for the articulation of the fibula; and at the fore part of the bone is a considerable tuberosity, of an inch and a half in length, to which the strong ligament of the rotula is fixed.

The body of the tibia is smaller than its extremities, and, being of a triangular shape, affords three surfaces. Of these, the external one is broad, and slightly hollowed by muscles above and below; the internal surface is broad and flat, and the posterior surface is narrower than the other two, and nearly cylindrical. This last has a slight ridge running obliquely across it, from the outer side of the upper end of the bone to about one-third of its length downwards. A little below this we observe a passage for the medullary vessels, which is pretty considerable, and slants obliquely downwards. Of the three angles which separate these surfaces, the anterior one, from its sharpness, is called the *spine*, or *shin*. This ridge is not straight, but describes a figure like an Italic *f*, turning first inwards, then outwards, and lastly inwards again. The external angle is more rounded and serves for the attachment of the interosseous ligament; and the internal one is more rounded still by the pressure of muscles.

The tibia enlarges again a little at its

lower extremity, and terminates in a pretty deep cavity, by which it is articulated with the uppermost bone of the foot. This cavity, in the recent subject, is lined with cartilage. Its internal side is formed into a considerable process, called *malleolus internus*, which, in its situation, resembles the styloid process of the radius. This process is broad, and of considerable thickness, and from it ligaments are extended to the foot. At its back part we find a groove, lined with a thin layer of cartilage, in which slide the tendons of the flexor digitorum longus, and of the tibialis posterior; and a little behind this is a smaller groove, for the tendon of the flexor longus pollicis. On the side opposite to the malleolus internus, the cavity is interrupted, and immediately above it is a rough triangular depression, which is furnished with cartilage, and receives the lower end of the fibula.

The whole of this lower extremity of the bone seems to be turned somewhat outwards, so that the malleolus internus is situated more forwards than the inner border of the upper extremity of the bone.

In the fœtus both ends of the tibia are cartilaginous, and become afterwards epiphyses.

TIBIAL ARTERIES. *Arteria tibiales.* The two principal branches of the popliteal artery: the one proceeds forwards, and is called the anterior tibial; the other backwards, and is called the posterior tibial; of which the external tibial, the fibular, the external and internal plantar, and the plantar arch, are branches.

TIBIALIS ANTICUS. (*Musculus tibialis anticus.*) *Tibio-sus-metatarsien* of Dumas. A flexor muscle of the foot, situated on the leg, which bends the foot by drawing it upwards and at the same time turns the toes inwards.

TIBIALIS GRACILIS. See *Plantaris*.

TIBIALIS POSTICUS (*Musculus tibialis posticus.*) *Tibio-tarsien* of Dumas. A flexor muscle of the foot, situated on the leg, which extends the foot, and turns the toes inwards.

TIC DOLOUREUX A painful affection of the nerves of the face, particularly of that branch of the fifth pair, which comes out of the infraorbital foramen.

TIGLIA GRANA. *Grana tili.* *Grana sigli* The grana tilia are seeds of a dark grey colour, the produce of *Troton tylium* of Linnæus, in shape very like the seed of the *ricinus communis*. They abound with an oil which is far more acrid and purgative than castor-oil.

TILBURY WATER. This is found at West Tilbury in Essex. It is an aperient and chalybeate now seldom used medicinally.

TILIA. (*Phœn, ulmus, the elm-tree.*) 1. The name of a genus of plants in the Linnæan system. Class, *Polyandria*. Order, *Monogynia*.

2. The pharmacopœial name of the lime or linden tree. *Tilia Europæa* of Linnæus. The flowers of this tree are supposed to possess anodyne and antispasmodic virtues. They have a moderately strong smell, in which their virtue seems to consist, and abound with a strong mucilage. They are in high esteem in France.

TILIA EUROPEA. The systematic name of the lime-tree. See *Tilia*.

TILIA GRANA. See *Tiglio Grana*.

TILMUS. (From *tyllan*, to pluck.) Floccitation or picking of bed-clothes, observable in the last stages of some disorders.

TIMAC. The name of a root imported from the East Indies, which is said to possess diuretic virtues, and therefore exhibited in dropsies. It is not known from what plant it is obtained.

TIN. *Stannum.* Jupiter of the alchemists. It has been much doubted whether this metal is found native. In the opinion of Kirwan there are sufficient authorities to determine the question in the affirmative. The native oxid of tin, or tin stone, occurs both massive and crystallized. Its colour is a dark brown, sometimes yellowish grey. When crystallized, it is somewhat transparent. The wood tin ore is a variety of the native oxid, termed so from its fibrous texture. This variety has hitherto been found only in Cornwall. It occurs in fragments which are generally round, and its colour is brown, sometimes inclining to yellow. Tin is also found mineralized by sulphur, associated always with a portion of copper, and often of iron. This ore is called *tin pyrites*. Its colour is yellowish grey. It has a metallic lustre, and a fibrous or lamellated texture; sometimes it exhibits prismatic colours. Tin is comparatively a rare metal, as it is not found in great quantity any where but in Cornwall or Devonshire; though it is likewise met with in the mines of Bohemia, Saxony, the island of Banca, the peninsula of Malacca, and in the East Indies.

Properties of tin.—Tin is of a brilliant white colour, though not quite so white as silver. It is one of the lightest of the metals; its specific gravity, when hammered, being 7.299. It is very fusible, melting at about 410° Fahr. By intense heat it is volatilized. It becomes oxidated by a moderate degree of heat. It easily bends and emits a noise, called the crackling of tin. It is exceedingly soft and ductile. It may be reduced to very thin leaves. *Tin foil*, or tin leaf, which is tin beat out, is about one-thousandth part of an inch thick. It has scarcely any sound.

It resists the action of the air. It crystallizes in rhombs formed of small octahedra. It unites, by fusion, with phosphorus and sulphur. It does not decompose water alone, or in the cold; but easily by means of many other bodies. It decomposes the sulphuric acid, and unites with the sulphureous acid. It decomposes the nitric acid, and is very soluble in muriatic acid. Nitromuriatic acid acts on it very readily. It is slightly soluble in the fluoric and boracic acids. Phosphoric acid attacks it by heat. Tartareous acid has no perceptible action on it. The oxalic and acetic acids dissolve it in small quantities. The alkalis appear to have some action on it by heat. It combines with sulphuret of potash by fusion. It unites with the greater number of the metals with facility. It decomposes muriate of mercury, and muriate of ammonia. It causes nitrate of potash to detonate. It takes fire spontaneously in oxygenated muriatic acid gas, and is capable of combining with two different proportions of oxygen.

Method of obtaining tin.—In order to obtain tin from its ore, a mere fusion of it, with pounded charcoal, is sufficient. Or, a certain quantity of the ore is first freed from sulphur and arsenic by torrefaction; after which it is mixed with equal parts of potash, one half of common resin, and two parts of black flux; the mixture is then fused in a crucible covered with charcoal, by means of which the metal is recovered in its metallic state. In order to obtain pure tin, let the tin obtained before be rapidly dissolved in strong nitric acid with heat. Thus some of the metals it may contain will be held in solution, and others become oxidated, but muriatic, or nitromuriatic acid will, on digestion, take up those oxids, and after sufficient abluion, leave that of tin, which may afterwards be reduced by mixing it when pulverised with double its weight of a flux formed of equal parts of pitch and borax, or resin and borax, and putting it into a covered crucible, lined with charcoal, which must be placed in a forge, and strongly heated for a quarter of an hour.

TINCÆ OS. (*Tinca*, a tench.) The mouth of the uterus is so called by some writers, from its resemblance to a tench's mouth.

TENCAL. See *Borax*.

TINCTORIUS. (From *tingo*, to die.) An epithet of a species of broom used by dyers: the *genista tinctoria* of Linnæus.

TINCTURA. (From *tingo*, to dye a tinctura.) A solution of any substance in spirit of wine. Rectified spirit of wine is the direct menstruum of the resins, and essential oils of vegetables, and totally extracts these active principles from sundry vegetable matters, which yield them to water not at all, or only in part. It

dissolves likewise the sweet saccharine matter of vegetables, and generally those parts of animal bodies in which their peculiar smell and taste reside.

The virtues of many vegetables are extracted almost equally by water and rectified spirit; but in the watery and spirituous tinctures of them there is this difference, that the active parts in the watery extractions are blended with a large proportion of inert gummy matter, on which their solubility in this menstruum in a great measure depends, while rectified spirit extracts them almost pure from gum. Hence, when the spirituous tinctures are mixed with watery liquors, a part of what the spirit had taken up from the subject generally separates and subsides, on account of its having been freed from that matter, which, being blended with it in the original vegetable, made it soluble in water. This, however, is not universal, for active parts of some vegetables when extracted by rectified spirits, are not precipitated by water, being almost soluble in both menstrua.

Rectified spirit may be tinged by vegetables of all colours, except blue: the leaves of plants, in general, will give out little of their natural colour to watery liquors, communicate to spirit the whole of their green tincture, which for the most part proves elegant, though not very durable.

Fixed alkaline salts deepen the colour of spirituous tinctures; and hence they have been supposed to promote the dissolving power of the menstruum, though this does not appear from experience; in the trials which have been made no more was found to be taken up in the deep-coloured tinctures than in the paler ones, and often not so much; if the alkali be added after the extraction of the tincture, it will heighten the colour as much as when mixed with the ingredients at first. The addition of these salts in making tinctures is not only needless but prejudicial, as they generally injure the flavour of aromatics, and superadd a quality sometimes contrary to the intention of the medicine.

Volatile alkaline salts, in many cases, promote the action of the spirits. Acids generally weaken it; unless when the acid has been previously combined with the vinous spirit into a compound of new qualities, called dulcified spirit.

TINCTURA ALOES. Tincture of aloë. "Take of the inspissated juice of spike aloë, half an ounce; extract of liquorice, an ounce and half; water, a pint; rectified spirit, four fluidounces." Macerate in a sand bath until the extract is dissolved, and then strain. This preparation possesses stomachic and purgative qualities, but should never be given where there is a tendency to hæmorrhoids. In choleric

cases and amenorrhœa it is preferred to other purges. The dose is from half a fluidounce to an ounce.

TINCTURA ALOES COMPOSITA. Compound tincture of aloe, formerly called *Elisir aloes*: *Elisir proprietas*. "Take of extract of spiked aloe, powdered, saffron, of each three ounces; tincture of myrrh, two pints." Macerate for fourteen days, and strain. A more stimulating compound than the former. It is a useful application to old indolent ulcers. The dose is from half a fluidrachm to two drachms.

TINCTURA ALOES VITRIOLATA. With the bitter infusion a drachm or two of this elegant tincture is extremely serviceable against gouty and rheumatic affections of the stomach and bowels, and also in the weaknesses of those organs which frequently attend old age.

TINCTURA ASSAFETIDÆ. Tincture of assafetida, formerly known by the name *tinctura fetida*. "Take of assafetida, four ounces; rectified spirit, two pints." Macerate for fourteen days and strain. Diluted with water, this is mostly given in all kinds of fits, by the vulgar. It is a useful preparation as an antispasmodic, especially in conjunction with vitriolated zinc. The dose is from half a fluidrachm to two drachms.

TINCTURA AURANTII. Tincture of orange-peel, formerly *tinctura corticis aurantii*. "Take of fresh orange-peel, three ounces; proof spirit, two pints." Macerate for fourteen days, and strain. A mild and pleasant stomachic bitter.

TINCTURA BENZOINI COMPOSITA. Compound tincture of benzoin, formerly known by the names of *tinctura benzoës composita*, and *balsamum traumaticum*. "Take of benzoin, three ounces; storax balsam, strained, two ounces; balsam of Tolu, an ounce; extract of spiked aloe, an ounce and half; rectified spirit, two pints." Macerate for fourteen days, and strain. This tincture is more generally applied externally to ulcers and wounds than given internally, though possessing expectorant, antispasmodic, and stimulating powers. Against coughs, spasmodic affections of the stomach and bowels, and diarrhœa, produced by ulcerations of those parts, it is a very excellent medicine. The dose when given internally is from half a fluidrachm to two drachms diluted.

TINCTURA CALUMBÆ. Tincture of calumba, formerly called *tinctura columbæ*. "Take of calumba root, powdered, two ounces and a half; proof spirit, two pints." Macerate for fourteen days, and strain. This tincture contains the active part of the root, and is generally given with the infusion of it, as a stomachic and adstringent.

TINCTURA CAMPHORÆ COMPOSITA. Compound tincture of camphor, formerly called *tinctura opis camphorata*; and *elixer pæregoricum*. "Take of camphor, two scruples; opium, dried and powdered, benzoic acid, of each a drachm; proof spirit, two pints." Macerate for fourteen days, and strain. The London college has changed the name of this preparation, because it was occasionally the source of mistakes under its old one, and tincture of opium was sometimes substituted for it. It differs also from the former preparation in the omission of the oil of aniseed, which was often complained of as disagreeable to the palate, and to which, as an addition, no increase of power could be affixed. The dose is from half a fluidrachm to half an ounce.

TINCTURA CANTHARIDIS. See *Tinctura Lyttæ*.

TINCTURA CAPSICI. Tincture of capicum. "Take of capsicum berries, an ounce; proof spirit, two pints." Macerate for fourteen days, and strain.

TINCTURA CARDAMOMI. Tincture of cardamom. "Take of cardamom-seeds, bruised, three ounces; proof spirit, two pints." Macerate for fourteen days, and strain. A powerful stimulating carminative. In spasm of the stomach, an ounce with some other diluted stimulant is given with advantage. The dose may vary according to circumstances, from half a drachm to an ounce and upwards.

TINCTURA CARDAMOMI COMPOSITA. Compound tincture of cardamom, formerly called *tinctura stomachica*. "Take of cardamom-seeds, carraway-seeds, cochineal, of each bruised, two drachms; cinnamon-bark, bruised, half an ounce; raisins, stoned, four ounces; proof spirit, two pints." Macerate for fourteen days, and strain. A useful and elegant carminative and cordial. The dose from half a fluidrachm to half an ounce and upwards.

TINCTURA CASCARILLÆ. Tincture of cascarrilla. "Take of cascarrilla-bark, powdered, four ounces; proof spirit, two pints." Macerate for fourteen days, and strain. A stimulating aromatic tonic, that may be exhibited in debility of the bowels and stomach, and in those cases of fever in which the Peruvian bark proves purgative. The dose from half a drachm to two drachms.

TINCTURA CASTOREI. Tincture of castor. "Take of castor, powdered, two ounces; rectified spirit, two pints." Macerate for fourteen days, and strain. A powerful stimulant and antispasmodic, mostly exhibited in hysterical affections in a dilute form. The dose is from half a fluidrachm to two drachms.

TINCTURA CATECHU. Tincture of catechu, formerly known by the name *tinctura japonica*. "Take of extract of catechu, cinnamon-bark, bruised, of each two ounces; proof spirit, two pints." Macerate for fourteen days, and strain. An aromatic adstringent, mostly given to the hysterical epilepsy of young women, which arises from irregularity in the menstrual discharge. The dose is from half a fluidrachm to two drachms.

TINCTURA CINCHONÆ. Tincture of cinchona. Formerly known by the name of *tinctura corticis peruviani simplex*. "Take of lance-leaved cinchona bark, powdered, seven ounces; proof spirit, two pints." Macerate for fourteen days, and strain. The dose is from a drachm to half a fluidounce. For its virtues see *Cinchona*.

TINCTURA CINCHONÆ COMPOSITA. Compound tincture of cinchona. "Take of lance-leaved cinchona bark, powdered, two ounces; orange-peel, dried, an ounce and half; serpentry-root, bruised, three drachms; saffron, a drachm; cochineal, powdered, two scruples; proof spirit, twenty fluidounces." Macerate for fourteen days, and strain. The dose is from one fluidrachm to half an ounce. For its virtues see *Cinchona*.

TINCTURA CINNAMOMI. Tincture of cinnamon. Formerly called *agua cinnamomi fortis*. "Take of cinnamon bark, bruised, three ounces; proof spirit, two pints." Macerate for fourteen days. The dose is from a fluidrachm to three drachms or more.

TINCTURA CINNAMOMI COMPOSITA. Compound tincture of cinnamon. Formerly called *tinctura aromatica*. "Take of cinnamon bark, bruised, six drachms; cardamom-seeds, bruised, three drachms; long pepper, powdered, ginger-root, sliced, of each two drachms; proof spirit, two pints." Macerate for fourteen days, and strain. The dose is from half a fluidrachm to two drachms or more.

TINCTURA DIGITALIS. Tincture of fox-glove. "Take of fox-glove leaves, dried, four ounces; proof spirit, two pints." Macerate for fourteen days, and strain. This tincture is introduced in the London Pharmacopœia as possessing the properties of the plant in a convenient, uniform, and permanent form; it is a saturated tincture, and in the same proportions has been long used in general practice. The dose is from ten to forty minims. For its virtues see *Digitalis*.

TINCTURA FERRI AMMONIATÆ. Tincture of ammoniated iron, formerly called *tinctura ferri ammoniacalis: tinctura storum martialis, tinctura muris mynaichti*. "Take of ammoniated iron, four ounces; proof spirit, a pint." Digest and strain. This is a most excellent chalybeate in all atonic

affections, and may be given with cinchona in the cure of dropsical and other cachectic diseases. The dose is from half a fluidrachm to two drachms.

TINCTURA FERRI MURIATIS. Tincture of muriate of iron. Formerly called *tinctura martis in spiritu salis: tinctura martis cum spiritu salis*: and lately known by the name of *tinctura ferri muriati*. "Take of subcarbonate of iron, half a pound; muriatic acid, a pint; rectified spirit, three pints." Pour the acid upon the carbonate of iron in a glass vessel, and shake it occasionally for three days. Set it by, that the fæces, if there be any, may subside; then pour off the solution, and add the spirit. Mr. Cline strongly recommends this 'in ischuria and many diseases of the kidneys and urinary passages. The dose is from ten to twenty drops. It is a good chalybeate, and serviceable against most diseases of debility without fever.

TINCTURA GENTIANÆ COMPOSITA. Compound tincture of gentian. Formerly called *tinctura amara*. "Take of gentian root, sliced, two ounces; orange-peel, dried, an ounce; cardamom-seeds, bruised, half an ounce; proof spirit, two pints." Macerate for fourteen days, and strain. The dose is from one fluidrachm to two drachms. For its virtues see *Gentiana*.

TINCTURA GUAIACI. Tincture of guaiacum. "Take of guaiacum gum resin, half a pound; rectified spirit, two pints." Macerate for fourteen days, and strain. This tincture, which possesses all the active parts of this peculiar vegetable matter, is now first introduced into the London Pharmacopœia. The dose is from one fluidrachm to two drachms. For its virtues see *Guaiacum*.

TINCTURA GUAIACI AMMONIATÆ. Ammoniated tincture of guaiacum. Formerly called *tinctura guaiacina volatilis*. "Take of guaiacum gum resin, powdered, four ounces; compound spirit of ammonia, a pint and half." Macerate for fourteen days, and strain. The dose is from one fluidrachm to two drachms.

TINCTURA HELLEBORI NIGRI. Tincture of black hellebore. Formerly called *tinctura melampodii*. "Take of black hellebore-root, sliced, four ounces; proof spirit, two pints." Macerate for fourteen days, and strain. The dose is from half a fluidrachm to a drachm. For its virtues consult *Helleborus niger*.

TINCTURA HUMULI. Tincture of hop. "Take of hops, five ounces; proof spirit, two pints." Macerate for fourteen days, and strain. Various modifications of the preparations of this bitter have lately been strongly recommended by Mr. Freke, (Observations on Humulus Lupulus,) and employed by many practitioners, who believe that it unites sedative and tonic powers,

and thus forms a useful combination. The dose is from half a fluidrachm to a drachm. See *Humulus*.

TINCTURA HYOSCIAMI. Tincture of henbane. "Take of henbane leaves, dried, four ounces; proof spirit, two pints." Macerate for fourteen days, and strain. That the henbane itself is narcotic is abundantly proved, that the same power is also found in its tincture is also certain, but to produce the same effects requires a much larger dose. In some of the statements made to the College of Physicians of London a different opinion has been given, and twenty-five drops have been considered as equivalent to twenty of tincture of opium; it does not produce costiveness, or the subsequent confusion of head which follows the use of opium, and will therefore be, even if its powers be weaker, of considerable use. The dose is from ten minims to one fluidrachm.

TINCTURA JALAPE. Tincture of jalap. Formerly called *tinctura jalapii*. "Take of jalap-root, powdered, eight ounces; proof spirit, two pints." Macerate for fourteen days, and strain. The dose is from one fluidrachm to half an ounce. For its virtues see *Jalapa*.

TINCTURA KINO. Tincture of kino. "Take of kino, powdered, three ounces; proof spirit, two pints." Macerate for fourteen days, and strain. All the adstringency of kino is included in this preparation. The dose is from half a fluidrachm to two drachms. See *Kino*.

TINCTURA LYTTE. Tincture of blistering fly. Formerly called *tinctura cantharidis*:—*tinctura cantharidum*. "Take of blistering flies, bruised, three drachms; proof spirit, two pints." Macerate for fourteen days, and strain. In the last edition of the London Pharmacopeia the colouring matter of the former preparation is omitted as useless, and that of the fly increased. It is a very acrid, diuretic, and stimulating preparation, which should always be administered with great caution from its known action on the parts of generation. In chronic eruptions on the skin, and dropsical diseases of the aged, it is often very useful when other medicines have been inert. The dose is from half a fluidrachm to two drachms.

TINCTURA MYRRHÆ. Tincture of myrrh. "Take of myrrh, bruised, three ounces; rectified spirit, twenty-two fluidounces; water, a pint and half." Macerate for fourteen days, and strain. The dose is from half a fluidrachm to a drachm. For its virtues see *Myrrha*.

TINCTURA OPII. Tincture of opium. "Take of hard opium, powdered, two ounces and a half; proof spirit, two pints." Macerate for fourteen days, and strain. The dose is from ten minims, or twenty

drops, to half a fluidrachm. For its virtues see *Opium*.

TINCTURA RHEI. Tincture of rhubarb. Formerly known by the names of *tinctura rhabarbari*, and *tinctura rhabarbari spiritosa*. "Take of rhubarb-root, sliced, two ounces; cardamom-seeds, bruised, an ounce and half; saffron, two drachms; proof spirit, two pints." Macerate for fourteen days, and strain. The dose is from half a fluidounce to an ounce and half. For its virtues see *Rhabarbarum*.

TINCTURA RHEI COMPOSITA. Compound tincture of rhubarb. Formerly called *tinctura rhabarbari composita*. "Take of rhubarb-root, sliced, two ounces; liquorice-root, bruised, half an ounce; ginger-root, sliced, saffron, of each two drachms; water, a pint; proof spirit, twelve fluidounces." Macerate for fourteen days, and strain. This is a mild stomachic aperient. The dose is from half a fluidounce to an ounce and half.

TINCTURA SCILLÆ. Tincture of squilla. "Take of squilla-root, fresh dried, four ounces; proof spirit, two pints." Macerate for fourteen days, and strain. The virtues of the squill (see *Scilla*) reside in this tincture, which is administered in doses of from twenty drops to a fluidrachm.

TINCTURA SENNÆ. Tincture of senna. Formerly called *elixir salutis*. "Take of senna-leaves, three ounces; caraway-seeds, three drachms; cardamom-seeds, bruised, a drachm; raisins, stoned, four ounces; proof spirit, two pints." Macerate for fourteen days, and strain. A carminative, aperient, and purgative in doses from two drachms to a fluidounce. See *Senna*.

TINCTURA SERPENTARIÆ. Tincture of serpentary. Formerly called *tinctura serpentariæ virginianæ*. "Take of serpentary root, three ounces; proof spirit, two pints." Macerate for fourteen days, and strain. This tincture possesses, in addition to the virtues of the spirit, those of the serpentaria. The dose is from half a fluidrachm to two drachms. See *Serpentaria*.

TINCTURA VALERIANÆ. Tincture of valerian. Formerly called *tinctura valerianæ simplex*. "Take of valerian-root, four ounces; proof spirit, two pints." Macerate for fourteen days, and strain. A useful antispasmodic in conjunction with others. The dose is from half a fluidrachm to two drachms. See *Valeriana*.

TINCTURA VALERIANÆ AMMONIATÆ. Ammoniated tincture of valerian. Formerly called *tinctura valerianæ volatilis*. "Take of valerian-root, four ounces; aromatic spirit of ammonia, two pints." Macerate for fourteen days, and strain. A strong antispasmodic and stimulating tincture. The dose is from half a fluidrachm to two drachms.

TINCTURA VERATRI. A very active alterative, recommended in the cure of epilepsy and cutaneous eruptions. Its administration requires great caution; the white hellebore being a powerful poison.

TINCTURA ZINGIBERIS. Tincture of ginger. "Take of ginger-root, sliced, two ounces; proof spirit, two pints." Macerate for fourteen days, and strain. A stimulating carminative. The dose is from a fluidrachm to three drachms.

Tincture of aloes. See *Tinctura aloes*.

Tincture of asafoetida. See *Tinctura asafoetida*.

Tincture of black hellebore. See *Tinctura hellebori nigri*.

Tincture of blistering fly. See *Tinctura lyttæ*.

Tincture of calumba. See *Tinctura calumbæ*.

Tincture of capsicum. See *Tinctura capsici*.

Tincture of cardamom. See *Tinctura cardamomi*.

Tincture of cascarella. See *Tinctura cascarielle*.

Tincture of castor. See *Tinctura castorei*.

Tincture of catechu. See *Tinctura catechu*.

Tincture of cinchona. See *Tinctura cinchonæ*.

Tincture of cinnamon. See *Tinctura cinnamomi*.

Tincture of fox-glove. See *Tinctura digitalis*.

Tincture of guaiacum. See *Tinctura guaiaci*.

Tincture of guaiacum, ammoniated. See *Tinctura guaiaci ammoniati*.

Tincture of ginger. See *Tinctura zinziberis*.

Tincture of henbane. See *Tinctura hyoscyami*.

Tincture of hops. See *Tinctura humuli*.

Tincture of jalap. See *Tinctura jalapii*.

Tincture of kino. See *Tinctura kino*.

Tincture of myrrh. See *Tinctura myrrhæ*.

Tincture of opium. See *Tinctura opii*.

Tincture of orange-peel. See *Tinctura aurantii*.

Tincture of rhubarb. See *Tinctura rhei*.

Tincture of senna. See *Tinctura sennæ*.

Tincture of serpentary. See *Tinctura serpentariæ*.

Tincture of squills. See *Tinctura scille*.

Tincture of valerian. See *Tinctura valerianæ*.

Tincture of valerian, ammoniated. See *Tinctura valerianæ ammoniata*.

Tincture, compound, of aloes. See *Tinctura aloes composita*.

Tincture, compound, of benzoin. See *Tinctura benzoini composita*.

Tincture, compound, of camphor. See *Tinctura camphoræ composita*.

Tincture, compound, of cardamom. See *Tinctura cardamomi composita*.

Tincture, compound, of cinnamon. See *Tinctura cinnamomi composita*.

Tincture, compound, of cinchona. See *Tinctura cinchonæ composita*.

Tincture, compound, of gentian. See *Tinctura gentianæ composita*.

Tincture, compound, of rhubarb. See *Tinctura rhubarbari composita*.

TINEA CAPITIS. (*Tinea*, from *teneo*, to hold.) The scald-head. A genus of disease in the class *locales* and order *diagnoses* of Cullen; characterised by small ulcers at the root of the hairs of the head, which produce a friable white crust.

Tinglass. See *Bismuth*.

TINNITUS AURIUM. See *Paracusia*.

TITANIUM. This is a lately discovered metal. It was first noticed by Macgregor as existing in the state of an oxid, mixed with iron, manganese, and silic, in a greyish black sand found in the vale of Menachan in Cornwall, and thence named menachanite, or oxid of titanium, combined with iron. It has since been discovered by Klaproth, in an ore named titanite, or oxid of titanium, combined with lime and silic. This ore is generally met with crystallized in four-sided prisms, not longer than a quarter of an inch. Its colour is a yellowish red, or blackish brown; it is opaque and of an imperfect lustre. It breaks with a foliated, uneven, or conchoidal fracture. It exists also in an ore called red schorl of Hungary, or red oxid of titanium. This ore, which is found generally crystallized in rectangular prisms, is of a brownish red colour; and its texture foliated. In all these ores titanium exists in the state of an oxid.

Properties of titanium.—Titanium has been only obtained in very small agglutinated grains. It is of a red-yellow and crystalline texture, brittle and extremely refractory. Its specific gravity is about 4.2; when broken with a hammer while yet hot from its recent reduction it shows a change of colours of purple, violet, and blue. In very intense heat it is volatilized. Most of the acids have a striking action on this metal. Nitric acid has little effect upon it. It is very oxidable by the muriatic acid. It is not attacked by the alkalies. Nitro-muriatic acid converts it into a white powder. Sulphuric acid when boiled upon it is partly decomposed. It is one of the most infusible metals. It does not combine with sulphur, but it may be united to phosphorus. It does not alloy with copper, lead, or arsenic, but combines with iron.

Method of obtaining titanium.—It is extremely difficult to reduce the oxid of titanium to the metallic state. However, the experiments of Klaproth, Hecht and Vauquelin, have proved its reducibility. According to the two latter, one part of oxid of titanium is to be melted with six of potash; the mass when cold is to be dissolved in water. A white precipitate will be formed which is carbonate of titanium. This carbonate is then made into a paste with oil, and the mixture is put into a crucible filled with charcoal powder and a little alumine. The whole is then exposed for a few hours to the action of a strong heat. The metallic titanium will be found in a blackish puffed-up substance, possessing a metallic appearance.

TITHYMALUS. (From *τιθος*, a dug, and *μαλός*, tender; so called from its smooth leaves and milky juice.) Spurge. Two plants are directed for medicinal purposes by this name. See *Tithymalus paralius* and *Esula minor*.

TITHYMALUS CYPARISSIUS. See *Esula minor*.

TITHYMALUS PARALIUS. Sea spurge. Every part of this plant, *Euphorbia paralius* of Linnæus, is violently cathartic and irritating, inflaming the mouth and fauces. It is seldom employed in the practice of this country; but where it is used vinegar is recommended to correct its irritating power.

TITHYMELEA. See *Thymelea*.

TITTILLICUM. (From *titillo*, to tickle; so called from its being easily tickled.) The arm-pit.

Toad-flax. See *Linaria*.

Tobacco. See *Nicotiana*.

Tobacco, English. See *Nicotiana minor*.

Tobacco, Virginian. See *Nicotiana*.

TOES. *Digiti pedis.* They consist of three distinct bones disposed in rows called phalanges, or ranks of the toes. The great toe has but two phalanges; the others have three ranks of bones, which have nothing particular, only the joints are made round and free, formed by a round head on one bone, and by a pretty deep hollow for receiving it, in the one above it.

Tolu balsam. See *Balsamum Tolutanum*.

TOLUIFERA BALSANUM. The systematic name of the tree which affords the Tolu balsam. See *Balsamum Tolutanum*.

TOLUTANUM BALSANUM. See *Balsamum Tolutanum*.

TOMEIUM. (From *τομω*, to cut.) An incision-knife.

TOMENTITIA. (From *tomentum*, a flock of wool; so called from its soft coat.) Cotton-weed.

TOMENTUM CEREBRI. (*Tomentum*, a flock of wool.) The small vessels that penetrate the cortical substance of the brain, from the pia mater, which, when

separated from the brain, and adhering to the pia mater, give it a flocky appearance.

TONIO SPASM. (*Spasmus tonicus*, *Tonus*, from *τύνω*, to pull or draw.) *Contractura a spasmo*. A rigid contraction of the muscles, without relaxation, as in trismus, tetanus, &c. See *Tetanus*.

TONICS. (*Medicamenta tonica*, from *τονω*, to strengthen.) Medicines which increase the tone of the muscular fibre; such as stimulants, adstringents, &c.

TONGUE. *Lingua.* A soft fleshy viscus, very moveable in every direction, situated inferiorly in the cavity of the mouth, and constituting the organ of taste. It is divided into a base, body, and back, an inferior surface, and two lateral parts. It is composed of muscular fibres, covered by a nervous membrane, on which are a great number of nervous papillæ, particularly at the apex and lateral parts; the rete mucosum, and epidermis. The arteries of the tongue are branches of the nine and labial. The veins empty themselves into the great lingual, which proceed to the external jugular. The nerves come from the eighth, ninth, and fifth pair. The use of this organ is for chewing, swallowing, sucking, and tasting. See also *Taste*.

TONSILS. *Tonsille.* *Amygdale Tonsiles.* *Tolles.* Two oblong, sub-oral glands, situated one on each side of the fauces, and opening into the cavity of the mouth by twelve or more large excretory ducts.

Tooth. See *Teeth*.

Tooth-ache. See *Odontalgia*.

TOPHUS (*Toph*, Hebrew.) *Ephoroma*. The concretion on the teeth or in the joints of gouty people. Also gravel.

TOPICA. (From *τοπος*, a place.) Medicines applied to a particular place.

TOPIARIA. A species of tumour in the skin of the head.

TORCULAR. (From *torqueo*, to twist.) The tourniquet; a bandage to check hæmorrhages after wounds or amputations.

TORCULAR HEROPHILI (*Torcular*, from *torqueo*, to twist.) *Lecheus Lenos*. The press of Herophilus. The place where the four sinuses of the dura mater meet together, first accurately described by Herophilus, the anatomist.

TORDYLUM OFFICINALE. (*Tordylus*, quasi *tortilium*, from *torqueo*, to twist; named from its tortuous branches.) The systematic name of the official *scellæ creticæ*. See *Scellæ creticæ*.

Tormentil. See *Tormentilla*.

TORMENTILLA. (From *tormenta*, pain; because it was supposed to relieve pain in the teeth) 1. The name of a genus of plants in the Linnæan system. Class, *Iconandria*. Order, *Monogynia*.

2. The pharmacopœial name of the *Asaphyllum*. *Consolida rubra*. *Convolvulus*.

termentil, or **upright septfoil**. *Tormentilla erecta* of Linnaeus:—*caule erectiusculo, foliis sessilibus*. The root is the only part of the plant which is used medicinally: it has a strong styptic taste, but imparts no peculiar sapid flavour: it has been long held in estimation as a powerful adstringent; and, as a proof of its efficacy in this way, it has been substituted for oak-bark in the tanning of skins for leather. Tormentil is ordered in the *pulvis creta compositus* of the London Pharmacopœia.

TORMENTILLA ERRECTA. The systematic name of the upright septfoil. See *Tormentilla*.

TORMINA. Gripes. Pains in the bowels.

TORPOR. A numbness, or deficient sensation.

TORTICOLLIS. (From *torqueo*, to twist, and *collum*, the neck.) The wry neck.

TORTURA OSSIS. The locked jaw.

TOTA BONA. See *Bonus Henricus*.

TOUCH. *Tactus*. The sensation by which we perceive any thing that is applied to the skin. The organ of touch is formed by the nervous papillæ, which are situated all over the skin, but more especially at the points of the fingers.

Touch-me not. See *Noli me tangere*.

Touch-wood. See *Agaricus*.

TOURNIQUET. (French, from *tourner*, to turn.) An instrument used for stopping the flow of blood into a limb.

TOXICARIA MACASARIENSIS. An Indian poison, obtained from a tree hitherto undescribed by any medical botanist, known by the name of Boas-upas; it is a native of South America. Concerning this plant various and almost incredible particulars have been related, both in ancient and modern times; some of them true, others probably founded on superstition. Rumphius testifies that he had not met with any other more dreadful produced from any vegetable. And he adds, that this poison, of which the Indians boast, was much more terrible to the Dutch than any warlike instrument. He likewise says, it is his opinion, that it is of the same natural order if not of the same genus as the cestrum.

TOXICODENDRUM. (From *τοξικον*, a poison, and *dendron*, a tree.) The poison-tree, which is so noxious that no insects ever come near it.

TOXICOLOGY. (*Toxicologia*, from *τοξον*, an arrow or bow; because the darts of the ancients were usually besmeared with some poisonous substance; and *λογος*, a discourse.) A dissertation on poisons. See *Poison*.

TOXICUM. (From *τοξον*, an arrow, which was sometimes poisoned.) Any deadly poison.

TOXITESIA. The artimesia or mugwort.

TRABECULA. (*Trabecula*, a small beam.) This word is mostly applied by anatomists to the small medullary fibres of the brain, which constitute the commissures.

TRACHEA. (So called from its roughness; from *τραχυς*, rough.) The wind-pipe. The trachea is a cartilaginous and membranous canal, through which the air passes into the lungs. Its upper part, which is called the larynx, is composed of five cartilages. The uppermost and smallest of these cartilages, is placed over the glottis or mouth of the larynx, and is called epiglottis, as closing the passage to the lungs in the act of swallowing. The sides of the larynx are composed of the two arytenoid cartilages, which are of a very complex figure, not easy to be described. The anterior and larger part of the larynx is made up of two cartilages, one of which is called thyroïdes or scutiformis, from its being shaped like a buckler; and the other cricoides or annularis, from its resembling a ring. Both these cartilages may be felt immediately under the skin, at the fore part of the thorax; and the thyroïdes, by its convexity, forms an eminence called the pomum adami, which is usually more considerable in the male than in the female subject.

All these cartilages are united to each other by means of very elastic ligamentous fibres; and are enabled by the assistance of their several muscles, to dilate or contract the passage of the larynx, and to perform that variety of motion which seems to point out the larynx, as the principal organ of the voice; for when the air passes out through a wound in the trachea, it produces no sound.

These cartilages are moistened by a mucus, which seems to be secreted by minute glands situated near them. The upper part of the trachea, and the cricoid and thyroid cartilages, are in some measure covered anteriorly by a considerable body, which is supposed to be of a glandular structure, and from its situation is called the thyroid gland, though its excretory duct has not yet been discovered, or its real use ascertained. The glottis is entirely covered by a very fine membrane, which is moistened by a constant supply of a watery fluid. From the larynx the canal begins to take the name of trachea or aspera arteria, and extends from thence as far down as the fourth or fifth vertebra of the back, where it divides into two branches, which are the right and left bronchial tube. Each of these bronchia ramifies through the substance of that lobe of the lungs, to which it is distributed, by an infinite number of branches, which are formed of cartilages separated from each other like those of the trachea, by an intervening membranous and ligamentary

substance. Each of these cartilages is of an angular figure; and as they become gradually less and less in their diameter, the lower ones are in some measure received into those above them, when the lungs, after being inflated, gradually collapse by the air being pushed out from them in expiration. As the branches of the bronchiz become more minute, their cartilages become more and more angular and membranous, till at length they become perfectly membranous, and at last become invisible. The trachea is furnished with fleshy or muscular fibres, some of which pass through its whole extent longitudinally, while the others are carried round it in a circular direction, so that by the contraction or relaxation of these fibres, it is enabled to shorten or lengthen itself, and likewise to dilate or contract the diameter of its passage. The trachea and its branches, in all their ramifications, are furnished with a great number of small glands which are lodged in their cellular substance, and discharge a mucous fluid on the inner surface of these tubes.

The cartilages of the trachea, by keeping it constantly open, afford a free passage to the air which we are obliged to be incessantly respiring; and its membranous part, by being capable of contraction or dilatation, enables us to receive and expel the air in a greater or less quantity, and with more or less velocity, as may be required in singing and declamation. This membranous structure of the trachea posteriorly, seems likewise to assist in the descent of the food, by preventing that impediment to its passage down the œsophagus, which might be expected, if the cartilages be complete rings. The trachea receives its arteries from the carotid and subclavian arteries, and its veins pass into the jugulars. Its nerves arise from the recurrent branch of the eighth pair, and from the cervical plexus.

TRACHELAGRA. (From *τραχάλος*, the throat, and *αγρα*, a seizure.) The gout in the neck.

TRACHELIUM. (From *τραχίλος*, the throat; so called from its efficacy in diseases of the throat.) The herb throat-wort.

TRACHELO. (From *τραχάλος*, the neck.) Names compounded of this word belong to muscles which are attached to the neck; as the

TRACHELOCELE. (From *τραχία*, the wind-pipe, and *κύη*, a tumour.) A wen or tumour upon the trachea.

TRACHELO-MASTOIDEUS. A muscle situated on the neck, which assists the complexus, but pulls the head more to one side. It is the *complexus minor seu mastoideus lateralis* of Winslow. *Trachelomastoidien* of Dumas. It arises from the transverse processes of the five inferior cervical vertebrae, where it is connected with the

transversalis cervicis and of the three superior dorsal, and it is inserted into the middle of the posterior part of the mastoid process.

TRACHELOPHRYA. (From *τραχάος*, the throat, and *φρυα*, a tumour.) A swelling of the bronchial gland.

TRACHELOS. (From *τραχός*, rough, because of the rough cartilages.) The wind-pipe.

TRACHEOTOMY. (*Tracheotomia*, from *τραχός*, rough, and *τομή*, to cut.) *Laryngotomia*. A synonym of bronchotomy. See *Bronchotomy*.

TRACHOMA. (From *τραχός*, rough.) An asperity in the internal superficies of the eyelid. The effects are a violent ophthalmia, and a severe pain, as often as the eyelid moves. The species are,

1. *Trachoma sabulosum*, from sand falling between the eye and the eyelid of persons travelling, blown by a high wind; this happens chiefly in sabulous situations, and may be prevented by spectacles for the purpose, or by guarding against the flight of sand by covering the eyes.

2. *Trachoma carunculorum*, which arises from caruncles, or fleshy verucæ, growing in the internal superficies of the eyelid. This species of the trachoma is called *morum palpebræ internæ*, because the tuberculous internal superficies appears of a livid red like a mulberry. Others call these caruncles *pladורות*.

3. *Trachoma herpeticum*, which are hard pustules in the internal superficies of the eyelids. This is also called *fyosis*, and *palpebra ficosa*, from its resemblance to granulated substances in a cut fig. With the Greeks it is nominated *atomablepharon*, or *proptoris*.

Tragacanth gum. See *Tragacantha*.

TRAGACANTHA. (From *τραγός*, a goat, and *ακανθα*, a thorn; so called from its pods resembling the goat's beard.) Goat's thorn. Milk-vetch. *Spina hirci*. *Astragalus aculeatus*. We are indebted to a French traveller of the name of Olivier for the discovery that the gum tragacanth of commerce, is the produce of a species of *astragalus* not before known. He describes it under the name of *astragalus verus*, being different both from *A. tragacantha* of Linnæus, and from the *A. gum-mifera* of Labillardiere. It grows in the North of Persia. Gum tragacanth, or gum dragant, (which is forced from this plant by the intensity of the solar rays, is concreted into irregular lumps or vermicular pieces, bent into a variety of shapes, and larger or smaller proportions, according to the size of the wound from which it issues,) differs from all other known gums in imparting to a very large quantity of water a thick and glutinous consistence. The demulcent qualities of this gum are to be considered as similar to those of gum arabic. It is seldom given alone, but frequently in

combination with more powerful medicines, especially in the form of troches, for which it is peculiarly well adapted: it gives name to an official powder, and is an ingredient in the compound powder of cerus.

TRAGICUS. (*Musculus tragicus*.) A proper muscle of the ear, which pulls the point of the tragus a little forward.

TRAGUM. (From *τραγος*, a goat; so named from its filthy smell.) Bastard dittany.

TRAGOCERUS. (From *τραγος*, a goat, and *κερας*, a horn; so named from the supposed resemblance of its leaves to the horn of a goat.) The aloë.

TRAGOPOGON. (From *τραγος*, a goat, and *πωγων*, a beard; so called because its downy seed, while enclosed in the calyx, resembles a goat's beard.) 1. The name of a genus of plants in the Linnæan system. *Class, Syngenesia. Order, Polygamia.*

2. The pharmacopœial name of the common goat's beard. The young stems of this plant, *Tragopogon pratense* of Linnæus, are eaten like asparagus, and are a pleasant and wholesome food. The root is also excellent, and was formerly used medicinally as a diuretic.

TRAGOPOGON PRATENÆ. The systematic name of the common goat's beard. See *Tragopogon*.

TRAGOPTHEUM. (From *τραγος*, a goat, and *πωγων*, wheat; so named from its beard.) Buck-wheat.

TRAGORCHIS. (From *τραγος*, a goat, and *ορχις*, a testicle; so named from the supposed resemblance of its roots to the testicles of a goat.) A species of orchis.

TRAGORISATUM. (From *τραγος*, a goat, and *οριζαντον*, marjoram; so called because goats are fond of it.) A species of wild marjoram.

TRAGOSKLINUM. (From *τραγος*, a goat, and *κλυον*, parsley; named from its hairy coat like the beard of a goat.) The burnet saxifrage was so called. See *Pimpinella*.

TRAGUS. (*Τραγος*, a goat; so called from its having numerous little hairs, or from its being hairy like the goat.) A small cartilaginous eminence of the auricle or external ear, placed anteriorly, and connected to the anterior extremity of the helix. It is beset with numerous little hairs, defending in some measure the entrance of the external auditory passage.

TRANS. (*Τρανς*.) *Raphe.* The line which divides the scrotum, and runs on to the anus.

TRANSFUSION. (*Transfusio*, from *transfundo*, to pour from one vessel into another.) The transmission of blood from one animal to another by means of a canula.

TRANSPIRATION. (*Transpiratio*, from *trans*, through, and *spiro*, to breathe.) A

synonym of perspiration. See *Perspiration*.

TRANSUDATION. The same as perspiration.

TRANSVERSALIS ABDOMINIS. A muscle situated on the anterior part of the abdomen. It arises internally or posteriorly from the cartilages of the seven lower ribs, being there connected with the intercostals and diaphragm, also from the transverse process of the last vertebra of the back from these of the four upper vertebrae of the loins, from the inner edge of the crista ilii and from part of Poupert's ligament, and it is inserted into the inferior bone of the sternum, and almost all the length of the linea alba. Its use is to support and compress the abdominal viscera.

TRANSVERSALIS ANTICUS PRINUS. See *Rectus capitis lateralis*.

TRANSVERSALIS CERVICIS. See *Longissimus dorsi*.

TRANSVERSALIS COLLI. A muscle, situated on the posterior part of the neck, which turns the neck obliquely backwards, and a little to one side.

TRANSVERSALIS DORSI. See *Multifidus spine*.

TRANSVERSALIS MAJOR COLLI. See *Longissimus dorsi*.

TRANSVERSALIS PEDIS. (*Musculus transversalis pedis*.) A muscle of the foot, which it contracts by bringing the great toe and the two outermost toes nearer each other.

TRANSVERSE SUTURE. *Sutura transversalis.* This suture runs across the face, and sinks down into the orbits, joins the bones of the skull to the bones of the face; but with so many irregularities and interruptions, that it can scarcely be recognised as a suture.

TRANSVERSO-SPINALES. See *Multifidus spine*.

TRANSVERSIS AURIS. (*Musculus transversus auris*.) A muscle of the external ear, which draws the upper part of the concha towards the helix.

TRANSVERSUS PERINÆ. (*Musculus transversus perinæ*.) A muscle of the organs of generation, which sustains and keeps the perinæum in its proper place.

TRANSVERSUS PERINÆI ALTER. *Inferior prostate* of Winslow. A small muscle occasionally found accompanying the former.

TRAPA NATANS. The systematic name of the plant which affords the nux aquatica. See *Tribulus aquaticus*.

TRAPEZIUM. (A four-sided figure; so called from its shape.) The first bone of the second row of the carpus.

TRAPEZIUS. (*Musculus trapezius*, from *τραπεζιος*, four-square; so named from its shape.) *Ocularis.* A muscle situated immediately under the integuments of the

posterior part of the neck and back. It arises, by a thick, round, and short tendon, from the lower part of a protuberance in the middle of the occipital bone backwards, and from the rough line that is extended from thence towards the mastoid process of the os temporis, and by a thin membranous tendon, which covers part of the complexus and splenius. It then runs downwards along the nape of the neck, and rises tendinous from the spinous processes of the two lowermost vertebrae of the neck, and from the spinous processes of all the vertebrae of the back, being inseparably united to its fellow, the whole length of its origin, by tendinous fibres, which, in the nape of the neck, form what is called *ligamentum colli*, or the cervical ligament. It is inserted fleshy into the broad and posterior half of the clavicle, tendinous and fleshy into one half of the acromion, and into almost all the spine of the scapula.

This muscle serves to move the scapula in different directions. Its upper descending fibres pull it obliquely upwards; its middle transverse ones pull it directly backwards; its inferior fibres, which ascend obliquely upwards, draw it obliquely downwards and backwards.

The upper part of the muscle acts upon the neck and head, the latter of which it draws backwards, and turns upon its axis. It likewise concurs with other muscles in counteracting the flexion of the head forwards.

TRAPEZOIDES OS. The second bone of the second row of the carpus; so called from its resemblance to the *trapezium* or quadrilateral geometrical figure.

TRAUMATIC. (From *τραυμα*, a wound.) Any thing relating to a wound.

Travellers, joy. See *Vitalba*.

Treacle. See *Thiaria*.

Treacle, mustard. See *Thlaspi*.

Trefail, marsh. See *Trifolium paludosum*.

TREMOR. An involuntary trembling of parts.

TREPAN. *Trephine.* An instrument used by surgeons to remove a portion of bone from the skull.

TREPHINE. See *Trepan*.

Triangularis. See *Sternocostalis* and *Depressor anguli oris*.

TRIBULUS AQUATICUS. (From *τριβη*, to vex, an instrument of war to be thrown in the way to annoy the enemy's horse: hence the name of an herb from its resemblance to this instrument.) *Nux aquatica.* The fruit of the *trapa nata* of Linnæus, of a quadrangular and somewhat oval shape, including a nut of a sweet farinaceous flavour, somewhat like that of the chesnut, which is apt to constipate the bowels, and produce disease; a poultice of these nuts

is said to be efficacious in resolving hard and indolent tumours.

TRICAUDALIS. (From *tres*, three, and *cauda*, a tail.) A muscle with three tails.

TRICEPS ADDUCTOR FEMORIS.

(*Triceps*, from *tres*, three, and *caput*, a head; having three heads.) Under this appellation are comprehended three distinct muscles. See *Adductor brevis, longus*, and *magnus femoris*.

TRICEPS AURIS. See *Refractentes auris*.

TRICEPS EXTENSOR CUBITI. This muscle, which occupies all the posterior part of the os humeri, is described as two distinct muscles by Douglas, and as three by Winslow. The upper part of its long head is covered by the deltoides: the rest of the muscle is situated immediately under the integuments.

It arises, as its name indicates, by three heads. The first, or long head, (the long head of the biceps externus, of Douglas: anconeus major, of Winslow, as it is called,) springs by a flat tendon of an inch in breadth, from the anterior extremity of the inferior costa of the scapula, near its neck, and below the origin of the *teres minor*. The second head, (the short head of the biceps externus of Douglas; anconeus externus of Winslow), arises by an acute tendinous and fleshy beginning, from the upper and outer part of the os humeri, at the bottom of its great tuberosity. The third head, (brachialis externus of Douglas; anconeus internus of Winslow), which is the shortest of the three, originates by an acute fleshy beginning, from the back part of the os humeri, behind the flat tendon of the latissimus dorsi. These three portions unite about the middle of the arm, so as to form one thick and powerful muscle, which adheres to the os humeri to within an inch of the elbow, where it begins to form a broad tendon, which after adhering to the capsular ligament of the elbow, is inserted into the upper and outer part of the olecranon, and sends off a great number of fibres, which help to form the fascia on the outer part of the fore-arm. The use of this muscle is to extend the fore-arm.

TRICHIASIS. (From *τριχ*, a hair.) *Trichia.* *Trichosis.* *Entropion.* *Dichiasis.* *Districhiasis.* *Capillitium.* *Distichia.* A disease of the eye-lashes, in which they are turned into towards the bulb of the eye.

TRICLISIVUS. (From *τριχ*, hair.) A species of fracture which appears like a hair, and is almost imperceptible.

TRICHOMA. (From *τριχ*, the hair.) The plaited hair. See *Phica polonica*.

TRICHOMANES. (From *τριχ*, hair, and *μανος*, thin, lax; so called because it resembles fine hair.) Common maiden-hair.

or spleen-wort. *Asplenium trichomanes* of Linnaeus :—*frondibus pinnatis, pinnis subrotundis crenatis*. This plant is admitted into the Edinburgh Pharmacopœia: the leaves have a mucilaginous, sweetish, subadstringent taste, without any particular flavour: they are esteemed useful in disorders of the breast, being supposed to promote the expectoration of tough phlegm, and to open obstructions of the viscera.

TRICHOMIS. See *Plica polonica*.

TRICHURIS. (From *τριχ*, a hair.) The long hair-worm. See *Worms*.

TRICUSPID VALVES. (*Valvula tricuspidæ*, from *tres*, three, and *cuspidæ*, a point; so called from their being three-pointed.) The name of the three valves situated at the origin of the aorta and pulmonary artery.

Trifol, water. See *Trifolium paludosum*.

TRIFOLIUM. (From *tres*, three, and *folium*, a leaf; so called because it has three leaves on each stalk.) The name of a genus of plants in the Linnaean system. Class, *Pentandria*. Order, *Monogynia*. Trefoil.

TRIFOLIUM ACETOSUM. The wood-sorrel was so called. See *Luzula*.

TRIFOLIUM AQUATICUM. See *Trifolium paludosum*.

TRIFOLIUM ARVENSE. Hare's-foot trefoil or lagopodium.

TRIFOLIUM AUREUM. Herb trinity; noble liver-wort.

TRIFOLIUM CABBALINUM. Melilotus.

TRIFOLIUM CÆRULUM. Sweet trefoil.

TRIFOLIUM FALCATUM. The Auricula muris. See *Pileocella*.

TRIFOLIUM FIBRINUM. See *Trifolium paludosum*.

TRIFOLIUM HEPATICUM. See *Hepatica nobilis*.

TRIFOLIUM MELILOTUS OFFICINALIS. The systematic name of the officinal melilot. See *Melilotus*.

TRIFOLIUM ODORATUM. See *Melilotus*.

TRIFOLIUM PALUDOSUM. *Trifolium aquaticum*. *Trifolium fibrinum*. *Mengyanthes*. Water-trefoil, or buck-bean. *Mengyanthes trifoliata* of Linnaeus :—*foliis ternatis*. The whole plant is so extremely bitter, that in some countries it is used as a substitute for hops, in the preparation of malt liquor.

It is sometimes employed in country places as an active eccoprotic bitter in hydropic and rheumatic affections. Cases are related of its good effects in some cutaneous diseases of the herpetic and seemingly cancerous kind.

TRIGEMINI. (*Trigeminus*, from *tres*, three, and *geminus*, double; three times double.) The fifth pair of nerves, which arise from the crura of the cerebellum, and are divided within the cavity of the cranium into three branches, viz. the *orbital*, *superior*, and *inferior maxillary*. The

orbital branch is divided into the frontal, lachrymal, and nasal nerves; the superior maxillary into the sphæno-palatine, posterior alveolar, and infra-orbital nerves; and the inferior maxillary into two branches, the internal lingual, and one more properly called the inferior maxillary.

TRIGONELLA FENUGRÆCUM. The systematic name of the fenugreek. See *Fenugreek*.

TRINITATIS HERBA. See *Hepatica nobilis*.

Trinity-herb. See *Hepatica nobilis*.

TRIPASTRUM APPELLIDIS. *Tripastrum archimædæ*. A surgical instrument for extending fractured limbs; so named because it resembled a machine invented by Apellides or Archimedes, for launching of ships, and because it was worked with three cords.

TRIQUETRA OSSICULA. (*Triquetrus*, from *tres*, three.) *Ossicula Wormiana*. The triangular-shaped bones, which are found mostly in the course of the lambdoidal suture.

TRISMUS. (From *τριζω*, to gnash.) Locked jaw. Spastic rigidity of the under jaw. *Capistrum* of Vogel. Dr. Cullen makes two species. 1. *Trismus nascentium*, attacking infants during the two first weeks from their birth. 2. *Trismus traumaticus*, attacking persons of all ages, and arising from cold or a wound. See *Tetanus*.

TRISTAGO. (*Quasi tristago*, from *tristis*, sad; because it dispels sadness.) The common germander is sometimes so called. See *Chamedrys*.

TRISTAGO PALUSTRIS. The water-germander was so called. See *Scordium*.

TRITOPHYA. (From *τριπλος*, tertian, and *φω*, of a like nature, or original.) It is an epithet of a fever much of a nature with a tertian, and taking its rise from it. Some call it a continued tertian. It is remittent or intermittent.

TRITOPHYA CAUSUS. The fever called *causus* by Hippocrates.

TRITEUS TERRAUS. The same as *Tri-teophya*.

TRITICUM. (From *tere*, to thresh from the husk.) See *Wheat*.

TRITORIUM. (From *trito*, to beat small.) A mortar. Also a glass for separating the oil from the water in distilling.

TROCAR. (Corrupted from *un trois quart*, French, a three quarters, from the three sides with which the point is made.) The name of an instrument used in tapping for the dropsy.

TROCHANTER. (From *τροχα*, to run; because the muscles inserted into them perform the office of running.) Two processes of the thigh-bone, which are distinguished into the greater and lesser. See *Femur*.

TROCHISCI ANTL. Starch lozenges are used in tickling coughs and acidities of the stomach and bowels.

TROCHISCI CACTI. These are exhibited in cardialgia, acidities of the prime viæ and diarrhoea.

TROCHISCI GLYCERYNI. A pectoral and demulcent lozenge.

TROCHISCI GLYCERYNI CUM OPIO. This lozenge possesses pectoral and anodyne qualities, but requires that the quantity be regulated, one grain being contained in a drachm.

TROCHISCI MAGNEŒI. Extremely serviceable in pyrosis and flatulent colic.

TROCHISCI NITRI. An attenuating diaphoretic, calculated to remove viscid phlegm arising from inflammatory angina.

TROCHISCI SULPHURIS. Aperient and antiscorbutic.

TROCHISUS. (Dim. of τροχός, a wheel.) A troch or round tablet. Troches and lozenges are composed of powders made up with glutinous substances into little cakes, and afterwards dried. This form is principally used for the more commodious exhibition of certain medicines, by fitting them to dissolve slowly in the mouth, so as to pass by degrees into the stomach; and hence these preparations have generally a considerable portion of sugar or other materials grateful to the palate. Some powders have likewise been reduced into troches, with a view to their preparation, though possibly for no very good reasons: for the moistening them and afterwards drying them in the air, must on this account be of greater injury, than any advantage accruing from this form can counterbalance.

General rules for making troches.

1. If the mass prove so glutinous as to stick to the fingers in making up, the hands may be anointed with any sweet or aromatic oil; or else sprinkled with powder or starch, or of liquorice, or with flour.

2. In order to thoroughly dry the troches, put them on an inverted sieve, in a shady airy place, and frequently turn them.

3. Troches are to be kept in glass vessels, or in earthen ones well glazed.

TROCHLEA. (Τροχλία, a pulley, from τροχός, to run.) A kind of cartilaginous pulley, through which the tendon of one of the muscles of the eye passes.

TROCHLEARIS. See *Obliquus superior oculi*.

TROCHLEATORES. The fourth pair of nerves are so called, because they are inserted into the musculus trochlearis of the eye. See *Pathetici*.

TROCHOIDES. (From τροχός, a wheel, and ὁμοί, resemblance.) *Axæ commissura*. A species of diarthrosis, or moveable connexion of bones, in which one bone rotates upon another; as the first cervical

vertebra upon the odontoid process of the second.

TROPEOLUM MAJUS. The systematic name of the Indian cress. See *Nasturtium indicum*.

TRUFFLE. *Lycoperdon tuber* of Linnaeus. *Tuber cibarium* of Dr. Withering. A solid fungus of a globular figure, which grows under the surface of the ground without any roots or the access of light, and attains a size from a pea to the largest potatoe. It has a rough blackish coat, and is destitute of fibres. Cooks are well acquainted with its use and qualities. It is found in woods and pastures in some parts of Kent, but is not very common in England. In France and Spain truffles are very frequent, and grow to a much larger size than they do here. In these places the peasants find it worth their while to search for them, and they train up dogs and swine for this purpose, who after they have been inured to their smell by their masters frequently placing them in their way, will readily scrape them up as they ramble the fields and woods.

TUBA EUSTACHIANA. *Tuba Aristotelica.* *Aqueductus.* *Aqueductus Fallopii.* *Moutus acicus.* *Palatinus ductus.* *Ductus auris palatinus.* The auditory tube. The Eustachian tube, so called because it was first described by Eustachius, arises in each ear from the anterior extremity of the tympanum by means of a bony semi-canal; runs forwards and inwards, at the same time becoming gradually smaller; and after perforating the petrous portion of the temporal bone terminates in a passage, partly cartilaginous and partly membranous, narrow at the beginning, but becoming gradually larger, and ending in a pouch behind the soft palate. It is through this orifice that the pituitary membrane of the nose enters the tympanum. It is always open, and affords a free passage for the air into the tympanum; hence persons hear better with their mouth open.

TUBA FALLOPIANA. *Tuba fallopi.* The Fallopian tube first described by Fallopius. The uterine tube. A canal included in two laminae of the peritonæum, which arises at each side of the fundus of the uterus, passes transversely and ends with its extremity turned downwards at the ovarium. Its use is to grasp the ovum, and convey the prolific vapour to it, and to conduct the fertilised ovum into the cavity of the uterus.

TUBERCLE. *Tuberculum.* A hard superficial tumour, circumscribed and permanent: or proceeding very slowly to supuration.

TUBEROULA QUADRIGEMINA. *Corpora quadrigemina.* *Eminentie quadrigeminae.* Four white oval tubercles of the

brain, two of which are situated on each side over the posterior orifice of the third ventricle and the aqueduct of Sylvius. The ancients called them testes and testes, from their supposed resemblance.

TUBERCULUM ANNULARE. The commencement of the medulla oblongata.

TUBERCULUM LOWERI. An eminence in the right auricle of the heart where the two veins meet; so called from Lower, who first described it.

TUBULI LACTIFERI. The ducts or tubes in the nipple, through which the milk passes.

TUMOR. A swelling.

TUMORES. (*Tumor*, from *tumeo*, to swell.) Tumours. An order in the class *locales* of Cullen's nosology, comprehending partial swellings without inflammation.

TUNBRIDGE WATER. Tunbridge Wells is a populous village in the county of Kent, which contains many chalybeate springs, all of which resemble each other very closely in their chemical properties. Two of these are chiefly used, which yield about a gallon in a minute, and therefore afford an abundant supply for the numerous invalids who yearly resort thither. The analysis of Tunbridge spring prove it to be a very pure water, as to the quantity of solid matter; and the saline contents (the iron excepted) are such as may be found in almost any water that is used as common drink. It is only as a chalybeate, and in the quantity of carbonic acid, that it differs from common water. Of this acid it contains one twenty-second of its bulk. The general operation of this chalybeate water is to increase the power of the secretory system in a gradual, uniform manner, and to impart tone and strength to all the functions; hence it is asserted to be of eminent service in irregular digestion; flatulency; in the incipient stages of those chronic disorders, which are attended with great debility; in chloris; and numerous other complaints incident to the female sex. The prescribed method of using the Tunbridge water, observes Dr. Saunders, is judicious. The whole of the quantity daily used, is taken at about two or three intervals, beginning at eight o'clock in the morning, and finishing about noon. The dose at each time varies from about one to three quarters of a pint; according to the age, sex, and general constitution of the patient, and especially the duration of the course, for it is found that these waters lose much of their effect by long habit.

TUNGSTEN. (*Tungsten*, Swed. ponderous stone.) This metal, which is never found but in combination, is by no means common. The substance known to mineralogists, under the name of tungsten, was, after some time, discovered to consist of

lime, combined with the acid of this metal. This ore is now called *tungstate of lime*, and is exceedingly scarce. It has been found in Sweden and Germany, both in masses and crystallized, of a yellowish white, or grey colour. It has a sparry appearance, is shining, of a lamellated texture, and semi-transparent. The same metallic acid is likewise found united to iron and manganese; it then forms the ore called Wolfram, or *tungstate of iron and manganese*. This ore occurs both massive and crystallized, and is found in Cornwall, Germany, France and Spain. Its colour is brownish black, and its texture foliated. It has a metallic lustre, and a lamellated texture; it is brittle and very heavy; it is found in solid masses in the state of layers interspersed with quartz. These two substances are therefore ores of the same metal.

Properties.—Tungsten appears of a steel grey colour. Its specific gravity is about 17.6. It is one of the hardest metals, but it is exceedingly brittle; and it is said to be almost as infusible as platina. Heated in the air it becomes converted into a yellow pulverulent oxid, which becomes blue by a strong heat or when exposed to light. Tungsten combines with phosphorus and sulphur, and with silver, copper, iron, lead, tin, antimony, and bismuth; but it does not unite with gold and platina. It is not attacked by sulphuric, nitric, or muriatic acids; nitro muriatic acids acts upon it very slightly. It is oxidable and acidifiable by the nitrates and super-oxigenated muriates. It colours the vitrified earths or the vitreous fluxes, of a blue or brown colour. It is not known what its action will be on water and different oxids. Its action on the alkalies is likewise unknown. It is not employed yet, but promises real utility, on account of its colouring property, as a basis for pigment, since the compounds it is said to form with vegetable colouring matters, afford colours so permanent, as not to be acted on by the most concentrated oxigenated muriatic acid, the great enemy of vegetable colours.

Methods of obtaining Tungsten.—The method of obtaining metallic tungsten is a problem in chemistry. Scheele, Bergman, and Gmelin, did not succeed in their attempts to procure it. Klaproth tried to reduce the yellow oxid of this metal with a variety of combustible substances, but without success. Messrs Ruprecht and Tondy say they have obtained this metal by using combustible substances alone; and by a mixture of combustible and alkaline matter.

The following process is recommended by Richter, an ingenious German chemist.

Let equal parts of tungstic acid and dried blood be exposed for some time to a red heat in a crucible; press the black powder which is formed, into another smaller crucible, and expose it again to a violent heat in a forge, for at least half an hour. Tungsten will then be found, according to this chemist, in its metallic state in the crucible.

TUNICA. (*A tuendo corpore*, because it defends the body.) A membrane or covering, as the coats of the eye, &c.

TUNICA ACINIFORMIS. The uvea or posterior of the lamella of the iris.

TUNICA ALBUGINEA OCULI. See *Conjunctive membrane*.

TUNICA ALBUGINEA TESTIS. See *Albuginea testis*.

TUNICA ARACHNOIDEA. See *Arachnoid membrane*.

TUNICA CELLULOSA RUTSCHII. The second coat of the intestines.

TUNICA CHOROIDEA. See *Choroid membrane*.

TUNICA CONJUNCTIVA. See *Conjunctive membrane*.

TUNICA CORNEA. See *Cornea*.

TUNICA FILAMENTOSA. The false or spongy chorion.

TUNICA RETINA. See *Retina*.

TUNICA VAGINALIS TESTIS. A continuation of the peritonæum through the inguinal ring, which loosely invests the testicle and spermatic cord.

TUNICA VILLOSA. The villous or inner folding coat of the intestines.

TUNSTAT. *Tunstas*. A salt formed by the combination of the tungstic acid, with different bases, as *tunstas of ammonia*.

Turbeth, mineral. See *Hydrargyrus vitriolatus*.

Turbeth root. See *Turpethum*.

TURBINATED BONES. (*Ossa turbinata*, from *turbo*, to sharpen at the top, shaped like a sugar-loaf.) The superior spongy portion of the ethmoid bone, and the inferior spongy bones, are so called by some writers.

TURBINATUM. The pineal gland.

Turbith. A cathartic Eastern bark; a species of cicely.

TURCICA SELLA. See *Sella turcica*.

Turneric. See *Curcuma*.

Turnhoof. A vulgar name of the ground-ivy. See *Hedera terrestis*.

Turnip. See *Rapa*.

Turnip, French. See *Rapus*.

TURPENTINE. The different turpentine employed medicinally are, the Chian or Cyprus turpentine, (see *Terebinthus vulgaris*), the common turpentine, (see *Terebinthina communis*), and the Venice turpentine, (see *Terebinthina veneta*). All these have been considered as hot, stimulating corroborants and detergents;

qualities which they possess in common. They stimulate the primæ viæ, and prove laxative; when carried into the blood-vessels they excite the whole system, and thus prove serviceable in chronic rheumatism and paralysis. Turpentine readily passes off by urine, which it imbues with a peculiar odour; also by perspiration and by exhalation from the lungs; and to these respective effects are ascribed the virtues it possesses in gravelly complaints, scurvy, and pulmonic disorders. Turpentine is much used in gleet and fluor albus, and in general with much success. The essential oil, in which the virtues of turpentine reside, is not only preferred for external use, as a rubefacient, but also internally as a diuretic and septic; the latter of which qualities it possesses in a very high degree. Formerly turpentine was much used as a digestive application to ulcers, &c. but, in the modern practice of surgery, it is almost wholly exploded.

Turpeth, mineral. See *Hydrargyrus vitriolatus*.

TURPETHUM, (from *turpeth*, Ind.) Turbeth. The cortical part of the root of a species of convolvulus, the *Convolvulus turpethum* of Linnaeus, brought from the East Indies, in oblong pieces, of a brown or ash colour on the outside and whitish within; the best is ponderous, not wrinkled, easy to break, and discovers to the eye a large quantity of resinous matter. When chewed, it at first imparts a sweetish taste, which is followed by a nauseous acrimony. It is considered as a purgative, liable to much irregularity of action.

TURPETHUM MINERALE. See *Hydrargyrus vitriolatus*.

TURUNDA. (*A terendo*, from its being rolled up.) A tent or suppository.

TUSSILAGO. (From *tussis*, a cough; because it relieves coughs.) 1. The name of a genus of plants in the Linnaean system. Class, *Syngenesia*. Order, *Polygamia superflua*.

2. The pharmacopœial name of what is also called *bechium*. *Bechium*. *Calcem equinum*. *Chamaëlie*. *Filius antipatris furfurella*. *Farfura*. *Tussilago vulgaris*. *Farfura bechium*. *Ungula caballina*. Colts-foot. *Tussilago farfura*: *scapo uniflora imbricata, foliis subcordatis angulatis denticulatis*. The sensible qualities of this plant are very inconsiderable; it has a rough mucilaginous taste, but no remarkable smell. The leaves have always been esteemed as possessing demulcent and pectoral virtues, and hence they have been exhibited in pulmonary consumptions, coughs, asthmas, and catarrhal affections. It is used as tea, or given in the way of infusion with liquorice-root or honey.

TUSSILAGO FARFARA. (*Farfara*, from *farfurus*, the white poplar; so called be-

cause its leaves resemble those of the white poplar.) The systematic name of the coltsfoot. See *Tussilago*.

TUSSILAGO PETASITES. (From *πετασος*, a hat; so named because its leaves are shaped like a hat.) The systematic name of the butter-bur. See *Petasites*.

TUSSIS. A cough. A sonorous concussion of the breast, produced by the violent and, for the most part, involuntary motion of the muscles of respiration. It is symptomatic of many diseases.

TUSSIS CONVULSIVA. See *Pertussis*.

TUSSIS EXANTHEMATICA. A cough attendant on an eruption.

TUSSIS FERINA. See *Pertussis*.

TUTIA. (Persian.) *Pompholyx Cadmia*. Tutty. A grey oxyd of zinc; it is generally formed by fusing lead, or mixed with blende, when it is incrustated in the chimneys of the furnace. Mixed with any common cerate, it is applied to the eye, in debilitated states of the conjunctive membrane.

TUTIA PREPARATA. Prepared tutty is often put into collyria, to which it imparts an adstringent virtue.

Tutty. See *Tutia*.

TYLOSIS. (From *τυλος*, a callous.) *Tylosis*. An induration or callous of the margin of the eye-lids.

TYMPANI MEMBRANA. See *Membrana tympani*.

TYMPANITES. (From *τυμπανον*, a drum; so called because the bell, is distended with wind, and sounds like a drum when struck.) *Tympany*. An elastic distention of the abdomen not readily yielding to pressure, and sounding like a drum, with costiveness and atrophy, but no fluctuation. Species: 1. *Tympanites intestinalis*, a lodgment of wind in the intestines, known by the discharge of wind giving relief. 2. *Tympanites abdominalis*, when the wind is in the cavity of the abdomen.

TYMPANUM. (*Τυμπανον*. A drum.) The drum or barrel of the ear. The hollow part of the ear in which are lodged the bones of the ear. It begins behind the membrane of the tympanum, which terminates the external auditory passage, and is surrounded by the petrous portion of the temporal bone. It terminates at the cochlea of the labyrinth, and has opening into it four foramina, viz. the orifice of the Eustachian tube and mastoid sinus, the fenestra ovalis, and rotunda. It contains the four ossicula auditus.

TYPHA. (From *τυφος*, a lake, because it grows in marshy places.) The name of a genus of plants in the Linnæan system. The cat's tail.

TYPHOMANIA. (From *τυφος*, to burn, and *μανια* delirium.) A complication of phrensy and lethargy with fever.

TYPHUS. (From *τυφος*, stupor.) A

species of continued fever, characterised by great debility, a tendency in the fluids to putrefaction and the ordinary symptoms of fever. It is to be readily distinguished from the inflammatory by the smallness of the pulse and the sudden and great debility which ensues on its first attack; and, in its more advanced stage, by the petechiæ, or purple spots, which come out on various parts of the body, and the fetid stools which are discharged; and it may be distinguished from a nervous fever by the great violence of all its symptoms on its first coming on.

The most general course that gives rise to this disease, is contagion, applied either immediately from the body of a person labouring under it, or conveyed in clothes or merchandise, &c.; but it may be occasioned by the effluvia arising from either animal or vegetable substances in a decayed or putrid state; and hence it is, that in low and marshy countries it is apt to be prevalent when intense and sultry heat quickly succeeds any great inundation. A want of proper cleanliness and confined air are likewise causes of this fever; hence it prevails in hospitals, jails, camps, and on board of ships, especially when such places are much crowded, and the strictest attention is not paid to a free ventilation and due cleanliness. A close state of the atmosphere, with damp weather, is likewise apt to give rise to putrid fever. Those of lax fibres, and who have been weakened by any previous debilitating cause, such as poor diet, long fasting, hard labour, continued want of sleep, &c. are most liable to it.

On the first coming on of the disease, the person is seized with languor, dejection of spirits, amazing depression and loss of muscular strength, universal weariness and soreness, pains in the head, back, and extremities and rigors; the eyes appear full, heavy, yellowish, and often a little inflamed; the temporal arteries throb violently, the tongue is dry and parched, respiration is commonly laborious, and interrupted with deep sighing; the breath is hot and offensive, the urine is crude and pale, the body is costive, and the pulse is usually quick, small, and hard, and now and then fluttering and unequal. Sometimes a great heat, load, and pain are felt at the pit of the stomach, and a vomiting of bilious matter ensues.

As the disease advances, the pulse increases in frequency, (beating often from 100 to 130 in a minute); there is vast debility, a great heat and dryness in the skin, oppression at the breast, with anxiety, sighing, and moaning; the thirst is greatly increased; the tongue, mouth, lips and teeth are covered over with a brown or black tenaceous fur; the speech is inarticulate, and scarcely intelligible; the pa-

ficient matters much, and delirium ensues. The fever continuing to increase still more in violence, symptoms of putrefaction shew themselves; the breath becomes highly offensive; the urine deposits a black and fetid sediment; the stools are dark, offensive, and pass off insensibly; hæmorrhages issue from the gums, nostrils, mouth, and other parts of the body; livid spots or petechiæ appear on its surface; the pulse intermits and sinks; the extremities grow cold; hiccough ensue; and death at last closes the tragic scene.

When this fever does not terminate fatally, it generally begins in cold climates to diminish about the commencement of the third week, and goes off gradually towards the end of the fourth, without any very evident crisis; but in warm climates it seldom continues above a week or ten days, if so long.

Our opinion, as to the event, is to be formed by the degree of violence in the symptoms, particularly after petechiæ, although in some instances recoveries have been effected under the most unpromising appearances. An abatement of febrile heat and thirst, a gentle moisture diffused equally over the whole surface of the body, loose stools, turbid urine, rising of the pulse, and the absence of delirium and stupor, may be regarded in a favourable

light. On the contrary, petechiæ, with dark, offensive, and involuntary discharges by urine and stool, fetid sweats, hæmorrhages, and hiccoughs, denote the almost certain dissolution of the patient.

The appearances usually perceived on dissection, are inflammations of the brain and viscera, but more particularly of the stomach and intestines, which are now and then found in a gangrenous state. In the muscular fibres there seems likewise a strong tendency to gangrene.

TYPHUS ÆGYPTIACUS. The plague of Egypt.

TYPHUS CARCERUM. The jail-fever.

TYPHUS CASTRENSIS. The camp-fever.

TYPHUS GRAVIOR. The severe species of typhus.

TYPHUS ICTERODES. Typhus with symptoms of jaundice.

TYPHUS MITIOR. The low fever.

TYPHUS NERVOSUS. The nervous fever.

TYPHUS PETECHIALIS. Typhus with purple spots.

TYRIASIS. *Tyriasis.* A species of leprosy in which the skin may be easily withdrawn from the flesh.

TYROSIS. (From *tyros*, to coagulate.) A disorder of the stomach from milk curdled in it.

U

ULCER. (*Ulcus*, from *ulcer*, a sore.)

A purulent solution of continuity of the soft parts of an animal body. Ulcers may arise from a variety of causes, as all those which produce inflammation, from wounds, specific irritations of the absorbents, from scurvy, cancer, the venereal or scrofulous virus, &c. The proximate or immediate cause is an increased action of the absorbents, and a specific action of the arteries, by which a fluid is separated from the blood upon the ulcerated surface. They are variously denominated; the following is the most frequent division: 1. *The simple ulcer*, which takes place generally from a superficial wound. 2. *The sinuous*, that runs under the integuments, and the orifice of which is narrow, but not callous. 3. *The fistulous ulcer*, or *fistula*, a deep ulcer, with a narrow and callous orifice. 4. *The fungous ulcer*, the surface of which is covered with fungous flesh. 5. *The gangrenous*, which is livid, fetid, and gangrenous. 6. *The scorbutic*, which depends on a scorbutic acrimony. 7. *The venereal*, arising from the venereal disease. 8. *The cancer-*

ous ulcer, or open cancer, (see *Cancer*.)

9. *The carious ulcer*, depending upon a carious bone. 10. *The inveterate ulcer*, which is of long continuance, and resists the ordinary applications. 11. *The scrofulous ulcer*, known by its having arisen from indolent tumours, its discharging a viscid, glary matter, and its indolent nature.

Ulcerated sore throat. See *Cynanche*.

ULMARIA. (From *ulmus*, the elm; so named because it has leaves like the elm.) *Regina prati.* *Barbara capræ.* Meadow-sweet. Queen of the meadows. This beautiful and fragrant plant is the *Spiræa ulmaria* of Linnaeus. The leaves are recommended as mild adstringents. The flowers have a strong smell, resembling that of May; they are supposed to possess antispasmodic and diaphoretic virtues, and as they are very rarely used in medicine, Linnaeus suspects that the neglect of them has arisen from the plant being supposed to be possessed of some noxious qualities, which it seemed to betray by its being left untouched by cattle. It may be observed, however, that the cattle also refuse the an-

golia and other herbs, whose innocence is apparent from daily experience.

ULMUS. 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Digymia*.

2. The pharmacopœial name of the common elm. *Ulmus campestris foliis duplicato-serratis, basi inæqualibus* of Linnæus. The inner tough bark of this tree, which is directed for use by the pharmacopœias, has no remarkable smell, but a bitterish taste, and abounds with a slimy juice, which has been recommended in nephritic cases, and externally as a useful application to burns. It is also highly recommended in some cutaneous affections allied to herpes and lepra. It is mostly exhibited in the form of decoction, by boiling four ounces in four pints of water to two pints; of which from four to eight ounces are given two or three times a day.

ULMUS CAMPESTRIS. The systematic name of the common elm. See *Ulmus*.

ULNA. (From *αλυσ*, the ulna, or cubit.) *Cubitus*. The larger bone of the fore-arm. It is smaller and shorter than the os humeri, and becomes gradually smaller as it descends to the wrist. We may divide it into its upper and lower extremities, and its body or middle part. At its upper extremity are two considerable processes, of which the posterior one and largest is named *olecranon*, and the smaller and anterior one the *coronoid* process. Between these two processes, the extremity of the bone is formed into a deep articulating cavity, which, from its semi-circular shape, is called the *greater sigmoid cavity*, to distinguish it from another, which has been named the *lesser sigmoid cavity*. The olecranon begins by a considerable tuberosity, which is rough, and serves for the insertion of muscles, and terminates in a kind of hook, the concave surface of which moves upon the pulley of the os humeri. This process forms the point of the elbow. The coronoid process is sharper at its extremity than the olecranon, but is much smaller, and does not reach so high. In bending the arm it is received into the fossa at the fore part of the pulley. At the external side of the coronoid process is the lesser sigmoid cavity, which is a small, semi-lunar, articulating surface, lined with cartilage, on which the round head of the radius plays. At the fore part of the coronoid process we observe a small tuberosity, into which the tendon of the brachialis internus is inserted. The greater sigmoid cavity, the situation of which we just now mentioned, is divided into four surfaces by a prominent line which is intersected by a small sinusity that serves for the lodgment of mucilaginous glands. The whole of this cavity is covered with cartilage. The body, or middle part of the ulna, is of a prismatic or triangular shape,

so as to afford three surfaces and as many angles. The external and internal surfaces are flat and broad, especially the external one, and are separated by a sharp angle, which, from its situation, may be termed the internal angle. This internal angle, which is turned towards the radius, serves for the attachment of the ligament that connects the two bones, and which is therefore called the *interosseous* ligament. The posterior surface is convex, and corresponds with the olecranon. The borders, or angles, which separate it from the other two surfaces, are somewhat rounded. At about a third of the length of this bone from the top, in its fore part, we observe a channel for the passage of vessels. The lower extremity is smaller as it descends, nearly cylindrical, and slightly curved forwards and outwards. Just before it terminates it contracts, so as to form a neck to the small head with which it ends. On the outside of this little head, answering to the olecranon, a small process, called the *styloid* process, stands out, from which a strong ligament is stretched to the wrist. The head has a rounded articulating surface, on its internal side, which is covered with cartilage, and received into a semi-lunar cavity formed at the lower end of the radius. Between it and the os cuneiforme, a moveable cartilage is interposed, which is continued from the cartilage that covers the lower end of the radius, and is connected by ligamentous fibres to the styloid process of the ulna. The ulna is articulated above with the lower end of the os humeri. This articulation is of the species called *ginglimus*, it is articulated also both above and below to the radius, and to the carpus at its lowest extremity. Its chief use seems to be to support and regulate the motions of the radius. In children, both extremities of this bone are first cartilaginous, and afterwards epiphyses, before they are completely united to the rest of the bone.

Ulnar artery. See *Cubital artery*.

Ulnar nerve. See *Cubital nerve*.

ULNARIS EXTERNUS. See *Extensor carpi ulnaris*.

ULNARIS INTERNUS. See *Flexor carpi ulnaris*.

UMBILICAL CORD. *Fundus umbilicalis*. *Funiculus umbilicalis*. The navel-string. A cord-like substance, of an intestinal form, about half a yard in length, that proceeds from the navel of the fœtus to the centre of the placenta. It is composed of a cutaneous sheath, cellular substance, one umbilical vein, and two umbilical arteries; the former conveys the blood to the child from the placenta, and the latter return it from the child to the placenta.

Umbilical hernia. See *Hernia umbilicalis*.

UMBILICAL REGION. *Regio umbil-*

bilicatis. The part of the abdominal parietes about two inches all round the navel.

UMBILICUS MARINUS. *Cotyledon marina*. *Androsace*. *Acetabulum marinum*. *Androsace methioli*. *Fungus petraeus marinus*. A submarine production found on rocks and the shells of fishes, about the coast of Montpelier, &c. It is said to be in the form of powder a useful antihelmetic and diuretic.

UNCIFORM BONE. (*Os unciforme*, from *uncus*, a hook, and *forma*, a likeness.) The last bone of the second row of the carpus or wrist; so named from its hook-like process, which projects towards the palm of the hand, and gives origin to the great ligament by which the tendons of the wrist are wound down. See *Carpal bones*.

UNGUENTUM. (From *ungō*, to anoint.) An ointment. The usual consistence of ointments is about that of butter. The following are among the best formulæ.

Ung. Cantharidis. See *Ceratum lyttæ*.

UNGUENTUM CETACEI. Ointment of spermaceti, formerly called *linimentum album*, and latterly *unguentum spermaceti*. "Take of spermaceti, six drachms; white wax, two drachms; olive oil, three fluid-ounces." Having melted them together over a slow fire, constantly stir the mixture until it gets cold. A simple emollient ointment.

UNGUENTUM CICUTÆ. Hemlock ointment. "Take of the fresh leaves of *cicuta*, and prepared hog's lard, of each four ounces." The *cicuta* is to be bruised in a marble mortar, after which the lard is to be added, and the two ingredients thoroughly incorporated by beating. They are then to be gently melted over the fire, and after being strained through a cloth, and the fibrous parts of the hemlock well pressed, the ointment is to be stirred till quite cold. To cancerous or scrophulous sores this ointment may be applied with a prospect of success.

UNGUENTUM ELEMI COMPOSITUM. Compound ointment of elemi, formerly called *linimentum urcei* and *unguentum e gummi elemi*. "Take of elemi, a pound; common turpentine, ten ounces; prepared suet, two pounds; olive oil, two fluid-ounces." Melt the elemi with the suet, then remove it from the fire, and immediately mix in the turpentine and oil, then strain the mixture through a linen cloth. Indolent ulcers, chilblains, chronic ulcers after burns, and indolent tumours, are often removed by this ointment.

UNGUENTUM HYDRARGYRI FORTIUS. Strong mercurial ointment, formerly called *unguentum ceruleum fortius*. "Take of purified mercury, two pounds; prepared lard, twenty-three ounces; prepared suet, an ounce." First rub the mercury with the suet and a little of the lard, until the globules disappear; then add the remain-

der of the lard, and mix. In very general use for mercurial frictions. It may be employed in almost all cases where mercury is indicated.

UNGUENTUM HYDRARGYRI MITIUS. Mild mercurial ointment, formerly called *unguentum ceruleum mitius*. "Take of strong mercurial ointment, a pound; prepared lard, two pounds." Mix. Weaker than the former.

UNGUENTUM HYDRARGYRI NITRATI. Ointment of nitrate of mercury, "Take of purified mercury, an ounce; nitric acid, two fluidounces; prepared lard, six ounces; olive oil, four fluidounces." First dissolve the mercury in the acid, then, while the liquor is hot, mix it with the lard and oil melted together. A stimulating and detergent ointment. *Tinea capitis*, *prophthalmia*, indolent tumours on the margin of the eye-lid, and ulcers in the urethra, are cured by its application.

UNGUENTUM HYDRARGYRI NITRATI MITIUS. Weaker only than the former.

UNGUENTUM HYDRARGYRI NITRICO-OXYDI. Ointment of nitric oxyd of mercury. "Take of nitric oxyd of mercury, an ounce; white wax, two ounces; prepared lard, six ounces." Having melted together the wax and lard, add thereto the nitric oxyd of mercury in very fine powder, and mix. A most excellent stimulating and escharotic ointment.

UNGUENTUM HYDRARGYRI PRÆCIPITATÆ ALBÆ. Ointment of white precipitate of mercury, formerly called *unguentum e mercurio precipitato albo*, and latterly *unguentum calcis hydrargyri albæ*. "Take of white precipitate of mercury, a drachm; prepared lard, an ounce and half." Having melted the lard over a slow fire, add the precipitated mercury, and mix. A useful ointment to destroy vermin in the head, and to assist in the removal of scald head, venereal ulcers of children, and cutaneous eruptions.

UNGUENTUM LYTTÆ. See *Ceratum lyttæ*.

UNGUENTUM OPHTHALMICUM. Ophthalmic ointment of Janin. "Take of prepared hogslard, half an ounce; prepared tutia, bole ammoniac, of each two drachms; white precipitate, one drachm." Mix. This celebrated ointment may be used for the same diseases of the eye and eye-lid as the *ung. hydrarg. nitrati*. It must be at first weakened with about twice its quantity of hogslard.

UNGUENTUM PICIS ARIDÆ. Pitch ointment, formerly called *unguentum basilicum nigrum vel tetrapharmacum*. "Take of pitch, yellow wax, yellow resin, of each nine ounces; olive oil, a pint." Melt them together, and strain the mixture through a linen cloth.

UNGUENTUM PICIS LIQUIDÆ. Tar ointment, formerly called *unguentum picis*;

unguentum e pice. "Take of tar, prepared suet, of each a pound." Melt them together, and strain the mixture through a linen cloth. The two preceding ointments are applicable to cases of tinea capitis, and some eruptive complaints; also to some kinds of irritable sores.

UNGUENTUM RESINÆ FLAVÆ. Yellow basilicon is in general use as a stimulant and detergent: it is an elegant and useful form of applying the resin.

UNGUENTUM SAMBUCI. Elder ointment, formerly called *unguentum sambucinum*. "Take of elder flowers, two pounds; prepared lard, two pounds." Boil the elder flowers in the lard until they become crisp, then strain the ointment through a linen cloth. A cooling and emollient preparation.

UNGUENTUM SULPHURIS. Sulphur ointment, formerly called *unguentum e sulphore*. "Take of sublimed sulphur, three ounces; prepared lard, half a pound." Mix. The most effectual preparation to destroy the itch. It is also serviceable in the cure of other cutaneous eruptions.

UNGUENTUM SULPHURIS COMPOSITUM. Compound sulphur ointment. "Take of sublimed sulphur, half a pound; white hellebore root, powdered, two ounces; nitrate of potash, a drachm; soft soap, half a pound; prepared lard, a pound and half." Mix. This preparation is introduced into the last London pharmacopeia as a more efficacious remedy for itch than common sulphur ointment. In the army, where it is generally used, the sulphur vivum, or native admixture of sulphur with various heterogeneous matters, is used instead of sublimed sulphur.

UNGUENTUM VERATRI. White hellebore ointment, formerly called *unguentum hellebori albi*. "Take of white hellebore root, powdered, two ounces; prepared lard, eight ounces; oil of lemons, twenty minimis." Mix.

UNGUENTUM ZINCI. Zinc ointment. "Take of the oxyd of zinc, an ounce; prepared lard, six ounces." Mix. A very useful application to chronic ophthalmia and relaxed ulcers.

UNGUIS. (From *ovξ*, a hook.) 1. The nail. The nails are horny laminae situated at the extremities of the fingers and toes.
2. An abscess or collection of pus between the lamellæ of the cornea transparents of the eye; so called from its resemblance to the lunated portion of the nail of the finger.

3. The lachrymal bone is so named from its resemblance to a nail of the finger. See *Lachrymal bone*.

UNGULA CABBALLINA. See *Tussilago*.

UNIONES. (*Unio*, pl. *uniones*, from *unus*, one; so called because there is never more than one found in the same shell, or according to others, for that many being

found in one shell not any one of them is like the other.) Pearls. See *Margarita*.

URACHUS. (From *uræ*, urine, and *χῶς*, to contain.) *Urinaculum*. The ligamentous cord that arises from the basis of the urinary bladder, which it runs along, and terminates in the umbilical cord. In the fetuses of brute animals, which the ancients mostly dissected, it is a hollow tube and conveys the urine to the allantoïd membrane.

URAGIUM. (From *uragos*, the hinder part of an army) The apex or extreme point of the heart.

URANISCU. (From *uragos*, the firmament; so called from its arch.) The palate.

URANIUM. This metal was discovered by Klaproth in the year 1789. It exists combined with sulphur, and a portion of iron, lead, and silex, in the mineral termed *Pechblende*, or *oxid of uranium*. Combined with carbonic acid it forms the *chalcite*, or *green mica*: and mixt with oxyd of iron, it constitutes the *uranitic ochre*. It is always found in the state of an oxyd with a greater or smaller portion of iron, or mineralized with sulphur and copper. The ores of uranium are of a blackish colour, inclining to a dark iron grey, and of a moderate splendor; they are of a close texture, and when broken present a somewhat uneven, and in the smallest particles a conchoidal surface. They are found in the mines of Saxony.

Properties of Uranium.—Uranium exhibits a mass of small metallic globules, agglutinated together. Its colour is a deep grey on the outside, in the inside it is a pale brown. It is very porous, and is so soft that it may be scraped with a knife. It has but little lustre. Its specific gravity is 6.440. It is more difficult to be fused than even manganese. When intensely heated with phosphate of soda and ammonia, or glacial phosphoric acid, it fuses with them into a grass-green glass. With soda or borax it melts only into a grey, opaque, scoriaceous bead. It is soluble in sulphuric, nitric, and muriatic acids. It combines with sulphur and phosphorus, and alloys with mercury. It has not yet been combined with other combustible bodies. It decomposes the nitric acid and becomes converted into a yellow oxyd. The action of uranium alone upon water, &c. is still unknown, probably on account of its extreme scarcity.

Method of obtaining Uranium.—In order to obtain uranium, the *pechblende* is first freed from sulphur by heat, and cleared from the adhering impurities as carefully as possible. It is then digested in nitric acid; the metallic matter that it contains is thus completely dissolved, while part of the sulphur remains undissolved, and part of it is dissipated under the form of sul-

phurated hydrogen gas. The solution is then precipitated by a carbonated alkali. The precipitate has a lemon-yellow colour when it is pure. This yellow carbonate is made into a paste with oil and exposed to a violent heat, bedded in a crucible containing and lined with charcoal.

Klaproth obtained a metallic globule 38 grains in weight, by forming a ball of 50 grains of the yellow carbonate with a little wax, and by exposing this ball in a crucible lined with charcoal to a heat equal to 170° of Wedgwood's pyrometer.

Richter obtained in a single experiment 100 grains of this metal, which seemed to be free from all admixture.

UNCOLA. (From *uraculus*, a small pitcher; so named from its uses in scouring glazed vessels.) The herb feverfew.

UREDE. (From *ure*, to burn.) An itching or burning sensation of the skin, which accompanies many diseases. The nettlerash is also so called.

URETER. (From *ure*, urine.) The membranous canal which conveys the urine from the kidney to the urinary bladder; at its superior part it is considerably the largest, occupying the greatest portion of the pelvis of the kidney; it then contracts to the size of a goose-quill, and descends over the *peas magnus* muscle and large crural vessels into the pelvis, in which it perforates the urinary bladder very obliquely. Its internal surface is lubricated with mucus to defend it from the irritation of the urine in passing.

URETERITIS. (From *ureter*, the ureter.) An inflammation of the ureter.

URETERICUS. (From *ureter*, the ureter, or *ureteritis*, an inflammation of the ureter.) Applied to an ischury, or suppression of urine, from an inflammation of the ureter.

URETEROLITHICUS. (From *ureter*, the ureter, and *lithon*, a stone.) Applied to an ischury from a stone in the ureter.

URETEROTROMBOIDES. (From *ureter*, the ureter, and *thrombos*, grumous blood, and *oides*, a likeness.) Applied to an ischury from grumous blood in the ureter.

URETEROPHLEGMATICUS. (From *ureter*, the ureter, and *phlegma*, phlegm.) Applied to a suppression of urine from pituitous matter in the ureter.

URETEROPUS. (From *ureter*, the ureter, and *pus*, pus.) Applied to an ischury from purulent matter in the ureter.

URETEROSTOMATICUS. (From *ureter*, the ureter, and *stoma*, a mouth.) Applied to a suppression of urine from an obstruction in the lower orifice of the ureter.

URETHRA. (From *ure*, the urine; because it is the canal through which the urine passes.) A membranous canal running from the neck of the bladder through the inferior part of the penis to the extremity of the glans penis, in which it opens

by a longitudinal orifice, called *meatus urinarius*. In this course it first passes through the prostate gland, which portion is distinguished by the name of the *prostatic urethra*; it then becomes much dilated, and is known by the name of the *bulbous part*, in which is situated a cutaneous eminence called the *caput gallinaginis* or *verumontanum*, around which are ten or twelve orifices of the excretory ducts of the prostate gland, and two of the spermatic vessels. The remaining part of the urethra contains a number of triangular mouths, which are the *lacunae*, or openings of the excretory ducts of the mucous glands of the urethra.

URETHELINTHICUS. (From *urethra*, the urethra, and *amphos*, worms.) Applied to an ischury from worms in the urethra.

URETHRITICUS. (From *urethrit*, an inflammation of the urethra.) Applied to a suppression of urine from an inflammation of the urethra.

URETHRITIS. (From *urethra*, the urethra.) An inflammation in the urethra. See *Gonorrhoea*.

URETHROCHYNODOS. (From *urethra*, the urethra, and *chyn*, a membrane.) Applied to an ischury from a membrane obstructing the urethra.

URETHROLITHICUS. (From *urethra*, the urethra, and *lithon*, a stone.) Applied to a suppression of urine from a stone in the urethra.

URETHROTROMBOS. (From *urethra*, the urethra, and *thrombos*, a grumous concretion.) A caruncle in the urethra.

URETHROPHLEGMATICUS. (From *urethra*, the urethra, and *phlegma*, phlegm.) Applied to an ischury from mucus obstructing the urethra.

URETHROPUS. (From *urethra*, the urethra, and *pus*, pus.) Applied to a suppression of urine from pus collected in the urethra.

URETICA. (From *ure*, urine.) Medicines which promote a discharge of urine.

URIAS. (From *ure*, urine.) The urethra.

URINE. (*Urina*; *ure*, from *urea*, to rush out.) The saline liquid, secreted in the kidneys, and dropping down from them, guttatum, through the ureters, into the cavity of the urinary bladder. The *secretory organ* is composed of the arterious vessels of the cortical substance of the kidneys, from which the urine passes through the uriniferous tubuli and renal papillae, into the renal pelvis: whence it flows drop by drop, through the ureters, into the cavity of the urinary bladder: where it is detained some hours, and at length, when *abundant*, eliminated through the urethra. The urine of an healthy man is divided in general into,

1. *Crude*, or that which is emitted one or two hours after eating; this is for the most part aqueous, and often vitiated by some foods, and,

2. *Cocted*, which is eliminated some hours after the digestion of the food, as that which is emitted in the morning after sleeping. This is generally in smaller quantity, thicker, more coloured, more acrid than at any other time. Of such cocted urine, the colour is usually citrine, and not unhandsome.

The degree of heat agrees with that of the blood; hence in atmospheric air it is warmer, as is perceived if the hand be washed with urine. The specific gravity is greater than water, and that emitted in the morning is always heavier than at any other time. The smell of fresh urine is not disagreeable. The taste is saltish and nauseous. The consistence is somewhat thicker than water. The quantity depends on that of the liquid drink, its diuretic nature, and the temperature of the air.

Changes of urine in the air. Preserved in an open vessel, it remains pellucid for some time, and at length there is perceived at the bottom a *nubecula* or little cloud, consolidated as it were from the gluten. This nubecula increases by degrees, occupies all the urine, and renders it opaque. The natural smell is changed into a putrid *cadaverous* one; and the surface is now generally covered with a *cuticle*, composed of very minute crystals. At length the urine regains its transparency, and the colour is changed from a yellow to a brown; the cadaverous smell passes into an *alkaline*, and a brown grumous sediment falls to the bottom, filled with white particles, deliquescent in the air, and so conglutinated as to form, as it were, little soft calculi.

Thus two sediments are distinguishable in the urine; the one white and gelatinous, and separated in the beginning; the other brown and grumous, deposited by the urine when putrid.

Spontaneous degeneration. Of all the fluids of the body, the urine first putrefies. In summer, after a few hours it becomes turbid, and sordidly black; then deposits a copious sediment, and exhales a fetor, like that of putrid cancers, which, at length becomes cadaverous. Putrid urine effervesces with acids, and if distilled, gives off, before water, an urinous volatile spirit.

The properties of healthy urine, are,

1. Urine reddens paper stained with turnsole and with the juice of radishes, and therefore contains an acid. This acid has been generally considered as the phosphoric, but Thénard has shewn that in reality it is the *acetic*.

2. If a solution of ammonia be poured into fresh urine, a white powder precipitates, which has the properties of *phosphate of lime*.

3. If the phosphate of lime precipitated from urine be examined, a little magnesia will be found mixed with it. Fourcroy and Vauquelin have ascertained that this is owing to a little *phosphate of magnesia* which urine contains, and which is decomposed by the alkali of lime employed to precipitate the phosphate of lime.

4. Proust informs us that *carbonic acid* exists in urine, and that its separation occasions the froth which appears during the evaporation of urine.

5. Proust has observed, that urine kept in new casks deposits small crystals, which effloresce in the air, and fall to powder. These crystals possess the properties of the *carbonate of lime*.

6. When fresh urine cools, it often lets fall a brick-coloured precipitate, which Scheele first ascertained to be crystals of *uric acid*. All urine contains this acid even when no sensible precipitate appears when it cools.

7. During intermitting fevers, and especially during diseases of the liver, a copious sediment of a brick-red colour is deposited from urine. This sediment contains the *rosacic acid* of Proust.

8. If fresh urine be evaporated to the consistence of a syrup, and muriatic acid be then poured into it, a precipitate appears which possesses the properties of *benzoic acid*.

9. When an infusion of tannin is dropped into urine, a white precipitate appears, having the properties of the combination of tannin and *albumen* or gelatine. Their quantity in healthy urine is very small, often indeed not sensible. Cruickshanks found that the precipitate afforded by tannin in healthy urine amounted to 1-240th part of the weight of the urine.

10. If urine be evaporated by a slow fire to the consistence of a thick syrup, it assumes a deep brown colour, and exhales a fetid ammoniacal odour. When allowed to cool, it concretes into a mass of crystals, composed of all the component parts of urine. If four times its weight of alcohol be poured into this mass, at intervals, and a slight heat be applied, the greatest part is dissolved. The alcohol which has acquired a brown colour is to be decanted off, and distilled in a retort in a sand heat till the mixture has boiled for some time and acquired the consistence of a syrup. By this time the whole of the alcohol has passed off, and the matter, on cooling, crystallizes in quadrangular plates, which intersect each other. This substance is *urea*, which composes 9-20ths of the urine, provided the watery part be excluded. It is this substance which characterises urine, and constitutes it what it is, and to which the greater part of the very singular phenomena of urine are to be ascribed.

11. According to Fourcroy and Vauquelin, the colour of urine depends upon the urea; the greater the proportion of urea the deeper the colour. But Proust has detected a *resinous matter* in urine similar to the resin of bile, and to this substance he ascribes the colour of urine.

12. If urine be slowly evaporated to the consistence of a syrup, a number of crystals make their appearance on its surface, these possess the properties of the *muriate of soda*.

13. The saline residuum which remains after the separation of urea from crystallized urine by means of alcohol, has been long known by the names of *fusible salt of urine*, and *microcosmic salt*. When these salts are examined, they are found to have the properties of phosphates. The rhomboidal prisms consist of *phosphate of ammonia*, united to a little *phosphate of soda*, the rectangular tables, on the contrary, are phosphate of soda united to a small quantity of phosphate of ammonia, urine then contains *phosphate of soda*, and *phosphate of ammonia*.

14. When urine is cautiously evaporated, a few cubic crystals are often deposited among the other salts, these crystals have the properties of *muriate of ammonia*.

15. When urine is boiled in a silver basin, it blackens the basin, and if the quantity of urine be large, small crusts of sulphuret of silver may be detached. Hence we see that urine contains *sulphur*.

Urine then contains the following substances :

1. Water.
2. Acetic acid.
3. Phosphate of lime.
4. Phosphate of magnesia.
5. Carbonic acid.
6. Carbonate of lime:
7. Uric acid.
8. Rosaic acid.
9. Benzoic acid.
10. Albumen.
11. Urea.
12. Resin:
13. Muriate of soda.
14. Phosphate of soda.
15. Phosphate of ammonia.
16. Muriate of ammonia.
17. Sulphur.

No liquor in the human body, however pure, is so variable, in respect to *quantity and quality*, as the urine; for it varies,

1. *In respect to age*: in the *fetus* it is inodorous, insipid, and almost aqueous; but as the *infant* grows, it becomes more acrid and fetid; and in *old age* more particularly so.

2. *In respect to drink*: it is secreted in greater quantity, and of a more pale colour, from cold and copious draughts.

It becomes green from an infusion of Chinese tea, or the use of the pulp of Cassia.

3. *In respect to food*: from eating the heads of asparagus, or olives, it contracts a peculiar smell; from the fruit of the opuntia, it becomes red; and from fasting turbid.

4. *In respect to medicines*: from the exhibition of rhubarb root, it becomes yellow; and from turpentine a violet colour.

5. *In respect to the time of the year*: in the winter the urine is more copious and aqueous; but in the summer, from the increased transpiration, it is more sparing, higher coloured, and so acrid that it sometimes becomes stranguous. The climate induces the same difference.

6. *In respect of the muscular motion of the body*: it is secreted more sparingly, and concentrated by motion; and is more copiously diluted, and rendered more crude by rest.

7. *In respect of the affection of the mind*: thus fright makes the urine pale.

Ure. The urine is an excrementitious fluid, like *lixivium*, by which the human body is not only liberated from the superfluous water, but also from the superfluous salts, and animal earth; and is defended from corruption.

Lastly, the *vis medicatrix nature* sometimes eliminates many morbid and acrid substances with the urine; as may be observed in fevers, dropsies, &c.

URINE, RETENTION OF. A want of the ordinary secretion of urine. In retention of urine there is none secreted; in a suppression, the urine is secreted but cannot be voided.

Urine, suppression of. See *Ischuria*.

URINA. See *Urine*.

Urinaculum. See *Urachus*.

URINARIA. (From *urina*, urine; so named from its diuretic qualities.) The herb dandelion. See *Taraxacum*.

URINÆ ARDOR. See *Dysuria*.

UROCRISIA. (From *uros*, urine, and *crisis*, to judge.) The judgment formed of diseases by the inspection of urine.

UROREXA. (From *uros*, the urine, and *exo*, to flow.) A discharge of the urine through the eroded perineum.

URSINA RADIX. The root of the plant called baldmoney. See *Meum athamanticum*.

UROSCOPIA. (From *uros*, the urine, and *scopia*, to inspect.) Inspection of urine, that a judgment of diseases may be made from its appearance.

URTICA. (*Ab urendo*; because it excites an itching and pustules like those produced by fire.) 1. The name of a genus of plants in the Linnæan system. Class, *Monocotyledon*. Order, *Tetrandria*. The nettle.

2. The pharmacopœial name of the common nettle. *Urtica dioica* of Linnæus. This plant is well known, and though generally despised as a noxious weed, has been long used for medical, culinary, and economical purposes. The young shoots in the spring possess diuretic and antiscorbutic properties, and are with these intentions boiled and eaten instead of cabbage greens.

URTICA DIOICA. The systematic name of the common stinging nettle. See *Urtica*.

URTICA MORTUA. See *Lamium album*.

URTICA PILULIFERA. The systematic name of the pillbearing nettle. See *Urtica romana*.

URTICA ROMANA. The plant which bears this name in the pharmacopœias is the *Urtica pilulifera* of Linnæus. The seed was formerly given against diseases of the chest, but is now deservedly forgotten. To raise an irritation in paralytic limbs the fresh plant may be employed as producing a more permanent sting than the common nettle.

URTICA URENS. The systematic name of a lesser nettle than the dioica, and possessing similar virtues.

URTICARIA. (From *urtica*, a nettle.) *Febris urticata.* *Uredo.* *Purpura urticata.* *Scarlatina urticata.* The nettle-rash. A species of exanthematous fever, known by pyrexia and an eruption on the skin like that produced by the sting of the nettle. The little elevations, called the nettle-rash, often appear instantaneously, especially if the skin be rubbed or scratched, and seldom stay many hours in the same place, and sometimes not many minutes. No part of the body is exempt from them; and where many of them rise together, and continue an hour or two, the parts are often considerably swelled, which particularly happens in the arms, face, and hands. These eruptions will continue to infest the skin, sometimes in one place and sometimes in another, for one or two hours together, two or three times a day, or perhaps for the greatest part of twenty-four hours. In some constitutions they last only a few days, in others many months.

URTICATIO. (From *urtica*, a nettle.) The whipping a paralytic or benumbed limb with nettles, in order to restore its feelings.

USNEA. *Muscus cranii humani.* This moss, *Lichen saxatilis* of Linnæus, when growing on the human skull, was formerly in high estimation, but now deservedly forgotten.

UTERARIA. (From *uterus*, the womb.) Medicines appropriated to diseases of the womb.

Uterine fury. See *Nymphomania*.

UTERUS. *Torrea.* *Matrix.* *Ager na-*

tura. *Hystera.* *Matra.* *Utriculus.* The womb. A spongy receptacle resembling a compressed pear, situated in the cavity of the pelvis, above the vagina, and between the urinary bladder and rectum.

The form of the uterus resembles that of an oblong pear flattened, with the depressed sides placed towards the ossa pubis and sacrum; but, in the impregnated state, it becomes more oval, according to the degree of its distention. For the convenience of description, and for some practical purposes, the uterus is distinguished into three parts. The fundus, the body, and the cervix; the upper part is called the fundus, the lower the cervix, the space between them, the extent of which is undefined, the body. The uterus is about three inches in length, about two in breadth at the fundus, and one at the cervix. Its thickness is different at the fundus and cervix, being at the former usually rather less than half an inch, and at the latter somewhat more; and this thickness is preserved throughout pregnancy, chiefly by the enlargement of the veins and lymphatics; there being a smaller change in the size of the arteries. But there is so great a variety in the size and dimensions of the uterus in different women, independent of the states of virginity, marriage, or pregnancy, as to prevent any very accurate mensuration. The cavity of the uterus corresponds with the external form; that of the cervix leads from the os uteri, where it is very small, in a straight direction, to the fundus, where it is expanded into a triangular form, with two of the angles opposed to the entrance into the Fallopian tubes; and at the place of junction between the cervix and the body of the uterus the cavity is smaller than it is in any other part. There is a swell, or fullness of all the parts towards the cavity, which is sometimes distinguished by a prominent line running longitudinally through its middle. The villous coat of the vagina is reflected over the os uteri, and is continued into the membrane which lines the cavity of the uterus. The internal surface of the uterus is corrugated in a beautiful manner, but the rugæ, or wrinkles, which are longitudinal, lessen as they advance into the uterus, the fundus of which is smooth. In the intervals between the rugæ are small orifices, like those in the vagina, which discharge a mucus, serving, besides other purposes, that of closing the os uteri very curiously and perfectly during pregnancy. The substance of the uterus, which is very firm, is composed of arteries, veins, lymphatics, nerves, and muscular fibres, curiously interwoven and connected together by cellular membrane. The muscular fibres are of a pale colour, and appear also

in their texture somewhat different from muscular fibres in other parts of the body. The arteries of the uterus are the spermatic and hypogastric. The spermatic arteries arise from the anterior part of the aorta, a little below the emulgents, and sometimes from the emulgents. They pass over the psoas muscles behind the peritonæum, enter between the two laminae or duplicatures of the peritonæum which form the broad ligaments of the uterus, and proceed to the uterus, near the fundus of which they insinuate themselves, giving branches in their passage to the ovaria and Fallopian tubes. The hypogastric arteries are on each side a considerable branch of the internal iliacs. They pass to the sides of the body of the uterus, sending off a number of smaller branches, which dip into its substance. Some branches also are reflected upwards to the fundus uteri, which anastomose with the spermatic arteries, and others are reflected downwards, supplying the vagina. The veins which reconvert the blood from the uterus are very numerous, and their size in the unimpregnated state is proportioned to that of the arteries; but their enlargement during pregnancy is such, that the orifices of some of them, when divided, will admit even of the end of a small finger. The veins anastomose in the manner of the arteries which they accompany out of the uterus, and then, having the same names with the arteries spermatic and hypogastric, the former proceeds to the vena cava on the right side, and on the left to the emulgent vein; and the latter to the internal iliac.

From the substance and surfaces of the uterus an infinite number of lymphatics arise, which follow the course of the hypogastric and spermatic blood-vessels. The first pass into the gland of the internal iliac plexus, and the other into the glands which are situated near the origin of the spermatic arteries. Of these Nuck first gave a delineation.

The uterus is supplied with nerves from the lower mesocolic plexus, and from two small flat circular ganglions, which are situated behind the rectum. These ganglions are joined by a number of small branches from the third and fourth sacral nerves. The ovaria derive their nerves from the renal plexus. By the great number of nerves these parts are rendered very irritable, but it is by those branches which the uterus receives from the intercostal, that the intimate consent between it and various other parts is chiefly preserved. The muscular fibres of the uterus have been described in a very different manner by anatomists, some of whom have asserted that its substance was chiefly muscular, with fibres running in transverse, orbicular,

or reticulated order, whilst others have contended that there were no muscular fibres whatever in the uterus. In the unimpregnated uterus, when boiled for the purpose of a more perfect examination, the former seems to be a true representation; and when the uterus is distended towards the latter part of pregnancy, these fibres are very thinly scattered; but they may be discovered in a circular direction, at the junction between the body and the cervix of the uterus, and surrounding the entrance of each Fallopian tube in a similar order. Yet it does not seem reasonable to attribute the time of labour to its muscular fibres only, if we are to judge of the power of a muscle by the number of fibres of which it is composed, unless it is presumed that those of the uterus are stronger than in common muscles. With respect to the glands of the uterus, none are discoverable dispersed through its substance upon the inner surface of the cervix; between the rugæ there are lacunæ which secrete mucus, and there are small follicles at the edge of the os uteri. These last are only observable in a state of pregnancy, when they are much enlarged. From the angles at the fundus of the uterus, two processes of an irregular round form originate, called, from the name of the first describer, the *Fallopian tubes*. They are about three inches in length, and, becoming smaller in their progress from the uterus, have an uneven, fringed termination, called the *fimbriæ*. The canal which passes through these tubes is extremely small at their origin, but it is gradually enlarged, and terminates with a patulous orifice, the diameter of which is about one third of an inch, surrounded by the fimbriæ. It is also lined by a very fine vascular membrane, formed into serpentine plicæ. Through this canal, the communication between the uterus and ovaria is preserved. The Fallopian tubes are wrapped in duplicatures of the peritonæum, which are called the broad ligaments of the uterus; but a portion of their extremities, thus folded, hangs loose on each side of the pelvis. From each lateral angle of the uterus, a little before and below the Fallopian tubes, the *round ligaments* arise, which are composed of arteries, veins, lymphatics, nerves, and a fibrous structure. These are connected together by cellular membrane, and the whole is much enlarged during pregnancy. They receive their outward covering from the peritonæum, and pass out of the pelvis through the ring of the external oblique muscle to the groin, where the vessels subdivide into small branches, and terminate at the mons veneris and contiguous parts. From the insertion of these ligaments into the groin, the reason appears why that part

generally suffers in all the diseases and affections of the uterus, and why the inguinal glands are in women so often found in a morbid or enlarged state. The duplicatures of the peritonæum, in which the Fallopian tubes and ovaria are involved, are called the *broad ligaments* of the uterus. These prevent the entanglement of the parts, and are conductors of the vessels and nerves as the mesentery is of those of the intestines. Both the round and broad ligaments alter their position during pregnancy, appearing to rise lower and more forward than in the unimpregnated state. Their use is supposed to be that of preventing the descent of the uterus, and to regulate its direction when it ascends into the cavity of the abdomen; but whether they answer these purposes may be much doubted. The use of the womb is for menstruation, conception, nutrition of the fœtus, and parturition. The uterus is liable to many diseases, the principal of which are *prolapsus uteri*, *procidentia uteri*, hydatids, dropsy of the uterus or tympanites uteri, moles, ulceration, &c.

UTERUS, RETROVERSION OF. By the term retroversion, such a change of the position of the uterus is understood, that the fundus is turned backwards and downwards upon its cervix, between the vagina and rectum, and the os uteri is turned forwards to the pubis, and upwards in proportion to the descent of the fundus, so that, by an examination *per vaginam*, it cannot be felt, or not without difficulty, when the uterus is retroverted. By the same examination there may also be perceived a large round tumour, occupying the inferior part of the cavity of the pelvis, and pressing the vagina towards the pubis. By an examination *per anum*, the same tumour may be felt, pressing the rectum to the hollow of the sacrum, and if both these examinations are made at the same time, we may readily discover that the tumour is confined between the vagina and rectum. Besides the knowledge of the retroversion which may be gained by these examinations, it is found to be accompanied with other very distinguishing symptoms. There is in every case, together with extreme pain, a suppression of urine; and by the continuance of this distention of the bladder the tumour formed by it in the abdomen often equals in size, and resembles in shape, the uterus in the sixth or seventh months of pregnancy; but it is necessary to observe, that the suppression of urine is frequently absolute only before the retroversion of the uterus, or during the time it is retroverted; for when the retroversion is completed, there is often a discharge of urine, so as to prevent an increase of the distention of the bladder,

though not in a sufficient quantity to remove it. There is also an obstinate constipation of the bowels, produced by the pressure of the retroverted uterus upon the rectum, which renders the injection of a clyster very difficult, or even impossible. But it appears that all the painful symptoms are chiefly in consequence of the suppression of urine; for none of those parts which are apt to sympathise in affections or diseases of the uterus are disturbed by its retroversion. The retroversion of the uterus has generally occurred about the third month of pregnancy, and sometimes after delivery it may likewise happen, where the uterus is, from any cause, enlarged to the size it acquires about the third month of pregnancy, but not with such facility as in the pregnant state, because the enlargement is then chiefly at the fundus. If the uterus is but little enlarged, or if it be enlarged beyond a certain time, it cannot well be retroverted; for, in the first case, should the cause of a retroversion exist, the weight at the fundus would be wanting to produce it; and in the latter the uterus would be raised above the projection of the sacrum, and supported by the spine.

UTRICARIA. (From *uter*, a bottle; so named from its appendages at the end of the leaves resembling bottles, to contain water.) A name of the nepenthes, or wonderful plant.

UTRICULUS. (Dim. of *uter*, a bottle; so called from its shape.) The womb.

UVA. (*Quasi uvula*, from its juice.) An unripe grape. A tumour on the eye resembling a grape.

UVA GRANA. Crane-berries. They are brought from New England, and are reckoned antiscorbutic.

UVA PASSA MAJOR. *Passula major*. The raisin. The dried fruit of the *Vitis vinifera* of Linnæus:—*foliis lobatis sinuatis nudis*. Raisins are prepared by immersing the fresh fruit into a solution of alkaline salt and soap-ley, made boiling hot, to which is added some olive-oil, and a small quantity of common salt, and afterwards drying them in the shade. They are used as agreeable, lubricating, acescent sweets in pectoral decoctions, and for obtunding the acrimony in other medicines, and rendering them grateful to the palate and stomach. They are directed in the *decoctum hordei compositum tinctura cœnæ*, and *tinctura cardamomi composita*.

UVA PASSA MINOR. *Passa corinthiaca*. The currants. The dried fruit of the *vitis corinthica*. Their virtues are similar to those of the *uva passa major*.

UVA URSI. Bear's whortle-berry. Bears' whorts, or bear's bilberries, called also *vaccaria vaccinia*. Trailing arbutus,

or bear-berry. *Arbutus uva ursi* of Linnaeus: —*canlibus procumbentibus, foliis integerrimis*. This plant, though employed by the ancients in several diseases, requiring adstringent medicines, had almost entirely fallen into disuse until the middle of the present century, when it first drew the attention of physicians as a useful remedy in calculous and nephritic complaints, which diseases it appears to relieve by its adstringent qualities.

UVEA. (From *uva*, an unripe grape.) The posterior lamina of the iris; so called because, in beasts, which the ancients chiefly dissected, it is of the colour of unripe grapes.

UVULA. (Dim. of *uva*, a grape.) *Columnella*. *Sion*. *Gargareon*. *Columna oris*.

Gurgulis interspectrum. The small conical fleshy substance hanging in the middle of the *velum pendulum palati*, over the root of the tongue. It is composed of the common membrane of the mouth, and a small muscle resembling a worm which arises from the union of the palatine bone, and descends to the tip of the uvula. It was called *Pulato staphilinus* by Douglas, and *Staphilinus epistaphilinus* by Winslow. By its contraction the uvula is raised up.

UVULARIA. (From *uvula*, because it cured diseases of the uvula.) The plant which bears this epithet is some pharmacopœias is the *Ruscus hypoglossum* of Linnaeus: it was formerly used against relaxation of the uvula, but now laid aside for more adstringent remedies.

V

VACCA. The cow.

VACCARIA. (From *vacca*, a cow; because it is coveted by cows.) The herb cow's-basil.

VACCINATION. The insertion of the matter to produce the cow-pox. See *Variola vaccine*.

VACCINIUM. (Quasi *baccinium*, from its berry.) The name of a genus of plants in the Linnæan system. Class, *Octandria*. Order, *Monogynia*.

VACCINIUM MYRTILLUS. The systematic name of the myrtle-berry. See *Myrtillus*.

VACCINIUM OXYCOCCUS. The systematic name of the cranberry-plant. See *Oxycoccus*.

VACCINIUM VITIS IDÆA. The systematic name of the red whortle-berry. See *Vitis idæa*.

VAGINA. A sheath. *Vagina uteri*. That canal which leads from the pudendum or external orifice to the uterus, is called the vagina. It is somewhat of a conical form, with the narrowest part downwards, and is described as being five or six inches in length, and about two in diameter. But it would be more proper to say, that it is capable of being extended to those dimensions; for in its common state, the os uteri is seldom found to be more than three inches from the external orifice, and the vagina is contracted as well as shortened. The vagina is composed of two coats, the first or innermost of which is villous, interspersed with many excretory ducts, and contracted into plicæ, or small transverse folds, particularly at the fore and back part, but, by child-bearing these are lessened or obliterated.

The second coat is composed of a firm membrane, in which muscular fibres are not distinctly observable, but which are endowed, to a certain degree, with contractile powers like a muscle. This is surrounded by cellular membrane, which connects it to the neighbouring parts. A portion of the upper and posterior part of the vagina is also covered by the peritonæum. The entrance of the vagina is constricted by muscular fibres, originating from the rami of the pubis, which run on each side of the pediculum, surrounding the posterior part, and executing an equivalent office, though they cannot be said to form a true sphincter.

The upper part of the vagina is connected to the circumference of the os uteri, but not in a straight line, so as to render the cavity of the uterus a continuation of that of the vagina. For the latter stretches beyond the former, and, being joined to the cervix, is reflected over the os uteri, which, by this mode of union, is suspended with protuberant lips in the vagina, and permitted to change its position in various ways and directions. When therefore these parts are distended and unfolded at the time of labour, they are continued into each other, and there is no part which can properly be considered as the precise beginning of the uterus or termination of the vagina.

The diseases of the vagina are, first, such an abbreviation and contraction as render it unfit for the uses for which it was designed: secondly, a cohesion of the sides in consequence of preceding ulceration.

thirdly, cicatrices after an ulceration of the parts: fourthly, excrescences: fifthly, fluor albus. This abbreviation and contraction of the vagina, which usually accompany each other, are produced by original defective formation, and they are seldom discovered before the time of marriage, the consummation of which they sometimes prevent. The curative intentions are to relax the parts by the use of emollient applications, and to dilate them to their proper size by sponge, or other tents, or, which are more effectual, by bougies gradually enlarged. But the circumstances which attend this disorder, are sometimes such as might lead us to form an erroneous opinion of the disease. A case of this kind, which was under Dr. Denman's care, from the strangury, from the heat of the parts, and the profuse and inflammatory discharge, was suspected to proceed from venereal infection; and with that opinion the patient had been put upon a course of medicine composed of quicksilver, for several weeks, without relief. When she applied to the Dr. he prevailed upon her to submit to an examination, and found the vagina rigid, so much contracted as not to exceed half an inch in diameter, and more than one inch and a half in length. The repeated, though fruitless attempts which had been made to complete the act of coition, had occasioned a considerable inflammation upon the parts, and all the suspicious appearances before mentioned. To remove the inflammation she was bled, took some gentle purgative medicines, used an emollient fomentation, and afterwards some unctuous applications; she was also advised to live separate from her husband for some time. The inflammation being gone, tents of various sizes were introduced into the vagina, by which it was distended, though not very amply. She then returned to her husband, and in a few months became pregnant. Her labour, though slow, was not attended with any extraordinary difficulty. She was delivered of a full-sized child, and afterwards suffered no inconvenience. Another kind of constriction of the external parts sometimes occurs, and which seems to be a mere spasm. By the violence or long continuance of a labour, by the morbid state of the constitution, or by the negligent and improper use of instruments, an inflammation of the external parts, or vagina, is sometimes produced in such a degree as to endanger a mortification. By careful management this consequence is usually prevented, but in some cases, when the constitution of the patient was prone to disease, the external parts have sloughed away, and in others, equal injury has been done to the vagina. But the effect of the inflammation is usually confined to the internal villous coat, which is sometimes

cast off wholly or partially. An ulcerated surface being thus left, when the disposition to heal has taken place, cicatrices have been formed of different kinds, according to the depth and extent of the ulceration; and there being no counteraction to the contractile state of the parts, the dimensions of the vagina become much reduced, or if the ulceration should not be healed, and the contractibility of the parts continue to operate, the ulcerated surfaces being brought together may cohere, and the canal of the vagina be perfectly closed.

Cicatrices in the vagina very seldom become an impediment to the connexion between the sexes; when they do, the same kind of assistance is required as was recommended in the natural contraction or abbreviation of the part; they always give way to the pressure of the head of the child in the time of labour, though in many cases with great difficulty. Sometimes the appearances may mislead the judgment; for the above author was called to a woman in labour, who was thought to have become pregnant; the hymen remained unbroken; but, on making very particular inquiry, he discovered that this was her second labour, and that the part which, from its form and situation, was supposed to be the hymen, with a small aperture, was a cicatrice, or unnatural contraction of the entrance into the vagina, consequent to an ulceration of the part after her former labour. Fungous excrescences arising from any part of the vagina or uterus, have been distinguished, though not very properly, by the general term polypus. See *Polypus*.

VAGINA OF NERVES. The outer covering of nerves. By some it is said to be a production of the pia mater only, and by others of the dura mater, because it agrees with it in tenacity, colour, and texture.

VAGINA OF TENDONS. A loose membranous sheath, formed of cellular membrane, investing the tendons, and containing an unctuous juice, which is secreted by the vessels of its internal surface. Ganglions are nothing more than an accumulation of this juice.

VAGINALIS TUNICA TESTIS. See *Tunica vaginalis testis*.

Vagum par. See *Par vagum*.

Valerian, celtic. See *Nardus celtica*.

Valerian, garden. See *Valeriana major*.

Valerian, great. See *Valeriana major*.

Valerian, lesser. See *Valeriana sylvestris*.

Valerian, wild. See *Valeriana sylvestris*.

VALERIANA. (From *Valerius*, who first particularly described it.) 1. The name of a genus of plants in the Linnæan system. Class, *Triandria*, Order, *Monogynia*. Valerian.

2. The pharmacopœial name of the wild valerian. *Valeriana minor*. Official valerian. *Valeriana officinalis* of Linnæus: *floribus triandris, foliis omnibus pinnatis*. The root of this plant has been long extolled as an efficacious remedy in epilepsy, which caused it to be exhibited in a variety of other complaints termed nervous, in which it has been found highly serviceable. It is also in very general use as an antispasmodic, and is exhibited in convulsive hysterical diseases. A simple and volatile tincture are directed in the pharmacopœias.

VALERIANA CELTICA. The systematic name of the celtic nard. See *Nardus celtica*.

VALERIANA MAJOR. *Phu*. The garden valerian. The root of this plant, *valeriana phu* of Linnæus, is said to be efficacious in removing rheumatism, especially the sciatica, and also inveterate epilepsies.

VALERIANA MINOR. See *Valeriana sylvestris*.

VALERIANA OFFICINALIS. The systematic name of the wild valerian. See *Valeriana sylvestris*.

VALERIANA PHU. The systematic name of the garden valerian. See *Valeriana major*.

VALERIANA SYLVESTRIS. See *Valeriana*.

VALLUM. (From *vallus*, a hedge stake; so called from the regular trench-like disposition of the hairs.) The eye-brows.

VALVA. (*Valva*; from *valveo*, to fold up.) A thin and transparent membrane situated within certain vessels, as arteries, veins, and absorbents, whose office appears to be to prevent the contents of the vessel from flowing back.

VALVE OF THE COLON. The end of the iliac portion of the small intestine enters the large one obliquely, and projects somewhat within it, so as to form a kind of valve, called from its discoverer the valve of Tulpus, also the valve of the cæcum.

Valves, semilunar. See *Semilunar valves*.

Valves, tricuspid. See *Tricuspid valves*.

Valves, triglochin. See *Tricuspid valves*.

VALVULA. (Dim. of *valva*.) A little valve.

VALVULA COLI. See *Intestines*.

VALVULA EUSTACHII. A membranous semilunar valve, which separates the right auricle from the inferior vena cava, first described by Eustachius.

VALVULA TULPII. See *Valves of the colon*.

VALVULÆ CONNIVENTES. The semilunar folds formed of the villous coat of the intestinum duodenum and jejunum. Their use appears to be to increase the surface of the intestines.

VALVULÆ MITRALES. See *Mitra valves*.

VALVULÆ SEMILUNARES. See *Semilunar valves*.

VALVULÆ TRIGLOCHINES. See *Tricuspid valves*.

VANELLOR. *Vanilla. Banilia. Bonilas. Aracus. Aromaticus.* A long, flattish pod, containing, under a wrinkled brittle shell, a reddish brown pulp, with small shining black seeds. The plant which affords this fruit is the *Epidendrum vanilla* of Linnæus:—*ascandens, foliis ovato-oblongis nervosis sessilibus caulinis, cirrhis spiræbibus*. Vanelloes have an unctuous aromatic taste, and a fragrant smell like that of some of the finer balsams heightened with musk. Although chiefly used as perfumes, they are said to possess aphrodisiac virtues.

Vanilla. See *Vanellor*.

VAPORARIUM. (From *vapor*, vapour.) A vapour-bath.

VARI. See *Jenthi*.

VARIA. (From *varius*, changeable.) The small-pox; also small red pimples in the face.

VARICELLA. (Dim. of *varia*, the small-pox; so called from its being changeable.) *Variola lymphatica.* The Chicken-pox. A genus of disease in the class *pyrexia* and order *exanthemata* of Cullen; known by moderate synocha; pimples bearing some resemblance to the small-pox, quickly forming pustules, which contain a fluid matter, and after three or four days from their first appearance, desquamate.

VARICOCELE. (From *varix*, a distended vein, and *cœla*, a tumour.) A swelling of the veins of the scrotum, or spermatic cord; hence it is divided into the *scrotal varicocele*, which is known by the appearance of livid and tumid veins on the scrotum; and *varicocele of the spermatic cord*, known by feeling hard vermiform vessels in the course of the spermatic cord.

Varicocele mostly arises from excessive walking, running, jumping, wearing of trusses, and the like, producing at first a slight uneasiness in the part, which if not remedied, continues advancing towards the loins.

VARIOLA. (From *varius*, changing colour, because it disfigures the skin.) The small pox. A genus of disease in the class *pyrexia* and order *exanthemata* of Cullen distinguished by synocha; eruption of red pimples on the third day, which on the eighth day contain pus, and drying, fall off in crusts.

It is a disease of a very contagious nature, supposed to have been introduced into Europe from Arabia, and in which there arises a fever, that is succeeded by a number of little inflammations in the skin, which proceed to suppuration, the matter formed thereby being capable of produc-

cising the disorder in another person. It makes its attack on people of all ages, but the young of both sexes are more liable to it than those who are much advanced in life; and it may prevail at all the seasons of the year, but in general is most prevalent in the spring and summer.

The small-pox is distinguished into the distinct and confluent, implying that in the former, the eruptions are perfectly separate from each other, and that in the latter, they run much into one another.

Both species are produced either by breathing air impregnated with the effluvia arising from the body of those who labour under the disease, or by the introduction of a small quantity of the variolous matter into the habit by inoculation; and it is probable that the difference of the small-pox is not owing to any difference in the contagion, but depends on the state of the person to whom it is applied, or on certain circumstances concurring with the application of it.

A variety of opinions have been entertained respecting the effect of the variolous infection on the fœtus in the uterus; a sufficient number of instances, however, has been recorded, to ascertain that the disease may be communicated from the mother to the child. In some cases, the body of the child, at its birth, has been covered with pustules, and the nature of the disease has been most satisfactorily ascertained by inoculating with matter taken from the pustules. In other cases, there has been no appearance of the disease at the time of the birth, but an eruption and other symptoms of the disease have appeared so early, as to ascertain that the infection must have been received previously to the removal of the child from the uterus.

Four different states, or stages, are to be observed in the small-pox: first, the febrile; second, the eruptive; third, the maturative; and fourth, that of the declination or scabbing. When the disease has arisen naturally, and is of the distinct kind, the eruption is commonly preceded by a redness in the eyes, soreness in the throat, pains in the head, back, and loins, weariness and faintness, alternate fits of chilliness and heat, thirst, nausea, inclination to vomit, and a quick pulse.

In some instances these symptoms prevail in a high degree, and in others they are very moderate and trifling. In very young children, startings and convulsion are apt to take place a short time previous to the appearance of the eruption, always giving great alarm to those not conversant with the frequency of the occurrence.

About the third or fourth day from the first seizure, the eruption shews itself in little red spots on the face, neck, and breast, and these continue to increase in

number and size for three or four longer, at the end of which time, they are to be observed dispersed over several parts of the body.

If the pustules are not very numerous, the febrile symptoms will generally go off on the appearance of the eruption, or they will become very moderate. It sometimes happens, that a number of little spots of an erysipelatous nature are interspersed amongst the pustules; but these generally go in again, as soon as the suppuration commences, which is usually about the fifth or sixth day, at which period, a small vesicle, containing an almost colourless fluid, may be observed upon the top of each pimple. Should the pustules be perfectly distinct and separate from each other, the suppuration will probably be completed about the eighth or ninth day, and they will then be filled with a thick yellow matter; but should they run much into each other, it will not be completed till some days later.

When the pustules are very thick and numerous on the face, it is apt about this time to become much swelled, and the eyelids to be closed up, previous to which, there usually arises a hoarseness, and difficulty of swallowing, accompanied with a considerable discharge of viscid saliva. About the eleventh day, the swelling of the face usually subsides, together with the affection of the fauces, and is succeeded by the same in the hands and feet, after which the pustules break, and discharge their contents; and then becoming dry, they fall in crusts, leaving the skin which they covered of a brown red colour, which appearance continues for many days. In those cases where the pustules are large, and are late in becoming dry and falling off, they are very apt to leave pits behind them; but where they are small, suppurate quickly, and are few in number, they neither leave any marks behind them, nor do they occasion much affection of the system.

In the confluent small pox, the fever which precedes the eruption is much more violent than in the distinct, being attended usually with great anxiety, heat, thirst, nausea, vomiting, and a frequent and contracted pulse, and often with coma or delirium. In infants, convulsive fits are apt to occur, which either prove fatal before any eruption appears, or they usher in a malignant species of the disease.

The eruption usually makes its appearance about the third day, being frequently preceded or attended with a rosy efflorescence, similar to what takes place in the measles; but the fever, although it suffers some slight remission on the coming out of the eruption, does not go off as in the distinct kind; on the contrary, it becomes increased after the fifth or sixth day, and

continues considerably throughout the remainder of the disease.

As the eruption advances, the face being thickly beset with pustules, becomes very much swelled, the eyelids are closed up, so as to deprive the patient of sight, and a gentle salivation ensues, which, towards the eleventh day, is so viscid as to be spit up with great difficulty. In children, a diarrhoea usually attends this stage of the disease instead of a salivation, which is to be met with only in adults. The vesicles on the top of the pimples are to be perceived sooner in the confluent small-pox than in the distinct; but they never rise to an eminence, being usually flattened; neither do they arrive to proper suppuration, as the fluid contained in them, instead of becoming yellow, turns to a brown colour.

About the tenth or eleventh day, the swelling of the face usually subsides, and then the hands and feet begin to puff up and swell, and about the same time the vesicles break, and pour out a liquor that forms into brown or black crusts, which, upon falling off, leave deep pits behind them that continue for life; and where the pustules have run much into each other, they then disfigure and scar the face very considerably.

Sometimes it happens that a putrescency of the fluids takes place at an early period of the disease, and shews itself in livid spots interspersed amongst the pustules, and by a discharge of blood by urine, stool, and from various parts of the body.

In the confluent small-pox, the fever, which, perhaps, had suffered some slight remission from the time the eruption made its appearance to that of maturation, is often renewed with considerable violence at this last-mentioned period, which is what is called the secondary fever, and this is the most dangerous stage of the disease. It has been observed, even amongst the vulgar, that the small pox is apt to appear immediately before or after the prevalence of the measles. Another curious observation has been made relating to the symptoms of these complaints, namely, that if, while a patient labours under the small-pox, he is seized with the measles, the course of the former is retarded till the eruption of the measles is finished. The measles appear, for instance, on the second day of the eruption of small-pox, the progress of this ceases, till the measles terminate by desquamation, and then it goes on in the usual way. Several cases are, however, recorded in the *Medical and Physical Journal*, as likewise in the third volume of the *Medical Commentaries*, in which a concurrence of the small-pox and measles took place without the progress of the former being retarded. The distinct small-pox is not attended with danger, except when it attacks pregnant women,

or approaches nearly in its nature to that of the confluent; but this last is always accompanied with considerable risk, the degree of which is ever in proportion to the violence and permanence of the fever, the number of pustules on the face, and the disposition to putrescency which prevails.

When there is a great tendency this way, the disease usually proves fatal between the eighth and eleventh day, but, in some cases, death is protracted till the fourteenth or sixteenth. The confluent small-pox, although it may not prove immediately mortal, is very apt to induce various morbid affections.

Both kinds of small-pox leave behind them a predisposition to inflammatory complaints, particularly to ophthalmia and visceral inflammations, but more especially of the thorax; and they not unfrequently excite scrophula into action which might otherwise have lain dormant in the system.

The regular swelling of the hands and feet upon that of the face subsiding, and its continuance for the due time, may be regarded in a favourable light.

The dissections which have been made of confluent small-pox, have never discovered any pustules internally on the viscera. From them it also appears that variolous pustules never attack the cavities of the body, except those to which the air has free access, as the nose, mouth, trachea, the larger branches of the bronchiz, and the outermost part of the meatus auditorius. In cases of prolapsus ani, they likewise frequently attack that part of the gut which is exposed to the air. They have usually shewn the same morbid appearances inwardly, as are met with in putrid fever, where the disease has been of the malignant kind. Where the febrile symptoms have run high, and the head has been much affected with coma or delirium, the vessels of the brain appear, on removing the cranium and dura mater, more turgid, and filled with a darker coloured blood than usual, and a greater quantity of serous fluid is found, particularly towards the base of the brain. Under similar circumstances, the lungs have often a darker appearance, and their moisture is more copious than usual. When no inflammatory affection has supervened, they are most usually found.

VARIOLA VACCINA. The cow-pox. Any pustulous disease affecting the cow, may be called the cow-pox: whether it arises from an over-distention of the udder, in consequence of a neglect in milking the cow, or from the sting of an insect, or any other cause. But the species which claims our particular attention, is that which was recommended to the world by Dr. Jenner, in the year 1798, as a substitute for the small-pox. This, which originates from the grease in the horse's heel,

is called the *genuine cow-pox*; all other kinds are *spurious*.

That the vaccine fluid, fraught with such unspeakable benefits to mankind, derives its origin from this humble source, however it may mortify human pride, or medical vanity, is confirmed by the observations and experiments of competent judges. For proofs of this assertion, the reader may consult the works of Dr. Jenner; the Medical and Physical Journal; and a treatise on the subject by Dr. Loy, of which an analysis is given in the Annals of Medicine for the year 1801; and Mr. King's work on this disease, which contains the whole mass of evidence that has appeared concerning it.

The genuine cow-pox appears on the teats of the cow, in the form of vesicles, of a blue colour approaching to livid. These vesicles are elevated at the margin, and depressed at the centre. They are surrounded with inflammation. The fluid they contain is limpid. The animals are indisposed; and the secretion of milk is lessened. Solutions of the sulphats of zinc and copper are a speedy remedy for these pustules; otherwise they degenerate into ulcers, which are extremely troublesome. It must, however, be recollected, that much of the obstinacy attending these cases is owing to the friction of the pustules, in consequence of milking. It is probable, that a solution of the superacetate of lead would be preferable to irritating applications.

Similar effects are produced in the hands of the milkers, attended with febrile symptoms, and sometimes with tumours in the axilla. Other parts, where the cuticle is abraded, or which are naturally destitute of that defence, are also liable to the same affection, provided active matter is applied. It even appears that, in some instances, pustules have been produced by the application of vaccine virus to the sound cuticle. One case of this kind may be found in a letter from Dr. Fowler, of Salisbury, to Dr. Pearson, published in the first work of Dr. Pearson on this subject.

The spurious cow-pox is white: and another criterion is, that both in the brute animal and in the human subject, when infected with the casual cow-pox, the sores occasioned by the genuine species are more difficult to heal, than those which are occasioned by the spurious kind. It is of the utmost importance to distinguish the genuine from the spurious sort, which is also in some degree infectious; since a want of such discrimination would cause an idea of security against the small-pox, which might prove delusive.

Dr. Jenner has elucidated one point of the first importance, relative to the genuine cow-pox itself. It had frequently been observed, that when this disorder prevailed in a farm, some of the persons

who contracted it by milking were rendered insusceptible of the small-pox, while others continued liable to that infection. This is owing to the different periods at which the disease was excited in the human subject; one person, who caught the disease while the virus was in an active state, is rendered secure from variolous contagion; while another who received the infection of the cow-pox when it had undergone a decomposition, is still susceptible of the small pox. This uncertainty of the prevention, the value of which is beyond all calculation, is probably the reason why it was not before introduced into practice.

From the violent opposition which vaccine inoculation has met with, in consequence of certain apparent failures in the casual way, it may be doubted whether the public would ever have adopted the practice, had not this fallacy been detected by Dr. Jenner. To him also we are indebted for another discovery of the first importance, namely, that the pustule excited in the human subject by vaccine matter, yields a fluid of a similar nature with that which was inserted. This experiment, so essential to the general propagation of the practice, and so happy in its result, was never before attempted. It was reserved to crown the labours of Dr. Jenner.

A considerable number of instances are on record, to prove that farriers and others who receive infection from the heel of a horse, are either partly or totally deprived of the susceptibility of the small-pox. When Dr. Jenner first published an account of his discoveries, this point was enveloped in some degree of obscurity. He then conceived, that the matter of grease was an imperfect preservative against the small-pox. This opinion was founded on the following circumstance: It had been remarked, that farriers either wholly escaped the small-pox, or had that distemper in a milder manner than other people. This, however, is easily reconcilable to reason, if we only suppose, that in some cases the infection is communicated when the virus possesses all its prophylactic virtue; and in others, when its specific quality is in some measure lost.

This variation in the effects produced by the virus of the horse, inclined Dr. Jenner to believe that it was modified, and underwent some peculiar alteration in the teats of the cow. He now concludes, that it is perfect when it excites the genuine disease in the cow; yet a considerable advantage is derived from its being transferred to the latter animal, the nipples of which furnish a more obvious and a more abundant source of this inestimable fluid than its original element the horse.

This theory, that the preservative against variolous contagion is perfect when it issues from the fountain-head, and comes immediately from the hands of Nature, is consonant with reason, and consistent with analogy. Thus one obstacle more to the universal adoption of the practice is removed.

Another point respecting vaccine inoculation, which has been much controverted, is the permanency of its effect. Instances have been known where persons have escaped the small-pox for a number of years, and yet have ultimately proved not insusceptible of its infection. When such persons had previously undergone the vaccine disease, their apparent security was erroneously ascribed to that cause; but we have not even a shadow of proof, that the cow-pox possesses in the least degree the property of a temporary prophylactic, since it appears not even to retard the eruption of the small-pox, where previous infection has been received.

By this remark, it is not meant to be asserted, that it never supercedes or modifies the small-pox, for we have great reason to believe that such beneficial effects often flow from vaccination; but where an eruption of the small-pox actually takes place after vaccine inoculation, the two diseases frequently co-exist, without retarding each other in the smallest degree. It is therefore contrary to all reason and analogy, to consider the cow-pox as a mere temporary preservative: it is nothing less than a perfect and permanent security against that terrible disease.

A number of cases are recorded by Dr. Jenner, and other authors who have written on this subject, in which persons who had received the cow-pox by casual infection, twenty, thirty, forty, and fifty years before, still continued insusceptible of variolous contagion, in whatever form it was applied.

As the cow-pox destroys the susceptibility of the small-pox, so the small-pox destroys that of the cow-pox. To this general rule, however, a few exceptions are said to have occurred. Certain it is, that a pustule has now and then been excited by the insertion of vaccine virus, in those who have had the small-pox, and that this pustule has been known to yield the genuing virus; but it is not equally certain that the pustule has been perfect in all respects. Possibly it may have been defective in point of size or duration; in respect to its areola, or the limpidity of its contents. That such a pustule has, in some instances, yielded effectual virus, is admitted; but this is no more than what has often happened, in cases where persons who have had the small-pox are a second

time submitted to that infection in the same form.

The artificial cow-pox in the human subject is much milder than the casual disease; and incomparably milder than the small-pox, even under the form of inoculation. It neither requires medicine nor regimen; it may be practised at any season of the year; and, not being infectious by effluvia, one person may be inoculated without endangering the life of another.

This affection produces no pestilential eruptions. When such attend vaccine inoculation, they are owing to some adventitious cause, such as the small-pox, which it is well known may co-exist with the cow-pox. The vaccine vesicle is confined to the parts where matter is inserted; it is therefore entirely a local and an inoculated disease. Nevertheless it is certain, that eruptions of other kinds in some instances attend vaccine inoculation; such as a nettle rash, or an eruption resembling a tooth rash, but rather larger than what is commonly called by that name.

Among other singularities attending the cow-pox, the mildness of the disease, under the form of inoculation, has been urged as an argument against the practice, the cause appearing to ordinary comprehensions, inadequate to the effect. This, it must be allowed, is the best apology that can be offered for scepticism on the point; but it will weigh but little when put into the scale against actual observation, and incontrovertible fact. The efficacy of the cow-pox as a safeguard against the small-pox, rests, perhaps, on more extensive evidence, and a more solid foundation, than any other axiom in the whole circle of medical science can boast.

That the cow-pox is not infectious by effluvia, is naturally concluded from its never being communicated from one person to another in the dairies; where the disease is casual, and appears under its worst form. The same inference may be drawn from its never spreading in a family, when only one person is inoculated at a time. To confirm this proposition more fully, the vaccine pustules have been ruptured, and persons who have never had the disorder have been suffered to inhale the effluvia several times a day, but to no purpose. This is no more than might be expected, in an affection where the pustulous appearance on the surface of the body is nearly local.

As to the constitutional indisposition, it is seldom considerable, unless there is a complication of this with some other distemper; and whenever any unfavorable symptoms appear they may in general be traced to some other cause. We have

dead great reason to believe, that no ill consequence ever arises from the cow-pox itself, unless from ignorance or neglect.

But notwithstanding the symptoms are so mild, they frequently occur at a very early period. A drowsiness, which is one of the most common attendants of the disease, is often remarked by the parents themselves, within forty-eight hours after the matter is inserted. In a majority of cases, a slight increase of heat is perceptible, together with an acceleration of the pulse, and other signs of pyrexia; but not in such a degree as to alarm the most timorous mother. Sometimes the patient is restless at nights; and now and then a case is met with, in which vomiting occurs, but in many cases, no constitutional indisposition whatever can be perceived. Even then, the cow-pox has never failed to prove an effectual preservative against the small-pox, provided the pustule has been perfect.

This being the grand criterion of the security of the patient, too minute an attention cannot be paid to its rise, progress, and decline. The best mode of inoculating is by making a very small oblique puncture in the arm, near the insertion of the deltoid muscle, with the point of a lancet charged with fluid matter. In order to render infection more certain, the instrument may be charged again, and wiped upon the puncture.

In places where the patient is likely to be exposed to variolous contagion, it is advisable to inoculate in more places than one, but, unless there is imminent danger of catching the small-pox, it is better not to make more than one puncture in each arm, lest too much inflammation should ensue.

The vaccine fluid may be taken for inoculation as soon as a vesicle appears; but if the vesicle is punctured at a very early period, it is more apt to be injured. When virus is wanting for inoculating a considerable number, it is better to let the pustule remain untouched till about the eighth day, by which time it has in general acquired a reasonable magnitude. After that day, if the pustule has made the usual progress, the matter begins to lose its virtue; but it may, in general, be used with safety, though with less certainty of producing infection, till the areola begins to be extensive.

The first sign of infection commonly appears on the third day. A small red spot rather elevated, may be perceived at the place where the puncture was made. Sometimes, however, the mark of infection having succeeded is not visible till a much later period. It may be retarded, or even entirely prevented, by any other disorder, such as dentition, or any com-

plaint attended with fever, or by extreme cold. Another frequent cause of a slow progress in the pustule, or a total failure of success, is debility. Sometimes it is impossible to discover any sign of infection for above a fortnight. In this respect the cow-pox is subject to the same laws, and liable to the same variation, as the small-pox.

When a considerable inflammation appears within two or three days after inoculation, there is reason to suspect that infection has not taken place; and if suppuration ensues, that suspicion ought, in general, to stand confirmed. Now and then, however, it happens, that after the spurious pustule, or, more properly speaking, the phlegmon, has run its course, which is within a few days, a vesicle begins to appear, bearing every characteristic of the genuine vaccine disease, and yielding a limpid and efficient virus for future inoculations. In this case the patient is as perfectly secured from all danger of the small-pox, as if no festering of the puncture had preceded. The occurrence of such a case, though rare, is worthy to be recorded; because some practitioners have concluded a spurious pustule to be a certain proof of failure.

The areola commonly begins to be extensive on the ninth day, and to decline about the eleventh or twelfth. At this period also the pustule begins to dry; the first sign of which is a brown spot in the centre. In proportion as this increases, the surrounding efflorescence decreases, till at length nothing remains but a circular scab, of a dark brown mahogany colour, approaching to black. Sometimes it resembles the section of a tamarind stone; and it often retains the depression in the centre, which characterises this disease before exsiccation takes place.

Instances have been known, where the vaccine pustule, though regular, and perfect in all other respects, has been totally destitute of areola; at least, where neither the medical practitioner, on visiting the patient, nor the attendants, have remarked any appearance of that symptom. In these cases, the patient has proved as insusceptible of variolous infection, as if the surrounding efflorescence had covered the whole arm. It must, however, be confessed, that we have no proof of the non-existence of an areola in these cases. It might have been trivial; it might have been transient; yet it might have been effectual. There is, however, greater reason to believe, that the surrounding efflorescence, though usually a concomitant circumstance, is not an essential requisite to the vaccine disease.

If by any accident the vesicle is ruptured, suppuration often ensues. In this case more attention than ordinary ought

to be paid to the progress, and to all the phenomena of the local affection; both on account of the uncertainty of success in the pustule, as a prophylactic; and the greater probability of tedious ulceration.

If there is room for the least doubt of the sufficiency of the first inoculation, a second ought to be performed without delay. This, if unnecessary, is seldom attended with inconvenience, and never with danger. Either no effect is produced, or a slight festering, which terminates in a few days. An exception occurs, but rarely, where a spurious, or, perhaps, even a genuine pustule, takes place, in those persons who are known to have had the cow-pox or the small-pox already; but this cannot be the least cause of alarm to any one who knows the benign character of the distemper.

Various topical applications, both stimulant and sedative, have been recommended, in order to allay the violence of inflammation. If the operation for the insertion of matter is not unnecessarily severe, nor the pustule irritated by friction or pressure, or other violence, no such applications are necessary. Nevertheless, if either the anxiety of the professional man, or the importunity of a tender parent, should demand a deviation from this general rule, any of the following remedies may be had recourse to. The pustule may be touched with very diluted sulphuric acid; which should be permitted to remain on the part half a minute, and then be washed off with a sponge dipped in cold water. This has been ignorantly, or artfully, called an escharotic; but any one who tries the application will soon discover, that its operation is mild and harmless.

To avoid cavil and misrepresentation, it is better to apply a saturnine lotion; either made with aq. litharg. comp. or cerusa acetata; compresses, dipped in such a lotion, may be applied at any time when inflammation runs high, and renewed as occasion requires.

If the pustule should chance to be broken, a drop of aq. litharg. acet. undiluted, may be applied as an esiccant; but if ulceration threatens to become obstinate, or extensive, a mild cataplasm is the best resource. In case the ulceration is only superficial, and not attended with immoderate inflammation, a bit of any adhesive plaister, spread on linen, will prove the most convenient dressing, and seldom fail of success. It will, in general, be unnecessary to renew it oftener than every other day.

These minute observations no one will despise, unless there be any person so ignorant as not to know that the care of the arm is almost the whole duty of the

medical practitioner in vaccine inoculation; and that nothing disgusts the public so much against the practice, as a sore arm, and the ill consequences which, from a neglect of that symptom, too often ensue.

When fluid virus cannot be procured, it is necessary to be cautious how it is preserved in a dry state. The most improper mode is that of keeping it on a lancet; for the metal quickly rusts, and the vaccine matter becomes decomposed. This method, however, is as likely to succeed as any, when the matter is not to be kept above two or three days. If the virus be taken on glass, care must be taken not to dilute it much; otherwise it will in all probability fail.

Cotton thread is a very commodious vehicle. If it is intended to be sent to any considerable distance, it ought to be repeatedly dipped in the virus. No particular caution is necessary with regard to the exclusion of air; nevertheless, as it can be done with so little trouble, and is more satisfactory to those who receive the matter, it is better to comply with the practice. On this account it may be enclosed in a glass tube, or in a tobacco-pipe sealed at each end, or between two square bits of glass, which may, if necessary, be also charged with the matter, and wrapped in gold-beater's skin.

Nothing is more destructive to the efficacy of cow-pox matter than heat: on this account it must not be dried near the fire, nor kept in a warm place. The advantage of inserting it in a fluid state is so great, that it is to be wished every practitioner would endeavour to keep a constant supply for his own use, by inoculating his patients in succession, at such periods as are most likely to answer that purpose.

The rapidity with which this practice now spreads in various parts of the globe, justifies our cherishing a hope, that it will ere long extinguish that most dreadful pestilence, and perpetual bane of human felicity, the small-pox.

VARIUS. (From *varus*, unequal, so called from the irregularity of its shape.) The cuboid bone is called os varium, from its irregular shape.

VARIX. (From *varus*, i. e. *obtusus*.) A dilatation of a vein. A genus of disease in the class *locales* and order *tumores* of Cullen; known by a soft tumour on a vein which does not pulsate. Varicose veins mostly become serpentine, and often form a plexus of knots, especially in the groins and scrotum.

VAS DEFERENS. (*Vas*, a vessel, and *deferens*, from *defere*, to convey.) A duct which arises from the epididymis, and passes through the inguinal ring in the spermatic cord into the cavity of the pelvis;

and terminates in the *vesiculæ seminales*. Its use is to convey the semen secreted in the testicle, and brought to it by the *epididymis* into the *vesiculæ seminales*.

VASA BREVIA. The arteries which come from the spleen, and run along the large arch of the stomach to the diaphragm.

VASA DEFERENTIA. See *Vas deferens*.

VASA VORTICOSA. The contorted vessels of the choroid membrane of the eye.

VASTUS EXTERNUS. (*Vastus*, so called from its size) A large, thick, and fleshy muscle situated on the outer side of the thigh: it arises, by a broad thick tendon, from the lower and anterior part of the great trochanter, and upper part of the *linea aspera*; it likewise adheres by fleshy fibres, to the whole outer edge of that rough line. Its fibres descend obliquely forwards, and after it has run four or five inches downwards, we find it adhering to the anterior surface and outer side of the *cruræus*, with which it continues to be connected to the lower part of the thigh, where we see it terminating in a broad tendon, which is inserted into the upper part of the patella laterally, and sends off an aponeurosis that adheres to the head of the tibia, and is continued down the leg.

VASTUS INTERNUS. This muscle, which is less considerable than the *vastus externus*, is situated at the inner side of the thigh, being separated from it by the *rectus*.

It arises tendinous and fleshy from between the fore-part of the *os femoris*, and the root of the lesser trochanter, below the insertion of the *psaos magnus*, and the *iliacus internus*; and from all the inner side of the *linea aspera*. Like the *vastus externus* it is connected with the *cruræus*, but it continues longer fleshy than that muscle. A little above the knee we see its outer edge uniting with the inner edge of the *rectus*, after which it is inserted tendinous into the upper part and inner side of the patella, sending off an aponeurosis which adheres to the upper part of the tibia.

VEIN. (*Vena*, from *venio*, to come, because the blood comes through it.) Veins are long membranous canals, which continually become wider, do not pulsate, and return the blood from the arteries to the heart. All veins originate from the extremities of arteries only, by anastomosis, and terminate in the auricles of the heart; e. g. the *vena cava* in the right, and the pulmonary veins in the left auricle. They are composed, like arteries, of three tunics or coats, which are much more slender than in the arteries, and are supplied with semilunar membranes or folds called valves.

Their use is to return the blood to the heart.

The blood is returned from every part of the body, except the lungs, into the right auricle, from three sources:

1. The *vena cava superior*, which brings it from the head, neck, thorax, and superior extremities.

2. The *vena cava inferior*, from the abdomen and inferior extremities.

3. The *coronary vein* receives it from the coronary arteries of the heart.

1. The *vena cava superior*. This vein terminates in the superior part of the right auricle, into which it evacuates the blood, from the right and left *subclavian veins*, and the *vena axylæ*. The right and left subclavian veins receive the blood from the head and upper extremities, in the following manner. The veins of the fingers, called *digitals*, receive their blood from the digital arteries, and empty it into,

The *cephalic of the thumb*, which runs on the back of the hand along the thumb, and evacuates itself into the external radial.

The *salvatorella*, which runs along the little finger, unites with the former, and empties its blood into the internal and external cubital veins. At the bend of the fore-arm are three veins, called the great cephalic, the basilic, and the median.

The *great cephalic* runs along the superior part of the fore-arm, and receives the blood from the external radial.

The *basilic* ascends on the under side, and receives the blood from the external and internal cubital veins, and some branches which accompany the brachial artery, called *venæ satellitum*.

The *median* is situated in the middle of the fore-arm, and arises from the union of several branches. These three veins all unite above the bend of the arm, and form

The *brachial vein*, which receives all their blood, and is continued into the axilla, where it is called

The *axillary vein*. This receives also the blood from the scapula, and superior and inferior parts of the chest, by the *superior and inferior thoracic vein*, the *vena muscularis*, and the *scapularis*.

The axillary vein then passes under the clavicle, where it is called the *subclavian*, which unites with the external and internal jugular veins, and the vertebral vein which brings the blood from the vertebral sinuses; it receives also the blood from the *mediastinal*, *pericardiac*, *diaphragmatic*, *thymic*, *internal mammary* and *laryngeal* veins, and then unites with its fellow, to form the *vena cava superior*, or, as it is sometimes called, *vena cava descendens*.

The blood from the external and in-

ternal parts of the head and face is returned in the following manner into the external and internal jugulars, which terminate in the subclavians.

The *frontal, angular, temporal, auricular, sublingual, and occipital* veins receive the blood from the parts after which they are named; these all converge to each side of the neck, and form a trunk, called the *external jugular vein*.

The blood from the brain, cerebellum, medulla oblongata, and membranes of these parts, is received into the lateral sinuses, or veins of the dura mater, one of which empties its blood through the foramen lacerum in basi crani into the *internal jugular*, which descends in the neck by the carotid arteries, receives the blood from the *thyroidal and internal maxillary veins*, and empties itself into the subclavians within the thorax.

The *vena azygos* receives the blood from the *bronchial, superior œsophageal, vertebral and intercostal veins*, and empties it into the superior cava.

2. *Vena cava inferior*. The *vena cava inferior* is the trunk of all the abdominal veins and those of the lower extremities, from which parts the blood is returned in the following manner. The veins of the toes, called the *digital veins*, receive the blood from the digital arteries, and form on the back of the foot three branches, one on the great toe called the *cephalic*, another which runs along the little toe, called the *vena saphena*, and on the back of the foot, *vena dorsalis pedis*; and on the sole of the foot they evacuate themselves into the *plantar veins*.

The three veins on the upper part of the foot coming together above the ankle, form the *anterior tibial*; and the plantar veins with a branch from the calf of the leg, called the *sural vein*, form the *posterior tibial*; a branch also ascends in the direction of the fibula, called the *peroneal vein*. These three branches unite before the ham, into one branch, the *subpopliteal vein*, which ascends through the ham, carrying all the blood from the foot: it then proceeds upon the anterior part of the thigh, where it is termed the *crural or femoral vein*, receives several muscular branches, and passes under Poupart's ligament into the cavity of the pelvis, where it is called the *external iliac*.

The arteries which are distributed about the pelvis evacuate their blood into the *external hemorrhoidal veins, the hypogastric veins, the internal pudendal, the vena magna ipsius penis, and obtratory veins*, all of which unite in the pelvis, and form the *internal iliac vein*.

The external iliac vein receives the blood from the external pudendal veins, and then unites with the internal iliac at

the last vertebra of the loins, and form the *vena cava inferior or ascendens*, which ascends on the right side of the spine, receiving the blood from the *sacral lumbar right spermatic veins*, and the *vena cava hepatica*; and having arrived at the diaphragm, it passes through the right foramen, and enters the right auricle of the heart, into which it evacuates all the blood from the abdominal viscera and lower extremities.

Vena cava hepatica. This vein ramifies in the substance of the liver, and brings the blood into the *vena cava inferior* from the branches of the *vena portæ*, a great vein which carries the blood from the abdominal viscera into the substance of the liver. The trunk of this vein, about the fissure of the liver in which it is situated, is divided into the hepatic and abdominal portions. The *abdominal portion* is composed of the *splenic, meseraic, and internal hemorrhoidal veins*. These three venous branches carry all the blood from the stomach, spleen, pancreas, omentum, mesentery, gall-bladder, and the small and large intestines, into the *sinus* of the *vena portæ*. The *hepatic portion* of the *vena portæ* enters the substance of the liver, divides into innumerable ramifications, which secrete the bile, and the superfluous blood passes into corresponding branches of the *vena cava hepatica*.

The *action of the veins*. Veins do not pulsate; the blood which they receive from the arteries flows through them very slowly, and is conveyed to the right auricle of the heart, by the contractility of their coats, the pressure of the blood from the arteries, called the *vis a tergo*, the contraction of the muscles, and respiration; and it is prevented from going backwards in the vein by the valves, of which there are a great number.

VERVEA DU GUACO. A plant which has the power of curing and preventing the bite of venomous serpents.

VELAMENTUM BOMBYCINUM. The interior soft membrane of the intestines.

VELUM PENDULUM PALATI. *Velum palatinum*. The soft palate. The soft part of the palate, which forms two arches, affixed laterally to the tongue and pharynx.

VENA AZYGOS. See *Azygos vein*.

VENA MEDINENSIS. See *Medinensis vena*.

VENA PORTÆ. (*Vena portæ*, a *portando*, because through it things are carried.) *Vena portarum*. The great vein, situated at the entrance of the liver, which receives the blood from the abdominal viscera, and carries it into the substance of the liver. It is distinguished into the *hepatic and abdominal portion*: the former is ramified through the substance of the liver, and carries the blood

destined for the formation of the bile, which is returned by branches to the trunk of the vena cava; the latter is composed of three branches; viz. the splenic, mesenteric, and internal hæmorrhoidal veins. See *Vena*.

VENÆ LACTEÆ. The lacteal absorbents were so called. See *Lacteals*.

Veneral disease. See *Gonorrhœa* and *Syphilis*.

VENTER. A term formerly applied to the larger circumscribed cavities of the body, as the abdomen and thorax.

VENTRICLE. A term given by anatomists to the cavities of the brain and heart. See *Cerebrum* and *Heart*.

VENTRICULUS PULMONARIS. The right ventricle of the heart.

VENTRICULUS SUCCENTURIATUS. That portion of the duodenum, which is surrounded by the peritoneum, is sometimes so large as to resemble a second stomach, and is so called by some writers.

VENUS. Copper was formerly so called by the chemists.

VERATRUM 1. The name of a genus of plants in the Linnæan system. Class, *Polygynia*. Order, *Monœcia*.

2. The pharmacopœial name of white hellebore. See *Helleborus albus*.

VERATRUM ALBUM. See *Helleborum albus*.

VERATRUM NIGRUM. See *Helleborus niger*.

VERBASCUM. (Quasi *barbascum*, from its hairy coat.) 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*. Mullien.

2. The pharmacopœial name of the *Candlea regia*: *Thapsus barbatus*: *Candelaria*: *Lunaria*: The *Verbascum nigrum* and *Verbascum thapsus* appear to be ordered indifferently by this name in the pharmacopœias. The flowers, leaves, and roots, are used occasionally as mild adstringents. The leaves possess a roughish taste, and promise to be of service in diarrhœas and other debilitated states of the intestines.

VERBASCUM NIGRUM. The systematic name of the black mullein. See *Verbascum*.

VERBASCUM THAPSUS. The systematic name of the yellow mullein. See *Verbascum*.

VERBENA. (Quasi *herbena*, a name of distinction for all herbs used in sacred rites.) 1. The name of a genus of plants in the Linnæan system. Class, *Dicandria*. Order, *Monogynia*.

2. The pharmacopœial name of what is also called *Verbenaca*. *Peristerium*. *Hierobastane cephalalgia*. *Herba sacra*. *Vervain*. This plant *Verbena officinalis* of Linnæus, is destitute of odour, and to the taste manifests but a slight degree of bitterness and adstringency. In former times the verbena seems to have been

held sacred, and was employed in celebrating the sacrificial rites; and with a view to this, more than the natural power of the plant, it was worn suspended about the neck as an amulet.

This practice, thus founded on superstition, was, however, in process of time, adopted in medicine; and therefore to obtain its virtues more effectually, the vervain was directed to be bruised before it was appended to the neck; and of its good effects thus used for inveterate headaches, Forestus relates a remarkable instance. In still later times it has been employed in the way of cataplasm, by which we are told the most severe and obstinate cases of cephalalgia have been cured, for which we have the authorities of Etmüller, Hartman, and more especially De Hean. Notwithstanding these testimonies in favour of vervain, it has deservedly fallen into disuse in Britain; nor has the pamphlet of Mr. Morley, written professedly to recommend its use in scrophulous affections, had the effect of restoring its medical character. This gentleman directs the root of vervain to be tied with a yard of white satin riband round the neck, where it is to remain till the patient recovers. He also has recourse to infusions and ointments prepared from the leaves of the plant, and occasionally calls in aid the most active medicines of the *Materia Medica*.

VERBENA FEMINA. The hedge mustard is sometimes so called. See *Erysimum*.

VERBENA OFFICINALIS. The systematic name of vervain. See *Verbena*.

VERDIGRIS. An impure subacetate of copper. It is prepared by stratifying copper plates with the husks of grapes, after the expression of their juice, and when they have been kept for some time imperfectly exposed to the air, in an apartment warm but not too dry, so as to pass to a state of fermentation, whence a quantity of vinegar is formed. The copper plates are placed in jars in strata, with the husks thus prepared, which are covered. At the end of twelve, fifteen, or twenty days these are opened: the plates have an efflorescence on their surfaces of a green colour and silky lustre: they are repeatedly moistened with water; and at length, a crust of verdigris is formed, which is scraped off by a knife, is put into bags, and dried by exposure of these to the air and sun. It is of a green colour, with a slight tint of blue.

In this preparation the copper is oxidised, probably by the atmospheric air, aided by the affinity of the acetic acid; and a portion of this acid remains in combination with the oxyde, not sufficient, however, to produce its saturation. When acted on by water, the acid with such a

portion of oxyde as it can retain in solution, are dissolved, and the remaining oxyde is left undissolved. From this analysis of it by the action of water, Proust inferred that it consists of 43 of acetate of copper, 27 of black oxyde of copper, and 30 of water, this water not being accidental, but existing in it in intimate combination.

Verdigris is used as a pigment in some of the processes of dyeing, and in surgery it is externally applied as a mild detergent in cleansing foul ulcers, or other open wounds. On account of its virulent properties, it ought not to be used as a medicine without professional advice; and in case any portion of this poison be accidentally swallowed, emetics should be first given, and afterwards cold water gently alkalized ought to be drank in abundance.

VERJUICE. An acid liquor prepared from grapes or apples, that are unfit to be converted into wine or cyder. It is also made from crabs. It is principally used in sauces and ragous, though it sometimes forms an ingredient in medicinal compositions.

VERMICULARIS. See *Illecebra*.

VERMIFORM PROCESS. *Protuberantia vermiformis*. The substance which unites the two hemispheres of the cerebellum like a ring, forming a process. It is called *vermiform*, from its resemblance to the contortions of worms.

VERMIFUGES. (*Vermifuga*, from *vermis*, a worm, and *fugo*, to drive away.) See *Anthelmintics*.

Vermilion. See *Cinnabar*.

VERMIS MORDICANS. *Vermis repens*. A species of herpetic eruption on the skin.

VERMIS TERRESTRIS. See *Earth-worm*.

VERONICA. 1. The name of a genus of plants in the Linnæan system. Class, *Diandria*. Order, *Monogynia*. Speedwell.

2. The pharmacopœial name of the *Veronica mas*. *Thea Germanica*. *Betonica pauli chamedryæ*. *Veronica officinalis* of Linnaeus; *spiciis lateralibus*; *pedunculatis foliis oppositis*; *caule procumbente* is not infrequent on dry barren grounds and heaths, as that of Hampstead, flowering in June and July. This plant was formerly used as a pectoral against coughs and asthmatic affections, but it is now justly forgotten.

VERRICULARIS TUNICA. The retina of the eye.

VERTEBRÆ. (From *verto*, to turn.) The spine is a long bony column, which extends from the head to the lower part of the trunk, and is composed of irregular bones which are called *vertebræ*.

The spine may be considered as being composed of two irregular pyramids, which are united to each other in that part of the loins where the last of the lumbar *vertebræ* is united to the os sacrum.

The *vertebræ*, which form the upper and longest pyramid, are called true *vertebræ*, and those which compose the lower pyramid, or the os sacrum and coccyx, are termed false *vertebræ*, because they do not in every thing resemble the others, and particularly because, in the adult state, they become perfectly immoveable, while the upper ones continue to be capable of motion. For it is upon the bones of the spine that the body turns, and their name has its derivation from the Latin verb *verto*, to turn, as observed above.

The true *vertebræ*, from their situations with respect to the neck, back, and loins, are divided into three classes, of cervical, dorsal, and lumbar *vertebræ*. We will first consider the general structure of all these, and then separately describe their different classes.

In each of the *vertebræ*, as in other bones, we may remark the body of the bone, its processes and cavities. The body may be compared to part of a cylinder cut off transversely; convex before, and concave behind, where it makes part of the cavity of the spine.

Each *vertebra* has commonly seven processes. The first of these is the *spinous* process, which is placed at the back part of the *vertebra*, and gives the name of spine to the whole of this *bony* canal. Two others are called *transverse* processes, from their situation with respect to the spine, and are placed on each side of the spinous process. The four others, which are called *oblique* processes, are much smaller than the other three. There are two of these on the upper and two on the lower part of each *vertebra*, rising from near the basis of the transverse processes. They are sometimes called *articular* processes, because they are articulated with each other; that is, the two superior processes of one *vertebra* are articulated with the two inferior processes of the *vertebra* above it: and they are called *oblique* processes, from their situation with respect to the processes with which they are articulated. These oblique processes are articulated to each other by a species of *ginglimus*, and each process is covered at its articulation with cartilage.

There is in every *vertebra*, between its body and apophyses, a foramen, large enough to admit a finger. These foramina correspond with each other through all the *vertebræ*, and form a long bony conduit, for the lodgment of the spinal marrow.

Besides this great hole, there are four notches on each side of every *vertebra* between the oblique processes and the body of the *vertebra*. Two of these notches are at the upper, and two at the lower part of the bone. Each of the superior notches, meeting with one of the superior notches of the *vertebra* below

forms a foramen; whilst the superior notches do the same with the inferior notches of the vertebra above it. These four foramina form passages for blood-vessels, and for the nerves that pass out of the spine.

The vertebrae are united together by means of a substance, compressible like cork, which forms a kind of partition between the several vertebrae. This intervertebral substance seems, in the fœtus, to approach nearly to the nature of ligaments; in the adult it has a great resemblance to cartilage. When cut horizontally, it appears to consist of concentric curved fibres, externally it is firmest and hardest; internally it becomes thinner and softer, till at length, in the centre, we find it in the form of a mucous substance, which facilitates the motions of the spine.

Genga, an Italian anatomist, long ago observed, that the change which takes place in these intervertebral cartilages, (as they are usually called) in advanced life, occasions the decrease in stature, and the stooping forwards, which are usually to be observed in old people. The cartilages then become shrivelled, and consequently lose, in a great measure, their elasticity. But, besides this gradual effect of old age, these cartilages are subject to a temporary diminution, from the weight of the body in an erect posture, so that people who have been long standing, or who have carried a considerable weight, are found to be shorter than when they have been long in bed. Hence we are taller in the morning than at night. This fact, though seemingly obvious, was not ascertained till of late years. The difference in such cases depends on the age and size of the subject; in tall, young people, it will be nearly an inch; but in older, or shorter persons, it will be less considerable.

Besides the connexion of the several vertebrae, by means of these cartilages, there are likewise many strong ligaments, which unite the bones of the spine to each other. Some of these ligaments are external, and others internal. Among the external ligaments, we observe one which is common to all the vertebrae, extending, in a longitudinal direction, from the fore part of the body of the second vertebra of the neck, over all the other vertebrae, and becoming broader as it descends towards the os sacrum, where it becomes thinner, and gradually disappears. This external longitudinal ligament, if we may so call it, is strengthened by other shorter ligamentous fibres, which pass from one vertebra to another, throughout the whole spine. The internal ligament, the fibres of which, like the external one, are spread

in a longitudinal direction, is extended over the back part of the bodies of the vertebrae, where they help to form the cavity of the spine, and reaches from the foramen magnum of the occipital bone to the os sacrum.

We may venture to remark, that all the vertebrae diminish in density and firmness of texture, in proportion as they increase in size, so that the lower vertebrae, though larger, are not so heavy in proportion as those above them. In consequence of this mode of structure, the size of the vertebrae is increased without adding to their weight; and this is an object of no little importance, in a part of the body, which, besides flexibility and suppleness, seems to require lightness as one of its essential properties.

In the fœtus, at the ordinary time of birth, each vertebra is found to be composed of three bony pieces, connected by cartilages which afterwards ossify. One of these pieces is the body of the bone; the other two are the posterior and lateral portions, which form the foramen for the medulla spinalis. The oblique processes are at that time complete, and the transverse processes beginning to be formed; but the spinous processes are totally wanting.

The cervical vertebrae, are seven in number, their bodies are smaller and of a firmer texture than the other bones of the spine. The transverse processes of these vertebrae are short, and forked for the lodgment of muscles; and, at the bottom of each of these processes, there is a foramen, for the passage of the cervical artery and vein. The spinous process of each of these vertebrae is likewise shorter than the other vertebrae, and forked at its extremity; by which means it allows a more convenient insertion to the muscles of the neck. Their oblique processes are more deserving of that name than either those of the dorsal or lumbar vertebrae. The uppermost of these processes are slightly concave, and the lowermost slightly convex. This may suffice for a general description of these vertebrae; but the first, second, and seventh, deserve to be spoken of more particularly. The first, which is called *Atlas*, from its supporting the head, differs from all the other vertebrae of the spine. It forms a kind of bony ring, which may be divided into its anterior and posterior arches, and its lateral portions. Of these, the anterior arch is the smallest and flattest; at the middle of its convex fore part we observe a small tubercle which is here what the body is in the other vertebrae. To this tubercle a ligament is attached, which helps to strengthen the articulation of the spine with the os occipitis. The back part of this anterior portion is concave,

and covered with cartilage, where it receives the odontoid process of the second vertebra. The posterior portion of the vertebra, or, more properly speaking, the posterior arch, is larger than the anterior one. Instead of a spinous process, we observe a rising, or tubercle, larger than that which we have just now described, on the fore part of the bone. The lateral portions of the vertebra project, so as to form what are called the transverse processes, one on each side, which are longer and larger than the transverse processes of the other vertebra. They terminate in a roundish tubercle, the end of which has a slight bend downwards. Like the other transverse processes, they are perforated at their basis, for the passage of the cervical artery. But, besides these transverse processes, we observe, both on the superior and inferior surface of these lateral portions of the first vertebra, an articulating surface, covered with cartilage, answering to the oblique processes in the other vertebra. The uppermost of these are oblong, and slightly concave, and their external edges rise somewhat higher than their internal brims. They receive the condylod processes of the os occipitis, with which they are articulated by a species of ginglymus. The lowermost articulating surfaces, or the inferior oblique processes, as they are called, are large, concave, and circular, and are formed for receiving the superior oblique processes of the second vertebra; so that the atlas differs from the rest of the cervical vertebra in receiving the bones, with which it is articulated both above and below.

In the fœtus we find this vertebra composed of five, instead of three pieces, as in the other vertebra. One of these is the anterior arch; the other four are the posterior arch and the sides, each of the latter being composed of two pieces. The transverse process, on each side, remains long in a state of epiphysis with respect to the rest of the bone. The second vertebra is called *dentata*, from the process on the upper part of its body, which has been, though perhaps improperly, compared to a tooth. This process, which is the most remarkable part of the vertebra, is of a cylindrical shape, slightly flattened, however, behind and before. Anteriorly it has a convex, smooth, articulating surface, where it is received by the atlas, as we observed in our description of that vertebra. It is by means of this articulation that the rotatory motion of the head is performed; the articulation of the os occipitis with the superior oblique processes of the first vertebra, allowing only a certain degree of motion backwards and forwards, so that when we turn the face either to the right or left, the

atlas moves upon this odontoid process of the second vertebra. But, as the face cannot turn a quarter of a circle, that is, to the shoulder, upon this vertebra alone, without being liable to injure the *medulla spinalis*, we find that all the cervical vertebra concur in this rotatory motion, when it is in any considerable degree; and indeed we see many strong ligamentous fibres arising from the sides of the odontoid process, and passing over the first vertebra, to the os occipitis, which not only strengthen the articulation of these bones with each other, but serve to regulate and limit their motion. It is on this account that the name of *moderators* has sometimes been given to these ligaments.

The transverse processes of the vertebra *dentata* are short, inclined downwards, and forked at their extremities. Its spinous process is short and thick. Its superior oblique processes a slightly convex, and somewhat larger than the articulating surfaces of the first vertebra, by which mechanism, the motion of that bone upon this second vertebra is performed with greater safety. Its inferior oblique processes have nothing singular in their structure.

The seventh vertebra of the neck differs from the rest chiefly in having its spinous process of a greater length, so that, upon this account, it has been sometimes called *vertebra prominens*.

The dorsal vertebra, which are twelve in number, are of a middle size, between the cervical and lumbar vertebra; the upper ones gradually losing their resemblance to those of the neck, and the lower ones coming nearer to those of the loins. The bodies of these vertebra are more flattened at their sides, more convex before, and more concave behind, than the other bones of the spine. Their upper and lower surfaces are horizontal. At their sides we observe two depressions, one at their upper, and the other at their lower edge, which, united with similar depressions in the vertebra above and below, form articulating surfaces, covered with cartilage, in which the heads of the ribs are received. These depressions, however, are not exactly alike in all the dorsal vertebra; for we find the head of the first rib articulated solely with the first of these vertebra, which has therefore the whole of the superior articulating surface within itself, independent of the vertebra above it. We may likewise observe a similarity in this respect in the eleventh and twelfth of the dorsal vertebra, with which the eleventh and twelfth ribs are articulated separately. Their spinous processes are long, flattened at the sides, divided at their upper and back part into two surfaces by a middle ridge, which is

received by a small groove in the inner part of the spinous process immediately above it, and connected to it by a ligament. Those spinous processes are terminated by a kind of round tubercle, which slopes considerably downwards, except in the three lowermost vertebræ, where they are shorter and more erect. Their transverse processes are of considerable length and thickness, and are turned obliquely backwards. Anteriorly they have an articulating surface, for receiving the tuberosity of the ribs, except in the eleventh and twelfth of the dorsal vertebræ to which the ribs are articulated by their heads only. In the last of these vertebræ the transverse processes are very short and thick, because otherwise they would be apt to strike against the lowermost ribs, when we bend the body to either side.

The *lumbar vertebræ*, the lowest of the true vertebræ, are five in number. They are larger than the dorsal vertebræ. Their bodies are extremely prominent, and nearly of a circular form at their fore part; posteriorly they are concave. Their intermediate cartilages are of considerable thickness, especially anteriorly, by which means the curvature of the spine forwards, towards the abdomen, in this part, is greatly assisted. Their spinous processes are short and thick, of considerable breadth, erect, and terminated by a kind of tuberosity. Their oblique processes are of considerable thickness; the superior ones are concave, and turned inwards; the inferior ones convex, and turned outwards. Their transverse processes are thin and long, except in the first and last vertebræ, where they are much shorter, that the lateral motions of the trunk might not be impeded. The inferior surface of all these vertebræ is slightly oblique, so that the fore part of the body of each is somewhat thicker than its hind part; but this is more particularly observable in the lowermost vertebra, which is connected with the *os sacrum*. Many anatomists describe the *os sacrum* and the *os coccygis* as when considering the bones of the spine, whilst others regard them as belonging more properly to the pelvis. These bones the reader may consult. It now remains to notice the uses of the spine. We find the spinal marrow lodged in this bony canal, secure from external injury. It defends the thoracic and abdominal viscera, and forms a pillar which supports the hand, and gives a general firmness to the whole trunk.

To give it a firm basis, we find the bodies of the vertebræ gradually increasing in breadth as they descend; and to fit it for a variety of motion, it is composed of a great number of joints, with an intermediate elastic substance, so that to great

firmness there is added a perfect flexibility.

We have already observed, that the lowermost and largest vertebræ are not so heavy in proportion as those above them; their bodies being more spongy, excepting at their circumference, where they are more immediately exposed to pressure; so that nature seems every where endeavouring to relieve us of an unnecessary weight of bone. But behind, where the spinal marrow is more exposed to injury, we find the processes composed of very hard bone; and the spinous processes are in general placed over each other in a slanting direction, so that a pointed instrument cannot easily get between them, excepting in the neck, where they are almost perpendicular and leave a greater space between them. Hence, in some countries, it is usual to kill cattle by thrusting a pointed instrument between the occiput and the atlas, or between the atlas and the second vertebra. Besides these uses of the vertebræ in defending the spinal marrow, and in articulating the several vertebræ, as is the case with the oblique processes, we shall find that they all serve to form a greater surface for the lodgment of muscles, and to enable the latter to act more powerfully on the trunk, by affording them a lever of considerable length.

In the neck, we see the spine projecting somewhat forwards, to support the head, which, without this assistance, would require a greater number of muscles. Through the whole length of the thorax it is carried in a curved direction backwards, and thus adds considerably to the cavity of the chest, and consequently affords more room to the lungs, heart, and large blood-vessels. In the loins, the spine again projects forwards, in a direction with the centre of gravity, by which means the body is easily kept in an erect posture; for otherwise we should be liable to fall forwards. But, at its inferior part, it again recedes backwards, and helps to form a cavity called the pelvis, in which the urinary bladder, intestinum rectum, and other viscera, are placed.

In a part of the body that is composed of so great a number of bones, and constructed for such a variety of motion, as the spine is, luxation is more to be expected than fracture; and this is very wisely guarded against in every direction, by the many processes that are to be found in each vertebra, and by the cartilages, ligaments, and other means of connexion, which we have described as uniting them together.

VERTEBRAL ARTERY. *Arteria vertebralis*. A branch of the subclavian, proceeding through the vertebræ to within the cranium, where, with its fellow, it

forms the basilar artery, the internal auditory, and the posterior artery of the dura mater.

VERTEX. The crown of the head.

VERTICALIA OSSA. See *Parietal bones*.

VERTICIS OS. See *Parietal bones*.

VERTIGO. Giddiness.

Vervain. See *Verbena*.

Vervain, female. See *Erysimum*.

VESANIÆ. (From *Vesanus*, a madman.)

The fourth order in the class *neuroses* of Cullen's nosological arrangement; comprehending diseases in which the judgment is unpaired, without either coma or pyrexia.

VESICA. (Dim. of *vas*, a vessel.) A bladder.

VESICA FELLIS. The gall-bladder. See *Gall-bladder*.

VESICA URINARIA. The urinary bladder. See *Urinary bladder*.

VESICATORIES. (*Vesicatoria*, from *vesica*, a bladder; because they raise a bladder.) See *Epiplastics*.

VESICLE (*Vesicula*, a diminutive of *vesica*, a bladder.) An elevation of the cuticle, containing a transparent watery fluid.

VESICULA FELLIS. The gall-bladder.

VESICULÆ DIVÆ BARBARÆ. The contagious small-pox.

VESICULÆ OINGIVARUM. The thrush.

VESICULÆ PULMONALES. The air cells which compose the greatest part of the lungs, and are situated at the termination of the bronchia.

VESICULÆ SEMINALES. Two membranous receptacles, situated on the back part of the bladder above its neck. The excretory ducts are called ejaculatory ducts. They proceed to the urethra, into which they open by a peculiar orifice at the top of the verumontanum. They have vessels and nerves from the neighbouring parts, and are well supplied with absorbent vessels, which proceed to the lymphatic glands about the loins. The use of the vesiculæ seminales is to receive the semen brought into them by the vasa deferentia, to retain, somewhat inspissate, and to excrete it *sub coitu* into the urethra, from whence it is propelled into the vagina uteri. See *Pemphigus*.

Vesicular fever. See *Pemphigus*.

VESTIBULUM. A round cavity of the internal ear, between the cochlea and semicircular canals, in which are an oval opening communicating with the cavity of the tympanum and the orifices of the semicircular canals. It is within this cavity and the semicircular canals, that the new apparatus, discovered by the celebrated neurologist Scarpa, lies. He has demonstrated membranous tubes, connected loosely by cellular texture, within the bony semicircular canals, each of which is dilated in the cavity of the vestibule into

an ampulla; it is upon these ampullæ, which communicate by means of an *abacus communis*, that branches of the portio molilis are expanded.

VERONICA CORNI. See *Betonica*.

VIBIX. (*Vibex*, sing. plu. *Vibices*.) The large purple spots which appear under the skin in certain malignant fevers.

VIBRISSÆ. (*Vibrissæ*, from *vibro*, to quaver.) Hairs growing in the nostrils. See *Capillus*.

VICHY WATER. Is obtained from the tepid mineral springs that arise in the vicinity of Vichy, in France. On account of its chalybeate and alkaline ingredients, it is taken internally, being reputed to be of great service in bilious colics, diarrhoeas, and in disorders of the stomach, especially such as arise from a relaxed or debilitated state of that organ.

These waters are likewise very useful when employed as a tepid bath, particularly in rheumatisms, sciatica, gout, &c. by combining the internal use with the external application, they have often effected a cure where other remedies had failed to afford relief.

VICIA FABA. The systematic name of the common bean plant. See *Bean*.

VICTORIALIS LONGA. This officinal is the *Allium victorialis* of Linnaeus. The root, which when dried loses its alliaceous smell and taste, is said to be efficacious in allaying the abdominal spasms of gravid females.

VIGILANCE. *Pervigilium.* Vigilance, when attended by anxiety, pain in the head, loss of appetite, and diminution of strength, is by Sauvage and Sagar considered as a genus of disease, and is called *agrypnia*.

VERRUCA. A wart.

VERRUCARIA. (From *Ferruca*, a wart; because it was supposed to destroy warts.) The herb turnsole.

VINCA. (From *vincio*, to bind; because of its said usefulness in making bands.) The name of a genus of plants in the Linnaean system. Class, *Pentandria*. Order, *Monogynia*. The herb periwinkle, or pervinckle.

VINCA MINOR. The systematic name of the lesser periwinkle. See *Vinca pervincka*. *Clematis daphnoides major*. This plant *Vinca minor* of Linnaeus, possesses bitter and adstringent virtues, and is said to be efficacious in stopping nasal hæmorrhages when bruised and put into the nose. Boiled it forms a useful adstringent gargle in common sore throat, and it is given by some in phthisical complaints.

VINCA PERVINCKA. The lesser periwinkle.

VISCETOXICUM. (From *vinco*, to overcome, and *toxicum*, poison; so named from its supposed virtues of resisting and expelling poisons.) *Hemidynamia*. *Ascle-*

pias. Swallow wort. Tame poison. The root of this plant *Asclepias vincetoxicum* of Linnæus, smells when fresh somewhat of valerian; chewed it imparts at first a considerable sweetness which is soon succeeded by an unpleasant subacid bitterness. It is given in some countries in the cure of glandular obstructions.

Vine. See *Vitis*.

Vine, white. See *Bryonia alba*.

Vine, wild. See *Bryonia alba*.

Vinegar. See *Acetum*.

Vinegar, distilled. See *Acetum distillatum*.

Vinegar, spirits of. See *Acetic acid*.

VINUM. See *Wine*.

VINUM ALOES. Wine of aloes. Formerly known by the names of *tinctura hieræ* and *tinctura sacra*. "Take of extract of spiked aloes, eight ounces; cannella bark, two ounces; wine, six pints; proof spirits, two pints." Rub the aloes into powder with white sand, previously cleansed from any impurities: rub the cannella-bark also into powder; and after having mixed these powders together, pour on the wine and spirit. Macerate for fourteen days, occasionally shaking the mixture, and afterwards strain. A stomachic purgative, calculated for the aged and phlegmatic, who are not troubled with the piles. The dose is from half a fluid-ounce to an ounce.

VINUM ANTIMONII. In small doses this proves alterative and diaphoretic, and a large dose emetic; in which last intention it is the common emetic for children.

VINUM ANTIMONII TARTARIZATI. See *Liquor antimonii tartarizati*.

VINUM FERRI. Wine of iron, formerly called *vinum chalybeatum*. "Take of iron filings, two ounces; wine, two pints." Mix, and set the mixture by for a month, occasionally shaking it; then filter it through paper. For its virtues see *Ferrium tartarizatum*.

VINUM IPECACUANHÆ. Wine of ipecacuanha. "Take of ipecacuanha-root, bruised, two ounces; wine, two pints." Macerate for fourteen days, and strain. The dose, when used as an emetic, from two fluidrachms to half an ounce.

VINUM OPII. Wine of opium, formerly known by the names of *laudanum liquidum Sydenhami*, and *tinctura thebaica*. "Take of extract of opium, an ounce; cinnamon-bark, bruised, cloves, bruised, of each a drachm; wine, a pint." Macerate for eight days, and strain. See *Opium*.

VIOLA. (From *io*, because it was first found in Ionia.) 1. The name of a genus of plants in the Linnæan system. Class, *Cyngenesia*. Order, *Monogynia*. The violet.

2. The pharmacopœial name of the *Violaria*. Sweet violet. *Viola odorata* of Linnæus:—*acaulis, foliis cordatis, stolonibus*

repentibus. The recent flowers of this plant are received into the catalogues of the *Materia Medica*. They have an agreeable sweet smell, and a mucilaginous bitterish taste. Their virtues are purgative or laxative, and by some they are said to possess an anodyne and pectoral quality. The official preparation of this flower is a syrup, which, to young children, answers the purpose of a purgative; it is also of considerable utility in many chemical inquiries, to detect an acid or an alkali; the former changing the blue colour to a red, and the latter to a green.

VIOLA CANINA. The dog violet. The root of this plant, *Viola canina* of Linnæus, possesses the power of vomiting and purging the bowels; with which intention a scruple of the dried root must be exhibited. It appears, though neglected in this country, worthy of the attention of physicians.

VIOLA IPECACUANHÆ. The plant which was supposed to afford the ipecacuanha-root.

VIOLA LUTEA. The wall-flower was so called. See *Cheirs*.

VIOLA ODORATA. The systematic name of the sweet violet. See *Viola*.

VIOLA PALUSTRIS. See *Pinguicula*.

VIOLA TRICOLOR. Harts-case. Pansies. This well known beautiful little plant grows in corn-fields, waste, and cultivated grounds, flowering all the summer months. It varies much by cultivation; and by the vivid colouring of its flowers often becomes extremely beautiful in gardens, where it is distinguished by various names. To the taste, this plant in its recent state is extremely glutinous, or mucilaginous, accompanied with the common herbaceous flavour and roughness. By distillation with water, according to Haase, it affords a small quantity of odorous essential oil, of a somewhat acrid taste. The dried herb yields about half its weight of watery extract, the fresh plant about one-eighth. Though many of the old writers on the *materia medica* represent this plant as a powerful medicine in epilepsy, asthma, ulcers, scabies, and cutaneous complaints, yet the *viola tricolor* owes its present character as a medicine to the modern authorities of Starick, Metzger, Haase, and others, especially as a remedy for the *crusta lactea*. For this purpose, a handful of the fresh herb, or half a drachm of it dried, boiled two hours in milk, is to be strained and taken night and morning. Bread, with this decoction, is also to be formed into a poultice and applied to the part. By this treatment it has been observed, that the eruption during the first eight days increases, and that the urine, when the medicine succeeds, has an odour similar to that of cats; but on continuing the use of the plant a sufficient time, this smell goes off, the scabs disappear, and the skin recovers

its natural purity. Instances of the successful exhibition of this medicine, as cited by these authors, are very numerous; indeed this remedy, under their management, seems rarely, if ever, to have failed. It appears, however, that Mursinna, Akermann, and Flennig, were less fortunate in the employment of this plant; the last of whom declares, that in the different cutaneous disorders in which he used it, no benefit was derived. Haase, who administered this species of violet in various forms, and large doses, extended its use to many chronic disorders; and from the great number of cases in which it proved successful, we are desirous of recommending it to a farther trial in this country.

It is remarkable that Bergius speaks of this plant as a useful mucilaginous purgative, and takes no notice of its efficacy in the crusta lactea, or in any other disease.

VIOLARIA. See *Viola*.

Violet, dog. See *Viola canina*.

Violet, sweet. See *Viola*.

Viper. See *Vipera*.

Viper-grass. See *Scorzonera*.

VIPERA. (*Quod vi pariat*; because it was thought that its young eat through the mother's bowels.) The viper, or adder. This viviparous reptile, *Coluber berus* of Linnæus, possesses the power of forming a poisonous fluid in little bags near its teeth. The flesh is perfectly innocent, and often taken by the common people against the king's evil, and a variety of disorders of the skin. Experience evinces it to be an inefficacious substance.

VIPERARIA. See *Serpentaria Virginiana*.

VIPERINA. (From *viper*, a snake; so called from the serpentine appearance of its roots.) See *Serpentaria virginiana*.

VIPERINA VIRGINIANA. See *Serpentaria virginiana*.

VIRGA AUREA. *Herba florea. Conyza coma aurea. Symphytum. Petrum. Elchrysium. Consolida saracenicæ.* Golden rod. The leaves and flowers of this plant, *Solidago virga aurea* of Linnæus, are recommended as aperients and corroborants in urinary obstructions, ulcerations of the kidneys and bladder, and it is said by some to be particularly useful in stopping internal hæmorrhages.

VIRGATA SUTURA. The sagittal suture of the skull.

VIRGINALE CLAUSTRUM. The hymen.

Virginian tobacco. See *Nicotiana*.

Virgins' bower, upright. See *Flammula jovis*.

Virgins' milk. A solution of gum benzoinum.

VIRUS. A synonym of contagion. See *Contagion*.

VIS CONSERVATRIX. See *Vis medicatrix*.

VIS ELASTICA. Elasticity.

VIS INERTIE. The propensity to rest inherent in nature.

VIS INSITA. This property is defined by Haller to be that power by which a muscle, when wounded, touched, or irritated, contracts, independent of the will of the animal that is the object of the experiment, and without its feeling pain. See *Irritability*.

VIS MEDICATRIX NATURE. *Vis conservatrix.* A term employed by physicians to express that healing power in an animated body, by which, when diseased, the body is enabled to regain its healthy actions.

VIS MORTUA. That property by which a muscle after the death of the animal, or a muscle immediately after having been cut out from a living body, contracts.

VIS NERVOSA. This property is considered by Whytt to be another power of the muscles by which they act when excited by the nerves.

VIS PLASTICA. That facility of formation which spontaneously operates in animals.

VIS A TERGO. Any impulsive power.

VIS VITÆ. The natural power of the animal machine in preserving life.

VISCUM. Mistletoe. *Viscum album* of Linnæus. This singular parasitical plant most commonly grows on apple-trees, also on the pear, hawthorn, service, oak, hazel, maple, ash, lime-tree, willow, elm, hornbeam, &c. It is supposed to be propagated by birds, especially by the field-fare and thrush, which feed upon its berries, the seeds of which pass through the bowels unchanged: and along with the excrement adhere to the branches of trees where they vegetate.

The mistletoe of the oak has, from the times of the ancient druids, been always preferred to that produced on other trees; but it is now well known that the *viscus quercus* differs in no respect from others.

This plant is the $\alpha\beta$ of the Greeks, and was in former times thought to possess many medicinal virtues; however, we learn but little concerning its efficacy from the ancient writers on the materia medica, nor will it be deemed necessary to state the extraordinary powers ascribed to the mistletoe by the crafty designs of druidical knavery. Both the leaves and branches of the plant have very little smell, and a very weak taste of the nauseous kind. In distillation they impregnate water with their faint unpleasant smell, but yield no essential oil. Extracts, made from them by water, are bitterish, roughish, and subsaline. The spirituous extracts of the wood has the greatest austerities, and that of the leaves the greatest bitterness. The berries abound with an extremely tenacious and most ungrateful sweet mucilage.

The viscous quercus obtained great reputation for the cure of epilepsy; and a case of this disease, of a woman of quality, in which it proved remarkably successful, is mentioned by Boyle. Some years afterwards its use was strongly recommended in various convulsive disorders by Colbach, who has related several instances of its good effects. He administered it in substance in doses of half a drachm, or a drachm, of the wood or leaves, or an infusion of an ounce. This author was followed by others, who have not only given testimony of the efficacy of the mistletoe in different convulsive affections, but also in those complaints denominated nervous, in which it was supposed to act in the character of a tonic. But all that has been written in favour of this remedy, which is certainly well deserving of notice, has not prevented it from falling into general neglect; and the colleges of London and Edinburgh have, perhaps not without reason, expunged it from their catalogues of the *materia medica*.

VISCUM ALBUM. The systematic name of the parasitical mistletoe. See *Viscum*.

VISCUS. Any organ or part which has an appropriate use, as the viscera of the abdomen, &c.

Vision. See *Sight*.

Vision, defective. See *Dysopia*.

VISUS DEFIGURATUS. See *Metamorphopsia*.

VITE ARBOR. See *Arbor vite*.

VITE LIGNUM. See *Guaiacum*.

Vital actions. See *Vital Functions*.

VITAL FUNCTIONS. Vital actions. Those actions of the body upon which life immediately depends, as the circulation of the blood, respiration, heat of the body, &c. See *Function*.

Vital principle. See *Life*.

VITALBA. Traveller's joy. This plant is common in our hedges, and is the *Clematis vitalba* of Linnæus;—*foliis pinnatis, folioli cordatis scandentibus*: its leaves when fresh produce a warmth on the tongue, and if the chewing is continued, blisters arise. The same effect follows their being rubbed on the skin. The plant has been administered internally to cure lues venerea, scrofula, and rheumatism. In France, the young sprouts are eaten, when boiled, as hoptops are in this country.

VITILIGO. (From *vitis*, to infect.) See *Alphus*.

VITIS. 1. The name of a genus of plants in the Linnæan system. Class, *Pentandria*. Order, *Monogynia*.

2. The pharmacopœial name of the *Vitis vinifera* of Linnæus:—*foliis lobatis sinuatis nudis*. Vine-leaves and the tendrils have an adstringent taste, and were formerly used in diarrhœas, hæmorrhages, and other disorders requiring refrigerant and styptic medicines. The juice or sap

of the vine, called lachryms, has been recommended in calculous disorders: and it is said to be an excellent application to weak eyes and specks of the cornea. The unripe fruit has a harsh, rough, sour taste; its expressed juice, called verjuice, was formerly much esteemed, but is now superseded by the juice of lemons; for external use, however, particularly in bruises and pains, verjuice is still employed, and considered to be a very useful application. See also *Uva passe, wine*, and *acetum*.

VITIS ALBA. See *Bryonia alba*.

VITIS ALTUS. St. Vitus's dance.

VITIS IDMEA. The red whortle-berry. The leaves of this plant, *Vaccinium vitis idæa* of Linnæus, are so adstringent as to be used in some places for tanning. They are said to mitigate the pain attendant on calculous diseases when given internally in the form of decoction. The ripe berries abound with a grateful acid juice; and are esteemed in Sweden as aperient, antiseptic, and refrigerant, and often given in putrid diseases.

VITIS SYLVESTRIS. White briony.

VITIS VINIFERA. The systematic name of the grape-tree. See *Vitis* and *wine*.

VITIS AGNUS CASTUS. The systematic name of the chaste tree. See *Agnus castus*.

VITRARIA. The pellytory of the wall.

VITREOUS HUMOUR. *Humour vitreus*. The pellucid body which fills the whole bulb of the eye behind the crystalline lens. The vitreous substance is composed of small cells which communicate with each other, and are distended with a transparent fluid.

Vitriol, acid of. See *Sulphuric acid*.

Vitriol, blue. See *Sulphus cupri*.

Vitriol, green. See *Ferri sulphas*.

Vitriol, Roman. See *Sulphas cupri*.

Vitriol, sweet spirit of. See *Spiritus ætheris vitriolici*.

Vitriol, white. See *Zinci sulphas*.

Vitriolated kali. See *Potasse sulphas*.

VITRIOLUM. (From *vitrum*, glass; so called from its likeness to glass. Hollandus says this word is fictitious, and composed from the initials of the following sentence: *Vade in terram rimundo, invenies, optimum lapidem veram medicinam.*) *Calcadinum*. *Calcatar*. *Calcatar*. *Calcanthos*. *Calcanthum*. *Calcitea*. *Vitriol*, or sulphate of iron.

VITRIOLUM ALBUM. See *Zinci sulphas*.

VITRIOLUM CÆRULEUM. See *Curpi sulphas*.

VITRIOLUM ROMANUM. See *Curpi sulphas*.

VITRIOLUM VIRIDE. See *Ferri sulphas*.

VITRUM. Glass.

VITRUM ANTIMONII. Glass of antimony: Antimony first calcined, then fused in a crucible.

VITRUM ANTIMONII SCRATUM. A di-

apboretic compound exhibited in the cure of dysenteries arising from checked perspiration.

VITRUM HYPOCLEPTICUM. A funnel to separate oil from water.

VOICE. *Vox.* The principal organ of the voice is the larynx; for, when it is injured, the air passes through the windpipe without yielding any sound.

VOISELLA. A probang, or instrument to remove bodies sticking in the throat.

VOLVULUS. (From *volvo*, to roll up) The iliac passion, or inflammation in the bowels, called twisting of the guts.

VOLVULUS TERRESTRIS. Small bindweed. The convolvulus minor.

VOMER. Named from its great resemblance to a plough-share. It is a slender thin bone separating the nostrils from each other, consisting of two plates much compressed together, very dense and strong, yet so thin as to be transparent; these two plates seem at every edge to separate from each other, and thus a groove is formed at every side.—1 This groove, on the upper edge, or, as it may be called, its base, is wide, and receives into it the projecting points of the ethmoid and sphenoid bones, and thus it stands very firmly and securely on the skull, and capable of resisting blows of considerable violence.—

2. The groove, upon the lower part, is narrower, and receives the rising line in the middle of the palate plate, where the bones join to form the palate suture. At the fore part it is united by a ragged surface, and by something like a groove, to the middle cartilage of the nose, and as the vomer receives the other bones into its grooves, it is, as it were, locked in on all sides, receiving support and strength from each, but more particularly from the thick and strong membrane which covers the whole, and which is so continuous as to resemble a periosteum, or rather

a continued ligament from its strength; thus the slender vomer possesses sufficient strength to advert from it all those evils which must inevitably have occurred, had it been less wisely or less strongly constructed.

VOMICA. (From *vomo*, to spit up; because it discharges a sanies.) An abscess of the lungs.

VOMICA LIQUORIS STRENI. Quick-silver.

VOMITING. *Vomitio.* A forcible ejection of food, or any other substance, from the stomach, through the oesophagus and mouth. It is either *idiopathic*, when arising from a cause in the stomach itself; or *symptomatic*, when originating from diseases seated in any other part of the body. Its immediate cause is an antiperistaltic, spasmodic, convulsive constriction of the stomach and pressure of the diaphragm and abdominal muscles. It is an effort of nature to expel whatever molests or is troublesome in the stomach.

Vomiting of blood. See *Hematemesis*.

VOMITUS CRUENTUS. See *Hematemesis*.

Voracious appetite. See *Bulimia*.

VOX ABSCESSA. Hoarseness, a loss of voice.

VULGAGO. The asarabacca was so called. See *Asarum*.

VULNERARIA. (From *vulnus*, a wound.) Medicines which heal wounds. A herb named from its uses in healing wounds.

VULNERARIA AQUA. Arguebusade.

VULNUS SCLOPETICUM. A gun-shot wound.

VULVA. (*Quasi vulva*, the aperture to the womb; or *quasi volta*, because the fetus is wrapped up in it.) The pudendum muliebre, or parts of generation proper to women; also a foramen in the brain.

VULVARIA. (From *vulva*, the womb; so named from its smell or use in disorders of the womb.) Stinking orach. See *Atriplex olide*.

W

WAKE rubin. See *Arum*.

Wall-flower. See *Cheiri*.

Wall-pellitory. See *Parietaria*.

Wall-pepper. See *Illecebra*.

Walnut. See *Juglans*.

WALTON WATER. A mineral spring near Tewkesbury in Gloucestershire, containing a small portion of iron dissolved in fixed air; of absorbent earth combined with hepatic air; of vitriolated magnesia, and muriated mineral alkali; but the pro-

portions of these constituent parts have not been accurately ascertained. Walton water is chiefly efficacious in obstructions and other affections of the glands.

WATER. *Aqua.* According to the opinion of modern chemists is a compound fluid, made up of two substances, neither of which can be exhibited separately, except in the gaseous form; and when æri-form, they are known, the one as hydrogen gas, or inflammable air; the other as

oxygen gas, or vital air. These gases, in the proportion of about three of hydrogen to eleven of oxygen, when united chemically, and reduced from the form of an air to that of a liquid, constitute the fluid, water.

It is transparent, without colour, smell, or taste; in every degree compressible; when pure, not liable to spontaneous change; liquid in the common temperature of our atmosphere, assuming the solid form at 32° Fahrenheit, and the gaseous at 212° , but returning unaltered to its liquid state on resuming any degree of heat between these points; capable of dissolving a greater number of natural bodies than any other fluid whatever, and especially those known by the name of the saline; performing the most important functions in the vegetable and animal kingdoms, and entering largely into their composition as a constituent part. The simple waters are the following:

1. *Distilled water.* This is the lightest of all others, containing neither solid nor gaseous substances in solution, is perfectly void of taste and smell, colourless and beautifully transparent, has a soft feel, and wets the fingers more readily than any other. It mixes uniformly with soap into a smooth opaline mixture, but may be added to a solution of soap in spirit of wine without injuring its transparency. The clearness of distilled water is not impaired by the most delicate chemical re-agents, such as lime-water, a solution of barytes in any acid, nitrated silver, or acid of sugar. When evaporated in a silver vessel it leaves no residuum; if preserved from access of foreign matter floating in the air, it may be kept for ages unaltered in vessels upon which it has no action, as it does not possess within itself the power of decomposition. As it freezes exactly at 32° of Fahrenheit, and boils at 212° under the atmospherical pressure of 29.8° inches, these points are made use of as the standard ones for their metrical division; and its specific weight being always the same under light pressure and temperature, it is employed for the comparative standard of specific gravity.

Pure distilled water can only be procured from water which contains no volatile matters that will rise in distillation, and continue still in union with the vapour when condensed. Many substances are volatile during distillation, but most of the gases, such as common air, carbonic acid, and the like, are incapable of uniting with water at a high temperature: other bodies, however, such as vegetable essential oil, and, in general, much of that which gives the peculiar odour to vegetable and animal matter, will remain in water after distillation. So the steam of many animal and vegetable decoctions has a certain flavour

which distinguishes it from pure water; and the aqueous exhalation from living bodies, which is a kind of distillation, has a similar impregnation.

To obtain distilled water perfectly pure, much stress was laid by former chemists on repeating the process a great number of times; but it was found by Lavoisier, that rain water once distilled, rejecting the first and last products, was as pure a water as could be procured by any subsequent distillations.

Distilled water appears to possess a higher power than any other as a solvent of all animal and vegetable matter, and these it holds in solution as little as possible altered from the state in which they existed in the body that yielded them. Hence the great practical utility of that kind of chemical analysis which presents the proximate constituent parts of these bodies, and which is effected particularly by the assistance of pure water. On the other hand, a saline, earthy, or otherwise impure water, will alter the texture of some of the parts, impair their solubility, produce material changes on the colouring matter, and become a less accurate analyser on account of the admixture of foreign contents.

Distilled water is seldom employed to any extent in the preparation of food, or in manufactures, on account of the trouble of procuring it in large quantities; but for preparing a great number of medicines, and in almost every one of the nicer chemical processes that are carried on in the liquid way, this water is an essential requisite. The only cases in which it has been used largely as an article of drink, have been in those important trials made of the practicability of procuring it by condensing the steam of sea water by means of a simple apparatus adapted to a ship's boiler; and these have fully shewn the ease with which a large quantity of fresh water of the purest kind, may be had at sea, at a moderate expense, whereby one of the most distressing of all wants may be relieved. There are one or two circumstances which seem to shew that water, when not already loaded with foreign matter, may become a solvent for concretions in urinary passages. At least, we know that very material advantage has been derived in these cases from very pure natural springs, and hence a course of distilled water has been recommended as a fair subject of experiment.

2. *Rain water*, is the next in purity to distilled water, is that which has undergone a natural distillation from the earth, and is condensed in the form of rain. This is a water so nearly approaching to absolute purity as probably to be equal to distilled water for every purpose except in the nicer chemical experiments. The fo-

reign contents of rain water appear to vary according to the state of the air through which it falls. The heterogeneous atmosphere of a smoky town will give some impregnation to rain as it passes through, and this, though it may not be at once perceptible on chemical examination, will yet render it liable to spontaneous change; and hence, rain water, if long kept, especially in hot climates, acquires a strong smell, becomes full of animalcula, and in some degree putrid. According to Margraff, the constant foreign contents of rain water appear to be some traces of the muriatic and nitric acids; but as this water is always very soft, it is admirably adapted for dissolving soap, or for the solution of alimentary or colouring matter, and it is accordingly used largely for these purposes. The specific gravity of rain water is so nearly the same as that of distilled water, that it requires the most delicate instruments to ascertain the difference. Rain, that falls in towns, acquires a small quantity of sulphat of lime and calcareous matter from the mortar and plaister of the houses.

3. *Ice and snow water.* This equals rain water in purity, and, when fresh melted, contains no air, which is expelled during freezing. In cold climates and in high latitudes, thawed snow forms the constant drink of the inhabitants during winter; and the vast masses of ice which float on the polar seas afford an abundant supply to the mariner. It is well known, that in a weak brine, exposed to a moderate freezing cold, it is only the watery part that congeals, leaving the unfrozen liquor proportionably stronger of the salt. The same happens with a dilute solution of vegetable acids, with fermented liquors, and the like; and advantage is taken of this property to reduce the saline part to a more concentrated form. Snow water has long lain under the imputation of occasioning those strumous swellings in the neck which deform the inhabitants of many of the Alpine valleys; but this opinion is not supported by any well-authenticated indisputable facts, and is rendered still more improbable, if not entirely overturned, by the frequency of the disease in Sumatra, where ice and snow are never seen, and its being quite unknown in Chili and in Thibet, though the rivers of these countries are chiefly supplied by the melting of the snow, with which the mountains are covered.

4. *Spring water.* Under this comprehensive class are included all waters that spring from some depth beneath the soil, and are used at the fountain head, or at least before they have run any considerable distance exposed to the air. It is obvious that spring water will be as various in its contents as the substances that compose the

soil through which it flows. When the ingredients are not such as to give any peculiar medical or sensible properties, and the water is used for common purposes, it is distinguished as a hard or soft spring, sweet or brackish, clear or turbid, and the like. Ordinary springs insensibly pass into mineral springs, as their foreign contents become more notable and uncommon; though sometimes waters have acquired great medical reputation from mere purity.

By far the greater number of springs are cold; but as they take their origin at some depth from the surface, and below the influence of the external atmosphere, their temperature is, in general, pretty uniform during every vicissitude of season, and always several degrees higher than the freezing point. Others, again, arise constantly hot, or with a temperature always exceeding the summer heat; and the warmth possessed by the water is entirely independent of that of the atmosphere, and varies little winter or summer.

One of the principal inconveniences in almost every spring water, is its hardness, owing to the presence of earthy salts, which, in by far the greater number of cases, are only the insipid substances, chalk and selenite, which do not impair the taste of the water; whilst the air which it contains, and its grateful coolness, render it a most agreeable, and generally a perfectly innocent, drink; though sometimes, in weak stomachs, it is apt to occasion an uneasy sense of weight in that organ followed by a degree of dyspepsia. The quantity of earthy salts varies considerably; but, in general, it appears that the proportion of five grains of these in the pint will constitute a hard water, unfit for washing with soap, and for many other purposes of household use or manufactures. The water of deep wells is always, *ceteris paribus*, much harder than that of springs which overflow their channel; for much agitation and exposure to air produce a gradual deposition of the calcareous earth; and hence spring water often incrusts to a considerable thickness the inside of any kind of tube through which it flows, as it arises from the earth. The specific gravity of these waters is also, in general, greater than that of any other kind of water, that of the sea excepted. Springs that overflow their channel, and form to themselves a limited bed, pass insensibly into the state of stream, or river water, and become thereby altered in some of their chemical properties.

5. *River water.* This is in general much softer and more free from earthy salts than the last, but contains less air of any kind; for, by the agitation of a long current, and in most cases a great increase of temperature, it loses common air and carbon

acids, and, with this last, much of the lime which it held in solution. The specific gravity thereby becomes less, the taste not so harsh, but less fresh and agreeable, and out of a hard spring is often made a stream of sufficient purity for most of the purposes where a soft water is required. Some streams, however, that arise from a clean siliceous rock, and flow in a sandy or stony bed, are from the outset remarkably pure. Such are the mountain lakes and rivulets in the rocky districts of Wales, the source of the beautiful waters of the Dee, and numberless other rivers that flow through the hollow of every valley. Switzerland has long been celebrated for the purity and excellence of its waters, which pour in copious streams from the mountains; and give rise to some of the finest rivers in Europe. An excellent observer and naturalist, the illustrious Haller, thus speaks of the Swiss waters, "vulgaribus aquis Helvetia super omnes fere Europæ regiones excoëllit. Nusquam liquidas illas aquas et crystalli similitas se rarihi obtulisse memini postquam ex Helvetia excessi. Ex scopulis enim nostræ per puros silices percolatæ nulla terra vitiantur." Some of them never freeze in the severest winter, the cause of which is probably, as Haller conjectures, that they spring at once out of a subterraneous reservoir so deep as to be out of the reach of frost, and during their short course when exposed to day they have not time to be cooled down from 53°, their original temperature, to below the freezing point.

Some river waters, however, that do not take their rise from a rocky soil, and are indeed at first considerably charged with foreign matter, during a long course, even over a rich cultivated plain, become remarkably pure as to saline contents, but often souled with mud, and vegetable or animal exuviz, which are rather suspended than held in true solution. Such is that of the Thames, which taken up at London at low water, is a very soft and good water, and, after rest and filtration, it holds but a very small portion of any thing that could prove noxious or impede any manufacture. It is also excellently fitted for sea store; but it here undergoes a remarkable spontaneous change. No water carried to sea becomes putrid sooner than that of the Thames. When a cask is opened after being kept a month or two, a quantity of inflammable air escapes, and the water is so black and offensive as scarcely to be borne. Upon racking it off, however, into large earthen vessels (oil jars are commonly used for the purpose), and exposing it to the air, it gradually deposits a quantity of black slimy mud, becomes clear as crystal, and remarkably sweet and palatable. The Seine has as high a reputation in France, and appears from accurate experi-

ments to be a river of great purity. It might be expected that a river which has passed by a large town, and received all its impurities, and been used by numerous dyers, tanners, hatters, and the like, that crowd to its banks for the convenience of plenty of water, should thereby acquire such a foulness as to be very perceptible to chemical examination for a considerable distance below the town; but it appears, from the most accurate examination, that where the stream is at all considerable, these kinds of impurity have but little influence in permanently altering the quality of the water, especially as they are for the most part only suspended and not truly dissolved; and, therefore, mere rest, and especially filtration, will restore the water to its original purity. Probably, therefore, the most accurate chemist would find it difficult to distinguish water taken up at London, from that procured at Hampton Court, after each has been purified by simple filtration.

6. *Stagnated waters.* The waters that present the greatest impurities to the senses, are those of stagnant pools, and low marshy countries. They are filled with the remains of animal and vegetable matter undergoing decomposition, and, during that process, becoming in part soluble in water, thereby affording a rich nutriment to the succession of living plants and insects which is supplying the place of those that perish. From the want of sufficient agitation in these waters, vegetation goes on undisturbed, and the surface becomes covered with conveer and other aquatic plants; and as these standing waters are in general shallow, they receive the full influence of the sun, which further promotes all the changes that are going on within them. The taste is generally vapid, and destitute of that freshness and agreeable coolness which distinguish spring water. However, it should be remarked, that stagnant waters are generally soft, and many of the impurities are only suspended, and therefore separable by filtration; and perhaps the unpalatableness of this drink has caused it to be in worse credit than it deserves, on the score of salubrity. The decidedly noxious effects produced by the air of marshes and stagnant pools, have been often supposed to extend to the internal use of these waters; and often, especially in hot climates, a residence near these places has been as much condemned on one account as on the other, and, in like manner, an improvement in health has been as much attributed to a change of water as of air.

Water-brash. See *Pyrosis*.

Water-cress. See *Nasturtium aquaticum*.

Water-dock. See *Hydrolythum*.

Water-lily, yellow. See *Iris palustris*.

Water-germander. See *Scordium*.

Water-hemp. See *Eupatorium*.

Water-lily, white. See *Nymphaea alba*.

Water-lily, yellow. See *Nymphaea lutea*.

Water-pump. See *Sium*.

Water-pepper. See *Hydrophiper*.

WATER RIZANTIA. A reed-like plant, *sizania aquatica* of Linnaeus, which grows in the swampy parts of Jamaica and Virginia. The Indians are exceedingly fond of its grain, and account it more delicious than rice.

Waters, mineral. See *Mineral Waters*.

Wux. See *Cera*.

WHEAT. The seeds of the *Triticum hybernium*, and *aestivum* of Linnaeus, are so termed. It is to these plants therefore we are indebted for our bread, and the various kinds of pastry. Wheat is first ground between mill-stones, and then sifted to obtain its farina or flour. The flour of wheat may be separated into its three constituent parts, in the following manner. The flour is to be kneaded into a paste with water in an earthen vessel, and the water continue pouring upon it from a cock which, as it falls upon the paste, takes up from it a very fine white powder, by means of which it acquires the colour and consistency of milk. This process is to be continued till the water run off clear, when the flour will be separated into three distinct parts: 1. A grey elastic matter that sticks to the hand, and on account of its properties has gained the name of the glutinous, or vegeto-animal part. 2. A white powder which falls to the bottom of the water, and is the *seculum* or starch. 3. A matter which remains dissolved in the water, and seems to be a sort of mucilaginous extracts.

Flour, from whatever species of corn obtained, is likewise disposed to vinous fermentation, on account of its saccharine contents, the aptitude for fermentation of these mealy seeds increases if they be first converted into malt; inasmuch as by this process, the gluten which forms the germ is separated, and the starchy part appears to be converted into saccharine matter. The making of malt, for which purpose barley and wheat are generally chosen, is as follows: The grains are put in the malting tub, and immersed in cold water, in a temperate and warm season, changing this fluid several times, especially in hot weather, and they are thus kept soaking till they be sufficiently soft to the touch. Upon this they are piled up in heaps on a roomy, clean, airy floor, where, by the heat spontaneously taking place, the vegetation begins, and the grains germinate. To cause the germination to go on uniformly, the heaps are frequently turned. In this state the vegetation is suffered to continue till the germs have about two-

thirds or three-fourths of the length of the corn. It is carried too far when the leafy germs have begun to sprout.

For this reason limits are set to the germination by drying the malt, which is effected by transferring it to the kiln, or by spreading it about in spacious airy lofts. Dried in the last way it is called air dried malt, in the first, kiln-malt. In drying this latter, care must be taken that it does not receive a burnt smell, or be in part converted into coal.

From this malt, beer is made by extraction with water and fermentation.

With this view a quantity of malt freed from its germs, and sufficient for one intended brewing, is coarsely bruised by grinding, and in the mash-tub first well mixed with some cold, then scalded with hot water, drawn upon it from the boiler. It is afterwards strongly and uniformly stirred. When the whole mass has stood quietly for a certain time, the extract, (mash,) or sweet wort, is brought into the boiler, and the malt remaining in the tub is once more extracted by infusion with hot water.

This second extract treated in like manner, is added to the first, and both are boiled together.

This clear decoction is now drawn off, and called boiled wort. To make the beer more fit for digestion, and at the same time to deprive it of its top great and unpleasant sweetness, the wort is mixed with a decoction of hops, or else these are boiled with it. After which it ought to be quickly cooled, to prevent its transition into acetous fermentation, which would ensue if it were kept too long in a high temperature.

On this account the wort is transferred into the cooler, where it is exposed with a large surface to cold air, and from this to the fermenting tub, that by addition of a sufficient portion of recent yeast it may begin to ferment. When this fermentation has proceeded to a due degree, and the yeast ceases to rise, the beer is conveyed into casks (casked), placed in cool cellars, where its finishes its fermentation, and where it is well kept and preserved, under the name of barrelled beer, with the precaution of filling up occasionally the vacancy caused in the vessels by evaporation; or the beer is bottled before it has done fermenting, and the bottles are stopped a little before the fermentation is completely over. By so doing the bottled beer is rendered sparkling. In this state it frequently bursts the bottles, by the disengagement of the carbonic acid gas which it contains, and it strongly froths, like Champaign, when brought into contact of air on being poured into another vessel.

Beer well prepared should be limpid and clear, possess a due quantity of spirit,

and excite no disagreeable sweet taste, and contain no disengaged acid. By these properties it is a species of vinous beverage, and is distinguished from wine, in the strict sense, and other liquors of that kind, by the much greater quantity of mucilaginous matter which it has received by extraction from the malted grains, but which also makes it more nourishing. Brown beer derives its colour from malt strongly roasted in the kiln, and its bitterish taste from the hops. Pale beer is brewed from malt dried in the air, or but slightly roasted, with but little or no hops at all.

Wheat, buck. See *Buck wheat*.

Wheat, eastern buck. See *Buck wheat, eastern*.

Wheat, Indian. See *Zea mays*.

WHEAT, TURKEY. The Turkey wheat is a native of America, where it is much cultivated, as it is also in some parts of Europe, especially in Italy and Germany. There are many varieties, which differ in the colour of the grain, and are frequently raised in our gardens by way of curiosity, whereby the plant is well known. It is the chief bread corn in some of the southern parts of America, but since the introduction of rice into Carolina, it is but little used in the northern colonies. It makes a main part too of the food of the poor people in Italy and Germany. This is the sort of wheat mentioned in the book of Ruth, where it is said that Boaz treated Ruth with parched ears of corn dipped in vinegar. This method of eating the roasted ears of Turkey wheat is still practised in the East; they gather in the ears when about half ripe, and having scorched them to their minds, eat them with as much satisfaction as we do the best flour bread.

In several parts of South America they parch the ripe corn, never making it into bread, but grinding it between two stones, mix it with water in a calabash, and so eat it. The Indians make a sort of drink from this grain, which they call *bici*. This liquor is very windy and intoxicating, and has nearly the taste of sour small beer: but they do not use it in common, being too lazy to make it often, and therefore it is chiefly kept for the celebration of feasts and weddings, at which times they mostly get intolerably drunk with it. The manner of making this precious beverage, is to steep a parcel of corn in a vessel of water, till it grows sour, then the old women, being provided with calabashes for the purpose, chew some grains of the corn in their mouths, and spitting it into the calabashes, empty them, spit it and all, into the sour liquor, having previously drawn off the latter into another vessel.

The chewed grain soon raises a fermentation, and when this ceases, the liquor is let off from the dregs, and set by till

wanted. In some of the islands in the South Sea, where each individual is his own lawgiver, it is no uncommon thing for a near relation to excuse a murderer, for a good drunken bout of ciri.

White swelling. See *Arthropoosis* and *Hydarthrus*.

WHITES. *Leucorrhœa. Fluor albus.* This disease is marked by the discharge of a thin white or yellow matter from the uterus and vagina, attended likewise with some degree of fever, smarting in making water, pains in the back and loins, anorexia and atrophy. In some cases, the discharge is of so acrid a nature, as to produce effects on those who are connected with the woman, somewhat similar to venereal matter, giving rise to excoriations about the glans penis and preputium, and occasioning a weeping from the urethra.

To distinguish leucorrhœa from gonorrhœa, it will be very necessary to attend to the symptoms. In the latter the running is constant, but in a small quantity; there is much ardor urine, itching of the pudenda, swelling of the labia, increased inclination to venery, and very frequently an enlargement of the glands in the groin; whereas in the former the discharge is irregular, comes away often in large lumps, and in considerable quantities, and is neither preceded by nor accompanied with any inflammatory affection of the pudenda.

Immoderate coition, injury done to the parts by difficult and tedious labours, frequent miscarriages, immoderate flowings of the menses, profuse evacuations, poor diet, and abuse of tea, and other causes giving rise to general debility, or to a laxity of the parts more immediately concerned, are those which usually produce the whites, vulgarly so called, from the discharge being commonly of that colour.

Fluor albus, in some cases, indicates that there is a disposition to disease in the uterus, or parts connected with it, especially where the quantity of the discharge is very copious, and its quality highly acrimonious. By some the disease has been considered as never arising from debility of the system, but as being always a primary affection of the uterus. Delicate women, with lax fibres, who remove from a cold climate to a warm one, are, however, very apt to be attacked with it, without the parts having previously sustained any kind of injury.

The disease shews itself by an irregular discharge from the uterus and vagina, of a fluid which, in different women, varies much in colour, being either of a white, green, yellow, or brown hue. In the beginning it is, however, most usually white and pellucid, and in the progress of the complaint acquires the various discolourations, and different degrees of acrimony,

from whence proceeds a slight degree of smarting in making water. Besides the discharge, the patient is frequently afflicted with severe and constant pains in the back and loins, loss of strength, failure of appetite, dejection of spirits, paleness of the countenance, chilliness, and languor. Where the disease has been of long continuance, and very severe, a slow fever, attended with difficult respiration, palpitations, faintings, and anasarcaous swellings of the lower extremities, often ensues.

A perfect removal of the disorder will at all times be a difficult matter to procure; but it will be much more so in cases of long standing, and where the discharge is accompanied with a high degree of acrimony. In these cases, many disorders, such as prolapsus uteri, ulcerations of the organ, atrophy and dropsy, are apt to take place, which in the end prove fatal.

Where the disease terminates in death, the internal surface of the uterus appears, on dissection, to be pale, flabby, and relaxed; and where organic affections have arisen, much the same appearances are to be met with as have been noticed under the head of menorrhagia. See *Leucorrhœa*.

Whortle-berry, bears. See *Uva urol.*

Whortle-berry, red. See *Vitis idœa*.

Widow-wail. See *Mezerium*.

Wild carrot. See *Daucus sylvestris*.

Wild cucumber. See *Cucumis agrestis*.

Wild nuxem. See *Nipus*.

Willow, crak. See *Salix*.

Willow, sweet. See *Myrtus brabantica*.

Willow, white. See *Salix*.

Willow-herb. See *Lynimachia purpurea*.

Willow-herb, rosebay. See *Rosebay willow-herb*.

Willow-leaved oak. See *Quercus phellos*.

WINE. *Vinum.* The fermented juice of the ripe fruit of the *Vitis vinifera* of Linnaeus:—*foliis lobatis sinuatis nudis*. The wines principally used in medicine are, the *vinum album hispanicum*, or mountain wine, *vinum canarium*, canary or sack wine, the *vinum rhenanum*, or Rhenish wine, and the *vinum rubrum*, or Port wine. On a chemical investigation, all wines consist chiefly of water, alcohol, a peculiar acid, the carbonic acid, tartar, and an adstringent gummi-resinous matter in which the colour of the red wine resides, and which is expressed from the husks of the grape. They differ from each other in the proportion of these ingredients, and particularly in that of alcohol, which they contain. The qualities of wines depend not only upon the difference of the grapes, as containing more or less of saccharine juice and the acid matter which accompanies it, but also upon circumstances attending the process of fermentation. New wines are liable to a strong degree of acescency when taken into the stomach, and

thereby occasion much flatulency and eructations of acid matter; heartburn and violent pains in the stomach from spasms are also often produced; and the acid matter, by passing into the intestines and mixing with the bile, is apt to occasion colics or excite diarrhoeas. Sweet wines are likewise more disposed to become acescent in the stomach than others: but as the quantity of alcohol which they contain is more considerable than appears sensibly to the taste, their acescency is thereby in a great measure counteracted. Red port, and most of the red wines, have an adstringent quality, by which they strengthen the stomach, and prove useful in restraining immoderate evacuations; on the contrary, those which are of an acid nature, as Rhenish, pass freely by the kidneys, and gently loosen the belly. But this, and perhaps all the thin or weak wines, though of an agreeable flavour, yet as containing little alcohol, are readily disposed to become acid in the stomach, and thereby to aggravate all arthritic and calculous complaints, as well as to produce the effects of new wine. The general effects of wine are, to stimulate the stomach, exhilarate the spirits, warm the habit, quicken the circulation, promote perspiration, and, in large quantities, to prove intoxicating, and powerfully sedative. In many disorders, wine is universally admitted to be of important service, and especially in fevers of the typhus kind, or of a putrid tendency; in which it is found to raise the pulse, support the strength, promote a diaphoresis, and to resist putrefaction; and in many cases it proves of more immediate advantage than the Peruvian bark. Delirium, which is the consequence of excessive irritability, and defective state of nervous energy, is often entirely removed by the free use of wine. It is also a well-founded observation, that those who indulge in the use of wine are less subject to fevers of the malignant and intermittent kind. In the putrid sore throat, in the small-pox, when attended with great debility and symptoms of putridity, in gangrenes, and in the plague, wine is to be considered as a principal remedy; and in almost all cases of languor, and of great prostration of strength, wine is experienced to be a more grateful and efficacious cordial than can be furnished from the whole class of aromatics.

Method of discovering whether wine has been adulterated with any metals prejudicial to the health.—The property which the sulphuret of potash and hepatic air, or sulphurated hydrogen, possess of precipitating lead in a black form, has been long ago made public; and this property has been employed to determine the quality of wines by means of the liquor probatorius Wirtembergensis, or Wirtemberg proving-liquor. But in trying

wines supposed to have been adulterated, this proof does more harm than service, because it precipitates iron of the same colour as the pernicious lead. Many wine-merchants of the greatest respectability rendered by these means suspected, have been ruined. There was wanting then a re-agent, which should discover in wine those metals only which are prejudicial to the health of man.

The following liquor, precipitates lead and copper in a black form, and arsenic of an orange colour, &c. but does not precipitate iron, which is not noxious, and rather salutary to the constitution, and frequently gets into wines by accident.

Method of preparing the proving liquor.—Mix equal parts of oyster-shells and crude sulphur in a fine powder, and put the mixture into a crucible; heat it in a wind furnace, and increase the fire suddenly, so as to bring the crucible to a white heat, for the space of fifteen minutes; pulverize the mass when it is cold, and preserve it in a bottle closely stopped. To prepare the liquor, put 120 grains of this powder and 120 grains of cream of tartar (acidulous tartrite of potash) into a strong bottle; fill the bottle with common water, which boil for an hour, and then let it cool; close the bottle immediately, and shake it for some time; after it has remained at rest to settle, decant the pure liquor, and pour it into small phials, capable of holding about an ounce each, first putting into each of them twenty drops of muriatic acid. They must be stopped very closely with a piece of wax, in which there is a small mixture of turpentine. One part of this liquor mixed with three parts of suspected wine, will discover, by a very sensible black precipitate, the least traces of lead, copper, &c. but will produce no effect upon iron, if it contains any of that metal. When the precipitate has fallen down, it may still be discovered whether the wine contains iron, by saturating the decanted liquor with a little salt of tartar, by which the liquor will immediately become black. Pure wines remain clear and bright after this liquor has been added to them.

Winter-bark. See *Winteranus cortex*.

Winter-cherry. See *Alkekengi*.

WINTERA AROMATICA. The systematic name of the winter bark-tree. See *Winteranus cortex*.

WINTERANUS CORTEX. *Winteranus cortex magellanicus.* The bark of the *Wintera aromatica pedunculis aggregatis terminalibus, pistillis quatuor*; it is very much allied in its properties to the *canella alba*. See *Canella alba*.

WINTERANUS SPURIUS. See *Canella alba*.

WHISPERING. A lowness of speech caused by uttering the words so feebly as

not to produce any vibration of the larynx.

Wolfs'bane. See *Aconitum*.

Wolfs'bane, blue. See *Aconitum*.

Womb. See *Uterus*.

Womb, inflammation of. See *Hysteritis*.

Wood-louse. See *Millipedes*.

Wood-sorrel. See *Luzula*.

Woody nightshade. See *Dnicamara*.

Worm-bark. See *Geoffrea*.

Wormgrass, perennial. See *Spigelia*.

Worm guinea. See *Dracunculus*.

Worm-ring. See *Herpes*.

Worm-seed. See *Santonicum*.

WORMS. *Vermes.* There are several kinds of animals which infest the human body. Their usual division is into those which inhabit only the intestinal canal, as the ascarides, &c.; and those which are found in other parts, as hydatids, &c. Such is the nature and office of the human stomach and intestines, that insects and worms, or their ovula, may not unfrequently be conveyed into that canal with those things, that are continually taken as food; but such insects, or worms, do not live long, and seldom, if ever, generate in a situation so different from their natural one. Besides these, there are worms that are never found in any other situation than the human stomach, or intestines, and which there generate and produce their species. Thus it appears that the human stomach and intestines are the seat for animalculæ, which are translated from their natural situation, and also for worms proper to them, which live in no other situation.

First Class.

This contains those which are generated and nourished in the human intestinal canal, and which there propagate their species.

Second Class.

Comprehends those insects or worms that accidentally enter the human primæ viz ab extra, and which never propagate their species in that canal, but are soon eliminated from the body; such are several species of *Scarabæi*, the *Lumbricus terrestris*, the *Fasciola*, the *Gordius intestinalis*, and others. The second class belongs to the province of natural history. The consideration of the first class belongs to the physician, which, from the variety it affords, may be divided into different orders, genera, and species.

Order I. Round worms.

Genus I. Intestinal ascarides.

Character. Body round, head obtuse, and furnished with three vesicles.

Species 1. Ascaris lumbricoides. The long round worm, or lumbricoid ascaris.

Character. When full grown, a foot in length. Mouth triangular.

II. Ascaris vermicularis. The thread or maw-worm.

Character. When full grown, half an inch in length, tail terminates in a fine point.

Genus II. Intestinal trichurides.

Character. Body round, tail three times the length of the body, head without vesicles.

Species. *Trichuris vulgaris*. The trichuris, or long thread-worm.

Character. The head furnished with a proboscis.

Order II. The flat worms.

Genus I. Intestinal tape-worm.

Character. Body flat and jointed.

Species I. *Tenia osculis marginalibus*. The long tape-worm.

Character. The oscula are situated upon the margin of the joints.

II. *Tenia osculis superficialibus*. The broad tape-worm.

Character. The oscula are placed upon the flattened surface.

These worms were all known to the ancients, the trichuris only excepted, and are mentioned in the works of Hippocrates, Galen, Celsus, Paulus Aegineta, and Pliny.

Worms may readily be distinguished by the following symptoms, viz. variable appetite, fetid breath, acid eructations and pains in the stomach, grinding of the teeth during sleep, picking of the nose, paleness of the countenance; sometimes dizziness, hardness and fulness of the belly; slimy stools, with occasional griping pains, more particularly about the navel, heat and itching about the anus; short dry cough; emaciation of the body; slow fever, with evening exacerbations and irregular pulse, and sometimes convulsive fits.

Wormwood. See *Absinthium vulgare*.

Wormwood, mountain. The *Artemisia glacialis* of Linnæus, which is common about the Alps, and similar in its virtues to the common wormwood.

Wormwood, Roman. See *Absinthium vulgare*.

Wormwood, sea. See *Absinthium maritimum*.

Wormwood, tartareum. See *Santonicum*.

Worm. An infusion of malt. This has been found useful in the cure of the scurvy.

Dr. Macbride, in his very ingenious experimental essays, having laid down as a principle, "that the cure of the scurvy depends on the fermentative quality in the remedies made use of," was led to inquire after a substance capable of being

preserved during a long sea voyage, and yet containing materials by which a fermentation might occasionally be excited in the bowels. Such an one appeared to him to be found in malt, which is well known to be the grain of barley, brought suddenly to a germinating state by heat and moisture, and then dried, whereby its saccharine principle is developed, and rendered easy of extraction by watery liquors. The sweet infusion of this he proposed to give as a dietetic article to scorbutic persons, excepting that it would ferment in their bowels, and give out its fixed air, by the antiseptic powers of which the strong tendency to putrefaction in this disease might be corrected.

It was some time before a fair trial of this proposed remedy could be obtained; and different reports were made concerning it. By some cases, however, published in a postscript of the second edition of the Doctor's work, in 1767, it appears that scorbutic complaints of the most dangerous kinds have actually been cured at sea by the use of wort. Its general effects were to keep the patients open, and to prove highly nutritious and strengthening. It sometimes purged too much, but this effect was easily obviated by the tincture thebaica. Other unquestionable cases of its success in this disease are to be seen in the London Medical Essays and Enquiries.

The use of wort has hence been adopted in other cases where a strong putrid disposition in the fluids appeared to prevail, as in cancerous and phagedenic ulcers; and instances are published in the fourth volume of the work above-mentioned of its remarkable good effects in these cases.

As the efficacy of the malt infusion depends upon its producing changes in the whole mass of fluids, it is obvious that it must be taken in large quantities for a considerable length of time, and rather as an article of diet than medicine. From one to four pints daily have generally been directed. The proportion recommended in preparing it, is one measure of ground malt to three equal measures of boiling water. The mixture must be well stirred, and left to stand, covered three or four hours. It should be made fresh every day.

Woundwort. See *Panax*.

Wrist, bones of. See *Carpal bones*.

X

XALAPPA. (From the province of Xalappa, in New Spain, whence it comes.) Jalap.

XANTHIUM. (From *ξανθος*, yellow; so named because it is said to make the hair yellow.) The lesser burdock. This herb *Xanthium strumarium* of Linnæus, was once esteemed in the cure of scrophula, but like most other remedies against this disease, proves ineffectual. The seeds are administered internally in some countries against erysipelas.

XANTHIUM STRUMARIUM. The systematic name of the lesser burdock. See *Xanthium*.

XERASIA. (From *ξηρος*, dry.) An excessive tenuity of the hairs similar to down.

XEROCOLLYRIUM. (From *ξηρος*, dry, and *χολυριον*, a collyrium.) A dry collyrium.

XEROMYRUM. (From *ξηρος*, dry, and *μυρον*, an ointment.) A dry ointment.

XEROPHTHALMIA. (*Ξηρος*, dry, and *οφθαλμια*, an inflammation of the eye.) A dry inflammation of the eye without discharge.

XIPHIUM. (From *ξίφος*, a sword; so named from the sword-like shape of its leaves.) Spurge wort.

XIPHOID. (*Xiphoides*, from *ξίφος*, a sword, and *ωδης*, likeness.) A term given by anatomists to parts which had some resemblance to an ancient sword, as the xiphoid cartilage.

Xiphoid cartilage. See *Cartilago eni-formis*.

XYLOALOE. See *Lignum aloes*.

XYLOBALSAMUM. See *Balsamum gileadense*.

Y

YAM. An esculent root, obtained principally from three species of *Dioscorea*, the *alata*, *bulbifera*, and *sativa*. They grow spontaneously in both Indies, and their roots are promiscuously eaten as the potatoe is with us. There is great variety in the colour, size, and shape of yams; some are generally blue or brown, round or oblong, and weigh from one pound to two. They are esteemed when dressed as being nutritious and easy of digestion, and are preferred to wheaten bread. Their taste is somewhat like the potatoe, but more luscious. The negroes, whose common food is yams, boil and mash them. They are also ground into flour, and made into bread and puddings.

When they are to be kept for some time, they are exposed upon the ground to the sun as we do onions, and when sufficiently withered they are put into dry sand in casks, and placed in a dry garret, where they remain often for many seasons without losing any of their primitive goodness.

Yarrow, common. See *Millefolium*.

Yawe. The African name for raspberry. See *Framboesia*.

Yellow fever. See *Febris continua*.

Yellow sunders. See *Santalum album*.

Yorkshire samite. See *Pinguicula*.

YPSILOGLOSSUS. (From *ψιλον*, the ypsiloid bone, and *γλωσσα*, the tongue.) A muscle originating in the ypsiloid bone, and terminating in the tongue.

YPSILOIDES. (From *ψ*, the Greek letter ypsilon, and *ωδης*, a likeness.) The os hyoides, so named from its likeness to the Greek letter *ψ* ypsilon.

YTTRIA. The heaviest of the earths. Its specific gravity is 4.842. It resembles glucine in several of its properties. It is smooth and insipid. It is infusible alone, but vitrifies with supersaturated borate of soda. It combines with the acids, and is precipitated from those solutions by ammonia and prussiate of potash. It is also precipitated by tannin. The precipitate is not soluble in acetous acid. As some of its salts are coloured, and its weight nearly approaches to that of metals, it is considered as the link which connects the metals with the earths. It differs from glucine in not being soluble in fixed alkalis, nor being precipitated by the succinates. Its attraction for the acids is also generally stronger than that of glucine, and its saline compounds have not the same saccharine taste. Its other properties have not yet been examined.

Z

SACCHARUM. See *Saccharum*.

SAFFRAN. (Arab.) Saffron.

SAIBAC. (Arab.) Quicksilver.

SARZA. An ancient and provincial name of the sarsaparilla.

SEA WAYS. The systematic name of the Indian wheat plant, a native of America, and cultivated in Italy and several parts of Europe, for its grain, which is ground for the same purposes as our wheat, to which it is very little inferior.

SEDOARIA. 1 The name of a genus of plants in the Linnean system. Class, *Monandria* Order, *Monogymia*. Zedoary.

2 The pharmacopœial name of the *Kempferia retunda* of Linneus, *foliis lanceolatis petiolatis*. The roots of this plant are brought to us in long pieces about the thickness of the little finger, two or three inches in length, bent, rough, and angular, or in roundish pieces about an inch in diameter, of an ash colour on the outside, and white within. They have an agreeable camphoraceous smell, and a bitterish aromatic taste. Though formerly much esteemed against rheumatic affections, they are at present thought to possess very little medicinal powers, although they have a place in the confectio aromatica of the London pharmacopœia.

SEDOARIA LONGA. The long roots of the *Zedoaria retunda* of Linneus.

SEDOARIA ROTUNDA. The systematic name of the zedoary plant. See *Zedoaria*.

Zedoary. See *Zedoaria*.

SERNA. An ulcerated impetigo. Lepra.

SIBETHUM. (From *Zobeth*, Arab.) Civetta. Civet. A soft unctuous odoriferous substance about the consistence of honey or butter, of a whiteish, yellowish, or brownish colour, sometimes blackish, contained in some excretory follicles near the anus of the *Fiberia zibetika* of Linneus. It has a grateful smell when diluted, and an unctuous subacid taste, and possesses stimulating, nervine, and antispasmodic virtues.

ZINC. (*Zincum*, Germ.) A metal found in nature combined with oxygen, carbonic acid, and sulphuric acid; and mineralised by sulphur. Native oxid of zinc is commonly called *calamine*. It occurs in a loose, and in a compact form, amorphous, of a white, grey, yellow, or brown colour, without lustre or transparency. Combined with carbonic acid, it is called *vitreous zinc ore*, or *native carbonate of zinc*. It is found in solid masses, sometimes in six-sided compressed prisms, both ends being covered with pentagons. Its colour

is generally greyish inclining to black. It is often transparent. *Sulphate of zinc* is found efflorescent in the form of stalactites or in rhombs. *Sulphuret of zinc*, or *blende*, is the most abundant ore. It is found of various colours; brown, yellow, hyacinth, black, &c. and with various degrees of lustre and transparency. This zinc ore is contaminated with iron, lead, argillaceous and siliceous earths, &c. It occurs both in amorphous masses and crystallized in a diversity of polygonal figures.

Properties of zinc.—Zinc on its fracture appears of a shining white colour, with a blueish tint. It possesses some degree of ductility, and may be extended when carefully passed between metallic cylinders in a flating mill. Its specific gravity is 7.190. It melts at 700° Fahr. By a further increase of heat it is volatilized without change. When melted in contact with the air, its surface becomes covered with an imperfect oxid; when heated a little above ignition it burns with a bright yellowish white flame, slightly tinged with green, and a white oxid is formed in light flakes, which are carried off by the rapid current of air in the burning metal. It undergoes very little alteration from the air, its surface becomes slightly tarnished. It is not acted on by water at the usual temperature of our atmosphere, but at that of ignition, it decomposes this fluid rapidly. It is oxidated and dissolved by the greater number of the acids. It has a very strong attraction for oxygen, and therefore precipitates the greater number of the metals from their acid solutions. All the alkalis when digested or boiled with zinc blacken its surface and dissolve a minute portion of it. It decomposes muriate of ammonia, sulphate of potash, and various other neutral salts. A mixture of nitrate of potash and zinc detonates with rapidity. Sulphur and zinc cannot be united by fusion. Gold, silver, platinum, and nickel form brittle compounds with zinc. It easily unites with mercury and tin. It does not combine with lead or bismuth. The most frequent and at the same time most useful combination of zinc is that with copper. It unites with great difficulty to arsenic, iron, and cobalt. It inflames in oxygenated muriatic acid gas, and fulminates by pressure or a blow with oxygenated muriate of potash. It is a very strong conductor of galvanism. The use of zinc in the arts is very considerable. In medicine the sulphate of zinc, or white vitriol, as it is called, is given internally

in the dose of from \mathfrak{Dj} to \mathfrak{Zss} , as a vomit. In small doses it cures dropsies, intermitting head-achs, and some nervous diseases, and is a powerful antispasmodic and tonic. A solution of white vitriol is also used to remove gleet, gonorrhoeas, and for cleaning foul ulcers.

Zinc, vitriolated. See *Zinci sulphus*.

ZINCUM. See *Zinc*.

ZINCUM CALCINATUM. See *Oxydum zinci*.

ZINCUM VITRIOLATUM. See *Zinci sulphas*.

ZINCUM VITRIOLATUM PURIFICATUM. See *Zinci sulphas*.

ZINGI. An ancient name of the stellated annised. See *Anisum stellatum*.

ZINGIBER. (*Zingiberaceae*, Indian.) *Zingiber album*. *Zingiber nigrum*. *Zingiber commune*. Ginger. *Amomum zingiber* of Linnaeus. The white and black ginger are both the produce of the same plant, the difference depending upon the mode of preparing them. Ginger is generally considered as an aromatic, and less pungent and heating to the system than might be expected from its effects upon the organ of taste. It is used as an antispasmodic and carminative. The cases in which it is more immediately serviceable are flatulent colics, debility, and laxity of the stomach and intestines; and in torpid and phlegmatic constitutions to excite brisker vascular action. It is seldom given but in combination with other medicines. In the pharmacopoeias it is directed in the form of a syrup and condiment, and in many compositions ordered as a subsidiary ingredient.

ZINGIBER ALBUM. The root of the *amomum zingiber* of Linnaeus is so termed when deprived of its radicles and sordes.

ZINGIBER COMMUNE. See *Zingiber*.

ZINGIBER NIGRUM. The root of the *Amomum zingiber* of Linnaeus is so called when suffered to dry with its radicles and the sordes which usually hang to it.

ZIRCON. An earth discovered in the year 1793, by Klaproth of Berlin, in the Zircon or Jargon, a gem first brought from the island of Ceylon, but also found in France, Spain, and other parts of Europe. Its colour is either grey, greenish, yellowish, reddish-brown, or purple. It has little lustre, and is nearly opaque. It is likewise found in another gem called the yacinth. This stone is of a yellowish red colour mixed with brown. It possesses lustre and transparency.

Properties.—It has a white colour, is exceedingly heavy, and rough or harsh to the touch like silice. It has neither taste nor odour, and is insoluble in water, but forms with it a kind of jelly. It melts with borax into a transparent colourless glass. It suffers in a charcoal crucible a pasty fusion by intense heat, and contracts in its dimensions, acquiring a

grey colour and scintillating hardness. In this state it is very hard and insoluble in acids. Its specific gravity is 4.3. Neither air nor combustible bodies act upon it. It is soluble in water, but retains while drying in the air a large quantity of it, which gives it the semi-transparency and appearance of a yellow jelly, or gum-arabic; it exhibits the same vitreous fracture. It unites with all the acids and forms salts, differing from those of the other bases by being decomposable by alumine, glucine, the alkalis, and by mere heat. It fuses with alumine and silice. It is insoluble even by boiling in a solution of alkalis, neither can it be fused with them by means of heat; but it is soluble in alkaline carbonates.

By these properties this earth may be distinguished from every other. It is still of no use in the arts.

Method of obtaining zircon.—Reduce the mineral to powder, mix it with three times its weight of potash, and fuse it in a crucible. Wash the obtained mass in distilled water, till the whole of the potash be extracted; then dissolve the residuum as far as possible in diluted muriatic acid. Boil the solution to precipitate any silice it may contain, filter it, and gradually add solution of potash. The zircon will now become precipitated. Wash it repeatedly in distilled water and dry it.

ZIZANIA AQUATICA. The systematic name of a reed whose grain is much esteemed. See *Water zizania*.

ZIZYPHUS. The jujubes were formerly so called. See *Jujuba*.

ZONA. (From *ζώνω*, to surround.) *Zoster*. The shingles.

ZOOLOGY. (*Zoologia*, from *ζωον*, an animal, and *λογος*, a discourse.) That part of natural history which treats of animals.

ZOONOMIA. (From *ζωον*, an animal, and *νομος*, a law.) The laws of organic life.

ZOOTOMY. (*Zootomia*. From *ζωον*, an animal, and *τομή*, to cut.) The dissection of animals.

ZOSTER. (From *ζώνω*, to gird.) A kind of erysipelas which goes round the body like a girdle.

ZUCHAR. (Arab.) Sugar.

ZYGOMA. (From *ζυγος*, a yoke; because it transmits the tendon of the temporal muscle like a yoke.) The cavity under the zygomatic process of the temporal bone, and os malle.

ZYGOMATIC PROCESS. An apophysis of the os jugale and another of the temporal bone are so called.

ZYGOMATIC SUTURE. *Sutura zygomatica*. The union of the zygomatic process of the temporal bone to the cheek bone.

ZYGOMATICUS MAJOR. This muscle arises from the cheek bone near the

zygomatic suture, taking a direction downwards and inwards to the angle of the mouth; it is a long slender muscle, which ends by mixing its fibres with the orbicularis oris and the depressor of the lip.

ZYGOMATICUS MINOR. This muscle arises a little higher up than the zygomaticus major, upon the cheek bone, but nearer the nose; it is much more slender than that muscle, and is often wanting. It is the zygomatic muscle that marks the face with that line which extends from the cheek bone to the corner of the mouth, which is particularly distinguishable in some persons. The zygomatic muscles pull the

angles of the mouth up as in laughter, and from, in this way, rendering the face distorted, it has obtained the name of distortor oris. The strong action of this muscle is more particularly seen in laughter, rage, or grinning.

ZITHOGALA. *Zithogala.* Beer and milk, which make together what we commonly call *posset-drink*; a term often to be met with in Sydenham.

ZZ. The ancients signify *Myrris* by these two letters, from *Myrris*, a name for it common amongst them; but the late writers use them only for the *Zinziber*, ginger.

Thursday Morning - Commenced
Continued till Wednesday
Morning - has subsided
building continues -
a large pile 21-22 to 505

THE END.

RETURN BIOLOGY LIBRARY
TO → 3503 Life Sciences Bldg. 642-2531

LOAN PERIOD 1 2 3

4 **1-MONTH--MONOGRAPH** 5 6

ALL BOOKS MAY BE RECALLED AFTER 7 DAYS
 Renewed books are subject to immediate recall

DUE AS STAMPED BELOW

MAY 1 1 1985		
DUE		
MAY 0 1 1985		
Subject to Recall Immediately		
DUE		
FEB 15 1991		
SUBJECT TO RECALL IMMEDIATELY		

UNIVERSITY OF CALIFORNIA, BERKELEY
 FORM NO. DD4, 12m, 12/80 BERKELEY, CA 94720

©s

Handwritten scribbles or marks at the top of the page.

Handwritten text, possibly a signature or a list of items, running diagonally across the upper left portion of the page.



